

Table of Radionuclides (Comments on evaluation)

M.-M. BÉ, V. CHISTÉ, C. DULIEU, M.A. KELLETT, X. MOUGEOT
E. BROWNE, C. BAGLIN
V.P. CHECHEV, A. EGOROV, N.K. KUZMENKO, V.O. SERGEEV
T. KIBÉDI
F.G. KONDEV
A. LUCA
M. GALÁN
X. HUANG, B. WANG
R.G. HELMER
E. SCHÖNFELD, R. DERSCH
V.R. VANIN, R.M. de CASTRO
A.L. NICHOLS
T.D. MACMAHON, A. PEARCE, A. ARINC
K.B. LEE
S.C. WU



Volumes 1 to 8, 2016

BUREAU INTERNATIONAL DES POIDS ET MESURES

Pavillon de Breteuil, F-92310 SÈVRES

Édité par le BIPM,
Pavillon de Breteuil
F-92312 Sèvres Cedex
France

ISBN-13 978-92-822-2264-5 (Vol. 8)

**TABLE DE RADIONUCLÉIDES
TABLE OF RADIONUCLIDES**

COMMENTS ON EVALUATIONS

Marie-Martine BÉ, Vanessa CHISTÉ, Christophe DULIEU, Mark KELLETT, Xavier MOUGEOT, Laboratoire National Henri Becquerel (LNHB), France;
Arzu ARINC, National Physical Laboratory (NPL), United Kingdom;
Valery CHECHEV, Nikolay KUZMENKO, Khlopin Radium Institute (KRI), Russia;
Tibor KIBÉDI, Australian National University (ANU), Australia;
Aurelian LUCA, National Institute for Physics and Nuclear Engineering "Horia Hulubei" (IFIN-HH), Romania;
Alan L. NICHOLS, Department of Physics, University of Surrey, United Kingdom.

Monographie BIPM-5 - Table of Radionuclides, Comments on evaluations, Volume 8

Marie-Martine BÉ, Vanessa CHISTÉ, Christophe DULIEU, Mark KELLETT, Xavier MOUGEOT, Laboratoire National Henri Becquerel (LNHB), France;
Arzu ARINC, National Physical Laboratory (NPL), United Kingdom;
Valery CHECHEV, Nikolay KUZMENKO, Khlopin Radium Institute (KRI), Russia;
Tibor KIBÉDI, Australian National University (ANU), Australia;
Aurelian LUCA, National Institute for Physics and Nuclear Engineering "Horia Hulubei" (IFIN-HH), Romania;
Alan L. NICHOLS, Department of Physics, University of Surrey, United Kingdom.

Sommaire

Depuis quelques années, un groupe composé d'évaluateurs spécialistes des données de décroissance radioactive s'est formé, avec l'objectif de réaliser une évaluation soignée et documentée de ces données pour des radionucléides intervenant dans de nombreuses applications. Ces évaluateurs se sont mis d'accord sur une méthodologie commune. Ce rapport inclut les commentaires sur les évaluations des radionucléides figurant dans le rapport Monographie BIPM-5, volume 8 :

^{41}Ca , ^{47}Sc , ^{52}Fe , ^{58}Co , ^{61}Cu , ^{63}Zn , ^{73}Se , ^{82}Rb , ^{82}Sr , ^{88}Y , ^{89}Zr , ^{93}Zr , $^{93\text{m}}\text{Nb}$, $^{94\text{m}}\text{Tc}$, ^{106}Ru , ^{106}Rh , ^{109}Cd , ^{127}Xe , ^{131}I , $^{131\text{m}}\text{Xe}$, ^{133}Ba , ^{138}La , ^{140}Ba , ^{140}La , ^{144}Ce , ^{144}Pr , $^{144\text{m}}\text{Pr}$, ^{148}Pm , $^{148\text{m}}\text{Pm}$, ^{151}Sm , ^{169}Er , ^{198}Au .

Summary

Over the past years, an informal collaboration of decay-data evaluators has been assembled with the goal of creating high-quality, well-documented evaluations of the decay data for a selected set of radionuclides that are of interest in various applications. This report includes, for each radionuclide, the evaluator's comments on how the evaluation was carried out for the radionuclides that are in the Monographie BIPM-5, volume 8:

^{41}Ca , ^{47}Sc , ^{52}Fe , ^{58}Co , ^{61}Cu , ^{63}Zn , ^{73}Se , ^{82}Rb , ^{82}Sr , ^{88}Y , ^{89}Zr , ^{93}Zr , $^{93\text{m}}\text{Nb}$, $^{94\text{m}}\text{Tc}$, ^{106}Ru , ^{106}Rh , ^{109}Cd , ^{127}Xe , ^{131}I , $^{131\text{m}}\text{Xe}$, ^{133}Ba , ^{138}La , ^{140}Ba , ^{140}La , ^{144}Ce , ^{144}Pr , $^{144\text{m}}\text{Pr}$, ^{148}Pm , $^{148\text{m}}\text{Pm}$, ^{151}Sm , ^{169}Er , ^{198}Au .

Monographie BIPM-5 - Table of Radionuclides, Comments on evaluations, volumes 1 to 7

^3H , ^7Be , ^{11}C , ^{13}N , ^{14}C , ^{15}O , ^{18}F , ^{22}Na , ^{24}Na , ^{32}P , ^{33}P , ^{35}S , ^{36}Cl , ^{37}Ar , ^{40}K , ^{41}Ar , ^{44}Sc , ^{44}Ti , ^{45}Ca , ^{46}Sc , ^{51}Cr , ^{54}Mn , ^{55}Fe , ^{56}Mn , ^{56}Co , ^{57}Co , ^{57}Ni , ^{59}Fe , ^{59}Ni , ^{60}Co , ^{63}Ni , ^{64}Cu , ^{65}Zn , ^{66}Ga , ^{67}Ga , ^{68}Ga , ^{68}Ge , ^{75}Se , ^{79}Se , ^{85}Kr , ^{85}Sr , ^{88}Y , ^{89}Sr , ^{90}Sr , ^{90}Y , $^{90\text{m}}\text{Y}$, $^{93\text{m}}\text{Nb}$, ^{99}Mo , ^{99}Tc , $^{99\text{m}}\text{Tc}$, ^{108}Ag , $^{108\text{m}}\text{Ag}$, ^{109}Pd , ^{109}Cd , ^{110}Ag , $^{110\text{m}}\text{Ag}$, ^{111}In , $^{123\text{m}}\text{Te}$, ^{123}I , ^{124}Sb , ^{125}Sb , ^{125}I , ^{127}Sb , ^{127}Te , $^{127\text{m}}\text{Te}$, ^{129}I , ^{131}I , $^{131\text{m}}\text{Xe}$, ^{132}Te , ^{133}I , ^{133}Xe , $^{133\text{m}}\text{Xe}$, ^{133}Ba , ^{134}Cs , $^{135\text{m}}\text{Xe}$, ^{137}Cs , ^{139}Ce , ^{140}Ba , ^{140}La , ^{141}Ce , ^{147}Nd , ^{147}Pm , ^{152}Eu , ^{153}Sm , ^{153}Gd , ^{154}Eu , ^{155}Eu , ^{159}Gd , ^{166}Ho , $^{166\text{m}}\text{Ho}$, ^{169}Yb , ^{170}Tm , ^{177}Lu , ^{182}Ta , ^{186}Re , ^{195}Au , ^{198}Au , ^{201}Tl , ^{203}Hg , ^{203}Pb , ^{204}Tl , ^{206}Hg , ^{206}Tl , ^{207}Tl , ^{207}Bi , ^{208}Tl , ^{209}Tl , ^{209}Pb , ^{209}Po , ^{210}Tl , ^{210}Pb , ^{210}Bi , ^{210}Po , ^{211}Pb , ^{211}Bi , ^{211}Po , ^{211}At , ^{212}Pb , ^{212}Bi , ^{212}Po , ^{213}Bi , ^{213}Po , ^{214}Pb , ^{214}Bi , ^{214}Po , ^{215}Bi , ^{215}Po , ^{215}At , ^{216}Po , ^{217}At , ^{217}Rn , ^{218}Po , ^{218}At , ^{218}Rn , ^{219}At , ^{219}Rn , ^{220}Rn , ^{221}Fr , ^{222}Rn , ^{223}Fr , ^{223}Ra , ^{224}Ra , ^{225}Ra , ^{225}Ac , ^{226}Ra , ^{227}Ac , ^{227}Th , ^{228}Ra , ^{228}Ac , ^{228}Th , ^{231}Th , ^{231}Pa , ^{232}Th , ^{232}U , ^{233}Th , ^{233}Pa , ^{234}Th , ^{234}Pa , $^{234\text{m}}\text{Pa}$, ^{234}U , ^{235}U , ^{236}U , ^{236}Np , $^{236\text{m}}\text{Np}$, ^{237}U , ^{237}Np , ^{238}U , ^{238}Np , ^{238}Pu , ^{239}U , ^{239}Np , ^{239}Pu , ^{240}Pu , ^{241}Pu , ^{241}Am , ^{242}Pu , ^{242}Am , $^{242\text{m}}\text{Am}$, ^{242}Cm , ^{243}Am , ^{243}Cm , ^{244}Am , $^{244\text{m}}\text{Am}$, ^{244}Cm , ^{245}Cm , ^{246}Cm , ^{252}Cf .

TABLE DE RADIONUCLÉIDES – COMMENTAIRES SUR LES ÉVALUATIONS

De nombreuses applications nécessitent la connaissance des données liées à la désintégration des radionucléides, telles que la période radioactive, les énergies et les intensités des divers rayonnements. Pour répondre aux demandes des utilisateurs, le Laboratoire National Henri Becquerel (LNE - LNHB, France) a, de 1982 à 1987, publié une table en quatre volumes [87Ta, 99Be]. Puis, en 1993 une coopération a été établie avec le Physikalisch-Technische Bundesanstalt (PTB, Allemagne) afin de reprendre cette étude et de la développer. En 1995, un nouveau groupe de travail international nommé Decay Data Evaluation Project (DDEP) s'est formé qui, en plus des deux laboratoires nationaux précédents, inclut : Idaho National Engineering and Environmental Laboratory (INEEL, USA), Lawrence Berkeley National Laboratory (LBNL, USA), Brookhaven National Laboratory (BNL, USA) et Khlopin Radium Institute (KRI, Russie).

Le propos de ce groupe est de fournir aux utilisateurs des données soigneusement évaluées qui puissent servir de référence. A cette fin, tous les membres du groupe suivent une méthodologie commune qui comprend :

- la lecture attentive de toutes les publications relatives à une grandeur ;
- une analyse statistique des données retenues ;
- le choix et l'utilisation des mêmes jeux de données, pour celles ayant fait l'objet d'études spécifiques par des spécialistes, tels les coefficients de conversion interne.

Tous ces points sont développés en détail dans le chapitre « Rules for evaluation and compilations ».

Par ailleurs, toutes les évaluations sont documentées et l'établissement des valeurs retenues explicité. Ce document est ensuite relu par deux membres du groupe.

Ce 8^{ème} volume regroupe les commentaires liés à l'évaluation des radionucléides suivants :

⁴¹Ca, ⁴⁷Sc, ⁵²Fe, ⁵⁸Co, ⁶¹Cu, ⁶³Zn, ⁷³Se, ⁸²Rb, ⁸²Sr, ⁸⁸Y, ⁸⁹Zr, ⁹³Zr, ^{93m}Nb, ^{94m}Tc, ¹⁰⁶Ru, ¹⁰⁶Rh, ¹⁰⁹Cd, ¹²⁷Xe, ¹³¹I, ^{131m}Xe, ¹³³Ba, ¹³⁸La, ¹⁴⁰Ba, ¹⁴⁰La, ¹⁴⁴Ce, ¹⁴⁴Pr, ^{144m}Pr, ¹⁴⁸Pm, ^{148m}Pm, ¹⁵¹Sm, ¹⁶⁹Er, ¹⁹⁸Au.

Ainsi que ceux précédemment publiés dans les volumes 1 à 7 :

³H, ⁷Be, ¹¹C, ¹³N, ¹⁴C, ¹⁵O, ¹⁸F, ²²Na, ²⁴Na, ³²P, ³³P, ³⁵S, ³⁶Cl, ³⁷Ar, ⁴⁰K, ⁴¹Ar, ⁴⁴Sc, ⁴⁴Ti, ⁴⁵Ca, ⁴⁶Sc, ⁵¹Cr, ⁵⁴Mn, ⁵⁵Fe, ⁵⁶Mn, ⁵⁶Co, ⁵⁷Co, ⁵⁷Ni, ⁵⁹Fe, ⁵⁹Ni, ⁶⁰Co, ⁶³Ni, ⁶⁴Cu, ⁶⁵Zn, ⁶⁶Ga, ⁶⁷Ga, ⁶⁸Ga, ⁶⁸Ge, ⁷⁵Se, ⁷⁹Se, ⁸⁵Kr, ⁸⁵Sr, ⁸⁸Y, ⁸⁹Sr, ⁹⁰Sr, ⁹⁰Y, ^{90m}Y, ^{93m}Nb, ⁹⁹Mo, ⁹⁹Tc, ^{99m}Tc, ¹⁰⁸Ag, ^{108m}Ag, ¹⁰⁹Pd, ¹⁰⁹Cd, ¹¹⁰Ag, ^{110m}Ag, ¹¹¹In, ^{123m}Te, ¹²³I, ¹²⁴Sb, ¹²⁵Sb, ¹²⁵I, ¹²⁷Sb, ¹²⁷Te, ^{127m}Te, ¹²⁹I, ¹³¹I, ^{131m}Xe, ¹³²Te, ¹³³I, ¹³³Xe, ^{133m}Xe, ¹³³Ba, ¹³⁴Cs, ^{135m}Xe, ¹³⁷Cs, ¹³⁹Ce, ¹⁴⁰Ba, ¹⁴⁰La, ¹⁴¹Ce, ¹⁴⁷Nd, ¹⁴⁷Pm, ¹⁵²Eu, ¹⁵³Sm, ¹⁵³Gd, ¹⁵⁴Eu, ¹⁵⁵Eu, ¹⁵⁹Gd, ¹⁶⁶Ho, ^{166m}Ho, ¹⁶⁹Yb, ¹⁷⁰Tm, ¹⁷⁷Lu, ¹⁸²Ta, ¹⁸⁶Re, ¹⁹⁵Au, ¹⁹⁸Au, ²⁰¹Tl, ²⁰³Hg, ²⁰³Pb, ²⁰⁴Tl, ²⁰⁶Hg, ²⁰⁶Tl, ²⁰⁷Tl, ²⁰⁷Bi, ²⁰⁸Tl, ²⁰⁹Tl, ²⁰⁹Pb, ²⁰⁹Po, ²¹⁰Tl, ²¹⁰Pb, ²¹⁰Bi, ²¹⁰Po, ²¹¹Pb, ²¹¹Bi, ²¹¹Po, ²¹¹At, ²¹²Pb, ²¹²Bi, ²¹²Po, ²¹³Bi, ²¹³Po, ²¹⁴Pb, ²¹⁴Bi, ²¹⁴Po, ²¹⁵Bi, ²¹⁵Po, ²¹⁵At, ²¹⁶Po, ²¹⁷At, ²¹⁷Rn, ²¹⁸Po, ²¹⁸At, ²¹⁸Rn, ²¹⁹At, ²¹⁹Rn, ²²⁰Rn, ²²¹Fr, ²²²Rn, ²²³Fr, ²²³Ra, ²²⁴Ra, ²²⁵Ra, ²²⁵Ac, ²²⁶Ra, ²²⁷Ac, ²²⁷Th, ²²⁸Ra, ²²⁸Ac, ²²⁸Th, ²³¹Th, ²³¹Pa, ²³²Th, ²³²U, ²³³Th, ²³³Pa, ²³⁴Th, ²³⁴Pa, ^{234m}Pa, ²³⁴U, ²³⁵U, ²³⁶U, ²³⁶Np, ^{236m}Np, ²³⁷U, ²³⁷Np, ²³⁸U, ²³⁸Np, ²³⁸Pu, ²³⁹U, ²³⁹Np, ²³⁹Pu, ²⁴⁰Pu, ²⁴¹Pu, ²⁴¹Am, ²⁴²Pu, ²⁴²Am, ^{242m}Am, ²⁴²Cm, ²⁴³Am, ²⁴³Cm, ²⁴⁴Am, ^{244m}Am, ²⁴⁴Cm, ²⁴⁵Cm, ²⁴⁶Cm, ²⁵²Cf.

Les données de décroissance radioactive de ces radionucléides peuvent être trouvées dans la Monographie BIPM-5 de la « Table de radionucléides », dans le CD-Rom NUCLÉIDE édité par le LNHB ou sur les pages web : <http://www.nucleide.org/NucData.htm>

TABLE OF RADIONUCLIDES – COMMENTS ON EVALUATIONS

Basic properties of radionuclides, such as half-life, decay mode and branchings, radiation energies and emission probabilities are commonly used in various research fields. To meet the demand for these data the LNHB produced a table that was published in four volumes [87Ta, 99Be] from 1982 to 1987. In 1993, a cooperative agreement was established between the Laboratoire National Henri Becquerel (LNE - LNHB, France) and the Physikalisch-Technische Bundesanstalt (PTB, Germany) to continue and expand this work. In 1995, a new international collaboration was formed, the Decay Data Evaluation Project (DDEP), which has the same objectives. Along with the evaluators from LNHB and PTB, this collaboration includes others from the Idaho National Engineering and Environmental Laboratory (INEEL, USA), the Lawrence Berkeley National Laboratory (LBNL, USA), the Brookhaven National Laboratory (BNL, USA) and the Khlopin Radium Institute (KRI, Russia). Its objective has been to provide carefully produced recommended values, which may eventually become standard data. With this goal in mind, the collaboration has adopted a uniform evaluation methodology that contains the following:

- a critical review of relevant publications;
- an accounting of all measured data;
- a uniform statistical analysis of the data;
- a presentation of values for quantities such as internal conversion coefficients, etc.;
- a review of evaluation by two other members of the collaboration.

These topics are described in detail in the chapter “Rules for evaluation and compilations”.

The evaluation of each individual radionuclide has a section (presented here) that describes the procedures used for deducing the recommended values. This documentation is included in order to establish the quality and completeness of each evaluation. It can also provide the basis for any future reevaluation by the DDEP or other groups.

This eighth volume contains the procedures and comments relevant to the evaluation for the following radionuclides:

^{41}Ca , ^{47}Sc , ^{52}Fe , ^{58}Co , ^{61}Cu , ^{63}Zn , ^{73}Se , ^{82}Rb , ^{82}Sr , ^{88}Y , ^{89}Zr , ^{93}Zr , $^{93\text{m}}\text{Nb}$, $^{94\text{m}}\text{Tc}$, ^{106}Ru , ^{106}Rh , ^{109}Cd , ^{127}Xe , ^{131}I , $^{131\text{m}}\text{Xe}$, ^{133}Ba , ^{138}La , ^{140}Ba , ^{140}La , ^{144}Ce , ^{144}Pr , $^{144\text{m}}\text{Pr}$, ^{148}Pm , $^{148\text{m}}\text{Pm}$, ^{151}Sm , ^{169}Er , ^{198}Au .

As well as those previously published in volumes 1 to 7:

^3H , ^7Be , ^{11}C , ^{13}N , ^{14}C , ^{15}O , ^{18}F , ^{22}Na , ^{24}Na , ^{32}P , ^{33}P , ^{35}S , ^{36}Cl , ^{37}Ar , ^{40}K , ^{41}Ar , ^{44}Sc , ^{44}Ti , ^{45}Ca , ^{46}Sc , ^{51}Cr , ^{54}Mn , ^{55}Fe , ^{56}Mn , ^{56}Co , ^{57}Co , ^{57}Ni , ^{59}Fe , ^{59}Ni , ^{60}Co , ^{63}Ni , ^{64}Cu , ^{65}Zn , ^{66}Ga , ^{67}Ga , ^{68}Ga , ^{68}Ge , ^{75}Se , ^{79}Se , ^{85}Kr , ^{85}Sr , ^{88}Y , ^{89}Sr , ^{90}Sr , ^{90}Y , $^{90\text{m}}\text{Y}$, $^{93\text{m}}\text{Nb}$, ^{99}Mo , ^{99}Tc , $^{99\text{m}}\text{Tc}$, ^{108}Ag , $^{108\text{m}}\text{Ag}$, ^{109}Pd , ^{109}Cd , ^{110}Ag , $^{110\text{m}}\text{Ag}$, ^{111}In , $^{123\text{m}}\text{Te}$, ^{123}I , ^{124}Sb , ^{125}Sb , ^{125}I , ^{127}Sb , ^{127}Te , $^{127\text{m}}\text{Te}$, ^{129}I , ^{131}I , $^{131\text{m}}\text{Xe}$, ^{132}Te , ^{133}I , ^{133}Xe , $^{133\text{m}}\text{Xe}$, ^{133}Ba , ^{134}Cs , $^{135\text{m}}\text{Xe}$, ^{137}Cs , ^{139}Ce , ^{140}Ba , ^{140}La , ^{141}Ce , ^{147}Nd , ^{147}Pm , ^{152}Eu , ^{153}Sm , ^{153}Gd , ^{154}Eu , ^{155}Eu , ^{159}Gd , ^{166}Ho , $^{166\text{m}}\text{Ho}$, ^{169}Yb , ^{170}Tm , ^{177}Lu , ^{182}Ta , ^{186}Re , ^{195}Au , ^{198}Au , ^{201}Tl , ^{203}Hg , ^{203}Pb , ^{204}Tl , ^{206}Hg , ^{206}Tl , ^{207}Tl , ^{207}Bi , ^{208}Tl , ^{209}Tl , ^{209}Pb , ^{209}Po , ^{210}Tl , ^{210}Pb , ^{210}Bi , ^{210}Po , ^{211}Pb , ^{211}Bi , ^{211}Po , ^{211}At , ^{212}Pb , ^{212}Bi , ^{212}Po , ^{213}Bi , ^{213}Po , ^{214}Pb , ^{214}Bi , ^{214}Po , ^{215}Bi , ^{215}Po , ^{215}At , ^{216}Po , ^{217}At , ^{217}Rn , ^{218}Po , ^{218}At , ^{218}Rn , ^{219}At , ^{219}Rn , ^{220}Rn , ^{221}Fr , ^{222}Rn , ^{223}Fr , ^{223}Ra , ^{224}Ra , ^{225}Ra , ^{225}Ac , ^{226}Ra , ^{227}Ac , ^{227}Th , ^{228}Ra , ^{228}Ac , ^{228}Th , ^{231}Th , ^{231}Pa , ^{232}Th , ^{232}U , ^{233}Th , ^{233}Pa , ^{234}Th , ^{234}Pa , $^{234\text{m}}\text{Pa}$, ^{234}U , ^{235}U , ^{236}U , $^{236\text{m}}\text{Np}$, ^{237}U , ^{237}Np , ^{238}U , ^{238}Np , ^{238}Pu , ^{239}U , ^{239}Np , ^{239}Pu , ^{240}Pu , ^{241}Pu , ^{241}Am , ^{242}Pu , ^{242}Am , $^{242\text{m}}\text{Am}$, ^{242}Cm , ^{243}Am , ^{243}Cm , ^{244}Am , $^{244\text{m}}\text{Am}$, ^{244}Cm , ^{245}Cm , ^{246}Cm , ^{252}Cf .

These evaluations may be found in the BIPM-5 Monographie, on the CD-Rom NUCLÉIDE published by the LNHB or in the web pages: <http://www.nucleide.org/NucData.htm>

A goal of the DDEP is to avoid future duplication of effort by disseminating these critically evaluated data with the hope that they will be included in many other collections of decay data.

REFERENCES

[87Ta] **Table de Radionucléides**, F. Lagoutine, N. Coursol, J. Legrand. ISBN 2-7272-0078-1 (LMRI, 1982-1987).

[85Zi] **W.L. Zijp**, Netherland Energy Research Foundation, ECN, Petten, The Netherlands, Rep. ECN-179.

[96He] **R.G. Helmer**, Proceedings of the Int. Symp. "Advances in alpha-, beta- and gamma-ray Spectrometry", St. Petersburg, September 1996, p. 71.

[96Be] **M.-M. Bé, B. Duchemin and J. Lamé**. Nucl. Instrum. Methods A369 (1996) 523 and Bulletin du Bureau National de Métrologie 110 (1998).

[99In] **Table de Radionucléides. Introduction, nouvelle version**. Introduction, revised version. Einleitung, überarbeitete Fassung. ISBN 2-7272-0201-6, BNM-CEA/LNHB BP 52, 91191 Gif-sur-Yvette Cedex, France.

[99Be] **M.-M. Bé, E. Browne, V. Chechev, R.G. Helmer, E. Schönfeld**. Table de Radionucléides, ISBN 2-7272-0200-8 and ISBN 2-7272-0211-3 (LNHB, 1988-1999).

[04Be] **M.-M. Bé, E. Browne, V. Chechev, V. Chisté, R. Dersch, C. Dulieu, R.G. Helmer, T.D. MacMahon, A.L. Nichols, E. Schönfeld**. Table of Radionuclides, Monographie BIPM-5, ISBN 92-822-2207-7 (set) and ISBN 92-822-2205-5 (CD), CEA/BNM-LNHB, 91191 Gif-sur-Yvette, France and BIPM, Pavillon de Breteuil, 92312 Sèvres, France.

and

M.-M. Bé, E. Browne, V. Chechev, V. Chisté, R. Dersch, C. Dulieu, R.G. Helmer, N. Kuzmenco, A.L. Nichols, E. Schönfeld. NUCLÉIDE, Table de Radionucléides sur CD-Rom, Version 2-2004, CEA/BNM-LNHB, 91191 Gif-sur-Yvette, France.

[06Be] **Marie-Martine BÉ, Vanessa CHISTÉ, Christophe DULIEU; Edgardo BROWNE, Coral BAGLIN; Valery CHECHEV, Nikolay KUZMENKO; Richard G. HELMER; Filip G. KONDEV; T. Desmond MACMAHON; Kyung Beom LEE**. *Table of Radionuclides, Monographie BIPM-5, vol. 3*, ISSN 92-822-2204-7 (set), ISBN 92-822-2218-7 (Vol. 3) and ISBN 92-822-2219-5 (CD), CEA/LNE-LNHB, 91191 Gif-sur-Yvette, France and BIPM, Pavillon de Breteuil, 92312 Sèvres, France.

[08Be] **Marie-Martine BÉ, Vanessa CHISTÉ, Christophe DULIEU; Edgardo BROWNE; Valery CHECHEV, Nikolay KUZMENKO; Filip G. KONDEV; Aurelian LUCA; Mónica GALÁN; Andrew PEARCE; Xiaolong HUANG**. *Table of Radionuclides, Monographie BIPM-5, vol.4*, ISBN 92-822-2230-6 (Vol. 4) and ISBN 92-822-2231-4 (CD), CEA/LNE-LNHB, 91191 Gif-sur-Yvette, France and BIPM, Pavillon de Breteuil, 92312 Sèvres, France.

[10Be] **Marie-Martine BÉ, Vanessa CHISTÉ, Christophe DULIEU, Xavier MOUGEOT, Edgardo BROWNE, Valery CHECHEV, Nikolay KUZMENKO, Filip G. KONDEV, Aurelian LUCA, Mónica GALAN, Arzu ARINC, Xiaolong HUANG, Alan NICHOLS**. Table of Radionuclides, Monographie BIPM-5, vol.5, ISBN 13 978-92-822-2234-8 (Vol. 5) et 13 978-92-822-2235-5 (CD-Rom), CEA/LNE-LNHB, 91191 Gif-sur-Yvette, France and BIPM, Pavillon de Breteuil, 92312 Sèvres, France.

Table of Radionuclides, Monographie BIPM-5, Commentaires, vol.5, ISBN 13 978-92-822-2235-5 (CD-Rom), CEA/LNE-LNHB, 91191 Gif-sur-Yvette, France and BIPM, Pavillon de Breteuil, 92312 Sèvres, France.

[11Be] **Marie-Martine BÉ, Vanessa CHISTÉ, Christophe DULIEU, Xavier MOUGEOT, Valery CHECHEV, Nikolay KUZMENKO, Filip G. KONDEV, Aurelian LUCA, Mónica GALÁN, Arzu ARINC, Xiaolong HUANG, B. WANG, Alan NICHOLS.** Table of Radionuclides, Monographie BIPM-5, vol.6, ISBN 13 978-92-822-2242-3 (Vol. 6) et 13 978-92-822-2243-0 (CD-Rom), CEA/LNE-LNHB, 91191 Gif-sur-Yvette, France and BIPM, Pavillon de Breteuil, 92312 Sèvres, France.

Table of Radionuclides, Monographie BIPM-5, Commentaires, vol.6, ISBN 13 978-92-822-2243-0 (CD-Rom), CEA/LNE-LNHB, 91191 Gif-sur-Yvette, France and BIPM, Pavillon de Breteuil, 92312 Sèvres, France.

[13Be] **Marie-Martine BÉ, Vanessa CHISTÉ, Christophe DULIEU, Xavier MOUGEOT, Valery CHECHEV, Filip G. KONDEV, Alan L. NICHOLS, Xiaolong HUANG, Baosong WANG.** Table of Radionuclides, Monographie BIPM-5, vol.7, ISBN 13 978-92-822-2248-5 (Vol. 7) et 13 978-92-822-2249-2 (CD-Rom), CEA/LNE-LNHB, 91191 Gif-sur-Yvette, France and BIPM, Pavillon de Breteuil, 92312 Sèvres, France.

Table of Radionuclides, Monographie BIPM-5, Commentaires, vol.7, ISBN 13 978-92-822-2249-2 (CD-Rom), CEA/LNE-LNHB, 91191 Gif-sur-Yvette, France and BIPM, Pavillon de Breteuil, 92312 Sèvres, France.

**AUTEURS POUR CORRESPONDANCE
AUTHOR'S MAIL ADDRESSES
ADRESSEN DER AUTOREN
AUTORES PARA CORRESPONDENCIA**

Toutes demandes de renseignements concernant les données recommandées et la façon dont elles ont été établies doivent être adressées directement aux auteurs des évaluations.

Information on the data and the evaluation procedures is available from the authors listed below.

Informationen über die Daten und Evaluationsprozeduren können bei den im folgenden zusammengestellten Autoren angefordert werden.

Todos los pedidos de información sobre datos recomendados y los métodos de evaluación utilizados, deben dirigirse directamente a los autores de las evaluaciones.

Dr. Arzu Arinc

National Physical Laboratory
Teddington,
Middlesex, TW11 OLM, United Kingdom
E-mail: Arzu.Arinc@npl.co.uk

Dr. Marie-Martine Bé (retired)

CEA/LNHB
91191 Gif-sur-Yvette, CEDEX, France
(Contact: Dr. Mark A. Kellett)

Dr. Valery P. Chechev

V.G. Khlopin Radium Institute
28, 2nd Murinsky Ave., 194021 St. Petersburg, Russia
Tel: 007 (812) 2473706
Fax: 007 (812) 2478095
E-mail: chechev@atom.nw.ru

Dr. Vanessa Chisté

CEA/LNHB
91191 Gif-sur-Yvette, CEDEX, France
Tel: 33-1-69-08-63-07
E-mail: vanessa.chiste@cea.fr

Dr. Mark A. Kellett

CEA/LNHB
91191 Gif-sur-Yvette, CEDEX, France
Tel: 33-1-69-08-27-76
Fax: 33-1-69-08-26-19
E-mail: mark.kellett@cea.fr

Dr. Tibor Kibédi

Department of Nuclear Physics
Australian National University
Canberra, Australia, ACT 0200
E-mail: Tibor.Kibedi@anu.edu.au

Dr. Nikolay K. Kuzmenko

V.G. Khlopin Radium Institute

28, 2nd Murinsky Ave., 194021 St. Petersburg, Russia

Tel: 007 (812) 2473706

Fax: 007 (812) 2478095

E-mail: akuzmenko@nk9433.spb.edu

Dr. Xavier Mougeot

CEA/LNHB

91191 Gif-sur-Yvette, CEDEX, France

Tel: 33-1-69-08-23-32

Fax: 33-1-69-08-26-19

E-mail: xavier.mougeot@cea.fr

Dr. Alan L. Nichols

Department of Physics

University of Surrey

Guildford GU2 7XH, United Kingdom

Tel: 44-1235-524077

E-mail: alanl.nichols@btinternet.com

RULES FOR EVALUATION AND COMPILATIONS

1. DATA SOURCES

Two main sources of data are used to obtain the recommended values:

- specific data evaluated from all available original publications (e.g., half-life),
- data already evaluated and compiled by specialists (e.g., Q-values); if a subsequent experimental study exists, the resulting measured value may be used, and its reference be included in a list of references for such a radionuclide.

2. EVALUATION RULES

All intermediate stages in the compilation and evaluation of a decay parameter are not presented in detail in order to avoid unnecessary complexity. The main stages comprise the following:

- critical analysis of published results and, if necessary, correction of these results to account for more recent values hitherto unavailable to the original experimentalists; as a rule, results without associated uncertainties are discarded, and the rejection of values is documented;
- data obtained through private communications are used only when all of the necessary information has been provided directly by the scientist who performed the measurements;
- adjustments may be made to the reported uncertainties;
- recommended values are deduced from an analysis of all measurements (or theoretical considerations), along with their standard deviations with a 1σ confidence level.

2.1. Evaluation of uncertainties

Definitions from “Guide to the expression of uncertainty in measurement” [1]:

Uncertainty (of measurement): parameter associated with the result of a measurement that characterizes the dispersion of the values that could reasonably be attributed to the measurand.

Standard uncertainty: uncertainty of the result of a measurement expressed as a standard deviation.

Type A evaluation (of uncertainty): method of evaluation of uncertainty by the statistical analysis of a series of observations.

Type B evaluation (of uncertainty): method of evaluation of uncertainty by means other than the statistical analysis of a series of observations.

The uncertainties given by authors are re-evaluated by combining the standard uncertainties σ_A and σ_B through the general law of variance propagation:

$$u_c = \sqrt{\sigma_A^2 + \sigma_B^2} \quad (1)$$

where u_c is the combined standard uncertainty,
 σ_A is the type A standard deviation, and
 σ_B is the type B standard uncertainty.

When the authors give insufficient information concerning their uncertainty calculations, the combined uncertainty u_c may be estimated by the evaluator, based on a knowledge of the measurement method(s).

2.2. Determination of the best value and associated uncertainty

(a) Results obtained by one author using one method

Sometimes only the final mean value and the combined standard uncertainty are given in the original publication. The following procedure is adopted if sufficient details are known.

For n individual values a_i ($i = 1 \dots n$), the best value is the arithmetical mean:

$$\bar{a} = \sum_{i=1}^n \frac{a_i}{n} \quad (2)$$

with type A standard deviation:

$$\sigma_A(\bar{a}) = \left[\frac{\sum_i (a_i - \bar{a})^2}{n(n-1)} \right]^{1/2} \quad (3)$$

If there are m contributions σ_{Bj} ($j = 1 \dots m$) to the type B standard uncertainty that are independent of each other:

$$\sigma_B(\bar{a}) = \left[\sum_{j=1}^m \sigma_{Bj}^2 \right]^{1/2} \quad (4)$$

Combined standard uncertainty:

$$u_c(\bar{a}) = \sqrt{\sigma_A^2(\bar{a}) + \sigma_B^2(\bar{a})} \quad (5)$$

Recommended value:

$$a = \bar{a} \pm u_c(\bar{a}) \quad (6)$$

(b) Results obtained by several authors employing the same method

For n individual values \bar{a}_i ($i = 1 \dots n$) having a standard deviation σ_{Ai} and a type B uncertainty σ_{Bi} , the best value is obtained by taking the mean weighted by the inverse of the variances.

$$\bar{a} = \frac{\sum_i (\bar{a}_i / \sigma_{Ai}^2)}{\sum_i (1 / \sigma_{Ai}^2)} \quad (7)$$

The associated values σ_A , σ_B are:

$$\sigma_A(\bar{a}) = \left[\sum_i (1 / \sigma_{Ai}^2) \right]^{-1/2} \quad (8)$$

$$\sigma_B(\bar{a}) = \sum_i (\sigma_{Bi})_{min} \quad \text{or} \quad \sigma_B(\bar{a}) = \sqrt{\sum_i (\sigma_{Bi})_{min}^2} \quad \text{or} \quad \sigma_B(\bar{a}) = (\sigma_B)_{min}$$

depending on the individual case, although $\sigma_B(\bar{a})$ cannot be less than the smallest σ_{Bi} .

σ_A and σ_B are combined quadratically to determine u_c :

$$u_c(\bar{a}) = \sqrt{\sigma_A^2(\bar{a}) + \sigma_B^2(\bar{a})} \quad (9)$$

and the recommended value is given by the expression:

$$a = \bar{a} \pm u_c(\bar{a}) \quad (10)$$

(c) Results obtained by different methods

When different measurement techniques have been applied, a weighted average is calculated using the combined uncertainties of the individual values as weights.

For n independent values a_i , each with a combined standard uncertainty u_{ci} , a weight p_i proportional to the inverse of the square of the individual u_{ci} can be assigned to each value.

$$a_w = \frac{\sum_{i=1}^n p_i a_i}{\sum_{i=1}^n p_i} \quad (11)$$

where the weights are $p_i = 1 / u_{ci}^2$.

An internal and an external uncertainty can be assigned to the mean value [2, 3]:

$$\sigma_{int}(a_w) = \left[\sum_i (1 / u_{ci}^2) \right]^{-1/2} \quad (12)$$

The internal variance $\sigma_{int}^2(a_w)$ is the expected uncertainty of the mean, based on the individual *a priori* variances u_{ci}^2 (by uncertainty propagation).

The external uncertainty is given by the equation:

$$\sigma_{ext}(a_w) = \left[\frac{\sum_i (a_i - a_w)^2 / u_{ci}^2}{(n-1) \sum 1 / u_{ci}^2} \right]^{1/2} \quad (13)$$

The external variance $\sigma_{ext}^2(a_w)$ includes the scatter of the data, and is based on the amount by which each a_i deviates from the mean when measured as a fraction of each given uncertainty u_{c_i} .

A measure of the consistency of the data is given by the ratio [2, 3]:

$$\sigma_{ext} / \sigma_{int} = \sqrt{\chi^2 / (n-1)} \quad (14)$$

If this ratio is significantly greater than unity, at least one of the input data most probably has an underestimated u_{c_i} which should be increased.

A critical value of $\chi^2 / (n-1)$ at 1 % confidence level is used as a practical test for discrepant data. The following table lists critical values of $\chi^2 / (n-1)$ for an increasing degree of freedom $\nu = n - 1$ [4].

v	critical $\chi^2/(n-1)$	v	critical $\chi^2/(n-1)$
1	6.6	12	2.2
2	4.6	13	2.1
3	3.8	14	2.1
4	3.3	15	2.0
5	3.0	16	2.0
6	2.8	17	2.0
7	2.6	18-21	1.9
8	2.5	22-26	1.8
9	2.4	27-30	1.7
10	2.3		
11	2.2	>30	$1 + 2.33\sqrt{2/\nu}$

If $\chi^2 / (n-1) \leq$ critical $\chi^2 / (n-1)$, the recommended value is given by:

$$a = a_w \pm \sigma_{int}(a_w) \quad (15)$$

If $\chi^2 / (n-1) >$ critical $\chi^2 / (n-1)$, the method of limitation of the relative statistical weight [3, 5] is recommended when there are three or more values; uncertainty of a value contributing more than 50 % to the total weight is increased to reduce its contribution to 50 % . The weighted and unweighted average and critical $\chi^2 / (n-1)$ are then recalculated:

if $\chi^2 / (n-1) \leq$ critical $\chi^2 / (n-1)$, the recommended value is given by:

$$a = a_w \pm (\text{the larger of } \sigma_{int}(a_w) \text{ and } \sigma_{ext}(a_w)) \quad (16)$$

if $\chi^2 / (n-1) >$ critical $\chi^2 / (n-1)$, the weighted or unweighted mean is chosen, depending on whether or not the uncertainties of the average values make them overlap with each other. If overlap occurs, the weighted average is recommended; otherwise the unweighted average is chosen. In either case, the uncertainty can be increased to cover the most accurate value.

Parameters evaluated according to these procedures and rules include half-lives, number of emitted particles, and some internal-conversion coefficients. All remaining data given in the tables of recommended data are generally taken from compilations.

2.3. Balanced decay schemes

All the probabilities for transitions and emitted radiations correspond to balanced decay schemes and permit the formulation of a fully consistent set of values. This balance implies the fulfillment of physical conservation principles as follows:

- The sum of the transition probabilities for all the transitions (α , β , ε) is equal to 1 (or 100 %); consequently, the sum of all the γ -ray transition probabilities (photons + internal conversion electrons) and all the (α , β , or ε) transitions feeding directly to the ground state is equal to 1 (or 100 %).
- For an excited nuclear level, the sum of the transition probabilities (α , β , γ , ε) feeding the level is equal to the sum of the transition probabilities depopulating this level;
- If the relative γ -ray emission probabilities $P(rel)_{\gamma_i}$ are known, the absolute emission probability $P(abs)_{\gamma_i}$ can be obtained from the equation:

$$P(abs)_{\gamma_i} = P(rel)_{\gamma_i} \times N \quad (17)$$

where N is the normalization factor, which may be determined from the equation:

$$N \sum_i P(rel)_{\gamma_i} (1 + \alpha_{i_i}) = 1 - B, \quad (18)$$

where α_{i_i} is the total conversion coefficient, and B , the (α , β , or ε) absolute branching to the ground state. The sum in equation (18) includes all the γ -ray transitions feeding the ground state.

3. COMPILATIONS

3.1. β and electron capture transitions

Depending on the individual radionuclide, the β -particle transition energies are either evaluated from experimental data (maximum β energies), or deduced from the atomic mass differences obtained from the tabulations of Audi and Wapstra [6] and the γ transition energies. The average β -particle energies are generally computed [7], and their $\log ft$ values as well as their ε/β^+ ratios are calculated using the tables of Gove and Martin [8].

Electron-capture transition energies are deduced from atomic mass differences and γ -ray transition energies. Capture probabilities P_K, P_L, \dots for allowed and non-unique first forbidden transitions can be calculated from equations where the ratios of the radial wave function components of the electron [9-11] and the corrective terms for exchange X^{LK} [12-16] are evaluated from tables.

3.2. γ -ray transitions

Internal conversion coefficients of pure multipolarity transitions are evaluated and compared with theoretical values that are interpolated from the tables of either Rösler *et al.* using a cubic spline method for $30 \leq Z \leq 104$ [17], or Band *et al.* [18]. The agreement of these theoretical values with experimental results is about 3 %.

Internal-conversion coefficients are calculated as described in Ref. [19] in order to include the effects of nuclear penetration in some M1 and E2 transitions.

Internal conversion coefficients for transitions with mixed multiplicities (e.g., M1 + E2) are calculated using tables of theoretical values using mixing ratios as shown below:

$$\alpha_i(\text{M1+E2}) = \alpha_i(\text{M1}) \frac{1}{1+\delta^2} + \alpha_i(\text{E2}) \frac{\delta^2}{1+\delta^2} \quad (19)$$

where $i = \text{K, L1, L2, } \dots \text{ T}$, refers to the individual atomic shell.

α_π coefficients for pair production are interpolated from theoretical values [20], with a precision between 5 % and 10 %.

3.3. Level spins and parities

Level spins and parities are usually from Nuclear Data Sheets [21].

3.4. Atomic shell constants

K-shell fluorescence yields ω_K and their uncertainties are taken from the evaluation of Bambynek *et al.* [22-24] with uncertainties ranging from 1 % ($Z > 35$) to 10 % ($Z = 5$), and from subsequent experimental results.

Mean L-shell fluorescence yields $\bar{\omega}_L$ are taken from the evaluation of Schönfeld and Janßen [25]. This evaluation includes both experimental [26-28] and theoretical values [29], and their uncertainties are equal to 4 % (for $Z > 29$).

Mean M-shell fluorescence yields $\bar{\omega}_M$ are obtained from the fitting of experimental data by Hubbell [28, 30].

Relative X-ray emission rates ($K\beta/K\alpha$) are taken from Schönfeld and Janßen [25], and $K\alpha_1/K\alpha_2$ from the theoretical values of Scofield [31]; uncertainties are assumed to be of the order of 1 %.

X-ray radiation energies are taken from the tables of Bearden [32].

Relative emission probabilities of K-Auger electron groups are deduced from the X-ray ratio, with uncertainties of the order of 3 % [25].

Energies of the K and L-Auger electrons are taken from Larkins [33].

The mean number of vacancies created in the L shell (from one K hole) n_{KL} and in the M shell (from one L hole) \bar{n}_{LM} are estimated from the preceding values.

3.5. m_0c^2 energy

m_0c^2 energy is defined as 510.998 902 (21) keV, as given by the CODATA Group [34].

REFERENCES

- [1] Guide to the expression of uncertainty in measurement, ISBN 92-67-20185-X ISO, Geneva, 1993.
- [2] WINKLER, G., "Data fitting and evaluation techniques for radioactive decay data", Proc. Symp. on Nuclear Data Evaluation Methodology, Brookhaven National Laboratory, USA, October 1992, World Scientific, Singapore (1993).
- [3] ZIJP, W.L., On the statistical evaluation of inconsistent measurement results illustrated on the example of the ^{90}Sr half-life, ECN Petten, Report ECN-179 (1985).
- [4] BEVINGTON, P.R., Data Reduction and Error Analysis for the Physical Sciences, Appendix C-4, McGraw-Hill (1969).
- [5] X-ray and gamma-ray standards for detector calibration, IAEA-TECDOC-619, Vienna (1991).
- [6] AUDI, G., WAPSTRA, A.H., The 1995 Atomic Mass Evaluation, Nucl. Phys. **A595** (1995) 4.
- [7] ENSDF – Logft program, ENSDF analysis and checking codes, Brookhaven National Laboratory, USA.
- [8] GOVE, N.B., MARTIN, M.J., Nucl. Data Tables **A10** (1971) 205.
- [9] SUSLOV, Y.P., Bull. Acad. Sci. USSR Phys. Ser. **33** (1969) 74.
- [10] BEHRENS, H., JANECKE, J., in Landolt-Bornstein: Numerical Data and Functional Relationships in Science and Technology, New Series Group 1, Vol. 4: Numerical Tables for Beta Decay and Electron Capture, Springer-Verlag, Berlin (1969).
- [11] MARTIN, M.J., BLICHERT-TOFT, P.H., Nucl. Data Tables **A8** (1970) 1.
- [12] BAHCALL, J.N., Phys. Rev. **129** (1963) 2683; *ibid.*, Phys. Rev. **132** (1963) 362; *ibid.*, Nucl. Phys. **71** (1965) 267.
- [13] FAESSLER, J.A., HUSTER, E., KRAFFT, O., KRAHN, F., Z. Phys. **238** (1970) 352.
- [14] SUSLOV, Y.P., Bull. Acad. Sci. USSR Phys. Ser. **34** (1970) 91.
- [15] VATAI, E., Nucl. Phys. **A156** (1970) 541.
- [16] SCHÖNFELD, E., Appl. Radiat. Isot. **49** (1998) 1353.
- [17] RÖSEL, F., FRIES, H.M., ALDER, K., PAULI, H.C., At. Data Nucl. Data Tables **21** (1978) 91.
- [18] BAND, I.M., TRZHASKOVSKAYA, M.B., NESTOR Jr., C.W., TIKKANEN, P.O., RAMAN, S., At. Data Nucl. Data Tables **81** (2002) 1.
- [19] PAULI, H.C., RAFF, U., A computer program for international conversion coefficients and particles parameters, Computer Physics Communications **9** (1975) 392.
- [20] SCHLUTER, P., SOFF, G., At. Data Nucl. Data Tables **24** (1979) 509.
- [21] NUCLEAR DATA SHEETS, National Nuclear Data Center, Brookhaven National Laboratory, USA.
- [22] BAMBYNEK, W., CRASEMANN, B., FINK, R.W., FREUND, H.-U., MARK, H., SWIFT, C.D., PRICE, R.E., VENUGOPALA RAO, P., Rev. Mod. Phys. **44** (1972) 716.
- [23] BAMBYNEK, W., Proc. Conf. on X-Ray and Inner Shell Processes in Atoms, Molecules and Solids, 20-24 August 1984, Leipzig, Germany.
- [24] W. BAMBYNEK, "Reorganization of atomic shells after radioactive decay", Proc. 2nd Int. Summer School, Low-level measurements of man-made radionuclides in the environment, June 1990, La Rabida, Huelva, Spain, World Scientific, Singapore (1991) 156-174.
- [25] SCHÖNFELD, E., JANSZEN, H., PTB report Ra-37 (1995).
- [26] MITCHELL, I.V., BARFOOT, K.M., Particle-induced X-Ray Emission Analysis, Application to Analytical Problems, Nucl. Sci. Appl. **1** (1981) 99-162.
- [27] COHEN, D.D., Nucl. Instrum. Meth. Phys. Res. **B22** (1987) 55.
- [28] HUBBELL, J.H., Bibliography and current status of K, L, and higher shell fluorescence yields for computation of photon energy absorption coefficients. NIST internal report 89-4144 (1989).
- [29] PURI, S., MEHTA, D., CHAND, B., SINGH, N., TREHAN, P.N., X-Ray Spectrometry **22** (1993) 358.
- [30] HUBBELL, J.H., TREHAN, P.N., SINGH, N., CHAND, B., MEHTA, D., GARG, M.L., GARG, R.R., SINGH, S., PURI, S., J. Phys. Chem. Ref. Data **23** (1994) 339.
- [31] SCOFIELD, J.H., Phys. Rev. **A9** (1974) 1041.
- [32] BEARDEN, J.A., Rev. Mod. Phys. **39** (1967) 78.
- [33] LARKINS, F.P., At. Data Nucl. Data Tables **20** (1977) 313.
- [34] MOHR, P.J., TAYLOR, B.N., Rev. Mod. Phys. **72** (2000) 351.

Table of contents

(Volumes 1 to 8 – All nuclides sorted by increasing mass number)

Mass	Nuclide	Page	Mass	Nuclide	Page	Mass	Nuclide	Page	Mass	Nuclide	Page
3	H-3	1	90	Sr-90	327	153	Gd-153	807	220	Rn-220	1127
7	Be-7	9	90	Y-90	329	154	Eu-154	813	221	Fr-221	1131
11	C-11	13	90	Y-90m	333	155	Eu-155	825	222	Rn-222	1139
13	N-13	17	93	Zr-93	337	159	Gd-159	833	223	Fr-223	1141
14	C-14	21	93	Nb-93m	339	166	Ho-166	839	223	Ra-223	1151
15	O-15	25	94	Tc-94m	343	166	Ho-166m	843	224	Ra-224	1163
18	F-18	29	99	Mo-99	369	169	Er-169	853	225	Ra-225	1167
22	Na-22	33	99	Tc-99	387	169	Yb-169	859	225	Ac-225	1171
24	Na-24	39	99	Tc-99m	393	170	Tm-170	865	226	Ra-226	1185
32	P-32	47	106	Ru-106	403	177	Lu-177	871	227	Ac-227	1191
33	P-33	51	106	Rh-106	405	182	Ta-182	879	227	Th-227	1197
35	S-35	53	108	Ag-108	425	186	Re-186	893	228	Ra-228	1209
36	Cl-36	55	108	Ag-108m	429	195	Au-195	899	228	Ac-228	1213
37	Ar-37	59	109	Pd-109	433	198	Au-198	903	228	Th-228	1249
40	K-40	63	109	Cd-109	443	201	Tl-201	913	231	Th-231	1259
41	Ar-41	69	110	Ag-110	453	203	Hg-203	917	231	Pa-231	1267
41	Ca-41	73	110	Ag-110m	459	203	Pb-203	921	232	Th-232	1289
44	Sc-44	79	111	In-111	471	204	Tl-204	927	232	U-232	1295
44	Ti-44	85	123	Te-123m	477	206	Hg-206	931	233	Th-233	1303
45	Ca-45	91	123	I-123	481	206	Tl-206	935	233	Pa-233	1315
46	Sc-46	93	124	Sb-124	491	207	Tl-207	943	234	Th-234	1329
47	Sc-47	97	125	Sb-125	525	207	Bi-207	947	234	Pa-234	1335
51	Cr-51	103	125	I-125	539	208	Tl-208	957	234	Pa-234m	1355
52	Fe-52	107	127	Sb-127	545	209	Tl-209	973	234	U-234	1367
54	Mn-54	111	127	Te-127	557	209	Pb-209	979	235	U-235	1373
55	Fe-55	117	127	Te-127m	565	209	Po-209	981	236	U-236	1387
56	Mn-56	121	127	Xe-127	573	210	Tl-210	985	236	Np-236	1391
56	Co-56	127	129	I-129	581	210	Pb-210	989	236	Np-236m	1397
57	Co-57	145	131	I-131	585	210	Bi-210	995	237	U-237	1401
57	Ni-57	155	131	Xe-131m	595	210	Po-210	999	237	Np-237	1407
58	Co-58	161	132	Te-132	599	211	Pb-211	1003	238	U-238	1417
59	Fe-59	167	133	I-133	603	211	Bi-211	1009	238	Np-238	1423
59	Ni-59	175	133	Xe-133	609	211	Po-211	1015	238	Pu-238	1429
60	Co-60	179	133	Xe-133m	615	211	At-211	1021	239	U-239	1441
61	Cu-61	183	133	Ba-133	619	212	Pb-212	1029	239	Np-239	1447
63	Ni-63	189	134	Cs-134	633	212	Bi-212	1033	239	Pu-239	1453
63	Zn-63	191	135	Xe-135m	641	212	Po-212	1041	240	Pu-240	1477
64	Cu-64	209	137	Cs-137	647	213	Bi-213	1043	241	Pu-241	1487
65	Zn-65	219	138	La-138	655	213	Po-213	1049	241	Am-241	1497
66	Ga-66	225	139	Ce-139	661	214	Pb-214	1053	242	Pu-242	1507
67	Ga-67	237	140	Ba-140	667	214	Bi-214	1061	242	Am-242	1513
68	Ga-68	249	140	La-140	673	214	Po-214	1075	242	Am-242m	1519
68	Ge-68	255	141	Ce-141	689	215	Bi-215	1079	242	Cm-242	1531
73	Se-73	257	144	Ce-144	695	215	Po-215	1089	243	Am-243	1539
75	Se-75	275	144	Pr-144	709	215	At-215	1095	243	Cm-243	1549
79	Se-79	285	144	Pr-144m	723	216	Po-216	1097	244	Am-244	1557
82	Rb-82	287	147	Nd-147	731	217	At-217	1101	244	Am-244m	1561
82	Sr-82	293	147	Pm-147	741	217	Rn-217	1105	244	Cm-244	1569
85	Kr-85	295	148	Pm-148	747	218	Po-218	1107	245	Cm-245	1577
85	Sr-85	299	148	Pm-148m	757	218	At-218	1109	246	Cm-246	1585
88	Y-88	303	151	Sm-151	769	218	Rn-218	1111	252	Cf-252	1591
89	Sr-89	311	152	Eu-152	775	219	At-219	1113			
89	Zr-89	315	153	Sm-153	795	219	Rn-219	1117			

Table of contents

(Volumes 1 to 8 – All nuclides sorted by alphabetical order)

Mass	Nuclide	Page	Mass	Nuclide	Page	Mass	Nuclide	Page	Mass	Nuclide	Page
225	Ac-225	1171	137	Cs-137	647	33	P-33	51	75	Se-75	275
227	Ac-227	1191	61	Cu-61	183	231	Pa-231	1267	79	Se-79	285
228	Ac-228	1213	64	Cu-64	209	233	Pa-233	1315	151	Sm-151	769
108	Ag-108	425	169	Er-169	853	234	Pa-234	1335	153	Sm-153	795
108	Ag-108m	429	152	Eu-152	775	234	Pa-234m	1355	82	Sr-82	293
110	Ag-110	453	154	Eu-154	813	203	Pb-203	921	85	Sr-85	299
110	Ag-110m	459	155	Eu-155	8235	209	Pb-209	979	89	Sr-89	311
241	Am-241	1497	18	F-18	29	210	Pb-210	989	90	Sr-90	327
242	Am-242	1513	52	Fe-52	107	211	Pb-211	1003	182	Ta-182	879
242	Am-242m	1519	55	Fe-55	117	212	Pb-212	1029	94	Tc-94m	343
243	Am-243	1539	59	Fe-59	167	214	Pb-214	1053	99	Tc-99	387
244	Am-244	1557	221	Fr-221	1131	109	Pd-109	433	99	Tc-99m	393
244	Am-244m	1561	223	Fr-223	1141	147	Pm-147	741	123	Te-123m	477
37	Ar-37	59	66	Ga-66	225	148	Pm-148	747	127	Te-127	557
41	Ar-41	69	67	Ga-67	237	148	Pm-148m	757	127	Te-127m	565
211	At-211	1021	68	Ga-68	249	209	Po-209	981	132	Te-132	599
215	At-215	1095	153	Gd-153	807	210	Po-210	999	227	Th-227	1197
217	At-217	1101	159	Gd-159	833	211	Po-211	1015	228	Th-228	1249
218	At-218	1109	68	Ge-68	255	212	Po-212	1041	231	Th-231	1259
219	At-219	1113	3	H-3	1	213	Po-213	1049	232	Th-232	1289
195	Au-195	899	203	Hg-203	917	214	Po-214	1075	233	Th-233	1303
198	Au-198	903	206	Hg-206	931	215	Po-215	1089	234	Th-234	1329
133	Ba-133	619	166	Ho-166	839	216	Po-216	1097	44	Ti-44	85
140	Ba-140	667	166	Ho-166m	843	218	Po-218	1107	201	Tl-201	913
7	Be-7	9	123	I-123	481	144	Pr-144	709	204	Tl-204	927
207	Bi-207	947	125	I-125	539	144	Pr-144m	723	206	Tl-206	935
210	Bi-210	995	129	I-129	581	238	Pu-238	1429	207	Tl-207	943
211	Bi-211	1009	131	I-131	585	239	Pu-239	1453	208	Tl-208	957
212	Bi-212	1033	133	I-133	603	240	Pu-240	1477	209	Tl-209	973
213	Bi-213	1043	111	In-111	471	241	Pu-241	1487	210	Tl-210	985
214	Bi-214	1061	40	K-40	63	242	Pu-242	1507	170	Tm-170	865
215	Bi-215	1079	85	Kr-85	295	223	Ra-223	1151	232	U-232	1295
11	C-11	13	138	La-138	655	224	Ra-224	1163	234	U-234	1367
14	C-14	21	140	La-140	673	225	Ra-225	1167	235	U-235	1373
41	Ca-41	73	177	Lu-177	871	226	Ra-226	1185	236	U-236	1387
45	Ca-45	91	54	Mn-54	111	228	Ra-228	1209	237	U-237	1401
109	Cd-109	443	56	Mn-56	121	82	Rb-82	287	238	U-238	1417
139	Ce-139	661	99	Mo-99	369	186	Re-186	893	239	U-239	1441
141	Ce-141	689	13	N-13	17	106	Rh-106	405	127	Xe-127	573
144	Ce-144	695	22	Na-22	33	217	Rn-217	1105	131	Xe-131m	595
252	Cf-252	1591	24	Na-24	39	218	Rn-218	1111	133	Xe-133	609
36	Cl-36	55	93	Nb-93m	339	219	Rn-219	1117	133	Xe-133m	615
242	Cm-242	1531	147	Nd-147	731	220	Rn-220	1127	135	Xe-135m	641
243	Cm-243	1549	57	Ni-57	155	222	Rn-222	1139	88	Y-88	303
244	Cm-244	1569	59	Ni-59	175	106	Ru-106	403	90	Y-90	329
245	Cm-245	1577	63	Ni-63	189	35	S-35	53	90	Y-90m	333
246	Cm-246	1585	236	Np-236	1391	124	Sb-124	491	169	Yb-169	859
56	Co-56	127	236	Np-236m	1397	125	Sb-125	525	63	Zn-63	191
57	Co-57	145	237	Np-237	1407	127	Sb-127	545	65	Zn-65	219
58	Co-58	161	238	Np-238	1423	44	Sc-44	79	89	Zr-89	315
60	Co-60	179	239	Np-239	1447	46	Sc-46	93	93	Zr-93	337
51	Cr-51	103	15	O-15	25	47	Sc-47	97			
134	Cs-134	633	32	P-32	47	73	Se-73	257			

³H – Comments on Evaluation by V.P. Chechev

The initial ³H decay data evaluation was done by Chechev in 1998 (1999Be). This current (revised) evaluation was carried out in April 2006. The literature available by April 2006 was included.

1. DECAY SCHEME

³H decays 100 % by β⁻-emission directly to the ground state of ³He.

2. NUCLEAR DATA

Q⁻ value is from 2003Au03.

The evaluated ³H half-life is based on the experimental data given in Table 1. This table has been taken from the paper of Lucas and Unterwiesinger (2000Lu17) which contains a comprehensive review and critical evaluation of the half-life of tritium.

Table 1. Experimental values of the ³H half-life (in years)

Reference	Author(s)	Measurement method	Half-life (years)	Stated uncertainty (years)	Meaning of the stated uncertainty	Comments
1936 McMillan	McMillan	Ionization current	>10	None	No uncertainty	Followed decay of radiation from irradiated beryllium for 4 months. OMITTED: limit only
1939 Alvarez	Alvarez and Cornog	Beta counting	0.41	0.11	Not given	One sample followed for 80 d. Chamber had diffusion losses. OMITTED: updated in 1940Alvarez
1940 Alvarez	Alvarez and Cornog	Beta counting	>10	None	No uncertainty	One sample followed for 5 months in new chamber. OMITTED: limit only
1940On01	O'Neal and Goldhaber	Beta counting	31	8	Not given	Counted tritium from irradiated lithium metal. OMITTED: outlier
1947Go08	Goldblatt <i>et al.</i>	Ionization current	10.7	2.0	Not given	Hydrogen + tritium in ionization chamber over 18 d. OMITTED: outlier
1947No01	Novick	Helium-3 collection	12.1	0.5	Not given	Two samples; accumulation times of 51 d and 197 d
1949Jenks	Jenks <i>et al.</i>	Helium-3 collection	12.46	0.20	Not given	Repeated measurements every two weeks until stable. OMITTED: updated in 1950Je60
1950Je60	Jenks <i>et al.</i>	Helium-3 collection	12.46	0.10	Probable error ^a	Four measurements over 206 d.
1951Jo15	Jones	Beta counting	12.41	0.05	Probable error ^a	Measurement of specific activity of tritium gas
1955Jo20	Jones	Helium-3 collection	12.262	0.004	Not given	Two samples; accumulation times of 578 d and 893 d
1958Po64	Popov <i>et al.</i>	Calorimetry	12.58	0.18	Not given	One sample; 21 measurements over 13 months
1963 Eichelberger	Eichelberger <i>et al.</i>	Calorimetry	12.355	0.010	Probable error ^a	Two samples measured over four years. OMITTED: updated in 1967Jo09

1966Merritt	Merritt and Taylor	Beta counting	12.31	0.13	Not given	Five gas counting measurements over 13 years
1967Jo09	Jordan <i>et al.</i>	Calorimetry	12.346	0.002	Probable error ^a	Five samples; 266 measurements over 6 years. OMITTED: updated in 1977RuZZ
1967Jo10	Jones	Helium-3 collection	12.25 12.31	0.08 0.42	99.7 % confidence limits	Two samples; accumulation times of 450 d to 800 d. Only the first value is usually quoted
1977RuZZ	Rudy and Jordan	Calorimetry	12.3232	0.0043	95 % confidence limits	Eight samples; 1353 measurements over 16 years
1980Un01	Unterweger <i>et al.</i>	Beta counting	12.43	0.05	1 standard uncertainty	Two sets of gas counting measurements 18 years apart. OMITTED: updated in 2000Unterweger
1987Bu28	Budick <i>et al.</i>	Bremsstrahlung counting	12.29	0.10	Not given	Two samples of tritium + xenon gas measured over 320 d. OMITTED: updated in 1991Bu13
1987O104	Oliver <i>et al.</i>	Helium-3 collection	12.38	0.03	1 standard uncertainty	Fifteen samples, each with accumulation times of 1 year to 2 years
1987Si01	Simpson	Beta counting	12.32	0.03	1 standard uncertainty	Tritium implanted in Si(Li) detector measured over 5.5 years
1988 Akulov	Akulov <i>et al.</i>	Helium-3 collection	12.279	0.033	1 standard uncertainty	Five series of measurements over 846 d
1991Bu13	Budick <i>et al.</i>	Bremsstrahlung counting	12.31	0.03	1 standard uncertainty	Two samples of tritium + xenon gas measured over 5.5 years
2000 Unterweger	Unterweger and Lucas	Beta counting	12.33	0.03	1 standard uncertainty	Three sets of gas counting measurements over 38 years

^a The probable error, PE, is the deviation from the population mean, μ , such that 50 % of the observations may be expected to lie between $\mu - PE$ and $\mu + PE$. For a normal distribution, the probable error can be converted to the standard deviation by multiplying by 1.4826.

As seen from Table 1 there are a number of measurements of the tritium half-life. Three of them stand out by their high precision (1955Jo20, 1967Jo09, 1977RuZZ). However, the uncertainties stated for the half-life in these works do not include an estimation of possible systematic errors. There are available newer measurements and discussions of the tritium half-life, so it is possible to estimate an "external" minimum uncertainty due to systematic effects (σ_{\min}) that should be added to the uncertainties stated in 1955Jo20, 1967Jo09 and 1977RuZZ. At that we can take into account the following circumstances:

a) The ³He collection result of 1955Jo20 has been obtained using only two points on each decay curve (for two samples). In the later work by the same method (1967Jo09) many experimental points were obtained on the decay curves (also for two samples) and the estimated systematic uncertainty made up 0.8 % for a 99.7 % confidence level.

b) The result of 1977RuZZ is a continuation of the measurements of 1967Jo09 for two tritide solids by calorimetric method for an additional 12 years. The difference of results of 1967Jo09 and 1977RuZZ proved to be 0.2 %, more than $5\sigma_{\exp}$ from 1977RuZZ and more than $10\sigma_{\exp}$ from 1967Jo09.

c) The comparative analysis of measurements of the radioactivity concentrations in several NBS tritiated-water standards over an 18-year period 1961 - 1978 (1980Un01) showed that for agreement of measurements (at given tritium half-life) their estimated standard errors (including a calorimetric method) should not be less 0.2 %.

Thus we have sufficient grounds for adding the "external" systematic error $\sigma_{\min} = 0.002 T_{1/2} (^3\text{H})$ into the uncertainties quoted in 1955Jo20, 1967Jo09 and 1977RuZZ. Lucas and Unterweger (2000Lu17) estimated the standard uncertainty of 1955Jo20 as 0.030 yr and that of 1977RuZZ as 0.025 yr.

Table 2 shows the modified set of half-life values, which has been formed from the original set by omitting the ten measurement results (see Comments in Table 1) and adjusting the uncertainties of 1955Jo20, 1977RuZZ and 1966Merritt. Latter was re-estimated in 2000Lu17.

Table 2. Selected measurement results for tritium half-life (in years)

Reference	Half-life	Measurement method	Comments on uncertainty
1947No01	12.1(5)	³ He collection	Author's stated uncertainty (ASU)
1950Je60	12.46(15)	³ He collection	ASU multiplied by 1.4826
1951Jo15	12.41(7)	Beta counting	Author's stated uncertainty
1955Jo20	12.262(30)	³ He collection	Uncertainty re-estimated in 2000Lu17
1958Po64	12.58(18)	Calorimetry	Author's stated uncertainty
1966Merritt	12.31(4)	Beta counting	Uncertainty re-estimated in 2000Lu17
1967Jo10	12.25(3)	³ He collection	Author's stated uncertainty
1977RuZZ	12.323(25)	Calorimetry	See text
1987Ol04	12.38(3)	³ He collection	Author's stated uncertainty
1987Si01	12.32(3)	³ H implanted into Si(Li)	Author's stated uncertainty
1988Akulov	12.279(33)	³ He collection	Author's stated uncertainty
1991Bu13	12.31(3)	Bremsstrahlung	Author's stated uncertainty
2000Unterweger	12.33(3)	Three sets of gas counting measurements over 38 years	Author's stated uncertainty

A weighted average for the final data set is 12.312 with an internal uncertainty of 0.010 and an external uncertainty of 0.013 and a reduced $\chi^2/\nu = 1.6$. An unweighted average is 12.33(3). Different statistical procedures from 1994Ka08 give the similar results: UINF, LWM, NORM – 12.312(10), PINF, BAYS and MBAYS – 12.312(13), IEXW – 12.314(14), RAJ – 12.311(10), CHV – 12.317(16). Lucas and Unterweger (2000Lu17) used three other statistical procedures including the method of determining the median and the estimated standard deviation of the median and adopted the value of 12.318(25).

The LWEIGHT computer program using the LWM procedure has led to the recommended value of 12.312(10).

The EV1NEW computer program (2000Ch01) has chosen the weighted average of 12.312 and recommended the smallest experimental uncertainty of 0.025 as a final uncertainty.

The adopted value of the ³H half-life is 12.312(25) years, or 4497(9) days.

It should be noted this half-life value has been evaluated for molecular tritium. The half-life of atomic tritium is less by ~0.26% (2004Ak16). See also 2005Ak04 for a bare triton half-life.

2.1. Tritium Beta End-Point Energy (E_{β}^0)

The tritium beta end-point energy depends upon the chemical state of the tritium in an experiment. The expression for E_{β}^0 of molecular tritium differs from that of a "bare" nucleus by the "chemical shift" $\Delta E = B(\text{RHe}^+) - B(\text{RT})$ (1985Ka21, 1989Re04) which is calculated taking into account the spectrum of

Comments on evaluation

final states (SFS). (Here the B values indicate electron binding energies for He+ ion and tritium atom, R indicates a chemical state).

For known ³He-³H atom mass difference (ΔMc^2) the tritium beta "end -point" energy measured in some experiment is :

$$E_{\beta}^0 = \Delta Mc^2 - E_{rec} - [B(He) - B(T)] + [B(RHe+) - B(RT)]$$

where E_{rec} is the helium recoil energy.

For tritium atom (nuclide) $E_{\beta}^0 = \Delta Mc^2 - 3.4 \text{ eV} - 64.3 \text{ eV} + \Delta E$ where $\Delta E = 40.82 \text{ eV}$.

With the recommended value of ΔMc^2 , the beta end-point energy for tritium nuclide is obtained by this way as 18563.6 eV. It is difficult to estimate the uncertainty of the ΔE calculation in 1985Ka21. Supposing it about the evaluated uncertainty of ΔMc^2 (Q value), we have E_{β}^0 (³H nuclide) = 18.564(2) keV.

For real forms of tritium sources in beta -spectrometry experiments the ³H end-point energies differ from the atomic value. For a molecular forms HT, CH₃T, valine the calculated E_{β}^0 makes 18572(2) eV. Below the measured end-point energies in some experiments are shown.

1987Bo07	Valine	18.579.4 ± 4 eV
1993Ba08	Molecular tritium	18.574.8 ± 0.6 eV
1993Su32	C ₁₄ H ₁₅ T ₆ O ₂ N ₃	18.578.3 ± 5.1 eV
1995St26	Gaseous tritium	18.568.5 ± 2.0 eV
2003Kr17	Gaseous tritium	18.570.5 eV

It should be noted that many works devoted to study of tritium beta -spectrum as it provided the most precise data of neutrino mass upper limit (see, for example, 2005Kr03, 2003Lo10, 2002Bo31 and references therein).

2.2. Average energy of beta particles of tritium per disintegration (<E_β>)

In Table 3 the available data of the <E_β> have been presented. The recommended value <E_β> has been obtained as the weighted average after corrections into the original results of the experiments and calculations. The calculation of the <E_β> with the LOGFT computer program using the adopted value Q⁻ = 18.591(1) keV gives 5.68 (±0.0011) keV.

Table 3. The available data of the tritium average beta energy (per disintegration, keV)

Reference	Method	Original	Re-estimated	Adopted
1950Je60	Calorimetry	5.69(4)	5.68(4) ^a	5.68(4)
1958Gr93	Calorimetry	5.57(1)	5.68(2) ^a	5.68(2)
1961Pi01	Calorimetry	5.73(3)	5.68(3) ^b	5.68(3)
1972Ma72	Calculation	5.7		5.7(1) ^d
1985Martin	Calculation	5.684(5)	5.680(5) ^c	5.68(1) ^d
1985Garcia	TDCR	5.70		5.70(2) ^d
1987Lagoutine, 1994Si21	Calculation	5.71(3)	5.70(3) ^c	5.70(3)
Recommended value 5.68(1) keV				

^a Corrected for the adopted tritium half-life of 12.312 y and heat output of 0.324(1) W/g

^b Corrected for the adopted tritium half-life of 12.312 y

^c Corrected for the adopted decay energy (Q⁻ = 18.591 keV)

^d Uncertainty attributed by the evaluator

3. REFERENCES

- 1936McMillan E.M. McMillan, Phys. Rev. 49(1936)875
[Half-life]
- 1939Alvarez L.W. Alvarez and L. Cornog, Phys. Rev. 56(1939)613
[Half-life]
- 1940Alvarez L.W. Alvarez and L. Cornog, Phys. Rev. 57(1940)248
[Half-life]
- 1940On01 R.D. O'Neal and M. Goldhaber, Phys. Rev. 58(1940)574
[Half-life]
- 1947Go08 M. Goldblatt, E.S. Robinson, and R.W. Spence, Phys. Rev. 72(1947)973
[Half-life]
- 1947No01 A. Novick, Phys. Rev. 72(1947)972
[Half-life]
- 1949Jenks G.H. Jenks, J.A. Ghormley, and F.H. Sweeton, Phys. Rev. 75(1949)701
[Half-life]
- 1950Je60 G.H. Jenks, F.H. Sweeton, and J.A. Ghormley, Phys. Rev. 80(1950)990
[Half-life, average beta energy]
- 1951Jo15 W.M. Jones, Phys. Rev. 83(1951)537
[Half-life]
- 1955Jo20 W.M. Jones, Phys. Rev. 100(1955)124
[Half-life]
- 1958Gr93 D.P. Gregory and D.A. Landsman, Phys. Rev. 109(1958)2091
[Average beta energy]
- 1958Po64 M.M. Popov et al., Atomnaya Energ.4(1958)269; J. Nucl. Energy 9(1959)190
[Half-life]
- 1961Pi01 W.L. Pillinger, J.J. Hentges, and J.A. Blair, Phys. Rev.121(1961)232
[Average beta energy]
- 1963Eichelberger J.F. Eichelberger, G.R. Grove, and L.V. Jones. Progress Report MLM -1160, US Department of Energy, Mound Laboratory, Miamisburg, Ohio, June 1963, p.5-6
[Half-life]
- 1966Merritt J.S. Merritt, J.G.V. Taylor, Chalk River Report AECL-2510(1966)
[Half-life]
- 1967Jo09 K.C. Jordan, B.C. Blanke, and W.A. Dudley, J. Inorg. Nucl. Chem. 29(1967)2129
[Half-life]
- 1967Jo10 P.M.S. Jones, J. Nucl. Materials 21(1967)239
[Half-life]
- 1972Ma72 J. Mantel, Intern. J. Appl. Rad. Isotopes 23(1972)407
[Average beta energy]
- 1977RuZZ C.R. Rudy and K.C. Jordan, In: MLM-2458, Monsanto Research Corporation, Miamisburg, Ohio(1977)
[Half-life]
- 1980Un01 M.P. Unterweger et al., Intern. J. Appl. Rad. Isotopes 31(1980)611
[Half-life]

- 1985Garcia E. Garcia-Torano and A. Gra u Malonda, *Comp. Phys. Commun.* 36(1985)307. See also 1994Si21
[Average beta energy]
- 1985Ka21 I.G. Kaplan, G.V. Smelov, and V.N. Smutny, *Phys.Lett.* 161B(1985)389
[Beta end-point energy and Q-value]
- 1985Martin M.J. Martin, In: *A handbook of radioactivity measurement procedures*, NCRP Report No 58 (1985), 2nd Edition, NCRP, Bethesda, Maryland, 368-373
[Average beta energy]
- 1987Bo07 S.D. Boris et al., *Phys.Rev.Lett.* 58, 2019 (1987); Erratum *Phys.Rev.Lett.* 61, 245 (1988)
[Beta end-point energy]
- 1987Bu28 B. Budick and Hong Lin, *Bull.Amer.Phys.Soc.*32 (1987)1063
[Half-life]
- 1987Lagoutine F. Lagoutine, N. Coursol, J. Legrand, *Table de Radionuclé ides*, ISBN-2-7272-0078-1 (LMRI, 1982-1987)
[Average beta energy]
- 1987Ol04 B.M. Oliver, H. Farrar IV, and M.M. Bretscher, *Intern. J. Appl. Rad. Isotopes* 38(1987)959
[Half-life]
- 1987Si01 J.J. Simpson, *Phys. Rev.* C35(1987)752
[Half-life]
- 1988Akulov Yu.A. Akulov, B.A. Mamyrin, L.V. Khabarin, V.S. Yudenich, and N.N. Ryazantseva, *Zh. Tekh. Fiz.* 14(1988)940, *Sov. Tech. Phys. Lett.* 14(1988)416
[Half-life]
- 1989Re04 A. Redondo and R.G.H. Robertson, *Phys. Rev.* C40(1989)368
[Beta end-point energy and Q-value]
- 1991Bu13 B. Budick, Jiansheng Chen, and Hong Lin, *Phys. Rev. Lett.* 67(1991)2630
[Half-life]
- 1993Ba08 H. Backe, H. Barth, J. Bonn et al., *Nucl. Phys.* A553(1991)313c
[Beta end-point energy]
- 1993Su32 H. Sun, D. Liang, S. Chen et al. *J. Chin. Nucl.Phys.* 15(1993)261
[Beta end-point energy]
- 1994Ka08 S.F. Kafala, T.D. MacMahon, and P.W. Gray, *Nucl. Instrum. Methods Phys. Res.* A339(1994)151
[Evaluation technique]
- 1994Si21 B.R.S. Simpson and B.R. Meyer, *Nucl. Instrum. Methods Phys. Res.* A339(1994)14
[Average beta energy]
- 1995St26 W. Stoeffl and D.J. Decman, *Phys. Rev. Lett.* 75(1995)3237
[Beta end-point energy]
- 1999BeM.-M. Bé, E.Browne, V.Chechev *et al.* In: *Table de Radionuclé ides*, CEA-ISBN 2-7272-0200-8, *Comments on Evaluations*, CEA-ISBN 2-7272-0211-3. 1999
[³H decay data evaluation-1998]
- 2000Ch01 V.P. Chechev and A.G. Egorov, *Appl. Radiat. Isot.* 52(2000)601
[Evaluation technique]
- 2000Lu17 L.L. Lucas and M.P. Unterweger, *J. Res. Natl. Inst. Stand. Technol.* 104(2000)541
[Half-life evaluation]

- 2000Unterweger M.P. Unterweger and L.L. Lucas, Appl. Radiat. Isot. 52(2000)527
[Half-life]
- 2002Bo31 J. Bonn, B. Bornschein, L. Bornschein et al., Prog. Part. Nucl. Phys. 48(2002)133
[Tritium beta-spectrum]
- 2003Au03 G. Audi, A.H. Wapstra, and C. Thibault, Nucl. Phys. A729(2003)337
[Q-value]
- 2003Kr17 Ch. Kraus, J. Bonn, B. Bornschein et al., Nucl. Phys. A721(2003)533c
[Beta end-point energy]
- 2003Lo10 V.M. Lobashev. Nucl. Phys. A719(2003)153c
[Tritium beta-spectrum]
- 2004Ak16 Yu.A. Akulov and B.A. Mamyurin. Phys. Lett. B600(2004)41
[Half-life]
- 2005Ak04 Yu.A. Akulov and B.A. Mamyurin, Phys. Lett. B610(2005)45
[Half-life]
- 2005Kr03 Ch. Kraus, B. Bornschein, L. Bornschein et al., Eur. Phys. J. C40(2005)447
[Tritium beta-spectrum]

⁷Be - Comments on Evaluation of Decay Data by R. G. Helmer

This evaluation was originally done in 1996 by R. G. Helmer and E. Schönfeld and minor editing was added in December 2000.

1. Decay Scheme

This decay scheme is complete since the only levels in ⁷Li below the decay energy are populated.

2. Nuclear Data

The Q value is from the mass evaluation in 1995Au04.

The adopted half-life is 53.22 (6) days.

The ⁷Be half-life has been observed to vary depending on the chemical form of the ⁷Be. Some of these measured variations are:

Reference	Chemical forms compared	$(\Delta T_{1/2} \times 10^4) / T_{1/2}$
1949Se20	Be - BeO	1.5 (9)
1953Kr16	Be - BeO	1.3 (5)
	BeO- BeF ₂	6.1 (6)
	Be - BeF ₂	7.4 (5)
1956Bo36	Be - BeF ₂	12 (1)
1970Jo21	BeO- BeF ₂	11.3 (6)
	BeO - BeBr ₂	14.7 (6)
	BeO- Be ₄ O(CH ₃ COO) ₆	-7.2 (6)
	BeO- Be(C ₅ H ₅) ₂	8.0 (7)
	BeO- Be(OH ₂) ₄	-3.7 (8)
	BeF ₂ - Be ₄ O(CH ₃ COO) ₆	-18.5 (8)
	Be(C ₅ H ₅) ₂ - Be(OH ₂) ₄	-11.7 (11)
	1999Hu20	BeO - Be(OH) ₂
	BeO - Be ²⁺ (OH ₂) ₄	-98.
1999Ra12	Be in Au - Be in Al ₂ O ₃	72 (7)

Excluding the much larger changes reported by 1999Hu20 and 1999Ra12, these measured changes range from 0.01% to 0.2%, or from 0.005 to 0.10 days, or 0.08 days, if the organic compounds are also omitted.

The adopted value of 53.22 (6) is from Limitation of Relative Statistical Weight (LRSW) (1985ZiZY, 1992Ra09) analysis of 53 (2) (1940Hi01), 52.93 (22) (1949Se20), 53.61 (17) (1953Kr16), 53.0 (4) (1956Bo36), 53.5 (2) (1957Wr37), 53.1 (3) (1965En01), 53.52 (10) (1970Jo21), 53.0 (3) (1974Cr05), 53.17 (2) (1975La16), 53.16 (1) (1982ChZF), 53.284 (4) (1982RuZV), and 53.12 (7) (1996Ja10). In this analysis the uncertainty of 1982RuZV value was increased from 0.004 to 0.0088 so that its relative

weight was reduced from 83 % to 50 %. The weighted average of these values is 53.225 with an internal uncertainty of 0.006, a reduced- χ^2 of 10.5, and an external uncertainty of 0.020. This uncertainty is increased by the LRSW method to 0.06 so that the most precise value of 53.284 is included; this uncertainty also includes the next most precise value of 53.16.

The chemical forms of the samples for which these half-lives were determined are: 1949Se20 Be metal or BeO and difference is not significant, 1953Kr16 Be metal, 1956Bo36 Be metal or BeF₂ and difference is not significant, 1970Jo21 average of data for BeF₂, BeO, and Be(C₅H₅)₂, and 1975La16 isolated Be atoms in aluminum matrix.

The adopted half-life is dominated by the values of 1975La16, 1982ChZF, and 1982RuZV which contribute 10 %, 39 %, and 50 % of the relative weight, respectively. The values of 1982ChZF and 1982RuZV differ by $\sim 10\sigma$ and contribute 3.8 and 4.1 to the reduced- χ^2 value of 10.5. Since these three values differ by 0.12 days and the chemical forms in the latter two cases are not known, the chemical variation data in the above table suggest that some of this difference may be due to chemical effects. This suggests that the adopted uncertainty of 0.06 days is reasonable for general use. In any case, the data on the chemical effects indicate that the adopted value can certainly be used for Be and BeO sources.

Values not used are 54.5 (J. F. Bonner as quoted in 1953Kr16, no uncertainty); and 54.3 (5) (1947BoAA as quoted in 1953Kr16, superseded by value of 1956Bo36); and 53.694 (6), 53.416 (6), and 54.226 (6) (1999Hu20). The values of 1999Hu20 have very small uncertainties and have very large variations, up to 1.5%, with chemical form which need to be confirmed. If this large shift and that of 1999Ra12 are correct, they would invalidate the uncertainty of our adopted value.

Also, the results of 2000Hu20 and 2000Li21 were obtained after this evaluation was completed, but these results would not change the adopted value.

Recent experiments have shown that the half-life of ⁷Be increases as much as 0.7% by imbedding this radionuclide in different matrices. The recommended value presented in this evaluation should be adequate for Be and BeO samples.

2.1 Electron-capture transitions

The adopted value for the electron capture to the 477-keV level is $P_{\epsilon}(477) = 10.44\%$ (4). This value is a weighted average of 10 (+20-7) (1938RuAA), 10.7 (20) (1949Wi13), 11.8 (12) (1949Tu06), 12.3 (6) (1951Di12), 10.35 (8) (1969TaZX), 10.47 (20) (1970MuZU), 10.42 (18) (1973Po10), 10.35 (8) (1974Go26), 10.10 (45) (1983Ba15), 10.61 (23) (1983Da14), 10.6 (5) (1983Do07), 10.9 (11) (1983Kn10), 10.7 (2) (1983Ma34), 9.8 (5) (1983No03), 11.4 (7) (1984Ev01), 10.61 (17) (1984Fi10), and 10.49 (7) (1984Sk01). This weighted average has an internal uncertainty of 0.039, a reduced- χ^2 of 1.35, and an external uncertainty of 0.045. The adopted value is dominated by the values of 1969TaZX, 1974Go24, and 1984Sk01 which contribute 23 %, 23 %, and 30 % of the relative weight, respectively. The largest contribution to the reduced- χ^2 is 0.6 from 1951Di12.

Values not used are 10.32 (16) (1962Ta11, superseded by 1969TaZX) and 10.5 (2) (W. Poenitz, 1966, superseded by 1973Po10).

The P_K and P_L values of 0.908 (12) and 0.092 (12) were calculated from the tables in 1998Sc28. The values from the LOGFT code are 0.97 and 0.03, which are different.

2.2 Gamma-ray transition

The γ -ray transition energy is computed from the γ -ray energy.

The internal-conversion coefficient is the measured value of 1964Kr04 and the mixing ratio was also determined by 1964Kr04. The theoretical values interpolated from the tables of 1976Ba63 are 7.73×10^{-7} for M1 and 2.96×10^{-6} for E2.

The gamma transition probability is :

Within its uncertainty, $P_\gamma(477) = I_\gamma(477) \times (1.0 + \alpha) = P_\epsilon(477)$

With $I_\gamma(477) = 10.44$ (4) % (c.f. § 2.1)

3. Atomic Data

The fluorescence yield is from the compilation of 1994Hu23.

4. Radiations

The conversion electron emission intensity is computed from $P_\gamma(477)$ and α_K .

The γ -ray energy is from the evaluation of 2000He14.

5. Main Production Modes

${}^6\text{Li}(d,n)$, ${}^{10}\text{B}(p,\alpha)$, and ${}^{12}\text{C}({}^3\text{He},2\alpha)$

6. References

- 1938RuAA L. H. Rumbaugh, R. B. Roberts, L. R. Hafstad, Phys. Rev. **54** (1938) 657 [P_ε]
 1940Hi01 J. E. Hill, Phys. Rev. **57** (1940) 567 [T_{1/2}]
 J. F. Bonner, Jr., report AECU-107, as quoted in 1953Kr16 [T_{1/2}]
 1947BoAA P. Bouchez, Daudel, Muxart, J. Phys. et Radium **8** (1947) 336, as quoted in 1953Kr16 [T_{1/2}]
 1949Se20 E. Segre, C. E. Wiegand, Phys. Rev. **75** (1949) 39; erratum Phys. Rev. **81** (1951) 284 [T_{1/2}]
 1949Tu06 C. M. Turner, Phys. Rev. **76** (1949) 148 [P_ε]
 1949Wi13 R. M. Williamson, H. T. Richards, Phys. Rev. **76** (1949) 614 [P_ε]
 1951Di12 J. M. Dickson, T. C. Randle, Proc. Phys. Soc. (London) **64A** (1951) 902 [P_ε]
 1953Kr16 J. J. Kraushaar, E. D. Wilson, K. T. Bainbridge, Phys. Rev. **90** (1953) 610 [T_{1/2}]
 1956Bo36 P. Bouchez, J. Tobaillem, J. Robert, R. Muxart, R. Mellet, P. Daudel, J. Phys. Rad. **17** (1956) 363 [T_{1/2}]
 1957Wr37 H. W. Wright, E. I. Wyatt, S. A. Reynolds, W. S. Lyon, T. H. Handley, Nucl. Sci. Eng. **2** (1957) 427 [T_{1/2}]
 1962Ta11 J. G. V. Taylor, J. S. Merritt, Can. J. Phys. **40** (1962) 926 [P_ε]
 1964Kr04 A. Kriester, Kernenergie **7** (1964) 748 [α]
 1965En01 J. B. A. England, B. L. Reece, Nucl. Phys. **72** (1965) 449 [T_{1/2}]
 1966PoAA W. Poenitz, J. Nucl. Energy **20** (1966) 825 [P_ε]
 1969TaZX J. G. V. Taylor, J. S. Merritt, report AECL-3512 (1969) [P_ε]
 1970MuZU M. Mutterer, Neutron Standards and Flux Normalization 452, AEC Symp. Series 23 (1970) [P_ε]

- 1970Jo21 H. W. Johlige, D. C. Aumann, H. J. Born, Phys. Rev. **C2** (1970) 1616 [$T_{1/2}$]
 1973Po10 W. P. Poenitz, A. Devolpi, Intern. J. Appl. Radiat. Isot. **24** (1973) 471 [P_{ϵ} , P_{γ}]
 1974Cr05 P. J. Cressy, Jr., Nucl. Sci. Eng. **55** (1974) 450 [$T_{1/2}$]
 1974Go26 I. W. Goodier, J. L. Makepeace, A. Williams, Intern. J. Appl. Radiat. Isot. **25** (1974) 373 [P_{ϵ} , P_{γ}]
 1975La16 F. Lagoutine, J. Legrand, C. Bac, Intern. J. Appl. Radiat. Isot. **26** (1975) 131 [$T_{1/2}$]
 1976Ba63 I. M. Band, M. B. Trzhaskovskaya, M. A. Listengarten, At. Data Nucl. Data Tables **18** (1976) 433 [α]
 1982ChZF P. Christmas, report NBS-SP-626 (1982) 100 & 198 [$T_{1/2}$]
 1982RuZV A. R. Rutledge, L. V. Smith, J. S. Merritt, report NBS-SP-626 (1982) 5 [$T_{1/2}$]
 1983Da14 C. N. Davids, A. J. Elwyn, B. W. Filippone, S. B. Kaufman, K. E. Rehm, J. P. Schiffer, Phys. Rev. C **28** (1983) 885 [P_{ϵ}]
 1983Ba15 D. P. Balamuth, L. Brown, T. E. Chapuran, J. Klein, R. Middleton, R. W. Zurmühle, Phys. Rev. C **27** (1983) 1724 [P_{ϵ}]
 1983Do07 T. R. Donoghue, E. Sugarbaker, M. Wiescher, T. C. Rinckel, K. E. Sale, C. P. Browne, E. D. Berners, R. W. Tarara, R. E. Warner, Phys. Rev. C **28** (1983) 875 [P_{ϵ}]
 1983Kn10 D. A. Knapp, A. B. McDonald, C. L. Bennett, Nucl. Phys. **A411** (1983) 195 [P_{ϵ}]
 1983Ma34 G. J. Mathews, R. C. Haight, R. G. Lanier, R. M. White, Phys. Rev. C **28** (1983) 879 [P_{ϵ}]
 1983No03 E. B. Norman, T. E. Chupp, K. T. Lesko, J. L. Osborne, P. J. Grant, G. L. Woodruff, Phys. Rev. C **27** (1983) 1728; erratum Phys. Rev. C **28** (1983) 1409 [P_{ϵ}]
 1984Ev01 H. C. Evans, I. P. Johnstone, J. R. Leslie, W. McLatchie, H.-B. Mak, P. Skensved, T. K. Alexander, Can. J. Phys. **62** (1984) 1139 [P_{ϵ}]
 1984Fi10 S. A. Fisher, R. L. Hershberger, Nucl. Phys. **A423** (1984) 121 [P_{ϵ}]
 1984Sk01 R. T. Shelton, R. W. Kavanagh, Nucl. Phys. **A414** (1984) 141 [P_{ϵ}]
 1985ZiZY W. L. Zijp, report ECN-179 (1985) [averaging]
 1992Ra09 M. U. Rajput, T. D. MacMahon, Nucl. Instr. Meth. **A312** (1992) 289 [averaging]
 1994Hu23 J. H. Hubbell, P. N. Trehan, Nirmal Singh, B. Chand, M. L. Garg, R. R. Garg, Surinder Singh, S. Puri, J. Phys. Chem. Ref. Data **23** (1994) 339 [ω_K]
 1995Au04 G. Audi, A. H. Wapstra, Nucl. Phys. **A595** (1995) 409 [Q]
 1996Ja10 M. Jaeger, S. Wilmes, V. Kölle, G. Staudt, P. Mohr, Phys. Rev. C **54** (1996) 423 [$T_{1/2}$]
 1998Sc28 E. Schönfeld, Appl. Radiat. Isot. **49** (1998) 1353 [P_L/P_K]
 1999Hu20 C.-H. Huh, Earth Plant. Sci. Lett. **171** (1999) 325 [$T_{1/2}$]
 1999Ra12 A. Ray, P. Das, S. K. Saha, S. K. Das, B. Sethi, A. Mookerjee, C. Basu Chauduri, G. Pari, Phys. Lett. B **455** (1999) 69 [$T_{1/2}$]
 2000He14 R. G. Helmer, C. van der Leun, Nucl. Instr. Meth. A **450** (2000) 35 [E_{γ}]
 2000Hu20 C. A. Huh, L. G. Liu, Journal of Radioanalytical and Nuclear Chemistry **246** (2000) 229 [$T_{1/2}$]
 2000Li21 L. G. Liu, C. A. Huh, Earth and Planetary Science Letters **180** (2000) 163 [$T_{1/2}$]

¹¹C – Comments on evaluation of decay data by V. Chisté and M. M. Bé

1) Decay Scheme

¹¹C disintegrates by β^+ emission (99.750(13)%) and electron capture (0.250(13)%) to the ground state of the stable nuclide ¹¹B.

2) Nuclear Data

The Q value (1982.5(9) keV) is from Audi and Wapstra evaluation (1995Au04), and has been calculated with the formula:

$$Q = M(A, Z) - M(A, Z - 1),$$

where M(A,Z) and M(A,Z-1) are the measured atomic masses of ¹¹C and ¹¹B, respectively.

E_{β^+} , calculated from this Q value ($E_{\beta^+} = 960.5(9)$ keV), is in agreement with a weighted average value of 959.8(5) keV, which was calculated from measured values (see **β^+ Transition and Electron Capture Transition**).

The measured ¹¹C half-life values (in minutes) are given below:

$T_{1/2}$

Reference	Value (min)
Smith (1941Sm11)	20.35 (8)
Solomon (1941So01)	20.5 (6)
Siegbahn (1944Si30)	20.0(4)
Dickson (1951Di12)	20.0 (1)
Kundu (1953Ku08)	20.74 (10)
Barber (1955Ba63)	20.26 (10)
Prokoshkin (1957Pr53)	20.8 (2)
Arnell (1958Ar15)	20.11 (13)
Kavanagh (1964Ka31)	20.34 (4)
Patterson (1965Pa10)	20.8 (4)
Awschalom (1969Aw02)	20.40 (4)
Hogstrom (1973Ho43)	19.8 (8)
Singh (1973SiYS)	20.0 (3)
Azuelos (1975Az01)	20.382 (20)
Behrens (1975Be28)	20.32 (12)

Evaluators calculated the weighted average of these 15 values using the Lweight program (version 3) as 20.369 min with an external uncertainty of 0.028 and a reduced χ^2 of 3.07. The value of Azuelos (1975Az01) has a relative statistical weight of 54%. Evaluators rejected Siegbahn's (1944Si30) value (quoted by Janecke (1960Ja12) and Raman (1978Ra21)), because they could not find the article, and therefore no details were available on how Siegbahn obtained such a value. For the remaining 14 values,

the largest contribution to the weighted average comes from the value of Azuelos (1975Az01), with a relative statistical weight of 57%. The program Lweight 3 has increased the uncertainty of the 1975Az01 value from 0,02 to 0,0231 in order to reduced its relative statistical weight to 50%. The adopted value is the weighted average : 20.370 min, with an external uncertainty of 0.029 min. The reduced χ^2 is 3.24.

b⁺ Transition and Electron capture transition

For the K/ β^+ ratio, the following values have been found in the literature:

Reference	Value (10 ⁻³)
Scobie (1957Sc02)	1.9(3)
Campbell (1967Ca21)	2.30 (+0.14;-0.11)

β^+ and electron capture probabilities have been calculated using the most recent value of K/ β^+ ratio measured by Campbell (1967Ca21), $P_K/P_{EC} = 0.9174(91)$ (See Section 2.2), and normalizing to a total probability ($P_{\beta^+} + P_{EC}$) of 100%. This leads to $P_{\beta^+} = 99.750(13)$ % and $P_{EC} = 0.250(13)$, respectively. The uncertainties were calculated through their propagation on the above formulas.

The experimental K/ β^+ ratio of Campbell is close to the theoretical values:

- a) 2.222 10⁻³ calculated with LOGFT program;
- b) 2.00 10⁻³ calculated by Scobie (1957Sc02);
- c) 2.18 10⁻³ calculated by Campbell (1967Ca21);
- d) 2.46 10⁻³ calculated by Vatai (1968Va23);
- e) 2.316 10⁻³ given by Fitzpatrick (1973Fi13);
- f) 2.11 10⁻³ given by Bambynek (1977Ba49);

Evaluators calculated a lg ft of 3.592 for this allowed transition. The value agrees with 3.599 suggested by Ajzenberg-Selove (1980Aj01, 1985Aj01 and 1990Aj01).

The partial sub shell capture probabilities given in Section 2.2 were calculated using the program EC Capture for an allowed transition.

The weighted mean of the β^+ end-point energy has been calculated (with the Lweight program, version 3) using the following measured values (in keV):

Reference	Values (keV)
Townsend (1940To03)	981(5)
Moore (1940Mo40)	1030(30)
Siegbahn(1944Si30)	993(1)
Richards (1950Ri07)	958(3)
Wong (1954Wo19)	968(8)
Campbell (1967Ca21)	958.2(14)
Fitzpatrick (1973Fi13)	960.2(10)
Azuelos (1975Az01)	960.0(10)
Behrens (1978Be28)	960.8(26)
Raman (1978Ra21)	960.1(11)

The weighted average of these 10 values is 967 keV with an uncertainty of 2.6 keV and a reduced χ^2 of 97. The values of 1944Si30, 1973Fi13 and 1975Az01 have a relative weight of 21%. The Townsend (1940To03), Moore (1940Mo40), Siegbahn (1944Si30) and Wong (1954Wo19) values have been rejected by the Lweight program, based on the Chauvenet's criterion. For the remaining 6 values, the largest contribution to the weighted average comes from the values of Fitzpatrick (1973Fi13) and Azuelos

(1975Az01), amounting to a statistical weight of 28%. The weighted average is 959.8 keV, with an internal uncertainty of 0,5 keV and a reduced χ^2 of 0,41. This value is in agreement with E_{β^+} (960.5(9) keV) deduced from the adopted Q value (1995Au04) in this evaluation.

3) Gamma-ray Emissions

The annihilation radiation emission probability ($I_{\gamma 511}$) is P_{β^+} (=99.750(13)%), multiplied by 2, without the correction factor for the annihilation-in-flight process in the medium. That is, $I_{\gamma 511} = 199.500(26)\%$.

References

- 1940Mo40 – B. L. Moore, Phys. Rev. 57 (1940) 355 [end-point energy].
 1940To03 – A. A. Townsend, Proc. Roy. Soc. (London) 177 (1940) 357 [end-point energy].
 1941Sm11 – J. H. C. Smith, D. B. Cowie, J. Appl. Phys. 12 (1941) 78 [$T_{1/2}$].
 1941So01 – A. K. Solomon, Phys. Rev. 60 (1941) 279 [$T_{1/2}$].
 1944Si30 – Siegbahn, Arkiv Mat. Astron. Fysik 30B (1944) 20 [$T_{1/2}$].
 1950Ri07 – H. T. Richards, R. V. Smith, Phys. Rev. 77 (1950) 752 [end-point energy].
 1951Di12 – J. M. Dickson, T. C. Randle, Proc. Phys. Soc. (London) 64A (1951) 902 [$T_{1/2}$].
 1953Ku08 – D. N. Kundu, T. W. Donaven, M. L. Pool, J. K. Long, Phys. Rev. 89 (1953) 1200 [$T_{1/2}$].
 1954Wo19 – C. Wong, Phys. Rev. 95 (1954) 765 [end-point energy].
 1955Ba63 – W. C. Barber, W. D. George, D. D. Reagan, Phys. Rev. 98 (1955) 73 [$T_{1/2}$].
 1957Pr53 – Iu. D. Prokoshkin, A. A. Tiapkin, Soviet. Phys. JETP 5 (1957) 148 [$T_{1/2}$].
 1957Sc02 – J. Scobie, G. M. Lewis, Phil. Mag. 2 (1957) 1089 [K/β^+ ratio].
 1958Ar15 – S. E. Arnell, J. Dubois, O. Almen, Nucl. Phys. 6 (1958) 196 [$T_{1/2}$].
 1960Ja12 – V. J. Janecke, Z. F. Naturf. 15A (1960) 593 [$T_{1/2}$].
 1964Ka31 – T. M. Kavanagh, J. K. P. Lee, W. T. Link, Can. J. Phys. 42 (1964) 1429 [$T_{1/2}$].
 1965Pa10 – J. R. Patterson, J. M. Poate, E. W. Tuttert, B. A. Robson, Proc. Phys. Soc. (London) 86 (1965) 1297 [$T_{1/2}$].
 1967Ca21 – J. L. Campbell, W. Leiper, K. W. Ledingham, R. W. P. Drever, Nucl. Phys. A96 (1967) 279 [K/β^+ ratio, end-point energy].
 1968Va23 – E. Vatai, Proc. Conf. Electron Capture and higher order processes in Nucl. Decay 2 (1968) 71 [K/β^+ ratio].
 1969Aw02 – M. Awschalom, F. L. Larsen, W. Schimmerling, Nucl. Inst. Meth. 75 (1969) 93 [$T_{1/2}$].
 1973Ho43 – K. R. Hogstrom, B. W. Mayes, L. Y. Lee, J. C. Allred, C. Goodman, G. S. Mutchler, C. R. Fletcher, G. C. Phillips, Nucl. Phys. A215 (1973) 598 [$T_{1/2}$].
 1973SiYS – J. Singh, Proc. Nucl. Phys. and Solid State Phys. Symp. 15B (1972) 1 [$T_{1/2}$].
 1973Fi13 – M. L. Fitzpatrick, K. W. D. Ledingham, J. Y. Gourlay, J. G. Lynch, J. Phys. A6 (1973) 713 [K/β^+ ratio, end-point energy].
 1975Az01 – G. Azuelos, J. E. Kitching, K. Ramavataram, Phys. Rev. C12 (1975) 563 [$T_{1/2}$, end-point energy].
 1975Aj01 – F. Ajzenberg-Selove, Nucl. Phys. A248 (1975) 1 [$T_{1/2}$, end point energy, Q, log ft].
 1975Be28 – H. Behrens, M. Kobelt, L. Szybisz, W. G. Thies, Nucl. Phys. A246 (1975) 317 [$T_{1/2}$, end-point energy].
 1977Az01 – G. Azuelos, J. E. Kitching, K. Ramavataram, Phys. Rev. C15 (1977) 1847 [$T_{1/2}$].
 1977Ba49 – W. Bambynek, H. Behrens, M. H. Chen, B. Crasemann, M. L. Fitzpatrick, K. W. D. Ledingham, H. Genz, M. Muttere, R. L. Intemann, Revs. Modern Phys. 49 (1977) 77 [Electron Capture].
 1978Ra21 – S. Raman, C. A. Houser, T. A. Walkiewicz, I. S. Towner, Atomic Data and Nucl. Data Tables 21 (1978) 567 [$T_{1/2}$, end point energy, Q, log ft].
 1980Aj01 – F. Ajzenberg-Selove, Nucl. Phys. A336 (1980) 1 [$T_{1/2}$, Q, log ft].
 1985Aj01 – F. Ajzenberg-Selove, Nucl. Phys. A433 (1985) 1 [$T_{1/2}$, Q, log ft].
 1990Aj01 – F. Ajzenberg-Selove, Nucl. Phys. A506 (1990) 1 [$T_{1/2}$, Q, log ft, end-point energy].
 1995Au04 – G. Audi, A.H. Wapstra, Nucl. Phys. A595 (1995) 409 [Q].
 2000Co21 – Codata Group, Revs. Modern Phys. 72 (2000) 351 [m_0c].

¹³N – Comments on evaluation of decay data by V. Chisté and M. M. Bé

1) Decay Scheme

¹³N disintegrates by β^+ emission (99,818 (13) %) and electron capture (0,182 (13) %) to the ground state of the stable nuclide ¹³C.

2) Nuclear Data

The Q value (2220,44 (27) keV) is from the evaluation of Audi and Wapstra (1995Au04), and has been calculated using the formula:

$$Q = M(A, Z) - M(A, Z - 1),$$

where M(A,Z) and M(A,Z-1) are the measured atomic masses of ¹³N and ¹³C, respectively.

The E $_{\beta^+}$ deduced from this Q value (E $_{\beta^+}$ = 1198,45 (27) keV) agrees with the weighted average value of 1199,00 (36) keV, deduced from measured values (see § **b⁺ Transition and Electron Capture Transition**).

The measured ¹³N half-life values (in minutes) are given below:

T_{1/2}

Reference	Value (min)
Ward (1939Wa35)	9,93 (3)
Siegbahn (1945Si02)	10,13 (10)
Cook (1948Co05)	10,2 (1)
Churchill(1953Ch34)	10,048 (32)
Wilkinson (1955Wi43)	10,08 (4)
Daniel (1957Da07)	9,960 (30)
Deineko (1957De22)	10,02 (10)
Norbeck (1957No17)	10,07 (6)
Arnell (1958Ar15)	9,960 (30)
King (1960Ki02)	9,93 (5)
Janecke (1960Ja12)	9,965 (5)
Ebrey (1965Eb03)	9,96 (2)
Bormann (1965Bo42)	10,05 (5)
Ritchie (1968Ri15)	9,963 (9)
Singh (1973SiYS)	10,0 (5)
Azuelos (1977Az01)	9,965(10)
Katoh (1989Ka08)	9,962 (20)

The weighted average has been calculated using the Lweight computer program (version 3).

The Siegbahn (1945Si02) and Cook (1948Co05) values have been shown to be outliers by the Lweight program, based on the Chauvenet's criterion. For the remaining 15 statistically consistent values, the largest contribution to the weighted average comes from the value of Janecke (1960Ja12), with statistical weight of 54 %. The reduced- χ^2 is 1,65.

The adopted value is the weighted average : 9,9670 min, with an uncertainty of 0,0037min.

2.1) β^+ Transition and Electron capture transition.

The β^+ and electron capture probabilities shown in Tables 2.1 and 2.2, respectively, have been deduced by using a K/β^+ ratio of $(1,68 \pm 0,12) \cdot 10^{-3}$ measured by Ledingham (1963Le06) and, normalizing to a total probability ($P_{\beta^+} + P_{EC}$) of 100%. This experimental K/β^+ ratio is close to the following theoretical values:

- a) $1,864 \cdot 10^{-3}$ calculated with LOGFT program;
- b) $1,939 \cdot 10^{-3}$ calculated by Fitzpatrick (1973Fi13);
- c) $1,800 \cdot 10^{-3}$ given by Bambynek (1977Ba49);
- d) $1,78 \cdot 10^{-3}$ given by Ledingham (1963Le06).

The uncertainties were estimated by standard error-propagation techniques.

The $lg ft$ value for β^+ transition (3,654) has been calculated with the program LOGFT for an allowed transition. This value agrees with 3,637 suggested by Ajzenberg -Selove (1981Aj01, 1986Aj01 and 1991Aj01).

The partial sub shell capture probabilities P_K and P_L were calculated for an allowed transition using the computer program EC-Capture.

A weighted average (1199,0(4) keV) of the β^+ end-point energy has been deduced (using the Lweight computer program, version 3) from the following measured values (in keV):

Reference	Values (keV)
Hornyak (1950Ho01)	1202 (5)
Grabowsky (1954Gr03)	1185 (25)
Daniel (1957Da07)	1190 (3)
Fitzpatrick (1973Fi13)	1198,5(9)
Raman (1978Ra21)	1198,7 (4)

The largest contribution (with an statistical weight of 81%) to the weighted average of these 5 values comes from the value of Raman (1978Ra21). The weighted average is 1199,00 keV, with an internal uncertainty of 0,36 and a reduced χ^2 of 2,2. This value agrees with E_{β^+} (1198,45(27) keV), which was deduced from the adopted Q value (1995Au04) in this evaluation.

3) Gamma-ray Emissions

The annihilation radiation emission intensity ($I_{\gamma 511}$) is $P_{\beta^+} (= 99,818 (13))$, multiplied by 2, without the correction factor for the annihilation-in-flight processus in the medium. That is, $I_{\gamma 511} = 199,636 (26) \%$.

4) Atomic Data

Atomic K-fluorescence yield (ω_K) is from Bambynek (1984Ba01).

References

- 1939Wa35 – W. Ward, Proc. Cambridge Phil. Soc. 35 (1939) 523 [$T_{1/2}$].
- 1945Si02 – S. Siegbahn, Arkiv F. Art. Mat. Fys. 32A (1945) 9 [$T_{1/2}$].
- 1948Co05 – C. S. Cook, L. M. Langer, C. Price Jr., M. B. Sampson, Phys. Rev. 74 (1948) 502 [$T_{1/2}$].
- 1950Ho01 – B. Hornyak, Phys. Rev. 77 (1950) 160 [end-point energy].
- 1952Aj01 – F. Ajzenberg, Rev. Mod. Phys. 24 (1952) 321 [$T_{1/2}$, end-point energy, log ft].
- 1953Ch34 – J. L. W. Churchill, W. M. Jones, S. E. Hunt, Nature 172 (1953) 460 [$T_{1/2}$].
- 1954Gr03 – N. Grabowsky, Bull. Acad. Pol. Sci. 2 (1954) 379 [end-point energy].
- 1955Wi43 – D. H. Wilkinson, Phys. Rev. 100 (1955) 32 [$T_{1/2}$].
- 1955Aj01 – F. Ajzenberg, Rev. Mod. Phys. 27 (1955) 27 [$T_{1/2}$, end-point energy, log ft].
- 1957No17 – E. Norbeck Jr., C. S. Littlejohn, Phys. Rev. 108 (1957) 754 [$T_{1/2}$].
- 1957De22 – S. Deineko, A. Ia. Taranov, A. K. Valter, Sov. Phys. - JETP 5 (1957) 201 [$T_{1/2}$].
- 1957Da07 – H. Daniel, U. Schmidt-Rohr, Z. Naturforsch 12A (1957) 750 [$T_{1/2}$].
- 1958Ar15 – S. E. Arnell, J. Dubois, O. Almen, Nucl. Phys. 6 (1958) 196 [$T_{1/2}$].
- 1958St30 – D. Strominger, J. L. Hollander, G. T. Seaborg, Rev. Mod. Phys. 30 (1958) 585 [$T_{1/2}$, end point energy, Q, log ft].
- 1960Ki02 – J. D. King, R. N. H. Haslam, R. W. Parsons, Can. J. Phys. 38 (1960) 231 [$T_{1/2}$].
- 1960Ja12 – V. J. Janecke, Z. F. Naturf. 15A (1960) 593 [$T_{1/2}$].
- 1963Le06 – K. W. D. Ledingham, J. A. Payne, R. W. P. Drever, Proc. Int. Conf. Role of Atomic Electrons in Nuclear Transformations 2 (1963) 359 [K/ β^+ ratio].
- 1965Eb03 – T. G. Ebrey, P. R. Gray, Nucl. Phys. 61 (1965) 479 [$T_{1/2}$].
- 1965Bo42 – M. Bormann, E. Fretwurst, P. Schehka, G. Wrege, Nucl. Phys. 63 (1965) 438 [$T_{1/2}$].
- 1968Va23 – E. Vatai, Proc. Conf. Electron Capture and higher order processes in Nucl. Decay 2 (1968) 71 [K/ β^+ ratio].
- 1968Ri15 – A. I. M. Ritchie, Nucl. Inst. Meth. 64 (1968) 181 [$T_{1/2}$].
- 1973SiYS – J. Singh, Proc. Nucl. Phys. and Solid State Phys. Symp. 15B (1972) 1 [$T_{1/2}$].
- 1973Fi13 – M. L. Fitzpatrick, K. W. D. Ledingham, J. Y. Gourlay, J. G. Lynch, J. Phys. A6 (1973) 713 [K/ β^+ ratio].
- 1976Aj01 – F. Ajzenberg-Selove, Nucl. Phys. A268 (1976) 1 [$T_{1/2}$, Q, log ft].
- 1977Az01 – G. Azuelos, J. E. Kitching, K. Ramavataram, Phys. Rev. C15 (1977) 1847 [$T_{1/2}$].
- 1977Ba49 – W. Bambynek, H. Behrens, M. H. Chen, B. Crasemann, M. L. Fitzpatrick, K. W. D. Ledingham, H. Genz, M. Muttere, R. L. Intemann, Revs. Modern Phys. 49 (1977) 77 [Electron Capture].
- 1978Ra21 – S. Raman, C. A. Houser, T. A. Walkiewicz, I. S. Towner, Atomic Data and Nucl. Data Tables 21 (1978) 567 [$T_{1/2}$, end point energy, Q, log ft].
- 1981Aj01 – F. Ajzenberg-Selove, Nucl. Phys. A360 (1981) 1 [$T_{1/2}$, Q, log ft].
- 1984Ba01 – W. Bambynek, Proc. X-ray and Inner-Shell Processes in Atoms, Molecules and Solids, Leipzig, Aug. 20-23, 1984. Edited by A. Meisel [Atomic Data]
- 1986Aj01 – F. Ajzenberg-Selove, Nucl. Phys. A449 (1986) 1 [$T_{1/2}$, Q, log ft].
- 1989Ka08 – T. Katoh, K. Kawade, H. Yamamoto, JAERI-M-089-083 (1989) [$T_{1/2}$]
- 1991Aj01 – F. Ajzenberg-Selove, Nucl. Phys. A523 (1991) 1 [$T_{1/2}$, Q, log ft].
- 1995Au04 – G. Audi, A.H. Wapstra, Nucl. Phys. A595 (1995) 409 [Q].
- 2000Co21 – Codata Group, Revs. Modern Phys. 72 (2000) 351 [m_0c].

**¹⁴C - Comments on evaluation of decay data
by M.M. Bé and V.P. Chechev**

This evaluation was completed in 1998, it was updated in January 2012 to include the most recent Q(β^-) update. The literature available by this date was included.

Nuclear Data

Half-life

In literature there are many measurements of the ¹⁴C half-life dating from 1946 to 1954 (Table 1). Mann *et al.* (1961) discussed the problem of spread of these measurement results from 4 700 to 7 200 years. They connect the divergence with very low enrichment of ¹⁴C (a few percentages) and a large systematic uncertainty arose from retention of a small quantity of carbon dioxide with a high specific activity during a gas dilution phase. Therefore, following Holden (1990Ho28) who evaluated the ¹⁴C half-life in 1990, we have omitted the measurement results before 1961 and considered only later measurements (Table 2). In all the latter works the number of ¹⁴C atoms has been determined by the mass-spectrometric method and the counting rate was measured by different methods as shown in Table 1.

Table 1: Results of ¹⁴C half-life measurements

NSR keynumber	Half-life of ¹⁴ C, years	Method
1946Re10	4 700 (400)	SA: GM; MS
1948No02	5 100 (200)	- " -
1948Ya02	7 200 (500)	- " -
1949Ha52	6 360 (200)	- " -
1949Jo07	5 589 (75)	- " -
1950En59	5 580 (90)	- " -
1950Mi10	6 360 (190)	- " -
1950Mi10	5 513 (165)	SA: PC ; MS
1951Ma30	5 370 (200)	SA: IC ; MS
1952Je11	6 030	SA: GM ; gas density
1954Ca41	5 900 (250)	SA: Cal ; gas density
1961Ma32	5 760 (50)	SA: PC ; MS
1961Wa16	5 780 (65)	SA: PC ; MS
1962O114	5 680 (40)	SA: PC ; MS
1964Hu09	5 745 (50)	SA: PC ; MS. 1961Ma32 value revised
1968Be47	5 660 (30)	SA: PC(GM) ; MS
1968Re13 +	5 736 (56)	SA: LS ; MS
1972Em01		

Usual designations:

SA - method of radionuclide specific activity determination, by mean of Geiger-Müller counter (GM), proportional counter (PC), calorimeter (Cal), ionization chamber (IC) or liquid scintillation counter (LS);
MS - determination of the number of atoms by the mass-spectrometric method.

Table 2: Selected measurement results and recommended value of ¹⁴C half-life

Year	Half-life of ¹⁴ C, <i>a</i>	Reference NSR keynumber
1961	5 780 (65)	1961Wa16
1962	5 680 (40)	1962Ol14
1964	5 745 (50)	1964Hu09
1968	5 660 (30)	1968Be47
1968	5 736 (56)	1968Re13/1972Em01
$\chi^2/n-1 = 1,2$; critical $\chi^2 = 3,3$		
Weighted average	5 697 (21) <i>a</i>	
Unweighted average	5 720 (22) <i>a</i>	
Recommended value	5 700 (30) <i>a</i>	

The adopted value of the ¹⁴C half-life is the weighted average of the five results listed in Table 2. Since they were all obtained by the same method of the specific activity measurement, the final uncertainty is taken as the lowest experimental uncertainty of the data set.

It should be noticed that Holden gave a similar evaluation of ¹⁴C, $T_{1/2}$ ($5\,715 \pm 30$ years), but he adopted the unweighted average of the same measurement results with addition to them of the average of three values obtained in 1949-1950.

From an analysis of fossil corals whose ages were determined via ²³⁴Th/²³⁴U/²³⁸U dating, a ¹⁴C half-life of “6 030” *a* should be expected (2007Ch**). A re-determination of the ¹⁴C half-life is required to improve radiocarbon-based researches.

Decay Energy and Characteristics of Electron Emission (β^-)

The ¹⁴C beta decay to the ground state level of ¹⁴Ni is expected to be allowed ($0^+ \rightarrow 1^+$). However it has been shown deviations in the shape of the ¹⁴C beta spectrum (2000Ku25, 1995Wi20). A summary of measured and predicted spectra is given in 2000Ku25.

The maximum energy of the β spectrum was deduced from the results of measurements, as listed below.

Table 3: Measured β end-point energy, E_0 .

Reference	E_0 (keV)	u_c	Remarks
Cook (1948Co10)	156,3	10	
Forster (1954Fo*)	155	1	
Smith (1975Sm02)	156,476	0,005	rf mass spectrometer
Sur (1991Su09)	155,74	0,08	¹⁴ C-doped Ge detector, taking into account anomalies in the β spectrum
Wietfeldt (1995Wi20)	155,95	0,22	¹⁴ C-doped Ge detector, taking into account anomalies in the β spectrum
Kuzminov (2000Ku25)	156,27	0,14	Wall-less proportional counter, taking into account anomalies in the β spectrum

It is noteworthy that the value reported by Smith (1975Sm02) is much more precise but also discrepant with the other results obtained by different methods.

The set of the four most precise values is discrepant with a $\chi^2/n-1 = 17$. Then the uncertainty of the Smith's value has been increased to 0,066 in order to reduce its weight to 50 %. The resulting weighted average with an expanded uncertainty to cover the most precise result is: 156,18 (30) keV.

This value is considerably less precise than the recommended value of 156,476 (4) keV given in Audi *et al.* (2003Au03).

On one hand, the weighted mean is only limited to values following ¹⁴C β⁻ decay and one value that comes from a direct mass-difference measurements using the rf technique; when the value recommended by Audi *et al.* (2003Au03/2011AuZZ) is deduced from the mass differences between ¹⁴C and ¹⁴N, determined using a robust least-squares procedure.

On the other hand, in that case the whole "robust least-squares procedure" in 2003Au03/2011AuZZ is dominated by the single ultra-precise mass-spectrometric value. And this exact ¹⁴C - ¹⁴N mass difference affects other masses, and not *vice versa*.

In this evaluation we will accept the Audi *et al.* recommendation, while following the Wietfeld's conclusion (1995Wi20): "We feel there is a significant problem in the ¹⁴C Q value and we hope that this will be resolved by future experiments".

The average energy per disintegration has been calculated, expecting an allowed form of β⁻-spectrum, by using the program BetaShape (2012Mo**) which includes the calculations of "exchange effects".

E _{max} (keV)	E _{mean} (keV)
156,18 (30)	49,1 (3)
156,476 (4)	49,16 (1)

References

- 1946Re10** A.F. Reid, J.R. Dunning. Phys.Rev. 70, (1946) 431 [T_{1/2}]
- 1948Co10** C.S. Cook, L.M. Langer, H.C. Price, Jr. Phys.Rev. 74 (1948) 548 [Q]
- 1948No02** L.D. Norris, M.G. Inghram. Phys.Rev. 73 (1948) 350 [T_{1/2}]
- 1948Ya02** L. Yaffe, J.M. Grunlund. Phys.Rev. 74 (1948) 696 [T_{1/2}]
- 1949Ha52** R.C. Hawkings, R.F. Hunter, W.B. Mann, W.H. Stephens. Can.J.Res. 27B (1949) 545 [T_{1/2}]
- 1949Jo07** W.M. Jones. Phys.Rev. 76 (1949) 885 [T_{1/2}]
- 1950En59** A.G. Engelkemeir, W.F. Libby, Rev.Sci.Inst. 21 (1950) 550 [T_{1/2}]
- 1950Mi10** W.W. Miller, R. Ballentine, W. Bernstein, L. Friedman, A.O.Nier, R.D.Evans. Phys.Rev. 77 (1950) 714 [T_{1/2}]
- 1951Ma30** G.G. Manov, L.F. Curtiss. J.Research Natl.Bur.Standards 46 (1951) 328 [T_{1/2}]
- 1952Je11** G.H. Jenks, H. Sweeton. Phys. Rev. 86 (1952) 803 [T_{1/2}]
- 1954Ca41** R.S. Caswell, J.M. Brabant, A. Schwebel. J.Res.Natl.Bur.Stand. 53 (1954) 27 [T_{1/2}]
- 1954Fo**** H.H. Forster, A. Oswald. Phys. Rev. 96,4 (1954) 1030 [Q]
- 1961Ma32** W.M. Mann, W.F. Marlow, E.E. Hughes, Appl. Rad.Isotopes 11 (1961) 57 [T_{1/2}]
- 1961Wa16** D.E. Watt, D. Ramsden, H.W. Wilson, Appl. Rad.Isotopes 11 (1961) 68 [T_{1/2}]
- 1962Ol04** I.V. Olsson, I. Karlen, A.H. Turnbull, N.J.D. Prosser, Ark.f.Fys. 22 (1962) 237 [T_{1/2}]
- 1964Hu09** E.E. Hughes, W.M. Mann, Appl. Rad.Isotopes 15 (1964) 97 [T_{1/2}]
- 1968Be47** F. Bella, M. Alessio, P. Fratelli, Nuovo Cim. 58B (1968) 232 [T_{1/2}]
- 1968Re22** S.A. Reynolds, ORNL-4343, p 76 (1968) [T_{1/2}]
- 1972Em01** J.F. Emery, S.A. Reynolds, E.I. Wyatt, G.I. Gleason. Nucl. Sci. Eng. 48 (1972)319 [T_{1/2}]
- 1975Sm02** L.G. Smith, A.H. Wapstra. Phys. Rev. C11, 4 (1975) 1392 [Q]
- 1990Ho28** N.E. Holden, Pure and Appl. Chem. 62 (1990) 941 [T_{1/2}]
- 1991Su09** B. Sur, *et al.* Phys. Rev. Lett. 66, 19 (1991) 2444 [Q]
- 1995Wi20** F.E. Wietfeldt, *et al.* Phys. Rev. C52, 2 (1995) 1028 [Q]
- 2000Ku25** V.V. Kuzminov, N.Ja. Osetrova. Phys. Atomic Nuclei 63, 7 (2000) 1292 [Q]
- 2003Au03** G. Audi, A.H. Wapstra, C. Thibault. Nucl. Phys. A729 (2003) 129 [Q]
- 2007Ch**** Tzu-Chien Chiu, *et al.* Quaternary Science Reviews 26 (2007) 18 [T_{1/2}]
- 2011AuZZ** G. Audi, W. Meng, <http://amdc.in2p3.fr/masstable/Ame2011int/mass.mas114> [Q_{β⁻}, Q_ε]
- 2012Mo**** X. Mougeot, M.-M. Bé, C. Bisch, M. Loidl. Evidence for the exchange effect in the beta decay of ²⁴¹Pu. Physical Review A86 (2012) 042506 [<Eβ>]

¹⁵O – Comments on evaluation of decay data by V. Chisté and M. M. Bé

1) Decay Scheme

¹⁵O disintegrates by β^+ emission (99,885 (6) %) and electron capture (0,115 (6) %) to the ground state of the stable nuclide ¹⁵N.

2) Nuclear Data

The Q value has been calculated using the formula:

$$Q = E_{\beta^+} + 2m_0c^2 = 2757,0 (13) \text{ keV}$$

where $E_{\beta^+} = 1735,0 (13) \text{ keV}$ is the weighted mean of the β^+ end-point energy (see **b⁺ Transition and Electron Capture**) and, $2m_0c^2 = 1021,9978 (42) \text{ keV}$ (2000Co21). The Q value calculated here is in agreement with the value of 2754,0 (5) from the Audi and Wapstra evaluation (1995Au04), which takes into account only Raman's value (1978Ra21, 1731,9 (7) keV) to determine the recommended Q value.

The measured ¹⁵O half-life values are, in seconds:

T_{1/2}

Reference	Value (sec)
McMillan (1935Mc02)	126 (5)
Brown (1950Br29)	118,0 (6)
Kline (1954Kl36)	123,4 (13)
Bashkin(1955Ba83)	121 (3)
Kistner (1957Ki22)	122 (5)
Penning (1957Pe12)	123,95 (50)
Kistner (1959Ki99)	124,1 (5)
Janecke (1960Ja12)	122,1 (1)
Nelson (1963Ne05)	122,6 (10)
Csikai (1963Cs02)	125 (2)
Vasil'ev (1963Va23)	114 (12)
Azuelos (1977Az01)	122,23(23)

The half-life weighted average has been calculated by the Lweight program (version 3).

The weighted average of all 12 values is 122,16 s with an internal uncertainty of 0,09 and a reduced $-\chi^2$ of 7,3. The value of 1960Ja12 has a relative weight of 76% and that of 1950Br29 contributes 4,4 to the reduced- χ^2 .

The evaluator has chosen to reject the McMillan (1935Mc02) and Csikai (1963Cs02), because they are far from the other values and with large uncertainties.

The Brown (1950Br29) and Vasil'ev (1963Va23) values have been rejected by the Lweight program, based on the Chauvenet's criterion. For the remaining 8 values, the largest contribution to the weighted average comes from the value of Janecke (1960Ja12), amounting to a statistical weight of 78% (reduced -

$\chi^2 = 4,01$). The program Lweight 3 has increased the uncertainty of the 1960Ja12 value from 0,1 to 0,186 in order to reduce its relative weight from 78% to 50%.

The adopted value is the weighted mean : $122,40$ s, with an uncertainty of $0,33$; or $2,041$ (6) min. The reduced- χ^2 is 3,2.

2.1) β^+ Transition and Electron capture

The β^+ and electron capture probabilities have been calculated taking into account a K/β^+ ratio of $(1,07 \pm 0,06) \cdot 10^{-3}$ measured by Leiper (1972Le06) and, normalizing to a total probability ($P_{\beta^+} + P_{EC}$) of 100%. The experimental K/β^+ ratio is close of its theoretical value ($= 0,99(1) \cdot 10^{-3}$) calculated with the LOGFT program. The uncertainties were calculated through their propagation on the above formulas.

The value of $\log ft$ of the β^+ transition (3,6) has been calculated with the program LOGFT for an allowed transition, in agreement with the value suggested by Ajzenberg -Selove, which is 3,637 (1981Aj01, 1986Aj01 and 1991Aj01).

The partial sub shell capture probabilities were calculated with the program EC-Capture for an allowed transition.

The weighted mean of the β^+ end-point energy has been calculated (with the Lweight program, version 3) using the following measured values (in MeV):

Reference	Values (MeV)
Fowler (1936Fo16)	1,7 (2)
Stephens (1937St03)	1,56 (20)
Perez-Mendez (1949Pe23), Brown (1950Br29)	1,683 (5)
Kington (1955Ki39)	1,735 (8)
Kistner (1957Ki22) (solid target)	1,723 (5)
Kistner (1957Ki22) (gaseous target)	1,736 (10)
Kistner (1959Ki99)	1,739 (2)
Raman (1978Ra21)	1,7319 (7)

The values given by Fowler (1936Fo16), Stephens (1937St03), Perez -Mendez (1949Pe23) and Kistner (1957Ki22 – solid target) were shown (by the Lweight program) to be statistically inconsistent with the other values (based on the Chauvenet's criterion), thus the evaluators rejected those 4 values. The largest contribution to the weighted average of the 4 remaining values comes from the value of Raman (1978Ra21), amounting to a statistical weight of 88% (reduced $\chi^2 = 3,8$). The program Lweight 3 has increased the uncertainty of the 1978Ra21 value from 0,0007 to 0,0019 in order to reduce its relative weight from 88% to 50%.

The adopted value is the weighted mean : $1735,0$ keV, with an external uncertainty of $1,3$ and a reduced- χ^2 of 2,2.

3) Gamma Emissions

The annihilation radiation emission probability ($I_{\gamma}(511)$), is P_{β^+} , or 99,885(6), multiplied by 2, without the correction factor for the annihilation-in-flight in the medium, that is $I_{\gamma}(511) = 199,770(12)\%$

4) Atomic Data

Atomic value (ω_K) is from Bambynek (1984Ba01).

References

- 1935Mc02 - E. McMillan, M. S. Livingston, Phys. Rev. 47 (1935) 452 [$T_{1/2}$].
- 1936Fo16 - W. A. Fowler, L. A. Delsasso, C. C. Lauritsen, Phys. Rev. 49 (1936) 561 [End-point energy].
- 1937St03 - W. E. Stephens, K. Djanab, T. W. Bonner, Phys. Rev. 52 (1937) 1079 [End-point energy].
- 1949Pe23 - V. Perez-Mendez, H. Brown, Phys. Rev. 76 (1949) 689 [End-point energy, ft value].
- 1950Br29 - H. Brown, V. P. Mendez, Phys. Rev. 78 (1950) 649 [$T_{1/2}$, end point energy].
- 1954Kl136 - R. M. Kline, D. J. Zaffarano, Phys. Rev. 96 (1954) 1620 [$T_{1/2}$].
- 1955Ba83 - S. Bashkin, R. R. Carlson, E. B. Nelson, Phys. Rev. 99 (1955) 107 [$T_{1/2}$].
- 1955Ki39 - J. D. Kington, J. K. Bair, H. O. Cohn, H. B. Willard, Phys. Rev. 99 (1955) 1393 [End-point energy].
- 1957Pe12 - J. R. Penning, F. H. Schmidt, Phys. Rev. 105 (1957) 647 [$T_{1/2}$].
- 1957Ki22 - O. C. Kistner, A. Schawarschild, B. M. Rustad, Phys. Rev. 105 (1957) 1339 [$T_{1/2}$, log ft, end-point energy].
- 1959Ki99 - O. C. Kistner, B. M. Rustad, Phys. Rev. 114 (1959) 1329 [$T_{1/2}$, end point energy].
- 1959Aj01 - F. Ajzenberg-Selove, T. Lauritsen, Nucl. Phys. 11 (1959) 1 [$T_{1/2}$, end point energy, Q, log ft].
- 1960Ja12 - V. J. Janecke, Z. F. Naturf. 15A (1960) 593 [$T_{1/2}$].
- 1963Ne05 - J. W. Nelson, E. B. Carter, G. E. Mitchel, R. H. Davis, Phys. Rev. 129 (1963) 1723 [$T_{1/2}$].
- 1963Cs02 - J. Csikai, G. Peto, Phys. Lett. 4 (1963) 252 [$T_{1/2}$].
- 1963Va23 - S. S. Vasil'ev, L. Ya. Shavtvalov, Bull. Acad. Sci. USSR 27 (1963) 1239 [$T_{1/2}$].
- 1970Aj01 - F. Ajzenberg-Selove, Nucl. Phys. A152 (1970) 1 [$T_{1/2}$, end point energy, Q, log ft].
- 1972Le06 - W. Leiper, R. W. P. Drever, Phys. Rev. C6 (1972) 1132 [K/β^+ ratio].
- 1973Fi13 - M. L. Fitzpatrick, K. W. D. Ledingham, J. Y. Gourlay, J. G. Lynch, J. Phys. A6 (1973) 713 [K/β^+ ratio].
- 1976Aj01 - F. Ajzenberg-Selove, Nucl. Phys. A268 (1976) 1 [$T_{1/2}$, Q, log ft].
- 1977Az01 - G. Azuelos, J. E. Kitching, K. Ramavataram, Phys. Rev. C15 (1977) 1847 [$T_{1/2}$].
- 1977Ba49 - W. Bambynek, H. Behrens, M. H. Chen, B. Crasemann, M. L. Fitzpatrick, K. W. D. Ledingham, H. Genz, M. Muttere, R. L. Intemann, Revs. Modern Phys. 49 (1977) 77 [Electron Capture].
- 1978Ra21 - S. Raman, C. A. Houser, T. A. Walkiewicz, I. S. Towner, Atomic Data and Nucl. Data Tables 21 (1978) 567 [$T_{1/2}$, end point energy, Q, log ft].
- 1981Aj01 - F. Ajzenberg-Selove, Nucl. Phys. A360 (1981) 1 [$T_{1/2}$, Q, log ft].
- 1984Ba01 - W. Bambynek, Proc. X-ray and Inner-Shell Processes in Atoms, Molecules and Solids, Leipzig, Aug. 20-23, 1984. Edited by A. Meisel [Atomic Data].
- 1986Aj01 - F. Ajzenberg-Selove, Nucl. Phys. A449 (1986) 1 [$T_{1/2}$, Q, log ft].
- 1991Aj01 - F. Ajzenberg-Selove, Nucl. Phys. A523 (1991) 1 [$T_{1/2}$, Q, log ft].
- 1995Au04 - G. Audi, A.H. Wapstra, Nucl. Phys. A595 (1995) 409 [Q].
- 2000Co21 - Codata Group, Revs. Modern Phys. 72 (2000) 351 [m_0c^2].

¹⁸F – Comments on evaluation of decay data by V. Chisté and M.M. Bé

1) Decay Scheme

¹⁸F disintegrates by β^+ emission (96.86(16)%) and electron capture (3.14(16)%) to the ground state of the stable nuclide ¹⁸O.

2) Nuclear Data

The Q value (1655.5 (6) keV) is from Audi and Wapstra (1995Au04), and has been calculated with the formula:

$$Q = M(A, Z) - M(A, Z - 1),$$

where M(A,Z) and M(A,Z-1) are the measured atomic masses of ¹⁸F and ¹⁸O, respectively.

E_{β^+} , calculated from this Q value ($E_{\beta^+} = 633.5(6)$ keV), is in agreement with a weighted average value of 633.2(3) keV, which was deduced from measured values (see **b⁺ Transition and Electron Capture Transition**).

The measured ¹⁸F half-life values (in minutes) are given below:

Reference	Value (min)
Snell (1937Sn14)	112 (4)
DuBridge (1938Br47)	107 (4)
Krishnan (1941Kr12)	112 (2)
Huber (1943Hu33)	115 (4)
Blaser (1949Bl30)	112 (1)
Jarmie (1955Ja12)	111 (1)
Bendel (1958Be08)	109.8 (12)
Markowitz (1958Ma12)	112 (1)
Carlson (1959Ca63)	109.70 (54)
Yule (1960Yu15)	110,2 (2)
Rayburn (1961Ra53)	111.0 (22)
Mahony (1962Ma15)	109.74 (21)
Beg (1963Be31)	109.6 (6)
Hofmann (1964Ho09)	110.5 (6)
Mahony (1964Ma07)	109.72 (6)
Ebrey (1965Eb02)	109.87 (12)
Bormann (1965Bo38)	111 (2)
Kavanagh (1969Ka17)	109.87 (12)
Hogstrom (1973Ho21)	95 (7)
Rutledge (1980Ru02)	109.71 (2)
Katoh (1989Ka01)	109.48 (8)
Schrader (2004Sc00)	109.748(21)

The only outliers values are 107 (4) min (1938Br47), 115 (4) min (1943Hu33) and 95 (7) min (1973Ho21), which contributed with a statistical weight of just $0.378 \cdot 10^{-5} \%$ (1973Ho21) to $0.116 \cdot 10^{-4} \%$ (1938Br47 and 1943Hu33) to the weighted average. Our recommended half-life is the weighted average of 109.728 (19) min, or 1.8288 (3) h ($\chi^2/\nu = 1.98$).

β^+ Transition and Electron capture transition

The β^+ and electron capture probabilities shown in Tables 2.1 and 2.2, respectively, have been deduced using a K/β^+ ratio of $(3.00 \pm 0.18) \cdot 10^{-2}$ measured by Drever (1956Dr02), $P_K/P_{EC} = 0.9267$ (48) (see Section 2.2) and, normalizing to a total probability ($P_{\beta^+} + P_{EC}$) of 100 %. This leads to $P_{\beta^+} = 96.86(19) \%$ and $P_{EC} = 3.14(19) \%$, respectively. The uncertainties were calculated through their propagation on the above formulas.

The experimental K/β^+ ratio of Drever is close to the theoretical values:

- a) $3.19 \cdot 10^{-2}$ calculated with LOGFT program;
- e) $3.31 \cdot 10^{-2}$ given by Fitzpatrick (1973Fi13);
- f) $3.14 \cdot 10^{-2}$ given by Bambynek (1977Ba49);

Using the LOGFT program evaluators calculated a $\lg ft$ of 3.57 for this allowed transition. This value agrees with 3.554 suggested by Ajzenberg-Selove (1972Aj01, 1978Aj01 and 1987Aj01).

The partial sub shell capture probabilities given in Section 2.2 were calculated using the program EC - Capture for an allowed transition.

The weighted mean of the β^+ end-point energy has been calculated (with the Lweight program, version 3) using the following measured values (in keV):

Reference	Values (keV)
Blaser (1949Bl30)	635 (15)
Ruby (1951Ru40)	649 (9)
Hofmann (1964Ho09)	635 (2)
Alburger (1970Al17)	632.9 (7)
Fitzpatrick (1973Fi13)	633.3 (3)

The weighted average of these 5 values is 633.2 keV with an internal uncertainty of 0.3 keV and a reduced χ^2 of 1.4. This value is in agreement with E_{β^+} (633.5 (6) keV) deduced from the adopted Q value (1995Au04) in this evaluation.

3) Gamma-ray Emissions

The annihilation radiation emission intensity ($I_{\gamma 511}$) is P_{β^+} (=96.86(19) %), multiplied by 2, without the correction factor for the annihilation-in-flight process in the medium. That is, $I_{\gamma 511} = 193.72(27) \%$.

References

- 1937Sn14 – A. H. Snell, Phys. Rev. 51 (1937) 143 [$T_{1/2}$].
- 1938Br47 – L. A. DuBridge, S. W. Barnes, J. H. Buck, C. V. Strain, Phys. Rev. 53 (1938) 447 [$T_{1/2}$].
- 1941Kr12 – R. S. Krishnan, Nature (London) 148 (1941) 407 [$T_{1/2}$].
- 1943Hu33 – O. Huber, O. Lienhard, P. Scherrer, H. Waffler, Helv. Phys. Acta 16 (1943) 33 [$T_{1/2}$].
- 1949Bl30 – J. P. Blaser, F. Boehm, P. Marmier, Phys. Rev. 75 (1949) 1953 [End-point energy, $T_{1/2}$].
- 1951Ru40 – L. Ruby, J. R. Richardson, Phys. Rev. 83 (1951) 698 [End-point energy].
- 1955Ja12 – N. Jarmie, Phys. Rev. 98 (1955) 41 [$T_{1/2}$].
- 1956Dr02 – R. W. P. Drever, A. Moljk, J. Scobie, Phil. Mag. 1 (1956) 942 [K/β^+ ratio].
- 1958Be08 – W. L. Bendel, J. McElhinney, R. A. Tobin, Phys. Rev. 111 (1958) 1297 [$T_{1/2}$].

- 1958Ko12 – J. Konijn, B. Van. Nooijen, H. L. Hagedoorn, A. H. Waspra, Nucl. Phys. 9 (1958) 296 [K/ β^+ ratio].
- 1958Ma12 – S. S. Markowitz, F. S. Rowland, Phys. Rev. 112 (1958) 1295 [$T_{1/2}$].
- 1959Aj01 – F. Ajzenberg-Selove, Nucl. Phys. 11 (1959) 1 [$T_{1/2}$, end point energy, Q, log ft].
- 1959Ca63 – C. H. Carlson, L. Singer, D. H. Service, W. D. Armstrong, Int. J. Appl. Rad. Isotopes 4 (1959) 210 [$T_{1/2}$].
- 1960Yu15 – H. P. Yule, A. Turkevich, Phys. Rev. 118 (1960) 1591 [$T_{1/2}$].
- 1961Ra53 – L. A. Rayburn, Phys. Rev. 122 (1961) 168 [$T_{1/2}$].
- 1962Ma15 – J. D. Mahony, S. S. Markowitz, UCRL 10624 (1963) [$T_{1/2}$].
- 1963Be31 – K. Beg, F. Brown, Int. J. Rad. Isotopes 14 (1963) 137 [$T_{1/2}$].
- 1964Ho09 – I. Hofmann, Acta. Phys. Austriaca 18 (1964) 309 [$T_{1/2}$].
- 1964Ma07 – J. D. Mahony, S. S. Markowitz, J. Inorg. Nucl. Chem. 26 (1964) 907 [$T_{1/2}$].
- 1965Eb02 – T. G. Ebrey, P. R. Gray, Nucl. Phys. 61 (1965) 479 [$T_{1/2}$].
- 1965Bo38 – M. Bormann, E. Fretwurst, P. Schehka, G. Wrege, H. Büttner, A. Lindner, H. Meldner, Nucl. Phys. 63 (1965) 438 [$T_{1/2}$].
- 1969Ka17 – R. W. Kavanagh, Nucl. Phys. A129 (1969) 172 [$T_{1/2}$].
- 1970Al17 – D. E. Alburger, D. H. Wilkinson, Phys. Lett. 32B (1970) 190 [End-point energy].
- 1972Aj01 – F. Ajzenberg-Selove, Nucl. Phys. A190 (1972) 1 [$T_{1/2}$, end point energy, Q, log ft].
- 1973Ho43 – K. R. Hogstrom, B. W. Mayes, L. Y. Lee, J. C. Allred, C. Goodman, G. S. Mutchler, C. R. Fletcher, G. C. Phillips, Nucl. Phys. A215 (1973) 598 [$T_{1/2}$].
- 1973Fi13 – M. L. Fitzpatrick, K. W. D. Ledingham, J. Y. Gourlay, J. G. Lynch, J. Phys. A6 (1973) 713 [K/ β^+ ratio, end-point energy].
- 1977Ba49 – W. Bambynek, H. Behrens, M. H. Chen, B. Crasemann, M. L. Fitzpatrick, K. W. D. Ledingham, H. Genz, M. Muttere, R. L. Intemann, Revs. Modern Phys. 49 (1977) 77 [Electron Capture].
- 1978Aj01 – F. Ajzenberg-Selove, Nucl. Phys. A300 (1978) 1 [$T_{1/2}$, Q, log ft].
- 1980Ru02 – A. R. Rutledge, L. V. Smith, J. S. Merritt, AECL 6692 (1980) 2 [$T_{1/2}$].
- 1987Aj01 – F. Ajzenberg-Selove, Nucl. Phys. A475 (1987) 1 [$T_{1/2}$, Q, log ft].
- 1989Ka01 – T. Katoh, K. Kawade, H. Yamamoto, JAERI-M 089-083 (1989) [$T_{1/2}$].
- 1995Au04 – G. Audi, A.H. Wapstra, Nucl. Phys. A595 (1995) 409 [Q].
- 2004Sc00 – H. Schrader, Appl. Rad. Isotopes 60 (2004) 317 [$T_{1/2}$].

²²Na - Comments on evaluation of decay data
M. Galán

No substantial differences with previous Helmer and Schönfeld ²²Na evaluation (1999BeZQ) are found. Only Q-value is changed and a new ε/β^+ experimental ratio (2009NA08) is available since 1997.

1) Decay Scheme

²²Na disintegrates by electron capture and β^+ emission to excited level of 1274-KeV in ²²Ne.

²²Na ground state has $J_\pi = 3^+$ from Helmer and Schönfeld evaluation (1997).

The level scheme is complete. A good agreement has been found between the total decay energy of 2843,0 (24) keV computed for this decay scheme by RADLST code and the Q value of 2843,02 (21) keV.

2) Nuclear Data

The Q value is from new value of 2009AuZZ: $Q_{\beta^+} = 2843,02$ (21) keV obtained from the most recent measurements of 2004Mu26 and 2008Mu05. Other: 2842,3 (4) (2003AU03).

The measured ²²Na half-life values, in years, are:

Reference	Value (a)	Comments
2002UN02, 1992UN01	2,6037 (3)	
1982RUZV	2,6018 (7)	
1980HO17	2,6019 (4)	
1965AN07	2,613 (11)	Rejected by Chauvenet's criterion
1965AN07	2,603 (1)	
1965AN07	2,602 (11)	
1961WY01	2,62 (2)	Rejected by Chauvenet's criterion
1957ME47	2,58 (3)	Rejected by Chauvenet's criterion
	Mean	Reduced χ^2
LWM	2,6029 (8)	3,32
NRM	2,6023 (3)	2,37
RT	2,6021 (3)	

1965AN07 reported a fourth value of 2,5917 (30) which has been omitted from the analysis as it is inconsistent with all of other values. The previous values of 2,6019 (3) in 1980RUZX (replaced by 1982RUZV) and that of 2,5775 (3) in 1982HOJZ (replaced by 1992UN01) have not been included.

The Lweight for Excel and AveTool computer codes have been used with these eight input values. The weighted mean of the Limitation of Relative Statistical Weight Method (LWM) was the same result in both codes. AveTool also estimates the weighted mean by two more methods: Normalised Residual Method (NRM) (1992JA06) and Rajeval Technique (RT) (1992RA08). Following the most conservative method of LWM the eight values have been considered.

As it was discussed by Helmer and Schönfeld in their previous ²²Na evaluation (see Comments on ²²Na evaluation, 1999BeZS), the value of 2002UN02 is inconsistent with the other recent values from 1982RUZV and 1980HO17 and one could exclude the values before the 70's.

The values in 1957ME47, 1961WY01 and 1965AN07 were rejected based on the Chauvenet's criterion. For the remaining values, the largest contribution to the weighted average comes from the value of Unterweger (2002UN02). The LWM method increased the uncertainty of this value 1,093 times in order to reduce its relative weight to 50 %. The final uncertainty is also expanded from 0,0004 to 0,0008 to include de most precise value of 2,6037.

The recommended value is the more conservative LWM mean, 2,6029 (8) a or 950,6 (3) d [1 a = 365,242 198 78 d (1999BeZQ)] with an internal uncertainty of 0,0002 and an external of 0,0004.

Level energy has been obtained from a least-squares fit to γ -ray energies (GTOL computer code).

2.1) Electron Capture and Positron Transitions

Many different ε/β^+ ratios for the 1274-keV level have been measured. They are reported in Table 1 and compared with theoretical estimations:

Reference	ε/β^+ (experimental)	ε/β^+ (theoretical)	Comments
1954KR**†	0,124 (12)		
1954SH**†	0,110 (6)	0,1135 (20)	
1954ZW**†		0,111	
1955AL**†	0,122 (10)		
1958KO75	0,109 (8)		
1959RA09	0,112 (4)		
1964WI04	0,1041 (7)		
1967LE07	0,1048 (7)	0,1138 (25) 0,100 (6)	omitting e ⁻ exchange correction with e ⁻ exchange correction
1968VA13	0,1042 (10)	0,1118 (25)	
1969MC06	0,1136 (97)		From K/ β^+ = 0,1050 (90). The factor 1/1,0816 from 1977BO10 was used.
1976MA38	0,1077 (6)		
1977BA48		0,1117 (4)	
1977BO10	0,1128 (57)		
1978FI11		0,1152 (3)	
1983BA41	0,1079 (3)		
1990KU11	0,1050 (29)	0,1116 (3)	
2009NA08	0,1084 (27)		

† References not appear in NSR database. Nomenclature has been added by evaluator.

As can be seen in Table 1, experimental results present important discrepancies and they do differ substantially from theoretical predictions. Firestone et al. (1978FI11) discussed further about the anomalous ε/β^+ in ²²Na.

Statistical analyses of the experimental values have been done. In the experimental dataset the LWM method rejected 1954KR01 and 1955AL01 values based on Chauvenet's criterion. The uncertainty of 1983BA41 was changed to reduce its relative weight to 50 %. For the 12 input values the weighted mean is 0,1068 with an internal uncertainty of 0,0002 and a external of 0,0005 and a reduced χ^2 of 2,25. The adopted value is 0,1068 (11) with an uncertainty increased to include the most precise value of 0,1079. If data before 1960 are rejected the LWM is 0,1067 (12) with expanded uncertainty and reduced χ^2 of 2,8.

Experimental data and theoretical estimations are found to differ up to 10 %.

The P_{β^+} and P_{ε} were derived as follows: with $\frac{P_{\varepsilon}(1274)}{P_{\beta^+}(1274)} = 0,1068(11)$ from experimental results

and with $\frac{P_{\beta^+}(1274)}{P_{\beta^+}(0)} = 1600(400)$ from 1953WR13, these ratios were introduced in the relationship

$100 = P_{\beta^+}(1274) + P_{\varepsilon}(1274) + P_{\beta^+}(0)$ neglecting the electron capture branching to the ground state.

Then one obtain, $P_{\beta^+}(0) = 0,056 (14) \%$.

Then, the LOGFT program (theory) was run considering $P_{\varepsilon+\beta^+}(1274) = 99,944 (14) \%$ and $P_{\varepsilon+\beta^+}(0) = 0,056 (14) \%$. The ε/β^+ for the ground state estimated by the code is $0,01782 (18)$. Thus one has:

$$100 = P_{\beta^+}(1274) + 0,1068(11) \times P_{\beta^+}(1274) + \frac{1}{1600(400)} \times P_{\beta^+}(1274) + 0,01782(18) \times \frac{1}{1600(400)} \times P_{\beta^+}(1274)$$

That gives:

$$P_{\beta^+}(1274) = 90,30 (9) \%$$

$$P_{\varepsilon}(1274) = 9,64 (9) \%$$

$$P_{\beta^+}(0) = 0,055 (14) \%$$

$$P_{\varepsilon}(0) = 0,00098 (25) \%$$

Using EC-Capture program we have: $P_K = 0,9233 (35)$ and $P_L = 0,0767 (35)$

2.2) γ -ray Transitions

Transition Probabilities

The γ -transition probability is $P_{\varepsilon^+}(1274) + P_{\beta^+}(1274) = 90,30 (9) + 9,64 (9) = 99,94 (13) \%$

Internal conversion coefficients

The internal conversion coefficients (ICC) have been calculated using the BrIcc computer code, which interpolated ICC values from tables of Band et al. (2002BA85). Associated uncertainties are 1,4 %. The theoretical value of $6,71 (9) \times 10^{-6}$ agrees with the value of $6,8 (4) \times 10^{-6}$ from the analysis of experimental data (1985HAZA).

The theoretical α_{π} (1979SC31) interpolated for this E2 transition is found to be $2,34 (3) \times 10^{-5}$.

3) Atomic Data

3.1) Atomic values (ω_k , ω_L and η_{KL}) are from 1996SC06.

3.1.1) X-Radiations, 3.1.2) Auger electrons

The X-ray and Auger electron emission probabilities have been deduced from γ -ray and conversion electron data by using the computer code EMISSION. Results were verified with the RADLST computer code. Differences between both codes were less than 4 %.

4) Electron Emissions

The β^+ and the electron capture emission probabilities are discussed above.

5) Photon Emissions

Energies

γ -ray energy 1274,537 (7) is from 2000HE14. The level energy has been computed to account for the recoil energy in the daughter nucleus.

γ -ray emissions

The absolute P_γ is evaluated from $P_{\gamma+ce}$ and the total internal conversion coefficient $\alpha = (\alpha_\pi + \alpha_T)$:

$$P_\gamma = \frac{P_{\gamma+ce}}{1 + \alpha} = \frac{99,94(13)}{1 + 3,01(4) \times 10^{-5}} = 99,94(13)\%$$

The annihilation radiation emission probability is taken to be 2 times P_{β^+} , that is 180,7 (2) % without the correction factor for the annihilation-in-flight.

Additional reference:

R.G. Helmer, E. Schönfeld (1999BeZS) Evaluation and comments on evaluation of ²²Na. Table des Radionucléides, CEA-ISBN 2-7272-0211-3 (1999).

References

- 1953WR13 B.T. Wright, Phys. Rev. 90 (1953) 159 [P_{β^+}]
 1954KR** W.E. Kreger, Phys. Rev. 96 (1954) 1554 [ϵ/β^+]
 1954SH** R. Sherr, R.H. Miller, Phys. Rev. 93 (1954) 1076 [ϵ/β^+]
 1954ZW** P.F. Zweifel, Phys. Rev. 96 (1954) 1572 [ϵ/β^+]
 1955AL** R.A. Allen, W.E. Burcham, K.F. Chakett, G.L. Munday, P. Reasbeck, Proc. Phys. Soc. 68 (1955) 681 [ϵ/β^+]
 1957ME47 W.F. Merrit, P.J. Campion, R.C. Hawkings, Can. J. Phys. 35 (1957) 16 [$T_{1/2}$]
 1958KO75 J. Konijn, B. Van Nooijen, H.L. Hagedoorn, A.H. Wapstra, Nucl. Phys 9 (1958) 296 [ϵ/β^+]
 1959RA09 M.K. Ramaswamy, Indian J. Phys 33 (1959) 285 [ϵ/β^+]
 1961WY01 E.I. Wyatt, S.A. Reynolds, T.H. Handley, W.S. Lyon, H.A. Parker, Nucl. Sci. Eng. 111 (1961) 74 [$T_{1/2}$]
 1964WI04 A. Williams, Nucl. Phys. 52 (1964) 324 [ϵ/β^+]
 1965AN07 S.C. Anspach, L.M. Cavallo, S.B. Garfinkel, J.M.R. Hutchinson, C.N. Smith, Report NP 15663 (1965) 1 [$T_{1/2}$]

- 1967LE07 H. Leutz, H. Wenninger, Nucl. Phys. A99 (1967) 55 [ϵ/β^+]
- 1968VA13 E. Vatai, D. Varga, J. Uchrin, Nucl. Phys. A116 (1968) 637 [ϵ/β^+]
- 1969MC06 M.F. McCann, K.M. Smith, J. Phys (London) A2 (1969) 392 [ϵ/β^+]
- 1970WA11 E. K. Warburton, G. T. Garvey, I. S. Towner, Ann. Phys. 57(1970)174 [P_{β^+}]
- 1971ME** J. S. Merritt, J. V. G. Taylor, Report AECL – 3912 (1971) [ϵ/β^+]
- 1973KA50 J. Kantele, M. Valkonen, Nucl. Instrum. Meth. 112(1973)501 [P_γ]
- 1976MA38 T.D. MacMahon, A.P. Baerg, Can. J. Phys. 54 (1976) 1433 [ϵ/β^+]
- 1977BA48 W. Bambynek, H. Behrens, M.H. Chen, B. Crasemann, M.L. Fitzpatrick, K.W.D. Ledingham, H.Genx, M. Mutterer, R.L. Intermann, Rev. Mod. Phys. 49 (1977) 77 [ϵ/β^+]
- 1977BO10 H.E. Bosch, J. Davidson, M. Davidson, L. Szbisz, Z. Phys A280 (1977) 321 [ϵ/β^+]
- 1978FI11 R.B. Firestone, Wm.C. McHarris, B.R. Holstein, Phys. Rev. C18 (1978) 2719 [ϵ/β^+]
- 1979SC31 P.Schluter, G.Soff, At.Data Nucl.Data Tables 24, 509 (1979) [α]
- 1980HO17 H. Houtermans, O. Molosevic, F. Reichel, Int. J. App. Radiat. Isot. 31 (1980) 153 [$T_{1/2}$]
- 1982RUZV A.R. Rutledge, L.V. Smith, J.S. Merrit, NBS – SP - 626 (1982) 5 [$T_{1/2}$]
- 1983BA41 A.P. Baerg, Can. J. Phys 61 (1983) 1222 [ϵ/β^+]
- 1985HaZA H.H. Hansen, European Appl. Res. Rept-Nucl. Sci. Technol. 6 (1985) 777 [α_K , α_T]
- 1990KU11 V. Kunze, W.D. Schmidt-Ott, H. Behrens, Z. Phys A 337 (1990) 169 [ϵ/β^+]
- 1991BAZS W. Bambynek, T. Barta, R. Jedlovsky, P. Christmas, N. Coursol, K. Debertain, R.G. Helmer, A.L. Nichols, F.J. Schima, Y.Yoshizawa. Report IAEA-TECDOC-619 (1991) [$T_{1/2}$, P_γ]
- 1992UN01 M.P. Unterweger, D.D. Hoppes, F.J. Schima, Nucl. Instrum. Meth. Phys. Res. A312 (1992) 349 [$T_{1/2}$]
- 1995SCZY E. Schönfeld, Report PTB 6.33-95-2 (1995) [P_K , P_L , P_M theory]
- 1996SC06 E. Schönfeld, H. Janssen, Nucl. Instrum. Meth. Phys. Res. A 369 (1996) 527 [atomic data]
- 1999BEZS M.-M. Bé, B. Duchemin, E. Browne, S.-C. Wu, V. Chechev, R. Helmer, E. Schönfeld. Table of Radionuclides: Comments on Evaluations. CEA-ISBN 2-7272-0211-3 (1999)
- 1999BEZQ M.-M. Bé, B. Duchemin, J. Lamé, C. Morillon, F. Piton, E. Browne, V. Chechev, R. Helmer, E. Schönfeld. Table of Radionuclides. CEA-ISBN 2-7272-0200-8 (1999)
- 2000HE14 R.G. Helmer, C. van der Leun, Appl. Radiat. Isot. 52 (2000) 601 [γ -ray energies]
- 2002BA85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor Jr., At. Data Nucl. Data Tables 81 (2002) 1 [ICC]
- 2002UN02 M.P. Unterweger, Nucl. Inst. Meth.A56 (2002) 125 [$T_{1/2}$]
- 2003AU03 G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)337 [Q value]
- 2004MU26 M. Mukherjee, A. Kellerbauer, D. Beck, K. Blaum, G. Bollen, F. Carrel, P. Delayahe, J. Dilling, S. George, C. Guenaut, F.Herfurth, A. Herlert, H.-J. Kluge, U. Koster, D. Lunney, S. Schwarz, L. Schweikhard, C. Yazidjian, Phys. Rev. Lett. 93 (2004) 150801 [Na mass excess]
- 2008MU05 M. Mukherjee, D. Beck, K. Blaum, G. Bollen, P. Delayahe, J. Dilling, S. George, C. Guenaut, F.Herfurth, A. Herlert, A. Kellerbauer, H.-J. Kluge, U. Koster, D. Lunney, S. Schwarz, L. Schweikhard, C. Yazidjian, Eur. Phys. J. A35 (2008) 31 [Na mass excess]
- 2009NA08 O. Nähle, K. Kossert, R. Klein, App. Radiat. Isot. 66 (2009) 865 [ϵ/β^+]
- 2009AUZZ G. Audi, W. Meng, D. Lunney, B. Pfeiffer, Priv. Comm. (2009) [Q value]

Comments of ²⁴Na Evaluation by R. G. Helmer and E. Schönfeld

1 Decay Scheme

The decay scheme is complete since the four levels populated in this decay are the only excited levels in ²⁴Mg below the decay energy.

The spins, parities, and half-lives of the excited levels are from the Endt evaluation 1990En08.

2 Nuclear Data

For the half-life, the following values are available (in hours):

14.90 (2)	1949Wi10, Wilson and Bishop (1949)	
15.10 (4)	1950Co69, Cobble and Atteberry (1950)	
14.97 (2)	1953Lo09, Lockett and Thomas (1953)	
14.90	1955To07, Tobailem (1955)	omitted - no uncertainty
14.959 (10)	1958Ca20, Campion and Merritt (1958)	
14.953 (13)	1960Wo07, Wolf(1960)	
15.05 (2)	1961Wy01, Wyatt et al. (1961)	superseded by 1972Em01
15.04 (5)	1962Mo21, Monahan et al. (1962)	
15.00 (2)	1968La10, Lagoutine et al. (1968)	superseded by 1982La25
15.16 (5)	1969Ke14, Kemeny (1969)	omitted - no background subtraction
15.030 (3)	1972Em01, Emery et al. (1972)	omitted - outlier
14.969 (12)	1974Ch25, Chakraborty (1974); average of 6 values with external uncertainty	
15.09 (6)	1976Ge06, Genz et al. (1976)	
15.010 (28)	1978Da21, Davis et al. (1978)	
14.9590 (12)	1980Ho17, Houtermans et al. (1980)	
14.964 (15)	1980Mu11, Muckenheim et al. (1980)	
14.965 (10)	1980RuZY, Rutledge et al. (1980)	superseded by 1982RuZY
14.965 (10)	1982RuZV, Rutledge et al. (1982)	
14.956 (3)	1982La25, Lagoutine, Legrand (1982);	originally $\sigma=0.008$ divided by 3
14.951 (3)	1982HoZJ, Hoppes et al. (1982)	superseded by 1992Un01
14.9575 (28)	1983Wa26, Walz et al. (1983)	
15.027 (2)	1989Ab05, Abzouzi et al. (1989)	omitted - outlier
14.90 (2)	1991Bo34, Bode et al. (1991)	
14.9512 (32)	1992Un01, Unterweger et al. (1992)	
14.86 (12)	1994Mi03, Mignonsin (1994)	

14.9574 (20) adopted value, LRSW weighted average

In the final weighted average, the values of 1972Em01 and 1989Ab05 have been omitted because they are outliers; both are over 30σ from the adopted value. If these values are included, the reduced- χ^2 value is about 80. For the 17 values included, the Limitation of Relative Statistical Weight, LRSW, method (1985ZiZY, 1992Ra08) increases the uncertainty of the value of 1980Ho17 from 0.001

to 0.0016 in order to reduce its relative weight from 73% to 50%. In addition to this relative weight, those of the values of 1982La25, 1983Wa34, and 1992Un01 are between 13 and 15%. For the final weighted average the internal uncertainty is 0.0012, the reduced- χ^2 value is 3.01, and the external uncertainty is 0.0020.

1974Ch25 have measured this half-life for solid NaCl and for an aqueous solution. No change of the half-life was observed, contrary to the report of 1969Ke14.

The Q_{β^-} value is taken from the 1995Au04 evaluation.

2.1 β^- Transitions

The energies are calculated from the Q_{β^-} value and the level energies. In the following list, nine values of the experimentally determined β^- end-point energy (in keV) for the transition to the 4122-keV level are compared with the value derived from Q value.

1394 (4)	1957Po36, Porter et al. (1957)
1389 (4)	1958Da10, Daniel (1958)
1389 (2)	1961De23, 1965De25, Depommier and Chabre (1961)
1395	1963Pa20, Paul et al. (1963)
1393 (3)	1964Le09, Lehmann (1964)
1394 (2)	1965Be24, Beekhuis and De Waard (1965)
1389.2 (5)	1969Bo48, Booij et al. (1969)
1389 (2)	1972Gi17, Gils et al. (1972)
1390 (1)	1976Ge06, Genz et al. (1976)
1392.94 (16)	$Q - E(4122)$

The measured and calculated probabilities (in %) of the β^- transitions are:

Level (keV)	1950Gr01 Grant(1950)	1951Tu12 Turner (1951)	Present evaluation
5236			0.057 (7)
4239			<0.002
4122	100	100	99.939 (8)
1368	<0.01	0.003	0.003 (2)
0			$<5 \times 10^{-10}$

The 4th forbidden β^- branch to the ground state has not been observed. From the experimental limit on the number of counts in the β^- spectrum above 4140 keV, 1951Tu12 give $\lg ft > 15.1$. The $\lg ft$ systematics of 1998Si17 lists four decays of this type with $\lg ft$ values of 22.5 to 24.3. Since this is a very small set of values, we have taken the lower limit of the ²⁴Na $\lg ft$ to be 20, which corresponds to $I_{\beta^-}(0) < 5 \times 10^{-10} \%$; this value is adopted.

The β^- branch to the 4238 level is a 2nd forbidden transitions and the $\lg ft$ systematics (1998Si17) give $\lg ft > 10.6$ which corresponds to $I_{\beta^-}(4238) < 0.002\%$; this value is adopted. This small value is supported by the adopted decay scheme for which the intensity of the 998-keV γ -ray feeding this level is more [0.00151(25)] than that depopulating it [0.00024(3) + 0.00084(10)]. An unobserved γ -ray of 116 keV could also depopulate this level.

No direct measurements are reported for the β^- transitions to the 4238- and 5236-keV levels. The adopted value for the transition to the 1368-keV level is based on the measurement of 1951Tu12 [Turner and Cavanagh (1951)] who gave no uncertainty. The adopted value for the transition to the 5236-keV level was calculated from probabilities of the two de exciting γ -rays and their internal and pair conversion.

The β^- branch to the 4122-keV level is 100% less the intensity of those to the levels at 0, 1368, 4238, and 5236 keV. The sum of the latter four is 0.061(8)%, so the former is 99.939(8)%.

2.2 Gamma Transitions

The transition probabilities of the 3866- and 4237-keV γ -rays are determined from the following measurements:

	3867 keV	4237 keV
1960Ar10, Artamonova <i>et al.</i> (1960)	0.09 (2)	0.0015 (5)
1962Mo21, Monahan <i>et al.</i> (1962)	0.075 (20)	0.008 (3)
1968Va06, van Klinken <i>et al.</i> (1968)	0.063 (6)	
1970Le12, Lebowitz (1970)	0.0489 (25)	<0.0033
1972Ra21, Raman <i>et al.</i> (1972)	0.061 (5)	0.00084 (10)
Adopted value	0.056(7)	0.00084(10)

For the 3866-keV γ -ray, the adopted value is the average of all five values, which gives an internal uncertainty of 0.0026, a reduced- χ^2 value of 2.46, and an external uncertainty of 0.0041, and the final uncertainty was expanded to include the most precise value. For the 4237-keV γ -ray, the value of 1972Ra21 is adopted as it is considered to be the most reliable and it is consistent with the limit of 1970Le12.

The 996- and 2869-keV γ -ray transitions are not observed in ²⁴Na decay, but their emission probabilities can be deduced from the relative probabilities in other decays or reactions. The transition probability of 996-keV γ -ray was calculated from the measured $P_\gamma(996)/P_\gamma(3866)$ ratio. For this ratio, the measured values are :

0.017 (5)	1972Me09, Meyer <i>et al.</i> (1972) from ²³ Na(p, γ)
0.019 (2)	1973Le15, Leccia <i>et al.</i> (1973) from ²³ Na(p, γ)
0.015 (3)	1975Bo43, Boydell <i>et al.</i> (1975) from ²³ Na(p, γ)
0.0260 (17)	1981Wa07, Warburton <i>et al.</i> (1981) from ²⁴ Al ϵ decay
0.030 (4)	1990En02, Endt <i>et al.</i> (1990) from ²³ Na(p, γ)
0.022 (4)	Adopted value

The adopted value is the weighted average value of 0.0222 with an internal uncertainty of 0.0011, a reduced- χ^2 of 4.6 and an external uncertainty of 0.0024. The LRSW method increases the final uncertainty to 0.004 to include the most precise value of 0.0260. With the above value of $P_\gamma(3866)$, we obtain $P_\gamma(996) = 0.00123(27)$.

The ratio $P_\gamma(2869)/P_\gamma(4237)$ ratio has been measured as follows:

0.30 (3)	1972Me09, Meyer <i>et al.</i> (1972) from ²³ Na(p, γ)
----------	--

0.30 (3)	1972Ra21, Raman <i>et al.</i> (1972) from ²⁴ Mg(n,n'γ)
0.299 (15)	1973Le15, Leccia <i>et al.</i> (1973) from ²³ Na(p,γ)
0.267 (7)	1973Br16, Branford (1973) from ²³ Na(p,γ)
0.299 (19)	1975Bo43, Boydell (1975) from ²³ Na(p,γ)
0.304 (19)	1981Wa07, Warburton <i>et al.</i> (1981) from ²⁴ Al ε decay

0.284 (7) Adopted value

The adopted value is the weighted average of all six values after the uncertainty for the 1973Br16 value was increased from 0.007 to 0.009 to reduce its relative weight from 63% to 50%. This average has an internal uncertainty of 0.006, a reduced-χ² of 1.37, and an external uncertainty of 0.007. With the above adopted value of 0.00084(10) for P_γ(4237), one obtains P_γ(2869) = 0.00024(3).

If there are no direct feeding the ground state by β⁻ decay or the unobserved γ transitions of 4122 and 5236 keV, T_γ(1368) = 100 - T_γ(4237) = 99.99916(10) where T_γ = P_γ (1.0 + α + α_π). Upper limits for transition intensities of the 4122- and 5236-keV γ-rays can be determined from the ratios measured by 1981Wa07: P_γ(4122)/P_γ(2754) < 0.00001, or P_γ(4122) < 0.001 and P_γ(5236)/P_γ(3867) < 0.004, so P_γ(5236) < 0.00023 and by 1972Ra21 and 1967En05 which give P_γ(4122) < 0.0009 and P_γ(5236) < 0.00002. If the 4122- and 5236-keV transitions have intensities equal to the latter upper limits, the value of T_γ(1368) would reduce from 99.99916 to 99.9983. Since it is unlikely that these two values will be at the limits, we have adopted the value of T_γ(1368) = 99.9990(3) and P_γ(1368) = 99.9935(5).

The 1114-keV transition between the 5236- and 4122-keV levels has not been observed in ²⁴Na decay. In the ²⁴Al decay, 1981Wa07 have found an upper limit of the ratio P_γ(1114)/P_γ(3867) < 0.007 which yields the value of P_γ(1114) < 0.0004.

The transition probability of the 2754-keV γ-ray is calculated from the balance condition T_γ(2754) = T_γ(1368) - [T_γ(2869) + T_γ(3867) + P_{β⁻}(1368)]. This yields T_γ(2754) = 99.9990(3) - 0.059(7) = 99.940(7)%, which gives P_γ(2754) = 99.872(8)%.

From the intensity balance at the 4238-keV level, for a possible depopulating γ-ray of 116 keV, P_γ(116) = 0.0004(3) + I_γ(4238). Since this γ-ray has not been observed, it is omitted from the scheme.

The internal-conversion coefficients are interpolated from the tables of theoretical values (Band *et al.*, 1976). The mixing parameters, δ, were based on the following information:

γ energy	1960Ba19	1963Br15	1973Le15	adopted
998			-5.1 (+8-12) or -0.47 (4)	-0.47 (4)
2869	+23 (9)		> 30	+23 (9)
3867		large	-0.21 (2) or >19	pure E2

The uncertainty of the interpolated conversion coefficients is assumed to be 3 %.

The internal-pair-formation coefficients (α_γ) for the 1368- and 2754-keV γ-rays have been interpolated from calculated values of 1979Sc31 and are in reasonable agreement with measured values which are:

1368 keV	2754 keV
	0.00116 (10) 1949Ra01
	0.00076 (19) 1950Mi82

Comments on evaluation

	0.00067 (10)	1951Cl50
0.00006 (1)	0.00071 (2)	1952Bl53
0.00003	0.00080	1952Sl52

In summary, the γ -ray photon and transition intensities are:

Energy (keV)	Transition (%)	Photon (%)
998	0.00151 (25)	
1114	<0.0004	
1368	99.9990 (3)	99.9935 (5)
2869	0.00024 (3)	
2754	99.940 (7)	99.872 (8)
3867	0.056 (7)	
4122	<0.0009	
4238	0.00084 (10)	
5236	<0.00002	

If P_γ is not given, it is equal to T_γ .

3 Atomic Data

The values for ω_K , the mean ω_L , and η_{KL} are taken from 1996Sc06.

3.1 X Radiation

The mean energies of the K_α radiations have been calculated from the wave lengths given by 1967Be65.

3.2 Auger Electrons

The mean energy of the KLL Auger electrons is taken from 1977La19.

4 Radiation Emission

4.1 Electron Emission

The energies and emission probabilities of the particles are the same as those given already in sect. 2.1. The energies of the electron from internal conversion and internal-pair formation are calculated from the γ -ray energies. The number of electrons per disintegration for various processes are calculated from the γ -ray emission probabilities, $\alpha_{\tau\gamma}$, α , and the atomic data.

4.2 Photon Emission

The energies of the two main γ -rays are from 2000He14. From the decay of ²⁴Na, the energies for the 3867- and 4238-keV γ -rays are 3867.5(3) from 1968Va06 and 1970Le12 and the 4237.4(10) keV from 1972Ra21. The energies of the 996- and 2869-keV γ -rays would then be calculated from the level energies. The adopted values for all four of these γ -rays have been taken from the decay of ²⁴Al (1981Wa07).

The number of photons per disintegration were calculated as described in sect. 2.2.

5 Main Production Modes

Taken from N. Coursol, Table de Radionucléides (1982).

References

- 1949Ra01 - E. R. Rae, Phil. Mag. **40**(149)1155 [α_π]
 1949Wi10 - R. Wilson, G. R. Bishop, Proc. Phys. Soc. (London) **62**(1949)457 [$T_{1/2}$]
 1950Co69 - J. W. Cobble, R. W. Atteberry, Phys. Rev. **80**(1950)917 [$T_{1/2}$]
 1950Gr01 - P. . Grant, Proc. Phys. Soc. (London) **63**(1950)1298 [P_{β^-}]
 1950Mi82 - W. Mims, H. Halban, R. Wilson, Nature **166**(150)1571 [α_π]
 1951Cl50 - M. R. Cleland, J. Townsend, A. L. Hughes, Phys. Rev. **84**(1951)298 [α_π]
 1951Tu12 - J. F. Turner, P. E. Cavanagh, Phil. Mag. **42**(1951)636 [P_{β^-}]
 1952Bl53 - S. D. Bloom, Phys. Rev. **88**(1952)312 [α_π]
 1952Sl52 - H. Slätis and K. Siegbahn, Arkiv f. Fysik **4**(1952)485 [α_π]
 1953Lo09 - E. E. Lockett, R. H. Thomas, Nucleonics **11**(1953)14 [$T_{1/2}$]
 1955To07 - J. Tobailem, J. Phys. Radium **16**(1955)48 [$T_{1/2}$]
 1957Po36 - F. T. Porter, F. Wagner, Jr., M. S. Freedman, Phys. Rev. **107**(1957)135 [$E_{\beta^{max}}$]
 1958Ca20 - P. J. Champion and J. S. Merritt, Can. J. Phys. **36**(1958)983 [$T_{1/2}$]
 1958Da10 - H. Daniel, Nucl. Phys. **8**(1958)191 [$E_{\beta^{max}}$]
 1960Ar10 - K. P. Artamonova, L. V. Gustova, Y. N. Podkapaev, O. V. Chubinskii, Soviet Phys. JETP **12**(1961)1109 [P_γ]
 1960Ba19 - R. Batchlor, A. J. Ferguson, H. E. Gove, A. E. Litherland, Nucl. Phys. **16**(1960)38 [δ]
 1960Wo07 - G. Wolf, Nukleonik **2**(1960)255 [$T_{1/2}$]
 1961De23, 1961De25 - P. Depommier, M. Chabro, J. Phys. Radium **22**(1961)656 and 674 [$E_{\beta^{max}}$]
 1961Wy01 - E. I. Wyatt, S. A. Reynolds, T. H. Handlcy, W. S. Lyon, H. A. Parkcr, Nucl. Sci. Eng. **11**(1961)74 [$T_{1/2}$]
 1961Mo09 - J. E. Monahan, S. Raboy, C. C. Trail, Nucl. Phys. **33**(1962)633 [P_γ]
 1962Mo21 - J. E. Monahan, S. Raboy, C. C. Trail, Nucl. Instr. Meth. **17**(1962)225 [$T_{1/2}$]
 1963Br15 - C. Broude, H. E. Gove, Ann. Phys. (New York) **23**(1963)71 [δ]
 1963Pa20 - H. Paul, F. P. Viehböck, P. Skarek, H. Baicr, I. Hoffmann, H. Wotke, Acta Phys. Austr. **16**(1963)278 [$E_{\beta^{max}}$]
 1964Le09 - J. Lehmann, J. Phys. **25**(1964)326 [$E_{\beta^{max}}$]
 1965Be24 - H. Beekhuis, H. De Waard, Nucl. Phys. **74**(1965)459 [$E_{\beta^{max}}$]
 1965Mu03 - G. Murray, R. L. Graham, J. S. Geiger, Nucl. Phys. **63**(1965)353 [E_γ]
 1965Re16 - J. J. Reidy, M. L. Wiedenbeck, Bull. Am. Phys. Soc. **10**(1965)1131, abstract SP2 [E_γ]
 1965Sp08 - E. Spring, Phys. Lett. **18**(1965)132 [α]
 1967Be65 - J. A. Bearden, Rev. Mod. Phys. **39**(1967)78 [E_X]
 1967En05 - P. M. Endt, C. van der Leun, Nucl. Phys. A **105**(1967)1 [P_γ]
 1968La10 - F. Lagoutine, Y. le Gallic, J. Lcgrand, Int. J. Appl. Rad. Isot. **19**(1968)475 [$T_{1/2}$]
 1968Va06 - J. van Klinken, F. Pleiter, H. T. Dijkstra, Nucl. Phys. A **112**(1968)372 [P_γ]
 1969Ke14 - P. Kemeny, Radiochem. Raccioanal. Letters **2**(1969)119 [$T_{1/2}$]
 1969Bo48 - H. M. W. Booij, E. A. Van Hoek, J. Blok, Nucl. Instr. and Meth. **72**(1969)40 [$E_{\beta^{max}}$]
 1970Le12 - J. Lebowitz, A. R. Sayres, C. C. Trail, B. Weber, P. L. Zirkind, Nuovo Cim. **65A**(1970)675 [E_γ, P_γ]
 1972Gi17 - H. J. Gils, D. Flothmann, R. Löhken, W. Wiesner, Nucl. Instr. Meth. **105**(1972)179 [$E_{\beta^{max}}$]
 1972Em01 - J. F. Emery, S. A. Reynolds, E. I. Wyatt, G. I. Gleason, Nucl. Sci. Eng. **48**(1972)319 [$T_{1/2}$]
 1972Me09 - M. A. Meyer, J. P. L. Reinecke, D. Reitmann, Nucl Phys. **A185**(1972)625 and Erratum Nucl.

- Phys. A196(1972)635 [E_γ, P_γ]
 1972Ra11 - F. Rahn, H. Camarda, G. Hacken, W. W. Havens, Jr., H. Liou, J. Rainwater, M. Slagowitz, S. Wynchank, Nucl. Sci. Eng. **47**(1972)372 [E_γ]
 1972Ra21 - S. Raman, N. B. Gove, J. K. Dicken, T. A. Walkiewicz, Phys. Lett. **40B**(1972)89 [E_γ, P_γ, P_β-]
 1973Br16 - D. Branford, Austral. J. Phys. **26**(1973)1995 [P_β-]
 1973Le15 - F. Leccia, M. M. Alconard, D. Castera, P. Hubert, P. Mennrath, J. Phys. **34**(1973)147 [P_γ, δ]
 1973Ra10 - S. Raman, N. B. Gove, Phys. Rev. C **7**(1973)1995 [P_β-]
 1974Ch25 - S. Chakraborty, Radiochem. Radioanal. Lett. **17**(1974)61 [T_{1/2}]
 1975Bo43 - S. G. Boydell, D. G. Sargood, Austral. J. Phys. **28**(1975)369 [P_γ]
 1976Ba63 - I. M. Band, M. B. Trzhaskovskaya, M. A. Listengarten, At. Data Nucl. Data Tables **18**(1976)433 [α]
 1976Ge06 - H. Genz, J. Reisberg, A. Richter, B. M. Schmitz, G. Schrieder, K. Werner, H. Behrens, Nucl. Instr. Meth. **134**(1976)309 [T_{1/2}, E_β^{max}]
 1977La19 - F. P. Larkins, Atom. Data Nucl. Data Tables **20**(1977)311 [E_{Auger}]
 1978Da21 - M. C. Davis, W. C. Bowman, J. C. Robertson, Intern. J. Appl. Radiat. Isot. **29**(1978) 331 [T_{1/2}]
 1979Gr01 - R. C. Greenwood, R. G. Helmer, R. J. Gehrke, Nucl. Instr. and Meth. **159**(1979)465 [E_γ]
 1979Sc31 - P. Schlüter, G. Soff, At. Data Nucl. Data Tables **24**(1979)509 [P_γ]
 1980Ho17 - H. Houtermans, O. Milosevic, F. Reichel, Intern. J. Appl. Radiat. Isot. **31**(1980)153 [T_{1/2}]
 1980Mu11 - W. Muckenheim, P. Rullhusen, F. Smend, M. Schumacher, Nucl. Instr. Meth. **173**(1980)403 [T_{1/2}]
 1980RuZY - A. R. Rutledge, L. V. Smith, J. S. Merritt, AECL-6692(1980) [T_{1/2}]
 1981Wa07 - E. K. Warburton, C. J. Lister, D. E. Alburger, J. W. Olness, Phys. Rev. **C23**(1981)1242 [P_γ]
 1982HoZJ - D. D. Hoppes, J. M. R. Hutchinson, F. J. Schima, M. P. Unterweger, NBS-SP-626 (1982)85 [T_{1/2}]
 1982La25 - F. Lagoutine, J. Legrand, Intern. J. Appl. Radiat. Isot. **33**(1982)711
 1982RuZV - A. R. Rutledge, L. V. Smith, J. S. Merritt, NBS-SP-626 (1982)5
 1983Wa26 - K. F. Walz, K. Debertin, H. Schrader, Intern. J. Appl. Radiat. Isot. **34**(1983)1191 [T_{1/2}]
 1985ZiZY - W. L. Zijp, report ECN FYS/RASA-85/19 (1985) [averages]
 1989Ab05 - A. Abzouzi, M. S. Antony, V. B. Ndocko Ndonguc, J. Radioanal. Nucl. Chem. **135**(1989)1 [T_{1/2}]
 1990En02 - P. M. Endt, et al., Nucl. Phys. **A510**(1990)209 [P_γ]
 1990En08 - P. M. Endt, Nucl. Phys. **A521**(1990)209; Errata and Addenda Nucl. Phys. **A529**(1991)763; and Errata Nucl. Phys. **A564**(1993)609 [J^π]
 1991Bo34 - P. Bode, M. J. J. Ammerlaan, M. Koese, Appl Radiat. Isot. **42**(1991)692 [T_{1/2}]
 1992Ra08 - M. U. Rajput, T. D. MacMahon, Nucl. Instr. Meth **A312**(1992)289 [T_{1/2}]
 1992Un01 - M. P. Unterweger, D. D. Hoppes, F. J. Schima, Nucl Instr. and Meth. **A312**(1992)349 [T_{1/2}]
 1994Hu23 - J. H. Hubbell, P. N. Trehan, Nirmal Singh, B. Chand, D. Mehta, M. L. Garg, R. R. Garg, Surinder Singh and S. Puri, J. Phys. Chem. Ref. Data **23**(1994)339 [ω_L]
 1994Mi03 - E. P. Mignonsin, Appl Radiat. Isot. **45**(1994)17 [T_{1/2}]
 1995Au04 - G. Audi, A. H. Wapstra, Nucl. Phys. **A595**(1995)409 [Q_β-]
 2000He14 - R. G. Helmer and C. van der Leun, Nucl. Instr. Meth. **A450**(2000)35 [γ-energies]

³²P – Comments on evaluation of decay data by V. Chisté and M. M. Bé

1) Decay Scheme

³²P disintegrates by β^- emission (100 %) to the ground state of the stable nuclide ³²S.

2) Nuclear Data

The Q value (1710,66 (21) keV) is from Audi and Wapstra evaluation (1995Au04), and has been calculated with the formula:

$$Q = M(A, Z) - M(A, Z + 1),$$

where M(A,Z) and M(A,Z+1) are the measured atomic masses of ³²P and ³²S, respectively.

This value is in agreement with a weighted average value of 1708 (7) keV, which was calculated from measured values of the β^- end-point energy (see **β^- Transition**).

The measured ³²P half-life values (in days) are given below:

Reference	$T_{1/2}$ Value (days)	Comments
Ambrosen (1934Am01)	17,5 (11)	Omitted from analysis
Preiswerk (1935Pr20)	15,0 (15)	"
Sizoo (1936Si10)	15,0 (1)	"
Newson (1937Ne19)	14,5 (3)	"
Capron (1938Ca08)	14,5 (3)	"
Cacciapuotu (1938Ca15)	14,30 (3)	
Mulder (1940Mu04)	14,07 (3)	Omitted, outlier
Klema (1948Kl06)	14,35 (5)	
Sinclair(1951Si26)	14,60 (5)	Omitted, outlier
Locket (1953Lo19)	14,50 (4)	Omitted, outlier
Bayly (1956Ba25)	14,30 (9)	
Anders (1957An57)	14,223 (30)	Original $Uc \times 2$
Daniel (1958Da08)	14,2 (3)	
Robert (1959Ro24)	14,55 (6)	Omitted, outlier
Marais (1961Ma01)	14,282 (20)	Original $Uc \times 2$
Goodier (1966Go17)	14,290 (28)	Original $Uc \times 2$
Pernaa (1969Pe16)	14,32 (1)	
Lagoutine (1969La28)	14,268 (42)	
Belyaev (1977Be21)	12 (2)	Omitted, outlier
Mudhole (1977Mu15)	14,35 (5)	
Precker (1979Pr36)	14,28 (4)	
Coursey (1994Co26)	14,26 (1)	

The first five and less precise historical values were omitted from analysis. In several cases original uncertainties have been enlarged to take into account systematic uncertainties in measurements.

The Mulder, Sinclair, Locket, Robert and Belyaev values have been shown to be outliers by the Lweight program, based on the Chauvenet's criterion. With the remaining 12 values, the weighted average is 14,284 d ; with an internal uncertainty of 0,006 d ; an external uncertainty of 0,01 and a reduced $-\chi^2$ of 2,89.

The adopted value is the weighted average : 14,284 d, with a final uncertainty expanded to include the most precise value of Coursey ((1994Co26), 14,26 (1) days) and is 0,036 d.

The large dispersion of the original set of data (reduced $-\chi^2 = 31,4$) is explained by the fact that ³²P is mainly produced by ³²S(n, γ)³²P reaction, then, resulting samples always contain ³³P as an impurity which could be not correctly taking into account.

β^- Transition transition

Evaluators calculated, with LOGFT program, a *lg ft* of 7,9 for this allowed transition. The value agrees with those suggested by Endt (1967En01, 1973En01, 1978En01 and 1990En01).

The weighted mean of the β^- end-point energy (or Q) has been calculated (with the Lweight program, version 3) using the following measured values (in keV):

Reference	Values (keV)
Lyman (1937Ly11)	1690 (24)
Newson (1937Ne19)	1590 (30)
Capron (1938Ca08)	1680 (50)
Siegbahn (1946Si07)	1712 (8)
Langer (1949La21)	1689 (10)
Marshaw (1950Ma28)	1708 (8)
Agnew (1950Ag05)	1718 (10)
Jensen (1952Je12)	1704 (8)
Antoneva (1954An18)	1712 (8)
Pohm (1956Po01)	1712 (6)
Ricci (1957Ri32)	1695 (15)
Daniel (1958Da08)	1705 (4)
Johnson (1958Jo12)	1711 (3)
Nichols (1961Ni22)	1707 (1)
Fehrentz (1961Fe15)	1705 (4)
Bosch (1963Bo36)	1706 (11)
Canthy (1966Ca31)	1697 (2)
Fishbeck (1968Fi17)	1710(2)
Flothmann (1969Fl25)	1701,2 (4)
Persson (1971Pe07)	1707 (4)
Booij (1971Bo06)	1706 (4)
Zemann (1971Ze02)	1711 (2)
Moore (1976Mo13)	1712,0 (8)
Greenwood (1993Gr10)	1710,0(30)
Kojima (2001Ko20)	1708 (2)

Evaluators calculated the weighted average of these 25 values using the Lweight program (version 3) as 1705,0 keV with an uncertainty of 3,8 and a reduced $-\chi^2$ of 9,6. The Lyman (1937Ly11), Newson (1937Ne19), Capron (1938Ca08), Langer (1949La21), Agnew (1950Ag05), Ricci (1957Ri32) and Canthy (1966Ca31) values have been shown to be outliers by the Lweight program, based on the Chauvenet's criterion. For the remaining 18 values, the weighted average is 1708,0 keV with an internal uncertainty

of 0,36 keV, an external uncertainty of 1,1 keV and a reduced $-\chi^2$ of 8,6. The final uncertainty is 7,0 keV (expanded so range includes the most precise value of Flothmann (1969F125)). This value is in agreement with the adopted Q value (1995Au04) in this evaluation.

References

- 1934Am01 - J. Ambrosen, Z. Phys. 91 (1934) 43 [Half-life].
 1935Pr20 - P. Preiswerk, H. Von Halban, Compt. Rend. 201 (1935) 722 [Half-life].
 1936Si10 - G. J. Sizoo, C. P. Koene, Physica 3 (1936) 1053 [Half-life].
 1937Ly11 - E. M. Lyman, Phys. Rev. 51 (1937) 1 [End-point energy].
 1937Ne19 - H. W. Newson, Phys. Rev. 51 (1937) 624 [Half-life, End-point energy].
 1938Ca08 - P. C. Capron, Physica 5 (1938) 882 [Half-life, End-point energy].
 1938Ca15 - N. B. Cacciapuoti, Nuovo Cimento 15 (1938) 213 [Half-life].
 1940Mu04 - D. Mulder, G. W. Hoeksema, G. J. Sizoo, Physica 7 (1940) 849 [Half-life].
 1946Si07 - K. Siegbahn, Phys. Rev. 70 (1946) 127 [End-point energy].
 1948Kl06 - E. D. Klema, A. O. Hanson, Phys. Rev. 73 (1948) 106 [Half-life].
 1949La21 - L. M. Langer, H. C. Price Jr, Phys. Rev. 76 (1949) 641 [End-point energy].
 1950Ag05 - H. M. Agnew, Phys. Rev. 77 (1950) 655 [End-point energy].
 1950Ma28 - S. D. Marshaw, J. J. L. Chen, G. L. Appleton, Phys. Rev. 80 (1950) 288 [End-point energy].
 1951Si26 - W. K. Sinclair, A. F. Holloway, Nature 167 (1951) 365 [Half-life].
 1952Je12 - E. N. Jensen, R. T. Nichols, J. Clement, A. Pohm, Phys. Rev. 85 (1952) 112 [End-point energy].
 1953Lo19 - E. E. Lockett, R. H. Thomas, Nucleonics 11 (1953) 14 [Half-life].
 1954An18 - H. M. Antoneva, Izv. Akad. Nauk. (Ser. Fiz.) 18 (1954) 93 [End-point energy].
 1956Ba25 - J. G. Bayly, Can. J. Research 28A (1956) 520 [Half-life].
 1956Po01 - A. V. Pohm, R. C. Waddell, E. N. Jensen, Phys. Rev. 101 (1956) 1315 [End-point energy].
 1957An57 - O. U. Anders, W. W. Wayne Meinke, Nucleonics 15 (1957) 68 [Half-life].
 1957Ri32 - R. A. Ricci, Physica 23 (1957) 693 [End-point energy].
 1958Da08 - H. Daniel, Nucl. Phys. 8 (1958) 191 [Half-life, End-point energy].
 1958Jo12 - O. E. Johnson, R. G. Johnson, L. M. Langer, Phys. Rev. 112 (1958) 2004 [End-point energy].
 1959Ro24 - J. Robert, Annales de Physique 4 (1959) 89 [Half-life].
 1961Ni22 - R. T. Nichols, R. E. McAdams, E. N. Jensen, Phys. Rev. 122 (1961) 172 [End-point energy].
 1961Fe15 - D. Fehrentz, H. Daniel, Nucl. Instr. Meth. 10 (1961) 185 [End-point energy].
 1961Ma01 - P. G. Marais, J. Deist, South African J. Agricultural Science 4 (1961) 627 [Half-life].
 1963Bo36 - H. E. Bosch, T. Urstein, Nucl. Instr. Meth. 24 (1963) 109 [End-point energy].
 1966Ca31 - M. J. Canty, W. F. Davidson, R. D. Connor, Nucl. Phys. 85 (1966) 317 [End-point energy].
 1966Go17 - I. W. Goodier, D. H. Pritchard, Int. J. Appl. Rad. Isotopes 17 (1966) 121 [Half-life].
 1967En01 - P. M. Endt, C. van der Leun, Nucl. Phys. A105 (1967) 1 [End-point energy, Half-life, Q, lg ft].
 1968Fi17 - H. J. Fischbeck, Phys. Rev. 173 (1968) 1078 [End-point energy].
 1969F125 - D. Flothmann, W. Wiesner, R. Lohken, H. Rebel, Z. Phys. 225 (1969) 164 [End-point energy].
 1969Pe16 - D. W. Perna, Int. J. Appl. Rad. Isotopes 20 (1969) 613 [Half-life].
 1969La28 - F. Lagoutine, J. Legrand, Y. Le Gallic, Int. J. Appl. Rad. Isotopes 20 (1969) 868 [Half-life].
 1971Bo06 - H. M. W. Booij, E. A. van Hoek, H. van der Molen, W. F. Slot, J. Blok, Nucl. Phys. A160 (1971) 337 [End-point energy].
 1971Pe07 - B. I. Persson, I. Plessner, Nucl. Phys. A167 (1971) 470 [End-point energy].
 1971Ze01 - H. Zemmann, Nucl. Phys. A175 (1971) 385 [End-point energy].
 1973En01 - P. M. Endt, C. van der Leun, Nucl. Phys. A214 (1973) 1 [End-point energy, Half-life, Q, lg ft].
 1976Mo13 - R. B. Moore, S. I. Hayakawa, D. M. Rehfield, Nucl. Instr. Meth. 133 (1976) 457 [End-point energy].
 1977Mu15 - T. S. Mudhole, Indian J. Pure and Appl. Phys. 15 (1977) 284 [Half-life].
 1977Be21 - B. N. Belyaev, S. S. Vasilenko, A. I. Egorov, A. I. Pautov, Izv. Akad. Nauk. (Ser. Fiz.) 41 (1977) 66 [Half-life].

- 1978En01 - P. M. Endt, C. van der Leun, Nucl. Phys. A310 (1978) 1 [End-point energy, Half-life, Q, lg ft].
- 1979Pr36 - J. Precker, K. Blansdorf, Atomkernenergie 34 (1979) 136 [Half-life, End-point energy].
- 1985Wa21 - A. H. Wapstra, Nucl. Phys. A432 (1985) 1 [End-point energy].
- 1990En01 - P. M. Endt, Nucl. Phys. A521 (1990) 1 [End-point energy, Half-life, Q, lg ft].
- 1993Gr10 - R. C. Greenwood, M. H. Putnam, Nucl. Instr. Meth. Phys. Res. A337 (1993) 106 [End-point energy].
- 1994Co26 - B. M. Coursey, J. M. Calhoun, J. Cessna, D. B. Golas, F. J. Schima, M. P. Unterweger, Nucl. Instr. Meth. Phys. Res. A339 (1994) 26 [Half-life].
- 1995Au04 - G. Audi, A. H. Wapstra, Nucl. Phys. A595 (1995) 409 [Q].
- 1996Sc33 - E. Schonfeld, H. Janßen, Nucl. Phys. Instr. Meth. Phys. Res. A369 (1996) 527 [Atomic data].
- 2001Ko20 - Y. Kojima, M. Shibata, H. Uno, K. Kawade, A. Taniguchi, Y. Kawase, K. Shizuma, Nucl. Instr. Meth. Phys. Res. A458 (2001) 656 [End-point energy].

³³P – Comments on evaluation of decay data by V. Chisté and M. M. Bé

1) Decay Scheme

³³P disintegrates by β^- emission (100 %) to the ground state of the stable nuclide ³³S.

2) Nuclear Data

The Q value (248,5 (11) keV) is from Audi and Wapstra evaluation (1995Au04), and has been calculated with the formula:

$$Q = M(A, Z) - M(A, Z + 1),$$

where M(A,Z) and M(A,Z+1) are the measured atomic masses of ³³P and ³³S, respectively.

Q, calculated with the formula, is in agreement with a weighted average value of 248,5 (10) keV, which the evaluators have calculated from measured values of the β^- end-point energy (see **b⁻ Transition**).

The measured ³³P half-life values (in days) are given below:

T_{1/2}

Reference	Value (days)
Sheline(1951Sh22)	25 (2)
Jensen (1952Je12)	24,8 (5)
Westermarck (1952We01)	25 (2)
Nichols (1954Ni06)	24,4 (2)
Westermarck (1954We03)	25,4 (2)
Russell (1958Ru07)	25 (1)
Fogelstrom-Fineman (1960Fo14)	25,2 (5)
Reynolds (1968Re20)	25,30 (5)
Lagoutine (1972La21)	25,56 (7)

Nichol's value (24,4 (2)) is an outlier (based on Chauvenet's criterion). The weighted average of the eight remaining values (excluding Nichol's value) is 25,383 days with an internal uncertainty of 0,040 days ($\chi^2 = 1,6$). Thus we recommend a half-life of 25,383 (40) d.

b⁻ Transition

Evaluators calculated, using the LOGFT program, a *lg ft* value of 5 for this allowed transition. This value agrees with those given by Endt (1967En01, 1973En01, 1978En01, 1990En01 and 1998En01).

The evaluators have calculated a weighted mean of the β^- end-point energy (or Q) from the following measured values (in keV):

Reference	Values (keV)
Sheline (1951Sh22)	270 (20)
Jensen (1952Je12)	260 (20)
Westermarck (1952We01)	246 (5)
Nichols (1954Ni06)	249 (2)
Elbek (1954El07)	252 (5)
Elbek (1954El08)	250 (5)
Westermarck (1954We03)	246 (5)
Russell (1958Ru07)	238 (5)
Polak (1984Po09)	248,3 (13)

Evaluators calculated the weighted average of these 9 values using the Lweight program (version 3) as 248,2 keV with an internal uncertainty of 1,0 and a reduced $-\chi^2$ of 0,87. The 2 values of Elbek (1954El07 and 1954El08) are independent measurements. The Sheline (1951Sh22), Jensen (1952Je12) and Russell (1958Ru07) values have been shown to be outliers by the Lweight program, based on the Chauvenet's criterion. For the remaining 6 values, the largest contributions to the weighted average come from the values of Polak (1984Po09), with a relative statistical weight of 59 %.

The weighted average of the six remaining input values is 248,5 keV with an internal uncertainty of 1,0 keV and a reduced $-\chi^2$ of 0,23. This value is in agreement with the adopted Q value (1995Au04) in this evaluation.

Atomic Data

Atomic values (ω_K and n_{KL}) are from (96Sc33).

References

- 1951Sh22 – P. K. Sheline, R. B. Holtzman, C. Y. Fan, Phys. Rev. 83(1951)215; Phys. Rev. 83(1951)919 [Half-life, End-point energy].
 1952Je12 - E. N. Jensen, R. T. Nichols, J. Clement, A. Pohm, Phys. Rev. 85(1952)112 [Half-life, End-point energy].
 1952We01 – T. Westermarck, Phys. Rev. 88(1952)573 [Half-life, End-point energy].
 1954Ni06 – R. T. Nichols, E. N. Jensen, Phys. Rev. 94(1954)369 [Half-life, End-point energy].
 1954El07 and 1954El08 – B. Elbek, K. O. Nielsen, O. B. Nielsen, Phys. Rev. 95(1954)96 [End-point energy].
 1954We03 – T. Westermarck, Arkiv Fysik 7(1954)87 [Half-life, End-point energy].
 1958Ru07 – J. E. Russell, Bull. Amer. Phys. Soc. 3(1958)61 [Half-life, End-point energy].
 1960Fo14 – I. Fogelstrom-Fineman, T. Westermarck, Acta Chem. Scan. 14(1960)2046 [Half-life].
 1967En01 – P. M. Endt, C. van der Leun, Nucl. Phys. A105(1967)1 [End-point energy, Half-life, Q, lg ft].
 1968Re03 – S. A. Reynolds, J. F. Emery, E. I. Wyatt, Nucl. Sci. Eng. 32(1968)46 [Half-life].
 1972La21 – F. Lagoutine, J. Legrand, C. Perrot, J. P. Brethon, J. Morel, Int. J. Appl. Rad. Isotopes 23(1972)219 [Half-life].
 1973En01 – P. M. Endt, C. van der Leun, Nucl. Phys. A214(1973)1 [End-point energy, Half-life, Q, lg ft].
 1978En01 – P. M. Endt, C. van der Leun, Nucl. Phys. A310(1978)1 [End-point energy, Half-life, Q, lg ft].
 1984Po09 – P. Polak, L. Lindner, Radiochimica Acta 35(1984)23 [End-point energy].
 1985Wa21 – A. H. Wapstra, Nucl. Phys. A432(1985)1 [End-point energy].
 1990En01 – P. M. Endt, Nucl. Phys. A521(1990)1 [End-point energy, Half-life, Q, lg ft].
 1995Au04 – G. Audi, A. H. Wapstra, Nucl. Phys. A595(1995)409 [Q].
 1996Sc33 – E. Schönfeld, H. Janßen, Nucl. Phys. Instr. Meth. Phys. Res. A369(1996)527 [Atomic data].
 1998En01 – P. M. Endt, Nucl. Phys. A633(1998)1 [End-point energy, Half-life, Q, lg ft].

**³⁵S - Comments on evaluation of decay data
by V.P. Chechev and M.M. Bé**

The first evaluation was completed by V. Chechev in 1998. It was updated in January 2012 to include the new $Q(\beta^-)$ value of 167.33 (3) keV (2011AuZZ), compared to 167.14 (8) keV, used in the original evaluation. The literature available by January 2012 was included.

Nuclear Data

Half-Life

In literature there are many measurements of the ³⁵S half-life. They are listed in Table 1.

Table 1. Measurements results and recommended value of ³⁵S half-life.

Reference	Value (d)	Uncertainty (d)	Remarks
1940Le**	88	5	GM, omitted
1941Ka**	88	3	GM, omitted
1943He**	87,1	1,2	GM
1949Ma**	88		Omitted
1952Ru23	80		Omitted
1958Se49	87,16	0,10	PC
1959Ca12	88,8	1,0	PC
1959Co56	86,35	0,17	PC
1961Wy01	89,0	0,5	PC
1961Oz01	87,1	0,9	Calorimetry
1965Fl02	87,9	0,3	PC
1968Wo06	87,39	0,10	4 π PC
1969La34	87,48	0,14	4 π PC, original Uc/3
1999Pa18	87,38	0,03	Omitted, β -spectrometer
$\chi^2/n-1 = 6,6$			
χ^2 crit. = 2,5			
UWM =	87,59		
LWM = (recommended)	87,25	0,15	Uc _{int} = 0,06 ; Uc _{ext} = 0,15

Conventional designations in the fourth column:

Measurement of counting rate decrease by Geiger-Müller counter (GM), proportional counter (PC), calorimeter (calorimetry), 4 π proportional counter (4 π PC).

The two values without uncertainty and the two oldest ones with high uncertainty were omitted from statistical analysis.

The value of Palermo *et al.* (1999Pa18) has been omitted because the measurement was carried out to check a source preparation process, only the statistical uncertainty was taken into account; in the publication, the uncertainty bars associated to each result, are significantly greater than the claimed uncertainty, moreover an impurity was observed in the source. It was then difficult to assess a real uncertainty and this value was rejected.

Decay Energy and Characteristics of Electron Emission (β^-)

The decay energy of ³⁵S has been adopted using the evaluations of Audi *et al.* (2011).

The end-point of the ³⁵S β^- -spectrum has been obtained from the correlation:

$E_{\beta^-} = Q_{\beta^-} - E_r$ where $E_r = 3$ eV is the maximum recoil energy of ³⁵Cl atom.

The average energy of the electrons per disintegration has been calculated for an allowed form of β^- - spectrum taking into account the adopted value of Q_{β^-} .

References

- | | | |
|----------|--|-------------------------------|
| 1940Le** | H. Levi, Nature 145 (1940) 588 | [T _{1/2}] |
| 1941Ka** | M.D. Kamen, Phys.Rev. 60 (1941) 537 | [T _{1/2}] |
| 1943He** | R.N. Hendricks et al., J.Phys. Chem. 47 (1943) 469 | [T _{1/2}] |
| 1949Ma** | W. Maurer, Z.Naturf. 4a (1949) 150 | [T _{1/2}] |
| 1952Ru23 | G. Rudstam, P.C. Stevenson, P.L. Folder, Phys.Rev. 87 (1952) 358 | [T _{1/2}] |
| 1958Se49 | H.H. Seliger, W.B. Mann and L.M. Cavallo, J.Research NBS 60 (1958) 447 | [T _{1/2}] |
| 1959Ca12 | J.P. Cali and L.F. Lowe, Nucleonics 17 (1959) 86 | [T _{1/2}] |
| 1959Co56 | R.D. Cooper and E.S. Cotton, Science 129 (1959) 1360 | [T _{1/2}] |
| 1961Wy01 | E.I. Wyatt et al., Nucl.Sci.Eng. 11 (1961) 74 | [T _{1/2}] |
| 1961Oz01 | Y. Ozias et al., Compt.rend. 253 (1961) 2944 | [T _{1/2}] |
| 1965Fl05 | K.F. Flynn, L.E. Glendenin and E.P. Steinberg, Nucl.Sci.Eng. 22 (1965) 416 | [T _{1/2}] |
| 1968Wo06 | E.J. Woodhouse and T.H. Norris, J.Inorg.Nucl.Chem. 301 (1968) 1373 | [T _{1/2}] |
| 1969La34 | F. Lagoutine et al., Intern.J.Appl.Rad.Isot. 20 (1969) 868 | [T _{1/2}] |
| 1999Pa16 | L.Palermo et al., Nucl. Instrum. Methods Phys. Res. A423 (1999) 337 | [T _{1/2}] |
| 2011AuZZ | G. Audi and W. Meng, private communication (2011) | [Q _{β⁻}] |

³⁶Cl - Comments on evaluation of decay data by M.-M. Bé and V.P. Chechev

This evaluation was completed in 1998, it was updated in January 2012. The literature available by this date was included. A new mean energy of the β decay is proposed.

1. Decay Scheme and total Decay Energy

³⁶Cl decay scheme is based on the measurements of Drever (1955Dr35) and Pierson (1967Pi04). The Q-values taken from Audi *et al.* (2011AuZZ) are based on many measurements.

2. Half-Life

The following values of the ³⁶Cl half-life in relation to β^- -decay to ³⁶Ar($T_{1/2\beta^-}$) presented in Table 1 have been considered .

Table 1. Results of ³⁶Cl \rightarrow ³⁶Ar decay half-life measurements

Reference NSRkeynumber	$T_{1/2}(\beta^-)$ ($10^5 a$)	Method and remarks
1949Re**	3,6	Microwave spectrometer, β G-M
1949Re**	2	Abundance by calculation from bombardment data
1949Wu15	4,4 (5)	Microwave spectrometer, β G-M
1955Ba93	3,08 (3)	Mass spectrometry, $4\pi\beta$ pc
1957Wr37	2,6 (4)	Cl(n, γ)Yield, β G-M.
1957Wr37	2,5 (4)	Mass spectrometry, β G-M. Same activity as above
1966Go07	3,10 (4)	Mass spectrometry, $4\pi\beta$ pc
1966Go07	3,06 (2)	Mass spectrometry, liq.scint. Same mass concentration as above

Wright *et al.* (1957Wr37) gave two results for ³⁶Cl half-life. The mass concentration of the samples was determined by two different methods, but the specific activity was determined only once and used to derive both half-life values, so, these values are not independent. Then, in this evaluation, the simple mean of the two results has been introduced for the statistical process.

Goldstein (1966Go07) published two results as well. However, in this work, they carried out one determination of the mass concentration and two separate measurements of the sample activity. Two half-life values were derived. Similarly, the simple mean is adopted, in this evaluation, with the highest experimental uncertainty because the author said that "he did not include any systematic error".

The values used for the statistical analysis are:

Reference	$T_{1/2}(\beta^-)$, ($10^5 a$)	
1949Wu15	4,4 (5)	
1955Ba93	3,08 (3)	
1957Wr37	2,55 (40)	
1966Go07	3,08 (4)	
χ^2 crit. = 3,8		

$\chi^2/n-1 = 2,9$		
UWM	3,28	
LWM	3,08	Int. $u_c = 0,024$; Ext. $u_c = 0,04$

The adopted value is: $3,08 (4) 10^5 a$.

From the basic relations:

$$T_{1/2} = \ln 2 / (\lambda_{ec} + \lambda_{\beta^-}) \text{ and } \lambda_{ec}/\lambda_{\beta^-} = P_{ec}/P_{\beta^-}$$

the total half-life of ³⁶Cl is obtained $T_{1/2}(\beta^-) \times 0,981 (1) = 3,02 (4) 10^5 a$.

3. Electron Capture

An experimental value of the ratio $(P_L/P_K)_{exp} = 0,112 (8)$ was measured in 1962Do07, and a theoretical value $(P_L/P_K)_T$ was calculated, assuming an allowed transition or 1st forbidden, from the tables of Schönfeld to be 0,095 (5) and, 0,0944 by using the LOGFT program. Theoretical and measured values are not consistent.

On one hand, there is only one measured value; on the other hand, this transition was shown to be of a non-unique second forbidden order.

However, the energy involved is high (1142 keV) so the difference between a 1st and a 2nd forbidden transition should be expected to be low. Then, the theoretical results are preferred and a conservative uncertainty of 5 % was applied.

The probability of the electron capture $P_{EC} = 1,9 (1) \%$ was deduced from the measured ratio $P_{EK}/P_{\beta^-} = 0,017 (1)$ (1955Dr35) and $P_{EK} = 0,904 (5) \times P_{EC}$.

4. β^+ Transition

The probability $P_{\beta^+} = 1,57 (30) 10^{-3} \%$ has been obtained by averaging the experimental data shown in Table 3.

Table 3. Measurement results for the probability of ³⁶Cl β^+ -decay (P_{β^+}), per 100 ³⁶Cl disintegrations.

Reference	$P_{\beta^+} (10^{-3}) \%$	Remarks
1962Do07	1,2 (5)	$P_{\beta^+} = 1,7 (1) \times 7(+3-1) 10^{-4} = 1,2 (+0,5-0,2) 10^{-3} \%$. The greatest uncertainty is adopted.
1962Be29, 1963Be38	2,3 (9)	
1965To05	1	No uncertainty, omitted
1967Pi04	1,66 (11)	Uncertainty increased to 0,40 to limit its weight
Recommended value	1,57 (30)	Reduced $\chi^2 = 0,6$; crit $\chi^2 = 4,6$

The recommended value P_{β^+} has been obtained by using the LWM procedure with the three input values from (1962Do07, 1963Be38 and 1967Pi04). The set of data is consistent then, the adopted value is the weighted mean with the internal uncertainty.

5. β^- Transition

The probability $P_{\beta^-} = 98,1 (1) \%$ was calculated from the balance relation:

$$P_{\beta^-} = 1 - P_{EC} - P_{\beta^+}$$

6. Atomic Data

The atomic constants ω_K , n_{KL} and relative emission probabilities of K X-ray components and K-Auger have been taken from Schönfeld (1996Sc06).

The energy values for Auger electrons have been taken from Larkins (1977La19).

7. Photon Emissions

The emission probabilities of K X-ray components and K-Auger electrons in sulfur were derived from the probability of the electron capture P_{EC} and the adopted values P_K and ω_K .

The emission probabilities of K X-ray components and Auger electrons of argon due to K-shell auto-ionization have been calculated using $P_{XK}(Ar)/P_{XK}(S) = 0,149$ (22) from 1976Lj03 and atomic constants.

The number of photons per 100 disintegrations for the annihilation radiation was deduced from the P_{β^+} value as determined above.

8. Beta Emissions

The end-point of the ³⁶Cl β^- spectrum has been obtained from $E_{\beta^-} = Q_{\beta^-} - E_r$ where $E_r = 18$ eV is the maximum recoil energy of the ³⁶Ar atom.

The end-point energy of the ³⁶Cl β^+ spectrum has been calculated as $E_{\beta^+} = Q_{\beta^+} - E_r - 1022,0$ keV, where $E_r = 2$ eV is the maximum recoil energy of ³⁶S atom.

Several papers report measurements or calculations of the ³⁶Cl β^- spectrum (see as examples: 1949Wu18, 1972Ma72, 1974Re11, 1993Sa24, 2004Kr10, 2005Gr41, 2008Ro31, etc.).

When using the program BetaShape (2011MoZU), which is based on theoretical considerations, for the mean β^- energy, we obtain:

- with the hypothesis of an allowed transition: 251 keV,
- with the hypothesis of a 2nd non-unique forbidden order, accepted as 1st unique forbidden: 278 keV,
- with the hypothesis of a 2nd non-unique forbidden order, accepted as 2st unique forbidden: 303 keV.

Recently, Rotzinger *et al.* (2008Ro31) measured the ³⁶Cl β^- spectrum by mean of a magnetic calorimeter, they kindly provided us their original recorded data, from which a mean β^- energy value of 316 keV has been derived.

This latter value is adopted, an uncertainty of 5 % is supposed, however it is difficult to estimate it correctly.

The β^+ transition is also of a 2nd non-unique forbidden order. Similarly the mean β^+ energy was calculated as 50 keV for an allowed transition and 58 keV for a 1st unique forbidden transition by the BetaShape program. The adopted value is 54 keV with an uncertainty which covers both hypothesis.

References

- 1949Re** S.A. Reynolds. Report ORNL 286 (1949) 219 [$T_{1/2}$]
- 1949Wu15 C.S. Wu, C.N. Townes, and L. Feldman, Phys.Rev. 76 (1949) 692 [$T_{1/2}$]
- 1949Wu18 C.S. Wu, L. Feldman. Phys.Rev. 76 (1949) 693 [β^- spectrum]
- 1955Ba93 R.M. Bartholomew et al., Can.J.Phys. 33 (1955) 43 [$T_{1/2}$]
- 1955Dr35 R.W.P. Drever, and A. Moljk, Phil.Mag. 46 (1955) 1337 [$P_{\epsilon_K}/P_{\beta^-}$]
- 1957Wr37 H.W. Wright et al., Nucl.Sci.Eng. 2 (1957) 427 [$T_{1/2}$]
- 1962Do07 P.W. Dougan et al., Phil.Mag. 7 (1962) 1223 [P_L/P_K]
- 1962Be29 D. Berenyi, Phys.Lett. 2 (1962) 332.
- 1963Ba38 D. Berenyi, Acta Phys.Acad.Sci.Hungary 16 (1963) 101. [P_{β^+}]
- 1965To05 W.R. Tolbert et al., Bull.Amer.Phys.Soc. 10 (1965) 589. [P_{β^+}]
- 1966Go07 G. Goldstein, J. Inorg.Nucl.Chem. 28 (1966) 937 [$T_{1/2}$]
- 1967Pi04 W.R. Pierson, Phys.Rev. 159 (1967) 951 [P_{β^+}]
- 1972Dz09 B.S. Dzhelepov, L.N. Zyryanova and Yu.P. Suslov. Beta Processes. Functions for Analysis of Beta Spectra and Electron Capture, Leningrad, "Nauka" Press, 1972 [$P_L/P_K, <E_{\beta^+}>$]
- 1972Ma72 J. Mantel, Int. J. Appl. Radiat. Isotop. 23 (1972) 407 [β^- spectrum]
- 1974Re11 M. Reich, H.M. Schupferling, Z.Phys. 271 (1974) 107. [β^- spectrum]
- 1976Lj03 A. Ljubicic, M. Jurcevicic, M. Vlatkovic, V.A. Logan, Phys.Rev. C13 (1976) 881 [$P_K(Ar), XK(Ar)/XK(S)$]

- 1977La19 F.P. Larkins, ADNDT 20 (1977) 313 [E (e_{AK}), I (e_{AK})]
- 1993Sa24 R. Sadler, H. Behrens, Z.Phys. A346 (1993) 25. [β^- spectrum]
- 1995ScZY E. Schönfeld, Report PTB-6.33-95-2 (1995) [P_K, P_L, P_M, P_N]
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl.Instr.Meth. Phys.Res. A369 (1996) 527 [P(K _{β})/P(K _{α}), ω_K , n_{KL}]
- 2004Kr10 A.A.Kriss, D.M.Hamby, Nucl.Instrum.Methods Phys.Res. A525, 553 (2004) [β^- spectrum]
- 2005Gr41 A. Grau Carles. Nucl.Instr.Meth. Phys.Res. A551 (2005) 312. [β^- spectrum]
- 2008Ro31 H. Rotzinger *et al.* J. Low Temp. Phys. 151 (2008) 1087. [β^- spectrum]
- 2011AuZZ G. Audi, W. Meng, <http://amdc.in2p3.fr/masstable/Ame2011int/mass.mas114> [Q _{β^-} , Q _{ϵ}]
- 2011MoZU X. Mougeot, M.-M. Bé ,V. Chisté, C. Dulieu, V. Gorozhankin, M. Loidl. Calculation of beta spectra for allowed and unique forbidden transitions. (2011) LSC 2010, advances in liquid scintillation spectrometry, 6-10 September 2010, p. 249, RadioCarbon, ISBN 978 0 9638314 7 7. [$\langle E_\beta \rangle$]

³⁷Ar - Comments on evaluation of decay data by V.P. Chechev

Evaluated in March 2012 with a literature cut-off by the same date.

1. Decay Scheme and Decay Energy

³⁷Ar disintegrates by 100 % electron capture (EC) transition to the ground state of the stable nuclide ³⁷Cl. The decay scheme is complete as there are no excited levels of ³⁷Cl below the EC decay energy Q^+ (1998En04).

Q^+ value has been taken from atomic mass adjustment (2003Au03, 2011AuZZ) based on the precise spectrometric measurement of the Q-value for (p, n) reaction on ³⁷Cl (1998Bo30).

2. Half-Life

The following values of the ³⁷Ar half-life presented in Table 1 were considered.

Table 1. Results of ³⁷Ar half-life measurements (in days)

Reference	Author(s)	Value	Comments
1944We**	Weimer et al.	34.1 (3)	Omitted; ³⁹ K (d,α), ³⁷ Cl (d,2n) and ³⁷ Cl (p,n), ionization chamber, t = 7 half-lives, possible source impurities, uncertainty strongly underestimated in an unknown amount
1952Mi**	Miskel and Perlman	35.0 (4)	Omitted; ⁴⁰ Ca (n,α), proportional counter, no details, possible source impurities
1959Ki41	Kiser and Johnston	34.30 (14)	Omitted; ⁴⁰ Ca (n,α), K Auger peak decay, proportional counter, t = 2 half-lives, possible source impurities, uncertainty strongly underestimated in an unknown amount
1965St09	Stoenner et al.	35.1 (1)	⁴⁰ Ca (n,α), 5 proportional counters, t = 5 half-lives, declared uncertainty includes possible systematic errors
1973Co26	Colomer and Gauvain	35.06 (9)	³⁷ Cl (p,n), 2 proportional counters, t = 5 and 3 half-lives; the authors reported the uncertainty of 0.18 d for 95% confidence level
1975Ki10	Kishore et al.	35.02 (2)	³⁷ Cl (p,n), several proportional counters, numerous decay curves (over 1 to 3 half-lives), 28 measurements
2001Re01	Renne and Norman	34.95 (4)	⁴⁰ Ca (n,α) and ⁴⁰ Ca (n,nα), ³⁷ Ar/ ³⁶ Ar mass spectrometry; the authors reported the uncertainty of 0.08 d for 95% confidence level

The four input values (from 1965St09, 1973Co26, 1975Ki10, and 2001Re01) have been adopted for the statistical processing. The weighted average for this data set is 35.011 with an internal uncertainty of 0.017 and external uncertainty of 0.019 ($\chi^2/\nu = 1.21$). The smallest experimental uncertainty reported is 0.02.

The recommended value of the ³⁷Ar half-life is **35.01 (2) days**.

3. Electron Capture

The energy of the electron-capture transition $3/2+ \rightarrow 3/2+$ in the decay of ³⁷Ar \rightarrow ³⁷Cl is equal to the adopted Q^+ value.

K, L, M-electron capture probabilities P_K , P_L , P_M were deduced using the EC-CAPTURE computer program (1998Sc28). It should be noted that according to theory L- and M-electron captures occur basically on L1 and M1 subshells (1972Dz09). Log ft value was calculated for the allowed electron-capture transition using the LOGFT computer code (NNDC Tools and Publications, Web programming: M. Emeric and A. Sonzogni, NNDC, Brookhaven National Laboratory).

4. Atomic Data, X-Ray and Auger Electron Emissions

Atomic values, ω_K , ω_L and n_{KL} are from Schönfeld and Janßen (1996Sc06).

The X-ray and Auger electron emission probabilities were calculated using the program EMISSION (2000Sc47). The calculation results including average energies per disintegration are given in Tables 2, 3.

Table 2. KX- and LX- rays in decay of ³⁷Ar

	Energy, keV	Number of photons per disintegration	Energy per disintegration, keV
X $K\alpha_2$	2.6208	0.0276 (7)	0.0723 (18)
X $K\alpha_1$	2.6224	0.0546 (14)	0.143 (4)
X $K\beta$	2.8156	0.0071 (4)	0.020 (1)
X $L\beta$	0.240	0.0020 (4)	0.00048 (10)
Total			0.236 (5)

Table 3. Auger electrons in decay of ³⁷Ar

	Energy, keV	Number of electrons per disintegration	Energy per disintegration, keV
L-Auger	0.17-0.26	1.665 (8)	0.38 (4)
K-LL	2.31 (7)	0.689 (6)	1.59 (5)
K-LM	2.57 (4)	0.119 (6)	0.306 (5)
K-MM	2.8	0.0051 (5)	0.014 (2)
Total			2.29 (6)

5. Internal Bremsstrahlung

The characteristics of the internal bremsstrahlung in the ³⁷Ar decay were calculated in 1986BrZQ (Table 4). The results of that calculation were taken for the evaluation of the average energy E_{IB} of internal bremsstrahlung per disintegration of ³⁷Ar: $E_{IB} = 0.139 (14)$ keV (2011Ch65). The relative uncertainty of E_{IB} was set as 10 % based on that the theoretical estimation of the internal bremsstrahlung intensity for allowed electron capture transitions (1972Dz09) agrees with experimental data within less than 15 %.

Table 4. Internal bremsstrahlung in decay of ³⁷Ar

Energy (keV)	Number of photons per disintegration	Energy (keV) per disintegration
10-20	$1.31 \cdot 10^{-6}$	$2.0 \cdot 10^{-5}$
20-40	$3.8 \cdot 10^{-6}$	$1.17 \cdot 10^{-4}$
40-100	$2.4 \cdot 10^{-5}$	$1.8 \cdot 10^{-3}$
100-300	$1.63 \cdot 10^{-4}$	0.034
300-600	$2.0 \cdot 10^{-4}$	0.087
600-814	$2.5 \cdot 10^{-5}$	0.0164
Total		0.139 (14)

6. Total energy of non-neutrino radiation per disintegration

Decay of ³⁷Ar is accompanied by emissions of X-rays, internal bremsstrahlung, Auger electrons and monochromatic neutrinos with energies of 811.05 (20), 813.60 (20), and 813.85 (20) keV. The neutrino energies were deduced from the value of decay energy $Q^+ = 813.87$ (20) keV (see Section 1) and the values of *K*, *L*, *M*-electron binding energies (Cl, *Z* = 17) (1977La19).

The total energy of non-neutrino radiation per disintegration releasing in the form of (*X* + Auger) emissions and internal bremsstrahlung is obtained from Tables 2–4 of 2.67 (7) keV. This value was specified using the calculation of (*X* + Auger) - energy releasing in each act of ³⁷Ar electron capture by the relation of $E = W_K P_K + W_L P_L + W_M P_M$, where W_J - binding energy of electron in *J*-shell, P_J - probability of electron capture in *J*-shell, *J* = *K*, *L*, *M*. The more accurate value of the energy of non-neutrino radiation per disintegration is 2.709 (16) keV per disintegration (2011Ch65).

7. References

- 1944We**** P.K. Weimer, J.D. Kurbatov, and M.L. Pool, Phys. Rev. 66 (1944) 209.
[Half-life]
- 1952Mi**** J.A. Miskel and M.L. Perlman, Phys. Rev. 87 (1952) 543.
[Half-life]
- 1959Ki41** R.W. Kiser and W.N. Johnston, J. Amer. Chem. Soc. 81 (1959) 1810.
[Half-life]
- 1965St09** R.W. Stoenner, O.A. Schaeffer, and S. Katoff, Science 148 (1965) 1325.
[Half-life]
- 1972Dz09** B.S. Dzhelepov, L.N. Zyryanova, Y.P. Suslov, Beta Processes. Functions for Analysis of Beta Spectra and Electron Capture, 'Nauka' Press, Leningrad (1972).
[Theoretical L- and M- electron captures]
- 1973Co26** J. Colomer and D. Gauvain, Int. J. Appl. Rad. Isot. 24 (1973) 391.
[Half-life]
- 1975Ki10** R. Kishore, R. Colle, S. Katoff, and J.B. Cumming, Phys. Rev. C12 (1975) 21.
[Half-life]

- 1977La19** F.P. Larkins, Atomic Data and Nuclear Data Tables 20 (1977) 313.
[Atomic electron binding energies]
- 1986BrZQ** E. Browne and R.B. Firestone, Table of Radioactive Isotopes. Ed. V.S. Shirley.
N.Y.: John Wiley and Sons (1986).
[Internal Bremsstrahlung]
- 1996Sc06** E. Schönfeld and H. Janßen, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527.
[Atomic Data]
- 1998Bo30** R. Bottger and H. Scholermann, Nucl.Phys. A642, 419 (1998)
[Q value]
- 1998En04** P.M. Endt, Nucl.Phys. A633, 1 (1998)
[³⁷Cl levels]
- 1998Sc28** E. Schönfeld, Appl. Radiat. Isot. 49 (1998) 1353.
[Calculation of P_K, P_L, P_M]
- 2000Sc47** E. Schönfeld and H. Janßen, Appl. Radiat. Isot. 52 (2000) 595.
[Calculation of X-ray and Auger electron emission probabilities]
- 2001Re01** P.R. Renne and E.B. Norman, Phys. Rev. C63 (2001) 047302.
[Half-life]
- 2003Au03** G. Audi, A.H. Wapstra, and C. Thibault, Nucl. Phys. A729, 337 (2003)
[Q value]
- 2011AuZZ** G. Audi, W. Meng, <http://amdc.in2p3.fr/masstable/Ame2011int/mass.mas114>
[Q value]
- 2011Ch65** V.P. Chechev, Phys. Atomic Nuclei 74 (2011) 1713
[Total energy of non-neutrino radiation per disintegration]

⁴⁰K - Comments on evaluation of decay data by X. Mougeot, R.G. Helmer

The initial evaluation was completed in 1998. This revised evaluation was done in 2009, taking into account the available literature by April 2009.

1 Decay Scheme

The decay scheme is complete since all of the levels in ⁴⁰Ar and ⁴⁰Ca below the decay energies are populated.

The J^π and half-life of the excited level are from 1990EN08 evaluation.

2 Nuclear Data

Q values are from Audi and Wapstra 2003 (2003AU03).

A full list of the half-life measurements available by April 2009, and the reasons why certain have been excluded by the evaluator, is given in Table 3.

Three types of measurements were carried out: $T_{1/2}(\beta^-)$ and $T_{1/2}(\text{EC}, 1460 \text{ keV})$ which are partial half-lives, and $T_{1/2}$ which is the total half-life. Branching ratios are needed to evaluate the ⁴⁰K half-life from these measurements: P_{β^-} for the ⁴⁰K→⁴⁰Ca transition, $P_{\text{ec},1460}$ for the ⁴⁰K→⁴⁰Ar²⁺(1460 keV) transition, P_{β^+} and $P_{\text{ec,gs}}$ for the ⁴⁰K→⁴⁰Ar⁰⁺(ground state) transition. So, $T_{1/2}(\beta^-)$ and $T_{1/2}(\text{EC}, 1460 \text{ keV})$ have been evaluated first and then, the branching ratios and the ⁴⁰K total half-life.

2.1 Partial half-lives

2.1.1 $T_{1/2}(\beta^-)$

Table 1: Partial measured β^- half-lives.

Reference	Partial $T_{1/2}(\beta^-)$ ($\times 10^9$ a)	Comments
1948Graf	1.48 (7)	
1948Hirzel	1.18 (19)	Excluded by LWEIGHT (Chauvenet's criterion)
1949Stout	1.29 (8)	
1950Smaller	1.76 (5)	Excluded by LWEIGHT (3σ criterion)
1951Delaney	1.24 (1)	Excluded by LWEIGHT (Chauvenet's criterion)
1951Good	1.46 (3)	
1955SU38	1.34 (3)	
1955KO21	1.36 (5)	
1956MC20	1.44 (1)	
1959KE26	1.46 (3)	
1960SA31	1.37 (4)	
1961GL07	1.400 (15)	
1962FL05	1.45 (40)	
1965BR25	1.36 (2)	
1965LE15	1.400 (2)	Uncertainty increased to 6.4×10^6 a by LWEIGHT
1966FE09	1.41 (2)	
1966Egelkraut	1.40 (7)	
1971Venkataramaiah	1.31 (6)	

The statistical processing was done using the LWEIGHT program. For $T_{1/2}(\beta^-)$, the program turned up three statistical outliers: 1948Hirzel (Chauvenet's criterion), 1950Smaller (3σ criterion), and 1951Delaney (Chauvenet's criterion). From the resulting discrepant data set, with a reduced- χ^2 value of 2.62, a weighted average was deduced. LWEIGHT increased the uncertainty of the most precise measurement (1965LE15) from 2 to 6.4×10^6 a in order to have a maximum contribution of 50 %. The second main contribution is 1956MC20 amounting for 20 %. Finally, this evaluation leads to:

$$T_{1/2}(b^-) = 1.407 (7) \times 10^9 \text{ a.}$$

2.1.2 $T_{1/2}(\text{EC}, 1460 \text{ keV})$

Table 2: Partial measured EC half-lives.

Reference	Partial $T_{1/2}(\text{EC}, 1460)$ ($\times 10^9$ a)	Comments
1947GL07	11 (2)	Excluded by LWEIGHT (Chauvenet's criterion)
1948Ahrens	11.6 (2)	
1950Sawyer	12 (1)	
1950Graf	12 (2)	
1953BU58	11.7 (5)	
1955SU38	13.4 (2)	
1955BA25	11.3 (5)	
1957WE43	11.7 (4)	
1960SA31	12.3 (6)	
1965LE15	12.2 (3)	
1966DeRuytter	12.2 (2)	
1966Egelkraut	11.8 (5)	

For the electronic capture (EC) part, all the partial half-lives, given in Table 2, were measured by detecting the 1460 keV gamma-ray in ⁴⁰Ar. In Table 3, a partial half-life for EC is listed, evaluated by 1956Wetherill: this evaluation used four measurements of the ⁴⁰Ar/⁴⁰K concentration ratio in young mica. Obviously, in this case, the total branching ratio of the ⁴⁰K → ⁴⁰Ar was determined. So, this result cannot be used to evaluate the partial $T_{1/2}(\text{EC}, 1460 \text{ keV})$.

The statistical processing was done using the LWEIGHT program. It turned up two statistical outliers: 1947GL07 and 1955SU38 (Chauvenet's criterion). A weighted average was adopted from the resulting consistent data set, with a reduced- χ^2 value of 0.87. The main contributions are 30 % for 1966DeRuytter and 1948Ahrens, and 13 % for 1965LE15. Finally, this evaluation gives:

$$T_{1/2}(\text{EC}, 1460 \text{ keV}) = 11.90 (11) \times 10^9 \text{ a.}$$

2.2 Branching ratios

The branching ratios were calculated following Helmer's method (1999BeZS). From the decay scheme:

$$P_{\text{ec},1460} + P_{b^+} + P_{b^-} + P_{\text{ec,gs}} = 1.$$

In order to calculate each branching ratio, the following quantities: $P_{\text{ec},1460}/P_{\beta^-}$, P_{b^+}/P_{β^-} and $P_{\text{ec,gs}}/P_{\beta^+}$ must be known.

The $P_{\text{ec},1460}/P_{\beta^-}$ ratio comes from the $T_{1/2}(\beta^-)/T_{1/2}(\text{EC}, 1460 \text{ keV})$ ratio. The partial half-lives evaluated above leads to: $P_{\text{ec},1460}/P_{\beta^-} = 0.1182 (12)$.

The β^+ transition of the ⁴⁰K is a difficult measurement, due to a very low intensity and the pair production which comes from the 1460 keV gamma-ray of ⁴⁰Ar. Few experiments were able to give more than an upper limit: 1959TI20 ($1.3 (7) \times 10^{-5}$), 1962EN01 ($1.12 (14) \times 10^{-5}$) and 1965LE15 ($1.5 (5) \times 10^{-5}$). The experimental set-up of 1962EN01 minimized the pair production. Following Helmer's choice, the most precise result is used in the present evaluation: $P_{b^+}/P_{\beta^-} = 1.12 (14) \times 10^{-5}$.

The $P_{ec,gs}/P_{\beta^+}$ ratio was calculated theoretically by Helmer, as described hereafter. The LOGFT program cannot calculate this ratio for this unique 3rd forbidden (3U) transition. But it can calculate the theoretical value for 1U and 2U transitions. For the former (1U), this ratio is 8.51 (9) and for the latter (2U), it is 45.20 (47). Making the assumption that the 3U ratio rises by the same factor (45.20/8.51), then $P_{ec,gs}/P_{\beta^+} = 240$. Following Helmer's choice, a value of **200 (100)** for $P_{ec,gs}/P_{\beta^+}$ was adopted in the present calculation.

The following branching ratios are then deduced:

$$P_{\beta^-} = 89.25 (17) \%, P_{ec,1460} = 10.55 (11) \%, P_{ec,gs} = 0.20 (10) \%, P_{\beta^+} = 0.00100 (12) \%$$

2.3 Total ⁴⁰K half-life

Table 3: Total half-lives used for the evaluation, determined from measurements and branching ratios.

Reference	Type of measurement	T _{1/2} (×10 ⁹ a)	Coefficient (%)	Total T _{1/2} (×10 ⁹ a)	Comments
1931Orban	Partial, EC 1460	0.5	-	-	Not used : no uncertainty
1947GL07	Partial, EC 1460	11 (2)	10.55 (11)	1.16 (21)	
1948Ahrens	Partial, EC 1460	11.6 (2)	10.55 (11)	1.224 (25)	
1948Graf	Partial, β ⁻	1.48 (7)	89.25 (17)	1.32 (6)	
1948Hirzel	Partial, β ⁻	1.18 (19)	89.25 (17)	1.05 (17)	Excluded by LWEIGHT (Chauvenet's criterion)
1949Stout	Partial, β ⁻	1.29 (8)	89.25 (17)	1.15 (7)	
1949Floyd	Total	1.54 (39)	100	1.54 (39)	Excluded by LWEIGHT (Chauvenet's criterion)
1950Sawyer	Partial, EC 1460	12 (1)	10.55 (11)	1.27 (11)	
1950Graf	Partial, EC 1460	12 (2)	10.55 (11)	1.27 (21)	
1950Faust	Total	1.14 (10)	100	1.14 (10)	
1950SA52	Total	1.27 (5)	100	1.27 (5)	
1950Spiers	Total	1.18	-	-	Not used : no uncertainty
1950Houtermans	Total	1.31 (7)	100	1.31 (7)	
1950Smaller	Partial, β ⁻	1.76 (5)	89.25 (17)	1.571 (45)	Excluded by LWEIGHT (3σ criterion)
1951Delaney	Partial, β ⁻	1.24 (1)	89.25 (17)	1.107 (9)	Excluded by LWEIGHT (Chauvenet's criterion)
1951Good	Partial, β ⁻	1.46 (3)	89.25 (17)	1.303 (27)	
1953BU58	Partial, EC 1460	11.7 (5)	10.55 (11)	1.23 (5)	
1955SU38	Partial, β ⁻	1.34 (3)	89.25 (17)	1.196 (27)	
1955SU38	Partial, EC 1460	13.4 (2)	10.55 (11)	1.414 (26)	Excluded by LWEIGHT (Chauvenet's criterion)
1955KO21	Partial, β ⁻	1.36 (5)	89.25 (17)	1.214 (45)	
1955BA25	Partial, EC 1460	11.3 (5)	10.55 (11)	1.19 (5)	
1956MC20	Partial, β ⁻	1.44 (1)	89.25 (17)	1.285 (9)	
1956Wetherill	Partial, EC and β ⁺	12.2 (6)	10.75 (15)	1.31 (7)	⁴⁰ Ar/ ⁴⁰ K in young mica
1957WE43	Partial, EC 1460	11.7 (4)	10.55 (11)	1.234 (44)	Direct measurement
1959KE26	Partial, β ⁻	1.46 (3)	89.25 (17)	1.303 (27)	
1960SA31	Partial, EC 1460	12.3 (6)	10.55 (11)	1.30 (6)	
1960SA31	Partial, β ⁻	1.37 (4)	89.25 (17)	1.223 (36)	
1961GL07	Partial, β ⁻	1.400 (15)	89.25 (17)	1.249 (14)	
1962FL05	Partial, β ⁻	1.45 (40)	89.25 (17)	1.29 (36)	
1965BR25	Partial, β ⁻	1.36 (2)	89.25 (17)	1.214 (18)	
1965LE15	Partial, EC 1460	12.2 (3)	10.55 (11)	1.287 (34)	
1965LE15	Partial, β ⁻	1.400 (2)	89.25 (17)	1.2495 (30)	


Reference	Type of measurement	T _{1/2} (×10 ⁹ a)	Coefficient (%)	Total T _{1/2} (×10 ⁹ a)	Comments
1966FE09	Partial, β-	1.41 (2)	89.25 (17)	1.258 (18)	Not used : erroneous uncertainty, see also 2001BE81
1966DeRuytter	Partial, EC 1460	12.2 (2)	10.55 (11)	1.287 (25)	
1966Egelkraut	Partial, EC 1460	11.8 (5)	10.55 (11)	1.24 (5)	
1966Egelkraut	Partial, β-	1.40 (7)	89.25 (17)	1.25 (6)	
1971Venkataramaiah	Partial, β-	1.31 (6)	89.25 (17)	1.17 (5)	
1972Gopal	Partial, β-	1.13 (6)	-	-	
1977CE04	Partial, EC 1460	12.30 (4)	-	-	Not used : erroneous uncertainty, see also 2001BE81
2004KO09	Total	1.248 (3)	100	1.2480 (30)	

In order to evaluate the ⁴⁰K half-life, each partial half-life was recalculated using the appropriate branching ratio. The corresponding uncertainty was also calculated.

The LWEIGHT program turned up five statistical outliers: four by Chauvenet’s criterion (1948Hirzel, 1949Floyd, 1951Delaney, 1955SU38 (EC, 1460)) and one by 3σ criterion (1950Smaller). A weighted average was adopted from the resulting consistent data set, with a reduced-χ² value of 1.62. The data used for the evaluation of the ⁴⁰K half-life can be seen in Figure 1. The two main contributions come from 1965LE15 (β-) and 2004KO09, each of them amounting by 43 %. The adopted value is: T_{1/2} = 1.2504 (25) × 10⁹ a. Since these measurements are not all independent, the adopted uncertainty is the most precise uncertainty on measurement: 3.0 × 10⁶ a, identical for 1965LE15 (β-) and 2004KO09.

The recommended value for the ⁴⁰K half-life is then: **T_{1/2} = 1.2504 (30) × 10⁹ a**, in good agreement with the evaluations by Helmer (1.265 (13) × 10⁹ a) (1999BeZS) and Chechev (1.258 (10) × 10⁹ a) (2001Chechev).

Figure 1: T_{1/2} measurements used for the present evaluation, recalculated with the branching ratios. The red ones are excluded by LWEIGHT.



2.4 Electron Capture Transitions

The evaluation of the branching ratios is described in Section 2.2. That is:

$$P_{ec,1460} = 10.55 \text{ (11) \%} \text{ and } P_{ec,gs} = 0.20 \text{ (10) \%}.$$

The $\log ft$ value for the 1U transition ($^{40}\text{K} \rightarrow ^{40}\text{Ar}^{2+}$) was computed using the LOGFT program:

$$\log ft = 11.55 \text{ (1)}.$$

LOGFT cannot calculate the $\log ft$ value for the 3U transition ($^{40}\text{K} \rightarrow ^{40}\text{Ar}^{gs}$). The evaluator chose the same method used in Section 2.2 to calculate the $P_{ec,gs}/\beta^+$ ratio.

$$\text{So, } \log ft \text{ (1U)} = 19.51 \text{ (5)} \text{ and } \log ft \text{ (2U)} = 20.41 \text{ (5)} \text{ and then, } \log ft \text{ (3U)} = 21.35 \text{ (10)}.$$

The P_K , etc. values were computed by the LOGFT program.

2.5 β^- Transitions

The β^- branching ratio is 89.25 (17) %, as deduced in Section 2.2. The average energy is from the LOGFT program.

The $\log ft$ value for this 3U transition ($^{40}\text{K} \rightarrow ^{40}\text{Ca}$) is calculated with the same method as previously, then $\log ft \text{ (3U)} = 20.58 \text{ (1)}$.

2.6 Gamma Transitions

The internal conversion coefficients were calculated using the BrIcc program (2008KI07) for the K, L and M shells. The total internal conversion coefficient is: $\alpha = 10.28 \text{ (15)} \times 10^{-5}$.

From the theoretical tables of 1979SC31, the internal pair formation coefficient is:

$$\alpha_{\pi}(1460, \text{E2}) = 7.3 \text{ (5)} \times 10^{-5}.$$

$$\text{So: } \alpha_T = \alpha + \alpha_{\pi}(1460, \text{E2}) = 17.6 \text{ (5)} \times 10^{-5}$$

3 Atomic Data (Ar, Z=10)

3.1 X Radiations and Auger electrons

The X-ray and Auger electron data were computed using the EMISSION program with the atomic data of Schönfeld and Janßen (1996SC06).

4 Radiation Emissions

4.1 Electron Emission

The β^+ and β^- intensities were evaluated as described above in Section 2.

4.2 Photon Emissions

No new measurement was carried out for the 1460 keV gamma-ray energy in ^{40}Ar since 1998. The adopted value was evaluated by Helmer (1999BeZS): $E_{\gamma} = 1460.822 \text{ (6) keV}$.

The gamma emission intensity is deduced from the electronic capture probability (see Section 2.2) and internal conversion coefficient (see Section 2.6):

$$I_{\gamma}(1460) = P_{EC}(1460) / [1 + \alpha_T] = 10.55 \text{ (11)} / 1.000176 \text{ (5) \%}.$$

So we have:

$$I_{\gamma}(1460) = 10.55 \text{ (11) \%}.$$

5. References

- 1931Orban - G. Orban, Sitzb. Akad. Wiss. Wien Abt. IIa, 140, 121 (1931) [$T_{1/2}$ EC]
 1947GL07 - E. Gleditsch, T. Graf, Phys. Rev. 72, 640 (1947) [$T_{1/2}$ EC]
 1948Ahrens - L.H. Ahrens, R.D. Evans, Phys. Rev. 74, 279 (1948) [$T_{1/2}$ EC]
 1948Graf - T. Graf, Phys. Rev. 74, 831 (1948) [$T_{1/2}$ β^-]
 1948Hirzel - O. Hirzel, H. Wäffel, Phys. Rev. 74 (1948) 1553 [$T_{1/2}$ β^-]

- 1949Stout - R.W. Stout, Phys. Rev. 75, 1107 (1949) [$T_{1/2}$ β^-]
- 1949Floyd - J.J. Floyd, L.B. Borst, Phys. Rev. 75, 1106 (1949) [$T_{1/2}$ Total]
- 1950Sawyer - G.A. Sawyer, M.L. Wiedenbeck, Phys. Rev. 76, 1535 (1950) [$T_{1/2}$ EC]
- 1950Graf - T. Graf, Rev. Sci. Inst. 21, 285 (1950) [$T_{1/2}$ EC]
- 1950Spiers - F.W. Spiers, Nature 165, 356 (1950) [$T_{1/2}$ Total]
- 1950Faust - W.R. Faust, Phys. Rev. 78, 624 (1950) [$T_{1/2}$ Total]
- 1950SA52 - G.A. Sawyer, M.L. Wiedenbeck, Phys. Rev. 79, 490 (1950) [$T_{1/2}$ Total]
- 1950Houtermans - F.G. Houtermans, O. Haxel, J. Heintze, Z. Physik 128, 657 (1950) [$T_{1/2}$ Total]
- 1950Smaller - B. Smaller, J. May, M. Freedman, Phys. Rev. 79, 940 (1950) [$T_{1/2}$ β^-]
- 1951Delaney - C.F.G. Delaney, Phys. Rev. 81, 158 (1951) [$T_{1/2}$ β^-]
- 1951Good - M.L. Good, Phys. Rev. 81, 891 (1951) [$T_{1/2}$ β^-]
- 1953BU58 - P.R. J. Burch, Nature 172, 361 (1953) [$T_{1/2}$ EC]
- 1955SU38 - A.D. Suttle, W.F. Libby, Anal. Chem. 22, 921 (1955) [$T_{1/2}$ β^- , EC]
- 1955KO21 - S. Kono, J. Phys. Soc. Japan 10, 495 (1955) [$T_{1/2}$ β^-]
- 1955BA25 - G. Backenstoss, K. Goebel, Z. Naturforsch. 10a, 920 (1955) [$T_{1/2}$ EC]
- 1956MC20 - A. McNair, R.N. Grover, H. W. Wilson, Phil. Mag. 1, 199 (1956) [$T_{1/2}$ β^-]
- 1956Wetherill - G.W. Wetherill, G.J. Wasserberg, L.T. Aldrich, G.R. Tilton, R.J. Hayden, Phys. Rev. 103, 987 (1956) [$T_{1/2}$ EC]
- 1957WE43 - G.W. Wetherill, Science 126, 545 (1957) [$T_{1/2}$ EC]
- 1959KE26 - W.H. Kelly, G.B. Beard, R.A. Peters, Nucl. Phys. 11, 492 (1959) [$T_{1/2}$ β^-]
- 1959TI20 - D.R. Tilley, L. Madansky, Phys. Rev. 116 (1959) 413 [β^+/β^- ratio]
- 1960SA31 - N.K. Saha, J.B. Gupta, Proc. Natl. Inst. Sci. India 26A (1960) 486 [$T_{1/2}$ β^- , EC]
- 1961GL07 - L.E. Glendenin, Ann. N.Y. Acad. Sci. 91, 166 (1961) [$T_{1/2}$ β^-]
- 1962EN01 - D.W. Engelkemeir, K.F. Flynn, L.E. Glendenin, Phys. Rev. 126 (1962) 1818 [β^+/β^- ratio]
- 1962FL05 - D.G. Fleishman, V.V. Glazunov, Sov. At. En. 12, 338 (1962) [$T_{1/2}$ β^-]
- 1965BR25 - G.A. Brinkman, A.H.W. Aten, Jr., J.Th. Veenboer, Physica 31, 1305 (1965) [$T_{1/2}$ β^-]
- 1965LE15 - H. Leutz, G. Schulz, H. Wenninger, Z. Phys. 187, 151 (1965) [$T_{1/2}$ β^- , EC, β^+/β^- ratio]
- 1966FE09 - I. Feuerhake, A. Hinzpeter, Naturwiss. 53, 272 (1966) [β^-]
- 1966DeRuytter - A.W. DeRuytter, A.H.W. Aten, Jr., A. Van Dulmen, C. Krol-Konig, E. Zuidema, Physica 32, 991 (1966) [$T_{1/2}$ EC]
- 1966Egelkraut - K. Egelkraut, H. Leutz, Physik Verhandl. 11, 67 (1966) [$T_{1/2}$ β^- , EC]
- 1967KI10 - J.D. King, N. Neff, H.W. Taylor, Nucl. Instr. Meth. **52** (1967) 349 [E_γ]
- 1970JA15 - J.F.W. Jansen, B.J. Meijer, P. Koldewijn, Radiochim. Acta **13** (1970) 171 [E_γ]
- 1971Venkataramaiah - P. Venkataramaiah, H. Sanjeevaiah, B. Sanjeevaiah, Ind. J. Pure Appl. Phys. 9, 133 (1971) [$T_{1/2}$ β^-]
- 1972Gopal - K. Gopal, H. Sanjeevaiah, B. Sanjeevaiah, Am. J. Phys. 40, 721 (1972) [$T_{1/2}$ β^-]
- 1977CE04 - A. Cesana, M. Terrani, Anal. Chem. 49, 1156 (1977) [$T_{1/2}$ EC]
- 1979HE13 - R.G. Helmer, R.J. Gehrke, R.C. Greenwood, Nucl. Instr. Methods **166** (1979) 547 [E_γ]
- 1979SC31 - P. Schluter, G. Soff, At. Data Nucl. Data Tables **24** (1979) 509 [α_π]
- 1990EN08 - P.M. Endt, Nucl. Phys. **A521** (1990) 1, Errata and Addenda Nucl. Phys. **A529** (1991) 763, Errata Nucl. Phys. **A564** (1993) 609 [J^π , $T_{1/2}$, excited levels]
- 1995SCZY - E. Schönfeld, report PTB-6.33-95-2 (1995) [EC probabilities]
- 1996SC06 - E. Schönfeld, H. Janßen, Nucl. Instr. Meth. **A369** (1996) 527 [ω_K , K x ray ratios, Auger e- ratios, atomic data]
- 1999BeZS - M.-M. Bé, B. Duchemin, E. Browne, S.-C. Wu, V. Chechev, R. Helmer, E. Schönfeld, report CEA, ISBN 2-7272-0211-3 (1999)
- 2001BE81 - F. Begemann, K.R. Ludwig, G.W. Lugmair, K. Min, L.E. Nyquist, P.J. Patchett, P.R. Renne, C.-Y. Shih, I.M. Villa, R.J. Walker, Geochim. Cosmochim. Acta 65 (2001) 111 [Uncertainties]
- 2001Chechev - V.P. Chechev, IAEA report (2001) INDC(CCP)-432 [$T_{1/2}$]
- 2003AU03 - G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. **A729** (2003) 337 [Q]
- 2004KO09 - K. Kossert, E. Günther, Appl. Rad. Isotop. 60 (2004) 459-464 [$T_{1/2}$ Total]
- 2008KI07 - T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr. Nucl. Instr. and Meth. **A589** (2008) 202-229 [BrICC]

**⁴¹Ar - Comments on evaluation of decay data
by V. Chisté and M. M. Bé**

This evaluation was completed in February 2010. Literature by February 2010 was included.

1 Decay Scheme

⁴¹Ar disintegrates 100 % by beta minus emissions to excited levels and to the ground state of ⁴¹K.

A good agreement was found between the effective Q value (2492 (7) keV) calculated from the decay scheme data and the adopted and recommended value from the mass adjustment of Audi (2003Au03).

2 Nuclear Data

The Q value is from the atomic mass evaluation Audi and Wapstra (2003Au03).

Spin and parities are from evaluation of P. M. Endt (1990En08).

Level energies and half-life for the 1293-keV excited level are from the mass-chain evaluation of J. A. Cameron and B. Singh (2001Ca59).

Experimental ⁴¹Ar half-life values (in min) are given in Table 1:

Table 1: Experimental values of ⁴¹Ar half-life.

Reference	Experimental value (min)	Comments
A. H. Snell (1936Sn01)	110 (1)	
E. Bleuler (1946Bl28)	109.4 (10)	
H. Brown (1950Br29)	107 (3)	Outlier.
W. Hälg (1951Ha78)	109.6 (4)	
A. Schwarzschild (1956Sc91)	111 (1)	Outlier.
H. Paul (1964Pa03)	109.8 (12)	
M. Bormann (1969Bo11)	103.5 (24)	Outlier.
F. Jundt (1971Ju04)	109 (2)	
A. R. Rutledge (1986Ru09)	109.32 (12)	
A. Abzouzi (1990Ab06)	109.640 (38)	
Recommended value	109.611 (38)	$\chi^2 = 1.13$

The values of H. Brown (1950Br29), Schwarzschild (1956Sc91) and Bormann (1969Bo11) have been shown to be outliers, based on the Chauvenet's criterion and thus were omitted in the final calculation. With the 7 remaining values (1936Sn01, 1946Bl28, 1951Ha78, 1964Pa03, 1971Ju04, 1986Ru09 and 1990Ab06), a weighted average was calculated using the LWEIGHT computer code (version 3). The largest contribution to the weighted average comes from the value of Abzouzi (1990Ab06), amounting to 89 %.

The adopted value is the weighted average of 109.611 min with an external uncertainty of 0.038 min. The reduced- χ^2 value is 1.13.

2.1 β^- Transitions

The maximum energies of the β^- transitions in the decay of ⁴¹Ar \rightarrow ⁴¹K have been obtained from the Q value (2003Au03) and the level energies given in Table 2 from J. A. Cameron (2001Ca59).

Table 2: ⁴¹K levels populated in the decay of ⁴¹Ar.

Level Number	Level energy (keV)	Spin and parity	Half-life
0	0	3/2 ⁺	
1	1293.64 (4)	7/2 ⁻	6.7 (5) ns
2	1677.0 (3)	7/2 ⁺	

The transition probability of the β transition to the ground state of ⁴¹K has been reported as (Table 3):

Table 3: Experimental values of β transition probability to the ground state of ⁴¹K.

Reference	Experimental value (%)	Comments
A. Schwarzschild (1956Sc91)	0.88	Not used: no uncertainty.
G. R. Kartashov (1961Ka19)	0.78 (2)	
H. Paul (1964Pa03)	0.82 (6)	
Recommended value	0.784 (19)	$\chi^2 = 0.4$

The adopted probability value of the β transition to the ground state is the weighted average of 0.784 % with an internal uncertainty of 0.019 %. The reduced- χ^2 value is 0.4.

For the 1293- and 1677-keV levels, the adopted β^- transition probabilities and the associated uncertainties were deduced from the γ transition probability balance at each level of the decay scheme (see **4.2 γ Emissions**).

The values of log ft and average β^- energies have been calculated with the program LOGFT for the allowed, unique 1st forbidden and 1st forbidden β^- transitions.

2.2 γ Transitions

The transitions probabilities were calculated using the γ -ray emission intensities and the relevant internal conversion coefficients (see **4.2 Gamma Emissions**).

Multipolarity and mixing ratio of 1293-keV γ -ray transition are from H. H. Eggenhuisen (1978Eg01): 1293-keV γ -ray: M2 + E3, with $\delta = 0.118$ (12)

The internal conversion coefficients (ICC's) for this γ -ray transition have been interpolated from theoretical values of I. M. Band (2002Ba85) using the BrIcc computer program (calculation for 'hole') (2008Ki07). The α_T theoretical value is compared with a measured value in Table 4. They are in agreement within the uncertainty limits.

Table 4: Experimental and recommended (calculated) values internal conversion coefficients.

	α_T (1293-keV)
G. R. Kartashov (1961Ka19)	6.8 (9) 10^{-5}
BrIcc program (recommended values)	7.44 (11) 10^{-5}

The internal pair formation coefficient α_π (1293 keV) is 4.92 (7) 10^{-6} (given by BrIcc computer program). So α_T , using to calculate the 1293-keV transition probability, is

$$\alpha_T = 4.92 (7) 10^{-6} + 7.44 (11) 10^{-5} = 7.93 (11) 10^{-5}$$

3 Atomic Data

Atomic values, ω_K , n_{KL} and the X-ray relative probabilities are from Schönfeld and Janßen (1996Sc06).

4 Emissions

4.1 K x-rays

The X-ray absolute intensities were deduced from the decay data using the EMISSION computer code.

4.2 Photon emissions

The energies of the γ -rays given in section 5.2 are from J. A. Cameron (2001Ca59).

The experimental relative γ -ray emission intensities in ⁴¹K have been obtained from all the available relative and absolute values.

The normalization factor to convert the relative emission intensities to absolute emission intensities is calculated using the formula:

$$N = \left(\frac{100 - P_\beta(0,0)}{(\sum(1 + \alpha_T)P_{rel})} \right) = \frac{100 - 0.784(19)}{(\sum(1 + \alpha_T)P_{rel})} = 0.99157 (20),$$

where the sum is over all the γ transitions to the ground state (1293- and 1677-keV) and α_T is the relevant coefficient. The uncertainty was calculated through the propagation on the formula given above.

The experimental γ -ray emission probabilities relative to 100 for the 1293-keV γ -ray are given in Table 5.

Table 5: Experimental data sets of the relative γ -ray emission intensities (%)

Reference / Energy (keV)	1293.64 (4)	1677.0 (3)
W. W. Pratt (1965Pr05)	100	5 (2) 10 ⁻²
F. Jundt (1971Ju04)	100	5.2 (5) 10 ⁻²
Evaluated value	100	5.19 (49) 10 ⁻²
χ^2		0.0094

The adopted values are the weighted means calculated by the LWEIGHT program (version 3).

The adopted relative and absolute γ -ray emission probabilities are given in Table 6.

Table 6: Recommended relative and absolute γ -ray intensities (%).

E γ (keV)	Relative γ -ray intensity (%)	Absolute γ -ray intensity (%)
1293.64 (4)	100	99.157 (20)
1677.0 (3)	5.19 (49) 10 ⁻²	5.15 (49) 10 ⁻²

6 References

- 1936Sn01 A. H. Snell, Phys. Rev. 49(1936)555 [Half-life].
- 1946B128 E. Bleuler, W. Bollmann, W. Zünti, Helv. Phys. Acta 19(1946)419 [Half-life].
- 1950Br29 H. Brown, V. Perez-Mendez, Phys. Rev. 78(1950)649 [Half-life].
- 1951Ha78 W. Hälg, Helv. Phys. Acta 24(1951)641 [Half-life].
- 1956Sc91 A. Schwarzschild, B. M. Rustad, C. S. Wu, Phys. Rev. 103(1956)1796 [Half-life, P_{β}].
- 1961Ka19 G. R. Kartashov, N. A. Burgov, A. V. Davydov, Bull. Acad. Sci. USSR, Phys. Ser. 25(1961)184 [P_{β} , α_T].
- 1962En05 P. M. Endt, C. van der Leun, Nucl. Phys. 34(1962)1 [Spin, parity].
- 1964Pa03 H. Paul, Acta Phys. Austriaca 18(1964)315 [Half-life, P_{β}].
- 1965Pr05 W. W. Pratt, Phys. Rev. 103(1965)B509 [Gamma-ray emission probability].
- 1969Bo11 M. Bormann, B. Lammers, Nucl. Phys. A130(1969)195 [Half-life].
- 1971Ju04 F. Jundt, E. Aslanides, A. Gallmann, E. K. Warburton, Phys. Rev. C4(1971)498 [Half-life, gamma-ray emission probability].
- 1978Eg01 H. H. Eggenhuisen, L. P. Ekström, G. A. P. Engelbertinik, H. J. M. Aarts, W. G. J. Langeveld, Nucl. Phys. A299(1978)175 [Mixing ratio].
- 1978En23 P. M. Endt, C. van der Leun, Nucl. Phys. A310(1978)1 [Spin, parity].
- 1986Ru09 A. R. Rutledge, L. V. Smith, J. S. Merritt, Appl. Rad. Isotop. 37(1986)1029 [Half-life].
- 1990En08 P. M. Endt, Nucl. Phys. A521(1990)1 [Spin, parity].
- 1990Ab06 A. Abzouzi, M. S. Antony, V. B. Ndocko Ndongué, D. Oster, J. Radioanal. Nucl. Chem. Lett. 145(1990)361 [Half-life].
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
- 2001Ca59 J. A. Cameron, B. Singh, Nucl. Data Sheets 94(2001)429 [Level energies, excited level half-life].
- 2002Ba85 I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor Jr., At. Data Nucl. Data Tables 81 (2002) 1 [ICC]
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr., Nucl. Instrum. Meth. Phys. Res. A589(2008)202 [Theoretical ICC].

⁴¹Ca - Comments on evaluation of decay data by X. Mougeot

This first evaluation was done in 2012, taking into account the available literature up to November 2012.

⁴¹Ca plays an important role in the long-term evaluation of the safety of final repositories for nuclear waste and is used to study the fine-scale chronology of the formation of the Solar System (2012JO04).

1 Decay Scheme

The decay scheme is complete since ⁴¹Ca can only decay through electron-capture to the ground state of ⁴¹K: $Q_e = 421,63$ (14) keV (from 2011AUZZ), and the first excited state of ⁴¹K is $1/2^+$ at 980,476 (6) keV (from 2001CA59).

The J^π of the ground states of both ⁴¹Ca ($7/2^-$) and ⁴¹K ($3/2^+$) are from the 2001CA59 evaluation. The electron-capture transition is therefore first forbidden unique.

2 Nuclear Data

The first measurement of the Q value was detailed in 1950RI59. The Q value adopted in this evaluation is from Audi *et al.* 2011 (2011AUZZ).

A full list of the half-life measurements available by November 2012, including the reasons why certain have been excluded by the evaluator, is given in Table 1.

Three types of measurements were carried out:

- 1) using accelerator mass spectrometry, $T_{1/2}({}^{41}\text{Ca})$ depends on $T_{1/2}({}^{36}\text{Cl})$;
- 2) using specific activity deduced from X-ray intensity measurements, $T_{1/2}({}^{41}\text{Ca})$ depends on neutron capture cross sections (σ_{40} and σ_{44}) and possibly on $T_{1/2}({}^{45}\text{Ca})$, $\omega_K(\text{K})$, P_K and P_L ;
- 3) using liquid scintillation counting (TDCR method) for the activity and thermal ionization mass spectrometry for the absolute isotopic composition, the uncertainty on $T_{1/2}({}^{41}\text{Ca})$ depends mainly on the uncertainty on P_K .

Thus, the available measurements have been recalculated using up-to-date values when it was necessary.

2.1 Half-life

In order to evaluate the ⁴¹Ca half-life, the measured half-life values were recalculated using up-to-date values of parameters. The corresponding uncertainties were also calculated. The following parameter values have been used:

- $N_A = 6,02214129$ (27) 10^{23} mol⁻¹ from 2008MO18.
- $\sigma_{40} = 0,41$ (2) b and $\sigma_{44} = 0,88$ (5) b from 2006MUZX.
- $I_{ab}({}^{40}\text{Ca}) = 0,9694$ (16) from 2011BE53.
- $\omega_K(\text{K}) = 0,143$ (4) from 1996SC06.
- LOGFT can calculate a first forbidden unique electron-capture transition. Using the Q value from 2011AUZZ, the following quantities have been calculated: $P_K = 0,894$ (9); $P_L = 0,0916$ (9); $P_{M+} = 0,01482$ (15). Thus, $P_L/P_K = 0,1025$ (10). A relative uncertainty of 1 % has been taken.
- $T_{1/2}({}^{36}\text{Cl}) = 3,02$ (4) 10^5 a from the latest DDEP evaluation (2012).
- $T_{1/2}({}^{45}\text{Ca}) = 162,64$ (11) d from the latest DDEP evaluation (2012).

Analysis of the publications

1962DR03 (Drouin *et al.*)

The source of ⁴¹Ca had been prepared ten years before by high neutron irradiation of a sample of

calcium carbonate. The number of ⁴¹Ca atoms was calculated as:

$$N_{41} = \frac{N_A}{40 \cdot 1000} \times I_{ab}(^{40}\text{Ca}) \times \sigma_{40} \times (8,07 \cdot 10^{20})$$

where the factor 1000 is used for converting grams to milligrams, and the last factor is the integrated neutron flux. With the parameter values given above, the new N_{41} value is: $4,83 (24) 10^{15}$ atoms·mg⁻¹. The decay rate is then:

$$DR = 594 (6) \times \frac{1}{0,86} \times \frac{1}{0,67} \times \frac{1}{\omega_K(K)} \times \left(1 + \frac{P_L}{P_K}\right) \times 2$$

where the first numerical term is the number of potassium K X-rays per min measured for 0,42 mg of Ca, the factor 2 corrects for the 50 % geometrical detection efficiency, 1/0,86 corrects for the self-absorption in the source (estimated to be 14 % in 1962DR03), and 1/0,67 corrects for the transmission of X-rays by the Auger electron absorber (estimated to be 67 % in 1962DR03).

Thus, the new value is: $DR = 1,59 (5) 10^4 \text{ min}^{-1}/(0,42 \text{ mg Ca})$. As:

$$T_{1/2} = \frac{N_{41} \cdot \ln 2}{DR/0,42}$$

the new half-life result is: $T_{1/2}(^{41}\text{Ca}) = 8,8 (5) 10^{10} \text{ min} = 1,68 (10) 10^5 \text{ a}$. The authors have estimated a relative uncertainty of about 15 %. Therefore, the final value is $T_{1/2}(^{41}\text{Ca}) = 1,68 (25) 10^5 \text{ a}$.

1966WA (Wahlin)

$T_{1/2}(^{41}\text{Ca}) = 1,00 (25) 10^5 \text{ a}$ was determined proportionally to $T_{1/2}(^{36}\text{Cl}) = 2,60 (8) 10^5 \text{ a}$. The up-to-date value of $T_{1/2}(^{36}\text{Cl})$ provides directly the up-to-date value of $T_{1/2}(^{41}\text{Ca}) = 1,16 (29) 10^5 \text{ a}$.

1974MA30 (Mabuchi *et al.*)

The authors gave a value of $T_{1/2}(^{41}\text{Ca}) = 1,03 (4) 10^5 \text{ a}$ using a neutron capture cross section on ⁴⁰Ca with a relative uncertainty of 10 %. The uncertainty on $T_{1/2}(^{41}\text{Ca})$ in 1974MA30 was clearly underestimated and has been recalculated by the evaluator to be $0,14 10^5 \text{ a}$. The half-life is calculated as follows:

$$T_{1/2}(^{41}\text{Ca}) = 108 (4) \times 5,29 \cdot 10^3 \times \frac{\sigma_{40}}{\sigma_{44}} \times T_{1/2}(^{45}\text{Ca}) \times \frac{1}{0,9750}$$

where the first numerical term is the ratio of measured activity of ⁴⁵Ca on measured activity of ⁴¹Ca, the second numerical term is the ratio (resulting from mass spectroscopy analysis) of the number of ⁴⁰Ca atoms on the number of ⁴⁴Ca atoms irradiated using a thermal neutron flux, and the last numerical term takes into account the time duration of irradiation and depends on experimental conditions. Thus, the up-to-date value is: $T_{1/2}(^{41}\text{Ca}) = 1,22 (10) 10^5 \text{ a}$.

1991PA10 (Paul *et al.*)

The authors gave a value of $T_{1/2}(^{41}\text{Ca}) = 1,01 (10) 10^5 \text{ a}$ derived from the specific activity determined from ⁴¹K X-ray measurements. They obtained: $S(^{41}\text{K}) = 23,98 (186) \text{ X-rays min}^{-1}\text{ng}^{-1}$.

As:

$$S(^{41}\text{Ca}) = \frac{S(^{41}\text{K})}{\omega_K P_K}$$

one has the new value: $S(^{41}\text{Ca}) = 188 (16) \text{ min}^{-1}\text{ng}^{-1}$, and the half-life is given by:

$$T_{1/2} = \frac{N_A \cdot \ln 2}{S(^{41}\text{Ca}) \cdot 41}$$

namely $T_{1/2}(^{41}\text{Ca}) = 1,03 (9) 10^5 \text{ a}$.

1991KL06 (Klein *et al.*)

The authors obtained a value of $T_{1/2}(^{41}\text{Ca}) = 1,03 (7) 10^5 \text{ a}$ which depends on the half-life of ³⁶Cl, taken as $T_{1/2}(^{36}\text{Cl}) = 3,01 (2) 10^5 \text{ a}$. As they measured a ratio $T_{1/2}(^{36}\text{Cl})/ T_{1/2}(^{41}\text{Ca}) = 2,91 (9)$, the updated value is $T_{1/2}(^{41}\text{Ca}) = 1,04 (7) 10^5 \text{ a}$. It is noteworthy that the authors estimated a final uncertainty two times larger, from $0,035 10^5 \text{ a}$ to $0,07 10^5 \text{ a}$, to take into account the dispersion of their measurements.

Evaluation


The half-life value and the reasons why some of them have been excluded by the evaluator are given in Table 1.

The LWEIGHT program excluded the result from 1962DR03, which was found to be an outlier by Chauvenet's criterion. A weighted average was adopted from the resulting consistent data set, with a reduced- χ^2 value of 1,43. The data used for the evaluation of the ⁴¹Ca half-life can be seen in Figure 1. The main contribution comes from 2012JO04, amounting to 91 %. The quality of this measurement leads the evaluator to keep this high contribution to the weighted average. The recommended value for the ⁴¹Ca half-life is then: $T_{1/2}({}^{41}\text{Ca}) = 1,002 (17) \times 10^5 \text{ a}$.

Table 1: Measured half-lives used for the evaluation.

Reference	Measured $T_{1/2}({}^{41}\text{Ca})$ (10^5 a)	Updated $T_{1/2}({}^{41}\text{Ca})$ (10^5 a)	Comments
1951BR94	1,2 (4)	-	Excluded: same authors as in 1962DR03
1953BR71	1,1 (3)	-	Excluded: same authors as in 1962DR03
1962DR03	0,75 (11)	1,68 (25)	Excluded by LWEIGHT (Chauvenet's criterion)
1966WA	1,00 (25)	1,16 (29)	
1974MA30	1,03 (14)	1,22 (10)	
1990FH13	1,03 (7)	1,04 (7)	Excluded: same authors as in 1991KL06
1991PA10	1,01 (10)	1,03 (9)	
1991KL06	1,03 (7)	1,04 (7)	
1992KU	1,01 (10)	1,03 (9)	Excluded: same authors as in 1991PA10
2012JO04	0,994 (15)	0,994 (15)	No updating required
Adopted	-	1,002 (17)	Weighted average using LWEIGHT

Figure 1: $T_{1/2}$ measurements used for the present evaluation. Blue crosses: measured. Red circles: updated.



2.2 Electron Capture Transition

⁴¹Ca can only decay through electron-capture to the ground state of ⁴¹K.

The LOGFT program can calculate this first forbidden unique electron-capture transition. The $\log ft$ value is: $\log ft(1U) = 10,53(3)$. LOGFT provides also the P_K , etc. values:

- $P_K = 0,894(9)$
- $P_L = 0,0916(9)$
- $P_{M+} = 0,01482(15)$
- $P_{L2}/P_{L1} = 0,00261(3)$
- $P_{L3}/P_{L1} = 0,0141(1)$

Thus: $P_L/P_K = 0,1025(10)$; $P_{L1} = 0,0901(9)$; $P_{L2} = 0,0002351(24)$; $P_{L3} = 0,001270(13)$. A relative uncertainty of 1 % has been taken by the evaluator.

3 Atomic Data (K, Z=19)

3.1 X Radiations and Auger electrons

The X-ray and Auger electron data were computed using the EMISSION program with the atomic data of Schönfeld and Janßen (1996SC06).

The authors of 1951SA31 measured $E_{XK}(^{41}\text{K}) = 3,40(15)$ keV. The authors of 1970SI24 measured $E_{XK}(^{41}\text{K}) = 3,17(10)$ keV. The adopted value from 1996SC06 is: $E_{XK}(^{41}\text{K}) = 3,31$ keV.

4 Radiation Emissions

4.1 Electron Emission

Internal ionization has been studied in 1988JA11. In this process, an orbital electron is ejected due to the sudden change of Coulomb potential and rearrangements in the electronic structure caused by the capture of another orbital electron by the nucleus. The energy released is statistically shared between the electron and the neutrino, giving rise to a continuous energy spectrum of emitted electrons that spans from 0 keV to the Q value of the transition. The integrated probability of electron ejection with an energy greater than 38 keV was found to be $1.7(2) \cdot 10^{-6}$ per K-capture.

4.2 Photon Emissions

In ordinary electron-capture decay, the energy is shared statistically between the neutrino and the nucleus. A second-order process, the internal bremsstrahlung, makes possible the emission of a photon together with the neutrino. The energy is thus shared between three bodies, giving rise to a continuous energy spectrum of photons that spans from 0 keV to the Q value of the transition. The probability of this process was measured in 1987HO16 to be: $P_{IB(1s)} = 2.5(3) \cdot 10^{-4}$ per K-capture.

Measurements of the photon spectrum can be found in 1973MY03, 1987HO16 and 1991PA10. Internal bremsstrahlung calculations can be found in 1992KA05 and 1973ZO03.

5. References

- | | |
|----------|--|
| 1950RI59 | H. T. Richards, R. V. Smith, C. P. Browne, Phys. Rev. 80, 524 (1950) [Q first measurement] |
| 1951BR94 | F. Brown, G. C. Hanna, L. Yaffe, Phys. Rev. 84, 1243 (1951) [Half-life] |
| 1951SA31 | V. L. Sailor, J. J. Floyd, Phys. Rev. 82, 960 (1951) [$E_{XK}(^{41}\text{K})$] |
| 1953BR71 | F. Brown, G. C. Hanna, L. Yaffe, Proc. Roy. Soc. (London) 220A, 203 (1953) [Half-life] |
| 1962DR03 | J. R. S. Drouin, L. Yaffe, Can. J. Chem. 40, 833 (1962) [Half-life] |
| 1966WA | L. Wahlin, UCOL-535-561 p.59, University of Colorado (1966) [Half-life] |
| 1970SI24 | B. Sitar, J. Chrapan, J. Oravec, K. Durcek, Jad. Energ. 16, 303 (1970) [$E_{XK}(^{41}\text{K})$] |
| 1973MY03 | B. Myslek, Z. Sujkowski, J. Zylicz, Nucl. Phys. A215, 79 (1973) [IB spectrum] |
| 1973ZO03 | B. A. Zon, Izv. Akad. Nauk SSSR, Ser. Fiz. 37, 1978 (1973); Bull. Acad. Sci. USSR, Phys. Ser. 37, No.9, 153 (1974) [IB calculations] |

- 1974MA30 H. Mabuchi, H. Takahashi, Y. Nakamura, K. Notsu, H. Hamaguchi, J. Inorg. Nucl. Chem. 36, 1687 (1974) [Half-life]
- 1987HO16 P. Hornshoj, T. Batsch, Z. Janas, M. Pfuetzner, A. Plochocki, K. Rykaczewski, Nucl. Phys. A472, 139 (1987) [IB spectrum]
- 1988JA11 Z. Janas, M. Pfuetzner, A. Plochocki, D. Seweryniak, Nucl. Phys. A486, 278 (1988) [Internal ionization]
- 1990FI13 D. Fink, J. Klein, R. Middleton, Nucl. Instrum. Methods Phys. Res. B52, 572 (1990) [Half-life]
- 1991PA10 M. Paul, I. Ahmad, W. Kutschera, Z. Phys. A340, 249 (1991) [Half-life]
- 1991KL06 J. Klein, D. Fink, R. Middleton, K. Nishiizumi, J. Arnold, Earth Planet. Sci. Lett. 103, 79 (1991) [Half-life]
- 1992KA05 L. Kalinowski, Z. Janas, M. Pfuetzner, A. Plochocki, P. Hornshoj, H. L. Nielsen, Nucl. Phys. A537, 1 (1992) [IB calculations]
- 1992KU W. Kutschera, I. Ahmad, M. Paul, Radiocarbon 34, 436 (1992) [Half-life, IB spectrum]
- 1996SC06 E. Schönfeld, H. Janßen, Nucl. Instr. Meth. **A369** (1996) 527 [ω_K , K x ray ratios, Auger e-ratios, atomic data]
- 2001CA59 J. A. Cameron, B. Singh, Nucl. Data Sheets 94, 429 (2001) [J^π of ground states, E_γ of first excited state]
- 2006MUZX S. F. Mughabghab, *Atlas of Neutron Resonances, 5 ed., Resonance Parameters and Thermal Cross Sections, Z = 1-100*, Elsevier, Amsterdam (2006) [Neutron capture cross sections]
- 2008MO18 P. J. Mohr, B. N. Taylor, D. B. Newell, Rev. Mod. Phys. 80, 633 (2008) [N_A]
- 2011AUZZ G. Audi, W. Meng, Atomic Mass Evaluation 2011, Private Communication (2011) [Q]
- 2011BE53 M. Berglund, M. E. Wieser, Pure Appl. Chem. 83, 397 (2011) [Isotopic abundance]
- 2012JO04 G. Jörg, Y. Amelin, K. Kossert, C. L. v. Gostomski, Geochimica et Cosmochimica Acta 88, 51-65 (2012) [Half-life]

⁴⁴Sc – Comments on evaluation of decay data

by E. Browne

The *Limitation of Relative Statistical Weights* ^[1] (LWM) method, used for averaging numbers throughout this evaluation, provided a uniform approach for the analysis of discrepant data. The uncertainty assigned to the recommended value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

⁴⁴Sc (T_{1/2} = 3.97 h) decays 94.27(5)% by β⁺, and 5.73(5)% by electron capture (Q(EC)=3653.3(19) keV (95Au04)^[2]) allowed transitions to levels at 1157.0 -, 2656.5-, and 3301.5-keV in ⁴⁴Ca (stable). A β⁺ transition from ⁴⁴Sc (J^π = 2⁺) to the ground state of ⁴⁴Ca (J^π = 0⁺) has not been observed. Such transition would be second-forbidden non unique, for which the systematic trend of log ft predicts a value > 10.6 (98Si17)^[4]. For ⁴⁴Sc this value corresponds to a β⁺ transition probability limit of < 0.005%. Therefore, I used no β⁺ feeding to the ground state, and normalized the decay scheme using the sum of the relative transition probabilities of the 1157.0-, 2656.4-, and 3301.3-keV gamma rays. This procedure produced a normalization factor N = (9.9875(3) × 10⁻⁴), as it will be shown in below.

Nuclear Data

The recommended half-life of ⁴⁴Sc, 3.97(4) h, is a weighted average (LWM, σ_{int}=0.01, χ²/ν= 8.0) of 3.927(8) h (69Ra16)^[5], 4.00(2) h (66Ta01)^[6], and 4.05(3) h ^[7]. Other values are: 4.04 h ^[8], 4.01 h ^[9], and 3.9 h ^[10], were not used because they have no uncertainties.

Gamma Rays

Tables Ia and Ib give gamma -ray energies and relative emission probabilities, respectively, reported by 90Me15^[11], 83Gu11^[12], 76Co06 ^[13], 74HeYW ^[14], 73Si05 ^[15], and 90Sc08 ^[16]. Recommended values (weighted averages (LWM)) are given on columns 5 and 7, respectively.

Table Ia - Gamma-Ray Energies

90Me15 ^[11] & 76Co06 ^[13] keV	83Gu11 ^[12] keV	74HeYW ^[14] keV	73Si05 ^[15] keV	Rec. Value keV	χ ² /ν
	646.55 (62)		646.5 (20)		
726.49	726.3 (15)		726.0 (15)		
	772.7 (12)		774		
1157.031 (15)	1157.015 (15)	1156.92 (15)	1156.9 (5)	1157.020 (15)	0.37
1499.489 (25)	1499.436 (15)	1499.20 (20)	1499.4 (3)	1499.460 (20)	1.3
2144.3 (1)	2144.43 (20)		2144.8 (8)	2144.33 (10)	0.34
2656.478 (30)	2656.435 (50)	2657.14 (20)	2656.4 (5)	2656.48 (7)	3.9
3301.3 (1)	3301.361 (55)	3301.6 (15)	3301.35 (6)	0.16	

Table Ib - Relative Gamma-Ray Emission Probabilities

Energy keV	90Me15 ^[11] & 76Co06 ^[13]	90Sc08 ^[16]	83Gu11 ^[12]	74HeYW ^[14]	73Si05 ^[15]	Rec. Value	χ^2/ν
646.5			0.040		0.043 (18)		
726.3	=0.014		0.053 (10)		0.051 (21)		
772.7	=0.0067*		0.062 (16)		0.041 (23)		
1157.020	1000(3)	1000 (1)	1000 (3) [#]	1000 (50)	1000 (3) [#]	1000 (3)	
1499.46	9.0 (2)	9.12 (15)	9.22 (37)	9.0 (10)	9.1 (4)	9.09 (15)	0.10
2144.33	0.02 (2)		0.035 (10) [#]		0.039 (7)	0.036 (7)	0.41
2656.48	1.11 (4)	1.15 (6)	1.11 (3)	1.4 (5)	1.3 (1)	1.12 (3)	0.98
3301.35	0.0064 (8)		0.016 (2)		0.018 (3)	0.017 (2) ^{&}	0.31

* From ⁴⁴K decay, relative to 9.0 for the emission probability of 1499-keV gamma ray.

Estimated by evaluator.

& Weighted average of 0.016(2) and 0.018(3).

The 726- and 772-keV gamma rays reported by 83Gu11^[12] and 73Si05^[15] were not observed by 90Me15^[11] and 76Co06^[13], who reported upper limits four and nine times lower, respectively, for their relative emission probabilities. Therefore, they probably do not belong to the decay of ⁴⁴Sc.

The 646-keV gamma ray was observed with about the same relative emission probability by both 83Gu11^[12] and 73Si05^[15]. These authors placed this gamma ray de-exciting a 3301-keV level, which is also de-excited by the 2144- and 3301-keV transitions. 90Me15^[11] and 76Co06^[13] did not report the 646-keV gamma ray. However, 76Co06^[13] have seen it in the β^- decay of ⁴⁴K. Table II shows the relative emission probabilities of the 646-, 2144-, and 3301-keV gamma rays, which de-excite the 3301-keV level, from both ⁴⁴Sc electron-capture and ⁴⁴K β^- decay.

Table II - Relative Emission Probabilities for the 646-, 2144-, and 3301-keV Gamma Rays from the 3301-keV Level

Energy keV	83Gu11 ^[12] P _{γ} From ⁴⁴ Sc EC Decay	73Si05 ^[15] P _{γ}	76Co06 ^[13] P _{γ} From ⁴⁴ K β^- Decay
646.5	0.040	0.043 (18)	1.5 (5)
2144.33	0.035	0.039 (7)	12.9 (8)
3301.35	0.016 (2)	0.018 (3)	5.5 (9)

R(646/2144)	1.1	1.1	0.12
R(2144/3301)	2.2	2.2	2.3

Table II shows that the ratio R(646/2144) is ten times lower from ⁴⁴K β^- decay than from ⁴⁴Sc electron-capture decay. Consequently, the 646-keV gamma-ray, observed from ⁴⁴K decay, does not de-excite the 3301-keV level, as 83Gu11 had suggested, and therefore, its existence is uncertain.

Multipolarities and Conversion Coefficients

A total measured conversion coefficient ^[17] $\alpha_t=6.3 (3)\times 10^{-5}$ for the 1157.020-keV gamma-ray suggests an E2 multipolarity for this gamma-ray. The 1499.46-keV gamma-ray has an M1+1.8 (4)% E2 multipolarity ($\delta = +0.137(7)$), determined in a $\gamma\gamma(\theta)$ measurement (68Wa21)^[3]. The theoretical conversion coefficients in Table 2.3 (Tables Section) for these transitions are from 76Ba63 ^[18]. Conversion coefficients for pair creation are theoretical values from 79Sc31^[30].

Absolute Emission Probabilities.

As mentioned before, the gamma-ray normalization factor *N* can be obtained as follows:

$$N = 1/[P_{\gamma(1157)}(1+\alpha_{1157}) + P_{\gamma(2256)} + P_{\gamma(3301)}] = 1/[1000 (3) (1 + 6.68 \times 10^{-5}) + 1.12 (3) + 0.064 (8)] = 9.9875(3) \times 10^{-4}$$

The internal pair conversion coefficients (from 79Sc31^[19]) for these gamma-rays are: $\alpha_{IP}(1157, E2) = 4.0 \times 10^{-6}$, $\alpha_{IP}(2256, E2) = 5.9 \times 10^{-4}$, and $\alpha_{IP}(3301, E2) = 9.0 \times 10^{-4}$. These coefficients were not included in the calculation shown above because their effect is negligible.

The fractional uncertainty in *N* should be added in quadrature to those in the relative emission probabilities. For the 1157.020 -keV gamma-ray, which dominates this normalization, the correct propagation of this uncertainty is as follows:

$$P_{\gamma(abs,1157)} = P_{\gamma(rel, 1157)} \times N = 1000 (3)/[1000 (3) (1 + 6.68 \times 10^{-5}) + 1.12 (3) + 0.064 (8)] = 1/ [1.0000668 + 1.18 (3)/1000 (3)] = 1/[1.0000668 + 0.00118 (3)] = 0.99875(3)$$

Notice that the fractional uncertainty of the relative emission probability is 0.3% , however, because of the effect of covariances, that in the absolute emission probability is just 0.003% . Table III shows the gamma-ray absolute emission probabilities.

Table III - Absolute Gamma-ray Emission Probabilities

Energy (keV)	$P_{\gamma}(\%)$
1157.020 (15)	99.875 (3)
1499.460 (20)	0.908 (15)
2144.33 (10)	$3.6 (7) \times 10^{-3}$
2656.48 (7)	0.112 (3)
3301.35 (6)	$1.7 (2) \times 10^{-3}$

Electron-Capture and b^+ Transitions

The electron-capture plus β^+ probabilities shown in the decay scheme have been deduced from gamma - ray transition intensity balances at each level. For the transition to the 1157 -keV level, the values of the individual β^+ and electron-capture probabilities (given in Table s 2.2 and 2.1, respectively) are based on the recommended $\epsilon/\beta^+ = 0.0499(5)$ ratio. This ratio is a weighted average of the experimental values 0.0499(5) (83Ba41)^[20] and 0.0497(23) (76St21)^[21]. Theory predicts 0.0489^[22].

Electron-capture probabilities to the various atomic sub -shells, i.e., P_K, P_L, P_{M+} in Table 2.1 are theoretical values (98Sc28)^[23] calculated with the computer program EC-CAPTURE^[24].

90Sc08^[16] measured the annihilation emission probability $P_{\gamma\pm}(511) = 1.88(3)$, which includes a 2.4% correction for positron annihilation -in-flight. I confirmed the value of this correction using the calculation procedure presented in Appendix D of the *Table of Radioactive Isotopes*^[25], as described below in Table IV.

Table IV - Annihilation-in-flight Correction Factor

E(bin) keV	$\langle\beta^+\rangle^*$ keV	β^+ (%) [#] %	$E_{avg}^{\&}$ keV	$P(E_{avg})^{\wedge}$ %	$\beta^+_{fl}^{\circledast}$ %
0-10	0.000434	0.0056	7.75		
10-20	0.0056	0.0355	15.77		
20-40	0.060	0.191	31.41		
40-100	1.12	1.50	74.67	0.5	0.0075
100-300	26.8	12.6	212.69	1.0	0.126
300-600	140.0	30.7	456.03	2.1	0.645
600-1300	418.0	48.6	860.0	3.4	1.652
1300-2497	10.8	0.80	1350.0	4.8	0.038
Total β^+ branching		94.0		Correction factor	2.47

*Average β^+ energy per decay

β^+ bin probability

& Average β^+ bin energy = $100 \langle\beta^+\rangle/\beta^+(\%)$

^ Positron annihilation-in-flight probability (from Fig.3, Appendix D, *Table of Radioactive Isotopes*)

⊙ Fraction (in %) of β^+ transitions that annihilate in flight = $0.01 \times \beta^+(\%) \times P(E_{avg})$

The final result, 2.47%, agrees with 2.4%, used by 90Sc08 [16].

Then, the β^+ probability is $P_{\beta^+}(1157) = 1.88(3)/2 = 0.940(15)$. The electron-capture probability, $P_{EC}(1157) = 0.9897(5) - 0.940(15) = 0.0497(15)$, although less accurate, is in agreement with the recommended value given in Table 2.1.

Levels half-life

The following half-life values: 2.61(14) ps (1157 -keV level), 30(3) fs (2656 -keV level), and 35 (18) fs (3301-keV level), shown on the level scheme, are from 90En08 [26].

Atomic Data

The X-ray and Auger-electron probabilities in Section 4 have been calculated with the computer program EMISSION^[27], using the gamma -ray and electron -capture data from Section 2, and atomic data from 96Sc06^[28]

Total Average Radiation Energy

The calculated (RADLST^[29]) total average radiation energy of 3653.3(25) keV (which includes all the radiations emitted by ⁴⁴Sc), agrees very well with $Q(EC) = 3653.3(19)$ keV (1995Au04 [2]) and confirms the self consistency of the ⁴⁴Sc decay scheme.

6 References

1. M. J. Woods and A. S. Munster, *Evaluation of Half-Life Data*, National Physical Laboratory, Teddington, UK, Rep. RS(EXT) 95, (1988) - 88WoZO
2. G. Audi and A. H. Wapstra, Nucl. Phys. **A595**, 409 (1995) - 95Au04 [Q(EC)]
3. H. K. Walter, A. Weitsch, H. J. Welke, Z. Physik **213**, 323 (1968) - 68Wa21 [Gamma-ray mult. Mixing, δ]

4. B. Singh, J. L. Rodriguez, S. S. M. Wong, and J. K. Tuli, Nucl. Data Sheets 84, 487 (1998) - 98Si17
[Log ft systematics]
5. H. Ravn, J. Inorg. Nucl. Chem. **31**, 1883 (1969) - 69Ra16
[$T_{1/2}$]
6. J. R. Tatarczuk, Phys. Rev. **143**, 818 (1966) - 66Ta16
[$T_{1/2}$]
7. D. R. Sachdev and L. Yaffe, Can. J. Chem. **47**, 1667 (1969)
[$T_{1/2}$]
8. L. A. Rayburn, Phys. Rev. **122**, 168 (1961) - 61Ra06
[$T_{1/2}$]
9. C. S. Khurana and H. S. Hans, Nucl. Phys. **28**, 560 (1961)
[$T_{1/2}$]
10. L. T. Dillman, J. J. Kraushaar, J. D. McCullen, Nucl. Phys. **42**, 383 (1963) - 63Di06
[$T_{1/2}$]
11. R. A. Meyer, Fizika (Zagreb) **22**, 153 (1990) - 90Me15
[E_{γ} , P_{γ}]
12. Guanjun Yuan et al., Nucl. Sci. Eng. **84**, 320 (1983) - 83Gu11
[E_{γ} , P_{γ}]
13. G. Coleman, R. A. Meyer, Phys. Rev. **C13**, 847 (1976) - 76Co06
[E_{γ} , P_{γ}]
14. R. L. Heath, Report ANCR-1000-2 (1974) - 74HeYW
[E_{γ} , P_{γ}]
15. J. J. Simpson, Nucl. Phys. **A203**, 221 (1973) - 73Si05
[E_{γ} , P_{γ}]
16. U. Schötzig, Nucl. Instrum. Methods Phys. Res. **A286**, 523 (1990) - 90Sc08
[P_{γ} , γ_{\pm} positron annihilation radiation]
17. J. W. Blue and E. Bleuer, Phys. Rev. **100**, 1324 (1955) - 55Bl23
[ICC]
18. I. M. Band et al., At. Data Nucl. Data Tables **18**, 433 (1976) - 76Ba63
[Theor. ICC]
19. P. Schluter and G. Soff, At. Data Nucl. Data Tables **24**, 509 (1979) - 79Sc31
[Theor. Int. Pair ICC]
20. A. P. Baerg, Can. J. Phys. **61**, 1222 (1983) - 83Ba41
[ϵ/β^+ ratio]
21. H. Stocker, A. P. Baerg, Can. J. Phys. **54**, 2396 (1976) - 76St21
[ϵ/β^+ ratio]
22. P. F. Zweifel, Phys. Rev. **107**, 329 (1957)
[Theor. ϵ/β^+ ratio]
23. E. Schönfeld, Appl. Radiat. Isot. **49**, 1353 (1998) - 98Sc28
[P_K , P_L , P_M]
24. E. Schönfeld, F. Y. Chu, and E. Browne, EC-CAPTURE, a computer program to calculate electron capture probabilities to atomic sub-shells, 1998.
25. E. Browne and R. B. Firestone, Table of Radioactive Isotopes, John Wiley and Sons, Inc., New York (1986) - 86BrZQ
[Positron Annih. in flight]
26. P. M. Endt, Nucl. Phys. **A521**, 1 (1990) - 90En08
[$T_{1/2}$ level]
27. H. Janßen and E. Schönfeld, EMISSION, a computer program for calculating X-ray and Auger-electron emission probabilities, 1998.
28. E. Schönfeld and H. Janßen, Nucl. Instrum. Meth. Phys. Res. **A369**, 527 (1996) - 96Sc06
[X-ray, ω_K]
29. The Program RADLST, Thomas W, Burrows, report BNL-NCS-52142, February 29, 1988.
30. P. Schluter and G. Soff, At. Data Nucl. Data Tables **24**, 509 (1979) - 79Sc31
[Internal Pair Conversion Coefficients]

⁴⁴Ti – Comments on evaluation of decay data

by E. Browne

Evaluation Procedures

The *Limitation of Relative Statistical Weights*^[1] (LWM) method, used for averaging numbers throughout this evaluation, provided a uniform approach for the analysis of discrepant data. The uncertainty assigned to the recommended value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

⁴⁴Ti ($T_{1/2} = 60.0$ y) decays 100% by electron capture ($Q(\text{EC}) = 267.5$ (19) keV) to excited levels at 67.9- and 146.2 keV only in ⁴⁴Sc ($T_{1/2} = 3.93$ h), which subsequently decays by $\text{EC} + \beta^+$ to ⁴⁴Ca (stable).

90Sc08 measured the relative emission probabilities of the 1157 -, 67.9- and 78.4-keV gamma rays from a ⁴⁴Ti - ⁴⁴Sc equilibrium source. Since the absolute emission probability of the 1157 -keV gamma ray from ⁴⁴Sc is well known (0.999)^[2], this measurement provided values for the absolute emission probabilities of the 67.9- and 78.4 keV gamma rays as well, thus normalizing the decay scheme of ⁴⁴Ti.

Nuclear Data

⁴⁴Ti is of considerable interest in astrophysics, since it is one of the few long-lived gamma-ray-emitting nuclides expected to be substantially produced during a supernova explosion. Moreover, the solar system abundance of ⁴⁴Ca is believed to have originated from the nucleosynthesis of ⁴⁴Ti and the subsequent decays. The characteristic 1157 -keV gamma ray from ⁴⁴Sc, which was observed from the young supernova remnant Cassiopeia A^[3], opened the possibility of deducing the mass of ⁴⁴Ti that was ejected in the explosion. For this calculation, however, it was needed (among other quantities) a reasonably precise knowledge of the ⁴⁴Ti half-life.

The recommended half-life of ⁴⁴Ti, 60.0 (11) y, is a weighted average (LWM, $\sigma_{\text{int}} = 0.5$, $\chi^2/\nu = 5.6$) of:

- 60.7 (12) y^[4] (method: decay of count rate),
- 59.0 (6) y (98Ah03^[5], method: decay of count rate),
- 60.3 (13) y (98Go05^[6], method: specific activity with beam fragmentation),
- 62 (2) y (98No06^[7], method: decay of count rate),
- 66.6 (16) y (90Al11^[8], method: decay of count rate), and
- 54.2 (21) y (83Fr27^[9], method: specific activity with accelerator mass spectroscopy).

The following results have not been included in the averaging:

- Preliminary results: 58 (10) y^[10] (method: specific activity with beam fragmentation), 39.0 (18) y^[11] (method: specific activity with beam fragmentation), and 63 (3) y (97No06^[12], method: decay of count rate).
- Older measurements: 48.2 (9) y (65Mo07^[13], method: specific activity), and 46.4 (17) y (65Wi05^[14], method: specific activity). These values significantly deviate from recent results, probably because of systematic errors.

Woosley and Diehl^[15] have recommended a half-life of 60 (1) y for ⁴⁴Ti, based on the 1998 values.

Gamma Rays

Energies

⁴⁴Ti emits gamma rays of 67.9 -, 78.4-, and a very weak one of 146.2 keV. The precise gamma -ray energies for the 67.9 - and 78.4-keV transitions given in Table 4.2 (and the values corrected for nuclear recoil, in Table 2.2) are weighted averages (LWM) of results from 63Kl06^[16], 67Ri06^[17], and 91We08^[18] (See Table I). Other: 88Al27^[19] (superseded by 91We08^[18]). The energy of 146.22 (3) keV for the 146 - keV is from level-energy differences. A measured value is: 147.0 (15) keV (67Ri06^[17]).

Table I - ⁴⁴Ti Gamma-ray Energies

	67.9 keV	78.4 keV
91We08 ^[18]	67.8679 (14)	78.3234 (10)*
67Ri06 ^[17]	67.85 (4)	78.38 (4)
63Kl06 ^[16]	67.85 (7)	78.44 (7)
Average	67.8679 (14)	78.36 (3)
χ^2/ν	0.13	1.3

* The uncertainty of 0.0010 was increased to 0.035 to reduce the statistical weight of this measurement from 99.9% to 50%. Original $\chi^2/\nu = 2.4$.

Emission Probabilities

The relative emission probabilities are average values (LWM) from 88Al27^[19], 90Sc08^[20], and 67Ri06^[17], as given in Table II below.

Table II - ⁴⁴Ti Relative Emission Probabilities

Energy keV	67Ri06 ^[17] P _γ (rel.)	88Al27 ^[19] P _γ (rel.)	90Sc08 ^[20] P _γ (rel.)	W. Average (LWM) P _γ (rel.)	χ^2/ν
67.8679 (14)	0.942 (15)*	0.981 (11)	0.960 (15)	0.965 (16) [@]	2.3
78.36 (3)	1.000 (11)*	1.000 (11)	1.000 (13)	1.000 (11) ^{&}	
146.22 (3)	0.0010 (3)	0.00093 (6)	0.00095 (3)	0.00095 (3)	0.05

* Original uncertainties of 0.005 seemed unrealistically low. Evaluator has increased these values.

& Uncertainty is the smallest of the individual values.

[@] Internal uncertainty $\sigma_{int}=0.011$

A factor to normalize relative to absolute emission probabilities was deduced as follows:

- N= 0.955 (15), from the average relative emission probabilities given in Table III column 2, the theoretical conversion coefficients from Section 2.2, and the condition that the total transition intensity to the ground state is 100%, as shown below.

$$[P_{\gamma}(67.8)(1 + \alpha_{68}) + P_{\gamma}(146)(1 + \alpha_{146})] N = 100\%$$

- N=0.974 (13), from the emission probability of the 78-keV gamma ray (0.974 (13)) relative to an absolute probability of 0.999 (1) for the 1157-keV gamma ray in the decay of ⁴⁴Sc in equilibrium with ⁴⁴Ti (90Sc08).

The (unweighted) average of these normalization factors is N_{avg}=0.964 (13) (smallest uncertainty from input values).

Table III gives recommended relative and absolute gamma-ray emission probabilities.

Table III - Recommended Relative and Absolute Gamma-Ray Emission Probabilities

E _γ (keV)	P _γ (rel.)*	P _γ (abs.) ^{&}
67.9	0.965 (16)	0.930 (15)
78.36 (3)	1.000 (11)	0.964 (11)
146.22 (3)	0.00095 (3)	0.00092 (3)

* From Table II, column 5.

[&] Values from column 2 multiplied by N_{avg}(=0.964 (13))

Multipolarities and Conversion Coefficients

The following experimental conversion coefficients: α_K = 0.123 (23) (67Ri06^[17]), α = 0.10 (5) (63Kl06^[16]) for the 67.9-keV gamma ray, and α_K = 0.031 (5) (67Ri06^[17]), α = 0.017 (8) (63Kl06^[16]) for the 78.4-keV gamma ray, suggest E1 and M1 multipolarities for the 67.9 - and 78.4-keV transitions, respectively. Spins of 0- (for the 146-keV level) and 1- (for the 67.9-keV level) require M1 multipolarity for the 78.4 -keV gamma ray. The evaluator has assigned from decay scheme (0 - to 2+)[M2] multipolarity to the 146-keV gamma ray.

Total conversion coefficients also may be deduced from the measured absolute gamma -ray emission probabilities of 90Sc08, by using 0.7 (3)% (88Al27, delayed -coincidence experiment) for the electron -capture feeding to the 67-keV level, and neglecting the very weak 146 -keV transition. These calculations are:

α(67.9) = [1.0/0.935 (15)] -1.0 = 0.069 (17); α(78.4) = [(1.0 - 0.007 (3))/0.974 (13)] -1.0 = 0.019 (14), which agree with the measured values. Where 0.935 (15) and 0.97 4 (13) (90Sc08) are the experimental absolute emission probabilities of the 67.9 - and 78.4-keV gamma rays, respectively. The absolute adopted emission probabilities were not used in this calculation because they are partially based on decay scheme considerations (that include the conversion coefficient of the 67.9-keV gamma ray.)

Table IV shows experimental and theoretical conversion coefficients for the 67.9 -, 78.4-, and 146-keV gamma rays.

Table IV - Conversion Coefficients

E _γ keV	α _T [@] From P _γ (%)	α _T Exp.	α _T [*] Theory	α _K Exp	α _K [*] Theory	Mult.
67.8679 (14)	0.069 (17)	0.10 (5) [#]	0.0845 (25)	0.123 (23) ^{&}	0.0766 (23)	E1
78.36 (3)	0.019 (14)	0.017 (8) [#]	0.032 (1)	0.031 (5) ^{&}	0.0273 (8)	M1
146.22 (3)			0.046 (1)		0.0414 (12)	M2

* Interpolated from ⁷⁶Ba63^[21]

From ⁶³Kl06^[16]

& From ⁶⁷Ri06^[17]

@ See text

The experimental conversion coefficients in Table IV are quite imprecise, therefore, the evaluator has adopted interpolated theoretical values as the recommended conversion coefficients. The interpolation was done with the computer program ICC^[22].

Electron-Capture Transitions

The EC probability to the 146-keV level is given by:

$$\epsilon(146) = [P_{\gamma}(78.4)+e(78.4) + P_{\gamma}(146)+e(146)] \times 100 = 99.5 (11)\% + 0.096 (3)\% = 99.6 (11)\%.$$

For the EC probability to the 0+ ground state of ⁴⁴Sc (0+ to 2+, second forbidden) a log ft >10.6 is expected from the systematic trend for second forbidden transitions (⁹⁸Si17), which corresponds to $\epsilon(0) < 0.04\%$. Using $\epsilon(0)=0.04\%$ and $\epsilon(146) = 99.6 (11)\%$ gives $\epsilon(67.4) = 0.4 (11)\%$. Experimental values for this quantity are 0.7 (3)% (⁸⁸Al27^[19]), and 1.9 (15)% (⁶⁷Ri06^[17]), both measured in γ -x ray coincidence experiments.

Electron-capture probabilities to the various atomic sub-shells, ie. P_K, P_L, P_{M+} in Table 2.1, are theoretical values (⁹⁸Sc28^[23]) calculated with the computer program EC-CAPTURE^[24].

Levels half-life

Table V shows the experimental half-life values for the 67.3 - and 78.4 keV levels, as well as their respective recommended (i.e., average) values.

Table V - ⁴⁴Sc Levels half-life

67.9 keV		78.4 keV	
153 (2) ns	(⁶⁷ Ri06 ^[17])	50 (3) μ s	(⁶³ Kl06 ^[16])
153 (1) ns	(⁶² Th12 ^[25])	49.5 (10) μ s	(⁶⁴ Br27 ^[27])
180 (20) ns	(⁵⁹ Cy90 ^[26])	51.2 (9)* μ s	(⁸⁸ Al27 ^[19])
166 (5) ns	(⁶³ Kl06 ^[16])		
155 (2) ns	(⁷⁵ Gu24 ^[28])		
154.8 (8) ns	(⁸⁸ Al27 ^[19])		
Avg.(LWM) = 154.2 (8) ns		Avg. (LWM) = 50.4 (7) μ s	
$\chi^2/\nu = 1.95$		$\chi^2/\nu = 0.77$	

* The uncertainty was increased from 0.3 ($\chi^2/\nu = 1.4$) to 0.9 to reduce its statistical weight from 91% to 50%.

Atomic Data

The X-ray and Auger-electron probabilities in Section 4 have been calculated using the gamma-ray and electron-capture data that are presented in Section 2, and using atomic data from ⁹⁶Sc06^[29].

Total Average Radiation Energy

Our calculated (RADLST^[30]) total average radiation energy of 268 (3) keV (which includes all the radiations emitted by ⁴⁴Ti), agrees very well with Q(EC) = 267.5 (19) keV (95Au04^[31]) and confirms the quality and completeness of the ⁴⁴Ti decay scheme.

References

1. M.J. Woods and A.S. Munster, *Evaluation of Half-Life Data*, National Physical Laboratory, Teddington, UK, Rep. RS(EXT) 95, (1988) - 88WoZO
2. Richard B. Firestone, Table of Isotopes, eighth edition, John Wiley & Sons, Inc., 1996.
[⁴⁴Sc ε+β⁺ decay, E_γ, I_γ]
3. A.F. Iyudin et al., *Astron. Astrophys.* **284**, L1 (1994)
4. F.E. Wietfeldt et al., *Phys. Rev.* **C59**, 528 (1999)
[T_{1/2}]
5. I. Ahmad et al., *Phys. Rev. Lett.* **80**, 2550 (1998) - 98Ah03
[T_{1/2}]
6. J. Gorres et al., *Phys. Rev. Lett.* **80**, 2554 (1998) - 98Go05
[T_{1/2}]
7. E.B. Norman et al., *Phys. Rev.* **C57**, 2010 (1998) - 98No06
[T_{1/2}]
8. D.E. Alburger and G. Harbottle, *Phys. Rev.* **C41**, 2320 (1990) - 90Al11
[T_{1/2}]
9. D. Frekers et al., *Phys. Rev.* **C28**, 1756 (1983) - 83Fr27
[T_{1/2}]
10. J. Meissner et al., in *Nuclei in the Cosmos III*, edited by Maurizio Busso and Claudia M. Raiteri, AIP Conf. Proc. No. 327 (AIP, New York, 1995), p. 303.
[T_{1/2}]
11. J. Meissner, Ph. D. thesis, University of Notre Dame, 1996.
[T_{1/2}]
12. E.B. Norman et al., in Proceedings of the International Conference on *Nuclei in the Cosmos IV*, *Nucl. Phys.* **A621**, 92c (1997) - 97No06
[T_{1/2}]
13. P.E. Moreland and D. Heymann, *J. Inorg. Nucl. Chem.* **27**, 493 (1965) - 65Mo07
[T_{1/2}]
14. J. Wing et al., *J. Inorg. Nucl. Chem.* **27**, 487 (1965) - 65Wi05
[T_{1/2}]
15. Stan Woosley and Roland Diehl, *Physics World* **11**, No. 7, 22 (1998)
[T_{1/2}]
16. J.K. Kliwer et al., *Nucl. Phys.* **49**, 328 (1963) - 63K106
[E_γ]
17. R.A. Ristinen and A.W. Sunyar, *Phys. Rev.* **153**, 1209 (1967) - 67Ri06
[E_γ, I_γ]
18. C. Wesselborg and D.E. Alburger, *Nuc. Instrum. Meth.* **A302**, 89 (1991) - 91We08
[E_γ]
19. D.E. Alburger and E.K. Warburton, *Phys. Rev.* **C38**, 1843 (1988) - 88Al27
[I_γ]

20. U. Schötzig, Nucl. Instrum. Meth. **A286**, 523 (1990) - 90Sc08
[I_γ]
21. I.M. Band et al., At. Data Nucl. Data Tables **18**, 433 (1976) - 76Ba63
[Theoretical internal conversion coefficients]
22. E. Yakusev and N. Coursol, ICC, a computer program to interpolate internal conversion coefficients, 1998.
23. E. Schönfeld, App. Rad. Isot. **49**, 1353 (1998) - 98Sc28
[P_K, P_L, P_M]
24. E. Schönfeld, F.Y. Chu, and E. Browne, EC-CAPTURE, a computer program to calculate electron capture probabilities to atomic sub-shells, 1998.
25. P. Thieberger, Arkiv Fysik **22**, 127 (1962) - 62Th12
[Level T_{1/2}]
26. E.W. Cybulska and L. Marquez, Nuovo Cimento **14**, 479 (1959) - 59Cy90
[Level T_{1/2}]
27. K. Bandi et al., Nucl. Phys. **59**, 33 (1964) - 64Br27
[Level T_{1/2}]
28. V.P. Gupta and D.K. Gupta, Indian J. Pure Appl. Phys. **13**, 334 (1975) - 75Gu24
[Level T_{1/2}]
29. E. Schönfeld and H. Janßen, Nucl. Instrum. Meth. Phys. Res. **A369**, 527 (1996) - 96Sc06
[X rays, ω_K]
30. *The Program RADLST*, Thomas W. Burrows, report BNL-NCS-52142, February 29, 1988.
31. G. Audi and A.H. Wapstra, Nucl. Phys. **A595**, 409 (1995) - 95Au04
[Q-value]
32. B. Singh, J.L. Rodriguez, S.S.M. Wong, and J.K. Tuli, Nucl. Data Sheets **84**, 487 (1998) - 98Si17
[Systematics of log *ft*]

**⁴⁵Ca - Comments on evaluation of decay data
by M.M. Bé**

This evaluation was completed in February 2012 with a literature cut-off by the same date.

Decay Scheme

⁴⁵Ca disintegrates by beta minus emission to the ⁴⁵Sc ground state mainly, with a very small branch of 0,002 % to the ⁴⁵Sc first excited level of 12,4 keV.

Level energies, spins and parities (J^π) were taken from the evaluation by Burrows (2008Bu01).

Nuclear Data

Half-life

The recommended value was derived from measurements listed in Table 1.

Table 1. Measurement results and recommended value of ⁴⁵Ca half-life.

Reference	Value (d)	Uncertainty (d)	Remarks
1952De01	163,5	4,0	GM, Stat. uc
1957Th**	153	2	
1959Ca12	167	3	PC
1961Wy01	165,1	0,7	PC
1965An07	162,63	0,11	4 π pc
1970Si24	163	2	PC
1994Lo04	162,67	0,25	LSC
$\chi^2/n-1 =$	0,34		
χ^2 crit. =	0,38		
UWM =	162,95		
LWM = (recommended)	162,64	0,11	U _{cint} = 0,10 ; U _{cext} = 0,018

Conventional designations in the fourth column:

Measurement of counting rate decrease by Geiger-Müller counter (GM), proportional counter (PC), 4 π proportional counter (4 π pc), Liquid Scintillation Counting (LSC).

Stat. uc: statistical uncertainty component.

Values from Thiry (1957Th**), Caswell (1959Ca12) and Wyatt (1961Wy01) have been found outliers due to Chauvenet's criterion. The set of the remaining four values is consistent, and then the adopted value is the weighted mean with the lowest of the experimental uncertainties.

Decay Energy and Characteristics of Electron Emissions

The decay energy of ⁴⁵Ca has been adopted from the evaluation of Audi *et al.* (2011AuZZ),

$Q_{\beta^-} = 258,0$ (7) keV.

It should be noted that in the previous evaluation (2003Au03) $Q_{\beta^-} = 255,8$ (8) keV. This difference seems due to a change of 2,5 keV in the ⁴⁵Sc mass.

Several papers report measurements of the ⁴⁵Ca β^- spectrum (1950Ma03, 1950Ke60, 1967Ha39); by using a Kurie plot, the nature of the transition $gs \rightarrow gs$ was determined to be allowed.

With this hypothesis, a mean beta energy of 76,8 keV was calculated with the BetaShape program (2011Mo**), which is based on theoretical considerations. It can be compared with the experimental value of 74,6 (30) keV obtained by Caswell (1952Ca10).

A weak β^- transition to the 12,4 keV level was deduced from the observation of K conversion electrons by Freedman *et al.* (1965Fr12). The ratio of the K conversion electrons over the total β emissions was measured being:

$$I(\text{ce}_K)/I\beta = 1,4 \times 10^{-5} (+8 -3)$$

The adopted value for this ratio is $I(\text{ce}_K)/I\beta = 1,7$ (6) $\times 10^{-5}$ in application of the method described in Audi (2003Au03, page 21) to experimental data with asymmetric uncertainties.

The internal conversion coefficients for a M2 transition were calculated with the BrIcc program in the “frozen orbital approximation” (Kibédi *et al.* - 2008Ki07), that is: $\alpha_T = 423$ (9) and $\alpha_K = 362$ (8).

Then, the probability of the β^- transition to the 12,4 keV level is $2,0$ (7) $\times 10^{-3}$ %.

Gamma-ray transition and emission

From the results above, the probability of the 12,4 keV gamma-ray transition is $2,0$ (7) $\times 10^{-3}$ % and the gamma-ray emission intensity is negligible.

References

- 1950Ma03 P.Macklin, et al. Phys. Rev. 77 (1950) 137. Beta spectrum
 1950Ke60 B.H.Ketelle. Phys. Rev. 80 (1950) 758. Beta spectrum
 1952Ca10 R.S. Caswell. Phys. Rev. 86,1 (1952) 82. Beta emission energies
 1952De01 C.F.G.Delaney, J.H.J.Poole. Phys. Rev. 89 (1953) 529. Half-life
 1957Th** H.Thiry. Bull. Soc. Royale Sci. Liège 25 (1957) 29. Half-life
 1959Ca12 J.P.Cali, L.F.Lowe. Nucleonics 17 (1959) 86. Half-life
 1961Wy01 E.I.Wyatt, S.A.Reynolds, T.H.Handley, W.S.Lyon, H.A.Parker. Nucl. Sci. Eng. 11 (1961) 74. Half-life
 1965An07 S.C. Anspach, L.M.Cavallo, S.B.Garfinkel, J.M.R Hutchinson, C.N.Smith. NP-15663 (1965). Half-life
 1965Fr12 M.S. Freedman, F.T.Porter, F.Wagner Jr.. Phys. Rev. B563 (1965) 140. Gamma-ray energies K-Conv. Elec. emission probabilities, K/L
 1967Ha39 F.Hareux. CEA Report R 3315 (1967). Beta spectrum
 1970Si24 B.Sitar, J.Chrapan, J.Oravec, K.Durcek. Jad. Energ. 16 (1970) 303. Half-life
 1994Lo04 J.M.Los Arcos, L.Rodriguez, M.Roteta, E.Garcia-Toraño. Nucl. Instrum. Methods Phys. Res. A339 (1994) 164. Half-life
 2003Au03 G.Audi, A.H.Wapstra, C.Thibault. Nucl. Phys. A729 (2003) 129. Q
 2008Ki07 T.Kibédi, W.Burrows, M.B.Trzhaskovskaya, P.M.Davidson, C.W.Nestor Jr.. Nucl. Instrum. Methods Phys. Res. A589 (2008) 202. ICC
 2011Mo** X.Mougeot, *et al.* RadioCarbon, P. Cassette Editor, ISBN 978 0 9638314 7 7 (2010) 249. Beta emission energy
 2011AuZZ G.Audi, W.Meng. Private communication (2011). Q_{β^-}

⁴⁶Sc - Comments on evaluation of decay data by R. G. Helmer

1 Decay Scheme

The only levels in ⁴⁶Ti below the decay energy are those populated in this ⁴⁶Sc β⁻ decay, so that portion of the decay scheme is complete. However, ⁴⁶Sc can also electron-capture decay, ε, to levels in ⁴⁶Ca with a decay energy of 1368 keV. The available levels are 0⁺ at 0 keV and 2⁺ at 1346 keV with ε branches that are 4th forbidden and 2nd forbidden, respectively. From systematics (1998Si17), the corresponding log *ft* limits are ≥ 22.5 and ≥ 10.6, and the deduced P_{ε+β+} limits are ≤ 1.0 × 10⁻¹² % and ≤ 2.5 × 10⁻⁶ %, respectively. Therefore, these ε branches are negligible.

The J^π values and half-lives for the excited levels are from Adopted Levels in Nuclear Data Sheets (2000Wu08).

2 Nuclear Data

Q value is from Audi and Wapstra 1995 (1995Au04).

The half-life values available are, in days:

85 (1)	(1940Wa01)	omitted from analysis
84.1 (3)	(1956Sc87)	omitted from analysis
83.89 (12)	(1957Ge07)	omitted from analysis
84.4 (2)	(1957Wr37)	omitted from analysis
83.80 (3)	(1965An07)	superseded by 1982HoZJ
84.34 (13)	(1974Cr05)	omitted as outlier
83.75 (3)	(1977MeZP)	superseded by 1980RuZY
83.819 (6)	(1980Ho17)	
83.79 (6)	(1980Ol03)	
83.752 (15)	(1980RuZY)	
83.79 (6)	(1982HoZJ)	superseded by 1992Un01
83.752 (15)	(1982RuZV)	same as 1980RuZY
83.73 (12)	(1983Wa26)	
83.83 (7)	(1992Un01)	
83.788 (22)	Adopted value	

This set of values is inconsistent which causes the adopted value to depend on the choice of the values used and the "averaging" method used. The values have decreased over time; the unweighted average of the four not superseded values before 1978 (1940Wa01, 1956Sc87, 1957Ge07, and 1957Wr37) is 84.18, whereas the same average for the five values after 1978 (1980Ho17, 1980Ol03, 1980RuZY, 1983Wa26, and 1992Un01) is 83.78. The values reported before 1960 were omitted from the analysis since it would have been difficult to determine the presence of a small amount of a longer-lived impurity with the spectroscopy methods then available.

The discrepancy among the values is illustrated by the values of 84.34(13) (1974Cr05),

83.819(6) (1980Ho17), and 83.752(15) (1980RuZY). The first two values differ by 0.52(13) and the last two by 0.067(16), or about 4σ in each case. The latter two values have the greatest weight in any weighted average, so the results will depend on how the analysis modifies their relative weight, and the first value will give the largest contribution to the χ^2 value. Of the remaining six values not superseded, that of 84.34(13) (1974Cr05) is considered an outlier and is omitted.

For the remaining five values not superseded, the following averages are obtained:

unweighted	83.784 (19)
weighted	83.810, $\sigma_{\text{int}}=0.006$, reduced- $\chi^2=4.46$, $\sigma_{\text{ext}}=0.013$
RAJEVAL	83.776 (20)
Normalized residuals	83.793 (16)
LRSW - weighted average	83.788, $\sigma_{\text{int}}=0.010$, reduced- $\chi^2=1.67$, $\sigma_{\text{ext}}=0.022$ and $\sigma_{\text{LRSW}}=0.031$

The RAJEVAL method (1992Ra08) increases both of the two smallest uncertainties, namely, 0.006 to 0.043 and 0.015 to 0.026, which causes the value of 1980RuZY to have the largest weight. The Normalized Residuals method (1992Ja06) also increases both of the two smallest uncertainties but by different amounts, namely, 0.006 to 0.022 and 0.015 to 0.028, which leaves the value of 1980Ho17 with the largest weight, but only by a small amount. In contrast, the Limitation of Relative Statistical Weight, LRSW, method (1985ZiZY, 1992Ra08) only increases the most precise uncertainty, namely that of 1980Ho17, from 0.006 to 0.014 in order to reduce its relative weight to 50% from its initial 84%. The LRSW method expands the final uncertainty to 0.031 in order to include the most precise value. [The LRSW method finally suggests the unweighted average of 83.96(14), but that choice is not accepted here.]

The results from the RAJEVAL, Normalized residuals, and LRSW methods all are in good agreement and the adopted value, 83.788(22) is taken as the latter value with its external uncertainty.

2.1 β^- Transitions

The β^- branch to the ground state of ^{46}Ti is 4th forbidden with an expected $\log ft \geq 22.5$ (1998Si17) and a corresponding $P_{\beta^-}(0) \leq 1 \times 10^{-11} \%$, the measured limit is $\leq 1 \times 10^{-4} \%$ (1954Ke04).

Similarly, for the 2nd forbidden decay to the 889 level, the expected $\log ft \geq 10.6$ which corresponds to $P_{\beta^-}(889) \leq 0.8 \%$. The measured I_{β^-} to this level are 0.096(1) (1954Ke04), 0.0036(7) (1956Wo09), ≤ 0.06 (1950Mo62), and ≤ 0.05 (1950So57). Some previous evaluators (e.g., 1986Al19) have assigned $I_{\beta^-}(889) = 0.0036(7)$ because it is consistent with the limits of 1950Mo62 and 1950So57. However, this evaluator has some reservations about the resulting precision for I_{β^-} (2009) and, therefore, has expanded the uncertainty and gives $I_{\beta^-}(889) = 0.004 \%$ (+364), which is consistent with the two limits and the value of 1956Wo09, and thus $I_{\beta^-}(2009) = 99.996(+4-36)$.

If symmetric uncertainties are required, as in ENSDF, for these quantities, $I_{\beta^-}(889) = 0.02(2)$ and $I_{\beta^-}(2009) = 99.98(2)$, adopted values.

The β^- average energies and $\log ft$ values are from LOGFT code.

2.2 Gamma Transitions

The J^π assignments are from the Adopted Levels in the Nuclear Data Sheets (2000Wu08) and these imply the two γ -rays have E2 multipolarities.

The internal-conversion coefficients were interpolated from the Band tables (1976Ba63).

The internal-pair-formation coefficient was interpolated from the theoretical values (1979Sc31) and is $IPFC(1120) = 0.0000022$ (4). This value is only about 2 % of the corresponding internal-conversion coefficient and, therefore, is negligible.

3 Atomic Data

The data are from 1996Sc06.

3.1 and 3.2

None

4 Radiation Emissions

4.1 Electron Emission

The emission intensities are calculated from the atomic data and the decay data.

4.2 Photon Emission

The γ -ray energies are from 2000He14 for the 889 and 1120 lines and the 2009 energy is the sum of these values corrected for nuclear recoil.

The relative γ -ray emission probability of the 2009-keV γ -ray is from 1980Fu07.

The emission probability of the 889 -keV γ -ray is $[100.0 - P_\gamma(2009)] / [1.0 + \alpha(889)] = 99.999987(10)/1.000167(5) = 99.9833(5)$ where the uncertainty is 5 ppm from the $(1.0 + \alpha)$ term.

That of the 1120 -keV γ -ray is $[I_{\beta}(2009) - P_\gamma(2009)] / [1.0 + \alpha(1120)] = 99.996(+4 - 36)/1.000095(3) = 99.986(+4-36)$, with symmetric uncertainties 99.98 (2). Here $\alpha(2009)$ has been neglected.

6 References

- 1940Wa01 - H. Walke, Phys. Rev. **57**(1940)163 [T_{1/2}]
 1950Mo62 - M. L. Moon, M. A. Waggoner, A. Roberts, Phys. Rev. **79**(1950)905 [I_β-]
 1950So57 - B. N. Sorensen, B. M. Dale, J. D. Kurbatov, Phys. Rev. **79**(1950)1007 [I_β-]
 1954Ke04 - G. L. Keister, F. H. Schmidt, Phys. Rev. **93**(1954)140 [I_β-]
 1956Sc87 - R. P. Schuman, M. E. Jones, A. C. McWherter, Jnorg. Nucl. Chem **3**(1956)160 [T_{1/2}]
 1956Wo09 - J. L. Wolfson, Can. J. Phys. **34**(1956)256 [I_β-]
 1957Ge07 - K. W. Geiger, Phys. Rev. **105**(1957)1539 [T_{1/2}]
 1957Wr37 - H. W. Wright, E. I. Wyatt, S. A. Reynolds, W. S. Lyon, T. H. Handley, Nuclear Sci. Eng. **2**(1957)427 [T_{1/2}]
 1965An07 - S. C. Anspach, L. M. Cavallo, S. B. Garfinkel, J. M. R. Hutchinson, C. N. Smith, NP-15663 (1965) [T_{1/2}]
 1974Cr05 - P. J. Cressy, Jr., Nucl. Sci. Eng. **55**(1974)450 [T_{1/2}]
 1976Ba63 - I. M. Band, M. B. Trzhaskovskaya, M. A. Listengarten, Atomic Data Nucl. Data Tables **18**(1976)433 [α]
 1977MeZP - J. S. Merritt, F. H. Gibson, AECL-5696 (1977) 40 [T_{1/2}]

- 1979Sc31 - P. Schluter, G. Soff, At. Data Nucl. Data Tables **24**(1979)509 [IPFC]
1980Fu07 - M. Fujishiro, Y. Satoh, K. Okamoto, T. Tsujimoto, Can. J. Phys. **58**(1980)1712 [P_{γ}]
1980Ho17 - H. Houtermans, O. Milosevic, F. Reichel, Int. J. Appl. Radiat. Isotop. **31**(1980)153 [$T_{1/2}$]
1980OI03 - J. B. Olomo, T. D. MacMahon, J. Phys.(London) **G6**(1980)367 [$T_{1/2}$]
1980RuZY - A. R. Rutledge, L. V. Smith, J. S. Merritt, AECL-6692 (1980) [$T_{1/2}$]
1882HoZJ - D. D. Hoppes, J. M. R. Hutchinson, F. J. Schima, M. P. Unterweger, NBS-SP-626 (1982) 85 [$T_{1/2}$]
1982RuZV - A. R. Rutledge, L. V. Smith, J. S. Merritt, NBS-SP-626 (1982) 5 [$T_{1/2}$]
1983Ru04 - A. R. Rutledge, L. V. Smith, J. S. Merritt, Nuk Instrum. Methods **206**(1983)211 [$T_{1/2}$]
1983Wa26 - K. F. Walz, K. Debertin, H. Schrader, Int. J. ApplRadiat. Isotop. **34**(1983)1191 [$T_{1/2}$]
1985ZiZY - W. L. Zijp, report ECN-179, Petten (1985) [analysis methodology]
1986Al19 - D. E. Alburger, Nucl. Data Sheets 49 (1986) 237 [evaluation]
1992Ra08 - M. U. Rajput, T. D. MacMahon, Nucl. Instr. Meth. **A312**(1992)289 [analysis methodology]
1992Un01 - M. P. Unterweger, D. D. Hoppes, F. J. Schima, Nucl. Instr. Meth. Phys. Res. **A312**(1992)349 [$T_{1/2}$]
1995Au04 - G. Audi, A. H. Wapstra, Nucl. Phys. **A595**(1995)409 [Q]
1996Sc06 - E. Schönfeld, H. Janßen, Nucl. Instr. Meth. **A369**(1996)527 [ω]
1998Si17 - B. Singh, J. L. Rodrigues, S. S. M. Wong, J. K. Tuli, Nucl. Data Sheets **84**(1998)487 [log ft systematics]
2000He14 - R. G. Helmer, C. van der Leun, Nucl. Instr. Meth. **A450**(2000)35 [E_{γ}]
2000Wu08 - S.-C. Wu, Nucl. Data Sheets **91**(2000)1 [J^{π} , multipolarities]

⁴⁷Sc - Comments on evaluation of decay data by X. Mougeot

This evaluation was completed in 2013, taking into account the available literature by November 2013. This radionuclide was first identified correctly by 1945HI04 who associated to ⁴⁷Sc the correct half-life and beta electron energies.

The study of the β^- decay of ⁴⁷Sc by 1986RE12 is the most precise and complete one. Notice also that it comes from a collaborative work of three metrological laboratories (CBNM, NPL, LNHB). Therefore, the final evaluated values adopted here are very similar to their results.

1 Decay Scheme

The decay scheme is complete for this β^- decay. The ground state of ⁴⁷Sc is 7/2⁻ (2007BU08). From $Q = 600,8$ (19) keV (2012WA38), two levels of ⁴⁷Ti are reachable (2007BU08): the ground state, 5/2⁻ and, the first excited state at 159 keV, 7/2⁻. The two beta transitions populating them are allowed.

Consistency of the decay scheme was checked in Section 4.2.2. The effective Q value resulting from total energy balance considerations is: $Q_{\text{eff}} = 601$ (35) keV, consistent with $Q = 600,8$ (19) keV from 2012WA38 adopted in this evaluation.

2 Nuclear Data

The Q value adopted in this evaluation, $Q = 600,8$ (19) keV, is from Wang *et al.* 2012 (2012WA38).

2.1 ⁴⁷Sc Half-life

A full list of the half-life values available by November 2013 is summarized in Table 1, including the experimental methods used. Values reported without uncertainty were not taken into account in the present evaluation.

The original value from 1968ME07 is 3,345 (3) d. In their paper, the authors gave all the measured values they obtained, reproduced in Table 2, from which they calculated a weighted average. The LWEIGHT program excludes the value 3,332 (3) d by Chauvenet's criterion and gives 3,3476 (17) d. The value 3,332 (3) d is therefore excluded of this evaluation. Dispersion of data was used by the evaluator to determine a systematic uncertainty, which was included to give: 3,348 (5) d. Nevertheless, the adopted value is provided by the arithmetical mean because the values are from the same authors using the same method. Including a systematic uncertainty, it was found 3,347 (5) d, consistent with the value provided by LWEIGHT.

The original value from 1969RA16 is 82,12 (9) h. It was just converted into days by the evaluator: 3,4217 (38) d.

The original value from 1986RE12 is 3,3492 (6) d, which comes from a weighted mean of six half-life measurements, given in Table 3: three from NPL using a ionization chamber, two from CBNM using Ge(HP), and one from CBNM using NaI(Tl) integral counting. An arithmetical mean was performed with the NPL values and the minimum uncertainty was chosen: 3,3484 (6) d. Identically, an arithmetical mean was determined from the measurements from CBNM: 3,3502 (13) d. The uncertainty was chosen to be the one from the arithmetical mean because CBNM used two different techniques.

Considering the values of Table 1 with uncertainties, the data set is discrepant. LWEIGHT recommends an unweighted average and expands the uncertainty so range includes the most precise value from 1986RE12: 3,3484 (6) d. Actually, there are two groups of consistent measurements: the ones obtained from beta measurements only, and the ones obtained using gamma measurements (including sometimes also beta measurements). This fact was highlighted by 1980MO26 but without giving an explanation. The evaluated half-life from the first group is 3,4219 (37) d. LWEIGHT excludes the value from 1956LI38 by Chauvenet's criterion, and adopts a weighted average with internal uncertainty. The evaluated half-life from the second group is 3,3485 (9) d. LWEIGHT excludes the values from 1955LY34, 1963HO17 and 1968BA33 by

Chauvenet’s criterion, and uses a weighted average with external uncertainty. The latter evaluated value was preferred by the evaluator because it includes the value from 1986RE12.

To conclude, the recommended half-life for the β^- decay of ⁴⁷Sc is: $T_{1/2} = 3,3485$ (9) d.

Table 1: ⁴⁷Sc half-life measurements

Reference	$T_{1/2}$ (d)	Method	Comments
1945HI04	3,40	β	Rejected: no uncertainty
1949KR12	3,43 (3)	β	
1953CH16	3,4	$\beta+\gamma$	Rejected: no uncertainty
1953CO44	3,40 (5)	β	
1953DU22	3,44	β	Rejected: no uncertainty
1953MA64	3,44 (5)	β	
1955LY34	3,45 (10)	$\beta + \gamma$	
1956LI38	3,3 (1)	β	
1959PO64	3,45 (6)	β	
1963HO17	3,38 (9)	γ	
1964MI07	3,40	γ	Rejected: no uncertainty
1968BA33	3,3 (1)	γ	
1968ME07	3,347 (5)	$\beta + \gamma$	Re-evaluated from 3,345 (3) d
1969RA16	3,4217 (38)	β	Calculated from 82,12 (9) h
1980MO26	3,341 (3)	γ	
1986RE12	3,3502 (13)	γ	Evaluated from the CBNM measurements
1986RE12	3,3484 (6)	$\beta + \gamma$	Evaluated from the NPL measurements
Evaluated	3,3485 (9)		From γ and $\beta + \gamma$ measurements

Table 2: ⁴⁷Sc half-life measurements from 1968ME07

Comments	$T_{1/2}$ (d)
	3,344 (3)
	3,347 (3)
	3,352 (3)
Excluded by Chauvenet’s criterion	3,332 (3)
	3,343 (6)
	3,351 (6)
Weighted average	3,345 (3)
Re-evaluated	3,347 (5)

Table 3: ⁴⁷Sc half-life measurements from 1986RE12

Comments	$T_{1/2}$ (d)
CBNM NaI(Tl) integral counting	3,3500 (6)
CBNM Ge(HP)	3,3480 (31)
CBNM Ge(HP)	3,3525 (15)
Evaluated	3,3502 (13)
NPL Ionization Chamber	3,3478 (10)
NPL Ionization Chamber	3,3491 (19)
NPL Ionization Chamber	3,3483 (6)
Evaluated	3,3484 (6)

2.2 Half-life of the first excited state of ⁴⁷Ti

The half-life of the first excited state of ⁴⁷Ti does not come from the β^- decay of ⁴⁷Sc. It was taken in 2007BU08: $T_{1/2}(159 \text{ keV}) = 210$ (6) ps.

2.3 Beta Transitions

Probabilities of the two beta transitions involved in the ⁴⁷Sc decay were determined by two methods in the literature: from direct β^- -electron measurements, or deduced from the absolute gamma-ray probability P_γ .

The β^- -electron measurements are given in Table 4. $P_{\beta 1}$ is associated with the ⁴⁷Sc ground state to the ⁴⁷Ti ground state transition, and $P_{\beta 2}$ is associated with the ⁴⁷Sc ground state to the 159 keV ⁴⁷Ti excited state transition. Using only the consistent values with uncertainties, LWEIGHT calculated a weighted average with

internal uncertainty and a found a reduced- χ^2 of 0,8 in both cases (P_{β_1} and P_{β_2} are correlated values). The resulting values are given in Table 4.

Table 4: ⁴⁷Sc beta transition probabilities from direct β -electron measurements. P_{β_1} is for the g.s. to g.s. transition, and P_{β_2} is for the g.s. to 159 keV state transition.

Reference	P_{β_1}	P_{β_2}
1953CH16	0,34 (4)	0,66 (4)
1955LY34	0,34 (3)	0,66 (3)
1955NI15	0,36	0,64
1956GR12	0,4	0,6
1956LI38	0,26	0,74
1964MI07	0,27 (4)	0,73 (4)
1986RE12	0,326 (14)	0,674 (14)
Evaluated	0,325 (12)	0,675 (12)

The absolute gamma-ray intensity was evaluated in Section 4.2.2: $I_\gamma(159 \text{ keV}) = 0,681$ (5). With the total internal conversion coefficient calculated in Section 2.4.2, it is found: $P_{\beta_2} = P_\gamma = I_\gamma(1 + \alpha_T) = 0,685$ (5). Thus: $P_{\beta_1} = 0,315$ (5). Notice that the measured $\alpha_T = 0,0045$ (3) from 1986RE12 leads to $P_{\beta_2} = 0,684$ (5) which is consistent and very close despite a discrepancy of about 30 % between the measured and calculated α_T .

The P_{β_1} and P_{β_2} values evaluated from direct β -electron measurements, or deduced from the absolute gamma-ray probability I_γ are consistent. In order to obtain a consistent decay scheme, the most precise values that come from I_γ were preferred. Therefore, the recommended values are:

$$P_{\beta_2} = 0,685 \text{ (5)} \text{ and } P_{\beta_1} = 0,315 \text{ (5)}.$$

These two beta transitions are allowed. Average energies and $\log ft$ values are from the LOGFT program.

2.4 Gamma Transition

The evaluated energy of the gamma transition is from 2007BU08 (see Section 4.2.1).

2.4.1 Mixing ratio and multipolarity

The mixing ratio of the first excited state of ⁴⁷Ti was taken from 2007BU08. It was evaluated using the half-life, B(E2) and the angular distribution of the emitted gamma-ray. The recommended mixing ratio is then: $\delta = -0,099$ (9). The corresponding multipolarity of the gamma transition is thus: **M1 + 0,97 (17) % E2**.

2.4.2 Internal conversion coefficients

The internal conversion coefficients of the gamma transition in ⁴⁷Ti were calculated with the BrIcc program (2008KI07) within the Frozen Orbitals approximation, using the mixing ratio from 2007BU08 given previously.

Notice the following measurement results: $\alpha_T = 0,0036$ (9) from 1953CH16 ; $\alpha_K/\alpha_L \sim 10$ from 1953CO44 ; $P_{eK} = 0,00277$ (14), $\alpha_K = 0,00406$ (21), $\alpha_K/\alpha_L \sim 10$, $\alpha_{L+} = 0,00041$ (10) and $\alpha_T = 0,0045$ (3) from 1986RE12.

3 Atomic Data

3.1 Fluorescence yields

The fluorescence yields are taken from 1996SC06.

3.2 X Radiations and Auger electrons

The X-ray and Auger electron data intensities are from Schönfeld and Janßen (1996SC06).

4 Gamma Emissions

4.2.1 Gamma-ray energies

The adopted energy of the first excited state of ⁴⁷Ti is from 2007BU08. Independent evaluation, described below, was carried out by the evaluator. Results agree very well. The one from 2007BU08 was preferred because of the completeness of their evaluation.

The gamma-ray energy measurements are given in Table 5. Values without uncertainty were not taken into account. The values from 1953CH16, 1955LY34 and 1955NI15 were excluded by the LWEIGHT program during the statistical process. The resulting data set is consistent and a weighted average was used with the internal uncertainty. The reduced- χ^2 is 0,05. The evaluated value of 159,382 (14) is very close to the value from 2007BU08.

Table 5: E_γ measurements of the levels of ⁴⁷Ti.
Final adopted value in red is from 2007BU08.

Reference	Energy (keV)
1953CH16	185 (7)
1953CO44	159,5
1955LY34	157 (7)
1955NI15	167 (2)
1956GR12	159 (2)
1956LI38	160 (5)
1964MI07	163
1967KO01	159,2 (5)
1969WO02	159,39 (5)
1972GEZG	159,381 (15)
Evaluated	159,382 (14)
2007BU08	159,373 (12)

4.2.2 Gamma-ray emission intensities

The measurement of the absolute intensity of the γ ray occurring in the ⁴⁷Sc decay requires the determination of the source activity. This was done only twice, in 1968BA33 and in 1986RE12, using a 4π β - γ coincidence method.

The value from 1968BA33 is $I_\gamma = 0,685$ (27).

The original value from 1986RE12 is $I_\gamma = 0,683$ (4), which comes from a weighted mean of four measurements, given in Table 6. An arithmetical mean was performed with the CBNM values and the minimum uncertainty was chosen: 0,6745 (40). Identically, the LNHB values led to: 0,6880 (40).

Then, LWEIGHT was used for these three values. From this consistent data set, LWEIGHT used a weighted average with its external uncertainty and found a reduced- χ^2 of 2,6. This evaluation leads to:

$$I_\gamma(159 \text{ keV}) = 0,681 (5).$$

Table 6: ⁴⁷Sc I_γ measurements.

Comments	I_γ	
CBNM Ge(HP)	0,678 (5)	} Arithmetical mean: 0,6745 (40)
CBNM Ge(HP)	0,671 (8)	
LNHB NaI(Tl)	0,687 (4)	} Arithmetical mean: 0,6880 (40)
LNHB Ge(HP)	0,689 (7)	
1968BA33	0,685 (27)	0,685 (27)
Evaluated		0,681 (5)

5. Electron Emissions

The β^- intensities were evaluated as described above in Section 2.3.

The energies of the conversion electrons were calculated from the gamma-transition energy given in Section 4.2.1 and the electron binding energies.

The emission intensities of the conversion electrons were calculated using the internal conversion coefficients (see Section 2.4.2). The emission intensities of K-Auger electrons were calculated using the transition probabilities given in Sections 2.3 and 2.4, the atomic data given in Section 3, and the internal conversion coefficients.

Consistency of the decay scheme was tested from total energy balance considerations. The resulting effective Q value, $Q_{\text{eff}} = 600,8$ (41) keV, is consistent with $Q = 600,8$ (19) keV from 2012WA38 adopted in this evaluation.

6. References

- 1945HI04 C.T. Hibdon, M.L. Pool, Phys. Rev. 67, 313 (1945) [First correct identification]
- 1949KR12 N.L. Krisberg, M.L. Pool, Phys. Rev. 75, 1693 (1949) [$T_{1/2}$]
- 1953CH16 L.S. Cheng, M.L. Pool, Phys. Rev. 90, 886 (1953) [$T_{1/2}$, P_β , ICC, E_γ]
- 1953CO44 J.M. Cork, J.M. LeBlanc, M.K. Brice, W.H. Nester, Phys. Rev. 92, 367 (1953) [$T_{1/2}$, ICC, E_γ]
- 1953DU22 J.E. Duval, M.H. Kurbatov, J. Am. Chem. Soc. 75, 2246 (1953) [$T_{1/2}$]
- 1953MA64 L. Marquez, Phys. Rev. 92, 1511 (1953) [$T_{1/2}$]
- 1955LY34 W.S. Lyon, B. Kahn, Phys. Rev. 99, 728 (1955) [$T_{1/2}$, P_β , E_γ]
- 1955NI15 R.T. Nichols, E.N. Jensen, Phys. Rev. 100, 1407 (1955) [P_β , E_γ]
- 1956GR12 W.E. Graves, S.K. Suri, Phys. Rev. 101, 1368 (1956) [P_β , E_γ]
- 1956LI38 L.J. Lidofsky, V.K. Fischer, Phys. Rev. 104, 759 (1956) [$T_{1/2}$, P_β]
- 1959PO64 A. Poularikas, R.W. Fink, Phys. Rev. 115, 989 (1959) [$T_{1/2}$]
- 1963HO17 S. Hontzeas, L. Yaffe, Can. J. Chem. 41, 2194 (1963) [$T_{1/2}$]
- 1964MI07 S.C. Misra, U.C. Gupta, N.P.S. Sidhu, Nucl. Phys. 51, 174 (1964) [$T_{1/2}$, P_β , E_γ]
- 1967KO01 J. Konijn, E.W.A. Lingeman, S.A. De Wit, Nucl. Phys. A 90, 558 (1967) [E_γ]
- 1968BA33 Z.T. Bak, P. Rieh, Int. J. Appl. Radiat. Isot. 19, 593 (1968) [$T_{1/2}$, P_γ]
- 1968ME07 J.W.T. Meadows, V.A. Mode, J. Inorg. Nucl. Chem. 30, 361 (1968) [$T_{1/2}$]
- 1969RA16 H. Ravn, J. Inorg. Nucl. Chem. 31, 1883 (1969) [$T_{1/2}$]
- 1969WO02 R.E. Wood, J.M. Palms, P. Venugopala Rao, Nucl. Phys. A 126, 300 (1969) [E_γ]
- 1972GEZG R.J. Gehrke, ANCR-1088, p. 392 (1972) [E_γ]
- 1980MO26 H. Mommsen, I. Perlman, J. Yellin, Nucl. Instrum. Methods 177, 545 (1980) [$T_{1/2}$]
- 1986RE12 D. Reher, H.H. Hansen, R. Vaninbroux, M.J. Woods, C.E. Grant, S.E.M. Lucas, J. Bouchard, J. Morel, R. Vatin, Int. J. Appl. Radiat. Isot. 37, 973 (1986) [$T_{1/2}$, P_β , ICC, P_γ]
- 1996SC06 E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. A 369 (1996) 527 [ω_K , K x ray ratios, Auger e-ratios, atomic data]
- 2007BU08 T.W. Burrows, Nucl. Data Sheets 108, 923 (2007) [ENSDF evaluation, mixing ratio, E_γ , J^π]
- 2008KI07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr. Nucl. Instrum. Meth. A 589 (2008) 202-229 [BrIcc]
- 2012WA38 M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer, Chin. Phys. C 36, 1603 (2012) [Q-value]

⁵¹Cr - Comments on evaluation of decay data by E. Schönfeld and R. G. Helmer

1 Decay scheme

The decay scheme is complete since there is only one excited level in ⁵¹V below the decay energy and it is populated in this decay.

The J^π and half-life of the excited level are from the 1997Zh09 evaluation.
See 1973De60 for a very complete evaluation of the nuclear and atomic data related to this decay.

2 Nuclear Data

Q value is from Audi and Wapstra (1995Au04).

The half-life data, in days, are as follows:

26.0	(10)	1940Wa02	
26.5	(10)	1940Wa02	
26		1948Ho04	
27		1948Mi12	
27.75	(30)	1952Ly17	
27.9	(2)	1956Ka33	
27.8	(1)	1956Sc87	
27.85	(2)	1957Ka65	
28.04	(16)	1957Ka65	
27.75	(30)	1957Wr37	
27.82	(20)	1963Ho17	
27.701	(6)	1964Ma56	
27.5		1965Sa09	
27.7	(2)	1967LaZZ	superseded by 1975La16
27.80	(51)	1968Bo25	
27.704	(3)	1969MeZV	superseded by 1982RuZV
27.679	(17)	1970WaAA	superseded by 1983Wa26
27.76	(15)	1972Em01	
28.1	(17)	1973ArZI	
27.721	(26)	1973LaAA	superseded by 1975La16
27.750	(9)	1973Vi13	
27.703	(8)	1974Ts01	
27.72	(3)	1975La16	
27.690	(5)	1980Ho17	
27.71	(1)	1982ChZF	
27.705	(12)	1982DeYX	superseded by 1983Wa26
27.73	(1)	1982HoZJ	superseded by 1992Un01
27.704	(3)	1982RuZV	
27.71	(3)	1983Wa26	
27.7010	(12)	1992Un01	
27.703	(3)	Adopted value	

Comments on evaluation

Three sets of half-life values were analyzed with the Limitation of Relative Statistical Weight, LRSW, method (1985ZiZY,1992Ra08) ; these sets had 21, 20, and 9 values. In all three cases the LRWS analysis increases the uncertainty of the 1992Un01 value from 0.0012 to 0.0021 in order to reduce its relative weight from 76% to 50%.

For all 21 values with uncertainties and not superseded, the LRSW weighted average is 27.7034 with an internal uncertainty of 0.0015, a reduced $-\chi^2 = 5.06$, and an external uncertainty of 0.0034. The largest contribution to this reduced- χ^2 is 2.7 from the first value from 1957Ka65. If this value is removed from the data, the remaining 20 values give an LRSW weighted average is 27.7026 with an internal uncertainty of 0.0015, a reduced- $\chi^2 = 2.49$, and an external uncertainty of 0.0024.

The third analysis was done with the nine values from the set of twenty which have uncertainties of ≤ 0.03 (namely, 1964Ma56, 1973Vi13, 1974Ts01, 1975La16, 1980Ho17, 1982ChZF, 1982RuZV, 1983Wa26, and 1992Un01). In this case the LRSW analysis gives a weighted average of 27.7025, an internal uncertainty of 0.0015, a reduced- χ^2 of 4.48, and an external uncertainty of 0.0032.

The adopted value of 27.703 (3) is consistent with all three of these results.

2.1 Electron Capture Transitions

The capture branching is determined from the $P_\gamma(320)$ value (see sec. 4.2).

The P_K etc. values from LOGFT and EC-CAPTURE codes agree quite well, namely

Level	P_K	LOGFT		EC-CAPTURE		
		P_L	P_{M+N}	P_K	P_L	P_M
0	0.892	0.0927	0.0154	0.8919 (17)	0.0934 (14)	0.0144 (6)
320	0.891	0.0935	0.0156	0.8910 (17)	0.0941 (14)	0.0145 (6)

The EC-CAPTURE values have been adopted.

2.3 Gamma Transitions

The internal-conversion coefficient of $\alpha = 0.00169$ (5) and $\alpha_K = 0.00154$ (3) are from the analysis of experimental data in 1985HaZA. These results are based on $\alpha = 0.00169$ (5) (1973Wi10) and α_K values of 0.00157 (8) (1969KaAA, as quoted in 1985HaZA), 0.00156 (8) (1970Ca17), 0.00146 (13) (1970Ri11), and 0.00153 (4) (1973Wi10). From $K/L = 11.3$ (6) and $L/M = 5.1$ (6) from 1969Dr01, one obtains $\alpha_L = 0.000136$ (8) and $\alpha_M = 0.000027$ (4). [An earlier evaluation by 1973De60 had available the latter three α_K values and deduced $\alpha_K = 0.00153$ (4) and from the above K/L and L/M ratios $\alpha = 0.00169$ (5).] Other measured values of α are 0.00162 (16) (1955Bu01), 0.0031 (2) (1955Es15), 0.0015 (2) 1956Of03 and 0.0016 (2) (1962Gu09) and those of α_K are 0.0029 (2) (1955Es15), 0.00138 (13) (1955Of01), 0.00146 (10) (1968Ri17, superseded by 1970Ri11), and 0.001527 (36) (1969WiAA, as quoted in 1985HaZA, superseded by 1973Wi10).

The mixing ratio, δ , deduced from these α_K and α_L and the conversion coefficients interpolated from the tables of 1976Ba63 is 0.40 (4). This compares reasonable well with the value of +0.465 (20) from the evaluation of 1997Zh09 which is based on the measured values of +0.43 (3) from (γ,γ') , +0.52 (7) from Coulomb excitation, and 0.49 (3) calculated from the adopted B(E2) and half-life values.

3, 3.1, and 3.2 X Radiations and Auger Electrons

Data are from 1996Sc06.

4.1 Electron Emissions

The data are from the γ -ray and atomic data in sec. 2.1, 2.2, and 3. A comparison of these intensities (in %) and those from RADLST gives :

	EMISSION	RADLST
L Auger	147.6 (10)	146.17 (16)
K Auger	66.4 (6)	66.32 (5)
K-320	0.0152 (3)	0.0166 (13)
L-320	0.00134 (8)	0.0016 (10)

The adopted values are from Emission.

4.2 Photon Emissions

The energy is from 2000He14.

The LRSW analysis of $9P_\gamma$ values gives the weighted average of 9.87% (5) with a reduced $\chi^2 = 0.96$. The input values are: 9.8 6 (1955Bu01), 9 1 (1955Co56), 9.72 15 (1953MeZZ), 10.20 63 (1965Dh01), 9.75 20 (average of 2 values of 1965Le24), 10.2 10 (1970Ri11), 9.85 9 (1980Sc07), 10.30 19 (1984Fi10), and 9.86 8 (1991Ba11). Others: \approx 2 (1940Wa02), 3 (1945Br02), 8 (1952Ly17), 21 (1952Ma49), 9.8 (1955Bi29), 7 (1955Co56), and 10.1 3 (1970ScAA, replaced by 1980Sc07). [From a set of five values, the evaluation of 1973De60 gives a result of 9.83% (14).]

The number of X rays was calculated, by the Emission program, from the γ -ray probabilities and atomic data in sec. 2.1, 2.2, and 3.

7. References

- 1940Wa02 - H. Walke, F. C. Thompson, J. Holt, Phys. Rev. **57** (1940) 171 [$T_{1/2}$]
 1945Br02 - H. Bradt, P. C. Gugelot, O. Huber, H. Medicus, P. Preiswerk, P. Scherrer, Helv. Phys. Acta **18** (1945) 252 [P_γ]
 1948Ho04 - H.H.Hopkins Jr., B.B.Cunningham, Phys.Rev. **73** (1948) 1406 [$T_{1/2}$]
 1948Mi12 - D.R.Miller, R.C.Thompson, B.B.Cunningham, Phys.Rev. **74** (1948) 347 [$T_{1/2}$]
 1952Ly17 - W.S.Lyon, Phys.Rev. **87** (1952) 1126 [$T_{1/2}$]
 1952Ma49 - D. Maeder, P. Preiswerk, A. Steinemann, Helv. Phys. Acta **25** (1952) 46 [P_γ , α_K]
 1955Bi29 - A.Bisi, E.Germagnoli, L.Zappa, Nuovo cimento **2** (1955) 1052 [P_γ]
 1955Bu01 - M. E. Bunker, J. W. Starner, Phys. Rev. **97** (1955) 1272 and **99** (1955) 1906 [P_γ , α , δ]
 1955Co56 - S. G. Cohen, S. Ofer, Phys. Rev. **100** (1955) 856 [Bremsstrahlung]
 1955Es15 - I. V. Estulin, E. M. Moiseeva, Sov. Phys. JETP **1** (1955) 463 [α , α_K]
 1955Of01 - Z. O'Friel, A. H. Huber, Phys. Rev. **99** (1955) 659, abstr. V5 [α_K]
 1956Ka33 - P.Kafalas, J.W.Irvine, Jr., Phys.Rev. **104** (1956) 703 [$T_{1/2}$]
 1956Of03 - Z. O'Friel, A. H. Huber, Phys. Rev. **101** (1956) 1076 [α]
 1956Sc87 - R. P. Schumann, M. E. Jones, A. C. Mewherter, J. Inorg. Chem. **3** (1956) 160 [$T_{1/2}$]
 1957Ka65 - G.M.Karavaev, S.A.Rusinova, Trudy Vsesoyuz.Nauch.- Issledovatel. Inst. Metrol. **30** (1957) 132 [$T_{1/2}$]
 1957Of07 - S. Ofer, R. Wiener, Phys. Rev. **107** (1957) 1639 [P_γ]
 1957Wr37 - H.W.Wright, E.I.Wyatt, S.A.Reynolds, W.S.Lyon, T.H.Handley, Nuclear Sci.and Eng**2** (1957) 427 [$T_{1/2}$]
 1962Fa02 - U.Fasoli, C.Manduchi, G.Zannoni, Nuovo cimento **23** (1962) 1126 [P_K , P_L]
 1962Gu09 - U. C. Gupta, M. G. Shanani, P. K. Srivastava, J. Sci. Industr. Res. **21B** (1962) 1 [α]
 1963Ho17 - S.Hontzeas, L.Yaffe, Can.J.Chem. **41** (1963) 2194 [$T_{1/2}$]

- 1963Kr02 - I.Y.Krause, Phys.Rev. **129** (1963) 1330 [γ mixing ratio]
 1963MeZZ - J. S. Merritt, J. G. V. Taylor, AECL-1778 (1963) 31 [P_γ]
 1964Ma56 - P. J. Marais, F. J. Haasbroek, J. H. M. Karsten. S. Afr. J. Agr. Sci. 7 (1964) 881 [$T_{1/2}$]
 1965Dh01 - K. C. Dhingra, U. C. Gupta, N. P. S. Sidhu, Current Sci. (India) 34 (1965) 504 [P_γ]
 1965Le24 - J. Legrand, report CEA-R-2813 (1965) [P_γ]
 1965Sa09 - S.R.Salisbury, R.A.Chalmers, Phys.Rev. **140** (1965) B305 [$T_{1/2}$]
 1966He07 - W.Heuer, Z.Physik **194** (1966) 224 [P_K, P_L]
 1967LaZZ - F.Lagoutine, Y.Le gallic, J.Legrand, Proc.Symp.Standardization of radionuclides, Vienna, Austria(1966), Intern.At.Energy Agency, Vienna, (1967) 603; CONF-661012 [$T_{1/2}$]
 1968Bo25 - M.Bormann, A.Behrend, I.Riehle, O.Vogel, Nucl.Phys. **A115** (1968) 309 [$T_{1/2}$]
 1968Ri17 - C. Ribordy, J. Kern, L. Schellenberg, O. Huber, Helv. Phys. Acta **41** (1968) 429 [α_K]
 1969Dr01 - O.Dragoun, C.Ribordy, O.Huber, Nucl.Phys. **A124** (1969) 337 [α]
 1969KaAA - J. W. Kane, Jr., Thesis, University of Alabama (1969) [α_K]
 1969MeZV - J. S. Merritt, J. G. V. Taylor, report AECL-3512, p.30 [$T_{1/2}$]
 1969WiAA - J. B. Willet, Thesis, Indiana University (1969) [α_K]
 1970Ca17 - H. C. Carter, J. H. Hamilton, Z. Phys. **235** (1970) 383 [α_K]
 1970Ho16 - R.N.Horoshko, D.Cline, P.M.S.Lesser, Nucl.Phys. **A149** (1970) 562 [γ mixing ratio]
 1970Ri11 - C. Ribordy, O. Huber, Helv. Phys. Acta **43** (1970) 345 [E_γ, α_K]
 1970ScAA - U. Schötzig, H. M. Weiss, K. F. Walz, Wissenschaftliche Abhandlungender Phys. -Techn. Bundesanstalt **22** (1970) 76 [P_γ]
 1970WaAA - K. F. Walz, U. Schötzig, Wissenschaftliche Abhandlungen der Phys-Techn. Bundesanstalt **22** (1970) 76 [$T_{1/2}$]
 1972Em01 - J.F.Emery, S.A.Reynolds, E.I.Wyatt, G.I.Gleason, Nucl.Sci.Eng. **48** (1972) 319 [$T_{1/2}$]
 1973ArZI - J.Araminowicz, J.Dresler, INR-1464 (1973) 14 [$T_{1/2}$]
 1973De60 - E. De Roost, F. Lagoutine, At. Energy Rev. **11** (1973) 642 [$\alpha, \delta, T_{1/2}, P_K, P_L$]
 1973LaAA - F. Lagoutine, to be published as quoted in 1973De60 [$T_{1/2}$]
 1973Vi11 - C. J. Visser, J. H. M. Karsten, F. J. Haasbroek, P. G. Marais, Agrochemophy sica **5** (1973) 15 [$T_{1/2}$]
 1973Wi10 - J. B. Willett, G. T. Emery, Ann. Phys. (New York) **78** (1973) 496 [α_K, δ]
 1974Ts01 - C. W. Tse, J. N. Mundy, W. D. McFall, Phys. Rev. C **10** (1974) 838 [$T_{1/2}$]
 1975Bo07 - G.L.Borchert, W.Scheck, K.P.Wieder, Z.Naturforsch. **30a** (1975) 274 [E_γ]
 1975La16 - F. Lagoutine, J. Legrand, C. Bac, Intern. J. Appl. Radiat. Isotop. **26** (1975) 131 [$T_{1/2}$]
 1976Ba63 - I. M. Band, M. B. Trzhaskovskaya. M. A. Listengarten, Atomic Data Nucl. Data Table**8** (1976) 433 [α_K, α_L]
 1978Kr19 - K.S.Krane, At.Data Nucl.Data Tables **22** (1978) 269 [δ]
 1978Mo22 - T.Morii, Nucl.Instrum.Methods **151** (1978) 489 [E_γ]
 1980Ho17 - H. Houtermans, O. Milosevic, F. Reichel, Intern. J. Appl. Radiat. Isotop. **31** (1980) 153 [$T_{1/2}$]
 1980RuZY - A. R. Rutledge, L. V. Smith, J. S. Merritt, report AECL-6692 (1980) [$T_{1/2}$]
 1980Sc07 - U. Schötzig, K. Debertin, K. F. Walz, Nucl. Instr. and Meth. **169** (1980) 43 [E_γ, P_γ]
 1982ChZF - P. Christmas, report NBS-SP-6262 (1982) 100 [$T_{1/2}$]
 1982DeYX - K.Debertin, U.Schotzig, K.F.Walz, NBS-SP-626 (1982) 101 [$T_{1/2}$]
 1982RuZV - A. R. Rutledge, L. V. Smith, J. S. Merritt, report NBS-SP-626 (1982) 5 [$T_{1/2}$]
 1983Wa26 - K. F. Walz, K. Debertin, H. Schrader, Intern. J. Appl. Radiat. Isot. **34** (1983) 1191 [$T_{1/2}$]
 1984Fi10 - S. A. Fisher, R. I. Hershberger, Nucl. Phys. **A423** (1984) 121 [P_γ, P_ϵ]
 1985HaZA - H. H. Hansen, European Appl. Res. Rept. Nucl. Sci. Technol. **6**, 4 (1985) 777; EUR 9478 EN [α, α_K]
 1985ZiZY - W. L. Zijp, report ECN-179, Petten (1985) [analysis methodology]
 1991Ba11 - T. Barta, L. Szücs, A. Zsinka, Appl. Radiat. Isot. **42** (1991) 490 [P_γ]
 1992Ra08 - M. U. Rajput, T. D. MacMahon, Nucl. Instr. Meth. **A 312** (1992) 289 [analysis methodology]
 1992Un01 - M. P. Unterweger, D. D. Hoppes, F. J. Schima, Nucl. Instr. and Meht. **A312** (1992) 349 [$T_{1/2}$]
 1995Au04 - G. Audi, A. H. Wapstra, Nucl. Phys. **A595** (1995) 409 [Q_ϵ]
 1995ScZY - E. Schönfeld, report PTB-6.33-95-2 (1995) [P_{ec}]
 1996Sc06 - E. Schönfeld, H. Janßen, Nucl. Instr. Meth. **A 369** (1996) 527 [ω]
 2000He14 - R..G. Helmer, C. van der Leun, Nucl. Instr. Meth. **A 450** (2000) 35 [E_γ]

**⁵²Fe – Comments on evaluation of decay data
by Aurelian Luca**

This evaluation was completed in October 2013. Includes all literature published before this date.

Decay Scheme

⁵²Fe decays 100 % by electron capture and β^+ to excited levels of ⁵²Mn. The isomer ^{52m}Mn is created within this decay chain (excitation energy of 377.7 keV, and half-life of 21.1 (2) min). The half-life of ^{52m}Mn was adopted from the evaluation of Huo *et al.* (2007), which is based on the experimental value of Juliano *et al.* (1959); another experimental half-life value is 21 (1) min, Katoh *et al.*, 1960.

Q_{EC} value for ⁵²Fe decay of 2375 (6) keV was adopted from Wang *et al.* (2012Wa38).

The spins, parities and level energies are adopted from the most recent mass-chain evaluation published for A=52 (2007Hu08). There is no information available about the spin and parity of the 1417.7-keV energy level of ⁵²Mn.

Half-life

Experimental half-lives used to calculate the recommended value are listed in the following table:

Reference	T _{1/2} (h)	uc (h)	Comments
1959Ju40	8.2	0.1	
1967Pa22	8.23	0.04	
1974Ro18	8.275	0.008	
Crit (χ^2)	4.6		
χ^2	0.88		
UWM	8.235	0.022	
LWM	8.273		
uc(WM)int. :	0.008		
uc(WM)ext. :	0.007		
Adopted:	8.273	0.008	

Although the half-life value measured by Saha and Farrer (1971Sa21) of 8.2 h was reported without any uncertainty, the evaluator has estimated an uncertainty of ± 0.2 h based on the reported counting statistics. However, because of a lack of information on the systematic uncertainty of the reported half-life, this particular measurement was not taken into account in the present evaluation. A half-life value given by Miller *et al.* (1948Mi12) of 7.8 hours was also set aside because the authors did not report the uncertainty, and an estimate could not be made from the published information.

The adopted data set is consistent and the recommended half-life is the weighted mean of the three listed values (LWM analysis, according to LWEIGHT4 computer program, version for MS Excel), with an uncertainty which is the maximum of the internal and external uncertainties.

Electron Capture and β^+ Transitions

All electron capture (EC) and β^+ energies were derived from the nuclear level energies and the Q value. Shell and sub-shells capture probabilities were calculated by means of the EC-Capture program (1998Sc28). The ground state of ⁵²Mn is only populated through the gamma-ray isomeric transition of 377.7 keV. There are two electron capture transitions feeding the excited states of 1417.7 and 546.4 keV, and only one β^+ transition with an energy of 807 (6) keV in competition with the electron capture process. This energy is in good agreement with the experimental value measured by Arbman and Svartholm (1956) of 0.804 (10) MeV. The probabilities of the two electron capture transitions (from the ground state of ⁵²Fe to the 2nd and 3rd excited states of ⁵²Mn) and the allowed β^+ transition (from the ground state of ⁵²Fe to the 2nd excited state of ⁵²Mn) were calculated from the decay scheme balance and the theoretical ratio (EC/ β^+) computed by the LOG FT program from the theoretical tables of Gove *et al.* (1971Go40). This theoretical ratio was 0.780 (21),

in agreement with the experimental value of 0.770 from 1959Ju40. The total (EC + β^+) transition probability to the excited state of 546.4 keV (⁵²Mn) is 99.9 (15) %. The LOG FT program was also used to calculate the log ft values for the EC and β^+ transitions.

The resulting 511-keV photon emission intensity is 112.2 (14) %.

Relative gamma emission probabilities, P_γ

Only one measurement of the gamma-ray energies and relative emission probabilities was found in the literature: Yaffe and Meyer (1977), 1977Ya08. Another measurement made by McIsaac and Gehrke (1972) gives a relative intensity of 3 (1) for the gamma emission with an energy of 355.6 (2) keV (intensity was 100 for the 168.77 (2) keV photons), and was not considered in the evaluation. The 377.749-keV gamma ray is the IT-decay process of Mn-52m directly to the ground state of Mn-52.

(i,f)	E_γ	$\Delta E_\gamma(\text{uc})$	Yaffe and Meyer (1977)	
			P_γ	$\Delta P_\gamma(\text{uc})$
(2,1)	168.689	0.008	1032	20
(1,0)	377.749	0.005	17.09	0.15
(3,1)	1039.928	0.019	0.99	0.04

A reference intensity of 1000 was adopted for the emission probability of the 1434.06 (1) keV gamma ray (this gamma transition follows the ⁵²Mn electron capture and β^+ transitions populating the nuclear levels of ⁵²Cr).

Internal conversion coefficients

The adopted ICC are the theoretical values calculated by the BrIcc program (2008Ki07). The gamma-ray transitions are [M1] (168.7 keV) and E4 (377.7 keV).

Normalization Factor

100 % of the transitions (β^+ , EC, γ - with the exception of the isomeric transition) in the decay of ⁵²Fe populate the first excited (isomeric) state of the ⁵²Mn daughter at 377.7 keV:

$$[P_{\gamma 168} \cdot (1 + \alpha_{T168})] + [P_{\gamma 1039} \cdot (1 + \alpha_{T1039})] = \frac{100}{N}$$

where $P_{\gamma 168}$ and $P_{\gamma 1039}$ are the relative emission probabilities of the 168.6-keV and 1039.9-keV gamma-rays, respectively, α_{T168} and α_{T1039} are the total internal conversion coefficients of the two transitions, and N is the normalization factor between the relative and absolute γ -ray probabilities.

A normalization factor (N) of 0.0961 (19) was deduced from the above. Using this factor and the relative γ probabilities adopted above, the absolute γ -ray emission probabilities were calculated.

Atomic Data

The fluorescence yield data, the relative K X-ray emission probabilities, the ratios P(KLX)/P(KLL) and P(KXY)/P(KLL) were taken from Schönfeld *et al.* (1996Sc06).

The Auger electron and X-ray absolute probabilities were calculated by the EMISSION program (2000Sc47) from the related decay data (γ emission probabilities, ICC, P_{EC} probabilities, etc.) as determined above.

Acknowledgement: the evaluator thanks Dr. Huang Xiaolong (China Nuclear Data Center, China Institute of Atomic Energy, Beijing, China) who provided several references of importance to the present evaluation.

References

- 1948Mi12** D.R. Miller, R.C. Thompson, B.B. Cunningham. Phys. Rev. 74 (1948) 347. [Half-life]
- 1956Ar33** E. Arberman, N. Svartholm. Ark. Fysik 10 (1956) 1. [Positron emission energy]
- 1959Ju40** J.O. Juliano, C.W. Kocher, T.D. Nainan, A.C.G. Mitchell. Phys. Rev. 113 (1959) 602.
[Half-life, Electron Capture/Beta plus ratio]
- 1960Ka20** T. Katoh, M. Nozawa, Y. Yoshizawa, Y. Koh. J. Phys. Soc. Jpn. 15 (1960) 2140.
[Half-life, Multipolarities]
- 1967Pa22** A. Pakkanen. Ann. Acad. Sci. Fenn. Series A, VI, 253 (1967) 25. [Half-life]
- 1971Go40** N.B. Gove, M.J. Martin. Nucl. Data Tables 10 (1971) 205. [EC/positron ratios, log ft]
- 1971Sa21** G.B. Saha, P.A. Farrer. Int. J. Appl. Radiat. Isot. 22 (1971) 495. [Half-life]
- 1972McYW** L.D. McIsaac, R.J. Gehrke. ANCR-1088 (1972) 384.
[Gamma ray energies, Gamma-ray relative emission probabilities]
- 1974Ro18** S.J. Rothman, N.L. Peterson, W.K. Chen, J.J. Hines, R. Bastar, L.C. Robinson, L.J. Nowicki,
J.B. Anderson. Phys. Rev. C9 (1974) 2272. [Half-life]
- 1977Ya08** R.P. Yaffe, R.A. Meyer. Phys. Rev. C16 (1977) 1581.
[Gamma ray energies, Gamma-ray relative emission probabilities]
- 1996Sc06** E. Schönfeld, H. Janssen. Nucl. Instrum. Methods Phys. Res. A369 (1996) 527.
[Atomic data]
- 1998Sc28** E. Schönfeld. Appl. Radiat. Isot. 49 (1998) 1353. [Fractional EC probabilities]
- 2000Sc47** E. Schönfeld, H. Janssen. Appl. Radiat. Isot. 52 (2000) 595. [P(X), P(Ae)]
- 2007Hu08** Huo Junde, Huo Su, Ma Chunhui. Nucl. Data Sheets 108 (2007) 773.
[Spin and Parity, Level energies, Half-life, Multipolarities]
- 2008Ki07** T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor Jr.
Nucl. Instrum. Methods Phys. Res. A589 (2008) 202. [Theoretical ICC]
- 2012Wa38** M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer.
Chin. Phys. C36 (2012) 1603. [Q-value]

**⁵⁴Mn - Comments on evaluation of decay data
by R. G. Helmer and E. Schönfeld**

1 Decay scheme

The decay scheme is complete since the only level in ⁵⁴Cr below the decay energy is populated in this decay. The β⁻ decay to ⁵⁴Fe is negligible.

The J^π and half-life of the excited level are from the 1993Hu04 evaluation.

2 Nuclear Data

Q value is from Audi and Wapstra 1995 (1995Au04) evaluation.

The half-life data, in days, are as follows:

291 (1)	1955Ba10	omitted from analysis
290 (6)	1956Ka33	omitted from analysis
278 (5)	1956Sc87	omitted from analysis
313.5 (7)	1961Wy01	
300	1964Be26	omitted from analysis
303 (1)	1964Ma14	omitted from analysis
311.9 (2)	1965An07	
311.9 (2)	1965An07	
312.6 (4)	1965An07	
314	1965Sa09	omitted from analysis
312 (5)	1968Ha47	
312.2 (3)	1968La10	quoted σ of 0.9 divided by 3
312.99 (5)	1968Zi01	quoted σ of 0.10 divided by 2, omitted from analysis
312.2 (9)	1969BoZX	
312.16 (11)	1973MeYE	superseded by 1982RuZV
315.40 (3)	1973Vi13	omitted from analysis
312.6 (8)	1974Cr05	
312.21 (5)	1979MeZY	superseded by 1980RuZY
312.21 (3)	1980RuZY	superseded by 1982RuZV
312.02 (4)	1982HoZJ	superseded by 1992Un01
312.21 (3)	1982RuZV	
312.19 (13)	1982RyZX	
312.15 (23)	1982RyZX	
312.028 (34)	1992Un01	
312.11 (5)	1997Ma75	
312.13 (3)	Adopted value	

The three values from before 1960 were omitted because it would have been difficult to determine the presence of impurities in the samples with the spectrometry methods available then. The two values without uncertainties were omitted. The quoted uncertainty for the value of 1968La10 was divided by 3 to convert it to a 1 σ value. The values of 1964Ma14, 1968Zi01, and 1973Vi13 were omitted since they are outliers; with the latter two both included the reduced- χ^2 is 21.7 and with only 1968Zi01 included, it is 7.4.

Adopted value of 312.13 (3) is from the Limitation of Relative Statistical Weight analysis (1985ZiZY, 1992Ra08) of the 13 remaining values. For this fit, the internal uncertainty is 0.020, the reduced- $\chi^2 = 2.06$, and the external uncertainty is 0.029. In this analysis, the three values from 1992Un01, 1982RuZV, and 1997Ma75 contribute 94% of the relative weight, and the latter two which are from the same laboratory contribute 60% of the relative weight.

2.1 and 2.2 Electron-Capture and β^+ Transitions

The unique 2nd forbidden $\epsilon + \beta^+$ transition to the ⁵⁴Cr ground state has not been observed, but an upper limit can be determined from the $\log ft$ systematics (1998Si17) as well as from searches for the positrons. From these $\log ft$ systematics, $\log f_{2ut} > 13.9$ which corresponds to $\epsilon + \beta^+$ branch of $< 0.0007\%$. The experimental limits on the β^+ intensity come from searches for the 511 keV annihilation radiation. These limits are $\leq 8 \times 10^{-5}\%$ (1968Be01), $\leq 4.4 \times 10^{-6}\%$ (1989Su08), and $\leq 5.7 \times 10^{-7}\%$ (1993Da20). From the latter value and the theoretical ϵ/β^+ ratio of 638(11), one has a capture probability of $\leq 0.0004\%$. Since this limit is lower than that from the $\log ft$ systematics, it is adopted.

The P_K etc. values for the branch to the 834keV level from the LOGFT and EGCAPTURE codes agree quite well, namely

	P_K	P_L	P_M
LOGFT	0.8895	0.0942	0.0163
EC-CAPTURE	0.8895 (17)	0.0950 (15)	0.0150 (16)

The EC-CAPTURE values have been adopted.

2.3 β^- Transitions

This unique 2nd forbidden β^- transition to the ⁵⁴Fe ground state has not been observed. A limit on its probability can be calculated from the $\log ft$ systematics (1998Si17) which give $\log f_{2ut} \geq 13.9$ and this corresponds to $I(\beta^-) \leq 0.0005\%$.

From cosmic-ray data and a model of galactic transport of cosmic rays, 1996Du15 deduce the partial half-life for β^- decay to be between 1×10^6 and 2×10^6 years, which corresponds to a β^- branch intensity between 0.00004% and 0.00009%.

2.4 Gamma Transitions and Internal-Conversion Coefficients

The α and α_K are from the analysis of the experimental data in 1985HaZA and, are based only on the data of 1966Ha07. The corresponding theoretical values interpolated from the tables of 1976Ba63 are 0.000252(8) and 0.000224(7) were α has been computed as $\alpha_K + 1.33 \times \alpha_L$.

3, 3.1 and 3.2 Atomic Data

Data are from 1996Sc06.

4.1 Electron Emissions

The data are deduced from the γ -ray probabilities and atomic data in sec. 2.1, 2.2, and 3.

A comparison of these intensities with those from the RADLIST code for this decay scheme is:

	Radlist	EMISSION
L Auger	143.3 (4)	143.0 (6)
K Auger	63.21 (12)	63.3 (5)
K-834	0.0224 (13)	0.0224 (11)
L-834	0.002199	0.00220 (13)

4.2 Photon Emissions

The energy is from the 2000He14 evaluation.

The γ -ray emission probability is computed as $I_{\epsilon}(834) / [1.0 + \alpha(834)] = 99.9997(3) / 1.000251(11) = 99.9746(11)$. The dominant component in the final uncertainty is from the uncertainty in α .

A comparison of the computed X-ray emission probabilities is:

	RADLST	EMISSION
K _{α2}	7.659 (15)	7.66 (13)
K _{α1}	15.04 (3)	15.0 (3)
K _{β}	3.056 (6)	3.05 (6)
K	25.76 (3)	25.7 (3)

And, the measured Cr K X ray emission probabilities include:

25.7 (4)	1963Ta19
24.3 (12)	1965Le21
25.14 (17)	1967Ba50
24.90 (53)	1967PeZZ
24.92 (17)	1968Ha47
24.4 (3)	1973KoAA
24.7 (9)	1973MuAA
25.93 (14)	1978Ma06
25.1 (7)	1980Co22

which are slightly lower than the calculated values, but generally are within the uncertainties.

6. References

- 1955Ba10 - E. W. Backofen, R. H. Herber, Phys. Rev. **97**(1955)743 [T_{1/2}]
- 1956Ka33 - P. Kafalas, J. W. Irvine, Jr., Phys. Rev. **104**(1956)703 [T_{1/2}]
- 1956Sc87 - R. P. Schuman, M. E. Jones, and A. C. McWherter, J. Inorg. Nucl. Chem**3**(1956)160 [T_{1/2}]
- 1961Wy01 - E. I. Wyatt, S. A. Reynolds, T.H. Handley, W. S. Lyons, H. A. Parker, Nucl. Sci. Eng. **11**(1961)74 [T_{1/2}]
- 1963Ta19 - J. G. V. Taylor, J. S. Merritt, Proc. Intern. Conf. Role of Atomic Electrons in Nuclear Transformations, Warsaw, vol. III, p. 465 (1963) [P_X]
- 1964Be26 - G. Ben-David, Nucl. Sci. Eng. **20**(1964)281 [T_{1/2}]
- 1964Ma14 - W. H. Marin, D. M. Clare, Nucl. Sci. Eng. **19**(1964)465 [T_{1/2}]
- 1965An07 - S. C. Anspach, L. M. Cavallo, S. B. Garfinkel, J. M. R. Hutchinson, C. N. Smith, report NP-15663(1965) [T_{1/2}]
- 1965Ta10 - J. G. V. Taylor, J. S. Merritt, report AECL-2501(1965)26 [T_{1/2}]
- 1965Le21 - K. F. Leistner, Atomkerenergie **10**(1965)311 [P_X]
- 1965Sa09 - S. R. Salisbury, R. A. Chalmers, Phys. Rev. **140**(1965)B305 [T_{1/2}]
- 1966Ha07 - J. H. Hamilton, S. R. Amtey, B. van Nooijen, A. V. Ramayya, J. J. Pinajian, Phys. Letters **19**(1966)682 [α]
- 1967Ba50 - W. Bambynek, Z. Phys. **206**(1967)66 [P_X]
- 1967PeZZ - M. Petel, H. Houtermans, "Standardization of Radionuclides", (IAEA, Vienna, 1967) 301 [P_X]
- 1968Be01 - D. Berenyi, D. Varga, B. Vasvari, E. Brucher, Nucl. Phys. **A106**(1968)248 [I_{β+}]
- 1968Ha47 - J. W. Hammer, Z. Phys. **216**(1968)355 [P_X]
- 1968La10 - F. Lagoutine, Y. le Gallic, J. Legrand, Intern. J. Appl. Radiat. Isot. **19**(1968)475 [T_{1/2}]
- 1968Zi01 - W. H. Zimmer, R. E. Dahl, Nucl. Sci. Eng. **32**(1968)132 [T_{1/2}]
- 1969BoZX - P. Bock, report KFK-1116 10/14/71 (1969) [T_{1/2}]
- 1973MeYE - J. S. Merritt, J. G. V. Taylor, report AECL-4657(1974)30 [T_{1/2}]
- 1973KoAA - A. A. Konstantinov, T. E. Sazonova, A. Konstantinov, Proc. Intern. Conf. Inner-Shell Ionization Phenomena and Future Applications, April 1972(1973) page 144 [P_X]
- 1973MuAA - A. Mukerji, Chin Lee, Proc. Intern. Conf. Inner -Shell Ionization Phenomena and Future Applications, April 1972 (1973) page 164 [P_X]
- 1973Vi13 - C. J. Visser, J. H. M. Karsten, F. J. Haasbroek, P. G. Marias, Agrochemop hysica **5**(1973)15 [T_{1/2}]
- 1974Cr05 - P. J. Cressy, Jr., Nucl. Sci. and Eng. **55**(1974)450 [T_{1/2}]
- 1976Ba63 - I. M. Band, M. B. Trzhaskovskaya, M. A. Listengarten, Atomic Data and Nuclear Data Tables **18**(1976)433 [α_K, α_L]
- 1978Ma06 - P. Magnier, J. Bouchard, M. Blondel, J. Legrand, J. -P. Perolat, R. Vatin, Z. Phys. **A284**(1978)383 or 389 [P_{1/2}]
- 1979MeZY - J. S. Merritt, A. R. Rutledge, L. V. Smith, F. H. Gibson, report NEANDC(CAN)-51/L (1979) 12 [T_{1/2}]
- 1980Co22 - D. D. Cohen, Nucl. Instr. Meth. **178**(1980)481 [P_X]
- 1980RuZV - A. R. Rutledge, L. V. Smith, J. S. Merritt, report AECL-6692 (1980) [T_{1/2}]
- 1982HoZJ - D. D. Hoppes, J. M. R. Hutchinson, F. J. Schima, M. P. Unterweger, report NBS-SP-626 (1982) 85 [T_{1/2}]
- 1982RuZV - A. R. Rutledge, L. V. Smith, J. S. Merritt, report NBS-SP-6262 (1982) 5 [T_{1/2}]
- 1982RyZX - A. Rytz, report NBS-SP-6262 (1982) 32 [T_{1/2}]
- 1985HaZA - H. H. Hansen, European App. Res. Reports, Nucl. Sci. and Technol. **6**, No. 4 (1985) 777; EUR 9478 EN [α]
- 1985ZiZY - W. L. Zijp, report ECN-179, Petten (1985) [analysis methodology]

Comments on evaluation

- 1989Su08 - B. Sur, K. R. Vogel, E. B. Norman, K. T. Lesko, R.-M. Larimer, E. Browne, Phys. Rev. **C39**(1989)1511 [$T_{1/2}$]
- 1992Ra08 - M. U. Rajput, T. D. Mac Mahon, Nucl. Instr. Meth. **A312**(1992)289 [analysis methodology]
- 1992Un01 - M. P. Unterweger, D. D. Hoppes, F. J. Schima, Nucl. Instr. Meth. **A312**(1992)349 [$T_{1/2}$]
- 1993Da20 - M. T. F. da Cruz, Y. Chan, A. Garcia, M. M. Hindi, G. Kenchian, R.M. Larimer, K. T. Lesko, E. B. Norman, R. G. Stokstad, F. E. Wietfeldt, I. Zlimen, PhysRev. **C48**(1993)31110 [ϵ , β^+]
- 1993Hu04 - J. Huo, H. Sun, W. Zhao, Q. Zhou, Nucl. Data Sheets **68**(1993)887 [J, $T_{1/2}$]
- 1995Au04 - G. Audi, A. H. Wapstra, Nucl. Phys. **A595**(1995)409 [Q]
- 1996Du15 - M. A. DuVernois, Phys. Rev. **C54**(1996)A2134 [$T_{1/2}$]
- 1997Ma75 - R. H. Martin, K. I. W. Burns, and J. V. G. Taylor, Nucl. Instr. Meth. **A390**(1997)267 [$T_{1/2}$]
- 1998Si17 - B. Singh, J. L. Rodriguez, S. S. M. Wong, J. K. Tuli, Nucl. Data Sheets **84**(1998)487 [$\log ft$ systematics]
- 2000He14 - R. G. Helmer, C. van der Leun, Nucl. Instr. Meth. **A450**(2000)35 [E_γ]

⁵⁵Fe - Comments on evaluation of decay data by M. M. Bé and V. Chisté

The initial evaluation was completed in April 1998. This revised evaluation was carried out in 2005, the literature available by December 2005 was taking into account.

1. Decay scheme

An Internal Bremsstrahlung electron capture spectrum was measured by **Isaac *et al.***, the intensity was found to be $3.24 (6) \times 10^{-5}$ relatively to K capture.

The J^π value and level energy are from **NDS 64,4** (1991). From other decay modes, the excited level energy has been determined to be 125.949 (10) keV.

2. Nuclear Data

- The Q value is from **Audi and Wapstra** (2003)
- The half-life values taking into account are, in days :

(1)	977.9	2.3	Lagoutine 1982 (DSA PC) ^a
(2)	1000.4	1.3	Houtermans 1980 (PC)
(3)	1009.0	1.7	Hoppes 1982 (PC, Si(Li))
(4)	996.8	6.0	Morel 1994 (Planar Ge)
(5)	995.0	3.0	Karmalitsyn 1998 (PC)
(6)	1003.5	2.1	Schötzig 2000 (Si(Li))
(7)	1005.2	1.4	Van Ammel 2006 (DSA PC)

^a (Method of measurement, PC = Proportional counter, DSA = Defined Solid Angle)

The (1) value is rejected because it is discrepant by Chauvenet's criterion.

With this value deleted, none of the other values has a relative weight greater than 50 %.

The Lweight calculation gives, for the six remaining values, a weighted mean value of 1003.4 d, with an external uncertainty of 1.7, an internal uncertainty of 0.7 and a reduced $-\chi^2$ of 5.4.

This set of value is inconsistent, the three values with lower uncertainties (2, 3 and 7) are not compatible within their uncertainty limits. No trend can be distinguished.

So, the external uncertainty has been expanded so range to include the most precise value of 1000.4 d.

The adopted value is $1003.4 (30) d$ or $2.747 (8) a$.

Other references not used in this evaluation due to their discrepancy or their great uncertainty comparing with the set of recent values above :

- 1037 (11) G.L. Brownell, C.J. Maletskos, Phys.Rev. 80 (1950) 1102
- 950 (7) R.P. Schuman et al., I.Inorg.Nuclear Chem. 3 (1956) 160
- 880 (44) J.S. Evens, R.A. Naumann, PPAD-2137-566 (1965) 10

2.1. Electron Capture transitions

- The EC transition energies are from $Q(\text{EC}) = 231.21$ (18) and from the individual level energies.
- The transition probabilities are deduced from the total gamma -ray transition probability balances at each level.
- The electron capture coefficients, for this allowed transition, were calculated by using the EC -capture program :

$$P_K = 0.8853$$
 (16) ; $P_L = 0.0983$ (13) ; $P_M = 0.0157$ (6) ; $P_N = 0.0006$ (2)

The LOGFT program gives :

$$P_K = 0.885$$
 (9) ; $P_L = 0.0974$ (10) ; $P_M = 0.0161$ (2) ; $P_{N+} = 0.00106$ (1)

Measurements were carried out by **Pengra et al.** :

$$P_K = 0.881$$
 (4) ; $P_L = 0.103$ (4) ; $P_{M+} = 0.0161$ (8)

Results from calculations and measurements are in good agreement, nevertheless the measured values are dependent on ω_K (= 0.314) and on the intensity of the $K\alpha$ X-ray (= 0.89). So, the recommended values are those of the EC-capture program.

- Several measurements or calculations were done to study the double K -shell ionization process. One can quote **Campbell et al.**; where the total probability for double vacancies in the K shell was found to be 1.3 (2) 10^{-4} , or **Kitahara et al.** where the probability for the ejection of another K electron during the K -capture decay was estimated to be 1.01 (27) 10^{-4} . As these phenomena have very small probabilities, these results are only quoted here as a matter of interest.

2.2. Gamma transitions

A weak gamma transition is deduced from the observation of a 126 keV gamma emission. The energy is derived from the level energy.

3. Atomic Data

Several data for ω_K are deduced from measurements :

- from **Smith**, $\omega_K = 0.320$ (3) ($P_K = 0.885$ (2))
- from **Konstantinov et al.**, $\omega_K = 0.312$ (3)
- from **Dobrilovic et al.**, $\omega_K = 0.322$ (5)
- from **Kuhn et al.**, $\omega_K = 0.310$ (23)
- from **Hubbell et al.**, $\omega_K = 0.321$ (7) (deduced from photoionization cross-section measurements)

A theoretical value was also calculated by **Chen** : $\omega_K = 0.323$.

These values are in good agreement (except **Konstantinov et al.** and **Khun et al.**) with the recommended value of $\omega_K = 0.321$ (5) from the semi-empirical fit of **Bambynek 1984**.

$\overline{\omega}_L$ and η_{KL} are from **Schönfeld et al.**

3.1.1. X Radiations

- The X-ray energies were obtained by conversion of the wavelength values from **Bearden** into energies with $1 \text{ \AA} = 1.000\,014\,81$ (92) 10^{-10} m.
- The emission intensities are calculated by the EMISSION program from PTB with ω_K , $\overline{\omega}_L$ and η_{KL} quoted above and, $K\beta/K\alpha = 0.1359$ (14), $K\alpha_2/K\alpha_1 = 0.5099$ (25) (**Schönfeld et al.**).
- With $P_K = 0.8853$ (16) for this allowed transition, and $\omega_K = 0.321$ (5) the total K X-ray emission intensity is then $P_K \times \omega_K = 0.284$ (5) which can be compared with the experimental values of 0.279 (8) (**Schötzig**) and of 0.283 (2) (**Smith**).

The value given by **Smith** was obtained in an international activity measurement exercise where six laboratories reported results for $P_K \times \omega_K$. The deduced weighted mean is in good agreement with the calculated value and has a better uncertainty. However, as pointed out by **Smith**, this uncertainty is probably underestimated. So, the value of $I_K = P_K \times \omega_K \times 100 = 28.4 (5) \%$ is adopted.

3.1.2. Auger Electrons

Complete measurements of the K Auger spectrum of manganese was performed by **Kovalik et al.**, they found for the relative intensities of the K Auger groups :

$$\text{KLM/KLL} = 0.26 (2)$$

$$\text{KMM/KLL} = 0.018 (2)$$

These values are in good agreement with the recommended values calculated with the EMISSION program:

$$\text{KLM/KLL} = 0.272 (3)$$

$$\text{KMM/KLL} = 0.0185 (4)$$

The energies were taken from **Larkins** or, for the missing lines, calculated from the electron binding energies. **Kovalik et al.** also measured the energies and found a good agreement for the KLM spectrum but observed discrepancies for the KLL and KMM groups.

4.2. Gamma emissions

A weak gamma emission superimposed on the intense inner -bremsstrahlung was observed by **Zlimen et al.** and interpreted as the deexcitation of the first excited state of Mn -55. The γ -ray energy is given as 126.0 (1) keV and the γ -ray intensity as $1.3 (1) \times 10^{-7} \%$.

From the level energy 125.949 (10) keV and with a recoil energy of 0.2 eV, the retained γ -ray energy is 125.949 (10) keV.

References

- J. G. **Pengra**, H. Genz, J. P. Renier, R. W. Fink. Phys. Rev. C5,6 (1972) 2007. PL/PK, PM/PL
- L. **Dobrilovic**, D. Bek-Uzarov, M. Simovic, K. Buraei, A. Milojevic. Proc. of the International Conference on Inner-shell Ionization Phenomena CONF-720404 (1973) 128. K fluorescence yield
- Tetsuo **Kitahara**, Sakae Shimizu. Phys. Rev. C11,3 (1975) 920. P(ionisation)
- F. P. **Larkins**. At. Data Nucl. Data Tables 20,4 (1977) 338. Auger Electrons
- M. H. **Chen**. Phys. Rev. A21-2 (1980) 436. K fluorescence yield
- H. **Houtermans**, O. Milosevic, F. Reichel. Int. J. Appl. Radiat. Isotop. 31 (1980) 153. Half-life
- U. **Kuhn**, H. Genz, W. Löw, A. Richter, H. W. Müller. Z. Phys. A - Atoms and Nuclei 300 (1981) 103. K fluorescence yield
- D. D. **Hoppes**, J. M. R. Hutchinson, F. J. Schima, M. P. Unterweger. NBS-Special publication 626 (1982) 85. Half-life
- F. **Lagoutine**, J. Legrand, C. Bac. Int. J. Appl. Radiat. Isotop. 33 (1982) 711. Half-life
- D. **Smith**. Nucl. Instrum. Methods 200 (1982) 383. PkWk
- W. **Bambynek**. A. Meisel Ed. Leipzig Aug. 20-23 (1984). K fluorescence yield
- A. A. **Konstantinov**, T. E. Sazonova, S. V. Sepman, E. A. Frolov. Metrologia 26 (1989) 205. K fluorescence yield
- M. C. P. **Isaac**, V. R. Vanin, O. A. M. Helene. Z. Phys. A. 335 (1990) 243. Beta emission energies
- A. **Kovalik**, V. Brabec, J. Novak, O. Dragoun, V. M. Gorozhankin, A. F. Novgorodov, Ts. Vylov. J. Elec. Spectro. Rel. Phenomena 50 (1990) 89. Auger electrons
- J. L. **Campbell**, J. A. Maxwell, W. J. Teesdale. Phys. Rev. C. 43,4 (1991) 1656. Double K capture probability

- I. **Zlimen**, E. Browne, Y. Chan, M. T. F. da Cruz, A. Garcia, R. -M. Larimer, K. T. Lesko, E. B. Norman, R. G. Stokstad, F. E. Wietfeldt. Phys. Rev. C. 46,3 (1992) 1136. Gamma Emission
- J. H. **Hubbell**, P. N. Trehan, Nirmal Singh, B. Chand, D. Mehta, M. L. Garg, R. R. Garg, Surinder Singh, S. **Puri**. J. Phys. Chem. Ref. Data 23-2 (1994) 339. K fluorescence yield
- J. **Morel**, M. Etcheverry, M. Vallée. Nucl. Instrum. Methods A339 (1994) 232. Half-life
- E. **Schönfeld**, H. Janssen. Report PTB Ra-37 (1995). L fluorescence yield, Kb/Ka
- N. I. **Karmalitsyn**, T. E. Sazonova, A. V. Zanevsky, S. V. Sepman. Int. J. Appl. Radiat. Isotop. 49,9 -11 (1998) 1363. Half-life
- U. **Schötzig**. Appl. Rad. Isotopes 53 (2000) 469. Half-life, X-ray emission intensities
- G. **Audi**, A. H. Wapstra. Nucl. Phys. A729, 1 (2003) 337 Q
- R. **Van Ammel**, S. Pommé, G. Sibbens. Appl. Rad. Isotopes 64 (2006) 1412. Half-life

Mn-56 – Comments on evaluation of decay data
by A. L. Nichols

Evaluated: November 1999
Re-evaluated: January 2004

Evaluation Procedures

Limitation of Relative Statistical Weight Method (LWM) was applied to average numbers throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

A reasonably simple and consistent decay scheme has been constructed from the gamma-ray measurements of 1967Au01, 1968Sh07, 1973Ar15, 1974Ti01, 1974Ho25 and 2004MiXX. Ten distinct gamma-ray emissions were identified with ⁵⁶Mn decay in these studies. An additional gamma ray at 3119.3 keV was identified by 1968Sh01, but this emission has been discarded due to a lack of evidence from the other studies.

Nuclear Data

The gamma-ray emissions of ⁵⁶Mn are reasonably well defined, and this radionuclide has suitable decay characteristics for use as a calibrant over the gamma-ray energy range 840 to 2550 keV.

Half-life

Half-life adopted from the evaluation of Woods for the IAEA -CRP: Update of X- and Gamma-ray Decay Data Standards for Detector Calibration. The measurements of 1968Sh07, 1971GoYM, 1972Em01, 1973La12, 1980RuZY, 1992An13 and 1994Ya02 were considered.

Reference	Half-life (days)
1968Sh07	0.10771(4)
1971GoYM	0.10742(33)
1972Em01	0.10779(25)
1973La12	0.107438(8)
1980RuZY	0.107350(33)
1992An13	0.107454(4) [§]
1994Ya02	0.1040(20) [*]
Evaluated value	0.107449(18)

[§] Uncertainty increased to ± 0.000008 to ensure weighting factor not greater than 0.50.

^{*} Method development study: removed from data set due to uncharacteristically large uncertainty.

Woods evaluation for IAEA-CRP (2004WoZZ): recommended half-life of 0.107449(19) days or 2.57878 (46) h (using above data set, but also excluding 1994Ya02 data), adopted for this evaluation.

Gamma Rays

Energies

A number of well-defined gamma-ray energies were adopted from the recommended standards of 2000He14. All other gamma-ray energies were calculated from the structural details of the proposed decay scheme and the nuclear level energies of 1999Hu04 (as derived from the energy measurements of 1973Ar15, 1974Ho25 and 1974Ti01). An additional gamma ray with an energy of 3119.3(5) keV was only detected by 1968Sh01, and has been discarded due to a lack of evidence in all of the other studies.

Emission Probabilities

Weighted mean relative emission probabilities were determined for all of the gamma rays assigned to the decay scheme, using the relevant data from the measurements of 1967Au01, 1968Sh07, 1973Ar15, 1974Ho25, 1974Ti01 and 2004MiXX. All gamma-ray emissions were expressed relative to the 846.7638 keV transition, which was arbitrarily assigned an uncertainty of 3% (100(3)%).

Gamma-ray Emission Probabilities: Relative to $P_g(846.7638 \text{ keV})$ of 100%

$E_g(\text{keV})$	P_g^{rel}						Recommended Values*
	1967Au01	1968Sh07	1973Ar15	1974Ho25	1974Ti01	2004MiXX	
846.7638(19) [†]	100(3)	100(3)	100(3)	100(3)	100(3)	100.000(103)	100(3)
1037.8333(24) [†]	-	-	0.06(1)	0.03(1)	0.040(5)	-	0.040(4) [§]
1238.2736(22) [†]	-	-	0.14(3)	0.13(1)	0.10(1)	0.097(2)	0.098(2) [§]
1810.726(4) [†]	30(3)	29.4(16)	28.6(15)	26.9(13)	27.5(8)	26.610(72)	27.2(4)
2113.092(6) [†]	17.4(17)	16.0(9)	16.0(8)	14.3(7)	14.5(4)	13.956(53)	14.4(3) [§]
2523.06(5) [‡]	1.10(15)	1.6(5)	1.14(5)	1.01(5)	1.00(3)	1.025(9)	1.03(2)
2598.438(4) [†]	-	-	0.026(5)	0.02(1)	0.019(2)	-	0.020(2)
2657.56(1) [‡]	0.60(10)	0.66(6)	0.71(4)	0.66(7)	0.66(2)	0.648(8)	0.652(7) [§]
2959.92(1) [‡]	0.31(6)	0.26(3)	0.30(2)	0.32(3)	0.31(1)	0.314(6)	0.311(5) [§]
3119.3(5) [#]	-	0.08(4)	-	-	-	-	-
3369.84(4) [‡]	0.22(5)	0.20(4)	0.15(2)	0.16(2)	0.17(1)	-	0.17(1)

[†] Energy adopted from 2000He14.

[‡] Energy calculated from the nuclear level energies specified by 1999Hu04.

[#] Energy from 1968Sh07, but transition not included in proposed decay scheme.

* Weighted mean values adopted using LWEIGHT, unless stated.

[§] Recommended values adopted from a combination of the normalised residuals and Rajeval methods (see 2004MaYY).

The normalisation factor for the gamma-ray emission probabilities was calculated from the proposed decay scheme via two routes:

(a) beta population of all ⁵⁶Fe nuclear levels derived from gamma -ray depopulation/population and summed, assuming β decay to ⁵⁶Fe ground state is zero (spin and parity considerations ($3^+ \rightarrow 0^+$)).

$$\begin{aligned} \text{for all nuclear levels populated by } \beta \text{ decay } \Sigma P_{\beta i} &= (101.163 \pm 1.479) \times NF = 100 \\ NF &= 0.989 (15) \end{aligned}$$

(b) population of ⁵⁶Fe ground state by gamma transitions, assuming β decay to ⁵⁶Fe ground state is zero.

$$\begin{aligned} \Sigma P_{\gamma i} (1 + \alpha_i) NF &= [P_{\gamma}(3369.84 \text{ keV}) + P_{\gamma}(2959.92 \text{ keV}) + P_{\gamma}(2657.62 \text{ keV}) + \\ P_{\gamma}(846.7638 \text{ keV}) (1 + \alpha_i)] \times NF &= 100 \\ 101.163(23) \times NF &= 100 \\ NF &= 0.9885(3) \end{aligned}$$

Hence, a normalisation factor of 0.9885(3) was adopted on the basis of the more accurate determination.

Multipolarities and Internal Conversion Coefficients

The nuclear level scheme specified by 1999Hu04 has been used to define the multipolarities of the gamma transitions on the basis of known spins and parities. Studies of the internal conversion coefficients of the some of these gamma transitions support the proposed transition types: (97%M1 + 3%E2) for the 1810.726 keV gamma rays (taken from 1989Co01); (99.96%M1 + 0.04%E2) and 100%E2 for the 1037.8333 and 1238.2736 keV gamma rays, respectively (taken from 1974Ho25).

Multipolarity Assignments

Reference	E _g (keV)	Multipolarity
1974Ho25	1037.83	99.96%M1 + 0.04%E2
	1238.27	E2
	1810.726(4)	96.5%M1 + 3.5%E2
	2113.092(6)	93.4%M1 + 6.6%E2
	2523.06(5)	94.1%M1 + 5.9%E2
	2598.438(4)	93.4%M1 + 6.6%E2
1989Co01	1810.726(4)	97%M1 + 3%E2
	2113.092(6)	96%M1 + 4%E2

Beta-particle Emissions

Energies

All beta-particle energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies of 1999Hu04 and the Q-value were used to determine the energies and uncertainties of the beta-particle transitions to the various levels.

Emission Probabilities

The beta-particle emission probabilities were calculated from the recommended gamma -ray emission probabilities and the theoretical internal conversion coefficients of 1976Ba63 (latter estimated by interpolation of the data). Log ft systematics can be applied to the beta -particle transition to the ground state of ⁵⁶Fe ($\Delta J=3, \Delta \pi = \text{no}$), with a lower limit for log ft of 13.9 (1998Si17), to give a beta-particle emission probability of < 0.0005 (set to zero).

Beta-particle Emission Probabilities

E _b (keV)	P _b
	Recommended Values*
250.2(3)	0.00020(2)
325.7(3)	0.0120(3)
572.6(3)	0.00040(4)
735.6(3)	0.145(3)
1037.9(3)	0.275(4)
1610.4(3)	0.00057(6)
2848.7(3)	0.566(7)

* Recommended emission probabilities derived from evaluated gamma-ray emission probabilities and theoretical internal conversion coefficients.

Atomic Data

The x-ray data have been calculated using the evaluated gamma-ray data, and the atomic data from 1996Sc06, 1998ScZM and 1999ScZX.

7 References

- 1967Au01 - R. L. Auble, W. C. McHarris, W. H. Kelly, The Decay Schemes of ⁵⁶Co and ⁵⁶Mn and their use as Calibration Standards, Nucl. Phys., A91(1967)225-237. [P_γ]
- 1968Sh07 - A. H. Sher, B. D. Pate, The Decay of ⁵⁶Co and ⁵⁶Mn, Nucl. Phys., A112(1968)85-96. [Half-life, P_γ]
- 1971GoYM - I. W. Goodier, M. J. Woods, A. Williams, Measurements of Nuclear Decay Schemes, Proc. Int. Conf. Chemical Nucl. Data, Canterbury, Editor: M. L. Hurrell (1971)175. [Half-life]
- 1972Em01 - J. F. Emery, S. A. Reynolds, E. I. Wyatt and G. I. Gleason, Half -Lives of Radionuclides – IV, Nucl. Sci. Eng., 48(1972)319-323. [Half-life]
- 1973Ar15 - G. Ardisson and C. Marsol, Niveaux de ⁵⁶Fe Peuplés dans la Désintégration de ⁵⁶Mn, Nucl. Phys., A212(1973)424-428. [P_γ]
- 1973La12 - F. Lagoutine and J. Legrand, Use of ⁵⁶Mn to Check Measuring Equipment, Nucl. Instrum. Meth., 112(1973)323. [Half-life]
- 1974Ho25 - S. Hofmann, Kernspektroskopische Untersuchungen zu den Termschemata der Kerne ⁵⁶Fe und ⁵⁶Co, Z. Phys., 270(1974)133-147. [P_γ]
- 1974Ti01 - K. G. Tirsell, L. G. Multhauf and S. Raman, Decays of ⁵⁸Mn, ⁵⁷Mn and ⁵⁶Mn, Phys. Rev., 10C(1974)785-794. [P_γ]
- 1976Ba63 - I. M. Band, M. B. Trzhaskovskaya and M. A. Listengarten, Internal Conversion Coefficients for Atomic Numbers Z = 30, At. Data Nucl. Data Tables, 18(1976)433. [ICC]
- 1980RuZY - A. R. Rutledge, L. V. Smith and J. S. Merritt, Decay Data for Radionuclides Used for the Calibration of X- and Gamma-Ray Spectrometers, AECL-6692(1980). [Half-life]
- 1989Co01 - S. P. Collins, S. A. Eid, S. A. Hamada, W. D. Hamilton and F. Hoyler, A Search for Mixed - symmetry States in the Mass A ~ 50 Region, J. Phys., G: Nucl. Part. Phys., 15(1989)321 -332. [Multipolarity]
- 1992An13 - M. S. Antony, D. Oster and A. Hachem, Precise Determination of Half -lives of ⁵⁶Mn, ¹⁹³Os and ^{197,199}Pt, J. Radioanal. Nucl. Chem. Letts., 164(1992)303-308. [Half-life]
- 1994Ya02 - T. Yassine and I. Othman, A Simple Method for the Rapid Assessment of the Half Lives of Short-Lived Radionuclides, Appl. Radiat. Isot., 45(1994)271-273. [Half-life]
- 1995Au04 - G. Audi and A. H. Wapstra, The 1995 Update to the Atomic Mass Evaluation, Nucl. Phys., A595(1995)409. [Q value]
- 1996Sc06 - E. Schönfeld and H. Janßen, Evaluation of Atomic Shell Data, Nucl. Instrum. Meth. Phys. Res., A369(1996)527-533. [X_K, X_L, Auger electrons]

- 1998ScZM - E. Schönfeld and G. Rodloff, Tables of the Energies of K-Auger Electrons for Elements with Atomic Numbers in the Range from $Z = 11$ to $Z = 100$, PTB Report PTB -6.11-98-1, October 1998. [Auger electrons]
- 1998Si17 - B. Singh, J. L. Rodriguez, S. S. M. Wong and J. K. Tuli, Nucl. Data Sheets, 84(1998)487. [log f_i]
- 1999Hu04 - Huo Junde, Nuclear Data Sheets for $A = 56$, Nucl. Data Sheets, 86(1999)315. [Nuclear structure, Energies]
- 1999ScZX - E. Schönfeld and G. Rodloff, Energies and Relative Emission Probabilities of K X-rays for Elements with Atomic Numbers in the Range from $Z = 5$ to $Z = 100$, PTB Report PTB -6.11-1999-1, February 1999. [X_K]
- 2000He14 - R. G. Helmer and C. van der Leun, Recommended Standards for γ -ray Energy Calibration (1999), Nucl. Instrum. Meth. Phys. Res., A450(2000)35-70. [E_γ]
- 2004MaYY - T. D. MacMahon, A. Pearce and P. Harris, Convergence of Techniques for the Evaluation of Discrepant Data, Appl. Radiat. Isot., 60(2004)275-281. [Statistical analyses]
- 2004MiXX - H. Miyahara, Y. Ogata, K. Fujiki, K. Katoh and N. Marnada, Highly -precise Measurements of the Relative Gamma-ray Intensities for ⁵⁶Mn and ⁷²Ga, Appl. Radiat. Isot., 60(2004)295-299. [P_γ]
- 2004WoZZ - M. J. Woods, Half -life Evaluations for IAEA -CRP on “Update of X-ray and Gamma -ray Decay Data Standards for Detector Calibration and Other Applications” (2004). [Half-life evaluation]

⁵⁶Co - Comments on evaluation of decay data by C.M. Baglin and T. D. MacMahon

This current evaluation was carried out in 2004. The literature available by September 2004 was included.

Evaluation Procedures

The *Limitation of Relative Statistical Weight* (LWM) [1985ZiZY] method, used almost exclusively for averaging numbers throughout this evaluation, provided a uniform approach for the analysis of discrepant data. In the few instances when an alternative technique was used, this fact has been noted. The uncertainty assigned in this evaluation to the recommended value is always greater than or equal to the smallest uncertainty in any of the experimental values used in the calculation.

1 Decay Scheme

⁵⁶Co decays 19.58 (11) % by positron (β^+) emission and 80.42 (11) % by electron capture (ϵ) to ⁵⁶Fe ($Q(\epsilon) = 4566.0$ (20) keV (2003Au03)). Altogether, 46 γ rays de-exciting 15 nuclear levels in ⁵⁶Fe have been reported. Except for the strong 847-keV transition, emission of conversion electrons is very low and negligible compared to that of γ rays (photons) because of the low atomic number ($Z=26$) of the daughter nucleus (⁵⁶Fe) and the high energy (> 700 keV) of the most intense γ -ray transitions. Consequently, neither conversion coefficients (most of them $< 2 \times 10^{-4}$) nor conversion electron energies and intensities have been tabulated in this evaluation. Pair production is also possible for transitions with $E_\gamma \geq 1022$ keV, but the internal-pair-formation coefficients (based on 1979Sc31) do not exceed 10^{-3} and are tabulated only for those transitions for which the coefficients exceed 4×10^{-4} or for which their omission would affect the deduced branching.

The evaluator has normalized the decay scheme assuming zero $\epsilon + \beta^+$ feeding from the 4^+ ⁵⁶Co parent to the 0^+ ⁵⁶Fe ground state. Then $\Sigma(I(\gamma + ce) \text{ to ground state}) = 100\%$. Based on the decay scheme, only the 847 γ , 2657 γ and 3370 γ feed the ground state. The 847 keV transition conversion coefficient is taken as $3.03(9) \times 10^{-4}$ (from Band *et al.*, 1976Ba63, assuming $\alpha = \alpha_K + 1.33 \alpha_L$ and a 3% uncertainty). The normalization factor N is then given by:

$$N = 100 / [I(847\gamma) (1 + \alpha(847\gamma)) + I(2657\gamma) + I(3370\gamma)]$$

Where: $I(847\gamma)$, $I(2657\gamma)$, $I(3370\gamma)$ are the relative values given in Table 2

$$= 100 / [100.0303 (9) + 0.0195 (20) + 0.0103 (8)]$$

$$= 100 / [100.0601 (23)]$$

$$= 0.999399 (23)$$

With this normalization, the probability of the 847 keV transition is : $P(847)(\gamma + ce) = 99.9702(23)\%$.

Electron-capture and β^+ transition probabilities to excited states in ⁵⁶Fe were determined from γ -ray transition intensity balance at each level and theoretical ϵ/β^+ ratios. It should be noted that the 2nd-forbidden transitions to the 2690 and 3370 levels, though weak, are probably overestimated since $\log ft$ values for these branches are significantly lower than expected from $\log ft$ systematics.

The evaluator has included level half-life data from the evaluation by Huo (1999Hu04) in the decay scheme drawing given here. The level energies shown in the drawing result from a least-squares adjustment of the γ -ray energies recommended in this evaluation.

2 Nuclear Data

The recommended value for the half-life of ⁵⁶Co is 77.236 (26) days, taken from the evaluation by Woods *et al.* (2004WoAA). This supersedes an earlier evaluation by two of these authors (2004Wo02) in which 77.20 (8) days ($\chi^2/\nu = 0.9$) was recommended. Measured values and their respective sources are:

Half-life (days)	Reference	Comments
77.2 (8)	1954Bu58	
77.3 (3)	1957Wr37	
78.76 (12)	1972Em01	statistical outlier
78.4 (5)	1974Cr05	statistical outlier
77.12 (10)	1977An13	
77.12 (7)	1978La21	
77.30 (9)	1989Al24	
77.08 (8)	1989Le17	
77.28 (4)	1989Sc17	
77.29 (3)	1990Al29	
77.210 (28)	1992Fu02	
77.29 (4)	1992Fu02	

The weighted average of all data published from 1977 onwards is 77.245 (23) days ($\chi^2/\nu = 2.2$), where the uncertainty shown is the external uncertainty (the internal uncertainty is 0.015 days).

$Q(\epsilon) = 4566.0$ (20) keV is adopted from 2003Au03.

2.1 b+ Transitions

The positron end-point energies, calculated from $E_{\beta^+} = Q(\epsilon) - E(\text{lev}) - 1022$, are the evaluator's values deduced using $Q(\epsilon) = 4566.0$ (20) keV (2003Au03) and level energies ($E(\text{lev})$) from the decay scheme. Absolute β^+ emission probabilities are from γ -ray intensity balance at each nuclear level and theoretical I_{β^+}/ϵ_i ratios. Note that the latter may not be reliable for the 2nd-forbidden branches.

2.2 Electron Capture Transitions

ϵ -transition energies, calculated from $E(\epsilon) = Q(\epsilon) - E(\text{lev})$, are evaluator's values deduced using $Q(\epsilon) = 4566.0$ (20) keV (2003Au03) and level energies ($E(\text{lev})$) from the decay scheme. Absolute ϵ transition probabilities are from γ -ray intensity balance at each nuclear level and theoretical I_{β^+}/ϵ_i ratios. These sum to 80.42(11)%, implying $I(\beta^+) = 19.58(11)\%$. Fractional atomic shell electron-capture probabilities (P_K, P_L, P_M) are evaluator's values calculated using the EC-CAPTURE computer program [2] for the relevant nuclear level energies.

3 Atomic Data

Emission probabilities are evaluator's values calculated using the EMISSION program (Version 3.04) [3], atomic data from 1996Sc06, and the γ -ray emission probabilities recommended here. The K X-ray and K-Auger electron energies are taken from Schönfeld and Rodloff [5] and [4], respectively; L X-ray and L-Auger electron energies are from Larkins [6].

4 Photon Emissions

4.1 Energies

γ -ray energies shown in boldface in Table 1 are from 2000He14. These values are based on a revised energy scale that uses the new adjusted fundamental constants and wave lengths deduced from an updated value of the lattice spacing of Si crystals [Cohen and Taylor [1]]. Helmer *et al.* (2000He14) fitted the adjusted γ -ray energy measurements for ⁵⁶Co to a level scheme, and deduced recommended γ -ray energy values from level-energy differences. Less precise energies are from 1990Me15, 1989Al25 (one transition only) and 1980St20. The latter authors adopted energies from the literature for the strongest transitions (shown in square brackets in Table 1) and made the general statement that the uncertainties in the other transition energies range from 0.05 keV to 0.8 keV; the evaluator has, therefore, assigned uncertainties of 0.8 keV to the four energies adopted from this study. The uncertainties in the γ -ray energies given in this evaluation are statistical only, as reported by authors. See Table 1.

4.2 Emission Probabilities

a. Relative intensities

Relative emission probability measurements are given in Table 2, panels a); panels b) show the results for several different analyses of those data along with the intensities recommended in the present evaluation. In cases where the authors indicated an uncertainty in the relative intensity of the 847keV reference line, that uncertainty was combined in quadrature with the statistical uncertainty for each of the other transitions prior to all analyses of the data.

The analysis of these data is complicated on account of two factors:

- (i) Discrepant data sets. Of the approximately 770 data points, successive runs of the program LWEIGHT identify a total of 87 statistical outliers based on the Chauvenet criterion; this seems an unusually large fraction. Most outliers, though by no means all, arise from the earlier measurements.
- (ii) The use by some authors of Ge detector efficiency calibration curves which are inadequate at the highest energies. This problem was first identified by McCallum and Coote (1975Mc07) and is discussed further by Baglin *et al.* in 2002Ba38.

One prescription for dealing with discrepant data is the limitation of relative statistical weight method proposed by Zijp (1985ZiZY) and incorporated in the program LWEIGHT. The program identifies a set of data as 'discrepant' whenever its reduced χ^2 -squared value exceeds the critical reduced χ^2 -squared value for the relevant number of data points. For those cases, it then increases the uncertainty for any datum whose statistical weight exceeds 50% until it no longer does so, then recalculates the weighted mean. If the weighted mean overlaps the unweighted mean, the weighted mean will be adopted. The uncertainty used is usually the internal uncertainty; however, the uncertainty will be expanded to include the most precise datum, if necessary, and the external uncertainty will be used if the internal uncertainty is less than the uncertainty in the most precise datum. Otherwise, the unweighted mean will be adopted; this does not seem to be a particularly useful number since it could so easily be skewed by the least reliable data.

Two additional techniques that might reasonably be applied to the analysis of these data are the Normalised Residuals (1991JaXX) and the Rajeval (1992Ra08) techniques. Both are iterative techniques which increase the uncertainties of any deviant data, but they use different prescriptions for identifying and adjusting the deviant data. The results of these analyses are also shown in Table 2.

Another logical approach would be to use the results from LWEIGHT after all statistical outliers have been eliminated from the dataset. Table 2 also gives the results from this analysis.

The second problem could be approached by considering data from only the eight experiments (2002MoZP, 2000Ra36, 1990Me15, 1980St20, 1978Ha53, 1977Ge12, 1974BoXX and 1971Si29) in which the detector efficiency has been *measured* (not extrapolated) up to at least the highest ⁵⁶Co transition energy (3611 keV). (Details of the efficiency calibrations for many measurements are sketchy at best, and some rely partially or totally on Monte Carlo calculations.) However, this approach greatly decreases the number of data points, so one should resort to this measure only at energies where significant problems are anticipated. The high precision

data from 1971Ca14, based on a linear extrapolation to high energy of a log(efficiency) versus log(energy) plot, have received considerable scrutiny in the literature, and 2002Ba38 deduced a multiplicative correction factor ($F = 1.116 - 0.155 E_\gamma + 0.0397 E_\gamma^2$, where E_γ is in MeV) to correct ⁶⁶Ga intensity data in 1971Ca14; this formula implies intensity correction factors of 0.98, 1.01 and 1.06, respectively, at $E_\gamma = 2.5, 3.0$ and 3.5 MeV. These factors apply equally to the ⁵⁶Co data from 1971Ca14 and to those from 1970Ph01 and 1974Ho25, all tied to the intensity scale in 1971Ca14. This situation suggests that data from only the eight selected references should be considered for $E_\gamma > 3000$ keV. However, although used only for $E_\gamma > 3000$ keV, the analysis of data from the selected references is shown in Table 2 for transitions of all energies, for the sake of completeness.

b. Absolute Intensities

Absolute emission probabilities are based on experimental results and decay β -scheme normalization arguments as follows:

- $I_{ce}(847\gamma, E2)/I_\gamma(847\gamma) = 3.03 (9) \times 10^{-4}$ (Theory (Band *et al.*, 1976Ba63), assuming $\alpha = \alpha_K + 1.33 \alpha_L$ and 3% uncertainty).
- No $\epsilon + \beta^+$ branch to ground state, so $\Sigma(I(\gamma + ce) \text{ to ground state}) = 100\%$.

The recommended absolute γ -ray emission probabilities are the relative values recommended in Table 2 multiplied by 0.999399 (23).

c. Annihilation radiation intensity

The 511-keV γ -ray intensity has not been experimentally determined but may be estimated from:

$$\begin{aligned} I(\gamma^+) &= 2 \times [100 - I(\epsilon) + I(\text{pair production})] \\ &= 2 \times [19.58 (11) + 0.024] \\ &= 39.21(22) \% \end{aligned}$$

4.3 Transition Multipolarities and Mixing Ratios

The transition multipolarities and mixing ratios have been taken directly from the evaluation by Huo (1999Hu04). Several additional transition multipolarities, deduced from the decay scheme, are shown enclosed by square brackets.

5 References

1. *The 1986 Adjustment of the Fundamental Physical Constants*, E.R. Cohen and B.N. Taylor, Rev. Mod. Phys. **59**, 1121 (1987).
2. *The Program EC-CAPTURE*, E. Schönfeld, F. Chu, and E. Browne. An interactive computer program for calculating electron capture probabilities P_K , P_L , P_M , and P_N to the K, L, M, and N atomic shells, respectively (1997).
3. *The Program EMISSION* (version 3.04 (2002)), E. Schönfeld and H. Janssen. A computer program for calculating emission probabilities of X-rays and Auger electrons emitted in nuclear disintegration processes.
4. *Tables of the energies of K-Auger electrons for elements with atomic numbers in the range from Z=11 to Z=100*, E. Schönfeld, G. Rodloff, Report PTB-6.11-98-1, October 1998.
5. *Energies and relative emission probabilities of K X-rays for elements with atomic numbers in the range from Z=5 to Z=100*, E. Schönfeld, G. Rodloff, Report PTB-6.11-1999-1, February 1999.
6. F.P. Larkins, Atomic Data and Nuclear Data Tables **20**, 313 (1977).

1954Bu58 - W.H. Burgus, G.A. Cowan, J.W. Hadley, W. Hess, T. Shull, M.L. Stevenson, H.F. York, Phys. Rev. **95**, 750 (1954) [Half-life].

1957Wr37 - H.W. Wright, E.I. Wyatt, S.A. Reynolds, W.S. Lyon, T.H. Handley, Nucl. Sci. Eng **2**, 427 (1957) [Half-life].

1965Pe18 - H. Pettersson, O. Bergman, C. Bergman, Ark. Fiz. **29**, 423 (1965) [Relative γ -ray emission

- probabilities].
- 1966Do07 - K.W. Dolan, D.K. McDaniels, D.O. Wells, Phys. Rev. **148**, 1151 (1966) [Relative γ -ray emission probabilities].
- 1966Hu17 - M.Huguet, H. Forest, C. Ythier, C. R. Acad. ScParis **263B**, 1342 (1966) [Relative γ -ray emission probabilities].
- 1966Sc01 - R. Schöneberg, M. Schumacher, A. Flammersfeld, Z. Physik **192**, 305 (1966) [Relative γ -ray emission probabilities].
- 1967Au01 - R.L. Auble, W.C. McHarris, W.H. Kelly, Nucl. Phys. **A91**, 225 (1967) [Relative γ -ray emission probabilities].
- 1967Ba75 - P.H. Barker, R.D. Connor, Nucl. Instrum. Methods **57**, 147 (1967) [Relative γ -ray emission probabilities].
- 1967Ch20 - C.Chasman, R.A. Ristinen, Phys. Rev. **159**, 915 (1967) [Relative γ -ray emission probabilities].
- 1968Sh07 - A.H. Sher, B.D. Pate, Nucl. Phys. **A112**, 85 (1968) [Relative γ -ray emission probabilities].
- 1969Ar04 - B.H. Armitage, A.T.G. Ferguson, G.C. Neilson, W.D.N. Pritchard, Nucl. Phys. **A133**, 241 (1969) [Relative γ -ray emission probabilities].
- 1969Au09 - G. Aubin, J. Barrette, M.Barrette, S. Monaro, Nucl. Instrum. Methods**76**, 93 (1969) [Relative γ -ray emission probabilities].
- 1969Sc09 - H.L. Scott, D.M. van Patter, Phys. Rev. **184**, 1111 (1969) [Relative γ -ray emission probabilities].
- 1970Ph01 - M.E. Phelps, D.G. Sarantites, W.G. Winn, Nucl. Phys. **A149**, 647 (1970) [Relative γ -ray emission probabilities].
- 1971Ca14 - D.C. Camp, G.L. Meredith, Nucl. Phys. **A166**, 349 (1971) [Relative γ -ray emission probabilities].
- 1971Ge07 - R.J. Gehrke, J.E. Cline, R.L. Heath, Nucl. Instrum. Methods **91**, 349 (1971) [Relative γ -ray emission probabilities].
- 1971Ge08 - A.-M. Genest, C. R. Acad. Sci. Paris **272**, 863 (1971) [Relative γ -ray emission probabilities].
- 1971Si29 - B.P. Singh, H.C. Evans, Nucl. Instrum. Methods **97**, 475 (1971) [Relative γ -ray emission probabilities].
- 1972Em01 - J.F. Emery, S.A. Reynolds, E.I. Wyatt, G.I. Gleason, Nucl. Sci. Eng. **48**, 319 (1972) [Half-life].
- 1972Pe20 - B.F. Peterman, S. Hontzeas, R.G. Rystephanick, Nucl. Instrum. Methods**104**, 461 (1972) [Relative γ -ray emission probabilities].
- 1974BoXX - S.G. Boydell, Doctoral Thesis, University of Melbourne, 1974 [Relative γ -ray emission probabilities].
- 1974Cr05 - P.J. Cressy, Nucl. Sci. Eng. **55**, 450 (1974) [Half-life].
- 1974Ho25 - S. Hofmann, Z. Physik **270**, 133 (1974) [Relative γ -ray emission probabilities].
- 1975Ka06 - T. Katou, Nucl. Instrum. Methods **124**, 257 (1975) [Relative γ -ray emission probabilities].
- 1975Mc07 - G.J. MacCallum, G.E. Coote, Nucl. Instrum. Methods **124**, 309 (1975) [Relative γ -ray emission probabilities].
- 1976Ba63 - I.M. Band, M.B. Trzhaskovskaya, M.A. Listengarten, At. Data Nucl. Data Tables**18**, 433 (1976) [Theoretical conversion coefficients].
- 1977An13 - M.E. Anderson, Nucl. Sci. Eng. **62**, 511 (1977) [Half-life].
- 1977Ge12 - R.J. Gehrke, R.G. Helmer, R.C. Greenwood, Nucl. Instrum. Methods**147**, 405 (1977) [Relative γ -ray emission probabilities].
- 1978Ha53 - M. Hautala, A.A. Anttila, J. Keinonen, Nucl. Instrum. Methods **150**, 599 (1978) [Relative γ -ray emission probabilities].
- 1978La21 - F. Lagoutine, J. Legrand, C. Bac, Intl. J. Appl. Radiat. Isotop. **29**, 269 (1978) [Half-life].
- 1979Sc31 - P. Schlütter and G. Soff, At. Data Nucl. Dat. Tables **24**, 509 (1979). [Internal pair conversion coefficients].
- 1980Sh28 - A.K. Sharma, R. Kaur, H.R. Verma, K.K. Suri, P.N. Trehan, Proc. Indian Natl. Sci. Acad. **46 A**, 181 (1980) [Relative γ -ray emission probabilities].
- 1980St20 - N.M. Stewart, A.M. Shaban, Z. Physik **A 296**, 165 (1980). Supersedes A.M. Shaban, N.M. Stewart, T.D MacMahon, Nucl. Instrum. Methods **165**, 109 (1979) [Relative γ -ray emission probabilities].
- 1980Yo05 - Y. Yoshizawa, Y. Iwata, T. Kaku, T. Katoh, J.-Z. Ruan, T. Kojima, Y. Kawada, Nucl. Instrum. Methods **174**, 109 (1980) [Precise relative γ -ray emission probabilities].
- 1982Gr10 - A. Grütter, Intl. J. Appl. Radiat. Isotop. **33**, 533 (1982) [Relative γ -ray emission probabilities].
- 1985ZiZY - W.L. Zijp, Report ECN FYS/RASA-**85/19** (1985) [Discrepant Data. Limited Relative Statistical

Weight Method].

1988Wa26 - G. Wang, E.K. Warburton, D.E. Alburger, Nucl. Instrum. Methods **A272**, 791 (1988) [Precise transition energies used in evaluation in reference 2000He14].

1989Al24 - D.E. Alburger, E.K. Warburton, Z. Tao, Phys. Rev. **C 40**, 2789 (1989) [Half-life].

1989Al25 - D.E. Alburger, E.K. Warburton, Z. Tao, Phys. Rev. **C 40**, 2891 (1989) [Relative γ -ray emission probability].

1989Le17 - K.T. Lesko, E.B. Norman, B. Sur, R.-M. Larimer, Phys. Rev. **C 40**, 445 (1989) [Half-life].

1989Sc17 - H. Schrader, Appl. Radiat. Isotop. **40**, 381 (1989) [Half-life].

1990Al29 - D.E. Alburger, C. Wesselborg, Phys. Rev. **C 42**, 2728 (1990) [Half-life].

1990Me15 - R.A. Meyer, Fizika **22**, 153 (1990). E_γ data presumed to supersede those in R.A. Meyer, T.N. Massey, Int. J. Appl. Radiat. Isotop. **34**, 1073 (1983) [Relative γ -ray emission probabilities and energies].

1991JaXX - M.F. James, R.W. Mills, D.R. Weaver, UKAEA Report, Winfrith Technology Centre, AE-RS-1082 (1991) [‘Normalised residuals’ technique for statistical analysis of data].

1992Ra08 - M.U. Rajput, T.D. MacMahon, Nucl. Instrum. Methods in Phys. Research **A312**, 289 (1992) [‘Rajeval’ technique for statistical analysis of data].

1992Fu02 - E. Funck, U. Schötzig, M.J. Woods, J.P. Sephton, A.S. Munster, J.C.J. Dean, P. Blanchis, B. Chauvenet, Nucl. Instrum. Methods Phys. Res. **A312**, 334 (1992) [Half-life].

1992ScZZ - U. Schötzig, H. Schrader, K. Debertin, Proc. Int. Conf. Nuclear Data for Science and Technology, Jülich, Germany (1992), p. 562 [Relative γ -ray emission probabilities].

1996Sc06 - E. Schönfeld, H. Janssen, Nucl. Instrum. Methods. Phys. Res. **A369**, 527 (1996) [Atomic data, X-rays, Auger electrons].

1999Hu04 - Junde Huo, Nuclear Data Sheets **86**, 315 (1999) [⁵⁶Co decay scheme].

2000He14 - R.G. Helmer, C. van der Leun, Nucl. Instrum. Methods. Phys. Res. **A450**, 35 (2000) [Precise evaluated γ -ray energies].

2000Ra36 - S. Raman, C. Yonezawa, H. Matsue, H. Iimura, N. Shinohara, Nucl. Instrum. Methods. Phys. Res. **A454**, 389 (2000) [Precise relative γ -ray emission probabilities].

2002Ba38 - C.M. Baglin, E. Browne, E.B. Norman, G.L. Molnár, T. Belgya, Zs. Révay, F. Szelecsenyi, Nucl. Instrum. Methods Phys. Res. **A481**, 365 (2002) [Intensity correction factor for 1971Ca14 data].

2002MoZP - G. Molnár, Zs. Révay, T. Belgya, **INDC(NDS)-437**, p. 23 (Appendix 4) (2002) [Precise relative γ -ray emission probabilities]

2003Au03 - G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. **A729**, 337 (2003) [Q values].

2004Wo02 - M.J. Woods, S.M. Collins, Applied Radiat. Isotop. **60**, 257 (2004) [Half-life evaluation].

2004WoAA - M.J. Woods, S.M. Collins, S.A. Woods, NPL Report CAIR 8 (January 2004) [Half -life evaluation].

Table 1. ⁵⁶Co Gamma-Ray Energies

2000He14	1990Me15	1989AI25	1980St20	Adopted
E _g (keV)	E _g (keV)	E _g (keV)	E _g (keV) ^a	E _g (keV)
	263.41 (10)		263.34	263.41 (10)
	411.38 (8)		410.94	411.38 (8)
	486.54 (11)		485.2	486.54 (11)
			655.0 (8) ^a	655.0 (8)
			674.7 (8) ^a	674.7 (8)
733.5085 (23)	733.72 (15)		733.6	733.5085 (23)
787.7391 (23)	787.88 (7)		787.77	787.7391 (23)
846.7638 (19)	846.772 (8)		[846.764]	846.7638 (19)
		852.78 (5)		852.78 (5)
896.503 (7)	896.56 (20)		896.55	896.503 (7)
977.363 (4)	977.485 (60)		977.39	977.363 (4)
996.939 (5)	997.33 (16)		996.48	996.939 (5)
1037.8333 (24)	1037.840 (6)		[1037.844]	1037.8333 (24)
	1089.03 (24)		1089.31	1089.03 (24)
1140.356 (7)	1140.28 (10)		1140.52	1140.356 (7)
1159.933 (8)	1160.08 (16)		1160.0	1159.933 (8)
1175.0878 (22)	1175.102 (6)		[1175.099]	1175.0878 (22)
	1198.78 (20)		1198.77	1198.78 (20)
1238.2736(22)	1238.282 (7)		[1238.287]	1238.2736(22)
	1272.2 (6)		1272.20	1272.2 (6)
1335.380 (29)	1335.56 (8)		1335.56	1335.380 (29)
1360.196 (4)	1360.215 (12)		[1360.206]	1360.196 (4)
	1442.75 (8)		1442.65	1442.75 (8)
	1462.34 (12)		1462.28	1462.34 (12)
1640.450 (5)	1640.54 (13)		1640.38	1640.450 (5)
1771.327 (3)	1771.351 (16)		[1771.350]	1771.327 (3)
1810.726 (4)	1810.714 (35)		[1810.722]	1810.726 (4)
1963.703 (11)	1963.99 (6)		[1963.714]	1963.703 (11)
2015.176 (5)	2015.181 (16)		[2015.179]	2015.176 (5)
2034.752 (5)	2034.755 (15)		[2034.159]	2034.752 (5)
2113.092 (6)	2113.185 (115)		[2113.107]	2113.092 (6)
2212.898 (3)	2212.96 (15)		[2212.921]	2212.898 (3)
	2276.36 (16)		2276.09	2276.36 (16)
	2373.7 (4)		2373.71	2373.7 (4)
	2523.86 (20)		2523.0	2523.0 (8) ^b
2598.438 (4)	2598.458 (13)		[2598.460]	2598.438 (4)
			2657.4 (8) ^a	2657.4 (8)
3009.559 (4)	3009.591 (22)		[3009.596]	3009.559 (4)
3201.930 (11)	3201.962 (16)		[3201.954]	3201.930 (11)
3253.402 (5)	3253.416 (15)		[3253.417]	3253.402 (5)
3272.978 (6)	3272.990 (15)		[3272.998]	3272.978 (6)
	3369.69 (30)		3369.97	3369.69 (30)
3451.119 (4)	3451.152 (17)		[3451.154]	3451.119 (4)
	3547.93 (6)		3548.27	3547.93 (6)
	3600.49 (40)		3600.85	3600.7 (4)
			3611.8 (8) ^a	3611.8 (8)

^a Authors took energies for the strongest lines from the literature (shown in square brackets) and stated that uncertainties varied from 0.05 to 0.8 keV for the others. The evaluator has conservatively assigned 0.8 keV to those lines whose energies are adopted in the present evaluation from this reference.

^b The datum from 1980St20 is adopted in preference to the more precise datum from 1990Me15 because the latter value fits its level-scheme placement poorly and is almost 1 keV higher than the γ -ray energy of 2522.88 (6) adopted in an evaluation (1999Hu04) which included information from sources other than ⁵⁶Co ϵ decay.

Table 2: ^{56}Co Relative Gamma-Ray Emission Probabilities[@], a) Experimental Data

Ref./Eg	263.4g	411.4g	486.5g	655.0g	674.7g	733.5g	787.7g	846.8g	852.8g	896.5g
65Pe18							1.04* (21)	100		
66Do07								100		
66Hu17								100		
66Sc01								100		
67Au01						0.10* (5)	0.4 (2)	100		
67Ba75								100		
68Sh07						0.13 (6)	0.2 (1)	100		
67Ch20							0.36 (5) (8)	100 (15) (0)		
69Ar04								100		
69Au09								100		
69Sc09							0.37 (4)	100		0.14* (4)
70Ph01	0.03 (1)		0.066 (6)			0.21 (4)	0.31 (6)	100		0.06 (1)
71Ca14	0.021 (4)	0.025 (5)	0.041 (7)			0.193 (3)	0.308 (8)	100		0.071 (4)
71Ge07								100		
71Ge08	0.05* (1)	0.024 (7)	0.050 (12)		0.03 (1)	0.18 (3)	0.28 (4)	100	0.04 (1)	0.08 (2)
71Si29 ^S							0.21 (6)	100		
72Pe20 ^d								100.0 (60) (0) 100.0 (56) (0) 100.0 (57) (0)		
74BoXX ^S								100		
74Ho25	0.020 (6)	0.025 (9)	0.07 (2)		0.03 (1)	0.165 (8)	0.29 (3)	100		0.062 (6)
75Ka06						0.219 (7)	0.311 (12)	100		0.089 (11)
77Ge12 ^S								100 (1) (0)		
78Ha53 ^S						0.143 (13)	0.34 (3)	100		0.077 (10)
80St20 ^S	0.022 (4)	0.031* (4)	0.069 (7)	0.038 (8)	0.038 (7)	0.195 (14)	0.320 (7)	100		0.063 (6)
80Sh28	0.031 ^c (9)	0.026 (8)	0.065 (11)		0.045 (20)	0.166 (12)	0.28 (1)	100		0.089 (13)
80Yo05			0.061 (10) (10)			0.193 (12) (12)	0.305 (13) (13)	100.0 (3) (0)		0.095 (18) (18)
82Gr10								100		
89Al25								100	0.050 (3)	
90Me15 ^S	0.022 (4)	0.025 (5)	0.055 (5)			0.20 (1)	0.31 (1)	100		0.070 (5)
92ScZZ						0.190 (7) (7)	0.315 (10) (10)	100.00 (26) (0)		0.086 (20) (20)
00Ra36 ^S								100		
02MoZP ^S								100.0 (2) (0)		

Table 2: ⁵⁶Co Relative Gamma-Ray Emission Probabilities[@]
b) Analysis

Eg	263.4g	411.4g	486.5g	655.0g	674.7g	733.5g	787.7g	846.8g	852.8g	896.5g
All Data										
# data points, N	7	6	8	1	4	13	17	33	2	12
$\chi^2/(N-1)$	1.5	0.31	1.7	N/A	0.31	4.2 ^b	2.0 ^b	N/A	0.92	1.4
I _γ : UWM	0.028 (4)	0.0260 (10)	0.060 (4)	-	0.036 (4)	0.176 (9)	0.350 (45)	100	0.045 (5)	0.082 (6)
I _γ : WM	0.0234 (20)	0.0269 (23)	0.0583 (27)	-	0.035 (5)	0.1909 (22)	0.309 (3)	100	0.049 (3)	0.0704 (22)
I _γ : LWM	= WM	= WM	= WM	-	= WM	0.176 (17) ^x	0.309 (11) ^x	100	= WM	= WM
I _γ : Norm Res	0.0234 (20)	0.0269 (23)	0.0583 (27)	-	0.035 (5)	0.1905 (37)	0.310 (4)	100	0.049 (3)	0.0704 (22)
I _γ : Rajeval	0.0227 (20)	0.0269 (23)	0.0602 (29)	-	0.035 (5)	0.1914 (24)	0.311 (4)	100		0.0704 (22)
Statistical Outliers Excluded			N/A	N/A	N/A				N/A	
# data points, N	6	5	-	-	-	12	16	33	-	11
$\chi^2/(N-1)$	0.36	0.01	-	-	-	4.3 ^b	1.4	N/A	-	1.3
UWM	0.0243 (20)	0.0250 (3)	-	-	-	0.182 (8)	0.307 (13)	100	-	0.077 (4)
WM	0.0223 (21)	0.0250 (28)	-	-	-	0.1911 (22)	0.309 (3)	100	-	0.0701 (22)
LWM	= WM	= WM	-	-	-	0.1909 (48) ^e	= WM	100	-	= WM
Selected Data										
# data points, N	2	2	2	1	2	3	4		0	3
$\chi^2/(N-1)$	0	0.88	2.7	N/A	0.43	6.6 ^b	1.5	N/A	N/A	0.83
I _γ : UWM	0.022 (0)	0.028 (3)	0.062 (7)	-	0.034 (4)	0.179 (18)	0.295 (29)	100	-	0.070 (4)
I _γ : WM	0.022 (3)	0.029 (3)	0.060 (4)	-	0.035 (6)	0.183 (7)	0.317 (6)	100	-	0.068 (4)
I _γ : LWM	= WM	= WM	= WM	-	= WM	0.183 (18) ^e	= WM	100	-	= WM
I _γ : Norm Res										
I _γ : Rajeval										
Adopted I_g	0.0234 (20)	0.0269 (23)	0.058 (3)	0.038 ^a (8)	0.035 (5)	0.191 (4)	0.310 (4)	100	0.049 (3)	0.0704 (22)
Source	All; WM	All; WM	All; WM	1980St220	All; WM	All; NR	All; NR	N/A	All; WM	All; WM

Table 2: ^{56}Co Relative Gamma-Ray Emission Probabilities (continued)[@], a) Experimental Data

Ref./Eg	977.4g	996.9g	1037.8g	1089.0g	1140.4g	1159.9g	1175.1g	1198.8g	1238.3g	1272.2g
65Pe18	1.73* (35)		14.1 (15)				2.1 (6)		66.8 (40)	
66Do07			12.4 (5)						71.2 (26)	
66Hu17			14.5 (15)				2.8* (5)		70.5 (70)	
66Sc01			14.0 (20)				1.4* (2)		66.3 (60)	
67Au01	1.36 (36)		12.8 (9)				2.4 (2)		69.5 (35)	
67Ba75	1.62* (10)		13.7 (8)				2.03* (14)		72.1 (50)	
68Sh07	1.01* (30)		12.1* (8)				2.2 (1)		70.2 (25)	
67Ch20	1.50* (23) (32)		14.0 (21) (30)		0.170 (26) (36)		1.60* (24) (34)		64 (10) (14)	
69Ar04	1.1 (1)		9.6* [†] (6)				1.9* (2)		69.6 (35)	
69Au09			13.08 (35)				1.73* (13)		68.3 (14)	
69Sc09					0.17 (3)					
70Ph01	1.35 (5)		14.0 (7)		0.24* (4)	0.11 (2)	2.25 (5)		68.5 (12)	
71Ca14	1.448 (14)	0.112 (6)	14.24 (14)	0.048 (9)	0.142 (9)	0.100 (9)	2.300 (25)	0.050 (7)	67.64 (68)	0.019 (1)
71Ge07			12.9 (5)				2.26 (23)		67.8 (15)	
71Ge08	1.42 (14)	0.13 (3)	14.4 (9)	0.04 (1)	0.16 (3)	0.11 (2)	2.29 (22)	0.06 (2)	69.6 (35)	0.024 (7)
71Si29 ^S	1.21* (6)		12.44 (31)				2.11 (5)			
72Pe20 ^d			13.45 (190) (206) 13.03 (172) (187) 12.72 (153) (169)				1.99* (27) (30) 2.18 (34) (36) 1.93* (25) (27)		70.9 (77) (88) 68.2 (72) (81) 66.9 (75) (84)	
74BoXX ^S			13.7 (6)				2.3 (1)		66.2 (10)	
74Ho25	1.37 (4)	0.17 (5)		0.07 (2)	0.13 (2)	0.078 (7)	2.25 (11)	0.028 (9)		0.022 (3)
75Ka06	1.386 (15)		13.922 (116)		0.107 (3)	0.095 (6)	2.180 (24)		66.37 (74)	
77Ge12 ^S	1.426 (15) (21)		14.04 (14) (20)				2.28 (2) (3)		66.4 (7) (10)	
78Ha53 ^S	1.38 (4)	0.170 (14)	13.5 (2)	0.06 (2)	0.117 (13)	0.08 (1)	2.11 (10)	0.044 (8)	65.1 (4)	0.035* (4)
80St20 ^S	1.41 (2)	0.092 (14)	14.11 (19)	0.050 (7)	0.125 (6)	0.074 (8)	2.30 (32)	0.04 (1)	68.47 (87)	0.038* (6)
80Sh28	1.38 (3)	0.11 (1)	14.06 (28)	0.075 (9)	0.11 (1)	0.079 (9)	2.22 (5)	0.035 (10)	67.59 (131)	0.022 (8)
80Yo05	1.435 (16) (16)	0.129 (14) (14)	14.16 (5) (7)	0.05 (3) (3)	0.131 (21) (21)	0.095 (14) (14)	2.241 (12) (14)	0.051 (9) (9)	66.06 (21) (29)	0.025 (8) (8)
82Gr10			13.85 (35)						65.8 (16)	
89Al25										
90Me15 ^S	1.440 (15)	0.112 (6)	14.0 (1)	0.05 (1)	0.15 (1)	0.10 (1)	2.28 (2)	0.05 (1)	67.6 (4)	0.020 (2)
92ScZZ	1.450 (15) (15)		14.18 (13) (13)		0.137 (5) (5)		2.289 (21) (21)		66.96 (60) (60)	0.024 (10) (10)
00Ra36 ^S			14.11 (22)				2.25 (4)		66.6 (10)	
02MoZP ^S	1.424 (6) (7)		14.07 (4) (5)				2.252 (9) (10)		66.20 (11) (17)	

Table 2: ⁵⁶Co Relative Gamma-Ray Emission Probabilities (continued)[@]
b) Analysis

Eg	977.4g	996.9g	1037.8g	1089.0g	1140.4g	1159.9g	1175.1g	1198.8g	1238.3g	1272.2g
All Data:										
# data points, N	20	8	30	8	13	10	29	8	29	9
$\chi^2/(N-1)$	2.7 ^b	3.0 ^b	4.5 ^b	1.3	5.3 ^b	1.6	2.8 ^b	0.92	1.8 ^b	3.1 ^b
I _γ ; UWM	1.39 (3)	0.128 (10)	13.51 (18)	0.055 (4)	0.145 (10)	0.092 (4)	2.15 (5)	0.045 (4)	67.84 (36)	0.0254 (22)
I _γ ; WM	1.423 (4)	0.116 (3)	14.018 (31)	0.054 (4)	0.1204 (21)	0.088 (3)	2.249 (6)	0.044 (3)	66.42 (12)	0.0206 (8)
I _γ ; LWM	1.423 (7) ^e	0.116 (6) ^e	13.51 (56) ^x	= WM	0.145 (38) ^x	= WM	2.15 (10) ^x	= WM	67.8 (16) ^x	0.025 (6) ^x
I _γ ; Norm Res	1.423 (7)	0.114 (4)	14.04 (5)	0.054 (4)	0.131 (4)	0.088 (3)	2.250 (9)	0.044 (3)	66.45 (16)	0.0205 (9)
I _γ ; Rajeval	1.425 (5)	0.113 (4)	14.055 (31)	0.051 (4)	0.133 (3)	0.088 (3)	2.254 (6)	0.044 (3)	66.44 (12)	0.0199 (8)
Statistical Outliers Excluded:		N/A		N/A		N/A		N/A	N/A	
# data points, N	14	-	28	-	11	-	21	-	-	7
$\chi^2/(N-1)$	1.7	-	2.6 ^b	-	4.0 ^b	-	1.6	-	-	0.36
UWM	1.406 (9)	-	13.70 (11)	-	0.137 (7)	-	2.240 (16)	-	-	0.0223 (8)
WM	1.424 (4)	-	14.03 (3)	-	0.1164 (23)	-	2.252 (6)	-	-	0.0196 (8)
LWM	= WM	-	13.70 (37) ^x	-	0.137 (30) ^x	-	= WM	-	-	= WM
Selected Data:										
# data points, N	6	3	8	3	3	3	8	3	7	3
$\chi^2/(N-1)$	3.1 ^b	9.1 ^b	4.9 ^b	0.12	2.8	2.1	1.9	0.25	4.4 ^b	8.5 ^b
I _γ ; UWM	1.382 (35)	0.125 (23)	13.75 (20)	0.053 (3)	0.131 (10)	0.085 (8)	2.24 (3)	0.045 (3)	66.65 (41)	0.031 (6)
I _γ ; WM	1.422 (6)	0.117 (5)	14.01 ((4)	0.0508 (55)	0.130 (5)	0.083 (5)	2.254 (8)	0.045 (5)	66.31 (14)	0.0242 (17)
I _γ ; LWM	1.422 (12) ^e	0.122 (21) ^e	13.98 (11) ^e	= WM	= WM	= WM	= WM	= WM	66.36 (36) ^e	0.028 (8) ^x
I _γ ; Norm Res										
I _γ ; Rajeval										
Adopted I_g	1.423 (7)	0.116 (6)	14.04 (5)	0.054 (4)	0.132 (4)	0.088 (3)	2.250 (9)	0.044 (3)	66.45 (16)	0.0202 (8)
Source	All; LWM	All; LWM	All; NR	All; WM	All; NR-Raj	All; WM	All; NR	All; WM	All; NR	All; NR-Raj

Table 2: ^{56}Co Relative Gamma-Ray Emission Probabilities (continued)[@], Experimental Data

Ref./Eg	1335.4g	1360.2g	1442.8g	1462.3g	1640.5g	1771.3g	1810.7g	1963.7g	2015.2g	2034.8g
65Pe18		4.0 (8)				16.2 (14)		0.75 (27)	4.1* (12)	9.2* (17)
66Do07		3.8 (3)				15.6 (13)			3.8* (7)	7.8 (10)
66Hu17		4.5 (7)				12.5* (13)	0.70* (14)	0.80 (15)	3.7* (6)	8.3 (15)
66Sc01		3.8 (4)				13.5* (14)		1.10* (15)	3.5* (4)	6.5* (8)
67Au01		4.5 (3)				16.1 (8)	0.4* (2)	0.59 (9)	2.7 (2)	7.4 (6)
67Ba75		4.8* (3)				16.9 (10)	1.3* (6)	1.1* (2)	2.93 (30)	7.37 (50)
68Sh07		4.2 (4)				16.7 (10)	0.5* (3)	0.63 (20)	2.9 (4)	7.7 (5)
67Ch20		4.0 (6) (8)				14.0* (21) (30)		0.68 (10) (14)	2.6 (4) (6)	6.6* (10) (14)
69Ar04		4.6 (3)				16.2 (10)		0.9* (2)	3.9* (3)	8.2 (5)
69Au09		4.15 (12)				14.95 (40)			2.78 (14)	7.56 (21)
69Sc09	0.12 (2)		0.23* (3)	0.12* (3)			0.65 (6)	0.63 (5)		
70Ph01	0.15* (2)	4.37 (13)	0.20 (2)	0.08 (2)	0.05 (2)	16.0 (5)	0.62 (6)	0.74 (3)	3.13 (10)	8.1 (2)
71Ca14	0.123 (3)	4.340 (45)	0.200 (8)	0.077 (1)	0.065 (9)	15.78 (16)	0.641 (8)	0.721 (15)	3.095 (31)	7.95 (8)
71Ge07		4.16 (21)				16.5 (8)			2.99 (20)	8.2 (6)
71Ge08	0.11* (2)	3.96 (40)	0.14* (2)	<0.02	0.07 (1)	14.9 (9)	0.55* (6)	0.67 (7)	2.83 (30)	7.7 (6)
71Si29 ^S		4.42 (8)					0.47* (6)	0.58 (5)	2.60 (12)	7.0* (3)
72Pe20 ^d		4.08 (51) (57) 4.4 (6) (6) 5.30* (78) (84)				15.36 (174) (197) 15.98 (180) (201) 14.55 (166) (186)			2.88 (42) (45) 2.28* (27) (30) 2.59 (45) (47)	6.25* (88) (96) 6.8* (8) (9) 6.85* (80) (89)
74BoXX ^S		4.4 (1)				15.9 (3)			3.1 (1)	7.8 (1)
74Ho25	0.120 (12)	4.35 (12)	0.177 (9)	0.065 (12)	0.063 (6)		0.63 (3)	0.71 (3)		
75Ka06	0.120 (3)	4.189 (52)	0.172 (4)	0.078 (3)	0.062 (3)	15.369 (241)	0.665 (23)	0.667 (21)	3.025 (72)	7.694 (146)
77Ge12 ^S		4.24 (4) (6)				15.65 (16) (22)	0.650 (7) (10)	0.724 (8) (11)	3.09 (5) (6)	7.95 (12) (14)
78Ha53 ^S	0.12 (2)	4.24 (15)	0.195 (10)		0.05 (1)	15.26 (15)	0.59* (3)	0.70 (2)	2.97 (3)	7.64 (6)
80St20 ^S	0.128 (6)	4.32 (6)	0.173 (7)	0.091 (13)	0.062 (7)	15.5 (4)	0.629 (13)	0.719 (15)	3.182 (66)	8.14 (17)
80Sh28	0.124 (10)	4.29 (8)	0.182 (11)	0.086 (3)	0.055 (9)	15.61 (30)	0.62 (2)	0.71 (2)	2.95 (6)	7.74 (2)
80Yo05	0.130 (6) (6)	4.265 (17) (21)	0.172 (7) (7)	0.084 (6) (6)	0.070 (11) (11)	15.49 (5) (7)	0.657 (23) (23)	0.707 (11) (11)	3.026 (14) (17)	7.766 (28) (36)
82Gr10		4.27 (15)				15.11 (38)			2.97 (11)	7.60 (19)
89Al25										
90Me15 ^S	0.125 (5)	4.33 (4)	0.20 (1)	0.077 (5)	0.06 (1)	15.70 (15)	0.64 (1)	0.720 (15)	3.08 (3)	7.89 (7)
92ScZZ	0.118 (6) (6)	4.29 (4) (4)	0.185 (7) (7)	0.065 (8) (8)	0.072 (12) (12)	15.48 (14) (15)	0.638 (8) (8)	0.724 (10) (10)	3.04 (5)(5)	7.90 (13) (13)
00Ra36 ^S		4.23 (7)				15.42 (25)			3.03 (5)	7.835 (120)
02MoZP ^S		4.22 (15) (15)				15.24 (8) (9)	0.641 (5) (5)	0.698 (3) (3)	2.976 (14) (15)	7.69 (3) (3)

Table 2: ^{56}Co Relative Gamma-Ray Emission Probabilities (continued)[@]
a) Analysis

E _g	1335.4g	1360.2g	1442.8g	1462.3g	1640.5g	1771.3g	1810.7g	1963.7g	2015.2g	2034.8g
All Data										
# data points, N	12	31	12	10	11	29	19	23	30	30
$\chi^2/(N-1)$	0.57	0.90	2.5 ^b	1.8	0.44	1.3	1.2	1.9 ^b	2.5 ^b	1.7
I _γ ; UWM	0.124 (3)	4.29 (5)	0.186 (6)	0.082 (5)	0.0617 (23)	15.43 (17)	0.64 (4)	0.738 (27)	3.06 (7)	7.64 (11)
I _γ ; WM	0.1229 (16)	4.283 (13)	0.1797 (23)	0.0779 (9)	0.0621 (21)	15.46 (4)	0.639 (3)	0.7030 (25)	3.015 (9)	7.746 (13)
I _γ ; LWM	= WM	= WM	0.180 (8) ^x	= WM	= WM	= WM	= WM	0.7060 (42) ^e	3.015 (39) ^x	=WM
I _γ ; Norm Res	0.1229 (16)	4.283 (13)	0.1797 (36)	0.0779 (9)	0.0621 (21)	15.46 (4)	0.639 (3)	0.7038 (37)	3.019 (14)	7.746 (18)
I _γ ; Rajeval	0.1229 (16)	4.283 (13)	0.1792 (25)	0.0774 (9)	0.0621 (21)	15.49 (4)	0.640 (3)	0.7094 (37)	3.025 (10)	7.744 (14)
Statistical Outliers Excluded					N/A					
# data points, N	10	29	10	9	-	26	12	20	24	23
$\chi^2/(N-1)$	0.45	0.80	2.3	1.7	-	1.2	0.43	1.6	2.3 ^b	1.6
UWM	0.1228 (12)	4.24 (4)	0.186 (4)	0.078 (3)	-	15.67 (11)	0.640 (4)	0.694 (12)	2.94 (4)	7.82 (5)
WM	0.1228 (16)	4.282 (13)	0.1799 (23)	0.0779 (9)	-	15.47 (4)	0.641 (3)	0.7028 (25)	3.014 (9)	7.748 (14)
LWM	= WM	= WM	= WM	= WM	-	= WM	= WM	= WM	2.94 (4) ^x	= WM
Selected Data										
# data points, N	3	8	3	2	3	7	6	6	8	8
$\chi^2/(N-1)$	0.12	0.89	3.1	1.0	0.50	2.0	115 ^b	2.9	4.7 ^b	3.5 ^b
I _γ ; UWM	0.1243 (23)	4.300 (28)	0.189 (8)	0.084 (7)	0.057 (4)	15.52 (9)	0.60 (3)	0.690 (22)	3.00 (6)	7.74 (12)
I _γ ; WM	0.126 (4)	4.309 (24)	0.185 (5)	0.079 (5)	0.059 (5)	15.40 (6)	0.590 (3)	0.7008 (28)	3.001 (11)	7.727 (23)
I _γ ; LWM	= WM	= WM	= WM	= WM	= WM	= WM	0.59 (5) ^x	= WM	3.006 (30) ^x	7.736 (48) ^e
I _γ ; Norm Res								0.701 (5)	2.999 (22)	7.727 (44)
I _γ ; Rajeval								0.713 (6)	2.997 (14)	7.713 (24)
Adopted I_g	0.1229 (16)	4.283 (13)	0.180 (4)	0.0779 (9)	0.0621 (21)	15.46 (4)	0.639 (3)	0.706 (4)	3.019 (14)	7.746 (13)
Source	All; WM	All; WM	All; NR	All; WM	All; WM	All; WM	All; WM	All; LWM	All; NR	All; WM

Table 2: ^{56}Co Relative Gamma-Ray Emission Probabilities (continued)[@] Experimental Data

Ref./E _g	2113.1g	2212.9g	2276.4g	2373.7g	2523.0g	2598.4g	2657.4g	3009.6g
65Pe18						17.4 (15)		1.3* (4)
66Do07						16.0* (27)		1.9* (8)
66Hu17	0.40 (9)	0.43 (9)	0.12 (3)	0.15* (3)	<0.03	20.0* (20)		1.25* (25)
66Sc01						17.4 (17)		1.5* (2)
67Au01	0.29 (5)					17.3 (9)		0.9 (2)
67Ba75	0.4 (1)	0.4 (1)				15.0* (13)		0.8 (3)
68Sh07	0.32 (15)	0.20* [†] (2)				17.0 (6)		1.0 (1)
67Ch20	0.56* (8) (12)	0.60* (9) (13)				14.0* (21) (30)		0.60* (9) (13)
69Ar04	0.3 (1)					18.7* (11)		0.9 (5)
69Au09						16.55 (44)		
69Sc09	0.32 (4)	0.46* (5)	0.14 (2)	0.11 (2)	0.09 (3)			
70Ph01	0.39 (3)	0.40 (3)	0.15 (2)	0.12 (2)	0.054 (15)	17.2 (4)		0.93 (6)
71Ca14	0.387 (4)	0.377 (10)	0.106 (5)	0.055 (12)	0.060 (5)	16.85 (17)		1.010 (11)
71Ge07						18.0* (9)		
71Ge08	0.26* (3)	0.28* (3)	0.10 (2)	0.08 (2)	0.07 (2)	16.5 (10)	~0.02	0.92 (10)
71Si29 ^S	0.34 (4)	0.30* (6)						1.55* (12)
72Pe20 ^d						15.65* (204) (224) 17.3 (22) (24) 14.44* (175) (193)		
74BoXX ^S						17.3 (4)		1.0 (2)
74Ho25	0.37 (2)	0.36 (2)	0.128 (8)	0.059 (12)	0.044 (10)		0.016 (5)	0.98 (9)
75Ka06	0.0.375 (17)	0.387 (18)	0.146 (7)			16.64 (22)		0.922 (29)
77Ge12 ^S	0.387 (8) (9)	0.406 (9) (10)				17.34 (26) (31)		1.06 (3) (3)
78Ha53 ^S	0.34 (2)	0.39 (2)	0.15 (2)	0.050 (6)	0.084 (9)	17.19 (15)	0.029 (4)	1.05 (3)
80St20 ^S	0.375 (14)	0.42 (2)	0.117 (9)	0.097 (12)	0.079 (11)	17.40 (38)	<0.05	0.84 (4)
80Sh28	0.35 (1)	0.35 (1)	0.115 (10)	0.079 (10)	0.14* [†] (1)	16.41 (33)	0.015 (3)	1.02 (2)
80Yo05	0.363 (7) (7)	0.389 (8) (8)	0.124 (7) (7)	0.083 (11) (11)	0.068 (11) (11)	16.96 (6) (8)	0.021 (6) (6)	
82Gr10								
89Al25								
90Me15 ^S	0.385 (5)	0.35 (1)	0.110 (5)	0.08 (1)	0.060 (5)	17.29 (15)		1.05 (1)
92ScZZ	0.376 (10) (10)	0.395 (14) (14)	0.128 (19) (19)	0.082 (22) (22)		17.26 (28) (28)		1.16 (3) (3)
00Ra36 ^S						17.1 (3)		
02MoZP ^S	0.372 (4) (4)	0.388 (4) (4)				16.82 (7) (8)		1.033 (11) (11)

Table 2: ⁵⁶Co Relative Gamma-Ray Emission Probabilities (continued)[@]

a) Analysis

Eg	2113.1g	2212.9g	2276.4g	2373.7g	2523.0g	2598.4g	2657.4g	3009.6g
All Data								
# data points, N	21	19	13	12	10	28	4	23
$\chi^2/(N-1)$	2.5 ^b	7.6 ^b	2.8 ^b	3.6 ^b	7.6 ^b	1.3	2.8	4.9 ^b
I_{γ} , UWM	0.365 (13)	0.383 (18)	0.126 (5)	0.087 (8)	0.075 (8)	16.89 (22)	0.020 (3)	1.07 (6)
I_{γ} , WM	0.3764 (21)	0.3795 (27)	0.1192 (24)	0.071 (3)	0.0687 (27)	16.97 (4)	0.0195 (20)	1.029 (5)
I_{γ} , LWM	0.376 (11) ^x	0.380 (8) ^x	0.119 (13) ^x	0.087 (37) ^x	0.069 (9) ^x	= WM	= WM	1.029 (21) ^x
I_{γ} , Norm Res	0.3761 (31)	0.385 (5)	0.1179 (36)	0.077 (6)	0.064 (4)	16.97 (4)	0.0184 (22)	1.030 (9)
I_{γ} , Rajeval	0.3756 (22)	0.387 (3)	0.1187 (28)	0.079 (4)	0.062 (3)	16.96 (4)	0.0168 (23)	1.029 (6)
Statistical Outliers Excluded			N/A				N/A	
# data points, N	19	14	-	11	9	20	-	17
$\chi^2/(N-1)$	1.9	2.8 ^b	-	3.3 ^b	1.7	1.2	-	4.4 ^b
UWM	0.360 (8)	0.389 (6)	-	0.081 (7)	0.068 (5)	17.06 (7)	-	0.975 (22)
WM	0.3769 (21)	0.3835 (27)	-	0.070 (3)	0.0631 (28)	16.96 (4)	-	1.028 (5)
LWM	= WM	0.384 (5) ^e	-	0.070 (20) ^x	= WM	= WM	-	0.975 (75) ^x
Selected Data								
# data points, N	6	6	3	3	3	7	1	7
$\chi^2/(N-1)$	1.9	4.4 ^b	2.0	7.8 ^b	3.4	2.2	N/A	7.5 ^b
I_{γ} , UWM	0.367 (9)	0.376 (18)	0.126 (12)	0.076 (14)	0.074 (7)	17.21 (7)	0.029 (4)	1.08 (8)
I_{γ} , WM	0.3770 (29)	0.386 (3)	0.113 (4)	0.064 (5)	0.067 (4)	17.01 (6)	-	1.039 (7)
I_{γ} , LWM	= WM	0.385 (9) ^e	= WM	0.068 (18) ^x	= WM	= WM	-	1.039 (19) ^e
I_{γ} , Norm Res	0.3770 (29)	0.389 (6)	0.113 (4)	0.080 (7)	0.072 (5)	17.13 (8)	-	1.043 (11)
I_{γ} , Rajeval	0.3773 (35)	0.390 (4)	0.112 (4)	0.082 (8)	0.080 (7)	17.20 (8)	-	1.043 (7)
Adopted I_{γ}	0.376 (3)	0.385 (5)	0.118 (4)	0.078 (6)	0.063 (4)	16.97 (4)	0.0195 (20)	1.039(19)
Source	All; NR	All; NR	All; NR	All; NR-Raj	All; NR-Raj	All; WM	All; WM	Sel; LWM

Table 2: ^{56}Co Relative Gamma-Ray Emission Probabilities (continued)[@], Experimental Data

Ref./E _g	3201.9g	3253.4g	3273.0g	3369.7g	3451.1g	3547.9g	3600.7g	3611.8g
65Pe18	3.2 (5)	8.5 (6)	1.5 (4)		0.95 (15)			
66Do07	2.9 (11)	5.8* (22)	1.2 (5)		0.7 (3)	0.2 (1)		
66Hu17	3.80* (45)	9.2* (9)	2.1 (4)		1.1 (2)	0.16 (3)	0.010 (5)	<0.005
66Sc01	3.4 (4)	8.3 (8)	1.9 (3)		0.7 (1)	0.21 (3)		
67Au01	3.4 (2)	7.8 (4)	1.5 (3)		0.87 (9)	0.15 (3)		
67Ba75	2.9 (3)	6.6 (6)	1.35 (20)		0.63* (15)	0.11* (5)		
68Sh07	2.8 (4)	7.3 (5)	1.5 (4)		0.83 (10)	0.15 (5)	0.02 (1)	
67Ch20	2.9 (4) (6)	7.2 (11) (15)	1.60 (24) (34)		0.72 (11) (15)	0.20 (3) (4)		
69Ar04	3.0 (2)	7.1 (4)	1.3 (1)		0.8 (1)	0.1* (1)		
69Au09	3.03 (14)	7.35 (21)	1.72 (13)		0.85 (7)			
69Sc09							0.024* (4)	0.007 (3)
70Ph01	3.10 (11)	7.5 (2)	1.72 (5)		0.89 (3)	0.18 (1)	0.014 (4)	0.011 (3)
71Ca14	3.03 (3)	7.390 (75)	1.755 (18)	0.011 (2)	0.875 (9)	0.178 (3)	0.015 (1)	0.0065 (10)
71Ge07	3.20 (35)	7.7 (9)	1.71 (25)		0.93 (20)	0.2 (1)		
71Ge08	2.81 (28)	7.0 (6)	1.69 (17)	0.015 (3)	0.82 (1)	0.15 (2)	0.014 (3)	0.007 (2)
71Si29 ^S			1.71 (9)		0.94 (2)	0.20 (3)		
72Pe20 ^d	2.86 (34) (38) 3.03 (36) (40) 2.55* (33) (36)	6.98 (86) (96) 7.4 (8) (9) 6.52 (78) (86)	- 1.57 (21) (23) 1.25 (20) (21)		0.98 (24) (25) 1.03 (14) (15) 0.84 (13) (14)			
74BoXX ^S	3.2 (1)	8.2 (4)	1.9 (1)		1.00 (4)	0.20 (2)		
74Ho25				0.008 (2)	0.89 (4)	0.178 (9)	0.016 (2)	0.008 (2)
75Ka06	3.067 (157)	7.45 (43)	1.697 (103)		0.936 (84)	0.164 (18)		
77Ge12 ^S	3.18 (10) (10)	7.79 (24) (25)	1.85 (6) (6)		0.93 (3) (3)	0.190 (6) (6)	0.0165 (7) (7)	0.0085 (4) (4)
78Ha53 ^S	3.24 (3)	7.97 (11)	1.84 (3)	0.010 (1)	0.95 (2)	0.196 (5)	0.012 (3)	0.005 (2)
80St20 ^S	3.03 (7)	7.60 (15)	1.815 (36)	0.011 (2)	0.90 (2)	0.196 (6)	0.015 (2)	0.010 (2)
80Sh28	3.04 (6)	7.52 (15)	1.77 (4)	0.007 (2)	0.90 (2)	0.19 (5)	0.015 (3)	0.007 (2)
80Yo05								
82Gr10								
89Al25								
90Me15 ^S	3.24 (3)	7.937 (65)	1.89 (2)	0.011 (2)	0.954 (10)	0.198 (5)	0.018 (1)	
92ScZZ	3.32 (7) (7)	8.13 (17) (17)	1.93 (4) (4)		0.973 (20) (20)	0.200 (5) (5)		
00Ra36 ^S	3.16 (6)	7.815 (160)	1.84 (4)		0.93 (3)	0.19 (1)		
02MoZP ^S	3.196 (17) (18)	7.85 (4) (4)	1.854 (12) (13)		0.94 (1) (1)	0.196 (2) (2)		

Table 2: ⁵⁶Co Relative Gamma-Ray Emission Probabilities (continued)[@], Analysis

Eg	3201.9g	3253.4g	3273.0g	3369.7g	3451.1g	3547.9g	3600.7g	3611.8g
All Data								
# data points, N	27	27	27	7	29	24	12	9
$\chi^2/(N-1)$	2.4 ^b	2.8 ^b	3.7 ^b	1.1	5.8 ^b	2.2 ^b	1.2	1.1
I _y ; UWM	3.10 (5)	7.55 (13)	1.68 (4)	0.0104 (10)	0.888 (19)	0.179 (6)	0.0158 (11)	0.0078 (6)
I _y ; WM	3.172 (11)	7.776 (27)	1.826 (8)	0.0100 (7)	0.905 (4)	0.1914 (13)	0.0162 (4)	0.0081 (3)
I _y ; LWM	3.10 (10) ^x	7.55 (30) ^x	1.68 (17) ^x	= WM	0.905 (30) ^x	0.179 (17) ^x	= WM	= WM
I _y ; Norm Res	3.188 (16)	7.82 (4)	1.838 (13)	0.0100 (7)	0.931 (7)	0.1934 (14)	0.0162 (4)	0.0081 (3)
I _y ; Rajeval	3.194 (12)	7.825 (28)	1.837 (9)	0.0100 (7)	0.932 (5)	0.1939 (14)	0.0162 (5)	0.0080 (4)
Statistical Outliers Excluded			N/A	N/A				N/A
# data points, N	25	25	-	-	28	22	11	-
$\chi^2/(N-1)$	2.4 ^b	2.9 ^b	-	-	5.8 ^b	2.2 ^b	0.99	-
UWM	3.089 (34)	7.56 (10)	-	-	0.897 (18)	0.185 (4)	0.0150 (8)	-
WM	3.173 (11)	7.775 (27)	-	-	0.905 (4)	0.1914 (13)	0.0161 (4)	-
LWM	3.09 (11) ^x	7.56 (29) ^x	-	-	0.905 (30) ^x	0.185 (11) ^x	= WM	-
Selected Data								
# data points, N	7	7	8	3	8	8	4	3
$\chi^2/(N-1)$	1.6	1.1	1.1	0.17	1.2	0.22	1.7	1.8
I _y ; UWM	3.178 (27)	7.88 (7)	1.837 (21)	0.0107 (3)	0.943 (10)	0.1958 (14)	0.0154 (13)	0.0078 (15)
I _y ; WM	3.205 (13)	7.868 (31)	1.856 (9)	0.0103 (8)	0.943 (6)	0.1957 (16)	0.0167 (5)	0.0084 (4)
I _y ; LWM	= WM	=WM	=WM	=WM	= WM	=WM	= WM	= WM
I _y ; Norm Res	3.205 (13)	7.868 (31)	1.856 (9)	0.0103 (8)	0.943 (6)	0.1957 (16)	0.0167 (5)	0.0084 (4)
I _y ; Rajeval	3.209 (13)	7.871 (31)	1.853 (10)		0.944 (6)	0.1957 (16)	0.0166 (6)	0.0085 (4)
Adopted I_g	3.205 (13)	7.87 (3)	1.856 (9)	0.0103 (8)	0.943 (6)	0.1957 (16)	0.0167 (5)	0.0084 (4)
Source	Sel; WM	Sel; WM	Sel; WM	Sel; WM	Sel; WM	Sel; WM	Sel; WM	Sel; WM

[@] Experimental data are listed along with the authors' statistical uncertainty in the least significant digits (given in parentheses). If two numbers are shown in parentheses, the second is the uncertainty after any uncertainty in the reference line (847) has been combined in quadrature with the former uncertainty. Note that reference codes are given with the leading two digits of the code omitted. In the 'Analysis' section, the following abbreviations have been used: UWM= unweighted mean; WM= weighted mean; LWM= values recommended by the program LWEIGHT; Norm Res = result from Normalised residuals analysis; Rajeval = result from Rajeval technique analysis; ~~N~~-Raj = mean of values from Normalised Residuals and Rajeval technique analyses, using the larger of the two uncertainties 'Sel' refers to data from eight selected references in

which the detector efficiency curves were well-characterised to at least 3600 keV (2002MoZP, 2000Ra36, 1990Me15, 1980St20, 1978Ha53, 1977Ge12, 1974BoXX and 1971Si29).

* This γ -ray intensity datum is identified by LWEIGHT as a statistical outlier based on the Chauvenet criterion.

^a Transition reported in one study only.

^b Exceeds critical value for $\chi^2/(N-1)$ so LWEIGHT considers the data in this dataset to be discrepant.

^c Reported as 0.310 in 1980Sh28, but this is clearly a typographical error; the value from the literature with which it is compared is also an order of magnitude too large.

^d 1972Pe20 took data using three different detectors (cylindrical, rectangular and trapezoidal), each calibrated using Monte Carlo calculations; data from these detectors are shown separately.

^e Weighted mean, external uncertainty recommended by LWEIGHT.

^f Datum rejected by Rajeval analysis.

^s Data from this reference included in 'selected data' analysis.

^x LWEIGHT has expanded the uncertainty to encompass the most precise datum.

⁵⁷Co - Comments on evaluation of decay data**by V. P. Chechev and N. K. Kuzmenko****1. Decay Scheme**

The 2nd forbidden electron capture (EC) transitions to the 3/2⁻ excited levels of 14,413 keV and 366,74 keV have not been observed, as well as the 2nd forbidden unique EC transition to the 1/2⁻ ground state of ⁵⁷Fe. From the log ft systematics the log ft of the 2nd forbidden transitions should be greater than 11,1 and 10,8, respectively, and for the 2nd forbidden unique transition, greater than 12,9. From these, the upper limits on the EC branch probabilities to the 14,413 keV level and ground state of ⁵⁷Fe are obtained as < 0,003 % and < 0,00035 %, and for the EC branch to the 366,74 keV level ≤ 0,002%. The calculations of the level probability balance in the decay scheme of ⁵⁷Co were made not taking into account the first two unobserved transitions. The EC branch probabilities to the levels of 136,47 keV, 366,74 keV and 706,42 keV were obtained from an probability balance of the gamma transitions.

2. Nuclear Data

Q value is from Audi and Wapstra (1995Au04).

There are available eight measurement results of the half-life of ⁵⁷Co (Table 1).

Table 1. Measurement results and evaluation of the half-life of ⁵⁷Co

Reference	Data set "1" $\chi^2=39,2$ $(\chi^2)_7^{0,05}=14,1$	Data set "2" $\chi^2=14,5$ $(\chi^2)_6^{0,05}=12,6$	Data set "3" $\chi^2=14,5$ $(\chi^2)_6^{0,05}=12,6$
1997Ma75	271,68(9)	271,68(9)	271,68(9)
1992Un01	272,11(26)	272,11(26)	272,11(26)
1983Wa26	271,84(4)	271,84(4)	271,84(4)
1981Va11	271,90(9)	271,90(9)	271,90(9)
1980Ho17	271,77(5)	271,77(5)	271,77(5)
1972La14	271,23(21)	271,23(21)	271,23(21)
1972Em01	269,8(4)	Omitted	Omitted
1965An07	271,65(13)	271,65(13)	271,65(13)
Evaluated value 271,80(5) d			

The value of 269,8(4) days from 1972Em01 was omitted on statistical considerations (because of a large contribution to χ^2 and also on the Chauvenet's criterion). This leads to the data set "2" of the seven values which coincides with the final data "3" as the LRSW method in statistical processing of the set "2" does not change the relative statistical weights.

The computer program EV1NEW 2000Ch01 has chosen the weighted mean of 271,80(5) days with the tS (or MBAYS) uncertainty as $(\chi^2)^{0,05}_{n-1} < \chi^2 < 10(\chi^2)^{0,05}_{n-1}$ (see evaluation technique in 2000Ch01). Other statistical procedures give, UWM-271,74(10), WM-271,80(3), CHV-271,83(7), UINF-271,80(4), PINF-271,80(4), BAYS-271,80(5), LWM-271,80(4), IEXW-271,75(8), NORM-271,80(4), RAJ-271,80(3). The computer program LWEIGHT leads to 271,80(3) days, the weighted mean with the internal uncertainty (the external uncertainty is 0,042). (The other evaluations of half-life of ⁵⁷Co see in 1990Ni03 and 1998Bh11).

The adopted value for the half-life of ⁵⁷Co is 271,80(5) days.

Half-life of excited levels in ⁵⁷Fe

The half-life of the excited levels (136 and 14 keV) have been evaluated being : **8,8(5)** ns [using 1989Ra17 and 1978AlZX] and **98,0(3)** ns [from 1961C111, 1965Ki03, 1967Ec05, 1969Ho28, 1978AlZX, 1995Ah04], respectively.

2.1. Electron Capture Transitions

The energies of the electron capture, ϵ , transitions have been calculated from the Q value and the level energies deduced from gamma transition energies.

The P_K , P_L and P_M values have been obtained from the tables of Schönfeld (1998Sc28). The experimental P_K values are available for $\epsilon_{0,2}$ EC transitions to the level of 136,47 keV: 0,885(9) in 1968Ru04 ; 0,87(2) in 1969Bo49 ; 0,922(10) in 1973 Mukerji and 0,89(4) in 1990Si03.

The electron capture probabilities of $\epsilon_{0,2}$, $\epsilon_{0,3}$ and $\epsilon_{0,4}$ have been calculated from the balance of the evaluated $P_{\gamma+ce}$ values for the 136,47 keV, 366,74 keV and 706,42 keV levels, respectively, assuming negligible EC transitions to the 14,4 keV level and the ground state of ⁵⁷Fe.

The calculated value of the sum of $P_{\gamma+ce}$ for the 4 gamma transitions to the ground state of ⁵⁷Fe is 99,996 (19) %.

2.2. Gamma Transitions and Internal Conversion Coefficients

The evaluated energies of gamma transitions are the energies of the gamma rays plus the recoil energy.

The probabilities of gamma transitions $P_{\gamma+ce}$ have been computed using the evaluated absolute gamma ray emission intensities and the total internal conversion coefficients (ICC). The ICC have been evaluated using the experimental information on the multipolarity admixture coefficients (see below) and the theoretical values from 1976Ba63.

The values of $\delta(E2/M1)$ have been adopted from the analysis of 1978Kr19 except for $\gamma_{2,1}$ which is obtained by weighting the 4 values of +0,120 from 1972Fo05, +0,116(1) from 1973Sc15, +0,1195(10) from 1975Co22 and +0,120(4) from 1972Kr15 (see also the evaluation of 1998Bh11). The weighted average of $\delta(E2/M1)$ for $\gamma_{2,1}$ is +0,1180(12).

The adopted values of $\delta(E2/M1)$ for other gamma transitions are 0,00223(18) for $\gamma_{1,0}$, +0,02 for $\gamma_{3,2}$, +0,083(5) for $\gamma_{4,3}$, +0,025(9) for $\gamma_{3,1}$, -0,45(5) for $\gamma_{3,0}$, +0,097(8) for $\gamma_{4,2}$ and -0,465(8) for $\gamma_{4,1}$.

There are many experimental values of ICC and the ratios of the fractional intensities of conversion electrons for $\gamma_{1,0}$, $\gamma_{2,1}$ and $\gamma_{3,0}$ which, with the exception of 1996Me11, support the adopted values of ICC:

$\gamma_{1,0}$ $\alpha_K=7,76(23)$, $\alpha_L=0,804(24)$ from 1976Ba63
 $\alpha_K=7,35(19)$ from 1985HaZA
 K:L:M+=100:9,59(13):1,48(15) from 1971Po05

$\gamma_{2,1}$ $\alpha_K=0,0214(12)$, K/L+=8,2(6) from 1967Ha06
 K:L:M+=100:9,0:1,5 from 1955Co31

$\gamma_{3,0}$ $\alpha_K=0,122(13)$, K/L+=8,6(5), $\alpha_T/\alpha_K=1,118(5)$ from 1967Ha06

There are 6 experimental values for the total ICC (α_T) of the low-energy gamma transition $\gamma_{1,0}$ (14,413 keV): 9,0(5) and 8,9(6) from 1965Ki03 ; 8,26(22) from 1965Mo22 ; 8,25(46) from 1966Sp06 ; 8,26(22) from 1968Ru04 and 8,19(18) from 1970Jo30. They can be compared to the adopted value of $\alpha_T=8,58(18)$.

3. Atomic Data

3.1. Fluorescence yields

The fluorescence yields are taken from 1996Sc06 (Schönfeld and Janßen).

3.2. X Radiations

The X-ray energies are based on the wavelengths in the compilation of 1967Be65 (Bearden).

The relative $K\beta/K\alpha$ emission probability is taken from 1998Be and 1997Lepy. They have shown that taking into account double-electron transitions with a simultaneous emission of a photon and Auger electron (the radiative Auger effect RAE) increases the value of $K\beta/K\alpha$ = from 0,1368(14) (1996Sc06) to 0,1419(19) (1998Be) or 0,1423(17) (1997Lepy). From these we have adopted $K\beta/K\alpha = 0,142(2)$.

The ratio $K\alpha_2/K\alpha_1$ is from 1996Sc06

3.3. Auger Electrons

The energies of Auger electrons are from 1977La19 (Larkins).

The ratios $P(KLX)/P(KLL)$ and $P(KXY)/P(KLL)$ are taken from 1996Sc06.

4. Photon Emissions

4.1 X-Ray Emissions

The total absolute emission intensity of KX-rays (P_{XK}) has been computed using the adopted value of ω_K , the evaluated total absolute emission probabilities (sums) of K conversion electrons (P_{ceK}) and K electron capture ($P_{\epsilon K}$).

The absolute emission intensities of the KX -ray components have been computed from the total P_{XK} using the relative probabilities from sect. 3.2.

Below the measured values of P_{Kα} and P_{XK} are compared to our calculated (evaluated) values:

	<i>Measured</i>		<i>Calculated</i>
	1989 Debertin	1994Ar22	(evaluated)
P_{Kα}, %	50,6(9)	50,1(5)	50,0(6)

	<i>Measured</i>			<i>Calculated</i>
	1968Ru04	1973 Mukerji	1978 Vylov	1989 Debertin (evaluated)
P_{XK}, %	56,9(8)	58,4(17)	55,3(15)	56,0(11) 57,1(9)

The total absolute emission intensity of LX -rays has been computed using absolute sums P_{CeL}, P_{CeK}, P_{EK}, P_{EL} and atomic data of section 3.1 (ω_K , $\bar{\omega}_K$, n_{KL}).

4.2. Gamma Emissions

The energies of the gamma rays $\gamma_{2,1}$ and $\gamma_{3,0}$ have been adopted from 1976Bo16 and 2000He14. The energies of other gamma rays have been obtained as the weighted means of measurement results listed in Table 2 or calculated from the decay scheme of ⁵⁷Co. The corrections to the revised energetic scale in 2000He14 (lowering the values by 5,80 ppm) do not change these values.

The evaluator has assumed no EC feeding to the ground and first excited states and used the total gamma -ray transition probabilities to these two states (except that for the 14,4 -keV transition) to normalize the decay scheme (using adopted relative photon intensities from Table 3, conversion coefficients from Section 2.2). This procedure has produced a normalization factor of 0,8551(6).

The absolute gamma ray emission intensity for $\gamma_{1,0}$ (14,413 keV) has been computed as follows: $P_\gamma(\gamma_{1,0}) = P_{\gamma+ce}(\gamma_{1,0}) / (1 + \alpha_T(\gamma_{1,0}))$, where $P_{\gamma+ce}(\gamma_{1,0}) = 87,57(16)$ comes from decay-scheme probability balance at the 14,4 -keV level, and $\alpha_T(\gamma_{1,0}) = 8,58$. The deduced value of $P_\gamma(\gamma_{1,0}) = 9,15(17)$ % can be compared with the experimental values, such as 9,5(2) % (1978Vylov), 9,54(12) % (1992ScZZ) and 9,16(15) % (1989Debertin). It agrees extremely well with the CRP experimental result from 1989 Debertin.

It should be noted also that the evaluated sum $P_\gamma(\gamma_{2,0}) + P_\gamma(\gamma_{1,0}) = 19,86(23)$ % agrees well with the measured value of 19,84(17)% in 1971Ko19.

Table 2 - Measured and adopted energies of gamma-rays in the decays of ⁵⁷Co → ⁵⁷Fe and ⁵⁷Mn → ⁵⁷Fe

	1965Ki03	1965Sp06	1970Gr13	1971Ko19	1972He42	1974Ti01 ^a	1976Bo16	1980Ve05	WM	Adopted
$\gamma_{1,0}$			14,408(5)		14,41247(29)	14,410(6)			-	14,41295(31) ^b
$\gamma_{2,1}$			122,07(3)	122,06(2)		122,063(4)	122,06065(12)		-	122,06065(12)
$\gamma_{2,0}$			136,473(4)	136,47(3)		136,473(4)	136,47356(29)		-	136,47356(29)
$\gamma_{3,2}$	229,8(10)	230,6(6)	230,4(5)	230,4(6)		230,25(4)		230,29(2)	230,27(3)	230,27(3)
$\gamma_{4,3}$	339,7(4)	339,7(5)	339,7(3)	339,68(28)		339,60(6)		339,54(18)	339,61(9)	339,67(3) ^b
$\gamma_{3,1}$	352,5(4)	352,4(5)	352,5(3)	352,23(27)		352,32(3)		352,36(1)	352,34(2)	352,34(2)
$\gamma_{3,0}$	366,8(5)	366,7(5)	336,8(4)	367,0(5)		366,73(4)		366,75(1)	366,74(3)	366,74(3) ^b
$\gamma_{4,2}$	570,0(4)	570,3(4)	570,1(3)	570,04(28)		569,93(5)		569,92(4)	569,94(4)	569,94(4)
$\gamma_{4,1}$	692,1(3)	692,1(3)	692,1(2)	692,44(6)		692,00(3)		692,03(2)	692,02(2)	692,01(2) ^b
$\gamma_{4,0}$	706,4(4)	706,8(4)	706,6(3)	706,46(34)		706,54(22)		706,40(20)	706,50(20)	706,42(2) ^b

a Experimental values from the decay of ⁵⁷Mn

b Calculated from decay scheme using the energies of $\gamma_{2,1}$, $\gamma_{2,0}$, $\gamma_{3,2}$, $\gamma_{3,1}$, $\gamma_{4,2}$

Table 3 - Relative emission probabilities of gamma rays in the decay of ⁵⁷Co

γ	E_γ	1965Ki03	1965Ma38	1971Ko19	1974 HeYW	1980Sc07 ^a	1982Gr10	Average	Adopted
$\gamma_{1,0}$	14			$1,14(5) \cdot 10^4$					$10,70(20)$ ^b
$\gamma_{2,1}$	122	10^5	10^5	10^5	10^5	10^5	10^5	10^5	100
$\gamma_{2,0}$	136	$1,25(8) \cdot 10^4$	$1,20(1) \cdot 10^4$	$1,30(4) \cdot 10^4$	$1,29(7) \cdot 10^4$	$1,236(9) \cdot 10^4$	$1,245(30) \cdot 10^4$	$1,253(18) \cdot 10^4$ ^c	$12,53(18)$
$\gamma_{3,2}$	230		0,2(2)	0,5(5)					$4(4) \cdot 10^{-4}$
$\gamma_{4,3}$	340		2,9(3)	4,5(4)					$0,0045(4)$ ^d
$\gamma_{3,1}$	352		2,0(2)	3,7(4)					$0,0037(4)$ ^d
$\gamma_{3,0}$	367		0,7(1)	1,5(4)					$0,0015(4)$ ^d
$\gamma_{4,2}$	570		16(1)	19,4(11)	10(10)			$18(2)$ ^e	$0,018(2)$
$\gamma_{4,1}$	692		188(5)	183(11)	190(30)			$186(7)$ ^f	$0,186(7)$
$\gamma_{4,0}$	706		5,5(6)	6,2(6)				$5,8(6)$ ^g	$0,0058(6)$

^a In 1980Sc07 the absolute gamma-ray emission probabilities are reported: $P_{\gamma_{2,0}(136)}=10,58(8)\%$ and $P_{\gamma_{2,1}(122)}=85,59(19)\%$. Their ratio is $0,1236(9)$.

^b Calculated as described in the text

^c The LWEIGHT program (version 1.2) has used an unweighted average and expanded the uncertainty so range includes the most precise value of 1980Sc07 . It is reasonable choice because of disagreement of the experimental values some uncertainties of which are only statistical.

^d Adopted from 1971Ko19.

^e LWEIGHT has used a weighted average and expanded the uncertainty so range includes the most precise value of 1965Ma38.

^f The method of Limitation of Relative Statistical Weights (LRSW) increased the uncertainty of 1965Ma38 to 10,3.

^g The experimental uncertainty is adopted as the uncertainty of the evaluated value.

5. Electron emissions

The energies of the conversion electrons have been calculated from the gamma transition energies given in sect. 2.2 and the electron binding energies.

The emission intensities of the conversion electrons have been calculated using the transition probabilities given in sect. 2.1 and 2.2, the atomic data given in sect. 3, and the internal conversion coefficients given in sect. 2.2.

The low energy electron spectrum from the decay of ⁵⁷Co has been analysed in 1997KoZJ using a combined electrostatic spectrometers. They obtained the following intensity ratios for the main spectrum components: (LMM+LXY) / KLL / KLLX / KMX / K-14,4 / L-14,4 / (M+N)-14,4 = 49,3(38): 59,6(23): 15,2(6): 1,2(2): 49,9(18): 5,1(3): 0,80(4). These values agree mainly with our evaluated data on electron emissions apart from the intensity of L Auger electrons. Perhaps, the latter is connected with difficulties of the electron spectrum measurement in the energy region of 0,6 - 0,7 keV. The discrepancy takes place also for the L/(M+N) and K/(M+N) ratios.

Also in 1997KoZJ $L_1/L_2 = 15,7(5)$, $L_1/L_3=39,3(16)$, $M_{2,3}/M_1=0,076(4)$ have been measured for the gamma transition $\gamma_{1,0}$ (14,4 keV).

References

1955Co31 J. M. Cork, M. K Brice, L. C Schmid, Phys. Rev. 99 (1955) 703.
(Relative conversion electron intensities)

1965An07 S. C. Anspach, L. M. Cavallo, S. B. Garfinkel, J. M. R. Hutchinson, C. N. Smith, In: NP-15663 1965. (Half-life)

1965Ki03 O. C. Kistner, A.W. Sunyar, Phys. Rev. 139 (1965) B295.
(Experimental ICC, gamma ray energies and relative emission probabilities)

1965Ma38 J. M. Mathiesen, J. P. Hurley, Nucl. Phys. 72 (1965) 475.
(Relative gamma ray emission probabilities)

1965Mo22 G. Moreau, G. Ambrosino, Compt. Rend. 261 (1965) 5438.
(Experimental ICC)

1966Sp06 E. H. Spejewski, Nucl. Phys. 82 (1966) 481.
(Experimental ICC and gamma ray energies)

1967Be65 J. A. Bearden, Revs. Modern Phys. 39 (1967) 78.
(X-ray energies)

1967Ha06 D. C. Hall, R. G. Albridge, Nucl. Phys. A91 (1967) 495.
(Experimental ICC)

1968Ru04 W. Rubinson, K. P. Gopinathan, Phys. Rev. 170 (1968) 969.
(Experimental ICC, P_K and P_{XK} values)

1969Bo49 H. E. Bosch, M. A. Farioli, N. Martin, M. C. Simon, Nuclear Instrum. Methods 73 (1969) 323. (Experimental P_K values)

1970Gr13 R. C. Greenwood, R. G. Helmer, R. J. Gehrke, Nuclear Instrum. Methods 77 (1970) 141.
(Gamma ray energies)

1970Jo30 D. P. Johnson, Phys. Rev. B1 (1970) 3551.
(Experimental ICC)

1971Ko19 J. Konijn, E. W. A. Lingeman, Nuclear Instrum. Methods 94 (1971) 389.
(Gamma ray energies and emission probabilities)

- 1971Po05 F. T. Porter, M. S. Freedman, Phys. Rev. C3 (1971) 2285.
(Relative conversion electron intensities)
- 1972Fo05 R. A. Fox, W. D. Hamilton, M. J. Holmes, Phys. Rev. C5 (1972) 853.
(Mixing ratios E2/M1 for gamma transitions)
- 1972He42 U. Heim, O. W. B. Schult, Z. Naturforsch.27a (1972) 1861.
(14,4 keV gamma ray energy)
- 1972Kr15 K. S. Krane, W. A. Steyert, Phys. Rev. C6 (1972) 2268.
(Mixing ratios E2/M1 for gamma transitions)
- 1972La14 F. Lagoutine, J. Legrand, C. Perrot, J. P. Brethon, J. Morel, Intern. J. Appl. Radiat. Isotop. 23 (1972) 219. (Half-life)
- 1973 Mukerji A. Mukerji, Lee Chin, In: Proc. of the Intern. Conf. on Inner-Shell Ionization Phenomena and future Applications, April 17-22, 1972, Conf-720404, USAEC-Technical Information Center, Oak Ridge, Tenn. (1973). (Experimental P_K and P_{XK} values)
- 1973Sc15 E. Schoeters, R. E. Silverans, L. Vanneste, Z. Phys. 260 (1973) 337.
(Mixing ratios E2/M1 for gamma transitions)
- 1974HeYW R. L. Heath, Gamma-Ray Spectrum Catalogue (1974), ANCR-1000-2.
(Relative gamma ray emission probabilities)
- 1974Ti01 K. G. Tirsell, L. G. Multhauf, S. Raman, Phys. Rev. C10 (1974) 785.
(Gamma ray energies)
- 1975Co22 E. J. Cohen, A. J. Becker, N. K. Cheung, H. E. Henrikson, Hyperfine Interact.1 (1975) 193. (Mixing ratios E2/M1 for gamma transitions)
- 1976Ba63 I. M. Band, M. B. Trzhaskovskaya, M. A. Listengarten, At. Data Nucl. Data Tables 18 (1976) 433. (Theoretical ICC)
- 1976Bo16 G. I. Borchert, Z. Naturforsch. 31a (1976) 387.
(Gamma ray energies)
- 1977La19 F. P. Larkins, At. Data Nucl. Data Tables 20 (1977) 311.
(Auger electron emission)
- 1978Kr19 K. S. Krane, At. Data Nucl. Data Tables 22 (1978) 269.
(Mixing ratios E2/M1 for gamma transitions)
- 1978 Vylov C. Vylov, Preprint JINR R6-10416, Dubna, 1977.
(Experimental P_{XK} and $P(\gamma_{14,4 \text{ keV}})$ values)
- 1980Ho17 H. Houtermans, O. Milosevic, F. Reichel, Intern. J. Appl. Radiat. Isotop. 31 (1980) 153.
(Half-life)
- 1980Sc07 U. Schötzig, K. Debertin, K. F. Walz, Nuclear Instrum. Methods 169 (1980) 43.
(Relative gamma ray emission probabilities)
- 1980Ve05 R. Vennink, J. Kopecky, P. M. Endt, P. W. M. Glaudemans, Nucl. Phys. A344 (1980) 421. (Gamma ray energies)
- 1981Va11 R. Vaninbroukx, G. Grosse, W. Zehner, J. Appl. Radiat. Isotop. 32 (1981) 589.
(Half-life)
- 1982Gr10 A. Grutter, Intern. J. Appl. Radiat. Isotop. 33 (1982) 533.
(Relative gamma ray emission probabilities)
- 1983Wa26 K. F. Walz, K. Debertin, H. Schrader, Intern. J. Appl. Radiat. Isotop. 34 (1983) 1191.
(Half-life)
- 1985HaZA H. H. Hansen, In: European App. Res. Rept. Nucl. Sci. Technol. 6, No 4 (1985) 777.
(Compilation of experimental ICC)
- 1989 Debertin K. Debertin, U. Schötzig, informal IAEA CRP paper GS/55 (1989), quoted in IAEA-TecDoc-619 (1991) (Experimental KX and $\gamma_{1,0}$ (14,4 keV) absolute emission probabilities)

Comments on evaluation

- 1990Ni03 A. L. Nichols, Nucl. Instrum. Methods Phys. Res. A286 (1990) 467.
(Half-life)
- 1990Si03 K. Singh, T. S. Gill, K. Singh, Appl. Radiat. Isot. 41 (1990) 333.
(Experimental P_K values)
- 1992ScZZ U. Schötzig, H. Schrader, K. Debertin. In: Proc. Inter. Conf. Nuclear Data for Science and Technology, Julich, Germany, 1992. P.562.
(Experimental $P_\gamma(14,4 \text{ keV})$ value)
- 1994Ar22 D. Arnold, G. Ulm, Nucl. Instrum. Methods Phys. Res. A339 (1994) 43.
(Experimental $K\alpha$ absolute emission probability)
- 1995Au04 G. Audi, A. H. Wapstra, Nucl. Phys. A595 (1995) 409.
(Q value)
- 1996Me11 R. Ya. Metskhvarishvili, Z. N. Miminoshvili, M. A. Elizbarashvili, L. V. Nekrasova, I. R. Metskhvarishvili, N. G. Khazaradze, N. M. Marchilashvili, I. V. Zhorzholiani, Yad. Fiz. 59, N o 5 (1996) 773,
Phys. Atomic Nuclei 59 (1996) 737.
(Relative conversion electron intensities and experimental ICC)
- 1996Sc06 E. Schönfeld, H. Janssen Nucl. Instrum. Methods Phys. Res. A369 (1996) 527.
(Atomic data)
- 1997 Lépy M.-C. Lépy, M.-M. Bé, J. Plagnard, In: CAARI'96 Conference proceedings AIP Press, 1067 (1997).(Experimental $K\beta/K\alpha$)
- 1997Ma75 R. H. Martin, K. I. W. Burns, J. G. V. Taylor, Nucl. Instrum. Methods Phys. Res. A390 (1997) 267. (Half-life)
- 1997KoZJ A. Kovalik, M. Rysavy, V. M. Gorozhankin, Ts . Vylou, D. V. Filosofov, M. A. Mahmoud, A. Minkova, N. Coursol, P. Casette, Ch. Briancon. Program and Thesis, Proc. 47th Ann. Conf. on Nucl. Spectrosc. Struct. At. Nuclei, Obninsk, 1997. P. 277.
(Relative low energy electron emission probabilities)
- 1998 Be M.- M. Bé, M.- C. Lépy, J. Plagnard, B. Duchemin, Applied Radiation Isotopes 49 (1998) 1367. (Experimental $K\beta/K\alpha$)
- 1998Bh11 M. R. Bhat, Nucl. Data Sheets 85 (1998) 415.
(Mixing ratios $E2/M1$ for gamma transitions)
- 1998Sc28 E. Schönfeld, Appl. Radiat. Isot. 49 (1998) 1353.
(Fractional electron capture probabilities P_K, P_L, P_M)
- 2000Ch01 V. P. Chechev, A. G. Egorov, Appl. Radiat. Isot. 52 (2000) 601.
(Evaluation technique)
- 2000He14 R. G. Helmer, C. van der Leun, Nucl. Instrum. Methods Phys. Res. A450 (2000) 35.
(Gamma ray energies)

⁵⁷Ni – Comments on evaluation of decay data by Shiu-Chin Wu

The *Limitation of Relative Statistical Weight* (1988WoZO) (LWM) method, used for averaging numbers throughout this evaluation, provided a uniform approach for the analysis of discrepant data. For two discrepant values, the method chooses the unweighted average. The uncertainty assigned to the recommended values was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

1. Decay Scheme

⁵⁷Ni decays by EC + β^+ to ⁵⁷Co states at 1377.65, 1504.81, 1757.58, 1919.55 and 2804.27 keV. The total β^+ branching has been measured by 1967Li08, 1962Ch20, 1958Ko60 and 1964Ru06. The weighted average of the results gives (45.9 ± 1.0) %, in agreement with the value of 43.5% predicted by theory [1; 1957Zw01].

2. Nuclear Data

The following values of the half-life of ⁵⁷Ni have been used to deduce a recommended value:

1	35.54(5) h	Dickens (1986)
2	35.65(5) h	Grutter (1982)
3	36.16(11) h	Rothman et al. (1974)
4	35.99(12) h	Ebrey and Gray (1965)
5	35.7(2) h	Rudstam (1964)
6	36.4(7) h	Friedlander et al. (1950)
7	35.7(10) h	Maienschein and Meem (1949)

The recommended half-life of ⁵⁷Ni, 35.9(3) h, is an average ($\chi^2/N-1=5.83$, LWM) of the seven values listed above. The LWM method changed the uncertainty of the averaged value from 0.1 h to 0.3 h, in order to overlap with the most precise value of 35.54 h. The value of 43.7(9) h by Rayburn (1961Ra06) differs from the average by about 8 σ , and was not included. Rudstam (1956Ru45) had previously reported a value of 37.6(5) h, which has been superseded by the more precise value of 35.7(2) h (1964Ru06) given above.

2.1 Electron Capture Transitions

Electron-capture energies given in Tables 2.2 have been deduced from the Q value and the level energies. EC + β^+ feedings to the levels are from gamma -ray emission probability balances. The electron -capture and positron emission probabilities to the individual levels are based on theoretical [1] β^+ /EC ratios. The fractional atomic shell electron -capture probabilities are theoretical values [1977Ba48] calculated with the EC-CAPTURE computer program [2]. EC decay to the ground state of ⁵⁷Co has not been observed. This transition would be 2nd forbidden non-unique, with a systematic *lg ft* value of 11.0 or higher. Its

corresponding probability, calculated with the LOGFT computer program [3], is less than 0.01%. Similarly, the EC decay to the 1st excited state has a probability of less than 0.001%.

2.2 Positrons Transitions

Electron-capture and β^+ end-point energies given in Tables 2.1 and 2.2 are equal to $Q_{EC} = 3264.2(26)$ keV (1995Au04) minus the individual level energies, and to the electron β^- -capture energies minus $2 m_0c^2$ (1022 keV), respectively.

2.3 Gamma Rays

Gamma-ray energies were measured with Ge(Li) detectors by Scardino *et al.* (1990Sc23); Rothman *et al.* (1974HeYW); Gatrousis *et al.* (1969Ga14); Lingeman *et al.* (1967Li08) and Piluso *et al.* (1966Pi01). The energies adopted here are the LWM averages, which are usually dominated by the values of 90Sc23.

Adopted	1990Sc23	1974HeYW	1969Ga14	1967Li08	1966Pi01	χ^2_R
127.164(3)	127.164(3)	127.192(25)	127.1(1)	127.6(5)**	127.2(1)	0.59
161.86(3)	161.86(3)		161.8(3)			0.04
304.1(1)	304.1(1)					
379.94(2)	379.94(2)		380.0(2)			0.09
541.9(1)	541.9(1)					
673.44(4)	673.44(4)		673.4(2)			0.04
696.0(4)	696.0(4)					
755.3(1)	755.3(1)					
906.98(5)	906.98(5)		906.8(3)			0.35
1046.54(14) [#]	1046.68(3)		1046.4(2)			0.98
1223.8(3) [#]	1224.00(4)		1223.5(4)			0.78
1279.99(6)	1279.99(6)					
1350.52(6)	1350.52(6)					
1377.62(4)	1377.63(3)	1377.59(4)	1377.6(2)	1378.0(5)	1378.1(2)	1.7
1603.28(6)	1603.28(6)					
1730.45(6)	1730.44(6)		1730.6(3)			0.27
1757.55(3)	1757.55(3)	1757.48(8)	1757.6(2)	1758.2(6)**	1757.7(2)	0.45
1897.0(5) [#]	1897.42(4)		1896.5(4)			2.6
1919.62(14)	1919.52(5)	1919.43(8)	1919.5(2)	1919.9(6)	1920.2(1)	11
2133.04(5)	2133.04(5)		2132.9(3)			0.21
2730.76(14)	2730.91(4)		2730.6(2)	2731(2)		0.61
2804.08(15)	2804.20(3)		2803.9(2)	2805.1(9)		1.2
3177.27(5)	3177.28(5)		3176.9(3)	3177.3(12)		0.78

** Statistical outlier, omitted.

[#] The LWM chose the unweighted average for these discrepant values.

Gamma-ray emission probabilities relative to that of the 1377.62 keV γ -ray measured with Ge(Li) detectors were reported by Scardino *et al.* (1990Sc23); Grutter (1982Gr10); Rothman *et al.* (1974HeYW); Gatrousis *et al.* (1969Ga14); Lingeman *et al.* (1967Li08) and Piluso *et al.* (1966Pi01). The LWM averages have been adopted here.

E_γ keV	Adopted	1990Sc23	1982Gr10	1974HeYW	1969Ga14	1967Li08	1966Pi01	χ^2_R
127.164(3)	19.8(6)	20.4(4)	20.3(2)	16.6(10)	20.0(6)	17.6(9)	15.0(9)	10
161.86(3)	0.025(3) [#]	0.0278(8)			0.022(11)			14
304.1(1)	0.0024(7)	0.0024(7)						
379.94(2)	0.089(7) [#]	0.082(2)			0.10(5)			4.2
541.9(1)	0.0045(6)	0.0045(6)						
673.44(4)	0.0600(18)	0.0601(18) ¹⁾			0.06(3)			0.38
696.0(4)	0.0011(8)	0.0011(8)						
755.3(1)	0.0066(8)	0.0066(8)						
906.98(5)	0.092(18) [#]	0.075(2)			0.110(6)			20
1046.54(14)	0.163(4)	0.164(4)			0.16(1)			0.20
1223.8(3)	0.094(16) [#]	0.077(3)			0.110(6)			18
1279.99(6)	0.0118(9)	0.0118(9)						
1350.52(6)	0.0024(12)	0.0024(12)						
1377.62(4)	100(2)	100	100	100	100	100	100	
1603.28(6)	0.0048(8)	0.0048(8)						
1730.45(6)	0.068(4) [#]	0.064(3) ²⁾			0.072(4)			2.5
1757.55(3)	7.5(5)	7.04(20)	7.63(20)	9.1(8)	7.7(2)	9.5(5)	6.9(3)	6.1
1897.0(5)	0.031(3) [#]	0.034(3)			0.028(14)			2.0
1919.62(14)	15.4(7)	15.0(3)	17.0(4)	18.9(12)	17.0(5)	22.4(11) ³⁾	14.7(2)	10
2133.04(5)	0.041(6) [#]	0.035(2) ²⁾			0.047(24)			13
2730.76(14)	0.024(4)	0.0243(6)			0.03(2)	0.015(2)		18
2804.08(15)	0.126(21)	0.120(4)			0.17(1)	0.088(9)		23
3177.27(5)	0.019(5)	0.0136(7)			0.024(1)	0.021(3)		21

¹⁾ The relative intensity of the 673.44 -keV γ -ray was listed in 1990Sc23 as 0.0601(15), and corrected as 0.0601(8) by Bhat (1992Bh05). However, a relative uncertainty of 1% for such a weak peak seems too low, it is probably a typographical error. We used 0.0601(18) here.

²⁾ As suggested by Bhat (1992Bh05), the intensity given in 1990Sc23 for the 1730.44 keV γ -ray (0.0614(3)) was changed to 0.064(3); and the uncertainty of the 2133.04 keV γ -ray (0.0350(2)) was increased by a factor of 10 here (possible typographical errors).

³⁾ Statistical outlier, omitted.

[#] The LWM chose the unweighted average for these discrepant values.

EC + β^+ feeding to the ground state of ^{57}Co has not been observed. A systematic $lg ft \geq 11.0$ for a second forbidden non-unique transition corresponds to $I_{\text{EC}} \leq 0.01\%$ for a possible EC transition to the ground state of ^{57}Co . Thus, we used the sum of the relative emission probabilities of the 1224.00 keV, 1377.63 keV, 1757.55 keV, 1897.42 keV, 1919.52 keV, 2133.04 keV, 2730.91 keV, 2804.20 keV and 3177.28 keV γ -rays to normalize the decay scheme. The 1377.62 keV gamma ray is the strongest transition, for which we used a fractional uncertainty of 2%, suggested by 1992Bh05. Similarly, for the first excited state at 1224 keV, a possible EC + β^+ transition would have a systematic $lg ft \geq 12.6$, which corresponds to an intensity $I_{\text{EC}} \leq 0.001\%$. Conversion coefficients used in these calculations are those of Band *et al.* [1976Ba63].

3. Atomic Data

The X-ray and Auger electron emission probabilities given in section 3 are values calculated by using the computer program EMISSION [4], the electron capture probabilities from section 2.2, and atomic data from 1996Sc06.

4. Radiation Emission

4.1 Electron Emission

The emission probabilities of the Auger electrons have been calculated here using the adopted nuclear and atomic electron capture transition data, and the program EMISSION [4]. The emission probabilities of conversion electrons were calculated using the adopted γ -ray emission probabilities and conversion coefficients (section 2.2).

4.2 Photon Emission

The emission probabilities of X-rays were calculated using the adopted nuclear and atomic electron capture transition data, and the program EMISSION [4]. The evaluation of the gamma-ray emission probabilities was discussed in section 2.3.

Total Average Radiation Energy

The total released average radiation energy (electron capture, neutrinos, nuclear recoil, photons and electrons) in the EC + β^+ decay of ⁵⁷Ni (calculated by using the computer program RADLST [5]) is 3264(32) keV. This value agrees well with 3264.2(26) keV from mass differences (1995Au04), and thus confirms the quality and completeness of the decay scheme.

References

- [1] - P. F. Zweifel, *Capture-Positron Branching Ratios*, Phys. Rev. **96**, 1572 (1954)
- [2] - Eckart Schönfeld, Frank Chu and Edgardo Browne, *EC-Capture*, a computer program to calculate electron-capture probabilities to various atomic shells for allowed and first-forbidden non-unique transitions. PTB, LBNL, Dec. 23, 1997.
- [3] - *The Program LOGFT*, National Nuclear Data Center, September 3, 1993.
- [4] - Herbert Janßen and Eckart Schönfeld, *EMISSION*, a computer program to calculate X-ray and Auger electron emission probabilities. PTB, Nov. 18, 1997.
- [5] - *The Program RADLST*, Thomas W. Burrows, report BNL-NCS-52142, February 29, 1988.
- 1949Ma38 F. Maienschein and J. L. Meem, Jr., *Coincidence Experiments in ⁶⁵Ni, ⁵⁷Ni, ¹¹⁰Ag and ¹¹⁴In*. Phys. Rev. **76**, 899 (1949)
- 1950Fr10 G. Friedlander, M. L. Perlman, D. Alburger, A. W. Sunyar, *Measurement of Absolute Electron Capture Rates with an Application to the Decay of ⁵⁷Ni*. Phys. Rev. **80**, 30 (1950)
- 1956Ru45 G. Rudstam, *Spallation of Medium Weight Elements*. Thesis, University of Uppsala (1956)
- 1957Zw01 P. F. Zweifel, *Allowed Capture-Positron Branching Ratios*, Phys. Rev. **107**, 329 (1957)

Comments on evaluation

- 1958Ko60 J. Konijn, H. L. Hagedoorn, B. van Nooijen, *Further Study on the Decay of ⁵⁷Ni*. Physica 24, 129 (1958)
- 1961Ra06 L. A. Rayburn, *14.4 MeV (n,2n) Cross Sections*. Phys. Rev. 122, 168 (1961)
- 1962Ch20 G. Chilosi, S. Monaro, R. A. Ricci, *Excited Levels in ⁵⁷Co from the Decay of ⁵⁷Ni*. Nuovo Cimento 26, 440 (1962)
- 1964Ru06 G. Rudstam, *Peripheral Reactions in Copper Induced by 19 GeV Protons*. Nucl. Phys. 56, 593 (1964).
- 1965Eb01 T. G. Ebrey, P. R. Gray, Precision, *Half-Life Measurements of Fourteen Positron-Emitting Nuclei*. Nucl. Phys. 61, 479 (1965).
- 1966Pi01 C. J. Piluso, D. O. Wells, D. K. McDaniels, *The Decay of ⁵⁶Ni and ⁵⁷Ni*. Nucl. Phys. 77, 193(1966).
- 1967Li08 E. W. A. Lingeman, J. Konijn, F. Diederix, B. J. Meijer, *The Decay of ⁵⁷Ni*. Nucl. Phys. A100, 136 (1967).
- 1969Ga14 C. Gatrousis, R. A. Meyer, L. G. Mann, J. B. McGrory, *Single-Particle and Core-Coupled States in ⁵⁷Co from the Decay of ⁵⁷Ni*. Phys. Rev. 180, 1052 (1969).
- 1974Ro18 S. J. Rothman, N. L. Peterson, W. K. Chen, J. J. Hines, R. Bastar, L. C. Robinson, L. J. Nowicki, J. B. Anderson, *Half-Lives of Nine Radioisotopes*. Phys. Rev. C9, 2272 (1974).
- 1976Ba63 I. M. Band, M. B. Trzhaskovskaya, M. A. Listengarten, *Internal Conversion Coefficients for Atomic Numbers Z < 30*. At. Data Nucl. Data Tables 18, 433 (1976).
- 1977Ba48 W. Bambynek, H. Behrens, M. H. Chen, B. Crasemann, M. L. Fitzpatrick, K. W. D. Ledingham, H. Genz, M. Mutterer and R. L. Intemann, *Orbital Electron Capture by the Nucleus*, Rev. Mod. Phys. **49**, 77 (1977)
- 1982Gr10 A. Grutter, *Decay Data of ⁵⁵⁻⁵⁸Co, ⁵⁷Ni, ^{60,61}Cu and ^{62,63}Zn*. Int. J. Appl. Radiat. Isotop. 33, 533 (1982).
- 1986Di02 J. K. Dickens, *Half Life of ⁵⁷Ni*. J. Radioanal. Nucl. Chem. 103, 273 (1986).
- 1988WoZO M. J. Woods and A. S. Munster, *Evaluation of Half-Life Data*. NPL RS(EXT)95 (1988).
- 1990Sc23 A. M. S. Scardino, O. Helene, P. R. Pascholati, V. R. Vanin, *Beta Decay of ⁵⁷Ni*. Z. Phys. A336, 313 (1990).
- 1992Bh05 M. R. Bhat, *Nuclear Data Sheets Update for A = 57*. Nucl. Data Sheets 67, 195 (1992).
- 1995Au04 G. Audi and A. H. Wapstra, *The 1995 Update to the Atomic Mass Evaluation*. Nucl. Phys. **A595**, 409 (1995).
- 1996Sc06 E. Schönfeld and H. Janßen, *Evaluation of Atomic Shell Data*, Nucl. Instr. Meth. in Phys. Res. **A369**, 527 (1996).

⁵⁸Co - Comments on evaluation of decay data by M.M. Bé

A first evaluation was completed in November 1993 with minor editing in October 1998 and in March 2002. This new evaluation was completed in August 2013 in accordance with the DDEP rules. Various quantities have been updated (Q value, ICC...) and some references published to this date included.

Decay Scheme

A comprehensive study of the ⁵⁸Co decay was published by Bambynek and Legrand (1973Ba67). ⁵⁸Co ($T_{1/2} = 70,85$ d) decays 100 % by electron capture and/or β^+ disintegrations to the two first excited levels in ⁵⁸Fe. The electron capture probabilities to upper levels in ⁵⁸Fe nucleus, if they exist, are of very weak intensity, since no gamma radiations have been observed from these levels. Disintegration by β^- transition to ⁵⁸Ni has never been observed.

Q^{EC} value for ⁶⁸Co decay was adopted from Wang *et al.* (2012Wa38). The spins, parities and level energies are based on the mass-chain evaluation of Nesaraja *et al.* (2010Ne01).

The proposed decay scheme is consistent since $Q_{calc} = 2307,9$ (27) keV, deduced from the evaluated average energies of all emissions, is identical to $Q_{\beta^-} = 2307,9$ (11) from the 2012 atomic mass evaluation of M. Wang *et al.* (2012Wa38).

Half-life

The experimental half-life values used to calculate the recommended value are listed in the following table:

References	$T_{1/2}$ (d)	uc (d)	Comments
Schumann, 1956Sc87	71,3	0,2	
Decowski, 1968De08	71,54	0,75	
Werner, 1972We17	71,1	0,2	
Araminowicz, 1973ArZI	71,83	6,12	
Lagoutine, 1975La16	70,78	0,033	Original uc divided by 3
Vaninbroukx, 1976Va30	70,81	0,033	Original uc divided by 3
Houtermans, 1980Ho17	70,916	0,03	Original uc multiplied by 2
Hoppes, 1982HoZJ	70,75	0,07	Superseded by 2012Fi12
Fitzgerald, 2012Fi12	70,77	0,11	
Crit (χ^2)	2,6		
$\chi^2 / (n-1)$	2,7		
UWM	71,131		
WM	70,845		
uc(WM)int. :	0,018		
uc(WM)ext. :	0,029		
Adopted	70,85	0,03	

Values given by Crisler *et al.* (1972Cr02) have not been taken into account because they were obtained as a mean to control another experiment and not as a specific study.

Uncertainties given by 1975La16 and 1976Va30 were divided by three because they are for a 99,7 % confidence level. But the unrealistic uncertainty of 0,015 claimed by 1980Ho17 was multiplied by two, then giving to the 1980Ho17 result a relative weight of 36 % comparable to 30 % for 1975La16 and 1976Va30.

The set of data is just consistent, the weighted mean is preferred with an uncertainty which is the lowest experimental uncertainty.

Electron Capture and Positron Emission

- The electron capture probability to the 1674 keV level is deduced from the decay scheme balance.
- The electron capture and the β^+ emission probabilities to the 810 keV level are deduced from the level balance and from the theoretical EC/ β^+ ratio of 5,61 (8). The resulting $I\beta^+$ intensity and the adopted EC/ β^+ ratio are in good agreement with the measured values as shown in the table below.
- The lg ft values, the fractional atomic shell electron capture probabilities and the average β^+ energy are calculated with the LOGFT program.

Measured values of $I\beta^+$ and ϵ / β^+ to the 811 keV level:

Reference	$I\beta^+$ (%)	ϵ / β^+
Good, 1946Go01	14,5 (5)	
Cook, 1956Co70		5,9 (2)
Grace, 1956Gr**	13,0 (13)	
Konijn, 1958Ko64	14,77 (30)	5,67 (14)
Ramaswamy, 1961Ra13	15,1 (5)	5,49 (18)
Biryukov, 1966Bi13	15,2 (2)	
Bambynek, 1968Ba49	15,00 (5)	
Williams, 1970Wi14		5,76 (4)
Goodier, 1971GoYM		5,76 (13)
Sykora, 1992Sy**		5,68 (21)
Adopted	14,94 (16)	5,61 (8)

The deduced value 14,94 (16) % for the $\beta_{+0,1}$ transition implies a 511 keV gamma emission of 29,88 (32) % which is in agreement with the measured values : 29,2 (1969GuZV) and 29,5 (3) (1974HeYW).

- A weak β^+ transition to the ground state was observed by Daniel (1958Da03), using a double focusing β spectrometer, with intensity 6,3 (+8,0 -5,4) 10^{-4} %, which was symmetrized to 8 (7) 10^{-4} %, following the method given in 1993Au03.

Gamma Rays

- The gamma ray energies are from Helmer (2000He14).

Relative γ ray intensities

- The measured 864 keV gamma ray intensities, relative to the 810 keV line, are:

References	$I_{\text{gr}}(\%)$	uc (%)	Comments
Frauenfelder, 1956Fr17	1,6	0,5	Omitted
MacArthur, 1962Ma33	1,2	0,06	Omitted
Malmskog, 1964Ma09	1,38	0,04	Omitted
Hill, 1965Hi12	0,77	0,04	
Bambynek, 1968Ba49	0,645	0,015	Superseded by Denecke
Ritter, 1968Ri03	0,81	0,03	
Dyer, 1972DyZY	0,7	0,02	
Denecke, 1973Ba67	0,69	0,02	As quoted in 1973Ba67
Legrand, 1973Ba67	0,69	0,02	As quoted in 1973Ba67
Heath, 1974HeYW	0,74		Given in the γ spectra catalogue, no information
Grütter, 1982Gr10	0,682	0,017	
Crit. χ^2	3,0		
Reduced χ^2	3,6		
UWM :	0,724		
WM :	0,704		
uc(WM)int. :	0,009		
uc(WM)ext. :	0,017		
Adopted :	0,704	0,022	Expanded

The set of the six adopted values (1965Hi12, 1968Ri03, 1972DyZY, Denecke, Legrand, 1982Gr10) is discrepant. The weighted mean with an expanded uncertainty to cover the most precise value is adopted.

- The 1674 gamma-ray emission intensities, relative to the 810 keV line, are:

References	$I_{\text{gr}}(\%)$	uc (%)	Comments
Frauenfelder, 1956Fr17	0,5	0,1	Omitted
Mac Arthur, 1962Ma33	0,66	0,05	Omitted
Malmskog, 1964Ma09	0,61	0,05	Omitted
Hill, 1965Hi12	0,68	0,05	
Bambynek, 1968Ba49	0,506	0,015	Superseded by Denecke
Ritter, 1968Ri03	0,57	0,03	
Dyer, 1972DyZY	0,49	0,03	
Denecke, 1973Ba67	0,527	0,015	As quoted in 1973Ba67
Legrand, 1973Ba67	0,525	0,013	As quoted in 1973Ba67
Heath, 1974HeYW	0,54		Given in the γ spectra catalogue, no information
Grütter, 1982Gr10	0,511	0,015	
Crit. χ^2	2,5		
Reduced χ^2	3,0		
UWM :	0,564		
WM :	0,531		
uc(WM)int. :	0,007		
uc(WM)ext. :	0,013		
Adopted :	0,531	0,013	

The set of the six adopted values (1965Hi12, 1968Ri03, 1972DyZY, Denecke, Legrand, 1982Gr10) is discrepant. The weighted mean with the external uncertainty is adopted.

Internal Conversion Coefficients

The γ transitions with energies 810 and 1674 keV respectively are of pure E2 character. The 864 keV transition is a M1 + E2 mixture.

Measured mixing ratio, δ , for the 864 keV transition:

References	δ	uc	Comments
1956Fr17	2,20	0,30	Co-58 decay, NaI-NaI, omitted
1962Ma33	1,5	+0,4 -0,5	Co-58 decay, NaI-NaI, omitted
1964Ma09	1,6	0,2	Co-58 decay, NaI-NaI, omitted
1966Ra22	1,56	0,04	Co-58 decay, NaI-NaI, omitted
1969Si13	1,46	0,07	Co-58 decay, NaI-NaI, omitted
1969Sc11	1,10	0,20	Co-58 decay, NaI-NaI, omitted
1969Fa05	0,57	0,06	Fe-57(n, γ), NaI – Ge(Li)
1971Si15	0,67	0,15	Co-58 decay, NaI – Ge(Li), original value: 0,61(+0,19 -0,10)
1972Fo05	0,94	+0,09 -0,08	Co-58 decay, NaI-NaI, omitted
1972Fo05	1,0	+0,6 -0,2	Co-58 decay, NaI – Ge(Li), omitted
1972Fo05	0,69	0,05	Co-58 decay, Ge(Li)
Adopted	0,643	0,041	

The adopted value is the weighted mean of the three results measured with a Ge(Li) detector: 1969Fa05, 1971Si15, and the last published value of 1972Fo05.

The result of 1971Si15 has been symmetrized following the method given in 1993Au03.

The adopted ICC(s) are the theoretical values interpolated by the BrIcc program (2008Ki07) from the tables of Band *et al.* (2002Ba85).

Absolute γ ray intensities

Considering the decay scheme, the β^+ transition (0,0008 %) being negligible, the sum of the two γ transitions populating the ground state must be equal to 100 %:

$$\sum_i I_{\gamma i} [1 + \alpha_{Ti}] = \frac{100 \%}{N}$$

where:

$I_{\gamma i}$ is the relative emission intensity of the 810 and 1674 keV rays and, α_{Ti} the related total internal conversion coefficient. N is a normalisation factor between the relative γ intensities and the absolute intensities.

$N = 0,9944$ (2).

Atomic Data

The adopted ω_K value of 0,355 (4) (1996Sc06) is in agreement with (0,352 (4)) from the measurement of Solé (1992So04).

The X-ray and Auger electron emission intensities are derived from the decay scheme parameters by using the program EMISSION.

The calculated K X-ray intensity of 26,8 (4) % is in good agreement with the measured K X-ray intensity of 25,96 (10) % (1968Ba49).

References

- 1946Go01 W.M.Good, D.Peaslee, M.Deutsch. Phys. Rev. 69 (1946) 313. Beta plus emission probability
- 1956Sc87 R.P.Schuman, M.E.Jones, A.C.McWherter. J. Inorg. Nucl. Chem. 3 (1956) 160. Half-life
- 1956Co70 C.S.Cook, F.M.Tomnovec. Phys. Rev. 104 (1956) 1407. Beta plus emission probabilities
- 1956Fr17 H.Frauenfelder. Phys. Rev. 103 (1956) 352. Gamma-ray emission probabilities, δ
- 1956Gr** M.A.Grace, G.A.Jones, J.O.Newton. Philos. Mag. 1 (1956) 363. Beta plus emission probability
- 1958Da03 H.Daniel. Z. Phys. 150 (1958) 144. Beta emission probabilities
- 1958Ko64 J.Konijn, H.L.Hagedoorn, H.Van Krugten, J.Slobben. Physica 24 (1958) 931. Beta plus emission probabilities
- 1961Ra13 M.K.Ramaswamy. Indian J. Phys. 35 (1961) 610. Beta plus emission probabilities
- 1962Ma33 D.MacArthur, R.Goodman, A.Artna, M.W.Johns. Nucl. Phys. 38 (1962) 106. Gamma-ray emission probabilities, δ
- 1964Ma09 S.Malmskog. Nucl. Phys. 51 (1964) 690. Gamma-ray emission probabilities
- 1965Hi12 M.W.Hill. Report BNWL-SA-315 (1965). Gamma-ray emission probabilities
- 1966Ra22 R.V.Rama Mohan, K.V.Reddy, B.B.V.Raju, S.Jnanananda. Indian J.Pure Appl.Phys. 4 (1966) 420. Mixing ratio
- 1966Bi13 E.I.Biryukov, E.G.Zaletskii, N.S.Shimanskaya. Bull. Acad. Sci. USSR 30 (1967) 514. Beta plus emission probabilities
- 1968Ba49 W.Bambynek, E.De Roost, E.Funck. Proceeding of the Conference on Electron Capture and Higher Order Processes in Nuclear Dec. (Budapest) (1968) 253. Gamma-ray emission probabilities, Beta+ emission probabilities, Elec. Capture probabilities
- 1968De08 P.Decowski, W.Grochulski, A.Marcinkowski, K.Siwiek, I.Sledzinska, Z.Wilhelmi. Nucl. Phys. A112 (1968) 513. Half-life
- 1968Ri03 J.C.Ritter, R.E.Larson, J.I.Hoover. Nucl. Phys. A110 (1968) 463. Gamma-ray emission probabilities
- 1969Sc11 U.Schötzig, H.Schrader, R.Stippler, F.Munnich. Z. Physik 222 (1969) 479. Mixing ratio
- 1969Si13 V.Singh, P.N.Tandon, S.H.Devare, H.G.Devare. Nucl. Phys. A137 (1969) 278. Mixing ratio
- 1969GuZV R.Gunnink, J.B.Niday, R.P.Anderson, R.A.Meyer. Report UCID-15439 (1969) . Gamma-ray emission probabilities
- 1969Fa05 U.Fanger, W.Michaelis, H.Schmidt, H.Ottmar. Nucl. Phys. A128 (1969) 641. Mixing ratio
- 1970Wi14 A.Williams. Nucl. Phys. A153 (1970) 665. Beta plus emission probabilities
- 1971Si15 N.C.Singhal, A.V.Ramayya, J.H.Hamilton, S.Raman. Z. Physik 245 (1971) 50. Mixing ratio
- 1971GoYM I.W.Goodier, M.J.Woods, A.Williams. Proc. Int. Conf. Chemical Nuc. Data, Canterbury, M.L. Hurrell Ed. (1971) 175. Beta plus emission probabilities
- 1972Cr02 D.F.Crisler, H.B.Eldridge, R.Kunselman, C.S.Zaidins. Phys. Rev. C5 (1972) 419. Half-life
- 1972DyZV N.C.Dyer, A.C.Rester, W.Croft, J.H.Hamilton. Proc. Int. Conf. Radioactivity in Nucl. Spectrosc., Nashville, Tenn. (1972) 1207. Gamma-ray energies and emission probabilities
- 1972Fo05 R.A.Fox, W.D.Hamilton, M.J.Holmes. Phys. Rev. C5 (1972) 853. Mixing ratio
- 1972We17 R.Werner, D.C.Santry. J. Nucl. Energy 26 (1972) 403. Half-life
- 1973Ba67 W.Bambynek, J.Legrand. Atomic Energy Rev. 11 (1973) 524. Gamma-ray energies and emission probabilities
- 1973ArZI J.Araminowicz, J.Dresler. Report INR-1464 (1973) 14. Half-life
- 1974HeYW R.L.Heath . Aerojet Nucl. Co. Report ANCR-1000-2 (1977). Gamma-ray energies and emission probabilities
- 1975La16 F.Lagoutine, F.Legrand, C.Bac. Int. J. Appl. Radiat. Isotop. 26 (1975) 131. Half-life
- 1976Va30 R.Vaninbrouck, G.Grosse. Int. J. Appl. Radiat. Isotop. 27 (1976) 727. Half-life
- 1980Ho17 H.Houtermans, O.Milosevic, F.Reichel. Int. J. Appl. Radiat. Isotop. 31 (1980) 153. Half-life (Pb-203).
- 1982Gr10 A.Grütter. Int. J. Appl. Radiat. Isotop. 33 (1982) 533. Gamma-ray energies and emission probabilities
- 1982HoZJ D.D. Hoppes, J.M.R.Hutchinson, F.J.Schima, M.P.Unterweger. NBS Special publication 626 (1982) 85, Half-life

- 1992Sy** I.Sykora. Rare Nuclear Processes: Proceedings of the 14th Europhysics Conference on Nuclear Physics, Bratislava, Czecho-Slovakia, 22-26 Oct. 1990 (1992) 141. Beta plus emission probabilities
- 1992So04 V.A.Solé. Nucl. Instrum. Methods A312 (1992) 303. K fluorescence yield
- 1996Sc06 E.Schönfeld, H.Janssen. Nucl. Instrum. Methods A369 (1996) 527. Atomic Data
- 2000He14 R.G.Helmer, C.Van der Leun. Nucl. Instrum. Methods A450 (2000) 35. Gamma-ray energies
- 2002Ba85 I.M.Band, M.B.Trzhaskovskaya, C.W.Nestor Jr., P.O.Tikkanen, S.Raman. At. Data Nucl. Data Tables 81 (2002) 1. Theoretical ICC
- 2003Au03 G.Audi, A.H.Wapstra, C.Thibault. Nucl. Phys. A729 (2003) 21
- 2008Ki07 T.Kibédi, T.W.Burrows, M.B.Trzhaskovskaya, P.M.Davidson, C.W.Nestor Jr. Nucl. Instrum. Methods Phys. Res. A589 (2008) 202. ICC
- 2010Ne01 C.D.Nesaraja, S.D.Geraedts, B.Singh. Nucl. Data Sheets 111 (2010) 897. Spin and Parity
- 2012Wa38 M.Wang, G.Audi, A.H.Wapstra, F.G.Kondev, M.MacCormick, X.Xu, B.Pfeiffer. Chinese Physics C36 (2012) 1603. Q
- 2012Fi12 R.Fitzgerald. J.Res.Natl.Inst.Stand.Technol. 117 (2012) 80. Half-life

⁵⁹Fe – Comments on evaluation of decay data by M.M. Bé and V. Chisté

1. Decay scheme

This decay scheme was well studied (Bérényi, Béraud, Collin, Ferguson, Heath, Pancholi, Metzger, Raman, etc.) so that the existence of beta transitions and the spin and parity of the ⁵⁹Co levels are clearly established. Some authors (Mukerji, Raman) carried out experiments in order to measure the weak β - branches. No clear evidence of a β -branching to the 1190 keV level was found, if this transition exists its branching ratio has an upper limit of 1×10^{-4} .

2. Nuclear Data

⁵⁹Fe half-life (in days)

Author	NSR	Value	Uc	Method
Metzger	52Me53	45.0	3.0	NaI
Keene	58Ke26	44.56	0.03	ionisation chamber
Pierroux	59Pi43	45.60	0.08	Electrometer à lames vibrantes
Fuschini	60Fu03	63.1	0.8	
Heath	60He06	45.0	5.0	NaI
Subba Rao	60Su10	46.5	1.0	
Wortman	63Wo01	45.0	3.0	
Emery	72Em01	44.5	0.2	NaI
Visser	73Vi13	44.75	0.04	NaI (s x 3)
Alstad	75Al02	45.3	0.3	Gas flow proportional counter
Houtermans	80Ho17	44.496	0.007	4 π - γ
Walz	83Wa26	44.53	0.07	4 π - γ ionisation chamber
Unterweger	92Un01	44.5074	0.0072	
Martin	97Ma75	44.472	0.008	4 π - γ ionisation chamber

The value from Fuschini was omitted due to its large deviation from the others.

The values from Subba Rao, Pierroux were rejected as outlier (Chauvenet' s criteria).

With this set of eleven remaining values, the reduced χ^2 is 6.4 and the Lweight program recommends the unweighted mean and expanded the uncertainty : 44.74 ± 0.24 .

With these eleven values the weighted mean and the external uncertainty are : 44.498 ± 0.011 .

Taking into account the most precise values (Keene, Visser, Houtermans, Walz, Unterweger and Martin) :

- the value from Visser was rejected as outlier;

- then the reduced $\chi^2 = 4$;

- the weighted mean is 44.495 with an external uncertainty of 0.008.

Regarding the fact that the four more recent measurements are compatible with this value and (for three of them) have a similar uncertainty, the recommended value is :

44.495 ± 0.008 d

Half-lives of ⁵⁹Co excited levels

Level 1100 keV

- Sidhu : ≤ 50 ps
- Béraud < 14 ps

Level 1291 keV (in ns)

Author	NSR	Value	Uc
Sidhu :	67Si01	0.60	0.05
Agarwal :	67Ag03	0.59	0.02
Béraud :	67Be60	0.575	0.011
Garg :	72Ga39	0.538	0.004
Green :	72Gr05	0.564	0.020
Arens :	71Ar07	0.564	0.005

The value from Chauhan (0.516 (6)) was not taken into account : it seems that the experiment is the same as those described in Garg *et al.*

For the six values above the reduced χ^2 is 5.45 and the critical $\chi^2 = 3$. Then, the uncertainty on the value given by Garg was increased by 1.08 in order to reduce its relative weight to 50 %. The reduced χ^2 is 5.10. This set of value is not consistent and the unweighted mean is adopted : **0.572 (34) ns**.

Level 1434 keV

Arens : 210 (20) ps

2.1 Beta Transitions

Beta transition energies

The adopted Q-value 1565.2 (6) keV is from Audi and Wapstra. It was determined from the measurements of Wortman and Metzger (see Table below)

The adopted energies and uncertainties of beta transitions are deduced from the Q-value and the levels energies and their uncertainties.

Measured beta energies are summarized in the following table :

keV	1565	475	273	132	85
Wortman	1573 \pm 3	475 \pm 3	273 \pm 5		
Berenyi		455 \pm 5	275 \pm 5		
Metzger	1560 \pm 8	462 \pm 3	271 \pm 3		
Mukerji	1566			132	85
Subba Rao	1580 \pm 20	470 \pm 6	280 \pm 6	150 \pm 10	
Raman	1575 \pm 20	461 \pm 10	268 \pm 10	128	80
<i>Evaluated</i>	1572 \pm 3	463.4 \pm 2.2	273.0 \pm 2.1	137 \pm 8	82.5 \pm 2.5
Adopted	1565.2 \pm 0.6	465.9 \pm 0.6	273.6 \pm 0.6	130.9 \pm 0.6	83.6 \pm 0.6

The 1565 keV transition is second forbidden non unique, with the shape factor given by Wortman (see below) the mean energy is 521 keV ; with the shape factor from Raman the mean energy is 584 keV ; these calculations were done with the SPEBETA program. In the Russian book Kolobachkin *et al.* the mean energy was calculated to be 523 keV.

Expecting a confirmation, the adopted value is 522 (2) keV.

Beta transition probabilities

The emission probabilities are calculated from gamma transition probability imbalance on each level. That was done for all the transitions, except for the weak 1565-keV to the ground state, the resulting values are in agreement with the experimental values (see Table below).

Taking into account the consistency of the decay scheme :

- the sum of all the transitions to the Co -59 ground state must be equal to 100 ; this leads to an intensity value of 0.12 (32) for the 1565 keV transition. This important uncertainty comes from the propagation of the uncertainties on the gamma transitions.
- the sum of all the beta transitions leaving from Fe-59 must be equal to 100 ; this gives a value of 0.13 (34) for the 1565 transition.

However, several authors measured this transition intensity and found values from 0.18 (4) % to 0.3 (1) % (Table below).

It must also be pointed out that the authors gave measured gamma emission probabilities after corrections, with a value of the $I_{\beta}(gs)$ taken as :

- 0.3% by Legrand, Béraud, Pancholi ;
- 0.18% by Miyahara.

From the previous remarks, it follows that the $I_{\beta}(gs)$ intensity is certainly greater than 0.10% (decay scheme) and less than 0.40% (experiments).

The adopted value is then : 0.25 (15) %.

Table : Measured I_{β}

Metzger (52Me53)			
1573 keV	$I_{\beta} = 0.3 (1)\%$	$\lg ft = 10.9$	
475 keV	$I_{\beta} = 54.8 (20)\%$	$\lg ft = 6.7$	
273 keV	$I_{\beta} = 44.9 (20)\%$	$\lg ft = 5.9$	
Wortman. (63Wo01) (No uncertainty given)			
1573 keV	$I_{\beta} = 0.30\%$	$\lg ft = 10.96$	shape factor $p^2 + 3.3 q^2$
475 keV	$I_{\beta} = 51.2\%$	$\lg ft = 6.74$	
273 keV	$I_{\beta} = 48.5\%$	$\lg ft = 5.92$	
Raman (74Ra13)			
1573 keV	$I_{\beta} = 0.18 (4)\%$	$\lg ft = 11.15 \pm 0.11$	shape factor $p^2 + 1.7 q^2$
475 keV	$I_{\beta} = 51 (3)\%$		
273 keV	$I_{\beta} = 47 (4)\%$		
(80-128)	$I_{\beta} = (1.4)\%$		
Berényi (60Be06) (No uncertainty given)			
1573 keV	$I_{\beta} < 0.5 \%$		
475 keV	$I_{\beta} = 55.4\%$	$\lg ft = 6.1$	
273 keV	$I_{\beta} = 44.6\%$	$\lg ft = 5.3$	

β - γ circular polarization asymmetry coefficients

Behrens (70BeZx) recommends :

For 466 β - 1099 γ : A= -0.164 (7)

For 273 β - 1292 γ : A= -0.15 (2)

2.2 Gamma transitions and internal conversion coefficients

1291 keV transition

Assuming a pure E2 transition, the theoretical ICC (from Band's tables) $a_T = 1.22 \cdot 10^{-4}$ is consistent with the experimental one from Metzger (52Me53) $\alpha_T = 1.35 (6) \cdot 10^{-4}$.

Other measurements :

Metzger (52Me53), $\alpha_K = 1.19 (6) \cdot 10^{-4}$

Hinman (53Hi02), $\alpha_T = 1.06 (16) \cdot 10^{-4}$

Collin (64Co34), $\alpha_T = 1.07 (8) \cdot 10^{-4}$

K.S.Krane *et al.* (1976Kr10) suggests a M3/E2 mixture of $\delta = -0.033 (30)$, that does not change the ICC value significantly.

1099 keV transition

Assuming a pure E2 transition, the theoretical ICC (from Band's tables) $a_T = 1.75 \cdot 10^{-4}$ is consistent with the experimental one from Metzger (52Me53) $\alpha_T = 1.87 (7) \cdot 10^{-4}$.

Other measurements :

Metzger(52Me53), $\alpha_K = 1.35 (6) \cdot 10^{-4}$

Hinman(53Hi02), $\alpha_T = 1.84 (27) \cdot 10^{-4}$

Collin(64Co34), $\alpha_T = 1.36 (10) \cdot 10^{-4}$

334 keV transition E2/M1

The measured values of the mixing ratio are the following :

Author	Delta
Pancholi	- 0.12 (6)
Eriksson	- 0.12 (4)
Arens	+ 0.05 + 0.03 - 0.07 or - 1.8 + 0.4 - 0.6
Adopted value	- 0.12 (6)
ICC (Band)	0.002 (1)

142 keV transition E2/M1

The measured values of the mixing ratio are the following :

Author	Delta
Pancholi	- 0.15 (6) < δ < 0.026
Eriksson	- 0.006 (12)
Arens	0.028 + 0.009 - 0.014 or - 1.78 + 0.15 - 0.20
Adopted value (from Krane 1977Kr13)	- 0.008 (7)
ICC (Band)	0.0160 (1)

192 keV transition E2/M1

The measured values of the mixing ratio are the following :

Author	Delta
Pancholi	- 0.22 (2)
Eriksson	0.21 (2)
Arens	- 0.21 (2) or $\delta > 14$
Bajaj	0.22 (2)
Collin	- 0.296 (23)
Adopted value	0.21 (1)
ICC (Band)	0.00899 (15)

Gamma emissionsGamma emission energies

The gamma emission energy of the following lines are from Helmer (2000He14) :

142.651 ± 0.002

192.349 ± 0.005

1099.245 ± 0.003

1291.590 ± 0.006

Others are from Pancholi.

Gamma emission intensities

Eight published papers describe measurements of the gamma emission intensities, all the values are given in absolute values.

Heath *et al.* do not give uncertainty, therefore these values are omitted.

The results given by Béraud *et al.* are with uncertainties of the order of 10%, they are not omitted but their relative weight is generally weak, as well as those of the values given by Mukerji *et al.*

J.Legrand *et al.* (70Le03), carried out β - γ coincidences measurements and deduced absolute values, assuming that the β branching to the ground state is 0.3%. The uncertainty adopted by Legrand is the sum of the statistical uncertainty assessed at 3σ and the systematic uncertainty at 1σ ; consequently, the standard deviation cannot be obtained dividing the original uncertainty by 3 and we divided the given uncertainties by 2 only.

Pancholi *et al.* (73Pa18), measured the relative values and normalized them such as $I(1099 + 1292 + 1481) = 99.7\%$, assuming $I\beta(gs) = 0.3\%$.

Miyahara *et al.* (1989Mi07), carried out activity measurements and deduced absolute values. This paper is the most recent one and gives the most precise values which contribute more than 50% in the adopted result for the two intense lines : 1099 and 1291 keV.

The following table summarizes all the values taken into account and the adopted results.

These different set of data are consistent, except for the original set of seven data for the 335 keV line where two values are outliers and are omitted (o). The adopted values are the weighted means.

keV	142	192	335	381
Mukerji	1.1 ± 0.16	3.3 ± 0.3	0.27 ± 0.03	
Legrand	0.98 ± 0.02	2.95 ± 0.04	0.24 ± 0.02	0.023 ± 0.002
Béraud	0.79 ± 0.8	2.50 ± 0.25	0.25 ± 0.05	0.022 ± 0.005
Collin	0.8 ± 0.2	2.8 ± 0.3	0.7 ± 0.3 ^(o)	
Miyahara	0.955 ± 0.030	2.851 ± 0.048	0.262 ± 0.016	
Ferguson	0.85 ± 0.15	2.4 ± 0.4	0.34 ± 0.07 ^(o)	
Pancholi	1.02 ± 0.04	3.08 ± 0.1	0.27 ± 0.01	0.018 ± 0.003
$\chi^{**2}/N-1(\text{critical})$	1.5 (2.8)	1.9 (2.8)	0.5	0.97
Adopted value	0.972 ± 0.015	2.918 ± 0.029	0.264 ± 0.007	0.0215 ± 0.0016

keV	1099	1291	1481
Mukerji	57.5 ± 3	42.4 ± 2.3	0.052 ± 0.006
Legrand	55.5 ± 0.8	44.1 ± 0.6	0.09 ± 0.01
Béraud	56.2 ± 5.6	43.5 ± 4.3	0.056 ± 0.012
Collin	56.5 ± 1.5	43.2 ± 1.5	
Miyahara	56.68 ± 0.22	42.99 ± 0.30	
Ferguson	56 ± 3	44 ± 3	
Pancholi	56.5 ± 1.5	43.2 ± 1.1	0.059 ± 0.006
$\chi^{**2}/N-1(\text{critical})$	0.4	0.5	3.6 (3.8)
Adopted value	56.59 ± 0.21	43.21 ± 0.25	0.0603 ± 0.0037

Angular correlation coefficients

Several authors determined the angular correlation coefficients. Some of them are summarized here as a matter of interest.

$$192\gamma - 1099\gamma, \frac{3}{2} \sqrt{(M1+E2)3/2} \sqrt{(E2)7/2} :$$

Author	NSR	A2	uc	A4	uc
Heath	60He06	0.024	0.005		
Rao	70Ra00	0.028	0.003	0.008	0.007
Arens	71Ar07	0.008	0.007		
Bajaj	72Ba**	0.008	0.004	0.004	0.008
Eriksson	73Er11	0.011	0.004	-0.003	0.004

$$335\gamma - 1099\gamma, \frac{1}{2} \sqrt{(M1+E2)3/2} \sqrt{(E2)7/2} :$$

Author	A2	uc	A4	uc
Rao	-0.043	0.003	-0.004	0.003
Arens	-0.064	0.011	-0.008	0.025
Eriksson	-0.040	0.010	-0.006	0.0006
Bajaj	-0.099	0.012		

$143\gamma - 1292\gamma, 1/2^-(M1+E2)3/2^-(E2)7/2^- :$

Author	A2	uc	A4	uc
Heath	- 0.069	0.005		
Rao	- 0.065	0.004	- 0.006	0.005
Arens	- 0.065	0.004		
Bajaj	- 0.070	0.005	0.014	0.015
Subrahmanyam	- 0.09	0.01		
Eriksson	- 0.070	0.003		

Conversion electrons

Conversion electron intensities were calculated from the gamma transition probabilities and the internal conversion coefficients.

Hinman(53Hi02) gives the ratio of the number of conversion electrons from the 1099 keV transition to the number of conversion electrons from the 1291 keV transition, to be equal to 1.91 (9).

There is a good agreement with the ratio (1.87) obtained from the calculated values in this evaluation.

References

- F. R. Metzger ; Phys. Rev. ; 88 (1952) 1360 ; T ICC, Half-life
 G. HINMAN ; D. BROWER, R. LEAMER ; Phys. Rev. ; 90 (1953) 370A ; T ICC
 J. P. KEENE ; L. A. MACKENZIE, C. W. GILBERT ; Phys. in Med. Biol. ; 2 (1958) 360 ; Half-life
 J. M. Ferguson ; Nucl. Phys. ; 12 (1959) 579 ; Gamma-ray emission probabilities
 A. Pierroux ; G. Guében, J. Govaerts ; Bull. Soc. Royale Sci. Liège ; 28, 7-8 (1959) 180 ; Half-life
 E. Fuschini et al ; Nuovo Cim. ; XVI,5 (1960) 1910 ; Half-life
 R. L. Heath ; C. W. Reich, D. G. Proctor ; Phys. Rev. ; 118,4 (1960) 1082 ; Half-life
 B. N. SUBBA RAO ; Proceeding of the Indian Acad. of Sciences ; 3A (1960) 130 ; Half-life
 D. E. Wortman ; L. M. Langer ; Phys. Rev. ; 131,1 (1963) 325 ; Half-life
 W. Collin et al ; Z. Physik ; 180 (1964) 143 ; Gamma-ray emission probabilities, mixing ratio
 Y. K. Agarwal ; C. V. K. Baba, S. K. Bhattacharjee ; Nucl. Phys. ; A99 (1967) 457 ; 1291 keV level half-life
 R. Béraud et al ; Comp. Rend. Acad. Sci. (Paris) ; 265-B (1967) 1354 ; Gamma-ray emission probabilities
 N. P. S. Sidhu ; U. C. Gupta ; Nucl. Phys. ; A91 (1967) 557 ; 1291 keV level half-life
 J. LEGRAND ; J. MOREL, C. CLEMENT ; Nucl. Phys. ; A142 (1970) 63 ; Gamma -ray emission probabilities
 I. ARENS ; H. J. KORNER ; Z. Phys. ; 242 (1971) 138 ; Mixing Ratio, Half-life
 M. I. Green ; P. F. Kenealy, G. B. Beard ; Nucl. Instrum. Methods ; 99 (1972) 445 ; 1291 keV level half-life
 R. K. Garg et al ; Z. Physik ; 257 (1972) 124 ; 1291 keV level half-life
 J. F. Emery ; S. A. Reynolds, E. I. Wyatt ; Nucl. Sci. Eng. ; 48 (1972) 319 ; Half-life
 S. C. PANCHOLI ; J. J. PINAJIAN, N. R. JOHNSON, A. KUMAR, S. K. SONI, M. M. BAJAJ, S. L. GUPTA, N. L. SAHA ; Phys. Rev. ; C8 (1973) 2277 ; Gamma-ray emission probabilities, Spin and Parity, Mixing Ratio
 C. J. Visser et al ; Agrochemophysica ; 5 (1973) 15 ; Half-life
 L. Eriksson ; L. Gidefeldt ; Physica Scripta ; 7 (1973) 169 ; Mixing ratio
 A. Mukerji ; D. Palazzo, J. D. Ullman ; Phys. Rev. ; C10,2 (1974) 949 ; Gamma-ray emission probabilities
 M. M. BAJAJ ; A. KUMAR, S. K. SONI, S. C. Pancholi, S. L. GUPTA, N. K. SAHA ; Proceeding of the nuclear physics and solid state physics symposium ; 14B,2 (1974) 375 ; Mixing ratio
 S. Raman et al ; Phys. Rev. ; C9,6 (1974) 2463 ; Beta emission probabilities and energies
 J. Alstad ; I. R. Haldorsen, A. C. Pappas, M. Skarestad ; J. Inorg. Nucl. Chem. ; 37 (1975) 873 ; Half-life
 K. S. Krane ; S. S. Roseblum, W. A. Steyert ; Phys. Rev. ; C 14,2 (1976) 653 ; 1291 keV Mixing ratio

K. S. Krane ; At. Data Nucl. Data Tables (19 (1977) 363 ; 142 keV Mixing ratio
H. HOUTERMANS ; O. MILOSEVIC, F. REICHEL ; Int. J. Appl. Radiat. Isotop. ; 31 (1980) 153 ; Half -
life
K. F. WALZ ; H. M. WEISS, K. DEBERTIN ; Int. J. Appl. Radiat. Isotop. ; 34 (1983) 1191 ; Half-life
H. Miyahara et al ; Appl. Rad. Isotopes ; 40,4 (1989) 343 ; Gamma-ray emission probabilities
G. Audi ; A. H. Wapstra ; Nucl. Phys. ; A595 (1995) 409 ; Q
R. H. Martin ; K. I. W. Burns, J. G. V. Taylor ; Nucl. Instrum. Methods ; A390 (1997) 267 ; Half-life
R. G. Helmer ; C. van der Leun ; Nucl. Instrum. Methods ; A450 (2000) 35 ; Gamma ray energies

Additional references

SPEBETA, a program to calculate beta spectra. P. Cassette. Note LPRI/92/307. CEA -LNHB, F-91191 Gif-sur-Yvette
Kolobachkin *et al.* Moscow Atomizdat, UDK 539 163(031), in Russian (1978)
M. M. Bajaj, A. KUMAR, S. K. SONI, S. C. Pancholi, S. L. GUPTA, N. K. SAHA. Report B.A.R.C. – 614(1972)41. Angular correlation coefficients. (72Ba**)
D. Berényi, G. Y. Mathé, T. Scharbert. Nucl. Phys. 14(1960)459. Beta emission energies and probabilities. (60Be06)
H. Behrens. Kerforschungszentrum Karlsruhe KFK 1249(1970)1. β - γ Asymmetry coefficients. (70BeZX)
S. D. Chauhan *et al.* Proceeding of the nuclear physics and solid state physics symposium 14B -2 (1972). 1291 keV level half-life. (72Ch**)
K. V. Ramana Rao *et al.* Proceeding of the nuclear physics and solid state physics symposium V2 (1970) 387. Angular correlation coefficients. (70****)
V. Subrahmanyam. J. Inorg. Nucl. Chem. 34(1972)3319. Angular correlation coefficients. (72Su06)

⁵⁹Ni - Comments on evaluation of decay data
M. Galán

1) Decay Scheme

⁵⁹Ni disintegrates mainly by electron capture (2nd forbidden non-unique) to the ground state of ⁵⁹Co. ⁵⁹Co ground state has $J_{\pi} = 7/2^{-}$ (2002BA42).

2) Nuclear Data

The Q value is from new values of 2009AuZZ: $Q_{\epsilon} = 1072,76$ (19) keV. Other: 1075,1 (13) keV (1976BE02).

Only one direct measurement has been performed for the ⁵⁹Ni half-life. The value of $7,6$ (0,5) $\times 10^4$ years given by 1981Ni08 from two samples by absolute activity measurement has been adopted as the recommended value.

Some other indirect measurements from nuclear reactions have been performed.

The measured ⁵⁹Ni half-life values, in years, are:

Reference ^a	Value (a) ($\times 10^4$)	Procedure	Comments
2008WAZW	9,7 (0,7)	⁶⁰ Ni(n,2n) ⁵⁹ Ni reaction	Corrected 1994Ru19 result with newest σ_{thermal} and K_{α} yield. E(n) = 17-19 MeV
1994RU19	10,8 (1,3)	⁵⁸ Ni(n, γ) ⁵⁹ Ni and ⁵⁴ Fe(n, γ) ⁵⁵ Fe reactions	E(n) = 14.8 MeV
1991NO08	29 (10)	⁶⁰ Ni(n,2n) ⁵⁹ Ni reaction	Authors used $\sigma = 104$ (25) mbarn measured by 1988BO31 (very poor statistics)
1981NI08	7,6 (0,5)	Absolute activity measurement	Recommended value
1956SA32	10,0 (2,5)	⁵⁸ Ni(n, γ) ⁵⁹ Ni reaction	E(n)= thermal. $T_{1/2}$ was not the purpose of the work
1951BR05	7,5 (1,3)	⁵⁸ Ni(n, γ) ⁵⁹ Ni and ⁵⁹ Co(n, γ) ⁶⁰ Co reactions	E(n)= thermal. Data used for these reactions $\sigma = 4.2$ and $\sigma = 34.5$ respectively
1951WI14	75	⁵⁸ Ni(n, γ) ⁵⁹ Ni reaction	No uncertainty given

^aEvaluator used the NSR (Nuclear Science References, Brookhaven Lab.) keynumbers.

The only absolute activity measurement has been performed by 1981NI08. This value was deduced from 2 samples prepared by means of the reaction ⁵⁹Co(p,n)⁵⁹Ni, purified by ion-exchange columns in order to remove ⁵⁸Co activity. 6,93 keV Co K_{α} X-rays were measured by a Xe filled proportional counter. The value of $7,6$ (5) $\times 10^4$ years estimated by 1981NI08 has been adopted as the recommended value as it comes from a direct measurement.

The five other measurements were all made in a very similar way. After neutron irradiation ⁵⁹Ni atoms are counted by mass spectrometry. Purification is performed to avoid iron or cobalt impurities. The induced activity in the samples has been counted via the Co K X-ray after the ⁵⁹Ni K capture.

2008WAZW corrected the half-life result given by 1994RU19 using the recently published thermal cross-section of ⁵⁸Ni (2004RA23) of 4,13 (5) barns instead of the old value of 4,6 (3) barns from 1981MUZQ. The ⁵⁹Ni half-life is reduced about 10 % but still higher than the recommended value. 2008WAZW analyzed the contribution to the uncertainty of the ⁵⁹Ni half-life mainly from the uncertainties in the cross-sections.

1991NO08 used the cross-section of ⁶⁰Ni(n,2n) measured by 1988BO30 but this value was obtained with very poor statistics (as mentioned by authors of 1991NO08).

The oldest value (with uncertainty) given by 1951BR05 was estimated using thermal-neutron cross-sections for ⁵⁸Ni(n,γ) and ⁵⁹Co(n,γ) of 4,17 and 34,5 barns respectively.

If a statistical analysis is made for the five indirect measurements, the Lweight code rejects 1991NO08 datum based on Chauvenet's criterion. For the other four values, the weighted mean with external uncertainty is $9,5 (6) \times 10^4$ years.

Due to the high discrepancy of the results depending on the technique used, new experimental direct measurements are needed.

2.1) Electron Capture and Positron Transitions

Experimental β^+ /K ratios are reported in Table 2 and compared with theoretical estimations:

Reference	β^+ /K		β^+ /EC	
	experimental	theoretical	experimental	theoretical
1991JA02	$4,2 (13) \times 10^{-7}$			
1976Be02			$1,5 \times 10^{-7}$	
Theory (Logft Code)		$1,75 \times 10^{-5}$		$1,55 \times 10^{-5}$

The LOGFT program (theory) was used with the recent published Q value from 2009AuZZ.

For a 2nd forbidden ε the Logft code gives a theoretical value of $P_{\beta^+} (1072,76) = 0,001 55 (4) \%$ and $P_{\varepsilon} (1072,76) = 99,998 44 (4) \%$ with a logft of 11,89 (3). Notice that the theoretical β^+/ε branching is in complete disagreement with the experimental results of $1,5 \times 10^{-7}$ (1976Be02) and $4,2 (13) \times 10^{-7}$ (1991Ja02). The reason for this discrepancy remains unknown.

The EC-Capture program gives: $P_K = 0,8870 (16)$, $P_L = 0,0966 (13)$, $P_M = 0,0156 (5)$, $P_N = 0,0008 (2)$

From the experimental value $\beta^+/K = 4,2 (13) \times 10^{-7}$ and assuming $K/EC = 0,8870 (16)$ then we have $\beta^+/EC = 3,7 (12) \times 10^{-7}$.

And from $\beta^+ + EC = 100$ we obtain:

$$P_{\beta^+} = 3,7 (12) \times 10^{-5} \%$$

$$P_{\varepsilon} = 99,999 96 (1) \%$$

Which are the adopted values.

3) Atomic Data

3.1) Atomic values (ω_k , ϖ_L and η_{KL}) are from 1996SC06.

3.1.1) X-Radiations, 3.1.2) Auger electrons

The X-ray and Auger electron emission probabilities have been estimated by using the computer code EMISSION. Results were verified with the RADLST computer code.

Good agreement has been found between the total decay energy of 1072,56 (19) keV computed for this decay scheme by RADLST code and the Q value of 1072,76 (19) keV.

4) Gamma emissions

The annihilation radiation emission probability ($I_{\gamma 511}$) is computed as $2 \times P_{\beta^+}$, without the correction factor for the annihilation-in-flight process in the medium, that is $I_{\gamma 511} = 7,4 (24) \times 10^{-5} \%$.

5) References

- 1949PO04 H.S. Pomerance, Phys. Rev. 76 (1949) 195
[σ_{thermal}]
- 1951BR05 A.R. Brosi, C.J. Borkowski, E.E. Conn, J.C. Griess, Phys. Rev. 81 (1951) 391 [$T_{1/2}$]
- 1951WI14 H.W. Wilson, Phys. Rev. 82 (1951) 548
[$T_{1/2}$]
- 1956SA32 B. Saraf. Phys. Rev. 102 (1956) 466
[$T_{1/2}$, inner Bremsstrahlung]
- 1976BE02 D. Bérenyi, G. Hock, A. Ménes, G. Székely, Cs. Ujhelyi, B.A. Zon. Nucl. Phys. A 256 (1976) 87
[I_{β} , Q, ε/β^+]
- 1981MUZQ S.F.Mughabghab, M.Divadeenam, N.E.Holden. Neutron Cross Sections, Vol.1, Neutron Resonance Parameters and Thermal Cross Sections, Part A, Z = 1-60, Academic Press, New York (1981).
[σ_{thermal}]
- 1981NI08 K. Nishiizumi, R. Gensho, M. Honda, Radiochim. Acta 29 (1981) 113
[$T_{1/2}$]
- 1988BO30 D.L. Bowers, L.R. Greenwood, J. Radioanal. Nucl. Chem. 123 (1988) 461
[σ_{thermal}]
- 1991JA02 Z. Janas, M. Pfützner, A. Plochocki, P. Hornshoj, H.L. Nielsen, Nucl. Phys. A524 (1991) 391
[K/β^+]
- 1991NO08 E.Nolte, T.Brunner, T.Faestermann, A.Gillitzer, G.Korschinek, D.Muller, B.Schneck, D.Weselka, V.N.Novikov, A.A.Pomansky, A.Ljubicic, D.Miljanic, H.Vonach, J.Phys.(London) G17 (1991) S355
[$T_{1/2}$]
- 1994RU19 W.Rühm, B.Schneck, K.Knie, G.Korschinek, L.Zerle, E.Nolte, D.Weselka, H.Vonach, Planet. Space Sci. 42 (1994) 227
[$T_{1/2}$]
- 1996SC06 E.Schönfeld, H. Janssen, Nucl. Instrum. Meth. A 369 (1996) 527
[atomic data]
- 2002BA42 C.M. Baglin, Nuclear Data Sheets 95 (2002) 215
[J, π]

- 2004RA23 S.Raman, X.Ouyang, M.A.Islam, J.W.Starner, E.T.Jurney, J.E.Lynn, G.Martinez-Pinedo, Phys. Rev. C 70 (2004) 044318
[σ_{thermal}]
- 2008WAZW A. Wallner, K. Knie, T. Faestermann, G. Korschinek, W. Kutschera, W. Rochow, G. Rugel, H. Vonach, Proc. Int. Conf. on Nuclear Data for Science and Technology, vol. 2, April 22-27, 2007, Nice (France)
[$T_{1/2}$]
- 2009AUZZ G.Audi, W. Meng, D. Lunney, B.Pfeiffer. Priv. Comm. (2009)
[Q value]

⁶⁰Co - Comments on evaluation of decay data by R. G. Helmer

This evaluation was originally completed in September 1996 with minor editing in February 1997 and post-review editing in January 2006 ; the literature available by January 2006 was included by M.-M.Bé (LNE-CEA/LNHB).

1 Decay scheme

In addition to the levels reported in this decay, there are levels in ⁶⁰Ni below the decay energy at 2284 keV (0+) and 2626 (3+). However, based on the limits on the β- branches to these levels (see sect. 2.1), this scheme is considered complete. The scheme is internally consistent since the total decay energy computed from the decay scheme is 2823.0 (5) keV compared to the Q value of 2823.07 (21) keV.

2 Nuclear Data

Q value is from Audi *et al.* (2003Au03).

The half-life values available, in days, are listed. If the value was published in years, it is shown here in years and also converted to days (365.242 days/year).

	Years	Days	Uc	Remarks
1940Li01	5,3 ± 0,7	1936	256	As quoted in 1963Go03 - Outlier (CHV)
1949Se20	5,08 ± 0,08	1855	29	As quoted in 1963Go03 - Outlier (3*S)
1950Br76	5,26 ± 0,17	1921	62	
1951Si25	5,25 ± 0,02	1917,5	7,3	As quoted in 1963Go03
1951To25	5,27 ± 0,07	1925	26	As quoted in 1963Go03
1953Ka21	5,21 ± 0,04	1903	15	Outlier (CHV)
1953Lo09	4,95 ± 0,04	1808	15	Omitted from analysis
1957Ge07	5,24 ± 0,03	1914	11	
1958Ke26	5,33 ± 0,04	1947	15	As quoted in 1965An07 - Outlier (CHV)
1963Go03	5,263 ± 0,003	1922,3	1,1	
1965An07	5,242 ± 0,008	1914,6	2,9	
1968La10	5,270 ± 0,007	1924,8	2,6	
1970Wa19	5,2719 ± 0,0011	1925,5	0,4	Replaced by 1983Wa26
1973Ha60	5,24 ± 0,21	1914	77	
1977Va30	5,283 ± 0,003	1929,6	1,1	
1980Ho17		1925,2	0,4	
1982HoZJ	5,282 ± 0,007	1929,2	2,6	Replaced by 1992Un01
1982RyZX		1924,8	1,0	
1982RyZX		1925,5	0,3	Omitted - unpublished result
1983Ru04		1925,02	0,47	
1983Wa26		1925,5	0,4	
1992Un01		1925,12	0,46	Replaced by 2002Un02
2002Un02		1925,20	0,25	
Adopted	5,2710 ± 0,0008	1925,21	0,29	

One input value (1949Se20) is outlier by 3 sigma, three others are outlier due to Chauvenet criterion (1940Li01, 1953Ka21, 1958Ke26).

For the remaining 14 values, the critical χ^2 is 2.1 ; the reduced χ^2 is 3 ; no value contributes over 50 % of the relative weight. The weighted average is 1925.21 with an internal uncertainty of 0.17 and an external uncertainty of 0.29.

2.1 β^- Transitions

In addition to the main decay to the $J^\pi = 4^+$ level at 2505 keV, there is the possibility of β^- decay from the 5^+ parent to the 0^+ levels at 0 and 2284 keV, the 2^+ levels at 1332 and 2158 keV, and the 3^+ level at 2626 keV.

The β^- decay to the 0^+ levels at 0 and 2284 keV are unique 4^{th} forbidden with expected $\log ft$ values (1973Ra10) > 23 and corresponding $P_{\beta^-} < 1 \times 10^{-10}\%$ and $< 1 \times 10^{-13}\%$, respectively. The decay to the 3^+ level at 2626 is 2^{nd} forbidden with an expected $\log ft > 11$ and a corresponding $P_{\beta^-} < 0.01\%$. This level decays mainly by γ 's of 467 and 1293 keV; the $P_{\gamma}(467)$ has been reported as $< 0.00023\%$ (1976Ca18) and $< 0.0004\%$ (1969Va20), which indicates $P_{\beta^-}(2626) < 0.001\%$.

The β^- decay to the 1332 level is unique 2^{nd} forbidden with an expected $\log ft \geq 12.8$ and a corresponding $P_{\beta^-} \leq 12\%$. The measured values are (in %): 0.15 (1) (1954Ke04), 0.010 (2) (56Wo09), 0.12 (61Ca05), and 0.08 (2) (1968Ha03). The average of 0.12% (3) is adopted.

The decay to the 2158-keV level is unique 2^{nd} forbidden with an expected $\log ft \geq 12.8$ and a corresponding $P_{\beta^-} \leq 0.02\%$. This branch is given as 0.000% (2) from 1969Ra23. (Value is given as 0.18% (3) in 1968Ha03, but this is apparently from a misinterpretation of the γ -ray spectrum.)

The decay to the 2505-keV level is then $100.0 - P_{\beta^-}(1332) - P_{\beta^-}(2158) = 0.12(3) - 0.000(2) = 99.88(3)\%$.

The β^- energies and $\log ft$ values are from the program LOGFT.

2.2 Gamma Transitions

The multiplicities are from the adopted γ data in the Nuclear Data Sheets (1993Ki10).

The total and K-shell internal-conversion coefficients, α and α_K , for the 1173- and 1332-keV γ rays are from the evaluation of the experimental measurements (1985HaZA) and the remaining values were interpolated from the Band tables (2002Ba85).

The internal-pair-formation coefficients were interpolated from the theoretical values of 1979Sc31 and are $\alpha_{\pi}(1173) = 0.000\ 006\ 2$ (7) and $\alpha_{\pi}(1332) = 0.000\ 034$ (4). The former is negligible since it is only about 5% of the corresponding α , but the latter is about 25% of the α , so it needs to be taken into account.

3 Atomic Data (Ni, Z=28)

The data are from Schönfeld and Janßen (1996Sc06).

3.1 and 3.2 X Radiation and Auger Electrons

The data were computed by RADLIST with the Schönfeld atomic data.

4 Radiation Emission

4.1 Electron Emission

Data were computed by the program RADLIST.

4.2 Photon Emissions

The γ -ray energies are from 2000He14 for the 1173-keV and 1332-keV lines and the others are deduced from the level energies resulting from a fit to the γ -ray energies. Besides the 1173 and 1332 values, the input to this fit included:

346.93 (7) from 1976Ca18 where the authors average their result and that of 1969Va20;
other: 346.95 (10) (1969Va20);

826.06 (3) from ⁵⁹Co(p, γ)⁶⁰Ni (1975Er05); others: 826.18 (20) (1969Va20) and 826.28 (9) (1976Ca18, but includes value of 1969Va20);

2158.57 (10) from ⁵⁹Co(p, γ) (1975Er05); others: 2158.8 (4) (1970Di01) from ⁶⁰Co decay and 2158.9 (2) (1969Ra07) and 2159.6 (8) (1969Ho22) from ⁶⁰Cu decay.

For the relative γ -ray emission probabilities, the following data were used.

Relative γ -ray emission probability

γ energy (keV) =	347	467	826	1173/1332	2158	2505
Reference						
1949Fl				100		<2. 5x10 ⁻⁵
1955Wb44	<0. 005			100	0. 0012 (2)	
1959Mb				100		- 4x10 ⁻⁵
1968Ha03			0. 19 (2)	100		
1969Ra23			<0. 02	100	<0. 002	
1969Va20	0. 0078 (12)	<0. 0004	0. 0055(47)	100		
1970Di 01	<0. 006		<0. 04	100	0. 0092 (16)	<4x10 ⁻⁵
1972Le14			0. 003 (2)	100	0. 0005 (2)	
1973Fu15				100	0. 0020 (13)	9(7) 10 ⁻⁶
1976Ca18	0. 00758 (50)	<0. 00023	0. 00762(80)	100. 0	0. 00111 (18)	
1977HaXC				100		<0. 001
1977Lo01	0. 0069 (10)	<0. 0012		100		
1978Fa03				100		<1. 0x10 ⁻⁵
1978Fu05				100		2. 0(4) x10 ⁻⁶
1988Se09				100		5. 2(20) 10 ⁻⁶
Adopted	0. 0075 (4)		0. 0076 (8)	100	0. 0012 (2)	2. 0(4) 10 ⁻⁶

These relative emission probabilities were normalized by requiring that the total β^- emission probability is 100%. For the 1332-keV γ ray, this means :

$$\begin{aligned}
 P_{\gamma}(1332) &= \{100.00 - P_{\gamma}(2158) \times [1+\alpha(2158)] - P_{\gamma}(2505) \times [1+\alpha(2505)]\} / [1.00+\alpha(1332)+\alpha_{\pi}(1332)] \\
 &= [100.00 - 0.0012(2) - 0.0000020(4)] / [1.000 + 0.000128(5) + 0.000034(4)] \\
 &= 99.9988(2) / 1.000162(6) = 99.9826(6)\%.
 \end{aligned}$$

In the evaluation 1991BaZS, this value is computed in the same fashion, but is given as 99.983 (6)% ; the origin of the larger uncertainty is not clear. Similarly, for the 1173-keV γ ray, this means :

$$\begin{aligned}
 P_{\gamma}(1173) &= \{P_{\beta^-}(2505) - P_{\gamma}(347) \times [1+\alpha(347)] - P_{\gamma}(2505) \times [1+\alpha(2505)]\} / [1.00+\alpha(1173)+\alpha_{\pi}(1173)] \\
 &= [99.88(3) - 0.0075(4) - 0.0000020(4)] / [1.000 + 0.000168(4) + 0.0000062(7)] \\
 &= 99.87(3) / 1.000174(4) = 99.85(3) \%.
 \end{aligned}$$

6 References

- 1940Li01 - J. J. Livingood, G. T. Seaborg, Rev. Mod. Phys. **12** (1940) 30 [T_{1/2}]
- 1949Se20 - E. Segrè, C. E. Weigand, Phys. Rev. **75** (1949) 39 [T_{1/2}]
- 1950Br76 - G. L. Brownell, C. J. Maletskos, Phys. Rev. **80** (1950) 1102 [T_{1/2}]
- 1951Si25 - W. K. Sinclair, A. F. Holloway, Nature **167** (1951) 365 [T_{1/2}]
- 1951To25 - J. Toboilem, Compt. rend. **233** (1951) 1360 [T_{1/2}]
- 1953Ka21 - J.Kastner, G.N.Whyte, Phys.Rev. **91** (1953) 332 [T_{1/2}]
- 1953Lo09 - E.E.Lockett, R.H.Thomas, Nucleonics **11**, No.3 (1953) 14 [T_{1/2}]
- 1954Ke04 - G.L.Keister, F.H.Schmidt, Phys.Rev. **93** (1954) 140 [I_β-]
- 1955Wo44 - J.L.Wolfson, Can.J.Phys. **33** (1955) 886 [P_γ]
- 1956Wo09.J.L.Wolfson,Can.J.Phys. **34** (1956) 256 [I_β]
- 1957Ge07 - K.W.Geiger, Phys.Rev. **105** (1957) 1593 [T_{1/2}]
- 1958Ke26 - J. P. Keene, L. A. Mackenzie, C. W. Gilbert, Phys. in Med. Biol. **2** (1958) 360 [T_{1/2}]
- 1961Ca05 - D.C.Camp, L.M.Langer, D.R.Smith, Phys.Rev. **123** (1961) 241 [I_β-]
- 1963Go03 - S.G.Gorbics, W.E.Kunz, A.E.Nash, Nucleonics **21**, No.1 (1963) 63 [T_{1/2}]
- 1965An07 - S.C.Anspach, L.M.Cavallo, S.B.Garfinkel, J.M.R.Hutchinson, C.N.Smith, NP-15663 (1965) [T_{1/2}]
- 1968Ha03 - H.H.Hansen, A.Spernal, Z.Physik **209** (1968) 111 [I_β-]
- 1968La10 - F.Lagoutine, Y.Le Gallic, J.Legrand, Intern.J.Appl.Radiation Isotopes **19** (1968) 475 [T_{1/2}]
- 1969Ho22 - E.J.Hoffman, D.G.Sarantites, Phys.Rev. **181** (1969) 1597 [E_γ]
- 1969Ra07 - F.Rauch, D.M.Van Patter, P.F.Hinrichsen, Nucl.Phys. **A124** (1969) 145 [E_γ]
- 1969Ra23 - S.Raman, Z.Physik **228** (1969) 387 [I_β-]
- 1969Va20 - J.R.Van Hise, D.C.Camp, Phys.Rev.Letters **23** (1969) 1248 [P_γ]
- 1970Di01 - W.R.Dixon, R.S.Storey, Can.J.Phys. **48** (1970) 483 [E_γ, P_γ]
- 1972Le14 - J.Legrand, C.Clement, Int.J.Appl.Radiat.Isotop. **23** (1972) 225 [P_γ]
- 1973Ha60 - G.Harbottle, C.Koehler, R.Withnell, Rev.Sci.Instrum. **44** (1973) 55 [T_{1/2}]
- 1973Ra10 - S.Raman, N.B.Gove, Phys.Rev. **C7** (1973) 1995 [log ft sys]
- 1975Er05 - B. Erlandsson, J. Lyttkens, A. Marcinkowski, Z. Phys. **A272** (1975) 67 [E_γ]
- 1976Ba63 - I. M. Band, M. B. Trzhaskovskaya, M. A. Listengarten, Atomic Data Nucl. Data Tables **18** (1976) 433 [α, α_K]
- 1976Ca18 - D.C.Camp, J.R.Van Hise, Phys.Rev. **C14** (1976) 261 [E_γ, P_γ]
- 1977Lo10 - M.A.Lone, C.B.Bigham, J.S.Fraser, H.R.Schneider, T.K.Alexander, A.J.Ferguson, A.B. McDonald, Nucl. Instrum. Methods **143** (1977) 331 [P_γ]
- 1976Va30 - R. Vaninbrouckx, G. Grosse, Intern. J. Appl. Radiat. Isop. **27** (1977) 727 [T_{1/2}]
- 1978Fa03 - H.Faust *et al.* J.Phys. (London) **G4** (1978) 247
- 1978Fu05 - M.Fujishiro, J.Nucl.Sci.Technol. **15** (1978) 237 [P_γ]
- 1979Sc31 - P.Schluter, G.Soff, At.Data Nucl.Data Tables **24** (1979) 509 [α_π]
- 1980Ho17 - H.Houtermans, O.Milosevic, F.Reichel, Int.J.Appl.Radiat.Isotop. **31** (1980) 153 [T_{1/2}]
- 1982RyZX - A. Rytz, NBS Special publication 626 (1982) 32 [T_{1/2}]
- 1983Ru04 - A.R.Rutledge, L.V.Smith, J.S.Merritt, Nucl.Instrum.Methods **206** (1983) 211 [T_{1/2}]
- 1983Wa26 - K.F.Walz, K.Deberlin, H.Schrader, Int.J.Appl.Radiat.Isotop. **34** (1983) 1191 [T_{1/2}]
- 1985HaZA - H.H.Hansen, European App.Res.Rept.Nucl.Sci.Technol. **6**, No.4 (1985) 777; EUR 9478 EN [α, α_K]
- 1988Se09 - S.Seuthe, H.W.Becker, C.Rolfs, S.Schmidt, H.P.Trautvetter, R. W. Kavanagh, F.B.Waanders, Nucl. Instrum. Methods **A272** (1988) 814 [P_γ]
- 1992Un01 - M.P.Unterweger, D.D.Hoppes, F.J.Schima, Nucl.Instrum.Methods Phys.Res. **A312** (1992) 349 [T_{1/2}]
- 1993Ki10 - M. M. King, Nucl.Data Sheets **48** (1986) 25 [J^π, multipolarities]
- 1995Au04 - G. Audi, A. H. Wapstra, Nucl. Phys. **A595** (1995) 409 [Q]
- 1996Sc06 - E. Schönfeld, H. Janßen, Nucl. Instr. Meth. **A 369** (1996) 527 (ω)
- 2000He14 - R. G. Helmer and C. van der Leun, Nucl. Instr. Meth. **A 450** (2000) 35 [E_γ]
- 2002Ba85 - I.M.Band, M.B.Trzhaskovskaya, C.W.Nestor, S.Raman. At. Data and Nucl. Data Tables **81**, 1&2 (2002) 1 [ICC]
- 2002Un02 - M.P. Unterweger. Applied Radiat. Isotopes **56** (2002) 125 [T_{1/2}]
- 2003Au03 - G. Audi, A. H. Wapstra, C.Thibault. Nucl. Phys. **A729**(2003)337 [Q]

⁶¹Cu – Comments on evaluation
by M.-M. Bé

This evaluation was completed in August 2013. Literature published to this date are included.

Decay Scheme

⁶¹Cu ($T_{1/2} = 3,366$ h) decays 100 % by electron capture and β^+ disintegrations to various excited levels and to the ground state of ⁶¹Ni.

Q^{EC} value for ⁶¹Cu decay was adopted from Wang *et al.* (2012Wa38).

The spins, parities and level energies are based on the mass-chain evaluation of Bhat (1999Bh04).

Half-life

The experimental half-life values used to calculate the recommended value are listed in the following table:

Reference	h	uc (h)	Comments
1969Ri04	3,408	0,012	Smallest of uncertainty measurement
1972Du09	3,26	0,05	
1973Ne02	3,39	0,04	
1982Gr10	3,333	0,01	Original uncertainty: 0,005
Crit (χ^2)	3,8		
$\chi^2 / (n-1)$	8,6		
UWM	3,348		
LWM	3,366		
uc(WM)int. :	0,008		
uc(WM)ext. :	0,023		
Adopted:	3,366	0,033	EXPANDED

The original uncertainty of 0,005 h given by Grütter (1982Gr10) was defined, in the article, as the statistical uncertainty only, at the 1σ level. It has been increased to 0,01 h, even if it is difficult to appreciate the real systematic component due to a lack of information in the publication.

The data set is discrepant, the adopted value is the weighted mean of the four values with an expanded uncertainty to cover the most precise value of 1982Gr10.

Values given by Berman (1954Be84), Martins *et al.* (1982Ma41) and Crisler *et al.* (1972Cr02) have not been taken into account because they were obtained as a mean to control another experiment and not as a specific study.

Electron Capture and β^+ transitions

All EC and β^+ energies were derived from the level energies and the Q value.

The shell and sub shells capture probabilities were calculated by mean of the EC-Capture program (1998Sc28).

To the ground state

In this decay scheme the ground state is fed by a transition where electron capture and β^+ emission are in competition, the determination of its probability is crucial to establish the decay scheme. Some attempts have

been made to determine the β^+ probability mainly from the measurement of the 511 keV photon emission in coincidence with other gamma emission. It must be noted that these results depend of other quantities and some of them were given without uncertainty. However, due to the importance of such a value, all the available results have been taken into consideration.

Nussbaum *et al.* (1956Nu02) investigated the Cu-61 decay by using coincidence techniques and obtained a β^+ probability to the ground state of 51 (5) %. They also measured the K conversion coefficients of the 67,4 and 282,8 keV transitions.

Bolotin *et al.* (1967Bo01) studied the decay scheme by using coincidence techniques and obtained a β^+ probability to the ground state of 50,7 %. Absolute γ ray intensities were given with an uncertainty of 5 %.

Schöneberg *et al.* (1967Sc02) determined the decay scheme from their measurement results and using the K conversion coefficients of the 67,4 and the 282,8 keV transitions measured by Nussbaum. Finally, they found a β^+ probability to the ground state of 51,7 % and a total 511 keV intensity of 126,1 (40) %. The other absolute γ ray intensities were given with an uncertainty of 3 – 4 %.

Béraud *et al.* (1967Be12) measured the β^+ in coincidence with other gamma rays, they adopted the probability of the 67,4 keV transition of Nussbaum *et al.* (1956Nu02). Finally, they obtained a β^+ probability to the ground state of 53,1 %. They determined the γ intensities with an uncertainty of 10 %.

Since three data have no uncertainty stated by the authors, the simple mean of the four values is adopted. An uncertainty which covers all values would lead to 51,6 (15) %, however considering the uncertainties on the various experimental results, an uncertainty of 5 % is adopted here. That is: β^+ to g.s. = 51,6 (25) %.

The theoretical ratios EC/ β^+ have been calculated by the Logft program from the theoretical tables of Gove *et al.* (1971Go40). The ratio for transitions to the ground state level is 0,3155 (33), then the total capture probability to the g.s. is 16,3 (8) %.

The total EC + β^+ to the g.s. is 67,9 (26) %.

To excited levels

The total EC + β^+ to each level of the decay scheme were derived from the P(ce+ γ) probability balance, then each component was obtained from the EC/ β^+ ratio.

Level	Energy (keV)	Spin	EC/ β^+	uc
0	0	3/2 ⁻	0,3155	0,0033
1	67,412	5/2 ⁻	0,3791	0,00394
2	282,9568	1/2 ⁻	0,7471	0,0079
3	656,012	1/2 ⁻	4,25	0,05
4	908,62	5/2 ⁻	38	0,06
5	1014,8	7/2 ⁻	199	5
6	1099,622	3/2 ⁻		
7	1132,332	5/2 ⁻		
8	1185,236	3/2 ⁻		
9	1609,639	5/2 ⁻		
10	1729,471	3/2 ⁻		
11	1997,7	5/2 ⁻		
12	2124	1/2 ⁻		

Other

The total β^+ emissions, as determined above, lead to a 511 keV photon emission of 123 (5) % which is in good agreement with the experimental value of Schöneberg *et al.* (1967Sc02) of 126,1 (40) %.

Relative gamma emission intensities, I_γ

Meyer (1978Me10) and Satyanarayana (1988Sa26) carried out relative measurements, the 282 keV gamma ray being the reference ray.

Nussbaum *et al.* (1956Nu02), Ritter (1969Ri04), Schöneberg (1967Sc02), Bolotin (1967), Béraud *et al.* (1967Be12) studied the decay scheme, they made use of coincidences techniques to determine the population of the various levels in Ni-61 and they published results for the gamma intensities. The absolute γ intensities and the intensities of the β^+ they obtained are correlated. The values of the β^+ intensities have already been used in this evaluation to obtain the branching ratio to the ground state, so, they have not been included in the determination of the relative gamma intensities.

The adopted relative gamma intensities are the mean of the values of Meyer (1978Me10) and Satyanarayana (1988Sa26). The adopted uncertainty is the standard deviation, but is never lower than the lowest experimental uncertainty.

Levels (i,f)	Energy (keV)		Meyer 1978		Satyan. 1988		UWM	uc ext.		Adopted relative I _γ	
	E _γ	E _γ (uc)	I _γ	I _γ (uc)	I _γ	I _γ (uc)	I _γ	I _γ (uc)	I _γ	I _γ (uc)	
(1,0)	67,412	0,003	31,5	1	34,7	1,1	33,1	1,6	33,1	1,6	
(7,5)	117,5				0,086	0,046	0,086	0,046	0,086	0,046	
	190,79	0,12	0,04	0,008					omitted		
(2,1)	215,55	0,18	0,036	0,009	0,176	0,058	0,106	0,021	0,106	0,058	
	276,688	0,053	0,21	0,02					omitted		
(2,0)	282,956	0,002	100		100	2,4	100		100	2,4	
(3,2)	373,05	0,005	17,2	0,5	17,6	0,4	17,4	0,195	17,4	0,4	
	443,5	0,1	0,03	0,01					omitted		
(8,3)	529,169	0,022	3,3	0,6	3,08	0,07	3,19	0,025	3,19	0,070	
	544,8				0,048	0,03	0,048	0,030	0,048 ^(*)	0,030	
(3,1)	588,605	0,009	9,6	0,1	9,57	0,17	9,585	0,013	9,585	0,1	
(4,2)	625,605	0,024	0,4	0,03	0,33	0,037	0,365	0,034	0,365	0,034	
	629,9	0,25	0,05	0,02					omitted		
(3,0)	656,008	0,004	85,3	2	88,34	1,52	86,82	1,465	86,82	1,520	
(9,4)	701,1	0,3	0,09	0,02					0,09 ^(w)	0,020	
(6,2)	816,692	0,013	2,89	0,06	2,52	0,06	2,705	0,185	2,705	0,185	
(10,4)	820,89	0,17	0,18	0,02					0,18 ^(w)	0,020	
(4,1)	841,211	0,017	1,98	0,08	1,755	0,054	1,8675	0,104	1,8675	0,104	
(8,2)	902,294	0,02	0,72	0,02	0,68	0,04	0,7	0,016	0,7	0,020	
(4,0)	908,631	0,017	9,7	0,3	9,03	0,168	9,365	0,286	9,365	0,286	
(5,1)	947,4	0,4	0,02	0,01	0,08	0,04	0,05	0,014	0,05	0,014	
(5,0)	1014,8	0,4	< 0,01		0,086	0,03	0,048		0,086	0,03	
(6,1)	1032,162	0,027	0,53	0,09	0,346	0,03	0,438	0,055	0,438	0,055	
(7,1)	1064,896	0,02	0,48	0,04	0,395	0,03	0,4375	0,041	0,4375	0,041	
(10,3)	1073,465	0,025	0,43	0,03	0,267	0,025	0,3485	0,080	0,3485	0,080	
(11,4)	1089,11		≤ 0,005						0,005 ^(w)		
(6,0)	1099,56	0,019	2,27	0,08	2,01	0,05	2,14	0,117	2,14	0,117	
(8,1)	1117,822	0,043	0,39	0,03	0,265	0,034	0,3275	0,062	0,3275	0,062	
(7,0)	1132,351	0,032	0,79	0,03	0,737	0,034	0,7635	0,026	0,7635	0,030	
(8,0)	1185,234	0,015	29,5	0,6	30,71	0,57	30,105	0,604	30,105	0,604	
(10,2)	1446,492	0,019	0,4	0,02	0,369	0,024	0,3845	0,015	0,3845	0,020	
(9,1)	1542,204	0,023	0,27	0,01	0,21	0,018	0,24	0,025	0,24	0,025	
(9,0)	1609,625	0,048	0,22	0,02	0,174	0,017	0,197	0,023	0,197	0,023	

Levels (i,f)	Energy (keV)		Meyer 1978		Satyan. 1988		UWM	uc ext.	Adopted relative I _γ	
	E _γ	E _γ (uc)	I _γ	I _γ (uc)	I _γ	I _γ (uc)	I _γ	I _γ (uc)	I _γ	I _γ (uc)
(10,1)	1662	0,019	0,42	0,02	0,433	0,024	0,4265	0,006	0,4265	0,020
(10,0)	1729,473	0,018	0,64	0,04	0,442	0,024	0,541	0,087	0,541	0,087
	1840,7	0,2	0,014	0,004					omitted	
(11,0)	1997,73	0,85	0,03	0,01	0,032	0,01	0,031	0,001	0,031	0,010
	2120				0,08		0,08		0,08 ^(*)	
(13,0)	2123,432	0,035	0,32	0,01	0,339	0,024	0,3295	0,007	0,3295	0,010

Some rays were observed only by Meyer (1978Me10) and not confirmed by Satyanarayana (1988Sa26), they were omitted. Similarly, some rays questionable (*) are not placed as well. But some rays (w) observed by Meyer were also observed by Wadsworth *et al.* (1977Wa03), they were adopted in this decay scheme.

Internal conversion coefficients

The adopted ICC are the theoretical values calculated by the BrIcc program (2008Ki07).

Most of the transitions are of M1+E2 type. The adopted mixing ratios are from the experimental study of Wadsworth *et al.* (1977Wa03) who carried out an extensive spectroscopy of the nucleus ⁶¹Ni.

For the 67 and 282 keV transitions, the K internal conversion coefficients were measured by Nussbaum *et al.* (1956Nu02) to be 0,11 and 0,004 respectively. These experimental values are in good agreement with the theoretical ones obtained by the BrIcc program for transitions of M1 character.

Normalisation Factor

For 100 % of ⁶¹Cu decays, 100 % of the transitions (β⁺, γ, EC) must populate the ground state of the daughter:

$$\sum_i I_{\gamma i} [1 + \alpha_{Ti}] = \frac{100\% - (P_{\beta^+} + P_{EC})}{N}$$

Where:

The sum $\sum_i I_{\gamma i} [1 + \alpha_{Ti}]$ is over the γ transitions feeding the ground state; $I_{\gamma i}$ is the relative emission intensity of the i^{th} gamma-ray, α_{Ti} is its total internal conversion coefficient and, $P_{\beta^+} + P_{EC}$ is the sum of β⁺ and EC – transitions populating the ground state. N is a normalisation factor between the relative γ intensities and the absolute values.

From above, the total EC + β⁺ to the g.s. is 67,9 (33) %, then the sum of the γ gamma transition probabilities to the g.s. must be equal to 32,1 (33) %. Therefore, a normalization factor of 12,0 (17) % is deduced.

Atomic Data

The fluorescence yield data, the relative K x-ray emission probabilities, the ratios P(KLX)/P(KLL) and P(KXY)/P(KLL) are from Schönfeld *et al.* (1996Sc06).

The Auger electrons and X-ray absolute intensities were calculated by the EMISSION program (2000Sc47) from the related decay data (γ intensities, ICC, P_{EC} probabilities, etc.) as determined above.

Data Consistency

The sum of all the energies involved in a decay (EC, γ, etc.) is 2238 (46) keV which can be compared with the value of 2237,5 (10) as adopted for the Q value.

References

- 1954Be84** A.I.Berman, K.L.Brown. Phys.Rev. 96, (1954) 83. Half-life
- 1956Nu02** R.H.Nussbaum, A.H.Wapstra, W.A.Bruil, M.J.Sterk, G.J.Nijgh, N.Grobben. Phys. Rev. 101 (1956) 905. Beta plus emission probabilities
- 1967Bo01** H.H.Bolotin, H.J.Fischbeck. Phys. Rev. 158 (1967) 1069. Beta plus emission probabilities
- 1967Sc02** R.Schöneberg, A.Flammersfeld. Z. Physik 200 (1967) 205. Beta plus emission probabilities
- 1967Be12** R.Béraud, I.Berkes, J.Daniere, M.Levy, G.Marest, R.Rougnny. Nucl. Phys. A99 (1967) 577. Beta plus emission probabilities
- 1969Ri04** J.C.Ritter, R.E.Larson. Nucl. Phys. A127 (1969) 399. Half-life
- 1971Go40** N.B. Gove, M.J. Martin. Nucl. Data Tables 10 (1971) 205. EC/positron ratios
- 1972Du09** G.H.Dulfer, B.O.ten Brink, T.J.Ketel, A.W.B.Kalshoven, H.Verheul. Z. Physik 251 (1972) 416. Half-life
- 1972Cr02** D.F.Crisler, H.B.Eldridge, R.Kunselman, C.S.Zaidins. Phys. Rev. C5 (1972) 419. Half-life
- 1973Ne02** D.A.Newton, S.Sarkar, L.Yaffe, R.B.Moore. J. Inorg. Nucl. Chem. 35 (1973) 361. Half-life
- 1977Wa03** R.Wadsworth, A.Kogan, P.R.G.Lornie, M.R.Nixon, H.G.Price, P.J.Twin. J. Phys. (London) G3 (1977) 35. Gamma rays
- 1978Me10** R.A.Meyer, A.L.Prindle, W.A.Myers, P.K.Hopke, D.Dieterly, J.E.Koops. Phys. Rev. C17 (1978) 1822. Half-life, gamma-ray energies and emission intensities
- 1982Ma41** M.N.Martins, E.Wolyne, M.C.A.Campos. Phys. Rev. C26 (1982) 1936. Half-life
- 1982Gr10** A.Grutter, . Int. J. Appl. Radiat. Isotop. 33 (1982) 533. Half-life
- 1988Sa26** G.Satyanarayana, N.Venkateswara Rao, G.S.Sri Krishna, M.V.S.Chandrasekhar Rao, S.Bhuloka Reddy, D.L.Sastry, S.N.Chintalapudi, V.V.Rao. Nuovo Cim. 99A (1988) 309. Gamma-ray emission intensities
- 1996Sc06** E. Schönfeld, H. Janssen. Nucl. Instrum. Methods Phys. Res. A369 (1996) 527. Atomic Data
- 1998Sc28** E. Schönfeld, .Appl. Radiat. Isot. 49 (1998) 1353. Fractional EC probabilities
- 1999Bh04** M.R.Bhat, .Nucl. Data Sheets 88 (1999) 3. Spin and Parity, level energies
- 2000Sc47** E. Schönfeld, H. Janssen. Appl. Radiat. Isot. 52 (2000) 595. P(X), P(Ae)
- 2008Ki07** T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor Jr. Nucl. Instrum. Methods Phys. Res. A589 (2008) 202. Theoretical ICC
- 2012Wa38** M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer. Chin. Phys. C36 (2012) 1603. Q-value

⁶³Ni - Comments on the Evaluation of Decay Data by K. B. Lee

This evaluation was completed in August 2005.

1. Decay Scheme

⁶³Ni disintegrates by β^- emission (100%) to the ground state of the stable nuclide ⁶³Cu.

2. Nuclear Data

The Q value (66.980 (15) keV) is from the measured value of Holzschuh (1999Ho09). This value is in agreement with 66.975 (15) keV from the atomic mass table of Audi et al. (2003Au03).

The measured ⁶³Ni half-life values are given below.

Reference	Values (years)	Comments
Brosi (1951Br)	85 (20)	Omitted from analysis
Wilson (1950Wi)	61	Omitted from analysis
McMullen (1956Mc)	125 (6)	Omitted from analysis
Horrocks (1962Ho)	93.9 (20)	Revised by Collé (1996Co25)
Barnes (1971Ba89)	101.21 (20)	Revised by Collé (1996Co25)
Collé (1996Co25)	101.06 (197)	

The first three older and less precise historical values were omitted from the analysis. The Horrocks (1962Ho) and Barnes (1971Ba89) values were revised by Collé (1996Co25) using more accurate nuclear data and thereby more rigorously calculated liquid scintillator detection efficiencies. The weighted average for the last three values is 98.7 years; with an internal uncertainty of 1.1 years; an external uncertainty of 2.4 years and a reduced- χ^2 of 4.38.

The evaluator's recommended value is the weighted average : 98.7 (24) years.

2.1 β^- Transitions

The evaluator has calculated (using the LOGFT program) a *log ft* of 6.7 for this allowed transition.

The various measured β^- end-point energy values (or Q-values) are summarized below.

Reference	Values (keV)	Comments
Preiss (1957Pr)	67.0 (5)	Omitted
Hsue (1966Hs01)	65.87 (15)	Omitted
Hetherington (1987He14)	66.946 (20)	Omitted
Kawakami (1992Ka29)	66.9451 (39)	Omitted
Ohshima (1993Oh2)	66.9459 (54)	Omitted
Ohshima (1993Oh2)	66.9433 (126)	Omitted
Holzschuh (1999Ho09)	66.980 (15)	Adopted value

Uncertainties given in 1993Oh2 include systematic values combined in quadrature with statistical uncertainties.

Holzschuh et al. (1990Ho09) pointed out that in the previous measurements of end-point energies the excitation of atomic electrons was not taken into account. That means that all the other values are biased by an amount of the order of the mean electron excitation energy (85 eV). Therefore the evaluator has recommended the value given in 1990Ho09. Besides, a second end-point energy given in 1993Oh2 was obtained under the assumption of the existence of a 17 keV neutrino.


3. Atomic Data

The fluorescence yield is from the compilation of 1996Sc33.

4. Radiations

The mean energy of beta particles has been computed as 17.434 (4) keV using the LOGFT program.

4. Main Production Modes



6. References

- 1951Br Brosi A. R., Borkowski C. J., Conn E. E. and Griess Jr J. C. , Phys. Rev. **82** (1951) 391-395 [$T_{1/2}$]
- 1950Wi Wilson H. W., Phys. Rev. **79** (1950) 1032-1033 [$T_{1/2}$]
- 1956Mc McMullen C. C., Pate B. D., Tomlinson R. H. and Yaffe L. , Can. J. Chem. **33** (1956) 1742-1746 [$T_{1/2}$]
- 1957Pr Ivor L. Preiss, R. W. Fink and B. L. Robinson, J. Inorg. Nucl. Chem. **4** (1957) 233-236 [End-point energy]
- 1962Ho Horrocks D. L. and Harkness A. L., Phys. Rev. **125** (1962) 1619-1620 [$T_{1/2}$]
- 1966Hs01 S. T. Hsue, L. M. langer, E. H. Spejewski and S. M. Tang, Nucl. Phys. **80** (1966) 657 [End-point energy]
- 1971Ba89 I.L. Barnes, S. B. Garfinkel and W. B. Mann, Int. J. Appl. Radiat. Isotop. **22** (1971) 777 [$T_{1/2}$]
- 1987He14 D. W. Hetherington, R. L. Graham, M. A. Lone, J. S. Geiger and G. E. Lee-Whiting, Phys. Rev. **C36** (1987) 1504 [End-point energy]
- 1992Ka29 H. Kawakami, S. Kato, T. Ohshima, C. Rosenfeld, H. Sakamoto, T. Sato, S. Shibata, J. Shirai, Y. Sugaya, T. Suzuki, K. Takahashi, T. Tsukamoto, K Ueno, K Ukai, S. Wilson and Y. Yonezawa, Phys. Lett. **287B** (1992) 45 [End-point energy]
- 1993Oh02 T. Ohshima, H, Sakamoto, T. Sato, J. Shirai, T. Tsukamoto, Y. Sugaya, K. Takahashi, T. Suzuki, C. Rosenfeld, S. Wilson, K Ueno, Y. Yonezawa and H. Kawakami, Phys. Rev. **D47** (1993) 4840 [End-point energy]
- 2003Au03 G. Audi, A. H. Wapstra and C. Thibault, Nucl.Phys. **A729** (2003) 337 [Q]
- 1996Sc33 E. Schönfeld and H. Janßen, Nucl. Phys. Instr. Meth. Phys. Res. A369 (1996) 527 [Atomic data]
- 1996Co25 R. Collé and B. Z. Zimmerman, Appl. Radiat. Isot. **47** (1996) 677 [$T_{1/2}$]
- 1999Ho09 E. Holzschuh, W. Kundig, L. Palermo, H. Stussi and P. Wenk, Phys. Lett. **451B** (1999) 247 [End-point energy]

**⁶³Zn – Comments on evaluation of decay data
by A. L. Nichols**

Evaluated: January/February 2013

Evaluation Procedures

Limitation of Relative Statistical Weight Method (LWM) and other analytical techniques were applied to obtain averaged data throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the *values* used to calculate the average.

Decay Scheme

⁶³Zn ($T_{1/2} = 38.33$ min) decays 100 % by electron capture/ β^+ decay ($Q(\text{EC}) = 3366.2$ (15) keV) to various excited nuclear levels and the ground state of ⁶³Cu (stable). A reasonably well-defined decay scheme was derived from the gamma-ray measurements of 1967De08, 1969Bo15, 1970Ki06, 1971GiZS, 1974KI02 and 1982Gr10, consisting of 20 EC/8 β^+ transitions and 64 gamma-ray emissions. Weighted mean relative emission probabilities were calculated for a majority of the main gamma rays, while equivalent data for the other gamma transitions were adopted primarily from the more comprehensive measurements of 1974KI02; all of these relative emission probabilities were defined in terms of the 669.93-keV gamma ray (100 %).

Nuclear Data

⁶³Zn decay is viewed as a potentially suitable beam monitor for the cyclotron production of β^+ emitters adopted in positron emission tomography (PET).

Half-life

The measurements of 1938St05, 1939De01, 1947Hu20, 1959Ri38, 1961Cu02, 1968Go10, 1969Bo11, 1969Bo15, 1972Cr02, 1974Co16, 1982Gr10 and 2000Le02 were adopted to give a weighted mean half-life of 38.33(10) minutes based on the limitation of relative statistical weight method (LWM).

Reference	Half-life (min)
1938St05	38.3 (5)
1939Bo05	38 [*]
1939De01	38.5 (8)
1947Hu20	38.3 (5)
1948Wa13	38.3 [*]
1959Ri38	38.1 (3)
1960Pr05	36 (2) [‡]
1961Cu02	38.4 (2)
1961Ra06	39.9 (8) [‡]
1961Va08	37.6 (3) [‡]
1965Pa18	39.0 (1) [‡]
1968Go10	38.4 (1)
1969Bo11	37.9 (2)
1969Bo15	38.5 (1)
1972Cr02	38.6 (2) [†]
1974Co16	38.0 (1)
1982Gr10	38.47 (5) [#]
2000Le02	38.1 (8)
Recommended value	38.33 (10) [§]

^{*} Uncertainty not specified – not included in weighted mean analysis of the data set.

[‡] Defined as an outlier, and therefore not adopted in the analytical procedures.

[†] Weighted mean of seven separate measurements (eighth measurement defined as an outlier), with an uncertainty of 0.2 adopted to align with the smallest uncertainty of the values used to calculate the average value.

[#] Only the statistical uncertainty was defined by the authors to be 0.05 at the 1 σ level – uncertainty has been increased by a factor of two (to 0.10) as a means of introducing an artificial systematic component.

[§] Recommended uncertainty has been adjusted from 0.05 to 0.10, in alignment with the smallest uncertainty of the values used to calculate the average value.

Limitation of relative statistical weight method (LWM), normalised residual method (NRM), Rajeval technique, bootstrap method, and Mandel-Paule approach were considered in the analysis of the data set.

Analytical method	Half-life (min)	$\chi^2/(N-1)$	$\chi^2/(N-1)_{\text{critical}}$
LWM	38.33 (5)*	2.14	2.25
NRM	38.37 (5)*	1.54	1.79
Rajeval	38.41 (5)*	0.91	–
Bootstrap	38.34 (10)	2.14	–
Mandel-Paule	null result	–	–

* Uncertainty increased from 0.05 to 0.10, in alignment with the smallest measured uncertainty of the values used to calculate the average value.

A half-life value of 38.33 (10) minutes is recommended, as quantified by the LWM analytical procedure.

Q value

Q_{EC} -value for ⁶³Zn EC decay of 3366.2 (15) keV was adopted from Wang *et al.* (2012Wa38).

Gamma-ray energies and emission probabilities

Energies

The well-defined nuclear level energies of 2001Ba27 were used to calculate the gamma transition energies and their uncertainties, and these data were adjusted to account for gamma recoil in the formulation of recommended gamma-ray emission energies and uncertainties. Greater confidence was placed on this approach because of the more wide-ranging origins of the gamma transition data even though the energies of a significant number of gamma-ray emissions have been directly measured by 1967De08, 1969Bo15, 1970Ki06, 1971GiZS and 1974Kl02.

Adopted energies, spins and parities for the nuclear levels of ⁶³Cu (2001Ba27).

Nuclear level number	Nuclear level energy (keV)	Spin and parity
0	0.0	3/2 –
1	669.93 (4)	1/2 –
2	962.02 (3)	5/2 –
3	1326.76 (5)	7/2 –
4	1412.16 (4)	5/2 –
5	1547.00 (5)	3/2 –
6	1860.63 (6)	7/2 –
7	2012.92 (11)	3/2 –
8	2062.45 (8)	(1/2) –
9	2081.32 (22)	5/2 (–)
10	2092.13 (10)	7/2 –
11	2336.54 (12)	5/2 –
12	2497.19 (9)	(3/2 –)
13	2511.06 (6)	1/2, 3/2, 5/2
14	2535.83 (7)	(5/2) –
15	2696.66 (13)	1/2 –, 3/2 –
16	2716.47 (9)	3/2 –, 5/2 –
17	2780.23 (21)	(1/2 –, 3/2 –)
18	2808.10 (8)	3/2 –
19	2857.9 (3)	(1/2 –, 3/2 –)
20	2888.9 (4)	1/2 –, 3/2, 5/2 –
21	3042.59 (8)	(5/2 –)
22	3101.4 (4)	1/2 –, 3/2 –

Gamma transition energies, and measured and recommended gamma-ray energies.

E _{TP} (keV) [#]	E _γ (keV)									
	1959Ha10	1959Ri38	1961Cu02	1961Va08	1967De08	1969Bo15	1970Ki06	1971GiZS	1974KI02	Recommended [§]
244.40 (22)	–	–	–	–	–	–	–	–	244.26 (50)	244.40 (22)
364.74 (6)	–	–	–	–	–	364.5 (20)*	–	–	365.22 (40)	364.74 (6)
443.70 (12)	–	–	–	–	–	–	–	–	443.13 (20)	443.70 (12)
450.14 (5)	–	–	–	–	–	450.0 (5)	449.8 (10)	450.0 (2)	449.93 (5)	450.14 (5)
475.91 (13)	–	–	–	–	–	–	–	–	475.8 (9)	475.91 (13)
–	–	511	511	detected	511	511.0	511	511.0	511.0	annihilation radiation
515.45 (9)	–	–	–	–	–	–	–	–	515.0 (10)*	515.45 (9)
534.32 (23)	–	–	–	–	–	–	–	–	533.8 (6)	534.32 (23)
584.98 (6)	–	–	–	–	–	–	–	585.2 (5)	584.82 (15)	584.98 (6)
624.34 (13)	–	–	–	–	–	–	–	624.1 (6)	624.26 (30)	624.34 (13)
669.93 (4)	670 (20)	670 (5)	669 (2) [†]	–	688 (1)	669.6 (2)	669.75 (20)	669.71 (10)	669.62 (5)	669.93 (4)
675.20 (9)	–	–	–	–	–	–	–	–	675.03 (60)	675.20 (9)
683.74 (17)	–	–	–	680 (10)	–	684.7 (17)	–	–	685.6 (6)	683.74 (17)
742.23 (6)	–	–	–	–	–	742.5 (5)	742.0 (10)	742.5 (5)	742.25 (10)	742.23 (6)
754.56 (23)	–	–	–	–	–	–	–	754.4 (7)	754.81 (80)	754.56 (23)
765.37 (11)	–	–	–	–	–	–	–	–	765.7 (5)	765.37 (11)
–	–	810 (10)	–	–	–	–	–	–	–	–
877.07 (6)	–	875 (10)	–	–	–	–	–	–	877.2 (8)	877.06 (6)
898.61 (7)	–	–	–	–	–	–	–	899.2 (6)	899.02 (40)	898.60 (7)
924.38 (13)	–	–	–	–	–	923.5 (10)	–	924.6 (6)	924.30 (50)	924.37 (13)
962.02 (3)	960 (30)	966 (6)	962 (2) [†]	970 (10)	961 (1)	961.9 (2)	962.1 (2)	962.14 (10)	962.06 (4)	962.01 (3)
988.83 (9)	–	–	–	–	–	–	–	–	989.6 (7)	988.82 (9)
1050.90 (11)	–	–	–	–	–	–	–	–	1048.78 (50)	1050.89 (11)
–	–	–	–	–	–	1087 (2)	–	–	–	–
–	–	1100 (20)	–	–	–	–	–	–	–	–
–	–	–	–	–	1116 (2)	–	–	–	–	⁶⁵ Zn decay
1123.67 (8)	–	–	–	–	–	1123.7 (3)	1123.6 (10)	1123.8 (6)	1123.72 (7)	1123.66 (8)
1130.11 (10)	–	–	–	–	–	–	–	1130.6 (5)	1130.67 (25)	1130.10 (10)
1149.66 (14)	–	–	–	–	–	1150 (2)	–	1149.6 (3)	1149.50 (16)	1149.65 (14)
1169.47 (10)	–	–	–	–	–	1168 (3)	–	–	1169.62 (30)	1169.46 (10)
–	–	–	–	–	–	1189 (3)	–	–	–	–
1209.07 (9)	–	–	–	–	–	1208 (2)	1209.1 (10)	1208.6 (8)	1208.78 (3)	1209.06 (9)
1233.23 (22)	–	–	–	–	–	–	–	–	1233.7 (5)	1233.22 (22)
–	–	1270 (20)	–	–	–	–	–	–	–	–
1326.76 (5)	–	1350 (20)	1330	1350 (20)	1327 (4)	1326.4 (3)	1327.1 (10)	1327.0 (4)	1327.03 (8)	1326.75 (5)

1342.99 (12)	–	–	–	–	–	–	1340.0 (10)	–	1341.7 (6)	1342.97 (12)
1374.52 (12)	–	–	–	–	–	1374.3 (3)	1374.4 (10)	1374.4 (3)	1374.47 (13)	1374.50 (12)
1389.71 (10)	–	–	–	–	?	–	–	1389.5 (6)	1389.66 (8)	1389.69 (10)
1392.52 (9)	–	–	–	–	1390 (4)	1391.5 (4)	1392.1 (10)	1392.3 (5)	1392.55 (8)	1392.50 (9)
1412.16 (4)	1440 (30)	1440 (30)	1420 (30)	1430 (20)	1412 (1)	1411.9 (2)	1412.1 (10)	1412.07 (20)	1412.08 (5)	1412.14 (4)
1445.7 (3)	–	–	–	–	–	–	–	–	1445.8 (4)	1445.7 (3)
1481.34 (9)	–	–	–	–	–	–	–	–	1479.08 (50)	1481.32 (9)
1547.00 (5)	–	1560 (30)	1550 (50)	1540 (20)	1548 (1)	1546.9 (2)	1546.6 (10)	1546.8 (6)	1547.04 (6)	1546.98 (5)
1573.81 (8)	–	–	–	–	1573 (4)	1574.0 (20)	1573.3 (10)	1573.4 (5)	1573.71 (20)	1573.79 (8)
1666.61 (13)	–	1670 (30)	–	–	–	–	–	–	1667.2 (6)	1666.59 (13)
1696.6 (10) ^Δ	–	–	–	–	1700 (4)	–	–	–	1696.6 (10)	1696.6 (10) ^Δ
1754.45 (9)	–	–	–	–	–	–	–	–	1754.89 (50)	1754.42 (9)
–	–	1800 (30)	–	–	–	1816.9 (10)	–	–	–	–
1827.26 (10)	–	–	1830 (40)?	–	1822 (4)	–	–	1825.4 (6)	1827.0 (5)	1827.23 (10)
1860.63 (6)	–	–	–	–	–	–	?	1860.9 (6)	1861.34 (30)	1860.60 (6)
1865.90 (8)	–	–	–	–	–	1865.3 (3)	1863.4 (10)	1865.7 (6)	1866.08 (30)	1865.87 (8)
1926.9 (4)	–	1900 (40)	–	–	–	1926.4 (20)	–	1926.8 (8)	1927.18 (70)	1926.9 (4)
2012.92 (11)	2000?	–	–	–	–	2012.0 (5)	2010.9 (10)	2012.0 (5)	2011.44 (50)	2012.89 (11)
2026.73 (14)	–	2040 (40)	2040 (40)	–	–	2027.2 (5)	2026.7 (10)	2026.9 (3)	2026.81 (30)	2026.70 (14)
2046.54 (10)	–	–	–	–	–	2048.6 (15)	–	2047.2 (8)	2046.44 (80)	2046.50 (10)
2062.45 (8)	–	–	–	–	2059 (4)	2062.0 (8)	2062.7 (10)	2062.3 (5)	2062.11 (30)	2062.41 (8)
2081.32 (22)	–	–	–	–	2082 (5)	2082.0 (5)	2082.1 (10)	2081.4 (7)	2081.44 (30)	2081.28 (22)
2092.13 (10)	–	–	–	–	–	–	–	2092.5 (8)	2092.64 (50)	2092.09 (10)
–	–	2140 (40)	–	–	–	2103.2 (30)	–	2103.3 (12)	–	–
2110.30 (21)	–	–	–	–	–	–	–	2110.6 (6)	2110.76 (50)	2110.26 (21)
2181.8 (7) ^Δ	–	–	–	–	–	–	–	–	2181.8 (7)	2181.8 (7) ^Δ
2188.0 (3)	–	–	–	–	–	–	–	–	2188.0 (7)	2188.0 (3)
2219.0 (4)	–	–	–	–	–	–	–	–	2219.91 (70)	2219.0 (4)
2336.54 (12)	2350 (50)	2330 (40)	2340 (40)	2300 (30)	2337 (2)	2336.5 (2)	2336.7 (10)	2336.8 (3)	2336.54 (30)	2336.49 (12)
–	–	–	–	–	2394 (4)	–	–	–	–	–
2497.19 (9)	–	–	–	–	2497 (4)	2497.7 (4)	2497.1 (10)	2497.5 (6)	2497.43 (40)	2497.14 (9)
2511.06 (6)	–	–	–	–	–	2512.0 (5)	2512.1 (10)	2512.5 (5)	2511.97 (50)	2511.01 (6)
2535.83 (7)	2550 (50)	2550 (50)	2550 (50)	–	2535 (2)	2535.9 (2)	2536.2 (10)	2536.2 (4)	2536.02 (30)	2535.78 (7)
–	–	–	–	–	2617 (6)	–	–	–	–	–
2696.66 (13)	–	2690 (50)	–	–	2694 (2)	2696.7 (2)	2696.7 (10)	2697.0 (4)	2696.57 (30)	2696.60 (13)
2716.47 (9)	–	–	–	–	–	2717.0 (5)	2716.8 (10)	2717.2 (5)	2716.89 (40)	2716.41 (9)
–	–	–	–	–	2767 (4)	–	–	–	–	–
2780.23 (21)	2750 (70)	2780 (60)	2770 (60)	–	2781 (4)	2780.1 (20)	2780.1 (10)	2780.6 (3)	2780.29 (40)	2780.16 (21)

2808.10 (8)	–	–	–	–	2805 (4)	2806.5 (4)	2807 (2)	2807.1 (6)	2806.62 (60)	2808.03 (8)
2857.9 (3)	–	–	–	–	2856 (4)	2857.4 (10)	2856 (2)	2857.8 (7)	2857.56 (80)	2857.8 (3)
–	–	–	–	–	2880 (4)	2882.1 (20) [‡]	–	–	–	–
2888.9 (4)	–	~ 2900	–	–	–	2891.1 (15)	2890 (2)	2889.5 (5)	2889.44 (80)	2888.8 (4)
3042.59 (8)	–	–	–	–	–	3044.9 (10)	3044 (2)	3044.0 (7)	3044.56 (80)	3042.51 (8)
–	–	–	–	–	–	3090.8 (2) [‡]	–	–	–	–
3101.4 (4)	–	–	3100 (80)	–	–	3100.9 (7)	3101 (2)	3100.3 (10)	3100.65 (80)	3101.3 (4)

[#] Determined from the nuclear level energies of 2001Ba27.

[§] Calculated by subtracting gamma recoil from gamma transition energy (E_{TP} (keV)).

^{*} Data derived from coincidence measurements.

[†] Energy determined from conversion-electron spectrum.

^Δ Unplaced in proposed decay scheme.

[‡] Inadequate data arising from insufficient statistics.

Emission Probabilities

Relative gamma-ray emission probabilities have been partially or fully determined in the measurements of 1967De08, 1969Bo15, 1970Ki06, 1971GiZS, 1974KI02 and 1982Gr10. While greater emphasis was placed on the measurements of 1971GiZS and 1974KI02, the equivalent studies of 1969Bo15, 1970Ki06 and 1982Gr10 proved to be significant in the weighted-mean analyses of specific gamma-ray emissions. All of the relative emission probabilities were suitably refined and quantified in terms of the emission probability of the 669.93-keV gamma ray (100.0 %).

The determination of both the N_γ(669.93 keV) / N_{β+} ratio and associated spectral correction by 1969Bo15 were of great importance in the calculation of the normalisation factor for the relative emission probabilities of the gamma rays and the resulting derivation of a consistent decay scheme:

$$N_{\gamma}(669.93 \text{ keV}) / N_{\beta+} = 9.14 (36) / 100 = 0.0914 (36)$$

In-flight annihilation and formation of ortho-positronium mitigates against immediate 100 % positron annihilation, and creates a shortfall in the resulting 511-keV detection of the order of 3.4 % that impacts in a similar manner with respect to the quantification of N_{β+}:

$$\text{correction to } N_{\gamma}(669.93 \text{ keV}) / N_{\beta+} \text{ constitutes a reduction of } 0.0914 (36) \times 0.034 = 0.00311 (12),$$

$$\text{and therefore } N_{\gamma}(669.93 \text{ keV}) / N_{\beta+} = 0.0914 (36) - 0.0031 = 0.0883 (35)$$

Published gamma-ray emission probabilities.

E _γ (keV)	P _γ									
	1959Ha10	1959Ri38	1961Cu02	1961Va08	1967De08 [#]	1969Bo15 [#]	1970Ki06 [#]	1971GiZS [*]	1974KI02 [*]	1982Gr10 [*]
244.40 (22)	–	–	–	–	–	–	–	–	0.065 (10)	–
364.74 (6)	–	–	–	–	–	0.02 (1) ⁰	–	–	0.14 (3)	–
443.70 (12)	–	–	–	–	–	–	–	–	0.20 (5)	–
450.14 (5)	–	–	–	–	–	0.27 (5)	0.2 (1)	2.7 (2)	2.88 (20)	–
475.91 (13)	–	–	–	–	–	–	–	–	0.07 (4)	–
511	–	200	200	intense	200	200	200	2188	–	–
515.45 (9)	–	–	–	–	–	–	–	–	0.26 (10) ⁰	–
534.32 (23)	–	–	–	–	–	–	–	–	0.06 (2)	–
584.98 (6)	–	–	–	–	–	–	–	0.22 (10)	0.40 (5)	–

624.34 (13)	–	–	–	–	–	–	–	0.18 (5)	0.17 (4)	–
669.93 (4)	intense	14.0 (8)	9.0 (6)	–	9.5 (5) (9.14 (36)	9.7 (2)	100	100	100
675.20 (9)	–	–	–	–	(–	–	–	0.18 (4)	–
683.74 (17)	–	–	–	detected	(0.041 (15)	–	–	0.05 (2)	–
742.23 (6)	–	–	–	–	–	0.079 (12)	0.07 (4)	0.76 (13)	0.82 (10)	–
754.56 (23)	–	–	–	–	–	–	–	0.33 (15)	0.08 (3)	–
765.37 (11)	–	–	–	–	–	–	–	–	0.08 (3)	–
810 (10) [†]	–	1.8 (1)	–	–	–	–	–	–	–	–
877.06 (6)	–	1.2 (1)	–	–	–	–	–	–	0.04 (2)	–
898.60 (7)	–	–	–	–	–	–	–	0.16 (6)	0.15 (3)	–
924.37 (13)	–	–	–	–	–	0.023 (8)	–	0.13 (6)	0.120 (24)	–
962.01 (3)	intense	10.0 (9)	6.7 (7)	detected	7.5 (5)	7.20 (20)	7.8 (2)	79.8 (32)	78.7 (40)	76.6 (19)
988.82 (9)	–	–	–	–	–	–	–	–	0.047 (13)	–
1050.89 (11)	–	–	–	–	–	–	–	–	0.054 (14)	–
1087 (2) [†]	–	–	–	–	–	0.029 (10)	–	–	–	–
1100 (20) [†]	–	0.70 (10)	–	–	–	–	–	–	–	–
1116 (2) [†]	–	–	–	–	detected	–	–	–	–	–
1123.66 (8)	–	–	–	–	–	0.14 (3)	0.16 (4)	1.3 (2)	1.35 (14)	–
1130.10 (10)	–	–	–	–	–	–	–	0.13 (4)	0.16 (3)	–
1149.65 (14)	–	–	–	–	–	0.027 (5)	–	0.21 (5)	0.23 (3)	–
1169.46 (10)	–	–	–	–	–	0.018 (4)	–	–	0.094 (20)	–
1189 (3) [†]	–	–	–	–	–	0.02 (1)	–	–	–	–
1209.06 (9)	–	–	–	–	–	0.014 (4)	0.03 (1)	0.17 (6)	0.15 (3)	–
1233.22 (22)	–	–	–	–	–	–	–	–	0.03 (1)	–
1270 (20) [†]	–	0.30 (5)	–	–	–	–	–	–	–	–
1326.75 (5)	–	0.41 (5)	< 0.05	detected	0.07 (4)	0.090 (11)	0.08 (2)	0.84 (6)	0.84 (5)	–
1342.97 (12)	–	–	–	–	–	–	0.06 (2)	–	0.03 (1)	–
1374.50 (12)	–	–	–	–	–	0.044 (8)	0.04 (2)	0.41 (6)	0.42 (3)	–
1389.69 (10)	–	–	–	–	0.17 (5) (0.16 (1) (0.17 (3) (0.44 (9)	0.52 (7)	–
1392.50 (9)	–	–	–	–	(((1.24 (12)	1.18 (18)	–
1412.14 (4)	modest	0.80 (7)	0.94 (15)	detected	0.95 (8)	0.80 (3)	0.93 (10)	9.3 (4)	9.08 (36)	8.85 (30)
1445.7 (3)	–	–	–	–	–	–	–	–	0.03 (1)	–
1481.32 (9)	–	–	–	–	–	–	–	–	0.02 (1)	–
1546.98 (5)	–	0.20 (5)	0.12 (3)	detected	0.14 (2)	0.13 (2)	0.17 (3)	1.6 (1)	1.49 (6)	–
1573.79 (8)	–	–	–	–	0.02 (1)	0.013 (4)	0.024 (10)	0.19 (7)	0.20 (2)	–
1666.59 (13)	–	0.60 (3)	–	–	–	–	–	–	0.017 (7)	–
1696.6 (1.0) ^Δ	–	–	–	–	0.03 (1)	–	–	–	0.024 (12)	–
1754.42 (9)	–	–	–	–	–	–	–	–	0.053 (12)	–

1816.9 (10) [†]	–	0.02 (1)	–	–	–	0.0028 (3)	–	–	–	–
1827.23 (10)	–	–	0.02 (1)?	–	0.02 (1)	–	–	0.11 (4)	0.051 (13)	–
1860.60 (6)	–	–	–	–	–	0.035 (3) (0.035 (20) (0.24 (5)	0.170 (24)	–
1865.87 (8)	–	–	–	–	–	((0.26 (5)	0.240 (26)	–
1926.9 (4)	–	0.08 (1)	–	–	–	0.006 (1)	–	0.082 (25)	0.070 (14)	–
2012.89 (11)	very weak	–	–	–	–	0.011 (2)	0.015 (5)	0.14 (3)	0.13 (2)	–
2026.70 (14)	–	0.12 (1)	0.14 (3)	–	–	0.060 (10)	0.08 (2)	0.78 (5)	0.68 (7)	–
2046.50 (10)	–	–	–	–	–	0.007 (1)	–	0.041 (14)	0.045 (13)	–
2062.41 (8)	–	–	–	–	0.04 (1)	0.032 (7)	0.04 (4)	0.41 (6)	0.42 (4)	–
2081.28 (22)	–	–	–	–	0.03 (2)	0.015 (4)	0.02 (1)	0.21 (5)	0.18 (2)	–
2092.09 (10)	–	–	–	–	–	–	–	0.057 (25)	0.03 (1)	–
2103.3 (12) [†]	–	0.06 (1)	–	–	–	0.003 (2)	–	0.069 (30)	–	–
2110.26 (21)	–	–	–	–	–	–	–	0.087 (22)	0.075 (15)	–
2181.8 (7) ^Δ	–	–	–	–	–	–	–	–	0.016 (10)	–
2188.0 (3)	–	–	–	–	–	–	–	–	0.02 (1)	–
2219.0 (4)	–	–	–	–	–	–	–	–	0.036 (10)	–
2336.49 (12)	weak	0.05 (1)	0.07 (2)	detected	0.080 (8)	0.095 (6)	0.09 (2)	0.99 (8)	0.91 (6)	–
2394 (4) [†]	–	–	–	–	0.010 (5)	–	–	–	–	–
2497.14 (9)	–	–	–	–	0.015 (5)	0.023 (2)	0.03 (2)	0.25 (3)	0.26 (3)	–
2511.01 (6)	–	–	–	–	–	0.013 (2)	0.009 (5)	0.12 (3)	0.12 (2)	–
2535.78 (7)	weak	0.08 (1)	0.06 (2)	–	0.07 (2)	0.077 (4)	0.09 (2)	0.80 (5)	0.81 (8)	–
2617 (6) [†]	–	–	–	–	0.005 (3)	–	–	–	–	–
2696.60 (13)	–	0.04 (1)	–	–	0.03 (1)	0.043 (8)	0.05 (2)	0.48 (4)	0.49 (5)	–
2716.41 (9)	–	–	–	–	–	0.012 (2)	0.016 (5)	0.140 (12)	0.16 (2)	–
2767 (4) [†]	–	–	–	–	0.009 (5)	–	–	–	–	–
2780.16 (21)	weak	0.03 (1)	0.04 (2)	–	0.010 (5)	0.018 (2)	0.018 (5)	0.180 (15)	0.19 (2)	–
2808.03 (8)	–	–	–	–	0.005 (2)	0.0027 (6)	0.005 (2)	0.039 (7)	0.05 (1)	–
2857.8 (3)	–	–	–	–	0.003 (1)	0.0033 (7)	0.004 (2)	0.032 (6)	0.04 (1)	–
2882.1 (20) [†]	–	–	–	–	0.0025 (10) (0.0010 (3) [‡]	–	–	–	–
2888.8 (4)	–	< 0.02	–	–	(0.0024 (5)	0.003 (1)	0.026 (3)	0.03 (1)	–
3042.51 (8)	–	–	–	–	< 0.004	0.0050 (10)	0.006 (1)	0.058 (10)	0.06 (1)	–
3090.8 (2) [†]	–	–	–	–	–	0.0004 (2) [‡]	–	–	–	–
3101.3 (4)	–	–	0.02 (1)	–	–	0.0008 (2)	0.002 (1)	0.009 (4)	0.007 (2)	–

[#] Emission probabilities expressed relative to P_γ for 511-keV annihilation radiation of 200 %.

^{*} Emission probabilities expressed relative to P_γ(669.93 keV) of 100 %.

^θ Data derived from coincidence measurements.

[†] Gamma emission not observed with confidence and regularity – not considered as identifiable with ⁶³Zn decay, and therefore not included in the proposed decay scheme.

^Δ Unplaced in proposed decay scheme.

[‡] Inadequate data arising from insufficient statistics.

Measured and recommended gamma-ray emission probabilities relative to P_γ(669.93 keV) of 100 %.

E _γ (keV)	P _γ ^{rel}						
	1967De08 [#]	1969Bo15 [#]	1970Ki06 [#]	1971GiZS	1974Kl02	1982Gr10	Recommended [†]
244.40 (22)	–	–	–	–	0.065 (10)	–	0.065 (10)
364.74 (6)	–	0.2 (1) ⁰	–	–	0.14 (3)	–	0.14 (3)
443.70 (12)	–	–	–	–	0.20 (5)	–	0.16 (5)
450.14 (5)	–	3.0 (6)	2 (1)	2.7 (2)	2.88 (20)	–	2.79 (20)
475.91 (13)	–	–	–	–	0.07 (4)	–	0.07 (4)
511	2105	2188	2062	2188	–	–	annihilation radiation
515.45 (9)	–	–	–	–	0.26 (10)⁰	–	0.26 (10)
534.32 (23)	–	–	–	–	0.06 (2)	–	0.06 (2)
584.98 (6)	–	–	–	0.22 (10)	0.40 (5)	–	0.40 (5)
624.34 (13)	–	–	–	0.18 (5)	0.17 (4)	–	0.14 (5)
669.93 (4)	100 (5) (100 (4)	100 (2)	100	100	100	100
675.20 (9)	(–	–	–	0.18 (4)	–	0.18 (4)
683.74 (17)	(0.45 (16)	–	–	0.05 (2)	–	0.05 (2)
742.23 (6)	–	0.86 (13)	0.7 (4)	0.76 (13)	0.82 (10)	–	0.82 (10)
754.56 (23)	–	–	–	0.33 (15)	0.08 (3)	–	0.19 (7)
765.37 (11)	–	–	–	–	0.08 (3)	–	0.08 (3)
877.06 (6)	–	–	–	–	0.04 (2)	–	0.04 (2)
898.60 (7)	–	–	–	0.16 (6)	0.15 (3)	–	0.11 (4)
924.37 (13)	–	0.25 (9)	–	0.13 (6)	0.120 (24)	–	0.121 (24)
962.01 (3)	79 (5)	78.8 (22)	80 (2)	79.8 (32)	78.7 (40)	76.6 (19)[§]	79.4 (20)
988.82 (9)	–	–	–	–	0.047 (13)	–	0.047 (13)
1050.89 (11)	–	–	–	–	0.054 (14)	–	0.054 (14)
1123.66 (8)	–	1.5 (3)	1.6 (4)	1.3 (2)	1.35 (14)	–	1.37 (14)
1130.10 (10)	–	–	–	0.13 (4)	0.16 (3)	–	0.16 (3)
1149.65 (14)	–	0.30 (6)	–	0.21 (5)	0.23 (3)	–	0.23 (3)
1169.46 (10)	–	0.20 (4)	–	–	0.094 (20)	–	0.094 (20)
1209.06 (9)	–	0.15 (4)	0.3 (1)	0.17 (6)	0.15 (3)	–	0.17 (4)
1233.22 (22)	–	–	–	–	0.03 (1)	–	0.03 (1)
1326.75 (5)	0.7 (4)	0.98 (12)	0.8 (2)	0.84 (6)	0.84 (5)	–	0.84 (5)
1342.97 (12)	–	–	0.6 (2)	–	0.03 (1)	–	0.03 (1)
1374.50 (12)	–	0.48 (9)	0.4 (2)	0.41 (6)	0.42 (3)	–	0.42 (3)
1389.69 (10)	1.8 (5) (1.8 (1) (1.8 (3) (0.44 (9)	0.52 (7)	–	0.52 (7)
1392.50 (9)	(((1.24 (12)	1.18 (18)	–	1.22 (12)
1412.14 (4)	10.0 (8)	8.8 (3)	9.6 (10)	9.3 (4)	9.08 (36)	8.85 (30)	9.0 (3)
1445.7 (3)	–	–	–	–	0.03 (1)	–	0.03 (1)
1481.32 (9)	–	–	–	–	0.02 (1)	–	0.02 (1)
1546.98 (5)	1.5 (2)	1.4 (2)	1.8 (3)	1.6 (1)	1.49 (6)	–	1.52 (6)
1573.79 (8)	0.2 (1)	0.14 (4)	0.25 (10)	0.19 (7)	0.20 (2)	–	0.19 (2)
1666.59 (13)	–	–	–	–	0.017 (7)	–	0.017 (7)
1696.6 (1.0) ^Δ	0.3 (1)	–	–	–	0.024 (12)	–	0.024 (12)
1754.42 (9)	–	–	–	–	0.053 (12)	–	0.053 (12)
1827.23 (10)	0.2 (1)	–	–	0.11 (4)	0.051 (13)	–	0.051 (13)
1860.60 (6)	–	0.38 (3) (0.36 (21) (0.24 (5)	0.170 (24)	–	0.14 (4)
1865.87 (8)	–	((0.26 (5)	0.240 (26)	–	0.244 (26)
1926.9 (4)	–	0.06 (1)	–	0.082 (25)	0.070 (14)	–	0.065 (14)
2012.89 (11)	–	0.12 (2)	0.15 (5)	0.14 (3)	0.13 (2)	–	0.13 (2)
2026.70 (14)	–	0.66 (11)	0.8 (2)	0.78 (5)	0.68 (7)	–	0.73 (5)
2046.50 (10)	–	0.08 (1)	–	0.041 (14)	0.045 (13)	–	0.043 (13)
2062.41 (8)	0.4 (1)	0.35 (8)[§]	0.4 (4)	0.41 (6)	0.42 (4)	–	0.42 (4)
2081.28 (22)	0.3 (2)[§]	0.16 (4)	0.2 (1)	0.21 (5)	0.18 (2)	–	0.18 (2)
2092.09 (10)	–	–	–	0.057 (25)	0.03 (1)	–	0.06 (3)
2110.26 (21)	–	–	–	0.087 (22)	0.075 (15)	–	0.079 (15)
2181.8 (7) ^Δ	–	–	–	–	0.016 (10)	–	0.016 (10)
2188.0 (3)	–	–	–	–	0.02 (1)	–	0.02 (1)
2219.0 (4)	–	–	–	–	0.036 (10)	–	0.036 (10)
2336.49 (12)	0.84 (8)	1.04 (7)	0.9 (2)	0.99 (8)	0.91 (6)	–	0.94 (6)
2497.14 (9)	0.16 (5)	0.25 (2)	0.3 (2)	0.25 (3)	0.26 (3)	–	0.25 (2)

2511.01 (6)	–	0.14 (2)	0.09 (5)	0.12 (3)	0.12 (2)	–	0.13 (2)
2535.78 (7)	0.7 (2)	0.84 (4)	0.9 (2)	0.80 (5)	0.81 (8)	–	0.82 (4)
2696.60 (13)	0.3 (1)	0.47 (9)	0.5 (2)	0.48 (4)	0.49 (5)	–	0.48 (4)
2716.41 (9)	–	0.13 (2)	0.16 (5)	0.140 (12)	0.16 (2)	–	0.146 (12)
2780.16 (21)	0.11 (6)	0.20 (2)	0.19 (5)	0.180 (15)	0.19 (2)	–	0.188 (15)
2808.03 (8)	0.05 (2)	0.030 (7)	0.05 (2)	0.039 (7)	0.05 (1)	–	0.044 (7)
2857.8 (3)	0.03 (1)	0.036 (8)	0.04 (2)	0.032 (6)	0.04 (1)	–	0.034 (6)
2888.8 (4)	0.026 (10)	0.026 (5)	0.03 (1)	0.026 (3)	0.03 (1)	–	0.026 (3)
3042.51 (8)	< 0.04	0.055 (11)	0.06 (1)	0.058 (10)	0.06 (1)	–	0.058 (10)
3101.3 (4)	–	0.009 (2)	0.02 (1)	0.009 (4)	0.007 (2)	–	0.008 (2)

[#] Emission probabilities adjusted in order to be expressed relative to $P_{\gamma}(669.93 \text{ keV})$ of 100 %.

[†] Weighted mean values adopted when judged appropriate.

⁰ Data derived from coincidence measurements.

[§] Defined as outlier, and therefore not adopted for consideration in the analytical procedures.

^Δ Unplaced in proposed decay scheme.

Weighted-mean analyses of the measured relative emission probabilities were carried out when judged appropriate. However, considerably more detail can be found within the gamma-ray measurements of Giesler (1971GiZS) and, in particular, Klaasse and Goudsmit (1974Kl02). Under these circumstances, the data of 1974Kl02 and sometimes 1971GiZS were directly adopted in preference to any semi-artificial weighting procedure. The relative emission probabilities of the 443.70-, 624.34-, 754.56-, 898.60-, 1860.60- and 2092.09-keV gamma rays were subsequently adjusted from 0.20 (5), 0.17 (4), 0.08 (3), 0.15 (3), 0.183 (24) and 0.03 (1) to 0.16 (5), 0.14 (5), 0.19 (7), 0.11 (4), 0.14 (4) and 0.06 (3), respectively, to achieve population-depopulation balances for the 1326.76-, 1860.63- and 2092.13-keV nuclear levels (shown as red in the table). Both the 1696.6- and 2181.8-keV gamma rays observed only by Klaasse and Goudsmit (1974Kl02) remain within the recommended data set, although they could not be placed in the proposed decay scheme.

The normalisation factor for the relative emission probabilities of the gamma rays can be directly determined from the spectral measurements of $P_{\gamma}^{abs}(669.93 \text{ keV})$ by 1961Cu02, 1967De08, 1969Bo15 and 1970Ki06.

	$P_{\gamma}^{abs}(669.93 \text{ keV})$	Comments
1961Cu02	9.0 (6)	3" x 3" NaI(Tl) detector, 8 % resolution
1967De08	9.5 (5)	5 x 20 mm ² Ge(Li) detector; defined as 688 (1) keV (presumably a misprint (668 keV)?)
1969Bo15	9.14 (36)	0.7 x 4.4 and 0.7 x 8.5 cm ² Ge(Li) detectors; Ge(Li)-NaI(Tl) coincidence system
1970Ki06	9.7 (2)	38 cm ³ Ge(Li) detector (FWHM of 3.5 keV for 1332 keV ⁶⁰ Co)

Inadequate spectral resolution is a significant issue in the measurements involving the NaI(Tl) detector system (1961Cu02), while the gamma-ray spectrum shown in detail by 1967De08 would appear to pose problems with respect to accurate peak-area analyses in the vicinity of 650 to 750 keV. Both 1969Bo15 and 1970Ki06 exhibit improved resolution over the gamma-ray energy range of 650 to 750 keV; however, these particular studies are not necessarily equivalent, and their reported $P_{\gamma}^{abs}(669.93 \text{ keV})$ values do not overlap and are effectively close to being discrepant. Therefore, as described in more detail below, the normalisation factor for the relative emission probabilities of the gamma rays has been preferably determined from the $N_{\gamma}(669.93 \text{ keV})/N_{\beta+}$ ratio (1969Bo15).

Multipolarities, and Internal Conversion and Internal-Pair Coefficients

The nuclear level scheme specified by Bai and Huo Junde has been used to define the multipolarities of the gamma transitions on the basis of known spins and parities (2001Ba27). Limited studies of the angular correlation coefficients, B(E2) values and mixing ratios support the proposed transition types: (M1 + E2) for the 364.74-, 450.14-, 669.93-, 962.01-, 1412.14- and 1546.98-keV gamma rays (1972Ro21, 1980Ku08 and 1998Si25). Other (M1 + E2) gamma transitions judged to be of reasonable importance in the derivation of the decay scheme were arbitrarily assigned multipolarities of (50 % M1 + 50 % E2) – i.e. gamma rays with energies of 443.70, 1123.66 and 1392.50 keV.

Recommended internal conversion coefficients have been determined from the frozen orbital approximation of Kibédi *et al.* (2008Ki07), based on the theoretical model of Band *et al.* (2002Ba85, 2002Ra45). A significant number of gamma transitions undergo decay via internal-pair formation, and the coefficient for this process has also been quantified in a few cases from the tabulations of 2008Ki07.

Gamma-ray emissions: multiplicities, and theoretical internal-conversion and internal-pair formation coefficients (frozen orbital approximation).

E_γ (keV)	Multiplicity	α_K	α_L	α_{M+}	$\alpha_{totalICC}$	α_{IPF}	α_{total}
244.40 (22)	(E2)	0.019 0 (3)	0.001 98 (3)	0.000 32	0.021 3 (3)	—	0.021 3 (3)
364.74 (6)	99.64% M1 + 0.36% E2 $\delta = -0.060(5)$ (1980Ku08)	0.001 84 (3)	0.000 184 (3)	0.000 026	0.002 05 (3)	—	0.002 05 (3)
443.70 (12)	(50% M1 + 50% E2) $\delta = 1.0$ (2)	0.001 77 (14)	0.000 179 (14)	0.000 031	0.001 98 (16)	—	0.001 98 (16)
450.14 (5)	98.7% M1 + 1.3% E2 $\delta = 0.115$ (10) (1980Ku08)	0.001 14 (4)	0.000 113 (5)	0.000 017	0.001 27 (5)	—	0.001 27 (5)
475.91 (13)	M1 + E2	—	—	—	—	—	—
515.45 (9)	(M1 + E2)	—	—	—	—	—	—
534.32 (23)	(M1 + E2)	—	—	—	—	—	—
584.98 (6)	M1 + E2	—	—	—	—	—	—
624.34 (13)	(E2)	—	—	—	—	—	—
669.93 (4)	98.8% M1 + 1.2% E2 $\delta = 0.11$ (2) (1980Ku08)	0.000 466 (7)	0.000 046 2 (7)	0.000 006 8	0.000 519 (8)	—	0.000 519 (8)
675.20 (9)	(M1 + E2)	—	—	—	—	—	—
683.74 (17)	M1 + E2	—	—	—	—	—	—
742.23 (6)	E2	0.000 512 (8)	0.000 051 1 (8)	0.000 007 4	0.000 571 (8)	—	0.000 571 (8)
754.56 (23)	(M1 + E2)	—	—	—	—	—	—
765.37 (11)	M1 + E2	—	—	—	—	—	—
877.06 (6)	M1 + E2	—	—	—	—	—	—
898.60 (7)	M1 + E2	—	—	—	—	—	—
924.37 (13)	M1 + E2	—	—	—	—	—	—
962.01 (3)	81.3% M1 + 18.7% E2 $\delta = -0.48$ (2) (1980Ku08)	0.000 226 (4)	0.000 022 3 (4)	0.000 003 1	0.000 251 (4)	—	0.000 251 (4)
988.82 (9)	(M1 + E2)	—	—	—	—	—	—
1050.89 (11)	M1 + E2	—	—	—	—	—	—
1123.66 (8)	(50% M1 + 50% E2) $\delta = 1.0$ (2)	0.000 171 (4)	0.000 016 9 (4)	0.000 002 5	0.000 190 (4)	0.000 001 41 (6)	0.000 192 (4)
1130.10 (10)	M1 + E2	—	—	—	—	—	—
1149.65 (14)	M1 + E2	—	—	—	—	—	—
1169.46 (10)	M1 + E2	—	—	—	—	—	—
1209.06 (9)	(M1 + E2)	—	—	—	—	—	—
1233.22 (22)	M1 + E2	—	—	—	—	—	—
1326.75 (5)	E2	0.000 126 8 (18)	0.000 012 51 (18)	0.000 001 81	0.000 141 1 (18)	0.000 034 6 (5)	0.000 175 7 (25)
1342.97 (12)	M1 + E2	—	—	—	—	—	—
1374.50 (12)	M1 + E2	—	—	—	—	—	—
1389.69 (10)	(E2)	—	—	—	—	—	—

Comments on evaluation

⁶³Zn

1392.50 (9)	(50% M1 + 50% E2) δ = 1.0 (2)	0.000 109 8 (19)	0.000 010 80 (19)	0.000 001 57	0.000 122 2 (19)	0.000 045 1 (16)	0.000 167 (4)
1412.14 (4)	63.4% M1 + 36.6% E2 δ = 0.76 (7) (1980Ku08)	0.000 105 5 (16)	0.000 010 38 (15)	0.000 001 50	0.000 117 4 (16)	0.000 048 2 (10)	0.000 166 (3)
1445.7 (3)	(E2)	—	—	—	—	—	—
1481.32 (9)	E2	—	—	—	—	—	—
1546.98 (5)	86.8% M1 + 13.2% E2 δ = 0.39 (7) (1980Ku08)	0.000 087 0 (13)	0.000 008 54 (13)	0.000 001 24	0.000 096 8 (13)	0.000 084 4 (16)	0.000 181 (3)
1573.79 (8)	(M1 + E2)	—	—	—	—	—	—
1666.59 (13)	E2	—	—	—	—	—	—
1754.42 (9)	M1 + E2	—	—	—	—	—	—
1827.23 (10)	(M1 + E2)	—	—	—	—	—	—
1860.60 (6)	E2	0.000 064 6 (9)	0.000 006 35 (9)	0.000 000 98	0.000 071 9 (9)	0.000 244 (4)	0.000 316 (5)
1865.87 (8)	(E2)	0.000 064 3 (9)	0.000 006 31 (9)	0.000 000 92	0.000 071 5 (9)	0.000 247 (4)	0.000 319 (5)
1926.9 (4)	(E2)	—	—	—	—	—	—
2012.89 (11)	M1 + E2	—	—	—	—	—	—
2026.70 (14)	M1 + E2	—	—	—	—	—	—
2046.50 (10)	M1 + E2	—	—	—	—	—	—
2062.41 (8)	(M1 + E2)	—	—	—	—	—	—
2081.28 (22)	(M1 + E2)	—	—	—	—	—	—
2092.09 (10)	E2	—	—	—	—	—	—
2110.26 (21)	M1 + E2	—	—	—	—	—	—
2188.0 (3)	M1 + E2	—	—	—	—	—	—
2219.0 (4)	M1 + E2	—	—	—	—	—	—
2336.49 (12)	M1 + E2	—	—	—	—	—	—
2497.14 (9)	(M1 + E2)	—	—	—	—	—	—
2511.01 (6)	[M1 + E2]	—	—	—	—	—	—
2535.78 (7)	(M1 + E2)	—	—	—	—	—	—
2696.60 (13)	M1 + E2	—	—	—	—	—	—
2716.41 (9)	M1 + E2	—	—	—	—	—	—
2780.16 (21)	M1 + E2	—	—	—	—	—	—
2808.03 (8)	M1 + E2	—	—	—	—	—	—
2857.8 (3)	M1 + E2	—	—	—	—	—	—
2888.8 (4)	M1 + E2	—	—	—	—	—	—
3042.51 (8)	M1 + E2	—	—	—	—	—	—
3101.3 (4)	M1 + E2	—	—	—	—	—	—

The normalisation factor for the relative emission probabilities of the gamma rays has been preferentially determined from the $N_{\gamma}(669.93 \text{ keV})/N_{\beta^+}$ ratio:

(a) $N_{\gamma}(669.93 \text{ keV}) / N_{\beta^+}$ ratio and associated spectral correction (1969Bo15)

$$N_{\gamma}(669.93 \text{ keV}) / N_{\beta^+} = 0.0914 (36) - 0.0031 = 0.0883 (35),$$

(b) $P_{EC}(\text{total})$ determined from balances of the relative transition probabilities calculated from the relative emission probabilities and total conversion coefficients of the γ rays populating-depopulating the nuclear levels of ⁶³Cu, and

(c) theoretical capture-to-positron ratios (ϵ/β^+) for the EC/ β^+ branches to the relevant nuclear levels of ⁶³Cu (1971Go40).

Knowing the relative $P_{EC}(\text{total})$ that populates each nuclear level by a combination of EC and β^+ decay, and their theoretical capture-to-positron ratios, relative P_{β^+} values can be determined:

Relative positron emission probabilities calculated from $P_{EC}(\text{total})$ and ϵ/β^+ ratios.

	Nuclear level energy (keV)	Calculated relative $P_{EC}(\text{total})$	ϵ/β^+ (1971Go40)	Relative P_{β^+}
9	2081.32 (22)	0.43 (8)	79.91	0.005
8	2062.45 (8)	1.90 (16)	61.55	0.030
7	2012.92 (11)	0.164 (3)	33.38	0.005
5	1547.00 (5)	1.239 (136)	1.459	0.504 (55)
4	1412.16 (4)	11.089 (400)	0.8551	5.978 (216)
2	962.02 (3)	75.018 (2012)	0.2397	60.513 (1623)
1	669.93 (4)	96.567 (184)	0.1320	85.307 (163)
0	0.0	X \rightarrow to be calculated	0.0465	X / 1.0465

Total relative positron emission probability ($P_{\beta^+}^{rel}$)

$$= \sum \text{relative } P_{\beta^+} = [(X / 1.0465) + 152.342 (1646)]$$

where X is the relative total EC transition probability ($P_{EC}(\text{total})$) directly to the ground state of ⁶³Cu.

Adopting the $N_{\gamma}(669.93 \text{ keV}) / N_{\beta^+}$ ratio and associated spectral correction (1969Bo15)

$$N_{\gamma}(669.93 \text{ keV}) / N_{\beta^+} = 0.0914 (36) - 0.0031 = 0.0883 (35),$$

this equation can be suitably re-arranged to accommodate the known relative transition probabilities and determine the value of X:

$$N_{\gamma}(669.93 \text{ keV}) = 100 \times F$$

$$N_{\beta^+} = [(X / 1.0465) + 152.342 (1646)] \times F$$

where F is the normalisation factor for the relative emission probabilities of the gamma rays. Thus, making these substitutions in the equation that quantifies the $N_{\gamma}(669.93 \text{ keV}) / N_{\beta^+}$ ratio:

$$\frac{100 \times F}{[(X / 1.0465) + 152.342 (1646)] \times F} = 0.0883 (35)$$

$$\frac{100}{0.0883 (35)} = \frac{X}{1.0465} + 152.342 (1646)$$

which can be re-arranged to give
$$X = [1132.503 (44890) - 152.342 (1646)] \times 1.0465$$

$$= 980.161 (44920) \times 1.0465 = 1025.738 (47009)$$

Direct γ population of the ground state of ⁶³Cu was calculated to sum to a relative transition probability of 194.864 (2027), and since the total population of this level can be expressed by the following equation:

$$\sum (\gamma \text{ transition probabilities direct to ground state}) + \text{total EC}/\beta^+ \text{ direct to ground state} = 100$$

$$194.864 (2027) \times F + 1025.738 (47009) \times F = 100$$

$$F = 100 / 1220.602 (47053) = 0.0819 (32)$$

When the relative EC/ β^+ transition probabilities are also summed to 100 %, a normalisation factor is obtained in full agreement with the above value:

$$\sum (\text{total EC}/\beta^+ \text{ transition probabilities populating all nuclear levels of } ^{63}\text{Cu}) = 100$$

$$194.93 (208) \times F + 1025.738 (47009) \times F = 100$$

$$F = 100 / 1220.668 (47055) = 0.0819 (32)$$

Therefore, a value of 0.082 (3) was adopted as the normalisation factor in order to determine the absolute transition and emission probabilities of the EC, β^+ particles and γ rays from their relative transition and emission probabilities.

EC/ β^+ Transitions

Energies

All EC/ β^+ energies were derived from the structural details of the proposed decay scheme. The nuclear level energies of 2001Ba27 and evaluated Q_{EC} -value of 3366.2 (15) keV (2012Wa38) were used to determine the recommended energies and uncertainties of the EC transitions and β^+ emissions.

Transition and Emission Probabilities

Total EC transition probabilities were derived for the population-depopulation imbalances of the relative emission probabilities of the gamma rays, their theoretical internal conversion and internal-pair formation coefficients, and a normalisation factor of 0.082 (3) for the gamma-ray emissions as calculated above. Component EC and β^+ transition and emission probabilities were determined from EC/ β^+ ratios (1971Go40), and $\log ft$ values and average E_{β^+} energies were derived by means of the LOGFT code. Fractional EC probabilities P_K , P_L , P_M and P_N were calculated by means of the EC-CAPTURE code (1998Sc28) as developed from the data tabulations of 1995ScZY.

EC decay to the ground state of Cu-63 can be determined by two routes: (a) from the normalisation factor and relative emission probabilities of all the gamma rays that populate the ground state directly, and (b) from the normalisation factor and calculated relative transition probabilities of all EC decays to the excited nuclear levels of Cu-63.

- (a) Normalisation factor and relative emission probabilities of gamma rays populating the ground state of Cu-63 directly

$$\begin{aligned} P_{\text{EC decay direct to ground state}} &= 100 \% - \sum (P_{\gamma} \text{ populating ground state of Cu-63}) \\ &= 100 - [194.864 (2027) \times F] \\ &= 100 - [194.864 (2027) \times 0.0819 (32)] \\ &= 100 - 16.0 (6) = 84.0 (6) \% \end{aligned}$$

- (b) Normalisation factor and relative transition probabilities of EC decay to excited nuclear levels of Cu-63

$$\begin{aligned} P_{\text{EC direct decay to ground state}} &= 100 \% - \sum (P_{\text{EC}} \text{ populating excited states of Cu-63}) \\ &= 100 - [194.93 (208) \times F] \\ &= 100 - [194.93 (208) \times 0.0819 (32)] \\ &= 100 - 16.0 (6) = 84.0 (6) \% \end{aligned}$$

Therefore, a value and uncertainty of 84.0 (0.6) % was adopted for the total P_{EC} transition directly to the ground state of Cu-63.

The emission probability for the 511-keV annihilation radiation was derived from the total positron emission probability of 92.8 (6) % as determined during the evaluation of the EC/ β^+ decay mode:

$$P_{\gamma^{\pm}} = 2 \times 92.8 (6) = 185.6 (0.9) \%$$

A consistent decay scheme was derived that consists of 20EC/8 β^+ transitions and 64 gamma-ray emissions, of which 84.0 (6) % of the EC/ β^+ decay occurs directly to the ground state of ⁶³Cu. Two observed gamma-ray emissions could not be placed in the proposed decay scheme, and there is also substantial 511-keV positron-annihilation radiation.

Recommended energies and transition probabilities of the EC/ β^+ decay of ⁶³Zn.

E_{EC} (keV)*	E_{β^+} (keV)	Av. E_{β^+} (keV)	P_{EC} (total)	ϵ/β^+ (theory)	P_{EC}	P_{β^+}	⁶³ Zn	⁶³ Cu	transition type	log ft	P_K	P_L	P_M	P_N
EC _{0,22} 264.8 (16)	–	–	0.0007 (2)	–	0.0007 (2)	–	3/2 –	1/2 –, 3/2 –	allowed	6.89	0.8802 (16)	0.1020 (13)	0.0168 (5)	0.0011 (1)
EC _{0,21} 323.6 (15)	–	–	0.0048 (8)	–	0.0048 (8)	–	3/2 –	(5/2 –)	(allowed)	6.24	0.8814 (16)	0.1010 (13)	0.0166 (5)	0.0010 (1)
EC _{0,20} 477.3 (16)	–	–	0.0104 (14)	–	0.0104 (14)	–	3/2 –	1/2 –, 3/2, 5/2 –	allowed	6.24	0.8831 (16)	0.0996 (13)	0.0163 (5)	0.0010 (1)
EC _{0,19} 508.3 (15)	–	–	0.0069 (12)	–	0.0069 (12)	–	3/2 –	(1/2 –, 3/2 –)	(allowed)	6.48	0.8833 (16)	0.0994 (13)	0.0163 (5)	0.0010 (1)
EC _{0,18} 558.1 (15)	–	–	0.0052 (10)	–	0.0052 (10)	–	3/2 –	3/2 –	allowed	6.68	0.8836 (16)	0.0992 (13)	0.0162 (5)	0.0010 (1)
EC _{0,17} 586.0 (15)	–	–	0.0298 (21)	–	0.0298 (21)	–	3/2 –	(1/2 –, 3/2 –)	(allowed)	5.97	0.8837 (16)	0.0991 (13)	0.0162 (5)	0.0010 (1)
EC _{0,16} 649.7 (15)	–	–	0.082 (7)	–	0.082 (7)	–	3/2 –	3/2 –, 5/2 –	allowed	5.62	0.8840 (16)	0.0988 (13)	0.0162 (5)	0.0010 (1)
EC _{0,15} 669.5 (15)	–	–	0.122 (6)	–	0.122 (6)	–	3/2 –	1/2 –, 3/2 –	allowed	5.47	0.8841 (16)	0.0988 (13)	0.0161 (5)	0.0010 (1)
EC _{0,14} 830.4 (15)	–	–	0.261 (14)	–	0.261 (14)	–	3/2 –	(5/2 –)	(allowed)	5.33	0.8846 (16)	0.0984 (13)	0.0161 (5)	0.0010 (1)
EC _{0,13} 855.1 (15)	–	–	0.011 (2)	–	0.011 (2)	–	3/2 –	1/2, 3/2, 5/2	[allowed]	6.73	0.8846 (16)	0.0983 (13)	0.0161 (5)	0.0010 (1)
EC _{0,12} 869.0 (15)	–	–	0.0247 (20)	–	0.0247 (20)	–	3/2 –	(3/2 –)	(allowed)	6.40	0.8846 (16)	0.0983 (13)	0.0161 (5)	0.0010 (1)
EC _{0,11} 1029.7 (15)	–	–	0.141 (9)	–	0.141 (9)	–	3/2 –	5/2 –	allowed	5.79	0.8849 (16)	0.0980 (13)	0.0160 (5)	0.0010 (1)
EC _{0,9} 1284.9 (15)	262.9 (15)	115.1 (6)	0.035 (7)	79.91	0.035 (7)	0.00043 (9)	3/2 –	5/2 (–)	(allowed)	6.59	0.8853 (16)	0.0978 (13)	0.0160 (5)	0.0010 (1)
EC _{0,8} 1303.8 (15)	281.8 (15)	123.0 (6)	0.155 (13)	61.55	0.153 (13)	0.0025 (2)	3/2 –	(1/2) –	(allowed)	5.96	0.8853 (16)	0.0978 (13)	0.0160 (5)	0.0010 (1)
EC _{0,7} 1353.3 (15)	331.3 (15)	143.6 (6)	0.0134 (3)	33.38	0.0130 (3)	0.00039 (2)	3/2 –	3/2 –	allowed	7.06	0.8853 (16)	0.0977 (13)	0.0160 (5)	0.0010 (1)
EC _{0,5} 1819.2 (15)	797.2 (15)	341.0 (7)	0.102 (11)	1.459	0.060 (7)	0.042 (4)	3/2 –	3/2 –	allowed	6.65	0.8856 (16)	0.0975 (13)	0.0159 (5)	0.0010 (1)
EC _{0,4} 1954.0 (15)	932.0 (15)	399.7 (7)	0.91 (3)	0.8551	0.42 (2)	0.49 (2)	3/2 –	5/2 –	allowed	5.87	0.8857 (16)	0.0974 (13)	0.0159 (5)	0.0010 (1)
EC _{0,2} 2404.2 (15)	1382.2 (15)	599.5 (7)	6.15 (16)	0.2397	1.19 (3)	4.96 (13)	3/2 –	5/2 –	allowed	5.601	0.8858 (16)	0.0973 (13)	0.0159 (5)	0.0010 (1)
EC _{0,1} 2696.3 (15)	1674.3 (15)	732.0 (7)	7.92 (2)	0.1320	0.92 (1)	7.00 (2)	3/2 –	1/2 –	allowed	5.813	0.8859 (16)	0.0972 (13)	0.0159 (5)	0.0010 (1)
EC _{0,0} 3366.2 (15)	2344.2 (15)	1041.9 (7)	84.0 (6)	0.0465	3.75 (5)	80.3 (6)	3/2 –	3/2 –	allowed	5.397	0.8860 (16)	0.0971 (13)	0.0158 (5)	0.0010 (1)
			Σ 99.985			Σ 92.8 (6)								

* Determined from the nuclear level energies of 2001Ba27 and Q-value of 3366.2(15) keV (2012Wa38).

Atomic Data

The x-ray and Auger-electron data have been calculated using the evaluated gamma-ray data, and atomic data from 1996Sc06, 1998ScZM and 1999ScZX. Both the x-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.02, 28 February 2012), as described in 2000Sc47. This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

K and L X-ray energies and emission probabilities of ⁶³Zn.

			Energy (keV)	Photons per 100 disint.	Relative probability
XL	(Cu)		0.811 – 1.022	0.095 8 (16)	5.68
	XL ₁	(Cu)	0.811	0.003 11 (8)	
	XL _α	(Cu)	0.929 – 0.930	0.054 7 (13)	
	XL _η	(Cu)	0.831	0.001 86 (5)	
	XL _β	(Cu)	0.949 – 1.022	0.036 1 (10)	
	XL _γ	(Cu)	0.952	0.000 052 (8)	
XK _α	XK _{α2}	(Cu)	8.027 92 (1)	0.865 (12)	51.3
	XK _{α1}	(Cu)	8.047 87 (1)	1.686 (22)	100
XK' _{β1}	XK _{β3}	(Cu)	8.905 41 (4)	0.355 (6)	21.1
	XK _{β1} "	(Cu)	8.905 39 (6)		
	XK _{β5}	(Cu)	8.977 1 (2)		

Auger-electron energies and emission probabilities of ⁶³Zn.

		Energy (keV)	Electrons per 100 disint.	Relative probability
e _{AK}	(Cu)		3.50 (5)	100
	KLL	6.731 – 7.059	2.69 (4)	
	KLX	7.746 – 8.064	0.749 (13)	
	KXY	8.739 – 8.982	0.052 0 (13)	
e _{AL}	(Cu)	0.68 – 0.80	9.30 (9)	346

Cu: ω_K = 0.454 (4); ω_L = 0.0097 (4); n_{KL} = 1.357 (4) were taken from 1996Sc06.

Electron energies were determined from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

Data Consistency

A Q_{EC}-value of 3366.2 (15) keV has been adopted from the atomic mass evaluation of Wang *et al.* (2012Wa38) while in the course of formulating the decay scheme of ⁶³Zn. This value has subsequently been compared with the Q-value calculated by summing the contributions of the individual emissions to the ⁶³Zn EC-decay process (i.e. EC/β⁺, electron, γ, etc.):

$$\text{calculated Q-value} = \sum (E_i \times P_i) = 3367 (14) \text{ keV}$$

Percentage deviation from the Q-value of Wang *et al.* is – 0.024 (42) %, which supports the derivation of a consistent decay scheme with a significant variant.

References

- 1938St05 C.V. Strain, Proton Induced Radioactivities, II. Nickel and Copper Targets, Phys. Rev. 54 (1938) 1021-1025. [Half-life]
- 1939Bo05 W. Bothe, W. Gentner, Die Wellenlängenabhängigkeit der Kernphotoeffekte; mit Anhang: Die radioaktiven Isotope des Selens, Z. Phys. 112 (1939) 45-64. [Half-life]
- 1939De01 L.A. Delsasso, L.N. Ridenour, R. Sherr, M.G. White, Artificial Radioactivity Produced by Protons, Phys. Rev. 55 (1939) 113-121. [Half-life]
- 1947Hu20 O. Huber, H. Medicus, P. Preiswerk, R. Steffen, Die radioaktive Umwandlung des Zn⁶³, Helv. Phys. Acta 20 (1947) 495-518. [Half-life]
- 1948Wa13 H. Wäffler, O. Hirzel, Relative Wirkungsquerschnitte für den (γ,n)-Prozess der Lithium-Gamma-Strahlung (Quantenergie hv = 17.5 MeV), Helv. Phys. Acta 21 (1948) 200-203. [Half-life]
- 1959Ha10 R.W. Hayward, E. Farrelly-Pessoa, D.D. Hoppes, R. van Lieshout, Gamme Radiation from ⁶³Zn, Nuovo Cimento 11 (1959) 153-155. [E_γ]
- 1959Ri38 R.A. Ricci, R.K. Girgis, R. van Lieshout, The Gamma Ray Spectrum of ⁶³Zn, Nuovo Cimento 11 (1959) 156-166. [Half-life, E_γ, P_γ]
- 1960Pr05 I.L. Preiss, R.W. Fink, New Isotopes of Cobalt : Activation Cross-sections of Nickel, Cobalt and Zinc for 14.8 MeV Neutrons, Nucl. Phys. 15 (1960) 326-336. [Half-life]
- 1961Ra06 L.A. Rayburn, 14.4-MeV (n,2n) Cross Sections, Phys. Rev. 122 (1961) 168-171. [Half-life]
- 1961Cu02 J.B. Cumming, N.T. Porile, Decay of Zn⁶³, Phys. Rev. 122 (1961) 1267-1274. [Half-life, E_γ, P_γ, P_{EC}]
- 1961Va08 S.S. Vasil'ev, No Hsieng Ch'ang, L.Ya. Shavtvalov, Investigation of the radiations from Zn⁶³, Sov. Phys. JETP 13 (1961) 331-332. [Half-life, E_γ, E_{β+}, P_{β+}]
- 1965Pa18 A. Paulsen, H. Liskien, Anregungsfunktionen der Reaktionen Ni⁵⁸(n,2n)Ni⁵⁷, Cu⁶⁵(n,2n)Cu⁶⁴ und Zn⁶⁴(n,2n)Zn⁶³ im Energiebereich von 12.6 bis 19.6 MeV, Nukleonik 7 (1965) 117-120. [Half-life]
- 1967De08 D. De Frenne, M. Dorikens, L. Dorikens-Vanpraet, J. Demuynck, The High-Energy Decay of ⁶³Zn, Nucl. Phys. A103 (1967) 203-208. [E_γ, P_γ]
- 1968Go10 J.D. Goss, F.L. Riffle, D.R. Parsignault, J.C. Harris, Precise Measurements of the Half-lives of ¹⁹Ne, ²³Mg, ^{26m}Al, ²⁷Si and ⁶³Zn, Nucl. Phys. A115 (1968) 113-119. [Half-life]
- 1969Bo11 M. Bormann, B. Lammers, Excitation Functions of (n,p) and (n,2n) Reactions for Some Isotopes of K, Mn, Zn and Cu, Nucl. Phys. A130 (1969) 195-208. [Half-life]
- 1969Bo15 I. Borchert, Der Zerfall des ⁶³Zn, Z. Phys. 223 (1969) 473-486. [Half-life, E_γ, P_γ]
- 1970Ki06 A. Kiuru, P. Holmberg, Decay of ⁶³Zn, Z. Phys. 233 (1970) 146-153. [E_γ, P_γ]
- 1971GiZS G.C. Giesler, Nuclear Spectroscopic Studies of Some Short-lived and Neutron Deficient Ga and Zn Isotopes, PhD thesis, Michigan State University (1971). [E_γ, P_γ, EC/β+ to ground state]

- 1971Gi03 G.C. Giesler, K.L. Kosanke, R.A. Warner, W.C. McHarris, Ge(Li)-Ge(Li) Sum Coincidence: A Bonus From Ge(Li)-Ge(Li) Megachannel Coincidence Experiments, Nucl. Instrum. Methods 93 (1971) 211-220. [γ - γ coincidence]
- 1971Go40 N.B. Gove, M.J. Martin, Log-*f* Tables for Beta Decay, Nucl. Data Tables 10 (1971) 205-237. [ϵ/β^+ ratios]
- 1972Cr02 D.F. Crisler, H.B. Eldridge, R. Kunselman, C.S. Zaidins, Reactions Induced by He³ Ions on Zn⁶⁴, Phys. Rev. C5 (1972) 419-430. [Half-life]
- 1972Ro21 R.L. Robinson, Z.W. Grabowski, Angular Correlations of γ Rays from Coulomb Excited ^{63,65}Cu, Nucl. Phys. A191 (1972) 225-232. [γ - γ angular correlation, δ]
- 1974Co16 R. Collé, R. Kishore, J.B. Cumming, Excitation Functions for (p,n) Reactions to 25 MeV on ⁶³Cu, ⁶⁵Cu and ¹⁰⁷Ag, Phys. Rev. C9 (1974) 1819-1830. [Half-life]
- 1974Kl02 A.A.C. Klaasse, P.F.A. Goudsmit, The Decay of ⁶³Zn, Z. Phys. 266 (1974) 75-82. [E_γ , P_γ]
- 1976Fu06 G.H. Fuller, Nuclear Spins and Moments, J. Phys. Chem. Ref. Data 5 (1976) 835-1092. [spin, magnetic dipole moment, electric quadrupole moment]
- 1977La19 F.P. Larkins, Semiempirical Auger-electron Energies for Elements $10 \leq Z \leq 100$, At. Data Nucl. Data Tables 20 (1977) 311-387. [Auger-electron energies]
- 1980Ku08 R.G. Kulkarni, D.P. Navalkele, Study of Negative Parity Levels in ⁶³Cu, Can. J. Phys. 58 (1980) 472-480. [γ - γ angular correlation, δ]
- 1982Gr10 A. Grütter, Decay Data of ⁵⁵⁻⁵⁸Co, ⁵⁷Ni, ^{60,61}Cu and ^{62,63}Zn, Int. J. Appl. Radiat. Isot. 33 (1982) 533-535. [Half-life, P_γ]
- 1995ScZY E. Schönfeld, Tables for the Calculation of Electron Capture Probabilities, PTB Report PTB-6.33-95-2, 1995. [P_K , P_L , P_M , P_N , P_O]
- 1996Sc06 E. Schönfeld, H. Janßen, Evaluation of Atomic Shell Data, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527-533. [ω_K , ω_L , K_β/K_α , $K_{\alpha 2}/K_{\alpha 1}$, KLX/KLL, KXY/KLL]
- 1998Sc28 E. Schönfeld, Calculation of Fractional Electron Capture Probabilities, Appl. Radiat. Isot. 49 (1998) 1353-1357. [P_K , P_L , P_M , P_N , P_O]
- 1998Si25 K.P. Singh, D.C. Tayal, H.S. Hans, Low-lying Levels in Cu and Zn Isotopes, Phys. Rev. C58 (1998) 1980-1985. [B(E2) values]
- 1998ScZM E. Schönfeld, G. Rodloff, Tables of the Energies of K-Auger Electrons for Elements with Atomic Numbers in the Range from $Z = 11$ to $Z = 100$, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX E. Schönfeld, G. Rodloff, Energies and Relative Emission Probabilities of K X-rays for Elements with Atomic Numbers in the Range from $Z = 5$ to $Z = 100$, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2000Le02 K.W.D. Ledingham, I. Spencer, T. McCanny, R.P. Singhal, M.I.K. Santala, E. Clark, I. Watts, F.N. Beg, M. Zepf, K. Krushelnick, M. Tatarakis, A.E. Dangor, P.A. Norreys, R. Allott, D. Neely, R.J. Clark, A.C. Machacek, J.S. Wark, A.J. Cresswell, D.C.W. Sanderson, J. Magill, Photonuclear Physics when a Multiterawatt Laser Pulse Interacts with Solid Targets, Phys. Rev. Lett. 84 (2000) 899-902. [Half-life]

- 2000Sc47 E. Schönfeld, H. Janßen, Calculation of Emission Probabilities of X-rays and Auger Electrons Emitted in Radioactive Disintegration Processes, *Appl. Radiat. Isot.* 52 (2000) 595-600. [P_X, P_{Ac}]
- 2001Ba27 E. Bai, Huo Junde, Nuclear Data Sheets for A = 63, *Nucl. Data Sheets* 92 (2001) 147-251. [nuclear levels]
- 2002Ba85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, Dirac–Fock Internal Conversion Coefficients, *At. Data Nucl. Data Tables* 81 (2002) 1-334. [ICC]
- 2002Ra45 S. Raman, C.W. Nestor, Jr., A. Ichihara, M.B. Trzhaskovskaya, How Good are the Internal Conversion Coefficients Now? *Phys. Rev. C* 66 (2002) 044312, 1-23. [ICC]
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr., Evaluation of Theoretical Conversion Coefficients using BrIcc, *Nucl. Instrum. Methods Phys. Res. A* 589 (2008) 202-229. [ICC]
- 2011StZZ N.J. Stone, Table of Nuclear Magnetic Dipole and Electric Quadrupole Moments, IAEA Report INDC(NDS)-0594, IAEA, Vienna, Austria (2011). [spin, magnetic dipole moment, electric quadrupole moment]
- 2012Wa38 M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer, The AME2012 Atomic Mass Evaluation, (I). Tables, Graphs and References, *Chin. Phys.* C36 (2012) 1603-2014. [Q-value]

⁶⁴Cu - Comments on evaluation of decay data by M.M. Bé and R.G. Helmer

This evaluation was completed in September 2010. To compare with the previous evaluation of R.G. Helmer in 2002, it includes several results published since this date (2002We02, 2007Qa02, 2010Wanke) and others obtained in the context of an Euramet exercise (2010 Bé).

Several procedures can be followed to determine the decay scheme of ⁶⁴Cu. In this evaluation we tried to introduce results coming from methods other than ionizing radiation measurements, in order to minimize the inherent correlation of the results.

1 Decay Scheme

The only levels in ⁶⁴Zn and ⁶⁴Ni below the decay energies are those populated in this decay, so the decay scheme is complete.

The J^π values and half-lives for the excited levels are from Adopted Levels in Nuclear Data Sheets (2007Si04).

2 Nuclear Data

Q values from 2003Au03 are 579.4 (7) keV for β^- decay and 1675.03 (20) keV for $\epsilon\beta^+$ decay.

The change in the half-life as a function of the chemical form or electron environment has been studied by several authors. These results are tallied after those used for the half-life evaluation.

The results of half-life measurements are listed below, in hours.

Not included in the evaluation			
	$T_{1/2}$	u_c	
(1935Am01)	10		omitted, no uncertainty
(1937He05)	12.5		omitted, no uncertainty
(1944Hu05)	11.9	1	omitted, same data as 1943Hu03
(1957Wr37)	12.87	0.05	superseded by 1972Em01
(1965He08)	13.9		omitted, no uncertainty
(1967Vi08)	12.8		omitted, no uncertainty
(1972WyZZ)	12.72	0.04	superseded by 1972Em01

The half-life values considered are, in hours:

	$T_{1/2}$	u_c	
(1936Va02)	12.8	0.1	
(1938Ri)	12.8	0.3	as cited in 1968Ke12
(1939Sa02)	12.8	0.3	as cited in 1968Ke12
(1943Hu03)	11.9	1	
(1950Ra62)	12.8	0.04	as cited in 1968Ke12

	T _{1/2}	u _c	
(1951Sc56)	12.74	0.07	
(1951Si91)	12.88	0.03	
(1955To07)	12.80	0.03	as cited in 1968Ke12
Rudstam	12.90	0.06	as cited in 1968Ke12
(1959Po64)	12.85	0.05	
(1965Pa18)	12.86	0.03	
(1966Fu14)	12.70	0.03	
(1966Li09)	12.86	0.03	
*(1968He20)	12.701	0.011	as cited in 1973De56
(1968Ke12)	12.80	0.04	
(1969Bo11)	12.65	0.17	
*(1972Em01)	12.715	0.007	
*(1972MeZM)	12.701	0.007	as cited in 1996Si12
(1973ArZI)	12.6	1	
*(1973De56)	12.699	0.002	
(1973Ne02)	12.82	0.04	
*(1974Ry01)	12.704	0.006	
*(1980RuZY, 1982RuZV)	12.701	0.003	
*(1989Ab22)	12.700	0.003	
*(2010Wanke)	12.705	0.005	
*(2010Bé - IFIN)	12.696	0.012	

The set of 25 unsuperseded values with uncertainties is inconsistent. The unweighted average is 12.76 (2) hours and the weighted average is 12.7025 with an internal uncertainty of 0.0013, a reduced- χ^2 of 6.3, and an external uncertainty of 0.003. It has been suggested that many of the older measurements give longer half-lives due to the presence of unidentified impurities. The value of 12.699 (2) used here for 1973De56 is the simple mean of their 22 measured values. The input value of 12.715 (7) is the evaluator's weighted average of the three values given in the paper of 1972Em01. The uncertainty given by 1989Ab22 has been increased to 0.003 to include systematic uncertainty components, but due to the very brief description of the process given in this paper, it is very difficult to assess them.

From the original set of 25 values, the most accurate ones (*) with uncertainties less than 0.012 hour have been accepted for statistical processing. In this set of nine values (1968He20, 1972Em01, 1972MeZM, 1973De56, 1974Ry01, 1980RuZY, 1989Ab22, 2010Wanke and 2010Bé - IFIN), the value of 1972Em01 was found outlier by the Chauvenet's criterion.

The adopted half-life is then the weighted average of the 8 remaining values. This average is 12.7003 with an internal uncertainty of 0.0013; an external uncertainty of 0.0007 and a reduced- χ^2 of 0.3.

As noted below, changes in this half-life of the order of 1 part in 10⁴ have been reported depending on the chemical form. Since these changes are comparable to the calculated uncertainty, the adopted uncertainty has been increased to 0.0020.

This half-life has been measured, and reported, many times primarily to identify the radionuclide observed, for example, in the process of cross section measurements. Some of these values, which are not included above are: 13 (1948Mi12); 12.8 (1950Ho26); and 13.8 (14), 13.6 (7), and 12.4 (17) (1972Cr02).

Since ⁶⁴Cu decays, in part, by electron capture, there have been several measurements of the variation

in the decay constant with the chemical form or atomic environment. The results from 1968 to 1975 are tallied in 1976Ha66 and given in the following table.

Reference and first author	Forms compared	$\Delta\lambda/\lambda \cdot 10^4$
1972Au Auric	Cu phtalocyanine in two forms	10.0 (16)
1972Em01 Emery	Cu metal	Cu(NO ₃) ₂ 15 (15)
1973Ha60 Harbottle	Cu metal	CuO 0 (3)
1973De56 Dema	Cu phtalocyanine in two forms	0.4 (20)
1974Je Jenschke	Cu metal	Cu(H ₂ O) ₆ SO ₄ 0.12 (9)
	Cu metal	Cu(H ₂ O) ₄ (NO ₃) ₂ 0.81 (10)
	Cu metal	Cu(2) 2.94
	Cu metal	Cu(3) 1.86
1974Jo17 Johnson	Cu phtalocyanine in two forms	1.4 (23)
	Cu phtalocyanine in two forms	3.7 (58)
	Cu metal	CuO 0.0 (23)
1975MaXN	Cu metal	Cu ₂ S 2.3 (10)
	Cu metal	CuInS ₂ 1.5 (10)
	Cu metal	Cu ₂ SnS ₃ 1.5 (10)
	CuInS ₂	Cu ₂ SnS ₃ 0 (1)
1979Eh01 Ehrhart	Cu metal	atom % Cu in Ag
		2 1.7 (3)
		5 1.6 (4)
		25 0.9 (4)
		50 0.7 (5)
1979Ko31 Koran	Cu metal	atom % Cu in Au
		2 3.1 (4)
		5 3.0 (4)
		25 1.4 (4)
		50 0.7 (5)
		75 -0.2 (9)

The earliest measurements gave larger values of $\Delta\lambda/\lambda$, but the values beginning in 1973 range from 0 to 0.000 37 (6). These values are similar in magnitude to the uncertainty of 1.5 parts in 10⁴ assigned to the adopted value. A set of measurements is also given in 1968Ke12, but the units of the results are not clear.

No dependence of the half-life with the temperature has been observed (2008Fa12).

2.1 β^- , β^+ and Electron Capture Transitions

The probabilities of the β^- , β^+ and ϵ branches were determined by a series of separate, but partially correlated, measurements by 1983Ch47, 1986Ka03 and 2007Qa02. These measurements included the β^- spectrum, β^+ spectrum, $4\pi\beta\text{-}\gamma$ coincidences, liquid scintillation counting, and X-, γ - ray spectrum. Then, in 1983Ch47 their analysis contained a least-squares fit to the various measured quantities and ratios of quantities, taking the covariances into account.

Another kind of investigation made by mass spectrometry measurements of the number of atoms of ⁶⁴Ni and ⁶⁴Zn produced in the decay of a ⁶⁴Cu sample (2002We02) led to the determination of the P _{β^-} branching ratio.

- β^- Transition

The published measured probabilities of the β^- transition are:

References	P_{β^-} (%)	uc (%)	Comments
1983Ch47	39.04	0.33	$4\pi\beta(\text{LS})-\gamma$ coincidence counting
1986Ka03	38.34	0.56	deduced
2002We02	38.06	0.3	Mass spectrometry
2007Qa02	38.4	1.2	2π PC – anti coincidence counting
$\chi^2/n - \text{crit } \chi^2$	1.6	3.8	
UWM	38.46	0.21	
WM	38.48	0.26	Adopted

From $P_{\beta^-} = 38.48$ (26) %, then $P(\beta^+ + \text{ec}) = 61.52$ (26) %.

- β^+ Transition

Two methods are possible to derive the $P(\beta^+)$ value:

- From published measured probabilities of the β^+ transition:

References	P_{β^+} (%)	uc (%)	Comments
1983Ch47	17.86	0.14	Ge(Li) spectrometry
1986Ka03	17.93	0.20	HPGe γ spectrometry
2007Qa02	17.8	0.4	$\gamma-\gamma$ coincidence counting
2010Wanke	17.56	0.16	HPGe γ spectrometry
2010Bé - CMI	17.69	0.19	HPGe γ spectrometry
2010Bé - LNHB	17.55	0.15	HPGe γ spectrometry
2010Bé – IFIN	17.65	0.60	HPGe γ spectrometry
$\chi^2/n - \text{crit } \chi^2$	0.7	2.8	
UWM	17.72	0.06	
WM	17.71	0.07	

- From theoretical calculations, using the LOGFT program, the ratio $P_{\text{ec}}/P(\beta^+)$ is: 2.485 (25), from the $P(\beta^+ + \text{ec}_{0,0}) = 61.05$ (26) % below, the $P(\beta^+)$ value is derived being $P(\beta^+) = 17.52$ (15) %.

The latter value has been obtained by an independent method and it is less correlated than the results of direct measurements. Moreover, it can be noted that the weighted mean of the four 2010 values, listed in the above table, of 17.59 (9) is very close to 17.52 (15). Thus, $P(\beta^+) = 17.52$ (15) % has been adopted in this evaluation.

- Electron Capture Transitions

The adopted $P_{\text{ec},1}$ value is deduced:

From $P_{g_{1345}} = 0.4744$ (33) % (§ 2.4), then $P_{\text{ec},1} = 0.4744$ (33) %,

and with $P(\beta^+ + \text{ec}) = 61.52$ (26) %, $P(\beta^+ + \text{ec}_{0,0}) = 61.05$ (26) %.

From the two values of $P(\beta^+)$ as determined above two $P_{ec,0}$ can be derived:

- With $P(\beta^+) = 17.71$ (7), $P(ec_{0,0}) = 43.34$ (27) %,
- With $P(\beta^+) = 17.52$ (15), $P(ec_{0,0}) = 43.53$ (20) %.

These values can be compared with the three experimental results obtained for the total P_{ec} (1983Ch47, 1986Ka03, 2007Qa02):

References	Total P_{ec}	uc (%)	Comments
1983Ch47	43.10	0.46	$4\pi\beta(LS)-\gamma$ coincidence counting
1986Ka03	43.73	0.52	$4\pi\beta\gamma(PC)-\gamma$ coincidence counting + HPGe γ spectrometry + $4\pi\beta-\gamma$ coincidence counting
2007Qa02	43.8	1.4	Si(Li) X-ray spectrometry
$\chi^2/n - \text{crit } \chi^2$	0.5	4.6	
UWM	43.54	0.22	$P(ec_{0,0}) = 43.07$ (33)
WM	43.40	0.33	$P(ec_{0,0}) = 42.93$ (33)

The unweighted and the weighted means above are consistent, within the uncertainty limits, with $P(ec_{0,0}) = 43.34$ (27) % calculated from experimental P_{β^+} values. This was expected since they were derived from the same sets of measurements.

The set of two values: **$P(\beta^+) = 17.52$ (15) % and $P(ec_{0,0}) = 43.53$ (20) %** has been adopted in this evaluation because it was derived from another different method (using theoretical $P_{ec}/P(\beta^+)$ ratio).

The average particle energies to the ⁶⁴Ni and ⁶⁴Zn ground states are 278.2 (9) and 190.7 (3) keV, respectively, and are from the LOGFT code. The log ft values to the ⁶⁴Ni ground state and level of 1345 keV are 4.973 (3) and 5.501 (6), respectively, and to the ⁶⁴Zn ground state - 5.302 (3), all of which are consistent with allowed transitions from the 1⁺ parent state.

2.2 Gamma Transitions

The J^π assignments are from the Adopted Levels in the Nuclear Data Sheets (2007Si04) and these imply the γ -ray has E2 multipolarity.

The internal-conversion coefficients (ICC) were interpolated from the tables of Band *et al.* (2002Ba85) by using the computer code BrIcc (2008Ki07) with the so called ‘‘Frozen orbital’’ approximation.

The internal-pair-formation coefficient was interpolated from the theoretical values and it is $IPFC(1345) = 0.000\ 039$.

From the ICC values and gamma ray emission intensity $I_{g1345} = 0.4743$ (33) % (§ 4.2), the 1345 keV gamma transition probability and electron capture probability to the first excited level in ⁶⁴Ni are deduced being: $P_{g1345} = P_{ec,1} = 0.4744$ (33) %.

3 Atomic Data

The data are from 1996Sc06.

4 Radiation Emissions

4.1 Electron Emissions

Auger electron emission intensities are deduced from the evaluated data set.

4.2 Photon Emissions

X-ray emission intensities are deduced from the evaluated data set.

The γ -ray energy is 1345.77 (6) keV from 1974HeYW.

The intensity of the 1345 keV gamma ray is deduced from the measured values:

Reference	$I_{\gamma 1345}$ (%)	uc (%)	Comments
1983Ch47	0.471	0.011	HPGe γ spectrometry
1986Ka03	0.487	0.020	HPGe γ spectrometry
2007Qa02	0.54	0.03	HPGe γ spectrometry
2010Wanke	0.474	0.005	HPGe γ spectrometry
2010Bé - CMI	0.476	0.006	HPGe γ spectrometry
2010Bé - LNHB	0.467	0.011	HPGe γ spectrometry
2010Bé- IFIN	0.481	0.017	HPGe γ spectrometry
$\chi^2/n - \text{crit } \chi^2$	0.24	3	
UWM	0.476	0.0029	
WM	0.4743	0.0033	Adopted

5 Various comparisons

The following tables summarize values of some ratios measured or deduced in the publications compared with those derived from the present data set. Both are in agreement within the uncertainty limits.

➤ $P_{\beta^-} / P_{\beta^+}$ ratio

Reference	$P_{\beta^-} / P_{\beta^+}$	uc	Remark
1946Br03	2.1		
1949Bo16	2.00	0.15	
1983Ch47	2.187	0.007	
1986Ka03	2.138	0.032	
$\chi^2/n - \text{crit } \chi^2$	2.2	6.6	
UWM	2.163	0.025	Value deduced from the present adopted data set: 2.196 (24)
WM	2.185	0.010	

➤ $I_{\gamma 1345} / P_{\beta^+}$ ratio

Reference	$I_{\gamma 1345} / P_{\beta^+}$	uc	Remark
1956Dz26	0.020 7		
1952Vl03	0.023	0.004	Omitted, outlier
1959Sc71	0.028 0	0.002 4	Omitted, outlier
1983Ch47	0.026 4	0.000 6	
1986Ka03	0.027 2	0.001 2	
2007Qa02	0.030 3	0.001 8	Omitted, outlier
2010Wanke	0.026 99	0.000 38	
2010Bé - CMI	0.026 91	0.000 45	
2010Bé - LNHB	0.026 6	0.000 7	
$\chi^2/n - \text{crit } \chi^2$	0.23	3.3	
UWM	0.026 82	0.000 14	Value deduced from the present adopted data set: 0.027 06 (30)
WM	0.026 84	0.000 24	

6 Main production modes

They are taken from: Table de Radionucléides, F; Lagoutine, N. Coursol, J. Legrand. ISBN 2 7272 0078-1

7 Other earlier publications not used in the evaluation

- H. von Bradt (1946Br03)

$$P_{\beta^-} / P_{\beta^+} = 2.1$$

- R. Bouchez (1949Bo16)

$$P_{\beta^-} / P_{\beta^+} = 2.00 \text{ (15)}$$

- Reynolds (1950Re51)

$$P(\beta^+ + ec) / P_{\beta^-} = 1.62 \text{ (11)}$$

$$P(ecK) / P(\beta^+) = 2.32 \text{ (28)}$$

- Vlaar (1952VI03)

$$I_{\gamma 1345} / P(\beta^+) = 0.023 \text{ (4)}$$

- Dzelepov *et al.* (1956Dz26)

$$I_{\gamma 1345} / P(\beta^+) = 0.0207$$

- Schmidt-Ott (1959Sc71)

$$I_{\gamma 1345} / P(\beta^+) = 0.0280 \text{ (24)}$$

8 References

- 1935Am01 E. Amaldi, O. D'Agostino, E. Fermi, B. Pontecorvo, F. Rasetti, E. Segre, Proc. Roy. Soc. (London) 149A(1935)522 [T_{1/2}]
- 1936Va02 S. N. Van Voorhis, Phys. Rev. 50(1936)895 [T_{1/2}]
- 1937He05 F. A. Heyn, Physica 4(1937)1224 [T_{1/2}]
- 1938Ridenour L. N. Ridenour, Phys. Rev. 53(1938)770 [T_{1/2}]
- 1939Sa02 R. Sagane, Phys. Rev. 55(1939)31 [T_{1/2}]
- 1943Hu03 O. Huber, O. Lienhard, H. Waffler, Helv. Phys. Acta 16(1943)226 [T_{1/2}]
- 1944Hu05 O. Huber, O. Lienhard, H. Waffler, Helv. Phys. Acta 17(1944)195 [T_{1/2}]
- 1946Br03 H. von Bradt *et al.* Phys. Acta 19(1946) 219 [P_{β⁻} / P_{β⁺}]
- 1948Mi12 D. R. Miller, R. C. Thompson, B. B. Cunningham, Phys. Rev. 74(1948)347 [T_{1/2}]
- 1949Bo16 R. Bouchez, G. Kayas, J. Phys. Radium 10, série 8 (1949) 110 [P_{β⁻} / P_{β⁺}]
- 1950Ho26 H. H. Hopkins, Phys. Rev. 77(1950)717 [T_{1/2}]
- 1950Ra62 E. Rabinowicz, Proc. Phys. Soc.(London) 63A(1950)1040 [T_{1/2}]
- 1950Re51 J.H. Reynolds, Phys. Rev. 79,5 (1950) 789 [P(ecK) / P(β⁺)]
- 1951Sc56 R. P. Schuman, A. Camilli, Phys. Rev. 84(1951)158 [T_{1/2}]
- 1951Si91 L. M. Silver, Can. J. Phys. 29(1951)59 [T_{1/2}]
- 1952VI03 H.T. Vlaar, Physica 18 (1952) 275 [I_{γ511} / I_{γ1345}]
- 1955To07 J. Tobailem, J. Phys. Radium 16(1955)48 [T_{1/2}]

- 1956Dz26 B.S. Dzelepov *et al.* Nuovo Cimento 3, Supp. 1, (1956) 49 [I_γ]
- 1957Wr37 H. W. Wright, E. I. Wyatt, S. A. Reynolds, W. S. Lyon, T. H. Handley, Nuclear Sci. and Eng. 2(1957)427 [T_{1/2}]
- 1959Sc71 W-D Schmidt-Ott, Z. Physik 154 (1959) 286 [I_{γ1345} / P(β⁺)]
- 1959Po64 A. Poularikas, R. W. Fink, Phys. Rev. 115(1959)989 [T_{1/2}]
- 1965He08 Z. He-Sung, N. S. Maltseva, V. N. Mekhedov, V. N. Rybakov, Soviet J. Nucl. Phys. 1(1965)132 [T_{1/2}]
- 1965Pa18 V. A. Paulsen, H. Liskien, Nukleonik 7(1965)117 [T_{1/2}]
- 1966Fu14 K. Fujiwara, O. Sueka, J. Phys. Soc. Japan 21(1966)1947 [T_{1/2}]
- 1966Li09 H. Liskien, A. Paulsen, Proc. Intern. Conf. Radiat. Meas. Nucl. Power, Berkeley, Engl., D. J. Littler, Ch., Editorial Panel, Inst. Phys. and the Physical. Society, London, Conf. Series No.2, (1966) p. 352 [T_{1/2}]
- 1967Vi08 G. P. Vinitzskaya, V. N. Levkovsky, V. V. Sokolsky, I. V. Kazachevsky, Sov. J. Nucl. Phys. 5(1967)839 [T_{1/2}]
- 1968He20 F. Heinrich, G. Philippin, Helv. Phys. Acta 41(1968)431 [T_{1/2}]
- 1968Ke12 P. Kemény, Rev. Roumaine Phys. 13(1968)901 [T_{1/2}]
- 1969Bo11 M. Bormann, B. Lammers, Nucl. Phys. A130(1969)195 [T_{1/2}]
- 1972Au P. Auric, J. I. Vargas, Chem. Phys. Lett. 15(1972)366 [T_{1/2}]
- 1972Cr02 D. F. Crisler, H. B. Eldridge, R. Kunselman, C. S. Zaidins, Phys. Rev. C5(1972)419 [T_{1/2}]
- 1972Em01 J. F. Emery, S. A. Reynolds, E. I. Wyatt, G. I. Gleason, Nucl. Sci. Eng. 48(1972)319 [T_{1/2}]
- 1972MeZM J. S. Merritt, J. G. V. Taylor, AECL-4257(1972) p. 25 [T_{1/2}]
- 1972WyZZ E. I. Wyatt, ORNL-4749(1972) p.61 [T_{1/2}]
- 1973ArZI J. Araminowicz, J. Dresler, INR-1464(1973) p.14 [T_{1/2}]
- 1973De56 I. Dema, G. Harbottle, Radiochem. Radioanal. Lett. 15(1973)261 [T_{1/2}]
- 1973Ha60 G. Harbottle, C. Koehler, R. Withnell, Rev. Sci. Instr. 44(1973)55 [T_{1/2}]
- 1973Ne02 D. A. Newton, S. Sarkar, L. Yaffe, R. B. Moore, J. Inorg. Nucl. Chem. 35(1973) 361 [T_{1/2}]
- 1974HeYW R. L. Heath, ANCR-1000-2(1974) [E_γ]
- 1974Je B. Jenschke, German Phys. Soc., Spring Conf.(1974) [T_{1/2}]
- 1974Jo17 J. A. Johnson, I. Dema, G. Harbottle, Radiochim. Acta 21(1974)196 [T_{1/2}]
- 1974Ry01 T. B. Ryves, K. J. Zieba, J. Phys.(London) A7(1974)2318 [T_{1/2}]
- 1976Ba63 I. M. Band, M. B. Trzhaskovskaya, M. A. Listengarten, Atomic Data Nucl. Data Tables **18** (1976) 433 [α]
- 1976Ha66 H. P. Hahn, H. J. Born, J. I. Kim, Radiochim. Acta 23(1976)23 [T_{1/2}]
- 1979Sc31 P. Schluter, G. Soff, At. Data Nucl. Data Tables **24** (1979)509 [IPFC]
- 1980RuZY A. R. Rutledge, L. V. Smith, J. S. Merritt, AECL-6692(1980) [T_{1/2}]
- 1982RuZV A. R. Rutledge, L. V. Smith, J. S. Merritt, NBS-SP-626(1982) p.5 [T_{1/2}]
- 1983Ch47 P. Christmas, S. M. Judge, T. B. Ryves, D. Smith, G. Winkler, Nucl. Instr. Meth. 215(1983)397 [P_{β-}, P_{β+}, P_γ, P_ε]

- 1986Ka03 Y. Kawada, Intern. J. Appl. Radiat. Isot. 37(1986)7 [P_{β^-} , P_{β^+} , P_{γ} , P_{ε}]
- 1989Ab22 A.Abzouzi, M.S.Antony, V.B.Ndocko Ndongue, J. Radioanal. Nucl. Chem. 135 (1989)455 [$T_{1/2}$]
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instr. Meth. **A369** (1996)527 [ω]
- 2002Ba85 I.M.Band, M.B.Trzhaskovskaya. At. Data. Nucl. Data Tables 88,1 (2002). [Theoretical ICC]
- 2002We02 G. Wermann, D. Alber, W. Pritzkow, G. Riebe, J. Vogl, W. Görner. Appl. Rad. Isotopes 56, 1-2 (2002) 145 [$\% \beta^-$]
- 2003Au03 G. Audi, A.H.Wapstra, C.Thibault, Nucl. Phys. **A729** (2003)337 [Q]
- 2007Qa02 S.M. Qaim, T.Bisinger, K.Hilgers, D.Nayak, H.H.Coenen, Radiochim. Acta 95 (2007) 67, [P_{β^-} , P_{β^+} , P_{γ} , P_{ε}]
- 2007Si04 B. Singh, Nucl. Data Sheets **108** (2007)197 [J^{π} . multipolarities]
- 2008Fa12 B.A. Fallin *et al.* Physical Review C78 (2008) 057301 [$T_{1/2}$]
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, and C.W.Nestor, Jr., Nucl. Instrum. Methods Phys. Res. **A589**, 202 (2008) [Theoretical ICC]
- 2010Wanke C. Wanke, K. Kossert, Ole J. Nähle, O. Ott. Appl. Radiat. Isot. (2010) doi: 10.1016/j.apradiso.2010.01.005, [P_{β^+} , P_{γ}]
- 2010Bé M.M. Bé *et al.* Euramet Project 1085, full report in preparation

⁶⁵Zn - Comments on evaluation of decay data by M.M. Bé, V. Chisté and R. G. Helmer

1 Decay scheme

This evaluation was originally completed in September 1996. New evaluation was completed in January 2005 taking into account results obtained as a part of a specific exercise dedicated to the ⁶⁵Zn activity and emission intensity measurements managed by the Euromet organization.

The decay scheme is complete since only two excited levels in ⁶⁵Cu below the decay energy are populated. Also, there is excellent agreement between the total decay energy of 1352.1 (19) keV computed from the evaluated decay scheme and the Q value of 1352.1 (3) keV.

2 Nuclear Data

Q = 1352.1 (3) value is from Audi *et al.* (2003Au03).

The measured ⁶⁵Zn half-life values, in days, are as follow:

245.0 (8)	1953To17	
243.5 (8)	1957Ge07	
246.4 (22)	1957Wr37	outlier
243.1 (7)	1965An07	replaced by 1982HoZJ
244.12 (12)	"	replaced by 1982HoZJ
242.78 (19)	"	omitted from analysis
243 (4)	1968Ha47	
258 (4)	1972Cr02	omitted from analysis
246 (5)	"	"
251 (6)	"	"
252 (6)	"	"
244.0 (2)	1972De24	replaced by 2004Va02
244.52 (7)	1973Vi13	Uncertainty given per 3 σ
244.3 (4)	1974Cr05	
243.75 (12)	1975La16	replaced by 2003Lu06
244.2 (1)	1982HoZJ	replaced by 1992Un01
243.97 (8)	1982DeYX	replaced by 1983Wa26
243.9 (3)	1983Wa26	replaced by 2004Sc04
244.16 (10)	1992Un01	(or 2002Un02)
244.15 (10)	2003Lu06	
243.66 (9)	2004Sc04	
243.8 (3)	2004Va02	

244.01 (9) Adopted

The four values of 1972Cr02 were omitted because they were not intended as $T_{1/2}$ measurements, but rather to determine the origin of certain γ -rays.

The very small uncertainty, 0.07 (3.3 σ), given by 1973Vi13 appears unrealistic when compared to the other quoted uncertainties at the same period of time, at least this uncertainty value should be increased. Moreover, this result is far from the mean and the published paper not detailed enough, so this result is omitted from analysis.

The value of 1957Wr37 was found outlier according to the Chauvenet's criterion.

As a rule, only one result per laboratory is retained in order to avoid possible correlation.

Then, the weighted average of the remaining eight values is 244.01 with an internal uncertainty of 0.05, an external uncertainty of 0.09 and, a reduced- χ^2 of 3.11 (the critical reduced- χ^2 is 2.60), no input value has more than 50% of the weight. The Lweight program suggested to expand the uncertainty to 0.31 in order to include the most precise value of 243.66 within its range.

But a small increased of the uncertainty given by 2004Sc04 from 0.09 to 0.11 leads to a reduced- χ^2 of 2.48 less than the critical one, then the Lweight program recommended the internal uncertainty as final uncertainty.

With these results in mind, the evaluator has chosen the weighted average and the external uncertainty.

2.1 Electron Capture Transitions

The ϵ branch to the 770-keV level is 2nd forbidden. From the log ft systematics (1973Ra10), the expected log ft value is > 11.0 and the corresponding $I_\epsilon(0)$ is $< 0.003\%$.

The P_K etc. values are computed from the Schönfeld tables (1995ScZY) for allowed transitions.

Level energy (keV) =	0	1115
P_K (S)	0.8853 (16)	0.8794 (17)
P_L (S)	0.0977 (15)	0.1027 (16)

The total branching ratios to each level were computed from the measured I_γ and adopted theoretical conversion coefficients.

The total branching ($\epsilon + \beta^+$) to the ground state is 49.77 (11) %. From the 511-keV gamma emission intensity measurements, the β^+ transition probability is deduced as 1.421 (7) % (see § Photon emissions).

The LOGFT program gives the theoretical ϵ/β^+ ratio as 34.03 (18). Using the ($\epsilon + \beta^+$) branching to the ground state as 49.77 (11) % ; the β^+ transition probability is then 1.42 (1)%. This value is in good agreement with the experimental observations.

From the LOGFT program, the theoretical ϵ_K/β^+ ratio is calculated as 29.86. This value can be compared with the corresponding experimental values of:

28.0 (32)	1953Pe14
30.3 (12)	1963Ta04
27.7 (15)	1968Ha47
31.3 (20)	1977Bo10
30.7 (11)	1984ScZP
30.3 (10)	1990Ku11

The measured 1115 γ/β^+ ratio is 35.1 (17) (1968Ha47).

For comparison with the adopted value for the β^+ transition probability of 1.421 (7)%, the measured values are :

	I_{β^+} (%)
1959Gl55	1.70 (10)
1962Be28	1.2 (3)
1963Ta04	1.40 (4)
1972De24	1.46 (2)

2.2 Gamma Transitions

The multiplicities are from the adopted γ -ray data (deduced from Coulomb excitation study and angular correlation data) in the journal Nuclear Data Sheets (1993Bh04).

The internal-conversion coefficients are interpolated from the tables of Band (2002Ba85). Mixing ratio of the 1115-keV transition is from Krane (1976Kr09). The ICC values for this high energy transition is very low so the influence of the uncertainty for the mixing ratio is negligible.

For the 1115-keV transition, the total and K-shell values of $1.85 (7) \times 10^{-4}$ and $1.66 (6) \times 10^{-4}$ respectively, evaluated by Hansen (1985HaZA) from measured values are in excellent agreement with the theoretical ones.

From the theoretical tables of 1979Sc31, the internal-pair-formation coefficients are $\alpha_{\pi}(1115, M1) = 1.2 \times 10^{-6}$ and $\alpha_{\pi}(1115, E2) = 1.6 \times 10^{-6}$, so $\alpha_{\pi}(1115) = 1.3 \times 10^{-6}$. This value is about 1% of the internal-conversion coefficient and therefore is negligible.

3 Atomic Data (Cu, Z=29)

Data are from 1996Sc06.

4.1 Electron Emission

The β^+ intensity to the ground state was deduced from the measured intensity of the 511-keV gamma ray.

4.2 Photon Emissions

The γ -ray energies are from the evaluation of Helmer *et al.* (2000He14) for the 1115-keV line where the values are on a scale on which the strong line from the decay of ¹⁹⁸Au is 411.80205 (17); from level energy differences for the 344-keV line; and from ⁶⁵Cu Adopted γ data in Nuclear Data Sheets (1993Bh04) and based on data from ⁶⁵Ni β^- decay for 770-keV line.

Photon emission intensities are deduced from the Euromet exercise results (2005Be**) and from other published values.

Absolute measured intensities of the 1115-keV line

	I (1115) (%)	Uc	
1959Gl55	51.3	3.0	
1960Go	46		
1963Ta04	50.7	0.5	
1966Ra21	51.3	1.5	
1968Ha47	52.4	1.0	Outlier
1972De24	50.75	0.10	Replaced by Euromet participant
1973Po10	49.3	0.8	
1982DeYX	50.39	0.26	replaced by 1990Sc08
1990Sc08	50.2	0.4	Replaced by Euromet participant
2003Lu06	49.76	0.21	Replaced by Euromet participant
Euromet-01	50.15	0.28	
Euromet-02	50.10	0.33	
Euromet-03	50.60	0.29	
Euromet-04	50.34	0.25	
Euromet-05	49.84	0.25	
Euromet-06	50.05	0.57	
Euromet-07	49.62	0.65	
Euromet-08	50.7	0.5	
Euromet-09	50.3	0.5	
Adopted	50.22	0.11	

The first part of the above Table lists the results published in various journals and the second part lists the values obtained as a part of the Euromet exercise (2005Be*).

The value from 1968Ha47 is omitted as outlier due to application of the Chauvenet's criterion. The results from 1972De24, 1990Sc08 and 2003Lu06 have been superseded by the results obtained by laboratories which have participated in the present Euromet exercise.

The LRSW analysis of the remaining 13 values gives a reduced χ^2 of 0.77 so the weighted mean of 50.22 and the internal uncertainty of 0.11, are adopted as final result.

344- and 770-keV Relative γ -ray emission intensities :

γ -ray energy (keV)	I(344)	I(770)	I(1115)
1960Ri06	≤ 0.5	≤ 1	100
1968St05	0.0060 (6)		100
Euromet-02	0.005067 (365)	0.005358 (439)	100
Euromet-09	0.00220 (86)	0.003 (17)	100
Adopted relative	0.005067 (365)	0.005358 (439)	100
Adopted absolute	0.00254 (18)	0.00269 (22)	50.22 (11)

The adopted relative values are those given by the participant 2 in the Euromet exercise. This participant activated Zinc (99.99 %) foil by thermal neutrons and obtained a Zn-65 activity of the order of 10 MBq, so he had a better counting statistic and then a better uncertainty.

511-keV photon emission

This particular emission is due to the annihilation of the β^+ positrons in the source and in the surrounding material (annihilation-in-flight). In γ -ray spectrometry, this phenomenon has the effect of removing, from the 511-keV peak, a fraction of the annihilation photons, the magnitude of this effect depends on the material in which the β^+ are stopped and then must be calculated by each experimentalist.

reference	Intensity (%)	Uc	Correction for annihilation, in %
1990Sc08	2.84*	0.04	0.5
Euromet-01	2.81 *	0.03	0.2
Euromet-02	2.841 *	0.027	Wider peak region
Euromet-03	2.75	0.017	
Euromet-04	3.00	0.018	
Euromet-05	2.848 *	0.020	0.34
Euromet-07	2.86	0.04	
Euromet-09	2.88 *	0.04	0.5

(*) taking annihilation-in-flight into account, magnitude given in the last column.

Reference 1990Sc08 is superseded by one of the Euromet participant. The weighted mean and standard uncertainty of the four values taking annihilation-in-flight into account, are : 2.842 ± 0.013 %.

The emission of additional 511-keV photons created by electron-positron pair creation is negligible (see § Gamma transitions).

X-ray emissions and Auger electron emissions

From the gamma-ray emission intensities, the internal conversion coefficients, the electron capture probabilities and electron capture sub shell probabilities, the X-ray and Auger electron emission intensities have been deduced.

Calculated K X-ray are compared with the measured values in the following table.

Reference	K α		K β		KX	
	Intensity	Uc	Intensity	Uc	Total	Uc
1963Ta19					40.0	0.6
1968Ha47					39.27	0.26
1968Ba**					38.66	0.17
1973Mu**					38.0	1.0
Euromet-05	32.1	1.6	4.50	0.023	36.6	1.6
Euromet-09	39	3.5	5.2	0.47	44.2	3.5
Weighted mean					38.87	0.22
Calculated	34.7	0.4	4.82	0.07	39.5	0.4

The weighted mean of the KX measured values (except Euromet-09 which is outlier) is lower than the calculated value deduced from the decay scheme. They barely agree within their uncertainty limits.

6 References

- 1953Pe14 - J. F. Perkins, S. K. Haynes, Phys. Rev. **92**(1953)687 [ϵ/β^+]
 1953To17 - J. Toballem, J. Phys. Radium **14**(1953)553 [$T_{1/2}$]
 1957Ge07 - K. W. Geiger, Phys. Rev. **105**(1957)1539 [$T_{1/2}$]
 1957Wr37 - H. W. Wright, E. I. Wyatt, S. A. Reynolds, W. S. Lyon, T. H. Handley, Nuclear Sci. Eng. **2**(1957)427 [$T_{1/2}$]
 1959Gl55 - G. I. Gleason, Phys. Rev. **113**(1959)287 [P_{β^+} , P_{γ}]
 1960Go - W. M. Good, W. C. Peacock, Bull. Amer. Phys. Soc. Abstract B4 (1960) 680 [P_{γ}]
 1960Ri06 - R. A. Ricci, G. Chilosi, G. Varcaccio, G. B. Vingiani, R. van Lieshout, Nuovo cimento **17**(1960)523 [P_{γ}]
 1962Be28 - D. Berenyi, Phys. Letters **3**(1962)142 [P_{β^+}]
 1963Ta04 - J. G. V. Taylor, J. S. Merritt, Phys. Can. **19**(1963) No. 3, 17, abstract 4.5 [P_{γ}]
 1963Ta19 - J. G. V. Taylor, J. S. Merritt, Proc. Int. Conf. role of atomic electrons in nuclear transformations, Warsaw, CONF-233 (1963) 465 [XK]
 1965An07 - S. C. Anspach, L. M. Cavallo, S. B. Garfinkel, J. M. R. Hutchinson, C. N. Smith, NP-15663(1965) [$T_{1/2}$]
 1966Ha07 - J. H. Hamilton, S. R. Amtey, B. van Nooijen, A. V. Ramayya, J. J. Pinajian, Phys. Letters **19**(1966)682 [α , α_K]
 1966Ra21 - P. S. Rao, Curr. Sci. **35**(1966)384 [P_{γ}]
 1968Ba** - W. Bambynek, D. Reher, Z. Physik **214**(1968)374 [XK]
 1968Ha47 - J. W. Hammer, Z. Physik **216**(1968)355 [$T_{1/2}$, P_{γ}]
 1968St05 - P. H. Stelson, Nucl. Phys. **A111**(1968)331 [P_{γ}]
 1972Cr02 - D. F. Crisler, H. B. Eldridge, R. Kunselman, C. S. Zaidins, Phys. Rev. **C5**(1972)419 [$T_{1/2}$]
 1972De24 - E. De Roost, E. Funck, A. Spornol, R. Vaninbrouckx, Z. Phys. **250**(1972)395 [ϵ/β^+ , P_{γ} , $T_{1/2}$]
 1973Po10 - W. P. Poenitz, A. Devolpi, Int. J. Appl. Radiat. Isotop. **24**(1973)471 [P_{γ}]
 1973Ra10 - S. Raman, N.B. Gove, Phys. Rev. C7 (1973) 1995 [lg ft]

- 1973Vi13 - C. J. Visser, J. H. M. Karsten, F. J. Haasbroek, P. G. Marais, *Agrochemophysica* **5**(1973)15 [T_{1/2}]
 1973Mu** - A.Mukerji, L.Chin, Atlanta Conf. Proc. AEA-CONF-720404 (1973) 164 [XK]
- 1974Cr05 - P. J. Cressy,Jr., *Nucl. Sci. Eng.* **55**(1974)450 [T_{1/2}]
 1975La16 - F. Lagoutine, J. Legrand, C. Bac, *Int. J. Appl. Radiat. Isotop.* **26**(1975)131 [T_{1/2}]
 1976Kr09 - K.S. Krane, S.S. Rosenblum, W.A. Steyert. *Phys. Rev. C* **14**, (1976) 650 [δ]
 1977Bo10 - H. E. Bosch, J. Davidson, M. Davidson, L. Szybisz, *Z. Phys.* **A280**(1977)321 [ε/β⁺]
 1979Sc31 - P. Schluter, G. Soff, *At. Data Nucl. Data Tables* **24**(1979)509 [α_π]
 1982DeYX - K. Debertin, U. Schötzig, K. F. Walz, NBS-SP-626(1982)101 [P_γ]
 1982HoZJ - D. D. Hoppes, J. M. R. Hutchinson, F. J. Schima, M. P. Unterweger, NBS-SP-626(1982)85 [T_{1/2}]
 1983Wa26 - K. F. Walz, K. Debertin, H. Schrader, *Int. J. Appl. Radiat. Isotop.* **34**(1983)1191 [T_{1/2}]
 1984ScZP - W.-D. Schmidt-Ott, J. Lauerwald, U. Bosch, H. Dornhofer, U. J. Schrewe, H. Behrens, 7th Proc. Intern. Conf. Atomic Masses Fund. Constants, Darmstadt-Seeheim (1984)210 [ε/β⁺]
 1985HaZA - H. H. Hansen, *European App. Res. Rept. Nucl. Sci. Technol.* **6**, No.4 (1985)777 [α, α_K]
 1990Ku11 - V. Kunze, W.-D. Schmidt-Ott, H. Behrens, *Z. Phys.* **A337**(1990)169 [ε/β⁺]
 1990Sc08 - U. Schötzig, *Nucl. Instrum. Methods Phys. Res.* **A286**(1990)523 [P_{β+}, P_γ]
 1991BaZS - W. Bambynek, T. Barta, R. Jedlovsky, P. Christmas, N. Coursol, K. Debertin, R. G. Helmer, A. L. Nichols, F. J. Schima, Y. Yoshizawa, report IAEA-TECDOC-619 (1991) [P_γ evaluation]
 1992Un01 - M. P. Unterweger, D. D. Hoppes, F. J. Schima, *Nucl. Instr. Meth.* **A312**(1992)349 [T_{1/2}]
 1993Bh04 - M. R. Bhat, *Nucl. Data Sheets* **69**(1993)209 [multipolarities, mixing ratios, J^π]
 1995ScZY - E. Schönfeld, report PTB-6.33-95-2 (1995) [P_K, P_L, P_M theory]
 1996Sc06 - E. Schönfeld, H. Janßen, *Nucl. Instr. Meth.* **A369**(1996)527 [ω_K, ω_L, Auger emis. prob.]
 2000He14 - R. G. Helmer and C. van der Leun, *Nucl. Instr. Meth.* **A450**(2000)35 [E_γ]
 2002Ba85 - I.M.Band, M.B.Trazhaskovskaya, C.W.Nestor, S.Raman. *At. Data and Nucl. Data Tables* **81**, 1&2 (2002) 1 [ICC]
 2002Un02 - M.P. Unterweger, *Applied Radiation Isotopes* **56** (2002) 125 [T_{1/2}]
 2003Lu06 - A. Luca, M.-N. Amiot, J. Morel, *Applied Radiation Isotopes* **58** (2003) 607 [T_{1/2}]
 2003Au03 - G. Audi, A. H. Wapstra, C.Thibault. *Nucl. Phys.* **A729**(2003)337 [Q]
 2004Sc04 - H. Schrader, *Applied Radiation Isotopes* **60** (2004) 317 [T_{1/2}]
 2004Va02 - R. Van Ammel, S. Pommé, G. Sibbens, *Applied Radiation Isotopes* **60** (2004) 337 [T_{1/2}]
 2005Be** - M.-M.Bé, Euromet 721, Report CEA R-6081. CEA, F-91191 Gif-sur-Yvette Cedex.
 2006Be** - M.-M.Bé, *Applied Radiation Isotopes* **64** (2006) 1396 [P_γ]

⁶⁶Ga – Comments on evaluation of decay data by E. Browne

1. Statistical Analysis of Data

The *Limitation of Relative Statistical Weight* (LWM) [1985ZiZY] method, used for averaging numbers throughout this evaluation, provided a uniform approach for the analysis of discrepant data. The uncertainty assigned in this evaluation to the recommended value is always greater than or equal to the smallest uncertainty in any of the experimental values used in the calculation.

2. Decay Scheme

⁶⁶Ga decays 56 (4) % by positron (β^+) emission and 44 (4) % by electron capture (ϵ) to ⁶⁶Zn ($Q(\epsilon) = 5175(3)$ keV (1995Au04)). About 140 γ -rays de-exciting 31 nuclear levels in ⁶⁶Zn are known. Emission of conversion electrons is very low and negligible compared to that of γ rays (photons) because of the low atomic number ($Z = 30$) of the daughter nucleus (⁶⁶Zn) and the high energy (> 1000 keV) of the most intense γ -ray transitions. Consequently, neither conversion coefficients (most of them $< 2 \times 10^{-4}$) nor a list of conversion electrons is given in this evaluation.

Evaluator has normalized the decay scheme using experimental results from 1960Sc06, decay scheme information, and theory. As expected from the spins and parities of ⁶⁶Ga (0+) and ⁶⁶Zn (0+), there is a significant $\epsilon + \beta^+$ feeding (51(4)%) to the ground state of ⁶⁶Zn. Electron-capture and β^+ transition probabilities to excited states in ⁶⁶Zn given in Section 2.1 are from γ -ray transition probability balance at each level and theoretical ϵ/β^+ ratios. The decay scheme shown here is that of 1998Bh02 with the addition of levels half-lives from 2002Ga20.

3. Nuclear Data

The recommended half-life of ⁶⁶Ga, 9.49(7) hours, is a weighted average (LWM, $\chi^2/\nu=2.9$) of 9.57(6) hours (1956Ru45), 9.50(10) hours (1959Ca15), and 9.33(8) hours (1964Ru06). Other values are: 9.45 hours (1950La55), and 9.35 hours (1967Va13).

$Q(\epsilon)=5175(3)$ keV is from 1995Au04.

4. Gamma Rays

Energies

γ -ray energies in Table 1 given in boldface are from 2000He14. These values are based on a revised energy scale that uses the new adjusted fundamental constants and wave lengths deduced from an updated value of the lattice spacing of Si crystals [Cohen and Taylor [1]]. Helmer and van der Leun (2000He14) fitted the adjusted γ -ray energies of ⁶⁶Ga to a level scheme, and deduced their recommended values from level-energy differences. Less precise energies are from 1993Al15 and 1994En02, but adjusted to those of 2000He14 using a least-squares procedure. Evaluator has considered the difference between these two energy scales to be a systematic adjustment that he applied to the recommended energies given here. Thus, the uncertainties in the γ -ray energies given in this evaluation are just statistical, as reported by authors. See Table 1.

Emission Intensities

The relative emission probabilities of the most intense γ rays (given in boldface) in Table 2 are values recommended in 2002Ba38 and in this evaluation. These are weighted averages (LWM) of results from Berkeley, Budapest, and of 2000Ra36. Some of the uncertainties given in 2002Ba38, however, may be smaller than those given here, which are always greater than or equal to the smallest uncertainty in any of the experimental values used in the calculation.

Relative emission probabilities of other γ rays are weighted averages (LWM) of values from 1970Ph01, 1971Ca14, and 1994En02, each corrected by evaluator for a systematic error in the detector efficiency above ~ 1100 keV. This error was caused by an inadequate extrapolation of the detector efficiency to higher energies, and affected its value by as much as 30% at 4806 keV (1975Mc07).

The correction factor ($F = 1.116 - 0.155 E_{\gamma}(\text{MeV}) + 0.0397 E_{\gamma}^2(\text{MeV})$) given in 2002Ba38 has been used here. Uncertainties in the recommended relative emission probabilities are only statistical and have been deduced from those given in the individual measurements (see Table 2).

Absolute emission intensities given here are based on experimental results and decay scheme normalization arguments as follows:

$$I_{\text{ce}}(1039 \gamma)/I_{\beta^+}(\text{gs}) = 2.08(10) \times 10^{-4} \quad (1960\text{Sc06})$$

$$I_{\beta^+}(\text{gs})/\Sigma I_{\beta^+} = 0.8697 \quad (1960\text{Sc06})$$

$$I_{\text{ce}}(1039 \gamma, E2)/I_{\gamma}(1039 \gamma) = 2.69(8) \times 10^{-4} \quad (\text{Theory, 1978Rö22}).$$

Therefore,

$$I_{\gamma}(1039 \gamma)/\Sigma I_{\beta^+} = 2.08(10) \times 10^{-4} \times 0.8697/2.69(8) \times 10^{-4} = 0.67(4).$$

Also $\Sigma I_{\beta^+}/\Sigma I_{\text{ei}} = 1.265$ from decay scheme and theoretical values of I_{β^+}/ϵ_i for each level. Using

$$\Sigma I_{\beta^+} + \Sigma I_{\text{ei}} = 100 \%, \text{ gives } \Sigma I_{\beta^+} = 55.8(24) \%, \text{ and}$$

$$I_{\gamma}(1039 \gamma) = 0.67(4) \times 55.8(24) = 37(3) \ %.$$

Absolute γ -ray emission intensities given in Section 5.2 are relative values multiplied by 0.37(3).

5. Positron (β^+) Transitions

Positron end-point energies given in section 2.1.1 ($E_{\beta^+} = Q(\epsilon) - E(\text{keV}) - 1022$) are evaluator's values deduced using $Q(\epsilon) = 5175(3)$ keV (1995Au04) and level energies ($E(\text{keV})$) from decay scheme. Absolute β^+ emission probabilities are from γ -ray intensity balance at each nuclear level and theoretical I_{β^+}/ϵ_i ratios.

6. Electron Capture (ϵ) Transitions

ϵ transition energies ($E(\epsilon) = Q(\epsilon) - E(\text{keV})$) are evaluator's values deduced using $Q(\epsilon) = 5175(3)$ keV (1995Au04) and level energies ($E(\text{keV})$) from decay scheme. Absolute ϵ transition probabilities are from γ -ray probability balance at each nuclear level and theoretical I_{β^+}/ϵ_i ratios. Fractional atomic shell electron-capture probabilities (P_K, P_L, P_M) are evaluator's values calculated using the EC-CAPTURE computer program [2] and the nuclear level energies presented here.

7. Atomic Data

The X-ray and Auger electron energies given in sections 3, 4 are from Schönfeld and Rodloff [4] and [5], respectively. Emission intensities are evaluator's values calculated using the EMISSION (Version V.3.04) [3] program, atomic data from 1996Sc06, and the recommended γ -ray emission intensities from section 5.2.

References

1. *The 1986 Adjustment of the Fundamental Physical Constants*, E.R. Cohen and B.N. Taylor, Rev. Mod. Phys. **59**, 1121 (1987).
 2. *The Program EC-CAPTURE*, E. Schönfeld, F. Chu, and E. Browne. An interactive computer program for calculating electron capture probabilities P_K , P_L , P_M , and P_N to the K, L, M, and N atomic shells, respectively (1997).
 3. *The Program EMISSION* (version 3.04 (2002)), E. Schönfeld and H. Janssen. A computer program for calculating emission probabilities of X-rays and Auger electrons emitted in nuclear disintegration processes.
 4. *Tables of the energies of K-Auger electrons for elements with atomic numbers in the range from Z=11 to Z=100*, E. Schönfeld, G. Rodloff, Report PTB-6.11-98-1, October 1998.
 5. *Energies and relative emission probabilities of K X-rays for elements with atomic numbers in the range from Z=5 to Z=100*, E. Schönfeld, G. Rodloff, Report PTB-6.11-1999-1, February 1999.
- 1960Sc06 - A. Schwarzschild, L. Grodzins, Phys. Rev. **119**, 276 (1960).
- 1970Ph01 - M. E. Phelps, D. G. Sarantites, W. G. Winn, Nucl. Phys. **A149**, 647 (1970).
(relative γ -ray emission probabilities)
- 1971Ca14 - D. C. Camp, G. L. Meredith, Nucl. Phys. **A166**, 349 (1971).
(relative γ -ray emission probabilities)
- 1975Mc07 - G. J. MacCallum, G. E. Coote, Nucl. Instrum. Methods **124**, 309 (1975).
(relative γ -ray emission probabilities)
- 1978Rö22 - F. Rösel, H. M. Friess, K. Alder, H. C. Pauli, At. Data. Nucl. Data Tables **21**, 92 (1978).
(theoretical conversion coefficients)
- 1985ZiZY - W. L. Zijp, Report ECN FYS/RASA-85/19 (1985).
(Discrepant Data Limited Relative Statistical Weight Method).
- 1993Al15 - C. Alderliesten, J. A. van Nie, A. P. Slok, P. M. Endt, Nucl. Instrum. Methods. Phys. Res. **A335**, 219 (1993).
(γ -ray energies)
- 1994En02 - P. M. Endt, C. Alderliesten, Nucl. Phys. **A575**, 297 (1994).
(γ -ray energies and relative emission probabilities)
- 1995Au04 - G. Audi, A. H. Wapstra, Nucl. Phys. **A595**, 409 (1995).
(Q_α)
- 1996Sc06 - E. Schönfeld, H. Janssen, Nucl. Instrum. Methods. Phys. Res. **A369**, 527 (1996).
(Atomic data, X-rays, Auger electrons)
- 1998Bh02 - M. R. Bhat, Nuclear Data Sheets 83, 789 (1998).
(⁶⁶Ga decay scheme)
- 2000He14 - R. G. Helmer, C. van der Leun, Nucl. Instrum. Methods. Phys. Res. **A450**, 35 (2000).
(Precise evaluated γ -ray energies)
- 2000Ra36 - S. Raman, C. Yonezawa, H. Matsue, H. Imura, N. Shinohara, Nucl. Instrum. Methods. Phys. Res. **A454**, 89 (2000).
(Precise relative γ -ray emission probabilities)
- 2002Ba38 - C. M. Baglin, E. Browne, E. B. Norman, G. L. Molnar, T. Belgya, Zs. Revay, F. Szelecsenyi, Nucl. Instrum. Meth. Phys. Res. **A481**, 365(2002).
(Precise relative γ -ray emission probabilities)
- 2002Ga20 - A. Gade, H. Klein, N. Pietralla, and P. von Brentano, Phys. Rev. C **65**, 054311 (2002)
(Levels half-life in ⁶⁶Zn)

Table 1. ⁶⁶Ga Gamma-Ray Energies

1993Al15, 1994En02	1993Al15 DE _g (keV)	2000He14 E _g (keV)	2000He14 DE _g (keV)	Fitted E _g (keV)
171.9	0.2			171.9 (2)
283.87	0.03			283.87 (3)
290.808	0.011			290.8105(11)
347.77	0.05			347.77 (5)
375.396	0.017			375.398 (17)
410.177	0.012			410.178 (12)
412.915	0.016			412.916 (16)
442.872	0.014			442.873 (14)
448.725	0.02			448.73 (2)
459.682	0.014			459.683 (14)
494.336	0.013			494.336 (13)
499.59	0.006			499.590 (6)
551.284	0.022			551.284 (22)
554.28	0.03			554.28 (3)
557.13	0.05			557.13(5)
562.241	0.01			562.241 (10)
578.54	0.019			578.540 (19)
600.789	0.021			600.788 (21)
653.569	0.014			653.568 (14)
658.57	0.03			658.57 (3)
670.252	0.014			670.251 (14)
680.56	0.1			680.56 (10)
<u>686.084</u>	0.007	686.080	0.006	686.080 (6)
705.033	0.015			705.031 (15)
708.36	0.05			708.36 (5)
718.97	0.05			718.97 (5)
723.17	0.05			723.17 (5)
749.68	0.1			749.68 (10)
763.64	0.03			763.64 (3)
796.21	0.05			796.21 (5)
800.13	0.05			800.13 (5)
<u>833.537</u>	0.003	833.5324	0.0021	833.5324 (21)
<u>853.046</u>	0.009	853.038	0.008	853.038 (8)
856.53	0.01			856.527 (10)
857.096	0.009			857.093 (9)
862.929	0.013			862.926 (13)
867.93	0.03			867.93 (3)
873.395	0.021			873.392 (21)
885	0.05			885.00 (5)
907.394	0.019			907.390 (19)
914.392	0.014			914.388 (14)
929.68	0.03			929.68 (3)
953.93	0.09			953.93 (9)
954.12	0.07			954.12 (7)
963.896	0.015			963.892 (15)
980.938	0.013			980.934 (13)
1008.593	0.012			1008.588 (12)
1010.962	0.019			1010.957 (19)
1015.086	0.018			1015.081 (18)
<u>1039.231</u>	0.006	1039.22	0.003	1039.220 (3)

1993Al15, 1994En02 E _g (keV)	1993Al15 DE _g (keV)	2000He14 E _g (keV)	2000He14 DE _g (keV)	Fitted E _g (keV)
1060.056	0.011			1060.051 (11)
1065.31	0.009			1065.305 (9)
1066.455	0.012			1066.450 (12)
1082.754	0.02			1082.75 (2)
1106.54	0.24			1106.53 (24)
1129.929	0.018			1129.923 (18)
1135.48	0.09			1135.47 (9)
<u>1147.9</u>	0.012	1147.896	0.010	1147.896 (10)
<u>1190.297</u>	0.008	1190.287	0.007	1190.287 (7)
1195.33	0.09			1195.32 (9)
1232.271	0.008			1232.264 (8)
1232.487	0.015			1232.480 (15)
1248.786	0.022			1248.779 (22)
1274.51	0.03			1274.50 (3)
1298.96	0.07			1298.95 (7)
1305.815	0.021			1305.807 (21)
<u>1333.12</u>	0.006	1333.112	0.005	1333.112 (5)
1356.112	0.009			1356.104 (9)
1356.328	0.015			1356.320 (15)
1357.258	0.012			1357.250 (12)
1409.36	0.24			1409.35 (24)
<u>1418.763</u>	0.006	1418.754	0.005	1418.754 (5)
1425.256	0.02			1425.25 (2)
1433.64	0.04			1433.63 (4)
<u>1458.67</u>	0.012	1458.662	0.012	1458.662 (12)
1468.98	0.05			1468.97 (5)
<u>1508.175</u>	0.011	1508.158	0.007	1508.158 (7)
1515.172	0.02			1515.162 (20)
1523.289	0.015			1523.279 (15)
1534.61	0.04			1534.60 (4)
1554.63	0.03			1554.62 (3)
1559.637	0.01			1559.627 (10)
1577.318	0.02			1577.308 (20)
1634.47	0.07			1634.46 (7)
1703.6	0.05			1703.59 (5)
1713.614	0.012			1713.602 (12)
<u>1740.918</u>	0.018	1740.904	0.016	1740.904 (16)
1787.45	0.09			1787.44 (9)
1797.95	0.09			1797.94 (9)
1868.118	0.02			1868.105 (20)
1872.753	0.006			1872.740 (6)
<u>1898.832</u>	0.009	1898.823	0.008	1898.823 (8)
<u>1918.341</u>	0.006	1918.329	0.005	1918.329 (5)
1927.97	0.04			1927.96 (4)
2009.643	0.016			2009.628 (16)
2026.031	0.025			2026.016 (25)
<u>2065.792</u>	0.008	2065.778	0.007	2065.778 (7)
2085.88	0.04			2085.86 (4)
2089	0.013			2088.985 (13)
<u>2173.334</u>	0.018	2173.319	0.015	2173.319 (15)
<u>2189.631</u>	0.009	2189.616	0.006	2189.616 (6)

Comments on evaluation

1993Al15, 1994En02	1993Al15	2000He14	2000He14	Fitted
E_g (keV)	DE_g (keV)	E_g (keV)	DE_g (keV)	E_g (keV)
<u>2213.19</u>	0.011	2213.181	0.009	2213.181 (9)
2265.86	0.24			2265.84 (24)
2292.188	0.013			2292.171 (13)
2341.691	0.011			2341.673 (11)
<u>2393.153</u>	0.01	2393.129	0.007	2393.129 (7)
<u>2422.544</u>	0.009	2422.525	0.007	2422.525 (7)
2433.826	0.018			2433.807 (18)
2467.99	0.07			2467.97 (7)
2492.44	0.03			2492.42 (3)
2537.11	0.05			2537.09 (5)
2588.573	0.013			2588.553 (13)
2631.46	0.09			2631.44 (9)
2698.94	0.05			2698.92 (5)
2713.75	0.05			2713.73 (5)
<u>2751.852</u>	0.006	2751.835	0.005	2751.835 (5)
<u>2780.12</u>	0.018	2780.095	0.016	2780.095 (16)
2785.7	0.3			2785.7 (3)
2802.8	0.5			2802.8 (5)
2843.153	0.016			2843.130 (16)
<u>2933.395</u>	0.017	2933.358	0.009	2933.358 (9)
<u>2977.12</u>	0.05	2977.083	0.043	2977.083 (43)
<u>2993.25</u>	0.04	2993.208	0.032	2993.208 (32)
<u>3046.697</u>	0.011	3046.684	0.009	3046.684 (9)
3085.4	0.4			3085.4 (4)
3212.526	0.019			3212.499 (19)
<u>3228.824</u>	0.009	3228.800	0.006	3228.800 (6)
3256.048	0.009			3256.021 (9)
3331.379	0.014			3331.351 (14)
<u>3380.882</u>	0.01	3380.850	0.006	3380.850 (6)
<u>3422.075</u>	0.012	3422.040	0.008	3422.040 (8)
<u>3432.343</u>	0.01	3432.309	0.007	3432.309 (7)
3738.13	0.05			3738.10 (5)
<u>3766.893</u>	0.018	3766.850	0.009	3766.850 (9)
3791.036	0.008			3791.004 (8)
3810.62	0.05			3810.59 (5)
<u>4085.875</u>	0.012	4085.853	0.009	4085.853 (9)
4295.224	0.01			4295.187 (10)
<u>4461.247</u>	0.013	4461.202	0.009	4461.202 (9)
<u>4806.06</u>	0.018	4806.007	0.009	4806.007 (9)
4865.91	0.04			4865.87 (4)
5005.62	0.23			5005.6 (3)

Y= A + BX and input energies (X) from 1994En02.

Table 2: ⁶⁶Ga Relative

Eg (keV)	1970Ph01		1971Ca14		Gamma-Ray 1994En02 lg	Emission 1994En02* lg(Corr.)	Intensities			Remarks	
	1970Ph01* lg	1970Ph01* lg(Corr.)	1971Ca14 lg	1971Ca14* lg(Corr.)			2000Ra00 lg	Berkeley lg	Budapest lg		Recomm. lg
171.9 (2)	0.028 (1)	0.028 (1)								0.028 (1)	O
283.87 (3)					0.0097 (21)	0.0097 (21)				0.0097 (21)	I
290.8105 (11)	0.150 (10)	0.150 (10)	0.131 (2)	0.131 (2)	0.146 (6)	0.146 (6)				0.133 (4)	A
347.77 (5)					0.0048 (15)	0.0048 (15)				0.0048 (15)	I
375.398 (17)					0.0058 (16)	0.0058 (16)				0.0058 (16)	I
410.178 (12)	0.300 (20)	0.300 (20)	0.172 (24)	0.172 (24)	0.177 (7)	0.177 (7)				0.177 (7)	I
412.916 (16)					0.0091 (13)	0.0091 (13)				0.0091 (13)	I
442.873 (14)					0.042 (3)	0.042 (3)				0.042 (3)	I
448.73 (2)	0.290 (10)	0.290 (10)	0.279 (58)	0.279 (58)						0.290 (10)	C
459.683 (14)	0.240 (10)	0.240 (10)	0.206 (35)	0.206 (35)						0.237 (10)	C
494.336 (13)					0.0152 (20)	0.0152 (20)				0.0152 (20)	I
499.590 (6)					0.013 (3)	0.013 (3)				0.013 (3)	I
551.284 (22)					0.0189 (16)	0.0189 (16)				0.0189 (16)	I
554.28 (3)					0.0122 (13)	0.0122 (13)				0.0122 (13)	I
557.13(5)					0.0166 (17)	0.0166 (17)				0.0166 (17)	I
562.241 (10)					0.0179 (17)	0.0179 (17)				0.0179 (17)	I
578.540 (19)	0.160 (10)	0.160 (10)	0.156 (20)	0.156 (20)						0.159 (10)	C
600.788 (21)					0.0365 (23)	0.0365 (23)				0.0365 (23)	I
653.568 (14)					0.0036 (12)	0.0036 (12)				0.0036 (12)	I
658.57 (3)					0.0203 (21)	0.0203 (21)				0.0203 (21)	I
670.251 (14)					0.0110 (18)	0.0110 (18)				0.0110 (18)	I
680.56 (10)					0.0040 (11)	0.0040 (11)				0.0040 (11)	I
686.080 (6)	0.690 (20)	0.690 (20)	0.645 (40)	0.645 (40)						0.681 (20)	C
705.031 (15)					0.0102 (11)	0.0102 (11)				0.0102 (11)	I
708.36 (5)					0.0234 (19)	0.0234 (19)				0.0234 (19)	I
718.97 (5)					0.0268 (20)	0.0268 (20)				0.0268 (20)	I
723.17 (5)					0.0093 (13)	0.0093 (13)				0.0093 (13)	I
749.68 (10)					0.0037 (11)	0.0037 (11)				0.0037 (11)	I
763.64 (3)					0.0240 (20)	0.0240 (20)				0.0240 (20)	I
796.21 (5)					0.0079 (17)	0.0079 (17)				0.0079 (17)	I
800.13 (5)					0.0027 (14)	0.0027 (14)				0.0027 (14)	I
833.5324 (21)	16.2 (7)	16.2 (7)	15.92 (17)	15.92 (17)			16.02 (24)	15.94 (14)	15.92 (6)	15.93 (6)	K
853.038 (8)			0.200 (5)	0.200 (5)	0.232 (12)	0.232 (12)				0.205 (5)	D
856.527 (10)			0.315 (10)	0.315 (10)	0.280 (12)	0.280 (12)				0.301 (17)	D

Comments on evaluation

⁶⁶Ga

Recomm. E _g (keV)	1970Ph01 lg	1970Ph01* lg(Corr.)	1971Ca14 lg	1971Ca14* lg(Corr.)	1994En02 lg	1994En02* lg(Corr.)	2000Ra00 lg	Berkeley lg	Budapest lg	Recomm. lg	Remarks
857.093 (9)					0.040 (12)	0.040 (12)				0.040 (12)	I
862.926 (13)					0.0410 (20)	0.0410 (20)				0.0410 (20)	I
867.93 (3)					0.0117 (14)	0.0117 (14)				0.0117 (14)	I
873.392 (21)					0.046 (3)	0.046 (3)				0.046 (3)	I
885.00 (5)					0.0051 (13)	0.0051 (13)				0.0051 (13)	I
907.390 (19)	0.300 (20)	0.300 (20)	<0.034 (10)		0.059 (4)	0.059 (4)				0.059 (4)	E
914.388 (14)	0.190 (10)	0.190 (10)	<0.030 (10)		0.073 (4)	0.073 (4)				0.073 (4)	E
929.68 (3)					0.0123 (15)	0.0123 (15)				0.0123 (15)	I
953.93 (9)					0.0027 (3)	0.0027 (3)				0.0027 (3)	I
954.12 (7)					0.0121 (17)	0.0121 (17)				0.0121 (17)	I
963.892 (15)					0.039 (3)	0.039 (3)				0.039 (3)	I
980.934 (13)	0.150 (20)	0.150 (20)	0.130 (5)	0.130 (5)						0.131 (5)	C
1008.588 (12)	0.183 (10)	0.183 (10)	0.138 (4)	0.138 (4)						0.160 (20)	C
1010.957 (19)					0.073 (4)	0.073 (4)				0.073 (4)	I
1015.081 (18)					0.033 (8)	0.033 (8)				0.033 (8)	I
1039.220 (3)	100	100	100	100	100	100	100.0 (16)	100.0 (9)	100.0 (3)	100.0 (3)	K
1060.051 (11)			0.033 (10)	0.033 (10)	0.043 (3)	0.043 (3)				0.042 (3)	F
1065.305 (9)					0.0063 (12)	0.0063 (12)				0.0063 (12)	I
1066.450 (12)					0.0064 (12)	0.0064 (12)				0.0064 (12)	I
1082.75 (2)					0.036 (2)	0.0358 (20)				0.0358 (20)	I
1106.53 (24)					0.0033 (10)	0.0033 (10)				0.0033 (10)	I
1129.923 (18)					0.0370 (21)	0.0367 (21)				0.0367 (21)	I
1135.47 (9)					0.0128 (13)	0.0128 (13)				0.0128 (13)	I
1147.896 (10)	0.22 (3)	0.22 (3)	0.211 (17)	0.211 (17)						0.212 (17)	C
1190.287 (7)	0.42 (4)	0.42 (4)	0.34 (1)	0.34 (1)						0.345 (19)	C
1195.32 (9)					0.0025 (9)	0.0025 (9)				0.0025 (9)	I
1232.264 (8)	1.14 (20)	1.12 (20)	1.38 (4)	1.36 (4)						1.35 (5)	C
1232.480 (15)	0.4 (2)	0.4 (2)	0.14 (4)	0.14 (4)						0.15 (5)	C
1248.779 (22)					0.0027 (9)	0.0027 (9)				0.0027 (9)	I
1274.50 (3)					0.0192 (15)	0.0189 (15)				0.0189 (15)	I
1298.95 (7)					0.0105 (12)	0.0103 (12)				0.0103 (12)	I
1305.807 (21)					0.0109 (12)	0.0107 (12)				0.0107 (12)	I
1333.112 (5)	3.28 (5)	3.21 (5)	3.25 (4)	3.18 (4)			3.17 (5)	3.20 (3)	3.171 (13)	3.175 (13)	K
1356.104 (9)	0.83 (30)	0.81 (30)	1.00 (10)	0.98 (10)						0.96 (10)	C
1356.320 (15)	0.3 (1)	0.29 (10)	0.35 (5)	0.34 (5)						0.33 (5)	C

Comments on evaluation

⁶⁶Ga

Recomm. E _g (keV)	1970Ph01 lg	1970Ph01* lg(Corr.)	1971Ca14 lg	1971Ca14* lg(Corr.)	1994En02 lg	1994En02* lg(Corr.)	2000Ra00 lg	Berkeley lg	Budapest lg	Recomm. lg	Remarks
1357.250 (12)	0.7 (2)	0.69 (20)	0.39 (10)	0.38 (10)						0.44 (13)	C
1409.35 (24)					0.0044 (18)	0.0043 (18)				0.0043 (18)	I
1418.754 (5)	1.65 (3)	1.61 (3)	1.700 (27)	1.659 (27)				1.640 (23)	1.659 (8)	1.657 (8)	M
1425.25 (2)					0.0167 (13)	0.0163 (13)				0.0163 (13)	I
1433.63 (4)					0.0050 (10)	0.0050 (10)				0.0050 (10)	I
1458.662 (12)	0.25 (7)	0.24 (7)	0.268 (6)	0.261 (6)						0.261 (6)	C
1468.97 (5)					0.0038 (10)	0.0037 (10)				0.0037 (10)	I
1508.158 (7)	1.48 (9)	1.44 (9)	1.520 (24)	1.478 (24)				1.503 (23)	1.496 (7)	1.497 (7)	M
1515.162 (20)					0.0172 (15)	0.0167 (15)				0.0167 (15)	I
1523.279 (15)					0.0152 (13)	0.0148 (13)				0.0148 (13)	I
1534.60 (4)					0.016 (4)	0.0155 (40)				0.016 (4)	I
1554.62 (3)					0.051 (3)	0.049 (3)				0.050 (3)	I
1559.627 (10)					0.061 (4)	0.059 (4)				0.059 (4)	I
1577.308 (20)					0.0111 (16)	0.0108 (16)				0.0108 (16)	I
1634.46 (7)					0.0098 (15)	0.0095 (15)				0.0095 (15)	I
1703.59 (5)					0.015 (5)	0.015 (5)				0.015 (5)	I
1713.602 (12)					0.068 (3)	0.066 (3)				0.066 (3)	I
1740.904 (16)	0.19 (4)	0.18 (4)	0.0800 (10)	0.0773 (10)						0.0773 (10)	G
1787.44 (9)					0.025 (2)	0.0240 (20)				0.0240 (20)	I
1797.94 (9)					0.0053 (14)	0.0051 (14)				0.0051 (14)	I
1868.105 (20)					0.0076 (15)	0.0073 (15)				0.0073 (15)	I
1872.740 (6)					0.064 (4)	0.062 (4)				0.062 (4)	I
1898.823 (8)	1.15 (3)	1.11 (3)	1.09 (4)	1.05 (4)				1.062 (23)	1.050 (8)	1.051 (8)	M
1918.329 (5)	5.65 (2)	5.45 (2)	5.625 (80)	5.427 (80)			5.33 (8)	5.44 (6)	5.360 (23)	5.368 (23)	K
1927.96 (4)					0.0063 (20)	0.0061 (20)				0.0061 (20)	I
2009.628 (16)					0.0086 (17)	0.0083 (17)				0.0083 (17)	I
2026.016 (25)					0.0073 (16)	0.0070 (16)				0.0070 (16)	I
2065.778 (7)	0.098 (16)	0.095 (16)	0.086 (4)	0.083 (4)						0.084 (4)	C
2085.86 (4)					0.006 (4)	0.0058 (40)				0.006 (4)	I
2088.985 (13)					0.032 (7)	0.031 (7)				0.031 (7)	I
2173.319 (15)	0.38 (3)	0.37 (3)	0.236 (12)	0.228 (12)						0.228 (12)	G
2189.616 (6)	15.0 (3)	14.5 (3)	15.06 (18)	14.56 (18)			14.54 (21)	14.50 (13)	14.39 (6)	14.42 (6)	K
2213.181 (9)	0.38 (5)	0.37 (5)	0.365 (12)	0.353 (12)						0.354 (12)	C
2265.84 (24)					0.0038 (14)	0.0037 (14)				0.0037 (14)	I
2292.171 (13)			0.110 (10)	0.107 (10)	0.047 (3)	0.046 (3)				0.046 (3)	H

Comments on evaluation

⁶⁶Ga

Recomm. E _g (keV)	1970Ph01 I _g	1970Ph01* I _g (Corr.)	1971Ca14 I _g	1971Ca14* I _g (Corr.)	1994En02 I _g	1994En02* I _g (Corr.)	2000Ra00 I _g	Berkeley I _g	Budapest I _g	Recomm. I _g	Remarks
2341.673 (11)					0.0089 (17)	0.0086 (17)				0.0086 (17)	I
2393.129 (7)	0.64 (2)	0.62 (2)	0.670 (20)	0.651 (20)						0.635 (20)	C
2422.525 (7)	5.06 (10)	4.93 (10)	5.16 (5)	5.023 (5)			5.12 (8)	5.15 (6)	5.072 (24)	5.085 (24)	K
2433.807 (18)					0.0206 (17)	0.0201 (17)				0.0201 (17)	I
2467.97 (7)					0.0234 (19)	0.0228 (19)				0.0228 (19)	I
2492.42 (3)			0.063 (6)	0.061 (6)	0.061 (4)	0.060 (4)				0.060 (4)	F
2537.09 (5)					0.014 (3)	0.014 (3)				0.014 (3)	I
2588.553 (13)			0.073 (7)	0.072 (7)	0.072 (4)	0.071 (4)				0.071 (4)	F
2631.44 (9)					0.008 (3)	0.008 (3)				0.008 (3)	I
2698.92 (5)					0.0101 (17)	0.0100 (17)				0.0100 (17)	I
2713.73 (5)					0.017 (5)	0.017 (5)				0.017 (5)	I
2751.835 (5)	60.9 (8)	60.3 (8)	61.2 (6)	60.6 (6)			61.2 (8)	61.5 (6)	61.34 (26)	61.35 (26)	K
2780.095 (16)	0.33 (2)	0.33 (2)	0.337 (8)	0.334 (8)						0.334 (8)	C
2785.7 (3)					0.0081 (14)	0.0080 (14)				0.0080 (14)	I
2802.8 (5)					0.0040 (11)	0.0040 (11)				0.0040 (11)	I
2843.130 (16)					0.0045 (9)	0.0045 (9)				0.0045 (9)	I
2933.358 (9)	0.57 (3)	0.57 (3)	0.574 (8)	0.576 (8)						0.576 (8)	C
2977.083 (43)			0.062 (6)	0.062 (6)						0.062 (6)	N
2993.208 (32)			0.084 (8)	0.085 (8)						0.085 (8)	N
3046.684 (9)	0.17 (2)	0.17 (2)	0.150 (6)	0.152 (6)						0.154 (6)	C
3085.4 (4)					0.0052 (13)	0.0053 (13)				0.0053 (13)	I
3212.499 (19)					0.0049 (10)	0.0050 (10)				0.0050 (10)	I
3228.800 (6)	3.85 (6)	3.96 (6)	3.96 (4)	4.08 (4)			4.06 (8)	4.07 (4)	4.087 (22)	4.082 (22)	K
3256.021 (9)	0.31 (3)	0.32 (3)	0.241 (5)	0.249 (5)	0.270 (14)	0.279 (14)				0.254 (10)	A
3331.351 (14)					0.0059 (8)	0.0061 (8)				0.0061 (8)	I
3380.850 (6)	3.68 (4)	3.85 (4)	3.78 (4)	3.95 (4)			3.96 (8)	3.99 (4)	3.950 (23)	3.960 (23)	K
3422.040 (8)	2.10 (9)	2.21 (9)	2.18 (4)	2.29 (4)				2.29 (3)	2.321 (16)	2.314 (16)	M
3432.309 (7)	0.73 (3)	0.77 (3)	0.740 (10)	0.778 (10)						0.777 (10)	C
3724.8 (10)			0.0060 (10)	0.0065 (10)						0.0065 (10)	N
3738.10 (5)			0.032 (3)	0.035 (3)	0.0353 (20)	0.0385 (20)				0.0374 (20)	F
3766.850 (9)	0.37 (2)	0.41 (2)	0.364 (14)	0.399 (15)						0.403 (15)	C
3791.004 (8)	2.63 (11)	2.89 (11)	2.675 (32)	2.940 (35)			2.96 (5)	2.96 (4)	2.929 (24)	2.941 (24)	K
3806.3 (10)			0.0060 (10)	0.0066 (10)						0.0066 (11)	N
3810.59 (5)			0.0210 (20)	0.0231 (22)	0.025 (3)	0.028 (3)				0.0248 (22)	F
3827.5 (8)			0.0170 (20)	0.0190 (22)						0.0190 (22)	N

Recomm.	1970Ph01	1970Ph01*	1971Ca14	1971Ca14*	1994En02	1994En02*	2000Ra00	Berkeley	Budapest	Recomm.	Remarks
E _g (keV)	I _g	I _g (Corr.)	I _g	I _g (Corr.)	I _g	I _g (Corr.)	I _g	I _g	I _g	I _g	
4085.853 (9)	2.91 (6)	3.33 (7)	3.07 (4)	3.52 (5)			3.38 (8)	3.42 (4)	3.455 (20)	3.445 (20)	K
4295.187 (10)	9.2 (2)	10.88 (24)	9.17 (11)	10.84 (13)			10.24 (26)	10.54 (15)	10.25 (7)	10.30 (8)	K, L
4461.202 (9)	1.84 (4)	2.23 (5)	1.875 (22)	2.277 (27)				2.20 (4)	2.275 (23)	2.26 (3)	M
4806.007 (9)	3.96 (6)	5.10 (6)	3.82 (4)	4.92 (4)			4.93 (11)	5.00 (7)	5.04 (3)	5.03 (3)	K
4865.87 (4)					0.0058 (5)	0.0075 (6)				0.0075 (6)	I
5005.6 (3)					0.0025 (3)	0.0033 (4)				0.0033 (4)	I

*γ-ray intensities (I_γ) corrected for a systematic inaccuracy in the detector efficiency curve above 1050 keV.

Correction factor $f = 1.116 - 0.155 E_{\gamma} (\text{MeV}) + 0.0397 E_{\gamma} \times E_{\gamma}$ (2002Ba38). Uncertainties are statistical values given by authors.

A: Weighted average of values from 1970Ph01, 1971Ca14, and 1994En02

B: Weighted average of values from 1971Ca14 and 1994En02. Value from 1970Ph01 is too high (peak may contain impurities).

C: Weighted average of values from 1970Ph01 and 1971Ca14.

D: Weighted average of values from 1970Ph01 and 1994En02.

E: From 1994En02. Value from 1970Ph01 is too high (peak may contain impurities).

F: Weighted average of values from 1971Ca14 and 1994En02.

G: From 1971Ca14. Value from 1970Ph01 is too high (peak may contain impurities).

H: From 1994En02. Value from 1971Ca14 is too high (peak may contain impurities).

I: From 1994En02.

K: Weighted average (in boldface) of values from 2000Ra36, from Berkeley, and from Budapest, as given in 2002Ba38 (except for the recommended uncertainties, which are never smaller than the smallest experimental uncertainty).

L: After correction for single-escape contribution from the 4806-keV line.

M; Weighted average (in boldface) of values from Berkeley and Budapest, as given in 2002Ba38

N: From 1971Ca14

O: Reported only by 1970Ph01.

⁶⁷Ga – Comments on evaluation of decay data by X. Mougeot and V.P. Chechev

The initial evaluation was completed in March 2000. This revised evaluation was done in 2011, taking into account the available literature by March 2011.

1. Decay Scheme

The spins and parities of the ground state of ⁶⁷Ga and of the levels of ⁶⁷Zn are from the evaluation of 2005HU18.

The main difficulty in the evaluation of the ⁶⁷Ga decay scheme is connected with the lack of measurements of the absolute intensity of the internal conversion electron component $P(\text{ce}_{1,0})$ from the 93 keV gamma transition (2000SI03). This value determines directly the probability $P(\varepsilon_{0,0})$ of the allowed electron capture transition to the ground state of ⁶⁷Zn. In many evaluations, including 1991BH06, $P(\varepsilon_{0,0})$ was adopted equal to zero. In a more recent evaluation (2005HU18), a value of 0,9 (9) % was adopted.

In this evaluation of the ⁶⁷Ga decay scheme, four measurements of $P(\text{ce}_{1,0})$ were taken into account: 1998AT04, 2000SI03, 2005YA01 and 2007BO. The analysis led to the evaluated value of $P(\text{ce}_{1,0}) = 0,3254$ (40) and to the probability of the electron capture transition to the ⁶⁷Zn ground state $P(\varepsilon_{0,0}) = 3,3$ (32) % (see Section 4.2.2). The large uncertainty of $P(\varepsilon_{0,0})$ mainly comes from the uncertainty of the evaluated mixing ratio δ (184 keV).

Among the adopted levels in 2005HU18, three levels are placed below the decay energy and are not taken into account in this evaluation: the $9/2^+$ at 604,48 (5) keV, the $7/2^-$ at 814,90 (6) keV and the $5/2^+$ at 979,85 (5) keV. These levels could be fed respectively by 3rd, 2nd and 1st transitions. As all the other electronic capture transitions are allowed, the branch intensities for these three levels must be much lower, and precisely the corresponding gamma-rays were never observed.

2. Nuclear Data

Q value is from 2003AU03: $Q^+(\text{}^{67}\text{Ga}) = 1000,8$ (12) keV.

2.1 ⁶⁷Ga half-life

The measured half-life values of ⁶⁷Ga are summarized in Table 1. The values from 1948HO04 and 1950HO26 were not used in the evaluation because no experimental uncertainty was reported. The values from 1982HOZJ and 2002UN02 are the same values as respectively 1978ME10 and 1992UN01, and were not used in the evaluation.

1972CR02 gives eight measurements using the same method. As this data set is discrepant, an unweighted mean was chosen, calculated using the LWEIGHT program: $T_{1/2}(\text{}^{67}\text{Ga}) = 3,38$ (9) d.


The measured half-life values used in this evaluation are summarized in Figure 1. The statistical processing was done using the LWEIGHT program. Four statistical outliers are excluded according to Chauvenet's criterion: 1938MA01, 1955TO27, 1964RU06 and 1972CR02, in red in Figure 1. A weighted average is 3,26125 (35) d from the resulting consistent data set, with a reduced- χ^2 value of 0,74. The statistical weights are 44 % for 1992UN01 and 20 % for 1980HO17, the two most precise measurements. As there are less than ten values, and as the most precise measurements use the same method, the smallest experimental uncertainty was preferred. Finally, the adopted value is:

$$\mathbf{T_{1/2}(\text{}^{67}\text{Ga}) = 3,2613 (5) d.}$$

Table 1: ⁶⁷Ga half-life measurements. The excluded values are crossed out.

Reference	T _{1/2} (⁶⁷ Ga) measurements	T _{1/2} (⁶⁷ Ga) in d	Comments
1938MA01	79 (2) h	3,29 (8)	Excluded by Chauvenet's criterion
1948HO04	83 h	3,46	Not used: no uncertainty
1948MC32	3,26 (2) d	3,26 (2)	
1950HO26	80 h	3,33	Not used: no uncertainty
1955TO27	77,9 (3) h	3,246 (13)	Excluded by Chauvenet's criterion
1964RU06	79,2 (5) h	3,300 (21)	Excluded by Chauvenet's criterion
1972CR02	78,5 (15) h 78,2 (12) h 84,7 (23) h 79,1 (14) h 69,7 (36) h 84,2 (11) h 90,7 (42) h 83,8 (44) h	3,27 (6) 3,26 (5) 3,53 (10) 3,30 (6) 2,90 (15) 3,508 (46) 3,78 (18) 3,49 (18)	
1972CR02	LWEIGHT	3,38 (9)	Unweighted mean, excluded by Chauvenet's criterion
1972LE37	78,26 (7) h	3,2608 (29)	
1978LA21	78,33 (10) h	3,2638 (42)	
1978ME10	3,261 (1) d	3,261 (1)	
1979DE42	3,2594 (12) d	3,2594 (12)	
1980HO17	3,2607 (8) d	3,2607 (8)	
1982HOZJ	3,261 (1) d	3,261 (1)	Not used: same as 1978ME10
1992UN01	3,2615 (5) d	3,2615 (5)	
2002UN02	3,2615 (5) d	3,2615 (5)	Not used: same as 1992UN01
2004DA05	3,2634 (16) d	3,2634 (16)	
2004SC04	3,2623 (15) d	3,2623 (15)	
Adopted		3,2613 (5) d	

Figure 1: T_{1/2} measurements used for the present evaluation. The red ones are excluded by LWEIGHT according to Chauvenet's criterion.



2.2 Half-lives of ⁶⁷Zn 93 keV and 184 keV metastable states

The measured half-life values of the first excited state of ⁶⁷Zn, at 93 keV, are given in Table 2. The value from 1953KE was not used because of the lack of uncertainty. The value from 1972LE37 comes from the same author as 1973LE18. 1973LE18 reports two consistent measurements, obtained with two independent methods. A weighted mean was chosen: $T_{1/2}({}^{67}\text{Zn}, 93 \text{ keV}) = 9,15 (14) \mu\text{s}$.

No outlier was found by the LWEIGHT program in the consistent data set. A weighted average was 9,002 (41) μs with a reduced- χ^2 value of 2,2. The statistical weight is 85 % for the most precise value from 1996HW03. For the same reasons as $T_{1/2}({}^{67}\text{Ga})$, the smallest experimental uncertainty was preferred. Finally, the adopted value, with its external uncertainty, is:

$$T_{1/2}({}^{67}\text{Zn}, 93 \text{ keV}) = 9,00 (4) \mu\text{s}.$$

Table 2: $T_{1/2}$ half-life measurements of the 93 keV level of ⁶⁷Zn.
The excluded values are crossed out.

Reference	$T_{1/2}$ in μs	Comments
1953KE	8,5	Not used: no uncertainty
1953ME52	9,5 (10)	
1957BU39	9,4 (3)	
1969IV02	8,8 (18)	
1971SU18	8,7 (1)	
1972LE37	9,10 (15)	Not used: same as 1973LE18
1973LE18	9,20 (20) 9,10 (20)	
1973LE18	9,15 (14)	Consistent set, same author, independent methods: weighted mean chosen
1975RO25	9,1 (4)	
1996HW03	9,01 (3)	
1998AT04	9,34 (20)	

The measured half-life values of the second excited state of ⁶⁷Zn, at 184 keV, are given in Table 3. The value from 1964AL28 is given with asymmetric uncertainties, and was symmetrised according to the method described in 2003AU02.

The LWEIGHT program found two statistical outliers, excluded according to Chauvenet's criterion: 1961HO05 and 1962RI09. A weighted average was 1,028 (13) ns in the resulting consistent data set, with a reduced- χ^2 value of 0,3. The statistical weight is 83 % for the most precise value from 1972EN08. For the same reasons as $T_{1/2}({}^{67}\text{Ga})$, the smallest experimental uncertainty was preferred. Finally, the adopted value, with its internal uncertainty, is:

$$T_{1/2}({}^{67}\text{Zn}, 184 \text{ keV}) = 1,028 (14) \text{ ns}.$$

Table 3: $T_{1/2}$ half-life measurements of the 184 keV level of ⁶⁷Zn.
The excluded values are crossed out.

Reference	$T_{1/2}$ in ns	Comments
1961HO05	1,45 (15)	Excluded by Chauvenet's criterion
1962RI09	2,2 (7)	
1964AL28	1,1 (5)	Symmetrised according to method of 2003AU02
1968LI02	1,01 (5)	
1972EN08	1,026 (14)	
1975RO25	1,06 (4)	

2.3 Electron Capture Transitions

The energies of the electron capture (ϵ) transitions were calculated from the Q value and the level energies deduced from gamma transition energies. The log ft values were computed using the LOGFT program.

The electron capture probabilities were calculated from the balance of the evaluated $P_{\gamma+ce}$ values taking into account the evaluated absolute intensity $P(\epsilon_{1,0}) = 0,3254$ (26) (see Section 4.2.2) that allows normalizing the total ground state gamma transition probability to 96,7 (31) per 100 disintegrations.

The experimental values of P_K are available for $\epsilon_{0,2}$ and $\epsilon_{0,3}$ from 1988BE55: $P_K(\epsilon_{0,2}) = 0,89$ (4) and $P_K(\epsilon_{0,3}) = 0,88$ (3). They were obtained using an old value of $\omega_K = 0,430$ (7).

The P_K , P_L , P_M and P_N values were computed using the EC-capture program. The P_{L2}/P_{L1} ratios were computed by the LOGFT program and were used to determine the P_{L1} and P_{L2} values.

2.4 Gamma Transitions and Internal Conversion Coefficients

The evaluated energies of gamma transitions are the energies of gamma-rays with adding the recoil energy (see Section 4.2.1).

2.4.1 Mixing ratios and multiplicities

The multiplicities of the 93 keV ($\gamma_{1,0}$) and 794 keV ($\gamma_{4,1}$) transitions are pure E2. For the 794 keV transition, the admixture of M3 is possible and was evaluated. The other gamma transitions have an M1 + E2 multipolarity. Only the absolute values of the mixing ratios were evaluated to deal with the inconsistency of the signs. The mixing ratios measurements are given in Table 4.

Table 4: Mixing ratios measurements of the gamma transitions in ⁶⁷Zn. The excluded values, by the evaluator or by Chauvenet's criterion during the statistical process, are crossed out. The mixtures are M1 + E2 for all the transitions, except for the 794 keV transition which has an E2 + M3 mixture.

Reference	91 keV	184 keV	209 keV	300 keV	393 keV	494 keV	703 keV	794 keV	887 keV
1961HO05		$\delta^2 = 0,15$							
1962RI09	$ \delta \leq 0,07$	0,51 (7)							
1964AL28		0,43 (8)							
1968LI02		0,41 (6)							
1969BO41			0,40 (15)						
1971SU18	-0,11 (5)	0,38 (3)		pure M1	0,5 (1)	0,18 (6)			
1973BA54			0,034 (21)	-0,181 (8)	0,043 (10)		0,090 (28)		
1974NI01		$\delta \begin{cases} > -0,8 \\ < -0,1 \end{cases}$	0,02 (4)	-0,21 (5)	0,11 (6)	2,2 (6)		0,04 (17)	
1975TH01	0,06 (5)	0,48 (11)	0,01 (20)	0,05 (7)	-0,02 (17)	-0,22 (20)		0,47 (11)	0,8 (6)
1975WE08	-0,15 (3) 2,6 (3)	-0,17 (7)	0,08 (5) -5,7 (20)	-0,11 (4) 2,3 (3)	0,09 (2) 3,2 (3)	0,06 (4) 2,8 (4)			
1978DU04		0,08 (4) -5,0 (8)	-0,10 (6) 3,6 (8)	0,20 (8) -3,1 (4)	-0,17 (8) -2,4 (3)	-0,17 (8) -1,7 (6)		-0,1 (1)	0,9 (3)
1978LO06						0,14 (3)		0,04 (4)	-0,96 (9)

Some values are given with asymmetric uncertainties: $\delta(184 \text{ keV})$ from 1964AL28 ; $\delta(300 \text{ keV})$ from 1973BA54 ; $\delta(794 \text{ keV})$ and $\delta(494 \text{ keV})$ from 1974NI01; $\delta(393 \text{ keV})$, $\delta(494 \text{ keV})$, $\delta(794 \text{ keV})$ and $\delta(887 \text{ keV})$ from 1975TH01. They were symmetrised according to the method described in 2003AU02.

The excluded values are crossed out in the table. For the 91 keV transition, the result from 1962RI09 rules out the second possibility from 1975WE08. For the 184 keV transition, the result from 1974NI01 rules out the second possibility from 1978DU04, and there is no uncertainty with the result of 1961HO05. For the 209 keV, 300 keV, 393 keV and 494 keV transitions, the second possibilities from 1975WE08 and 1978DU04 are ruled out by comparison with the other results. For the 300 keV transition, the result from 1971SU18 is ruled out by comparison with the other results. For the 494 keV transition, the second possibility from 1974NI01 is ruled out by comparison with the most precise measurement from 1978LO06.

The other values that are crossed out were excluded according to Chauvenet's criterion during the statistical processing with the LWEIGHT program: 1969BO41 for 209 keV and 1971SU18 for 393 keV. All the data set are consistent except for the 184 keV and 794 keV transitions which are discrepant. The existence of only one measurement from 1973BA54 for the 703 keV transition should be underlined. A weighted average was used for each gamma transition. The adopted values are given in Table 5. Internal uncertainty was chosen by the LWEIGHT program for the red adopted values, external uncertainty for the green ones and expanded uncertainty for the blue one.

Table 5: Adopted mixing ratios and the corresponding multipolarity.
Internal uncertainty was chosen by LWEIGHT for the red adopted values,
external uncertainty for the green ones and expanded uncertainty for the blue one.

	Adopted $ \delta $	Multipolarity	Comments
91 keV	0,123 (25)	M1 + 1,5 (6) % E2	Consistent, weighted average
184 keV	0,31 (7)	M1 + 8,8 (36) % E2	Discrepant, weighted average
209 keV	0,042 (17)	M1 + 0,18 (14) % E2	Consistent, weighted average
300 keV	0,178 (10)	M1 + 3,07 (33) % E2	Consistent, weighted average
393 keV	0,051 (16)	M1 + 0,26 (16) % E2	Consistent, weighted average
494 keV	0,110 (34)	M1 + 1,2 (7) % E2	Consistent, weighted average
703 keV	0,090 (28)	M1 + 0,8 (5) % E2	Only one measurement
794 keV	0,09 (11)	E2 + 0,8 (19) % M3	Discrepant, weighted average
887 keV	0,95 (9)	M1 + 47,4 (47) % E2	Consistent, weighted average

2.4.2 Internal conversion coefficients

The internal conversion coefficients measurements of the gamma transitions of ⁶⁷Zn are summarized in Table 6.

The adopted values are calculated with the BrIcc program (2008KI07) and can be seen in Table 7. They were calculated using the mixing ratios evaluated previously. These values agree satisfactorily with the measured ones. For the 794 keV transition, the two possible multipolarities were tested and the calculated internal conversion coefficients agree well inside the uncertainties. The possible admixture of M3 is kept.

Table 6: Internal conversion coefficients measurements of the gamma transitions of ⁶⁷Zn.

Reference	91 keV	93 keV	184 keV	209 keV	300 keV	393 keV	494 keV	887 keV	
1938AL02	$\alpha_K \sim 0,07$								
1953KE	$\alpha_K = 0,074$	$\alpha_K = 0,63$	$\alpha_K = 0,011$	$\alpha_K = 0,029$	$\alpha_K = 0,0029$	$\alpha_K = 0,0019$			
1953ME52	$\alpha_T = 0,54 (5)$								
1966FR12	$\alpha_K = 0,77 (8)$		$\alpha_K = 0,0156 (10)$	$\alpha_K = 0,0075 (7)$	$\alpha_K = 0,00337 (30)$	$\alpha_K = 0,00192 (15)$	$\alpha_K = 0,00119 (15)$	$\alpha_K = 0,00034 (7)$	
1969LI04	$\alpha_K = 0,066 (10)$ $\alpha_{L1} = 0,0069 (11)$ $\alpha_{L2,3} = 0,00069 (30)$								
1988BE55			$\alpha_K = 0,89 (4)$				$\alpha_K = 0,883 (28)$		

Table 7: BrIcc calculations for the internal conversion coefficients of the gamma transitions of ⁶⁷Zn.

	α_T	α_K	α_{Ltot}	Comments
91 keV	0,091 (6)	0,081 (5)	0,008 7 (7)	
93 keV	0,854 (12)	0,748 (11)	0,092 2 (13)	
184 keV	0,016 9 (21)	0,015 1 (19)	0,001 58 (20)	
209 keV	0,009 01 (14)	0,008 06 (13)	0,000 827 (13)	
300 keV	0,003 88 (6)	0,003 48 (6)	0,000 354 (6)	
393 keV	0,001 93 (3)	0,001 728 (25)	0,000 174 8 (25)	
494 keV	0,001 149 (18)	0,001 030 (16)	0,000 103 8 (17)	
703 keV	0,000 524 (8)	0,000 470 (7)	0,000 047 0 (7)	
794 keV E2+M3	0,000 54 (6)	0,000 48 (5)	0,000 049 (6)	Adopted
794 keV pure E2	0,000 523 (8)	0,000 469 (7)	0,000 047 3 (7)	
887 keV	0,000 354 (7)	0,000 318 (6)	0,000 031 8 (6)	

3. Atomic Data

3.1 Fluorescence yields

The fluorescence yields are taken from 1996SC06.

3.2 X Radiations and Auger electrons

The X-ray energies are based on the wave lengths in the compilation of 1967BE65 (Bearden). The energies of Auger electrons are from the SAISINUC software (see also 1977LA19 (Larkins) and 1987Table (Table de Radionucléides)).

The X-ray and Auger electron probabilities were computed using the EMISSION program with the atomic data from 1996SC06.

4. Photon Emissions

4.1 X-Ray Emissions

The total absolute emission probabilities of the K and L X-rays were computed using the EMISSION program with the atomic data from 1996SC06, the evaluated electron capture probabilities and the evaluated conversion electron probabilities.

The authors of 1979DE42 measured the following ratios: $P(XK_{\alpha})/P(184\text{keV}) = 2,37 (5)$ and $P(XK_{\beta})/P(184\text{keV}) = 0,331 (7)$. The absolute intensity measurements from 2005YA01 are: $P(XK_{\alpha 2}) = 17,1 (8) \%$, $P(XK_{\alpha 1}) = 32,3 (14) \%$, $P(XK_{\beta 1,3}) = 6,44 (28) \%$ and $P(184 \text{ keV}) = 21,4 (9) \%$. The relative uncertainties were increased to 4,4 % (see Section 4.2.2). This leads to the consistent ratio $P(XK_{\alpha})/P(184 \text{ keV}) = 2,31 (12)$ and the discrepant ratio $P(XK_{\beta})/P(184 \text{ keV}) = 0,301 (18)$.

With the previous calculations and the evaluated value of $P(184 \text{ keV})$ (see Section 4.2.2), these ratios become: $P(XK_{\alpha})/P(184 \text{ keV}) = 2,39 (8)$ and $P(XK_{\beta})/P(184 \text{ keV}) = 0,338 (14)$. They are summarized in Table 8.

Table 8: Comparison of the $P(XK_{\alpha})/P(184 \text{ keV})$ and $P(XK_{\beta})/P(184 \text{ keV})$ ratios.

	1979DE42	2005YA01	Evaluated
$P(XK_{\alpha})/P(184 \text{ keV})$	2,37 (5)	2,31 (12)	2,39 (8)
$P(XK_{\beta})/P(184 \text{ keV})$	0,331 (7)	0,301 (18)	0,338 (14)

4.2 Gamma Emissions

4.2.1 Gamma-ray energies

The gamma-ray energy measurements are given in Table 9. The excluded values are crossed out. The values from 1978ME10 come from the same author as 1990ME15. The most recent data set was preferred, even if it corresponds to the less precise measurements of the two publications. The other values that are crossed out were excluded according to Chauvenet's criterion during the statistical process by the LWEIGHT program. All the data set are consistent and a weighted average was used each time. The adopted values are in red or in green when respectively the internal or the external uncertainties were chosen. The reduced- χ^2 are also mentioned.

Table 9: E_{γ} measurements of γ -rays in ⁶⁷Zn.

The excluded values are crossed out. The values from 1978ME10 come from the same author as 1990ME15. The other values were excluded according to Chauvenet's criterion. Internal uncertainty was chosen by LWEIGHT for the red adopted values, external uncertainty for the green ones.

Reference	$E_{\gamma 2,1}$ (keV)	$E_{\gamma 1,0}$ (keV)	$E_{\gamma 2,0}$ (keV)	$E_{\gamma 3,2}$ (keV)	$E_{\gamma 3,1}$ (keV)
1958CH08	91,22 (4)	93,26 (4)	184,46 (27)		
1966FR12	91,275 (20)	93,317 (20)	184,595 (40)	208,96 (6)	300,24 (7)
1969RA15	91,26 (10)	93,25 (10)	184,53 (10)	208,95 (10)	300,22 (10)
1971SU18	91 (2)	93 (2)	184 (2)	208 (2)	299 (2)
1974AR22		93,2 (2)	184,0 (2)		
1974HEYW	91,31 (5)	93,32 (2)	184,56 (2)	208,93 (2)	300,18 (2)
1977AB02			184 (1)		
1978DU04		93,3 (5)	184,63 (3)	208,91 (4)	300,24 (5)
1978ME10	91,266 (5)	93,311 (5)	184,577 (10)	208,951 (10)	300,219 (10)
1990ME15	91,237 (35)	93,291 (30)	184,569 (30)	208,970 (30)	300,230 (25)
Adopted	91,263 (15)	93,307 (12)	184,577 (17)	208,939 (15)	300,232 (21)
χ^2	0,74	0,54	1,7	0,48	0,02

Reference	E _{γ3,0} (keV)	E _{γ4,3} (keV)	E _{γ4,2} (keV)	E _{γ4,1} (keV)	E _{γ4,0} (keV)
1966FR12	393,65 (6)	494,31 (10)	703,6 (2)	794,7 (2)	888,0 (2)
1969RA15	393,60 (10)				
1971SU18	393 (2)	493 (2)	701 (2)	794 (2)	886 (2)
1974HEYW	393,47 (3)	494,19 (8)		794,49 (20)	887,68 (15)
1977AB02	393 (3)				884 (12)
1978DU04	393,54 (5)	494,1 (6)	703,2 (3)	794,39 (8)	887,67 (7)
1978ME10	393,529 (10)	494,169 (15)	703,110 (15)	794,386 (15)	887,693 (15)
1990ME15	393,539 (25)	494,132 (30)	703,08 (5)	794,38 (5)	887,664 (40)
Adopted	393,528 (20)	494,143 (28)	703,11 (8)	794,400 (41)	887,676 (33)
χ²	1,5	1,3	2,5	0,67	0,91

The energies of the gamma transitions were then deduced by adding the recoil energy of the nucleus. The proton and neutron masses come from 2008MO18. The mass excess of ⁶⁷Zn comes from 2003AU02. The largest recoil energy is for the 887 keV transition, with a value less than 6,3 eV.

4.2.2 Gamma-ray emission probabilities

The measurements of gamma-ray emission probabilities are summarized in Table 10. The excluded values are crossed out. 1953KE was not used because of the lack of uncertainty. The intensity of the doublet from 1967VR03 is not useful. In 1975TH01, the intensities are normalized by level. With the 184 keV emission as reference, only the measurement of the 91 keV intensity can be used. In 1978LO06, the reference is the complete decay of the 888 keV level. As there is no value for the 184 keV emission, these values were not used. The values from 1978ME10 come from the same author as 1990ME15. The most recent data set was preferred.

Table 10: Measurements of gamma-ray emission probabilities from the decay of ⁶⁷Ga. The values that are crossed out were excluded by the evaluator. The energies are in keV. Values from 1969LI04, 2005YA01 and 2007BO are absolute emission probabilities measurements. The relative uncertainties of the values from 2005YA01 were increased up to 4,4 %.

Reference	91	93	184	209	300	393	494	703	794	887
1953KE	2,7	63,9	29,6	±	20,2	4,9	0,4		0,2	0,4
1953ME52	7,0 (5)	93,0 (5)	44,1 (30)	3,0 (8)	27,5 (10)	9,7 (10)				
1966FR12	1,5 (4)	73 (7)	23,1 (16)	2,50 (25)	16,2 (16)		0,100 (15)	0,015 (2)	0,06 (1)	0,160 (16)
1967VR03	γ_{2,1}+γ_{1,0} 229 (20)		100	10,9 (5)	75,6 (50)	20,4 (12)	0,24 (3)	0,05 (1)	0,23 (2)	0,58 (6)
1969LI04	3,27 (45)	38,4 (38)	23,7 (27)							
1969RA15	155 (11)	360 (25)	1000 (70)	2,4 (3)	15,7 (12)	4,3 (4)				
1974HEYW	13 (1)	100 (5)	62 (3)	7,1 (4)	50 (3)	14 (1)	3,7 (3)		0,15 (2)	0,43 (4)
1975TH01	13,1 (4)		86,9 (25)	9,6 (3)	70,3 (21)	20,1 (6)	23,8 (11)	4,8 (4)	18,4 (7)	53,2 (16)
1978LO06							25 (2)	5 (2)	21 (2)	49 (2)
1978ME10	80,0 (3)	1000 (3)	552 (2)	62,8 (3)	448 (1)	125,4 (5)	1,83 (2)	0,292 (9)	1,37 (5)	3,88 (2)
1979DE42	15,0 (5)	185 (6)	100,0 (11)	11,35 (13)	79,9 (9)	22,0 (3)	0,322 (7)	0,060 (5)	0,251 (7)	0,712 (11)
1990ME15	30 (2)	366 (14)	217 (9)	24 (1)	166 (4)	45 (1)	0,7 (1)	0,10 (1)	0,53 (3)	1,49 (5)
2005YA01	3,11 (14)	38,8 (17)	21,4 (9)	2,46 (11)	16,6 (7)	4,6 (2)				
2007BO	3,11 (3)	38,61 (35)	21,13 (10)	2,396 (13)	16,74 (8)	4,642 (25)	0,0657 (33)		0,0565 (24)	0,1522 (35)

The value from 1966FR12 for the 91 keV emission is given with asymmetric uncertainties, and was symmetrised according to the method described in 2003AU02. 2005YA01 and 2007BO measured absolute emission probabilities. The uncertainties from 2005YA01 are too low. They should be at least about 4,4 % according to the authors, and they were expanded.

In this evaluation, the data were normalized to the 184 keV emission, used as reference. They are given in Table 11. The crossed out values were excluded according to Chauvenet’s criterion during the statistical process by LWEIGHT. All the data sets are consistent and a weighted average was used each time. Red adopted values stand for internal uncertainty, green ones stand for external uncertainty. The reduced- χ^2 are also mentioned.

Table 11: Measurements of gamma-ray emission probabilities used in this evaluation, normalized to the 184 keV emission used as reference.

The values that are crossed out were excluded according to Chauvenet’s criterion during the statistical process by LWEIGHT.

Red values stand for internal uncertainty, green ones stand for external uncertainty.

Reference	91 keV	93 keV	184 keV	209 keV	300 keV	393 keV	494 keV	703 keV	794 keV	887 keV
1953ME52	15,9 (16)	211 (14)	100	6,8 (19)	62,4 (48)	22,0 (27)				
1966FR12	6,5 (18)	316 (37)	100	10,8 (13)	70 (8)		0,43 (7)	0,065 (10)	0,260 (47)	0,69 (8)
1967VR03			100	10,9 (5)	76 (5)	20,4 (12)	0,24 (3)	0,05 (1)	0,23 (2)	0,58 (6)
1969LI04	13,8 (25)	162 (24)	100							
1974HEYW	21,0 (19)	161 (11)	100	11,5 (9)	81 (6)	22,6 (19)	6,0 (6)		0,242 (34)	0,69 (7)
1975TH01	15,1 (6)		100							
1979DE42	15,0 (5)	185 (6)	100	11,35 (18)	79,9 (13)	22,00 (39)	0,322 (8)	0,060 (5)	0,251 (8)	0,712 (14)
1990ME15	13,8 (11)	169 (10)	100	11,1 (7)	76,5 (37)	20,7 (10)	0,32 (5)	0,046 (5)	0,244 (17)	0,687 (37)
2005YA01	14,53 (6)	181 (8)	100	11,5 (5)	77,7 (34)	21,3 (11)				
2007BO	14,72 (16)	182,7 (19)	100	11,34 (8)	79,2 (5)	21,97 (16)	0,31 (16)		0,267 (11)	0,720 (17)
Adopted	14,74 (14)	181,8 (19)	100	11,33 (7)	79,22 (48)	21,92 (14)	0,318 (12)	0,0539 (42)	0,252 (6)	0,712 (10)
χ^2	0,34	1,3	-	0,20	0,32	0,61	2,4	1,8	0,67	0,21

These evaluated relative emission probabilities were used with the absolute emission probability of the 93 keV level P(93 keV) to calculate the absolute gamma-ray emission probabilities. P(93 keV) was determined using the total internal conversion coefficient $\alpha_T(93 \text{ keV})$ calculated with the BrIcc program in Section 2.4.2, and the evaluated value of P($ce_{1,0}$).

Four values of P($ce_{1,0}$) were used in this evaluation: 0,3285 (40) from 1998AT04; 0,3198 (40) from 2000SI03; 0,331 (15) from 2005YA01; 0,330 (6) from 2007BO. The first value was recalculated by 2000SI03. The second value is an unweighted mean of the two results 0,3213 (14) and 0,3182 (27). The authors of 2000SI03 precise that the uncertainties are only 1σ statistical uncertainties. The final uncertainty of P($ce_{1,0}$) was increased to be at least the second lowest uncertainty, the one from 1998AT04. The values from 2005YA01 and 2007BO were calculated with the measured absolute intensities and with $\alpha_T(93 \text{ keV})$. The relative uncertainty of 2005YA01 was increased to 4,4 %, as explained above.

No outlier was found by the LWEIGHT program in the consistent data set. A weighted average was 0,3254 (26) with a reduced- χ^2 value of 1,1. The statistical weight is 40 % for the two most precise values from 1998AT04 and 2000SI03. As there are only four useful values, the smallest experimental uncertainty was preferred. Finally, the adopted value, with its external uncertainty, is:

$$P(\epsilon_{1,0}) = 0,3254 (40).$$

With $\alpha_T(93 \text{ keV}) = 0,854 (12)$ (see Table 7 in Section 2.4.2), the absolute intensity of the 93 keV gamma-ray transition was found to be:

$$P(93\text{keV}) = 0,381 (7).$$

With the absolute gamma-ray emission probabilities and the adopted total internal conversion coefficient of each transition, the transition probabilities $P_{\gamma+ce}$ were determined. They are given in Table 12. The resulting probabilities of the electron capture transitions were found to be: $P(\epsilon_{0,4}) = 0,280 (8) \%$; $P(\epsilon_{0,3}) = 23,60 (47) \%$; $P(\epsilon_{0,2}) = 22,3 (27) \%$; $P(\epsilon_{0,1}) = 50,5 (17) \%$. It leads to the probability of the electron capture to the ground state of ⁶⁷Zn: $P(\epsilon_{0,0}) = 3,3 (32) \%$. The large uncertainty of $P(\epsilon_{0,0})$ mainly comes from the uncertainty of the evaluated mixing ratio $\delta(184 \text{ keV})$.

Table 12: The absolute gamma-ray emission probabilities, calculated with the evaluated value of $P(\epsilon_{1,0})$ and the adopted $\alpha_T(93 \text{ keV})$. The transition probabilities were calculated from the P_γ and the α_T of each transition.

Transition	Relative intensity	P_γ (%)	$P_{\gamma+ce}$ (%)
$\gamma_{2,1}$ (91 keV)	14,74 (14)	3,09 (7)	3,37 (24)
$\gamma_{1,0}$ (93 keV)	181,8 (19)	38,1 (7)	70,6 (16)
$\gamma_{2,0}$ (184 keV)	100	20,96 (44)	21,3 (27)
$\gamma_{3,2}$ (209 keV)	11,33 (7)	2,37 (5)	2,40 (6)
$\gamma_{3,1}$ (300 keV)	79,22 (48)	16,60 (37)	16,67 (45)
$\gamma_{3,0}$ (393 keV)	21,92 (14)	4,59 (10)	4,60 (12)
$\gamma_{4,3}$ (494 keV)	0,318 (12)	0,0666 (29)	0,0667 (31)
$\gamma_{4,2}$ (703 keV)	0,0539 (42)	0,0113 (9)	0,0113 (9)
$\gamma_{4,1}$ (794 keV)	0,252 (6)	0,0528 (17)	0,053 (6)
$\gamma_{4,0}$ (887 keV)	0,712 (10)	0,1492 (38)	0,1493 (48)

5. Electron Emissions

The energies of the conversion electrons were calculated from the gamma-transition energies given in Section 4.2.1 and the electron binding energies.

The emission probabilities of the conversion electrons were calculated using the internal conversion coefficients given in Section 2.4.2. The emission probabilities of K-Auger electrons were calculated using the transition probabilities given in Sections 2.3 and 2.4, the atomic data given in Section 3, and the internal conversion coefficients given in Section 2.4.2.

6. References

- 1938AL02 L.W. Alvarez, Phys. Rev. 54, 486 (1938) [ICC]
 1938MA01 W.B. Mann, Phys. Rev. 54, 649 (1938) [$T_{1/2}$]
 1948HO04 H.H. Hopkins Jr., B.B. Cunningham, Phys. Rev. 73, 1406 (1948) [$T_{1/2}$]
 1948MC32 D.A. McCown, L.L. Woodward, M.L. Pool, Phys. Rev. 74, 1311 (1948) [$T_{1/2}$]
 1950HO26 H.H. Hopkins, Phys. Rev. 77, 717 (1950) [$T_{1/2}$]
 1953KE B.H. Ketelle, A.R. Brosi, F.M. Porter, Phys. Rev. 90, 567 (1953) [$T_{1/2}$, P_γ , ICC]
 1953ME52 W.E. Meyerhof, L.G. Mann, H.I. West Jr., Phys. Rev. 92, 758 (1953) [$T_{1/2}$, P_γ , ICC]
 1955TO27 J. Tobailem, Ann. Phys. 10, 783 (1955) [$T_{1/2}$]
 1957BU39 A.J. Bureau, C.L. Hammer, Phys. Rev. 105, 1006 (1957) [$T_{1/2}$]
 1958CH08 E.L. Chupp, J.W.M. Dumond, F.J. Gordon, R.C. Jopson, H. Mark, Phys. Rev. 109, 2036 (1958) [E_γ]

- 1961HO05** R.E. Holland, F.J. Lynch, Phys. Rev. 121, 1464 (1961) [$T_{1/2}$, mixing ratios]
- 1962RI09** R.C. Ritter, P.H. Stelson, F.K. McGowan, R.L. Robinson, Phys. Rev. 128, 2320 (1962) [$T_{1/2}$, mixing ratios]
- 1964AL28** D.G. Alkhozov, V.D. Vasilev, G.M. Gusinskii, I.K. Lemberg, V.A. Nabichvrishvili, Bull. Acad. Sci. USSR, Phys. Ser. 28, 1575 (1965) [$T_{1/2}$, mixing ratios]
- 1964RU06** G. Rudstam, Nucl. Phys. 56, 593 (1964) [$T_{1/2}$]
- 1966FR12** M.S. Freedman, F.T. Porter, F. Wagner, Phys. Rev. 151, 886 (1966) [E_γ , P_γ , ICC]
- 1967BE65** J.A. Bearden, Rev. Mod. Phys. 39, 78 (1967) [E_x]
- 1967VR03** J. Vrzal, B.S. Dzheleпов, J. Liptak, L.N. Moskvina, V.P. Prikhodtseva, B.M. Sabirov, V.G. Tishin, J. Urbanets, L.G. Tsaritsyna, Bull. Acad. Sci. USSR, Phys. Ser. 31, 1687 (1968) [P_γ]
- 1968LI02** R.M. Lieder, M. Fleck, K. Killig, M. Forker, K.-H. Speidel, E. Bodenstedt, Nucl. Phys. A 106, 389 (1968) [$T_{1/2}$, mixing ratios]
- 1969BO41** E. Bozek, R. Broda, J. Golczewski, A.Z. Hryniewicz, R. Kulesa, M. Rybicka, S. Szymczyk, W. Walus, Acta Phys. Polon. 36, 1065 (1969) [Mixing ratios]
- 1969IV02** E.A. Ivanov, A. Iordachescu, G. Pascovici, Rev. Roum. Phys. 14, 317 (1969) [$T_{1/2}$]
- 1969LI04** A. Li-Scholz, H. Bakhru, Phys. Rev. 177, 1629 (1969) [P_γ , ICC]
- 1969RA15** S. Raman, J.J. Pinajian, Nucl. Phys. A 131, 393 (1969) [E_γ , P_γ]
- 1971SU18** V. Sutela, L. Tukia, Ann. Acad. Sci. Fenn., Ser. A VI, No.364 (1971) [$T_{1/2}$, E_γ , mixing ratios]
- 1972CR02** D.F. Crisler, H.B. Eldridge, R. Kunselman, C.S. Zaidins, Phys. Rev. C 5, 419 (1972) [$T_{1/2}$]
- 1972EN08** H. Engel, P. John, B. Reuse, Z. Naturforsch. 27a, 1368 (1972) [$T_{1/2}$]
- 1972LE37** V.E. Lewis, M.J. Woods, I.W. Goodier, Int. J. Appl. Radiat. Isotop. 23, 279 (1972) [$T_{1/2}$]
- 1973BA54** C. Bargholtz, L. Eriksson, L. Gidefeldt, Z. Phys. 263, 89 (1973) [Mixing ratios]
- 1973LE18** V.E. Lewis, D. Smith, A. Williams, Metrologia 9, 14 (1973) [$T_{1/2}$]
- 1974AR22** P.O. Aronsson, B.E. Johansson, J. Rydberg, G. Skarnemark, J. Alstad, B. Bergersen, E. Kvale, M. Skarestad, J. Inorg. Nucl. Chem. 36, 2397 (1974) [E_γ]
- 1974HEYW** R.L. Heath, ANCR-1000-2 (1974) [E_γ , P_γ]
- 1974NI01** A. Nilsson, Z.P. Sawa, Phys. Scr. 9, 83 (1974) [Mixing ratios]
- 1975RO25** S. Roodbergen, H. Visser, W. Molendijk, H.S. Bedet, H. Verheul, Z. Phys. A 275, 45 (1975) [$T_{1/2}$]
- 1975TH01** M.J. Throop, Y.T. Cheng, D.K. McDaniels, Nucl. Phys. A 239, 333 (1975) [P_γ , mixing ratios]
- 1975WE08** S.A. Wender, J.A. Cameron, Nucl. Phys. A 241, 332 (1975) [Mixing ratios]
- 1977AB02** A.L. Abu-Ghazaleh, I. M. Naqib, G. Brown, J. Phys. (London) G 3, 253 (1977) [E_γ]
- 1977LA19** F.P. Larkins, At. Data Nucl. Data Tables 20, 311 (1977) [E_{Ae}]
- 1978DU04** R. Duffait, A. Charvet, R. Chery, Phys. Rev. C 17, 2031 (1978) [E_γ , mixing ratios]
- 1978LA21** F. Lagoutine, J. Legrand, C. Bac, Int. J. Appl. Radiat. Isotop. 29, 269 (1978) [$T_{1/2}$]
- 1978LO06** P.R.G. Lornie, A. Kogan, G.D. Jones, M.R. Nixon, H.G. Price, R. Wadsworth, P.J. Twin, J. Phys. (London) G 4, 923 (1978) [P_γ , mixing ratios]
- 1978ME10** R.A. Meyer, A.L. Prindle, W.A. Myers, P.K. Hopke, D. Dieterly, J.E. Koops, Phys. Rev. C 17, 1822 (1978) [$T_{1/2}$, E_γ , P_γ]
- 1979DE42** K. Debertain, W. Pessara, U. Schotzig, K.F. Walz, Int. J. Appl. Radiat. Isotop. 30, 551 (1979) [$T_{1/2}$, P_x , P_γ]
- 1980HO17** H. Houtermans, O. Milosevic, F. Reichel, Int. J. Appl. Radiat. Isotop. 31, 153 (1980) [$T_{1/2}$]
- 1982HOZJ** D.D. Hoppes, J.M.R. Hutchinson, F.J. Schima, M.P. Unterweger, NBS-SP-626, p.85 (1982) [$T_{1/2}$]
- 1987Table** F. Lagoutine, N. Coursol, J. Legrand, Table de Radionucléides, CEA-LMRI, ISBN-2-7272-0078-1 (1982-1987) [E_{Ae}]

- 1988BE55** R.B. Begzhanov, K.Sh. Azimov, N.A. Ilkhamdzhanov, R.A. Magrupov, Sh.A. Mirakhmedov, A. Mukhammadiev, M. Narzikulov, Bull. Acad. Sci. USSR, Phys. Ser. 52, No.11, 88 (1988) [Experimental P_K values]
- 1990ME15** R.A. Meyer, Fizika (Zagreb) 22, 153 (1990) [E_γ , P_γ]
- 1991BH06** M.R. Bhat, Nucl. Data Sheets 64, 875 (1991) [Decay scheme]
- 1992UN01** M.P. Unterweger, D.D. Hoppes, F.J. Schima, Nucl. Instrum. Methods Phys. Res. A 312, 349 (1992) [$T_{1/2}$]
- 1996HW03** H.Y. Hwang, C.B. Lee, T.S. Park, H.J. Kim, Nucl. Instrum. Methods Phys. Res. A 383, 447 (1996) [$T_{1/2}$]
- 1996SC06** E. Schönfeld, H. Janssen, Nucl. Instrum. Methods Phys. Res. A 369, 527 (1996) [Atomic Data]
- 1998AT04** M.R.P. Attie, M.F. Koskinas, M.S. Dias, K.A. Fonseca, Appl. Radiat. Isot. 49, 1175 (1998) [$T_{1/2}$, P_{ce}]
- 2000SI03** B.R.S. Simpson, T.P. Ntsoane, Appl. Radiat. Isot. 52, 551 (2000) [P_{ce}]
- 2002UN02** M.P. Unterweger, Appl. Radiat. Isot. 56, 125 (2002) [$T_{1/2}$]
- 2003AU02** G. Audi, O. Bersillon, J. Blachot, A.H. Wapstra, Nucl. Phys. A 729, 3 (2003) [Symmetrization of asymmetric uncertainties]
- 2003AU03** G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A 729, 337 (2003) [Q]
- 2004DA05** M.A.L. da Silva, M.C.M. de Almeida, C.J. da Silva, J.U. Delgado, Appl. Radiat. Isot. 60, 301 (2004) [$T_{1/2}$]
- 2004SC04** H. Schrader, Appl. Radiat. Isot. 60, 317 (2004) [$T_{1/2}$]
- 2005HU18** J. Huo, X. Huang, J.K. Tuli, Nucl. Data Sheets 106, 159 (2005) [Decay scheme]
- 2005YA01** P. Yalcin, Y. Kurucu, Appl. Radiat. Isot. 62, 63 (2005) [P_γ , P_{ce}]
- 2007BO** C. Bobin, J. Bouchard, C. Hamon, M.G. Iroulart, J. Plagnard, Appl. Radiat. Isot. 65, 757 (2007) [P_γ , P_{ce}]
- 2008KI07** T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr. Nucl. Instr. and Meth. A589 (2008) 202-229 [BrIcc]
- 2008MO18** P.J. Mohr, B.N. Taylor, D.B. Newell, Rev. Mod. Phys. 80, 633 (2008) [m_p , m_n]

⁶⁸Ga - Comments on evaluation
by M.-M. Bé and E. Schönfeld

This evaluation was completed in 1996, it was reviewed in November 2011. The literature available by this date was included.

1 Decay Scheme

The decay scheme of Ga-68 is taken from Vo *et al.* (1994Vo15) who discovered 5 very weak transitions between already known levels in the decay scheme established by Lange *et al.* (1973La01). From other excitation modes, the existence of levels with energy (J^π): 2370,3 (15); 2417,44 (6) (4)⁺; 2510,2 (15); 2750,38 (8) keV (3⁻) was shown, however radiations originating from these levels were not observed in the Electron Capture decay of Ga-68 (1994Vo15). Transitions to the 2417 and 2510 keV levels would be third and second forbidden and therefore if they exist, their intensities would be very weak. The values of the half-lives of the excited states of ⁶⁸Zn are taken from Burrows (2002Bu29).

2 Nuclear Data

The Q^+ value is from Audi *et al.* (2003Au03), adopted from Slot (1972Sl03) measurement, $Q^+ = 2921,1$ (12) keV.

A value of 2912 (10) keV was also measured by Kojima *et al.* (2001Ko07).

The following values of the half-life of ⁶⁸Ga have been taken into consideration:

Reference	$T_{1/2}$ (min)	Remarks
M. L. Perlman	68,0	omitted
G. L. Gleason (1960Gl04)	67,7 (3)	
L. A. Rayburn (1961Ra06)	69,2 (14)	outlier
T. G. Ebrey (1965Eb01)	68,33 (9)	Coin. Count. NaI(Tl), statistical uncertainty only
M. Borman (1965Bo42)	68,2 (1)	
J. M. Ootukalam (1971Oo01)	68,5 (5)	NaI, brief note, statistical uncertainty only
Smith and Williams (1971Sm02)	67,80 (8)	IC
Iwata <i>et al.</i> (1983Iw02)	67,629 (24)	Ge(Li)
Luca <i>et al.</i> (2012Lu*)	67,87 (10)	IC
Adopted	67,83 (20)	$\chi^2_{\text{crit}} = 2,8$; $\chi^2 = 11$

The set of 7 values used in the averaging process is not consistent with a reduced χ^2 of 11. The limitation of relative statistical weight procedure has then increased the Iwata's uncertainty to 0,05 in order to reduce the relative weight of this value to 50 %.

Therefore, the resulting (and adopted) weighted mean is 67,83 min with an expanded uncertainty of 0,20 to cover the most precise value.

2.1, 2.2 Electron Capture Transitions and β^+ Transitions

The sum of the EC + β^+ transition probabilities were deduced from the sum of the gamma transition probabilities populating and depopulating each level of the decay scheme. The EC/ β^+ ratios were calculated by using the Logft program, a relative uncertainty of 5 % was assumed. For level 0, the theoretical value is quite close to the experimental values of 0,10 (2) (1959Ra04) ; for level 1, it lies between 1,28 (12) and 1,63 (11), experimentally determined by Ramaswamy (1959Ra04) and Sykora (1992Sy**) respectively.

The individual EC and β^+ probabilities have then been derived.

	P(EC+ β^+) %	EC/ β^+	P(EC) %	P(β^+) %
Level 0	96,62 (3) %	0,102 (5)	8,94 (41)	87,68 (41)
Level 1	3,00 (3) %	1,51 (7)	1,80 (5)	1,20 (4)
Level 2	0,0338 (23) %	127,5 (64)	0,0335 (23)	0,00026 (2)

From the values above, the sum of P(β^+) amounts for 88,88 (41) %, it corresponds to a 511 keV photon intensity of 177,8 (8) %, this result can be compared with experimental values.

Several authors measured the 511 keV photon emission, I_{511} , relatively to the 1077 emission:

Reference	I_{511}/I_{1077}	u_c	
Horen (1959Ho85)	5460	600	
Craseman (1956Cr29)	2880	340	Omitted
Carter (1968Ca15)	5930	600	
Ramaswamy (1959Ra04)	3900	420	Outlier
Schönfeld (1994Sc44)	5537	52	Uc increased to 220 to limit its weight to 50 %
Luca (2012Lu*)	5569	264	
Weighted mean ; internal uncertainty	5570	160	With $I_{g1077} = 3,235 (30) \%$, the total I_{511} is 180 (5) %.

Moreover, two authors measured directly the I_{511} in absolute values: Schönfeld (1994Sc44), 178,29 (22) % and Luca (2012Lu*), 181 (6) %.

All these results are consistent within the uncertainty limits.

The energies are derived from the Q value and the level energies. The fractional probabilities for EC are calculated using the "Tables for Calculation of Electron Capture" (E. Schönfeld, PTB-Laboratory report 6.33-95-2 (1995)). These values are based on wave functions of Mann and Waber (1973) with exchange and overlap corrections of Bahcall and Vatai; see W. Bambynek et al., *Rev. Mod. Phys.* 49(1977)77. Note that the sum $P_K + P_L + P_M$ is not equal to 1 because of a very small fraction of capture from the N shell.

2.3 Gamma Transitions

The level differences (as well as the gamma-ray energies in Section 4.2) are taken from a compilation of Helmer (1997HeZZ) which takes into account also data from sources other than the Ga-68 decay.

The transition probabilities were derived from the emission intensities and the conversion coefficients. The conversion coefficients are interpolated from Band's tables by using the program BrIcc with the "frozen orbital approximation" (Kibédi *et al.* 2008Ki07). The adopted mixing ratios are derived from the experimental results as summarized below. For the 1261 keV transition where the values are discrepant the adopted value comes from the γ - γ directional correlation measurements of Vo *et al.* (1994Vo15). Eight transitions are pure E2 because the initial or the final level has $J = 0$.

Experimental mixing ratios δ :

E γ (keV)	483	805	938	1261	1744
Reference					
1960Ra06				-1,8 (2)	
1962Ko01		+4 (+3 -2)			
1963Ta03				-2,25 (30)	
1971Ot01		-1,45 (15)		-0,21 (+6 -4)	0,24 (13)
1973La01		-1,46 (14)		0,14 (4)	0,29 (5)
1994Vo15	-0,12 (16) -1,7 (9)	-1,55 (5)	-0,7 (3)	-0,15 (2)	0,27 (2)
Adopted	1,0 (5) ^e	-1,53 (5) ^w	-0,7 (3)	-0,15 (2) ^v	0,272 (18) ^w
M1 / E2 (%)	50 (25)	70 (1)	33 (8)	2,2 (1)	6,9 (1)

^e : estimated ; ^w : weighted mean ; ^v : 1994Vo15.

3 Atomic Data

All these data are taken from E. Schönfeld and H. Janssen, *Nucl. Instr. and Methods in Phys. Res. A* 369(1996)527.

3.1 X Radiations

The energies are based on the wavelengths compiled by J. A. Bearden, *Rev. Mod. Phys.* 39(1967)78. The relative probabilities for K α radiation are based on $P(K\beta)/P(K\alpha)$ and $P(K\alpha_2)/P(K\alpha_1)$ values as given in the above cited paper of Schönfeld and Janßen (1994). The relative probabilities for L quanta are derived from the corresponding absolute values (Section 4.2) setting $P(K\alpha_1) = 100$.

3.2 Auger Electrons

The energies of KLL and KLX Auger electrons are taken from the paper of F. P. Larkins, *Atomic Data and Nuclear Data Tables* 20 (1977) 313. The relative emission probabilities of K Auger electrons are taken from the above cited paper of Schönfeld and Janßen (1994). The relative emission probabilities of L Auger electrons are derived from the corresponding absolute probabilities (Section 4.1) setting $P(KLL) = 100$.

4. Electron Emissions

The energies of the Auger electrons are the same as above. The energies of the conversion electrons are calculated from the energies of the gamma rays and the corresponding electron binding energies. The emission intensities of the Auger electrons are calculated from the transition probabilities of the EC transitions (2.1) and gamma transitions (2.3) using the atomic data given in Sections 3 and 3.2, the fractional electron capture probabilities and the conversion coefficients. The emission intensities of conversion electrons are calculated from the transition probabilities and conversion coefficients given in Section 2.3.

5. Photon Emissions

The X ray energies are the same as in 3.1. The energies of the gamma rays are taken from Helmer (1997HeZZ). The emission intensities of X rays were calculated from the decay scheme data. Schönfeld *et al.* (1994Sc44) measured the ratio R of K x-ray intensities following the decay of Ge-68 and Ga-68, they obtained $R = 9,57 (8)$. From the evaluation of Ge-68 decay data and the present one, a value $R = 9,36 (26)$ is derived. They are in agreement within the uncertainty limits.

To determine the relative gamma-ray emission intensities the following values have been considered:

	Vaughan 1969Va16	Carter 1968Ca15	Lange 1973La01	Vo 1994Vo15	Schönfeld 1994Sc44	Luca 2012Lu*	$\chi^2/n-1$	Adopted
227	-	-	-	0,0037 (15)				0,0037 (15)
483	-	-	-	0,0082 (9)				0,0082 (9)
579	0,7 (1) *	1,1 (2)	1,00 (12)	1,05 (15)	1,14 (15)	1,35 (30) *	0,2	1,06 (7)
683	-	-	-	0,0097 (6)		-		0,0097 (6)
806	2,2 (2) *	2,8 (2)	2,95 (12)	2,81 (14)	2,90 (31)	2,68 (34)	0,3	2,87 (8)
939	-	-	-	0,0055 (5)		-		0,0055 (5)
1077	100	100	100	100	100	100		
1166	-	-	-	0,0005 (3)		-		0,0005 (3)
1261	3,1 (2)	2,9 (2)	3,00 (7)	2,75 (14)	3,06 (31)	2,60 (28)	1	2,95 (6)
1744	0,5 (1)	0,28 (4)	0,30 (4)	0,295 (15)		-	1,4	0,297 (16)
1883	4,8 (3)	4,1 (4)	4,33 (12)	4,6 (2)	3,86 (59)	3,94 (42)	1,1	4,39 (10)
2338	<0,1	0,04 (2)	0,050 (6)	0,031 (3)		-	4	0,035 (5)
2821	-	-	0,015 (2)	0,0139 (11)				0,0144 (11)

* Omitted from statistical processing

Where there are only values from Vo *et al.*, these results have been adopted unchanged. In all other cases weighted means were calculated by using the Lweight program, the adopted uncertainty being the highest of the internal or external uncertainty.

Two published papers report absolute measurements of the gamma intensities, for the 1077 keV emission, they are:

- Schönfeld *et al.* (1994): 3,22 (3) %

- Luca *et al.* (2012): 3,25 (11) %

From these two results, an absolute value of the 1077 keV intensity is adopted which is the simple mean with the lowest experimental uncertainty, that is: $I_{g1077} = 3,235 (30) \%$.

All the other absolute intensity values were derived from I_{g1077} and the relative values as determined above.

A possible 1656 keV transition was observed only indirectly via conversion electrons by Slot *et al.* (1972Si03) ($P_{ce(1656)}/P_{ce(1077)} = 0,010 (2)$).

The absolute emission intensity of the annihilation photon (having 511 keV energy) is deduced from the decay scheme data and the theoretical EC/ β^+ ratios as described in § 2.1

7. References

- 1956Cr29** B.Crasemann, et al. Phys. Rev. 102,5 (1956) 1344 . Beta plus emission probabilities
1959Ho85 D.J.Horen. Phys. Rev. 113, 2 (1959) 572 . Beta plus emission probabilities
1959Ra04 M.K.Ramaswamy. Nucl. Phys. 10 (1959) 205 . Electron Capture/Beta plus Ratio
1960Ra22 M.K.Ramaswamy, P.Jastram. Nucl. Phys. 16 (1960) 113. Angular correl. 1077/1261 keV
1960Ra06 M.K.Ramaswamy, P.S.Jastram. Nuclear Phys. 16 (1960) 113 [δ]
1962Ko01 S.Kono J.Phys.Soc.Japan 17 (1962) 907 [δ]
1963Ta03 H.W.Taylor, R.McPherson Can.J.Phys. 41 (1963) 554 [δ]
1965Bo42 M.Bormann, et al.. Nucl. Phys. 63 (1965) 438. half-life
1965Eb01 T.G.Ebrey, et al.. Nucl. Phys. 61 (1965) 479. half-life
1968Ca15 H.K.Carter, J.H.Hamilton, A.V.Ramayya, J.J.Pinajian. Phys. Rev. 174 (1968) 1329.
Gamma-ray emission intensities
1969Va16 K.Vaughan, et al.. Nucl. Phys. A132 (1969) 561. Gamma-ray emission intensities
1971Oo01 J.M.Oottukulam, et al.. American. J. Phys. 39 (1971) 221. Half-life
1971Sm02 D.Smith, A.Williams. Int. J. Appl. Radiat. Isotop. 22 (1971) 615. Half-life

- 1971Ot01** H.Ottmar, N.M.Ahmed, U.Fanger, D.Heck, W.Michaelis, H.Schmidt Nucl.Phys. A164 (1971) 69 [δ]
- 1972Sl03** W.F.Slot, G.H.Dülfer, H.Van der Molen, A.Verheuil. Nucl. Phys. A186 (1972) 28. Beta emission energies, T ICC
- 1973La01** J.Lange, J.H.Hamilton, P.E.Little, D.L.Haffox, D.C.Morton, L.C.Whitblock, J.J.Pinajian. Phys. Rev. C7 (1973) 177. E2/M1 Mixing Ratios, Gamma emission probabilities
- 1977Kr17** K.S.Krane. At. Data. Nucl. Data Tables 20 (1977) 211. E2/M1 Mixing Ratios
- 1983Iw02** Y.Iwata, M.Kawamoto, Y.Yoshizawa. Int. J. Appl. Radiat. Isotop. 34 (1983) 1537. Half-life
- 1992Sy**** I.Sykora. Acta. Phys.Univ.Comen. 33 (1992) 25. Electron Capture/Beta plus Ratio
- 1994Sc44** E.Schönfeld, U.Schötzig, E.Günther, H.Schrader. Appl. Rad. Isotopes 45 (1994) 955. Electron Capture/Beta plus Ratio,
- 1994Vo15** D.T.Vo, W.H.Kelley, F.K.Wohn, J.C.Hill, J.P.Vary, M.A.Deleplanque, F.S.Stephens, R.M.Diamond, J.R.B.Oliveira, A.O.Machiavelli, J.A.Becker, E.A.Henry, M.J.Brinkman, M.A.Stover, J.E.Draper. Phys. Rev. C50 (1994) 1713. Gamma emission energies and probabilities, Mixing Ratios
- 2002Bu29** Nucl. Data Sheets 97 (2002) 1. Spin and parity, Half-life excited levels, Prod. modes
- 2001Ko07** Y.Kojima *et al.* Nucl. Instrum. Methods Phys. Res. A458 (2001) 656 [Q]
- 2003Au03** G.Audi, A.H.Wapstra, C.Thibault. Nucl. Phys. A729 (2003) 337. [Q]
- 2012Lu14** A. Luca *et al.* Appl. Rad. Isotopes, 70,9 (2012) 1876. Gamma-ray intensities

**⁶⁸Ge - Comments on evaluation
by M.-M. Bé and E. Schönfeld**

This evaluation was completed in 1996, it was updated to include the most recent Q value evaluation in December 2011. The literature available by this date was included.

Decay Scheme

The first excited state in Ga-68 is at 175 keV. As can be seen from the Q value an energy of this amount is not available. Thus, the decay scheme of Ge-68 is complete.

Nuclear Data

The $Q^+ = 106,9$ (24) keV is from 2011AuZZ, which supersedes the value of 106 (6) keV (2003Au03).

The following values of the half-life of ⁶⁸Ge have been taken into account:

Reference	$T_{1/2}$ (d)	
H. H. Hopkins, Jr.	250	omitted
G. Rudstam	288 (6)	
B. Crasemann <i>et al.</i>	275 (20)	
Waters <i>et al.</i> 1981	270,82 (27)	
Schönfeld <i>et al.</i> 1994	270,99 (19)	
Adopted	270,95 (26)	χ^2 crit = 3,8 ; $\chi^2 = 2,8$

The set of four data is consistent, the weighted mean is adopted. Due to their large uncertainties, the values of Rudstam and Crasemann have practically no weight. The external uncertainty of 0,26 d is adopted, whereas the internal uncertainty is 0,15 d .

Electron Capture Transition

The energy is derived from the Q value. The fractional probabilities for EC are calculated using the "Tables for Calculation of Electron Capture" (E. Schönfeld, PTB-Laboratory report 6.33-95-2 (1995)). These values are based on wave functions of Mann and Waber (1973) with exchange and overlap corrections of Bahcall and Vatai; see W. Bambynek *et al.*, *Rev. Mod. Phys.* 49(1977)77.

Atomic Data

All these data are taken from E. Schönfeld and H. Janssen, *Nucl. Instr. and Methods in Phys. Res. A* 369(1996)527.

X Radiations

The energies are based on the wavelengths compiled by J. A. Bearden, *Rev. Mod. Phys.* 39(1967)78. The relative probabilities for K_{α} radiation are based on $P(K_{\beta})/P(K_{\alpha})$ and $P(K_{\alpha 2})/P(K_{\alpha 1})$ values as given in the paper of Schönfeld and Janßen (1994).

Auger Electrons

The energies of KLL and KLX Auger electrons are taken from the paper of F. P. Larkins, (Atomic Data and Nuclear Data Tables 20 (1977) 313).

The relative emission probabilities of K Auger electrons are taken from the paper of Schönfeld and Janßen (1994). The relative emission probabilities of L Auger electrons are derived from the corresponding absolute probabilities setting $P(\text{KLL}) = 100$.

Radiation Emissions**Electron Emissions**

The energies of the Auger electrons are the same as above. The emission intensities are calculated from the transition probability of the EC transition, the fractional electron capture probabilities and by using the atomic data given in sections above.

Photon Emissions

The X ray energies are the same as described above. The emission intensities are calculated from the transition probability of the EC transition, the fractional electron capture probabilities and by using the atomic data given in sections above.

Main Production Mode

Zn-66(α , 2n)Ge-68

References

- 1950Ho**** H.H.Hopkins Jr. Phys. Rev. 77 (1950) 717. Half-life
1956Ru45 G.Rudstam. Thesis U. Uppsala (1956). Half-life
1956Cr25 B.Crasemann, D..E.Rehfuss, H.T.Easterday. Phys.Rev. 102 (1956) 1344. Half-life
1981Wa26 S.L.Waters, G.R.Forse, P.L.Horlock, M.J.Woods. Int. J. Appl. Radiat. Isotop. 32 (1981) 757. Half-life
1994Sc44 E.Schönfeld, U.Schötzig, E.Günther, H.Schrader. Int. J. Appl. Radiat. Isotop. 45 (1994) 955. Half-life, Electron Capture/Beta plus Ratio, Gamma emission and annihilation probabilities
2002Bu29 T.W.Burrows. Nucl. Data Sheets 97 (2002) 1. Spin and parity
2003Au03 G.Audi, A.H.Wapstra, C.Thibault. Nucl. Phys. A729 (2003) 337. Q value
2011AuZZ G.Audi, W.Meng. private communication (2011) Q

**⁷³Se – Comments on evaluation of decay data
by A. L. Nichols**

Evaluated: October/November 2013 and March 2014

Evaluation Procedures

Limitation of Relative Statistical Weight Method (LWM) and other analytical techniques were applied to obtain averaged data throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the *values* used to calculate the average.

Decay Scheme

⁷³Se ($T_{1/2} = 7.10$ h) decays 100 % by electron capture/ β^+ decay ($Q(\text{EC}) = 2725$ (7) keV) to various excited nuclear levels and the ground state of ⁷³As. A reasonably well-defined decay scheme was derived from the gamma-ray measurements of 1969Ma21, 1970Me20 (1970MeZZ) and 1980Te01(1978TeZY), consisting of 16 EC/6 β^+ transitions and 53 gamma-ray emissions. Eighteen gamma-ray emissions were originally unplaced in the decay scheme, but only seven of these transitions have been successfully incorporated into the proposed decay scheme during the course of the current study. Weighted mean relative emission probabilities were calculated for the main gamma rays, while equivalent data for the other gamma transitions were adopted primarily from the more comprehensive measurements of 1980Te01. All of these relative emission probabilities have been defined in terms of the 360.866-keV gamma ray (100 %).

Nuclear Data

⁷³Se decay is viewed as a potentially suitable β^+ emitter for application in positron emission tomography (PET).

Half-life of ⁷³Se

The measurements of 1957Be46, 1960Ri09, 1967Ra08, 1969Ma21 and 1976Bo19 were adopted to give a weighted mean half-life of 7.10 (9) hours based on the limitation of relative statistical weight method (LWM).

Reference	Half-life (h)
1948Co07	7.1*
1951Sc70	7.1*
1957Be46	7.1 (3)
1960Ri09	7.1 (2)
1967Ra08	7.0 (1)
1969Ma21	7.18 (9)
1976Bo19	7.08 (15)
Recommended value	7.10 (9)[§]

* Uncertainty not specified – not included in weighted mean analysis of the data set.

§ Recommended uncertainty has been adjusted from 0.06 to 0.09, in alignment with the smallest uncertainty of the values used to calculate the average value.

Limitation of relative statistical weight method (LWM), normalised residual method (NRM), Rajeval technique, bootstrap method, and Mandel-Paule approach were considered in the analysis of the data set.

Analytical method	Half-life (h)	$\chi^2/(N-1)$	$\chi^2/(N-1)_{\text{critical}}$
LWM	7.10 (6)*	0.45	3.32
NRM	7.10 (6)*	0.45	2.37
Rajeval	7.10 (6)*	0.45	–
Bootstrap	7.09 (3)	0.45	–
Mandel-Paule	null result	–	–

* Uncertainty increased from 0.06 to 0.09, in alignment with the smallest measured uncertainty of the values used to calculate the average value.

A half-life value of 7.10 (9) hours is recommended, as quantified by the LWM analytical procedure.

Half-life of daughter ⁷³As ground state

A half-life value of 80.30 (6) days for the ground state of daughter ⁷³As was adopted from 2004Si08.

Q-value

Q_{EC}-value for ⁷³Se EC/β⁺ decay of 2725 (7) keV was adopted from Wang *et al.* (2012Wa38).

Gamma-ray energies and emission probabilitiesEnergies

The well-defined nuclear level energies of 2004Si08 were used to calculate the gamma transition energies and their uncertainties, and these data were adjusted to account for gamma recoil in the formulation of recommended gamma-ray emission energies and uncertainties. Greater confidence was placed on this overall method of assignment because of the more wide-ranging origins of the gamma transition data in 2004Si08, even though the energies of a significant number of gamma-ray emissions have been directly measured by 1969Ma21, 1970Me20 and 1980Te01. The 1961.8-, 2434.1- and 2476.1-keV levels are notable exceptions to this approach – the energies of these particular nuclear levels of ⁷³As were determined from the energies of observed gamma rays that possess the potential to populate already known nuclear levels after depopulating proposed nuclear levels that can be closely identified with the energies of other particular levels listed by 2004Si08. Short-lived half-lives for the 67.039- and 427.906-keV nuclear levels were adopted from weighted-mean analyses of the measurements of 1956Ha10, 1963Bo16, 1963Bo26, 1966Ol04, 1969Iv02, 1969Ma21, 1972ReZN and 1977KeZY.

Adopted energies, spins and parities for the nuclear levels of ⁷³As (2004Si08).

Nuclear level number	Nuclear level energy (keV)	Spin and parity	Half-life
0	0.0	3/2 –	80.30 (6) d ⁷³ As
1	67.039 (8)	5/2 –	4.97 (7) ns
2	427.906 (21)	9/2 +	5.75 (20) μs
3	510.055 (17)	(5/2) +	
4	993.766 (12)	(7/2) –	
5	1037.13 (3)	(13/2) +	
6	1178.052 (21)	(7/2) –	
7	1275.14 (7)	(7/2) +	
8	1293.09 (10)	(11/2) +	
9	1293.37 (3)	(7/2) +	
10	1328.89 (5)	(7/2, 9/2) +	
11	1850.59 (5)	(9/2) +	
12	1910.13 (12)	(9/2 +, 11/2)	
13	1961.8 (2)*	(3/2, 5/2, 7/2)	
14	1975.41 (11)	(7/2, 9/2, 11/2 +)	
15	2180.66 (10)	(7/2, 9/2 +)	
16	2311.63 (6)	(7/2, 9/2 +)	
17	2434.1 (4)#	3/2+, 5/2+	
18	2476.1 (2)†	(11/2, 15/2)	
19	2482.87 (23)	(7/2, 9/2 +)	
20	2584.09 (11)	(7/2, 9/2 –)	

* Energy of nuclear level derived from observed 1451.7-keV gamma-ray depopulation, with spin assigned originally by 2004Si08 for a 1962.23 (17)-keV nuclear level.

Energy of nuclear level derived from observed 2006.2-keV gamma-ray depopulation, with spin and parity assigned originally by 2004Si08 for a 2436 (10)-keV nuclear level.

† Energy of nuclear level derived from observed 1439.0- and 2048.2-keV gamma-ray depopulation, with spin assigned originally by 2004Si08 for a 2475.49 (20)-keV nuclear level.

Half-life measurements and recommended values of the 67.039- and 427.906-keV nuclear levels of ⁷³As.

Level (keV)	Half-life								
	1956Ha10	1963Bo16	1963Bo26	1966Ol04	1969Iv02	1969Ma21	1972ReZN	1977KeZY	Recommended#
67.039 (ns)	< 5	6.1 (4)	5.0 (1)	4.95 (7)	–	–	–	–	4.97 (7) ns
427.906 (μs)	6.0 (2)	5.0 (4)	–	–	6 (1)	5.6 (3)	5.8 (2)	5.66 (23)	5.75 (20) μs

Data analysis based on limitation of relative statistical weight method (LWM), normalised residual method (NRM), Rajeval technique, and bootstrap method.

Gamma transition energies, and measured and recommended gamma-ray energies.

E_{TP} (keV) [#]	E_{γ} (keV)					
	1960Ri09	1967Ra08	1969Ma21	1970Me20	1980Te01	Recommended [§]
67.039 (8)	65 (2)	66.9(5)	67.0 (1)	67.0 (1)	67.07 (10)	67.039 (8)
360.867 (23)	360 (5)	360.4 (5)	361.1 (1)	360.9 (1)	361.2 (3)	360.866 (23)
427.906 (21)	–	–	–	–	428.3 (3)	427.905 (21)
443.016 (19)	–	–	443(1)	442.5 (8)	442.3 (3)	443.015 (19)
510.055 (17)	–	–	–	509.5 (8)	509.4	510.053 (17)
–	–	–	511.0	511.0	511.0	annihilation radiation
557.50 (11)	–	–	556.9 (8)	–	557.9 (5)	557.50 (11)
575.45 (9)	–	–	–	–	575.9 (5)	575.45 (9)
600.3 (3) ^Δ	–	–	–	–	600.33 (29)	600.3 (3) ^Δ
609.22 (4)	–	–	–	–	609.17 (19)	609.22 (4)
682.04 (11)	–	–	683*	682.0 (8)	682.25 (20)	682.04 (11)
700.27 (13)	–	–	700.5 (8)	699.9 (10)	700.0 (5)	700.27 (13)
765.09 (7)	–	–	765.0 (9)	764.4 (8)	765.07 (12)	765.09 (7)
783.32 (4)	–	–	783.7 (7)	782.9 (8)	783.7 (3)	783.32 (4)
793.0 (5) ^Δ	–	–	–	–	793.0 (5)	793.0 (5) ^Δ
813.46 (6)	–	–	–	–	813.38 (29)	813.46 (6)
818.84 (5)	–	–	819.0 (3)	818.4 (8)	818.65 (15)	818.84 (5)
847.23 (7)	–	–	847.5 (5)	847.2 (8)	847.16 (17)	847.22 (7)
856.82 (5) [Ⓛ]	–	–	–	–	857.0 (3)	856.81 (5) [Ⓛ]
865.18 (10) [¶] {	–	–	865.4 (2)	864.9 (5)	865.09 (12)	} 865.17 (10) [¶]
865.46 (3) [¶] {	–	–	–	–	–	} 865.45 (3) [¶]
873.00 (12)	–	–	–	–	872.56 (27)	872.99 (12)
887.29 (10)	–	–	–	–	887.46 (18)	887.28 (10)
900.98 (5)	–	–	901.2 (4)	901.3 (5)	900.73 (10)	900.97 (5)
926.727 (14)	–	–	–	–	926.19 (15)	926.721 (14)
930.09 (15) ^Δ	–	–	–	–	930.09 (15)	930.09 (15) ^Δ
968.0 (2) [‡]	–	–	969 (1)	968.1 (8)	968.1 (1)	968.0 (2) [‡]
982.74 (8)	–	–	983.6 (4)	982.5 (8)	982.75 (13)	982.73 (8)
993.766 (12)	–	–	–	–	993.52 (12)	993.759 (12)
1002.61 (10)	–	–	–	–	1001.9 (2)	1002.60 (10)
1018.26 (7)	–	–	–	–	1018.52 (13)	1018.25 (7)
1036.49 (9)	–	–	–	–	1036.38 (9)	1036.48 (9)
1111.013 (23)	–	–	1111.0 (4)	1110.8 (5)	1110.64 (6)	1111.004 (23)
1153.98 (24)	–	–	–	–	1153.9 (3)	1153.97 (24)
1159.0 (4) [‡]	–	–	–	–	1158.2 (4)	1159.0 (4) [‡]
1208.10 (7)	–	–	–	–	1207.9 (3)	1208.09 (7)
1215.4 (8) ^Δ	–	–	–	1215.4 (8)	1215.4 (8)	1215.4 (8) ^Δ
1226.33 (3)	–	–	–	–	1226.6 (9)	1226.32 (3)
1249.9 (2) ^Δ	–	–	–	–	1249.9 (2)	1249.9 (2) ^Δ
1275.14 (7)	–	–	–	–	1274.39 (21)	1275.13 (7)
1308.95 (13)	–	–	–	–	1308.9 (2)	1308.94 (13)
1317.86 (6)	–	–	–	–	1317.75 (21)	1317.85 (6)
1323.81 (20) ^Δ	–	–	–	–	1323.81 (20)	1323.81 (20) ^Δ
1340.54 (5)	–	–	1341.0 (5)	1340.7 (8)	1340.50 (7)	1340.53 (5)
1406.04 (11)	–	–	–	–	1406.3 (4)	1406.03 (11)
1422.68 (6)	–	–	1422.9 (2)	1422.5 (5)	1422.68 (7)	1422.67 (6)
1439.0 (2) [▲]	–	–	–	–	1439.10 (17)	1439.0 (2) [▲]
1451.7 (2) [‡]	–	–	–	–	1451.62 (2) [*]	1451.7 (2) [‡]
1482.22 (6)	–	–	1480 (1)	1482.3 (10)	1482.29 (12)	1482.20 (6)
1547.50 (11)	–	–	1548.1 (6)	1548.0 (8)	1547.45 (12)	1547.48 (11)
1670.61 (10)	–	–	–	–	1670.81 (16)	1670.59 (10)
1738.4 (5) ^Δ	–	–	–	–	1738.4 (5)	1738.4 (5) ^Δ
1752.75 (10)	–	–	–	–	1752.88 (20)	1752.73 (10)
1801.58 (6)	–	–	–	–	1801.36 (14)	1801.56 (6)
1847.8 (3) ^Δ	–	–	–	–	1847.82 (26)	1847.8 (3) ^Δ
1883.72 (6)	–	–	–	–	1883.85 (15)	1883.69 (6)

1889.57 (20) ^Δ	–	–	–	–	1889.57 (20)	1889.57 (20) ^Δ
1972.82 (23)	–	–	–	–	1973.4 (4)	1972.79 (23)
2006.2 (4) [‡]	–	–	–	–	2006.2 (4)	2006.2 (4) [‡]
2023.9 (3) ^Δ	–	–	–	–	2023.86 (26)	2023.9 (3) ^Δ
2048.2 (2) [▲]	–	–	–	–	2048.1 (8)	2048.2 (2) [▲]
2054.96 (23)	–	–	–	–	2053.8 (6)	2054.93 (23)
2156.18 (11)	–	–	–	–	2156.04 (14)	2156.15 (11)
2170.5 (3) ^Δ	–	–	–	–	2170.50 (26)	2170.5 (3) ^Δ
2517.05 (11)	–	–	–	–	2517.31 (25)	2517.00 (11)

[#] Determined from the nuclear level energies of 2004Si08.

[§] Calculated by subtracting gamma recoil from gamma transition energy (E_{TP} (keV)).

^Δ Unplaced in proposed decay scheme.

[¶] Unresolved doublet derived from proposed decay scheme.

^{*} Reported uncertainty believed to be in error when compared with defined uncertainties of all other equivalent data – modified to 1451.62(20) keV.

^Φ Proposed gamma-ray depopulation of 1850.59-keV nuclear level of ⁷³As.

[†] Proposed gamma-ray depopulation of 1961.8-keV nuclear level of ⁷³As.

[‡] Proposed gamma-ray depopulation of 2434.1-keV nuclear level of ⁷³As.

[▲] Proposed gamma-ray depopulation of 2476.1-keV nuclear level of ⁷³As.

Emission Probabilities

All of the relative emission probabilities were suitably refined and quantified in terms of the emission probability of the 360.866-keV gamma ray (100.0 %). Relative gamma-ray emission probabilities have been partially or fully determined in the measurements of 1960Ri09, 1967Ra08, 1969Ma21, 1970Me20 and 1980Te01, although much greater emphasis was placed on the studies of 1969Ma21, 1970Me20 and 1980Te01 when undertaking weighted-mean analyses of the appropriate and most significant relative emission probabilities. Considerably more detail can be found within the gamma-ray measurements of ten Brink *et al.* (1980Te01), and therefore the data of 1980Te01 were directly adopted for many of the low-intensity gamma-ray emissions in preference to any semi-artificial weighting procedure. Eighteen previously unplaced gamma-ray emissions were re-considered from the point of view of the known nuclear levels of ⁷³As and the possible observation of emissions from other radionuclides that might be generated during the various irradiations (i.e. Ga, Ge, As and other Se isotopes). A number of these gamma-ray emissions can be incorporated into the proposed decay scheme: 856.81, 968.0, 1159.0, 1439.0, 1451.7, 2006.2 and 2048.2 keV. Despite these efforts, eleven observed gamma rays remain unplaced in the proposed decay scheme, although they are contained within the recommended data set: 600.3, 793.0, 930.09, 1215.4, 1249.9, 1323.81, 1738.4, 1847.8, 1889.57, 2023.9 and 2170.5 keV.

Previously unplaced gamma transitions (2004Si08).

E_{γ} (keV)	p_{γ}^{rel}	Assignment	Comments
600.3 (3)	0.021 (3)	–	Only observed by 1980Te01. Possible to assign to 1178.06 → 577.90 and/or 993.78 → 393.43, but lack of evidence of depopulating 577.9-keV, and 393.4- and 309.1-keV γ -ray emissions, respectively, as specified by 1997So08.
793.0 (5)	0.066 (2)	–	Only observed by 1980Te01.
856.81 (5)	0.024 (6)	1850.59 → 993.766	Only observed by 1980Te01.
930.09 (15)	0.005 (1)	–	Only observed by 1980Te01.
968.0 (2)	0.012 (5)	1961.8 → 993.766	Observed by 1969Ma21, 1970Me20 and 1980Te01.
1159.0 (4)	0.003 (1)	2434.1 → 1275.14	Only observed by 1980Te01.
1215.4 (8)	0.065 (10)	–	Observed by 1970Me20 and 1980Te01.
1249.9 (2)	0.004 (1)	–	Only observed by 1980Te01.
1323.81 (20)	0.007 (1)	–	Only observed by 1980Te01.
1439.0 (2)	0.002 (1)	2476.1 → 1037.13	Only observed by 1980Te01.
1451.7 (2)	0.006 (2)	1961.8 → 510.055	Only observed by 1980Te01.
1738.4 (5)	0.002 (1)	–	Only observed by 1980Te01.
1847.8 (3)	0.008 (1)	–	Only observed by 1980Te01.
1889.57 (20)	0.003 (1)	–	Only observed by 1980Te01.
2006.2 (4)	0.002 (1)	2434.1 → 427.906	Only observed by 1980Te01.
2023.9 (3)	0.002 (1)	–	Only observed by 1980Te01.
2048.2 (2)	0.001 (1)	2476.1 → 427.906	Only observed by 1980Te01.
2170.5 (3)	0.002 (1)	–	Only observed by 1980Te01.

Extensive γ singles, γ - γ coincidence and conversion-electron spectral measurements by Sohler *et al.* (1997So08) of the prompt gamma-ray emission induced by the ⁷³Ge(p, $n\gamma$) reaction provided additional support towards defining the detailed structure of the ⁷³Se decay scheme.

Nine possible unresolved doublets have been proposed in the studies of 1997So08 that required further assessment to ensure their justifiable adoption or rejection:

- 317-keV doublet rejected – no 317-keV gamma rays observed in the ⁷³Se EC decay studies,
- 376-keV doublet rejected – no 376-keV gamma rays observed in the ⁷³Se EC decay studies,
- 484-keV doublet rejected – no 484-keV gamma rays observed in the ⁷³Se EC decay studies,
- 510-keV doublet – identified as the gamma-ray transition between the 510.055 → 0.0 keV nuclear levels in the ⁷³Se decay scheme; other possibility of a gamma-ray transition between the 577.9 → 67.04 keV nuclear levels rejected because no 577.9-keV gamma emission was observed, although specified to be the major depopulating transition from a 577.9-keV nuclear level by 1997So08,
- 600-keV doublet – both gamma-ray transitions between the 1178.0 → 577.9 and 993.8 → 393.4 keV nuclear levels rejected because of a lack of evidence of any depopulating 577.9-keV, and 393.4- and 309.1-keV gamma-ray emissions, respectively, as specified by 1997So08,
- 643-keV doublet rejected – no 643-keV gamma rays observed in the ⁷³Se EC decay studies,
- 724-keV doublet rejected – no 724-keV gamma rays observed in the ⁷³Se EC decay studies,
- 865-keV doublet accommodated as the two gamma transitions between the 1293.37 → 427.906 and 1293.09 → 427.906 keV nuclear levels,
- 994-keV doublet – identified as the gamma-ray transition between the 993.8 → 0.0 keV nuclear levels in the ⁷³Se decay scheme; other possibility of a gamma-ray transition between the 1077.8 → 84.3 keV nuclear levels rejected because there is no gamma-ray evidence for the existence of depopulating 1077.8- and 84.3-keV nuclear levels in ⁷³Se EC decay.

The undetermined relative emission probability of the 510.053-keV γ ray was calculated to be 0.27 (3) % from the relative emission probability ratio of the 510.053- and 443.015-keV γ rays depopulating the 510.055-keV nuclear level of ⁷³As, as determined by 1997So08 from their studies of the ⁷³Ge(p,n γ) reaction to be 214.1 (270) % and 41.1 (41) %, respectively. Another problem is posed by the 865-keV gamma-ray doublet which arises from the depopulation of both the 1293.37- and 1293.09-keV nuclear levels of ⁷³As, as detected in the γ - γ coincidence studies of the ⁷³Ge(p,n γ)⁷³As reaction by 1997So08. Gamma-ray spectroscopy studies of this region by 1969Ma21, 1970Me20 and 1980Te01 were only interpreted in terms of one 865-keV gamma transition and a single 1293-keV nuclear level, while 1997So08 undertook γ - γ coincidence measurements in which an 865-keV doublet of 865.17/865.45 keV was demonstrated to exist of unresolved structure. Problems arose in the evaluation when attempting to achieve a satisfactory population-depopulation balance for the 1293.37-keV level involving the 865.45-keV gamma ray and relative intensity determined by 1997So08, which also impacted inevitably on the 865.17-keV gamma-ray depopulation of the 1293.09-keV level. The relative emission probability of the 865.45-keV gamma ray was calculated to be 0.02 (1) % from the relative emission probability ratio of the 865.45- and 783.32-keV gamma rays depopulating the 1293.37-keV nuclear level of ⁷³As, as determined by 1997So08, while the relative emission probability of the 865.17-keV gamma ray depopulating the 1293.09-keV nuclear level of ⁷³As was defined as 0.540 (17) - 0.02 (1) = 0.52 (2) %.

Absolute total (EC + β^+) decay to the 67.039-keV was determined from the positron spectroscopy studies of 1951Sc70, 1956Ha10 and 1968Ak03. A weighted-mean value of 1.07 ± 0.20 was adopted for the relative P_{β^+} of the $\beta_{0,1}^+$ transition as derived from the positron emission measurements of 1951Sc70 and 1956Ha10 and 1968Ak03 with estimated uncertainties. The reasonably comprehensive gamma-ray spectroscopy studies of 1969Ma21, 1970Me20 and 1980Te01 support an extremely significant and dominant population of the 427.906-keV nuclear level of ⁷³As by direct EC/ β^+ decay. An EC/ β^+ ratio of 0.522 can be assigned to this allowed 2297-keV EC transition/1275-keV β^+ emission as determined from the tabulations of Gove and Martin (1971Go40). Both the equivalent relative P_{EC} and total $P_{(EC + \beta^+)}$ were calculated on the basis of the dominance of the 2297-keV EC/ β^+ transition. Furthermore, absolute total $P_{(EC + \beta^+)}$ values of 98.8 (2) % and 1.2 (2) % can be determined for the ($EC_{0,2} + \beta_{0,2}^+$) plus all lowly-populated higher-energy levels, and ($EC_{0,1} + \beta_{0,1}^+$), respectively.

Measured and calculated positron emission and EC transition probabilities (1951Sc70, 1956Ha10, 1968Ak03).

E _{β+} (keV)	Relative P _{β+}				ε/β (theory) 1971Go40	Relative P _{EC}	Relative total P _(EC+β+)	Absolute total P _(EC+β+) (%)
	1951Sc70	1956Ha10	1968Ak03 [#]	Recommended relative P _{β+}				
250 ?	1.3	–	–)	0.522)))
750 ?	11.8	–	8 → 8.8) 100) 52.2) 152.2) 98.8 (2)
1275 β _{0,2} ⁺	100	100	91.3 → 100))))
1636 β _{0,1} ⁺	1.4 → 1.4 (2)	~ 1 → 1.0 (2)	0.7 → 0.8 (2)	1.07 (20)	0.741	0.79	1.86 → 1.9 (2)	1.2 (2)
			Σ 100				Σ 154.1 ≡ 100	Σ 100

[#] Described as the intensities of partial β+ transitions.

A normalisation factor for the relative gamma-ray emission and EC/β⁺ transition probabilities was calculated by assigning a value of 100 % to the total gamma and EC/β⁺ transition probabilities directly populating the 67.039- and 0.0-keV nuclear levels of ⁷³As, assuming zero EC/β⁺ decay directly to the ground state based on spin-parity considerations (9/2+ → 3/2– constitutes a third forbidden non-unique transition (3, yes)):

$$\sum_{67.039\text{ keV}}^{\gamma\text{ population}} TP_{\gamma} + \sum_{0.0\text{ keV}}^{\gamma\text{ population}} TP_{\gamma} + \sum (P_{EC_{0,1}} + P_{\beta_{0,1}^+}) = 100$$

and excluding the 67.039-keV gamma transition between the two nuclear levels. Thus:

$$101.590 (20) F + 0.363 (34) F + 1.2 (2) = 100$$

where F is the normalisation factor for the relative gamma-ray emission and EC/β⁺ transition probabilities.

$$F = 98.8 (2) / 101.953 (39)$$

$$F = 0.9691 \pm 0.0020$$

Measured and recommended gamma-ray emission probabilities relative to P_γ(360.866 keV) of 100 %.

E _γ (keV)	P _γ ^{rel}	1960Ri09 [#]					Recommended [*]	
		1960Ri09 [#]	1967Ra08	1969Ma21	1970Me20	1980Te01	Recommended [*]	
γ _{1,0} 67.039 (8)	105 (10) → 66 (9)	76 (8)	73 (7)	73.0 (74)	72 (8)	73 (7) ↔ 81 (8) ⁺		
γ _{2,1} 360.866 (23)	160 (15) → 100	100	100	100	100	100		
γ _{2,0} 427.905 (21)	–	–	–	–	0.080 (15)	0.080 (15)		
γ _{3,1} 443.015 (19)	–	–	0.04 (1)	0.06 (1)	0.052 (3)	0.052 (3)		
γ _{3,0} 510.053 (17)	–	–	–	–	–	0.27 (3) [§]		
γ [±] 511	200 → 125 (12)	–	126 (3)	128 (13)	–	126 (3) → 133 (8) ^θ annihilation radiation		
γ _{11,8} 557.50 (11)	–	–	0.04 (1)	–	0.055 (2)	0.054 (2)		
γ _{11,7} 575.45 (9)	–	–	–	–	0.150 (7)	0.150 (7)		
γ _{-1,1} 600.3 (3) ^Δ	–	–	–	–	0.021 (3)	0.021 (3)		
γ _{5,2} 609.22 (4)	–	–	–	–	0.051 (4)	0.051 (4)		
γ _{14,9} 682.04 (11)	–	–	0.02 (1)	0.09 (4)	0.020 (2)	0.020 (2)		
γ _{14,7} 700.27 (13)	–	–	0.03 (1)	0.04 (1)	0.046 (2)	0.045 (2)		
γ _{7,3} 765.09 (7)	–	–	0.10 (3)	0.14 (2)	0.131 (2)	0.131 (2)		
γ _{9,3} 783.32 (4)	–	–	0.06 (2)	0.06 (2)	0.060 (2)	0.060 (2)		
γ _{-1,2} 793.0 (5) ^Δ	–	–	–	–	0.066 (2)	0.066 (2)		
γ _{11,5} 813.46 (6)	–	–	–	–	0.009 (1)	0.009 (1)		
γ _{10,3} 818.84 (5)	–	–	0.02 (1)	0.04 (2)	0.038 (2)	0.037 (2)		
γ _{7,2} 847.22 (7)	–	–	0.06 (2)	0.08 (1)	0.083 (6)	0.081 (6)		
γ _{11,4} 856.81 (5) ^θ	–	–	–	–	0.024 (6)	0.024 (6)		
γ _{8,2} 865.17 (10) [¶]	–	–	} 0.49 (3)	} 0.55 (6)	} 0.540 (17)	} 0.540 (17)	0.52 (2) [§]	
γ _{9,2} 865.45 (3) [¶]	–	–	}	}	}	}	0.02 (1) [§]	
γ _{12,5} 872.99 (12)	–	–	–	–	0.039 (7)	0.039 (7)		
γ _{15,9} 887.28 (10)	–	–	–	–	0.011 (8)	0.011 (8)		
γ _{10,2} 900.97 (5)	–	–	0.12 (3)	0.15 (2)	0.139 (2)	0.139 (2)		
γ _{4,1} 926.721 (14)	–	–	–	–	0.004 (1)	0.004 (1)		
γ _{-1,3} 930.09 (15) ^Δ	–	–	–	–	0.005 (1)	0.005 (1)		
γ _{13,4} 968.0 (2) [†]	–	–	0.010 (5)	0.02 (1)	–	0.012 (5)		

$\gamma_{16,10}$	982.73 (8)	–	–	0.02 (1)	0.03 (1)	0.035 (1)	0.035 (1)
$\gamma_{4,0}$	993.759 (12)	–	–	–	–	0.005 (1)	0.005 (1)
$\gamma_{15,6}$	1002.60 (10)	–	–	–	–	0.004 (1)	0.004 (1)
$\gamma_{16,9}$	1018.25 (7)	–	–	–	–	0.055 (2)	0.055 (2)
$\gamma_{16,7}$	1036.48 (9)	–	–	–	–	0.015 (1)	0.015 (1)
$\gamma_{6,1}$	1111.004 (23)	–	–	0.18 (5)	0.19 (2)	0.207 (2)	0.207 (2)
$\gamma_{19,10}$	1153.97 (24)	–	–	–	–	0.005 (1)	0.005 (1)
$\gamma_{17,7}$	1159.0 (4) [‡]	–	–	–	–	0.003 (1)	0.003 (1)
$\gamma_{7,1}$	1208.09 (7)	–	–	–	–	0.004 (1)	0.004 (1)
$\gamma_{-1,4}$	1215.4 (8) ^Δ	–	–	0.07 (1)	0.06 (1)	–	0.065 (10)
$\gamma_{9,1}$	1226.32 (3)	–	–	–	–	0.003 (2)	0.003 (2)
$\gamma_{-1,5}$	1249.9 (2) ^Δ	–	–	–	–	0.004 (1)	0.004 (1)
$\gamma_{7,0}$	1275.13 (7)	–	–	–	–	0.007 (1)	0.007 (1)
$\gamma_{20,7}$	1308.94 (13)	–	–	–	–	0.004 (1)	0.004 (1)
$\gamma_{16,4}$	1317.85 (6)	–	–	–	–	0.006 (1)	0.006 (1)
$\gamma_{-1,6}$	1323.81 (20) ^Δ	–	–	–	–	0.007 (1)	0.007 (1)
$\gamma_{11,3}$	1340.53 (5)	–	–	–	0.09 (2)	0.071 (2)	0.071 (2)
$\gamma_{20,6}$	1406.03 (11)	–	–	–	–	0.002 (1)	0.002 (1)
$\gamma_{11,2}$	1422.67 (6)	–	–	0.11 (3)	0.14 (2)	0.140 (5)	0.139 (5)
$\gamma_{18,5}$	1439.0 (2) [▲]	–	–	–	–	0.002 (1)	0.002 (1)
$\gamma_{13,3}$	1451.7 (2) [†]	–	–	–	–	0.006 (2)	0.006 (2)
$\gamma_{12,2}$	1482.20 (6)	–	–	0.008 (5)	0.03 (1)	0.023 (1)	0.023 (1)
$\gamma_{14,2}$	1547.48 (11)	–	–	0.03 (1)	0.04 (1)	0.032 (1)	0.032 (1)
$\gamma_{15,3}$	1670.59 (10)	–	–	–	–	0.005 (1)	0.005 (1)
$\gamma_{-1,7}$	1738.4 (5) ^Δ	–	–	–	–	0.002 (1)	0.002 (1)
$\gamma_{15,2}$	1752.73 (10)	–	–	–	–	0.011 (1)	0.011 (1)
$\gamma_{16,3}$	1801.56 (6)	–	–	–	–	0.020 (5)	0.020 (5)
$\gamma_{-1,8}$	1847.8 (3) ^Δ	–	–	–	–	0.008 (1)	0.008 (1)
$\gamma_{16,2}$	1883.69 (6)	–	–	–	–	0.031 (2)	0.031 (2)
$\gamma_{-1,9}$	1889.57 (20) ^Δ	–	–	–	–	0.003 (1)	0.003 (1)
$\gamma_{19,3}$	1972.79 (23)	–	–	–	–	0.001 (1)	0.001 (1)
$\gamma_{17,2}$	2006.2 (4) [‡]	–	–	–	–	0.002 (1)	0.002 (1)
$\gamma_{-1,10}$	2023.9 (3) ^Δ	–	–	–	–	0.002 (1)	0.002 (1)
$\gamma_{18,2}$	2048.2 (2) [▲]	–	–	–	–	0.001 (1)	0.001 (1)
$\gamma_{19,2}$	2054.93 (23)	–	–	–	–	0.003 (1)	0.003 (1)
$\gamma_{20,2}$	2156.15 (11)	–	–	–	–	0.005 (1)	0.005 (1)
$\gamma_{-1,11}$	2170.5 (3) ^Δ	–	–	–	–	0.002 (1)	0.002 (1)
$\gamma_{20,1}$	2517.00 (11)	–	–	–	–	0.005 (1)	0.005 (1)

[#] Emission probabilities expressed relative to P_γ for 511-keV annihilation radiation of 200 %.

^{*} Emission probabilities expressed relative to $P_\gamma(360.866\text{ keV})$ of 100 %.

⁺ Weighted-mean relative emission probability can be either adjusted in an arbitrary manner from 73 (7) % to 81 (8) % if assigned multipolarity of 100 % M1, or modified to (M1 + 4.3 % E2) multipolarity to maintain relative emission probability of 73 (7) % to ensure satisfactory depopulation of the 67.039-keV nuclear level to the ground state of ⁷³As.

[§] Calculated from relative emission probability ratio of 510.053- and 443.015-keV γ rays depopulating the 510.055-keV nuclear level of ⁷³As as determined by 1997So08 from studies of ⁷³Ge(p, γ) reaction.

^o 511-keV annihilation radiation: weighted-mean relative emission probability of 126 (3) % compares with a value of 133 (1) % calculated from the total absolute positron emission probability and normalisation factor - uncertainty increased artificially to derive a more acceptable relative emission probability of 133 (8) %.

^Δ Unplaced in proposed decay scheme.

[†] Unresolved doublet derived from defined decay scheme.

[§] Relative emission probability of 865.45-keV γ ray calculated to be 0.02 (1) % from relative emission probability ratio of 865.45- and 783.32-keV γ rays depopulating 1293.37-keV nuclear level of ⁷³As (relative emission probabilities of 18.1 (151) and 69.78 (21), respectively, from 1997So08); as a consequence, the relative emission probability of 865.17-keV γ ray depopulating the 1293.09-keV nuclear level of ⁷³As was defined as 0.540 (17) – 0.02 (1) → 0.52 (2) %.

^o Gamma ray depopulates proposed 1850.59-keV nuclear level of ⁷³As.

[†] Gamma ray depopulates proposed 1961.8-keV nuclear level of ⁷³As.

[‡] Gamma ray depopulates proposed 2434.1-keV nuclear level of ⁷³As.

[▲] Gamma ray depopulates proposed 2476.1-keV nuclear level of ⁷³As.

The measured relative emission probability of the 67.039-keV gamma ray proved to be particularly problematic from the point of view of the derived depopulation of the 67.039 nuclear level to the ground state of ⁷³As. Based on a calculated relative transition probability of 102.83 (21) and 100 % M1 total internal conversion coefficient of 0.272 (4), a relative emission probability of 80.8 (3) % was calculated in comparison to a weighted-mean value of 73 (7) % from the measured data:

$$TP_{\gamma}^{rel}(67.039 \text{ keV}) = \sum_{67.039 \text{ keV}}^{\gamma \text{ population}} TP_{\gamma}^{rel} + \text{total relative } P_{EC_{0,1}}$$

$$= 101.590 (20) + 1.24 (21) = 102.83 (21)$$

Therefore:

$$P_{\gamma}^{rel}(67.039 \text{ keV}) = \frac{TP_{\gamma}(67.039 \text{ keV})}{(1 + \text{total } ICC_{67.039 \text{ keV}})} = \frac{102.83 (21)}{1.272 (4)} = 80.8 (3)$$

Thus, a relative emission probability of 81 (8) % could be adopted for the 67.039-keV gamma ray, whereby the recommended uncertainty could be brought into a sensible degree of alignment with the measured data and their uncertainties of 1969Ma21, 1970Me20 and 1980Te01.

An alternative procedure would be to modify the total internal conversion coefficient to a value of 0.409 (14) in order to achieve good agreement between the weighted-mean relative emission probability of 73 (7) % and the calculated relative transition probability of 102.83 (21) for the 67.039-keV gamma transition:

$$P_{\gamma}^{rel}(67.039 \text{ keV}) = \frac{TP_{\gamma}^{rel}(67.039 \text{ keV})}{(1 + \text{total } ICC_{67.039 \text{ keV}})}$$

$$\text{and therefore } (1 + \text{total } ICC_{67.039 \text{ keV}}) = \frac{TP_{\gamma}^{rel}(67.039 \text{ keV})}{P_{\gamma}^{rel}(67.039 \text{ keV})} = \frac{102.83 (21)}{73 (7)} = 1.409 (14)$$

$$\text{total } ICC(67.039 \text{ keV}) = 0.409 (14), \text{ which defines a calculated mixing ratio } (\delta) \text{ of } 0.211 (10).$$

Further studies of this gamma-ray emission are merited to improve understanding and confidence in this important feature of the decay scheme of ⁷³Se.

Multipolarities, and Internal Conversion and Internal-Pair Coefficients

The nuclear level scheme specified by Singh has been used to define the multipolarities of the gamma transitions on the basis of known spins and parities (2004Si08). Various γ - γ coincidence studies have resulted in the determination of a limited number of transition types and mixing ratios: 100 % M1 for the 67.039-keV gamma ray, (99.88 % M2 + 0.12 % E3) for the 360.866-keV gamma ray, and 100 % E3 for the 427.905-keV gamma ray (1963Bo26, 1967Ra08, 1969Ma21, 1970Me20, 1970Qu03, 1980Te01 and 1988Be39). While various E1 and E2 gamma transitions could also be identified, mixing ratios for all of the low-intensity (M1 + E2) gamma rays have not been defined and quantified. The weighted-mean emission probability and resulting transition probability of the substantial 67.039-keV gamma emission is in some disagreement with the rest of the proposed decay-scheme data. As described above, this problem can be addressed by either arbitrarily increasing the relative emission probability from 73 (7) % to 81 (8) % for a 100 % M1 transition, or modifying the mixing ratio to 0.211 (10) to give a multipolarity of (95.7 % M1 + 4.3 % E2). However, the weighted mean value of the measured relative emission probabilities has been adopted to be 73 (7) %, with the internal conversion coefficients defined for a 100 % M1 transition (α_{total} of 0.272 (4)).

Recommended internal conversion coefficients have been determined from the frozen orbital approximation of Kibédi *et al.* (2008Ki07), based on the theoretical model of Band *et al.* (2002Ba85, 2002Ra45). A significant number of gamma transitions undergo decay via internal-pair formation, and the coefficient for this process has also been quantified in a few cases from the tabulations of 2008Ki07.

Gamma-ray emissions: multipolarities, and theoretical internal-conversion (frozen orbital approximation) and internal-pair formation coefficients.

E_γ (keV)	Multipolarity	α_K	α_L	α_{M+}	α_{totalICC}	α_{IPF}	α_{total}
67.039 (8)	100 % M1 (1967Ra08, 1970Qu03, 1988Be39)	0.241 (4)	0.026 4 (4)	0.004 6	0.272 (4)	–	0.272 (4)
	or 95.7 % M1+ 4.3 % E2 δ = 0.211 (10)	0.356 (12)	0.045 4 (19)	0.007 6	0.409 (14)	–	0.409 (14)
360.866 (23)	99.88 % M2 + 0.12 % E3 δ = – 0.035 (10) (1963Bo26, 1970Qu03, 1988Be39)	0.011 65 (17)	0.001 286 (18)	0.000 214	0.013 15 (19)	–	0.013 15 (19)
427.905 (21)	E3	0.01195 (17)	0.001 397 (20)	0.000 223	0.013 57 (19)	–	0.013 57 (19)
443.015 (19)	(E1)	0.000 926 (13)	0.000 095 5 (14)	0.000 015 5	0.001 037 (15)	–	0.001 037 (15)
510.053 (17)	(E1)	0.000 650 (10)	0.000 067 0 (10)	0.000 011	0.000 728 (11)	–	0.000 728 (11)
557.50 (11)	(M1 + E2)	–	–	–	–	–	–
575.45 (9)	(M1 + E2)	–	–	–	–	–	–
600.3 (3) ^Δ	–	–	–	–	–	–	–
609.22 (4)	(E2)	0.001 258 (18)	0.000 132 7 (19)	0.000 021 3	0.001 412 (20)	–	0.001 412 (20)
682.04 (11)	(E1)	0.000 330 (5)	0.000 033 8 (5)	0.000 005 2	0.000 369 (6)	–	0.000 369 (6)
700.27 (13)	(M1 + E2)	–	–	–	–	–	–
765.09 (7)	(M1 + E2)	–	–	–	–	–	–
783.32 (4)	(M1 + E2)	–	–	–	–	–	–
793.0 (5) ^Δ	–	–	–	–	–	–	–
813.46 (6)	(E2)	0.000 573 (8)	0.000 059 7 (9)	0.000 009 3	0.000 642 (9)	–	0.000 642 (9)
818.84 (5)	(M1 + E2)	–	–	–	–	–	–
847.22 (7)	(M1 + E2)	–	–	–	–	–	–
856.81 (5)	(E1)	0.000 203 (3)	0.000 020 7 (3)	0.000 003 3	0.000 227 (4)	–	0.000 227 (4)
865.17 (10)	(M1 + E2)	–	–	–	–	–	–
865.45 (3)	(M1 + E2)	–	–	–	–	–	–
872.99 (12)	(E2)	0.000 479 (7)	0.000 049 9 (7)	0.000 008 1	0.000 537 (8)	–	0.000 537 (8)
887.28 (10)	(M1 + E2)	–	–	–	–	–	–
900.97 (5)	(M1 + E2)	–	–	–	–	–	–
926.721 (14)	(M1 + E2)	–	–	–	–	–	–
930.09 (15) ^Δ	–	–	–	–	–	–	–
968.0 (2)	–	–	–	–	–	–	–
982.73 (8)	(M1 + E2)	–	–	–	–	–	–
993.759 (12)	(E2)	0.000 350 (5)	0.000 036 3 (5)	0.000 005 7	0.000 392 (6)	–	0.000 392 (6)
1002.60 (10)	(E1)	0.000 148 4 (21)	0.000 015 17 (22)	0.000 002 4	0.000 166 0 (24)	–	0.000 166 0 (24)
1018.25 (7)	(M1 + E2)	–	–	–	–	–	–
1036.48 (9)	(M1 + E2)	–	–	–	–	–	–

Comments on evaluation

⁷³Se

1111.004 (23)	(M1 + E2)	-	-	-	-	-	-
1153.97 (24)	(M1 + E2)	-	-	-	-	-	-
1159.0 (4)	-	-	-	-	-	-	-
1208.09 (7)	(E1)	0.000 105 0 (15)	0.000 010 72 (15)	0.000 001 78	0.000 117 5 (15)	0.000 052 5 (8)	0.000 170 0 (24)
1215.4 (8) ^Δ	-	-	-	-	-	-	-
1226.32 (3)	(E1)	0.000 102 3 (15)	0.000 010 43 (15)	0.000 001 67	0.000 114 4 (15)	0.000 064 3 (9)	0.000 179 (3)
1249.9 (2) ^Δ	-	-	-	-	-	-	-
1275.13 (7)	(M2)	0.000 394 (6)	0.000 040 8 (6)	0.000 006 8	0.000 442 (6)	0.000 004 16 (6)	0.000 446 (7)
1308.94 (13)	(E1)	0.000 091 2 (13)	0.000 009 30 (13)	0.000 001 53	0.000 102 0 (13)	0.000 120 3 (17)	0.000 222 (4)
1317.85 (6)	(E1)	0.000 090 2 (13)	0.000 009 19 (13)	0.000 001 50	0.000 100 8 (13)	0.000 126 1 (18)	0.000 227 (4)
1323.81 (20) ^Δ	-	-	-	-	-	-	-
1340.53 (5)	(E2)	0.000 180 (3)	0.000 018 5 (3)	0.000 003 0	0.000 201 (3)	0.000 037 6 (6)	0.000 239 (4)
1406.03 (11)	(M1 + E2)	-	-	-	-	-	-
1422.67 (6)	(M1 + E2)	-	-	-	-	-	-
1439.0 (2)	-	-	-	-	-	-	-
1451.7 (2)	-	-	-	-	-	-	-
1482.20 (6)	(M1 + E2)	-	-	-	-	-	-
1547.48 (11)	(M1 + E2)	-	-	-	-	-	-
1670.59 (10)	(M1 + E2)	-	-	-	-	-	-
1738.4 (5) ^Δ	-	-	-	-	-	-	-
1752.73 (10)	(M1 + E2)	-	-	-	-	-	-
1801.56 (6)	(M1 + E2)	-	-	-	-	-	-
1847.8 (3) ^Δ	-	-	-	-	-	-	-
1883.69 (6)	(M1 + E2)	-	-	-	-	-	-
1889.57 (20) ^Δ	-	-	-	-	-	-	-
1972.79 (23)	(M1 + E2)	-	-	-	-	-	-
2006.2 (4)	-	-	-	-	-	-	-
2023.9 (3) ^Δ	-	-	-	-	-	-	-
2048.2 (2)	-	-	-	-	-	-	-
2054.93 (23)	(M1 + E2)	-	-	-	-	-	-
2156.15 (11)	(E1)	0.000 041 3 (6)	0.000 004 19 (6)	0.000 000 69	0.000 046 1 (6)	0.000 739 (11)	0.000 785 (11)
2170.5 (3) ^Δ	-	-	-	-	-	-	-
2517.00 (11)	(M1 + E2)	-	-	-	-	-	-

^Δ Unplaced in proposed decay scheme.

Recommended gamma-ray energies, relative and absolute emission probabilities, and transition probabilities.

	E_γ (keV)	P_γ^{rel}	P_γ^{abs}	Transition probability (%)
$\gamma_{1.0}$	67.039 (8)	73 (7) 100 % M1 \leftrightarrow (95.7 % M1 + 4.3 % E2)	70.7 \rightarrow 71 (7) 100 % M1 \leftrightarrow (95.7 % M1 + 4.3 % E2)	90 (9) 100 % M1 \leftrightarrow 100.0 (95.7 % M1 + 4.3 % E2)
$\gamma_{2.1}$	360.866 (23)	100	96.91 (20)	98.18 (20)
$\gamma_{2.0}$	427.905 (21)	0.080 (15)	0.078 (14)	0.079 (14)
$\gamma_{3.1}$	443.015 (19)	0.052 (3)	0.050 (3)	0.050 (3)
$\gamma_{3.0}$	510.053 (17)	0.27 (3)	0.26 (3)	0.26 (3)
γ^+	511	126 (3) \rightarrow 133 (8)	129 (8)	–
$\gamma_{11.8}$	557.50 (11)	0.054 (2)	0.052 (2)	0.052 (2)
$\gamma_{11.7}$	575.45 (9)	0.150 (7)	0.146 (7)	0.146 (7)
$\gamma_{-1.1}$	600.3 (3) ^{Δ}	0.021 (3)	0.020 (3)	0.020 (3)
$\gamma_{5.2}$	609.22 (4)	0.051 (4)	0.049 (4)	0.049 (4)
$\gamma_{14.9}$	682.04 (11)	0.020 (2)	0.019 (2)	0.019 (2)
$\gamma_{14.7}$	700.27 (13)	0.045 (2)	0.044 (2)	0.044 (2)
$\gamma_{7.3}$	765.09 (7)	0.131 (2)	0.127 (2)	0.127 (2)
$\gamma_{9.3}$	783.32 (4)	0.060 (2)	0.058 (2)	0.058 (2)
$\gamma_{-1.2}$	793.0 (5) ^{Δ}	0.066 (2)	0.064 (2)	0.064 (2)
$\gamma_{11.5}$	813.46 (6)	0.009 (1)	0.009 (1)	0.009 (1)
$\gamma_{10.3}$	818.84 (5)	0.037 (2)	0.036 (2)	0.036 (2)
$\gamma_{7.2}$	847.22 (7)	0.081 (6)	0.078 (6)	0.078 (6)
$\gamma_{11.4}$	856.81 (5)	0.024 (6)	0.023 (6)	0.023 (6)
$\gamma_{8.2}$	865.17 (10)	0.52 (2)	0.50 (2)	0.50 (2)
$\gamma_{9.2}$	865.45 (3)	0.02 (1)	0.02 (1)	0.02 (1)
$\gamma_{12.5}$	872.99 (12)	0.039 (7)	0.038 (7)	0.038 (7)
$\gamma_{15.9}$	887.28 (10)	0.011 (8)	0.011 (8)	0.011 (8)
$\gamma_{10.2}$	900.97 (5)	0.139 (2)	0.135 (2)	0.135 (2)
$\gamma_{4.1}$	926.721 (14)	0.004 (1)	0.004 (1)	0.004 (1)
$\gamma_{-1.3}$	930.09 (15) ^{Δ}	0.005 (1)	0.005 (1)	0.005 (1)
$\gamma_{13.4}$	968.0 (2)	0.012 (5)	0.012 (5)	0.012 (5)
$\gamma_{16.10}$	982.73 (8)	0.035 (1)	0.034 (1)	0.034 (1)
$\gamma_{4.0}$	993.759 (12)	0.005 (1)	0.005 (1)	0.005 (1)
$\gamma_{15.6}$	1002.60 (10)	0.004 (1)	0.004 (1)	0.004 (1)
$\gamma_{16.9}$	1018.25 (7)	0.055 (2)	0.053 (2)	0.053 (2)
$\gamma_{16.7}$	1036.48 (9)	0.015 (1)	0.015 (1)	0.015 (1)
$\gamma_{6.1}$	1111.004 (23)	0.207 (2)	0.201 (2)	0.201 (2)
$\gamma_{19.10}$	1153.97 (24)	0.005 (1)	0.005 (1)	0.005 (1)
$\gamma_{17.7}$	1159.0 (4)	0.003 (1)	0.003 (1)	0.003 (1)
$\gamma_{7.1}$	1208.09 (7)	0.004 (1)	0.004 (1)	0.004 (1)
$\gamma_{-1.4}$	1215.4 (8) ^{Δ}	0.065 (10)	0.063 (10)	0.063 (10)
$\gamma_{9.1}$	1226.32 (3)	0.003 (2)	0.003 (2)	0.003 (2)
$\gamma_{-1.5}$	1249.9 (2) ^{Δ}	0.004 (1)	0.004 (1)	0.004 (1)
$\gamma_{7.0}$	1275.13 (7)	0.007 (1)	0.007 (1)	0.007 (1)
$\gamma_{20.7}$	1308.94 (13)	0.004 (1)	0.004 (1)	0.004 (1)
$\gamma_{16.4}$	1317.85 (6)	0.006 (1)	0.006 (1)	0.006 (1)
$\gamma_{-1.6}$	1323.81 (20) ^{Δ}	0.007 (1)	0.007 (1)	0.007 (1)
$\gamma_{11.3}$	1340.53 (5)	0.071 (2)	0.069 (2)	0.069 (2)
$\gamma_{20.6}$	1406.03 (11)	0.002 (1)	0.002 (1)	0.002 (1)
$\gamma_{11.2}$	1422.67 (6)	0.139 (5)	0.135 (5)	0.135 (5)
$\gamma_{18.5}$	1439.0 (2)	0.002 (1)	0.002 (1)	0.002 (1)
$\gamma_{13.3}$	1451.7 (2)	0.006 (2)	0.006 (2)	0.006 (2)
$\gamma_{12.2}$	1482.20 (6)	0.023 (1)	0.022 (1)	0.022 (1)
$\gamma_{14.2}$	1547.48 (11)	0.032 (1)	0.031 (1)	0.031 (1)
$\gamma_{15.3}$	1670.59 (10)	0.005 (1)	0.005 (1)	0.005 (1)
$\gamma_{-1.7}$	1738.4 (5) ^{Δ}	0.002 (1)	0.002 (1)	0.002 (1)
$\gamma_{15.2}$	1752.73 (10)	0.011 (1)	0.011 (1)	0.011 (1)
$\gamma_{16.3}$	1801.56 (6)	0.020 (5)	0.019 (5)	0.019 (5)
$\gamma_{-1.8}$	1847.8 (3) ^{Δ}	0.008 (1)	0.008 (1)	0.008 (1)
$\gamma_{16.2}$	1883.69 (6)	0.031 (2)	0.030 (2)	0.030 (2)
$\gamma_{-1.9}$	1889.57 (20) ^{Δ}	0.003 (1)	0.003 (1)	0.003 (1)

$\gamma_{19,3}$	1972.79 (23)	0.001 (1)	0.001 (1)	0.001 (1)
$\gamma_{17,2}$	2006.2 (4)	0.002 (1)	0.002 (1)	0.002 (1)
$\gamma_{-1,10}$	2023.9 (3) ^Δ	0.002 (1)	0.002 (1)	0.002 (1)
$\gamma_{18,2}$	2048.2 (2)	0.001 (1)	0.001 (1)	0.001 (1)
$\gamma_{19,2}$	2054.93 (23)	0.003 (1)	0.003 (1)	0.003 (1)
$\gamma_{20,2}$	2156.15 (11)	0.005 (1)	0.005 (1)	0.005 (1)
$\gamma_{-1,11}$	2170.5 (3) ^Δ	0.002 (1)	0.002 (1)	0.002 (1)
$\gamma_{20,1}$	2517.00 (11)	0.005 (1)	0.005 (1)	0.005 (1)

^Δ Unplaced in proposed decay scheme.

Energies and emission probabilities of noteworthy internal conversion electrons.

		Energy (keV)	Electrons per 100 disint.
ec _{1,0T}	(As)	55.172 – 67.037	19 (11)
ec _{1,0K}	(As)	55.172 (8)	17 (10)
ec _{1,0L}	(As)	65.513 – 65.716	1.8 (10)
ec _{1,0M+}	(As)	66.836 – 67.037	0.32 (18)
ec _{2,1T}	(As)	349.00 – 360.86	1.27 (30)
ec _{2,1K}	(As)	349.00 (3)	1.13 (25)
ec _{2,1L}	(As)	359.34 – 359.54	0.12 (3)
ec _{2,1M+}	(As)	360.66 – 360.86	0.021 (5)

EC/ β^+ Transitions

Energies

All EC/ β^+ energies were derived from the structural details of the proposed decay scheme. The nuclear level energies of 2004Si08 and evaluated Q_{EC} -value of 2725 (7) keV (2012Wa38) were used to determine the recommended energies and uncertainties of the EC transitions and β^+ emissions.

Transition and Emission Probabilities

A large majority of the EC transition probabilities were derived for the population-depopulation imbalances of the relative emission probabilities of the gamma rays, their theoretical internal conversion and internal-pair formation coefficients. A value of 0.9691 (20) was adopted as the normalisation factor in order to determine the absolute transition and emission probabilities of the EC and β^+ particles from their resulting relative transition and emission probabilities. Component EC and β^+ transition and emission probabilities were determined from EC/ β^+ ratios (1971Go40), and $\log ft$ values and average E_{β^+} energies were derived by means of the LOGFT code. Fractional EC probabilities P_K , P_L , P_M and P_N were calculated by means of the EC-CAPTURE code (1998Sc28) as developed from the data tabulations of 1995ScZY. Possible EC transitions directly to the 1293.37, 1037.13-, 993.766- and 510.055-keV nuclear levels of ⁷³As were assumed to be zero, based on gamma population-depopulation balances and/or spin-parity considerations, while EC decay directly to the ground state was assigned a value of zero from a spin-parity assessment alone (third forbidden non-unique (3, yes)).

The absolute emission probability of the 511-keV annihilation radiation was derived from the total absolute positron emission probability of 64.6 (4) % as determined during the evaluation of the EC/ β^+ transitions/emissions:

$$\text{absolute } P_{\gamma}^{\pm} = 2 \times 64.6 (4) = 129.2 (8) \% \rightarrow 129 (1) \%$$

$$[\text{relative } P_{\gamma}^{\pm} = \frac{\text{absolute } P_{\gamma}^{\pm}}{F} = \frac{129.2 (8)}{0.9691 (20)} = 133.3 (9) \% \rightarrow 133 (1) \%$$

where F is the normalisation factor for the gamma-ray emission probabilities. Recommended uncertainties have been artificially increased:

$$\text{relative } P_{\gamma}^{\pm} = 133 (8) \%, \text{ and therefore } \text{absolute } P_{\gamma}^{\pm} = 129 (8) \%$$

to ensure full coverage of the range of measured relative emission probabilities for the annihilation radiation (i.e. 125 (12) %, 126 (3) % and 128 (13) % \rightarrow 133 (8) %).

A decay scheme was derived that consists of 16EC/ $6\beta^+$ transitions and 53 gamma-ray emissions. Substantial 511-keV annihilation radiation was also observed. Eleven additional gamma-ray emissions of reasonably low intensity could not be placed in the proposed decay scheme.

Recommended energies and transition probabilities of the EC/ β^+ decay of ⁷³Se.

E_{EC} (keV)*	E_{β^+} (keV)	Av. E_{β^+} (keV)	P_{EC} (total)	ϵ/β^+ (theory) 1971Go40	P_{EC}	P_{β^+}	⁷³ Se	⁷³ As	transition type	log ft	P_K	P_L	P_M	P_N	
EC _{0,20}	141 (7)	–	–	0.0155 (20)	–	0.0155 (20)	–	9/2 +	(7/2, 9/2 –)	(1 st forbidden non-unique)	6.2	0.8646 (19)	0.1136 (16)	0.0199 (5)	0.0020 (2)
EC _{0,19}	242 (7)	–	–	0.0087 (20)	–	0.0087 (20)	–	9/2 +	(7/2, 9/2 +)	allowed	7.0	0.8723 (16)	0.1072 (13)	0.0186 (4)	0.0018 (2)
EC _{0,18}	249 (7)	–	–	0.0029 (10)	–	0.0029 (10)	–	9/2 +	(11/2, 15/2)	(allowed)	7.5	0.8726 (16)	0.1070 (13)	0.0186 (4)	0.0018 (2)
EC _{0,17}	291 (7)	–	–	0.0048 (19)	–	0.0048 (19)	–	9/2 +	3/2 +, 5/2 +	[2 nd forbidden non-unique]	7.4	0.8740 (16)	0.1058 (13)	0.0184 (4)	0.0018 (1)
EC _{0,16}	413 (7)	–	–	0.157 (6)	–	0.157 (6)	–	9/2 +	(7/2, 9/2 +)	(allowed)	6.2	0.8764 (16)	0.1038 (13)	0.0180 (4)	0.0018 (1)
EC _{0,15}	544 (7)	–	–	0.030 (8)	–	0.030 (8)	–	9/2 +	(7/2, 9/2 +)	(allowed)	7.1	0.8778 (15)	0.1027 (13)	0.0178 (4)	0.0017 (1)
EC _{0,14}	750 (7)	–	–	0.094 (3)	–	0.094 (3)	–	9/2 +	(7/2, 9/2, 11/2 +)	(allowed)	6.9	0.8789 (15)	0.1018 (13)	0.0176 (4)	0.0017 (1)
EC _{0,13}	763 (7)	–	–	0.017 (5)	–	0.017 (5)	–	9/2 +	(3/2, 5/2, 7/2)	(allowed)	7.7	0.8790 (15)	0.1017 (13)	0.0176 (4)	0.0017 (1)
EC _{0,12}	815 (7)	–	–	0.060 (7)	–	0.060 (7)	–	9/2 +	(9/2 +, 11/2)	(allowed)	7.2	0.8792 (15)	0.1016 (13)	0.0175 (4)	0.0017 (1)
EC _{0,11}	874 (7)	–	–	0.433 (11)	–	0.433 (11)	–	9/2 +	(9/2) +	(allowed)	6.4	0.8794 (15)	0.1014 (12)	0.0175 (4)	0.0017 (1)
EC _{0,10}	1396 (7)	374 (7)	164 (3)	0.132 (3)	37.50	0.129 (3)	0.0034 (2)	9/2 +	(7/2, 9/2) +	(allowed)	7.3	0.8804 (15)	0.1006 (12)	0.0173 (4)	0.0017 (1)
EC _{0,8}	1432 (7)	410 (7)	179 (3)	0.452 (20)	26.30	0.435 (19)	0.017 (2)	9/2 +	(11/2) +	(allowed)	6.8	0.8804 (15)	0.1006 (12)	0.0173 (4)	0.0017 (1)
EC _{0,7}	1450 (7)	428 (7)	187 (3)	0.006 (2)	22.39	0.0057 (19)	0.0003 (1)	9/2 +	(7/2) +	(allowed)	8.7	0.8804 (15)	0.1006 (12)	0.0173 (4)	0.0017 (1)
EC _{0,6}	1547 (7)	525 (7)	228 (3)	0.195 (2)	10.59	0.178 (2)	0.017 (1)	9/2 +	(7/2) –	(1 st forbidden non-unique)	7.3	0.8805 (15)	0.1005 (12)	0.0173 (4)	0.0017 (1)
EC _{0,2}	2297 (7)	1275 (7)	555 (3)	97.24 (20)	0.522	33.3 (5)	63.9 (5)	9/2 +	9/2 +	allowed	5.36	0.8810 (15)	0.1001 (12)	0.0172 (4)	0.0017 (1)
EC _{0,1}	2658 (7)	1636 (7)	745 (3)	1.2 (2)	0.741	0.51 (9)	0.69 (11)	9/2 +	5/2 –	1 st forbidden unique	8.7 ^{lu}	0.8811 (15)	0.1000 (12)	0.0172 (4)	0.0017 (1)
				Σ 100.0479		Σ 35.3806	Σ 64.6277								
						Σ 100.0083									

* Determined from the nuclear level energies of 2004Si08 and Q-value of 2725(7) keV (2012Wa38).

Atomic Data

The x-ray and Auger-electron data have been calculated using the evaluated gamma-ray data, and atomic data from 1996Sc06, 1998ScZM and 1999ScZX. Both the x-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.02, 28 February 2012), as described in 2000Sc47. This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

The inability to obtain a sound and satisfactory balance with respect to the population-depopulation of the 67.039-keV nuclear level of ⁷³As could have arisen as a consequence of either a poorly quantified emission probability for the 67.039-keV gamma ray when identified to be a 100 % M1 transition, or a suitably well-defined emission probability for the 67.039-keV gamma ray with inappropriate ICC data. A weighted-mean absolute emission probability of 71 (7) % has been adopted for the 67.039-keV gamma ray with a multipolarity of 100 % M1 rather than (95.7 % + 4.3 % E2). Such a difference impacts on the X-ray and Auger-electron data as shown below:

K and L X-ray energies and emission probabilities of ⁷³Se.

Energy (keV)			Multipolarity of 67.039-keV gamma transition			
			100 % M1		(95.7 % M1 + 4.3 % E2) XX	
			Photons per 100 disint.	Relative probability	Photons per 100 disint.	Relative probability
XL	(As)	1.120 – 1.524	1.05 (3)	6.48	1.24 (4)	6.56
	XL ₁	(As) 1.120	0.0270 (10)		0.0316 (14)	
	XL _α	(As) 1.282	0.613 (21)		0.72 (3)	
	XL _η	(As) 1.155	0.0152 (7)		0.0178 (9)	
	XL _β	(As) 1.317 – 1.388	0.397 (17)		0.466 (23)	
	XL _γ	(As) 1.524	0.00230 (9)		0.00268 (11)	
XK _α	XK _{α2}	(As) 10.50814 (9)	8.3 (3)	51.2	9.7 (5)	51.3
	XK _{α1}	(As) 10.54380 (9)	16.2 (6)	100	18.9 (9)	100
XK' _{β1}	XK _{β3}	(As) 11.7204 (5)	}		}	
	XK _{β1}	(As) 11.7263 (2)	} 3.70 (14)	22.8	} 4.31 (21)	22.8
	XK _{β5}	(As) 11.821 (1)	}		}	
XK' _{β2}	XK _{β2}	(As) 11.8643 (3)	0.140 (7)	0.86	0.163 (10)	0.86

Auger-electron energies and emission probabilities of ⁷³Se.

Energy (keV)			Multipolarity of 67.039-keV gamma transition			
			100 % M1		(95.7 % M1 + 4.3 % E2)	
			Electrons per 100 disint.	Relative probability	Electrons per 100 disint.	Relative probability
e _{AK}	(As)		21.0 (8)		24.4 (12)	
	KLL	8.746 – 9.149	15.7 (6)	100	18.3 (9)	100
	KLX	10.114 – 10.541	4.91 (19)	31.3	5.7 (3)	31.1
	KXY	11.460 – 11.862	0.384 (17)	2.45	0.447 (24)	2.44
e _{AL}	(As)	0.90 – 1.23	65.3 (15)	416	76.5 (22)	418

As: ω_K = 0.575 (4); ω_L = 0.0155 (5); n_{KL} = 1.232 (4) were taken from 1996Sc06.

Electron energies were determined from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

Data Consistency

A Q_{EC} -value of 2725 (7) keV has been adopted from the atomic mass evaluation of Wang *et al.* (2012Wa38) while in the course of formulating the decay scheme of ⁷³Se. This value has subsequently been compared with the Q-value calculated by summing the contributions of the individual emissions to the ⁷³Se EC-decay process (i.e. EC/ β^+ , electron, γ , etc.):

$$\text{calculated Q-value} = \sum (E_i \times P_i) = 2718 (16) \text{ keV}$$

Percentage deviation from the Q-value of Wang *et al.* is 0.3 (6) %, which supports the derivation of a reasonably consistent decay scheme with a large variant despite known inadequacies identified with the gamma depopulation of the first excited state of ⁷³As (i.e. 67.039-keV nuclear level).

References

- 1948Co07 W.S. Cowart, M.L. Pool, D.A. McCown, L.L. Woodward, Artificially Radioactive Se⁷³ and Se⁷⁵, Phys. Rev. 73 (1948) 1454-1457. [Half-life]
- 1951Sc70 F.R. Scott, The Radioactivity of Se⁷³, Phys. Rev. 84 (1951) 659-665. [E β^+ , P β^+ , E γ , P γ , E_{ce}, P_{ce}, E_{Ae}, P_{Ae}]
- 1955Ha85 R.W. Hayward, D.D. Hoppes, The Disintegration of Se-73, Phys. Rev. 98 (1955) 1172, MA2. [E β^+ , log *ft*]
- 1956Ha10 R.W. Hayward, D.D. Hoppes, The Disintegration of Se-73, Phys. Rev. 101 (1956) 93-97. [E β^+ , P β^+ , E γ , P γ , E_{ce}, P_{ce}, Half-lives (67- and 428-keV nuclear levels)]
- 1957Be46 J. Beydon, R. Chaminade, M. Crut, H. Faraggi, J. Olkowsky, A. Papineau, Étude de la Transmutation du Cuivre par L'Azote et l'Oxygène, Nucl. Phys. 2 (1956/57) 593-618. [Half-life, E γ]
- 1957Ku57 M.Ia. Kuznetsova, V.N. Mekhedov, Procedure for Measuring the Activity of Nuclei Undergoing K-capture, Izvest. Akad. Nauk SSSR, Ser. Fiz. 21 (1957) 1020-1024; Columbia Tech. Trans. 21 (1958) 1021-1025. [K capture]
- 1960Ku06 T. Kuroyanagi, Decays of Photoreaction Products of Selenium, J. Phys. Soc. Japan 15 (1960) 2179-2187. [β^+ decay]
- 1960Ri09 R.A. Ricci, R. van Lieshout, H.J. van den Bold, Scintillation Spectrometry Study of the ⁷³Se Isomeric Pair, Physica 26 (1960) 1014-1020. [Half-life, E γ , P γ]
- 1963Bo16 H.H. Bolotin, Double Isomerism in As⁷³, Phys. Rev. 131 (1963) 774-777. [Half-life (67- and 428-keV nuclear levels)]
- 1963Bo26 E. Bodenstein, G. Strube, W. Engels, H. Blumberg, R.-M. Lieder, E. Gerdau, Phys. Lett. 6 (1963) 290-291. [Half-life (67-keV nuclear level), δ]
- 1966Ol04 B. Olsen, L. Boström, Numerical Analysis of the Time Spectrum of Delayed Coincidences, II, Nucl. Instrum. Methods. 44 (1966) 65-72. [Half-life (67-keV nuclear level)]
- 1967Ra08 P.V. Rao, R.W. Fink, Gamma Decay of Se^{73m,g} and Se^{81m,g} Isomeric Pairs, Phys. Rev. 154 (1967) 1028-1032. [Half-life, E γ , P γ , α_{tot}]
- 1968At04 A.H.W. Aten Jr., J.C. Kapteyn, Decay of ^{73m}Se, ⁷¹Se and ⁷⁰Se, Radiochim. Acta 9 (1968) 48-49. [67- and 361-keV gamma rays, ^{73m}Se]

- 1968Ak03 M.P. Akhmed, K.A. Baskova, S.S. Vasil'ev, L.Ya. Shavtvalov, Study of the Radiation of ${}_{34}\text{Se}^{73}$, *Yad. Fiz.* 8 (1968) 240-246; *Sov. J. Nucl. Phys.* 8 (1969) 138-141.
[E_{β+}, P_{β+}, log *ft*, P_γ]
- 1968Iv02 E.A. Ivanov, A. Iordăchescu, G. Pascovici, Nuclear Isomers with Half-lives in the 2-30 μs Range Excited by Pulsed-beam Proton Irradiations, *Rev. Roum. Phys.* 13 (1968) 879-885.
[Half-life (428-keV nuclear level)]
- 1969Iv02 E.A. Ivanov, A. Iordăchescu, G. Pascovici, New Nuclear Isomers in the Microsecond Region, *Rev. Roum. Phys.* 14 (1969) 317-319. [Half-life (428-keV nuclear level)]
- 1969Ma21 K.W. Marlow, A. Faas, The Radioactive Decay of ${}^{73g}\text{Se}$ and ${}^{73m}\text{Se}$, *Nucl. Phys.* A132 (1969) 339-352. [Half-life, E_γ, P_γ, Half-life (428-keV nuclear level)]
- 1970Me20 R.D. Meeker, A.B. Tucker, Decay of ${}^{73m,g}\text{Se}$, ${}^{75m}\text{Ge}$ and ${}^{77m}\text{Ge}$, *Nucl. Phys.* A157 (1970) 337-357. [E_γ, P_γ]
- 1970MeZZ R.D. Meeker, Decay of ${}^{73m,g}\text{Se}$, ${}^{75m}\text{Ge}$ and ${}^{77m,g}\text{Ge}$, PhD thesis, Iowa State University (1970). [E_γ, P_γ]
- 1970Qu03 D. Quitmann, J.M. Jaklevic, Measurement of the Magnetic Moments of the Microsecond-isomers in ${}^{73}\text{As}$ and ${}^{206}\text{Pb}$, *Z. Naturforsch.* 25a (1970) 975-976. [A₂, A₄, Multipolarity]
- 1971Go40 N.B. Gove, M.J. Martin, Log-*f* Tables for Beta Decay, *Nucl. Data Tables* 10 (1971) 205-237. [e/β⁺ ratios]
- 1972ReZN E. Recknagel, Perturbed Angular Distribution Following Nuclear Reactions, Summer School on Interaction of Radiation with Matter, July 1971, Predeal, Romania, Hahn-Meitner Institute report HMI-B-115 (1972). [Half-life (428-keV nuclear level)]
- 1974Be54 R.R. Betts, S. Mordechai, D.J. Pullen, B. Rosner, W. Scholz, Study of ${}^{71,73,75,77}\text{As}$ with Ge(³He,d) Reactions, *Nucl. Phys.* A230 (1974) 235-252. [⁷³As nuclear levels]
- 1975Va03 P. van der Merwe, E. Barnard, J.A.M. de Villiers, J.G. Malan, An Investigation of ${}^{73}\text{As}$ Energy Levels by Means of (p,n) and (p,nγ) Reactions, *Nucl. Phys.* A240 (1975) 273-292. [⁷³As nuclear levels from ${}^{73}\text{Ge}(p,n)$ and (p,nγ) reactions]
- 1976Bo19 M. Bormann, H.-K. Feddersen, H.-H. Hölscher, W. Scobel, H. Wagener, (n,2n) Anregungsfunktionen für ${}^{54}\text{Fe}$, ${}^{70}\text{Ge}$, ${}^{74}\text{Se}$, ${}^{85}\text{Rb}$, ${}^{86,88}\text{Sr}$, ${}^{89}\text{Y}$, ${}^{92}\text{Mo}$, ${}^{204}\text{Hg}$ im Neutronenenergiebereich 13-18 MeV, *Z. Phys.* A277 (1976) 203-210. [Half-life]
- 1976Sc13 M. Schrader, H. Reiss, G. Rosner, H.V. Klapdor, Investigation of the Level Schemes of ${}^{73,75,77}\text{As}$ via the (³He,d) Reaction, *Nucl. Phys.* A263 (1976) 193-209. [⁷³As nuclear levels]
- 1977KeZY T.J. Ketel, In-beam Measurements on Nuclear Magnetic Dipole Moments of Isomeric States in ${}^{73}\text{As}$, ${}^{112,114,115,117}\text{Sb}$ and ${}^{139}\text{Pr}$, PhD thesis, Vrije Universiteit, Amsterdam (1977). [Half-life (428-keV nuclear level)]
- 1977La19 F.P. Larkins, Semiempirical Auger-electron Energies for Elements 10 ≤ Z ≤ 100, *At. Data Nucl. Data Tables* 20 (1977) 311-387. [Auger-electron energies]
- 1978TeZY B.O. ten Brink, Properties of the Nuclei ${}^{69}\text{Ge}$, ${}^{69,70,71,72,73}\text{As}$ Investigated by Gamma Spectroscopic Methods, PhD thesis, Vrije Universiteit, Amsterdam (1978). [E_γ, P_γ]
- 1980Te01 B.O. ten Brink, P. van Nes, C. Hoetmer, H. Verheul, On the Structure of ${}^{71}\text{As}$ and ${}^{73}\text{As}$, *Nucl. Phys.* A338 (1980) 24-44. [E_γ, P_γ]

- 1988Be39 I. Berkes, R. Hassani, M. Massaq, Ground State Spin and Magnetic Moment of ⁷³Se, Phys. Rev. C38 (1988) 2329-2331. [Ground state spin, magnetic moment, δ]
- 1992Sc21 Th. Schaefer, E. Lohmann, R. Vianden, The Quadrupole Moment of the 66-keV $5/2^-$ State of ⁷³As, Z. Phys. – Hadrons and Nuclei A343 (1992) 279-281. [Quadrupole moment (67-keV nuclear level)]
- 1995ScZY E. Schönfeld, Tables for the Calculation of Electron Capture Probabilities, PTB Report PTB-6.33-95-2, 1995. [P_K, P_L, P_M, P_N, P_O]
- 1996Sc06 E. Schönfeld, H. Janßen, Evaluation of Atomic Shell Data, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527-533. [$\omega_K, \omega_L, K_\beta/K_\alpha, K_{\alpha 2}/K_{\alpha 1}, KLX/KLL, KXY/KLL$]
- 1997So08 D. Sohler, Zs. Podolyák, J. Gulyás, T. Fényes, A. Algora, Zs. Dombrádi, S. Brant, V. Paar, Structure of the ⁷³As Nucleus, Nucl. Phys. A618 (1997) 35-54. [⁷³As nuclear levels from ⁷³Ge(p,n γ) reaction, γ branches]
- 1998Sc28 E. Schönfeld, Calculation of Fractional Electron Capture Probabilities, Appl. Radiat. Isot. 49 (1998) 1353-1357. [P_K, P_L, P_M, P_N, P_O]
- 1998ScZM E. Schönfeld, G. Rodloff, Tables of the Energies of K-Auger Electrons for Elements with Atomic Numbers in the Range from Z = 11 to Z = 100, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX E. Schönfeld, G. Rodloff, Energies and Relative Emission Probabilities of K X-rays for Elements with Atomic Numbers in the Range from Z = 5 to Z = 100, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2000Sc47 E. Schönfeld, H. Janßen, Calculation of Emission Probabilities of X-rays and Auger Electrons Emitted in Radioactive Disintegration Processes, Appl. Radiat. Isot. 52 (2000) 595-600. [P_X, P_{Ae}]
- 2002Ba85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, Dirac–Fock Internal Conversion Coefficients, At. Data Nucl. Data Tables 81 (2002) 1-334. [ICC]
- 2002Ra45 S. Raman, C.W. Nestor, Jr., A. Ichihara, M.B. Trzhaskovskaya, How Good are the Internal Conversion Coefficients Now? Phys. Rev. C66 (2002) 044312, 1-23. [ICC]
- 2004Si08 Balraj Singh, Nuclear Data Sheets for A = 73, Nucl. Data Sheets 101 (2004) 193-323. [Nuclear levels]
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr., Evaluation of Theoretical Conversion Coefficients using BrIcc, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202-229. [ICC]
- 2012Wa38 M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer, The AME2012 Atomic Mass Evaluation, (I). Tables, Graphs and References, Chin. Phys. C36 (2012) 1603-2014. [Q-value]

⁷⁵Se - Comments on evaluation of decay data by V. Chisté and M. M. Bé

1 Decay Scheme

⁷⁵Se disintegrates 100 % by electron capture to excited levels and to the ground state of ⁷⁵As.

A good agreement was found between the effective Q value (860 (18) keV) calculated from the decay scheme data and the adopted and recommended value from the mass adjustment of Audi (2003Au03).

2 Nuclear Data

The Q value is from Audi and Wapstra (2003Au03).

Level energies, spins and parities are from the mass-chain evaluation of A. R. Farhan and B. Singh (1999Fa05).

Experimental ⁷⁵Se half-life values (in days) are given in Table 1:

Table 1: Experimental values of ⁷⁵Se half-life.

Reference	Experimental value (d)	Comments
H. N. Friedlander (1947Fr08)	115 (5)	Outlier
W. S. Cowart (1948Co07)	127 (7)	Outlier
J. M. Cork (1950Co58)	128	Not used: no uncertainty.
H. M. Wright (1957Wr37)	119.9 (6)	Outlier
H. T. Easterday (1960Ea02)	120.4 (2)	Outlier
F. Lagoutine (1975La16)	118.45 (25)	Outlier
M. J. Martin (1976MaZW)	120 (1)	Outlier
H. Houtermans (1980Ho17)	119.779 (4)	
U. Schötzig (1980Sc07)	119.76 (5)	
D. D. Hoppes (1982HoZJ)	119.80 (7)	
A. Iwahara (1994Iw04)	119.0 (5)	Outlier
M. He (2002He19)	115.0 (117)	Not used
M. P. Unterweger (2002Un02)	119.809 (66)	
Recommended value	119.781 (24)	$\chi^2 = 0.14$

The value in 2002He19 was omitted because this value is just a verification of how good their experimental set-up was. The first 6 values (1947Fr08, 1948Co07, 1957Wr37, 1960Ea02, 1975La16, 1976MaZW) and the Iwahara value (1994Iw04) have been shown to be outliers, based on the Chauvenet's criterion and thus were omitted in the final calculation. With the 4 remaining values (1980Ho17, 1980Sc07, 1982HoZJ and 2002Un02), a weighted average was calculated using the LWEIGHT computer code (version 3). The largest contribution to the weighted average comes from the value of Houtermans (1980Ho17), amounting to 63 %. The LWEIGHT increases the uncertainty of 1980Ho17 value from 0.004 to 0.034 in order to reduce its relative weight from 63 % to 50 %.

The adopted value is the weighted average of 119.781 d with an internal uncertainty of 0.024 d. The reduced- χ^2 value is 0.14.

2.1 Electron Capture Transitions

The energies of the electron capture transitions have been obtained from the Q(EC) value (2003Au03) and the level energies given by A. R. Farhan (1999Fa05).

The adopted electron capture transition probabilities and associated uncertainties were deduced from the γ transition probability balance at each level in the decay scheme.

The partial sub-shell capture probabilities given in this section were calculated using the computer program EC-Capture.

2.2 γ Transitions

The γ transition probabilities were deduced using the γ -ray emission intensities and the relevant internal conversion coefficients (see **5.2 Gamma Emissions**).

Table 2 shows the multiplicities (no mixing ratios given) of γ transitions, deduced from the conversion coefficient (1999Fa05) analysis.

Table 2: Multiplicities of γ transitions.

Multiplicity	E_γ (keV)
E1	121.1155 (11), 136.0001 (6), 400.6572 (8),
[E2]	80.9365 (15), 373.61 (24), 556.90 (18), 821.56 (18)
E2	96.7340 (9)
E3	303.9236 (10)
[M1, E2]	249.3 (3), 419.1 (4), 468.6 (4), 542.02 (18), 617.8 (4)
M1 (+E2)	14.8847 (13)
M2 + (E3)	24.3815 (14)

For (M1 + E2) γ transitions (66-, 198-, 264-, 279- and 572-keV), the mixing ratios (δ) are given in Table 5, they were deduced by comparison between experimental values of K internal conversion coefficients and the theoretical values calculated using the BrIcc computer code (2008Ki07).

Since the γ transitions with $E_\gamma = 121$ - and 136-keV were determined to be pure E1, their α_K coefficients can be interpolated from theoretical values and then used to deduce the α_K coefficient of the 264-keV γ -ray which has been used as the reference line in all the measurements. Then:

$$\alpha_k(264) = \frac{\alpha_k(136) \times I_\gamma(136)}{I_{CEK}(136)}$$

and

$$\alpha_k(264) = \frac{\alpha_k(121) \times I_\gamma(121)}{I_{CEK}(121)} \quad (\text{see Table 4})$$

where:

- I_{CEK} is the weighted average of the experimental values of the relative conversion electron intensities shown in Table 3a (2nd and 3rd columns, respectively);
- I_γ is the weighted average of the experimental values of the relative γ -ray emission intensities given in Table 8 (see **5.2 Gamma Emission**).

Table 3a: Experimental and recommended values of relative conversion electron intensities (I_{CEK}) and photon (I_γ) intensities for (M1 + E2) γ -rays.

Energy (keV) Reference	121	136	66	198	264	279	572
1955Sc09	173 (14)	420 (34)	68 (10)	6.4 (9)	100	53 (7)	
1959Me76		377 (20)			100	53.6 (16)	
1960De06	180 (12)	450 (30)	80 (12)	6.8 (10)	100	63 (5)	
1960Gr03	154 (5)	384 (13)	73.7 (49)	7.30 (37)	100	49.2 (33)	0.055 (22)
1961Ed02	187 (15)	378 (30)	99 (12)	7 (1)	100	53 (5)	
1965Br19	167 (26)	520 (70)		7.0 (12)	100	52 (7)	
1970Pa25	174 (17)	399 (32)	72.3 (10)	7.36 (41)	100	52.5 (23)	0.0099 (9)
2005Ra29	169.91 (27)	377.94 (41)	88.47 (20)	6.41 (5)	100	42.93 (22)	0.0103 (34)
Recommended I_{CEK}	169.88 (42)	377.95 (41)	81 (8)	6.44 (7)	100	53.2 (12)	0.0100 (13)
χ^2	2.4	0.08	7.5	1.9		0.05	2.1
Recommended I_γ	28.7 (6)	97.8 (34)	1.792 (34)	2.48 (10)	100	42.36 (6)	0.06165 (49)
χ^2	4.16	5.08	6.07	4.43		0.51	1.43

Table 3b: Experimental and recommended values of relative conversion electron intensities I_{CEK} and photon (I_γ) intensities for additional (M1 + E2) γ -rays.

Energy (keV) Reference	24	80	96	303	400	419	556	617
1955Sc09		~ 8.1 (7)*	~ 720 (60)*	15.6 (13)	3.6 (5)			
1959Me76				15.4 (9)	3.6 (4)			
1960De06		14 (3)	940 (60)	16 (1)	3.8 (3)			
1960Gr03	1250 (150)	7 (1)	645 (32)	16.1 (9)	3.76 (28)			
1961Ed02			753 (60)	17.2 (17)	3.8 (3)			
1965Br19			750 (110)	17.0 (26)				
1970Pa25			724 (19)	16.6 (5)	3.71 (4)	0.006 6 (7)		0.000 85 (9)
2005Ra29	1010 (1)	4.04 (4)	502 (1)	16.4 (8)	3.98 (4)	0.012 0 (4)	0.009 5 (32)	0.001 1 (4)
Recommended I_{CEK}	1010.0 (16)	5.9 (18)	610 (110)	16.16 (29)	3.84 (13)	0.006 8 (9)	0.009 5 (32)	0.000 86 (9)
χ^2	2.6	6.2	12	0.45	3.9	1.8		0.37
Recommended I_γ	0.046 (11)	0.0161 (9)	5.71 (12)	2.2267 (44)	19.384 (36)	0.020 6 (11)	0.004 7 (2)	0.007 71 (9)
χ^2	4.56	1.13	13.53	0.80	1.5	5.03		0.179

* Not used

Table 4: Determination of α_{K264} .

Energy (keV)	I_{CEK} (%)	I_γ (%)	α_K (by BrIcc)	α_{K264}
121	169.88 (42)	28.7 (6)	0.037 2 (6)	0.006 28 (17)
136	377.95 (41)	97.8 (34)	0.026 3 (4)	0.006 81 (26)
			Adopted	0.006 44 (24)

The adopted α_K coefficient for the 264-keV γ transition is 0.006 44, weighted average with an external uncertainty of 0.000 24 and a reduced- χ^2 of 2.87.

Table 5 shows the final results of experimental α_K (deduced using $\alpha_{K264} = 0.006 44 (24)$), together with mixing ratios δ (deduced from a comparison between experimental (column 2) and theoretical (column 5, calculated with the BrIcc computer code) α_K values.

Table 5: Recommended conversion coefficients and mixing ratios.

E_γ (keV)	I_{CEK}/I_γ	α_K experimental (= $(\alpha_{K264} * I_{\text{CEK}}/I_\gamma)$)	δ (mixing ratio)	α_K theoretical (given by BrIcc)	Multipolarities
24	22 (5) 10 ³	141 (34)		165.4 (24)	M2 + (E3)
66	45.2 (45)	0.291 (31)	0.121 (33)	0.29 (3)	M1 + E2
80	370 (110)	2.4 (7)		1.486 (21)	[E2]
96	107 (19)	0.69 (13)		0.772 (11)	E2
198	2.60 (11)	0.016 7 (9)	0.315 (39)	0.016 7 (9)	M1 + E2
264	1	0.006 44 (24)	-0.10 (7)	0.006 46 (25)	M1 + E2
279	1.256 (28)	0.008 09 (35)	-0.578 (44)	0.008 1 (4)	M1 + E2
303	7.26 (13)	0.046 7 (19)		0.046 9 (7)	E3
400	0.198 (7)	0.001 28 (6)		0.001 202 (17)	E1
419	0.330 (47)	0.002 13 (31)		0.003 0 (10)	[M1,E2]
556	2.0 (7)	0.013 0 (44)		0.001 628 (25)	[E2]
572	0.162 (21)	0.001 04 (14)	0.19 (1)	0.001 04 (1)	M1 + E2
617	0.112 (12)	0.000 72 (8)		0.001 03 (18)	[M1,E2]

Then, for all γ transitions, the adopted detailed and total internal conversion coefficients (ICC) and associated uncertainties have been obtained using the BrIcc computer program with “the frozen orbital approximation” (2008Ki07).

3 Atomic Data

Atomic values, ω_K , ω_L and n_{KL} and X-ray and Auger electron relative probabilities are from Schönfeld and Janßen (1996Sc06).

4 Electron emissions

The conversion electron emission probabilities were deduced from ICC values and γ -ray emission probabilities.

5 Photon Emissions

5.1 X-rays

The X-ray absolute intensities were deduced from the decay data using the EMISSION computer code and are compared in Table 6 with measured values found in the literature. A good agreement has been found between the experimental and our values deduced from decay scheme.

Table 6: Experimental and recommended (calculated) values of X-ray absolute intensities.

	1966Ra09*	1970Pa25*	1974Ca29*	1983Si25*	1992Sc09*	1996Sa22	2000Zhan ^g	Recommended
K x-ray	55.5 (14)	53.1 (15)	53.5 (29)	57.5 (13)	54.7 (11)	58.3 (14)	55.6 (12)	56.0 (13)
K α x-ray				49.2 (13)	47.6 (11)		48.3 (10)	48.4 (13)
K β x-ray				8.25 (24)	7.13 (17)		7.3 (2)	7.58 (25)

*Using normalization factor of 0.5875 (19) (see Table 7, **5.2 Gamma Emissions**)

5.2 Gamma emissions

The energies of the γ -rays given in section 5.2 are from A. F. Farhan (1999Fa05).

The experimental relative γ -ray emission intensities from ⁷⁵Se have been obtained from all the available relative and absolute values. The normalization factor to convert relative γ -ray emission probabilities to absolute values is from a weighted average of measured absolute values for the 264-keV γ -ray absolute intensity.

Table 7: Experimental 264-keV absolute gamma-ray emission intensities.

References	Experimental values
Y. Yoshizawa (1983Yo03)	0.580 (9)
U. Schötzig (1992Sc09)	0.5950 (54)
H. Miyahara (1994Mi22)	0.5870 (17)
Recommended value	0.5875 (19), $\chi^2 = 1.35$

The experimental γ -ray emission probabilities relative to 100 for the 264-keV γ -ray are given in Table 8.

Our recommended relative and absolute γ -ray emission probabilities are given in Table 9.

The adopted values are the weighted means calculated by the LWEIGHT program (version 3) with the following restrictions:

*: Discrepant data set, omitted from analysis.

@: data set already taken into account in 1987JeZZ, then these references have been omitted from the analysis.

μ : the experimental value has been shown to be an outlier value by the Lweight program.

Table 8: Experimental data sets of the relative γ -ray emission intensities (%) (cont. next page).

Energy (keV)	14	24	66	80	96	121	136 198 249 264			
Reference										
1955Sc09			1.8 (1)		6.6 (15)	28 (5)	94 (12)		100	
1958Va02			2.1 (8)		5.8 (6)	24.5 (30)	76 (5)	3.6 (4)	100	
1959Vo30									100	
1960De06						28 (5)	86 (15)		100	
1960Gr03			1.53 (15)		5.5 (3)	27.9 (13)	96 (5)	2.6 (2)	100	
1961Ed02			1.63 (6)		5.57 (18)	28.0 (6)	95.5 (18)	2.4 (1)	100	
1965Br19						30 (10)	130 (40)		100	
1966Ra09			1.64 (5)		5.53 (16)	27.8 (8)	94.9 (20)	2.28 (5)	100	
1969Ra12			1.4 (4)		4.83 (10)	29.2 (29)	96.0 (96)	2.25 (23)	100	
1970Pa25		0.044 (6)	1.72 (4)	0.015 (3)	5.12 (10)	27.7 (5)	95.0 (18)	2.38 (7)	100	
1970Na14			1.54 (8)		5.43 (16)	28.5 (9)	94.0 (28)	2.78 (14)	100	
1971Ge07			1.77 (1)		5.6 (5)	28.2 (14)	98.3 (46)	2.43 (12)	100	
1971Pr07		0.032 (10)								
1973Su10*			0.97 (6)		4.7 (2)	25.4 (12)	90.3 (28)	2.5 (1)	100	
1973Te06			2.0 (5)		5.0 (5)	25.8 (25)	94.6 (82)	2.2 (2)	100	
1973Th07	0.034 (6)	0.063 (8)	1.50 (15)	0.011 (3)	5.4 (4)	26.7 (30)	95.7 (70)	2.59 (20)	100	
1974Ca29		0.036 (4)							100	
1977Ge12			1.86 (11)		5.90 (35)	29.8 (13)	102.0 (30)	2.53 (11)	100	
1978Pr08		0.065 (8)	1.46 (20)	0.012 (4)	5.22 (20)	27.1 (40)	95.5 (60)	2.5 (4)	100	
1983Yo03					5.78 (17)	29.24 (32)	99.2 (10)	2.51 (4)	100	
1984Si06		0.052 (9)	1.91 (3)	0.014 (4)		29.96 (26)	102.5 (10)	2.52 (6)	100	
1987JeZZ - n°1		0.045 (6)	1.850 (31)		5.93 (8)	29.23 (19)	99.9 (5)	2.518 (16)	100	
1987JeZZ - n°2		0.127 (12) ^u	1.82 (7)		5.68 (19)	29.1 (9)	96.3 (28)	2.52 (9)	100	
1987JeZZ - n°3			1.76 (9)		6.13 (22)	27.9 (9)	94.6 (30)	2.25 (9)	100	
1987JeZZ - n°4			1.95 (6)		6.47 (19)	29.2 (5)	99.9 (14)	2.568 (35)	100	
1987JeZZ - n°5						29.3 (7)	99.9 (13)	2.48 (6)	100	
1987JeZZ - n°6			1.78 (7)		5.41 (16)	28.5 (7)	95.9 (27)	2.38 (6)	100	
1987JeZZ - n°7			2.00 (18)		5.13 (33)	30.0 (13)	99.5 (40)	2.53 (10)	100	
1987JeZZ - n°8			1.860 (22)		5.790 (42)	28.65 (18)	98.2 (6)	2.509 (20)	100	
1987JeZZ - n°9			1.960 (41)		5.63 (5)	28.96 (18)	99.9 (6)	2.581 (16)	100	
1987JeZZ - n°10		0.0446 (20)	1.910 (25)		5.91 (7)	29.16 (33)	99.7 (11)	2.534 (28)	100	
1987JeZZ - n°11			1.940 (34)		5.88 (8)	29.43 (32)	100.4 (11)	2.514 (28)	100	
1987JeZZ - n°12			1.88 (1)		5.830 (22)	29.31 (11)	101.2 (3)	2.586 (11)	100	
1987JeZZ - n°13			1.950 (24)		5.91 (6)	29.24 (29)	99.4 (12)	2.500 (35)	100	
1990An07 [@]			1.962 (29)		5.93 (9)	29.24 (41)	100.0 (17)	2.50 (5)	100	
1990Me15		0.0460 (46)	1.87 (9)	0.0190 (41)	5.71 (35)	29.8 (15)	100 (6)	2.54 (24)	100	
1990Wa09 [@]			1.960 (49)		5.91 (12)	29.1 (6)	99.5 (20)	2.50 (6)	100	
1992Sc09 [@]		0.0446 (20)	1.910 (25)		5.91 (7)	29.16 (33)	99.7 (11)	2.534 (28)	100	
1994Bh07*	0.003 (2)	0.056 (6)	1.912 (3)	0.013 (4)		30.1 (9)	102.3 (11)	2.51 (8)	100	
1994Mi22					5.779 (45)	29.76 (19)	100.2 (6)	2.555 (20)	100	
1997Lo10						28.05 (27)	98.41 (36)	2.58 (7)	100	
2005Ra29	0.035 (1)	0.035 (1)	1.79 (1)	0.017 (1)	5.10 (4)	27.40 (22)	94.1 (8)	2.42 (2)	0.0067 (2)	100
Evaluated	0.035 (1)	0.046 (11)	1.792 (34)	0.0161 (9)	5.71 (12)	28.7 (6)	97.8 (34)	2.48 (10)	0.0067 (2)	100
χ^2	0.027	4.56	6.07	1.13	13.53	4.16	5.08	4.43		

Energy (keV)	279	303	373	400 418		468	542 557		572	617 821	
Reference											
1955Sc09	45.7 (40)	2.0 (5)		24.8 (25)							
1958Va02	52 (5)			28 (2)							
1959Vo30	44.1 (44)	3.2 (12)		22.7 (15)					0.068 (46)		
1960De06	42.5 (20)	2.15 (30)		23 (2)							
1960Gr03	41.0 (25)	2.5 (3)		22.3 (23)					0.18 (6)		
1961Ed02	42.2 (6)	2.29 (14)		19.5 (6)							
1965Br19	53 (15)										
1966Ra09	43.0 (9)	2.39 (5)		22.3 (5)	0.0322 (6)				0.0636 (13)	0.00777 (15)	
1969Ra12	41.3 (41)	2.06 (21)		19.2 (19)	0.020 (3)				0.053 (8)	0.0076 (10)	
1970Pa25	42.0 (8)	2.19 (7)		20.4 (5)	0.023 (2)	0.0010 (5)			0.063 (2)	0.0075 (2)	
1970Na14	41.9 (13)	2.20 (11)		19.5 (6)							
1971Ge07	43.2 (22)	2.31 (12)		19.6 (12)							
1971Pr07						0.00054 (18)					0.000216 (10)
1973Su10*	42.5 (15)	2.20 (8)		19.0 (6)	0.0140 (16)				0.054 (3)	0.0075 (31)	
1973Te06	40.0 (22)			19.6 (7)							
1973Th07	42.1 (8)	2.11 (30)		18.0 (4)	0.017 (3)				0.048 (5)	0.059 (7)	
1974Ca29											
1977Ge12	42.4 (18)	2.21 (7)		19.1 (6)							
1978Pr08	42.6 (8)	2.3 (4)	0.0042 (4)	18.8 (6)	0.018 (4)	0.00062 (10)	0.00022 (4) ^u	0.00006 (2) ^u	0.050 (4)	0.0062 (8)	0.00028 (2)
1983Yo03	42.43 (29)	2.234 (20)		19.42 (16)	0.0231 (21)				0.0634 (29)	0.0078 (21)	
1984Si06	42.4 (4)										
1987JeZZ - n°1	42.53 (23)	2.248 (13)		19.27 (13)	0.0206 (7)				0.0602 (20)	0.0072 (7)	
1987JeZZ - n°2	43.9 (13)	2.25 (7)		19.7 (6)	0.024 (9)				0.0625 (26)	0.0067 (10)	0.0016 (12) ^u
1987JeZZ - n°3	42.2 (13)	2.21 (8)		19.1 (6)							
1987JeZZ - n°4	42.6 (6)	2.091 (27)		19.41 (24)							
1987JeZZ - n°5	42.6 (9)	2.24 (5)		19.50 (42)							
1987JeZZ - n°6	42.4 (9)	2.23 (6)		19.17 (39)	0.0102 (32)				0.0580 (41)	0.0076 (6)	0.00030 (15)
1987JeZZ - n°7	42.6 (16)	2.24 (8)		19.5 (7)	0.0154 (11)				0.0590 (34)	0.0080 (6)	0.0013 (7) ^u
1987JeZZ - n°8	42.48 (31)	2.234 (19)		19.60 (14)					0.0610 (18)		
1987JeZZ - n°9	42.36 (22)	2.224 (12)		19.79 (10)					0.0617 (14)	0.0063 (18)	
1987JeZZ - n°10	42.5 (5)	2.242 (25)		19.49 (22)	0.0196 (12)				0.0610 (11)	0.0078 (5)	
1987JeZZ - n°11	42.4 (5)	2.220 (27)		19.08 (23)	0.0217 (5)				0.0603 (9)	0.0077 (3)	
1987JeZZ - n°12	42.25 (7)	2.219 (9)		19.36 (4)	0.0247 (13)				0.067 (2)	0.0108 (12)	
1987JeZZ - n°13	42.69 (37)	2.239 (22)		19.51 (17)					0.064 (3)		
1990An07[@]	42.7 (5)	2.238 (31)		19.51 (24)					0.064 (5)		
1990Me15	42.2 (21)	2.23 (11)		19.5 (10)	0.0180 (31)				0.0600 (42)	0.0077 (6)	0.000220 (23)
1990Wa09[@]	42.4 (9)	2.25 (5)		20.19 (43)	0.0209 (5)				0.0589 (12)	0.0076 (2)	
1992Sc09[@]	42.5 (5)	2.242 (25)		19.49 (22)	0.0196 (12)				0.0610 (11)	0.0078 (5)	
1994Bh07*	42.55 (10)										
1994Mi22	42.78 (25)	2.239 (18)		19.31 (12)							
1997Lo10	43.63 (29)	2.199 (11)		18.84 (16)					0.066 (3)		
2005Ra29	43.07 (34)	2.27 (2)	0.0044 (2)	20.13 (16)	0.035 (1)	0.0036 (2) ^u	0.00074 (1)	0.0047 (2)	0.062 (1)	0.0078 (2)	0.0015 (2) ^u
Evaluated	42.36 (6)	2.2267 (44)	0.00436 (18)	19.384 (36)	0.0206 (11)	0.00061 (9)	0.00074 (1)	0.0047 (2)	0.06165 (49)	0.00771 (9)	0.00028 (14)
χ^2	0.51	0.80	0.2	1.55	5.03	0.38			1.43	0.179	2.85

Table 9: Recommended relative and absolute γ -ray intensities (%).

E_{γ} (keV)	Relative γ -ray intensity (%)	Absolute γ -ray intensity (%)	E_{γ} (keV)	Relative γ -ray intensity (%)	Absolute γ - ray intensity (%)	E_{γ} (keV)	Relative γ -ray intensity (%)	Absolute γ -ray intensity (%)
14	0.035 (1)	0.020 6 (6)	198	2.48 (10)	1.46 (6)	418	0.020 6 (11)	0.012 1 (6)
24	0.046 (11)	0.027 (6)	249	0.006 7 (2)	0.003 94 (12)	468	0.000 61 (9)	0.000 36 (5)
66	1.792 (34)	1.053 (20)	264	100	58.75 (19)	542	0.000 74 (1)	0.000 435 (6)
80	0.016 1 (9)	0.009 5 (5)	279	42.36 (6)	24.89 (9)	557	0.004 7 (2)	0.002 76 (12)
96	5.71 (12)	3.35 (7)	303	2.226 7 (44)	1.3082 (50)	572	0.061 65 (49)	0.036 22 (31)
121	28.7 (6)	16.86 (36)	373	0.004 36 (18)	0.002 56 (11)	617	0.007 71 (9)	0.004 53 (5)
136	97.8 (34)	57.7 (20)	400	19.384 (36)	11.388 (42)	821	0.000 28 (14)	0.000 134 (8)

6 References

- 1947Fr08 H. N. Friedlander, L. Seren, S. H. Turkel, Phys. Rev. 72(1947)23 [Half-life].
- 1948Co07 W. S. Cowart, M. L. Pool, D. A. McCown, L. L. Woodward, Phys. Rev. 73(1948)1454 [Half-life].
- 1950Co58 J. M. Cork, W. C. Rutledge, C. E. Branyan, A. E. Stoddard, J. M. Le Blanc, Phys. Rev. 79(1950)889 [Half-life].
- 1955Sc09 A. W. Schardt, J. P. Welker, Phys. Rev. 99(1955)810 [Gamma-ray energies and emission intensities].
- 1957Wr37 H. W. Wright, E. I. Wyatt, S. A. Reynolds, W. S. Lyon, T. H. Handley, Nucl. Sci. Eng. 2(1957)427 [Half-life].
- 1959Me76 F. R. Metzger, W. B. Todd, Nucl. Phys. 10(1959)220 [K conversion electron intensity].
- 1960Ea02 H. T. Easterday, R. L. Smith, Nucl. Phys. 20(1960)155 [Half-life].
- 1960Gr03 E. P. Grigoriev, A. V. Zolotavin, Nucl. Phys. 14(1960)443 [K conversion electron intensity].
- 1960De06 M. de Cröes, G. Bäckström, Ark. Fysik 16(1960)567 [K conversion electron intensity].
- 1961Ed02 W. F. Edwards, C. J. Gallagher, Nucl. Phys. 26(1961)649 [K conversion electron intensity].
- 1965Br19 D. R. Brundrit, S. K. Sen, Nucl. Phys. 68(1965)287 [K conversion electron intensity].
- 1966Ra09 P. V. Rao, D. K. McDaniels, B. Crasemann, Nucl. Phys. 81(1966)09 [Gamma-ray emission intensities].
- 1969Pa05 T. Paradellis, S. Hontzeas, Nucl. Phys. A131(1969)378 [Gamma-ray emission intensities].
- 1970Pa25 T. Paradellis, S. Hontzeas, Can. J. Phys. 48(1970)2254 [Gamma-ray emission intensities].
- 1970Pr07 W. W. Pratt, Nucl. Phys. A170(1971)223 [Gamma-ray emission intensities].
- 1973Su10 V. Sutela, Ann. Acad. Sci. Fen. AVI 407(1973) [K conversion electron intensity].
- 1973Th07 R. N. Thomas, R. V. Thomas, J. Phys. (London) A6(1973)1037 [Gamma-ray emission intensities].
- 1974Ca29 J. L. Campbell, J. Phys. (London) A7(1974)1451 [Gamma-ray emission intensities].
- 1975La19 F. Lagoutine, J. Legrand, C. Bac, Int. J. Appl. Radiat. Isotop. 26(1975)131 [Half-life].
- 1976MaZW M. J. Martin, Report ORNL 5114 (1976) [Half-life].
- 1977Kr13 K. S. Krane, At. Data Nucl. Data Tables 19(1977)363 [Mixing ratio].
- 1977Pr08 R. Prasad, Can. J. Phys. 55(1977)2036 [Gamma-ray emission intensities and energies].
- 1977Ge12 R. J. Gehrke, R. G. Helmer, R. C. Greenwood, Nucl. Instrum. Meth. 147(1977)405 [Gamma-ray emission energies].

- 1980Sc07 U. Schötzig, K. Debertin, K. F. Walz, Nucl. Instrum. Meth. 169(1980)43 [Half-life, gamma-ray emission intensities].
- 1980Ho17 H. Houtermans, O. Milosevic, F. Reichel, Int. J. Appl. Radiat. Isotop. 31(1980)153 [Half-life].
- 1982HoZJ D. D. Hoppes, J. M. R. Hutchinson, F. J. Schima, M. P. Unterweger, NBS 626(1982)85 [Half-life].
- 1983Si25 K. Singh, R. Mittal, M. L. Hasiza, H. S. Sahota, Indian J. Phys. 57A(1983)127 [X-ray intensities].
- 1983Yo03 Y. Yoshizawa, Y. Iwata, T. Katoh, J. -Z. Ruan, Y. Kawada, Nucl. Instrum. Meth. 212(1983)249 [Gamma-ray emission intensities].
- 1984Si02 K. Singh, H. S. Sahota, J. Phys. (London) G10(1984)241 [Gamma-ray emission intensities].
- 1987JeZZ R. Jedlovszki, T. Barta, M. Csikos, Gy. Horvath, L. Szücs, A. Szinka, Report OHM-GS 32(1987) [Gamma-ray emission intensities].
- 1990An07 A. Andai, T. Barta, L. Szücs, Nucl. Instrum. Meth. Phys. Res. A286(1990)457 [Gamma-ray emission intensities].
- 1990Je01 R. Jedlovszki, L. Szücs, A. Szörényi, Nucl. Instrum. Meth. Phys. Res. A286(1990)462 [Gamma-ray emission intensities].
- 1990Me15 R. A. Meyer, Fisika 22(1990)153 [Gamma-ray emission intensities].
- 1990Wa03 X. L. Wang, Y. Wang, Nucl. Instrum. Meth. Phys. Res. A286(1990)460 [Gamma-ray emission intensities].
- 1991BaZS W. Bambynek, T. Barta, R. Jedlovszki, P. Christmas, N. Coursol, K. Debertin, R. G. Helmer, A. L. Nichols, F. J. Schima, Y. Yoshizawa, Report IAEA TECDOC 619(1991) [Half-life, gamma-ray emission intensities].
- 1992Sc09 U. Schötzig, Nucl. Instrum. Meth. Phys. Res. A312(1992)141 [Gamma-ray emission intensities].
- 1994Mi22 H. Miyahara, H. Matumoto, C. Mori, N. Takeuchi, T. Genka, Nucl. Instrum. Meth. Phys. Res. A339(1994)203 [Gamma-ray emission intensities].
- 1994Iw04 A. Iwahara, I. P. A. Salati, R. Poledna, C. J. da Silva, L. Tauhata, Nucl. Instrum. Meth. Phys. Res. A339(1994)381 [Half-life].
- 1996Sa22 T. E. Sazonova, A. V. Zanevsky, S. V. Sepman, Nucl. Instrum. Meth. Phys. Res. A369(1996)421 [X-ray intensities].
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
- 1997Lo10 L. C. Longoria, J. S. Benitez, Appl. Rad. Isotopes 48(1997)1069 [Gamma-ray emission intensities].
- 1999Fa05 A. R. Farhan, B. Singh, Nucl. Data Sheets 86(1999)785 [Level energies, spin and parity].
- 2000Zhang Q.-S. Zhang, L. Yin-ming, Y. Chang, C. Yan, W. Li, At. Energ. Sci. Tech. (Chine) 34(2000)422 [X-ray intensities].
- 2002Un02 M. P. Unterweger, Appl. Rad. Isotopes 56(2002)125 [Half-life].
- 2002He19 M. He, S. Jiang, L. Diao, S. Wu, C. Li, Nucl. Instrum. Meth. Phys. Res. B194(2002)393 [Half-life].
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].
- 2005Ra29 D. R. Rao, K. V. Sai, M. Sainath, K. Venkataramaniah, Eur. Phys. J. A26(2005)41 [Gamma-ray emission intensities].
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr. , Nucl. Instrum. Meth. Phys. Res. A589(2008)202 [Theoretical ICC].

⁷⁹Se - Comments on evaluation of decay data by M. M. Bé and V. Chisté

This evaluation was completed in January 2006.

1. Decay scheme

The J^π value and level energy are from NDS 70,3 (1993).

2. Nuclear Data

- The Q value is from Audi *et al.* (2003)
- Published values of the half life are, in years :

Historical values		<i>a</i>	
1949	Parker <i>et al.</i>	$\leq 6.5 \times 10^4$	Report ORNL- 499, p.45
1951	Glendenin	$\geq 7 \times 10^6$	Radiochemical studies : The fission products, C.D. Coryell, N. Sugarman, New-York, McGraw Hill (1951) 596
Revised value			
1993	B. Singh	$\leq 6.5 \times 10^5$	NDS 70,3 p. 452
Measured Values			
1995	Yu Runlan, Guo Jingru <i>et al.</i>	$4.8 (4) \times 10^5$	J. Radioanalytical and Nuclear Chemistry, Articles, 196,1 p. 165
1997	Jiang Songsheng, Guo Jingru <i>et al.</i>	$1.1 (2) \times 10^6$	Nucl. Instr. Methods B123, p 405
2000	Ming He, Shan Jiang <i>et al.</i>	$1.24 (19) \times 10^5$	Nucl. Instr. Methods B172, p 177
2002	Songsheng Jiang, Ming He <i>et al.</i>	$2.95 (38) \times 10^5$	Nucl. Instr. Methods A489, p 195 or Chin. Phys. Lett. 18 (2001) 746
2002	Ming He, Songshen Jiang <i>et al.</i>	$2.80 (36) \times 10^5$	Nucl. Instr. Methods B194, p 393
2006	Bienvenu, <i>et al.</i>	$3.77 (19) \times 10^5$	To be published
Adopted		$3.56 (40) \times 10^5$	

Assessments of the Se-79 half-life were done in the years 49-50 (Parker, Glendenin) and a value of $6.5 \times 10^4 a$ was accepted by the various tables and chart of isotopes.

In 1993, due to inconsistencies in the measured and calculated fission yields of ⁷⁹Se for an irradiated fuel from a reactor, the calculations of Parker were reviewed (Singh) and a new value of $6.5 \times 10^5 a$ (i.e. one order of magnitude more) was deduced. Hence, in 1995 a Chinese team carried out the first measurement of this half-life by the means of a radiochemical method, they obtained $4.8 (4) \times 10^5 a$. However, and since this date, the same team, using the same ⁷⁹Se source published various results (see table above), the highest being $1.1 (2) \times 10^6 a$ (1997), and the last $2.80 (36) \times 10^5 a$ (2002).

Only one value (the last) will be used in this evaluation.

In NDS 96,1 (2002) B. Singh adopts the result of 2.96 (38) from the Chinese team.

In 2006, an independent result was published by P. Bienvenu *et al.* confirming the range 10^5 a for this half-life. In this study, the concentration of ⁷⁹Se was measured using ICP-MS coupled with Electro-Thermal Vaporisation to eliminate potential isobaric interferences and, the activity was measured using LSC after gamma ray spectrometry to check the contribution of residual radioactive contaminants.

In this evaluation, the adopted value is the weighted mean of the last Chinese value (NIM B194) and of the Bienvenu *et al.* value. They are in the same range but not consistent so, the adopted uncertainty is the external uncertainty.

2.1 b emission

⁷⁹Se is a pure beta minus emitter which disintegrates directly to the ground state level of ⁷⁹Br, no gamma rays are emitted.

The end-point energy is deduced from the Q value. The mean beta energy was calculated for a 1st forbidden unique transition.

References

- G.W. Parker *et al.* ORNL-499 (1949) 45
- L.E. Glendenin, MDDC-1694-C (1948) and Radiochemical studies : The fission products, C.D. Coryell, N. Sugarman, New-York, McGraw Hill (1951) 596
- B. Singh. O.W. Hermann, quoted in Nuclear Data Sheets 70,3 (1993) 452
- B. Singh, Nuclear Data Sheets 96,1 (2002) 30
- Yu Runlan *et al.* J. Radioanalytical and Nuclear Chemistry, Articles, 196,1 (1995) 165
- Jiang Songsheng *et al.* Nucl. Instr. Methods B123 (1997) 405
- Ming He *et al.* Nucl. Instr. Methods B172 (2000) 177
- Jiang Songsheng, Ming He, *et al.* Nucl. Instr. Methods A489 (2002) 195
- Ming He, Songshen Jiang *et al.* Nucl. Instr. Methods B194 (2002) 393
- G. Audi, A. H. Wapstra. Nucl. Phys. A729, 1 (2003) 337 Q
- P. Bienvenu, P. Cassette, G. Andreoletti, M. -M. Bé, J. Comte, M. -C. Lépy. Appl. Rad. Isotopes 65 (2007) 355

⁸²Rb - Comments on evaluation by M.-M. Bé

This evaluation was completed in January 2015. All literature available by this date was included.

1 Decay Scheme

The decay scheme of Rb-82 is taken from Meyer *et al.* (1983Me08) who observed 46 γ rays in the decay of Rb-82 to Kr-82, whereas only the most intense rays were studied in the other published papers. A statistical analysis process was adopted when several values existed, while all of the other γ rays as given by Meyer *et al.* were accepted.

Rb-82 is produced in the EC decay of Sr-82 ($T_{1/2} = 25,347 (17) d$).

2 Nuclear Data

Q^+ value of 4403 (3) keV is from 2012Wa38.

The following measured values of the half-life of ⁸²Rb have been taken into consideration:

Reference	$T_{1/2}$ (min)	Uc (min)
1953Kr10	1,267	0,083
1953Li27	1,25	0,03
1978Gr17	1,273	0,002
1987Wo01	1,2575	0,0005
Chi2 crit	3,8	
Chi2	10,1	
UWM	1,2618	
LWM	1,2652	
uc(WM)int	0,0014	
uc(WM)ext	0,0045	
Adopted	1,2652	0,0045

This set of data is discrepant, and the uncertainty in the half-life given by 1987Wo01 has been increased to 0,0020 in order to limit the weight of this particular value to 50 %.

2.1 and 2.2 Electron Capture Transitions and β^+ Emissions

The sums of the EC + β^+ transition probabilities were deduced from the sum of the gamma transition probabilities populating and depopulating each level of the decay scheme. The EC/ β^+ ratios were calculated using the Logft program, and a relative uncertainty of 5 % was assumed. Individual EC and β^+ probabilities were derived from these data.

From the values listed below, the sum of $P(\beta^+)$ amounts to 95,45 (30) %, which corresponds to a 511-keV photon intensity of 190,9 (6) %. This result and the ratio I_{511}/I_{776} can be compared with experimental values.

Reference	$I\beta^+$ (%)	u_c	I_{511}/I_{776}	u_c
1962Sa10	96,1	–	14,3	2,2
1967Vr07	95,4	–	14,52	0,90
1969Ra06	94,6	–	14,13	1,0
1970Gr01	–	–	14,08	0,8
1987Ho06	96,0	0,2	12,886	0,034
1987Ju01	95,87	0,15	12,67	0,15
Adopted	95,45	0,30	12,71	0,17

The energies are derived from the Q value and the level energies. The fractional probabilities for EC are calculated using the "Tables for Calculation of Electron Capture" (E. Schönfeld, PTB Laboratory report 6.33-95-2 (1995) - 1995ScZY). These values are based on the wave functions determined by J.B. Mann and J.T. Waber (*Atomic Data* **5** (1973) 201), with the exchange and overlap corrections of J.N. Bahcall and E. Vatai (see 1977Ba48; J.N. Bahcall, *Phys. Rev.* **132** (1963) 362; *Nucl. Phys.* **71** (1965) 267; E. Vatai, *Nucl. Phys.* **A156** (1970) 541).

2.3 Gamma Transitions

The transition probabilities were derived from the emission intensities and the conversion coefficients, but in most cases the internal conversions are weak and can be neglected. When given, the conversion coefficients have been interpolated from tables of Band (2002Ba85) by using the BrIcc program with the "frozen orbital approximation" (Kibédi *et al.*, 2008Ki07).

3. Atomic Data

All atomic data are taken from E. Schönfeld and H. Janssen, *Nucl. Instrum. Methods Phys. Res.* A369 (1996)527 (1996Sc06).

4. Electron Emissions

The energies of the conversion electrons are calculated from the energies of the γ rays and the corresponding electron binding energies.

Emission intensities for the Auger electrons and X-rays were derived from the decay scheme parameters by means of the EMISSION program.

5. Photon Emissions

The following data have been considered in the determination of the γ -ray emission intensities relative to a value of 100 % for the intensity of the 776-keV γ ray:

Reference	E (keV)					
	696,86 (5)	698,37 (5)	711,2 (1)	1395,14 (3)	1474,88 (1)	1703,19 (4)
Vrzal (1967Vr07)	–	1,35 (8)	0,36 (5)	3,98 (15)	0,63 (3)	0,42 (3)
Raman (1969Ra06)	0,430(22)	1,04 (5)	0,35 (9)	3,53 (18)	0,61 (3)	0,37 (9)
Graeffe (1970Gr01)	–	1,28 (10)	0,31 (5)	3,8 (20)	0,89 (10) ^(o)	0,48 (10)
Hrynkiewicz (1970Hr02)	0,22 (7) ^(o)	1,0 (1)	0,23 (6)	3,78 (40)	0,68 (7) ^(o)	0,41 (10)
Meyer (1983Me08)	0,51 (2)	0,99 (3)	0,38 (2)	3,51 (2)	0,59 (2)	0,335 (4)
Hoppes (1987Ho06)	–	1,54 (2) ^(o)	0,470(21)	3,52 (3)	0,601 (14)	0,330 (12)
Chi2	6,9	5,5	4,8	2,4	0,4	2,2
Chi2 crit	6,6	3,3	3,0	3,0	3,8	3,0
UWM	0,470	1,132	0,35	3,687	0,6078	0,3908
WM	0,470	1,062	0,400	3,5212	0,6025	0,3362
uc int	0,015	0,026	0,013	0,0162	0,0100	0,0037
uc ext	0,040	0,062	0,029	0,0251	0,0066	0,0055
LWM	0,47	1,06	0,400	3,521	0,602	0,336
Adopted	0,47 (4)	1,06 (7)	0,400 (29)	3,521 (25)	0,602 (14)^(m)	0,336 (6)

Reference	E (keV)			
	2167,59 (4)	2410,26 (5)	2479,65 (4)	2940,0 (3)
Vrzal (1967Vr07)	0,34 (3)	0,15 (2)	0,22 (5)	–
Raman (1969Ra06)	0,25 (6)	0,12 (3)	0,220 (55)	–
Graeffe (1970Gr01)	0,32 (5)	0,20 (5)	0,28 (6)	0,08 (3)
Hrynkiewicz (1970Hr02)	0,35 (8)	0,17 (4)	0,28 (7)	0,07 (2)
Meyer (1983Me08)	0,28 (1)	0,156 (8)	0,27 (1)	0,021 (4) ^(u)
Hoppes (1987Ho06)	–	–	–	–
Chi2	1,3	0,6	0,4	2,5
Chi2 crit	3,3	3,3	3,3	4,6
UWM	0,308	0,1592	0,254	0,057
WM	0,2872	0,1546	0,2672	0,0470
uc int	0,0092	0,0070	0,0094	0,0118
uc ext	0,0103	0,0054	0,0063	0,0186
LWM	0,287	0,155	0,267	0,047
Adopted	0,287 (1)	0,155 (8)^(m)	0,267 (10)^(m)	0,047 (19)

^(o) Omitted from statistical processing.

^(m) Minimum experimental uncertainty.

^(u) Uncertainty increased to limit the weighting to 50 %.

All other rays listed in the tables and decay scheme are from Meyer *et al.* (1983Me08). The γ rays with energies of 466,9 and 2508,9 keV given in Meyer are of uncertain origin, and have not been included in the proposed decay scheme.

Two E0 transitions with very weak probabilities of the order of $(10^{-5} - 10^{-6})$ %, as deduced by 1985Ze03 from the measurement of conversion electrons with a mini-orange spectrometer, were also not included in the tables.

Several measured values of the absolute intensity of the 776 keV γ -ray exist, obtained mainly by means of coincidence techniques:

Reference	I γ (%)	Uncertainty (%)	
1962Sa10	13	–	Coincidence techniques
1967Vr07	13,2	–	Coincidence techniques
1969Ra06	13,4	–	Coincidence techniques
1987Ho06	14,9	0,4	Coincidence techniques
1987Ju01	15,13	0,27	Original uncertainty = 0,18
2012Gr03	14,93	0,37	Ion implantation of ⁸² Sr
Chi2	0,16		
Chi2 crit	4,61		
UWM	14,987		
WM	15,024		
uc int	0,19		
uc ext	0,08		
LWM	15,02		
Adopted	15,02	0,19	WM and internal uncertainty

There are only three measurements in which the uncertainties are quoted. The uncertainty given by 1987Ju01 was increased to limit the weighting of this value to 50 %.

The absolute emission intensity of the 511-keV annihilation photon has been deduced from the decay scheme data and the theoretical EC/ β^+ ratios, as described in Section 2.1

6. References

- 1953Kr10** P.Kruger, N.Sugarman. Phys. Rev. 90 (1953) 158. Half-life
1953Li27 L.M.Litz, S.A.Ring, W.R.Balkwell. Phys. Rev. 92 (1953) 288. Half-life
1962Sa10 M.Sakai, H.Ikegami, T.Yamazaki. J. Phys. Soc. Japan 17 (1962) 1087. Gamma-ray emission intensities
1967Vr07 J.Vrzal, B.S.Dzhelepov, A.G.Dmitriev, N.N.Zhukovskii, J.Liptak, L.N.Moskvin, J.Urbanets, L.G.Tsaritsyna. Bull. Acad. Sci. USSR, Phys. Ser. 31 (1968) 1701. Gamma-ray emission intensities
1969Ra06 S.Raman, J.J.Pinajian. Nucl. Phys. A125 (1969) 129. Gamma-ray emission intensities
1970Gr01 G.Graeffe, S.Vaisala, J.Heinonen. Nucl. Phys. A140 (1970) 161. Gamma-ray emission intensities
1970Hr02 A.Z.Hryniewicz, J.Styczen, W.Walus, R.Broda. Acta Phys. Polonica A38 (1970) 501. Gamma-ray emission intensities
1977Ba48 W. Bambynek, H.Behrens, M.H.Chen, B.Crasemann, M.L.Fitzpatrick, K.W.D.Ledingham, H.Genz, M.Mutterer, R.L.Intemann, Rev. Mod. Phys. 49 (1977) 77 (Erratum. Rev. Mod. Phys. 49 (1977) 961) EC transition probabilities
1978Gr17 P.M.Grant, B.R.Erdal, R.E.Whipple, R.J.Daniels, H.A.O'Brien Jr. Phys. Rev. C18 (1978) 2799. Half-life
1983Me08 R.A.Meyer, J.F.Wild, K.Eskola, M.E.Leino, S.Vaisala, K.Forssten, U.Kaup, A.Gelberg. Phys. Rev. C27 (1983) 2217. Gamma-ray emission intensities
1985Ze03 A.Zemel, T.Hageman, J.J.Hamill, J.van Klinken. Phys. Rev. C31 (1985) 1483. Gamma transitions
1987Ho06 D.D.Hoppes, B.M.Coursey, F.J.Schima, D.Yang. Appl. Radiat. Isotopes 38 (1987) 195. Half-life
1987Ju01 S.M.Judge, M.J.Woods, S.L.Waters, K.R.Butler. Appl. Radiat. Isotopes 38 (1987) 185. Gamma-ray emission intensities

- 1987Wo01** M.J.Woods, S.M.Judge, S.E.M.Lucas. Appl. Radiat. Isotopes 38 (1987) 191. Gamma-ray emission intensities
- 1995ScZY** E.Schönfeld. PTB Report PTB-6.33-95-2, Braunschweig, Germany (1995). EC transition probabilities
- 1996Sc06** E.Schönfeld, H.Janssen. Nucl. Instrum. Methods Phys. Res. A369 (1996) 527. Atomic Data
- 1998Sc28** E.Schönfeld. Appl. Radiat. Isotopes 49 (1998) 1353. Atomic Data
- 2000Sc47** E.Schönfeld, H.Janssen. Appl. Radiat. Isotopes 52 (2000) 595. Atomic Data
- 2002Ba85** I.M.Band, M.B.Trzhaskovskaya, C.W.Nestor, Jr., P.O.Tikkanen, S.Raman. At. Data. Nucl. Data Tables 81 (2002) 1. Theoretical ICC
- 2008Ki07** T.Kibédi, T.W.Burrows, M.B.Trzhaskovskaya, P.M.Davidson, C.W.Nestor Jr. Nucl. Instrum. Methods Phys. Res. A589 (2008) 202. Theoretical ICC
- 2012Gr03** C.J.Gross, K.P.Rykaczewski, D.W.Stracener, M.Wolinska-Cichocka, R.L.Varner, D.Miller, C.U.Jost, M.Karny, A.Korgul, S.Liu, M.Madurga. Phys. Rev. C85 (2012) 024319. Gamma-ray emission intensities
- 2012Wa38** M.Wang, G.Audi, A.H.Wapstra, F.G.Kondev, M.MacCormick, X.Xu, B.Pfeiffer. Chin. Phys. C36 (2012) 1603. Q

⁸²Sr - Comments on evaluation by M.-M. Bé

This evaluation was completed in December 2014. All literature available by this date was included.

1. Decay Scheme

Sr-82 decays by electron capture to the ground state of Rb-82 ($T_{1/2} = 1,2652$ min). There is no decay to the excited level of Rb-82m ($T_{1/2} = 6,47$ h).

2. Nuclear Data

Q_{EC} -value of 178 (7) keV for Sr-82 EC decay is taken from 2012Wa38.

The following measured values for the half-life of ⁸²Sr have been taken into account:

Reference	$T_{1/2}$ (d)	Comments
1953Kr10	25,5 (5)	
1953Li27	27	omitted
1958Sa20	25,0 (4)	
1978Gr17	25,55 (15)	
1987Ju02	25,342 (41)	Uncertainty calculated as the combination, in quadrature, of the statistic and systematic uncertainties given in the text.
1987Ho06	25,36 (3)	Superseded by 2009Pi02
2009Pi02	25,346 (17)	Weighted mean of two values
Adopted	25,347 (17)	χ^2 crit = 3,3 ; $\chi^2 = 0,7$

The set of five data and their uncertainties is consistent, and their weighted mean value has been adopted along with the lowest experimental uncertainty.

3. Electron Capture Transition

The energy is derived from the Q_{EC} -value, and the fractional probabilities for EC were calculated using the "Tables for Calculation of Electron Capture" (1998Sc28).

4. Atomic Data

The fluorescence yield data, the relative K X-ray emission probabilities, the ratios $P(KLX)/P(KLL)$ and $P(KXY)/P(KLL)$ are from Schönfeld *et al.* (1996Sc06).

The Auger electrons and X-ray absolute intensities were calculated by the EMISSION program (2000Sc47) from the related decay data (P_{EC} sub shell probabilities, etc.) as determined above.

5. Main Production Mode

Rb-85(p, 4n)Sr-82
Nat. Rb(p, xn)Sr-82

6. References

- 1953Kr10** P.Kruger, N.Sugarman. Phys. Rev. 90 (1953) 158. Half-life
1953Li27 L.M.Litz, S.A.Ring, W.R.Balkwell. Phys. Rev. 92 (1953) 288. Half-life
1958Sa20 V.Sangiust. Nuovo Cim. 9 (1958) 446. Half-life
1978Gr17 P.M.Grant, B.R.Erdal, R.E.Whipple, R.J.Daniels, H.A.O'Brien Jr. Phys. Rev. C18 (1978) 2799. Half-Life
1987Ho06 D.D.Hoppes, B.M.Coursey, F.J.Schima, D.Yang. Appl. Radiat. Isotopes38 (1987) 195. Half-life
1987Ju02 S.M.Judge, A.M.Privitera, M.J.Woods. Appl. Radiat. Isotopes38 (1987) 193. Half-life
1996Sc06 E.Schönfeld, H.Janssen. Nucl. Instrum. Methods Phys. Res. A369 (1996) 527. Atomic Data
1998Sc28 E.Schönfeld. Appl. Radiat. Isotopes49 (1998) 1353. Atomic Data
2000Sc47 E.Schönfeld, H.Janssen. Appl. Radiat. Isotopes52 (2000) 595. Atomic Data
2009Pi02 L.Pibida, R.Fitzgerald, M.Unterweger, M.M.Hammond, D.Golas. Appl. Radiat. Isotopes67 (2009) 636. Half-life
2012Wa38 M.Wang, G.Audi, A.H.Wapstra, F.G.Kondev, M.MacCormick, X.Xu, B.Pfeiffer. Chin. Phys. C36 (2012) 1603. Q

⁸⁵Kr – Comments on evaluation of decay data by V. Chisté and M. M. Bé

This evaluation was completed in July 2003 and the half life value has been updated in May 2004.

1) Decay Scheme

⁸⁵Kr disintegrates by β^- emission to the ⁸⁵Rb ground state (99.562(10)%) and to the second excited level at 513.998(5) keV (0.438(10)%). The decay scheme is based mainly on the measurements of the 514 keV γ -emission intensity (see § 4. Radiation Emission, 4.2 Gamma Ray Emissions).

2) Nuclear Data

The Q value is from Audi and Wapstra (1995Au04)

Level energies, spins and parities are from R. A. Meyer (1980Me06).

The measured ⁸⁵Kr half-life values are, in years:

$T_{1/2}$

Reference	Value (a)	Comments
Thode (1948Th06)	9.4 (4)	
Turner (1953Tu22)	10.57 (14)	
Wanless (1953Wa17)	10.27 (18)	
Lerner (1963Le07)	10.76 (2)	
Anspach (1965An07)	10.75 (3)	
Johnston (1974Jo12)	10.714 (57)	
Walz (1983Wa15)	10.702(8)	Superseded by 2003Sc49
Unterweger (1992Un03)	10.7720(38)	Superseded by 2002Un04
Eberszkorn (1996Er06)	10.757 (49)	
Unterweger (2002Un04)	10.7756(33)	
Schrader (2003Sc49)	10.724(7)	

Evaluators calculated the weighted average of these 9 values using the Lweight program (version 3) as 10.750 years with an external uncertainty of 0.011 and a reduced $-\chi^2$ of 6.34. Evaluators rejected the Thode (1948Th06), Turner (1953Tu22) and Wanless (1953Wa17) values based on the Chauvenet's criterion. For the remaining 6 values, the largest contribution to the weighted average comes from the value of Unterweger (2002Un04), amounting to 79%. The program Lweight 3 increased the uncertainty for the 2002Un04 value from 0.0033 to 0.0064 in order to reduce its relative weight from 79% to 50%.

The adopted value is the weighted mean : 10.752 a, with an uncertainty of 0.023 (expanded so range includes the most precise value of Unterweger (2002Un04)) and a reduced- χ^2 of 6.

2.1) β^- Transitions

The β^- probabilities and the associated uncertainties have been deduced from γ transition probability balance at each level of the decay scheme, i. e., $P_{\beta}(0,0) = 99.562(10)\%$ and $P_{\beta}(0,2) = 0.438(10)\%$. The values of $\log ft$ have been calculated with the program LOGFT for the Allowed and 1st Unique Forbidden transitions.

2.2) Gamma Transitions

Probabilities

The transition probabilities have been calculated from the gamma emission intensities and the internal conversion coefficients (see § 4.2) **Gamma Ray Emissions**).

Mixing ratios and internal conversion coefficients

The adopted δ ($= 0.072(4)$) for the 151 keV γ -transition and the gamma transition multipolarities of the 362 keV ((E3)) and of the 513 keV (M2, from ⁸⁵Sr ground state decay) were adopted from Sievers (1991Si01).

The theoretical internal conversion coefficients (table 1) have been interpolated from values in 1978Ro22 using the ICC Computer Code (program Icc99v3a – GETICC dialog).

Table 1:

E_g (keV)	Multipolarity	Value of α_K	Value of α_L	Value of α_T
151.18 (3)	M1 + 0.52(4)% E2	0.0430(13)	0.00485(14)	0.0488(14)
362.81 (3)	(E3)	0.0292(9)	0.0040(1)	0.0340(10)
513.998 (5)	M2	0.00635(19)	0.00072(2)	0.00721(21)

For the 151 keV γ -transition, the α_T is calculated as follows:

$$\alpha_T(M1) * \%(M1) + \alpha_T(E2) * \%(E2) = (0.00479(14) * 0.9948(4)) + (0.213(6) * 0.0052(4)) = 0.0488(14)$$

Calculations of ICC uncertainties for transitions:

* For the all transitions, uncertainties in α_T , α_K and α_L calculated values with ICC Computer Code (program Icc99v3a) are taken to be 3% .

3) Atomic Data

Atomic values (ω_K , ω_L and n_{KL}) are from Schönfeld (1996Sc33).

The X-ray and Auger probabilities are calculated by Emission program.

4) Radiation emissions

4.2) Gamma ray emissions

Gamma ray energies (in keV) are from R. A. Meyer (1980Me06).

Emission probability values are deduced from measured values of the 514 keV absolute γ -emission intensity in Table 2 and using values relative to 514-keV transition for the other gamma-rays (1980Me06) shown in Table 3.

Table 2:

Reference	514 keV γ -emission intensity (%)	Comments
Geiger (1961Ge19)	0.46 (4)	
Eastwood (1964Ea01)	0.431(17)	
Denecke (1967De05)	0.435 (13)	
Weighted Average (Lweight 3)	0.435 (10)	Reduced- $\chi^2 = 0.22$

Table 3:

Energy (keV)	Relative γ -emission intensity measured by R. A. Meyer (1980Me06) (%)	Absolute γ -emission intensity (%)
151	0.0005 (3)	0.0000022(13)
362	0.0005 (1)	0.00000218(44)
514	100	0.435(10)

With these values shown in table 3, and the values of α_T calculated using the ICC Computer Code (table 1, section 2.2), evaluators deduced the γ -transition probability (table 4).

Table 4:

Energy (keV)	Transition probability (%)
151	0.0000023(14)
362	0.00000225(45)
514	0.438(10)

References

- 1948Th06 – H. G. Thode, *Nucleonics* 3(1948)14 [Half-life].
 1953Tu22 – J. F. Turner, *AERE – N/R* 1254(1953) [Half-life].
 1953Wa17 – R. K. Wanless, H. G. Thode, *Can. J. Phys.* 31(1953)517 [Half-life].
 1961Ge19 – K. W. Geiger, J. S. Merritt, J. G. V. Taylor, *Nucleonics* 19(1961)97 [514 γ -Branching ratio].
 1963Le07 – J. Lerner, *J. Inorg. Nucl. Chem.* 25(1963)749 [Half-life].
 1964Ea01 – T. A. Eastwood, F. Brown, R. D. Werner, *Can. J. Phys.* 42(1964)218 [514 γ -Branching ratio].
 1965An07 – S. C. Anspach, L. M. Cavallo, S. B. Garfinkel, J. M. R. Hutchinson, C. N. Smith, *NP* – 15663(1965) [Half-life].
 1967De05 – B. Denecke, E. de Roost, A. Spornol, R. Vaninbrouckx, *Nucl. Sci. Eng.* 28(1967)305 [514 γ -Branching ratio].
 1971Ho05 – D. J. Horen, *Nucl. Data Sheets* B5(1971)131 [Multipolarities, mixing ratio, half-life, J^π].
 1974Jo12 – J. W. Johnston, *BNWL – B* – 369(1974) [Half-life].
 1978Ro22 – F. Rösel et al, *At. Data Nucl. Data Tables* 21(1978)91 [ICC].
 1980Me06 – R. A. Meyer, J. E. Fontanilla, N. L. Smith, C. F. Smith, R. C. Ragaini, *Phys. Rev.* C21(1980)2590 [Energy and emission probability].
 1980Si01 – H. Sievers, *Nucl. Data Sheets* 30(1980)501 [Multipolarities, mixing ratio, half-life, J^π].
 1983Wa15 – K. F. Walz, K. Debertin, H. Schrader, *Int. J. Appl. Rad. Isotopes* 34(1983)1191 [Half-life].
 1991Si01 – H. Sievers, *Nucl. Data Sheets* 62(1991)271 [Multipolarities, mixing ratio, half-life, J^π].
 1992Un03 – M. P. Unterweger, D. D. Hoppes, F. J. Schima, *Nucl. Instr. Meth. Phys. Research* A312(1992)349 [Half-life].
 1995Au04 – G. Audi, A. H. Wapstra, *Nucl. Phys.* A565(1995)1 [Q-value].
 1996Er06 – L. Erbeszkorn, Á. Szörényi, J. Vágvolgyi, *Nucl. Instr. Meth. Phys. Research* A369(1996)463 [Half-life].
 1996Sc33 – E. Schönfeld, H. Janßen, *Nucl. Instr. Meth. Phys. Research* A369(1996)527 [Atomic Data].
 2002Un04 – M. P. Unterweger, *Appl. Rad. Isotopes* 56(2002)125 [Half-life].
 2003Sc49 – H. Schrader, *Appl. Rad. Isotopes* 60, 2-3 (2004) 317 [Half-life].

⁸⁵Sr - Comments on evaluation of decay data by E. Schönfeld, R. Dersch

1 Decay Scheme

The decay scheme is taken from Torti et al. (1972) and Meyer et al. (1980). A level at 951 keV which is depopulated by four gamma transitions (see Section 4.3) was observed by Barnard et al. (1973) in n, γ reactions. An EC transition to this level in the ⁸⁵Sr disintegration would be second forbidden. An upper limit of $3 \cdot 10^{-7}$ was estimated for this transition. The existence of EC transitions to the levels at 281 keV (unique third forbidden) and 151 keV (third forbidden) is also questionable.

Below the Q_{EC} value there are also levels at 919,7 keV (possibly two levels, $1/2^-$ or $3/2^-$ and $5/2^-$, populated in the decay of 68 min ⁸⁵Sr^m and several reactions) and 731,822 keV ($3/2^-$, populated in the decay of 4 h ⁸⁵Kr^m and several reactions). EC transitions from ⁸⁵Sr ground state to these levels would be both 3rd forbidden, γ rays from these levels have not been observed in the decay of ⁸⁵Sr.

The main transitions in the EC decay of ⁸⁵Sr are the EC transition populating the 514 keV level of ⁸⁵Rb and the γ transition of 514 keV depopulating this level. Besides these transitions there is an EC transition to the 869 keV level which is mainly depopulated by 869 keV γ rays.

The half-lives of the excited levels were taken from Sievers (1991). The half-life of the 514 keV level was determined by Siekman (1956), Löbner (1964), Miller *et al.* (1972) and Walz and Weiß (1976). Sievers took the value of Miller et al. which claims to be the most accurate one.

2 Nuclear Data

The following values of the half-life of ⁸⁵Sr have been considered ($T_{1/2}$ in d):

1	66	Dubridge and Marshall (1940)
2	65,0(7)	Herrmann and Strassmann (1956)
3	64,0(2)	Wright et al. (1957)
4	63,9(27)	Sattler (1962)
5	65,19(13)	Anspach et al. (1965)
6	66,6(6)	Grotheer et al. (1969)
7	64,93(22)	Emery et al. (1972)
8	64,68(23)	Lagoutine et al. (1972)
9	65,0(49)	Araminowicz and Dressler (1972)
10	65,0(50)	Vatai et al. (1974)
11	64,84(3)	Merritt and Gibson (1976); replaced by value 13
12	64,84(1)	Thomas (1978)
13	64,845(9)	Rutledge et al. (1980)
14	64,856(7)	Houtermans et al. (1980)
15	64,851(6)	Hoppes et al. (1982); replaced by value 17
16	64,85(14)	Walz et al. (1983)
17	64,8530(81)	Unterweger et al. (1992)
18	64,847(3)	unweighted mean of 12, 13, 14, 16, 17
19	64,850(7)	LWM (0,004 (int), 0,003 (ext), reduced χ^2 0,46), uncertainty enlarged to the uncertainty of the most accurate single value for the same five values

Values 1 - 11 are only of historical interest. They were not included in the averaging procedure.

The Q_{EC} value was taken from Audi and Wapstra (1995).

2.1 Electron capture Transitions

The main EC transition $\epsilon_{0,3}$ to the 514 keV level in ⁸⁵Rb is allowed ($\lg ft = 6,2$). A transition leading directly to the ground state ($\epsilon_{0,0}$) is unique 1st forbidden. The transition probability of this transition was estimated by Yoshizawa and Inoue (1991) by using the average $\lg ft$ value (according to Gove and Martin (1971)) of $9,47 \pm 0,17$ for seven neighbouring nuclei with uncertainty of 2σ . Their result is 0,8(4)%. The probability for the EC transition $\epsilon_{0,4}$ is deduced from the probabilities of the depopulating γ ray transitions. Concerning EC transitions $\epsilon_{0,2}$ and $\epsilon_{0,1}$ see Section 1. The data for the population and depopulation of the 151 keV level are discrepant as $P_{\gamma+ce}(4,1) + P_{\gamma+ce}(3,1) + P_{\gamma+ce}(2,1)$ is larger than $P_{\gamma+ce}(1,0)$. This can be explained (for example) by a too small value for $P_{\gamma+ce}(1,0)$. Moreover, it supports the assumption that an EC transition to the first excited level of ⁸⁵Rb at 151 keV does not exist.

Double K shell ionization was found by Schupp and Nagy (1984) $6,0(5) 10^{-5}$ per disintegration.

2.2 Gamma Transitions

The transition probability of 0,8(4)% for the EC transition directly feeding the ground state of ⁸⁵Rb yields for $P_{\gamma+ce}(514 \text{ keV}) = 99,2(4)\%$. Furthermore, with the total conversion coefficient of the 514 keV transition $I_{\gamma}(514) = 98,5(4)\%$. The transition probabilities of the other gamma transitions are derived from the measured emission probabilities (Sect. 4.2).

The conversion coefficients are interpolated from the tables of Rösler et al. (1978). The main transition $\gamma_{3,0}$ is assumed to have pure M2 multipolarity. The conversion coefficients of the other transitions have little influence on the balancing procedure because the emission probabilities of the assigned transitions are very small.

3 Atomic data

The atomic data are taken from Schönfeld and Janßen (1996).

3.1 X Radiation

The energies are based on the wavelengths of Bearden (1967). The relative probabilities are taken from Schönfeld and Janßen (1996).

3.2 Auger electrons

The energies are taken mainly from Larkins (1977). The relative probabilities are taken from Schönfeld and Janßen (1996).

4 Radiation Emission

4.1 Electron emission

The energies of the Auger electrons are the same as above. The energies of the conversion electrons are calculated from the transition energy and the binding energies. The number of Auger electrons per disintegration are calculated using the above mentioned atomic shell data and the program EMISSION. The number of conversion electrons related to the 514 keV γ -transition are calculated from the transition probability and the conversion coefficients.

4.1 X-ray emission

For the total K X-ray emission intensity, it was found three measured values :

Comments on evaluation

1	59,59(35)	Grotheer et al. (1969)
2	58,6(3)	Bambynek and Reher (1970)
3	58,66(47)	Thomas (1978)
4	59,04(34)	Weighted mean
5	58,95(32)	Unweighted mean
6	59,2 (6)	calculated from P_e , P_K , ω_K , P_{g+ce} This is the adopted value.

4.2 Photon Emission

The accuracy of the γ ray energy of the main line has improved during the last years, in keV :

1	514,0	Sattler (1962), Vartanov (1966)
2	513,98(3)	Legrand et al. (1968)
3	513,998	Ragaini et al (1972), Meyer et al. (1980)
4	514,009(12)	Helmer et al. (1978)
5	514,0076(22)	Kumahora et al. (1983)
6	514,00492(50)	Chang et al. (1993)
7	514,0048(22)	Helmer and van der Leun (2000), evaluation

The γ ray energies of the other transitions are taken from Sievers (1991).

From the balance of the decay scheme $P_{\gamma+ce}$ (514 keV) is calculated to be 99,2(4)%.

The ratio of the emission probabilities of the 869 keV and the 514 keV transitions were determined to be:

1	$1,7 \cdot 10^{-4}$	Sattler (1962)
2	$1,0(2) \cdot 10^{-4}$	Vartanov et al. (1966)
3	$1,4(2) \cdot 10^{-4}$	Vatai et al. (1974)
4	$1,154(63) \cdot 10^{-4}$	Pratt (1977)
5	$1,25(5) \cdot 10^{-4}$	Thomas (1978)
6	$1,25(5) \cdot 10^{-4}$	Meyer et al. (1980)
7	$1,23(3) \cdot 10^{-4}$	LWM of values 2 - 6

With the above-mentioned $I_\gamma(514) = 98,5(4) \%$ this yields $I_\gamma(869) = 0,0121(4) \%$.

Barnard *et al.* (1973) have observed in (n,n' γ) measurements a level at 951,3 keV in ^{85}Rb which is depopulated by the following gamma transitions: 951,3 keV (86 %), 800,2 keV (9 %), 670,3 keV (4 %) and 437,7 keV (1 %). If this level with the populated in the ^{85}Sr decay, the corresponding EC transition is second forbidden ($9/2^+ \rightarrow 5/2^+$; $\lg ft > 11,2$; transition energy 114(4) keV). Meyer *et al.* (1980) observed a 951 keV gamma ray in two spectra with high counting statistics and estimated an upper limit of $3 \cdot 10^{-7}$ for the emission probability of these gamma rays.

Levels at 731,9 keV ($3/2^-$) and 921 keV ($1/2^-$, $3/2^-$) in ^{85}Rb have not been found to be populated in the studies of the ^{85}Sr decay carried out by Meyer *et al.* (1980).

A level in ^{85}Rb at 281 keV, found by Barnard *et al.* (1973), is depopulated according to Meyer *et al.* (1980) by 129,8 keV gamma rays with an emission probability of $< 5 \cdot 10^{-3}$. As this is an upper limit the existence of this transition is not sure. Therefore, the population and depopulation of this level is given in the above decay scheme by dashed lines.

The gamma ray emission intensities in Table 5.2 and the corresponding values of the transition probabilities $P_{\gamma+ce}$ given in Table 2.2 are from Meyer *et al.* (1980) (129,8/151,1/355,0/362,8 keV) whereas the value for the 717,8 keV gamma rays is from Jerbic -Zorc (1990). The origin of the values for the 514 keV and 869 keV gamma rays were already explained above.

5 Main Production Modes

The main production modes are taken from Sievers (1991).

6 References

- L. A. Dubridge, J. Marshall, *Phys. Rev.* 58 (1940) 7
 [$T_{1/2}$]
 G. Herrmann, F. Strassmann, *Z. Naturforschg.* 11a (1956) 946
 [$T_{1/2}$]
 J. G. Siekman, *Nuclear Physics* 2 (1956/57) 254
 [$T_{1/2}$ 514 keV level]
 H. W. Wright, E. L. Wyatt, S. A. Reynolds, W. S. Lyon, T. H. Handley, *Nucl. Sci. Eng.* 2 (1957) 427
 [$T_{1/2}$]
 M. K. Ramaswamy, B. A. Bishara, P. S. Jastram, *Bull. Am. Phys. Soc.* 7 (1962) 341
 [P_{γ}]
 A. R. Sattler, *Phys. Rev.* 127 (1962) 854
 [$T_{1/2}$, P_{γ}]
 K. E. G. Löbner, *Nuclear Physics* 58 (1964) 49
 [$T_{1/2}$ 514 keV level]
 S. C. Anspach L. M. Cavallo, S. B. Garfinkel, J. M. R. Hutchinson, C. N. Smith, *NBS Misc. Publ.* 260-9 (1965)
 [$T_{1/2}$]
 J. Legrand, J. P. Boulanger, J. P. Brethon, *Nucl. Phys.* A107 (1968) 177
 [E_{γ}]
 M. McDonnell, M. K. Ramaswamy, *Nuclear Physics* A127 (1969) 531
 [Q_{EC}]
 H.-H. Grotheer, J. W. Hammer, K.-W. Hoffmann, *Z. Physik* 225 (1969) 293
 [$T_{1/2}$]
 I. F. Bubb, S. I. H. Naqui, J. L. Wolfson, *Nucl. Phys.* A167 (1971) 252
 [P_{γ}]
 J. F. Emery, S. A. Reynolds, E. I. Wyatt, G. I. Gleason, *Nucl. Sci. Eng.* 48 (1972) 319
 [$T_{1/2}$]
 J. Araminowicz, R. Dresler, *Report INR-1464* (1973) 14
 [$T_{1/2}$]
 R. C. Ragaini, C. F. Smith, R. A. Meyer, *Bull. Am. Phys. Soc.* 17 (1972) 444
 [levels]
 F. Lagoutine, J. Legrand, C. Perrot, J. P. Brethon, J. Morel, *Int. J. Appl. Rad. and Isot.* 23 (1972) 219
 [$T_{1/2}$]
 J. F. Emery, S. A. Reynolds, E. I. Wyatt, *Nucl. Science and Eng.* 48 (1972) 319
 [$T_{1/2}$]
 R. P. Torti, V. M. Cottles, V. R. Dave, J. A. Nelson, R. M. Wilenzick, *Phys. Rev.* 6C (1972) 1686
 [E_{γ} decay scheme]
 P. D. Bond, G. J. Kumbartzki, *Nucl. Phys.* A205 (1973) 239
 [J^{π} , P_{γ}]
 J. S. Merritt and F. H. Gibson, *AECL-5315* (1976) 37
 [$T_{1/2}$]
 R. G. Helmer, R. C. Greenwood, R. J. Gehrke, *Nucl. Instr. and Meth.* 155 (1978) 189
 [E_{γ}]
 Y. Yoshizawa, *Jaeri-M* 7567 (1978)
 [P_{γ}]
 G. Schupp, H. J. Nagy, *Phys. Rev.* 29 (1984) 1414
 [double K shell ionization]

For other references see Chapter “References” in the Table Part.

**⁸⁸Y - Comments on evaluation of decay data
by V.P. Chechev and N.K. Kuzmenko**

The first DDEP evaluation of ⁸⁸Y decay data was done by E. Schönfeld in 1998 with minor update in 2004 (2004BeZR). The current evaluation has been completed in June 2015 with a literature cut-off by the same date.

1. DECAY SCHEME

The structure of the adopted decay scheme of ⁸⁸Y is based on the ENSDF evaluation by McCutchan and Sonzogni NDS (2014Mc01). ⁸⁸Y disintegrates by the electron capture and β⁺-transition to the ⁸⁸Sr excited levels. Below the Q-value of 3622.6 keV there are two additional levels at 3486.6 and 3522.8 keV. They are not shown in the decay scheme because they are not populated in the disintegration of ⁸⁸Y. Up to now these levels were observed only in other disintegration processes, for example in the decay of ⁸⁸Rb (17.77 min).

An EC or β⁺ transition to the ground state of ⁸⁸Sr were also not observed. This is due to the high order of forbiddenness of such transition (4⁻ → 0⁺). Thus, the decay scheme of ⁸⁸Y adopted here is complete.

The spin, parity and half-life of the ⁸⁸Sr excited levels were adopted from the evaluation by 2014Mc01.

2. NUCLEAR DATA

Q⁺ - value is from 2012 mass evaluation of Wang et al. (2012Wa38).

The recommended half-life of ⁸⁸Y is based on the experimental results given in Table 1.

Table 1. Experimental values of the ⁸⁸Y half-life (in days)

No	Author(s) and year	Reference	T _{1/2}	Method and comments
1	DuBridge and Marshall (1940)	1940Du09	105 (5)	Ionization chamber; <i>omitted</i> (too large uncertainty)
2	Peacock and Jones (1948)	1948Pe13	104	Ionization chamber; <i>omitted</i> (no uncertainty)
3	Ramaswamy and Jastram (1960)	1960Ra20	105	Ionization chamber; <i>omitted</i> (no uncertainty)
4	Wyatt <i>et al.</i> (1961)	1961Wy01	108.1 (3)	4πγ ionization chamber; <i>omitted</i> (on the Chauvenet's criterion)
5	Anspach <i>et al.</i> (1965)	1965An07	106.52 (3)	4πγ ionization chamber; <i>omitted</i> (superseded by 13)
6	Anspach <i>et al.</i> (1965)	1965An07	106.67 (3)	4πγ ionization chamber; <i>omitted</i> (superseded by 13)

7	Grotheer <i>et al.</i> (1969)	1969Gr12	108.4 (9)	NaI(Tl)-detectors; <i>omitted</i> (on the Chauvenet's criterion)
8	Lagoutine <i>et al.</i> (1975)	1975La16	106.6 (4)	$4\pi\gamma$ ionization chamber; <i>omitted</i> (superseded by 18)
9	Bormann <i>et al.</i> (1976)	1976Bo19	107.1 (14)	$4\pi\gamma$ ionization chamber
10	Konstantinov <i>et al.</i> (1977)	1977Ko**	107.15 (65)	$4\pi\gamma$ ionization chamber
11	Houtermans <i>et al.</i> (1980)	1980Ho17	106.612 (50)	$4\pi\gamma$ ionization chamber; original uncertainty of 0.014 has been increased to 0.050 to ensure that the "weight" of this measurement would not exceed of the "weights" (26%) of two most recent values (18 and 20).
12	Debertin <i>et al.</i> (1982)	1982DEYX	106.64 (8)	Ge(Li) detector; <i>omitted</i> (superseded by 14)
13	Hoppes <i>et al.</i> (1982)	1982HOZJ	106.64 (5)	$4\pi\gamma$ ionization chamber; <i>omitted</i> (superseded by 15)
14	Walz <i>et al.</i> (1983)	1983Wa26	106.66 (6)	$4\pi\gamma$ ionization chamber
15	Unterweger <i>et al.</i> (1992)	1992Un01	106.626 (44)	$4\pi\gamma$ ionization chamber; <i>omitted</i> (superseded by 17)
16	Martin <i>et al.</i> (1997)	1997Ma75	106.65 (13)	$4\pi\gamma$ ionization chamber
17	Unterweger (2002)	2002Un02	106.63 (4)	4π ionization chamber; <i>omitted</i> (superseded by 19)
18	Amiot <i>et al.</i> (2005)	2005Am03	106.63 (5)	$4\pi\gamma$ ionization chamber
19	Fitzgerald (2012)	2012Fi12	106.62 (4)	$4\pi\gamma$ ionization chamber; <i>omitted</i> (superseded by 20)
20	Unterweger and Fitzgerald (2014)	2014Un01	106.63 (5)	$4\pi\gamma$ ionization chamber
Recommended value			106.63 (5) d	LWM

Values 1-3 from very early measurements (1940-1960) have been omitted as they are much less accurate. Values 5, 6, 8, 12, 13, 15, 17, 19 were not used because they were replaced ultimately by later results of the same laboratory.

Values 4, 7 have been rejected by the LWEIGHT computer program based on the Chauvenet's criterion. An unweighted average of the remained seven values is 106.78 (9) d. A weighted average is 106.632 d. The LWEIGHT program using the limitation of relative statistical weight method (LWM) has chosen the weighted average with the internal uncertainty of 0.025 d. The external uncertainty is 0.011 d. The ratio of the reduced $\chi^2 / (\chi^2)_{\text{crit}}$ is 0.2/2.8. The smallest experimental uncertainty of 0.05 d has been adopted for the uncertainty of the recommended value.

Thus, the recommended value of ^{88}Y half-life is **106.63 (5) days**. It can be compared with an earlier evaluation of 106.626 (21) (2004BeZR).

The measurements with the largest contributions to the weighted average are listed in Table 1a.

Table 1a. The measured values having the largest contributions to the weighted average of ^{88}Y half-life.

	Houtermans <i>et al.</i> (1980)	Walz <i>et al.</i> (1983)	Amiot <i>et al.</i> (2005)	Unterweger and Fitzgerald (2014)
T_{1/2} (d)	106.612 (50)	106.66 (6)	106.63 (5)	106.63 (5)
Weight	26%	18%	26%	26%

2.1. Electron Capture and β^+ Transitions

The electron capture and the β^+ emission probabilities, P_{e^+} and P_{β^+} , have been deduced from the transition intensity balance for ^{88}Sr each level (Table 2). The energies, spins, parities and half-lives of levels have been adopted from 2014Mc01.

Table 2. ^{88}Sr levels populated in the ^{88}Y decay

Level	Energy (keV)	Spin and parity	Half-life	P_{e^+} (%)	P_{β^+} (%)
0	0	0+	Stable	-	-
1	1836.090 (8)	2+	0.154 (8) ps	5.7 (3)	0.21 (1)
2	2734.137 (8)	3-	0.70 (5) ps	94.3 (3)	-
3	3218.489 (22)	2+	0.154 (10) ps	0.023 (4)	-
4	3584.784 (19)	5-	0.14 (4) ns	0.048 (18)	-

A positron transition to the ground state was not observed. However, sufficient energy for a positron transition is available for a transition to the 1836 keV level. The emission probability of these positrons was determined to be 0.203 (16) % by Barkov *et al.* (1974BaYZ) and that agrees well with the adopted value, 0.21 (1) %, calculated from the theoretical EC/ β^+ ratio of 25.6 (8) for an unique first forbidden transition interpolated from the table of Gove and Martin (1971Go40).

The maximum beta energy of the β^+ spectrum was found by Antonewa *et al.* (1979An36) to be 764.6 (15) keV corresponding to a Q value of 3622.6 (15) keV.

The $\log ft$ values, the fractional atomic shell electron capture probabilities and the average β^+ energy have been calculated with the LOGFT code.

2.2. Gamma Transitions and Internal Conversion Coefficients

The gamma-ray transition probabilities $P_{\gamma+ce}$ were calculated from the gamma ray emission probabilities, the total conversion coefficients, and the adopted internal pair creation coefficients.

Table 3. Comparison of the theoretical and experimental internal conversion coefficients for two most intense gamma-lines.

	$\alpha_K \times 10^{-4}$	$\alpha_T \times 10^{-4}$	K/L+M+...	
898 keV	2.73 (4)	3.07 (5)	8.01 (16)	Theory, BrIccFO adopted value
	E1(+M2)			
	$\delta = -0.002$ (9)			
	2.5 (3)	2.8 (3)	8.0 (2)	1971Al06
	3.01 (21)	3.45 (24)	7.0 (5)	1966Ha07
	3.4 (7)		1952Me50	
1836 keV	1.45 (2)	1.63 (2)	7.80 (16)	Theory, BrIccFO adopted value
	E2			
		1.7 (4)		1971Al06
	1.24 (16)	1.40 (16)	7.8 (3)	1966Ha07

Adopted ICC(s) are theoretical values interpolated by the BrIcc computer program based on BrIccFO approximation (2008Ki07). The multipolarities and mixing ratios δ have been taken from 2014Mc01 (Adopted Levels and Gammas). The mixing ratio parameter for the 898 keV transition was evaluated by Müller (1988) in 1988Mu09 to be $\delta = -0.002$ (9), i.e. this transition is an almost pure E1 transition.

The adopted internal pair creation coefficients, α_π , have been calculated with the BrIcc computer program. The experimental values were determined by Allan (1971Al06) as follows: $\alpha_\pi = 0.00023$ (3) for gamma-ray with energy of 1836 keV and $\alpha_\pi = 0.00033$ (5) for gamma-ray with energy of 2734 keV and can be compared with the adopted theoretical values of 0.000230 (4) and 0.000440 (7), respectively.

3. ATOMIC DATA

SAISINUC software has been used to determine the atomic data (fluorescence yields, X-ray energies and relative probabilities, and Auger electrons energies and relative probabilities).

4. ELECTRON AND POSITRON EMISSIONS

The energies of the conversion electrons have been obtained from the gamma-ray transition energies and the electron binding energies.

The absolute emission probabilities of the conversion electrons have been deduced using recommended P_{γ} and ICC values.

The absolute emission probabilities of K and L Auger electrons have been calculated using the EMISSION computer program [2000Sc47].

The number of electron-positron pairs per 100 disintegrations have been obtained using the adopted α_{π} values.

β^{+} average energies have been calculated using the LOGFT computer program.

5. PHOTON EMISSIONS

5.1. X-ray Emissions

The absolute emission probabilities of Sr KX- and LX- rays have been calculated using the EMISSION computer program [2000Sc47]. The adopted total KX-ray emission probability of 60.5 (5) % can be compared with the measured value of 61.5 (6) % (1994Ko34).

5.2. Gamma ray emissions

The energies of gamma rays in ^{88}Sr have been calculated from the level energies taking into account the recoil energy.

The measured relative intensities and the adopted values obtained with the LWEIGHT code are listed in Table 4 (see also 2004BeZR).

Table 4. Experimental and adopted relative gamma-ray intensities in decay of ^{88}Y

N		$\gamma_{850.6}$	$\gamma_{898.0}$	$\gamma_{1382.4}$	$\gamma_{1836.1}$	$\gamma_{2734.1}$	$\gamma_{3218.4}$
1	Peelle (1960Pe23)	-	94.0 (7)	-	100	0.597 (25)	-
2	Shastry and Bhattacharyya (1964Sh16)	-	91*	3*	100	0.97*	0.03*
3	Sakai <i>et al.</i> (1966Sa08)	-	-	-	100	0.63 (4)	0.0095 (3)*
4	Schötzig <i>et al.</i> (1973Sc40)	-	94.9 (5)*	-	100	-	-
5	Ardisson <i>et al.</i> (1974Ar12)	0.066 (13)	92.0 (7) ^a	0.021 (6)	100	0.724 (70)	0.0071 (20)
6	Heath (1974HeYW)	-	92.1*	-	100	0.54 (9)	0.007*
7	Antoneva <i>et al.</i> (1979An36)	0.030 (4)	92.2 (37)	0.014 (3)	100	-	-

N		$\gamma_{850.6}$	$\gamma_{898.0}$	$\gamma_{1382.4}$	$\gamma_{1836.1}$	$\gamma_{2734.1}$	$\gamma_{3218.4}$
8	Yoshizawa <i>et al.</i> (1980Yo05)	-	94.4 (3)	-	100	-	-
9	Debertin <i>et al.</i> (1982DeYX)	-	95.2 (5)* ^b	-	100	-	-
10	Hoppes <i>et al.</i> (1982HoZJ)	-	94.9 (4)	-	100	-	-
11	Schötzig (1990Sc08)	-	94.8 (9)	-	100	-	-
	Adopted	0.048 (18)	94.3 (3)	0.016 (3)	100	0.612 (25)	0.0071 (20)

*indicates the value omitted from the current analysis

^a adopted in 1974Ar12 from the measurement by Jardine (1971Ja21)

^b 1977 measurement result by Debertin *et al.*

The experimental values 2, 6 from Shastry and Bhattacharyya (1964Sh16) and Heath (1974HeYW), respectively, marked by asterisks were not taken into account because leak of uncertainties.

Adopted values:

- for **γ_{850} keV**: the evaluated value has been obtained by LWM with values 5 and 7. The LWEIGHT code has increased the original uncertainty of 1979An36 to 0.013 to ensure the relative weight of this measurement would not exceed 50%; the ratio of the reduced $\chi^2 / (\chi^2)_{\text{crit}}$ is 3.8/6.6.
- for **γ_{898} keV**: the values 2 and 6 have not been taken into account because of the lack of uncertainties and the values 4 and 9 are superseded by value 11. The value of the relative intensity of 92.2 (37) for this gamma line, referred to as Antoneva *et al.* (1979An36), has been calculated by the evaluators from the measured ratio $P(\text{ce}_K; \gamma_{1,0}) / P(\text{ce}_K; \gamma_{2,1}) = 0.576$ (20) in 1979An36 (where $P(\text{ce}_K; \gamma) = \alpha_K \times P\gamma$ is the probability of K-conversion electron emission) by using the current theoretical values of α_K . The evaluated value has been obtained by LWM with values 1, 5, 7, 8, 10 and 11. The ratio of the reduced $\chi^2 / (\chi^2)_{\text{crit}}$ is 2.8/3.0.
- for **γ_{2734} keV**: the evaluated value has been obtained by LWM with values 1, 3, 5 and 6. The ratio of the reduced $\chi^2 / (\chi^2)_{\text{crit}}$ is 1.2/3.8.
- for **γ_{3522} keV**: upper limit of $P\gamma$ (3522 keV)/ $P\gamma$ (1836 keV) < 0.001 has been taken from Ardisson *et al.* (1974Ar12);
- for **γ_{484} keV**: upper limit of $P\gamma$ (484 keV)/ $P\gamma$ (1836 keV) < 9×10^{-4} has been taken from Antoneva *et al.* (1979An36).

The normalization factor (**0.99346 (25)**) to convert the adopted relative gamma ray intensities to absolute emission probabilities has been deduced from the gamma ray transition

intensity balance for the ground state assuming no direct electron capture or beta plus feeding to the ground state.

5. ENERGY CONSERVATION

The total average energy of 3622 (7) keV per decay, calculated from the current evaluated data, corresponds well to the available energy of 3622.6 (15) keV (Q) from the atomic mass tables (2012Wa38), confirming the consistency of the decay scheme and the reliability of this evaluation.

6. REFERENCES

- 1940Du09** L. A. DuBridge and J. Marshall, Phys. Rev. 58 (1940) 7. [T_{1/2}]
- 1948Pe13** W. C. Peacock, J. W. Jones, Report AECD, 1812 (1948). [P_{β+}, T_{1/2}]
- 1952Me50** F.R.Metzger, H.C.Amacher, Phys.Rev. 88, 147 (1952). [Experimental ICC]
- 1960Pe23** R. W. Peelle, report ORNL, 3016 (1960), 110. [P_γ]
- 1960Ra20** M. K. Ramaswamy and P. S. Jastram, Nucl. Phys. 19 (1960) 243. [T_{1/2}]
- 1961Wy01** E. I. Wyatt, S. A. Reynolds, T. H. Handley, W. S. Lyon, H. A. Parker, Nucl. Sci. Eng. 11, 74 (1961). [T_{1/2}]
- 1964Sh16** S. Shastry and R. Bhattacharyya, Nucl. Phys. 55 (1964) 397. [P_γ]
- 1965An07** S. C. Anspach, L. M. Cavallo, S. B. Garfinkel, J. M. R. Hutchinson, C. N. Smith, Report NP-15663 (1965). [T_{1/2}]
- 1966Sa08** M. Sakai, T. Yamazaki, J. M. Hollander, Nucl. Phys. 84 (1966) 302. [P_ε, P_γ]
- 1966Ha07** J. H. Hamilton, S. R. Amtey, B. van Nooijen, A. V. Ramayya, J. J. Pinajian, Phys. Lett. 19 (1966) 682. [Experimental ICC]
- 1969Gr12** H. H. Grotheer, J. W. Hammer, K. W. Hoffmann, Z. Phys. 225 (1969) 293. [P_{β-}]
- 1971Al06** C. J. Allan, Nucl. Instrum. Methods 91 (1971) 117. [α_π]
- 1971Go40** N. B. Gove and M. J. Martin, Nuclear Data Tables 10 (1971) 205. [P_ε/P_{β+}]
- 1971Ja21** L. J. Jardine, Nucl. Instrum. Methods 96 (1971) 259. [P_γ]
- 1973Sc40** U. Schötzig, K. Debertin, H.M. Weiss, PTB Mitt. 83, 307 (1973) [P_γ]
- 1974Ar12** G. Ardisson, S. Laribi, C. Marsol, Nucl. Phys A223 (1974) 616. [P_γ]
- 1974BaYZ** A. V. Barkow, W. M. Winogradow, A. W. Solotawin, W. M. Makarow, T. M. Usypko, Program and abstracts of 24. Conference on nuclear spectroscopy and nuclear structure, Kharkov, 29 Jan. -1 Feb.1974, AN SSSR, Moscow, 1974, 58. [β⁺, α_π]
- 1974HeYW** R. L. Heath, Gamma-ray Spectrum Catalogue, USAEL, Rep. ANCR 1000-2 (1974). [E_γ, P_γ]
- 1975La16** F. Lagoutine, J. Legrand, C. Bac, Int. J. Appl. Radiat. Isot. 26 (1975) 131. [T_{1/2}]
- 1976Bo19** M. Bormann, H.-K. Feddersen, H.-H. Hölscher, W. Scobel and H. Wagner, Z. Physik A277 (1976) 203. [T_{1/2}]
- 1977Ko**** A. A. Konstantinov, T. E. Sazonowa and S. W. Sepman, Conference: 27. Annual conference on nuclear spectroscopy and nuclear structure, Tashkent, USSR, 22-25 March 1977, p.6. [T_{1/2}]

- 1979An36** N. M. Antoneva, V. M. Vindogradov, E. P. Grigorev, P. P. Dmitriev, A. V. Zolotavin, G. S. Katykhin, N. Krasnov, V. N. Makarov, Bull. Ac. Sci.USSR, Phys. Ser. 43 (1979) 155. [P β^+ , α_π , P γ]
- 1980Ho17** H. Houtermans, O. Milosevic, F. Reichel, Int. J. Appl. Radiat. Isot. 31 (1980) 153. [T $_{1/2}$]
- 1980Yo05** Y. Yoshizawa, Y. Iwata, T. Kaku, T. Katoh, J. Ruan, T. Kojima, Y. Kawada, Nucl. Instr. and Meth.174 (1980) 109. [P γ]
- 1982DeYX** K. Debertain, U. Schötzig, K. F. Walz, National Bureau of Standards, Washington, D.C., Report NBS-SP- 626 (1982) 101. [T $_{1/2}$, P γ]
- 1982HoZJ** D. D. Hoppes, J. M. R. Hutchinson, F. J. Schima, M. P. Unterweger, National Bureau of Standards, Washington, D.C., Report NBS-SP-626 (1982) 85. [T $_{1/2}$, P γ]
- 1983Wa26** K. F. Walz, K. Debertain and H. Schrader, Int. J. Appl. Radiat. Isot. 34 (1983) 1191. [T $_{1/2}$]
- 1988Mu09** H.-W. Muller, Nucl. Data Sheets 54 (1988) 1.
[Gamma-ray multipolarities and mixing ratios]
- 1990Sc08** U. Schötzig, Nucl. Instrum. Methods Phys. Res. A286 (1990) 523. [P γ].
- 1992Un01** M.P. Unterweger, D.D. Hoppes, F.J. Schima, Nucl. Instrum. Methods Phys. Res. A312 (1992) 349. [T $_{1/2}$]
- 1994Ko34** A. A. Konstantinov, T. E. Sazonova, S. V. Sepman, A. V. Zanevsky, N. I. Karmalitsyn, Nucl. Instrum. Methods Phys. Res. A339 (1994) 200. [P $\kappa\kappa$]
- 1997Ma75** R. H. Martin, K. I. W. Burns, J. G. V. Taylor, Nucl. Instrum. Methods Phys. Res. A390 (1997) 267. [T $_{1/2}$]
- 2000Sc47** E. Schönfeld and H. Janßen. Appl. Radiat. Isot. 52 (2000) 595.
[X-ray and Auger electron emission probabilities]
- 2002Un02** M.P. Unterweger, Appl. Radiat. Isot. 56, 125 (2002). [T $_{1/2}$]
- 2004BeZR** M.-M. Bé, V. Chiste, C. Dulieu, E. Browne, V. Chechev, N. Kuzmenko, R. Helmer, A. Nichols, E. Schönfeld, and R. Dersch, *Table of Radionuclides (Vol. 1 – A = 1 to 150)*. Bureau International des Poids et Mesures, 2004. ⁸⁸Y. P. 153-160. [Previous ⁸⁸Y decay data evaluation by DDEP collaboration]
- 2005Am01** M. N. Amiot, J. Bouchard, M.-M. Bé, J. B. Adamo, Appl. Rad. Isotop. 62 (2005) 11. [T $_{1/2}$]
- 2008Ki07** T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor Jr., Nucl. Instrum. Methods Phys. Res. A589 (2008) 202.
[BrIcc computer program]
- 2012Fi12** R. Fitzgerald. J. Res. Natl. Inst. Stand. Technol. 117 (2012) 80. [T $_{1/2}$]
- 2012Wa38** M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer, Chin. Phys. C36 (2012) 1603. [Q]
- 2014Mc01** E. A. McCutchan, A. A. Sonzogni, Nucl.Data Sheets 115, 135 (2014).
[Decay scheme structure, ⁸⁸Sr adopted levels and gammas, gamma-ray multipolarities]
- 2014Un01** M.P. Unterweger, R. Fitzgerald, Appl. Radiat. Isot. 87 (2014) 92. [T $_{1/2}$]

⁸⁹Sr – Comments on evaluation of decay data by E. Schönfeld

This evaluation was completed by E. Schönfeld (PTB) in November 1999.
The half-life evaluation was up-dated by M.-M. Bé (LNHB) in November 2002.

1 Decay Scheme

Below the Q -value there are no other levels of ⁸⁹Y. Thus, the decay scheme is complete. Spins and parities of the levels and $\lg ft$ values are taken from Sievers (1989). The half -life of the isomeric level at 909 keV was determined by Yule (1967) to be 16,06(4) s and by Durrani and Köhler (1966) to be 15,91 (17) s. The weighted mean is 16,05 (4) s. Earlier determinations were carried out by Swann and Metzger (1955) and Sattler (1962). The excited levels of ⁸⁹Y were studied by Robinson *et al.* (1969).

2 Nuclear Data

For the half-life evaluation the following measurements, carried out since 1954, were considered ($T_{1/2}$ in d):

Reference	Value (days)	Uncertainty	Comments
Herrmann (1954)	50,4	0,5	Superseded by the 2 nd value
Herrmann and Strassmann (1955)	50,5	0,2	
Kjelberg and Papas (1956)	51	1	Omitted, outlier
Osmond and Overs (1959)	50,36	0,18	
Sattler (1952)	53,6	0,4	Omitted, outlier
Marsden and Yaffee	50		Omitted, no uncertainty
Flynn <i>et al.</i>	52,7	0,5	Omitted, outlier
Anspach <i>et al.</i> (1965)	50,70	0,19	
Anspach <i>et al.</i> (1965)	50,52	0,04	Original uncertainty = 0,03
Baba <i>et al.</i> (1971)	50,55	0,09	
Lagoutine <i>et al.</i> (1972)	50,75	0,25	Superseded by Amiot
Amiot <i>et al.</i> (2003)	50,65	0,05	
Recommended value	50,57	0,03	Weighted mean

Four values have been omitted from the analysis, the uncertainty on the second Anspach value (Anspach *et al.*, 1965) has been multiplied by 1,33 in order to reduce its relative weight to 50 % in the calculation of the weighted mean and because it seems optimistic when compared with the other data. The set of six values taken into account in this analysis has a reduced $-\chi^2$ of 1,2. Finally, the adopted value (half -life, uncertainty) is the weighted mean and the external uncertainty.

The Q -value is taken from Audi and Wapstra (1995).

2.1 b- Transitions

The shape of the unique 1st forbidden β spectrum of ⁸⁹Sr was measured by Wohn and Talbert (1970). They found the end-point energy to be 1488(4) keV. The shape corrected $\lg ft$ was calculated by these authors to be 8,36. Earlier, the maximum beta end-point energy was determined to be 1463(5) keV by Bisi *et al.* (1955). This value is too small compared with the result of Wohn and Talbert and the larger value taken from the compilation of Audi and Wapstra (1995) which is the here adopted one.

Internal bremsstrahlung accompanying the first forbidden beta decay of ^{89}Sr was measured by Babu et al. (1987), Sayibaba et al. (1987), Basha et al. (1991) and Dhaliwal et al. (1994). Sayibaba et al. carried out their measurements with a HPGe detector and a multichannel analyzer along with a standard geometrical set-up. Their results are satisfactorily accounted for by the KUB theory. Basha et al. compared also their measurements with the theoretical spectra. Dhaliwal et al. measured the spectra using an extrapolation procedure with a beta stopper method. Their results are in agreement with the Lewis and Ford theory in the whole energy region covered by the present measurements and do not favour the KUB and Nilsson theories beyond a photon energy of 400 keV.

2.2 Gamma Transition

The energy of the gamma rays following the ^{89}Sr β^- decay was measured by Merritt et al. (1982) to be 909,12(7) keV whereas Sievers gives 908,96(4) keV as unweighted average from several (n, γ)-reactions and from the decay of ^{89}Zr ($T_{1/2} = 78,4$ h). In the present evaluation 909,0(1) keV is adopted. The transition probability of the gamma transition is calculated from the gamma ray emission probability of the 909 keV transition (see section 4.2) and the conversion coefficient of this transition. The conversion coefficients are interpolated from the tables of Rösel et al. (1978).

3 Atomic Data

The atomic data are taken from Schönfeld and Janßen (1996).

3.1 X Radiation

The energies are based on the wavelengths of Bearden (1967). The relative probabilities are taken from Schönfeld and Janßen (1996).

3.2 Auger Electrons

The energies of the Auger electrons are taken mainly from Larkins (1977). The ratios $P(\text{KLX})/P(\text{KLL})$ and $P(\text{KXY})/P(\text{KLL})$ are taken from Schönfeld and Janßen (1996).

4 Radiation Emission

4.1 Electron Emission

The energies and emission probabilities of the β^- particles correspond to the data given already in Section 2.1. The number of conversion electrons per disintegration has been calculated using the gamma ray emission probability P_γ and the conversion coefficient as given in Section 2.2. The emission probabilities of the Auger electrons have been calculated with the PTB program EMISSION using the atomic data as given in Section 3.

4.2 Photon Emissions

The gamma ray emission intensity, per one disintegration, was found to be:

1	$9,71(24) \cdot 10^{-5}$	Merritt et al. (AECL)	1980	replaced by value 3
2	$9,65(29) \cdot 10^{-5}$	Hoppes et al. (NBS)	1980	
3	$9,54(16) \cdot 10^{-5}$	Merritt et al. (AECL)	1982	
4	$9,61(13) \cdot 10^{-5}$	Schötzig (PTB)	1990	
5	$9,555(34) \cdot 10^{-5}$	Schima (NIST)	1998	
6	$9,56(6) \cdot 10^{-5}$	adopted value	1999	

Value 1 is replaced by value 3, value 6 is the LWM of values 2, 3, 4 and 5. The reduced χ^2 of this set is 0,19.

The emission probabilities of K-X rays are very small. This is caused by the small values of $P_{\gamma+ce}$ and α_K . Lyon and Rickard (1955) were the first who detected these weak gamma rays.

The number of emitted KX rays due to K-shell internal-ionization probabilities in nuclear beta decay were measured in comparison to the absolute beta decay rate by Hansen and Parthasaradhi (1974). Their experimental

result is $8,6 (7) 10^{-4}$ quanta per decay. The contribution of K conversion of the 909 keV γ -transition is only $5,1 10^{-7}$ per decay.

5 Main Production Modes

The production mode are taken from Sievers (1989).

6 References

- G. Herrmann, *Z. f. Elektrochemie* 58 (1954) 626
 [$T_{1/2}$]
 W. S. Lyon and R. R. Rickard, *Phys. Rev.* 100 (1955) 112
 [$T_{1/2}$]
 C. P. Swann and F. R. Metzger, *Phys. Rev.* 100 (1955) 1329
 [$T_{1/2}$ isomeric level]
 A. Bisi, S. Terrani and L. Zappa, *Il Nuovo Cimento II* (1955) 1297
 [E_{β}]
 G. Herrmann and F. Strassmann, *Z. Naturforschg.* 10A (1955) 146
 [$T_{1/2}$]
 A. Kjelberg and A. C. Papas, *Nucl. Phys.* 1 (1956) 322
 [$T_{1/2}$]
 R. G. Osmond, N. J. Owers, *J. inorg. Nucl. Chem.* 9 (1959) 96
 [$T_{1/2}$]
 A. R. Sattler, *Nucl. Phys.* 36 (1962) 648
 [$T_{1/2}$]
 D. A. Marsden and L. Yaffee, *Can. J. Chem.* 43 (1965) 249
 [$T_{1/2}$]
 K. F. Flynn, L. G. Glendenin and E. E. Steinberg, *Nucl. Sci. Eng.* 22 (1965) 416
 [$T_{1/2}$]
 S.C. Anspach, L.M. Cavallo, S.B. Garfinkel, J.M.R. Hutchinson, C.N. Smith, *NP-15663* (1965)
 [$T_{1/2}$]
 J. A. Bearden, *Rev. Mod. Phys.* 39 (1967) 78
 [$E(KX)$]
 E. L. Robinson, R. C. Hagenauer and E. Eichler, *Nucl. Phys.* A123 (1969) 471
 [E_{γ} , P_{γ}]
 F. K. Wohn and W. L. Talbert, Jr., *Nucl. Phys.* A146 (1970) 33
 [E_{β}]
 S. Baba, H. Baba and H. Natsume, *J. Inorg. Nucl. Chem.* 33 (1971) 589
 [$T_{1/2}$]
 F. Lagoutine, J. Legrand, C. Perrot, J. P. Brethon and J. Morel, *Int. J. Appl. Rad. Isot.* 23 (1972) 219
 [$T_{1/2}$]
 H. H. Hansen and K. Parthasaradhi, *Phys. Rev.* C9 (1974) 1143
 [P_{XK}]
 F. P. Larkins, *Atomic Data and Nuclear Data Tables* 20 (1977) 313
 [$E(KLL)$, $E(KLX)$]
 F. Rösler, H. M. Fries, K. Alder, H. C. Pauli, *At. Data Nucl. Data Tables* 21 (1978) 91
 [α_t]
 J. S. Merritt, K. M. Ophel, A. R. Rutledge and L. V. Smith, *AECL 7102* (1980) 32
 [P_{γ}]
 J. S. Merritt, A. R. Rutledge and L. V. Smith, *Int. J. Appl. Radiat. Isot.* 33 (1982) 77
 [E_{γ} , P_{γ}]
 T. Sayibaba, K. Narasimha Murty and C. R. Rao, *Il Nuovo Comento*, 97A (1987) 365
 [Bremsstrahlung]
 B. R. S. Babu, P. Venkataramaiah and H. Sanjeeviah, *Nucl. Instr. Meth.* A255 (1987) 96
 [Bremsstrahlung]
 H. Sievers, *Nuclear Data Sheets* 58 (1989) 351
 [production modes, spins, parities]
 U. Schötzig, *Nucl. Instr. Methods* A286 (1990) 523
 [P_{γ}]

- A. M. Basha, E. I. Khalil, M. Hussein and H. Ragab, *Indian J. Phys.* 65A (1991) 120
[Bremsstrahlung]
- A. S. Dhaliwal, M. S. Powar and M. Singh, *J. Phys. G Nucl. Part. Phys.* 20 (1994) 135
[Bremsstrahlung]
- G. Audi, A. H. Wapstra, *Nucl. Phys.* A595 (1995) 409
[Q]
- E. Schönfeld and H. Janßen, *Nucl. Instr. Methods* A369 (1996) 527
[ω_K , ω_L , K_β/K_α , n_{KL}]
- F. J. Schima, *Appl. Radiat. Isot.* 49 (1997) 1359
[P_γ]
- M. N. Amiot, J. Bouchard, M.-M. Bé, J. B. Adamo. To be published (2004)
[$T_{1/2}$]

**⁸⁹Zr – Comments on evaluation of decay data
by A. L. Nichols**

Evaluated: May/June 2014

Evaluation Procedures

Limitation of Relative Statistical Weight Method (LWM) and other analytical techniques were applied to obtain averaged data throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

⁸⁹Zr ($T_{1/2} = 78.42$ h) undergoes 100 % electron capture/ β^+ decay ($Q(\text{EC}) = 2832.8$ (28) keV) to various excited nuclear levels and the ground state of ⁸⁹Y. A reasonably well-defined decay scheme was derived from the gamma-ray measurements of [1968Hi12](#), [1969Ro02](#), [1971Ar18](#) and [1979Ba46](#), consisting of five EC transitions, one β^+ emission and five gamma rays. Weighted mean relative emission probabilities were calculated for the gamma rays. All of these relative emission probabilities have been defined in terms of the 871.094-keV gamma ray (100 %).

Nuclear Data

⁸⁹Zr decay is viewed as a potentially suitable β^+ emitter for application in positron emission tomography (PET).

Half-life of ⁸⁹Zr

The measurements of [1940Du05](#), [1953Sh48](#), [1960Ha26](#), [1961Ra06](#), [1964Va03](#), [1969Ro02](#) and [1984Sk01](#) were adopted to give a weighted mean half-life of 78.42 (13) hours based on the limitation of relative statistical weight method (LWM).

Reference	Half-life (h)
1938Sa01	70*
1940Du05	78 (1)
1940Sa08	68 (2)†
1951Hy24	77*
1951Sh24	79.3*
1953Ka11	78*
1953Sh48	79 (2)
1960Ha26	79.0 (5)
1961Ra06	79.4 (16)
1962Ho10	79*
1964Va03	78.43 (8)†
1969Ro02	78.0 (2)
1984Sk01	78.62 (17)
Recommended value	78.42 (13)#

* Uncertainty unspecified – not included in weighted mean analysis of the data set.

† Defined as an outlier, and therefore not adopted in the analytical procedures.

‡ Uncertainty increased from ± 0.08 to 0.13 to reduce weighting below 0.50.

Recommended uncertainty has been adjusted from ± 0.09 to 0.13, in alignment with the smallest uncertainty of the values used to calculate the average value.

A small change has been observed in the half-life of ⁸⁹Zr when bound at the lattice site of the Ti ion in BaTiO₃ as the crystal undergoes a phase transition at a temperature (T_c) of 120 °C ([1970Ga03](#)). This experimental study has been further assessed in terms of modelling the impact of changes in the chemical bonding and ligand field strength of the BaTi(Zr)O₃ lattice ([1973Le13](#)). Theoretical estimates of the relative change of the decay constant quantified in terms of the expression $\Delta\lambda_{\text{EC}}/\lambda_{\text{EC}}$ (where $\lambda = \ln 2/t_{1/2}$) were of the order of 0.0015 and 0.0065 compared with the experimental observations of 0.00080 (3). Measured data represent observed changes in the EC-decay half-life of ⁸⁹Zr of the order of ~ 0.08 % (no more than 0.06 h).

Half-lives of ^{89m}Y and ^{89m}Zr metastable states

The measurements of 1955Sw92, 1962Br42, 1966Du07, 1967Yu01, 1968Bo52 and 1995ItZY were adopted to give a weighted mean half-life for ^{89m}Y of 15.84 (18) seconds, based on the limitation of relative statistical weight method (LWM).

Reference	^{89m} Y half-life (s)
1951Go42	14 (2) [‡]
1955Sw92	16.1 (3)
1962Br42	16.3 (13)
1966Du07	15.91 (17)
1967Yu01	16.06 (4)
1968Bo52	15.78 (11)
1995ItZY	15.663 (5) [†]
Recommended value	15.84 (18)[#]

[‡] Defined as an outlier, and therefore not adopted in the analytical procedures.

[†] Uncertainty increased from ± 0.005 to 0.04 to reduce weighting to 0.50.

[#] Weighted average adopted, with the recommended uncertainty increased to include the precise value of 15.66 (4) s.

The measurements of 1951Sh89, 1953Sh48, 1964Va03, 1969Ro02 and 1992KaZM were adopted to give a weighted mean half-life for ^{89m}Zr of 4.170 (19) minutes based on the limitation of relative statistical weight method (LWM).

Reference	^{89m} Zr half-life (min)
1940Du05	4.5 [*]
1951Sh89	4.4 (1)
1953Ka11	4.25 [*]
1953Sh48) 4.5 (5)
) 4.40 (4)
) 4.4 (3)
1964Va03	4.18 (1)
1969Ro02	4.16 (6)
1992KaZM	4.145 (9) [†]
Recommended value	4.170 (19)

^{*} Uncertainty unspecified – not included in weighted mean analysis of the data set.

[†] Uncertainty increased from ± 0.009 to 0.010 to reduce weighting to no more than 0.50.

Q-value

Q_{EC}-value for ⁸⁹Zr EC/β⁺ decay of 2832.8 (28) keV was adopted from Wang *et al.* (2012Wa38).

Gamma-ray energies and emission probabilities

Energies

The well-defined nuclear level energies of 2013Si03 were used to calculate the gamma transition energies and their uncertainties, and these data were adjusted to account for gamma recoil in the formulation of recommended gamma-ray emission energies and uncertainties. Greater confidence was placed on this overall method of assignment because of the more wide-ranging origins of the gamma transition data in 2013Si03.

Adopted energies, spins, parities and half-lives of the nuclear levels of ⁸⁹Y (2013Si03).

Nuclear level number	Nuclear level energy (keV)	Spin and parity	Half-life
0	0.0	1/2 –	stable ⁸⁹ Y
1	908.97 (3)	9/2 +	15.84(18) s ^{89m} Y
2	1744.74 (18)	5/2 –	0.62(14) ps [*]
3	2529.80 (20)	7/2 +	0.08(3) ps [†]
4	2566.55 (15)	11/2 +	
5	2622.1 (3)	9/2 +	0.21(10) ps [*]

^{*} Determined from Doppler shift attenuation measurements of the ⁸⁷Rb(α,2nγ)⁸⁹Y reaction by 1992Fu04.

[†] Determined from Doppler shift attenuation measurements of the ⁸⁹Y(p,p'γ)⁸⁹Y reaction by 1973BeYD.

Gamma transition energies, and measured and recommended gamma-ray energies.

E _{TP} (keV) [#]	E _γ (keV)										
	1953Sh48	1961Mo12	1964Va03	1968Dr02	1968Hi12	1969GuZV	1969Ro02	1971Ar18	1974HeYW	1992Fu04	Recommended [§]
–	–	511	511	511	511.0	511	–	–	511.006	–	annihilation radiation
908.97 (3)	920 (10)	915	908	908 (2)	909.1 (1)	909.1 (1)	909.2 (1)	908 (1)	909.14 (7)	909.1 (1)	908.97 (3)
1620.83 (20)	–	–	1621 (9)	1617 (2)	1620.6 (7)	–	1620.8 (2)	1620 (1)	1620.76 (19)	1620.9 (5)	1620.81 (20)
1657.58 (15)	–	–	–	1655 (2)	1656.9 (7)	–	1657.3 (2)	1657 (1)	1657.26 (19)	1657.5 (2)	1657.56 (15)
1713.1 (3)	–) 1750 (20)	1705 (4)	1712 (2)	1713.0 (6)	1713.4 (2)	1712.9 (8)	1712 (1)	1712.8 (4)	1713.3 (4)	1713.1 (3)
1744.74 (18)	–)	–	1744 (2)	1744.4 (7)	–	1744.5 (2)	1744 (1)	1744.47 (19)	1744.9 (3)	1744.72 (18)

[#] Determined from the nuclear level energies of 2013Si03.

[§] Calculated by subtracting gamma recoil from gamma transition energy (E_{TP} (keV)).

Measured and recommended gamma-ray emission probabilities relative to P_γ(908.97 keV) of 100 %.

E _γ (keV) [#]	P _γ ^{rel}									
	1961Mo12	1964Va03	1968Dr02 [*]	1968Hi12	1969GuZV [*]	1969Ro02	1971Ar18	1974HeYW [†]	1979Ba46	Recommended [‡]
γ [±] 511	44.5 (15)	44.2 (9)	–	44.3 (20)	0.47 → 47.5	–	–	45.1 (6)	–	46.0 (6) [§] annihilation radiation
γ _{1,0} 908.97 (3)	100	100	100	100	0.99 → 100	100	99.86 → 100	99.871 (3) → 100	100	100
γ _{3,1} 1620.81 (20)	–	0.12 (1)	0.077	0.068 (9)	–	0.071 (7)	0.08 (1)	0.072 (5)	0.075 (5)	0.075 (5) [#]
γ _{4,1} 1657.56 (15)	–	–	0.098	0.105 (5)	–	0.10 (1)	0.12 (1)	0.107 (4)	0.108 (5)	0.107 (5) [†]
γ _{5,1} 1713.1 (3)) 0.4 (1)	0.98 (3)	0.77	0.760 (10)	0.0067 → 0.68	0.77 (7)	0.86 (5)	0.763 (13) → 0.764 (13)	0.731 (13)	0.752 (10) ^Δ
γ _{2,0} 1744.72 (18))	–	0.096	0.127 (4)	–	–	0.14 (1)	0.129 (3)	0.118 (4)	0.124 (4) ⁰

^{*} Unquantified uncertainties – not included in weighted mean analysis of the data set.

[†] Described as obtained from the 1998 version of ENSDF when prepared as an electronic spectral database in 1998 - not included in weighted mean analysis of the data set.

[‡] Emission probabilities expressed relative to P_γ(908.97 keV) of 100 %.

[§] Relative emission probability of 46.0 (6) % for the 511-keV annihilation radiation was calculated from the total absolute positron emission probability of 22.8 (3) % and normalisation factor of 0.9903 (2).

[#] Recommended uncertainty has been adjusted from ± 0.004 to 0.005, in alignment with the smallest uncertainty of the values used to calculate the average value.

[†] Recommended uncertainty has been adjusted from ± 0.003 to 0.005, in alignment with the smallest uncertainty of the values used to calculate the average value.

^Δ Recommended uncertainty has been adjusted from ± 0.008 to 0.010, in alignment with the smallest uncertainty of the values used to calculate the average value.

⁰ Recommended uncertainty has been adjusted from ± 0.003 to 0.004, in alignment with the smallest uncertainty of the values used to calculate the average value.

Gamma-ray emissions: multiplicities, and theoretical internal-conversion (frozen orbital approximation) and internal-pair formation coefficients.

E _γ (keV)	Multipolarity	α _K	α _L	α _{M+}	α _{ICtotal}	α _{IPF}	α _{total}
908.97 (3)	100%M4	0.007 43 (11)	0.000 906 (13)	0.000 174	0.008 51 (12)	–	0.008 51 (12)
1620.81 (20)	M1 + E2	–	–	–	–	–	–
1657.56 (15)	M1 + E2	–	–	–	–	–	–
1713.1 (3)	M1 + E2 1984HaZC, 1985HaZI	–	–	–	–	–	–
1744.72 (18)	E2	0.000 172 2 (25)	0.000 018 6 (3)	0.000 003 6	0.000 194 4 (25)	0.000 188 (3)	0.000 382 (6)

Emission Probabilities

All of the relative emission probabilities were quantified in terms of the emission probability of the 908.97-keV gamma ray (100.0 %), and the gamma-ray measurements of [1968Hi12](#), [1969Ro02](#), [1971Ar18](#) and [1979Ba46](#) were used to derive weighted-mean relative emission probabilities.

The normalisation factor for the relative gamma-ray emission and EC/ β^+ transition probabilities can be calculated via two related routes:

(a). A total value of 100 % can be assigned to the gamma transition probabilities directly populating the 0.0-keV nuclear level of ⁸⁹Y, assuming zero EC/ β^+ decay directly to the ground state on the basis of spin-parity considerations (9/2+ \rightarrow 1/2- constitutes a third forbidden unique transition (4, yes)):

$$\sum_{0.0 \text{ keV}}^{\gamma \text{ population}} TP_{\gamma} = 100$$

Thus: $100.975 (13) F = 100$

where F is the normalisation factor for the relative gamma-ray emission probabilities.

$$F = 100 / 100.975 (13) = 0.99034 \pm 0.00013$$

(b). Gamma population-depopulation of the various nuclear levels of ⁸⁹Y provide a means of determining the relative EC/ β^+ transition probabilities from the relative γ -ray transition probabilities. Imbalances for each proposed nuclear level of Y-89 (at energies of 2622.1, 2566.5, 2529.8, 1744.7 and 908.97 keV) can be calculated on the basis of the evaluated relative gamma-ray emission probabilities, and their adopted total internal conversion coefficients. These population-depopulation imbalances in the relative gamma-transition probabilities constitute relative EC (EC + positron) transition probabilities of 0.752 (10) F, 0.107 (5) F, 0.075 (5) F, 0.124 (4) F and 99.917 (17) F to give a total of 100.975 (18) F in which the uncertainties of the EC_{0,2}, EC_{0,3}, EC_{0,4} and EC_{0,5} transitions are correctly included once in the summation. The total relative EC transition probability can be set to 100 % EC decay, assuming zero EC/ β^+ decay directly to the ground state based on spin-parity considerations (third forbidden unique):

$$\sum_0^{EC/\beta^+ \text{ populations}} TP_{EC} = 100$$

Thus: $100.975 (18) F = 100$

where F is the normalisation factor for the calculated relative EC/ β^+ transition probabilities (and relative gamma-ray emission probabilities).

$$F = 100 / 100.975 (18) = 0.99034 \pm 0.00018$$

On the basis of the above, a normalization factor of (0.9903 \pm 0.0002) was adopted for the relative gamma-ray emission and EC/ β^+ transition probabilities.

Multipolarities and Internal Conversion Coefficients

The nuclear level scheme specified by Singh has been used to define the multipolarities of the gamma transitions on the basis of known spins and parities ([2013Si03](#)). Only the significant emission probability of the 908.97-keV gamma ray merits the determination of the internal conversion coefficients for this 100 % M4 transition, although such data have also been derived for the 1744.72-keV E2 gamma-ray emission. Recommended internal conversion coefficients have been determined from the frozen orbital approximation of Kibédi *et al.* ([2008Ki07](#)), based on the theoretical model of Band *et al.* ([2002Ba85](#), [2002Ra45](#)), while the internal-pair formation coefficient for the 1744.72-keV E2 gamma transition was quantified from the tabulations of [2008Ki07](#).

Energies and emission probabilities of noteworthy internal conversion electrons.

		Energy (keV)	Electrons per 100 disint.
ec _{1,0 T}	(Y)	891.93 – 908.97	0.84 (3)
ec _{1,0 K}	(Y)	891.93 (3)	0.73 (3)
ec _{1,0 L}	(Y)	906.60 – 906.89	0.089 (3)
ec _{1,0 M+}	(Y)	908.58 – 908.97	0.017 (1)

Comments on evaluation

Recommended gamma-ray energies, relative and absolute emission probabilities, and transition probabilities

E_γ (keV)	P_γ^{rel} (%)	P_γ^{abs} (%)	Transition probability (%)
γ^\pm 511 annihilation radiation	46.0 (6) ^s	45.6 (6)	–
$\gamma_{1,0}$ 908.97 (3)	100	99.03 (2)	99.87 (2)
$\gamma_{3,1}$ 1620.81 (20)	0.075 (5)	0.074 (5)	0.074 (5)
$\gamma_{4,1}$ 1657.56 (15)	0.107 (5)	0.106 (5)	0.106 (5)
$\gamma_{5,1}$ 1713.1 (3)	0.752 (10)	0.745 (10)	0.745 (10)
$\gamma_{2,0}$ 1744.72 (18)	0.124 (4)	0.123 (4)	0.123 (4)

^s Relative emission probability of 46.0 (6) % for the 511-keV annihilation radiation was calculated from the total absolute positron emission probability of 22.8 (3) % and normalisation factor of 0.9903 (2).

EC/ β^+ transitions

Energies

All EC/ β^+ energies were derived from the structural details of the proposed decay scheme. The nuclear level energies of [2013Si03](#) and evaluated Q_{EC} -value of 2832.8 (28) keV ([2012Wa38](#)) were used to determine the recommended energies and uncertainties of the EC transitions and β^+ emissions.

Transition and Emission Probabilities

⁸⁹Zr undergoes 100 % EC/ β^+ decay. All resulting EC transition probabilities were derived from the population-depopulation imbalances of the relative emission probabilities of the gamma rays, their theoretical internal-conversion and internal-pair formation coefficients. However, an EC transition directly to the ground state of ⁸⁹Zr was assumed to be zero, based on spin-parity considerations alone (third forbidden unique (4, yes)). A value of 0.9903 (2) was adopted as the normalisation factor in order to determine the absolute transition and emission probabilities of the EC and β^+ particles from their resulting relative transition and emission probabilities. Component EC and β^+ transition and emission probabilities were determined from EC/ β^+ ratios ([1971Go40](#)), and $\log ft$ values and average E_{β^+} energies were derived by means of the LOGFT code. Fractional EC probabilities P_K , P_L , P_M and P_N were calculated by means of the EC-CAPTURE code ([1998Sc28](#)) as developed from the data tabulations of [1995ScZY](#).

Measurements by [1953Sh48](#), [1961Mo12](#), [1964Va03](#), [1968Hi12](#) and [1979Ba46](#) have been made to determine or provide a means of calculating the EC/ β^+ ratio for the 1924-keV EC transition. All relevant EC/ β^+ data for ⁸⁹Zr are summarised in the table below.

EC/ β^+ ratio for the 1924-keV EC transition of ⁸⁹Zr.

Reference	ϵ/β^+ (1924-keV EC transition)
Gove and Martin (1971Go40), theory	3.36 (5)
derived from 1953Sh48 , and quoted by Bambynek <i>et al.</i> (1977Ba48)	4 (1)
1961Mo12 , as adjusted and quoted by Bambynek <i>et al.</i> (1977Ba48)	3.54 (14) → 3.48 (15)
1964Va03 , and quoted by Bambynek <i>et al.</i> (1977Ba48)	3.43 (10)
derived from 1968Hi12 , and quoted by Bambynek <i>et al.</i> (1977Ba48)	3.47 (21)
Baillie <i>et al.</i> (1979Ba46)	3.634 (55)

The absolute emission probability of the 511-keV annihilation radiation was derived from the total absolute positron emission probability of 22.8 (3) % as determined during the evaluation of the EC/ β^+ transitions/emissions:

$$absolute P_\gamma^\pm = 2 \times 22.8 (3) = 45.6 (6) \%$$

$$[relative P_\gamma^\pm = \frac{absolute P_\gamma^\pm}{F} = \frac{45.6(6)}{0.9903(2)} = 46.0 (6) \%]$$

where F is the normalisation factor for the gamma-ray emission probabilities.

A decay scheme was derived that consists of five EC transitions, one β^+ emission and five gamma rays. Furthermore, 511-keV annihilation radiation was also observed, with an absolute emission probability of 45.6 (6) %.

Recommended energies and transition probabilities of the EC/ β^+ decay of ⁸⁹Zr.

E_{EC} (keV) *	E_{β^+} (keV)	Av. E_{β^+} (keV)	P_{EC} (total)	ϵ/β^+ (theory) 1971Go40	P_{EC}	P_{β^+}	⁸⁹ Zr	⁸⁹ Y	transition type	log ft	P_K	P_L	P_M	P_N
EC _{0,5} 211 (3)	–	–	0.745 (10)	–	0.745 (10)	–	9/2 +	9/2 +	allowed	6.18	0.8575 (17)	0.1165 (13)	0.0223 (5)	0.0036 (2)
EC _{0,4} 266 (3)	–	–	0.106 (5)	–	0.106 (5)	–	9/2 +	11/2 +	allowed	7.25	0.8615 (16)	0.1134 (13)	0.0216 (5)	0.0035 (2)
EC _{0,3} 303 (3)	–	–	0.074 (5)	–	0.074 (5)	–	9/2 +	7/2 +	allowed	7.52	0.8632 (16)	0.1120 (13)	0.0213 (4)	0.0035 (2)
EC _{0,2} 1088 (3)	–	–	0.123 (4)	–	0.123 (4)	–	9/2 +	5/2 –	first forbidden unique	9.09 ^{lu}	0.8677 (15)	0.1082 (12)	0.0208 (4)	0.0033 (2)
EC _{0,1} 1924 (3)	902 (3)	395.7 (14)	98.95 (2)	3.36	76.2 (3)	22.8 (3)	9/2 +	9/2 +	allowed	6.152	0.8731 (15)	0.1041 (12)	0.0196 (4)	0.0032 (2)
			Σ 99.998		Σ 77.248	Σ 22.8								
					Σ 100.048									

* Determined from the nuclear level energies of [2013Si03](#) and Q-value of 2832.8 (28) keV ([2012Wa38](#)).

Atomic Data

The X-ray and Auger-electron data have been calculated using the evaluated gamma-ray data, and atomic data from [1996Sc06](#), [1998ScZM](#) and [1999ScZX](#). Both the X-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.02, 28 February 2012), as described in [2000Sc47](#). This program incorporates atomic data from [1996Sc06](#), along with the adoption of a capture probability ratio P_{L1}/P_{L2} from [1997ScZY](#) and P_{L3} of zero, and the evaluated gamma-ray data.

K and L X-ray energies and emission probabilities of ⁸⁹Zr.

			Energy (keV)	Photons per 100 disint.	Relative probability
XL	(Y)		1.686 – 2.347	2.36 (5)	8.74
	XL ₁	(Y)	1.686	0.0568 (15)	
	XL _α	(Y)	1.920 – 1.923	1.48 (4)	
	XL _η	(Y)	1.762	0.0234 (7)	
	XL _β	(Y)	1.996 – 2.078	0.778 (19)	
	XL _γ	(Y)	2.153 – 2.347	0.01331 (24)	
XK _α	XK _{α2}	(Y)	14.8829 (1)	14.08 (13)	52.1
	XK _{α1}	(Y)	14.9585 (1)	27.01 (20)	100
XK' _{β1}	XK _{β3}	(Y)	16.7259 (7))	25.1
	XK _{β1}	(Y)	16.7381 (5)) 6.78 (8)	
	XK _{β5}	(Y)	16.880 (2))	
XK' _{β2}	XK _{β2}	(Y)	17.0156 (9)) 0.94 (4)	3.48
	XK _{β4}	(Y)	17.0362 (12))	

Auger-electron energies and emission probabilities of ⁸⁹Zr.

		Energy (keV)	Electrons per 100 disint.	Relative probability
e _{AK}	(Y)		19.4 (3)	
	KLL	12.205 – 12.784	13.72 (21)	100
	KLX	14.238 – 14.956	5.16 (9)	37.6
	KXY	16.251 – 17.034	0.484 (11)	3.53
e _{AL}	(Y)	1.27 – 1.89	79.5 (7)	579

Y: $\omega_K = 0.716$ (4); $\omega_L = 0.0289$ (7); $n_{KL} = 1.081$ (4) were taken from [1996Sc06](#).

Electron energies were determined from electron binding energies tabulated by Larkins ([1977La19](#)) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

Data Consistency

A Q_{EC} -value of 2832.8 (28) keV has been adopted from the atomic mass evaluation of Wang *et al.* ([2012Wa38](#)) for the EC decay of ⁸⁹Zr. This value has subsequently been compared with the Q-value calculated by summing the contributions of the individual emissions to the ⁸⁹Zr EC-decay process (i.e. EC/ β^+ , electron, γ , etc.):

$$\text{calculated Q-value} = \sum (E_i \times P_i) = 2834 (7) \text{ keV}$$

Percentage deviation from the Q-value of Wang *et al.* is 0.04 (28) %, which supports the derivation of a reasonably consistent decay scheme albeit with a significant variant.

References

- 1938Sa01** R. Sagane, S. Kojima, G. Miyamoto, M. Ikawa, Preliminary Report on the Radioactivity Produced in Y, Zr and Mo, Phys. Rev. 54 (1938) 542-543. [Half-life]
- 1940Du05** L.A. DuBridge, J. Marshall, Radioactive Isotopes of Sr, Y and Zr, Phys. Rev. 58 (1940) 7-11. [⁸⁹Zr and ^{89m}Zr half-lives]
- 1940Sa08** R. Sagane, S. Kojima, G. Miyamoto, M. Ikawa, Artificial Radioactivity Induced in Zr and Mo, Phys. Rev. 57 (1940) 1179-1180. [Half-life]
- 1951Go42** M. Goldhaber, E. der Mateosian, G. Scharff-Goldhaber, A.W. Sunyar, M. Deutsch, N.S. Wall, Isomeric State of Y⁸⁹ and the Decay of Zr⁸⁹, Phys. Rev. 83 (1951) 985-986. [Half-life, EC decay]
- 1951Hy24** E.K. Hyde, G.D. O'Kelley, Radiochemical and Spectrometer Studies of Several Neutron-Deficient Zirconium Isotopes and Their Decay Products, Phys. Rev. 82 (1951) 944-951. [Half-life, E_γ]
- 1951Sh24** K. Shure, M. Deutsch, Radiations from Zr⁸⁹, Phys. Rev. 82 (1951) 122. [Half-life]
- 1951Sh89** F.J. Shore, W.L. Bendel, R.A. Becker, 4.4-minute Radiations from Zr⁸⁹, Phys. Rev. 83 (1951) 688-689. [^{89m}Zr half-life]
- 1953Ka11** L. Katz, R.G. Baker, R. Montalbetti, The Photoneutron Cross Sections of Rb⁸⁷, Zr⁹⁰ and Mo⁹², Can. J. Phys. 31 (1953) 250-261. [⁸⁹Zr and ^{89m}Zr half-lives]
- 1953Sh48** F.J. Shore, W.L. Bendel, H.N. Brown, R.A. Becker, Radiations from Zr⁸⁹, Phys. Rev. 91 (1953) 1203-1211. [⁸⁹Zr and ^{89m}Zr half-lives, P_γ(909 keV)/P_{β+} ratio]
- 1955Sw92** C.P. Swann, F.R. Metzger, Production of Y^{89m}, Ba^{137m} and Hg^{199m} by Inelastic Neutron Scattering, Phys. Rev. 100 (1955) 1329-1333. [Half-life]
- 1957Ku57** M.I. Kuznetsova, V.N. Mekhedov, Procedure for Measuring the Activity of Nuclei Undergoing K-capture, Izvest. Akad. Nauk SSSR, Ser. Fiz. 21 (1957) 1020-1024. [K-capture]
- 1960Ha26** J.H. Hamilton, L.M. Langer, W.G. Smith, Evidence for Small Deviations in the Allowed Positron Spectrum of Zr⁸⁹, Phys. Rev. 119 (1960) 772-776. [Half-life]
- 1961Mo12** S. Monaro, G.B. Vingiani, R. van Lieshout, Aspects of the Decay of ²⁷Mg and ⁸⁹Zr, Physica 27 (1961) 985-993. [Half-life, E_γ, P_γ, EC/β⁺ ratio]
- 1961Ra06** L.A. Rayburn, 14.4-MeV (n,2n) Cross Sections, Phys. Rev. 122 (1961) 168-171. [Half-life]
- 1962Br42** E.T. Bramlitt, R.W. Fink, Absolute Activation Cross Sections for Reactions of Niobium with 14.5-MeV Neutrons, J. Inorg. Nucl. Chem. 24 (1962) 1321-1325. [^{89m}Y half-life]
- 1962Ho10** D.A. Howe, L.M. Langer, D. Wortman, Search for A 9/2⁺ Level at 1.5 MeV in Y⁸⁹, Nucl. Phys. 37 (1962) 476-481. [Gamma-ray emissions feeding 909-keV level]
- 1964Aw02** Y. Awaya, Y. Tendow, Decay of Zr⁸⁶ and Gamma Rays of Zr⁸⁷ and Zr⁸⁹, J. Phys. Soc. Japan. 19 (1964) 606-613. [1700(100)-keV γ ray]

- 1964Va03** D.M. van Patter, S.M. Shafroth, Decay of 78.4 h Zr⁸⁹ and 4.18 min Zr^{89m}, Nucl. Phys. 50 (1964) 113-135. [⁸⁹Zr and ^{89m}Zr half-lives, E_γ, P_γ, E_{EC}, P_{EC}, P_{β+}, EC/β⁺ ratio]
- 1966Du07** S.A. Durrani, W. Köhler, Average Cross Sections and Half-lives of Au-197m and Y-89m Excited by Inelastic Scattering of Reactor Neutrons, Trans. Am. Nucl. Soc. 9 (1966) 479-481. [^{89m}Y half-life]
- 1967Yu01** H.P. Yule, Half-lives of Some Short-Lived Radioactive Isotopes, Nucl. Phys. A94 (1967) 442-448. [^{89m}Y half-life]
- 1968Bo52** P. Bornemisza-Pausperl, P. Hille, Über (n,n'γ)-Reaktionen an ⁷⁹Br, ⁸⁹Y, ¹⁷⁶Yb und ¹⁹⁷Au im Zusammenhang mit der Frage des (n,2n)-Reaktionsmechanismus, Osterr. Akad. Wiss. Math-Naturw. Kl. Sitzber. Abt. II, 176 (1968) 227-243. [^{89m}Y half-life]
- 1968Dr02** J.E. Draper, J.A. McCray, Levels in ⁸⁹Y Populated by Decay of ⁸⁹Zr, Nucl. Phys. A120 (1968) 234-240. [⁸⁹Zr and ^{89m}Zr half-lives, E_γ, P_γ]
- 1968Hi10** P.F. Hinrichsen, S.M. Shafroth, D.M. van Patter, Energy Levels in ⁸⁹Y, Phys. Rev. 172 (1968) 1134-1138. [⁸⁹Y nuclear levels]
- 1968Hi12** P.F. Hinrichsen, Decay of 78.4 h ⁸⁹Zr, Nucl. Phys. A118 (1968) 538-544. [E_γ, P_γ, P_{EC}, P_{β+}, P_γ(909 keV)/P_γ(511 keV) ratio]
- 1969GuZV** R. Gunnink, J.B. Niday, R.P. Anderson, R.A Meyer, Gamma-ray Energies and Intensities, Lawrence Radiation Laboratory report UCID-15439 (1969). [E_γ, P_γ]
- 1969Ro02** E.L. Robinson, R.C. Hagenauer, E. Eichler, Decay of ⁸⁹Zr and ^{89m}Zr and Levels of ⁸⁹Y, Nucl. Phys. A123 (1969) 471-476. [⁸⁹Zr and ^{89m}Zr half-lives, E_γ, P_γ]
- 1970Ga03** St. Gagneux, P. Huber, H. Leuenberger, P. Nyikos, Änderung der Zerfallskonstante von ⁸⁹Zr in BaTiO₃, Helv. Phys. Acta 43 (1970) 39-58. [⁸⁹Zr half-life variation]
- 1971Ar18** R. Arlt, N.G. Zaitseva, B. Kracik, M.G. Loshchilov, G. Musiol, Chan Thanh Minh, H. Strusny, Decay of ⁸⁹Zr and ⁸⁷Zr, Bull. Acad. Sci. USSR, Phys. Ser. 35 (1972) 52-59. [E_γ, P_γ, P_{EC}, P_{β+}]
- 1971Go40** N.B. Gove, M.J. Martin, Log-f Tables for Beta Decay, Nucl. Data Tables 10 (1971) 205-237. [ε/β⁺ ratios]
- 1973BeYD** W. Beens, Lifetimes of Excited States in Nuclei with Mass Number Around A = 88 by Means of the Döppler Shift Attenuation Method, PhD thesis, Vrije Universiteit, Amsterdam (1973). [half-lives of ⁸⁹Y nuclear levels]
- 1973Le13** H. Leuenberger, Interpretation of the Change in Half-life of ⁸⁹Zr in BaTi(Zr)O₃ at the Upper Phase Transition (T_c - 120°C), Physica 64 (1973) 621-624. [theoretical ⁸⁹Zr half-life variation]
- 1974HeYW** R.L. Heath, Gamma-ray Spectrum Catalogue, Ge and Si Detector Spectra, 4th edition, Aerojet Nuclear Corporation report ANCR-1000-2 (1974); updated 1998, and available at: www.inl.gov/gammaray/catalogs/catalogs.shtml [Half-life, E_γ, P_γ]
- 1977Ba48** W. Bambynek, H. Behrens, M.H. Chen, B. Crasemann, M.L. Fitzpatrick, K.W.D. Ledingham, H. Genz, M. Mutterer, R.L. Intemann, Orbital Electron Capture by the Nucleus, Rev. Mod. Phys. 49 (1977) 77-221; Erratum, Rev. Mod. Phys. 49 (1977) 961-962. [ε/β⁺ ratios]

- 1977La19** F.P. Larkins, Semiempirical Auger-electron Energies for Elements $10 \leq Z \leq 100$, At. Data Nucl. Data Tables 20 (1977) 311-387. [Auger-electron energies]
- 1979Ba46** A.D. Baillie, K.W.D. Ledingham, J.G. Lynch, M. Campbell, Electron Capture to Positron Emission Ratios in the Allowed Decay of ⁸⁹Zr, J. Phys. G: Nucl. Phys. 5 (1979) 1433-1444. [P_{EC}/P_{β^+} , $P_{KX}/P_{\gamma}(909 \text{ keV})$, ϵ/β^+ ratio, E_{γ} , P_{γ}]
- 1984HaZC** J.J. Hamill, A. Zemel, Conversion-electron Spectroscopy in the Region Near ⁹⁰Zr, KVI 1983 Annual Report (1984) 18-19. [1713-keV γ transition between 2622.1-908.97 keV levels of ⁸⁹Y]
- 1984Sk01** R.T. Skelton, R.W. Kavanagh, Measurement of the ⁷Be Branching Ratio, Nucl. Phys. A414 (1984) 234-240. [Half-life, P_{EC}]
- 1985HaZI** J. Hamill, A.J. Hoeven, J. van Klinken, V.A. Wichers, A. Zemel, Conversion-electron Spectroscopy Near ⁹⁰Zr, KVI 1984 Annual Report (1985) 18-19. [1713-keV γ transition between 2622.1-908.97 keV levels of ⁸⁹Y]
- 1992Fu04** L. Funke, G. Winter, J. Döring, L. Käubler, H. Prade, R. Schwengner, E. Will, Ch. Prochostov, W. Andrejtscheff, L.G. Kostova, P.O. Lipas, R. Wirowski, Neutron-core Excitations in the N = 50 Nucleus ⁸⁹Y, Nucl. Phys. A541 (1992) 241-265. [E_{γ} , P_{γ} , half-lives of ⁸⁹Y nuclear levels]
- 1992KaZM** K. Kawade, H. Yamamoto, A. Tanaka, A. Hosoya, T. Katoh, T. Iida, A. Takahashi, Measurement of Beta-decay Half-lives of Short-lived Nuclei, Proc. 1991 Symposium on Nuclear Data, 28-29 November 1991, JAERI report JAERI-M-92-027, Eds. M. Baba, T. Nakagawa (1992) 364-368. [^{89m}Zr half-life]
- 1995ItZY** S. Itoh, M. Yasuda, H. Yamamoto, T. Iida, A. Takahashi, K. Kawade, Measurement of Beta-decay Half-lives of Short-lived Nuclei by Using High-rate Spectroscopy Amplifier, Proc. 1994 Symposium on Nuclear Data, 17-18 November 1994, JAERI report JAERI-Conf-95-008, Eds. M. Kawai, T. Fukahori (1995) 185-188. [^{89m}Y half-life]
- 1995ScZY** E. Schönfeld, Tables for the Calculation of Electron Capture Probabilities, PTB Report PTB-6.33-95-2, 1995. [P_K , P_L , P_M , P_N , P_O]
- 1996Sc06** E. Schönfeld, H. Janßen, Evaluation of Atomic Shell Data, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527-533. [ω_K , ω_L , K_{β}/K_{α} , $K_{\alpha 2}/K_{\alpha 1}$, KLX/KLL , KXY/KLL]
- 1997La20** S. Lahiri, B. Mukhopadhyay, N.R. Das, Simultaneous Production of ⁸⁹Zr and ^{90,91m,92m}Nb in α -particle Activated Yttrium and their Subsequent Separation by HDEHP, Appl. Radiat. Isot. 48 (1997) 883-886. [⁸⁹Y(α ,p3n)⁸⁹Zr mode of production]
- 1998Sc28** E. Schönfeld, Calculation of Fractional Electron Capture Probabilities, Appl. Radiat. Isot. 49 (1998) 1353-1357. [P_K , P_L , P_M , P_N , P_O]
- 1998ScZM** E. Schönfeld, G. Rodloff, Tables of the Energies of K-Auger Electrons for Elements with Atomic Numbers in the Range from Z = 11 to Z = 100, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX** E. Schönfeld, G. Rodloff, Energies and Relative Emission Probabilities of K X-rays for Elements with Atomic Numbers in the Range from Z = 5 to Z = 100, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2000Sc47** E. Schönfeld, H. Janßen, Calculation of Emission Probabilities of X-rays and Auger Electrons Emitted in Radioactive Disintegration Processes, Appl. Radiat. Isot. 52 (2000) 595-600. [P_X , P_{Ae}]

- 2002Ba85** I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, Dirac–Fock Internal Conversion Coefficients, *At. Data Nucl. Data Tables* 81 (2002) 1-334. [ICC]
- 2002Ra45** S. Raman, C.W. Nestor, Jr., A. Ichihara, M.B. Trzhaskovskaya, How Good are the Internal Conversion Coefficients Now? *Phys. Rev. C* 66 (2002) 044312, 1-23. [ICC]
- 2008Ki07** T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr., Evaluation of Theoretical Conversion Coefficients using BrIcc, *Nucl. Instrum. Methods Phys. Res. A* 589 (2008) 202-229. [ICC]
- 2012Wa38** M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer, The AME2012 Atomic Mass Evaluation, (I). Tables, Graphs and References, *Chin. Phys. C* 36 (2012) 1603-2014. [Q-value]
- 2013Si03** B. Singh, Nuclear Data Sheets for A = 89, *Nucl. Data Sheets* 114 (2013) 1-208. [Nuclear levels]

⁹⁰Sr - Comments on evaluation of decay data by V. Chisté

This evaluation was completed in 2005. The literature available by August 2005 was included.

1 Decay Scheme

⁹⁰Sr disintegrates by β^- emission to the fundamental level of ⁹⁰Y ($T_{1/2} = 2.6684 (13) \text{ d}$). The decay scheme and level spins and parities are from the evaluation of E. Browne (1997Br34).

2 Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2003Au03).

The ⁹⁰Sr half-life has been evaluated from the following data (in days):

1950Po67	7270 (110)
1955Wi15	10117 (146)
1958An40	10702 (584)
1965FI01	10227 (146)
1965FI01	10410 (329)
1965An07	10527 (51)
1978La21	10282 (13)
1983Ra09	10589 (92)
1989Ko57	10665 (37)
1992ScZZ	10513 (14)
1994Ma50	10561 (14)
1996Wo06	10495 (4)
2004Sc49	10557 (11)
Adopted	10522 (27) d or 28.80 (7) y

The half-life experimental values of 1950Po67 (7270 (110) d), 1955Wi15 (10117 (146) d), 1978La21 (10282 (13) d), 1983Ra09 (10589 (92) d) are rejected by the evaluator following the recommendation given by 1996Wo06.

The half-life weighted average has been calculated by LWEIGHT computer program (version 3).

The evaluator has chosen to take into account the nine values with associated uncertainty for the calculation. One of them (10227 (146) d) from Flynn (1965FI01) is rejected by the LWEIGHT computer program, based on the Chauvenet's criterion. The largest contribution to the weighted average comes from the value of Woods (1996Wo06) amounting to 76 %. The LWEIGHT program has increased the uncertainty of the 1996Wo06 value from 4.0 to 7.1 in order to reduce its relative weight from 76 % to 50 %.

The recommended value is the weighted average of 10522 d (28.80 (7) y), with an uncertainty of 27 d (expanded so range includes the most precise value of Woods (1996Wo06)). The reduced χ^2 value is 8.

2.1 b- Transitions

The maximum energy of the β^- transition in the decay of ⁹⁰Sr to ground state in ⁹⁰Y has been adopted from the Q value of 2003Au03 ($E_{\beta^-} = Q = 545.9$ (14) keV), and is in agreement with the experimental value of 546.0 (16) keV, measured with a magnetic β -ray spectrometer (1983Ha15).

The lg ft value (9.3) for the 546keV unique first forbidden transition and mean energy value (196 (1) keV) have been calculated with the Logft computer program (version 7.2a).

For measured first forbidden shape factors, see 1964Da16 and 1983Ha35.

3 Atomic Data

Atomic values, ω_K , $\bar{\omega}_L$ and n_{KL} , are from Schönfeld and Janßen (1996Sc33).

4 References

- 1950Po67 – R.I. Powers, A.F. Voigt, Phys. Rev. 79(1950)175 [$T_{1/2}$].
 1955Wi15 – D.M. Wiles, R.H. Tomlinson, Can. J. Phys. 33(1955)133 [$T_{1/2}$].
 1958An40 – M.P. Anikina, R.N. Ivanov, G.M. Kukavadze, B.V. Ershler, Atomnaya Energ. 4(1958)198; J. Nucl. Energ. 9(1959)167 [$T_{1/2}$].
 1964Da16 – H. Daniel, G.T. Kaschl, H. Schmitt, K. Springer, Phys. Rev. 136(1964)B1240 [β^- shape factor].
 1965Fl01 – K.F. Flynn, L.E. Gleindenin, A.L. Harkness, E.P. Steinberg, J. Inorg. Nucl. Chem. 27(1965)21 [$T_{1/2}$].
 1965An07 – S.C. Anspach, L.M. Cavallo, S.B. Garfinkel, J.M.R. Hutchinson, C.N. Smith, N. P- 15663 (1965) [$T_{1/2}$].
 1978La21 – F. Lagoutine, J. Legrand, C. Bac, Int. J. Appl. Radiat. Isotop. 29(1978)269 [$T_{1/2}$].
 1983Ha35 – H.H. Hansen, Int. J. Appl. Radiot. Isotop. 34(1983)1241 [β^- shape factor].
 1983Ra09 – H. Ramthun, Nucl. Instrum. Meth. 207(1983)445 [$T_{1/2}$].
 1989Ko57 – A.E. Kochin, M.G. Kuzmina, I.A. Sokolova, P.L. Merson, Metrologia 26(1989)203 [$T_{1/2}$].
 1992ScZZ – U. Schötzgig, H. Schrader, K. Debertain, Proc. Inter. Conf. Nucl. Data for Science and Technology, Julich (1992)562 [$T_{1/2}$].
 1994Ma50 – R.H. Martin, K.I.W. Burns, J.G.V. Taylor, Nucl. Instrum. Meth. Phys. Res. A339(1994)158 [$T_{1/2}$].
 1996Sc33 – E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
 1996Wo06 – M.J. Woods, S.E.M. Lucas, Nucl. Instrum. Meth. Phys. Res. A369(1996)534 [$T_{1/2}$].
 1997Bro34 – E. Browne, Nucl. Data Sheets 82(1997)420 [Spin, parity, energy level].
 2003Au03 – G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].
 2004Sc49 – H. Schrader, Appl. Rad. Isotopes 60(2004)317 [$T_{1/2}$].

⁹⁰Y - Comments on evaluation of decay data by V. Chisté

This evaluation was completed in 2005. Updated version in November 2006 and the literature available by this date included.

1 Decay Scheme

⁹⁰Y disintegrates by β^- emission mainly (99.983 %) to the stable ⁹⁰Zr ground state level. The decay scheme and level energies, spins and parities are based on the evaluation of E. Browne (1997Br34).

A weak beta branch occurs to the 1760 keV excited level which decays by a E0 gamma transition. This 0⁺0⁺ transition undergoes with the emission of two particles materialized by the emission of two gamma, or an electron-positron pair, or internal conversion.

2 Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2003Au03).

The half-life of the ⁹⁰Y ground state has been evaluated from the following data (in hours) :

1937Po07	57.6 (24)
1937St08	60.5 (20)
1938Sa01	66 (3)
1940Sa02	66 (2)
1946Bo01	61 (1)
1954Ch29	64.60 (43)
1955Sa27	64.029 (24)
1955Vo03	64.24 (30)
1956He77	64.8 (2)
1957Pe05	63.97 (10)
1961He09	64.10 (8)
1963Vo02	63.74 (10)
1966Ri01	64.06 (11)
1967Bi02	64.6 (8)
1968La10	64.21 (8)
1969Gr38	63.46 (13)
2004Ko18	64.053 (20)

Adopted **64.041 (31) h** or **2.6684 (13) d**

The weighted average has been calculated with LWEIGHT computer program (version 3).

The evaluator has chosen to take into account the twelve most precise values for the calculation, since the 50's: 1954Ch29, 1955Sa27, 1955Vo03, 1956He77, 1957Pe05, 1961He09, 1963Vo02, 1966Ri01, 1967Bi02, 1968La10, 1969Gr38 and 2004Ko18. The evaluator's choice is supported by the fact that in preliminary calculation with LWEIGHT program, the 1937P07, 1937St08 and 1946Bo01 values have been rejected based on the Chauvenet's criterion.

With the data set of twelve values, the largest contribution to the weighted average comes from the value of Kossert amounting to 51%. The LWEIGHT program has increased the uncertainty of the 2004Ko18 value

from 0.020 to 0.0202 in order to reduce its relative weight from 51 % to 50 % .

The weighted average of 64.041 h and the external uncertainty of 0.031 is the half-life adopted value. The reduced- χ^2 value is 4.7.

2.1 b⁻ Transitions

The maximum energy of the β^- transitions in the decay of ⁹⁰Y to excited states in ⁹⁰Zr has been calculated from the relation of

$$E_{\beta^-} = Q_{\beta^-}(\text{from 2003Au03}) - E_{\text{level in Zr-90}}(\text{from 1997Br34})$$

In the case of the transition $\beta_{0,0}^-$ (to the ground state), many experimental values of E_{β^-} have been found in literature (measured with β^- -ray spectrometer), as shown in the following table (Table 1). It can be noted that the evaluated value, 2279.8 (17) keV, is in agreement with all experimental values.

Table 1: Experimental and adopted energy of the $\beta_{0,0}^-$ transition

Reference	E_{β^-} (keV)
T. Yuasa and J. Laberrigue-Frolov (1957Yu06)	2265 (5)
O.E. Johnson et al. (1958Jo33)	2261 (3)
R.T. Nichols et al. (1961Ni02)	2271 (2)
S. André and P. Depommier (1964An12)	2268 (2)
L.M. Langer et al. (1964La13)	2273 (5)
H. Daniel et al. (1964Da16)	2284 (5)
P.G. Hansen et al. (1966Ha15)	2275 (5)
P. Riehs (1966Ri01)	2280 (5)
T. Nagarajan et al. (1971Na09)	2288 (3)
H. Hansen (1983Ha35)	2279.5 (29)
C. Greenwood and M.H. Putnam (1993Gr17)	2274.8 (30)
Adopted value	2279.8 (17)

For the probabilities of the β^- transitions, the available published data are given in Table 2:

Table 2: Measured and adopted probabilities of β^- transitions in %.

Populated level (keV)	1961La07	1970Va09	1976Gr16	Adopted values
ground state	99.9885 (15)	99.977 (9)		99.983 (6)
1760.72	0.0115(15)	0.023(9)		0.017 (6)
2186.282			0.0000014 (3)	0.0000014 (3)

For the ground state and 1760.72-keV β^- transitions, the adopted values are the weighted averages of the two values given with uncertainties.

The lg ft values have been calculated with the LOGFT program (version 7.2a).

2.2 g Transitions

The 1760- and 2186-keV γ -ray transition probabilities are 0.017 (6) % and 0.0000014 (3) %, respectively. These values come directly from the evaluated β^- transition probabilities and adopted decay scheme.

Multipolarities of these γ -ray transitions are from 1997Br34.

The internal conversion coefficients (α_T , α_K and α_L) for 2186-keV γ -ray transition has been calculated using the ICC Computer Code (program Icc99v3a – GETICC dialog). The adopted values have been interpolated from the new tables of Band et al.(2002Ba85). The uncertainties in α_T , α_K and α_L have been estimated as 3 %.

The intensity of the conversion electrons was measured by Legrand (1972) being $1,3 (7) \times 10^{-2} \%$.

3 Atomic Data

Atomic values, ω_K , ω_L and n_K , are from Schönfeld and Janßen (1996Sc33).

5 Photon emissions

5.1 g-ray Emissions

The 2186-keV γ -ray absolute emission probability has been deduced from the total ($\gamma+ce$) transition probability of 0.0000014 (3) % (§ 2.2) and the theoretical α_T (2002Ba85) for a E2 transition.

The ratio of two photon decay $P_{\gamma\gamma}$, occurring during the $0^+ - 0^+$ gamma transition, to the sum of internal-pair decay $P_{e^+e^-}$ and internal-conversion decay P_{ic} : $P_{\gamma\gamma} / (P_{e^+e^-} + P_{ic})$ is 0,040 (5) (1997Br34).

The number of positrons (leading to the emission of the 511 keV annihilation peak) is : 31,9 (5) 10^{-4} per 100 beta decays (R.G.Selwyn). Other values : 36 (5) $\times 10^{-4}$ (1956Gr21) and 34 (4) (1961La07).

X-ray emissions aren't given in the table file. $IK\alpha$ was measured by Legrand (1972) being $3,7 (5) \times 10^{-4} \%$.

6 References

- 1937Po07 – M.L. Pool, J.M. Cork, R.L. Thornton, Phys. Rev. 52(1937)239 [$T_{1/2}$].
 1937St08 – D.W. Stewart, J.L. Lawson, J.M. Cork, Phys. Rev. 52(1937)901 [$T_{1/2}$].
 1938Sa01 – R. Sagane, S. Kojima, G. Miyamoto, M. Ikawa, Phys. Rev. 54(1938)970 [$T_{1/2}$].
 1940Sa02 – R. Sagane, S. Kojima, G. Miyamoto, M. Ikawa, Phys. Rev. 57(1940)1179 [$T_{1/2}$].
 1946Bo01 – W. Bothe, Z. Naturforsch. 1(1946)173 [$T_{1/2}$].
 1954Ch29 – A. Chetham-Strode Jr., E.M. Kinderman, Phys. Rev. 93(1954)1029 [$T_{1/2}$].
 1955Sa27 – M.L. Salutsky, H.W. Kirby, Anal. Chem. 27(1955)567 [$T_{1/2}$].
 1955Vo03 – H.L. Volchok, J.L. Kulp, Phys. Rev. 97(1955)102 [$T_{1/2}$].
 1956He77 – G. Herrmann, F. Strassmann, Z. Naturforsch. 11a(1956)946 [$T_{1/2}$].
 1956Gr21 – J.S. Greenberg, M. Deutsch, Phys. Rev. 102,2 (1956) 415 [$P_{e^+e^-}$].
 1957Pe05 – D.F. Peppard, G.W. Mason, S.W. Moline, J. Inorg. Nucl. Chem. 5(1957)141 [$T_{1/2}$].
 1957Yu06 – T. Yuasa, J. Laberrig ue-Frolow, Journal de Physique et le Radium 18(1957)498 [End -point energy].
 1961La07 – H.Langhoff, H.H. Hennies. Z. Physik 164 (1961) 166. [$P_{e^+e^-}$].
 1958Jo33 – O.E. Johnson, R.G. Johnson, L.M. Langer, Phys. Rev. 112(1958)2004 [End-point energy].
 1961He09 – R.L. Heath, J.E. Cline, C.W. Reich, E.C. Yates, E.H. Turk, PhysRev. 123(1961)903 [$T_{1/2}$, γ -ray emission intensity].
 1961La07 – H. Langhoff, H.H. Hennies, Z. Physik 164(1961)161 [Branching ratio].
 1961Ni02 – R.T. Nichols, R.E. McAdams, E.N. Jensen, Phys. Rev. 122(1961)172 [End-point energy].
 1962Ne02 – M. Nessin, T.H. Kruse, K.E. Eklund, Phys. Rev. 125(1962)639 [α_π].
 1963Vo02 – H.R. von Gunten, W. Scherle, H. Hugli, Nucl. – Med. (Stuttgart) 3(1963)417 [$T_{1/2}$].
 1964An12 – S. André, P. Depommier, J. Physique 25(1964)673 [End-point energy].
 1964La13 – L.M. Langer, E.H. Spejewski, D.E. Wortman, Phys. Rev. 135(1964)B581 [End-point energy].
 1964Da16 – H. Daniel, G.Th. Kaschl, H. Schmitt, K. Springer, Phys. Rev. 136(1964)B1240 [End-point energy].
 1966Ri01 – P. Riehs, Nucl. Phys. 75(1966)381 [$T_{1/2}$, End-point energy].
 1967Bi02 – J.K. Bienlein, G. Grof, W. Kreische, W. Lampert, G. Loos, Nucl. Phys. A92(1967)549 [$T_{1/2}$].
 1968La10 – F. Lagoutine, Y. Le Gallic, J. Legrand, Int. J. Appl. Radiat. Isotopes 19(1968)475 [$T_{1/2}$].
 1969Gr38 – V.P. Groll, F. Grass, K. Buchtela, Radiochem. Acta 12(1969)152 [$T_{1/2}$].
 1970Va09 – J.C. Vanderleeden, P.S. Jastram, Phys. Rev. C1(1970)1025 [Branching ratio].

- 1971Na09 – T. Nagarajan, M. Ravindranath, K.V. Reddy, *Nuovo Cim.* 2A(1971)662 [End-point energy].
- 1972Le** – J.Legrand *et al.* Proc. Int. Conf. Inner shell Ionization Phenom. And future applications, CONF-720404, Vol. 3, (1972) 2167 [IX_K]
- 1973Ra10 – S. Raman, N.B. Gove, *Phys. Rev.* C7(1973)1995 [I_γ].
- 1973Ha18 – A. Hanser, *Nucl. Instrum. Meth.* 107(1973)187 [I_γ].
- 1974Kl06 – A. Kluge, K. Kroth, F.J. Schröder, W. Thomas, H. Toschinski, C. Günther, *Nucl. Phys.* A224(1974)1 [I_γ].
- 1976Gr16 – H.C. Griffin, *Radiochem. Radioanal. Lett.* 27(1976)353 [Branching ratio].
- 1978Ra05 – G.N. Rao, C. Günther, *Phys. Rev.* C17(1978)1266 [I_γ].
- 1983Ha35 – H.H. Hansen, *Int. J. Appl. Radiat. Isot.* 34(1983)1241 [End-point energy].
- 1993Gr17 – C. Greenwood, M.H. Putnam, *Nucl. Instrum. Meth. Phys. Res.* A337(1993)106 [End-point energy].
- 1996Sc33 – E. Schönfeld, H. Janßen, *Nucl. Instrum. Meth. Phys. Res.* A369(1996)527 [Atomic data].
- 1997Br34 – E. Browne, *Nucl. Data Sheets* 82(1997)421 [E_{level}, spin, parity and multipolarity].
- 2002Ba85 – I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, *Atomic Data and Nuclear Data Tables* 81(2002)1 [α].
- 2003Au03 – G. Audi, A.H. Wapstra, C. Thibault *Nucl. Phys.* A729(2003)129 [Q].
- 2004Ko18 – K. Kossert, H. Schrader, *Applied Radiation and Isotopes* 60(2004)741 [T_{1/2}].
- 2006Se** - R.G. Selwyn *et al.* *Applied Radiation and Isotopes*, 65 (2007) 318 [P_{e+e-}]

⁹⁰Y^m - Comments on evaluation of decay data by V. Chisté

This evaluation was completed in 2005. The literature available by August 2005 was included.

1 Decay Scheme

⁹⁰Y^m disintegrates 99.9981 (2) % through isomeric transition to the ⁹⁰Y ground state and 0.0019 (2) % by β⁻ emission to the 2318 keV excited state in ⁹⁰Zr. The decay scheme, level energies, spins and parities and half-lives of excited states are based on the evaluation of E. Browne (1997Br34).

2 Nuclear Data

The Q value in the decay of ⁹⁰Y^m → ⁹⁰Zr (2961.8 (17) keV) has been calculated from the following relation:

$$Q(^{90}\text{Y}^{\text{m}} \rightarrow ^{90}\text{Zr}) = Q(^{90}\text{Y} \rightarrow ^{90}\text{Zr}) + Q(^{90}\text{Y}^{\text{m}} \rightarrow ^{90}\text{Y})$$

Both latter values are from the atomic mass evaluation of Audi *et al.* (2003Au03).

The experimental ⁹⁰Y^m half-life values (in hours) are given in Table 1:

Table 1: Experimental values of ⁹⁰Y^m half-life

Reference	Value (h)
Carter-Waschek and Linder (1961Ca12)	3.2 (1)
Heath et al.(1961He09)	3.14 (10)
Haskin and Vandenbosch (1961Ha17)	3.19 (6)
Abecasis et al.(1962Ab03)	3.15 (5)
Grench et al.(1967Gr02)	3.19 (1)
Anthony et al.(1992An19)	3.244 (5)
Adopted	3.19 (6)

The weighted average has been calculated with the LWEIGHT computer program (version 3).

The evaluator has chosen to take into account the seven values with associated uncertainties for the statistical processing. The largest contribution to the weighted average comes from the value of Anthony (1992An19) amounting to 79 %. The LWEIGHT program has increased the uncertainty for the 1992An19 value from 0.005 to 0.010 in order to reduce its relative weight from 79 % to 50 %.

The recommended value is the weighted average of 3.19 h with a final uncertainty of 0.06, expanded to include the most precise value of Anthony (1992An19, 3.244 (5) h). The reduced-χ² value is 3.5.

2.1 b- Transitions

The maximum energy of the β⁻ transition in the decay of ⁹⁰Y^m → ⁹⁰Zr has been calculated from the relation:

$$E_{\beta^-} = Q(^{90}\text{Y}^{\text{m}} \rightarrow ^{90}\text{Zr}, \text{ from } 2003\text{Au03}) - E_{\text{level in Zr-90}}(\text{from } 1997\text{Br34}) = 642.9 (17) \text{ keV.}$$

The *lg ft* of 9.6 and mean energy of 231.7 (7) keV have been calculated with the LOGFT computer program for the 642-keV unique first forbidden transition.

The 642-keV β^- transition probability is deduced from the ratio $I_{\gamma}(2319 \text{ keV})/ I_{\gamma}(479 \text{ keV})$ given by H. C. Griffin (1976Fr16). The value of this ratio has been recalculated by the evaluator with the adopted photon branching ratio (see **5.2 g-ray Emission**).

2.2 g Transitions

For the $^{90}\text{Y}^m \rightarrow ^{90}\text{Y}$ and $^{90}\text{Y}^m \rightarrow ^{90}\text{Zr}$ branching, the transition probabilities have been calculated using gamma ray intensities and the internal conversion coefficients (see **5.2 g-ray emissions**).

Multipolarities of γ -ray transitions in both decays of $^{90}\text{Y}^m$ are from 1997Br34:

202-keV γ -ray : M1 + E2, $\delta = -0.04$ (4)

479-keV γ -ray : M4 (+ E5)

682-keV γ -ray : E5

2319-keV γ -ray (from $^{90}\text{Y}^m \rightarrow ^{90}\text{Zr}$): E5

The internal conversion coefficients (ICC's) have been calculated using the Icc99v3a computer program (GETICC dialog). The adopted values have been interpolated from new tables of Band et al.(2002Ba85). The uncertainties of internal conversion coefficients have been estimated as 3 %.

3 Atomic Data

Atomic values are from 1996Sc33.

4 Electron Emissions

The Auger electrons emission probabilities have been calculated from γ -ray and conversion electron data by using the EMISSION computer program. The Auger electrons emission probabilities of ^{90}Zr aren't given in the table file, because they are negligible (of the order of 10^{-7}).

5 Photon Emissions

5.1 X-ray Emissions

The X-ray emission probabilities have been calculated from γ -ray and conversion electron data by using the EMISSION computer program. The X-ray emission probabilities of ^{90}Zr aren't given in the table file, because they are negligible (of the order of 10^{-7}).

5.2 g-ray Emissions

The relative emission probabilities measured in the isomeric decay of $^{90}\text{Y}^m$ are given in Table 2. The 479keV line as been taken as 100 %.

Table 2: Relative γ -ray emission probabilities measured in the isomeric decay of $^{90}\text{Y}^m$, in %.

Energy (keV)	Heath (1961He09)	Hanser (1973Ha18)	Raman (1973Ra10)	Kluge (1974K106)	Griffin (1976Gr17)	Rao (1978Ra05)	Evaluated Values
202.53	104.99 (44)	107.2 (4)	103.7 (33)	none	none	none	106.1 (11)
682.04	< 0.01	none	none	0.40 (8)	0.34 (5)	0.35 (3)	0.352 (24)

For each γ -ray, the evaluated relative γ -ray emission probabilities are weighted averages (calculated with the LWEIGHT computer program, version 3) of the three values measured with uncertainties.

The normalization factor to convert the relative emission probabilities to the absolute emission probabilities has been calculated from the intensity balance at the ⁹⁰Y ground state. As β⁻ branching in the ⁹⁰Y^m is negligible (1976Gr16), the normalization factor is:

$$\text{Normalization factor} = \frac{100\%}{[(1 + a_T(202))P_{rel}(202)] + [(1 + a_T(682))P_{rel}(682)]}$$

From the theoretical α_T and the evaluated relative emission probabilities of the 202- and 682-keV γ -rays (Table 2), the normalization factor becomes **0.915 (9) %**. The uncertainty was calculated through the propagation on the formula given above.

The 479-keV transition probability is given by:

$$P_{(\gamma+ce)}(682 \text{ keV}) + P_{(\gamma+ce)}(479 \text{ keV}) = 100 \%$$

Taking into account the evaluated normalization factor, the theoretical α_T and the evaluated relative emission probability of the 682-keV γ -rays (Table 2), then $P_{(\gamma+ce)}(682 \text{ keV}) = 0.329 (23) \%$ and, therefore, $P_{(\gamma+ce)}(479 \text{ keV}) = 99.671 (23) \%$.

The evaluated relative and absolute emission intensities for the 202-, 479- and 682-keV γ -rays are given in Table 3:

Table 3: Evaluated relative and absolute γ -ray emission intensities.

Energy (keV)	Relative emission intensity (%)	Absolute emission intensity (%)
202.53	106.1 (11)	97.1 (14)
479.53	99.4 (10)	90.97 (24)
682.04	0.352 (24)	0.322 (22)

From the 479-keV γ -ray absolute emission intensity value (Table 3) and the value of $I_\gamma(2319 \text{ keV}) / I_\gamma(479 \text{ keV}) = 2.1 (2) \cdot 10^{-5}$, as given by Griffin (1976Gr16), then $I_\gamma(2319 \text{ keV}) = 0.0019 (2) \%$.

6 References

- 1961Ca12 – C. Carter-Waschek, B. Linder, Nucl. Phys. 27(1961)415; Erratum Nucl. Phys. 31(1962)351 [T_{1/2}].
 1961Ha17 – L. Haskin, R. Vandenbosch, Phys. Rev. 123(1961)184 [T_{1/2}].
 1961He09 – R.L. Heath, J.E. Cline, C.W. Reich, E.C. Yates, E.H. Turk, Phys. Rev. 123(1961)903 [T_{1/2}, I_γ].
 1962Ab03 – S. Abecasis, H. Bosch, M.C. Caracoche, A. Mococho, H. Vignau, Rev. Union Mat. Arg., Assoc. Fis. Arg. 21(1962)104; Nucl. Sci. Abstr. 17(1963)3732, Abstr. 28327 [T_{1/2}].
 1967Gr02 – H.A. Grench, K.L. Coop, H.O. Menlove, F.J. Vaughn, Nucl. Phys. A94(1967)157 [T_{1/2}].
 1973Ha18 – A. Hanser, Nucl. Instrum. Meth. 107(1973)187 [I_γ].
 1973Ra10 – S. Raman, N.B. Gove, Phys. Rev. C7(1973)1995 [I_γ].
 1974Kl06 – A. Kluge, Nucl. Phys. A224(1974)1 [I_γ].
 1976Gr16 – H.C. Griffin, Radiochem. Radioanal. Lett. 27(1976)353 [P_β].
 1978Ra05 – G.N. Rao, C. Günther, Phys. Rev. C17(1978)1266 [I_γ].
 1992An19 – M.S. Anthony, D. Oster, A. Hachem, J. Radioanal. Nucl. Chem. 166(1992)63 [T_{1/2}].
 1996Sc33 – E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
 1997Br34 – E. Browne, Nucl. Data Sheets 82(1997)379 [E_{level}, spin, parity, multipolarity, T_{1/2}].
 2002Ba85 – I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, Atomic Data and Nuclear Data Tables 81(2002)1 [α].
 2003Au03 – G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].

⁹³Zr - Comments on evaluation of decay data

by M.A. Kellett

This evaluation was completed in September 2013. Literature cutoff date: September 2013.

Limitation of Relative Statistical Weights Method (LWM) was applied to average the decay data when appropriate (unless otherwise stated) by use of the LWEIGHT program and/or the Excel implementation of same. All uncertainties are given as the combined uncertainty to one standard deviation.

The ⁹³Zr samples used in the various measurements were produced either by chemical extraction from irradiated uranium fuel (⁹³Zr is a major fission product), or via activation of the zircaloy cladding of modern PWR fuel.

Impurities and/or interfering nuclei include ⁹⁵Zr, ^{93m}Nb (the decay daughter), ⁹³Mo and ⁵⁵Fe (this latter particularly for liquid scintillation measurements). The presence of ⁹⁵Zr is easily dealt with as it has a comparatively short half-life (~ 64 d) and so extracted samples are kept to allow the activity of this nuclide to decay away. Radiochemical separation and mass spectrometry were performed before measurements in almost all of the published reports in order to remove these impurities.

The most important complication in measuring the decay properties for this nuclide are linked to it being a pure beta emitter and also a severe lack of knowledge of the decay scheme, particularly the beta branching fraction to the metastable state and ground state of the daughter, ⁹³Nb. Observation of the 30.77 keV γ ray from ^{93m}Nb, following β^- -decay of ⁹³Zr, only becomes possible once equilibrium is reached, i.e. > 80 a.

1 Decay Scheme

⁹³Zr decays 100 % via two β^- -transitions to ^{93m}Nb (16.12 (15) a) and ⁹³Nb (stable).

The overall decay scheme presented is complete and consistent. Its overall consistency is verified by the comparison between Q_{calc} (90 (6) keV) and Q_{β^-} (90,3 (15) keV) from the 2012 atomic mass evaluation of M. Wang *et al.* (2012Wa38). The level energy of ^{93m}Nb, along with the spins and parities, are taken from 2004BeZQ.

2 Nuclear Data**2.1. Half-life**

The experimental half-life values of ⁹³Zr are given in Table 1, resulting in a recommended value of **1,61 (6) $\times 10^6$ a**.

Table 1: Experimental values of ⁹³Zr half-life (in $\times 10^6$ a)

Reference	$T_{1/2}$ values (10^6 a)	Uncertainty
1950St90 [†]	1,5 – 8,5	
1952G1** [†]	0,93	0,05
1953G131 [†]	0,95	0,05
1972FIZM ^a	1,53	0,1
2010Ya01 ^b	1,13	0,11
2010Ca01	1,64	0,06
Recommended^c	1,61	0,06

LWEIGHT reduced $\chi^2 = 0,89$ (critical $\chi^2 = 6,63$)

[†] Superseded by 1972FIZM

^a Private communication from K.F. Flynn (ANL) to D. Kocher (ORNL) claiming to include their “best information” concerning ⁹³Zr/⁹³Nb, so supersedes 1950St90, 1952G1** and 1953G131

^b The LSC measurement method is questionable (definition of quenching factor, use of ⁶³Zn spectral shape for normalisation, etc.), and so this value has been excluded from the dataset

^c Recommended value is the weighted mean of 1972FIZM and 2010Ca01, where the value of 2010Ca01 amounts to 73 % of the total weight, due to its far lower uncertainty. This uncertainty is adopted for the recommended value.

The recommended half-life for ^{93m}Nb is **16,12 (15) a**, taken from 2004BeZQ.

2.2. Beta Transitions

Energies and Emission Probabilities

Beta-particle energies were determined from the difference between the adopted Q-value and the final level energies of the daughter ⁹³Nb. A level energy of 30,77 (2) keV has been adopted from 2004BeZQ. Beta-particle transition probabilities were taken from 2010Ca01, who used an iterative approach to look at the in-growth of ^{93m}Nb over a three year period, determined from X-ray measurements, and hence deduced the beta transition probabilities to ^{93m}Nb (73 (5) %) and ⁹³Nb (27 (5) %) respectively. Note the uncertainty of 5 % on each is from the text of the report, not Figure 7, which shows 6 %. It is noteworthy that these values are significantly different from those given in 1972FIZM (≥ 95 % and ≤ 5 %, respectively), but due to the earlier measurements only quoting limits, and that the later measurements used an iterative fitting approach, it is felt they are more reliable.

References

- 1950St90 E.P.Steinberg, L.E.Glendenin, Phys.Rev. 78 (1950) 624
[Half-life; Beta branching; Beta emission energies]
- 1952Gl** L.E.Glendenin, E.P.Steinberg, ANL-4833 (1952) 89
[Half-life; Beta branching; Beta emission energies; Nb-93m half-life; Conversion electrons; X-rays]
- 1953Gl131 L.E.Glendenin, E.P.Steinberg, ANL-5000 (1953) 55
[Half-life; Beta branching; Conversion electrons; X-rays]
- 1972FIZM K.F.Flynn, Priv. Comm. 1972
[Half-life; Beta branching; Conversion electrons; X-rays]
- 2004BeZQ M.M.Bé, V.Chisté, C.Dulieu, E.Browne, V.Chechev, N.Kuzmenko, R.Helmer, A.Nichols, E.Schönfeld, R.Dersch, Monographie BIPM-5 Vol. 1(2004)
[Nb-93 levels; Nb-93m half-life, IT emission]
- 2008DuZX C.Dulieu, M.M.Bé, V.Chisté, Proc. Int. Conf. on Nuclear Data for Science and Technology, 22-27 April 2007, Nice, France (2008) 97
[SAISINUC software]
- 2010Ca01 P.Cassette, F.Chartier, H.Isnard, C.Frechou, I.Laszak, J.P.Degros, M.M.Bé, M.C.Lépy, I.Tartès, Appl.Radiat.Isot. 68 (2010) 122
[Half-life, Beta branching]
- 2010Ya01 J.Yang, S.Zhang, Y.Ding, F.Shu, J.Zhang, Radiochim.Acta 98 (2010) 59
[Half-life]
- 2012Wa38 M.Wang, G.Audi, A.H.Wapstra, F.G.Kondev, M.MacCormick, X.Xu, B.Pfeiffer, Chin. Phys. C36 (2012) 1603
[Q]

^{93m}Nb – Comments on evaluation of decay data by V.P. Chechev, N.K. Kuzmenko and M.A. Kellett

This evaluation was originally completed in May 2001 and was updated in October 2013 by M.A. Kellett (CEA/LNE-LNHB) to include new internal conversion coefficients and thus the associated atomic data (X-ray emissions, conversion- and Auger-electrons). No new references have been published during the intervening period.

1 Decay scheme

The ^{93m}Nb decay scheme is very simple. It includes the single 30.77 keV gamma transition with the well-established multipolarity of M4 (1972Ko59, 1997Ba13).

2 Nuclear Data

Q(IT) value is the energy of the isomeric transition to the ground state of ⁹³Nb (1977Mo07).

There are available seven measurements of the ^{93m}Nb half-life, in years:

Half-life	Reference	Comment
~ 4	1954Sc74	Omitted (approximate)
13.6 (3)	1965Fl02	Omitted (low precision)
11.4 (9)	1976He**	Omitted (low precision)
16.4 (4)	1977Ll01	Omitted (preliminary)
15.3 (13)	1980Va**	Omitted (preliminary)
16.10 (19)	1981Ll01	
16.13 (15)	1983Va25	
16.12 (15)	Recommended value (weighted mean)	

The measurement result of 1954Sc74 was omitted as only approximate. The 1977Ll01 and 1980Va** values, measured by Lloret and Vaninbroux, respectively, were only preliminary results. They were obtained from observations over relatively short periods – in both cases the measurements continued over only about four or so years. Consequently only the final values of 1981Ll01 and 1983Va25 have been used by the evaluator for statistical processing. The lower values of 1965Fl02 and 1976He** were omitted as being less precise and they disagree with the two more precise measurements of 1981Ll01 and 1983Va25.

Averaging of these latter values gives a weighted mean of 16.12 with an internal uncertainty of 0.12 and an external uncertainty of 0.01. As the measurement method was the same in both cases, the minimum input uncertainty of 0.15 has been chosen for the final uncertainty of the weighted mean. Thus, the evaluated ^{93m}Nb half-life is **16.12 (15) years**.

2.1 Gamma Transition and Internal Conversion Coefficients.

The energy of the gamma transition, 30.77 (2) keV, has been taken from the 1977Mo07 measurement. The 1972FlZM measurement value of 30.4 (3) keV is significantly less accurate.

The multipolarity of the gamma transition, M4, is determined confidently from measured subshell ratios:

$$K/(L+M) = 0.18 (2) (1964Ho08),$$

$$K/L = 0.21 (2) (1964Ho08),$$

$$K/(L+M+\dots) = 0.19 (2) (1982Re09),$$

$$L/(M+N+\dots) = 3.8 (4) (1982Re09).$$

The internal conversion coefficient (α_K) is obtained by interpolation of the ICC tables using the BrIcc code of Kibédi (2008Ki07). The relative uncertainty of α_K has been adopted as 1.5 % in accordance with the available estimations of the reliability of the calculations of the theoretical ICC with a pure multipolarity (see 2008Ki07). The α_K adopted value of $2.60 (4) 10^4$ agrees well with the measured value of $2.58 (15) 10^4$ (1976Ju04) and the evaluated value of $2.63 (6) 10^4$ (1987La**), but disagrees with the value of $1.7 (3) 10^4$ calculated in (1977Mo07) from the measured ratio $P_\gamma/P_{XK} = 8 (1) 10^{-5}$.

The adopted value of α_K is supported by the measurement result of $2.4 (9) 10^4$ obtained by the quite different method – investigation of "electron bridge" in ^{93m}Nb decay (1999ZhZY).

The evaluated α_L , α_M , α_T are also theoretical values for M4 multipolarity.

3 Atomic Data

3.1 Fluorescence yields

The fluorescence yields are taken from 1996Sc06 (Schönfeld and Janßen).

3.2 X Radiations

The X-ray energies are based on the wave lengths in the compilation of 1967Be65 (Bearden). The relative K X-ray emission probabilities are taken from 1999ScZX.

3.3 Auger Electrons

The energies of Auger electrons are from 1977La19 (Larkins) and 1987La** (Lagoutine, Table de Radionucléides).

The ratios $P(KLX)/P(KLL)$ and $P(KXY)/P(KLL)$ are taken from 1996Sc06.

4 Photon Emissions

4.1 X-Ray Emissions

The total K X-ray absolute emission probability computed with use of the ICC α_T , α_K and the K-fluorescence yield $\omega_K = 0.751 (4)$ is 11.54 (26) per 100 disintegrations. It is consistent with the measurements of 11.6 (4) (1978Ba**, 1980Va**) and 11.5 (3) (1983Va25). (See these references also in 1991BaZS). Other measured values include 10.7 (3) (1982Al**), 11.04 (28) (1985Ge**), and 11.12 (22) (1990Co17). The adopted value of the total K X-ray absolute emission probability is thus 11.54 (26).

The absolute emission probabilities of the K X-ray components have been computed from P_{XK} using the relative probabilities from 1996Sc06.

The total L X-ray absolute emission probability has been computed with use of the ICC α_L and the atomic data of $\omega_L = 0.0347 (9)$, $n_{KL} = 1.045 (4)$ from 1996Sc06.

4.2 Gamma Emissions

The energy of the gamma ray, 30.77 (2) keV, is from the 1977Mo07.

The absolute emission probability of the gamma ray, $5.91 (9) 10^{-4}$ per 100 disintegrations, is computed from the decay scheme using the ICC α_T .

5 Electron Emissions

The energies of the conversion electrons have been calculated from the gamma-transition energies given in 2.1 and the electron binding energies using the computer program EMISSION (v3.10, 28-Jan-2003) described in 2000Sc47.

The total emission probability of the conversion electrons, 99.999 409 (9), has been obtained as $P_{(ec1,0T)} = 100 - P_\gamma$ (per 100 disintegrations). The emission probabilities of the K-, L-, M-, NO-conversion electrons have been calculated using the conversion coefficients given in 2.1.

The values of the emission probabilities of K-Auger electrons have been calculated using the gamma transition probability given in 2.1, the atomic data given in 3, and the conversion coefficients given in 2.1 also using the computer program EMISSION (v3.10, 28-Jan-2003) described in 2000Sc47.

6 References

- 1954Sc74 R.P.Schuman, Phys. Rev. 96 (1954) 121
(Half-life)
- 1964Ho08 K.Hohmuth, G.Muller, J.Schinthmeister, Nucl. Phys. 52 (1964) 590
(ICC subshell ratios)
- 1965Fl02 K.F.Flynn, L.E.Glendenin, E.P.Steinberg, Nucl. Sci. Eng. 22 (1965) 416
(Half-life)
- 1967Be65 J.A.Bearden, Rev. Mod. Phys. 39 (1967) 78
(X-ray energies)
- 1972FIZM K.F.Flynn, Priv. Comm. (1972)
(Gamma-ray energies)
- 1972Ko59 D.C.Kocher, Nucl. Data Sheets 8 (1972) 527
(Multipolarity)
- 1976He** F.Hegeudues, Report EUR 5667E I (1976) 757
(Half-life)
- 1976Ju04 M.Jurcevic, A.Ljubivic, D.Rendic, Fizika (Zagreb) 8 (1976) 81
(K ICC)
- 1977La19 F.P.Larkins, At. Data Nucl. Data Tables 20 (1977) 313
(Auger electron energies)
- 1977Ll01 R.L.Lloret, Radiochem. Radioanal. Lett. 29 (1977) 165
(Half-life)
- 1977Mo07 J.Morel, J.-P.Pérolat, N.Coursol, C.R. Acad. Sci. (Paris) B284 (1977) 223
(X-ray and gamma-ray energies and emission probabilities, K ICC)
- 1978Ba** W.Bambynek, D.Reher, R.Vaninbroux, Proc. Int. Conf. on Neutron Physics and Nuclear Data for Reactors and other Applied Purposes, Harwell, September 1978, OECD Nuclear Energy Agency, Paris (1978) 778
(K X-ray emission probability)
- 1980Va** R.Vaninbroux, Liquid Scintillation Counting (D.L.Horrockx, E.L.Alpen Eds.) Academic Press, New York, Vol.1 (1980) 43
(K X-ray emission probability)

- 1981Ll01 R.Lloret, Radiochem. Radioanal. Lett. 50 (1981) 113
(Half-life)
- 1982Al** W.G.Alberts, R.Hollnagel, K.Knauf, W.Pessara, Proc. 4th ASTM-EURATOM Symposium on Reactor Dosimetry (F.B.K.Kam Ed.), NUREG/CP-0029, Gaithersburg 1 (1982) 433
(K X-ray emission probability)
- 1982Re09 D.Reher, Int. J. Appl. Radiat. Isotop. 33 (1982) 537
(ICC subshell ratios, multipolarity)
- 1983Va25 R.Vaninbroukx, Int. J. Appl. Radiat. Isotop. 34 (1983) 1211
(Half-life, K X-ray emission probability)
- 1985Ge** R.J.Gehrke, J.W.Rogers, J.D.Baker, Proc. 5th ASTM-EURATOM Symposium on Reactor Dosimetry, Geesthacht, FRG, 24-28 September 1984, Dordrecht 1 (1985) 319
(K X-ray emission probability)
- 1987La** F.Lagoutine, N.Coursol, J.Legrand, Table de Radionucléides. ISBN-2-7272-0078-1. LMRI, 1982-1987, BP 52, 91191 Gif-sur-Yvette Cedex, France
(ICC, multipolarity)
- 1990Co17 B.M.Coursey, R.Vaninbroukx, D.Reher, J.M.R.Hutchinson, P.A.Mullen, K.Debertin, R.J.Gehrke, J.W.Rogers, L.D.Koepfen, D.Smith, Nucl. Instrum. Methods Phys. Res. A290 (1990) 537
(K X-ray emission probability)
- 1991BaZS W.Bambynek, T.Barta, R.Jedlovsky, P.Christmas, N.Coursol, K.Debertin, R.G.Helmer, A.L.Nichols, F.J.Schima, Y.Yoshizawa, TECDOC-619, IAEA. X-ray and gamma-ray standards for detector calibration. Vienna (1991)
(K X-ray emission probability)
- 1996Sc06 E.Schönfeld, H.Janßen, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527
(Atomic data)
- 1997Ba13 E.Baglin, Nucl. Data Sheets 80 (1997) 1
(Decay scheme)
- 1999ScZX E.Schönfeld, G.Rodloff, Report PTB-6 11-1999-1 (1999)
(Atomic data)
- 1999ZhZY V.A.Zheltonozhsky, A.G.Zelinsky, Yu.M.Shevchenko, E.G.Shemchuk, Program and Thesis, Proc.49th Ann. Conf. Nucl. Spectrosc. Struct. At Nuclei, Dubna (1999) 100
(Electron, X-ray and gamma-ray emission probabilities, K ICC)
- 2000Sc47 E.Schönfeld, H.Janssen, Appl. Radiat. Isot. 52 (2000) 595
(X-ray and Auger Electron emission probabilities)
- 2008Ki07 T.Kibédi, T.W.Burrows, M.B.Trzhaskovskaya, P.M.Davidson, C.W.Nestor Jr, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202
(Theoretical ICCs)

**^{94m}Tc – Comments on evaluation of decay data
by A. L. Nichols**

Evaluated: March/April 2014

Evaluation Procedures

Limitation of Relative Statistical Weight Method (LWM) and other analytical techniques were applied to obtain averaged data throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

^{94m}Tc ($T_{1/2} = 51.9$ min) undergoes 100% electron capture/ β^+ decay ($Q(\text{EC}) = 4332$ (5) keV) to various excited nuclear levels and the ground state of ⁹⁴Mo. A reasonably well-defined decay scheme was derived from the gamma-ray measurements of 1969Ba09 and 1986AgZX, and γ - γ coincidence studies of 2003Fr02, consisting of 16 EC/ β^+ transitions and 52 gamma-ray emissions. While nineteen gamma-ray emissions were originally defined as unplaced in the decay scheme, twelve of these transitions have been successfully incorporated into the proposed decay scheme during the course of the current study, while the 1022-keV gamma emission was identified as the (511 + 511 keV) sum peak. Thus, six of the observed gamma rays remain unplaced. Weighted mean relative emission probabilities were calculated for a significant number of the gamma rays, although some of the recommended values arose solely from either 1969Ba09 or 1986AgZX. All of these relative emission probabilities have been defined in terms of the 871.094-keV gamma ray (100%).

A further sixteen gamma rays have been observed/proposed by Fransen *et al.* (2003Fr02), but an inability to derive and quantify their absolute emission probabilities has prevented their satisfactory insertion and adoption in the decay scheme and data file. If these additional gamma transitions could have been suitably introduced into the ^{94m}Tc decay scheme, they would have impacted to a small degree on the EC/ β^+ decay data.

Nuclear Data

^{94m}Tc decay is viewed as a potentially suitable β^+ emitter for application in positron emission tomography (PET).

Half-life of ^{94m}Tc

The measurements of 1948Mo19 (enriched ⁹⁴Mo(d,2n) ^{94m}Tc), 1950Me21 (⁹⁴Mo(p,n) ^{94m}Tc) and 1962Mo06 (enriched ⁹⁴Mo(d,2n) ^{94m}Tc) were adopted to give a weighted mean half-life of 51.9 (10) minutes based on the limitation of relative statistical weight method (LWM). There is no evidence for the production of ⁹⁴Ru (half-life of 52.1 (6) minutes), or potential complication of transient equilibrium that would be associated with such an occurrence. Furthermore, unrelated studies involving the ⁹⁴Mo(d,2n) reaction that focus on the resulting gamma ray emissions have shown that only Tc activities are produced, without the presence of the 367-, 525- and 892-keV gamma rays of ⁹⁴Ru (1969Ba09). All half-life measurements of ^{94m}Tc as produced via the ⁹⁴Mo(d,2n) and ⁹⁴Mo(p,n) reactions are judged to be valid, while under certain circumstances (e.g. adoption of ⁹²Mo(α ,2n)⁹⁴Ru(EC) ^{94m}Tc and ⁹⁴Mo(α ,4n)⁹⁴Ru(EC) ^{94m}Tc as modes of preparation), the presence of parent ⁹⁴Ru with an extremely similar half-life of 52.1 (6) minutes will complicate data analyses.

Reference	Half-life (min)
1948Mo19	50 (2)
1950Me21) 52.5 (15)
)	51.5 (10)
1962Mo06	53 (2)
Recommended value	51.9 (10)*

* Recommended uncertainty has been adjusted from ± 0.8 to 1.0, in alignment with the smallest uncertainty of the values used to calculate the average value.

Half-life of ⁹⁴Tc ground state

The measurements of 1962Mo06, 1963Ma21 and 1965Ba48 were adopted to give a weighted mean half-life of 293 (1) minutes based on the limitation of relative statistical weight method (LWM).

Reference	Half-life (min)
1962Mo06	270 (12)
1963Ma21	293 (1)
1965Ba48	295 (10)
Recommended value	293 (1)

Half-life of parent ⁹⁴Ru

⁹⁴Ru undergoes 100% EC decay through the population of the 443(3)- and 968(3)-keV nuclear levels of ⁹⁴Tc and the 367-, 525- and 892-keV gamma transitions to populate the ^{94m}Tc metastable level at 76 (3) keV. No other gamma-ray emissions are identified with the EC decay of ⁹⁴Ru. With parent ⁹⁴Ru (half-life of 52.1 (6) min) and ^{94m}Tc as a daughter radionuclide (half-life of 51.9 (10) min), the determination of these almost identical half-lives necessitated additional investigation to ensure these two separate values are valid and not a consequence of transient equilibrium being established between parent and daughter prior to measurements.

Reference	Half-life (min)
1967Ei01	53 ± 1
1968Bo27	51.8 ± 0.6
Recommended value	52.1 ± 0.6*

* Recommended uncertainty has been adjusted from ± 0.5 to 0.6, in alignment with the smallest uncertainty of the values used to calculate the average value.

Eichler et al. (1967Ei01) produced ⁹⁴Ru by means of the ⁹⁴Mo(α,4n)⁹⁴Ru and ⁹²Mo(α,2n)⁹⁴Ru with 58- and 32-MeV α particles, respectively. This material was purified by distillation and precipitation, and the only contaminant activities were observed to be ⁹⁵Ru and ^{94m}Tc. Little to no positron emission was observed to occur in the ⁹⁴Ru decay process on the basis of the 511-keV annihilation radiation, which only increased substantially with the in-growth of ^{94m}Tc. A ⁹⁴Ru half-life of (53 ± 1) min was measured by following the decay of only the 367.2(5)-, 525(1)- and 891.2(5)-keV gamma rays, which constitute the primary decay signatures of ⁹⁴Ru (and are not attributed to ^{94m}Tc); other peaks in the spectrum were assigned to ^{94m}Tc and ⁹⁵Ru.

Boswell and McGee (1968Bo27) also produced ⁹⁴Ru by irradiation of natural Mo foil with 80-100 μA beam of 40-MeV α particles to induced the ⁹⁴Mo(α,4n)⁹⁴Ru and ⁹²Mo(α,2n)⁹⁴Ru reactions. Subsequent purification included dissolution of the target and distillation of ruthenium tetraoxide (RuO₄), followed by removal of Tc daughters by a milking procedure with a growth period of 20 minutes between each of six milkings. Decay of each milking sample was followed for seven days by means of a NaI(Tl) well-type detector. Corrected yields of ⁹⁴Tc were determined and plotted against time of separation to give a half-life for parent ⁹⁴Ru of (51.8 ± 0.6) min. The gamma-ray spectrum emitted from the decay of an isolated ruthenium fraction was also determined by means of NaI(Tl) and Ge(Li) detectors – both the 367- and 890-keV gamma rays assigned to ⁹⁴Ru were determined to decay with a half-life of (53 ± 1) min.

Q-value

Q_{EC}-value for ^{94m}Tc EC/β⁺ decay of 4332 (5) keV was adopted from Wang *et al.* (2012Wa38).

Gamma-ray energies and emission probabilitiesEnergies

The well-defined nuclear level energies of 2006Ab37 were used to calculate the gamma transition energies and their uncertainties, and these data were adjusted to account for gamma recoil in the formulation of recommended gamma-ray emission energies and uncertainties. Greater confidence was placed on this overall method of assignment because of the more wide-ranging origins of the gamma transition data in 2006Ab37. The 1573.76-, 1741.65-, 3331.74- and 3534.32-keV levels are exceptions to this approach – the energies of these particular nuclear levels of ⁹⁴Mo were determined from previously unplaced gamma rays that possess the potential to depopulate nuclear levels derived from the γ-γ coincidence studies of 2003Fr02.

Adopted energies, spins and parities for the nuclear levels of ⁹⁴Tc and ⁹⁴Mo (2003Fr02, 2006Ab37).

Nuclear level number	Nuclear level energy (keV)	Spin and parity	Half-life
0	0.0	7 +	293 (1) min ⁹⁴ Tc
1	76 (3)	(2) +	51.9 (10) min ^{94m} Tc
0	0.0	0 +	stable ⁹⁴ Mo
1	871.098 (16)	2 +	
2	1573.76 (4)*	4 +	
3	1741.65 (15)*	0 +	
4	1864.31 (5)	2 +	
5	2067.35 (6)	2 +	
6	2294.79 (16)	4 +	
7	2393.02 (6)	2 +	
8	2423.45 (9)	6 +	
9	2533.87 (12)	3 –	
10	2610.57 (16)	5 –	
11	2739.91 (7)	1 +	
12	2805.04 (19)	3 +	
13	2869.90 (8)	2 +	
14	2872.40 (11)	6 +	
15	2955.55 (13)	(8 +)	
16	2965.41 (6)	3 +	
17	3128.66 (7)	1 +	
18	3163.29 (19)	(3 +)	
19	3165.77 (9)	6 +	
20	3331.74 (17)*	(3 +)	
21	3339.54 (17)	6 +	
22	3400.83 (17)		
23	3447.6 (4)	(1, 2 +)	
24	3511.86 (14)	1 (+)	
25	3534.32 (9)*	2 +	
26	3792.87 (15)	2 +	
27	3892.16 (7)	(2 +)	

* Nuclear level assignment based originally on studies of other related nuclear processes, but introduced in the proposed ^{94m}Tc decay scheme to explain observations of previously unplaced gamma-ray emissions.

Emission Probabilities

All of the relative emission probabilities were suitably quantified in terms of the emission probability of the 871.094-keV gamma ray (100.0%). Sufficiently detailed decay data measurements were only reported in gamma-ray spectroscopic studies by 1969Ba09 and 1986AgZX, and γ - γ coincidence studies of 2003Fr02. When appropriate, weighted-mean analyses have been undertaken for the recommendation of the most significant gamma-ray emission probabilities. Although lacking necessary descriptive detail, the measurements of Ageev *et al.* (1986AgZX) and Fransen *et al.* (2003Fr02) are the most substantial, and therefore their data were adopted for many of the low-intensity gamma-ray emissions in preference to any semi-artificial weighting procedure. This approach was occasionally fortified with data supportive towards the emission of particular gamma rays by Sugiyama and Kikuchi (1976Su04), as taken from their nuclear-level studies of ⁹⁴Mo populated by the ⁹⁴Mo(n,n' γ)⁹⁴Mo reaction. The comprehensive studies of 2003Fr02 focused on γ intensity ratios from individual nuclear levels of ⁹⁴Mo rather than the determination of absolute γ -ray emission probabilities, and therefore not all of these observed transitions could be suitably quantified in the desired manner – while listed in the tabulations below, these particular γ rays are not included in the decay-data file because their absolute emission probabilities remain unknown. Nineteen previously unplaced gamma-ray emissions were re-considered from the point of view of the known nuclear levels of ⁹⁴Mo. A number of these gamma-ray emissions can be incorporated into the proposed decay scheme: 875.1, 998.2, 1499.0, 1670.1, 1757.9, 1769.9, 2027.5, 2257.5, 2664.1, 2869.9, 3400.8 and 3640.6 keV. Despite these efforts, six observed gamma rays remain unplaced in the proposed decay scheme, although they are contained within the recommended data set: 1037.2, 1357.4, 3065.6, 3085.8, 3640.6 and 4136.2 keV.

Gamma transition energies, and measured and recommended gamma-ray energies.

E _{TP} (keV) [#]	E _γ (keV)								
	1962Mo06	1964Ha29	1968Ar06	1968Ka25 [‡]	1969Ba09	1976Su04	1986AgZX*	2003Fr02	Recommended [§]
293.37 (14)	–	–	–	293 (1)	–	–	–	293.4 (1)	293.37 (14)
325.67 (9)	–	–	–	–	–	–	–	325.7 (3)	325.67 (9)
358.3 (3)	–	–	–	–	–	–	–	358.0 (5)	358.3 (3)
448.95 (14)	–	–	–	449.0 (10)	–	–	–	449.0 (1)	448.95 (14)
466.52 (13)	–	–	–	–	–	466.7 (8)	–	466.4 (3)	466.52 (13)
–	511	511	–	–	–	–	–	–	annihilation radiation
528.71 (8)	–	–	–	–	–	–	–	528.7 (3)	528.71 (8)
532.10 (16)	–	–	–	532.0 (10)	–	–	–	532.1 (1)	532.10 (16)
669.56 (13)	–	–	–	–	–	–	–	669.6 (2)	669.56 (13)
672.56 (9)	–	–	–	–	–	–	–	672.0 (7)	672.56 (9)
702.66 (4)	–	–	–	704.0 (8)	–	702.5 (5)	–	702.63 (10)	702.66 (4) ⁰
721.03 (17)	–	–	–	–	–	721.5 (5)	–	721.0 (2)	721.03 (17)
742.32 (13)	–	–	–	743.0 (15)	–	–	–	742.2 (2)	742.32 (13)
802.55 (10)	–	–	–	–	–	–	–	802.6 (2)	802.55 (10)
849.69 (10)	–	–	–	851.0 (10)	–	–	–	849.7 (1)	849.69 (10)
870.55 (22)	(((((871.4 (5)	((870.55 (22) [†]
871.098 (16)	870 (5) (870.7 (7) (870.9(2) (872.5 (10) (871.05 (7) (871.4 (5)	871.0 (3) (871.09 (10) (871.094 (16)
875.60 (9)	(((((–	875.1 (3)	875.5 (2)	875.60 (9)
898.06 (9)	–	–	–	–	–	–	–	898.1 (1)	898.06 (9)
916.09 (19)	–	–	–	917.0 (10)	–	–	–	916.2 (1)	916.09 (19)
940.73 (20)	–	–	–	–	–	–	–	–	940.72 (20)
960.11 (13)	–	–	–	–	–	961.5 (10)	–	960.1 (3)	960.10 (13)
993.21 (5)	–	–	993.1 (5)	–	993.19 (9)	993.1 (8)	993.4 (3)	993.1 (1)	993.20 (5)
998.26 (17)	–	–	–	–	998.1 (10)	–	998.2 (3)	998.2 (2)	998.25 (17)
1005.59 (9)	–	–	–	–	1005.8 (3)	–	1006.1 (3)	1005.5 (1)	1005.58 (9)
–	–	–	–	–	–	–	1022.2 (3)	–	(511 + 511) keV sum peak
1037.2 (3) ^Δ	–	–	–	–	–	–	1037.2 (3)	1036.8 (2)	1037.2 (3) ^Δ
1061.31 (9)	–	–	–	–	–	–	–	1061.1 (5)	1061.30 (9)
1101.10 (8)	–	–	–	–	–	–	1101.3 (3)	1101.1 (1)	1101.09 (8)
1196.25 (6)	–	–	1196.2 (7)	–	1196.4 (3)	–	1196.4 (3)	1196.2 (1)	1196.24 (6)
1231.28 (19)	–	–	–	–	–	–	–	–	1231.27 (19)
1264.35 (9)	–	–	–	–	1264.9 (4)	–	1264.7 (3)	1264.3 (1)	1264.34 (9)
1357.4 (15) ^Δ	–	–	–	–	1357.4 (15)	–	–	–	1357.4 (15) ^Δ
1391.65 (7)	–	–	–	–	–	1391.4 (10)	–	1391.6 (1)	1391.64 (7)
1399.85 (16)	–	–	–	–	–	–	–	1399.9 (2)	1399.84 (16)
1423.69 (16)	–	–	–	–	–	–	–	1423.7 (3)	1423.68 (16)
1467.43 (18)	–	–	–	–	–	–	–	1467.3 (3)	1467.42 (18)

Comments on evaluation

^{94m}Tc

1499.14 (9)	-	-	-	-	-	-	1499.0 (3)	1499.1 (1)	1499.13 (9)
1521.92 (6)	1530 (10)	1530 (70)	1522.0 (3)	1510 (2)	1522.1 (2)	1522.3 (10)	1521.7 (3)	1521.8 (1)	1521.91 (6)
1536.52 (18)	-	-	-	-	-	-	-	1536.5 (2)	1536.51 (18)
1592.01 (10)	-	-	-	1593 (2)	-	-	-	1592.0 (1)	1592.00 (1)
1662.77 (12)	-	-	-	-	-	1663.1 (10)	-	1662.7 (3)	1662.75 (12)
1670.01 (10)	-	-	-	-	-	-	1670.1 (3)	1670.0 (1)	1669.99 (10)
1757.98 (17)	-	-	-	1767 (2)	1757.9 (10)	-	1757.9 (3)	1758.0 (2)	1757.96 (17)
1770.21 (21)	-	-	-	-	-	-	1769.9 (3)	1770.4 (2)	1770.19 (21)
1824.81 (9)	-	-	-	-	-	-	-	1824.9 (3)	1824.79 (9)
1864.31 (5)	1870 (20) (1870 (70) (-	1868 (1864.2	1864.5 (10)	1864.4 (3)	1864.3 (2)	1864.29 (5)
1868.81 (7)	((1868.8 (3)	(1868.68 (8)	1869 (2)	1868.6 (3)	1868.8 (1)	1868.79 (7)
1928.56 (16)	-	-	-	-	1928.8 (20)	-	1928.2 (3)	1928.5 (2)	1928.54 (16)
1933.94 (19)	-	-	-	-	-	-	-	-	1933.92 (19)
1998.80 (8)	-	-	-	-	-	-	-	1998.9 (2)	1998.78 (8)
2027.85 (9)	-	-	-	-	-	-	2027.5 (3)	2027.9 (2)	2027.83 (9)
2067.35 (6)	-	-	-	-	2067.4 (5)	-	2066.7 (3)	2067.4 (1)	2067.33 (6)
2094.31 (6)	-	-	-	-	-	-	-	2094.3 (1)	2094.28 (6)
2257.56 (7)	-	-	-	-	-	-	2257.5 (3)	2257.6 (1)	2257.53 (7)
2292.19 (19)	-	-	-	-	-	-	2292.2 (3)	2292.2 (2)	2292.16 (19)
2393.02 (6)	2410 (20)	2410 (100)	2393.0 (7)	-	2393.2 (4)	-	2392.6 (3)	2393.1 (1)	2392.99 (6)
2460.64 (17)	-	-	-	-	-	-	-	2460.8 (8)	2460.61 (17)
2529.73 (17)	-	-	2530.0 (10)	-	2529.8 (3)	2530.9 (20)	2529.6 (3)	2529.7 (3)	2529.69 (17)
2576.5 (4)	-	-	-	-	2577.2 (20)	2577.9 (20)	2577.2 (3)	2576.5 (5)	2576.5 (4)
2640.76 (14)	-	-	-	-	2641.6 (15)	-	2640.6 (3)	2640.7 (3)	2640.72 (14)
2663.22 (9)	-	-	-	-	2664.1 (20)	-	-	2663.2 (2)	2663.18 (9)
2739.91 (7)	2730 (20)	2740 (100)	2740.0 (7)	2740	2740.1 (3)	2741.8 (20)	2739.8 (3)	2739.9 (1)	2739.87 (7)
2869.90 (8)	-	-	-	-	-	-	2869.9 (3)	2870.0 (2)	2869.85 (8)
3021.06 (7)	-	-	-	-	3021.6 (10)	-	3020.9 (3)	3021.0 (1)	3021.01 (7)
3065.7 (3) ^Δ	-	-	-	-	-	-	3065.6 (3)	-	3065.6 (3) ^Δ
3085.9 (3) ^Δ	-	-	-	-	-	-	3085.8 (3)	-	3085.8 (3) ^Δ
3128.66 (7)	3200 (30)	3200 (100)	3129.2 (7)	-	3129.1 (5)	-	3128.6 (3)	3128.5 (2)	3128.60 (7)
3400.83 (17)	-	-	-	-	-	-	3400.8 (3)	-	3400.76(17)
3447.6 (4)	-	-	-	-	-	-	3447.0 (3)	3447.5 (10)	3447.5 (4)
3511.86 (14)	-	-	-	-	3512.5 (15)	-	3511.8 (3)	3511.6 (2)	3511.79 (14)
3534.32 (9)	-	-	-	-	-	-	-	3534.0 (4)	3534.25 (9)
3640.7 (3) ^Δ	-	-	-	-	-	-	3640.6 (3)	-	3640.6 (3) ^Δ
3792.87 (15)	-	-	3793.5 (10)	-	3793.1 (15)	-	3792.5 (3)	3792.3 (10)	3792.79 (15)
3892.16 (7)	-	-	-	-	3892.7 (25)	-	3891.7 (3)	3891.6 (10)	3892.07 (7)
4136.3 (3) ^Δ	-	-	-	-	-	-	4136.2 (3)	-	4136.2 (3) ^Δ

Determined from the nuclear level energies of 2006Ab37.

‡ All gamma-ray energies listed as reported by 1968Ka25, but only 872.5 (10)-, 1868- and 2740-keV gamma rays were identified with the EC/ β^+ decay of ^{94m}Tc – as defined in their proposed decay schemes, all other gamma rays were assigned to EC/ β^+ decay of ^{94}Tc .

* All uncertainties defined as ± 0.3 keV.

§ Calculated by subtracting gamma recoil from gamma transition energy (E_{TP} (keV)).

⁰ Introduced as depopulating (1573.76 – 871.098)-keV transition from proposed 1573.76-keV nuclear level, although unobserved in published ^{94m}Tc gamma-ray measurements.

[†] Introduced as depopulating (1741.65 – 871.098)-keV transition from proposed 1741.65-keV nuclear level, although unobserved in published ^{94m}Tc gamma-ray measurements.

^Δ Unplaced in proposed decay scheme.

Previously unplaced gamma transitions (2006Ab37).

E_γ (keV)	P_γ^{rel}	Assignment	Comments
875.1 (3)	1.1 (3)	2739.91 → 1864.31	Observed by 1986AgZX, and supported by the studies of 2003Fr02.
998.2 (3)	0.25 (2)	2739.91 → 1741.65	Observed by 1969Ba09 and 1986AgZX, and supported by the studies of 2003Fr02.
1022.2 (3)	0.029 (15)	N/A	(511 + 511) keV sum peak.
1037.2 (3)	0.047 (15)	–	Only observed by 1986AgZX.
1357.4 (15)	0.20 (8)	–	Only observed by 1969Ba09.
1499.0 (3)	0.071 (12)	3892.16 → 2393.02	Observed by 1986AgZX, and supported by the studies of 2003Fr02.
1670.1 (3)	0.039 (2)	3534.32 → 1864.31	Observed by 1986AgZX, and supported by the studies of 2003Fr02.
1757.9 (3)	0.16 (2)	3331.74 → 1573.76	Observed by 1969Ba09 and 1986AgZX, and supported by the studies of 2003Fr02.
1769.9 (3)	0.027 (7)	3511.86 → 1741.65	Observed by 1986AgZX, and supported by the studies of 2003Fr02.
2027.5 (3)	0.022 (4)	3892.16 → 1864.31	Observed by 1986AgZX, and supported by the studies of 2003Fr02.
2257.5 (3)	0.061 (5)	3128.66 → 871.098	Observed by 1986AgZX, and supported by the studies of 2003Fr02.
2664.1 (20)	0.070 (2)	3534.32 → 871.098	Observed by 1969Ba09, and supported by the studies of 2003Fr02.
2869.9 (3)	0.017 (2)	2869.90 → 0.0	Observed by 1986AgZX, and supported by the studies of 2003Fr02.
3065.6 (3)	0.012 (4)	–	Only observed by 1986AgZX.
3085.8 (3)	0.017 (4)	–	Only observed by 1986AgZX.
3400.8 (3)	0.005 (2)	3400.83 → 0.0	Only observed by 1986AgZX.
3447.0 (3)	0.006 (1)	3447.6 → 0.0	Observed by 1986AgZX, and supported by the studies of 2003Fr02.
3640.6 (3)	0.007 (2)	–	Only observed by 1986AgZX.
4136.2 (3)	0.007 (1)	–	Only observed by 1986AgZX.

Another point of note involves summed gamma peaks that could also be potentially observed:

- unplaced 1022.2-keV gamma transition is believed to be the sum peak of (511 + 511);
- placed 1864.79-keV gamma could partially contain or fully be the sum peak of (871.09 + 993.20);
- placed 2392.99-keV gamma could partially contain the sum peak of (871.09 + 1521.91);
- placed 2739.87-keV gamma could partially contain the sum peak of (871.09 + 1868.79).

Multiple assignments of this nature are very difficult to address when experimental details are lacking. Therefore, these possible summing effects have been ignored, other than defining the 1022.2-keV emission as a (511 + 511) sum peak.

The existence of the 1864.29-keV γ ray is uncertain, being effectively unobserved by 1969Ba09 (assigned relative emission probability ≤ 0.25), but detected by 1986AgZX (relative emission probability of 0.41 (4)). Extensive γ -ray studies by 2003Fr02 indicate the depopulation of the 1864.31-keV nuclear level of ⁹⁴Mo by means of 993.20- and 1864.29-keV γ transitions. Under these circumstances, the relative emission probabilities of these two γ rays as determined by 2003Fr02 have been adopted to determine a relative emission probability of 0.24 (3) % for the previously ill-defined 1864.29-keV γ ray.

As proposed, population of the 1741.65-keV nuclear level by the 998.25- and 1770.19-keV γ rays requires depopulation by an unobserved 870.55-keV γ transition. Similarly, population of the 1573.76-keV nuclear level by the 1757.96-keV γ ray implies depopulation by an unobserved 702.66-keV γ transition. Both of these ill-defined depopulating γ rays have been added to the proposed decay scheme, with relative emission probabilities based on the values of two (and one) transition probabilities populating the original nuclear levels. One implication of the above is that the 871-keV γ emission constitutes an unresolved doublet (870.55 and 871.094 keV) – however, the relative emission probability of the postulated 870.55-keV γ ray is so small at 0.28 (3) % compared with the main 871.094-keV γ ray of 100% (99.72%) as to be of negligible impact on calculations of all of the absolute γ -ray emission probabilities.

The studies of Fransen *et al.* are particularly noteworthy from the point of view of their comprehensive nature when compared with all other previous measurements (2003Fr02). Their experimental studies included photon scattering experiments, γ - γ coincidence measurements of the EC/ β^+ decay of ^{94m}Tc, and in-beam γ - γ coincidence studies of the emissions from the ⁹¹Zr(α ,n)⁹⁴Mo and ⁹⁴Mo(n,n' γ) reactions. Much quantitative detail was derived from these studies concerning γ emissions and their mixing ratios, along with the identification of 57 nuclear levels of ⁹⁴Mo, of which seventeen were newly observed. Only nuclear level data of direct relevance to their γ - γ coincidence measurements of the EC/ β^+ decay of ^{94m}Tc are given below in tabulated form, and assisted greatly in the derivation of a more comprehensive decay scheme. Unfortunately, the relative γ -ray emission probabilities derived by 2003Fr02 apply only in terms of depopulating groups of γ transitions for a particular nuclear level – therefore, absolute emission

probabilities for the full γ -ray decay cannot be determined from this source. Some of these parameters can be calculated in conjunction with the equivalent 1969Ba09/1986AgZX singles measurements, but this approach was not possible when direct γ equivalence data were unavailable (as was found to be the case for the 293.37-, 448.95-, 466.52-, 532.10-, 669.56-, 702.66-, 721.03-, 742.32-, 849.69-, 870.55-, 916.09-, 940.72-, 960.10-, 1231.27-, 1423.68-, 1592.00-, 1662.75- and 1933.97-keV γ rays). With such a significant lack of emission probability data, these particular γ transitions have not been included in the proposed decay scheme, although relevant nuclear levels have been noted.

Nuclear levels of ⁹⁴Mo and γ transitions of relevance to the EC/ β^+ decay of ^{94m}Tc – as listed below, I_γ are sets of relative intensities as applied only to individual groups of γ emissions depopulating each specific nuclear level (2003Fr02).

E_x (keV)	J_i^π	J_f^π	E_γ (keV)*	I_γ (%)	$\delta(J_i^\pi \rightarrow J_f^\pi)$	P_γ^{rel} (%)*
871.09 (10)	2 ₁ ⁺	0 ₁ ⁺	871.09 (10)	100.0	0	100
1573.72 (14)	4 ₁ ⁺	2 ₁ ⁺	702.63 (10)	100.0	0.00 (4)	–
1741.65 (15) [†]	0 ₂ ⁺	2 ₁ ⁺	870.55 (22)	100.0	–	–
1864.3 (1)	2 ₂ ⁺	0 ₁ ⁺	1864.3 (2)	10.3 (10)	0	0.24 (3) / 0.24 (3)
		2 ₁ ⁺	993.1 (1)	100.0 (10)	– 2.0 (10)	2.35 (19)
2067.4 (1)	2 ₃ ⁺	0 ₁ ⁺	2067.4 (1)	15.1 (7)	0	0.13 (1) / 0.12 (1)
		2 ₁ ⁺	1196.2 (1)	100.0 (7)	0.15 (4)	0.76 (7)
2294.7 (2)	4 ₂ ⁺	2 ₁ ⁺	1423.7 (3)	13.3 (2)	0.08 (8)	–
		4 ₁ ⁺	721.0 (2)	100.0 (2)	0.03 (4)	–
2393.1 (1)	2 ₄ ⁺	0 ₁ ⁺	2393.1 (1)	11.11 (22)	0	0.52 (5) / 0.53 (4)
		2 ₁ ⁺	1521.8 (1)	100.0 (20)	– 0.12 (3)	4.76 (30)
		2 ₂ ⁺	528.7 (3)	0.719 (33)	–	0.034 (2)
		2 ₃ ⁺	325.7 (3)	0.61 (4)	–	0.029 (2)
2423.4 (2)	6 ₁ ⁺	4 ₁ ⁺	849.7 (1)	100.0	– 0.04 (5)	–
2533.8 (3)	3 ₁ [–]	2 ₁ ⁺	1662.7 (3)	100.0 (22)	0.03 (7)	–
		4 ₁ ⁺	960.1 (3)	81.3 (31)	0.00 (2)	–
		2 ₂ ⁺	669.6 (2)	31.9 (13)	– 0.03 (13)	–
		2 ₃ ⁺	466.4 (3)	57.3 (10)	0.00 (3)	–
2610.5 (2) ?	5 ₁ [–]	4 ₁ ⁺	1036.8 (2) ?	100.0	0.00 (4)	–
2739.9 (1)	1 ₁ ⁺	0 ₁ ⁺	2739.9 (1)	65.4 (13)	0	3.69 (26) / 3.82 (21)
		2 ₁ ⁺	1868.8 (1)	100.0 (20)	– 0.12 (2)	5.84 (30)
		0 ₂ ⁺	998.2 (2)	4.44 (10)	0	0.23 (2) / 0.26 (2)
		2 ₂ ⁺	875.5 (2)	24.4 (5)	– 0.10 (2)	0.84 (20) / 1.42 (8)
		2 ₃ ⁺	672.0 (7)	3.02 (48)	–	0.18 (3)
2805.0 (3) [†]	3 ₁ ⁺	2 ₁ ⁺	1933.9 (4)	75.7 (30)	–0.66 (14) or – 1.7 ^{+0.4} _{–0.5}	–
		4 ₁ ⁺	1231.2 (3)	100.0 (53)	7.6 ^{+4.7} _{–3.0}	–
		2 ₂ ⁺	940.7 (4)	62.9 (39)	2.3 ^{+0.7} _{–0.5}	–
2870.0 (2)	2 ₅ ⁺	0 ₁ ⁺	2870.0 (2)	17.3 (5)	0	0.022 (7) / 0.017 (2)
		2 ₁ ⁺	1998.9 (2)	13.1 (6)	1.3 ^{+1.4} _{–0.4}	0.0131 (6)
		2 ₂ ⁺	1005.5 (1)	100.0 (36)	– 0.05 (4)	0.10 (3)
		2 ₃ ⁺	802.6 (2)	26.2 (15)	–	0.0262 (15)
2872.4 (2)	6 ₂ ⁺	6 ₁ ⁺	449.0 (1)	100.0	0.14 (6)	–
2955.5 (3)	(8 ₁ ⁺)	6 ₁ ⁺	532.1 (1)	100.0	– 0.03 (5)	–
2965.3 (2)	3 ₂ ⁺	2 ₁ ⁺	2094.3 (1)	36.9 (14)	1.1 ^{+1.0} _{–0.4}	0.0166 (6)
		4 ₁ ⁺	1391.6 (1)	63.0 (24)	– 0.08 (6)	0.0284 (11)
		2 ₂ ⁺	1101.1 (1)	100.0 (23)	– 0.09 (6)	0.045 (15)
		2 ₃ ⁺	898.1 (1)	23.0 (12)	2.0 ^{+1.2} _{–0.6} or 0.39 (25)	0.0104 (5)
3128.6 (2)	1 ₂ ⁺	0 ₁ ⁺	3128.5 (2)	100.0 (3)	0	1.42 (10)
		2 ₁ ⁺	2257.6 (1)	4.29 (10)	0.74 ^{+0.21} _{–0.17}	0.056 (18) / 0.061 (5)
		2 ₂ ⁺	1264.3 (1)	18.27 (37)	– 0.08 (3)	0.21 (2) / 0.26 (2)
		2 ₃ ⁺	1061.1 (5)	1.16 (11)	–7.0 ^{+3.0} _{–20.0} or – 0.57 (16)	0.017 (2)
3163.3 (3)	(3 ₄ ⁺)	2 ₁ ⁺	2292.2 (2)	100.0 (13)	0.17 (4)	0.053 (18)
		3 ₁ ⁺	358.0 (5)	16.7 (13)	– 0.35 (12)	0.0089 (7)
3165.8 (2)	6 ₃ ⁺	4 ₁ ⁺	1592.0 (1)	100.0 (35)	– 0.01 (6)	–
		6 ₁ ⁺	742.2 (2)	29.4 (11)	0.15 (7)	–
		6 ₂ ⁺	293.4 (1)	79.4 (25)	0.18 (5)	–
3331.7 (3)	(3 ⁺)	2 ₁ ⁺	2460.8 (8)	7.4 (11)	–	0.012 (2)
		4 ₁ ⁺	1758.0 (2)	100.0 (55)	– 0.10 (3)	0.16 (2)
		2 ₂ ⁺	1467.3 (3)	47.9 (29)	0.3 ^{+2.9} _{–0.2}	0.077 (5)
3339.6 (3)	6 ₄ ⁺	6 ₁ ⁺	916.2 (1)	100.0	– 0.02 (7)	–
3400.8 (2) [†]	–	0 ₁ ⁺	3400.76 (17)	–	–	0.005 (2)
		2 ₁ ⁺	2529.7 (3)	100.0 (8)	–	0.36 (4)

		2 ₂ ⁺	1536.5 (2)	4.2 (8)	–	0.015 (3)
3447.6 (5)	(2)	0 ₁ ⁺	<u>3447.5 (10)</u>	5.1 (3)	0	<u>0.005 (2)</u> / 0.006 (1)
		2 ₁ ⁺	<u>2576.5 (5)</u>	100.0 (3)	–1.9 ^{+0.5} _{–0.6} or – 0.08 (10)	<u>0.12 (2)</u>
3511.7 (2)	1 ₃ ⁽⁺⁾	0 ₁ ⁺	<u>3511.6 (2)</u>	100.0 (11)	0	<u>0.067 (7)</u>
		2 ₁ ⁺	<u>2640.7 (3)</u>	51.6 (13)	–	<u>0.037 (9)</u> / 0.035 (4)
		0 ₂ ⁺	<u>1770.4 (2)</u>	48.6 (92)	0	<u>0.020 (8)</u> / 0.033 (7)
3534.3 (2)	2 ₈ ⁺	0 ₁ ⁺	3534.0 (4)	5.09 (55)	0	0.0036 (4)
		2 ₁ ⁺	<u>2663.2 (2)</u>	100.0 (23)	– 0.3 (2)	<u>0.07 (6)</u> → 0.070 (2)
		2 ₂ ⁺	<u>1670.0 (1)</u>	56.1 (20)	0.15 (19)	<u>0.037 (12)</u> / 0.04 (3) → 0.039 (2)
3792.8 (3)	2 ₉ ⁺	0 ₁ ⁺	<u>3792.3 (10)</u>	77.8 (20)	0	<u>0.054 (8)</u> / 0.062 (16)
		2 ₂ ⁺	<u>1928.5 (2)</u>	100.0 (40)	–	<u>0.08 (2)</u>
		2 ₄ ⁺	<u>1399.9 (2)</u>	54.9 (30)	–	0.044 (3)
3892.2 (2)	(2 ⁺)	0 ₁ ⁺	<u>3891.6 (10)</u>	17.4 (9)	–	<u>0.014 (2)</u> / 0.016 (3)
		2 ₁ ⁺	<u>3021.0 (1)</u>	100.0 (24)	–	<u>0.093 (15)</u>
		2 ₂ ⁺	<u>2027.9 (2)</u>	22.3 (10)	–	<u>0.025 (6)</u> / 0.021 (4)
		2 ₃ ⁺	1824.9 (3)	25.8 (10)	–	0.024 (1)
		2 ₄ ⁺	<u>1499.1 (1)</u>	79.4 (22)	–	<u>0.062 (20)</u> / 0.074 (12)

* Underscored entries for E_γ and P_γ^{rel} denote previously observed/determined data; other P_γ^{rel} data have been calculated from relevant ratios of I_γ and P_γ^{rel} as tabled.

† While not specified in the studies of 2003Fr02, the proposed 1741.65-keV nuclear level arises from the observed population of this level by the 998.2- and 1770.4-keV γ transitions.

‡ Although not identified in the EC/β⁺ decay studies of ^{94m}Tc by 2003Fr02, the 2805.0- and 3400.8-keV nuclear levels have been introduced as a consequence of the observation of the 358.0-keV γ transition that populates the 2805.0-keV level (2003Fr02), and the 2529.7 and 3400.76-keV γ transitions that depopulate the 3400.8-keV level (1969Ba09, 1986AgZX).

Measured and recommended gamma-ray emission probabilities relative to P_γ(871.094 keV) of 100%.

E _γ (keV)	P _γ ^{rel} *							
	1962Mo06	1964Ha29	1965Ba48	1968Ar06	1968Ka25 [†]	1969Ba09	1986AgZX	2003Fr02 [†]
γ _{19,14} 293.37 (14)	–	–	–	–	–	–	–	unquantified
γ _{7,5} 325.67 (9)	–	–	–	–	–	–	–	0.029 (2)
γ _{18,12} 358.3 (3)	–	–	–	–	–	–	–	0.0089 (7)
γ _{14,8} 448.95 (14)	–	–	–	–	–	–	–	unquantified
γ _{9,5} 466.52 (13)	–	–	–	–	–	–	–	unquantified
γ [±] 511	144 (12) → 158 (13)	–	160	–	–	–	–	–
γ _{7,4} 528.71 (8)	–	–	–	–	–	–	–	0.034 (2)
γ _{15,8} 532.10 (16)	–	–	–	–	–	–	–	unquantified
γ _{9,4} 669.56 (13)	–	–	–	–	–	–	–	unquantified
γ _{11,5} 672.56 (9)	–	–	–	–	–	–	–	0.18 (3)
γ _{2,1} 702.66 (4)	–	–	–	–	–	–	–	unquantified
γ _{6,2} 721.03 (17)	–	–	–	–	–	–	–	unquantified
γ _{19,8} 742.32 (13)	–	–	–	–	–	–	–	unquantified
γ _{13,5} 802.55 (10)	–	–	–	–	–	–	–	0.0262 (15)
γ _{8,2} 849.69 (10)	–	–	–	–	–	–	–	unquantified
γ _{3,1} 870.55 (22)	–	–	–	–	–	–	–	–
γ _{1,0} 871.094 (16)	91 → 100	91 → 100	100	100	100 (5)	100	100	unquantified
γ _{11,4} 875.60 (9)	–	–	–	–	–	–	0.84 (20)	1.42 (8)
γ _{16,5} 898.06 (9)	–	–	–	–	–	–	–	0.0104 (5)
γ _{21,8} 916.09 (19)	–	–	–	–	–	–	–	unquantified
γ _{12,4} 940.72 (20)	–	–	–	–	–	–	–	–
γ _{9,2} 960.10 (13)	–	–	–	–	–	–	–	unquantified
γ _{4,1} 993.20 (5)	–	–	–	2.6	–	2.35 (30)	2.35 (19)	[2.35 (19)]
γ _{11,3} 998.25 (17)	–	–	–	–	–	0.25 (8)	0.23 (2)	0.26 (2)
γ _{13,4} 1005.58 (9)	–	–	–	–	–	0.16 (8)	0.094 (30)	[0.10 (3)]
2γ [±] 1022	–	–	–	–	–	–	0.029 (15)	–
γ 1037.2 (3) ^Δ	–	–	–	–	–	–	0.047 (15)	unquantified
γ _{17,5} 1061.31 (9)	–	–	–	–	–	–	–	0.017 (2)
γ _{16,4} 1101.09 (8)	–	–	–	–	–	–	0.045 (15)	[0.045 (15)]
γ _{5,1} 1196.24 (6)	–	–	–	0.9	–	0.80 (10)	0.74 (7)	[0.76 (7)]
γ _{12,2} 1231.27 (19)	–	–	–	–	–	–	–	–
γ _{17,4} 1264.34 (9)	–	–	–	–	–	0.23 (8)	0.21 (2)	0.26 (2)
γ 1357.4 (15) ^Δ	–	–	–	–	–	0.20 (8)	–	–
γ _{16,2} 1391.64 (7)	–	–	–	–	–	–	–	0.0284 (11)
γ _{26,7} 1399.84 (16)	–	–	–	–	–	–	–	0.044 (3)
γ _{6,1} 1423.68 (16)	–	–	–	–	–	–	–	unquantified
γ _{20,4} 1467.42 (18)	–	–	–	–	–	–	–	0.077 (5)
γ _{27,7} 1499.13 (9)	–	–	–	–	–	–	0.062 (20)	0.074 (12)

Comments on evaluation

^{94m}Tc

$\gamma_{7,1}$	1521.91 (6)	10 (2) → 11 (2)	10 (2) → 11 (2)	38	5.9	–	4.80 (30)	4.72 (33)	[4.76 (30)]
$\gamma_{22,4}$	1536.51 (18)	–	–	–	–	–	–	–	0.015 (3)
$\gamma_{19,2}$	1592.00 (10)	–	–	–	–	–	–	–	unquantified
$\gamma_{9,1}$	1662.75 (12)	–	–	–	–	–	–	–	unquantified
$\gamma_{25,4}$	1669.99 (10)	–	–	–	–	–	–	0.037 (12)	0.039 (2)
$\gamma_{20,2}$	1757.96 (17)	–	–	–	–	–	0.16 (4)	0.16 (2)	[0.16 (2)]
$\gamma_{24,3}$	1770.19 (21)	–	–	–	–	–	–	0.020 (8)	0.033 (7)
$\gamma_{27,5}$	1824.79 (9)	–	–	–	–	–	–	–	0.024 (1)
$\gamma_{4,0}$	1864.29 (5)	–	–	–	–	unquantified (≤ 0.25	0.41 (4)	0.24 (3)
$\gamma_{11,1}$	1868.79 (7)	9 (2) → 10 (2)	9 (2) → 10 (2)	21	5.8	(6.10 (30)	5.42 (38)	[5.84 (30)]
$\gamma_{26,4}$	1928.54 (16)	–	–	–	–	–	0.09 (5)	0.08 (2)	[0.08 (2)]
$\gamma_{12,1}$	1933.97 (19)	–	–	–	–	–	–	–	–
$\gamma_{13,1}$	1998.78 (8)	–	–	–	–	–	–	–	0.0131 (6)
$\gamma_{27,4}$	2027.83 (9)	–	–	–	–	–	–	0.025 (6)	0.021 (4)
$\gamma_{5,0}$	2067.33 (6)	–	–	–	–	–	0.09 (3)	0.13 (1)	0.12 (1)
$\gamma_{16,1}$	2094.28 (6)	–	–	–	–	–	–	–	0.0166 (6)
$\gamma_{17,1}$	2257.53 (7)	–	–	–	–	–	–	0.056 (18)	0.061 (5)
$\gamma_{18,1}$	2292.16 (19)	–	–	–	–	–	–	0.053 (18)	[0.053 (18)]
$\gamma_{7,0}$	2392.99 (6)	1.4 (5) → 1.5 (5)	< 1.4 (5) → < 1.5 (5)	–	0.6	–	0.50 (20)	0.52 (5)	0.53 (4)
$\gamma_{20,1}$	2460.61 (17)	–	–	–	–	–	–	–	0.012 (2)
$\gamma_{22,1}$	2529.69 (17)	–	–	–	0.7	–	0.33 (8)	0.37 (4)	[0.36 (4)]
$\gamma_{23,1}$	2576.5 (4)	–	–	–	–	–	0.13 (5)	0.12 (2)	[0.12 (2)]
$\gamma_{24,1}$	2640.72 (14)	–	–	–	–	–	very weak	0.037 (9)	0.035 (4)
$\gamma_{25,1}$	2663.18 (9)	–	–	–	–	–	0.07 (6)	–	0.070 (2)
$\gamma_{11,0}$	2739.87 (7)	5.0 (15) → 5.5 (17)	5.0 (15) → 5.5 (17)	2.2	3.8	unquantified	3.74 (35)	3.66 (26)	3.82 (21)
$\gamma_{13,0}$	2869.85 (8)	–	–	–	–	–	–	0.022 (7)	0.017 (2)
$\gamma_{27,1}$	3021.01 (7)	–	–	–	–	–	0.08 (6)	0.094 (15)	[0.093 (15)]
γ	3065.6 (3) ^Δ	–	–	–	–	–	–	0.012 (4)	–
γ	3085.8 (3) ^Δ	–	–	–	–	–	–	0.017 (4)	–
$\gamma_{17,0}$	3128.60 (7)	2.2 (8) → 2.4 (9)	2.2 (8) → 2.4 (9)	3.3	1.2	–	1.47 (15)	1.39 (10)	[1.42 (10)]
$\gamma_{22,0}$	3400.76 (17)	–	–	–	–	–	–	0.005 (2)	–
$\gamma_{23,0}$	3447.5 (4)	–	–	–	–	–	–	0.005 (2)	0.006 (1)
$\gamma_{24,0}$	3511.79 (14)	–	–	–	–	–	0.06 (2)	0.068 (7)	[0.067 (7)]
$\gamma_{25,0}$	3534.25 (9)	–	–	–	–	–	–	–	0.0036 (4)
γ	3640.6 (3) ^Δ	–	–	–	–	–	–	0.007 (2)	–
$\gamma_{26,0}$	3792.79 (15)	–	–	–	0.1	–	0.05 (2)	0.054 (5)	0.062 (16)
$\gamma_{27,0}$	3892.07 (7)	–	–	–	–	–	0.016 (10)	0.014 (2)	0.016 (3)
γ	4136.2 (3) ^Δ	–	–	–	–	–	–	0.007 (1)	–

Measured and recommended gamma-ray emission probabilities relative to P_γ(871.094 keV) of 100% (continued).

E _γ (keV)	P _γ ^{rel} *	Assessed and/or calculated [†]		Recommended [*]
		1969Ba09/1986AgZX	1969Ba09/1986AgZX/ 2003Fr02	
γ _{19,14} 293.37 (14)	–	–	–	unknown
γ _{7,5} 325.67 (9)	–	–	–	0.029 (2)
γ _{18,12} 358.3 (3)	–	–	–	0.0089 (7)
γ _{14,8} 448.95 (14)	–	–	–	unknown
γ _{9,5} 466.52 (13)	–	–	–	unknown
γ [±] 511	–	–	–	147 (1) ⁰ annihilation radiation
γ _{7,4} 528.71 (8)	–	–	–	0.034 (2)
γ _{15,8} 532.10 (16)	–	–	–	unknown
γ _{9,4} 669.56 (13)	–	–	–	unknown
γ _{11,5} 672.56 (9)	–	–	–	0.18 (3)
γ _{2,1} 702.66 (4)	0.19 (2) ⁺	–	–	0.19 (2) ⁺
γ _{6,2} 721.03 (17)	–	–	–	unknown
γ _{19,8} 742.32 (13)	–	–	–	unknown
γ _{13,5} 802.55 (10)	–	–	–	0.0262 (15)
γ _{8,2} 849.69 (10)	–	–	–	unknown
γ _{3,1} 870.55 (22)	0.28 (3) [§]	–	–	0.28 (3) [§]
γ _{1,0} 871.094 (16)	100	(100)	–	100
γ _{11,4} 875.60 (9)	0.84 (20)	–	1.1 (3)	1.1 (3)
γ _{16,5} 898.06 (9)	–	–	–	0.0104 (5)
γ _{21,8} 916.09 (19)	–	–	–	unknown
γ _{12,4} 940.72 (20)	–	–	–	unknown
γ _{9,2} 960.10 (13)	–	–	–	unknown
γ _{4,1} 993.20 (5)	2.35 (19)	–	–	2.35 (19)
γ _{11,3} 998.25 (17)	0.23 (2)	–	0.25 (2)	0.25 (2)
γ _{13,4} 1005.58 (9)	0.10 (3)	–	–	0.10 (3)
2γ [±] 1022	0.029 (15) (511 + 511) keV sum peak	–	–	0.029 (15) (511 + 511) keV sum peak
γ 1037.2 (3) ^Δ	0.047 (15)	–	–	0.047 (15)
γ _{17,5} 1061.31 (9)	–	–	–	0.017 (2)
γ _{16,4} 1101.09 (8)	0.045 (15)	–	–	0.045 (15)
γ _{5,1} 1196.24 (6)	0.76 (7)	–	–	0.76 (7)
γ _{12,2} 1231.27 (19)	–	–	–	unknown
γ _{17,4} 1264.34 (9)	0.21 (2)	–	0.23 (2)	0.23 (2)
γ 1357.4 (15) ^Δ	0.20 (8)	–	–	0.20 (8)
γ _{16,2} 1391.64 (7)	–	–	–	0.0284 (11)
γ _{26,7} 1399.84 (16)	–	–	–	0.044 (3)
γ _{6,1} 1423.68 (16)	–	–	–	unknown
γ _{20,4} 1467.42 (18)	–	–	–	0.077 (5)

Comments on evaluation

^{94m}Tc

$\gamma_{27.7}$	1499.13 (9)	0.062 (20)	0.071 (12)	0.071 (12)
$\gamma_{7.1}$	1521.91 (6)	4.76 (30)	–	4.76 (30)
$\gamma_{22.4}$	1536.51 (18)	–	–	0.015 (3)
$\gamma_{19.2}$	1592.00 (10)	–	–	unknown
$\gamma_{9.1}$	1662.75 (12)	–	–	unknown
$\gamma_{25.4}$	1669.99 (10)	0.037 (12)	0.039 (2)	0.039 (2)
$\gamma_{20.2}$	1757.96 (17)	0.16 (2)	–	0.16 (2)
$\gamma_{24.3}$	1770.19 (21)	0.020 (8)	0.027 (7)	0.027 (7)
$\gamma_{27.5}$	1824.79 (9)	–	–	0.024 (1)
$\gamma_{4.0}$	1864.29 (5)	0.24 (3) ^s	0.24 (3)	0.24 (3) ^s
$\gamma_{11.1}$	1868.79 (7)	5.84 (30)	–	5.84 (30)
$\gamma_{26.4}$	1928.54 (16)	0.08 (2)	–	0.08 (2)
$\gamma_{12.1}$	1933.97 (19)	–	–	unknown
$\gamma_{13.1}$	1998.78 (8)	–	–	0.0131 (6)
$\gamma_{27.4}$	2027.83 (9)	0.025 (6)	0.022 (4)	0.022 (4)
$\gamma_{5.0}$	2067.33 (6)	0.13 (1)	0.12 (1)	0.12 (1)
$\gamma_{16.1}$	2094.28 (6)	–	–	0.0166 (6)
$\gamma_{17.1}$	2257.53 (7)	0.056 (18)	0.061 (5)	0.061 (5)
$\gamma_{18.1}$	2292.16 (19)	0.053 (18)	–	0.053 (18)
$\gamma_{7.0}$	2392.99 (6)	0.52 (5)	0.53 (4)	0.53 (4)
$\gamma_{20.1}$	2460.61 (17)	–	–	0.012 (2)
$\gamma_{22.1}$	2529.69 (17)	0.36 (4)	–	0.36 (4)
$\gamma_{23.1}$	2576.5 (4)	0.12 (2)	–	0.12 (2)
$\gamma_{24.1}$	2640.72 (14)	0.037 (9)	0.035 (4)	0.035 (4)
$\gamma_{25.1}$	2663.18 (9)	0.07 (6)	–	0.070 (2)
$\gamma_{11.0}$	2739.87 (7)	3.69 (26)	3.75 (21)	3.75 (21)
$\gamma_{13.0}$	2869.85 (8)	0.022 (7)	0.017 (2)	0.017 (2)
$\gamma_{27.1}$	3021.01 (7)	0.093 (15)	–	0.093 (15)
γ	3065.6 (3) ^A	0.012 (4)	–	0.012 (4)
γ	3085.8 (3) ^A	0.017 (4)	–	0.017 (4)
$\gamma_{17.0}$	3128.60 (7)	1.42 (10)	–	1.42 (10)
$\gamma_{22.0}$	3400.76 (17)	0.005 (2)	–	0.005 (2)
$\gamma_{23.0}$	3447.5 (4)	0.005 (2)	0.006 (1)	0.006 (1)
$\gamma_{24.0}$	3511.79 (14)	0.067 (7)	–	0.067 (7)
$\gamma_{25.0}$	3534.25 (9)	–	–	0.0036 (4)
γ	3640.6 (3) ^A	0.007 (2)	–	0.007 (2)
$\gamma_{26.0}$	3792.79 (15)	0.054 (5)	0.055 (5)	0.055 (5)
$\gamma_{27.0}$	3892.07 (7)	0.014 (2)	0.015 (2)	0.015 (2)
γ	4136.2 (3) ^A	0.007 (1)	–	0.007 (1)

^s Emission probabilities expressed relative to P_γ(871.094 keV) of 100%.

^A As reported by 1968Ka25, only 872.5(10)-, 1868- and 2740-keV gamma rays were identified with the EC/β⁺ decay of ^{94m}Tc – all other gamma rays were assigned to EC/β⁺ decay of ⁹⁴Tc.

[†] P_γ^{rel} data of 2003Fr02 are only based on the relative depopulation of individual nuclear levels of ⁹⁴Mo, and are not relative to P_γ(871.094 keV) of 100%, nor expressed per 100 decays of ^{94m}Tc – P_γ^{rel} data cannot always be quantified in the form required, as denoted by “unquantified” or placing specific values in parentheses [] when they have been adopted directly from the data of 1969Ba09 and 1986AgZX in order to calculate lesser gamma transitions depopulating the same nuclear level.

‡ Derived by various means: calculated gamma depopulation of particular nuclear levels based entirely on their known population sum; weighted mean of either direct gamma-ray spectroscopy measurements (1969Ba09 and 1986AgZX) and combination of various gamma-ray studies (1969Ba09, 1986AgZX and 2003 Fr02).

^o Relative emission probability of 147 (1) % for the 511-keV annihilation radiation was calculated from the total absolute positron emission probability of 69.2 (4) % and normalisation factor of 0.9404 (21).

⁺ As observed by 1976Su04 and 2003Fr02, population of the 1573.76-keV nuclear level by the 1391.64- and 1757.96-keV γ rays with relative emission probabilities of 0.027 (7) and 0.16 (2) %, respectively, implies depopulation by a 702.66-keV γ transition with a relative emission probability of 0.19 (2) %.

[§] Population of the 1741.65-keV nuclear level by the 998.25- and 1770.19-keV γ rays requires depopulation by an 870.55-keV γ transition as observed by 1976Su04, with a sum relative emission probability of 0.28 (3) %.

[§] Existence of 1864.29-keV γ ray and presence within the proposed decay scheme are questionable (compare 1969Ba09 with 1986AgZX) - relative emission probability calculated to be 0.24 (3) % from the relative emission probability ratio of 993.20-keV γ ray depopulating the same 1864.31-keV nuclear level of ^{94}Mo (relative emission probabilities from 2003Fr02 of 10.3(10) and 100.0 (10) for the 1864.29- and 993.20-keV γ rays, respectively).

^Δ Unplaced in proposed decay scheme.

Comments on evaluation

The normalisation factor for the relative gamma-ray emission and EC/ β^+ transition probabilities can be calculated via two related routes:

(a). A total value of 100% can be assigned to the gamma transition probabilities directly populating the 0.0-keV nuclear level of ^{94}Mo , assuming zero EC/ β^+ decay directly to the ground state on the basis of spin-parity considerations ((2) $^+$ \rightarrow 0 $^+$ constitutes a second forbidden non-unique transition (2, no)):

$$\sum_{0.0 \text{ keV}}^{\gamma \text{ population}} TP_{\gamma} = 100$$

Thus: $106.34 (24) * F = 100$

where F is the normalisation factor for the relative gamma-ray emission and EC/ β^+ transition probabilities.

$$F = 100 / 106.34 (24) = 0.9404 \pm 0.0021$$

(b). Gamma population-depopulation of the various nuclear levels of ^{94}Mo provide a means of determining the relative EC/ β^+ transition probabilities from the relative γ -ray transition probabilities. Assuming zero EC/ β^+ decay directly to the ground state based on spin-parity considerations (second forbidden non-unique), the calculated relative EC/ β^+ transition probabilities sum to 100%:

$$\sum_0^{EC/\beta^+ \text{ populations}} TP_{EC} = 100$$

Thus: $106.35 (77) * F = 100$

where F is the normalisation factor for the relative gamma-ray emission and EC/ β^+ transition probabilities

$$F = 100 / 106.35 (77) = 0.940 \pm 0.007$$

Therefore, a normalization factor of (0.9404 \pm 0.0021) was adopted for the relative gamma-ray emission and EC/ β^+ transition probabilities.

Multipolarities, and Internal Conversion and Internal-Pair Coefficients

The nuclear level scheme specified by Abriola and Sonzogni has been used to define the multipolarities of the gamma transitions on the basis of known spins and parities (2006Ab37). A significant number of transition types and mixing ratios have been determined from the γ - γ coincidence measurements of Fransen *et al.* (2003Fr02): (89.1%M1 + 10.9%E2) for the 358.3-keV gamma ray, (99.0%M1 + 1.0%E2) for the 875.60- and 1757.96-keV gamma rays, (20%M1 + 80%E2) for the 898.06- and 993.20-keV gamma rays, (99.75%M1 + 0.25%E2) for the 1005.58-keV gamma ray, (75.5% + 24.5%E2) for the 1061.31-keV gamma ray, (99.20%M1 + 0.80%E2) for the 1101.09-keV gamma ray, (97.80%M1 + 2.20%E2) for the 1196.24- and 1669.99-keV gamma rays, (99.36%M1 + 0.64%E2) for the 1264.34- and 1391.64-keV gamma rays, (91.7% + 8.3%E2) for the 1467.42- and 2663.18 -keV gamma rays, (98.58%M1 + 1.42%E2) for the 1521.91- and 1868.79-keV gamma rays, (37.2%M1 + 62.8%E2) for the 1998.78-keV gamma ray, (45.2%M1 + 54.8%E2) for the 2094.28-keV gamma ray, (64.6%M1 + 35.4%E2) for the 2257.53-keV gamma ray, (97.2 %M1 + 2.8%E2) for the 2292.16-keV gamma ray, and provisional selection of (21.7%M1 + 78.3%E2) for the 2576.5-keV gamma ray. The 998.25-, 2739.87- and 3128.60-keV gamma rays were defined as 100%M1, and the 325.67-, 528.71-, 672.56- and 802.55-keV gamma rays were arbitrarily assigned multipolarities of (50%M1 + 50%E2), while various E2 gamma transitions were also identified.

Recommended internal conversion coefficients have been determined from the frozen orbital approximation of Kibédi *et al.* (2008Ki07), based on the theoretical model of Band *et al.* (2002Ba85, 2002Ra45). A significant number of gamma transitions undergo decay via internal-pair formation, and the coefficient for this process has also been quantified in a few cases from the tabulations of 2008Ki07.

Gamma-ray emissions: multipolarities, and theoretical internal-conversion (frozen orbital approximation) and internal-pair formation coefficients.

E_γ(keV)	Multipolarity	α_K	α_L	α_{M+}	α_{totalICC}	α_{IPF}	α_{total}
293.37 (14)	96.9%M1 + 3.1%E2 δ = + 0.18 (5) 2003Fr02	0.0128 (3)	0.00149 (4)	0.00031	0.0146 (3)	–	0.0146 (3)
325.67 (9)	50%M1 + 50%E2 δ = 1.0 (2)	0.0128 (8)	0.00156 (11)	0.00034	0.0147 (9)	–	0.0147 (9)
358.3 (3)	89.1%M1 + 10.9%E2 δ = – 0.35 (12) 2003Fr02	0.0080 (4)	0.00093 (5)	0.00027	0.0092 (4)		0.0092 (4)
448.95 (14)	98.1%M1 + 1.9%E2 δ = + 0.14 (6) 2003Fr02	0.00442 (7)	0.000503 (8)	0.000107	0.00503 (8)	–	0.00503 (8)
466.52 (13)	E1	0.001544 (22)	0.0001729 (25)	0.000036	0.001753 (25)	–	0.001753 (25)
528.71 (8)	50%M1 + 50%E2 δ = 1.0 (2)	0.00325 (8)	0.000378 (11)	0.000082	0.00371 (9)	–	0.00371 (9)
532.10 (16)	(E2)	0.00346 (5)	0.000412 (6)	0.000078	0.00395 (6)	–	0.00395 (6)
669.56 (13)	E1	0.000666 (10)	0.0000740 (11)	0.000015	0.000755 (11)	–	0.000755 (11)
672.56 (9)	50%M1 + 50%E2 δ = 1.0 (2)	0.00176 (3)	0.000201 (4)	0.000039	0.00200 (3)	–	0.00200 (3)
702.66 (4)	E2	0.001608 (23)	0.000187 (3)	0.000035	0.00183 (3)	–	0.00183 (3)
721.03 (17)	99.91%M1 + 0.09%E2 δ = + 0.03 (4) 2003Fr02	0.001457 (21)	0.0001636 (23)	0.0000334	0.001654 (24)	–	0.001654 (24)
742.32 (13)	97.8%M1 + 2.2%E2 δ = + 0.15 (7) 2003Fr02	0.001365 (20)	0.0001533 (22)	0.0000317	0.001550 (22)	–	0.001550 (22)
802.55 (10)	50%M1 + 50%E2 δ = 1.0 (2)	0.001146 (16)	0.0001301 (19)	0.0000269	0.001303 (19)	–	0.001303 (19)
849.69 (10)	E2	0.000997 (14)	0.0001141 (16)	0.000023	0.001134 (16)	–	0.001134 (16)
870.55 (22)	E2	0.000940 (4)	0.0001075 (15)	0.0000225	0.001070 (15)	–	0.001070 (15)
871.094 (16)	E2	0.000939 (14)	0.0001073 (15)	0.0000217	0.001068 (15)	–	0.001068 (15)
875.60 (9)	99.0%M1 + 1.0%E2 δ = – 0.10 (2) 2003Fr02	0.000945 (14)	0.0001056 (15)	0.0000214	0.001072 (15)	–	0.001072 (15)
898.06 (9)	(20%M1 + 80%E2) δ = + 2.0 (+1.2, -0.6) [or 0.39 (25)] 2003Fr02	0.000877 (13) 0.000891 (13)	0.0000996 (14) 0.0000998 (14)	0.0000204 0.0000212	0.000997 (15) 0.001012 (15)	– –	0.000997 (15) 0.001012 (15)
916.09 (19)	99.96%M1 + 0.04%E2 δ = – 0.02 (7) 2003Fr02	0.000856 (12)	0.0000956 (13)	0.0000194	0.000971 (14)	–	0.000971 (14)
940.72 (20)	16%M1 + 84%E2	0.000786 (12)	0.0000892 (13)	0.0000188	0.000894 (13)	–	0.000894 (13)

Comments on evaluation

^{94m}Tc

	$\delta = + 2.3 (+0.7, -0.5)$ 2003Fr02						
960.10 (13)	E1	0.000316 (5)	0.0000349 (5)	0.0000071	0.000358 (5)	–	0.000358 (5)
993.20 (5)	20%M1 + 80%E2 $\delta = - 2.0 (10)$ 2003Fr02	0.000696 (13)	0.0000786 (13)	0.0000164	0.000791 (15)	–	0.000791 (15)
998.25 (17)	100%M1	0.000710 (10)	0.0000792 (11)	0.0000167	0.000806 (12)	–	0.000806 (12)
1005.58 (9)	99.75%M1 + 0.25%E2 $\delta = - 0.05 (4)$ 2003Fr02	0.000699 (10)	0.0000779 (11)	0.0000161	0.000793 (12)	–	0.000793 (12)
1037.2 (3) ^A	–	–	–	–	–	–	–
1061.31 (9)	(75.5%M1 + 24.5%E2) $\delta = - 7.0 (+3.0, -20.0)$ [or – 0.57 (16)] 2003Fr02	0.000595 (9) 0.000616 (10)	0.0000672 (10) 0.0000688 (10)	0.0000138 0.0000142	0.000676 (10) 0.000699 (11)	– –	0.000676 (10) 0.000699 (11)
1101.09 (8)	99.20%M1 + 0.80%E2 $\delta = - 0.09 (6)$ 2003Fr02	0.000576 (8)	0.0000640 (9)	0.0000130	0.000653 (9)	0.000000492 (8)	0.000653 (10)
1196.24 (6)	97.80%M1 + 2.20%E2 $\delta = + 0.15 (4)$ 2003Fr02	0.000483 (7)	0.0000536 (8)	0.0000114	0.000548 (7)	0.00000577 (9)	0.000553 (8)
1231.27 (19)	1.7%M1 + 98.3%E2 $\delta = + 7.6 (+4.7, -3.0)$ 2003Fr02	0.000431 (6)	0.0000483 (7)	0.0000097	0.000489 (7)	0.0000125 (20)	0.000502 (7)
1264.34 (9)	99.36%M1 + 0.64%E2 $\delta = - 0.08 (3)$ 2003Fr02	0.000431 (6)	0.0000478 (7)	0.0000092	0.000488 (6)	0.00001516 (22)	0.000503 (7)
1357.4 (15) ^A	–	–	–	–	–	–	–
1391.64 (7)	99.36%M1 + 0.64%E2 $\delta = - 0.08 (6)$ 2003Fr02	0.000353 (5)	0.0000391 (6)	0.0000079	0.000400 (5)	0.0000404 (6)	0.000441 (7)
1399.84 (16)	M1 + E2	–	–	–	–	–	–
1423.68 (16)	E2	0.000319 (5)	0.0000355 (5)	0.0000075	0.000362 (5)	0.0000577 (8)	0.000419 (6)
1467.42 (18)	91.7%M1 + 8.3%E2 $\delta = + 0.3 (+2.9, -0.2)$ 2003Fr02	0.000316 (15)	0.0000350 (15)	0.0000070	0.000358 (16)	0.000061 (10)	0.000419 (9)
1499.13 (9)	M1 + E2	–	–	–	–	–	–
1521.91 (6)	98.58%M1 + 1.42%E2 $\delta = - 0.12 (3)$ 2003Fr02	0.000295 (5)	0.0000326 (5)	0.0000064	0.000334 (5)	0.0000776 (11)	0.000411 (6)
1536.51 (18)	–	–	–	–	–	–	–
1592.00 (10)	E2	0.000255 (4)	0.0000284 (4)	0.0000066	0.000290 (4)	0.0001198 (17)	0.000410 (6)
1662.75 (12)	E1	0.0001198 (17)	0.00001311 (19)	0.0000027	0.0001356 (17)	0.000369 (6)	0.000505 (7)
1669.99 (10)	97.80%M1 + 2.20%E2	0.000245 (4)	0.0000270 (4)	0.0000055	0.000277 (4)	0.000132 (3)	0.000410 (6)

	$\delta = + 0.15$ (19) 2003Fr02						
1757.96 (17)	99.0%M1 + 1.0%E2 $\delta = - 0.10$ (3) 2003Fr02	0.000221 (3)	0.0000244 (3)	0.0000050	0.000251 (4)	0.0001677 (24)	0.000418 (6)
1770.19 (21)	(M1 + E2)	-	-	-	-	-	-
1824.79 (9)	(M1 + E2)	-	-	-	-	-	-
1864.29 (5)	E2	0.000189 (3)	0.0000209 (3)	0.0000041	0.000214 (3)	0.000241 (4)	0.000455 (7)
1868.79 (7)	98.58%M1 + 1.42%E2 $\delta = - 0.12$ (2) 2003Fr02	0.000196 (3)	0.0000216 (3)	0.0000044	0.000222 (3)	0.000215 (3)	0.000438 (7)
1928.54 (16)	M1 + E2	-	-	-	-	-	-
1933.97 (19)	69.7%M1 + 30.3%E2 $\delta = - 0.66$ (14) [or - 1.7 (+0.4, -0.5)] 2003Fr02	0.000182 (3) 0.000178 (3)	0.0000200 (3) 0.0000197 (3)	0.0000040 0.0000043	0.000206 (3) 0.000202 (3)	0.000253 (5) 0.000266 (5)	0.000459 (7) 0.000468 (7)
1998.78 (8)	37.2%M1 + 62.8%E2 $\delta = + 1.3$ (+1.4, -0.4) 2003Fr02	0.000168 (3)	0.0000186 (3)	0.0000044	0.000191 (3)	0.000293 (9)	0.000484 (10)
2027.83 (9)	(M1 + E2)	-	-	-	-	-	-
2067.33 (6)	E2	0.0001562 (22)	0.00001722 (25)	0.00000358	0.0001770 (22)	0.000338 (5)	0.000515 (8)
2094.28 (6)	45.2%M1 + 54.8%E2 $\delta = + 1.1$ (+1.0, -0.4) 2003Fr02	0.000155 (3)	0.0000171 (3)	0.0000039	0.000176 (3)	0.000336 (11)	0.000512 (11)
2257.53 (7)	64.6%M1 + 35.4%E2 $\delta = + 0.74$ (+0.21, -0.17) 2003Fr02	0.0001356 (20)	0.00001491 (22)	0.00000309	0.0001536 (20)	0.000407 (8)	0.000561 (9)
2292.16 (19)	(97.2%M1 + 2.8%E2) $\delta = + 0.17$ (4) 2003Fr02	0.0001330 (19)	0.00001461 (21)	0.00000300	0.0001506 (19)	0.000411 (6)	0.000562 (8)
2392.99 (6)	E2	0.0001203 (17)	0.00001322 (19)	0.00000278	0.0001363 (17)	0.000496 (7)	0.000633 (9)
2460.61 (17)	(M1 + E2)	-	-	-	-	-	-
2529.69 (17)	-	-	-	-	-	-	-
2576.5 (4)	(21.7%M1 + 78.3%E2) $\delta = - 1.9$ (+0.5, -0.6) [or - 0.08 (10)] 2003Fr02	0.0001061 (15) 0.0001072 (15)	0.00001164 (17) 0.00001176 (17)	0.00000246 0.00000243	0.0001202 (16) 0.0001214 (16)	0.000574 (11) 0.000542 (8)	0.000694 (12) 0.000664 (10)
2640.72 (14)	(M1 + E2)	-	-	-	-	-	-
2663.18 (9)	91.7%M1 + 8.3%E2 $\delta = - 0.3$ (2) 2003Fr02	0.0001009 (15)	0.00001106 (16)	0.00000224	0.0001142 (15)	0.000585 (10)	0.000699 (11)
2739.87 (7)	100%M1	0.0000959 (14)	0.00001051 (15)	0.00000217	0.000109 (14)	0.000616 (9)	0.000725 (11)
2869.85 (8)	E2	0.0000881 (13)	0.00000964 (14)	0.00000196	0.0000997 (13)	0.000717 (10)	0.000816 (12)

Comments on evaluation

^{94m}Tc

3021.01 (7)	(M1 + E2)	-	-	-	-	-	-
3065.6 (3) ^Δ	-	-	-	-	-	-	-
3085.8 (3) ^Δ	-	-	-	-	-	-	-
3128.60 (7)	100%M1	0.0000758 (11)	0.00000829 (12)	0.00000172	0.000086 (11)	0.000785 (11)	0.000871 (13)
3400.8(3)	-	-	-	-	-	-	-
3447.5 (4)	-	-	-	-	-	-	-
3511.79 (14)	(M1 + E2)	-	-	-	-	-	-
3534.25 (9)	E2	0.0000625 (9)	0.00000682 (10)	0.00000148	0.0000708 (9)	0.000994 (14)	0.001065 (15)
3640.6 (3) ^Δ	-	-	-	-	-	-	-
3792.79 (15)	E2	0.0000559 (8)	0.00000609 (9)	0.00000121	0.0000632 (8)	0.001086 (16)	0.001149 (16)
3892.07 (7)	-	-	-	-	-	-	-
4136.2 (3) ^Δ	-	-	-	-	-	-	-

^Δ Unplaced in proposed decay scheme.

Recommended gamma-ray energies, relative and absolute emission probabilities, and transition probabilities.

E_γ (keV)	origins	P_γ^{rel}	origins	P_γ^{abs}	Transition probability (%)
$\gamma_{19,14}$ 293.37 (14)	2003Fr02	unknown	–	–	–
$\gamma_{7,5}$ 325.67 (9)	2003Fr02	0.029 (2)	2003Fr02	0.027 (2)	0.027 (2)
$\gamma_{18,12}$ 358.3 (3)	2003Fr02	0.0089 (7)	2003Fr02	0.0084 (7)	0.0085 (7)
$\gamma_{14,8}$ 448.95 (14)	2003Fr02	unknown	–	–	–
$\gamma_{9,5}$ 466.52 (13)	1976Su04/2003Fr02	unknown	–	–	–
γ^+ 511	annihilation radiation	147 (1)	annihilation radiation	138 (1)	–
$\gamma_{7,4}$ 528.71 (8)	2003Fr02	0.034 (2)	2003Fr02	0.032 (2)	0.032 (2)
$\gamma_{15,8}$ 532.10 (16)	2003Fr02	unknown	–	–	–
$\gamma_{9,4}$ 669.56 (13)	2003Fr02	unknown	–	–	–
$\gamma_{11,5}$ 672.56 (9)	2003Fr02	0.18 (3)	2003Fr02	0.17 (3)	0.17 (3)
$\gamma_{2,1}$ 702.66 (4)	1976Su04/2003Fr02	0.19 (2) ⁺	–	0.18 (2)	0.18 (2)
$\gamma_{6,2}$ 721.03 (17)	1976Su04/2003Fr02	unknown	–	–	–
$\gamma_{19,8}$ 742.32 (13)	2003Fr02	unknown	–	–	–
$\gamma_{13,5}$ 802.55 (10)	2003Fr02	0.0262 (15)	2003Fr02	0.0246 (14)	0.0246 (14)
$\gamma_{8,2}$ 849.69 (10)	2003Fr02	unknown	–	–	–
$\gamma_{3,1}$ 870.55 (22)	1976Su04	0.28 (3) ^s	–	0.26 (3)	0.26 (3)
$\gamma_{1,0}$ 871.094 (16)	1969Ba09/1976Su04/1986AgZX/2003Fr02	100	1969Ba09/1986AgZX/2003Fr02	94.04 (21)	94.14 (21)
$\gamma_{11,4}$ 875.60 (9)	1986AgZX/2003Fr02	1.1 (3)	1986AgZX/2003Fr02	1.0 (3)	1.0 (3)
$\gamma_{16,5}$ 898.06 (9)	2003Fr02	0.0104 (5)	2003Fr02	0.0098 (5)	0.0098 (5)
$\gamma_{21,8}$ 916.09 (19)	2003Fr02	unknown	–	–	–
$\gamma_{12,4}$ 940.72 (20)	2003Fr02	unknown	–	–	–
$\gamma_{9,2}$ 960.10 (13)	1976Su04/2003Fr02	unknown	–	–	–
$\gamma_{4,1}$ 993.20 (5)	1969Ba09/1976Su04/1986AgZX/2003Fr02	2.35 (19)	1969Ba09/1986AgZX	2.21 (18)	2.21 (18)
$\gamma_{11,3}$ 998.25 (17)	1969Ba09/1986AgZX/2003Fr02	0.25 (2)	1969Ba09/1986AgZX/2003Fr02	0.24 (2)	0.24 (2)
$\gamma_{13,4}$ 1005.58 (9)	1969Ba09/1976Su04/1986AgZX/2003Fr02	0.10 (3)	1969Ba09/1986AgZX	0.09 (3)	0.09 (3)
2 x γ^+ 1022	(511 + 511) keV sum peak	0.029 (15)	1986AgZX	0.027 (14)	–
$\gamma_{-1,1}$ 1037.2 (3) ^A	1976Su04/1986AgZX	0.047 (15)	1986AgZX	0.044 (14)	0.044 (14)
$\gamma_{17,5}$ 1061.31 (9)	2003Fr02	0.017 (2)	2003Fr02	0.016 (2)	0.016 (2)
$\gamma_{16,4}$ 1101.09 (8)	1976Su04/1986AgZX/2003Fr02	0.045 (15)	1986AgZX	0.042 (14)	0.042 (14)
$\gamma_{5,1}$ 1196.24 (6)	1969Ba09/1976Su04/1986AgZX/2003Fr02	0.76 (7)	1969Ba09/1986AgZX	0.71 (7)	0.71 (7)
$\gamma_{12,2}$ 1231.27 (19)	1976Su04/2003Fr02	unknown	–	–	–
$\gamma_{17,4}$ 1264.34 (9)	1969Ba09/1986AgZX/2003Fr02	0.23 (2)	1969Ba09/1986AgZX/2003Fr02	0.22 (2)	0.22 (2)
$\gamma_{-1,2}$ 1357.4 (15) ^A	1969Ba09	0.20 (8)	1969Ba09	0.19 (8)	0.19 (8)
$\gamma_{16,2}$ 1391.64 (7)	1976Su04/2003Fr02	0.0284 (11)	2003Fr02	0.0267 (10)	0.0267 (10)
$\gamma_{26,7}$ 1399.84 (16)	2003Fr02	0.044 (3)	2003Fr02	0.041 (3)	0.041 (3)
$\gamma_{6,1}$ 1423.68 (16)	2003Fr02	unknown	–	–	–
$\gamma_{20,4}$ 1467.42 (18)	2003Fr02	0.077 (5)	2003Fr02	0.072 (5)	0.072 (5)
$\gamma_{27,7}$ 1499.13 (9)	1986AgZX/2003Fr02	0.071 (12)	1986AgZX/2003Fr02	0.067 (11)	0.067 (11)
$\gamma_{7,1}$ 1521.91 (6)	1969Ba09/1976Su04/1986AgZX/2003Fr02	4.76 (30)	1969Ba09/1986AgZX	4.48 (28)	4.48 (28)
$\gamma_{22,4}$ 1536.51 (18)	2003Fr02	0.015 (3)	2003Fr02	0.014 (3)	0.014 (3)

Comments on evaluation

^{94m}Tc

γ _{19,2}	1592.00 (10)	2003Fr02	unknown	–	–	–
γ _{9,1}	1662.75 (12)	1976Su04/2003Fr02	unknown	–	–	–
γ _{25,4}	1669.99 (10)	1986AgZX/2003Fr02	0.039 (2)	1986AgZX/2003Fr02	0.037 (2)	0.037 (2)
γ _{20,2}	1757.96 (17)	1969Ba09/1986AgZX/2003Fr02	0.16 (2)	1969Ba09/1986AgZX	0.15 (2)	0.15 (2)
γ _{24,3}	1770.19 (21)	1986AgZX/2003Fr02	0.027 (7)	1986AgZX/2003Fr02	0.025 (6)	0.025 (6)
γ _{27,5}	1824.79 (9)	2003Fr02	0.024 (1)	2003Fr02	0.023 (1)	0.023 (1)
γ _{4,0}	1864.29 (5)	1969Ba09/1976Su04/1986AgZX/2003Fr02	0.24 (3) [§]	2003Fr02	0.23 (3)	0.23 (3)
γ _{11,1}	1868.79 (7)	1969Ba09/1976Su04/1986AgZX/2003Fr02	5.84 (30)	1969Ba09/1986AgZX	5.49 (28)	5.49 (28)
γ _{26,4}	1928.54 (16)	1969Ba09/1986AgZX/2003Fr02	0.08 (2)	1969Ba09/1986AgZX	0.075 (19)	0.075 (19)
γ _{12,1}	1933.97 (19)	1976Su04/2003Fr02	unknown	–	–	–
γ _{13,1}	1998.78 (8)	2003Fr02	0.0131 (6)	2003Fr02	0.0123 (6)	0.0123 (6)
γ _{27,4}	2027.83 (9)	1986AgZX/2003Fr02	0.022 (4)	1986AgZX/2003Fr02	0.021 (4)	0.021 (4)
γ _{5,0}	2067.33 (6)	1969Ba09/1976Su04/1986AgZX/2003Fr02	0.12 (1)	1969Ba09/1986AgZX/2003Fr02	0.11 (1)	0.11 (1)
γ _{16,1}	2094.28 (6)	2003Fr02	0.0166 (6)	2003Fr02	0.0156 (6)	0.0156 (6)
γ _{17,1}	2257.53 (7)	1986AgZX/2003Fr02	0.061 (5)	1986AgZX/2003Fr02	0.057 (5)	0.057 (5)
γ _{18,1}	2292.16 (19)	1976Su04/1986AgZX/2003Fr02	0.053 (18)	1986AgZX	0.050 (17)	0.050 (17)
γ _{7,0}	2392.99 (6)	1969Ba09/1986AgZX/2003Fr02	0.53 (4)	1969Ba09/1986AgZX/2003Fr02	0.50 (4)	0.50 (4)
γ _{20,1}	2460.61 (17)	2003Fr02	0.012 (2)	2003Fr02	0.011 (2)	0.011 (2)
γ _{22,1}	2529.69 (17)	1969Ba09/1976Su04/1986AgZX/2003Fr02	0.36 (4)	1969Ba09/1986AgZX	0.34 (4)	0.34 (4)
γ _{23,1}	2576.5 (4)	1969Ba09/1976Su04/1986AgZX/2003Fr02	0.12 (2)	1969Ba09/1986AgZX	0.11 (2)	0.11 (2)
γ _{24,1}	2640.72 (14)	1969Ba09/1986AgZX/2003Fr02	0.035 (4)	1986AgZX/2003Fr02	0.033 (4)	0.033 (4)
γ _{25,1}	2663.18 (9)	1969Ba09/2003Fr02	0.070 (2)	1969Ba09/2003Fr02	0.066 (2)	0.066 (2)
γ _{11,0}	2739.87 (7)	1969Ba09/1976Su04/1986AgZX/2003Fr02	3.75 (21)	1969Ba09/1986AgZX/2003Fr02	3.53 (20)	3.53 (20)
γ _{13,0}	2869.85 (8)	1986AgZX/2003Fr02	0.017 (2)	1986AgZX/2003Fr02	0.016 (2)	0.016 (2)
γ _{27,1}	3021.01 (7)	1969Ba09/1986AgZX/2003Fr02	0.093 (15)	1969Ba09/1986AgZX	0.087 (14)	0.087 (14)
γ _{-1,3}	3065.6 (3) ^Δ	1986AgZX	0.012 (4)	1986AgZX	0.011 (4)	0.011 (4)
γ _{-1,4}	3085.8 (3) ^Δ	1986AgZX	0.017 (4)	1986AgZX	0.016 (4)	0.016 (4)
γ _{17,0}	3128.60 (7)	1969Ba09/1986AgZX/2003Fr02	1.42 (10)	1969Ba09/1986AgZX	1.34 (9)	1.34 (9)
γ _{22,0}	3400.76 (17)	1986AgZX	0.005 (2)	1986AgZX	0.005 (2)	0.005 (2)
γ _{23,0}	3447.5 (4)	1986AgZX/2003Fr02	0.006 (1)	1986AgZX/2003Fr02	0.006 (1)	0.006 (1)
γ _{24,0}	3511.79 (14)	1969Ba09/1986AgZX/2003Fr02	0.067 (7)	1969Ba09/1986AgZX	0.063 (7)	0.063 (7)
γ _{25,0}	3534.25 (9)	2003Fr02	0.0036 (4)	2003Fr02	0.0034 (4)	0.0034 (4)
γ _{-1,5}	3640.6 (3) ^Δ	1986AgZX	0.007 (2)	1986AgZX	0.007 (2)	0.007 (2)
γ _{26,0}	3792.79 (15)	1969Ba09/1986AgZX/2003Fr02	0.055 (5)	1969Ba09/1986AgZX/2003Fr02	0.052 (5)	0.052 (5)
γ _{27,0}	3892.07 (7)	1969Ba09/1986AgZX/2003Fr02	0.015 (2)	1969Ba09/1986AgZX/2003Fr02	0.014 (2)	0.014 (2)
γ _{-1,6}	4136.2 (3) ^Δ	1986AgZX	0.007 (1)	1986AgZX	0.007 (1)	0.007 (1)

[†] Population of the 1573.76-keV nuclear level by the 1391.64- and 1757.96-keV γ rays with relative emission probabilities of 0.027 (7) and 0.16 (2) %, respectively, implies depopulation by a 702.66-keV γ transition observed by 1976Su04 and 2003Fr02, with a relative emission probability of 0.19 (2) %.

[§] Population of the 1741.65-keV nuclear level by the 998.25- and 1770.19-keV γ rays requires depopulation by an 870.55-keV γ transition observed by 1976Su04, with a sum relative emission probability of 0.28 (3) %.

^Δ Unplaced in proposed decay scheme.

EC/β⁺ TransitionsEnergies

All EC/β⁺ energies were derived from the structural details of the proposed decay scheme. The nuclear level energies of 2006Ab37 and evaluated Q_{EC}-value of 4332 (5) keV (2012Wa38) were used to determine the recommended energies and uncertainties of the EC transitions and β⁺ emissions.

Transition and Emission Probabilities

Gamma-ray studies of the EC decay of ⁹⁴Ru have revealed 100% feeding to ^{94m}Tc, and a lack of evidence for subsequent ^{94m}Tc IT decay to the ground state (≤ 0.1% relative to the 871.094-keV gamma-ray emission of ^{94m}Tc (1967Ei01)). On the basis of these findings, ^{94m}Tc has been assumed to undergo 100% EC/β⁺ decay.

A large majority of the EC transition probabilities were derived from the population-depopulation imbalances of the relative emission probabilities of the gamma rays, their theoretical internal-conversion and internal-pair formation coefficients. A value of 0.9404 (21) was adopted as the normalisation factor in order to determine the absolute transition and emission probabilities of the EC and β⁺ particles from their resulting relative transition and emission probabilities. Component EC and β⁺ transition and emission probabilities were determined from EC/β⁺ ratios (1971Go40), and log f t values and average E_{β⁺} energies were derived by means of the LOGFT code. Fractional EC probabilities P_K, P_L, P_M and P_N were calculated by means of the EC-CAPTURE code (1998Sc28) as developed from the data tabulations of 1995ScZY.

Possible EC transitions directly to the 3339.54-, 3165.77-, 2955.55-, 2872.40-, 2610.57-, 2423.45-, 2294.79-, 1741.65- and 1573.76-keV nuclear levels of ⁹⁴Mo were assumed to be zero, based on gamma population-depopulation balances and/or spin-parity considerations, while EC decay directly to the ground state was assigned a value of zero from a spin-parity assessment alone (second forbidden non-unique (2, no)). Although both the 2805.04- and 2533.87-keV nuclear levels of ⁹⁴Mo may undergo EC and γ population and γ depopulation, these possibilities have not been confidently confirmed, and therefore direct EC transition probabilities of zero were assigned (despite these possible transitions being allowed (1, no) and first forbidden non-unique (1, yes), respectively).

The absolute emission probability of the 511-keV annihilation radiation was derived from the total absolute positron emission probability of 69.2 (4) % as determined during the evaluation of the EC/β⁺ transitions/emissions:

$$\text{absolute } P_{\gamma}^{\pm} = 2 \times 69.2 (4) = 138.4 (8) \% \rightarrow 138 (1) \%$$

$$[\text{relative } P_{\gamma}^{\pm} = \frac{\text{absolute } P_{\gamma}^{\pm}}{F} = \frac{138.4 (8)}{0.9404 (21)} = 147.2 (9) \% \rightarrow 147 (1) \%]$$

where F is the normalisation factor for the gamma-ray emission probabilities.

A decay scheme was derived that consists of 16 EC/7 β⁺ transitions and 52 gamma-ray emissions. Substantial 511-keV annihilation radiation was also observed. A further sixteen gamma rays have been observed/proposed by Fransen *et al.* (2003Fr02), but an inability to derive and quantify their absolute emission probabilities has prevented their satisfactory insertion and adoption in the decay scheme and data file – if these additional gamma transitions could have been suitably introduced into the ^{94m}Tc decay scheme, they would have impacted to a small degree on the EC/β⁺ decay data. Finally, although six other gamma-ray emissions of reasonably low intensity could not be placed in the proposed decay scheme, they have been included in the final data set because their absolute emission probabilities have been measured.

Recommended energies and transition probabilities of the EC/ β^+ decay of ^{94m}Tc.

E_{EC} (keV)*	E_{β^+} (keV)	Av. E_{β^+} (keV)	P_{EC} (total)	ϵ/β^+ (theory) 1971Go40	P_{EC}	P_{β^+}	^{94m} Tc	⁹⁴ Mo	transition type	log ft	P_K	P_L	P_M	P_N	
EC _{0,27}	440 (5)	–	–	0.212 (13)	–	0.212 (13)	–	(2) +	(2 +)	(allowed)	5.6	0.8620 (15)	0.1121 (12)	0.0220 (4)	0.0039 (2)
EC _{0,26}	539 (5)	–	–	0.169 (20)	–	0.169 (20)	–	(2) +	2 +	(allowed)	5.9	0.8639 (15)	0.1106 (11)	0.0216 (4)	0.0039 (2)
EC _{0,25}	798 (5)	–	–	0.106 (3)	–	0.106 (3)	–	(2) +	2 +	(allowed)	6.4	0.8664 (15)	0.1086 (11)	0.0212 (4)	0.0038 (2)
EC _{0,24}	820 (5)	–	–	0.121 (10)	–	0.121 (10)	–	(2) +	1 (+)	(allowed)	6.4	0.8666 (15)	0.1085 (11)	0.0212 (4)	0.0038 (2)
EC _{0,23}	884 (5)	–	–	0.118 (19)	–	0.118 (19)	–	(2) +	(1, 2 +)	(allowed)	6.4	0.8669 (14)	0.1082 (11)	0.0211 (4)	0.0038 (2)
EC _{0,22}	931 (5)	–	–	0.36 (4)	–	0.36 (4)	–	(2) +		[allowed]	6.0	0.8672 (14)	0.1080 (11)	0.0211 (4)	0.0038 (2)
EC _{0,20}	1000 (5)	–	–	0.234 (20)	–	0.234 (20)	–	(2) +	(3 +)	(allowed)	6.3	0.8675 (14)	0.1078 (11)	0.0210 (4)	0.0038 (2)
EC _{0,18}	1169 (5)	147 (5)	–	0.058 (17)	5888	0.058 (17)	–	(2) +	(3 +)	(allowed)	7.0	0.8681 (14)	0.1073 (11)	0.0209 (4)	0.0037 (2)
EC _{0,17}	1203 (5)	181 (5)	–	1.63 (9)	2344	1.63 (9)	–	(2) +	1 +	(allowed)	5.57	0.8682 (14)	0.1072 (11)	0.0209 (4)	0.0037 (2)
EC _{0,16}	1367 (5)	345 (5)	157.5 (22)	0.094 (14)	160	0.093 (14)	0.00058 (9)	(2) +	3 +	(allowed)	6.9	0.8686 (14)	0.1069 (11)	0.0208 (4)	0.0037 (2)
EC _{0,13}	1462 (5)	440 (5)	198.5 (22)	0.15 (3)	61.66	0.15 (3)	0.0024 (5)	(2) +	2 +	(allowed)	6.8	0.8688 (14)	0.1067 (11)	0.0208 (4)	0.0037 (2)
EC _{0,11}	1592 (5)	570 (5)	254.3 (22)	10.5 (4)	23.44	10.1 (4)	0.427 (21)	(2) +	1 +	(allowed)	5.03	0.8690 (14)	0.1066 (11)	0.0207 (4)	0.0037 (2)
EC _{0,7}	1939 (5)	917 (5)	404.8 (22)	4.9 (3)	4.41	4.0 (2)	0.91 (6)	(2) +	2 +	(allowed)	5.60	0.8694 (14)	0.1062 (11)	0.0207 (4)	0.0037 (2)
EC _{0,5}	2265 (5)	1243 (5)	548.7 (23)	0.56 (8)	1.595	0.34 (5)	0.22 (3)	(2) +	2 +	(allowed)	6.8	0.8697 (14)	0.1060 (11)	0.0206 (4)	0.0037 (2)
EC _{0,4}	2468 (5)	1446 (5)	639.6 (23)	0.80 (19)	0.974	0.39 (9)	0.41 (10)	(2) +	2 +	(allowed)	6.82	0.8699 (14)	0.1059 (11)	0.0206 (4)	0.0037 (2)
EC _{0,1}	3461 (5)	2439 (5)	1094.4 (24)	80.0 (5)	0.1908	12.8 (1)	67.2 (4)	(2) +	2 +	(allowed)	5.606	0.8704 (14)	0.1055 (11)	0.0205 (4)	0.0037 (2)
				Σ 100.012		Σ 30.881	Σ 69.16998								
						Σ 100.05098									

* Determined from the nuclear level energies of 2006Ab37 and Q-value of 4332 (5) keV (2012Wa38).

Atomic Data

The x-ray and Auger-electron data have been calculated using the evaluated gamma-ray data, and atomic data from 1996Sc06, 1998ScZM and 1999ScZX. Both the X-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.02, 28 February 2012), as described in 2000Sc47. This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

K and L X-ray energies and emission probabilities of ^{94m}Tc.

			Energy (keV)	Photons per 100 disint.	Relative probability
XL	(Mo)		2.016 – 2.831	1.198 (22)	10.6
	XL ₁	(Mo)	2.016	0.0279 (8)	
	XL _α	(Mo)	2.290 – 2.293	0.750 (18)	
	XL _η	(Mo)	2.120	0.0100 (3)	
	XL _β	(Mo)	2.395 – 2.518	0.389 (10)	
	XL _γ	(Mo)	2.623 – 2.831	0.0208 (4)	
XK _α	XK _{α2}	(Mo)	17.3745 (2)	5.93 (11)	52.4
	XK _{α1}	(Mo)	17.47954 (2)	11.31 (19)	100
XK _{β1}	XK _{β3}	(Mo)	19.5904 (2))	26.3
	XK _{β1}	(Mo)	19.6085 (2)) 2.97 (6)	
	XK _{β5}	(Mo)	19.774 (5))	
XK _{β2}	XK _{β2}	(Mo)	19.9653 (7)) 0.457 (18)	4.04
	XK _{β4}	(Mo)	19.998)	

Auger-electron energies and emission probabilities of ^{94m}Tc.

		Energy (keV)	Electrons per 100 disint.	Relative probability
e _{AK}	(Mo)		6.28 (15)	100
	KLL	14.172 – 14.855	4.37 (11)	
	KLX	16.592 – 17.478	1.74 (5)	
	KXY	18.990 – 19.996	0.172 (5)	
e _{AL}	(Mo)	1.48 – 2.25	29.8 (4)	682

Mo: ω_K = 0.767 (4); ω_L = 0.0381 (9); n_{KL} = 1.029 (4) were taken from 1996Sc06.

Electron energies were determined from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

Data Consistency

A Q_{EC}-value of 4332 (5) keV has been adopted from the atomic mass evaluation of Wang *et al.* (2012Wa38) while in the course of formulating the decay scheme of ^{94m}Tc. This value has subsequently been compared with the Q-value calculated by summing the contributions of the individual emissions to the ^{94m}Tc EC-decay process (i.e. EC/β⁺, electron, γ, etc.):

$$\text{calculated Q-value} = \sum (E_i \times P_i) = 4331 (17) \text{ keV}$$

Percentage deviation from the Q-value of Wang *et al.* is 0.02 (42) %, which supports the derivation of a reasonably consistent decay scheme with a large variant.

References

- 1948Mo19 E.E. Motta, G.E. Boyd, Characterization of Tc Activities Produced by Deuteron Bombardment of Separated Mo Isotopes, Phys. Rev. 74 (1948) 220-222. [Half-life]
- 1950Me21 H. Medicus, P. Preiswerk, P. Scherrer, Untersuchungen Über den Radioaktiven Zerfall bei Isotopen des Technetiums, Helv. Phys. Acta 23 (1950) 299-346. [Half-life, E_{β+}, P_{β+}, P_{EC}]
- 1962Mo06 S. Monaro, G.B. Vingiani, R.A. Ricci, R. van Lieshout, A New Technetium Activity: 4.5 hour ^{94m}Tc, Physica 28 (1962) 63-69. [Half-lives (⁹⁴Tc, ^{94m}Tc), E_γ, P_γ, E_{β+}, P_{β+}, P_{EC}]
- 1963Ma21 J.M. Matuszek, Jr., T.T. Sugihara, The Decay of 293 min Tc⁹⁴, Nucl. Phys. 42 (1963) 582-589. [Half-life of ground state ⁹⁴Tc]
- 1964Ha29 J.H. Hamilton, K.E.G. Löbner, A.R. Sattler, R. van Lieshout, The Energy Levels in ⁹⁴Mo Populated by the Decays of ⁹⁴Tc and ^{94m}Tc, Physica 30 (1964) 1802-1812. [E_γ, P_γ, E_{β+}, P_{cc}, ICC]
- 1965Ba48 K.A. Baskova, S.S. Vasil'ev, M.A. Khamo-Leila, L.Ya. Shavtvalov, Investigation of the Radiation from Tc⁹⁴, Tc⁹⁶ and Zr⁸⁷, Bull. Acad. Sci. USSR, Phys. Ser. 29 (1966) 2094-2102. [P_γ, Half-life of ground state ⁹⁴Tc]
- 1967Ei01 E. Eichler, G. Chilosi, N.R. Johnson, The Decay of ⁹⁴Ru to Levels in ⁹⁴Tc, Phys. Lett. 24B (1967) 140-142. [⁹⁴Tc nuclear levels, ^{94m}Tc decay mode, ⁹⁴Ru half-life]
- 1968Ar06 N.K. Aras, E. Eichler, G.G. Chilosi, Decay of ⁹⁴Tc Isomers into the Levels of ⁹⁴Mo, Nucl. Phys. A112 (1968) 609-624. [E_γ, P_γ]
- 1968Bo27 G.G.J. Boswell, T. McGee, The Decay of Ruthenium 94, J. Inorg. Nucl. Chem. 30 (1968) 1139-1143. [⁹⁴Ru half-life]
- 1968Ka25 G.S. Katykhin, M.K. Nikitin, Yu.N. Podkopaev, J. Vrzal, J. Liptak, Gamma-ray Spectra of ^{93,93m}Tc and ^{94,94m}Tc, Bull. Acad. Sci. USSR, Phys. Ser. 32 (1969) 739-744. [E_γ]
- 1969Ba09 J. Barrette, A. Boutard, S. Monaro, Energy Levels of ⁹⁴₄₂Mo Excited in the Decay of the ^{94m,g}Tc Isomeric Pair, Can. J. Phys. 47 (1969) 995-1003. [E_γ, P_γ]
- 1971Go40 N.B. Gove, M.J. Martin, Log-*f* Tables for Beta Decay, Nucl. Data Tables 10 (1971) 205-237. [ε/β⁺ ratios]
- 1976Su04 Y. Sugiyama, S. Kikuchi, Study of ⁹⁴Mo Through the (n,n'γ) Reaction, Nucl. Phys. A264 (1976) 170-187. [Nuclear levels, E_γ]
- 1977La19 F.P. Larkins, Semiempirical Auger-electron Energies for Elements 10 ≤ Z ≤ 100, At. Data Nucl. Data Tables 20 (1977) 311-387. [Auger-electron energies]
- 1986AgZX V.A. Ageev, V.S. Belyavenko, I.N. Vishnevsky, V.A. Zheltonozhsky, Decay of ^{94m}Tc, Program and Theses 36th Ann. Conf. Nucl. Spectrosc. Struct. At. Nuclei, Kharkov (1986) 77. [E_γ, P_γ]
- 1995ScZY E. Schönfeld, Tables for the Calculation of Electron Capture Probabilities, PTB Report PTB-6.33-95-2, 1995. [P_K, P_L, P_M, P_N, P_O]
- 1996Sc06 E. Schönfeld, H. Janßen, Evaluation of Atomic Shell Data, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527-533. [ω_K, ω_L, K_β/K_α, K_{α2}/K_{α1}, KLX/KLL, KXY/KLL]
- 1998Sc28 E. Schönfeld, Calculation of Fractional Electron Capture Probabilities, Appl. Radiat. Isot. 49 (1998) 1353-1357. [P_K, P_L, P_M, P_N, P_O]

- 1998ScZM E. Schönfeld, G. Rodloff, Tables of the Energies of K-Auger Electrons for Elements with Atomic Numbers in the Range from $Z = 11$ to $Z = 100$, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX E. Schönfeld, G. Rodloff, Energies and Relative Emission Probabilities of K X-rays for Elements with Atomic Numbers in the Range from $Z = 5$ to $Z = 100$, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2000Sc47 E. Schönfeld, H. Janßen, Calculation of Emission Probabilities of X-rays and Auger Electrons Emitted in Radioactive Disintegration Processes, Appl. Radiat. Isot. 52 (2000) 595-600. [P_X, P_{Ae}]
- 2002Ba85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, Dirac-Fock Internal Conversion Coefficients, At. Data Nucl. Data Tables 81 (2002) 1-334. [ICC]
- 2002Ra45 S. Raman, C.W. Nestor, Jr., A. Ichihara, M.B. Trzhaskovskaya, How Good are the Internal Conversion Coefficients Now? Phys. Rev. C66 (2002) 044312, 1-23. [ICC]
- 2003Fr02 C. Fransen, N. Pietralla, Z. Ammar, D. Bandyopadhyay, N. Boukharouba, P. von Brentano, A. Dewald, J. Gableske, A. Gade, J. Jolie, U. Kneissl, S.R. Leshner, A.F. Lisetskiy, M.T. McEllistrem, M. Merrick, H.H. Pitz, N. Warr, V. Werner, S.M. Yates, Comprehensive Studies of Low-spin Collective Excitations in ⁹⁴Mo, Phys. Rev. C67 (2003) 024307, 1-19. [Nuclear levels, E_γ, P_γ, δ]
- 2006Ab37 D. Abriola, A.A. Sonzogni, Nuclear Data Sheets for $A = 94$, Nucl. Data Sheets 107 (2006) 2423-2578. [Nuclear levels]
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr., Evaluation of Theoretical Conversion Coefficients using BrIcc, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202-229. [ICC]
- 2012Wa38 M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer, The AME2012 Atomic Mass Evaluation, (I). Tables, Graphs and References, Chin. Phys. C36 (2012) 1603-2014. [Q-value]

⁹⁹Mo - Comments on evaluation of decay data
 by C. Morillon*, M. M. Bé*, V. Chechev**, A. Egorov**
 * CEA-BNM/LNHB, 91191 Gif sur Yvette France
 ** Khlopin Radium Institute, Saint Petersburg, Russia

This evaluation was completed in December 2000 with minor editing in September 2001. Updated half-life value in 2004.

1- DECAY SCHEME

Molybdenum 99 disintegrates to the technetium 99 excited levels by beta minus transitions. The 1205 keV (3/2-) and 1321 keV (1/2-) levels could be fed by non-unique 1st forbidden β⁻ decays. From lg ft systematic and with lg ft ≥ 8, the β⁻ branches to 1205 keV and 1321 keV levels, if they exist, would be expected ≤ 0,010% and ≤ 0,00014%, respectively. Forbiddenness of other possible β⁻-transitions is still greater. Therefore, all of these unobserved branches can be considered negligible.

Unlike the decay scheme of Peker based mainly on Goswamy (1992Go22), we have not found any justification for placing β⁻ transition to the 534 keV level. The P_{γ+ce} balance for this level has led to the evaluated probability of β⁻ transition of the order of 0,0010(10) %. Also because of the significant lg ft, the attribution of 3/2+ to the 534 keV level seems to be unlikely.

Apart from that, in comparison with 1994Pe15 we have shown a β⁻-transition feeding the 1072 keV level. The spin and parity of this level are not defined exactly. Other J^π values are from Peker.


For this evaluation Mo-99 and Tc-99m are considered being in equilibrium. Therefore, the ratio of their activities is 1,1.

2- NUCLEAR DATA

Q⁻ is from Audi and Wapstra 1995 (95Au04).

- The measured **half-life** values are, in hours :

66,0(1)	Seiler (1947Se25)	²³⁵ U(n,f) ic
66,00(15)	Gunn <i>et al.</i> (1957Gu62)	²³⁵ U(n,f),Mo(n,γ) pc
66,96(9)	Wright <i>et al.</i> (1957Wr37)	⁹⁸ Mo(n,γ)
67,2(2)	Protopopov <i>et al.</i> (1958Pr71)	²³⁵ U(n,f) GM
65,6(2)	Newman (1961Ne15)	²³⁵ U(n,f) pc
66,7(1)	Crowther and Eldridge (1965 Cr03)	⁹⁸ Mo(n,γ) well scin
65,98(10)	Baldwin (1967Ba37)	Mo(n,γ) from 2 meas. pc + scin
66,69(6)	Reynolds <i>et al.</i> (1968Re04)	²³⁵ U(n,f) ic
66,5(2)	Baba <i>et al.</i> (1971Ba28)	²³⁸ U(p,f)
66,02(1)	Emery <i>et al.</i> (1972Em01)	²³⁵ U(n,f)
66,16(30)	Dickens (1979Di07)	
65,945(3)	Houtermans <i>et al.</i> (1980Ho17)	ic
65,924(6)	Hoppes <i>et al.</i> (1982HoZJ)	ic
	Unterweger <i>et al.</i> (1992Un01)	
65,942(12)	Walz <i>et al.</i> (1983Wa26)	Superseded by 2003Sc49
66,02(9)	Butsev <i>et al.</i> (1998)	⁹⁸ Mo(n,γ)
65,85(10)	Butsev <i>et al.</i> (1998)	²³⁵ U(n,f)
64,92(25)	Butsev <i>et al.</i> (1998)	181Ta(12C,x) ⁹⁹ Mo
65,974(14)	Schrader <i>et al.</i> (2003Sc49)	ic



Looking at the graphical representation given above, it appears that 5 values are $\geq 66,5$ h and 12 are in the range $> 65,5$ and $< 66,5$. The five high values are rejected of the statistical treatment (Chauvenet's criterion). The last value given by V.S. Butsev (1998) has also been rejected : $^{181}\text{Ta}(^{12}\text{C},x)^{99}\text{Mo}$ is an exotic reaction, and the result is clearly outlier.

When processing the 17 values, the LWEIGHT program has detected 1957Wr37, 1958Pr71, 1961Ne15, 1965Cr03, 1968Re04, 1971Ba28, 1979Di07 and 2 values of 1998Bu (65,85(10) and 64,92(25)) to be outliers, based on Chauvenet's criterion. The Limited Relative Statistical Weight method increases the uncertainty for the 1980Ho17 value from 0,003 to 0,00481 and used the unweighted mean of 65,983(38) with the large uncertainty that does not correspond to the most accurate measured values (1980Ho17, 1982HoZJ or 1992Un01 and 2003Sc49).

With the set of the 5 most recent values (1972Em01, 1980Ho17, 1982HoZJ or 1992Un01, 1998Bu01 and 2003Sc49), the Limited Relative Statistical Weight method increases the uncertainty for the 1980Ho17 value from 0,003 to 0,00482 and used the weighted mean of 65,949 (14), where 0,014 is the external uncertainty, the reduced- χ^2 is 10,4.

The adopted value is 65,949(14) h or 2,7479(6) d.

- The measured half-life values of the 140,5 keV level of Tc-99 are in ns:

0,277	(14)	STEINER <i>et al.</i> (1969St04)
0,160	(20)	MCDONALD (1971Do02)
0,205	(4)	ALFTER <i>et al.</i> (1993Al23)
0,237	(14)	SHENOY <i>et al.</i> (1973Sh21)

The value of Steiner (1969) given here, is from the original article ; the NDS value (1994Pe15) from the same reference is very different : 0,192 ns.

The value of 0,160(20) ns from J.McDonald (1971) is very far from the others and is not taken into account.

The values from Alfter and Shenoy were determined by using the Moessbauer effect.

The uncertainty on the Alfter *et al.* (1993) value was increased 2,47 times by LRSW.

Reduced- $\chi^2 = 8,94$

LWEIGHT has used the weighted average and the external uncertainty.

The adopted value from the LWEIGHT program is : **0,221(20) ns**

- The measured half-life value of the 181 keV level is **3,61(7) ns** (McDonald (1971))
- The values of the level energies are from NDS 73,1.

2.1 BETA-MINUS TRANSITIONS

The energies of β^- -transitions have been computed from the Q value and the adopted level energies. The probabilities of β^- -transitions have been obtained from the $P_{\gamma+ce}$ balance for each level based on the P_γ normalization factor of 0,1212(15) (see section 4.2.3).

The sum of all the beta transition probabilities leaving the molybdenum must be equal to 100 %; this leads to a probability of 82,1(15)% for the beta transition feeding the 142 keV level, taking into account the gamma transitions feeding this level.

The measured energies and probabilities of some β^- -transitions are given below for comparison with calculated data:

	Measured ^a		Calculated	
	Energy, keV	Probability (%)	Energy, keV	Probability (%)
$\beta^-_{0,12}$	245	0,2	228,1(10)	0,011 (1)
$\beta^-_{0,9}$	450(10)	14	436,6(10)	16,45 (30)
$\beta^-_{0,4}$	840(5)	2	848,1(10)	1,18 (3)
$\beta^-_{0,2}$	1214(1)	80(2)	1214,5(10)	82,1 (15)

^a Nagarajan (1971Na01) except P($\beta^-_{0,2}$) for which unweighted mean of six experimental results quoted in Kholnov (1982KhZW) is given.

2.2 - GAMMA TRANSITIONS and INTERNAL CONVERSION COEFFICIENTS

The evaluated energies of the gamma transitions are the sums of the energies of gamma rays and the recoil energy.

2.2.1- INTERNAL CONVERSION COEFFICIENTS

The ICC have been evaluated using experimental information for the multipolarity admixture coefficients and the theoretical values from 1978Ro22 (Rösel *et al.*) and 1976Ba63 (Band *et al.*) (for $\gamma_{2,1}$).

The relative uncertainties of ICC were adopted to be 2%, for pure multiplicities. The ICC uncertainties for mixed multiplicities were evaluated by taking into account the uncertainties of the respective multipolarity admixture coefficients given in the referenced papers.

The internal conversion coefficients adopted in this evaluation are the theoretical values deduced from the Rösel et al. (1978Ro22) tables. They have been compared with experimental values.

Transition 3-1 : 40,584 keV

Internal Conversion Coefficients α_T

Some authors measured the mixing ratio δ :

δ	First author and NSR code	Transition	α_T (Rösel <i>et al.</i>)
-0,008 (8)	GARDULSKI (1974Ga01)	M1 + 0,0064%E2	3,80
0,03 (3)	SINGH (1982Si16)	M1 + 0,09%E2	3,87
-0,119 (8)	ALFTER (1993Al23)	M1 + 1,4%E2	4,18
	MCDONALD (1971Mc02)	M1 + 1,4(2)%E2	4,18(13) (adopted)

The E2 admixture of 1,4(2) % for $\gamma_{3,1}(40,6 \text{ keV})$ has been adopted from 1971Mc02. The $\gamma(\theta)$ precise measurement of 1993Al23 confirmed this value ($\delta = -0,119(8)$) and rejected the 0,0064 % value of 1974Ga01 which was adopted in Peker's evaluation (1994Pe15). This increases the total ICC for $\gamma_{3,1}$ from 3,76 to 4,18 and improves the intensity balance for the 140,5 keV and 181,1 keV levels.

Internal Conversion Coefficients a_K

a_K	Transition	First author and year
3,2 (2)	M1 transition	Ranakumar (1969)
3,7 (5)	M1 transition	Bashandy (1969Ba03)
3,27 (19)	Weighted average, external uncertainty	LWEIGHT ($\chi^2 = 0,86$)
Adopted: 3,50 (8)	M1+1,4(2)%E2	Rösel <i>et al.</i> (with the adopted admixture)

Internal Conversion Coefficients a_L

From the measurement of the K/L ratio of the conversion electron emission probabilities and, with $\alpha_K = 3,50(8)$, the α_L value is deduced :

K/L	a_L	First author and year
9,3 (20)	0,38(8)	RAVIER (1961)
8,3 (9)	0,42(5)	BASHANDY (1969Ba03)
	0,41(4)	LWEIGHT ($\chi^2 = 0,18$)
Adopted:	0,560 (13)	Rösel <i>et al.</i> for M1+1,4(2)%E2

Transition 1-0 : 140,511 keV

Internal Conversion Coefficients a_T

Experimental measurements :

0,118 (8)	AMTEY <i>et al.</i> (1966)
0,113 (6)	DICKENS and LOVE (1980)
0,122 (5)	VUORINEN (1969)
0,118 (3)	LEGRAND <i>et al.</i> (1973)
0,1181(23)	LWEIGHT (reduced- $\chi^2 = 0,44$ weighted average and internal uncertainty)
Adopted: 0,119(3)	Rösel <i>et al.</i> (1978) for M1+3,2(3)%E2

Dickens and Love (1980) have determined α_T from the α_K value given by Gardulski and Wiedenbeck (1974) and the K/L/MN values reported by Hager and Selzer and by Medsker (NDS - 12-4 - 1974).

α_T was evaluated by Vuorinen (1969) from measurements of conversion electrons in coincidence with fluorescence X-rays.

Multipolarity

There are a significant number of measurements. However most authors gave different values with and without large uncertainties: these multipolarities make it possible to calculate the total internal conversion coefficients. We have assigned a 5% uncertainty to α_T :

/d/	Transition	a_T (Rösel)	
0,31 (2)	M1 + 8,25% E2	0,132(7)	SINGH and SAHOTA (1982Si16)
0,178 (12)	M1 + 3,1% E2	0,119(6)	ALFTER <i>et al.</i> (1993Al23)
	M1 + 4%(2) E2	0,121(6)	MCDONALD <i>et al.</i> (1971Mc02)
	M1+<3%E2		VOINOVA <i>et al.</i> (1971Vo06)
0,194(30)	M1+E2		VUORINEN (1969Vu03)
	M1+<8%E2		VAN EIJK <i>et al.</i> (1968Va14),calculated from ICCK
	M1+9%(5)E2	0,134(7)	VAN EIJK <i>et al.</i> (1968), calculated from K/L ratio
	M1+2,8%E2	0,118(6)	COOK <i>et al.</i> (1969 Co18)
	M1+7(3)%E2	0,129(7)	MEYER (1974)
	M1+1,4%E2	0,114(6)	DICKENS and LOVE (1980Di16)
	M1+6,5(40)E2	0,128(7)	AGEEV <i>et al.</i> (1969Ag04)
0,118(6)	M1+1,4(2)%E2	0,114(6)	GARDULSKI and WIEDENBECK (1974Ga01)
	M1+2,8(3)%E2	0,118(6)	GEIGER (1968GeZW)
	M1+9%E2		SIMONITS <i>et al.</i> (1982Si15)
	M1+E2		AMTEY <i>et al.</i> (1966Am04)
	M1		BASHANDY (1969Ba54)
		0,120(2)	LWEIGHT (reduced- $\chi^2= 1,16$), weighted average and external uncertainty= 0,0015
0,186 (8)	M1 + 3,2(3)%E2	0,119 (3)	Adopted (Rösel <i>et al.</i>)

From each determination of the multipolarity of the transition, the Rösel theoretical internal coefficient was calculated. From the set of the 10 deduced ICC values the LWEIGHT program recommends a weighted mean of 0,120(2). The value obtained is very close to that obtained by considering the 4 experimental values for α_T (see table above).

Internal Conversion Coefficients α_K

Experimental values:

0,096 (6)	VOINOVA <i>et al.</i> (1971Vo06)
0,093 (6)	VOINOVA <i>et al.</i> (1971Vo06)
0,102 (7)	VAN EIJK <i>et al.</i> (1968Va14)
0,094 (8)	VUORINEN (1969Vu03)
0,102 (5)	DICKENS and LOVE (1980Di16)
0,096 (3)	LWEIGHT ($\chi^2=0,35$; weighted average and internal uncertainty)
0,104 (3)	Rösel <i>et al.</i> (1978) (adopted)

- α_K was measured by Voinova *et al.* (1971) with a spectrometer which provided simultaneous measurement of conversion electrons and γ -ray spectra.
- Van Eijk *et al.*(1968) calculated ICCK from measurements of the 140,5 keV gamma-ray emission probability (P_γ) relative to the gamma-ray emission probability of the 661,6 keV gamma transition in decay of Cs-137 and from measurements of the conversion electron emission probability P_{ce} of the 140,5 keV K-conversion line relative to the conversion electron emission probability of the 661,6 keV K-conversion line in decay of Cs-137. With $P_{ceK} = 6,84(19)$; $P_\gamma = 6,00(35)$; $\alpha_K(661,6 \text{ keV}) = 0,0896(15)$ (Helmer in BÉ 1999 (1999BeZQ)), the value becomes 0,102(7).
- Vuorinen (1969) evaluated the internal conversion coefficient α_K by measuring the electron conversion emissions following the conversion of the 140 keV gamma ray in coincidence with fluorescence X-rays.

- α_K given by Dickens and Love (1980) was computed from the tables of Hager and Seltzer for a M1 transition and a 1,4% E2 admixture. An 5% uncertainty assigned to α_K reflects the added uncertainty to the usual 3% assignment due to the rapid change of α_K with admixture. This value is not taken into account in our calculations.

Internal Conversion Coefficients α_L

From each measurement of the K/L ratios of the conversion electron emission probabilities, and with $\alpha_K = 0,104(3)$, a value for α_L is deduced :

K/L	α_L	
8,1 (5)	0,0125(8)	BASHANDY(1969Ba03)
7,70 (30)	0,0132(7)	VAN EIJK <i>et al.</i> (1968Va14)
8,3 (3)	0,0122(6)	RAVIER <i>et al.</i> (1961Ra04)
7,63 (32)	0,0133(7)	BRAHMAVAR (1968)
7,8 (3)	0,0130(6)	GEIGER (1968GeZW)
	0,0128(3)	LWEIGHT has used the weighted average and the internal uncertainty. Reduced- $\chi^2 = 0,52$
Adopted	0,0129 (4)	Rösel <i>et al.</i> (1978)

Transition 2-0: 142,683 keV

Internal Conversion Coefficients α_T

For a M4 transition the theoretical value from Rösel is : **40,9(8)**.

Internal Conversion Coefficients α_K

- The 2 following values were calculated from experimental data and given by the authors :
 23 (6) Van Eijk *et al.* (1968)
 30 (3) Bashandy (1969Ba54)

Van Eijk *et al.* (1968) calculated the K ICC value from the ratios of $K(142,7)/K(140,5) = 0,072(32)$ and $I_\gamma(142,7)/I_\gamma(140,5) = 0,00030(6)$ after correction for $\alpha_K(661,6 \text{ keV, Cs-137}) = 0,0896(15)$

Bashandy (1969) calculated the K ICC from internal conversion spectra and photon emission probabilities $I_\gamma(142)/I_\gamma(140) = 0,00030(6)$

- The following α_K coefficients are calculated from the $K(142,7)/K(140,5)$ ratio given by the authors and based on the ratio $I_\gamma(142,7)/I_\gamma(140,5) = 0,00030(6)$ given by Van Eijk *et al.* (1968) and on $\alpha_K(140,5) = 0,104(3)$.

$K(142,7)/K(140,5)$	$\alpha_K(142,7)$	
0,072(4)	24 (6)	AMTEY <i>et al.</i> (1966Am04)
0,0746(12)	25 (6)	GEIGER (1968GeZW)
0,075 (8)	26 (6)	AGEEV <i>et al.</i> (1969Ag04)

If we take into account the ratio $I_\gamma(142,7)/I_\gamma(140,5) = 0,00021(3)$ given by Dickens and Love (1980Di16), with $\alpha_K(140,5) = 0,104(3)$, the same calculations give higher results for $\alpha_K(142,7)$:

$K(142,7)/K(140,5)$	$\alpha_K(142,7)$	
0,072(4)	34 (6)	AMTEY <i>et al.</i> (1968)
0,0746 (12)	36 (5)	GEIGER (1968)
0,075 (8)	36 (7)	AGEEV <i>et al.</i> (1969)

If we have taken into account all the six possible data, the weighted average, with the external uncertainty, calculated by LWEIGHT is 29,5(18) (reduced- $\chi^2=0,87$)

The **adopted** theoretical K conversion coefficient, for a M4 transition, is : **29,3(6)** (Rösel *et al.* (1978)).

Internal Conversion Coefficients α_L

From the measurement of the ratio of the conversion electron intensities (BASHANDY and IBRAHIEM), with $\alpha_K = 29,3(6)$, α_L can be deduced. This value is close to the adopted theoretical value:

K/L	α_L		
2,9 (5)	10,1 (18)	M4 transition	BASHANDY and IBRAHIEM
Adopted:	9,35 (20)	M4 transition	Rösel <i>et al.</i> (1978)

Transition 3-0 : 181,094 keV

Internal Conversion Coefficients α_T

0,140(5) DICKENS and LOVE (1980Di16)
 GARDULSKI and WIENBECK (1974Ga01) measured a low multipole mixing ratio of 0,002(7) for a M3/E2 transition.

For a E2 transition, the theoretical value is : **0,149(3)** (Rösel *et al.* (1978))

Internal Conversion Coefficients α_K

0,13(3)		RAVIER <i>et al.</i> (1961)
0,127(11)*	E2 ≤ 12%M1	VAN EIJK <i>et al.</i> (1968)
0,133(20)	E2 transition	BASHANDY (1969Ba54)
0,12(1)		VOINOVA <i>et al.</i> (1972)
0,125(7)		LWEIGHT (reduced- $\chi^2 = 0,16$, weighted average and the internal uncertainty)
0,125 (3)	E2 transition	Rösel <i>et al.</i> (adopted)

(*) value corrected for $\alpha_K(661\text{keV Cs-137})=0,0896(15)$ (Helmer in Bé 1999)

Internal Conversion Coefficients α_L

From the measurement of ratio K/L of conversion electron intensities, with $\alpha_K = 0,125(3)$, α_L can be deduced:

K/L	α_L	Transition	
4,9 (1)	0,025(6)		RAVIER <i>et al.</i> (1961)
6,8 (7)	0,0184(20)		BASHANDY (1969Ba03)
Adopted:	0,0191 (4)	E2	Rösel <i>et al.</i> (1978)

Transition 4-2 : 366,422 keV

Internal Conversion Coefficients α_T

0,0081 (2)		DICKENS and LOVE (1980)
0,00915 (18)	M1 transition	Rösel <i>et al.</i> (1978) (adopted)

Internal Conversion Coefficients a_K

0,0072 (10)		BASHANDY (1969Ba54)
0,00802(16)	M1 transition	Rösel <i>et al.</i> (1978) (adopted)

Transition 13-7 : 380,13 keV**Internal Conversion Coefficients a_K**

0,009 (1)	M1+E2	BASHANDY (1969Ba54)
0,0091 (7)	M1+63(22)%E2	Rösel <i>et al.</i> (1978) (adopted)

- From the value of Bashandy (1969Ba54), it can be deduced a M1+63%E2 transition and multipole mixing ratio $\delta = 1,3(6)$.

Transition 14-7 : 410,27 keV**Internal Conversion Coefficients a_K**

0,0060 (8)		BASHANDY (1969Ba54)
0,0065 (2)	M1+20(3)%E2	Rösel <i>et al.</i> (1978) (adopted)

Transition 9-4 : 411,492 keV**Internal Conversion Coefficients a_K**

0,0030 (5)	E1 transition	BASHANDY (1969Ba54)
0,00226(5)	E1 transition	Rösel <i>et al.</i> (1978) (adopted)

Transition 12-6 : 457,60 keV

The E2 admixture of 72(55) % has been adopted from the evaluation of Kholnov (1982KhZW).

Internal Conversion Coefficients a_K

0,0054 (6)		BASHANDY (1969Ba54)
0,0054 (4)	M1+72(55)%E2	Rösel <i>et al.</i> (1978) (adopted)

Transition 6-2 : 528,790 keV**Internal Conversion Coefficients a_K**

0,0050 (6)	E2 transition	BASHANDY (1969Ba54)
0,00375(8)	E2 transition	Rösel
0,00331(7)	M1 transition	Rösel <i>et al.</i> (1978) (adopted)

Transition 8-1: 621,771 keV**Internal Conversion Coefficients α_K**

0,0020 (4)		BASHANDY (1969Ba54)
0,00227 (5)	M1 transition	Rösel <i>et al.</i> (1978) (adopted)

Transition 9-3 :739,503 keV**Internal Conversion Coefficient α_K**

0,0016 (4)	M1 or E2	BASHANDY(1969Ba54)
0,00154 (40) *		VAN EIJK et a.l. (1968)
0,00151 (3)	E2+7,6%M1	Rösel <i>et al.</i> (1978) (adopted)

*value corrected for $\alpha_K(661\text{keV Cs-137}) = 0,0896(15)$ (Helmer in BÉ 1999)

The multipole mixing ratio : $\delta = 3,58(20)$ measured by Gardulski and Wiedenbeck (1974), leads to an E2 + 7,2% M1 transition.

Singh and Sahota (1982) indicated an E2 + 8,0(1)%M1 multipolarity.

Transition 9-2 : 777,924 keV**Internal Conversion Coefficient α_K**

0,0005 (1)		BASHANDY (1969Ba54)
0,000518 (10)	E1 transition	Rösel <i>et al.</i> (1978) (adopted)

Transition 10-3 : 822,976 keV**Internal Conversion Coefficient α_K**

0,0004 (1)		BASHANDY(1969Ba54)
0,0004 (1)	E1+1%M2 transition	SINGH (1982)
0,000461(9)	E1 transition	Rösel <i>et al.</i> (1978) (adopted)

For an E1+1%M2 transition, the theoretical value would be higher than the experimental values and we do not accept this type of transition.

Transition 13-3 : 960,759 keV**Internal Conversion Coefficient**

Based on $\alpha_K = 0,0024(5)$ Bashandy deduced a M2 multipolarity. From the decay scheme Singh gave a M2 + E3 multipolarity. This is not consistent with the adopted spins and parities which lead to a M1+E2 transition. For a M1 transition, $\alpha_T = 0,00097$ from the Rösel tables.

Transition 13-1 : 1001,348 keV**Internal Conversion Coefficient**

Based on $\alpha_K = 0,0018(3)$ Bashandy deduced a M2+E3 multipolarity. This is not consistent with the adopted spins and parities which lead to a E2+M3 transition. For a E2 transition, $\alpha_T = 0,00083$ from the Rösler tables.

2.2.2 GAMMA TRANSITION PROBABILITIES

The gamma transition probabilities have been calculated from the gamma emission probabilities and the internal conversion coefficients for the transitions occurring above the 142 keV level.

The total gamma and beta transition probabilities populating the 142 keV level is : 87,65(19)%. Within the Tc-99m decay, the 2,17 keV gamma transition probability (from the level 2 to the level 1) is deduced to be : 99,0(4)%; the 142 keV gamma transition probability is evaluated to be : 1,0(1) % and the 140 keV gamma transition probability is 99,0(4)%.

So, the transition probabilities are deduced to be : 86,8(19)% and 0,88(6)% for the 2,17 keV and the 142 keV, respectively. Taking into account the level balance, the 140 keV transition probability is deduced to be 92,1(19) %.

3. Atomic Data**3.1. Fluorescence yields**

- ω_K is from Bambynek (1984)
- ω_L , η_{KL} , η_{LM} are from Schönfeld and al.(1996)
- ω_M is from Hubbell and al. (1994)

3.2. X Radiations

The X-ray energies are based on the wave lengths in the compilation of 1967Be65 (Bearden). The relative K X-ray emission $K\beta/K\alpha$ and $K\alpha_2/K\alpha_1$ probabilities are taken from 1996Sc06.

3.3. Auger Electrons

The energies of Auger electrons are from 1977La** (Larkins). The ratios $P(KLX)/P(KLL)$ and $P(KLY)/P(KLL)$ are taken from 1996Sc06.

4. Photon Emissions**4.1. X-Ray Emissions**

The total absolute emission probability of K X-rays (P_{XK}) has been computed using the adopted value of ω_K and the evaluated total absolute emission probability of K conversion electrons (P_{ceK}). The absolute emission probabilities of the K X-ray components have been computed from P_{XK} using the relative probabilities from 1996Sc06.

The measured values of the total absolute emission probability of K X-rays ($P_{XK} \times 100$) are given below in comparison with the calculated (adopted) value:

Dickens and Love	Goswamy	Calculated (adopted)
11,3(5)	11,5(4)	11,2(2)

Above agreement of the measured and calculated values shows concord between the evaluated data for ⁹⁹Mo including the gamma -ray emission probabilities, gamma -multipolarity admixtures, ICC α_K and the fluorescence yield ω_K .

The total absolute emission probability of L X-rays has been computed using total absolute sums P_{ceL} , P_{ceK} , and atomic data of section 3 (ω_K , ω_L , η_{KL}).

M X-ray and Auger spectra have been investigated in Gerasimov. The influence of the chemical state on the K X-ray intensity has been studied in Yoshihara (1981Yo08).

4.2. GAMMA RAY EMISSIONS

4.2.1 GAMMA RAY ENERGIES

The γ -ray energies of $\gamma_{2,1}$ (2,17 keV), $\gamma_{3,1}$ (40,6 keV) and $\gamma_{1,0}$ (140,5 keV) are taken from Gerasimov (1981Ge05), Gardulski (1972Ga37) and Helmer (2000He14), respectively. These values are based on the most accurate measurements with the electrostatic spectrometer ($E_{\gamma_{2,1}}$, see also Lacasse (1971La12)) and curved-cristal spectrometer ($E_{\gamma_{3,1}}$ and $E_{\gamma_{1,0}}$, see also Helmer (1981He15)). The energies of $\gamma_{2,0}$ (142,7 keV), $\gamma_{3,0}$ (181,1 keV), $\gamma_{7,0}$ (761,7 keV) and $\gamma_{11,0}$ (1072,2 keV) have been computed from the Q value and the adopted energies of other gamma transitions using gamma cascades in the decay scheme. The energy of $\gamma_{15,4}$ (689,6 keV) is taken from 1969Co18 (this γ -ray was seen also by Goswamy *et al.* (1992Go22) but was defined as some contamination in the source). All other gamma -ray energies have been adopted from the recent measurements with large volume Ge(Li) and high-purity Ge detectors by R.A. Meyer (1990Me15).

4.2.2 GAMMA RAY RELATIVE EMISSION PROBABILITIES

Several authors measured the relative emission probabilities to the emission probability of 739 keV line, and others to the emission probability of the 140,5 keV line.

In this evaluation the 739 keV line is taken as the reference line rather than the 140 keV line because the 739 line is not a part of the Tc-99m decay scheme, and the measurements carried out relative to this line, are more recent.

Measurements relative to the 140,5 keV line have been taken into account by converting the data so that they are relative to the 739 keV line.

The available experimental values for the γ -ray relative emission probabilities are given in Table 1. Where necessary, these data (including uncertainties) have been converted by the evaluators to values relative to the $\gamma_{9,3}$ (739,5 keV) taken as 100. Some old references differ widely far from more recent studies and are not included in the statistical processing.

The adopted (evaluated) values are displayed in last column of Table 1. Reasons for adopting specific data are given in Table 2 which includes the following designations :

R indicates that the value was rejected due to Chauvenet criteria.

n is the number of values taken into account, WM is the weighted mean, s and S are the internal and external uncertainties of WM, respectively;

" χ^2 -table" is $(\chi^2)^{0,05}_{n-1}$, "reduced χ^2 -set" is $\chi^2/(n-1)$ for the given data set; s_{min} is the minimum experimental uncertainty for the given data set, tS is the external uncertainty multiplied by the Student's factor *t*, "MBAYS" is the uncertainty from a modified Bayesian analysis.

The doublet $\gamma_{14,7}+\gamma_{9,4}$ (410-411 keV) has been calculated as two different lines because several authors were able to distinguish separated values.

For the doublet $\gamma_{7,3}+\gamma_{8,3}$ (580-581 keV) several authors measured only one line, except Meyer (see Table 1).

For the doublet $\gamma_{12,4}+\gamma_{8,1}$ (620-622 keV) the emission intensity was computed for the two combined lines in order to take into account most of the measurements, and then these lines were separated by using the intensity ratio for components deduced from the measurements of Meyer of 0,09(3).

Table 1. Experimental and evaluated values for γ -ray relative emission probabilities

	keV	Van Eijk	Cook	Gehrke Heath	Morel	Dickens 1980	Yang 1980	Singh	Chen Da	Meyer 1990	Goswamy 1992	Evaluated
$\gamma_{3,1}$	40,58		6,9(8)	4,6(18) <i>R</i>	5,9(15)	8,68(27)		7,7(6)		8,6(5)	8,49(25)	8,43(20)
$\gamma_{1,0}$	140,5	649(25)	704(45)	730(49)	743(19)	747(12)	759(20)	686(49)	752(28)	755(26)	739(11)	739(11)
$\gamma_{2,0}$	142,6	0,195(40)				0,149(25)				0,189(11)		0,174(14)
$\gamma_{9,7}$	158,8		0,10(3)	0,095(30)	0,112(15)			0,11(4)		0,139(8)	0,156(6)	0,12(4)
$\gamma_{6,4}$	162,4			0,073(22)	0,067(15)			0,078(13)		0,097(5)	0,098(5)	0,094(5)
$\gamma_{3,0}$	181	48,7(23)	49,9(34)	49,6(42)	49,1(16)	50,1(7)		49,8(33)	48,7(13)	50,3(17)	49,4(8)	49,6(7)
$\gamma_{10,7}$	242,3		0,0070(25)					0,0118(44)		0,0117(17)	0,021(4)	0,0114(28)
$\gamma_{9,6}$	249	0,039(20)	0,05(2)					0,04(3)		0,024(3)	0,032(4)	0,0285(30)
$\gamma_{4,2}$	366,4	10,6(8)	10,7(6)	10,0(9)	9,8(3)	9,52(32)		9,8(8)		9,92(25)	9,82(15)	9,85(15)
$\gamma_{13,7}$	380,1	0,071(20)	0,07(2)	0,058(15)	0,045(15)			0,07(2)		0,075(3)	0,086(7)	0,075(4)
$\gamma_{5,2}$	391,7	0,016(4)									0,026(5)	0,021(5)
$\gamma_{14,7}$	410,3	0,010(5)						0,009(9)		0,016(4)		0,013(3)
$\gamma_{9,4}$	411,5	0,18(2)	0,13(2)	0,36(4) <i>R</i>	0,134(23)			0,14(2)		0,120(6)		0,133(10)
$\gamma_{12,6}$	457,6	0,039(20)	0,08(2)					0,04(2)		0,056(5)	0,067(5)	0,061(5)
$\gamma_{10,5}$	469,6		0,0060(15) <i>R</i>							0,022(4)	0,022(4)	0,022(4)
$\gamma_{6,2}$	528,8	0,39(5)	0,49(5)	0,36(4)	0,43(6)			0,44(4)		0,447(15)	0,47(2)	0,446(15)
$\gamma_{11,5}$	537,8		0,0100(25)					0,009(3)		0,013(5)	0,027(5)	0,012(4)
$\gamma_{7,3}$	580,5	0,026(7)						0,021(8)		0,036(4)	0,026(4)	0,0294(31)
$\gamma_{8,3}$	581,3									0,008(4)		0,008(4)
$\gamma_{12,4}^+$	620	0,21(3)	0,217(22)	0,19(6)	0,30(4)			0,26(2)		0,232(11)	0,24(4)	0,236(11)
$\gamma_{8,1}$	621,7											

	keV	Van Eijk	Cook	Gehrke Heath	Morel	Dickens 1980	Yang 1980	Singh	Chen Da	Meyer 1990	Goswamy 1992	Evaluated
$\gamma_{15,4}$	689,6		0,0035(15)									0,0035(15)
$\gamma_{9,3}$	739,5	100	100	100	100	100	100	100	100	100	100	100
$\gamma_{7,0}$	761,8		0,019(5)							0,0092(8)	0,033(3)	0,019(11)
$\gamma_{9,2}$	777,9	35,1(24)	34,9(20)	35,8(30)	35,5(10)	35,8(9)		34,8(19)		35,3(12)	35,1(5)	35,3(5)
$\gamma_{10,3}$	822,9	1,04(8)	1,11(8)	1,09(10)	1,09(5)	1,09(5)		1,10(7)		1,06(4)	1,10(2)	1,09(2)
$\gamma_{10,2}$	861,2		0,006(2)	0,015(6)				0,005(3)			0,006(3)	0,006(2)
$\gamma_{13,3}$	960,722	0,78(7)	0,78(6)	0,80(8)	0,76(4)	0,84(4)		0,79(6)		0,76(4)	0,78(2)	0,78(2)
$\gamma_{12,2}$	986,4	0,013(5)	0,014(4)	0,016(4)						0,0108(9)	0,012(4)	0,0112(8)
$\gamma_{13,1}$	1001	0,045(13)	0,036(16)	0,027(4)	0,052(15)			0,045(12)		0,033(1)	0,045(4)	0,035(3)
$\gamma_{15,3}$	1017	0,006(3)									0,005(2)	0,0055(21)
$\gamma_{15,2}$	1056,2		0,008(2)					0,007(3)		0,0083(9)	0,0089(7)	0,0085(7)
$\gamma_{11,0}$	1072,2							0,010(4)				0,010(4)

Table 2. Results of data statistical processing on relative γ -ray emission probabilities

	n	WM	s	S	c ²		Final uncertainty and type
					table	set	
$\gamma_{3,1}$	6	8,43	0,16	0,20	14,07	1,82	0,20 (S)
$\gamma_{1,0}$	10	739	5,7	7,6	18,31	2	11 (<i>S_{min}</i>)
$\gamma_{2,0}$	3	0,174	0,014	0,014		1	0,014 (S)*
$\gamma_{9,7}$	6	0,12 ^d	0,0047	0,0078	11,07	3	0,04 (S)
$\gamma_{6,4}$	5	0,094	0,0033	0,0042	9,49	1,6	0,005 (<i>S_{min}</i>)
$\gamma_{3,0}$	10	49,6	0,42	0,20	16,92	0,13	0,7 (<i>S_{min}</i>)
$\gamma_{10,7}$	4	0,0114	0,0014	0,0024	7,82	2,96	0,0028 (<i>tS</i>)
$\gamma_{9,6}$	5	0,0285	0,0027	0,0026	9,49	0,9	0,0030 (<i>S_{min}</i>)*
$\gamma_{4,2}$	9	9,85	0,11	0,08	15,51	0,58	0,15 (<i>S_{min}</i>)
$\gamma_{13,7}$	7	0,075	0,0037	0,0042	12,59	1,3	0,004 (S)*
$\gamma_{5,2}$	2	0,021	0,0035	0,005		2	0,005 (S)*
$\gamma_{14,7}$	3	0,013	0,003	0,002		0,56	0,003 (S)
$\gamma_{9,4}$	5	0,133	0,007	0,01		1,81	0,01 (S)*
$\gamma_{12,6}$	5	0,061	0,0034	0,0040	9,49	1,4	0,005 (<i>S_{min}</i>)
$\gamma_{10,5}$	3	0,022 ^b					0,004 ^b
$\gamma_{6,2}$	7	0,446	0,010	0,012	12,59	1,1	0,015 (<i>S_{min}</i>)
$\gamma_{11,5}$	4	0,012	0,0017	0,0032	7,82	3,6	0,0038 (<i>tS</i>)
$\gamma_{7,3}$	4	0,0294	0,0025	0,0031		1,6	0,0031 (S)
$\gamma_{8,3}$	1	0,008					0,004
$\gamma_{12,4} + \gamma_{8,1}$	7	0,236	0,0083	0,0085	12,59	1	0,011 (<i>S_{min}</i>)*
$\gamma_{15,4}$		0,0035 ^c					0,0015 ^c
$\gamma_{9,3}$		100					
$\gamma_{7,0}$	3	0,019	0,0018	0,0077	5,99	18	0,011 (MBAYS)*
$\gamma_{9,2}$	9	35,3	0,34	0,17	15,51	0,2	0,5 (<i>S_{min}</i>)
$\gamma_{10,3}$	8	1,09	0,015	0,0063	14,07	0,1	0,02 (<i>S_{min}</i>)*
$\gamma_{10,2}$	4	0,006	0,0014	0,0012	7,82	0,6	0,002 (<i>S_{min}</i>)
$\gamma_{13,3}$	9	0,78	0,014	0,0083	15,51	0,08	0,02 (<i>S_{min}</i>)
$\gamma_{12,2}$	5	0,0112	0,0015	0,0008	9,49	0,44	0,0008 (S)
$\gamma_{13,1}$	7	0,035	0,0017	0,0026	12,59	1,9	0,0028 (<i>tS</i>)*
$\gamma_{15,3}$		0,0055 ^d					0,0021 ^d
$\gamma_{15,2}$	4	0,0085	0,00056	0,00025	7,82	0,22	0,0007 (<i>S_{min}</i>)*
$\gamma_{11,0}$		0,010 ^e					0,004 ^e

^a Adopted from Goswamy (1992Go22)

^b Adopted from Meyer (1990Me15) and 1992Go22 (the same values)

^c Adopted from Cook (1969Co18)

^d Unweighted average

^e Adopted from Singh (1982Si16)

* LRSW increased an uncertainty for one of the values(1992Go22 or 1990Me15).

All values for relative γ -ray emission probabilities are given for the equilibrium mixture ⁹⁹Mo + ⁹⁹Tc^m.

For $\gamma_{2,0}$ (142,7 keV) the following measured intensity ratios of $\gamma_{2,0}/\gamma_{1,0}$ (140,5 keV) have been used: 3,0(6)·10⁻⁴ (Van Eijk), 2,0(2) ·10⁻⁴ (Ageev), 2,0(3) ·10⁻⁴ (Dickens, 1980Di16), 2,50(9) ·10⁻⁴ (Meyer, 1990Me15). The weighted average of these values is 2,29(16) ·10⁻⁴ with an external uncertainty; in terms of the $\gamma_{9,3}$ (739,5 keV) a relative intensity of 0,169(12) is obtained.

For $\gamma_{11,0}$ (1072,2 keV) the relative γ -ray emission probability is taken from Singh (1982Si16).

4.2.3 GAMMA RAY ABSOLUTE EMISSION PROBABILITIES

Several absolute measurements of the emission intensity of the 739 keV line are available to give a consistent set of data.

Emission 9 - 3 : 739,500(17) keV**Absolute measurement : photon emission per 100 decays**

11,9 (3)	Chen Da - 1985 (Ge(Li) gamma spectrometer) (measured)
12,3 (3)	Simonits (1981)
12,14 (22)	Dickens and Love(1980) (calculated)
12,00 (33)	Meyer (Fizika - 22 - p153 (1990))

Lweight has used the weighted average and the internal uncertainty. Reduced- $\chi^2=0,45$

Adopted absolute g emission probability: 12,12(15)%

This absolute γ -ray emission probability can be compared with the value obtained by considering the balance of the decay scheme. The γ -ray absolute emission probabilities P_γ have been computed using relative ($\gamma+ce$)-probabilities (relatively to the 739,5 keV gamma ray) and the ⁹⁹Tc ground state intensity balance, which assumes no β -feeding to the g.s. and the 140,5 keV level as confirmed by the high degree of forbiddenness. The P_γ intensity of the 739 keV line has been deduced to be 12,18(17)% taking into account the correlation $\Sigma P_\beta=1$ and the factor of 1,100 for the gamma transitions in Tc-99m.

All the absolute gamma ray emission probabilities are given per 100 disintegrations of Mo⁻⁹⁹ (in equilibrium with Tc-99m) taking into account the correction factor of 1,100 for $\gamma_{2,1}(2,17 \text{ keV})$, $\gamma_{2,0}(142,7 \text{ keV})$ and $\gamma_{1,0}(140,5 \text{ keV})$ intensities.

It should be noted that Singh and Sahota (1982Si16) have reported nine controversial γ -rays at energies of 38,4; 163,4; 319,8; 321,0; 352,9; 599,6; 721,7; 940 and 1082,0 keV. These γ -rays have not been confirmed by Goswamy *et al.* (1992Go22) and are not placed in the decay scheme; neither are the 344,6 keV γ -ray observed by Cook *et al.* (1969Co18) and the 89 ,4; 455,84; 490,53 keV γ -rays observed by Meyer (1990Me15).

5. Electron Emissions

The energies of the conversion electrons have been calculated from the gamma⁻transition energies given in 2.2 and the electron binding energies. The emission probabilities have been calculated using the conversion coefficients given in 2.2. and the gamma emission probabilities.

Many measurements of conversion electron spectra for ⁹⁹Mo in equilibrium with ⁹⁹Tc^m have been made (1968Va14, 1969Ag04, 1969Ba03, 1969Ba54, 1969Ra01, 1971La12, 1971Vo06, 1973Le29, 1981Ge05). However the computed values of the conversion electron energies and emission probabilities are more accurate.

The values of the emission probabilities of K-Auger electrons have been calculated using the gamma transition probabilities given in 2.1 and 2.2, the atomic data given in 3. and the conversion coefficients given in 2.2.

Experimental Auger spectra can be found in 1981Ge05.

BETA-MINUS EMISSIONS

The β^- transition energies are derived from the level energies.

T. NAGARAJAN (1971Na01) analysed the β spectrum of Mo-99. This study revealed four β groups with end points :

	Energy keV	Transition probability
$\beta_{0,2}$	1214(1)	84
$\beta_{0,4}$	840(5)	2
$\beta_{0,9}$	450(10)	14
$\beta_{0,12}$	245	<0,2

No evidence was found for a β group with endpoint higher than 1214 keV.

These values are in a rough agreement with those established by considering the balance of the decay scheme (paragraph 2.1).

REFERENCES

- J. A. SEILER, Report ANL-4000(1947)119 [Half-life]
 S. R. GUNN, H. G. HICKS, H. B. LEVY, P. C. STEVENSON, Phys. Rev. 107(1957)1642 [Half-life]
 H. W. WRIGHT, E. I. WYATT, S. A. REYNOLDS, W. S. LYON, T. H. HANDLEY, Nucl. Sci. Eng. 2(1957) 427 [Half-life]
 A. N. Protopopov , G. M.Tolmachev et al., J. Nucl. Energy 10A(1959)80 [Half-life]
 J. RAVIER, P. MARGUIN, A. MOUSSA, J. Phys. Radium 22(1961)249 [K/L]
 R. D. Newman, Priv. Comm. (1961) [Half-life]
 P. CROWTHER, Nucl. Phys. 66(1965)472 [Half-life, Gamma emission probabilities, T ICC, K ICC]
 S. R. AMTEY, J. H. HAMILTON, M. J. ZENDER, Phys. Lett. 23-10(1966)581 [K ICC, L ICC, M ICC]
 J. A. Bearden, Rev. Mod. Phys., 39,1(1967)78 [X-Rays energies]
 M. N. BALDWIN, Nucl. Sci. Eng. 30(1967)144 [Half-life]
 C. W. E. VAN EIJK, B. VAN NOOIJNEN, F. SCHUTTE , Nucl. Phys. A121(1968)440 [Gamma-ray energies, K-Conv. Elec. , Gamma-ray emission probabilities]
 J. S. GEIGER, AECL-3166 PR-P-79(1968)29 [Gamma-ray energies, K-Conv. Elec., Gamma emission probabilities]
 S. M. BRAHMAVAR, J. H. Hamilton, J. J. Pinajian, Quoted in Nucl. Phys. A121(1968)440 [ICC]
 S. A. Reynolds, J. F. Emery, E. I. Wyatt, Nucl. Sci. Eng. 32(1968)46 [Half-life]
 N. RANAKUMAR, R. W. FINK, P. V. RAO, Nucl. Phys. A217(1969)683 [K ICC]
 W. B. COOK, L. SCHELLENBERG, M. W. JOHNS, Nucl. Phys. A139(1969)277 [Gamma-ray energies, Gamma-ray emission probabilities]
 E. BASHANDY, Z. Naturforsch 24A(1969)1893 [K ICC]
 V. A. AGEEV et al., Bull. Ac. Sc. USSR, Phys. Ser. 33(1969)1183 [Conv. Elec. emission probabilities]
 A. VUORINEN, Ann. Acad. Sci. Fenn. Ser. A VI.(1969)311 [T ICC]
 E. BASHANDY, N. IBRAHIEM, Z. Phys. 219(1969)337 [K ICC, K/L]
 P. STEINER, E. Gerdau, W. Hautsch, D. Streenken, Z. Physik 221(1969)281 [Level half-life]
 N. A. VOINOVA et al., Bull. Ac. Sc. URSS, Phys. Ser. 35(1971)794 [K ICC]
 J. Mc DONALD , A. BACKLIN, S. G. MALMSKOG, Nucl. Phys. A162(1971)365 [Mean half-life, Multipolarity mixing ratio]
 T. NAGARAJAN, M. RAVINDRANATH, K. V. REDDY, Phys. Rev. C3(1971)247 [Beta emission energies]
 W. M. Lacasse, J. H. Hamilton, Nucl. Phys. A171(1971)641 [Gamma ray energies]
 S. BABA, H. BABA, H. NATSUME, J. Inorg. Nucl. Chem. 33(1971)589 [Half-life]
 P. L. GARDULSKI, M. L. WIEDENBECK, Nucl. Instrum. Methods 105(1972)169 [Gamma-ray energies]
 N. A. VOINOVA, A. I. EGOROV, Y. V. KALINICHEV, A. G. SERGEEV, Bull. Ac. Sc. URSS, Phys. Ser. 35(1972)794 [K ICC]
 J. F. EMERY, S. A. REYNOLDS, E. Y. WYATT, G. I. GLEASON, Nucl. Sci. Eng. 48(1972)319 [Half-life]

- G. K. SHENOY, G. ABSTREITER, G. M. KALVI US, K. SCHWOCHAU, K. H. LINSE, J. Phys. A (London) 6(1973)L144 [Level half-life]
- J. LEGRAND et al., Report CEA-R-4427(1973) [Gamma-ray emission probabilities, Conv. Elec. emission probabilities]
- R. J. GEHRKE, L. D. Mac ISSAC, R. L. HEATH, Report ANCR-1129(1973)23 [Gamma-ray energies, Gamma-ray emission probabilities]
- P. L. GARDULSKI, M. L. WIEDENBECK, Phys. Rev. C9(1974)262 [Spin and Parity, Mixing Ratio]
- R. L. HEATH, Report ANCR-1002(1974) [Gamma-ray energies, Gamma-ray emission probabilities]
- R. A. MEYER, Report UCRL-76207(1974) [Spin and Parity]
- F. P. Larkins, At. Data Nucl. Data Tables 20,4(1977)338 [Auger Electrons]
- J. MOREL, J. P. PEROLAT, Report CEA-N-2043(1978) [Gamma-ray emission probabilities]
- J. K. DICKENS, T. A. LOVE, Radiochem. Radioanal. Letters 39(1979) 107 [Gamma-ray emission probabilities]
- H. HOUTERMANS, D. MILOSEVIC, F. REICHEL, Int. J. Appl. Radiat. Isotop. 31(1980)153 [Half-life]
- J. K. DICKENS, T. A. LOVE, Nucl. Instrum. Methods 175(1980)535 [Gamma-ray emission probabilities]
- Yang Chunxiang et al., Chin. J. Nucl. Phys. 2(1980)41 [Gamma-ray emission probabilities]
- R. G. HELMER, A. J. CAFFREY, R. J. GEHRKE, R. C. GREENWOOD, Nucl. Instrum. Methods 188(1981)671 [Gamma-ray energies]
- K. Yoshihara, A. Hibino, I. Yamoto, H. Kaji, Radiochem. Radioanal. Lett. 48(1981)303 [KX-ray intensities]
- A. SIMONITS, J. Radioanal. Nucl. Chem. 67(1981) [Gamma-ray emission probabilities]
- V. N. Gerasimov et al., Yadern. Fiz. 34(1981)3 [Gamma ray energies]
- K. SINGH, H. S. SAHOTA, J. Phys. Soc. Jap. 51,12(1982)153 [Multipole mixing ratio]
- D. D. HOPPE, J. M. R. HUTCHINSON, F. J. SCHIMA, M. P. UNTERWE GER, NBS-SP-626(1982)85 [Half-life]
- Y. V. KHOLNOV, Handbook, Energoizdat, Moscou (1982) [Beta emission probabilities]
- K. F. WALZ, K. DEBERTIN, H. SCHRADER, Appl. Rad. Isot. 34(1983)1191 [Half-life]
- W. Bambynek, X-84 Proc. X-Ray and Inner-Shell Processes in Atoms, Molecules and Solids, A. Meisel Ed., Leipzig Aug. 20-23(1984) [K fluorescence yield]
- CHEN DA, ZHANG QIAOLIAN, CHANG YONGFU, IEEE-Transactions on Nuclear Science, Vol.32- n°1 (1985)153 [Gamma-ray emission probabilities]
- R. A. MEYER, Fizika 22(1990)153 [Gamma-ray energies, Gamma-ray emission probabilities]
- J. GOSWAMY, B. CHAND, D. MEHTA, N. SINGH, P. N. TREHAN, Appl. Rad. Isotopes 43(1992)1467 [Gamma-ray emission probabilities]
- I. ALFTER, E. BODENSTEDT, B. HAMER, W. KNICHEL, R. MÜSSELER, R. SAJOK, T. SCHAEFER, J. SCHÜTH, R. VIANDEN, Z. Physik A347(1993)1 [Level half-life]
- L. K. PEKER, Nucl. Data Sheets 73,1(1994)1 [Spin and Parity]
- J. H. Hubbell, P. N. Trehan, Nirmal Singh, B. Chand, D. Mehta, M. L. Garg, R. R. Garg, Surinder Singh, S. J. Puri, Phys. Chem. Ref. Data 23-2(1994)339 [M fluorescence yield]
- G. AUDI, A. H. Wapstra, Nucl. Phys. A595(1995)409 [Q]
- E. SCHÖNFELD, H. Janssen, Nucl. Instrum. Methods A369(1996)527 [L fluorescence yield]
- V. S. BUTSEV, G. L. BUTSEVA, D. CHULTEM, P. I. GOLUBEV, D. KISS, E. J. LANGROCK, R. A. ZULKARNEV, Heavy Ion Physics 8(1998)227 [Half-life]
- M.-M. BÉ, E. Browne, V. Chechev, R. Helmer, E. Schönfeld, Table de Radionucléides, ISBN 2 7272 0200 8, CEA, F-91191 Gif sur Yvette ; (1999) [T ICC (Cs-137)]
- R. G. HELMER, C. van der Leun, Nucl. Instrum. Methods A450(2000)35 [Gamma ray energies]
- H. SCHRADER, Appl. Rad. Isotopes, 60(2004)317 [Half-life]

⁹⁹Tc - Comments on evaluation of decay data by X. Mougeot

This evaluation was done in 2010, taking into account the available literature by March 2010.

1 Decay Scheme

The decay scheme is complete since all of the levels in ⁹⁹Ru below the decay energies are populated. The J^π of the ground and excited levels are from the evaluation of Muller et al. (1986Mu09).

2 Nuclear Data

The Q value is from 2003AU03: $Q(\beta^-) = 293,8 (14)$ keV. Measurements of the end-point of the main β transition are given in Table 1.

Table 1: Measured end-points of the main β transition.

Reference	E_{\max} (keV)	Uncertainty (keV)
1947MO15	320	-
1950KE02	300	10
1951TA05	290	4
1952FE16	290	4
1960BO08	290	10
1966SN02	294	4
1974RE11	293	2

The evaluation of the ⁹⁹Tc half-life is described in the next section. Table 2 summarizes the measurements and their methodology.

Although the β -decay of ⁹⁹Tc is practically 100 % from its $9/2^+$ ground state to the $5/2^+$ ground state of ⁹⁹Ru, 1973LE10 and 1974EN02 observed a β -decay with a very small intensity to the $3/2^+$ first excited state of ⁹⁹Ru. Thus, its energy, half-life and the multipolarity of the de-exciting γ -ray are evaluated. The β -branching ratio is evaluated next.

An analysis of the published form factors of the main β^- transition is presented here. With the described limitations, an evaluation was carried out and the mean energy of the spectrum was calculated.

2.1 ⁹⁹Tc half-life

The measured half-life values of ⁹⁹Tc are given in Table 2 together with the experimental methods that were used. The value from 1947MO15 was not used in the evaluation because an experimental uncertainty was not reported. The value from 1960BO08 is a more recent one from the same authors.

Table 2: ⁹⁹Tc half-life measurements.

Reference	$T_{1/2}$ ($\times 10^5$ a)	Uncertainty ($\times 10^5$ a)	Method	Observations
1947MO15	9,4	-	Aluminium absorption	Not used: no uncertainty
1951FR05	2,12	0,04	Aluminium absorption	
1960BO08	2,15	0,05	Aluminium absorption	Same authors as 1947MO15
1966GO10	2,14	0,05	Liquid Scintillation Counting	
1984CO30	2,111	0,012	Liquid Scintillation Counting	

The statistical processing was done using the LWEIGHT program. A weighted average was adopted here from the resulting consistent data set, with a reduced- χ^2 value of 0,29. The statistical weight is 82 % for 1984CO30, the most recent and precise measurement. Finally, the adopted value, with its internal uncertainty, is:

$$T_{1/2} = 2,115 (11) \times 10^5 \text{ a.}$$

2.2 γ transition : first excited state of ⁹⁹Ru

2.2.1 Energy

The measured energies of the first excited state of ⁹⁹Ru are given in Table 3 with the experimental methods used.

Table 3: Measurements of the energy of the first excited state of ⁹⁹Ru.

Reference	Energy (keV)	Uncertainty (keV)	Method
1967MO20	89,36	0,40	⁹⁹ Rh decay, γ Ge(Li)
1970AN12	89,6	0,5	⁹⁹ Rh, γ Ge(Li)
1971LE20	89,4	1,0	⁹⁸ Mo(α ,3n) ⁹⁹ Ru, γ Ge(Li)
1973LE10	89,7	0,4	⁹⁹ Tc decay, β Si(Li)
1974EN02	89,5	0,2	⁹⁹ Tc decay, γ Si(Li)

The statistical processing was done using the LWEIGHT program. A weighted average was adopted from the resulting consistent data set, with a reduced- χ^2 value of 0,10. The statistical weight is 59 % for 1974EN02, the most precise measurement. Finally, this evaluation gives:

$$E_{\gamma}({}^{99}\text{Ru}) = 89,52 (15) \text{ keV.}$$

2.2.2 $T_{1/2}({}^{99}\text{Ru}, 89 \text{ keV})$

The measured half-life values of the first excited state of ⁹⁹Ru are given in Table 4 together with the experimental methods used. The original uncertainty of 1972GU01, not explained in detail in the article, seems to be underestimated. 1973BE72 used the same method, with nearly the same statistics, and reported an uncertainty of 0,6. The uncertainty of 0,1 given by 1972GU01 seems to be only that from the data fitting. Thus, the evaluator decided to increase the uncertainty of 1972GU01 from 0,1 to 0,6.

Table 4: Measurements of the half-life of the first excited state of ⁹⁹Ru.

Reference	$T_{1/2}$ (ns)	Uncertainty (ns)	Method	Observation
1964BO28	19,7	0,4	⁹⁹ Rh, γ spectro.	Uncertainty increased from 0,1 to 0,6
1965KI01	20,0	1,0	⁹⁹ Rh, γ spectro.	
1965MA27	20,7	0,3	⁹⁹ Rh, γ spectro.	
1972GU01	20,5	0,6	⁹⁹ Rh, γ Ge(Li)	
1973BE72	21,04	0,6	⁹⁹ Rh, γ Ge(Li)	
1974EN02	18,9	1,0	⁹⁹ Tc decay, γ Si(Li)	

The statistical processing was done using the LWEIGHT program. A weighted average was adopted here from the resulting consistent data set, with a reduced- χ^2 value of 1,52. The statistical weight is 45 % for 1965MA27. These authors gave some details on their estimation of the uncertainty, and there is no reason to believe it was underestimated. Finally, this evaluation gives:

$$T_{1/2}({}^{99}\text{Ru}, 89 \text{ keV}) = 20,36 (25) \text{ ns.}$$

2.2.3 Multipolarity

The γ transition from the $3/2^+$ first excited state to the $5/2^+$ ground state of ⁹⁹Ru is M1+E2. Measurements were carried out to obtain the mixing ratio $\delta^2 = E2/M1$. They are summarized in Table 5 with the experimental methods used. Only two measurements were used for the evaluation because most of the publications are from the same author. Only the most recent one, which is also the most precise, was included. The value from 1973BE72 was not used because the experimental uncertainty was not reported.

The statistical processing was done using the LWEIGHT program. A weighted average was adopted here from the resulting consistent data set, with a reduced- χ^2 value of 2,69. The statistical weight is 94 % for 1976KI02, the most precise measurement. Finally, this evaluation gives:

$$\delta^2({}^{99}\text{Ru}, 89 \text{ keV}) = 2,45 (6).$$

Then $\delta = -1,56 (2)$, and the multipolarity is:

$$\text{M1} + 71,0 (5) \% \text{ E2.}$$

Table 5: Measurements of the multipolarity mixing ratio of the first excited state of ⁹⁹Ru. The values from 1973Gibb and 1976KI02 were the only ones that were used for the evaluation.

Reference	$\delta^2 = E2/M1$	Uncertainty	Method	Observation
1964KI01	~ 2	-	Ru-99 Mössbauer γ transition	Not used: no uncertainty
1965KI01	2,4	0,9	Ru-99 Mössbauer γ transition	Same author as 1964KI01
1966KI02	2,7	0,6	Ru-99 Mössbauer γ transition	$\delta < 0$, same author as 1964KI01
1972Wagner	2,7	0,6	Ru-99 Mössbauer γ transition	Coming from 1966KI02
1973BE72	2,57	-	⁹⁹ Rh, γ Ge(Li)	Not used: no uncertainty
1973Gibb	2,72	0,17	Ru-99 Mössbauer γ transition	
1976KI02	2,43	0,04	Ru-99 Mössbauer γ transition	$\delta = -1,56 (2)$, same author as 1964KI01

2.2.4 Branching ratio

1973LE10 and 1974EN02 inferred a small β^- transition from the $9/2^+$ ground state of ⁹⁹Tc to the $3/2^+$ 89 keV level of ⁹⁹Ru, by detecting a de-exciting γ -ray. Thus, this β^- transition is second unique forbidden, whereas the main transition is second non-unique forbidden.

The authors reported the number of photons detected per decay: $6,5 (1,5) \times 10^{-6}$ for 1973LE10 and $4,9 (1,7) \times 10^{-6}$ for 1974EN02. Next, they used a total internal conversion coefficient $\alpha_T = 1,5$ calculated from 1968HA52 to determine the corresponding total γ -ray transition probability, and thus the β^- branching.

The absolute γ -ray intensity was evaluated. The statistical processing was done using the LWEIGHT program. A weighted average was adopted from the resulting consistent data set, with a reduced- χ^2 value of 0,50. The statistical weight is 56 % for 1973LE10. Finally, this evaluation gives:

$$I_{\text{abs}}(^{99}\text{Ru}, 89 \text{ keV}) = 5,8 (11) \times 10^{-4} \text{ \%}$$

The total conversion coefficient α_T was calculated using the BrIcc program (2008KI07): $\alpha_T = 1,495 (25)$. Thus the β^- branching is equal to $I_{\text{abs}}(1 + \alpha_T)$. Finally, this evaluation gives:

$$P_{\beta 0,1} = 1,45 (30) \times 10^{-3} \text{ \%}$$

2.3 β^- transition

The branching of the main β^- transition is practically 100 %. A small contribution to the first excited state of ⁹⁹Ru exists, with a branching of $1,45 (30) \times 10^{-3} \text{ \%}$, as deduced in Section 2.2.4.

The main β^- transition is from the $9/2^+$ ground state of ⁹⁹Tc to the $5/2^+$ ground state of ⁹⁹Ru. This is a second forbidden non-unique transition, thus one can expect a form factor as given below (1976Behrens):

$$C(W) = A(W)q^2 + B(W)\lambda_2 p^2 + D(q^4 + 10/3\lambda_2 q^2 p^2 + \lambda_3 p^4),$$

where q is the linear momentum of the neutrino, p the linear momentum of the electron, and W is the normalised energy of the electron. Measurements show that the following form factor for a first unique forbidden transition gives a good description of the measured energy spectrum:

$$C(W) = q^2 + \lambda p^2.$$

The determination of the form factor is highly dependent on the calculated spectrum used for the comparison with experimental data. Consequently, the form factor depends on the hypothesis made and the data used for the calculation: Coulomb corrections, screening correction due to electron cloud, finite nuclear size correction, radiative corrections, end-point energy, and nature of the transition. The form factor is generally determined by a comparison with a calculated allowed spectrum.

Table 6: Measurements of the form factor of the main β^- transition.

Reference	λ	Uncertainty	E_{max} (keV)	Energy range (keV)	Method	Observation
1951TA05	~ 1	-	290 (4)	150 - end-point	Mag. spectro.	Not used: no uncertainty
1952FE16	0,50	0,13	292 (3)	60 - end-point	Mag. spectro.	Recalculated by 1966Lipnik
1966SN02	0,49	0,04	294 (4)	50 - 280	Plastic scint.	Recalculated by 1976Behrens
1974RE11	0,54	0,02	293 (2)	55 - 250	Si(Li)	

Experimental data are summarized in Table 6 with the energy range of their validity, and the experimental methods used. The value from 1951TA05 was not used because no experimental uncertainty was reported. The value from 1952FE16 was calculated by 1966Lipnik in the correct form. The value from 1966SN02 was recalculated by 1976Behrens using more recent tables for the Fermi function, leading to an increase of the uncertainty from 0,011 to 0,04. The statistical processing was done using the LWEIGHT

program. A weighted average was adopted from the resulting consistent data set, with a reduced- χ^2 value of 0,65. The statistical weight is 78 % for 1974RE11, the most recent and precise measurement. Finally, this evaluation gives:

$$C(W) = q^2 + 0,529 (18) p^2.$$

It should be underline the main difficulty of the evaluation of form factors: the authors of the published data did not describe in detail all the possible sources of distortion of the measured spectra and their contributions. Obviously, this is a difficult task. In some articles, it is clear that all these problems were not taken into account. The temptation could be great to adjust some known parameters within the uncertainty range to obtain a result close to the previous published ones.

Resulting from the violation parity, electrons emitted in nuclear β -decay are longitudinally spin-polarized. If the decaying neutron is influenced by the nuclear structure in which it is embedded, the value of the polarization may be altered. The authors of 1990GA13 measured the longitudinal electron polarization, and they suggested that the decaying neutron is not influenced more than 3,3 % by the nuclear structure. This could explain why a form factor of a first unique forbidden transition is sufficient to describe a second non-unique forbidden transition. One can note the usual approximation in the theoretical calculation of β spectra: a non-unique forbidden transition is treated as a unique forbidden transition with the same variation of the total angular momentum. It means that a second non-unique forbidden transition is treated as a first unique forbidden transition.

The mean energy of the β spectrum was calculated with the Q value and the form factor given previously. The calculation is based on the analytical approach developed by N.B. Gove and M.J. Martin (1971GO40) and it includes the following correction terms: Coulomb corrections (1961RO33), screening correction due to electron cloud (1954GO69), finite nuclear size correction (1980Dillman), and radiative corrections (1982Behrens). The uncertainty is estimated by the product of E_{mean} with the uncertainty on λ . The result is:

$$E_{\text{mean}} = 94,6 (17) \text{ keV}.$$

The log ft value for the main transition (second non-unique forbidden) has been calculated with the LOGFT program: $\log ft (^{99}\text{Tc} \rightarrow ^{99}\text{Ru}^{\text{gs}}) = 12,323 (7)$. In the same way, for the second unique forbidden transition to the first excited state of ^{99}Ru : $\log ft (^{99}\text{Tc} \rightarrow ^{99}\text{Ru}^*) = 15,82 (9)$.

For the sake of completeness, we mention some publications on K-shell auto-ionization probabilities accompanying the β decay of ^{99}Tc : 1967ST36, 1972WA32, 1974HA12, and 1980LA02, for experimental studies, and 1977IS05, for theoretical studies. The emitted β particle can ionize the electron cloud of the daughter nucleus, ^{99}Ru , distorting the β spectrum. This phenomenon is negligible in almost all applications, since its probability is about 0,05 % per emitted β .

3 Atomic Data (Ru, Z=44)

3.1 X Radiations and Auger electrons

The X-ray and Auger electron data were computed using the EMISSION program with the atomic data of Schönfeld and Janßen (1996SC06).

4 Radiation Emissions

4.1 Electron Emission

The β^- intensities were evaluated as described above in Section 2.

4.2 Photon Emission

The details of the photon emission evaluation are in Section 2. ^{99}Ru decays from its first excited state at 89,52 (15) keV, with a half-life of 20,36 (25) ns, and a γ -ray multipolarity of $M1 + 71,0 (5) \% E2$. The absolute γ -ray emission intensity is evaluated as 5,8 (11) $\times 10^{-4} \%$, leading to a β^- branching to $^{99}\text{Ru}^*(89 \text{ keV})$ of $1,45 (30) \times 10^{-3} \%$.

5. References

- 1947MO15 E.E. Motta, G.E. Boyd, Q.V. Larson, Phys. Rev. 72, 1270 (1947) [E_{\max} , $T_{1/2}$]
- 1950KE02 B.H. Ketelle, J.W. Ruch, Phys. Rev. 77, 565 (1950) [E_{\max}]
- 1951FR05 S. Fried, A.H. Jaffey, N.F. Hall, L.E. Glendenin, Phys. Rev. 81, 741 (1951) [$T_{1/2}$]
- 1951TA05 S.I. Taimuty, Phys. Rev. 81, 461 (1951) [E_{\max} , Form Factor]
- 1952FE16 L. Feldman, C.S. Wu, Phys. Rev. 87, 1091 (1952) [E_{\max} , Form Factor]
- 1954GO69 R.H. Good, Jr., R.H. Good, Phys. Rev. 94, 931 (1954) [β spectrum calculation]
- 1960BO08 G.E. Boyd, Q.V. Larson, E.E. Motta, J. Am. Chem. Soc. 82, 809 (1960) [E_{\max} , $T_{1/2}$]
- 1961RO33 M.E. Rose, Relativistic Electron Theory, John Wiley and Sons, Inc., New York (1961) [β spectrum calculation]
- 1964BO28 E. Bodenstedt, C. Gunther, J. Radloff, W. Engels, W. Delang, M. Forker, H. Luig, Phys. Lett. 13, 330 (1964) [$T_{1/2}$ γ]
- 1964KI01 O.C. Kistner, R. Segnan, Bull. Am. Phys. Soc. 9, No.4, 396, BC13 (1964) [Multipolarity]
- 1965KI01 O.C. Kistner, S. Monaro, A. Schwarzschild, Phys. Rev. 137, B23 (1965) [$T_{1/2}$ γ , Multipolarity]
- 1965MA27 E. Matthias, S.S. Rosenblum, D.A. Shirley, Phys. Rev. 139, B532 (1965) [$T_{1/2}$ γ]
- 1966GO10 G. Goldstein, J.A. Dean, J. Inorg. Nucl. Chem. 28, 285 (1966) [$T_{1/2}$]
- 1966KI02 O.C. Kistner, Phys. Rev. 144, 1022 (1966) [Multipolarity]
- 1966Lipnik P. Lipnik, J.W. Sunier, Phys. Rev. 145, 746 (1966) [Form Factor]
- 1966SN02 R.E. Snyder, G.B. Beard, Phys. Rev. 147, 867 (1966) [E_{\max} , Form Factor]
- 1967MO20 G.A. Moss, D.K. McDaniels, Phys. Rev. 162, 1087 (1967) [E_{γ}]
- 1967ST36 P. Stephas, B. Crasemann, Phys. Rev. 164, 1509 (1967) [K-shell autoionization]
- 1968HA52 R.S. Hager, E.C. Seltzer, Nucl. Data A 4, 1 (1968) [α_T]
- 1970AN12 N.M. Antoneva, E.P. Grigorev, L.F. Protasova, Bull. Acad. Sci. USSR, Phys. Ser. 34, 771 (1971) [E_{γ}]
- 1971GO40 N.B. Gove, M.J. Martin, Nucl. Data Tables A 10, 205 (1971) [β spectrum calculation]
- 1971LE20 C.M. Lederer, J.M. Jaklevic, J.M. Hollander, Nucl. Phys. A 169, 489 (1971) [E_{γ}]
- 1972GU01 D.K. Gupta, C. Rangacharyulu, R. Singh, G.N. Rao, Nucl. Phys. A 180, 311 (1972) [$T_{1/2}$ γ]
- 1972Wagner F.E. Wagner, B.D. Dunlap, G.M. Malvius, H. Schaller, R. Felscher, H. Spieler, Phys. Rev. Lett. 28, 530 (1972) [Multipolarity]
- 1972WA32 R.L. Watson, E.T. Chulick, R.W. Howard, Phys. Rev. C 6, 2189 (1972) [K-shell autoionization]
- 1973BE72 R.B. Begzhanov, D.A. Gladyshev, K.S. Azimov, M. Narzikulov, K.T. Teshabaev, Russ. Phys. J. 16, 9 (1973) 1258 [$T_{1/2}$ γ , Multipolarity]
- 1973Gibb T.C. Gibb, R. Greatrex, N.N. Greenwood, P. Kaspi, J. Chem. Soc., Dalton Trans., 1253 (1973) [Multipolarity]
- 1973LE10 J. Legrand, J. Morel, Phys. Rev. C 8, 366 (1973) [Branching Ratio, E_{γ} , I_{abs} γ]
- 1974EN02 C.E. Engelke, J.D. Ullman, Phys. Rev. C 9, 2358 (1974) [Branching Ratio, E_{γ} , $T_{1/2}$ γ , I_{abs} γ]
- 1974HA12 H.H. Hansen, K. Parthasaradhi, Phys. Rev. C 9, 1143 (1974) [K-shell autoionization]
- 1974RE11 M. Reich, H.M. Schupferling, Z. Phys. 271, 107 (1974) [E_{\max} , Form Factor]
- 1976Behrens H. Behrens, L. Szybisz, ZAED Phys. Data, 6-1 (1976) [Form Factor]
- 1976KI02 O.C. Kistner, A.H. Lumpkin, Phys. Rev. C 13, 1132 (1976) [Multipolarity]
- 1977IS05 Y. Isozumi, S. Shimizu, T. Mukoyama, Nuovo Cim. 41 A, 359 (1977) [K-shell autoionization]
- 1980Dillman L.T. Dillman, EDISTR, ORNL/TM-6689 (1980) [β spectrum calculation]
- 1980LA02 C.E. Laird, P.C. Hummel, H.-C. Liu, Phys. Rev. C 21, 723 (1980) [K-shell autoionization]
- 1982Behrens H. Behrens, W. Bühring, Electron radial wave functions and nuclear beta-decay, Oxford Science Publications (1982) [β spectrum calculation]
- 1984CO30 B.M. Coursey, J.A.B. Gibson, M.W. Heitzmann, J.C. Leak, Int. J. Appl. Radiat. Isotop. 35, 1103 (1984) [$T_{1/2}$]
- 1986MU09 H.-W. Muller, D. Chmielewska, Nucl. Data Sheets 48, 663 (1986) [J^{π}]
- 1990GA13 R. Gauder, S. Fuchs, A. Hilscher, K.-W. Hoffmann, E. Lehmann, R. Sadler, Z. Phys. A 336, 53 (1990) [Longitudinal electron polarization]
- 1996SC06 E. Schönfeld, H. Janßen, Nucl. Instr. Meth. **A369** (1996) 527 [ω_K , K x ray ratios, Auger e-ratios, atomic data]
- 2003AU03 G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A 729, 337 (2003) [Q]
- 2008KI07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr. Nucl. Instr. and Meth. A589 (2008) 202-229 [BrICC]

⁹⁹Tc^m - Comments on evaluation of decay data
by C. Morillon*, M. M. Bé*, V. Chechev, A. Egorov****

This evaluation was completed in December 2000. The half-life has been updated in January 2004.

1. DECAY SCHEME

Tc-99m mainly decays to the ground level of Tc-99.

Very weak beta minus transitions to the ground and two excited levels of Ru-99 have been observed. The J^π values and the level energies are from Peker(1994Pe15).

2. NUCLEAR DATA

Q_{IT} (⁹⁹Tc^m) from the 142,7 keV level energy
 Q (⁹⁹Tc^m) from Audi and Wapstra (1995)

2.1 HALF-LIFE

- The measured half-life values are, in hours:

1	6,13(5)	CROWTHER and ELDRIDG	(1965)	1965Cr03	rejected
2	6,006(7)	GOODIER and WILLIAMS	(1966)	1966Go22	
3	6,014 (4)	VUORINEN	(1969)	1969Vu03	
4	6,031 (12)	LEGRAND et al.	(1970)	1970Le07	
5	6,007 (2)	SANTRY and BOWES	(1989)	1989Sa**	
6	6,03 (13)	DECOMBAZ et al.	(1972)	1972De76	
7	6,02 (1)	EMERY et al.	(1972)	1972Em01	
8	6,049 (35)	EMERY et al.	(1972)	1972Em01	rejected
9	6,02 (3)	MEYER	(1974)	1974Me**	
10	6,008 (4)	RUTLEDGE et al.	(1980)	1980RuZY	TcO ₄ Na
11	6,006 (2)	HOUTERMANS et al.	(1980)	1980Ho17	No precision
12	6,0072 (10)	AYRES and HIRSHFELD	(1982)	1982Ay**	Normal saline solution
13	6,0170(19)	AYRES and HIRSHFELD	(1982)	1982Ay**	Acid solution
	6,0062 (7)	WALZ et al.	(1983)	1983Wa26	Superseded by 2003Sc49
14	6,020(2)	KOLTSOV et al.	(1998)	1998Ko**	TcO ₄ Na
15	6,0058(12)	SCHRADER	(2004)	2004Sc49	TcO ₄ Na
16	6,0071(21)	Da SILVA et al.	(2004)	2004Si04	TcO ₄ Na

The chemical medium probably has an influence on the half -life. Changes in the half -life values have been observed with the modification of external environment or chemical composition (influence on internal conversion of electrons of 2,17 keV transition in external shells : Mazaki (1980Ma03) , Koltsov, and others).

Comparisons of the decay constant of Tc -99m in different chemical environments were made. In the following table λ₀ is the decay constant for Tc-99m in the form of pertechnetate (TcO₄).

Author	Type of source	Source pair	Relative variation of decay constant, %
Koltsov	Sulfide	$[\lambda_0 - \lambda (\text{Tc}_2\text{S}_7)] / \lambda_0$	0,14 (8)
Koltsov	Silver	$[\lambda_0 - \lambda (\text{Ag})] / \lambda_0$	0,35 (7)
Koltsov	Gold	$[\lambda_0 - \lambda (\text{Au})] / \lambda_0$	0,25 (7)
Mazaki	Sulfide	$[\lambda_0 - \lambda (\text{Tc}_2\text{S}_7)] / \lambda_0$	0,32 (7)
Mazaki	Sulfide - metal	$[\lambda (\text{Tc}_2\text{S}_7) - \lambda (\text{Metal})] / \lambda (\text{Metal})$	0,056 (3)
Ayres		Acid solution – Normal saline	0,16

If we consider the set of 16 measured values given in the table above, where :

- Emery *et al.* (1972) and Ayres and Hirshfeld (1982) measured the half -life of Tc-99m by 2 different methods or 2 media: both values were taken into account. (NB : the experiment and results described by Ayres and Hirshfeld are the same as those described by Hoppes *et al.* in NBS-SP-626 (1982) 85 and by Unterweger *et al.* in NIM A312 (1992) 349) ;
- the value of Crowther and Eldridge (1965) and the second value of Emery *et al.* (1972) are rejected due to the Chauvenet criterion.

With the set of 14 remaining values, LWEIGHT recommended the unweighted average (Reduced $\chi^2 = 5,3$) and expanded the uncertainty to include the most precise value of 6,0072 (Ayres *et al.* 1983). This leads to 6,014 (7) h.

With the 7 most recent values (from 10 to 16) (>1980), the LWEIGHT program derived the weighted mean and expanded the uncertainty: the recommended value is 6,0089 (19) h. (Reduced- $\chi^2 = 10,2$).

Nevertheless, the most commonly used chemical composition is sodium pertechnetate (TcO_4Na) in a physiological saline solution, this solution is chemically stable. This is the result of the way of production of ⁹⁹Tc^m for medical purposes. The metallic matrix have been made for very specific studies and do not correspond to a general use.

Then, taking into consideration the most recent values obtained from a (TcO_4Na) solution, i.e. values 10 – 12 – 14 – 15 – 16 ; the value 14 (Koltsov) is outlier, omitting it the weighted mean is 6,006 7 (7) with the internal uncertainty, the reduced χ^2 is 0,32.

Conclusions :

- Due to the fact that the pertechnetate solution is a stable solution and the most commonly used, *the adopted half-life is : 6,006 7 (10) h*, uncertainty of the most precise measurement value.
- Uncertainty should be enlarged to 0,009, to take into account a possible chemical effect of 0,15% for other solutions, then the half life would be : 6,007 (9) h.

DECAY Tc-99m to Tc-99

- Measured half-life of the 140,5 keV level in ns:

0,277 (14)	STEINER <i>et al.</i> (1969St04)
0,160 (20)	MCDONALD <i>et al.</i> (1971Mc02)
0,205 (4)	ALFTER <i>et al.</i> (1993A123)
0,237 (14)	SHENOY <i>et al.</i> (1973Sh21)

The value of Steiner *et al.*(1969) is from the original article; the NDS value from the same reference has been adjusted to 0,192 ns.

The value of 0,160(20) ns from McDonald *et al.* (1971) deviates far from the others and is not taken into account.

The Steiner *et al.*(1969) and Shenoy *et al.*(1973) values were determined using the Mössbauer effect. The uncertainty in the Alfter *et al.* (1993) value was increased 2,47 times by LWEIGHT.

Reduced- $\chi^2 = 8,94$

LWEIGHT has used the weighted average and the external uncertainty.

The adopted value is : **0,221(20) ns**

· **Level energy of technetium 99**

The values of the level energies are from NDS 73,1

Level 2 142,6833 (11)

Level 1 140,5108 (10)

2.2 GAMMA TRANSITIONS and INTERNAL CONVERSION COEFFICIENTS

The energies of the gamma transitions are derived from the energies of the gamma rays, taking recoil into account. The probabilities of gamma transitions $P_{\gamma+ce}$ have been computed using the evaluated absolute gamma-ray emission probabilities and the total internal conversion coefficients (ICC).

INTERNAL CONVERSION COEFFICIENTS

The ICC have been evaluated using the experimental information of the multipolarity admixture coefficients and the theoretical values from Röseler *et al.* and Band *et al.* (for $\gamma_{2,1}$ 2,17 keV).

For pure multiplicities the uncertainties on the ICC values are adopted to be 2%. For mixed multiplicities the uncertainties of ICC were evaluated taking into account the uncertainties of respective multipolarity admixture coefficients.

The ICC adopted values are compared with the measured values, and are, generally, in good agreement.

Transition 2-1: 2,17 keV

No experimental value has been found. Band theoretical values (1976Ba63):

$\alpha_T = 1,35 (4) 10^{10}$ and $\alpha_M = 1,19 (3) 10^{10}$

Transition 1-0: 140,511 keV

Total Internal Conversion Coefficient α_T

Experimental measurements :

0,118 (8)	AMTEY <i>et al.</i> (1966)
0,113 (6)	DICKENS and LOVE (1980)
0,122 (5)	VUORINEN (1969)
0,118 (3)	LEGRAND <i>et al.</i> (1973Le29)
0,1181(23)	LWEIGHT (reduced $\chi^2 = 0,44$; weighted average and internal uncertainty)
Adopted: 0,119(3)	Rosel <i>et al.</i> for M1+3,3(3)%E2

Dickens and Love (1980) determined α_T from the α_k value given by Gardulski and Wiedenbeck (1974) and the K/L/MN values reported by Hager and Selzer and by Medsker (NDS 12-4 - 1974)

α_T was evaluated by Vuorinen (1969) from measurements of conversion electrons in coincidence with fluorescence X-rays.

Multipolarity

Large number of measurements have been made. However, most of the authors gave different values without, or with a large uncertainty. These multiplicities permit the calculation of the total internal conversion coefficients, to which we have assigned a 5% uncertainty:

/d/	Transition	a_T (Rösel)	
0,31 (2)	M1 + 8,25% E2	0,132(7)	SINGH and SAHOTA (1982Si16)
0,178 (12)	M1 + 3,1% E2	0,119(6)	ALFTER (1993Al23)
	M1 + 4%(2) E2	0,121(6)	MCDONALD <i>et al.</i> (1971Mc02)
	M1+<3%E2		VOINOVA <i>et al.</i> (1972Vo06)
0,194(30)	M1+3,8%E2		VUORINEN (1969Vu03)
	M1+<8%E2		VAN EIJK <i>et al.</i> (1968Va14) calculated from ICCk
	M1+9%(5)E2	0,134(7)	VAN EIJK <i>et al.</i> (1968) calculated from K/L ratio
	M1+2,8%E2	0,118(6)	COOK <i>et al.</i> (1969Co18)
	M1+7(3)%E2	0,129(7)	MEYER (1974)
	M1+1,4%E2	0,114(6)	DICKENS and LOVE (1980Di16)
	M1+6,5(40)E2	0,128(7)	AGEEV <i>et al.</i> (1969Ag04)
0,118(6)	M1+1,4(2)%E2	0,114(6)	GARDULSKI and WIEDENBECK (1974Ga01)
	M1+2,8(3)%E2	0,118(6)	GEIGER (1968GeZW)
	M1+9%E2		SIMONITS <i>et al.</i> (1981Si15)
	M1+E2		AMTEY <i>et al.</i> (1966Am04)
	M1		BASHANDY (1969Ba54)
		0,120(2)	LWEIGHT (reduced- $\chi^2= 1,16$), weighted average and external uncertainty= 0,002
0,186 (8)	M1+ 3,2(3)%E2	0,119(3)	Adopted (Rösel <i>et al.</i>)

From each determination of the multipolarity of the transition, the Rösel theoretical internal coefficient was calculated. From the set of the 10 deduced ICC values the LWEIGHT program recommends a weighted mean of 0,120(2). The value is very closed to that obtained by considering the 4 experimental values for α_T (see table above).

Internal Conversion Coefficients a_K

Experimental values:

0,096(6)	VOINOVA <i>et al.</i> (1971Vo06)
0,093 (6)	VOINOVA <i>et al.</i> (1971Vo06)
0,102 (7)	VAN EIJK <i>et al.</i> (1968Va14)
0,094 (8)	VUORINEN (1969Vu03)
0,102 (5)	DICKENS and LOVE (1980Di16)
0,096 (3)	LWEIGHT ($\chi^2=0,35$; weighted average and internal uncertainty)
0,104 (3)	Rösel <i>et al.</i> (1978) (adopted)

- α_K was measured by Voinova *et al.* (1971) with a spectrometer which provided simultaneous measurement of conversion electrons and γ -ray spectra.

- Van Eijk *et al.*(1968) calculated α_K from measurements of the 140,5 keV gamma -ray emission probability (P_γ) relative to the gamma -ray emission probability of the 661,6 keV gamma transition in the decay of Cs-137, and from measurements of the conversion electron emission probability P_{ce} of the 140,5 keV K-conversion line relative to the conversion electron emission probability of the 661,6 keV K-conversion line in the decay of Cs-137: $P_{ceK} = 6,84(19)$; $P_\gamma = 6,00(35)$; $\alpha_K(661,6 \text{ keV}) = 0,0896 (15)$ (Helmer in 1999BeZQ).
- Vuorinen (1969) evaluated the internal conversion coefficient α_K by measuring the electron conversion emissions following the conversion of the 140 keV gamma -ray in coincidence with fluorescence X-rays.
- α_K given by Dickens and Love (1980) was computed from the tables of Hager and Seltzer for a M1 transition and a 1,4% E2 admixture. An 5% uncertainty assigned to α_K reflects the added uncertainty to the usual 3% due to the rapid change of α_K with admixture. This value is not taken into account in our calculations.

Internal Conversion Coefficients α_L

α_L can be deduced from measurements of the K/L ratio of the conversion electron emission probabilities, and with $\alpha_K = 0,104(3)$:

K/L	α_L	
8,1 (5)	0,0125(8)	BASHANDY(1969Ba03)
7,70 (30)	0,0132(7)	VAN EIJK <i>et al.</i> (1968Va14)
8,3 (3)	0,0122(6)	RAVIER <i>et al.</i> (1961Ra04)
7,63 (32)	0,0133(7)	BRAHMAVAR (1968)
7,8 (3)	0,0130(6)	GEIGER (1968 GeZW)
	0,0128(3)	LWEIGHT has used the weighted average and the internal uncertainty. Reduced- $\chi^2 = 0,52$
Adopted	0,0129(4)	Rösel <i>et al.</i> (1978)

Transition 2-0: 142,683 keV

Internal Conversion Coefficients α_T

For a M4 transition the theoretical value from Rösel is : **40,9(8)**.

Internal Conversion Coefficients α_K

- The two following values were calculated from experimental data, and listed by the authors:
 23 (6) VAN EIJK *et al.* (1968)
 30 (3) BASHANDY (1969Ba54)

Van Eijk *et al.* (1968) calculated the K ICC from the ratios of $K(142,7)/K(140,5) = 0,072(32)$ and $I_\gamma(142,7)/I_\gamma(140,5) = 0,00030(6)$, after correction for $\alpha_K(661,6 \text{ keV, Cs-137}) = 0,0896(15)$

Bashandy (1969) calculated the K ICC from internal conversion spectra and photon emission probabilities $I_\gamma(142)/I_\gamma(140) = 0,00030(6)$

- The following α_K coefficients are calculated from the $K(142,7)/K(140,5)$ ratio given by the authors, based on the ratio $I_\gamma(142,7)/I_\gamma(140,5) = 0,00030(6)$ [Van Eijk (1968)] and $\alpha_K(140,5) = 0,104(3)$.

$K(142,7)/K(140,5)$	$\alpha_K(142,7)$	
0,072(4)	24 (6)	AMTEY (1966Am04)
0,0746(12)	25 (6)	GEIGER (1968GeZW)
0,075 (8)	26 (6)	AGEEV <i>et al.</i> (1969Ag04)

If we take into account the ratio $I_{\gamma}(142,7)/I_{\gamma}(140,5) = 0,00021(3)$ given by Dickens and Love (1980Di16), with $\alpha_K(140,5) = 0,104(3)$ the same calculations give higher results for $\alpha_K(142,7)$:

K(142,7)/K(140,5)	$\alpha_K(142,7)$	
0,072(4)	34 (6)	AMTEY (1966)
0,0746 (12)	36 (5)	GEIGER (1968)
0,075 (8)	36 (7)	AGEEV <i>et al.</i> (1969)

If we take into account all the six possible data, the weighted average, with the external uncertainty, calculated by LWEIGHT is 29,5(18) (reduced- $\chi^2 = 0,87$)

The **adopted** theoretical K conversion coefficient, for a M4 transition, is : **29,3(6)** (Rösel *et al.* (1978)).

Internal Conversion Coefficients α_L

From the measurement of the ratio of the conversion electron intensities, with $\alpha_K = 29,3(6)$, it can be deduced that α_L (BASHANDY and IBRAHIEM) is closed to the adopted theoretical value:

K/L	α_L		
2,9 (5)	10,1 (18)	M4 transition	BASHANDY and IBRAHIEM (1969Ba03)
Adopted:	9,35 (20)	M4 transition	RÖSEL <i>et al.</i> (1978)

3. ATOMIC DATA

3.1. FLUORESCENCE YIELDS

The fluorescence yields are taken from Schönfeld and Janßen (96Sc06).

3.2. X RADIATIONS

The X-ray energies are based on the wavelengths given by Bearden and were converted into energy with $1\text{Å} = 1,00001481(92) \cdot 10^{-10}\text{m}$

The emission intensities are calculated with the EMISSION program from PTB. No experimental data have been found.

3.3. AUGER ELECTRONS

The energies of Auger electrons are from 1977La** (Larkins).

The ratios P(KLX)/P(KLL) and P(KLY)/P(KLL) are taken from 1996Sc06.

4. PHOTON EMISSIONS

4.1. X-RAY EMISSIONS

The absolute emission probabilities of K X-rays (P_{XK}) have been computed using the adopted value of ω_K , the evaluated internal conversion coefficients and the emission probabilities.

4.2. GAMMA RAY EMISSIONS

4.2.1 GAMMA RAY ENERGIES

The γ -ray energies of $\gamma_{2,1}(2,17\text{ keV})$ and $\gamma_{1,0}(140,5\text{ keV})$ are taken from Gerasimov *et al.* (1981Ge05) and Helmer (2000He14), respectively. These values are based on the most accurate measurements with an

electrostatic spectrometer ($E_{\gamma_{2,1}}$, see also 1971La12 – Lacasse and Hamilton) and curved-crystal spectrometer ($E_{\gamma_{1,0}}$, see also 1981He15 – Helmer *et al.*). The energy of $\gamma_{2,0}$ (142,7 keV) has been computed as the sum of the adopted energies of $\gamma_{2,1}$ (2,17 keV) and $\gamma_{1,0}$ (140,5 keV).

4.2.2 GAMMA RAY EMISSION INTENSITIES

140,511 keV (1,0)

Absolute values (per 100 decays)

88,20 (26)	Chen Da (1985)
87,30(21)	Simonits <i>et al.</i> (1981Si15)
88,75 (14)	Rutledge <i>et al.</i> (1980Ru20)
87,2 (5)	Dickens and Love (1980Di16) (calculated)
88,0 (24)	Legrand <i>et al.</i> (1973Le29)

LWEIGHT has been used to derive the weighted average and expand the uncertainty so that the range includes the most precise value of 88,75(14). This leads to the average of 88,4(4) % (reduced- $\chi^2 = 2,24$). Omitting the calculated value of Dickens and Love (1980) and the value of Simonits (1981) from statistical considerations, we have a weighted average of 88,5 % with an external uncertainty of 0,2. LWEIGHT has increased the uncertainty of Rutledge *et al.* (1980) to 0,258. Reduced- $\chi^2 = 1,14$. The **adopted** value is : **88,5(2)%**

142,675 keV (2,0)

Relative measurements of the $\gamma_{1,0}$ (140,5 keV) line are not precise: from 0,00020(3) of Dickens *et al.*(1980) to 0,00030(6) of Van Eijk *et al.* (1969).

The ratio of $I_{\gamma+ce}(142,7)/I_{\gamma+ce}(140,5)$ from the ⁹⁹Mo+⁹⁹Tc^m evaluation for the “slow” component of the 140,5 keV transition is 0,0097(7), corresponding to $I_{\gamma}(142,7)/I_{\gamma}(140,5) = 0,00026(2)$ and $P_{\gamma}(142,7) = \mathbf{0,023(2)\%}$ (**adopted value**).

5. ELECTRON EMISSIONS

The energies of the conversion electrons have been calculated from the gamma-transition energies given in 2.2 and the electron binding energies. Emission probabilities have been calculated using the conversion coefficients given in 2.2. and the adopted gamma emission probabilities.

Measurements of conversion electron spectra for ⁹⁹Tc^m (in equilibrium with ⁹⁹Mo) have been made in many studies (Van Eijk -1968Va14, Ageev-1969Ag04, Bashandy-1969Ba03, Bashandy-1969Ba54, Ravier-1961Ra01, Lacasse-1971La12, Voinova-1971Vo06, Legrand-1973Le29, Gerasimov-1981Ge05). However, the computed values of the conversion electron energies and emission probabilities are more accurate.

The values of the emission probabilities of K-Auger electrons have been calculated using the transition probabilities given in 2.1 and 2.2, the atomic data given in 3. and the conversion coefficients given in 2.2.

Experimental Auger spectra can be found in 1981Ge05 (Gerasimov *et al.*).

Tc-99m to Ru-99 b- DECAY

From Alburger *et al.* (1980Al02) the total transition probability of the β -transition is: 0,0037(6)%

2- NUCLEAR DATA

Level energy of Ru-99

The values of the level energies are from Peker (NDS 73,1)

Level 2	322,38 (6)
Level 1	89,68 (5)

2.1- b-TRANSITIONS

Only Alburger *et al.* (1980) have totally studied the beta decay of Tc-99m. The lg ft values were calculated by Singh *et al.* (1998) and derived from measurements by Alburger *et al.* (1980):

Transition	Energy	lg ft Singh <i>et al.</i>	lg ft Alburger <i>et al.</i>	Nature
0-0	434,8 (26)	9,4	9,39(11)	unique first-forbidden
0-1	346,7(20)	8,7	8,66(8)	first-forbidden
0-2	113,8 (20)	8,50	7,79(3)	first-forbidden

The adopted values of lg ft and average beta energies have been calculated using the LOGFT program and the level energies from ENSDF.

2.2 GAMMA TRANSITIONS and INTERNAL CONVERSION COEFFICIENTS

Multipolarity

Transition 322 keV	M1+(E2)
Transition 233 keV	(M1+E2)
Transition 89 keV	29%M1+E2 ($\delta = -1,56(2)$ measured by Kistner (1976Ki02))

Internal Conversion Coefficients

No experimental data have been found in the known literature. The Rösler tables were used to deduce theoretical coefficients :

keV	a _T	a _K	a _L	a _M
322,4	0,01747	0,01519		
232,8	0,0478	0,0412		
89,6	1,492	1,171	0,270	0,0512

3. ATOMIC DATA

The fluorescence yields taken from 96Sc06 (Schönfeld and Janßen) are: $\omega_K = 0,796(4)$, $\omega_L = 0,0453(11)$, $n_{KL} = 1,000(4)$

4. PHOTON EMISSIONS

4.1 X-RAY EMISSIONS

The emission intensities are very low and have not been calculated.

4.2 GAMMA EMISSIONS

Energy, keV	Relative emission probability	Absolute emission intensity	Author(s)
322	0,97*10 ⁻⁶ (15)	0,96*10 ⁻⁴ (6)	Jones and Griffin (1970Jo24)
	1,10*10 ⁻⁶ (10)		Decombaz <i>et al.</i> (1972De76)
	1,13*10 ⁻⁶ (9)		Alburger <i>et al.</i> (1980A102)
	1,09*10 ⁻⁶ (6)		LWEIGHT reduced- $\chi^2 = 0,42$ weighted mean and internal uncertainty
232	0,95*10 ⁻⁷ (17)	0,84*10 ⁻⁵ (15)	Alburger <i>et al.</i> (1980)
		1,04*10 ⁻³ (20)	deduced from the level balance
89			

For the 322 keV and the 232 keV gamma -rays, the measured emission probabilities are relative to the 140,5 keV emission probability. The absolute emission probabilities are deduced from the adopted absolute emission probability of the 140,5 keV gamma-ray: 88,5(2) %.

For the 89 keV line, no experimental value is available.

The 89 keV level is mainly fed by the beta transition from Tc -99m. With a beta transition probability of $2,6(5) \times 10^{-3}$ and $\alpha_T = 1,49(5)$, the absolute emission probability is : $1,04(20) \times 10^{-3}$.

5. ELECTRON EMISSIONS

For the 434,8 and 346,7 keV β^- transitions, the energies and transition probabilities were measured by Alburger (1980).

For the third β^- transition of 113,8 keV, no direct experimental data was found.

The energy is estimated by Alburger *et al.* (1980), and the absolute transition probability is derived from 3 experimental and relative values :

$$\begin{array}{ll} P_{\gamma}(322)/P_{\gamma}(140,5) = 1,10(6) \times 10^{-6} & \text{Decombaz } et al.(1972) \\ P_{\gamma}(322)/P_{\gamma}(140,5) = 0,97(15) \times 10^{-6} & \text{Jones and Griffin (1970Jo24)} \\ P_{\gamma}(322)/P_{\gamma}(140,5) = 1,113(9) \times 10^{-6} & \text{Alburger } et al.(1980) \end{array}$$

The weighted mean of γ emission probability relative to the 140 keV -line calculated by Alburger *et al.* (1980) is: $1,10(6) \times 10^{-6}$.

The gamma transitions probabilities are calculated from the gamma emission probabilities and the internal conversion coefficients :

$$\begin{array}{l} P_{\gamma}(322) = P_{\gamma}(322) \times (1 + \alpha_T(322)) \\ P_{\gamma}(322) = 1,10 \times 10^{-6} \times P_{\gamma}(140,5) \\ P_{\gamma}(322) = 1,10 \times 10^{-6} \times 88,5 \times 1,0175 = 0,99 \times 10^{-4} \end{array}$$

As the level 0 is feeding by 93% of the transitions starting from the 322 keV-level, the probability of the 322-keV β transition can be deduced : $0,99 \times 10^{-6}/0,93 = \mathbf{1,06(6) \times 10^{-4}}$.

References

- K. T. Bainbrige, M. Goldhaber, E. Wilson. Phys. Rev. 90 (1953) 430 ; Half-life
 J. RAVIER, P. MARGUIN, A. MOUSSA. J. Phys. Radium 22 (1961) 249 ; K/L
 P. CROWTHER, J. S. Eldridge. Nucl. Phys. 66 (1965) 472 ; Half-life
 S. R. AMTEY. Phys. Lett. 23-10 (1966) 581 ; M ICC, multipolarity
 I. W. GOODIER, A. WILLIAMS. Nature 210 (1966) 614 ; Half-life
 C. W. E. Van EIJK , B. VAN NOOIJNEN, F. SCHUTTE. Nucl. Phys. A121 (1968) 440 ;Gamma-ray energies, Conv. Elec. emission probabilities
 S. M. BRAHMAVAR, J. H. Hamilton, J. J. Pinajian. Quoted in Nucl. Phys. A121 (1968) 440 ; ICC
 J. S. GEIGER. AECL-3166 PR-P-79 (1968) 29 ;Gamma-ray energies, K-Conv. Elec. emission probabilities
 P. STEINER, E. GERDAU, W. HAUTSCH, D. STEENKEN. Z. Phys. 221 (1969) 281 ; Half-life
 W. B. COOK, L. Schellenberg, M. W. Johns. Nucl. Phys. A139 (1969) 277 ; Multipole mixing ratio
 P. STEINER. Z.Phys. 221 (1969) 281 ; Half-life
 E.BASHANDY, N. Ibrahiem. Z. Phys. 219 (1969) 337 ; K/L ratio
 V. A. AGEEV et al. Izv. Akad. Nauk SSSR, Ser. Fiz. 33 (1969) 1279 ; Conv. Elec. emission probabilities
 A. VUORINEN. Ann. Acad. Sci. Fenn. Ser. A VI (1969) 311 ; ICC, ; Half-life
 E. BASHANDY. Z. Naturforsch 24A (1969) 1893 ; K ICC, Multipolarity
 J. LEGRAND, F. Lagoutine, J. P. Brethon. Int. J. Appl. Rad. Isot. 21 (1970) 139 ; Half-life
 J. D. JONES, H. C. GRIFFIN. Radiochem. Radioanal. Letters 4-6 (1970) 381 ; Beta emission probabilities, Gamma-ray energies

- J. Mc DONALD, A. BÄCKLIN, S. G. MALMSKOG. Nucl. Phys. A162) 365 ; Multipolarity, Half-life, Gamma-ray emission probabilities
- W. M. LACASSE, J. H. HALMILTON. Nucl. Phys. A171 (1971) 641 ; M ICC/N ICC, Gamma-ray energies
- N. A. VOINOVA, A. I. EGOROV, Yu. V. KALINICHEV, A. G. SERGEEV. Bull. Ac.Sc URSS, Phys. Ser. 35 (1972) 794 ; K ICC, K/L, Multipolarity
- J. F. EMERY. Nucl. Sci. Eng. 48-3 (1972) 319 ; Half-life
- M. DECOMBAZ, J. J. GOSTELY, P. LERCH. Radiochem. Radioanal. Letters 10 (1972) 119 ; Gamma emission probabilities
- G. K. SHENOY. J. Phys. (London) A6 (1973) L144 ; Half-life
- J. Legrand, M. Blondel, P. Magnier, C. Perrot, J. P. Brethon. Report CEA-R-4427 (1973) ; Gamma-ray emission probabilities, Conv. Elec. emission probabilities
- R. A. MEYER. Report UCRL-76207 (1974) ; Half-life
- P. L. GARDULSKI, M. L. WIEDENBECK. Phys. Rev. C9,1 (1974) 262 ; Multipole mixing ratio
- I. M. Band, M. B. Trzhaskovskaya, M. A. Listengarten. Atomic Data and Nuclear Data Tables 18 (1976) 433 ; K and L-shell internal conversion coefficients
- O. C. KISTNER, A. H. LUMPKIN. Phys. Rev. C13 (1976) 1132 ; Multipolarities
- H. HOUTERMANS, D. MILOSEVIC, F. REICHEL. Int. J. Appl. Radiat. Isotop. 31 (1980) 153 ; Half-life
- D. E. ALBURGER, P. RICHARDS, T. H. KU. Phys. Rev. C21 (1980) 705 ; Beta emission probabilities, Beta emission energies
- H. MAZAKI, S. KAKIUCHI, T. MUKOYAMA, M. MATSUI. Phys. Rev. C21 (1980) 344 ; Half-life
- J. K. DICKENS, T. A. Love. Nucl. Inst. Meth. 175 (1980) 535 ; Gamma-ray emission probabilities, X-ray emission probabilities, K ICC, T ICC
- A. R. RUTLEDGE, L. V. SMITH, J. S. MERRIT. Report AECL-6692 (1980) ; Half-life
- R. G. HELMER, A. J. CAFFREY, R. J. GEHRKE, R. C. GREENWOOD. Nucl. Instrum. Methods 188 (1981) 151 ; Gamma-ray energies
- V. N. GERASIMOV, A. G. ZELENKOV, V. M. KULAKOV, V. A. PCHELIN, A. A. SOLDATOV. Sov. J. Nucl. Phys. 34-1 (1981) 1 Multipolarity, Gamma ray energies
- A. Simonits, L. Moens, F. De Corte, A. De Wispelaere, J. Hoste. J. Radioanal. Chem. 67 (1981) 61 ; Gamma-ray emission probabilities
- K. Singh, H. S. Sahota. J. Phys. Soc. Jap. 51-12 (1982) 3766 ; Multipole mixing ratio
- R. L. AYRES, A. T. HIRSCHFELD. Int. J. Appl. Rad. Isot. 33-10 (1982) 835 ; Half-life
- K. F. WALZ, K. Debertin, H. Schrader. Int. J. Appl. Rad. Isot. 34-8 (1983) 1191 ; Half-life
- CHEN DA. IEEE-Transactions on Nuclear Science 32-1 (1985) 71 ; Gamma-ray emission probabilities
- D. C. SANTRY, G. C. BOWES. Health Physics 57-4 (1989) 673 ; Half-life
- I. ALFTER, Z. Phys. A347 (1993) 1 Nuclear levels ; Half-life
- L. K. PEKER. Nucl. Data Sheets 73,1 (1994) 1 ; Level energies, Spin and Parity
- G. Audi, A. H. Wapstra. Nucl. Phys. A 595 (1995) 409 ; Q
- E. Schönfeld, H. Janssen. Nucl. Instrum. Methods A369 (1996) 527 ; Fluorescence yields
- B. Singh, J. L. Rodriguez, S. S. M. Wong, J. K. Tuli. Nucl. Data Sheets 84,3 (1998) ; lg ft
- V. KOLTSOV, L. G. Mashirov, D. N. Suglovov. Bull. Acad. Sci. USSR, Phys. Ser. 62,5 (1998) 789 ; Half-life
- M.-M. BÉ, E. BROWNE, V. CHECHEV, R. HELMER, E. SCHÖNFELD. Table de Radionucléides, ISBN 2 7272 0200 8, CEA, F-91191 Gif sur Yvette (1999) ; T ICC (Cs-137)
- R. G. HELMER, C. van DER LEUN. Nucl. Instrum. Methods Phys. Res. A450 (2000) 35 ; Gamma ray energies
- H. SCHRADER. Appl. Rad. Isotopes 60(2004)317 ; Half-life.
- M. A. L. da SILVA, M. C. M. de ALMEIDA, C. J. da SILVA, J. U. DELGADO. Appl. Rad. Isotopes 60(2004)301 ; Half-life.

**¹⁰⁶Ru – Comments on evaluation of decay data
by A. Arinc**

Evaluation completed: February 2012

Literature cut-off date: September 2011

Evaluation procedure

Weighted mean analyses were applied to determine recommended values throughout the evaluation when the data were in statistical agreement. When the data were not in statistical agreement, the Limitation of Relative Statistical Weights (LRSW) was used. Uncertainties were expanded to match the minimum input uncertainty where appropriate.

Decay scheme

¹⁰⁶Ru disintegrates by beta minus emission to the ground state of ¹⁰⁶Rh. The spin and parity of ¹⁰⁶Rh ($J^\pi=1+$) are taken from the mass-chain evaluation of De Frenne (2008De09).

The Q^- value of 39.40 (21) keV is taken from the evaluation of Wang et al. (2012Wa38).

Half-life

The experimental half-life values used for calculating the recommended value are given in Table 1. The half-life value from Walz (1983Wa26) was omitted from the analysis as a more recent publication was available from the same laboratory (Schrader, 2004Sc04).

The AveTool computer code was used to calculate the average using three statistical methods: Limitation of Relative Statistical Weights (LRSW), Normalised Residual Methods (NRM) and the Rajeval Technique (RT).

The data set is discrepant with a reduced- χ^2 of 12.1 (@ 99 % confidence level) on the weighted mean (WM) which is larger than the critical reduced- χ^2 of 2.80. The recommended value of 371.5 (21) days is calculated using LRSW technique and was chosen as it includes the most precise value from Houtermans (1980Ho17).

Table 1. Experimental half-life values of ¹⁰⁶Ru.

Reference	Half-life (days)	Comments
1956Sc87	366.6 (9)	
1957Me47	373 (4)	Published as 1.02 (1) year
1960Ea02	365.8 (17)	
1961Wy01	368.0 (18)	
1965Fl02	371 (1)	
1980Ho17	373.59 (15)	
1983Wa26	371.7 (15)	Not included in averaging
2004Sc04	370.5 (6)	
UWM	369.7 (11)	
WM	371.47 (30)	reduced- $\chi^2 = 12.1$
LRSW	371.5 (21)	
NRM	370.3 (7)	
RT	370.4 (5)	
Recommended value	371.5 (21)	

Beta particles

The maximum beta-particle emission energy was derived from the $Q(\beta^-) = 39.40 (21)$ keV value from Wang et al. (2012Wa38).

The mean beta-particle energy and $\log ft$ values were calculated from the LOGFT code.

References

- 1950Ag01 H.M. Agnew, Phys. Rev. 77 (1950) 655.
[E_β]
- 1956Sc87 R.P. Schuman, M.E. Jones, A.C. McWherter, J. Inorg. Nuclear Chem. 3 (1956) 160.
[Half-life]
- 1957Me47 W.F. Merritt, P.J. Campion, R.C. Hawkings, Can. J. Phys. 35 (1957) 16.
[Half-life]
- 1960Ea02 H.T. Easterday, R.L. Smith, Nuclear Phys. 20 (1960) 155.
[Half-life]
- 1961Wy01 E.I. Wyatt, S.A. Reynolds, T.H. Handley, W.S. Lyon, H.A. Parker, Nucl. Sci. Eng. 11 (1961) 74.
[Half-life]
- 1965Fl02 K.F. Flynn, L.E. Glendenin, E.P. Steinberg, Nucl. Sci. Eng. 22 (1965) 416.
[Half-life]
- 1980Ho17 H. Houtermans, O. Milosevic, F. Reichel, Int. J. Appl. Radiat. Isotop. 31 (1980) 153.
[Half-life]
- 1983Wa26 K.F. Walz, K. Debertin, H. Schrader, Int. J. Appl. Radiat. Isotop. 34 (1983) 1191.
[Half-life]
- 2004Sc04 H. Schrader, Appl. Radiat. Isot. 60 (2004) 317.
[Half-life]
- 2008De09 D. De Frenne, A. Negret, Nucl. Data Sheets 109 (2008) 943.
[Spin, parity]
- 2012Wa38 M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer, Chinese Physics C36 (2012) 1603.
[Q value]

**¹⁰⁶Rh – Comments on evaluation of decay data
by A. Arinc**

Evaluation completed: March 2013

Literature cut-off date: December 2012

Evaluation procedure

Weighted mean analyses were applied to determine recommended values throughout the evaluation when the data were in statistical agreement. When the data were not in statistical agreement, the Limitation of Relative Statistical Weights (LRSW) was used. Uncertainties were expanded to match the minimum input uncertainty where appropriate.

Decay scheme

¹⁰⁶Rh is the daughter of ¹⁰⁶Ru and disintegrates by beta minus emission to various excited levels and the ground state of ¹⁰⁶Pd. The spins, parities, multipolarities, mixing ratios and level energies of ¹⁰⁶Pd are based on the mass-chain evaluation of De Frenne and Negret (2008De09).

The Q(β⁻) value of 3546 (5) keV is taken from the evaluation of Wang *et al.* (2012Wa38).

Half-life

The experimental half-life values used for calculating the recommended value are given in Table 1. The half-life values from 1946Se30 and 1950Gl05 were omitted from the analysis as no uncertainty was given.

The data set is discrepant with a reduced-χ² of 10.9 on the Weighted Mean (WM) which is larger than the critical reduced-χ² of 6.63. The recommended value of 30.1 (3) seconds is the Unweighted Mean (UWM) of 1966Mi10 and 1969KoZW.

Table 1. Experimental half-life values of ¹⁰⁶Rh.

Reference	Half-life (seconds)	Comments
1946Se30	40	no uncertainty
1950Gl05	30	no uncertainty
1966Mi10	30.36 (15)	
1969KoZW	29.80 (8)	
UWM	30.1 (3)	
WM	29.92 (23)	reduced-χ ² = 10.9
Recommended value	30.1 (3)	

In the majority of circumstances ¹⁰⁶Rh will be encountered in equilibrium with ¹⁰⁶Ru therefore an accurate knowledge of the half-life is not required; however for applications where the ¹⁰⁶Rh is unsupported by ¹⁰⁶Ru (for example decay heats), new measurements of the half-life would be beneficial as the available data are inconsistent and in excess of 40 years old.

Gamma rays

When more than one publication was available per laboratory only the latest published data were used (e.g. 1972Ma71 and 1973Ma35; 1960Ro12 and 1965Ro09). Published data with no reported uncertainty were not used for averaging process (i.e. 1971Az02, 1957Fi50).

Energies

The recommended values were derived from the differences in level energies as adopted from the mass-chain evaluation of 2008De09, and were compared to the experimental values calculated from the weighted mean (calculated with LWEIGHT4 code) of 1982Ka10, 1977Ok02, 1969St03, 1973Ma35, 1975Hs02, 1965Ro09, 1967Fo09, 1967Vr05 and 1968Ha35. When there was a disagreement between the calculated and evaluated gamma-ray emission energies, the latter were used. Although a few recent publications were available (1976Sh25, 1983Ku03 and 1993Ch32) on the precise measurement of the 511.85 keV gamma-ray emission energy, the recommended value of 511.8534 (23) keV was taken from Helmer and van der Leun (2000He14). The data set from Rao and Fink (1967Ra11) was not taken into account as many of the values in this publication were excluded by Chauvenet's criterion as outliers. In the publication of Forest *et al.* (1967Fo09) only the values up to 2614 keV were used as this was the last energy calibration point available to the user based upon the information given, and any energies above this point will have been based upon extrapolation. In the publication of Hsue *et al.* (1975Hs02), only the use of a standard gamma calibration source was mentioned, which usually does not cover gamma-ray emissions as high as 3.3 MeV. The values from 1975Hs02 above 2000 keV are generally lower and with smaller associated uncertainties compared with the rest of the values reported by other authors, which may imply that an extrapolation of the energies was performed rather than a more precise interpolation. Therefore, the evaluator has decided to increase the uncertainties of 1975Hs02 above 2000 keV to a minimum of 0.3 keV to be in line with other reported uncertainties.

Adopted nuclear levels of ¹⁰⁶Pd can be seen in Table 2.

The experimental results and recommended values can be seen in Table 1 of Appendix 1.

Table 2. Adopted nuclear levels of ¹⁰⁶Pd.

Nuclear level number	Nuclear level energy (keV)	Spin and parity
0	0	0 +
1	511.8547 (23) ^a	2 +
2	1128.02 (3)	2 +
3	1133.76 (4)	0 +
4	1229.30 (4)	4 +
5	1557.68 (4)	3 +
6	1562.25 (3)	2 +
7	1706.44 (5)	0 +
8	1909.37 (16)	(1 +, 2 +)
9	2001.48 (5)	0 +
10	2242.48 (5)	2 +
11	2278.11 (9)	0 +
12	2308.82 (5)	2 +
13	2439.10 (7)	2 +
14	2484.66 (20)	(1 -)
15	2500.31 (8)	2 -
16	2624.40 (5)	0 +
17	2705.30 (8)	(1 +)
18	2717.59 (21)	
19	2783.74 (21)	2 +
20	2820.97 (9)	2 +
21	2828.29 (9)	0 +
22	2877.92 (7)	0 +
23	2902.48 (10)	2 +
24	2917.86 (8)	2 +
25	2968.68 (21)	3 -
26	3037.32 (17)	1, 2
27	3054.97 (9)	1 +
28	3083.91 (18)	0
29	3163.7 (3)	(1, 2+)
30	3221.37 (25)	0 +
31	3249.9 (5)	2 +
32	3252.0 (4)	2 +
33	3273.5 (7)	1, 2
34	3299.2 (7)	
35	3320.5 (3)	0 +
36	3376.6 (9)	
37	3401.9 (6)	

^a) Derived from Helmer and van der Leun (2000He14) gamma-ray emission energy of 511.8534 (23) keV.

Probabilities

The gamma-ray emission probabilities were calculated as weighted means (calculated with LWEIGHT4 code) of the measurements of 1982Ka10, 1977Ok02, 1975Hs02, 1975De17, 1973Ma35, 1969Od01, 1696St03, 1968Ha35, 1967Vr05, 1967Ra11, 1967Fo09, 1966Ov01, 1962Am03, 1960Se07 and 1960Ro12. In some of the earlier papers (1960Ro12, 1960Se07, 1962Am03 and 1966Ov01) the high energy gamma-ray emission probabilities were impossible to place due to large differences between the energies measured by the authors and the adopted energies. In the publication of Hsue *et al.* (1975Hs02), the reported annihilation contribution of

5.1 % on the most intense peak at 511.85 keV was disputed by Kaur *et al.* (1982Ka10) as being almost negligible. The annihilation contribution is dependent on the specifications of the gamma-ray spectrometry system and cannot be verified without precise details on the equipment and shielding used. Therefore the evaluator has decided to increase the reported uncertainties of 1975Hs02 by 5.1 %.

Values were initially normalised to an intensity for the 511.85 keV gamma-ray emission of 1000.

The experimental results and recommended values can be seen in Table 2 of Appendix 1.

Internal conversion coefficients and E0 transition probability

Theoretical internal conversion coefficients (ICCs) were calculated using the BrIcc code (Kibédi *et al.*, 2008Ki07) with the “frozen orbital” approximation, which uses interpolated values of Band *et al.* (2002Ba85).

The transition at 1133.75 keV is pure E0 and absolute measurements for this transition were not available; the transition probability cannot be calculated using the balance of the decay scheme because the beta probabilities are not well known for the levels concerned. Using the ratio of the K-conversion electron intensity of the E0 transition to that of the E2 transition at 621.9 keV, $q^2_{\text{K}}(\text{E0/E2}) = 0.162$ (7) from Kibédi and Spear (2005Ki02) and the following data for the 621.9 keV line:

Relative gamma-ray emission probability (P_{γ}): 481 (5)

ICC for K shell (α_{K}): 0.002 82 (40)

The calculated relative K-conversion intensity for the 1133.7 keV gamma ray is:

$$= P_{\gamma(621)} \times \alpha_{\text{K}(621)} \times q^2_{\text{K}}(\text{E0/E2 of 1133 to 621})$$

$$= 481 (5) \times 0.00282 (40) \times 0.162 (7) = 0.22 (3)$$

Using the electronic factors for E0 calculated by BrIcc and assuming proportionality between the electronic factors and the conversion on each subshell, this gives a total relative conversion intensity for the 1133.7 keV transition of 0.25 (3).

Normalisation factor

Five absolute experimental values reported for the strongest gamma ray line at 511.85 keV are given in Table 3.

The AveTools computer code was used to calculate the average using three statistical methods: Limitation of Relative Statistical Weights (LRSW), Normalised Residual Methods (NRM) and the Rajeval Technique (RT).

Table 3. Absolute experimental values for the gamma ray line at 511.85 keV.

Reference	Absolute $P_{\gamma_{1,0}}$ (%)	Used	Comments
1953Ka47	20.5	no	No uncertainty
1966Ov01	21.0 (20)	no	Later publication 1975Ge06 available
1969Od01	20.6 (6)	yes	
1975De17	20.47 (23)	yes	
1975Ge06	22.0 (11)	yes	Uncertainty quoted @ 95 % confidence
UWM	21.0 (5)		
WM	20.7 (4)		
LRSW	20.9 (5)		
NRM	20.63 (21)		
RT	20.52 (21)		
Recommended value	20.52 (23)		Minimum input uncertainty used

The recommended value was taken from the Rajeval Technique and the minimum input uncertainty from 1975De17 was used. Using the absolute $P_{\gamma_{1,0}}$ value calculated above we can derive a normalisation factor of 0.020 52 (23).

The theoretical value obtained from the balance of the decay scheme to the ground state, taking the beta emission efficiency $\beta_{0,0}$ from 1992Gr21 of 79.1 (16) %, is 0.0202 (16) which is in good agreement with the value calculated from the experimental data.

Beta particles

Energies

The values of the maximum beta-particle energies were derived from the $Q(\beta^-) = 3546 (5)$ keV value from Wang *et al.* (2012Wa38) and the adopted nuclear level energies given in Table 2. The experimental data for the maximum beta-particle emission energies to the ground state ($\beta_{0,0 \max}$) is in good agreement with the calculated data and can be found in Table 4.

The mean beta-particle energies and $\log ft$ values were calculated with the LOGFT code.

Table 4. Experimental values for the maximum beta-particle emission energies to the ground state and first excited state of ¹⁰⁶Pd.

Reference	$\beta_{0,0 \max}$ (MeV)	$\beta_{0,1 \max}$ (MeV)
1947Pe07	3.55 (10)	
1958Gr07	3.55 (1)	
1966JoZZ	3.55 (100)	3.04 (7)
1963Ke13	3.55	3.07
1969Be74	3.52 (12)	
Recommended value	3.546 (5)	3.034 (5)

Emission probabilities

The theoretical beta-particle probabilities were calculated from the P($\gamma+ce$) balance using the GTOL software and are given in Table 5.

Table 5. Beta-particle emission energies, probabilities, transition type and log *ft* values of ¹⁰⁶Rh.

Transition	E _{βmax} (keV)	P _{β} (%)	Transition type	log <i>ft</i>
$\beta_{0,0}$	3546 (5)	78.80 (24)	Allowed	5.18
$\beta_{0,1}$	3034 (5)	8.2 (3)	Allowed	5.87
$\beta_{0,2}$	2418 (5)	0.608 (21)	Allowed	6.58
$\beta_{0,3}$	2412 (5)	9.82 (15)	Allowed	5.37
$\beta_{0,4}$	2317 (5)	0.005 1 (5)	Unique 2 nd Forbidden	11.0
$\beta_{0,6}$	1984 (5)	1.67 (3)	Allowed	5.79
$\beta_{0,7}$	1840 (5)	0.066 4 (10)	Allowed	7.06
$\beta_{0,8}$	1637 (5)	0.002 77 (21)	(Allowed)	8.24
$\beta_{0,9}$	1545 (5)	0.448 (9)	Allowed	5.93
$\beta_{0,10}$	1304 (5)	0.037 2 (8)	Allowed	6.72
$\beta_{0,11}$	1268 (5)	0.043 (5)	Allowed	6.62
$\beta_{0,12}$	1237 (5)	0.043 0 (7)	Allowed	6.57
$\beta_{0,13}$	1107 (5)	0.020 8 (5)	Allowed	6.71
$\beta_{0,14}$	1061 (5)	0.000 93 (15)	(1 st Forbidden)	7.99
$\beta_{0,15}$	1046 (5)	0.028 4 (6)	1 st Forbidden	6.48
$\beta_{0,16}$	922 (5)	0.090 (3)	Allowed	5.78
$\beta_{0,17}$	841 (5)	0.010 6 (4)	(Allowed)	6.56
$\beta_{0,18}$	828 (5)	0.000 23 (12)	-	-
$\beta_{0,19}$	762 (5)	0.001 17 (8)	Allowed	7.36
$\beta_{0,20}$	725 (5)	0.009 0 (3)	Allowed	6.40
$\beta_{0,21}$	718 (5)	0.007 31 (19)	Allowed	6.47
$\beta_{0,22}$	668 (5)	0.026 2 (9)	Allowed	5.81
$\beta_{0,23}$	644 (5)	0.007 60 (18)	Allowed	6.29
$\beta_{0,24}$	628 (5)	0.018 3 (7)	Allowed	5.87
$\beta_{0,25}$	577 (5)	0.000 22 (4)	Unique 1 st Forbidden	7.82
$\beta_{0,26}$	509 (5)	0.002 2 (3)	-	-
$\beta_{0,27}$	491 (5)	0.010 1 (5)	Allowed	5.76
$\beta_{0,28}$	462 (5)	0.002 78 (13)	-	-
$\beta_{0,29}$	382 (5)	0.000 70 (5)	(Allowed)	6.55
$\beta_{0,30}$	325 (5)	0.004 02 (13)	Allowed	5.56
$\beta_{0,31}$	296 (5)	0.000 086 (16)	Allowed	7.09
$\beta_{0,32}$	294 (5)	0.000 21 (4)	Allowed	6.70
$\beta_{0,33}$	272 (5)	0.000 049 (14)	-	-
$\beta_{0,34}$	247 (5)	0.000 082 (21)	-	-
$\beta_{0,35}$	226 (5)	0.000 87 (8)	Allowed	5.71
$\beta_{0,36}$	169 (5)	0.000 025 (9)	-	-
$\beta_{0,37}$	144 (5)	0.000 012 5 (19)	-	-

Table 6. Published values for the beta-particle emission probability to the ground state of ¹⁰⁶Pd.

Reference	$\beta_{0,0}$ (%)
1947Pe07	82 (2)
1958Gr07	90
1960Se07	70
1962Am03	67.2
1963Ke13	63
1966JoZZ	90
1992Gr21	79.1 (16)
Recommended value	78.80 (24)

The most recent and most precise measurement of the beta-particle emission probability to the ground state of ¹⁰⁶Pd reported by Greenwood *et al.* (1992Gr21) using total absorption gamma-ray spectrometry (TAGS) and the value derived theoretically from the singles gamma studies agree within the stated uncertainties.

Atomic data

The values of ω_K , ω_L , n_{KL} and relative probabilities of the X-ray and Auger emissions were derived from Schönfeld and Janßen (1996Sc06).

The energies and relative emission probabilities of the X-ray and Auger electrons have been calculated using the computer code EMISSION. A summary of the results is given in Tables 7 and 8. There are no known publications of measured data with which these values can be compared.

Table 7. Calculated L X-ray emission energies and probabilities.

L Xray	Energy, /keV	Calculated value
L λ	2.50	0.000 174 (5)
L α	2.83 – 2.84	0.004 68 (11)
L η	2.66	0.000 060 8 (17)
L β	2.99 – 3.17	0.002 70 (8)
L γ	3.25 – 3.55	0.000 242 (6)
LX total		0.007 85 (14)

Table 8. Calculated K X-ray emission energies and probabilities.

K Xray	Energy, /keV	Calculated value
K α_2	21.02	0.031 0 (5)
K α_1	21.18	0.058 6 (9)
K β_1'	23.8 – 24.0	0.016 08 (29)
K β_2'	24.30 – 24.34	0.002 73 (10)
KX total		0.108 5 (17)

References

- 1946Se30 W. Seelmann-Eggebert, *Naturwiss.* 33 (1946) 279.
[Half-life]
- 1947Pe07 W.C. Peacock, *Phys. Rev.* 72 (1947) 1049.
[E_β, P_β]
- 1950GI05 L.E. Glendenin, E.P. Steinberg, *National Nuclear Energy Series NNEs* 9 (1950) 793.
[Half-life]
- 1953Ka47 B. Kahn, W.S. Lyon, *Phys. Rev.* 92 (1953) 902.
[E_γ, P_γ]
- 1957Fi50 E.I. Firsov, A.A. Bashilov, *Izvest. Akad. Nauk SSSR, Ser. Fiz.* 21 (1957) 1633;
Columbia Tech. Transl. 21 (1958) 1619.
[E_γ, P_γ]
- 1958Gr07 E.P. Grigorev, A.V. Zolotavin, I.I. Kuzmin, E.D. Pavlitskaia, *Izvest. Akad. Nauk SSSR, Ser. Fiz.* 22 (1958) 194; *Columbia Tech. Transl.* 22 (1959) 191.
[E_β, P_β]
- 1960Ro12 R.L. Robinson, F.K. McGowan, W.G. Smith, *Phys. Rev.* 119 (1960) 1692.
[E_γ, P_γ]
- 1960Se07 O.J. Segaert, J.L. Demuynck, *Nucl. Phys.* 16 (1960) 492.
[E_γ, P_γ]
- 1962Am03 S.Y. Ambiyev, R.P. Sharma, *Nucl. Phys.* 29 (1962) 657.
[E_γ, P_γ, P_β]
- 1963Ke13 T.J. Kennett, G.L. Keech, *Nucl. Instrum. Methods* 24 (1963) 142.
[E_β, P_β]
- 1965Ro09 R.L. Robinson, P.H. Stelson, F.K. McGowan, J.L.C. Ford, Jr., W.T. Milner, *Nucl. Phys.* 74 (1965) 281.
[E_γ]
- 1966JoZZ A.S. Johnston, G.M. Julian, *Bull. Am. Phys. Soc.* 11 (1966) No.3, 409.
[E_β, P_β]
- 1966Mi10 V. Middelboe, *Nature* 211 (1966) 283.
[Half-life]
- 1966Ov01 V.V. Ovechkin, T.K. Ragimov, D.F. Rau, *Yadern. Fiz.* 4 (1966) 683; *Soviet J. Nucl. Phys.* 4 (1967) 484.
[E_γ, P_γ]
- 1967Fo09 H. Forest, M. Huguet, C. Ythier, *C. R. Acad. Sci., Paris* 264B (1967) 1614.
[E_γ, P_γ]
- 1967Ra11 P.V. Rao, R.W. Fink, *Nucl. Phys.* A103 (1967) 385.
[E_γ, P_γ]
- 1967Vr05 J. Vrzal, E.P. Grigorev, A.V. Zolotavin, J. Liptak, V.O. Sergeev, J. Urbanets, *Izvest. Akad. Nauk SSSR, Ser. Fiz.* 21 (1967) 696; *Bull. Acad. Sci. USSR, Phys. Ser.* 31 (1968) 692.
[E_γ, P_γ]
- 1968Ha35 J. Hattula, E. Liukkonen, *Ann. Acad. Sci. Fennicae, Ser. A VI*, No.274 (1968).
[E_γ, P_γ]
- 1969Be74 E. Beck, *Nucl. Instrum. Methods* 76 (1969) 77.
[E_β]
- 1969KoZW Y. Kobayashi, *Japan Atomic Energy Research Institute report JAERI-1178* (1969) 21.
[Half-life]
- 1969Od01 P. Odru, *Radiochim. Acta* 12 (1969) 64.
[P_γ]

- 1969St03 Kl.-D. Strutz, H.-J. Strutz, A. Flammersfeld, Z. Phys. 221 (1969) 231.
[E_γ, P_γ]
- 1971Az02 T. Azuma, Y. Sato, Annual Rep. Radiat. Center Osaka Prefect. 12 (1971) 28.
[E_γ, P_γ]
- 1972Ma71 C. Marsol, O. Rahmouni, G. Ardisson, C. R. Acad. Sci., Paris 275B (1972) 805.
[E_γ, P_γ]
- 1973Ma35 C. Marsol, G. Ardisson, Rev. Roum. Phys. 18 (1973) 1101.
[E_γ, P_γ]
- 1975De17 K. Debertain, U. Schotzig, K.F. Walz, H.M. Weiss, Ann. Nucl. Energy 2 (1975) 37.
[P_γ]
- 1975Ge06 A.M. Geidelman, V.V. Ovechkin, D.F. Rau, P.I. Fedotov, Y.V. Kholnov, Izvest. Akad. Nauk SSSR, Ser. Fiz. 39 (1975) 555; Bull. Acad. Sci. USSR, Phys. Ser. 39 (1975) No.3, 76.
[P_γ]
- 1975Hs02 S.T. Hsue, H.H. Hsu, F.K. Wohn, W.R. Western, S.A. Williams, Phys. Rev. C12 (1975) 582.
[E_γ, P_γ]
- 1976Sh25 K. Shizuma, H. Inoue, Y. Yoshizawa, Nucl. Instrum. Methods 137 (1976) 599.
[E_γ]
- 1977Ok02 K. Okano, Y. Kawase, S. Yamada, J. Phys. Soc. Japan 43 (1977) 381.
[E_γ, P_γ]
- 1982Ka10 R. Kaur, A.K. Sharma, S.S. Sooch, N. Singh, P.N. Trehan, J. Phys. Soc. Japan 51 (1982) 23.
[E_γ, P_γ]
- 1983Ku03 H. Kumahora, H. Inoue, Y. Yoshizawa, Nucl. Instrum. Methods 206 (1983) 489.
[E_γ]
- 1992Gr21 R.C. Greenwood, D.A. Struttman, K.D. Watts, Nucl. Instrum. Methods Phys. Res. A317 (1992) 175.
[P_β]
- 1993Ch32 T. Chang, S. Wang, H. Wang, B. Meng, Nucl. Instrum. Methods Phys. Res. A325 (1993) 196.
[E_γ]
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527.
[X_K, X_L, Auger electrons]
- 2000He14 R.G. Helmer, C. van der Leun, Nucl. Instrum. Methods Phys. Res. A450 (2000) 35.
[E_γ]
- 2002Ba85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor Jr., P.O. Tikkanen, S. Raman, At. Data Nucl. Data Tables 81 (2002) 1.
[ICC]
- 2005Ki02 T. Kibédi, R.H. Spear, At. Data Nucl. Data Tables 89 (2005) 77.
[E0 transition]
- 2008De09 D. De Frenne, A. Negret, Nucl. Data Sheets 109 (2008) 943.
[Spin, parity, energy level, multipolarity]
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor Jr., Nucl. Instrum. Methods Phys. Res. A589 (2008) 202.
[Theoretical ICC]
- 2012Wa38 M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer, Chinese Physics C36 (2012) 1603.
[Q value]

Appendix 1: Experimental and recommended gamma-ray emission energies and probabilities.

Table 1. Experimental and recommended gamma-ray emission energies (keV).

	1982Ka10	1977Ok02	1975Hs02 ^a	1973Ma35	1969St03	1968Ha35	1967Fo09 ^b	1967Ra11 ^c	1967Vr05	1965Ro09	Evaluated	Recommended
Y _{6,3}	428.40 (20)	428.4 (2)	428.39 (22)	428.3 (3)	427.9 (5)	427.7 (10)	-	-	-	-	428.35 (20)	428.49 (5)
Y _{6,2}	434.25 (21)	434.2 (3)	434.3 (5)	434.1 (5)	435.0 (15)	-	-	-	-	-	434.23 (21)	434.23 (4)
Y _{9,6}	439.17 (27)	439.5 (5)	-	439.6 (10)	-	-	-	-	-	-	439.3 (3)	439.23 (6)
Y _{1,0}	511.83 (8)	511.85 (10)	511.80 (15)	511.8 (8)	511.8 (2)	511.9 (3)	512.0 (5)	511.6 (5)	511.8 (2)	511.6 (5)	-	511.8534 (23)^d
Y _{7,2}	578.3 (2)	578.4 (2)	578.2 (9)	578.3 (3)	578.0 (4)	578.0 (15)	-	-	-	-	578.31 (20)	578.42 (6)
Y _{2,1}	616.22 (9)	616.2 (3)	616.33 (17)	616.1 (2)	616.2 (3)	616.0 (6)	616.0 (7)	616.6 (5)	-	616.2 (9)	616.22 (9)	616.16 (3)
Y _{3,1}	621.93 (6)	621.84 (10)	622.2 (3)	621.7 (2)	621.8 (3)	622.0 (3)	622.4 (5)	622.8 (5)	622.0 (2)	622.2 (5)	621.91 (6)	621.90 (4)
Y _{10,6}	680.25 (14)	680.2 (2)	680.6 (4)	680.1 (3)	680.1 (6)	-	-	-	-	-	680.21 (14)	680.23 (6)
Y _{10,5}	684.8 (2)	684.7 (2)	684.8 (8)	684.7 (4)	684.3 (8)	-	-	-	-	-	684.75 (20)	684.80 (6)
Y _{17,9}	702.8 (10)	-	-	-	-	-	-	-	-	-	702.8 (10)	702.8 (10)
Y _{11,6}	715.9 (2)	715.9 (2)	715.7 (5)	715.8 (3)	716.2 (3)	-	-	-	-	-	715.92 (20)	715.86 (9)
Y _{4,1}	717.4 (2)	717.3 (2)	717.3 (5)	717.3 (5)	717.1 (3)	716.6 (10)	-	717.6 (5)	-	-	717.34 (20)	717.44 (4)
Y _{12,5}	751.3 (2)	751.1 (6)	751.7 (8)	751.4 (10)	749.6 (15)	749 (2)	-	-	-	-	751.26 (20)	751.26 (20)
Y _{9,2}	873.49 (5)	873.5 (1)	873.73 (9)	873.1 (2)	873.1 (3)	873.7 (4)	873.5 (6)	873.4 (5)	-	873.5 (7)	873.52 (5)	873.46 (6)
Y _{15,5}	942.6 (4)	939.6 (20)	-	942.7 (10)	-	940 (3)	-	-	-	-	942.5 (4)	942.63 (9)
Y _{5,1}	1045.6 (6)	1045.8	1044.8 (15)	1046.5	1045.7	-	-	-	-	-	1045.5 (6)	1045.82 (4)
Y _{6,1}	1050.41 (6)	1050.4 (1)	1050.47 (7)	1050.3 (3)	1050.1 (3)	1050.3 (4)	1050.1 (6)	1049.8 (5)	-	1049.8 (7)	1050.42 (6)	1050.39 (3)
Y _{16,6}	1062.14 (5)	1062.2 (1)	1062.22 (8)	1062.0 (3)	1062.1 (5)	1062.7 (10)	1061.5 (8)	1061.8 (7)	-	-	1062.16 (5)	1062.14 (6)
Y _{10,3}	1108.7 (1)	1108.7 (5)	1108.66 (19)	1108.5 (3)	1108.9 (3)	-	-	-	-	-	1108.69 (10)	1108.71 (6)
Y _{10,2}	1114.48 (5)	1114.5 (3)	1114.69 (10)	1114.4 (3)	1114.6 (3)	1114 (2)	-	1114.0 (9)	-	-	1114.52 (5)	1114.45 (6)

	1982Ka10	1977Ok02	1975Hs02 ^a	1973Ma35	1969St03	1968Ha35	1967Fo09 ^b	1967Ra11 ^c	1967Vr05	1965Ro09	Evaluated	Recommended
Y _{2,0}	1128.07 (5)	1128.1 (2)	1128.21 (6)	1128.0 (2)	1128.0 (3)	1128.0 (4)	1127.8 (6)	1127.8 (5)	-	1127.4 (7)	1128.07 (5)	1128.01 (3)
Y _{3,0}	-	-	-	-	-	-	-	-	-	-	-	1133.75 (4)
Y _{11,2}	1150.2 (2)	1150.2 (2)	1150.21 (24)	1150.4 (5)	1150.1 (15)	1148 (2)	-	-	-	-	1150.20 (20)	1150.08 (9)
Y _{18,5}	1159.9 (2)	-	-	-	1159.7 (20)	-	-	-	-	-	1159.80 (20)	1159.90 (21)
Y _{12,2}	1180.73 (8)	1180.8 (2)	1180.86 (9)	1180.5 (3)	1180.6 (3)	1180 (2)	-	1179.8 (5)	-	-	1180.77 (8)	1180.79 (6)
Y _{7,1}	1194.54 (5)	1194.6 (1)	1194.68 (6)	1194.4 (3)	1194.3 (3)	1194.5 (6)	1194.2 (8)	1192.6 (5)	-	-	1194.59 (5)	1194.58 (5)
Y _{13,4}	1209.8 (2)	1209.7 (3)	1210.2 (7)	1209.7 (10)	-	-	-	-	-	-	1209.79 (20)	1209.79 (8)
Y _{20,6}	1258.8 (2)	1258.8 (3)	-	1258.3 (10)	1257.0 (20)	-	-	-	-	-	1258.77 (20)	1258.71 (9)
Y _{21,6}	1266.0 (2)	1266.1 (3)	1266.4 (7)	1266.0 (5)	1266.6 (15)	-	-	-	-	-	1266.05 (20)	1266.03 (9)
Y _{13,3}	1305.2 (2)	1305.3 (3)	1305.2 (8)	1305.3 (5)	1305.6 (10)	-	-	-	-	-	1305.24 (20)	1305.33 (8)
Y _{22,6}	1315.7 (2)	1315.7 (3)	1315.43 (24)	1315.7 (5)	1316.8 (15)	1315.4 (10)	-	-	-	-	1315.61 (20)	1315.66 (8)
Y _{24,6}	1355.7 (3)	1356.0 (3)	-	1355.5 (5)	1355.0 (6)	-	-	-	-	-	1355.7 (3)	1355.60 (9)
Y _{24,5}	1360.2 (3)	1360.1 (3)	1360.4 (4)	1360.1 (3)	1359.9 (6)	-	-	-	-	-	1360.2 (3)	1360.17 (9)
Y _{15,2}	1372.3 (3)	1372.3 (3)	1371.7 (3)	1372.0 (4)	-	1372 (4)	-	-	-	-	1372.1 (3)	1372.28 (9)
Y _{8,1}	1397.6 (2)	1397.7 (3)	1397.66 (15)	1397.1 (3)	1397.3 (4)	1396 (2)	-	-	-	-	1397.56 (15)	1397.51 (16)
Y _{9,1}	1489.6 (6)	1489.7 (6)	1489.46 (18)	1489.1 (5)	1488.9 (6)	1489.5 (15)	-	-	-	-	1489.42 (18)	1489.61 (5)
Y _{16,2}	1496.33 (13)	1496.6 (2)	1496.22 (8)	1496.1 (3)	1496.4 (3)	1497.5 (6)	1498.0 (12)	1495.5 (5)	1497 (1)	-	1496.30 (8)	1496.37 (6)
Y _{27,5}	1498.8 (2)	-	1498.71 (16)	1498.2 (8)	-	-	-	-	-	-	1498.73 (16)	1498.73 (16)
Y _{6,0}	1562.25 (6)	1562.2 (1)	1562.20 (6)	1561.9 (2)	1562.0 (3)	1562.4 (6)	1562.0 (10)	1561.6 (5)	-	1562.5 (11)	1562.21 (6)	1562.24 (3)
Y _{17,3}	1572.4 (2)	1572.3 (6)	1572.7 (3)	1572.4 (5)	1574.4 (9)	-	-	-	-	-	1572.47 (20)	1572.47 (20)

	1982Ka10	1977Ok02	1975Hs02 ^a	1973Ma35	1969St03	1968Ha35	1967Fo09 ^b	1967Ra11 ^c	1967Vr05	Evaluated	Recommended
Y _{17,2}	1577.2 (2)	1576.9 (8)	1577.1 (5)	1577.0 (7)	1579.8 (10)	-	-	-	-	1577.16 (20)	1577.27 (9)
Y _{20,3}	1687.4 (3)	1687.6 (10)	-	1687.9 (10)	1686.1 (20)	-	-	-	-	1687.4 (3)	1687.20 (10)
Y _{20,2}	1693.2 (3)	1693.0 (4)	-	1693.5 (8)	-	1695 (3)	~1694.5	-	-	1693.2 (3)	1693.2 (3)
Y _{10,1}	1730.5 (2)	1730.5 (3)	1730.6 (3)	1730.1 (3)	1730.0 (15)	-	~1730.8	-	-	1730.44 (20)	1730.44 (20)
Y _{11,1}	1766.25 (5)	1766.2 (2)	1766.25 (7)	1766.0 (2)	1766.2 (3)	1766.4 (15)	1767.7 (20)	1765.3 (5)	1765 (1)	1766.24 (5)	1766.24 (9)
Y _{23,2}	1774.5 (3)	1774.4 (7)	1774.2 (8)	1773.9 (5)	1775.0 (15)	-	-	-	-	1774.4 (3)	1774.44 (10)
Y _{24,3}	1784.1 (3)	1783.8 (7)	-	1784.5 (10)	1784.0 (20)	-	-	-	-	1784.1 (3)	1784.08 (9)
Y _{12,1}	1796.94 (9)	1796.9 (1)	1796.77 (7)	1796.6 (2)	1796.7 (3)	1797.0 (15)	1798.1 (20)	1795.7 (5)	1796 (1)	1796.83 (7)	1796.95 (5)
Y _{28,4}	1855.0 (2)	-	1854.8 (10)	1854.4 (5)	1854.8 (4)	-	-	-	-	1854.89 (20)	1854.89 (20)
Y _{26,2}	1909.3 (2)	1909.4 (2)	1909.3 (5)	1909.5 (5)	1909.4 (8)	-	~1909.2	-	-	1909.36 (20)	1909.28 (17)
Y _{13,1}	1927.22 (9)	1927.3 (2)	1926.96 (8)	1927.0 (3)	1927.2 (3)	1927.0 (15)	1927.8 (15)	1925.3 (5)	-	1927.09 (8)	1927.23 (7)
Y _{28,2}	1954.6 (4)	1954.9 (8)	-	1955.9 (10)	1956.9 (20)	-	-	-	-	1954.9 (4)	1954.9 (4)
Y _{14,1}	1973.5 (10)	1973.3 (8)	-	-	-	-	-	-	-	1973.4 (8)	1973.4 (8)
Y _{15,1}	1988.44 (8)	1988.6 (1)	1988.09 (7)	1988.1 (3)	1988.5 (3)	1989.0 (15)	1989.1 (15)	1986.3 (5)	-	1988.32 (23)	1988.44 (8)
Y _{30,2}	2093.3 (4)	2093.0 (4)	2092.5 (12)	2092.8 (10)	2094.1 (15)	-	-	-	-	2093.1 (4)	2093.33 (25)
Y _{16,1}	2112.54 (6)	2112.6 (2)	2112.1 (3) ^a	2112.4 (3)	2112.7 (2)	2113.6 (10)	2112.8 (20)	2111.0 (5)	-	2112.54 (6)	2112.52 (5)
Y _{35,3}	2185.7 (5)	-	-	-	-	-	-	-	-	2185.7 (5)	2185.7 (5)
Y _{17,1}	2193.2 (1)	2193.3 (2)	2192.7 (3) ^a	2193.0 (3)	2193.2 (6)	2194 (2)	2193.9 (22)	2189.3 (5)	-	2193.17 (10)	2193.17 (10)
Y _{10,0}	2242.4 (1)	2242.4 (2)	2242.0 (3) ^a	2242.4 (5)	2242.4 (8)	2243 (2)	2242.3 (20)	2241.4 (10)	-	2242.40 (10)	2242.45 (5)
Y _{19,1}	2271.9 (2)	2271.8 (3)	2271.5 (3)	2271.4 (5)	2271.7 (5)	2278 (3)	2271.9 (30)	-	-	2271.75 (20)	2271.86 (21)

	1982Ka10	1977Ok02	1975Hs02 ^a	1973Ma35	1969St03	1968Ha35	1967Fo09 ^b	1967Ra11 ^c	Evaluated	Recommended
Y _{20,1}	2309.0 (1)	2309.3 (2)	2308.5 (3) ^a	2308.4 (8)	2308.7 (4)	2310 (2)	2309.9 (20)	2307.8 (9)	2308.99 (10)	2309.09 (9)
Y _{21,1}	2316.4 (1)	2316.5 (2)	2315.9 (3) ^a	2316.6 (3)	2317.6 (5)	2317 (2)	2316.3 (30)	2314.8 (9)	2316.39 (10)	2316.41 (9)
Y _{22,1}	2366.04 (7)	2366.1 (2)	2365.6 (3) ^a	2365.8 (3)	2366.4 (3)	2366.4 (15)	2366.5 (20)	2366.9 (5)	2366.03 (7)	2366.04 (7)
Y _{23,1}	2390.6 (1)	2390.6 (2)	2390.0 (3) ^a	2390.7 (3)	2390.8 (3)	2389 (3)	2391.4 (20)	2390.3 (5)	2390.58 (10)	2390.60 (10)
Y _{24,1}	2405.96 (9)	2406.0 (2)	2405.5 (3) ^a	2406.0 (3)	2406.0 (3)	2406 (2)	2406.5 (20)	2406.6 (5)	2405.94 (9)	2405.98 (8)
Y _{13,0}	2439.1 (1)	2439.1 (2)	2438.6 (3) ^a	2439.0 (3)	2438.5 (3)	2439 (3)	2440.8 (20)	2437.3 (5)	2439.01 (10)	2439.07 (7)
Y _{25,1}	2456.8 (2)	2455.9 (15)	-	2457.3 (10)	-	-	-	-	2456.80 (20)	2456.79 (21)
Y _{14,0}	2484.6 (2)	2484.8 (3)	2484.0 (3)	2484.6 (5)	2485.7 (10)	-	2486.0 (20)	-	2484.54 (20)	2484.63 (20)
Y _{26,1}	2525.2 (6)	2525.2 (10)	2525.4 (10)	2525.4 (10)	2525.2 (20)	-	-	-	2525.3 (6)	2525.43 (17)
Y _{27,1}	2542.7 (1)	2542.8 (3)	2542.9 (3) ^a	2542.8 (3)	2543.4 (3)	2544 (2)	2544.3 (20)	2544.9 (6)	2542.79 (10)	2542.79 (10)
Y _{28,1}	2571.1 (2)	2571.2 (3)	2570.7 (3)	2571.2 (3)	2571.2 (3)	2573 (5)	2571.5 (30)	2571.4 (6)	2571.16 (20)	2571.16 (20)
Y _{29,1}	2651.4 (2)	2651.7 (3)	2651.0 (4)	2651.0 (5)	2651.6 (7)	2650 (5)	2653.1 (30) ^b	2650.4 (6)	2651.39 (20)	2651.39 (20)
Y _{17,0}	2705.1 (3)	2705.3 (3)	2705.1 (3)	2705.3 (5)	2706.6 (3)	-	-	-	2705.2 (3)	2705.26 (8)
Y _{30,1}	2709.5 (3)	2709.6 (3)	2708.7 (3) ^a	2709.2 (5)	2709.4 (3)	2708 (3)	2710.1 (15) ^b	2709.0 (5)	2709.3 (3)	2709.48 (25)
Y _{32,1}	2740.1 (4)	2740.5 (4)	-	2739.5 (10)	2740.5 (6)	-	-	-	2740.3 (4)	2740.1 (4)
Y _{34,1}	2787.3 (7)	2788.5 (5)	-	2788.5 (10)	-	-	-	-	2788.2 (5)	2788.2 (5)
Y _{35,1}	2809.0 (3)	2809.2 (3)	2808.4 (5)	2809.1 (5)	2809.5 (6)	-	2812.1 (18) ^b	2814.4 (8)	2809.1 (3)	2809.1 (3)
Y _{20,0}	2821.1 (3)	2821.3 (3)	2820.3 (3)	2821.0 (5)	2821.2 (4)	2825 (5)	2823.0 (18) ^b	2826.6 (8)	2821.2 (3)	2821.2 (3)
Y _{36,1}	-	2865.0 (10)	-	-	-	-	-	-	2865.0 (10)	2865.0 (10)
Y _{23,0}	2902.5 (8)	2902.7 (5)	-	2901.8 (10)	2904.1 (15)	-	-2898.7 ^b	-	2902.6 (5)	2902.6 (5)

	1982Ka10	1977Ok02	1975Hs02 ^a	1973Ma35	1969St03	1968Ha35	1967Fo09 ^b	1967Ra11 ^c	Evaluated	Recommended
Y _{24,0}	2917.9 (3)	2918.0 (3)	2917.1 (3)	2917.3 (5)	2917.9 (8)	2920 (5)	2919.7 (18) ^b	2918.0 (8)	2917.6 (3)	2917.6 (3)
Y _{26,0}	3037.4 (3)	3037.3 (3)	3036.5 (3)	3036.6 (5)	3036.8 (6)	3038 (2)	3039.7 (20) ^b	3042.0 (8)	3037.0 (3)	3037.3 (3)
Y _{27,0}	3055.0 (4)	3054.9 (3)	3053.5 (9)	3055.1 (5)	3055.6 (10)	-	3057.2 (30) ^b	3058.0 (10)	3055.0 (3)	3055.0 (3)
Y _{29,0}	3164.7 (10)	3165.4 (13)	-	3165.3 (25)	3161.5 (20)	-	-	-	3164.6 (10)	3164.6 (10)
Y _{31,0}	3249.8 (5)	3249.8 (5)	-	3251.8 (15)	-	-	~3252 ^b	-	3249.9 (5)	3249.8 (5)
Y _{33,0}	3273.4 (7)	3273.3 (5)	-	3275.5 (15)	-	-	-	-	3273.5 (5)	3273.4 (7)
Y _{36,0}	-	3375.9 (14)	-	-	-	-	-	-	3375.9 (14)	3375.9 (14)
Y _{37,0}	-	3401.8 (9)	-	-	-	-	-	-	3401.8 (9)	3401.8 (9)

^{a)} For data above 2000 keV, the uncertainty was increased to a minimum of 0.3 keV.
^{b)} Only data up to the last energy calibration point of 2614 keV were taken into account in evaluation.
^{c)} Data set not used as majority of values were rejected as outliers by Chauvenet's criterion.
^{d)} Data taken from Helmer and van der Leun (2000He14).

Table 2. Experimental and recommended relative gamma-ray emission probabilities.

E _γ (keV)	1982Ka10	1977Ok02	1975Hs02 ^a	1975De17	1973Ma35	1969St03	1969Od01	1968Ha35	1967Vr05	1967Ra11	1967Fo09	1966Ov01	1962Am03	1960Ro12	1960Se07	Recommended
428.49 (5)	3.46 (11)	3.6 (4)	3.12 (24)	-	3.48 (20)	4.5 (5)	-	1.4 (5)	-	-	-	-	-	-	-	3.43 (11)
434.23 (4)	0.99 (10)	0.9 (2)	1.18 (8)	-	0.77 (8)	2.3 (6)	-	-	-	-	-	-	-	-	-	0.97 (21)
439.23 (6)	0.62 (10)	0.46 (20)	-	-	0.50 (8) ^b	-	-	-	-	-	-	-	-	-	-	0.54 (8)
511.8534 (23)	1000	1000	1000 (65)	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000 (50)	1000 (40)	1000	1000
578.42 (6)	0.41 (4)	0.46 (3)	0.4 (3)	-	0.46 (4)	0.41 (4)	-	0.64 (2)	-	-	-	-	-	-	-	0.44 (3)
616.16 (3)	37.2 (33)	35.9 (72)	43 (6)	35.9 (7)	38 (2)	33.7 (27)	32 (2)	35 (7)	34 (4)	35 (3)	50 (10)	-	-	49 (13)	-	35.6 (7)
621.90 (4)	486.8 (60)	474 (24)	515 (38)	487 (6)	476 (23)	476 (15)	477 (5)	480 (10)	485 (10)	485 (10)	473 (6)	530	440 (50)	470 (30)	530 (20)	481 (5)
680.23 (6)	0.54 (3)	0.45 (6)	0.44 (8)	-	0.54 (5)	0.36 (6)	-	-	-	-	-	-	-	-	-	0.50 (3)
684.80 (6)	0.27 (1)	0.24 (4)	0.31 (22)	-	0.28 (3)	0.12 (3)	-	-	-	-	-	-	-	-	-	0.269 (10)
702.8 (10)	0.014 (9)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.014 (9)
715.86 (9)	0.49 (2)	0.48 (8)	0.34 (15)	-	0.46 (6)	0.75 (7) ^c	0.69 (20) ^c	0.80 (10) ^c	-	<0.9 ^c	-	-	-	-	-	0.484 (20)
717.44 (4)	0.32 (2)	0.33 (7)	0.34 (15)	-	0.39 (6)	-	-	-	-	-	-	-	-	-	-	0.327 (20)
751.26 (20)	0.053 (11)	0.044 (15)	0.06 (4)	-	0.12 (4)	0.09 (9)	-	0.12 (3)	-	-	-	-	-	-	-	0.059 (11)
873.46 (6)	21.5 (3)	20.9 (13)	23.7 (17)	20.9 (4)	25.0 (12)	20.2 (8)	23 (1)	19.0 (12)	20 (1)	21 (1)	18.0 (25)	-	17.1 (20)	18 (2)	35 (5)	21.2 (3)
942.63 (9)	0.028 (7)	0.026 (10) ^d	-	-	~0.05	-	-	0.12 (4)	-	-	-	-	-	-	-	0.029 (9)
1045.82 (4)	0.65 (8)	-	0.59 (18)	-	-	-	-	-	-	-	-	-	-	-	-	0.64 (8)
1050.39 (3)	76.4 (15)	72.7 (44)	84 (6)	71.1 (9)	73.5 (40)	70.6 (28)	73.5 (30)	72 (4)	80 (5)	75 (4)	63 (10)	-	-	68 (4)	90 (10)	72.6 (9)
1062.14 (6)	1.57 (2)	1.43 (4)	1.67 (14)	-	1.45 (10)	1.20 (9)	1.7 (6)	1.2 (2)	1.1 (2)	1.3 (1)	2.0 (3)	-	-	-	-	1.48 (9)
1108.71 (6)	0.29 (1)	0.24 (7)	0.26 (4)	-	0.22 (2)	0.25 (3)	-	-	-	-	-	-	-	-	-	0.273 (14)
1114.45 (6)	0.58 (1)	0.51 (9)	0.55 (5)	-	0.56 (5)	0.40 (6)	0.62 (25)	0.45 (10)	0.64	1.2 (4)	-	-	-	-	-	0.572 (13)

E _γ (keV)	1982Ka10	1977Ok02	1975Hs02 ^a	1975De17	1973Ma35	1969St03	1969Od01	1968Ha35	1967Vr05	1967Ra11	1967Fo09	1966Ov01	1960Ro12	Recommended
1128.01 (3)	19.8 (3)	19.2 (4)	22.0 (16)	18.6 (4)	19.3 (9)	18.7 (8)	21.5 (10)	19 (2)	20.6	20 (1)	16.0 (25)	-	24 (3)	19.4 (3)
1133.75 (4) ^e	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1150.08 (9)	0.15 (1)	0.132 (8)	0.17 (3)	-	0.14 (2)	<0.02	-	0.32 (10)	-	-	-	-	-	0.140 (8)
1159.90 (21)	0.008 (6)	-	-	-	-	0.02 (1)	-	-	-	-	-	-	-	0.011 (6)
1180.79 (6)	0.71 (1)	0.68 (4)	0.76 (6)	-	0.69 (4)	0.68 (7)	1.1 (3)	0.60 (5)	0.83	0.9 (1)	-	-	-	0.704 (10)
1194.58 (5)	2.81 (2)	2.64 (10)	3.11 (23)	-	2.60 (13)	2.56 (13)	2.7 (2)	2.6 (2)	2.9	2.8 (2)	2.3 (3)	-	-	2.791 (24)
1209.79 (8)	0.022 (5)	0.015 (4)	0.027 (14)	-	0.020 (4)	-	-	-	-	-	-	-	-	0.019 (4)
1258.71 (9)	0.028 (4)	0.036 (7)	-	-	0.038 (8)	0.048 (24)	-	-	-	-	-	-	-	0.032 (4)
1266.03 (9)	0.051 (5)	0.058 (8)	0.06 (3)	-	0.052 (10)	0.033 (16)	-	-	-	-	-	-	-	0.053 (5)
1305.33 (8)	0.065 (7)	0.051 (6)	0.040 (23)	-	0.043 (8)	0.14 (6)	-	-	-	-	-	-	-	0.053 (6)
1315.66 (8)	0.17 (1)	0.162 (12)	0.16 (3)	-	0.11 (1)	0.18 (6)	0.25 (10)	0.13 (10)	-	-	0.20 (6)	-	-	0.147 (23)
1355.60 (9)	0.031 (7)	0.017 (6)	-	-	0.058 (10)	0.031 (15)	-	-	-	-	-	-	-	0.029 (12)
1360.17 (9)	0.109 (6)	0.077 (6)	0.095 (24)	-	0.14 (2)	0.058 (10)	-	-	-	-	-	-	-	0.090 (19)
1372.28 (9)	0.101 (9)	0.091 (7)	0.137 (22)	-	0.10 (2)	-	-	0.27 (10)	-	-	-	-	-	0.097 (7)
1397.51 (16)	0.13 (1)	0.122 (10)	0.169 (23)	-	0.17 (2)	0.15 (2)	-	0.27 (10)	-	-	-	-	-	0.135 (10)
1489.61 (5)	0.06 (3)	0.087 (14)	0.090 (13)	-	0.19 (4)	0.08 (2)	-	0.2 (1)	0.15	-	-	-	-	0.090 (13)
1496.37 (6)	1.09 (3)	1.30 (8)	1.19 (9)	-	1.11 (8)	1.25 (6)	1.6 (1)	1.5 (5)	1.4	1.50 (15)	0.99 (10)	-	-	1.17 (8)
1498.73 (16)	0.33 (2)	-	0.37 (4)	-	0.25 (6)	-	-	-	-	-	-	-	-	0.331 (20)
1562.24 (3)	8.0 (1)	7.66 (38)	8.9 (7)	7.26 (10)	7.4 (4)	7.1 (3)	7.8 (4)	8.6 (10)	8	8.1 (5)	7.2 (8)	8.1 (3)	6.3 (5)	7.6 (4)
1572.47 (20)	0.09 (1)	0.083 (9)	0.086 (23)	-	0.13 (2)	-	-	-	-	-	-	-	-	0.090 (9)

E _γ (keV)	1982Ka10	1977Ok02	1975Hs02 ^a	1973Ma35	1969St03	1969Od01	1968Ha35	1967Vr05	1967Ra11	1967Fo09	1966Ov01	1960Ro12	1960Se07	Recommended
1577.27 (9)	0.052 (9)	0.042 (8)	0.056 (24)	0.06 (1)	0.08 (3)	-	-	-	-	-	-	-	-	0.051 (8)
1687.20 (10)	0.027 (6)	0.010 (8)	-	0.05 (1)	0.05 (2)	-	-	-	-	-	-	-	-	0.027 (8)
1693.2 (3)	0.032 (6)	0.035 (10)	-	0.058 (17)	-	0.10 (5)	0.06 (3)	-	-	0.084 (17)	-	-	-	0.040 (7)
1730.44 (20)	0.109 (6)	0.090 (9)	0.100 (17)	0.12 (1)	0.07 (2)	0.10 (5)	-	-	-	0.077 (15)	-	-	-	0.102 (6)
1766.24 (9)	1.68 (3)	1.34 (12)	1.62 (13)	1.20 (6)	1.13 (6)	1.3 (1)	1.4 (2)	1.5	1.3 (1)	1.1 (2)	2.6 (6)	1.9 (3)	2.0 (7)	1.46 (22)
1774.44 (10)	0.062 (11)	0.045 (4)	0.042 (22)	0.060 (18)	0.030 (15)	-	-	-	-	-	-	-	-	0.046 (4)
1784.08 (9)	0.021 (6)	0.0055 (15)	-	0.02 (1)	0.025 (15) ^f	-	-	-	-	-	-	-	-	0.021 (6)^g
1796.95 (5)	1.36 (2)	1.22 (7)	1.48 (11)	1.27 (6)	1.23 (7)	1.35 (10)	1.1 (2)	1.4	1.3 (1)	0.99 (20)	-	-	-	1.334 (22)
1854.89 (20)	0.061 (4)	-	0.043 (23)	0.12 (3)	0.04 (2)	-	-	-	-	-	-	-	-	0.061 (5)
1909.28 (17)	0.070 (5)	0.060 (5)	0.068 (22)	0.040 (4)	0.033 (8)	-	-	-	-	<0.058	-	-	-	0.052 (12)
1927.23 (7)	0.75 (2)	0.66 (4)	0.80 (6)	0.71 (11)	0.64 (5)	0.63 (6)	0.70 (15)	0.65	0.64 (6)	0.74 (8)	-	-	-	0.716 (20)
1954.9 (4)	0.011 (6)	0.009 (2)	-	0.015 (5)	-	-	-	-	-	-	-	-	-	0.009 9 (20)
1973.4 (8)	0.009 (5)	0.008 (2)	-	-	-	-	-	-	-	-	-	-	-	0.008 1 (20)
1988.44 (8)	1.28 (2)	1.18 (7)	1.41 (11)	1.23 (6)	1.07 (7)	1.0 (2)	1.3 (2)	1.2	1.2 (1)	1.25 (20)	1.4 (6)	1.0 (2)	-	1.255 (21)
2093.33 (25)	0.018 (4)	0.013 (3)	0.022 (16)	0.012 (3)	0.03 (3)	-	-	-	-	-	-	-	-	0.014 (3)
2112.52 (5)	1.71 (3)	1.69 (9)	1.95 (16)	1.75 (15)	1.56 (11)	1.7 (2)	1.7 (2)	1.6	1.7 (1)	1.8 (3)	1.5 (3)	1.3 (2)	3 (1)	1.71 (3)
2185.7 (5)	0.012 (3)	-	-	-	-	-	-	-	-	-	-	-	-	0.012 (3)
2193.17 (10)	0.24 (1)	0.229 (13)	0.26 (3)	0.25 (2)	0.27 (3)	0.5 (2)	0.24 (6)	0.4	0.23 (3)	0.26 (4)	-	-	-	0.241 (10)
2242.45 (5)	0.101 (3)	0.086 (6)	0.105 (12)	0.09 (1)	0.070 (15)	0.077 (10)	0.13 (2)	-	<0.3	0.083 (12)	-	-	-	0.095 (4)
2271.86 (21)	0.067 (4)	0.052 (4)	0.051 (11)	0.045 (5)	0.070 (14)	0.063 (10)	0.08 (3)	-	-	0.068 (10)	-	-	-	0.057 (4)

E _γ (keV)	1982Ka10	1977Ok02	1975Hs02 ^a	1973Ma35	1969St03	1969Od01	1968Ha35	1967Vr05	1967Ra11	1967Fo09	1960Ro12	1960Se07	Recommended
2309.09 (9)	0.276 (7)	0.252 (13)	0.306 (26)	0.28 (2)	0.29 (2)	0.35 (5)	0.30 (6)	0.3	0.29 (4)	0.30 (4)	-	-	0.280 (7)
2316.41 (9)	0.312 (7)	0.287 (15)	0.348 (27)	0.29 (3)	0.28 (2)	0.31 (5)	0.30 (6)	0.3	0.29 (4)	0.28 (3)	-	-	0.303 (7)
2366.04 (7)	1.14 (3)	1.08 (6)	1.25 (9)	1.12 (10)	1.04 (8)	1.1 (1)	1.4 (3)	0.95	1.1 (1)	1.35 (15)	1.7 (2)	-	1.13 (3)
2390.60 (10)	0.319 (7)	0.30 (2)	0.337 (27)	0.30 (3)	0.32 (3)	0.39 (4)	0.34 (8)	0.26	0.33 (4)	0.38 (4)	-	-	0.321 (7)
2405.98 (8)	0.71 (2)	0.68 (4)	0.81 (6)	0.60 (6)	0.60 (5)	0.78 (5)	0.72 (15)	0.58	0.70 (7)	0.85 (9)	-	-	0.705 (20)
2439.07 (7)	0.225 (6)	0.214 (11)	0.253 (25)	0.25 (2)	0.19 (2)	0.22 (5)	0.25 (8)	0.2	0.27 (3)	0.29 (3)	-	0.3 (1)	0.226 (6)
2456.79 (21)	0.014 (2)	0.010 (2)	-	0.006 (3)	-	-	-	-	-	-	-	-	0.010 9 (21)
2484.63 (20)	0.044 (3)	0.036 (4)	0.05 (1)	0.030 (3)	0.026 (6)	0.059 (15)	-	-	-	0.062 (15)	-	-	0.037 (7)
2525.43 (17)	0.010 (2)	0.0028 (14)	0.015 (8)	0.005 (2)	0.004 (4)	-	-	-	-	-	-	-	0.005 2 (15)
2542.79 (10)	0.145 (4)	0.139 (8)	0.158 (14)	0.14 (1)	0.12 (1)	0.10 (2)	0.15 (5)	0.14	0.12 (2)	0.17 (3)	-	-	0.141 (4)
2571.16 (20)	0.071 (3)	0.066 (4)	0.068 (12)	0.050 (5)	0.066 (7)	0.053 (10)	0.07 (3)	0.06	0.66 (15)	0.090 (13)	-	-	0.065 (3)
2651.39 (20)	0.034 (2)	0.032 (4)	0.030 (7)	0.030 (5)	0.033 (6)	-	~ 0.02	-	0.53 (8)	0.039 (15)	-	-	0.033 0 (20)
2705.26 (8)	0.123 (5)	0.125 (20)	0.137 (22)	0.10 (1)	0.15 (2)	-	-	-	-	-	-	-	0.121 (6)
2709.48 (25)	0.183 (5)	0.185 (20)	0.211 (24)	0.18 (2)	0.17 (2)	0.19 (5)	0.31 (9) ^h	0.2	0.35 (4) ^h	0.39 (5) ^h	0.3 (1) ^h	0.3 (1) ^h	0.182 (5)
2740.1 (4)	0.012 (2)	0.010 (2)	-	0.009 (2)	0.010 (3)	-	-	-	-	-	-	-	0.010 3 (20)
2788.2 (5)	0.0041 (20)	0.0039 (10)	-	0.004 (1)	-	-	-	-	-	-	-	-	0.004 0 (10)
2809.1 (3)	0.034 (2)	0.029 (2)	0.033 (8)	0.028 (3)	0.020 (4)	-	-	-	0.04 (1)	0.055 (16)	-	-	0.030 1 (20)
2821.2 (3)	0.059 (2)	0.057 (4)	0.074 (12)	0.060 (6)	0.051 (6)	0.12 (5)	0.05 (3)	0.045	0.07 (2)	0.070 (14)	-	-	0.058 6 (20)
2865.0 (10)	-	0.0007 (4)	-	-	-	-	-	-	-	-	-	-	0.000 7 (4)
2902.6 (5)	0.0032 (10)	0.0028 (10)	-	0.003 (1)	0.006 (2)	-	-	-	-	<0.0102	-	-	0.003 2 (10)

E _γ (keV)	1982Ka10	1977Ok02	1975Hs02 ^a	1973Ma35	1969St03	1968Ha35	1967Vr05	1967Ra11	1967Fo09	Recommended
2917.6 (3)	0.045 (2)	0.046 (5)	0.046 (7)	0.050 (5)	0.048 (5)	0.05 (2)	0.06	0.06 (2)	0.067 (13)	0.046 0 (20)
3037.0 (3)	0.050 (2)	0.053 (6)	0.065 (9)	0.052 (6)	0.048 (5)	0.06 (2)	0.05	0.07 (2)	0.076 (11)	0.051 4 (20)
3055.0 (3)	0.017 (2)	0.014 (2)	0.019 (9)	0.025 (3)	0.014 (7)	-	-	0.023 (6)	0.023 (7)	0.017 6 (20)
3164.6 (10)	0.0014 (7)	0.00029 (13)	-	~ 0.001	0.002 (2)	-	-	-	-	0.001 1 (6)^g
3249.8 (5)	0.0051 (13)	0.0039 (8)	-	0.004 (1)	-	-	-	-	<0.015	0.004 2 (8)
3273.4 (5)	0.0025 (11)	0.0020 (7)	-	0.003 (1)	-	-	-	-	-	0.002 4 (7)
3375.9 (14)	-	0.00055 (10)	-	-	-	-	-	-	-	0.000 55 (10)
3401.8 (9)	-	0.00061 (9)	-	-	-	-	-	-	-	0.000 61 (9)

^{a)} Uncertainty of 5.1 % added in quadrature to published data.

^{b)} Sum of 437.5 keV and 439.6 keV gamma-ray emission probabilities.

^{c)} Sum of 715.86 keV and 717.44 keV gamma-ray emission probabilities.

^{d)} Gamma ray only observed in coincidence spectra.

^{e)} E0 transition, no gamma ray observed.

^{f)} Sum of 1784.0 keV and 1789.0 keV gamma-ray emission probabilities.

^{g)} Recommended data was calculated using Rajeval Technique as the two most precise data are not in agreement.

^{h)} Sum of 2705.26 keV and 2709.48 keV gamma-ray emission probabilities.

¹⁰⁸Ag – Comments on evaluation of decay data by V. Chisté and M. M. Bé

The full decay data evaluation was completed in 2005. The literature available by January 2005 was included.

1. Decay Scheme

¹⁰⁸Ag disintegrates by electron capture (2,19 (14) %) and β^+ emission (0,283 (20) %) to excited states of ¹⁰⁸Pd and by β^- emission (97,53 (14) %) to excited states of ¹⁰⁸Cd .

2. Nuclear Data

The Q values are from the 2003Au03 evaluation.

Level energies, spin, parities and half -life of excited states are from J. Blachot (2000BI04, see also 1982Ha37).

The half-life of the ¹⁰⁸Ag ground state has been determined from the following data (in minutes):

1958Gu31	2,43 (5)
1960Wa10	2,42 (2)
1965Eb38	2,41 (2)
1971Jo07	2,38 (3)
1974HeYW	2,41 (1)
1974Ry01	2,37 (1)
1991Yamamoto	2,353 (9)
Adopted	2,382 (11)

The half-life weighted average has been calculated by Lweight program (version 3).

The evaluator has chosen to take into account the seven values with associated uncertainty for the calculation. The largest contributions to the weighted average come from values of Head (1974HeYW), Ryves (1974Ry01) and Yamamoto (1991Yamamoto) (25 %, 25 % and 31 %, respectively).

The weighted average value is 2,382 min with a reduced $-\chi^2$ value of 4,35. The external uncertainty is 0,011 min. Then, the adopted value is 2,382 (11) min.

2.1 β^- transition

The maximum energy of the β^- transitions in the decay of ¹⁰⁸Ag to excited states in ¹⁰⁸Cd is calculated from:

$$E_{\beta^-} = Q(\text{from 2003Au03}) - E_{\text{level in Cd-108}}(\text{from 2000BI04})$$

For the probabilities of the β^- transitions, the published data are (table 1):

Table 1: β^- transition measured intensity values in %.

Populated Level	1953Pe16	1960Wa10	1962Fr02	1965Fr01
β^- ¹⁰⁸ Cd ground state	97,3	93,8	95,0 (3)	95,9 (3)
β^- ¹⁰⁸ Cd 632 keV	0,8	1,9	1,73 (10)	1,75 (10)

For the β^- ¹⁰⁸Cd ground state transition, the values given by 1953Pe16 and 1960Wa10 have no uncertainties and the other two values are from the same author; the evaluators have chosen the most recent value published by L. Frevert (1965Fr01). This value, 95,9 (3) %, is important to determine the decay-scheme normalization factor (see **Gamma Ray Transition and Emission**).

For the β^- transition to the ¹⁰⁸Cd 632 keV level, the adopted value (1,63 (26) %), consistent with the Frevert value (1,75 (10) %) (table 1) has been deduced from the decay scheme balance.

The total β^- branching ratio was deduced taking into account that gamma -ray adopted relative emission intensities (see **4.1 Gamma Emissions**), the normalization factor (see **4.1 Gamma Emissions**) and the $I_{\beta^+, \epsilon}$ (g.s.) = 2,01 (12) % (see **2.3 Electron capture transition**):

$$I_{\beta^+, \epsilon} = I_{\beta^+, \epsilon}(\text{g.s.}) + N * [I_{\gamma}(433 \text{ keV}) + I_{\gamma}(931 \text{ keV}) + I_{\gamma}(1441 \text{ keV}) + I_{\gamma}(1539 \text{ keV})]$$

$$I_{\beta^+, \epsilon} = 2,01(12) \% + 0,0046(7) * [100 + 0,105(8) + 0,585(28) + 0,205(14)] = 2,47 (14)\%$$

$$\text{And } I_{\beta^-} = 100 - 2.47(14) \% = 97,53 (14) \%$$

The lg ft values have been calculated by Logft program (version 7.2a).

2.2 β^+ transition

The maximum energy of the β^+ transitions in the decay of ¹⁰⁸Ag is calculated by the same way as for the β^- transition.

For the probability of β^+ transition to the ground state, the published data are (table 2):

Table 2: β^+ transition probability measured values in %.

Level Populated	1953Pe16	1960Wa10	1962Fr02	1965Fr01
β^+ ¹⁰⁸ Pd ground state	0,14	0,36	0,28 (2)	0,28 (2)

From the total of 0,283 (20) % (2 transitions: to the 433 -keV level and to the ground state) β^+ transition decaying by this mode, 0,28 (2)%, measured by Frevert (1965Fr01) go directly to the ground state. Most of the remaining 0,0026 (3) % (2000Bl04 and 1982Ha37) populate the 433 -keV level (from theoretical ratio ϵ/β^+) (this electron -capture transition to the 433 -keV level hasn't been measured by Frevert (1965Fr01)).

2.3 Electron capture transition

Some values for the electron capture branching ratio (in %) have been found in the literature, as shown in the following table:

Populated Level	1953Pe16	1960Wa10	1962Fr02	1965Fr01
EC ¹⁰⁸ Pd ground state	1,5	3,35	2,49 (25)	1,73 (12)
EC ¹⁰⁸ Pd 433 keV level	0,06	0,18	0,19 (3)	0,19 (3)
EC ¹⁰⁸ Pd 1052 keV level	0,22	0,42	0,26 (3)	0,27 (3)

For the ground state, the adopted value is the most recent measurement of Frevert (1965Fr01). For the other levels, the electron-capture probabilities have been deduced from the imbalance at each level of the decay scheme. It can be noted that for the levels at 433 keV and 1052 keV the adopted electron capture branchings of 0,19 (8) % and 0,243 (39) %, respectively, are consistent with the Frevert measured values.

P_K , P_L , P_M values have been calculated for allowed electron -capture transitions in the decay of ¹⁰⁸Ag to the excited states in Pd-108 using the EC-Capture computer program.

2.4 Gamma transitions

Probabilities

The transitions probabilities have been calculated from the gamma -ray emission intensities and the internal conversion coefficients (see **Gamma ray emission**).

Multipolarity and internal conversion coefficients

For the 433 - ([E2]), 633 -(E2) and 1441 -keV ([E2]) gamma -ray transitions, multipolarities are from J. Blachot (2000Bl04, see also 1982Ha37)

The internal conversion coefficients (α_T , α_K and α_L) for these transitions have been calculated using the ICC Computer Code (program Icc99v3a – GETICC dialog). The adopted values have been interpolated from the new tables of Band (2002Ba85).

Their uncertainties are taken as 3% of the calculated values with the ICC computer code.

3. Atomic data

Atomic values, ω_K , ω_L and η_{KL} , are from Schönfeld (1996Sc33).

The X-ray and Auger electrons emission probabilities are calculated from the data set values by using the program EMISSION.

4. Photon Emissions

4.1 Gamma Emissions

The measured relative emission intensities are given in table 3, they are relative to the 433 -keV gamma ray taken as 100. Energy values are in keV.

Table 3: Measured relative gamma emission intensities in %.

Energy (keV)	Okano et al. (1971Ok01)	Singhal (1973Si02)	Adopted values
383,13 (16)	none	0,18 (6)	0,18 (6)
388,36 (7)	none	0,37 (12)	0,37 (12)
433,938 (5)	100	100	100
497,13 (12)	0,25 (9)	0,45 (11)	0,33 (7)
618,86 (5)	54,1 (24)	52,4 (26)	53,3 (18)
632,98 (5)	355,1 (14,9)	349,6 (175)	353 (11)
880,26 (10)	0,65 (3)	0,64 (5)	0,647 (26)
931,07 (12)	0,091(16)	0,11 (1)	0,105 (8)
1007,22 (5)	2,71 (11)	2,79 (14)	2,74 (9)
1106,01 (7)	0,26 (2)	0,33 (3)	0,282 (17)
1441,15 (5)	0,56 (4)	0,61 (4)	0,585 (28)
1539,94 (7)	0,20 (2)	0,21 (2)	0,205 (14)

The adopted values are the weighted averages of the two values given with uncertainties. One set of values, N. D. Johnson (1971Jo07), was not taken into account by the evaluator because the measured relative emission probabilities were relative to that of the 633 keV gamma ray and not to that of the 433 keV gamma ray as done by the other authors (normalization could introduce an overestimation of uncertainties).

The normalization factor to convert the relative emission intensities to absolute emission intensities is calculated with the formula:

$$\text{Normalization} = \frac{100 - I_{\beta^-}(\text{g.s.}) - I_{\beta^+,e}(\text{g.s.})}{(\sum(1 + a_T)P_{rel})}$$

where the sum is to be done over all the gamma transitions to the ground state, and:
 $I_{\beta^-}(\text{g.s.}) = 95,9$ (3) % and $I_{\beta^+,e}(\text{g.s.}) = 2,01$ (12) %. (see explanations above)

From the theoretical α_T and the evaluated relative emission intensities (table 3), the calculated normalization factor is 0,0046 (7). The uncertainties were propagated on the above formula. Absolute emission intensities are given in table 4.

Table 4: Absolute emission intensities for the γ -rays in the decay of the ¹⁰⁸Ag (in %).

Energy (keV)	Relative Emission intensity	Absolute emission intensity
383,13 (16)	0,18 (6)	0,00083 (30)
388,36 (7)	0,37 (12)	0,0017 (6)
433,938 (5)	100	0,46 (7)
497,13 (12)	0,33 (7)	0,00152 (40)
618,86 (5)	53,3 (18)	0,245 (39)
632,98 (5)	353 (11)	1,62 (26)
880,26 (10)	0,647 (26)	0,00298 (48)
931,07 (12)	0,105 (8)	0,00048 (8)
1007,22 (5)	2,74 (9)	0,0126 (20)
1106,01 (7)	0,282 (17)	0,00130 (22)
1441,15 (5)	0,585 (28)	0,00269 (44)
1539,94 (7)	0,205 (14)	0,00094 (16)

5. References

- 1953Pe16 – M. L. Perlman, W. Bernstein, R. B. Schwartz, Phys. Rev. 92(1953)1236 [Branching ratio].
 1958Gu31 – G. Gueben, Inst. Inter. Sci. Nucl. Monographie n°. 2 (1958) [T_{1/2}].
 1960Wa10 – M. A. Wahlgren, W. W. Meinke, Phys. Rev. 118(1960)181 [T_{1/2}].
 1962Fr02 – L. Frevert, Z. Phys. 169(1962)456 [P_β].
 1965Eb38 – T. B. Ebrey, P. R. Gray, Nucl. Phys. 61(1965)479 [T_{1/2}].
 1965Fr01 – L. Frevert, R. Schöneberg, A. Flammersfeld, Z. Phys. 182(1965)439 [P_β].
 1971Jo07 – N. D. Johnson, J. H. Hamilton, A. F. Fluk, N. R. Johnson, Z. Phys. 243(1971)395 [T_{1/2}, E_γ, I_γ].
 1971Ok01 – K. Okano, Y. Kawase, S. Uehara, T. Hayashi, Nucl. Phys. A164(1971)545 [E_γ, I_γ].
 1973Si02 – N. C. Singhal, N. R. Johnson, E. Eichler, Phys. Rev. C7(1973)774 [E_γ, I_γ].
 1974HeYW – R. L. Heath, ANCR-1000-2(1974) [T_{1/2}].
 1974Ry39 – T. B. Ryves, K. J. Zieba, J. Phys. (London) A7(1974)2318 [T_{1/2}].
 1982Ha37 – R. L. Haese, F. E. Bertrand, B. Harmatz, M. J. Martin, Nucl. Data Sheets 37(1982)289 [Energy level, multipolarity, spin, branching ratio].
 1991Yamamoto – H. Yamamoto, K. Kawade, T. Katoh, A. Hosoya, M. Shibata, A. Osa, T. Iida, A. Takahashi, Proc. of Int. Conf. ‘Nucl. Data for Science and Technology’ (1991), p. 565 [T_{1/2}].
 1996Sc33 – E. Schönfeld, H. Janssen, Nucl. Instrum. Meth. Phys. Res. A369 (1996)527 [Atomic data].
 2000Bl04 – J. Blachot, Nucl. Data Sheets 91(2000)135 [Energy level, multipolarity, spin, branching ratio].
 2002Ba85 – I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr., P. O. Tikkanen, S. Raman, Atomic Data and Nuclear Data Tables 81(2002)1 [α].
 2003Au03 – G. Audi, A. H. Wapstra, Nucl. Phys. A729(2003)129 [Q].

¹⁰⁸Ag^m – Comments on evaluation of decay data by V. Chisté and M. M. Bé

The full decay data evaluation was completed in 2005. The literature available by January 2005 was included.

1. Decay Scheme

¹⁰⁸Ag^m disintegrates 90.9 (6) % by electron capture to the 1771 keV excited state in Pd -108, and by 9.1(6)% through isomeric transitions (two gamma-rays in cascade) in ¹⁰⁸Ag.

2. Nuclear Data

The Q value (= 2031 (6) keV) is from the 2003Au03.

Level energies, spin and parities are from J. Blachot (2000Bl04).

The measured ¹⁰⁸Ag^m half-life values are, in years :

1969Ha07	127 ± 7
1969Vo06	310 ± 132
1992Sc25	418 ± 15
2004Sc49	438 ± 9

The evaluators have chosen as their recommended value the most recent result from Schrader (2004Sc49) who followed the decay by using a ionisation chamber for about 20 years.

2.1 Electron capture transition

For the 260 keV electron capture transition, the adopted value has been deduced from the decay -scheme balance at the 1771-keV level.

P_K, P_L, P_M have been calculated for allowed electron capture transition in the decay of ¹⁰⁸Ag^m to the 1771-keV excited state in Pd-108 using the EC Capture computer program.

2.4 Gamma transitions

Probabilities

The transition probabilities have been calculated using the gamma-ray emission intensities and the relevant internal conversion coefficients (see **Gamma ray Emission**)

Multipolarity and internal conversion coefficients

The multipolarities for the 30- (M4) and 79-keV gamma-ray transitions (E1) in ¹⁰⁸Ag, and the 433-([E2]), 614- (E2) and 722-keV (E2) gamma-ray transitions in ¹⁰⁸Pd have been taken from J. Blachot (2000Ba04, see also 1982Ha37).

The internal conversion coefficients (α_T , α_K and α_L) for these gamma-ray transitions have been interpolated from the tables of Band (2002Ba85) using the ICC Computer Code (program Icc99v3a – GETICC dialog). Their uncertainties are taken to be 3%.

3. Atomic data

Atomic values for ω_K , ω_L and η_{KL} , are from Schönfeld (1996Sc33).

The X-ray and Auger electron emission probabilities have been calculated from the data set values by using the program EMISSION.

4. Photon Emissions**4.1 Gamma-ray Emissions**

The energy of the 433-, 614- and 722-keV gamma-ray lines are from Helmer et al. (2000He14).

The measured relative emission intensities are given in table 1, they are relative to the 433-keV gamma ray taken as 100. Energy values are in keV.

Table 1: Measured relative emission intensities, in %.

Energy (keV)	Kistner (1966Ki03)	Kracíková (1968Kr23)	Hamilton (1971Ha31)	Heath (1974HeYW)	Weighted Average values
γ in ¹⁰⁸ Ag					
30.309 (8)	none	none	none	none	none
79.131 (3)	7.3 (8)	8.3 (9)	none	none	7.7 (6)
γ in ¹⁰⁸ Pd					
433.938 (4)	100	100	100	100 (5)	100
614.276 (4)	103 (3)	105 (10)	99.3 (20)	100 (5)	100.5 (16)
722.907 (10)	102 (2)	102 (10)	100.4 (20)	100 (5)	100.8 (16)

Adopted values are weighted averages (calculated by the Lweight program, version 3) of the four values measured with uncertainties. The normalization factor to convert the relative emission intensities to absolute emission intensities is calculated with the formula:

$$\text{Normalization} = \frac{100}{[(1 + a_T(433))P_{rel}(433)] + [(1 + a_T(79))P_{rel}(79)]}$$

where the 79- and 433-keV gamma-ray transitions populate the ground state level of ¹⁰⁸Ag and ¹⁰⁸Pd, respectively.

From the theoretical α_T and the relative evaluated emission intensities of the 79- and 433-keV gamma-rays (table 1), the normalization factor becomes 0.901 (6). The uncertainty was calculated through the propagation on the formula given above. Absolute emission intensities are given in table 2.

Table 2: Absolute emission intensities for the γ -rays, in %.

Energy (keV)	Relative Emission intensity	Absolute emission intensity
79.131 (3)	7.7 (6)	6.9 (5)
433.938 (4)	100	90.1 (6)
614.276 (4)	100.5 (16)	90.5 (16)
722.907 (10)	100.8 (16)	90.8 (16)

The 30-keV transition probability in the decay of $^{108}\text{Ag}^m \rightarrow ^{108}\text{Ag}$ is equal to 9.1 (6) % (from decay scheme transition probability balance).

Energy (keV)	Transition probability (%)	Absolute emission intensity (%)
30.309 (8)	9.1 (6)	0.0000215 (18)

The 30-keV absolute emission intensity has been deduced from the total transition probability and the theoretical α_T (Band *et al.*, 2002) for a M4 transition.

5. References

- 1960Wa07 – M. A. Wahlgren, W. W. Meinke, Phys. Rev. 118(1960)181 [Branching ratio].
 1965Jo38 – J. H. Hamilton, J. F. W. Jansen, P. F. A. Goudsmit, A. R. Sattler, Nucl. Phys. 61(1965)257 [Electron conversion].
 1966Ki03 – O. C. Kistner, A. W. Sunyar, Phys. Rev. 143(1966)918 [Emission probabilities].
 1968Jo04 – J. H. Hamilton, A. V. Ramayya, Bull. Am. Phys. Soc. 13(1968)249 [Energy values].
 1968Kr01 – T. I. Kracikova, B. Kracik, Czech. J. Phys. B18(1968)143 [Emission probabilities].
 1969Ha23 – G. Harbottle, Radiochim. Acta 13(1969)132 [Half-life].
 1969Vo06 – H. Vonach, M. Hille, Z. Phys. 227(1969)381 [Half-life].
 1971Ha31 – J. H. Hamilton, S. M. Brahmavar, J. B. Gupta, R. W. Lide, P. H. Stelson, Nucl. Phys. A172(1971)139 [Emission probabilities].
 1974HeYW – R. L. Heath, ANCR-1000-2(1974) [T_{1/2}].
 1975Mo09 – T. Morii, T. Saito, Nucl. Instrum. Meth. 131(1975)197 [Energy values].
 1982Ha37 – R. L. Haese, F. E. Bertrand, B. Harmatz, M. J. Martin, Nucl. Data Sheets 37(1982)289 [Energy level, multipolarity, spin, branching ratio].
 1982Ma11 – H. Maria, J. Dalmaso, G. Ardisson, Nucl. Instrum. Meth. 195(1982)621 [Energy value].
 1992Sc25 – U. Schötzig, H. Schrader, K. Debertin, Julich Conf., Nucl. Data for Science and Technology (1992)562 [Half-life].
 1996Sc33 – E. Schönfeld, H. Janssen, Nucl. Instrum. Meth. Phys. Res. A369 (1996)527 [Atomic data].
 2000Bi04 – J. Blachot, Nucl. Data Sheets 91(2000)135 [Energy level, multipolarity, spin, branching ratio].
 2000He14 – R. G. Helmer, C. van der Leun, Nucl. Instrum. Meth. Phys. Res. A450(2000)35 [Energy].
 2002Ba85 – I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor Jr., P. O. Tikkanen, S. Raman, Atomic Data and Nuclear Data Tables 81(2002)1 [α].
 2003Au03 – G. Audi, A. H. Wapstra, Nucl. Phys. A729(2003)129 [Q].
 2004Sc49 – H. Schrader, Appl. Rad. Isotopes 60(2004)317 [Half-life].

¹⁰⁹Pd - Comments on evaluation of decay data by A. L. Nichols

Evaluated: January 2007/March 2009

A.1. Evaluation Procedure

Limitation of Relative Statistical Weight Method (LWM) was applied to average the measured decay data when appropriate (see below).

Decay Scheme

A reasonably comprehensive decay scheme was constructed from the gamma-ray studies of 1968Gr02, 1968Be22, 1969Sc12, 1970Bo37, 1975El10 and 1978Pr08. Other earlier studies involved the use of low-resolution NaI(Tl) detectors, and these data have been set aside from consideration in this particular evaluation [1962Br15, 1962Ec02]. The gamma-ray emission probabilities were expressed in terms of the emission probability of the 647.3-keV gamma ray (100 %), and weighted mean data were derived as appropriate.

Most of the beta decay goes directly to the 88.034-keV metastable state of ¹⁰⁹Ag (half-life of 39.7 s). Hence, the resulting 88.03360-keV gamma ray dominates the decay scheme.

A.2. Nuclear Data

¹⁰⁹Pd undergoes beta decay to ¹⁰⁹Ag, including population of the 88.034-keV nuclear level (¹⁰⁹Ag^m, half-life of 39.7 s) that undergoes 100 % gamma decay to the stable ground state of ¹⁰⁹Ag.

Half-life (¹⁰⁹Pd)

The recommended half-life has been determined from the measurements of Gueben and Govaerts (1958Gu09), Starner (1959St28), Brandhorst and Cobble (1962Br15), Bormann *et al.* (1970Bo22), Gindler and Glendenin (1977Gi11), Chatterjee and Baliga (1983Ch42) and Abzouzi *et al.* (1990Ab06). A value of 13.58 (12) hours was derived in terms of LWM, with the uncertainty extended to include the most precise measurement of 13.7012 hours.

Half-life measurements (¹⁰⁹Pd)

Reference	Half-life (h)
1958Gu09	13.99 ± 0.16 13.20 ± 1.66
1959St28	13.45 ± 0.01
1962Br15	13.47 ± 0.01
1970Bo22	13.67 ± 0.07
1977Gi11	13.427 ± 0.014
1983Ch42	13.85 ± 0.17
1990Ab06	13.7012 ± 0.0024
Recommended value	13.58 ± 0.12

Half-life (¹⁰⁹Ag^m)

The recommended half-life has been determined from the measurements of Helmholtz (1941He03), Wiedenbeck (1945Wi11), Bradt *et al.* (1945Br06, 1946Br07, 1947Br05), Wolicki *et al.* (1951Wo15), Middelboe (1967Mi11), Abrams and Pelekis (1967Ab07), and Cottrell (1973Co10). A value of 39.7 (2) seconds was derived in terms of LWM.

Half-life measurements (¹⁰⁹Ag^m)

Reference	Half-life (s)
1941He03	40 ± 3
1945Wi11	40.4 ± 0.2
1947Br05	39.2 ± 0.2
1951Wo15	40 ± 1
1967Mi11	39.80 ± 0.18
1967Ab07	39.3 ± 0.3
1973Co10	35 ± 5
Recommended value	39.7 ± 0.2

Gamma RaysEnergies

Gamma-ray transition energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies of 2006Bi02 were adopted, and used to determine the energies of the gamma-ray transitions between the populated-depopulated levels, apart from the 88.03360-keV gamma ray which was adopted from 2000He14.

Emission Probabilities

Specific features of the earlier decay data studies of ¹⁰⁹Pd were considered and adopted to varying degrees during the course of assembling a reasonably consistent decay scheme (1953Av25, 1953Nu04, 1954Mo38, 1957Ma16, 1957Wa05, 1962Ec02, 1967Bi08, 1968BaZY and 1970Fo01). Relative emission probabilities and their uncertainties were determined from the measurements of 1968Gr02, 1968Be22, 1969Sc12, 1970Bo37, 1975Ei10, 1978Pr08 and 1983Ch42 normalized to the 647.3-keV gamma ray (100 %). These seven data sets were in reasonably good agreement with each other, although some difficulties occurred as noted below.

The 44.7-keV gamma ray has only been observed and quantified by 1968Gr02, 1968Be22 and 1970Bo37. Equivalent relative emission probabilities for the high-intensity 88.0336-keV gamma ray varied by as much as 25 %, and all of these measurements possess rather large uncertainties of ± 9 % to 10 % - not a particularly satisfactory situation for such an important gamma transition. A few gamma-ray emissions of questionable origin were noted by 1975Ei10 (in particular), 1978Pr08 and 1983Ch42. Thus, the 114.2-, 400.7-, 500.6-, 565.1- and 787.1-keV gamma rays have only been observed by 1975Ei10. The 327.2-, 395.6-, 609.8- and 869.5-keV gamma rays are also of doubtful origin.

Gamma-ray emission probabilities as measured and reported.

transition	E _γ (keV)	P_{γ}^{rel}								
		1962Ec02 [#]	1968BaZY	1968Be22	1968Gr02	1969Sc12	1970Bo37	1975E110	1978Pr08	1983Ch42
γ _{2,1} (Ag)	44.7 (1)	-	-	4.5 (14)	3.6 (11)	-	4.8 (5)	-	-	-
γ _{1,0} (Ag)	88.03360 (103)	4.35 (90)	-	11600 (1160)	8900 (800)	3850 (350)	11700 (1150)	11568 (995)	14600 (1300)	16252 (2194)
γ _{4,3} (Ag)	103.8 (2)	-	-	1.0 (3)	2.2 (7)	1.0 (2)	1.9 (2)	2.2 (4)	2.8 (5)	1.59
γ _{14,6} (Ag)	114.2 (9) ?	-	-	-	-	-	-	0.20 (6)	-	-
γ _{16,11} (Ag)	134.2 (2)	-	3.2	3.7 (11)	3.2 (10)	1.4 (3)	4.0 (4)	3.8 (4)	6.1 (9)	8.0 (13)
γ _{16,10} (Ag)	145.1 (2)	-	2.5	2.5 (8)	2.7 (8)	1.2 (2)	3.1 (3)	2.8 (3)	3.8 (8)	4.3 (18)
γ _{7,4} (Ag)	286.7 (3)	-	-	-	-	0.15 (4)	-	0.660 (72)	0.5 (1)	0.84 (11)
γ _{10,4} (Ag)	309.1 (3)	-	-	9 (1)	-	5.0 (15)	11.0 (9)	13.30 (14)	20 (6)	29.8 (36)
γ _{3,0} (Ag)	311.4 (1)	0.046 (9)	100	85 (9)	100	34 (3)	91 (8)	100	124 (6)	140.4 (152)
γ _{4,1} (Ag)	327.2 (2) ?	-	-	-	-	-	-	-	-	0.52 (5)
γ _{7,3} (Ag)	390.5 (2)	0.010 ?	2.3	3.0 (9)	2.5 (5)	1.0 (2)	3.2 (3)	3.2 (4)	3.6 (3)	3.6 (4)
γ _{9,3} (Ag)	395.6 (3)	-	-	-	-	0.07 (3)	-	0.60 (19)	0.27 (5)	0.46 (13)
γ _{23,6} (Ag)	400.7 (6) ?	-	-	-	-	-	-	0.20 (7)	-	-
γ _{10,3} (Ag)	413.0 (2)	0.016 (3)	47 (complex)	22 (2)	26 (8)	7 (1)	23 (2)	23.50 (23)	29 (2)	32.9 (37)
γ _{4,0} (Ag)	415.2 (2)	-	-	45 (5)	23 (7)	11.3 (10)	45.0 (42)	35.2 (3)	42 (3)	46.0 (50)
γ _{11,3} (Ag)	423.9 (2)	-	1.7	3.8 (11)	1.8 (4)	1.0 (2)	3.9 (3)	3.1 (3)	3.5 (3)	3.8 (4)
γ _{15,4} (Ag)	447.6 (4)	0.005	1.8	3.3 (10)	2.6 (6)	0.88 (20)	3.3 (3)	2.46 (30)	3.5 (3)	3.6 (4)
γ _{16,4} (Ag)	454.3 (3)	-	0.9	2.5 (8)	-	0.56 (25)	2.3 (2)	1.80 (23)	1.7 (2)	1.7 (2)
γ _{20,4} (Ag)	496.9 (10)	-	-	0.2 (1)	-	-	0.15 (3)	-	0.31 (6)	0.33 (9)
γ _{14,3} (Ag)	500.6 (6) ?	-	-	-	-	-	-	0.15 (3)	-	-
γ _{15,3} (Ag)	551.4 (3)	0.006 (1) ?	1.9	2.6 (8)	1.5 (5)	0.65 (15)	2.5 (3)	2.1 (3)	2.7 (2)	2.6 (3)
γ _{16,3} (Ag)	558.1 (2)	-	6.4	9.8 (10)	6.2 (8)	2.6 (3)	9.6 (9)	8.70 (95)	9.9 (7)	9.7 (10)
γ _{6,2} (Ag)	565.1 (5) ?	-	-	-	-	-	-	0.35 (4)	-	-
γ _{11,2} (Ag)	602.6 (2)	0.003	21	34 (3)	21.5 (20)	8.5 (5)	34 (3)	28.1 (20)	35 (2)	33.2 (37)
γ _{6,1} (Ag)	609.8 (4) ?	-	-	-	-	0.15 (7)	-	0.60 (15)	-	-
γ _{10,1} (Ag)	636.3 (1)	-	31	41 (4)	27 (3)	10.6 (5)	41.0 (38)	32.5 (3)	42 (3)	41.5 (45)
γ _{11,1} (Ag)	647.3 (1)	0.031 (6)	64	100	65 (5)	26 (2)	100	81.2 (80)	100	100
γ _{7,0} (Ag)	701.9 (2)	0.004 (1)	9.2	15 (2)	9.2 (10)	3.3 (3)	14.7 (12)	11.20 (12)	13.5 (18)	12.6 (14)

transition	E _γ (keV)	P _γ ^{rel}								
		1962Ec02 [#]	1968BaZY	1968Be22	1968Gr02	1969Sc12	1970Bo37	1975El10	1978Pr08	1983Ch42
γ _{9.0} (Ag)	707.0 (2)	-	3.8	6.9 (7)	4.5 (5)	1.7 (2)	7.1 (7)	5.8 (6)	6.3 (9)	5.6 (7)
γ _{10.0} (Ag)	724.4 (1)	-	-	1.2 (4)	-	0.20 (5)	1.1 (1)	0.8 (2)	0.4 (1)	0.27 (7)
γ _{16.2} (Ag)	736.7 (2)	-	4.4	7.8 (8)	5.0 (6)	1.8 (2)	7.7 (7)	6.1 (7)	6.8 (9)	6.4 (7)
γ _{19.2} (Ag)	778.3 (5)	-	-	4 (1)	-	1.6 (5)	4.0 (4)	4.7 (6)	7.3 (25)	9.6 (13)
γ _{16.1} (Ag)	781.4 (1)	0.010 (2)	34	49 (5)	33 (3)	11.7 (12)	50 (4)	40.0 (35)	48 (3)	50.5 (56)
γ _{23.3} (Ag)	787.1 (3) ?	-	-	-	-	-	-	0.070 (4)	-	-
γ _{19.1} (Ag)	823.0 (4)	-	0.5	0.8 (2)	-	0.20 (3)	0.5 (1)	1.2 (3)	0.77 (11)	0.66 (8)
γ _{15.0} (Ag)	862.8 (2)	-	0.4	0.6 (2)	< 0.5	0.14 (3)	0.3 (1)	0.40 (15)	0.66 (11)	0.68
γ _{16.0} (Ag)	869.5 (1) ?	-	-	-	-	-	-	-	0.21 (6) ?	-
γ _{23.2} (Ag)	965.8 (3)	-	-	0.10 (3)	-	-	< 0.1	0.3 (1)	0.25 (4)	0.28
γ _{23.1} (Ag)	1010.5 (2)	-	-	0.10 (3)	-	-	< 0.1	0.200 (66)	0.11 (4)	0.12 (4)

[#] NaI(Tl) detectors were used with a lack of spectral resolution – this data set was discarded.

Relative gamma-ray emission probabilities – re-normalised for weighted mean analysis.

E _γ (keV)	P _γ ^{rel}								
	1968BaZY [§]	1968Gr02	1968Be22	1969Sc12	1970Bo37	1975El10	1978Pr08	1983Ch42	Recommended
44.7 (1)	-	5.5 (17)	4.5 (14)	-	4.8 (5)	-	-	-	4.8 (5)
88.03360 (103)	-	13690 (1230)*	11600 (1160)*	14810 (1350)	11700 (1150)*	14250 (1230)	14600 (1300)	16252 (2194)*	14540 (750)
103.8 (2)	-	3.4 (11)*	1.0 (3)*	3.8 (8)*	1.9 (2)*	2.7 (5)	2.8 (5)	1.5*	2.8 (4)
114.2 (9) ?	-	-	-	-	-	0.25 (8)	-	-	0.25 (8)
134.2 (2)	5	4.9 (15)	3.7 (11)*	5.4 (12)	4.0 (4)	4.7 (5)	6.1 (9)*	8.0 (13)*	4.4 (3)
145.1 (2)	3.9	4.2 (12)	2.5 (8)	4.6 (8)	3.1 (3)	3.4 (4)	3.8 (8)	4.3 (18)	3.3 (2)
286.7 (3)	-	-	-	0.58 (15)	-	0.81 (9)	0.5 (1)	0.84 (11)	0.70 (5)
309.1 (3)	-	-	9 (1)*	19 (6)	11.0 (9)*	16.4 (2)	20 (6)	29.8 (36)*	16.4 (2)
311.4 (1)	156	154 (12)*	85 (9)*	131 (12)	91 (8)*	123 (12)	124 (6)	140.4 (152)*	125 (5)
327.2 (2) ?	-	-	-	-	-	-	-	0.52 (5)	0.52 (5)
390.5 (2)	3.6	3.8 (8)	3.0 (9)*	3.8 (8)	3.2 (3)*	3.9 (5)	3.6 (3)	3.6 (4)	3.7 (2)
395.6 (3)	-	-	-	0.27 (12)	-	0.74 (23)*	0.27 (5)	0.46 (13)*	0.27 (5)

E _γ (keV)	P _γ ^{rel}								
	1968BaZY [§]	1968Gr02	1968Be 22	1969Sc12	1970Bo37	1975El10	1978Pr08	1983Ch42	Recommended
400.7 (6) ?	-	-	-	-	-	0.25 (9)	-	-	0.25 (9)
413.0 (2)	73(complex)	40 (12)*	22 (2)	27 (4)	23 (2)	28.94 (28)	29 (2)	32.9 (37)	27 (2)
415.2 (2)		35 (11)	45 (5)	43 (4)	45.0 (42)	43.3 (4)	42 (3)	46.0 (50)	43.3 (4)
423.9 (2)	2.7	2.8 (6)*	3.8 (11)	3.8 (8)	3.9 (3)	3.8 (4)	3.5 (3)	3.8 (4)	3.7 (2)
447.6 (4)	2.8	4.0 (9)	3.3 (10)	3.4 (8)	3.3 (3)	3.03 (37)	3.5 (3)	3.6 (4)	3.4 (2)
454.3 (3)	1.4	-	2.5 (8)	2.2 (10)	2.3 (2)	2.22 (28)	1.7 (2)	1.7 (2)	2.0 (1)
496.9 (10)	-	-	0.2 (1)	-	0.15 (3)*	-	0.31 (6)	0.33 (9)	0.29 (5)
500.6 (6) ?	-	-	-	-	-	0.18 (4)	-	-	0.18 (4)
551.4 (3)	3.0	2.3 (8)	2.6 (8)	2.5 (6)	2.5 (3)	2.6 (4)	2.7 (2)	2.6 (3)	2.6 (2)
558.1 (2)	10	9.5 (12)	9.8 (10)	10 (1)	9.6 (9)	10.7 (12)	9.9 (7)	9.7 (10)	9.9 (4)
565.1 (1) ?	-	-	-	-	-	0.43 (5)	-	-	0.43 (5)
602.6 (2)	33	33.0 (31)	34 (3)	33 (2)	34 (3)	34.6 (25)	35 (2)	33.2 (37)	34 (1)
609.8 (4) ?	-	-	-	0.6 (3)	-	0.74 (19)	-	-	0.7 (2)
636.3 (1)	48	42 (5)	41 (4)	41 (2)	41.0 (38)	40.0 (4)	42 (3)	41.5 (45)	40.1 (4)
647.3 (1)	100	100	100	100	100	100	100	100	100
701.9 (2)	14.4	14.2 (15)	15 (2)	12.7 (12)	14.7 (12)	13.79 (14)	13.5 (18)	12.6 (14)	13.8 (2)
707.0 (2)	5.9	6.9 (8)	6.9 (7)	6.5 (8)	7.1 (7)	7.1 (7)	6.3 (9)	5.6 (7)*	6.8 (3)
724.4 (1)	-	-	1.2 (4)	0.8 (2)	1.1 (1)	1.0 (3)	0.4 (1)*	0.27 (7)*	1.0 (1)
736.7 (2)	6.9	7.7 (9)	7.8 (8)	6.9 (8)	7.7 (7)	7.5 (9)	6.8 (9)	6.4 (7)	7.2 (3)
778.3 (5)	-	-	4 (1)*	6.2 (19)	4.0 (4)*	5.8 (7)	7.3 (25)	9.6 (13)*	5.9 (6)
781.4 (1)	53	51 (5)	49 (5)	45 (5)	50 (4)	49 (4)	48 (3)	50.5 (56)	49 (2)
787.1 (3) ?	-	-	-	-	-	0.086 (5)	-	-	0.086 (5)
823.0 (4)	0.8	-	0.8 (2)	0.77 (12)	0.5 (1)*	1.5 (4)*	0.77 (11)	0.66 (8)	0.72 (6)
862.8 (2)	0.6	< 0.8	0.6 (2)	0.54 (12)	0.3 (1)*	0.49 (18)	0.66 (11)	0.68*	0.59 (7)
869.5 (1) ?	-	-	-	-	-	-	0.21 (6) ?	-	0.21 (6)
965.8 (3)	-	-	0.10 (3)*	-	< 0.1	0.4 (1)	0.25 (4)	0.28*	0.27 (4)
1010.5 (2)	-	-	0.10 (3)	-	< 0.1	0.25 (8)	0.11 (4)	0.12 (4)	0.12 (2)

[§] Uncertainties were not assigned to the intensity measurements – this data set was discarded.

* Data were not used in the weighted mean analysis process (LWM) – some of these data lack quantified uncertainties, while other data deviate considerably from the majority of equivalent data from other sources.

Gamma-ray emissions: recommended energies, relative emission probabilities, multiplicities and theoretical internal conversion coefficients (frozen orbital approximation).

E_γ (keV)	P_γ^{rel}	Multipolarity	α_K	α_L	α_{M+}	α_{tot}
44.7 (1)	4.8 (5)	M1 + E2 $\delta = 0.533$	5.69 (9)	2.69 (5)	0.62 (2)	9.00 (15)
88.033 60 (103)	14 540 (750)	E3	11.41 (16)	12.06 (17)	2.86 (4)	26.33 (40)
103.8 (2)	2.8 (4)	M1 + E2 $\delta = -0.045$	0.329 (6)	0.041 (1)	0.009	0.379 (7)
114.2 (9) ?	0.25 (8)	(M1 + E2)	-	-	-	-
134.2 (2)	4.4 (3)	M1 + E2 ($\delta = 0.15$)	0.165 8 (25)	0.021 2 (4)	0.005 (1)	0.192 (3)
145.1 (2)	3.3 (2)	(M1 + E2) $\delta = 0.132$	0.132 6 (20)	0.016 70 (25)	0.003 7	0.153 (2)
286.7 (3)	0.70 (5)	M1 + E2 ($\delta = 0.199$)	0.021 6 (3)	0.002 64 (4)	0.000 56	0.024 8 (4)
309.1 (3)	16.4 (2)	(E1)	0.005 91 (9)	0.000 697 (10)	0.000 163	0.006 77 (10)
311.4 (1)	125 (5)	M1 + E2 $\delta = -0.22$	0.017 49 (25)	0.002 13 (3)	0.000 48	0.020 1 (3)
327.2 (2) ?	0.52 (5)	E1	0.005 09 (8)	0.000 599 (9)	0.000 131	0.005 82 (9)
390.5 (2)	3.7 (2)	M1 + E2 $\delta = 0.19$	0.009 80 (14)	0.001 178 (17)	0.000 262	0.011 24 (16)
395.6 (3)	0.27 (5)	(E1)	0.003 12 (5)	0.000 366 (6)	0.000 084	0.003 57 (5)
400.7 (6) ?	0.25 (9)	(M1 + E2)	-	-	-	-
413.0 (2)	27 (2)	(E1 (+ M2)) $\delta = 0.18$	0.003 66 (7)	0.000 442 (8)	0.000 098	0.004 20 (8)
415.2 (2)	43.3 (4)	E2	0.009 44 (14)	0.001 257 (18)	0.000 283	0.010 98 (16)
423.9 (2)	3.7 (2)	E1 (+ M2) $\delta = -0.27$	0.004 36 (7)	0.000 536 (9)	0.000 124	0.005 02 (8)
447.6 (4)	3.4 (2)	M1 + E2 $\delta = -0.16$	0.006 98 (10)	0.000 833 (12)	0.000 187	0.008 00 (12)
454.3 (3)	2.0 (1)	E1	0.002 22 (4)	0.000 259 (4)	0.000 051	0.002 53 (4)
496.9 (10)	0.29 (5)	M1 + E2 ($\delta = 0.20$)	0.005 41 (8)	0.000 644 (10)	0.000 146	0.006 2 (1)
500.6 (6) ?	0.18 (4)	(E1)	0.001 756 (25)	0.000 205 (3)	0.000 049	0.002 01 (3)
551.4 (3)	2.6 (2)	M1 + E2 $\delta = -0.28$	0.004 20 (6)	0.000 500 (7)	0.000 12	0.004 82 (7)
558.1 (2)	9.9 (4)	E1 (+ M2) $\delta = -0.26$	0.002 07 (4)	0.000 249 (4)	0.000 061	0.002 38 (4)
565.1 (5) ?	0.43 (5)	(E2)	0.003 86 (6)	0.000 489 (7)	0.000 111	0.004 46 (7)
602.6 (2)	34 (1)	E2	0.003 24 (5)	0.000 407 (6)	0.000 093	0.003 74 (6)
609.8 (4) ?	0.7 (2)	(M1 + E2)	-	-	-	-
636.3 (1)	40.1 (4)	(E2)	0.002 81 (4)	0.000 350 (5)	0.000 07	0.003 23 (5)
647.3 (1)	100	M1 + E2	-	-	-	-
701.9 (2)	13.8 (2)	M1 + E2 $\delta = 0.029$	0.002 39 (4)	0.000 280 (4)	0.000 06	0.002 73 (4)
707.0 (2)	6.8 (3)	(E1)	0.000 807 (12)	0.000 093 3 (13)	0.000 020 7	0.000 921 (13)
724.4 (1)	1.0 (1)	(E1)	0.000 766 (11)	0.000 088 5 (13)	0.000 019 5	0.000 874 (13)
736.7 (2)	7.2 (3)	E2	0.001 93 (3)	0.000 236 (4)	0.000 044	0.002 21 (4)
778.3 (5)	5.9 (6)	M1 + E2	-	-	-	-
781.4 (1)	49 (2)	M1 + E2	-	-	-	-
787.1 (3) ?	0.086 (5)	(E1)	0.000 644 (9)	0.000 074 3 (11)	0.000 016 7	0.000 735 (11)
823.0 (4)	0.72 (6)	M1 + E2	-	-	-	-
862.8 (2)	0.59 (7)	E2	0.001 313 (19)	0.000 158 3 (23)	0.000 038 7	0.001 51 (2)
869.5 (1) ?	0.21 (6)	M2 (+ E3)	0.003 72 (6)	0.000 453 (7)	0.000 097	0.004 27 (6)
965.8 (3)	0.27 (4)	-	-	-	-	-
1010.5 (2)	0.12 (2)	-	-	-	-	-

Much of the lower-energy gamma-ray data of 1968Gr02, 1968Be22, 1970Bo37 and 1983Ch42 deviated significantly from the studies of 1969Sc12, 1975E110 and 1978Pr08 (particularly below 400 keV) and after careful consideration of the individual data sets, some of these measurements were set aside and not included in the eventual weighted mean analyses. Despite these problems, every effort has been made to incorporate all of the gamma-ray data within a reasonably comprehensive decay scheme. One result of this effort is the introduction of two relatively poorly defined nuclear levels at 697.8 (5/2+) and 812.0 (3/2+) keV, primarily to accommodate the 114.2-, 500.6-, 565.1- and 609.8-keV gamma rays. Additional low-intensity gamma transitions were also incorporated into the proposed decay scheme, including the 327.2-, 400.7-, 787.1- and 869.5-keV gamma rays.

Multipolarities and Internal Conversion Coefficients

The nuclear level scheme specified by Blachot (2006B102) has been used to define the multipolarities of the gamma transitions on the basis of known spins and parities. Somewhat disparate mixing ratios were obtained by 1970Ro14, 1975E110, 1977Bo04 and 1978Pr08 based on angular correlation measurements, and these data were used to determine the assignments and internal conversion coefficients of the 103.8-, 145.1-, 311.4-, 390.5-, 413.0-, 423.9-, 447.6-, 551.4-, 558.1- and 701.9-keV gamma rays. Recommended internal conversion coefficients were determined from the theoretical tabulations of Band *et al.* (2002Ba85, 2002Ra45) by means of the methodology of Kibedi *et al.* (2008Ki07) in which the frozen orbital approximation was adopted. Finally, the theoretical internal conversion coefficients and mixing ratio of the 44.7-keV (M1 + E2) gamma transition were derived from the population-depopulation balance of the 132.74-keV nuclear level (with no populating beta transition).

A normalization factor of 0.000 252 (14) was calculated from the internal conversion coefficients and relative emission probabilities of the gamma-ray transitions populating the ¹⁰⁹Ag ground state directly, assuming that there is no direct beta feeding as implied from the spins and parities derived for the ¹⁰⁹Pd (5/2+) and ¹⁰⁹Ag (1/2-) ground states:

$$\sum P_{\gamma+ce}^{rel} = 100\%$$

$$397\ 572\ (21\ 307)\ F = 100$$

$$F = 0.000\ 251\ 53 \pm 0.000\ 013\ 51\ [= 0.000\ 252 \pm 0.000\ 014]$$

Beta-particle Emissions

Energies and emission probabilities

The beta-particle energies were calculated from the structural detail of the proposed decay scheme. Nuclear level energies adopted from Blachot (2006B102) and a Q_{β^-} value of 1116.1 ± 2.0 keV from Audi *et al.* (2003Au03) were used to determine the energies and uncertainties of the beta-particle transitions. Beta-particle emission probabilities were calculated from the relative gamma-ray emission probabilities, the associated normalization factor and the theoretical internal conversion coefficients derived from Kibedi *et al.* (2008Ki07). Direct beta population of the 132.74-keV nuclear level and ground state of ¹⁰⁹Ag were assumed to be zero on the basis of spin and parity considerations (5/2+ to 9/2+ (2nd forbidden non-unique), and 5/2+ to 1/2- (1st forbidden unique), respectively).

Beta-particle Emission Probability per 100 Disintegrations of ¹⁰⁹Pd.

Transition	E _β (keV)	P _β	Transition type	logft
$\beta_{0,23}^-$	17.6 ± 2.0	0.000 18 ± 0.000 03	(allowed)	6.22
$\beta_{0,20}^-$	204.0 ± 2.2	0.000 074 ± 0.000 014	1 st forbidden non-unique	9.87
$\beta_{0,19}^-$	205.1 ± 2.0	0.001 66 ± 0.000 17	allowed	8.53
$\beta_{0,16}^-$	246.6 ± 2.0	0.019 4 ± 0.000 9	allowed	7.72
$\beta_{0,15}^-$	253.3 ± 2.0	0.001 67 ± 0.000 10	1 st forbidden non-unique	8.82
$\beta_{0,14}^-$	304.1 ± 2.1	0.000 108 ± 0.000 024	(allowed)	10.3
$\beta_{0,11}^-$	380.8 ± 2.0	0.033 4 ± 0.001 5	allowed	8.096
$\beta_{0,10}^-$	391.8 ± 2.0	0.020 4 ± 0.000 9	(allowed)	8.351
$\beta_{0,9}^-$	409.1 ± 2.0	0.001 78 ± 0.000 12	(allowed)	9.47
$\beta_{0,7}^-$	414.2 ± 2.0	0.004 60 ± 0.000 21	1 st forbidden non-unique	9.08
$\beta_{0,6}^-$	418.3 ± 2.0	0.000 16 ± 0.000 07	(allowed)	10.55
$\beta_{0,4}^-$	700.9 ± 2.0	0.006 3 ± 0.000 2	1 st forbidden non-unique	9.73
$\beta_{0,3}^-$	804.7 ± 2.0	0.019 1 ± 0.002 2	1 st forbidden non-unique	9.46
$\beta_{0,1}^-$	1028.1 ± 2.0	99.891 ± 0.003	allowed	6.134 (5)

Σ 99.999 832

A.3. Atomic Data

The X-ray and Auger electron data have been calculated using the evaluated gamma-ray data, and the atomic data from 1996Sc06, 1998ScZM and 1999ScZX.

References

- 1940A101 L.W. ALVAREZ, A.C. HELMHOLZ, E. NELSON, Isomeric silver and the Weizsäcker theory, Phys. Rev. 57 (1940) 660-661. [¹⁰⁹Ag^m half-life]
- 1941He03 A.C. HELMHOLZ, Long-lived radioactive Cd from deuteron bombardment of Ag, Phys. Rev. 60 (1941) 160, 11. [¹⁰⁹Ag^m half-life]
- 1945Br06 H. BRADT, P.C. GUGELOT, O. HUBER, H. MEDICUS, P. PREISWERK, P. SCHERRER, Die metastabilen Zustände der Silberkerne Ag¹⁰⁷ und Ag¹⁰⁹, Helv. Phys. Acta 18 (1945) 256-258. [¹⁰⁹Ag^m half-life]
- 1945Wi11 M.L. WIEDENBECK, The nuclear excitation of silver and cadmium, Phys. Rev. 67 (1945) 92-97. [¹⁰⁹Ag^m half-life]
- 1946Br07 H. BRADT, P.C. GUGELOT, O. HUBER, H. MEDICUS, P. PREISWERK, P. SCHERRER, R. STEFFEN, Die Silberkerne Ag¹⁰⁷ und Ag¹⁰⁹, Helv. Phys. Acta 19 (1946) 218-219. [¹⁰⁹Ag^m half-life]
- 1947Br05 H. BRADT, P.C. GUGELOT, O. HUBER, H. MEDICUS, P. PREISWERK, P. SCHERRER, R. STEFFEN, Die metastabilen Zustände der Silberkerne Ag¹⁰⁷ und Ag¹⁰⁹, Helv. Phys. Acta 20 (1947) 153-165. [¹⁰⁹Ag^m half-life, ICC]

- 1951Wo15 E.J. WOLICKI, B. WALDMAN, W.C. MILLER, The nuclear excitation of Ag¹⁰⁷ and Ag¹⁰⁹ by X-rays, Phys. Rev. 82 (1951) 486-488. [¹⁰⁹Ag^m half-life]
- 1953Av25 P. AVIGNON, État isomérique de ¹⁰⁹Ag, J. Phys. Radium 14 (1953) 636-637. [α_K (88 keV)]
- 1953Nu04 R.H. NUSSBAUM, A.H. WAPSTRA, N.F. VERSTER, H. CERFONTAIN, On the radioactive decay of ¹¹²Pd and ¹¹²Ag, Physica, 19 (1953) 385-390. [α_K (88 keV)]
- 1954Mo38 J. MOREAU, Étude de la radioactivité de ¹⁰⁹₄₆Pd, J. Phys. Radium 15 (1954) 380-381. [ICC ratios (88 keV)]
- 1957Ma16 R.L. MACKLIN, N.H. LAZAR, W.S. LYON, Neutron activation cross sections with Sb-Be neutrons, Phys. Rev. 107 (1957) 504-508. [P_γ (88-keV)]
- 1957Wa05 A.H. WAPSTRA, W. VAN DER EIJK, The decay of ¹⁰⁹Cd, ¹⁰⁹Ag and ¹⁰⁹Pd, Nucl. Phys. 4 (1957) 325-329; Erratum, Nucl. Phys. 4 (1957) 695. [α_K (88 keV)]
- 1958Gu09 G. GUEBEN, J. GOVAERTS, La méthode d'analyse par activation en utilisant les neutrons d'une source Ra-Be, Inst. Interuniv. Sci. Nucléaires, Bruxelles, Monographie No. 2 (1958). [¹⁰⁹Pd half-life]
- 1959St28 J.W. STARNER, Decay of Pd¹⁰⁹ and Pd^{109m}, Bull. Am. Phys. Soc. 4, No. 2 (1959) 99, L2. [¹⁰⁹Pd half-life, E γ , P γ]
- 1962Br15 H.W. BRANDHORST, Jr., J.W. COBBLE, Decay of Ru¹⁰⁵, Pd¹⁰⁹, and Rb⁸⁶, Phys. Rev. 125 (1962) 1323-1328. [¹⁰⁹Pd half-life, E γ , P γ , P β]
- 1962Ec02 S.F. ECCLES, Gamma ray spectroscopy of ¹⁰⁷Cd, ¹⁰⁹Pd and ¹¹¹Pd, Physica 28 (1962) 251-261. [E γ , P γ]
- 1967Mi11 V. MIDDELBOE, Some accurate half-life determinations, Mat. Fys. Medd. Dan. Vid. Selsk. 35, No. 8 (1966). [¹⁰⁹Ag^m half-life]
- 1967Ab07 I.A. ABRAMS, L.L. PELEKIS, Excitation of metastable levels in nuclei by the γ rays of ^{116m}In, Program and Theses, Proc. 17th All-Union Conf. Nucl. Spectrosc. Struct. At. Nuclei, Kharkov (1967) 30. [¹⁰⁹Ag^m half-life]
- 1967BI08 J.L. BLACK, W. GRUHLE, The excited states of ¹⁰⁷Ag and ¹⁰⁹Ag, Nucl. Phys. A93 (1967) 1-30. [spin, parity, γ - γ coincidence]
- 1968BaZY W.E. BARNES, H.T. EASTERDAY, Decay of Pd¹⁰⁹, Oregon State University Progress Report RLO-1062-681 (1968) 8-10. [E γ , P γ]
- 1968Be22 G. BERZINS, M.E. BUNKER, J.W. STARNER, Ge(Li)-Ge(Li) coincidence studies of the decay of ¹⁰⁹Pd, Nucl. Phys. A114 (1968) 512-528. [E γ , P γ]
- 1968Gr02 G. GRAEFFE, G.E. GORDON, Decay of 13.5 h ^{109g}Pd to levels of ¹⁰⁹Ag, Nucl. Phys. A107 (1968) 67-80. [E γ , P γ]
- 1969Sc12 W.C. SCHICK, Jr., W.L. TALBERT, Jr., Gamma-ray decay schemes of ^{109g}Pd, ^{111g}Pd and ^{111m}Pd, Nucl. Phys. A128 (1969) 353-387. [E γ , P γ]
- 1970Bo22 M. BORMANN, H.H. BISSEM, E. MAGIERA, R. WARNEMUNDE, Total cross sections and isomeric cross-section ratios for (n, 2n) reactions in the energy region 12-18 MeV, Nucl. Phys. A157 (1970) 481-496. [¹⁰⁹Pd half-life]
- 1970Ba37 E. BASHANDY, Contributions to the decays of ¹⁰⁹Pd and ¹⁹⁹Pt, Z. Phys. 236 (1970) 130-143. [E γ , P γ]

- 1970Fo01 J.L.C. FORD, Jr., R.L. ROBINSON, P.H. STELSON, T. TAMURA, CHEUK-YIN WONG, Elastic and inelastic proton scattering from ¹⁰⁹Ag, Nucl. Phys. A142 (1970) 525-544. [spin, parity]
- 1970Ro14 R.L. ROBINSON, F.K. MCGOWAN, P.H. STELSON, W.T. MILNER, Coulomb excitation of ^{107,109}Ag, Nucl. Phys. A150 (1970) 225-246. [nuclear levels, BR_γ, δ]
- 1973Co10 C.W. COTTRELL, Transitions, with change in parity, produced in ^{107,109}Ag by Coulomb excitation, Nucl. Phys. A204 (1973) 161-171. [¹⁰⁹Ag^m half-life]
- 1975E110 F. EL-BEDEWI, Z. MILIGY, H. HANAFI, Nuclear states of ¹⁰⁹Ag and ¹¹¹Ag, Acta Phys. 38 (1975) 153-177. [E_γ, P_γ, δ]
- 1977Bo04 H.E. BOSCH, V.M. SILBERGLEIT, M. DAVIDSON, J. DAVIDSON, Determination of mixing ratios for some cascades corresponding to ¹⁰⁹Ag, Can. J. Phys. 55 (1977) 175-179. [spin, δ]
- 1977Gi11 J.E. GINDLER, L.E. GLENDENIN, The half-lives of ¹⁰⁹Pd and ¹¹²Pd, Inorg. Nucl. Chem. Lett. 13 (1977) 95-99. [¹⁰⁹Pd half-life]
- 1978Pr08 I. PROCHÁZKA, T.I. KRACÍKOVÁ, V. JAHELKOVÁ, Z. HONS, M. FIŠER, J. JURŠÍK, The decay of ^{109g}Pd, Czech. J. Phys. B28 (1978) 134-140. [E_γ, P_γ, P_K, α_K]
- 1983Ch42 M.B. CHATTERJEE, B.B. BALIGA, Decay of ^{109g}Pd to the levels of ¹⁰⁹Ag, Fizika (Zagreb) 15 (1983) 273-282. [E_γ, P_γ]
- 1990Ab06 A. ABZOUZI, M.S. ANTONY, V.B. NDOCKO NDONGUE, D. OSTER, Re-determination of several half-lives, J. Radioanal. Nucl. Chem. 145 (1990) 361-368. [¹⁰⁹Pd half-life]
- 1996Sc06 E. SCHÖNFELD, H. JANßEN, Evaluation of atomic shell data, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527-533. [X_K, X_L, Auger electrons]
- 1998ScZM E. SCHÖNFELD, G. RODLOFF, Tables of the energies of K-Auger electrons for elements with atomic numbers in the range from Z = 11 to Z = 100, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX E. SCHÖNFELD, G. RODLOFF, Energies and relative emission probabilities of K X-rays for elements with atomic numbers in the range from Z = 5 to Z = 100, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2000He14 R.G. HELMER, C. VAN DER LEUN, Recommended standards for γ-ray energy calibration (1999), Nucl. Instrum. Methods Phys. Res. A450 (2000) 35-70. [E_γ]
- 2002Ba85 I.M. BAND, M.B. TRZHASKOVSKAYA, C.W. NESTOR, Jr., P.O. TIKKANEN, S. RAMAN, Dirac-Fock internal conversion coefficients, At. Data Nucl. Data Tables 81 (2002) 1-334. [ICC]
- 2002Ra45 S. RAMAN, C.W. NESTOR, Jr., A. ICHIHARA, M.B. TRZHASKOVSKAYA, How good are the internal conversion coefficients now? Phys. Rev. C66 (2002) 044312, 1-23. [ICC]
- 2003Au03 G. AUDI, A.H. WAPSTRA, C. THIBAULT, The AME2003 atomic mass evaluation (II). Tables, graphs and references, Nucl. Phys. A729 (2003) 337-676. [Q-value]
- 2006B102 J. BLACHOT, Nuclear data sheets for A = 109, Nucl. Data Sheets 107 (2006) 355-506. [nuclear levels]
- 2008Ki07 T. KIBÉDI, T.W. BURROWS, M.B. TRZHASKOVSKAYA, P.M. DAVIDSON, C.W. NESTOR, Jr., Evaluation of theoretical conversion coefficients using BrIcc, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202-229. [ICC]

**¹⁰⁹Cd - Comments on evaluation of decay data
by M.M. Bé, E. Schönfeld**

This evaluation was first completed by E. Schönfeld in 1996, and was completely reviewed by M.M. Bé in November 2014. New half-life and γ intensity measurement results and improved calculations of the internal conversion coefficients were included.

1 Decay Scheme

The observed transition in the decay of ¹⁰⁹Cd is an allowed EC transition $\varepsilon_{0,1}$ to the 88 keV level in ¹⁰⁹Ag. If there is an EC branch to the ground state of ¹⁰⁹Ag, it would have $\Delta J = 2$ with no change of parity (2006B102), so it would be 2nd forbidden. From the paper of S. Raman et al. (1973Ra10) it is then expected to have a $\lg ft$ greater than 11 and this corresponds to an EC branch of less than 0,005%.

Below the decay energy of ¹⁰⁹Cd there is a level at 132,74 (11) keV in ¹⁰⁹Ag, $J^\pi = 9/2^+$, $T_{1/2} = 2,60$ (12) ns. This level has been observed in the decay of ¹⁰⁹Pd but not in the decay of ¹⁰⁹Cd. If this EC branch exists, it is 2nd forbidden with an expected $\lg ft$ greater than 11 and an emission probability of less than 0,0003%.

2 Nuclear Data

The Q_{EC} value of **215,5 (18) keV** is taken from the atomic mass evaluation of Wang *et al.* (2012Wa38).

In the past, some discrepancies in the Q_{EC} values were observed: 183,9 keV was derived from internal bremsstrahlung measurements (Gopinathan *et al.* (1968Go25)); 201 (3) keV from $P(L)/P(K) = 0,193$ (3) (Goedbloed (1968GoZY), Goedbloed *et al.* (1970Go39)) measured experimentally; 220 (3) keV from $P(L, M, N)/P(K) = 0,227$ (2) (average from Leutz *et al.* (1965Le06), Goedbloed (1968GoZY), Goedbloed *et al.* (1970Go39)) measured experimentally.

From the present accepted Q_{EC} value of 215,5 (18) keV, the P_K , P_L , etc. have been derived (§2.1) and used to calculate (with the atomic data) the X-ray intensities. The ratio of the X-ray over the γ ray intensity was found to be consistent with the experimental values (§5), that support the adopted Q_{EC} value.

The following values of the half-life have been considered ($T_{1/2}$, in days):

Reference	$T_{1/2}$ (d)	Comments
Gum and Pool (1950Gu54)	470 (8)	Omitted
Leutz <i>et al.</i> (1965Le06)	453 (2)	Omitted
East and Murphy (1968Ea01)	459 (6)	Omitted
Reynolds <i>et al.</i> (1968Re04)	450 (5)	Omitted
Vaninbroukx <i>et al.</i> (1981Va11)	461,9 (3)	
Lagoutine and Legrand (1982La25)	463,1 (4)	Original uncertainty 3σ divided by 2
Hoppes <i>et al.</i> (1982HoZJ)	463,2 (6)	Superseded by 1992Un01
Unterweger <i>et al.</i> (1992Un01)	463,26 (63)	Superseded by 2002Un02
Martin <i>et al.</i> (1997Ma75)	460,2 (4)	Original uncertainty multiplied by 2
Unterweger (2002Un02)	463,26 (63)	Same value as in 1992Un01, Superseded by 2012Fi12
Schrader (2004Sc04)	459,6 (17)	
Van Ammel <i>et al.</i> (2011Va02)	462,29 (30)	Weighted mean of two results
Fitzgerald (2012Fi12)	462,6 (7)	Superseded by 2014Un01

Unterweger and Fitzgerald (2014Un01)	462,3 (8)	
Fenwick	462,40 (20)	<i>Not yet published, measurement in progress – not included</i>
Chi2 crit.:	3,0	
Chi2 :	6,2	
UWM	461,57	
WM	461,93	
uc(WM)int. :	0,17	
uc(WM)ext. :	0,41	
Adopted	461,9 (4)	

The uncertainty given by Lagoutine and Legrand (1982La25) was calculated as the sum of the systematic components and three times the statistical component, thus it cannot be simply divided by three. It was divided by two only, that gives it a weight comparable to the other values published at the same period of time.

With the original uncertainty of 0,2 d given by Martin (1997Ma75) the set of the six accepted data is strongly not consistent with a reduced χ^2 of 13,6 and this value weighs for 45%, thus it has been multiplied by two. Then, the values of 1981Va11 and 2011Va02 amounts for 37% and those of 1982La25 and 1997Ma75 for 17%.

In spite of, the set of the six data is not consistent, the three most recent values ranging from 459,6 d (2004Sc04) to 462,3 d (2014Un01).

The LWEIGHT program has used the weighted mean and external uncertainty.

Makaryunas and Makaryunene (1984Ma**) searched for a chemical alteration of the probability of EC by the ¹⁰⁹Cd nucleus. Metallic Cd, CdS and CdTe were measured. No significant change ($\Delta\lambda/\lambda < 1 \cdot 10^{-4}$) has been found from a 1000 d measurement with a NaI(Tl) detector equipped with a Be window and collimation.

Half-life (^{109m}Ag)

The recommended half-life has been determined from the measurements of Alvarez (1940Al01), Wiedenbeck (1945Wi11), Bradt *et al.* (1947Br05), Wolicki *et al.* (1951Wo15), Middelboe (1967Mi11), Abrams and Pelekis (1967Ab07), Cottrell (1973Co10) and Yoshida *et al.* (2000Yo07).

The values of 1973Co10 and 2000Yo07 have been omitted as outliers due to Chauvenet's criterion. The adopted value is the weighted mean of the six remaining values, *i.e.* 39,7 (2) seconds.

Half-life measurements (^{109m}Ag)

Reference	Half-life (s)	Comments
1940Al01	40 (2)	
1941He01	40	Same as 1940Al01
1945Wi11	40,4 (2)	
1947Br05	39,2 (2)	
1951Wo15	40 (1)	
1967Mi11	39,80 (18)	
1967Ab07	39,3 (3)	
1973Co10	35 (5)	Outlier
2000Yo07	38,0 (12)	Outlier
Recommended value	39,7 (2)	

2.1 Electron Capture Transitions

Available experimental values of P_K , P_L , P_M are listed below. They are compared with the theoretical values calculated by the EC-CAPTURE program (1998Sc28) for an allowed transition and $Q = 215,5$ (18) keV; the resulting values are in good agreement with the measured ones.

P_K	P_L	P_{M+}	P_L/P_K	P_{LMN}/P_K	Reference
			0,28 (3)		Der Mateosian (1953De26)
			0,32 (4)		Bertolini et al. (1954Be41)
0,805 (27)				0,24 (4)	Wapstra and van der Eijk (1957Wa05)
0,814 (2)	0,159	0,027	0,195 (5)	0,228 (3)	Leutz et al. (1965Le06)
0,778 (25)	0,184	0,038	0,237 (15)	0,332 (15) ^(O)	Moler and Fink (1965Mo06)
0,794 (25)	-	-	-	0,26 (4)	Durosinmi-Etti <i>et al.</i> (1966Du01)
0,816 (2)	0,157 (5)	0,027	0,193 (3) ^(LW)	0,226 (3)	Goedbloed <i>et al.</i> (1970Go39)
0,780 (15)	-	-	-	0,282	Plch <i>et al.</i> (1979Pl04)
0,815 (2)			0,199 (8)	0,227 (2)	Weighted mean
0,812 (3)	0,150 (3)	0,0385 (10)	0,183 (9)	0,231 (10)	Theory

^(O) outlier; ^(LW) limited weight

2.2 Gamma Transitions

The conversion coefficients of this E3 transition have been extensively measured. The main studies are summarized below.

Experimentally and theoretically determined conversion coefficients are compiled in the following tables:

	α_K	α_T	α_K/α_L	$\alpha_K/(\alpha_L+\alpha_M+\alpha_N)$	Reference
1	12,4 (10)	-		0,85 (2)	Brunner <i>et al.</i> (1953Br73)
2	10,3 (5)	-	-	-	Wapstra and van der Eijk (1957Wa05)
3	-	-	0,95 (3)	-	Boyd <i>et al.</i> (1964Bo12)
4	11,0 (3)	24,7 (5)	-	-	Leutz <i>et al.</i> (1965Le06)
5	11,3 (4)	24,2 (14)	-	-	Sen and Durosinmi-Etti (1965Se08)
6	12,7 (9)	29,4 (25)	0,94	0,76 (2)	Foin <i>et al.</i> (1968Fo03)
7	-	-	-	0,76 (2)	Planskoy (1969Pl08)
8	10,6 (5)	-	-	-	Bashandy (1970Ba37)
9	-	25,4 (5)	-	-	Legrand <i>et al.</i> (1973Le29)
10	11,4 (3)	26,4 (4)	0,933	0,760	Dragoun <i>et al.</i> (1976Dr07)
11	9,6 (2)	-	-	-	Prochazka <i>et al.</i> (1978Pr**)
12	11,4 (3)	26,4 (3)	-	-	Plch <i>et al.</i> (1979Pl04)
13	-	26,21 (14)	-	-	Ballaux <i>et al.</i> (1988Ba60)
14	-	26,67 (9)	-	-	Ratel (1994Ra37)

15		26,30 (25)			Kossert <i>et al.</i> (2006Ko27)
	11,28 (12)	26,62 (9)	0,913	0,736	Weighted mean of experimental values
16		26,46 (8)			The α_t value corresponds to the I_γ as determined in §4.2
17	11,41 (16)	26,3 (4)	0,946 (19)	0,765 (14)	BRICC Frozen Orbital approximation

	Brenner and Perlman (1972Br02)	Davidonis <i>et al.</i> (1980Da23)	Dragoun <i>et al.</i> (1976Dr07)	Shevelev <i>et al.</i> (1978Sh08)		Theory: BRICC Frozen Orbital approximation
α_{L1}			0,63 (13)			0,891 (13)
α_{L2}			5,48 (18)			5,16 (8)
α_{L3}			6,11 (20)			6,02 (9)
α_M			2,40 (8)			2,47 (4)
$L_1 / L_2 / L_3$	0,132 (8) / 0,830 (20) / 1	0,148 (7) / 0,86 (2) / 1	0,103 (22) / 0,897(42) / 1	0,159 (13) / 0,860 (20) / 1		0,148 (3) / 0,857 (18) / 1
K/L/M/N				0,98 (5) / 1 / 0,20 (1) / 0,050 (5)		0,946 (19) / 1 / 0,205 (4) / 0,032 (1)

The most recent values, from 1976 to 2006, of α_T and α_K are generally in agreement with the theoretical values within the uncertainty limits. Also, the experimental and theoretical ratios $L_1 / L_2 / L_3$ and K/L/M/N are most often in agreement, which supports the improvements made in the theoretical calculations (Band and Trzhaskovskaya (2002Ba85)) for such E3 transitions since the previous evaluation.

As α_T and I_γ are closely connected, further experimental values can be found in papers which are dealing with the determination of I_γ (§4.2).

Double K-shell vacancy creation in the decay of ¹⁰⁹Cd has been measured by van Eijk and Wijnhorst (1977Va05): $P_{KK}(\text{IC}) = 2,8 (7) \cdot 10^{-5}$ per K internal conversion. In a later paper, van Eijk *et al.* (1979Va04), determined the probability $P_{KK}(\text{IC})$ of double K-shell vacancy creation per K internal conversion of the 88 keV E3 transition in the decay of ^{109m}Ag, by means of a K_α -X-ray-K-X-ray coincidence experiment on ¹⁰⁹Pd, to be $13,0 (11) \cdot 10^{-5}$. From a similar experiment on ¹⁰⁹Cd, the probability $P_{KK}(\text{EC})$ of double K-shell vacancy production per K-electron capture decay of ¹⁰⁹Cd has been determined to be $1,02 (36) \cdot 10^{-5}$. Nagy *et al.* (1975Na01) found the probability that a double K-shell vacancy is formed per K-shell internal conversion to be $1,53 (24) \cdot 10^{-4}$. Horvat and Ilakovac (1985Ho06) measured the decay of the double-K-shell vacancy state in ^{109m}Ag, the probability of creation of double K-shell vacancies per ¹⁰⁹Cd decay was determined to be $6,07 (12) \cdot 10^{-5}$.

The energy shift of the hyper satellite Ag $K_{\alpha 1}^H$ -X-ray line was found to be 532 (6) eV.

Martin *et al.* (1975Ma32) measured ratios of L subshell conversion electrons.

Ilakovac *et al.* (1988II01) searched for double photon decay of the ¹⁰⁹Ag metastable state at 88 keV and found an experimental upper limit of the relative transition probability $P_{\gamma\gamma}/P_\gamma < 6 \cdot 10^{-7}$ using a pair of Ge detectors and a fast-slow coincidence system.

3 Atomic data

The atomic data, $P(K_{\beta})/P(K_{\alpha})$ and $P(K_{\alpha_2})/P(K_{\alpha_1})$, $P(KLX)/P(KLL)$ and $P(KXY)/P(KLL)$, etc. have been taken from Schönfeld and Janßen (1996Sc06).

4 Electron Emissions

The conversion electron energies were calculated from the transition energy and the binding energies of the electrons of the corresponding shells. The number of conversion and Auger electrons per 100 disintegrations is based on the decay scheme parameters as given in Section 2.1 and 2.2 and on the atomic data as given in Section 3.

5. Photon Emissions

5.1 X-ray Emissions

The total KX- ray emission intensity is (assuming there is no EC transition to the ground state):

$$I_{KX} = \omega_K \{P_K + [\alpha_K/(1 + \alpha_L)]\} \times 100 \% = 102,2 (9) \%$$

The measured values of the ratio I_{KX} / I_{γ} are summarized below and compared with the value deduced from the decay scheme parameters.

	I_{KX} / I_{γ}	Reference
1	33,8 (7) ^(O)	Wapstra and van der Eijk (1957Wa05) - outlier
2	26,2 (6)	Leutz <i>et al.</i> (1965Le06)
3	22,2 (6) ^(O)	Jansen and Wapstra (1966Ja01) - outlier
4	29,1 (10)	Freedman <i>et al.</i> (1966Fr12)
5	30 (4)	Foin <i>et al.</i> (1968Fo03)
6	26,2 (5)	Campbell and McNelles (1972Ca16)
7	27,0 (3)	Dragoun <i>et al.</i> (1976Dr07) – superseded by 1979PI04
8	27,3 (6)	Plch <i>et al.</i> (1979PI04)
9	27,34 (27)	Hoppes and Schima (1982HoZF)
10	27,9 (4)	Egorov <i>et al.</i> (1989Eg**)
11	27,7 (5)	Unweighted mean
12	27,25 (29)	Weighted mean; reduced- $\chi^2 = 2,4$; Crit. $\chi^2 = 2,8$
13	27,92 (45)	Present evaluation using the above equation together with the adopted values of ω_K , P_K , α_K , α_T and I_{γ}

In the set of data above, the results 1 and 2 are outliers due to Chauvenet's criterion. The value 7 is superseded by 8. The remaining set of seven results is consistent with a reduced χ^2 of 2,4. The unweighted and weighted means (values 11 and 12, respectively) are in agreement within the uncertainty limits with the value 13 obtained in this evaluation.

5.2 Gamma Emissions

	E_{γ} (keV)	Reference
1	88,008 (42)	Freedman <i>et al.</i> (1966Fr12)
2	88,041 (87)	Schima and Hutchinson (1967Sc22)
3	88,05 (5)	Libert (1967Li10)
4	88,033 (42)	Pierson and Marsh (1967Pi05)
5	88,09 (3)	Foin <i>et al.</i> (1968Fo03)

6	88,21 (3)	Furuta and Rhodes (1968Fu05)
7	88,036 (8)	Heath (1969HeZY)
8	88,036 (8)	Greenwood <i>et al.</i> (1970Gr13)
9	88,035 (6)	Raaside (1970Ra37)
10	88,035 (4)	Morii (1978Mo22)
11	88,0336 (10)	Helmer and van der Leun (2000He14), adopted here.

The γ ray energy is taken from Helmer and van der Leun (2000He14).

The number of X photons per 100 disintegrations is based on P_K , P_L , P_M as given in Section 2.1, α_K , α_L as given in Section 2.2 and the atomic data as given in Section 3.

The following values for the number of γ photons per 100 disintegrations have been taken into account:

	I_γ (%)	correspond. α_i	Comments on I_γ results
1	3,65 (4)	26,4 (3)	Plch <i>et al.</i> (1979Pl04) – superseded by [2]
2	3,62 (2)	26,82 (14)	Plch (UVVVR, 1988) – simple mean of two values deduced from activity measurements
4	3,65 (3)	26,40 (23)	Gostely (IER, 1988)
5	3,59 (11)		Konstantinov (IMM, 1988)
6	3,70 (6)	26,0 (5)	Park <i>et al.</i> (KSRI, 1988)
7	3,60 (2)	26,78 (8)	Chauvenet (LMRI, 1988) – original uncertainty $\times 2$
8	3,57 (10)	27,0 (8)	Woods and Smith (NPL, 1988)
9	3,65 (8)	26,4 (6)	Szörenyi <i>et al.</i> (OMH, 1988)
10	3,675 (18)	26,21 (15)	Ballaux (NBS, 1988) and 1988Ba60
11	3,66 (5)	26,3 (4)	Hino and Kawada (ETL, 1988) and 1989Hi01
12	3,68 (4)	26,2 (5)	Funck and Schötzig (PTB, 1989), Schötzig <i>et al.</i> (1992ScZZ) – superseded by [14]
13	3,65 (2)	26,4 (4)	Nedovesov (1989Ne**) based on Plch measurements - omitted
14	3,663 (33)	26,30 (25)	Kossert <i>et al.</i> (2006Ko27)
A	3,641 (11)		Weighted mean of values 2–11 and 14 Reduced $\chi^2 = 1,2$; crit. $\chi^2 = 2,4$
15	3,89 (7)	24,7 (5)	Leutz <i>et al.</i> (1965Le06) from α_i
16	3,97 (21)	24,2 (14)	Sen and Durosini-Etti (1965Se08); from α_i
17	3,29 (25)	29,4 (25)	Foin <i>et al.</i> (1968Fo03); from α_i
18	3,79 (7)	25,4 (5)	Legrand <i>et al.</i> (1973Le29) ; from α_i
19	3,65 (5)	26,4 (4)	Dragoun <i>et al.</i> (1976Dr07); from α_i
B	3,74 (6)		Weighted mean of values 15–19; Reduced $\chi^2 = 3,2$; crit. $\chi^2 = 3,3$
C	3,641 (10)	26,46 (8)	Weighted mean of 2–11, 14 and 19 Reduced $\chi^2 = 1,1$; crit. $\chi^2 = 2,3$.
D	3,66 (5)	26,3 (4)	From theoretical ICC - Adopted

The first set of data (1 - 14) lists the I_γ results obtained by direct measurements of the total activity of a ¹⁰⁹Cd solution by means of absolute methods and the photon emission rate by gamma spectrometry counting. All these results, except the value 14, were obtained in the framework of an international comparison of activity measurements (1994Ra37), some of them (10, 11, 12) were also published elsewhere. When two results were given (Plch, 1988) the simple mean was adopted. The original

uncertainty of the value 7 has been multiplied by two; this value then has a weight comparable to the others. The weighted mean of values 2–11 and 14 is given on line A of the above table.

The second set of data (15 – 19) lists the I_γ results derived from measurements of the internal conversion coefficients. The set of five values is just consistent with a reduced χ^2 of 3,2. The weighted mean is listed on line B.

When processing all the available data (1 – 19) the values (15 -18) were found to be outliers due to Chauvenet's criterion. The set of the eleven remaining values is consistent with a reduced χ^2 of 1,1; the weighted mean is given on line C.

An I_γ value has been derived on line D, from the theoretical value of the total internal coefficient for an E3 transition interpolated from Band *et al.* tables (2002Ba85) using the BRICC program with the “frozen orbital approximation” (Kibédi *et al.* 2008Ki07). The I_γ and ICC values listed on line D are in good agreement with the experimental values obtained on line C.

It is noteworthy that the I_γ determined in various studies are dependent on the efficiency of the γ detector at 88 keV, but there are only a few γ rays available to establish an efficiency curve in this energy region. For this reason, and in order to propose an independent value, the I_γ derived from the theoretical ICC is adopted here, that is: $I_\gamma = 3,66$ (5) %.

6 Main Production Modes

Taken from the “Table de Radionucléides”, LMRI, 1982:

Cd-108(n, γ)Cd-109, σ : 1,1 (3) barns
Possible impurities: Ag-110m

Ag-109(p,n)Cd-109
Possible impurities: none

7 References

- 1940Al01** L.W.Alvarez, A.C.Helmholz, E.Nelson. *Phys. Rev.* 57 (1940) 660. Isomeric Silver and the Weizsacker Theory [$T_{1/2}$ Isomeric level]
- 1941He01** A.C. Helmholz. *Phys. Rev.* 60 (1941) 415. Energy and Multipole Order of Nuclear γ -Rays [$T_{1/2}$ Isomeric level]
- 1945Wi11** M.L.Wiedenbeck. *Phys. Rev.* 67 (1945) 92. The Nuclear Excitation of Silver and Cadmium [$T_{1/2}$ Isomeric level]
- 1947Br05** H.Bradt, P.C.Gugelot, O.Huber, H.Medicus, P.Preiswerk, P.Scherrer, R.Steffen
Helv. Phys. Acta 20 (1947) 153. Die metastabilen Zustände der Silberkerne Ag107 und Ag109 [$T_{1/2}$ Isomeric level]
- 1950Gu54** J. R. Gum and M. L. Pool, *Phys. Rev.* 80 (1950) 315 [$T_{1/2}$]
- 1951Wo15** E.J.Wolicki, B.Waldman, W.C.Miller. *Phys. Rev.* 82 (1951) 486. The Nuclear Excitation of Ag107 and Ag109 by X-Rays [$T_{1/2}$ Isomeric level]
- 1953Br73** J. Brunner, O. Huber, R. Joly and D. Maeder, *Helv. Phys. Acta* 26 (1953) 588 [$\alpha_K, \alpha_K/(\alpha_L + \alpha_M + \alpha_N)$]
- 1953De26** E. Der Mateosian *Phys. Rev.* 92 (1953) 938 [P_L/P_K]
- 1954Be41** G. Bertolini, A. Bisi, E. Lazzarini and L. Zappa, *Nuovo Cimento* 11 (1954) 539 [P_L/P_K]
- 1957Wa05** A. H. Wapstra and W. van der Eijk, *Nucl. Phys.* 4 (1957) 325 [$I_{KX}/I_\gamma, P_K, P_{LMN}/P_K, \alpha_K$]
- 1964Bo12** H. W. Boyd, J. H. Hamilton, A. R. Sattler and P. F. A. Goudsmit, *Physica* 30 (1964) 124 [α_K/α_L]

- 1965Le06** H.Leutz, K.Schneckenberger, H.Wenninger. *Nucl. Phys.* 63 (1965) 263. [$T_{1/2}$, I_{KX}/I_γ , P_L/P_K , ICC, Q(EC)]
- 1965Mo06** R.B.Moler, R.W.Fink. *Phys. Rev.* 2B (1965) B282. [P_K , P_L , P_M , P_{LMN}/P_K]
- 1965Se08** S.K.Sen, I.O.Durosinmi-Etti. *Phys. Lett.* 18 (1965) 144. [$\alpha_{K,T}$]
- 1966Ja01** J.W.F. Jansen and A. H. Wapstra, *Internal Conversion Processes* Nashville (ed. J. H. Hamilton; Academic Press, New York) (1966) p. 237. [I_{KX}/I_γ]
- 1966Fr12** M. S. Freedman, F. T. Porter and F. Wagner, *Phys. Rev.* 151 (1966) 886 [E_γ , I_{KX}/I_γ]
- 1966Du01** I.O.Durosinmi-Etti, D.R.Brundrit, S.K.Sen. *International Conversion Processes*, Ed. Hamilton, Acad. Press, New York (1966) 201. [P_K , P_{LMN}/P_K]
- 1967Ab07** I.A. Abrams, L.L. Pelekis. Program and Theses, Proc. 17th All Union Conf. Nucl. spectroscopy and Struct. At. Nuclei, Kharkov, p. 30 (1967). Excitation of Metastable Levels in Nuclei by the γ -Rays of ^{116m}In. [$T_{1/2}$ Isomeric level]
- 1967Li10** J. Libert, *Nucl. Phys.* A 102 (1967) 477 [E_γ]
- 1967Mi11** V.Middelboe. Kgl. Danske Videnskab. Selskab, Mat.-Fys. Medd. 35, No.8 (1966) Some Accurate Half-Life Determinations. [$T_{1/2}$ Isomeric level]
- 1967Pi05** W. R. Pierson and R. H. Marsh, *Nucl. Phys.* A 104 (1967) 511 [E_γ]
- 1967Sc22** F.J. Schima and J. M. R. Hutchinson, *Nucl. Phys.* A 102 (1967) 667 [E_γ]
- 1968Ea01** L.V. East and H. M. Murphy, Jr., *Nucl. Phys.* A 107 (1968) 382 [$T_{1/2}$]
- 1968Fo03** K. C. Foin, A. Gizon and J. Oms, *Nucl. Phys.* A 113 (1968) 241 [I_{KX}/I_γ , $\alpha_{K,T}$, E_γ]
- 1968Fu05** T. Furuta and J. R. Rhodes, *Intern. J. Appl. Rad. Isotopes* 19 (1968) 483 [E_γ]
- 1968Go25** K.P.Gopinathan, W.Rubinson. *Bull. Am. Phys. Soc.* 13, 11 (1968) 1452. (Q(EC))
- 1968GoZY** W.Goedbloed. Proc. Conf. Electron capture, Higher order processes. In *Nucl. Decays*, Debrecen, Hungary, D.Berenyi, Ed. Eötvös Lorand, Phys. Soc. Budapest, vol. 1 (1968) 92. [Q(EC)]
- 1968Re04** S.A.Reynolds, J.F.Emery, E.I.Wyatt. *Nucl. Sci. Eng.* 32 (1968) 46. [$T_{1/2}$]
- 1969HeZY** R. L. Heath, Proc. Int. Conf. on Radioactivity in Nucl. Spectroscopy, Nashville, USA (1969) [E_γ]
- 1969Pi08** B. Planskoy, *Nucl. Instr. Meth.* 73 (1969) 205 [$\alpha_K/\alpha_L + \alpha_M + \alpha_N$]
- 1970Gr13** R. C. Greenwood, R. G. Helmer and R. J. Gehrke, *Nucl. Instr. and Meth.* 77 (1970) 141 [E_γ]
- 1970Ra37** D. E. Raeside, *Nucl. Instr. and Methods* 87 (1970) 7 [E_γ]
- 1970Ba37** E. Bashandy, *Z. Phys.* 236 (1970) 130 [α_K]
- 1970Go39** W.Goedbloed, S.C.Goverse, C.P.Gerner, A.Brinkman, J.Blok. *Nucl. Instrum. Methods* 88 (1970) 197. [P_K , P_L , P_M , P_{LMN}/P_K , Q(EC)]
- 1972Br02** D. S. Brenner and M. L. Perlman, *Nucl. Phys.* A 181 (1972) 207 – 216 [$\alpha_{L1/L2/L3}$]
- 1972Ca16** J.L. Campbell, L.A. McNelles. *Nucl. Instrum. Methods* 98 (1972) 433. [I_{KX}/I_γ]
- 1973Co10** C.W.Cottrell. *Nucl.Phys.* A204 (1973) 161. Transitions, with Change in Parity, Produced in 107,109Ag by Coulomb Excitation. [$T_{1/2}$ Isomeric level]
- 1973Le29** J. Legrand, M. Blondel and P. Magnier, *Nucl. Instr. and Methods* 112 (1973) 101 [α_L]
- 1973Ra10** S. Raman, *Phys. Rev.* C 7 (1973) 1995 [$lg ft$]
- 1975Ma32** B. Martin, D. Merkert and J. L. Campbell, *Z. Physik* A 274 (1975) 15 [$I_{ec(L1/L2/L3)}$]
- 1975Na01** H. J. Nagy, G. Schupp, R. R. Hurst, *Phys. Rev.* C 11 (1975) 205 [P_{KK}]
- 1976Dr07** O.Dragoun, V.Brabec, M.Rysavy, J.Plch, J.Zderadicka. *Physik* A279 (1976) 107. [Int. Conv. Coefficients, I_{KX}/I_γ , I_γ]
- 1977Va05** C. W. E. van Eijk and J. Wijnhorst, *Phys. Rev.* C 15 (1977) 1068 [P_{KK} (IC)]
- 1978Sh08** G. A. Shevelev, A. G. Troitskaya and V. M. Kartashov, *Izv. Akad. Nauk SSSR, Ser. Fiz* 42 (1978) 211. [$\alpha_{L1/L2/L3}$]
- 1978Mo22** T. Morii, *Nucl. Instr. and Methods* 151 (1978) 489 [E_γ]
- 1978Pr**** I. Prochazka, T. I. Kracikova, V. Jahelkova, Z. Hons, M. Friser, and J. Jursik, *Czech. J. Phys.* B28 (1978) 134 [α_K]
- 1979Pi04** J.Plch, P.Dryak, J.Zderadicka, E.Schönfeld, A.Szörenyi. *Czech. J. Phys.* B29 (1979) 1071. [I_{KX}/I_γ , P_K , $\alpha_{K,T}$, I_γ]

- 1979Va04** C. W. E. van Eijk, J. Wijnhorst, M. A. Popelier, *Phys. Rev. C* 19 (1979) 1047 [P_{KK} (IC)]
- 1980Da23** R. I. Davidonis, R. K. Zhirgulyavichyus, R. A. Kalinauskas, V. I. Kerskulis, K. V. , *Izv. Akad. Nauk SSSR, Ser. Fiz.* 44 (1980) 1060 [$\alpha_{L1/L2/L3}$]
- 1981Va11** R. Vaninbrouckx, G. Grosse, W. Zehner. *Int. J. Appl. Radiat. Isot.* 32 (1981) 589. [$T_{1/2}$]
- 1982HoZJ** D.D.Hoppes, J.M.R.Hutchinson, F.J.Schima, M.P.Unterweger. NBS-Special Publ. 626 (1982) 85 - 99. [$T_{1/2}$, I_{KX}/I_{γ} , I_{γ}]
- 1982La25** F.Lagoutine, J.Legrand. *Int. J. Appl. Radiat. Isotop.* 33 (1982) 711. [$T_{1/2}$]
- 1984Ma**** K. V. Makaryunas and E. K. Makaryunene, *Izv. Akad. Nauk SSSR, Ser. Fiz.* 48 (1984) 23-27. [$T_{1/2}$]
- 1985Ho06** H. Horvat, K. Ilakovac, *Phys. Rev. A* 31 (1985) 1543 [P_{KK} (IC)]
- 1988Ba60** C. Ballaux, B.M. Coursey, D.D. Hoppes. *Appl.Radiat.Isot.* 39 (1988) 1131. ^{109}Pd and ^{109}Cd Activity Standardization and Decay Data. [α_{T} , I_{γ}]
- 1988Pi01** K. Ilakovac, G. Jerbic-Zorc, M. Bozin, R. Posic, W. Horvat, *Fizika (Zagreb)* 20 (1988) 91. [$P_{\gamma\gamma}$]
- 1989Hi01** Y. Hino, Y. Kawada. *Appl. Radiat. Isot.* 40 (1989) 79. [I_{γ}]
- 1989Eg**** A.G. Egorov, Y.S. Egorov, V.G. Nedovesov, G.E. Shchukin, K.P. Yakovlev. 39th Conf. on Nucl. Spectroscopy and Atomic Nucleus Structure, Tashkent, USSR 18-21 April 1989, Lo. Nauka, Leningrad (1989) 505. [I_{KX}/I_{γ}]
- 1989Ne**** V.G. Nedovesov, V.P. Chechev, E.S. Checheva. *Measuring Technics* 6 (1989) 52. [I_{γ}]
- 1992ScZZ** U.Schötzig, H.Schrader, K.Debertin. Proc. Int. Conf. Nuclear Data for Science and Technology, Jülich 1991 (1992) 562. [I_{γ}]
- 1992Un01** M.P. Unterweger, D.D. Hoppes, F.J. Schima, *Nucl. Instr. Meth.* A312 (1992) 349. [$T_{1/2}$]
- 1994Ra37** G. Ratel. *Nucl. Instrum. Methods Phys. Res.* A345 (1999) 289. [α_{T} , I_{γ}]
- 1996Sc06** E. Schönfeld, H. Janssen. *Nucl. Instrum. Methods Phys. Res.* A369 (1996) 527. [Atomic Data]
- 1997Ma75** R.H.Martin, K.I.W.Burns, J.G.V.Taylor. *Nucl. Instrum. Methods* A390 (1997) 267. [$T_{1/2}$]
- 1998Sc28** E. Schönfeld. *Appl. Radiat. Isot.* 49 (1998) 1353. [Fractional EC probabilities]
- 2000He14** R.G.Helmer, C.Van der Leun. *Nucl. Instrum. Methods* A450 (2000) 35. [E_{γ}]
- 2000Yo07** E.Yoshida, T.Kobayashi, Y.Kojima, K.Shizuma. *Nucl. Instrum. Methods Phys. Res.* A449 (2000) 217. Half-Lives of Isomeric Levels of ^{107m}Ag , ^{109m}Ag and ^{103m}Rh Photoactivated by ^{60}Co γ -Ray Irradiation [$T_{1/2}$ Isomeric level]
- 2002Ba85** I.M. Band, M.B. Trzhaskovskaya. *At. Data. Nucl. Data Tables* 88 (2002) 1. Theoretical ICC
- 2002Un02** M.P. Unterweger, *Appl. Radiat. Isot.* 56, 125 (2002). [$T_{1/2}$]
- 2004Sc04** H.Schrader. *Appl. Radiat. Isot.* 60 (2004) 317. Half-life measurements with ionization chambers. A study of systematic effects and results. [$T_{1/2}$]
- 2006Bl02** J.Blachot. *Nucl. Data Sheets* 107 (2006) 355 [spin and parity]
- 2006Ko27** K.Kossert, H.Janssen, R.Klein, M.K.H.Schneider, H.Schrader. *Appl.Radiat.Isot.* 64 (2006) 1031. Activity standardization and nuclear decay data of ^{109}Cd . [I_{γ}]
- 2008Ki07** T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor Jr. *Nucl. Instrum. Methods Phys. Res.* A589 (2008) 202. [Theoretical ICC]
- 2011Va02** R.Van Ammel, S.Pommé, J.Paepen, G.Sibbens. *Appl.Radiat.Isot.* 69 (2011) 785. Measurement of the ^{109}Cd half-life. [$T_{1/2}$]
- 2012Fi12** R.Fitzgerald. *J.Res.Natl.Inst.Stand.Technol.* 117 (2012) 80. [$T_{1/2}$]
- 2012Wa38** M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer. *Chin. Phys.* C36 (2012) 1603. [Q-value]
- 2014Un01** M.P.Unterweger, R.Fitzgerald. *Appl.Radiat.Isot.* 87 (2014) 92. Update of NIST half-life results corrected for ionization chamber source-holder instability. [$T_{1/2}$]

¹¹⁰Ag – Comments on evaluation of decay data by R. G. Helmer

1) Decay Scheme

The β^- emission to ¹¹⁰Cd from the ¹¹⁰Ag ground state occurs in 99,70% (6) of the decays and the remaining 0,30% (6) is by electron capture to ¹¹⁰Pd.

2) Q values and half-lives

The Q values from the 1995Au04 evaluation for the decay of the ¹¹⁰Ag ground state are 2892,2 (16) keV for the β^- decay and 892 (11) keV for the electron-capture decay.

The half-life of the ¹¹⁰Ag ground state has been determined from the following data (in seconds):

1935Am01	22	omitted, no uncertainty
1938Po03	22	omitted, no uncertainty
1938Re04	23	omitted, no uncertainty
1944F101	24	omitted, no uncertainty
1946Hi06	24,5 (3)	
1954Bo39	24 (2)	
1957Se19	24,2 (12)	
1962Ma38	24,42 (14)	
1967Yu01	24,93 (22)	
1970Va08	24,7 (7)	
Adopted	24,56 (11)	

The adopted value is the weighted average of the six values with uncertainties, and the reduced- χ^2 value is 0,82, so the values are consistent.

3) g-ray data

The energies for the γ -rays from the decay of ¹¹⁰Ag (24 s) were determined as shown in Table 1. The precise energies from the ¹¹⁰Ag^m (249 d) isomer decay are adopted where appropriate.

Table 1. γ -ray energies from the β^- decay of ¹¹⁰Ag (24 s).

1970Va08	1972Ka34 ^a	Adopted ^b
	295,3 (1)	295,3 (2)
657,8 (2)	657,6 (1)	657,7600 (11) ^c
815,5 (3)	815,5 (1)	815,5 (2)
817,8 (12)	818,2 (1)	818,0244 (18) ^c
	1074,0 (1)	1074,0 (2)
1125,9 (3)	1125,8 (1)	1125,699 (20) ^d
1186,4 (7)	1186,3 (1)	1186,3 (2)

1421,8 (13)	1421,4 (1)	1421,5 (2)
1475,8 (13)	1475,8 (1)	1475,7792 (23) ^c
1630,0 (12)	1629,9 (1)	1629,9 (2)
1674,2 (9)	1674,3 (1)	1674,3 (2)
1783,3 (13)	1783,6 (7)	1783,46 (3) ^d
	2004,4 (2)	2004,4 (2)

^a The author's uncertainties are quoted to 0,01 keV, but the energies are only given to 0,1 keV, so the last digit in the uncertainty is of no use.

^b For energies from 1972Ka34 and 1970Va08, a minimum uncertainty of 0,2 keV has been used for the adopted value.

^c From evaluation of 2000He14,

^d From adopted value from ¹¹⁰Ag^m decay.

The relative emission probabilities of the γ -rays from the decay of ¹¹⁰Ag (24 s) were determined from the measurements in Table 2 :

Table 2: Relative emission probabilities of the γ -rays from the decay of ¹¹⁰Ag (24 s)

E _{γ} (keV)	1970Va08	1972Ka34	Adopted
295		0,17 (3)	0,17 (3)
657	100,	100,	100,
815	0,79 (12)	0,85 (2)	0,85 (2)
818	0,10 (9)	0,20 (1)	0,20 (1)
1074		0,02 (1)	0,02 (1)
1125	0,36 (3)	0,34 (1)	0,34 (1)
1186	0,056 (2)	0,06 (1)	0,06 (1)
1421	0,044 (30)	0,05 (1)	0,05 (1)
1475	0,11 (5)	0,08 (1)	0,08 (1)
1629	0,048 (30)	0,05 (1)	0,05 (1)
1674	0,15 (6)	0,16 (1)	0,16 (1)
1783	0,17 (9)	0,10 (1)	0,10 (1)
2004		0,08 (1)	0,08 (1)

The normalization of the relative emission probabilities for the γ -rays from the decay of ¹¹⁰Ag (24 s) depends on the probability of the β branch to the ground state of ¹¹⁰Cd and the fact that 0,30(6)% of the decays are by electron capture to ¹¹⁰Pd (1961Fr01). The intensity of the β branch to the ¹¹⁰Cd ground state can be obtained from the ratio of the emission probabilities for the branches to the 657 -keV level and the ground state, $I_{\beta}(657)/I_{\beta}(0)$, as deduced from the decomposition of the β^- spectrum. However, the following results for this ratio are very inconsistent.

	$I_{\beta^-}(657)/I_{\beta^-}(0)$
1962Ka07	0,14 (5)
1963Da03	0,21
1963Fr07	0,0465 (25)
1967Mo12	0,070 (22)
Adopted	0,047 (4)

The adopted value is the weighted average of the three values with uncertainties. For this average the internal uncertainty is 0,0025 and the external uncertainty is 0,0038. Although the reduced- χ^2 value is 2,30, this does not necessarily imply an inconsistent set since one has only three values. If one does consider it an inconsistent set and applies the Limitation of Relative Statistical Weight rule (1985ZiZY, 1992Ra08) of reducing the relative weight of the 1963Fr07 value from 98% to 50%, the weighted average becomes 0,064 with an internal uncertainty of 0,014, a reduced $-\chi^2$ value of 1,6, and an external uncertainty of 0,018.

From this β^- branching ratio, the 0,30 (6)% electron $-\text{capture}$, and 0,1% β^- branching to higher energy levels, the branch to the ground state is 95,1(4) % and that to the 657 -keV level is 4,5(4) %. The emission probability of the 657 -keV γ -ray is then 4,6 (4) % of the decays of the ground state including both the direct and indirect feeding.

Table 3: Absolute emission probabilities for the γ -rays from the decay of the ¹¹⁰Ag ground state.

E_{γ}	P_{γ} (%)
295	0,0078 (16)
657	4,6 (4)
815	0,039 (4)
818	0,0092 (9)
1074	0,0009 (5)
1125	0,0156 (14)
1186	0,0028 (5)
1421	0,0023 (5)
1475	0,0037 (6)
1629	0,0023 (5)
1674	0,007 (1)
1783	0,0046 (8)
2004	0,0037 (6)

The γ -ray multiplicities and mixing ratios were taken from the 2000De11 evaluation and are as follows:

E1: 1421 -keV
 E2: 657, 815, 1074, 1186,1475, 1783, 2004 -keV
 M1+E2: 818 [d = - 1,36 (7)] ; 1125 [d = + 0,33 (8)]
 E2(+M1): 1629 [d = + 0,06 (3)]
 (E1): 295 -keV

4) Atomic data

From the EMISSION code and the decay data, the following information was obtained.

Quantity	Pd (Z=46)	Cd (Z=48)
ω_K	0,820(4)	0,842(4)
ω_L average	0,0536 (13)	0,0632 (16)
n_{KL}	0,975 (4)	0,953 (4)
$K_{\alpha 2}/K_{\alpha 1}$	0,5293 (25)	0,5317 (25)
K_{β}/K_{α}	0,2099 (17)	0,2151 (18)

Due to the high energy of the strong transitions, the Auger electrons are negligible and no related data are included here.

The K X-ray emission probabilities are calculated as follows:

From the decay of ¹¹⁰Ag (24 s), the Pd X-rays per 100 decays of parent:

$K_{\alpha 2}$	0,060 (12)
$K_{\alpha 1}$	0,114 (23)
K_{β}	0,037 (8)

and the Cd X-rays per 100 decays of parent:

$K_{\alpha 2}$	0,00322 (28)
$K_{\alpha 1}$	0,0061 (6)
K_{β}	0,00200 (18)

5) β^- decay intensities

The β^- decay intensities for the decay of the ¹¹⁰Ag ground state are simply deduced from the above data and the γ -ray probability balances.

6) References

- 1935Am01 - E. Amaldi, O. D'Agostino, E. Fermi. B. Pontecorvo, R. Rasetti, and E. Segrè, Proc. Roy. Soc. (London) 149A(1935)522 [$T_{1/2}$ as cited in 1962Ma38].
 1938Po03 - M. L. Pool, Phys. Rev. 53(1938)116 [$T_{1/2}$].
 1938Re04 - H. Reddemann, Naturwiss. 26(1938)124 [$T_{1/2}$ as cited in 1962Ma38].
 1944Fl01 - A. Flammersfeld, Naturwiss. 32(1944)36 [$T_{1/2}$ as cited in 1962Ma38].
 1946Hi06 - O. Hirzel, H. Wäffler, Helv. Phys. Acta 19(1946)214 [$T_{1/2}$ as cited in 1953Ho01 and/or 1962Ma38].
 1954Bo39 - F. I. Boley, Phys. Rev. 94(1954)1078 [$T_{1/2}$].
 1957Se19 - M. L. Seghal, Indian J. Phys. 31(1957)630 [$T_{1/2}$ as cited in 1962Ma38].
 1962Ka07 - T. Katoh, Y. Yoshizawa, Nucl. Phys. 32(1962)5 [E_{β} , I_{β} , E_{γ} , α_K , Mult].
 1962Ma38 - S. Malmskog, J. Konijn, Nucl. Phys. 38(1962)196 [$T_{1/2}$].
 1963Da03 - H. Daniel, O. Mehling, D. Schotte, Zeits. f. Phys. 172(1963)202 [E_{β} , P_{β}].

- 1963Fr07 - L. Frevert, P. H. Heckmann, A. Flammersfeld, Zeits. f. Phys. 175(1963)221 [E_β , P_β].
 1963Su07 - T. Suter, P. Reyes-Suter, W. Scheuer, Nucl. Phys. 47(1963)251 [E_γ , I_{e^-}].
 1964Ne05 - W. B. Newbolt, J. H. Hamilton, Nucl. Phys. 530(1964)353 [E_γ , I_{e^-} , α_K , Mult].
 1965Fr01 - L. Frevert, R. Schöneberg, A. Flammersfeld, Zeits. f Phys. 182(1965)439 [I_e].
 1967Mo12 - J. A. Moragues, P. Reyes-Suter, T. Suter, Nucl. Phys. A99(1967)652 [E_β , I_β , E_γ , I_γ].
 1967Yu01 - H. P. Yule, Nucl. Phys. A94(1967)442 [$T_{1/2}$].
 1970Kr03 - K. S. Krane, R. M. Steffen, Phys. Rev. C2(1970)724 [δ].
 1970Su03 - S. P. Sud, P. C. Mangal, P. N. Trehan, Aust. J. Phys. 23(1970)87 [δ].
 1970Va08 - J. R. Van Hise, M. C. Kelley, R. G. Lanier, N. R. Johnson, Phys. Rev. C1(1970)8161 [$T_{1/2}$, E_γ , I_γ].
 1972Ka34 - Y. Kawase, K. Okano, S. Uehara, T. Hayashi, Nucl. Phys. A193(1972)204 [E_γ , I_γ].
 1973Ga10 - P. L. Gardulski, M. L. Wiedenbeck, Phys. Rev. C7(1973)2080 [δ].
 1973Jo08 - P. D. Johnston, N. J. Stone, Nucl. Phys. A206(1973)273 [δ].
 1978Wa07 - G. W. Wang, A. J. Becker, L. M. Chirovsky, J. L. Groves, C. S. Wu, Phys. Rev. C18(1978)476 [δ].
 1979Sc31 - P. Schlüter, G. Soff, Atomic Data Nuclear Data Tables 24(1979)509 [α_π].
 1979Ve03 - H. R. Verma, A. K. Sharma, R. Kaur, K. K. Suri, P. N. Trehan, J. Phys. Soc. Japan 47(1979)16 [E_γ , I_γ , δ].
 1985ZiZY - W. L. Zijp, Report ECN FYS/RASA-85/19 (1985) [averages].
 1992Ra08 - M. U. Rajput, T. D. MacMahon, Nucl. Instr. Meth. A312(1992)289 [averages].
 1993Ka37 - V. M. Kartashov, A. I. Oborovsky, A. G. Troitskaya, Bull. Russ. Acad. Sci. 57(1993)1554 [I_{e^-}].
 1993Ki18 - L. L. Kiang, P. K. Teng, G. C. Kiang, W. S. Chang, P. J. Tu, J. Phys. Soc. Japan 62 (1993)888 [E_γ , I_γ , δ].
 1995Au04 - G. Audi, A. H. Wapstra, Nucl. Phys. A595(1995)409 [Q].
 2000De11 - D. DeFrenne, E. Jacobs, Nucl. Data Sheets 89(2000)481 [J^π , multipolarities, δ].
 2000He14 - R. G. Helmer, C. van der Leun, Nucl. Instr. Meth. A450(2000)35 [E_γ].

**¹¹⁰Ag^m – Comments on evaluation of decay data
by R. G. Helmer**

1) Decay Scheme

The β^- decay of the ¹¹⁰Ag^m (249 d) isomer to levels in ¹¹⁰Cd occurs in 98,64(8) % of the decays and the remaining 1,36(8) % is by an isomeric transition to the ¹¹⁰Ag ground state (24 s). The β^- emission to ¹¹⁰Cd from the ground state occurs in 99,70(6) % of the decays and the remaining 0,30(6) % is by electron capture to ¹¹⁰Pd. The comments on the decay ¹¹⁰Ag (24 s) ground state are provided under that decay.

2) Q values and half-lives

The Q values from the 1995Au04 evaluation for the decay of the ¹¹⁰Ag ground state are 2892,2 (16) keV for the β^- decay so the decay energy for the β^- decay of the ¹¹⁰Ag^m (249 d) isomer is then 3009,8 (16) keV.

The half-life of the ¹¹⁰Ag^m isomeric state has been determined from the following data (in days):

1938Li07	225 (20)	omitted, large uncertainty
1950Gu54	270	omitted, no uncertainty
1976WaZH	249,78 (4)	superseded by 1983Wa26
1980Ho17	249,74 (5)	
1983Wa26	249,79 (2)	
Adopted	249,78 (2)	

The adopted value is the weighted average of the last two values, and the reduced- χ^2 value is 0,86.

3) g-ray data

Several of the γ -rays from the decay of the isomer ¹¹⁰Ag^m (249 d) have precisely measured energies; these values were taken from the evaluation 2000He14 and are on a scale for which the energy of the strong line from the decay of ¹⁹⁸Au is 411,80205(17) keV. The other energies were determined as shown in Table 1 from the data of 1979Ve03, 1981Ma09, 1990Me15, and 1993Ki18. In order to provide a set of energies consistent with those of 2000He14, the values 1990Me15 were adjusted by additive amounts of 0 to 15 eV as shown in the table. No additional uncertainty was assigned for these adjustments. The values of the remaining references were not adjusted.

Table 1. γ -ray energies (keV)

1979Ve03	1981Ma09 ^a	1993Ki18	1990Me15	1990Me15 adjusted & rounded	2000He14	Adopted
			116,485 (46)	116,48 (5)		116,48 (5)
120,4 (2)	120,3 (1)	120,2 (2)	120,226 (26)	120,23 (3)		120,23 (3)
133,3 (2)	133,4 (1)	133,2 (1)	133,333 (7)			133,333 (7)
219,2 (2)	219,4 (1)	219,4 (1)	219,348 (8)			219,348 (8)

1979Ve03	1981Ma09 ^a	1993Ki18	1990Me15	1990Me15 adjusted & rounded	2000He14	Adopted
221,0 (1)	221,0 (1)	221,1 (2)	221,079 (10)			221,079 (10)
229,3 (2)	229,4 (1)	229,4 (3)	229,423 (23)			229,423 (23)
	264,4 (1)	264,1 (3)	264,254 (58)	264,25 (6)		264,25 (6)
266,9 (2)	267,0 (1)	267,0 (3)	266,913 (12)			266,913 (12)
	341,4 (1)	340,9 (5)	341,2 (2)			341,3 (2)
	356,4 (1)	356,5 (2)	356,43 (10)			356,43 (10)
360,7 (2)	360,0 (1)	360,2 (5)	360,228 (75)	360,23 (8)		360,23 (8)
365,54 (10)	365,4 (1)	365,3 (1)	365,450 (10)	365,448 (10)		365,448 (10)
387,2 (2)	387,1 (1)	387,1 (6)	387,075 (9)	387,073 (9)		387,073 (9)
397,1 (2)	396,8 (1)	396,5 (6)	396,897 (23)	396,895 (23)		396,895 (23)
	409,6 (1) ^d	409,6 (4)	409,330 (45)	409,33 (5)		409,4 (5)
446,87 (5)		446,8 (2)	446,808 (8)		446,812 (3)	446,812 (3)
466,9 (2)	466,9 (1)	465,8 (7)	467,029 (36)	467,03 (4)		467,03 (4)
493,8 (2)	493,0 (1)	493,6 (1)	493,432 (91)	493,43 (9)		493,43 (10)
554,8 (2)	544,5 (1)	544,9 (5)	544,555 (45)	544,55 (5)		544,55 (5)
	572,7 (1)	573,1 (7)	573,0 (4)			572,8 (2)
	603,1 (1)	603,1 (4)	603,065 (90)	603,06 (9)		603,08 (10)
620,45 (5)		620,4 (1)	620,362 (1)		620,3553 (17)	620,3553 (17)
626,24 (5)	626,1 (1)	626,4 (2)	626,262 (10)	626,258 (10)		626,258 (10)
	630,6 (1)	630,7 (4)	630,626 (55)	630,62 (6)		630,62 (6)
	648,2 (10)	647,8 (4)				647,8 (4)
657,75 (5)		657,7 (2)	657,766 (5)		657,7600 (11)	657,7600 (11)
	666,1 (2)	667,1 (1)				666,6 (5)
	676,6 (1)		676,58 (10)			676,58 (10)
677,72 (5)		677,6 (1)	677,623 (7)		677,6217 (12)	677,6217 (12)
687,10 (5)			687,005 (11)		687,0091 (18)	687,0091 (18)
706,74 (5)			706,688 (8)		706,6760 (15)	706,6760 (15)
	708,3 (1)	708,6 (5)	708,133 (20)	708,128 (20)		708,128 (20)
	714,9 (1)	715,0 (3)				714,9 (1)
744,35 (5)			744,279 (8)		744,2755 (19)	744,2755 (18)
763,98 (5)			763,947 (8)		763,9424 (17)	763,9424 (17)
	774,8 (1)	774,6 (1)	774,8 (2)			774,70 (10)
818,00 (5)			818,037 (8)		818,0244 (18)	818,0244 (18)
884,65 (5)			884,037 (8)		884,6781 (13)	884,6781 (13)

1979Ve03	1981Ma09 ^a	1993Ki18	1990Me15	1990Me15 adjusted & rounded	2000He14	Adopted
937,55 (5)			937,505 (13)		937,485 (3)	937,485 (3)
957,3 (2)	957,4 (1)	957,6 (7)	957,368 (85)	957,35 (9)		957,35 (10)
997,12 (5)	997,2 (1)	997,2 (4)	997,258 (15)	997,243 (15)		997,243 (15)
1019,0 (2)	1019,1 (1)	1018,8 (5)	1018,893 (50)	1018,88 (5)		1018,95 (8)
	1050,1 (3)	1051,8 (6)				1050,5 (5)
1085,7 (1)	1085,5 (1)	1085,3 (4)	1085,462 (14)	1085,447 (14)		1085,447 (14)
1117,7 (2)	1117,5 (1)	1117,2 (3)	1117,474 (28)	1117,46 (3)		1117,46 (3)
1125,7 (2)	1125,6 (1)	1125,6 (4)	1125,714 (20)	1125,699 (20)		1125,699 (20)
1163,5 (2)	1163,1 (2)	1163,1 (3)	1163,159 (75)	1163,14 (8)		1163,14 (8)
1165,6 (2)	1164,5 (2)	1165,2 (8)	1164,959 (85)	1164,94 (9)		1164,94 (9)
	1186,7 (1)	1186,5 (2)	1186,7 (2)			1186,7 (1)
1251,2 (2)	1251,0 (1)	1251,2 (3)	1251,057 (42)	1251,04 (4)		1251,04 (4)
1300,0 (2)	1300,1 (1)	1300,3 (4)	1300,03 (12)	1300,02 (12)		1300,05 (10)
1334,53 (10)	1334,4 (1)	1334,3 (3)	1334,341 (17)	1334,326 (17)		1334,326 (17)
1384,47 (5)			1384,305 (8)		1384,2931 (20)	1384,2931 (20)
	1421,1 (1)	1420,9 (5)	1420,081 (50)	1420,07 (5)		1420,07 (5)
	1465,6 (1)	1465,6 (1)				1465,6 (1)
1475,80 (5)			1475,305 (12)		1475,7792 (23)	1475,7792 (23)
1505,05 (5)			1505,039 (8)		1505,0280 (20)	1505,0280 (20)
1562,37 (5)			1562,305 (9)		1562,2940 (18)	1562,2940 (18)
	1572,3 (2)		1572,4 (2)			1572,4 (2)
1592,8 (1)	1593,0 (2)	1593,1 (4)	1592,672 (95)	1592,66 (10)		1592,80 (15)
	1630,0 (2)	1630,0 (1)	1629,692 (63)	1629,68 (6)		1629,75 (15)
	1698,5 (2)	1698,9 (1)				1698,8 (2)
1775,6 (2)	1775,4 (1)	1775,4 (2)	1775,422 (39)	1775,41 (4)		1775,41 (4)
1783,4 (2)	1783,6 (1)	1783,4 (2)	1783,480 (30)	1783,46 (3)		1783,46 (3)
1903,9 (2)	1903,4 (1)	1904,1 (8)	1903,530 (35)	1903,52 (4)		1903,52 (4)
	2004,6 (1)	2003,8 (8)	2004,74 (10)	2004,72 (10)		2004,65 (10)

^a The uncertainties of 0,1 keV are from a general statement and not specific to each γ -ray.

^d Reported to be a doublet.

The relative γ -ray intensities for the decay of $^{110}\text{Ag}^m$ (249 d) are given in Table 2. The adopted values are the weighted averages computed by the Limitation of Relative Statistical Weight method (1985ZiZY, 1992Ra09) and take into account the measurements from 1976De, 1977Ge12, 1979Ve03, 1980Ro22, 1980Yo05, 1981Ma09, 1990Me15, and 1993Ki18.

The γ -ray energies in Table 2 that are flagged with a "c" are from the evaluation 2000He14 and are considered especially suitable for energy calibration.

Table 2. Relative γ -ray intensities for $^{110}\text{Ag}^m$ decay

Energy (keV)	1969Br03 1972Ph04 ^a	1976De	1977Ge12	1979Ve03	1980Ro22	1980Yo05	1981Ma09	1990Me15	1993Ki18	LRSW average	χ_R^2 if > 1,0	σ_{int}	σ^{ext}	σ_{LWM}
116,48 (5)	isomeric decay							0,085 (3)						
120,23 (3)	<0,15			0,17 (3)			0,18 (1)	0,19 (1)	0,66(1) ^e	0,179 (9)				
133,333(7)	0,9 (2)			0,86 (13)			0,80 (5)	0,77 (3)	0,78 (2)	0,780 (16)				
219,348(8)	1,3 (3)			0,80 (6)			0,77 (5)	0,70 (2)	0,81 (1) ⁱ	0,76 (5)	5,8	0,013	0,030	0,046
221,079 (10)	1,1 (3)			0,80 (11)			0,74 (5)	0,72 (1)	0,67 (3)	0,716 (10)	1,1	0,009	0,010	
229,423 (23)	0,32 (15)			0,19 (5)			0,11 (1)	0,128 (8) ⁱ	0,22 (3)	0,126 (14)	4,7	0,007	0,014	
264,25 (6)							0,070 (7)	0,059 (5)	0,11 (3)	0,064(6)	2,0	0,004	0,006	
266,913 (12)	0,5 (1)			0,65 (6)			0,37 (2)	0,43 (1) ⁱ	0,53 (4)	0,43 (4)	9,5	0,012	0,037	
341,3 (2)							0,06 (3)	0,022 (4)	0,13 (9)	0,023 (5)	1,5	0,004	0,005	
356,43(10)							0,06 (3)	0,045 (3)	0,04 (2)	0,045 (3)				
360,23 (8)				0,14 (2)			0,11 (5)	0,035(7) ⁱ	0,09 (5)	0,08 (5)	5,4	0,012	0,028	0,048
365,448 (10)	1,1 (2)			1,27(14)		0,91 (19)	0,92 (5)	1,02 (8)	1,10 (12)	0,98 (5)	1,8	0,038	0,050	
387,073(9)	0,43 (9)			0,54 (13)		0,8 (4)	0,54 (3)	0,55 (1)	0,61 (24)	0,549 (9)				
396,895 (23)	0,36 (8)			0,68 (12)		0,6 (3)	0,35 (2)	0,43 (1) ⁱ	0,30 (10)	0,39 (4)	3,8	0,014	0,027	0,036
409,4 (5)							0,08 (4)	0,068 (7)	0,01 (4)	0,067 (7)	1,1	0,007	0,007	
446,812 (3) ^c	35 (2)		38,6 (4)	41,8 (6) ^e	39,0 (12)	39,55 (28)	39 (2)	38,9 (6)	38,22 (12) ⁱ	38,7 (5)	2,9	0,15	0,25	0,48
467,03 (4)				0,35 (5)			0,26 (2)	0,26 (5)	0,21 (5)	0,264 (19)	1,4	0,016	0,019	
493,43(10)				0,06 (2)			0,10 (2)	0,11 (1)	0,13 (4)	0,101 (11)	1,8	0,008	0,011	
544,55 (5)				0,10 (2)			0,19 (1)	0,22 (1)	0,15 (6)	0,19 (3)	9,8	0,007	0,021	0,027
572,8 (2)							0,19 (1)	0,13 (3)	0,14 (6)	0,183 (13)	2,1	0,009	0,013	

Energy (keV)	1969Br03 1972Ph04 ^a	1976De	1977Ge12	1979Ve03	1980Ro22	1980Yo05	1981Ma09	1990Me15	1993Ki18	LRSW average	χ_R^2 if > 1,0	σ_{int}	σ^{ext}	σ_{LWM}
603,08(10)							0,20 (3)	0,042 (9) ⁱ	0,30 (12)	0,12 (8)	8,2	0,021	0,059	0,081
620,3553 (17) ^c	29 (2)		29,3 (3)	29,5 (4)	31,4 (13)	29,65 (19)	28,0 (14)	29,4 (5)	28,00 (15) ⁱ	28,8 (8)	10,1	0,10	0,32	0,8
626,258 (10)	1,85 (20)			2,2 (2)		2,28 (14)	2,3 (1)	2,48 (4)	2,10 (3) ⁱ	2,27 (17)	12,7	0,025	0,09	0,17
630,62 (6)							0,30 (2)	0,40 (1) ⁱ	0,30 (8)	0,35 (5)	6,6	0,014	0,035	0,050
647,8 (4)							0,19 (4)		0,186 (4)	0,185 (5)	1,6	0,004	0,005	
657,7600 (11) ^c	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000				
666,6 (5)							0,16 (2) ⁱ		0,43 (5)	0,30 (14)	14,6	0,035	0,14	
676,58(10)								1,5 (1)						
677,6217 (12) ^c	122 (7)		113,1(11)	111 (2)	112,6 (29)	110,9 (8)	112 (6)	112 (2)	112,6 (11)	111,9 (5)				
687,0091 (18) ^c	74 (6)		68,5 (7)	75,8 (14) ^e	69,0 (27)	68,0 (6)	67 (3)	68,5 (5) ⁱ	69,2 (21)	68,3 (3)				
706,6760 (15) ^c	172 (7)	175 (10)	176,7 (18)	175,4 (20)	176,2 (22)	176,6 (10)	174 (7)	172,8 (5) ⁱ	176,9 (26)	174,6 (7)	1,9	0,5	0,6	
708,128 (20)							2,0 (2)	2,9 (2)	2,4 (3)	2,4 (5)	5,1	0,11	0,29	0,46
714,9 (1)							0,09 (2)		0,17 (6)	0,098 (24)	1,6	0,019	0,024	
744,2755 (18) ^c	44 (4)		49,2 (5)	52,3 (8)	49,5 (16)	50,00 (27)	48,0 (25)	49,3 (8)	50,2 (14)	49,9 (3)	2,0	0,21	0,31	
763,9424 (17) ^c	240 (8)	237 (2)	236,0 (24)	243,7 (30)	237,4 (31)	235,5 (9)	243 (12)	236 (3)	239,1 (53)	236,4 (7)	1,1	0,70	0,74	
774,70 (10)							0,03 (2)	0,02 (1)	0,092 (4) ⁱ	0,06 (3)	15,4	0,006	0,025	0,035
818,0244 (18) ^c	78 (3)		77,3 (8)	80,5 (10)	77,4 (17)	77,6 (4)	79 (4)	77,1 (5)	78,8 (18)	77,7 (4)	1,7	0,27	0,35	
845,8 (1)							0,10 (3)		0,10 (2)	0,10 (2)				
884,6781 (13) ^c	796 (20)	775 (5)	769 (8)	811 (10)	780 (10)	767,6 (26)	800 (40)	771 (10)	706,6 (12) ⁱ	784 (12)	13,3	1,5	5,3	12,5
927,6 (1)							0,065 (10)		0,067 (8)	0,063 (6)				
937,483 (3) ^c	365 (11)	366 (3)	362,2 (36)	380 (4)	369 (4)	363,1 (12) ⁱ	374 (18)	363 (6)	376 (8)	365,7 (26)	2,7	1,2	1,9	2,6

Energy (keV)	1969Br03 1972Ph04 ^a	1976De	1977Ge12	1979Ve03	1980Ro22	1980Yo05	1981Ma09	1990Me15	1993Ki18	LRSW average	χ_R^2 if > 1,0	σ_{int}	σ^{ext}	σ_{LWM}
957,35(10)				0,28 (5)			0,11 (1)	0,08 (1)	0,14 (5)	0,099 (19)	6,2	0,007	0,017	0,019
997,243 (15)	1,4 (2)			1,6 (1)		1,42 (5)	1,4 (1)	1,32 (4)	1,33 (10)	1,36 (4)	1,8	0,033	0,043	
1018,95(8)	0,3 (1)			0,17 (5)			0,15 (1)	0,15 (1)	0,08 (5)	0,149 (7)				
1050,5 (5)							0,08 (1)		0,08 (6)	0,08 (1)				
1085,447 (14)	0,58 (8)			0,95 (10)		0,66 (12)	0,74 (4)	0,71 (2)	0,81 (24)	0,76 (4)	1,2	0,035	0,0371	
1117,46(3)	0,39 (7)			0,55 (20)		0,41 (6)	0,52 (3)	0,52 (1)	0,38 (20)	0,517 (9)				
1125,699 (20)	0,26 (6)			0,35 (10)		0,38 (8)	0,34 (2)	0,30 (2)	0,22 (21)	0,322 (14)				
1163,14(8)				1,5 (1)			0,54 (5) ⁱ	0,79 (7)	1,0 (4)	0,78 (24)	23,4	0,04	0,19	0,24
1164,94(9)				0,96 (10) ^e			0,42 (5)	0,50(5)	0,47 (4)	0,46 (3)				
1186,7 (1)								0,015 (5)	0,0170 (5)	0,0170 (5)				
1251,04(4)	0,58 (19)			0,52 (5)		0,24 (7)	0,31 (2)	0,26 (1) ⁱ	0,25 (2)	0,28 (3)	7,5	0,0090	0,0260	
1300,05 (10)				0,20 (2)		0,25 (8)	0,19 (1)	0,21 (1)	0,22 (11)	0,200 (7)				
1334,326 (17)	1,55 (20)			1,8 (1)		1,49 (6)	1,40 (7)	1,49 (5) ⁱ	1,55 (33)	1,50 (5)	2,8	0,03	0,05	
1384,2931 (20) ^c	277 (8)	261 (2)	257,0 (26)	277,9 (30)	271 (5)	256,6 (8) ⁱ	278 (14)	261 (5)	276,6 (26)	262 (5)	12,8	0,8	2,9	5,0
1420,07(5)						0,39 (3)	0,27 (2)	0,24 (2)	0,37 (9)	0,28 (4)	6,2	0,013	0,032	0,041
1465,6 (1)							0,019 (2)							
1475,7792 (23) ^c	45,0 (20)		42,1 (4)	44,8 (6)	44,9 (12)	42,22 (17) ⁱ	45 (2)	42,4 (8)	45,7 (13)	42,7 (5)	4,6	0,20	0,43	0,5
1505,0280 (20) ^c	148 (4)	139 (1)	138,4 (14)	145,2 (16)	147,0 (29)	137,8 (5) ⁱ	151 (7)	140,1 (19)	149,2 (28)	139,4 (16)	6,1	0,45	1,1	1,6
1562,2940 (18) ^c	13,3 (6)		12,50(13) ⁱ	13,2 (2)	14,0 (8)	10,87 (7)	13,0 (7)	12,6 (6)	13,5 (4)	12,8 (3)	3,4	0,11	0,21	0,30
1572,4 (2)								0,012 (3)						
1592,80 (15)				0,4 (1)		0,221 (13)	0,20 (2)	0,22 (1)	0,34 (18)	0,219 (8)	1,2	0,007	0,0081	

Energy (keV)	1969Br03 1972Ph04 ^a	1976De	1977Ge12	1979Ve03	1980Ro22	1980Yo05	1981Ma09	1990Me15	1993Ki18	LRSW average	χ_R^2 if > 1,0	σ_{int}	σ^{ext}	σ_{LWM}
1629,75 (15)						0,061 (11)	0,036 (4)	0,046 (5)	0,11 (5)	0,042 (5)	2,6	0,003	0,005	
1698,8 (2)							0,019 (2)		0,012 (4)	0,018 (3)	2,4	0,002	0,003	
1775,41(4)				0,067(10)		0,067 (11)	0,076 (4)	0,063 (4)	0,07 (6)	0,069 (3)	1,4	0,0026	0,0031	
1783,46(3)				0,085 (30)		0,103 (11)	0,110 (6)	0,092 (3)	0,07 (4)	0,107 (5)				
1903,52(4)				0,20 (2)		0,158 (15)	0,18 (1)	0,16 (1)	0,15 (2)	0,169 (7)	1,5	0,006	0,007	
2004,65 (10)							0,012(1) ⁱ	0,011 (2)	0,028 (4)	0,013 (4)	7,7	0,0013	0,0035	

a The values from these two articles, by the same authors, are for comparison and were not used in the calculated averages.

c γ -ray energy is from the 2000He14 evaluation and is useful for energy calibrations.

e Value was not used in the calculation of the average.

i The published uncertainty, which is given, was increased in the LRSW analysis to reduce the relative weight to 50 %.

The mixing ratios for the M1+E2 γ -rays have been evaluated in this work (from references 1962Ka07, 1963Su07, 1964Ne05, 1970Kr03, 1973Jo08, 1978Wa07, 1979Ve03, 1980Ru03, 1990Ke02, and 1993Ki18). The results are very similar to those in the most recent ENSDF evaluation (2000De11), so those from ENSDF have been used. From the measurements of 1979Ve03, mixing ratios for M3 contributions to predominantly E2 transitions are quoted in ENSDF. The δ (M3/E2) values that do not include 0,0 in their uncertainties are those of 763 and 1562 keV γ -rays; both are $\delta = -0,10 (+2-3)$. Although the conversion coefficients are small, the high precision of the relative γ -ray intensities makes them significant; for example, $\alpha_{(657)} = 0,00318$.

The normalization of the relative emission probabilities for the γ -rays from the decay of ¹¹⁰Ag^m (249 d) is determined by requiring that the sum of the γ -ray transition intensities to the ground states of ¹¹⁰Cd and ¹¹⁰Ag be 100 % of the decays of the isomeric state. However, the 657 keV γ -ray occurs in both the direct β^- decay and that which follows the isomeric decay. Since 4,6(4) % the ground state decays lead to the 657-keV γ ray, the intensity of the isomeric decay is reduced by this fraction in computing the intensity feeding the ground states.

Then, in the units of Table 2, one has $I_{\gamma(116)}[1+\alpha_{(116)}][0,954] + I_{\gamma(657)}[1+\alpha_{(657)}] + I_{\gamma(1475)} + I_{\gamma(1783)} = 0,085[169][0,954] + 1000[1,003] + 42,7 + 0,107$. If an uncertainty of 5 % is assigned to $\alpha_{(116)}$, this sum is 1059,5 (9), so the normalization factor for the γ -ray intensities in Table 2 is 0,09438 (8).

The resulting intensity of the isomeric decay branch is then $0,085[0,09438][169] = 1,36$ with an uncertainty of 0,08 and that of the β^- decay is 98,64 (8) %. This gives the 657 keV photon intensity of 94,38 (8) per 100 decays of the isomeric state.

The isomeric decay of ¹¹⁰Ag^m (249 d) occurs via an M4 γ -ray of 116,48 (5) keV with $\alpha = 168$ [i.e., $P_{\gamma} = 0,0080$ (4)] followed by an E1 γ -ray of 1,113 keV energy. The γ -rays following the β^- decay of the ground state are all very weak due to the small isomeric decay branch (1,36 %) and the large β^- branch to the ground state (95,1 %). Also, the 4,6 % branch to the 657 level is already included in Table 2. Therefore, the remaining γ -rays following the β^- decay of the ground state are neglected.

The γ -ray multipolarities and mixing ratios were taken from the 2000De11 evaluation and are as follows:

E1: 603, 1421-keV

E1(+M2): 409 [$\delta = -0,029(23)$]; 997 [$\delta = -0,30(46)$]; 1117 [$\delta = +0,021(44)$]; 1300 [$\delta = +0,0(1)$]

E2: 626, 657, 884, 1085, 1334, 1475, 1592, 1783, 2004

(E2): 467; 774

M1(+E2): 120 [$\delta = -0,13(33)$]

M1+E2: 446 [$\delta = -0,38(2)$]; 544; 620 [$\delta = -0,50(4)$]; 677 [$\delta = 0,36(2)$]; 687 [$\delta = -1,76(6)$]; 706 [$\delta = -1,42$ (7)]; 708 [$\delta = -0,15(9)$]; 818 [$\delta = -1,36(7)$]; 957 [$\delta = -0,9(7)$]; 1018 [$\delta = -0,56(35)$]; 1125 [$\delta = +0,33(8)$]; 1163 [$\delta = -0,03(+6-9)$]; 1164 [$\delta = +0,0(3)$]; 1384 [$\delta = -0,44(2)$]; 1505 [$\delta = -1,21(4)$]; 1629 [$\delta = +0,06(3)$]; 1697; 1775

E2(+M3): 744 [$\delta = -0(+16-10)$]; 937 [$\delta = -0,07(+7-3)$]; 1562 [$\delta = -0,10(+2-3)$]

M3+E2: 763 [$\delta = -0,10 (+2-3)$]

4) Atomic data

From the EMISSION code and the decay data, the following information was obtained.

Quantity	Ag (Z=47)	Cd (Z=48)
ω_K	0,831 (4)	0,842 (4)
ω_L average	0,0583 (14)	0,0632 (16)
n_{KL}	0,964 (4)	0,953 (4)
$K_{\alpha 2}/K_{\alpha 1}$	0,5305 (25)	0,5317 (25)
K_{β}/K_{α}	0,2125 (17)	0,2151 (18)

Due the high energy of the strong transitions, the Auger electrons are negligible and no related data are included here.

The K X-ray emission probabilities are calculated as follows:

For the decay of ¹¹⁰Ag^m (249 d), Ag KX-rays per 100 decays of parent

$K_{\alpha 2}$	0,198 (12)
$K_{\alpha 1}$	0,372 (22)
K_{β}	0,121 (7)

Cd KX-rays per 100 decays of the parent

$K_{\alpha 2}$	0,153 (9)
$K_{\alpha 1}$	0,288 (16)
K_{β}	0,095 (6)

5) β^- decay intensities

The β^- decay intensities for the decay of the ¹¹⁰Ag ground state are simply deduced from the above data and the γ -ray intensity balances. Since the spin of the isomeric state is large, namely 6, there are several β^- decay branches for which the $\log ft$ systematics (1998Si17) given lower limits on the intensities than can be derived from the intensity balances. These data are given in Table 3

Table 3. Data used to deduce β^- decay intensities and $\log ft$ values.

Level(keV)	J^π	$\Delta J, \Delta \pi$	$\log ft$ limit	I_β from $\log ft$ limit	I_β from intensity balance	I_β adopted	$\log ft$
0	0 ⁺	6,no			1,3 (4)	0	
657	2 ⁺	4,no	>22	<10 ⁻¹⁰	-1,2 (12)	0	
1475	2 ⁺	4,no	>22	<10 ⁻¹⁰	0,08 (8)	0	
1522	4 ⁺	2,no	>10,6	<6	0,8 (13)	<2	>11
1783	2 ⁺	4,no	>22	<10 ⁻¹¹	0,0156 (23)	0	
2078	3 ⁻	3,yes	>16,5	<10 ⁻⁶	0,002 (8)	<10 ⁻⁶	>16,5
2162	3 ⁺	3,no	>13,9	<0,0004	-0,01 (19)	<0,0004	>13,9
2220	4 ⁺	2,no	>10,6	<0,6	0,06 (9)	<0,15	>11,2

Level(keV)	J ^π	ΔJ,Δπ	logft limit	I _β from logft limit	I _β from intensity balance	I _β adopted	logft
2250	4 ⁺	2,no	>10,6	<0,6	0,06 (5)	0,06 (5)	11,5
2287	2 ⁺	4,no	>22	<2x10 ⁻¹²	0,0040 (5)	0	
2356	(1 ⁺ ,2 ⁺)	4 or 5, no	>22	<10 ⁻¹²	0	0	
2433	3 ⁺	3,no	>13,9	<0,0001	-0,008 (6)	0	
2479	6 ⁺	0,no			30,8 (3)	30,8 (3)	8,282
2539	5 ⁻	1,yes			0,060 (4)	0,060 (4)	10,82
2561	4 ⁺	2,no	>10,6	<0,1	-0,003 (7)	<0,005	>11,8
2659	5 ⁻	1,yes			0,031 (4)	0,031 (4)	10,67
2662					0	0	
2705	4 ⁺	2,no	>10,6	<0,03	0,006 (23)	<0,029	>10,5
2707	4 ⁺	2,no	>10,6	<0,03	-0,010 (7)	0	
2793	4 ⁺	2,no	>10,6	<0,03	-0,013 (7)	0	
2842	5 ⁻	1,yes			0,0252 (10)	0,0252 (10)	9,73
2876	6 ⁺	0,no			0,392 (18)	0,392 (18)	8,23
2926	5 ⁺	1,no			67,5 (6)	67,5 (6)	5,36

6) References

- 1938Li07 - J. J. Livingood, G. T. Seaborg, Phys. Rev. 54 (1938) 88 [T_{1/2}]
1950Gu54 - J. R. Gum, M. L. Pool, Phys. Rev. 80 (1950) 315 [T_{1/2}]
1963Su07 - T. Suter, P. Reyes-Suter, W. Scheuer, Nucl. Phys. 47 (1963) 251 [E_γ, I_{e-}]
1964Ne05 - W. B. Newbolt, J. H. Hamilton, Nucl. Phys. 53 (1964) 353 [E_γ, I_{e-}, α_K, Mult]
1964Sc06 - J. Schintlmeister, L. Werner, Nucl. Phys. 51 (1964) 383 [E_β, I_⊗, I_{e-}]
1969Br03 - S. M. Brahmavar, J. H. Hamilton, A. V. Ramayya, E. F. Zganjar, C. E. Bemis Jr., Nucl. Phys. A125 (1969) 217 [E_γ, I_γ]
1970Kr03 - K. S. Krane, R. M. Steffen, Phys. Rev. C2 (1970) 724 [δ]
1970Su03 - S. P. Sud, P. C. Mangal, P. N. Trehan, Aust. J. Phys. 23 (1970) 87 [δ]
1972Ph04 - G. B. Philips, S. M. Brahmavar, J. H. Hamilton, T. Kracikova, Nucl. Phys. A182 (1972) 606 [E_γ, I_γ]
1973Ga10 - P. L. Gardulski, M. L. Wiedenbeck, Phys. Rev. C7 (1973) 2080 [δ]
1973Jo08 - P. D. Johnston, N. J. Stone, Nucl. Phys. A206 (1973) 273 [δ]
1974Pr07 - W. W. Pratt, J. Inorg. Nucl. Chem. 36 (1974) 1199 [E_γ, I_γ]
1976De - K. Debertain, U. Schötzig, K. F. Walz, H. M. Weiss, Proc. ERDA Symposium on X- and Gamma-ray Sources and Applications, Ann. Arbor. (1976) 59 [I_γ]
1976WaZH - K. F. Walz, H. M. Weiss, K. Debertain, Priv. Comm. (Octobre 1976) [T_{1/2} as cited in Nuclear Data Sheets 38 (1983) 545]
1977Ge12 - R. J. Gehrke, R. G. Helmer, R. C. Greenwood, Nucl. Instr. Meth. 147 (1977) 405 [I_γ]
1978Ke14 - J. Kern, S. Schwitz, Nucl. Instr. Meth. 151 (1978) 549 [E_γ]
1978Wa07 - G. W. Wang, A. J. Becker, L. M. Chirovsky, J. L. Groves, C. S. Wu, Phys. Rev. C18 (1978) 476 [δ]

- 1979Co14 – E. J. Cohen, H. R. Andrews, T. F. Knott, F. M. Pipkin, D. C. Santry, Phys. Rev. C20 (1979) 847 [δ]
- 1979Ve03 – H. R. Verma, A. K. Sharma, P. Kaur, K. K. Suri, P. N. Trehan, J. Phys. Soc. Japan 47 (1979) 16 [E_γ , I_γ , δ]
- 1979Sc31 - P. Schlüter, G. Soff, Atomic Data Nuclear Data Tables 24 (1979) 509 [α_π]
- 1980Ba58 – V. V. Babenko, I. N. Vishnevskii, V. A. Zheltonozhskii, V. P. Svyato, V. V. Trishin, Bull. Acad. Sci. (USSR), Phys. Ser. 44,#5, (1980) 132 [$\gamma\gamma(\theta)$, δ]
- 1980Ho17 – H. Houtermans, O. Milosevic, F. Reichel, Intern. J. Appl. Radiat. Isot. 31 (1980) 153 [$T_{1/2}$]
- 1980Ro22 - W. M. Roney, Jr., W. A. Seale, Nucl. Instr. Meth. 171 (1980) 389 [I_γ]
- 1980Ru03 - W. D. Ruhter, D. C. Camp, Nucl. Instr. Meth. 173 (1980) 489 [δ]
- 1980Yo05 – Y. Yoshizawa, Y. Iwata, T. Katu, T. Katoh, J.-Z. Ruan, T. Kojima, Y. Kawada, Nucl. Instr. Meth. 174 (1980) 109 [I_γ]
- 1981Ma09 - G. Mallet, J. Phys. Soc. Japan 50 (1981) 384 [E_γ , I_γ]
- 1981Ma25 - G. Mallet, J. Dalmasso, H. Maria, G. Ardisson, J. Phys. G – Nucl. Phys. 7 (1981) 1259 [scheme]
- 1983Me17 - R. A. Meyer, T. N. Massey, Intern. J. Appl. Radiat. Isot. 34 (1983) 1073 [E_γ]
- 1983Wa26 – K. F. Walz, K. Debertain, H. Schrader, Inter. J. Appl. Radiat. Isot. 34 (1983) 1191 [$T_{1/2}$]
- 1985ZiZY - W. L. Zijp, Report ECN FYS/RASA-85/19 (1985) [averages]
- 1988Kr03 - K. S. Krane, N. S. Schulz, Phys. Rev. C37 (1988) 747 [δ]
- 1990Me15 - R. A. Meyer, Fizika 22 (1990) 153 [E_γ , I_γ]
- 1991Ba63 – I. M. Band, M. B. Trzhaskovskaya, Bull. Acad. Sci. (USSR) , Phys. Ser. 55,#11 (1991) 39 [α]
- 1992Gr18 – R. C. Greenwood, R. G. Helmer, M. A. Lee, M. H. Putnan, M. A. Oates, D. A. Strttrmann, K. D. Watts, Nucl. Instr. Meth. A314 (1992) 514 [I_β]
- 1992Ra08 - M. U. Rajput, T. D. MacMahon, Nucl. Instr. Meth. A312 (1992) 289 [averages]
- 1993Ki18 – L. L. Kiang, P. K. Teng, G. C. Kiang, W. S. Chang, P. J. Tu, J. Phys. Soc. Japan 62 (1993) 888 [E_γ , I_γ , δ]
- 1995Au04 - G. Audi, A. H. Wapstra, Nucl. Phys. A595 (1995) 409 [Q]
- 1998Si17 – B. Singh, J. L. Rodriguez, S. S. M. Wong, J. K. Tuli, Nucl. Data Sheets 84 (1998) 487 [logft systematics]
- 2000De11 - D. DeFrenne, E. Jacobs, Nucl. Data Sheets 89 (2000) 481 [J^π , multipolarities, δ]
- 2000He14 - R. G. Helmer, C. van der Leun, Nucl. Instr. Meth. A450 (2000) 35 [E_γ]

¹¹¹In - Comments on evaluation of decay data by V.P. Chechev.

The initial ¹¹¹In decay data evaluation was done by V.P. Chechev in 1998 (1999Be). This current (revised) evaluation has been carried out in March 2006. The literature available by March 2006 has been included.

1 Decay Scheme

Transitions to the ground state and the excited level of 245 keV of ¹¹¹Cd have not been observed. Limits on the electron capture branches to these levels can be deduced from the log ft systematics of 1998Si17. The transitions to the levels at 0 and 245 keV are 4th and 2nd forbidden with expected log ft's of > 22 and > 10.6, respectively. The corresponding electron capture branch limits are < 1.0×10⁻¹⁴ % and < 5×10⁻⁴ %, respectively (2003B110).

The upper limit of 0.01 % has been found for the electron capture branch to the excited level of 396 keV by Meyer and Landrum (1972MeZD).

2 Nuclear Data

Q_{EC} value is from 2003Au03.

The evaluated ¹¹¹In half-life is based on the experimental data given in Table 1.

Table 1. Experimental values of the ¹¹¹In half-life (in days)

<u>Reference</u>	<u>Author(s)</u>	<u>Value</u>	<u>Comments</u>
1949He06	Helmholz <i>et al.</i>	2.84 (3)	
1957Ma26	Maier	2.81 (1)	
1968Li08	Liskien	2.84 (11)	
1968Sm08	Smend <i>et al.</i>	2.96 (8)	
1972Em01	Emery <i>et al.</i>	2.83 (1)	
1972Gu19	Gureev <i>et al.</i>	2.84	Uncertainty is not quoted
1978La21	Lagoutine <i>et al.</i>	2.802 (1)	Quoted uncertainty, corresponding to 99.7 % confidence level, has been reduced by a factor 3
1980Ho17	Houtermans <i>et al.</i>	2.8071 (15)	
1982HoZY	Hoppes <i>et al.</i>	2.8048 (5)	Replaced by 1992Un01
1983Wa26	Walz <i>et al.</i>	2.8049 (5)	
1986Ru09	Rutledge <i>et al.</i>	2.8048 (1)	
1992Un01	Unterweger <i>et al.</i>	2.80477 (53)	Cited also in 2002Un02
2004Sc04	Schrader	2.8063 (7)	

The value of 1972Gu19 has been omitted because of the absence of an estimated uncertainty. The value of 1982HoZY has been omitted as it is replaced in 1992Un01. The value of 1968Sm08 has been omitted as outlier using the Chauvenet's criterion. Hence the eleven values have been used for the statistical data processing.

The uncertainty of 1986Ru09 was increased to 0.00030 to adjust weights according to the LRSW method. A weighted average for the final data set is 2.8049 with an internal uncertainty of 0.00021 and an external uncertainty of 0.00034 and a reduced $\chi^2/\nu = 2.5$. An unweighted average is 2.815 (5).

Different statistical procedures (1994Ka08) give the following results: UINF, PINF and NORM- 2.8049 (3), LWM - 2.815 (10), IEXW - 2.805 (13), RAJ - 2.8049 (2), BAYS and MBAYS - 2.8049 (4).

The adopted value of the ¹¹¹In half-life is 2.8049 (4) days.

The evaluated half-life of the metastable level of 396 keV (^{111m}Cd) is based on the experimental results given in Table 2.

Table 2. Experimental values of the ^{111m}Cd half-life (in minutes)

Reference	Author(s)	Value
1945Wi11	Wiedenbeck	48.7 (3)
1948Ho37	Hole	50 (2)
1949He06	Helmholz et al.	48.6 (3)
1968Bo28	Bornemisza-Pauspertl et al.	49.4 (7)
1987Ne01	Nemeth et al.	48.54 (5)
1997We13	Wen et al.	48.30 (15)

The uncertainty of 1987Neo1 was increased to 0.12 to adjust weights according to the Limitation of Relative Statistical Weight (LRSW) method. A weighted average for the final data set is 48.50 with an internal uncertainty of 0.085 and an external uncertainty of 0.082 and a reduced $\chi^2/\nu = 0.93$. An unweighted average is 48.9 (3).

Different statistical procedures (1994Ka08) give the following results: IEXW, LWM, MBAYS, NORM and UINF – 48.50 (9), PINF – 48.50 (8), RAJ – 48.51(9), BAYS – 48.50 (11).

The adopted value of the ¹¹¹In half-life is 48.50 (9) minutes.

2.1 Electron Capture Transitions

The electron capture transition energies have been calculated from Q_{EC} value and the ¹¹¹Cd level energies given in Table 3 from 2003Bi10. The electron capture transition probability $P_{0,2} = 5 (5) 10^{-3}$ has been evaluated taking into account the observed upper limit of 1×10^{-2} (1972MeZD). The fractional electron capture probabilities P_K, P_L, P_M have been calculated using the LOGFT computer program.

Table 3. ¹¹¹Cd levels populated in the ¹¹¹In ϵ -decay

Level number	Energy, keV	Spin and parity	Half-life	Probability of EC-transition (x100)
0	0.0	1/2 ⁺	Stable	$< 1.0 \times 10^{-14}$
1	245.35 (4)	5/2 ⁺	84.5 ns	$< 5 \times 10^{-4}$
2	396.16 (5)	11/2 ⁻	48.50 min	0.005 (5)
3	416.63 (5)	7/2 ⁺	0.12 ns	99.995 (5)

2.2 g Transitions

The energies of γ -ray transitions are virtually the same as the γ -ray energies because nuclear recoil is negligible. The γ -ray transition probabilities have been calculated from the γ -ray emission probabilities and the evaluated total internal conversion coefficients (α_T).

The evaluated α_T values for $\gamma_{1,0}$ (245 keV) and $\gamma_{3,1}$ (171 keV) gamma-ray transitions have been obtained from the sets of 5 data including theoretical values (Table 4). The values of $\alpha_K, \alpha_L, \alpha_M$ have been calculated from the evaluated α_T using the theoretical ratios $\alpha_K/\alpha_L/\alpha_M/\alpha_{NO}$. The relative uncertainties of $\alpha_K, \alpha_L, \alpha_M$ have been taken as 2 %.

The theoretical α_T has been used for the E3 $\gamma_{2,1}$ (151 keV) gamma-ray transition (see also 1973Pathak).

Table 4. Experimental, theoretical and evaluated values of the total internal conversion coefficients (α_T)

	1956St64	1966Sp04	1975Sh29	1985Ka29	Theory (2006Ra03)	Evaluated
$\gamma_{1,0}$ (245 keV)	0.0621 (15)	0.0618 (15)	0.0634 (30)	0.0620 (7)	0.0637 (9)	0.0625 (7)
$\gamma_{3,1}$ (171 keV)	0.099 (3)	0.100 (3)	0.124 (6)	0.1018 (13)	0.1068 (15)	0.1036 (24)

The theoretical α_T values have been calculated using the BRICC computer program (2006Ra03).

The gamma-ray transition multiplicities have been adopted from measurements of 1956St54 and 1974Kr03. The gamma-ray multipolarity mixing ratio $\delta(E2/M1)$ of the $\gamma_{3,1}$ (171 keV)-transition has been evaluated using the following data:

0.146(3)	Steffen (1956St64)
0.141(3)	Budz-Jorgensen (1973)
0.145	Kreische and Lampert (1974Kr03)
0.144(3)	Weighted average of 1956St04 and 1973Budz-Jorgensen

The adopted value of 0.144 (3) corresponds to an E2 admixture of 2.07 (9) %.

3 Atomic Data

3.1. Fluorescence yields

The fluorescence yield data ω_K , $\overline{\omega}_L$, n_{KL} are from 1996Sc06 (Schönfeld and Janßen).

3.2. X Radiations

The energy values for X-rays have been calculated from the wavelengths given by Bearden (1967Be65). The relative emission probabilities of KX ray components have been taken from 1996Sc06.

3.3. Auger Electrons

The energies of Auger electrons are from 1977La19 (Larkins) and Table of Isotopes. The ratios $P(KLX)/P(KLL)$, $P(KXY)/P(KLL)$ are taken from 1996Sc06.

4 Electron Emissions

The energies of the conversion electrons have been calculated from the gamma transition energies and the electron binding energies. The emission probabilities of conversion electrons have been deduced from the evaluated $P(\gamma)$ and ICC values.

The total absolute emission probability of K Auger electrons has been calculated with the EMISSION computer program using the adopted $\omega_K = 0.842$ (4).

The absolute total emission probability of L Auger electrons has been calculated with the EMISSION computer program using the adopted $\overline{\omega}_L = 0.0632$ (16).

Experimental data on conversion electrons (1951Mc61, 1966Sp04, 1975Sh29) and Auger electrons (2005Ya03) are concordant with the adopted values

5 Photon Emissions

5.1 X-ray Emissions

The absolute emission probabilities of Cd KX-rays have been calculated with the EMISSION computer program using the adopted values of P_K and ω_K (Cd).

The absolute emission probabilities of Cd LX-rays have been calculated with the EMISSION computer program using the adopted values of P_L , ω_L (Cd), P_K , ω_K (Cd), n_{KL} (Cd).

5.2 g-ray Emissions

The energy of $\gamma_{2,1}$ -ray (151 keV) has been taken from 1975Sh29.

The energy of the $\gamma_{3,1}$ -ray (171 keV) has been evaluated using the experimental results given below:

172.1 (5)	McGinnis (1951Mc11) - Omitted from data processing
171.29 (3)	Sparrman et al. (1966Sp04)
171.20 (10)	Heath (1974HeYW)
171.28 (3)	Shevelev et al. (1975Sh29)
171.28 (3)	Weighted average (adopted value)

The energy of the $\gamma_{1,0}$ -ray (245 keV) has been evaluated using the experimental results given below:

246.6 (7)	McGinnis (1951Mc11) - Omitted from data processing
245.35 (4)	Sparrman et al.(1966Sp04)
245.27 (10)	Heath(1974HeYW)
245.35 (4)	Shevelev et al. (1975Sh29)
245.35 (4)	Weighted average (adopted value)

The absolute emission probabilities of $\gamma_{2,1}$ (151 keV), $\gamma_{3,1}$ (171 keV) and $\gamma_{1,0}$ (245 keV) gamma rays have been calculated using the below relations:

$$P\gamma_{2,1} (\times 100) = 99.995 (5) / (1 + \alpha_T (\gamma_{2,1}))$$

$$P\gamma_{3,1} (\times 100) = 0.005 (5) / (1 + \alpha_T (\gamma_{3,1}))$$

$$P\gamma_{1,0} (\times 100) = 100 / (1 + \alpha_T (\gamma_{1,0})).$$

In 1975Sh29 the latter value has been estimated as ~ 0.003 .

The relative intensity of $\gamma_{1,0} / \gamma_{3,1}$ from 0.90 to 0.97 has been measured with an accuracy not better than 3 % in the above works. This accuracy is considerably worse in comparison with the calculation from the decay scheme using α_T values.

6 References

- 1945Wi11 - M. L. Wiedenbeck, Phys. Rev. 67(1945)92 [$T_{1/2}$ (^{111m}Cd)].
 1948Ho37 - N. Hole, Arkiv. Mat. Astron. Fysik 36A(1948)N09 [$T_{1/2}$ (^{111m}Cd)].
 1949He06 - A. S. Helmholtz, R. W. Hayward, C. L. McGinnis, Phys. Rev. 75(1949)1469A. See also 1951Mc11 [$T_{1/2}$ (^{111m}Cd), $T_{1/2}$ (¹¹¹In)].
 1951Mc11 - C. L. McGinnis, Phys. Rev. 81(1951)734 [$T_{1/2}$ (^{111m}Cd), E_γ , I_γ , Ice].
 1956St64 - R. H. Steffen, Phys. Rev. 103(1956)116 [$\delta(E2/M1)$ of $\gamma_{3,1}$ -transition].
 1957Ma26 - A. Maier, Helv. Phys. Acta 30(1957)611 [$T_{1/2}$ (¹¹¹In)].
 1966Sp04 - P. Sparrman, A. Marrelius, T. Sundstrom, H. Petterson, Z. Phys. B192(1966)439 [E_γ , I_γ , ICC].
 1967Be65 - J. A. Bearden, Rev. Mod. Phys. 39(1967)78 [E_X].
 1968Bo28 - P. Bornemisza-Pauspertl, J. Karolyi, G. Peto, ATOMKI Kozlemen 10(1968)112 [$T_{1/2}$ (^{111m}Cd)].
 1968Li08 - H. Liskien, Nucl. Phys. A118(1968)379 [$T_{1/2}$ (¹¹¹In)].

- 1968Sm08 - F. Smend, W. Weirauch, W.-D. Schmidt-Ott, Z. Phys. 214(1968)437 [$T_{1/2}$ (¹¹¹In)].
- 1972Em01 - J. F. Emery et al., Nucl. Sci. Eng. 48(1972)319 [$T_{1/2}$ (¹¹¹In)].
- 1972Gu19 - S. E. Gureev, T. Islamov, V. S. Usachenko, Izv. Akad. Nauk. SSSR, Ser. FizMat. 1(1972)87 [$T_{1/2}$ (¹¹¹In)].
- 1972MeZD - R. A. Meyer, J. H. Landrum, Bull. Am. Phys. Soc. 17(1972)906 [$P_{E0,2}$].
- 1973Budz-Jorgensen - C. Budz-Jorgensen, Phys. Rev. B8(1973)5411 [$\delta(E2/M1)$ of $\gamma_{3,1}$ -transition].
- 1973Pathak - B. P. Pathak, S. K. Mukherjee, Radiochem. Radioanal. Lett. 15(1973)187 [α_T of $\gamma_{2,1}$ -transition].
- 1974HeYW - R. L. Heath, ANCR-1000-2 (1974) [E_γ , I_γ].
- 1974Kr03 - W. Kreische, W. Lampert, Z. Phys. 266(1974)51 [$\delta(E2/M1)$ of $\gamma_{3,1}$ -transition].
- 1975Sh29 - G. A. Shevelev, A. T. Troytskaya, V. M. Kartashov, Izv. Akad. Nauk. SSSR, Ser. Fiz. 39(1975)2038 [E_γ , I_γ , ICC].
- 1977La19 - F. P. Larkins, Atomic Data and Nuclear Data Tables 20(1977)313 [E_{eAK} , E_{eAL}].
- 1978Ro22 - F. Rösler, H. M. Fries, K. Alder, H. C. Pauli, Atomic Data and Nuclear Data Tables 21(1978)92 [Theoretical ICC].
- 1978La21 - F. Lagoutine, J. Legrand, C. Bac, Int. J. Appl. Radiat. Isotopes 29(1978)269 [$T_{1/2}$ (¹¹¹In)].
- 1980Ho17 - H. Houtermans, O. Milosevic, F. Reichel, Int. J. Appl. Radiat. Isotopes 31(1980)153 [$T_{1/2}$ (¹¹¹In)].
- 1982HoZY - D. D. Hoppes et al., NBS Special Publication 626(1982)85 [$T_{1/2}$ (¹¹¹In)].
- 1983Wa26 - K. F. Walz, K. Debertin, H. Schrader, Int. J. Appl. Radiat. Isotopes 34(1983)1191 [$T_{1/2}$ (¹¹¹In)].
- 1985Ka29 - Y. Kawada, Y. Hino, Nucl. Instrum. Methods A241(1985)199 [ICC α_T].
- 1986Ru09 - A. R. Rutledge, L. V. Smith, J. S. Merritt, Int. J. Appl. Radiat. Isotopes 37(1986)1029 [$T_{1/2}$ (¹¹¹In)].
- 1987Ne01 - Zs. Nemeth, L. Lakosi, I. Pavlicsek, A. Veres, Int. J. Appl. Radiat. Isot. 38(1987)63 [$T_{1/2}$ (^{111m}Cd)].
- 1992Un01 - M. P. Unterweger, D. D. Hoppes, F. J. Schima, Nucl. Instrum. Meth. in Phys. Res. A312(1992)349 [$T_{1/2}$ (¹¹¹In)].
- 1994Ka08 - S. F. Kafala, T. D. MacMahon, P. W. Gray, Nucl. Instrum. Meth. Phys. Res. A339(1994)151 [Evaluation technique].
- 1996Sc06 - E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [$P(K\beta)/P(K\alpha)$, ω_K , ω_L , n_{KL}].
- 1997We13 - Xiao-qiong Wen et al., Nucl. Instrum. Meth. Phys. Res. A379 (1997) 478 [$T_{1/2}$ (^{111m}Cd)].
- 1998Si17 - B. Singh, J. L. Rodriguez, S. S. Wong, J. K. Tuli, Nucl. Data Sheets 84 (1998) 487 [$lg ft$].
- 1999Be - M.-M. Bé, B. Duchemin, J. Lame, C. Morillon, F. Piton, E. Browne, V. Chechev, R. Helmer, E. Schönfeld. Table de Radionucléides, CEA-ISBN 2-7272-0200-8. 1999. Comments on Evaluations, CEA-ISBN 2-7272-0211-3. 1999 [¹¹¹In decay data evaluation-1998].
- 2003Au03 - G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)3 [Q value].
- 2003Bl10 - J. Blachot, Nuclear Data Sheets 100 (2003) 179 [¹¹¹Cd level scheme and energies].
- 2004Sc04 - H. Schrader, Applied Radiation and Isotopes 60 (2004) 317 [$T_{1/2}$ (¹¹¹In)].
- 2005Ya03 - E. A. Yakushev et al., Applied Radiation and Isotopes 62 (2005) 451 [Auger electrons].
- 2006Ra03 - S. Raman, M. Ertugrul, C. W. Nestor, Jr., M. B. Trzhaskovskaya, At. Data Nucl. Data Tables 92(2006)207 [Theoretical ICC].

¹²³Te^m - Comments on evaluation of decay data by M. M. Bé and V. Chisté

This evaluation was completed in October 1993 and has been updated in September 2002. Several measurements of the gamma emission intensity and of the total internal conversion coefficient of the 159-keV line were carried out. The decay scheme has been constructed mainly from these measurements.

Nuclear Data

- Spins and parities are from the LPRI "Table de Radionucléides" [1]-
- The half-life value is the weighted average of : 11 9,7(3) (Emery 1970 – 1970EmZY) and 119,2(1) (Coursey 1992 – 1992Co11) ; its uncertainty is the internal uncertainty.

Gamma Transitions

- 88-keV gamma transition

For this M4 transition, the various theoretical conversion coefficients differ by about 5%. They are compared with measured values in the following table :

	Th. value Band 2002 – (2002Ba85)	Th. Value Rösel 1978 – (1978Ro22)	Exp. Value Kalinauskas 1969 – (1968Ka20)	Exp. Value Raman 1973 – (1973Ra32)	Exp. value Chu 1964 – (1964Ch18)
α_T	1099	1151	1000 (70)	1080 (40)	
α_K	463	483			455 (9)
α_L	493	517			482 (14)
α_M	118	124			

Values interpolated from the new Band *et al.* tables (2002Ba85), have been adopted following the recommendations of Gorozhankin (2002) [3].

The transition probability has been deduced from the decay scheme balance at the 159-keV level.

- 247-keV gamma transition

The conversion coefficients, for this E5 transition, were calculated using the new tables of Band *et al.* (2002Ba85) as suggested by Gorozhankin [2, 3]. The theoretical α_T (7,75 (30)) agrees with the measured value (8,1(4)) given by Raman (1973Ra32).

The transition probability has been deduced using this theoretical value for α_T and the gamma emission intensity (see below).

- 159-keV gamma transition

For the 159-keV gamma transition, the following values of the mixing ratio squared δ^2 have been found in the literature :

Reference	d ²	a _T
Goldberg <i>et al.</i> – (1955Go25)	0,013(1)	1,919 10 ⁻¹
Fagg <i>et al.</i> – (1955Fa40)	0,0034(20)	1,905 10 ⁻¹
Chu <i>et al.</i> – (1964Ch08)	0,0067(11)	1,909 10 ⁻¹
Gupta <i>et al.</i> – (1966Gu02)	0,011(8)	1,916 10 ⁻¹
Alkhazov <i>et al.</i> – (1964Al28)	0,004(5)	1,906 10 ⁻¹
Törnkvist <i>et al.</i> – (1969To02)	0,0119(9)	1,917 10 ⁻¹
Krane – (1977Kr13)	0,01232 (47) (adopted value)	1,918 10 ⁻¹

The internal conversion coefficients were calculated by ICC Computer Code [2] by interpolation of the Rösel tables (1978Ro22).

Elsewhere, the following measurements of the α_T coefficients were carried out :

Chu1964 (1964ch08)	0,1964 (74)
Hatch1966 (1966Ha03)	0,1979 (54)
Janssen1992 (1999Ja15)	0,1932 (46)
Janssen1992 (1999Ja15)	0,1895 (13)

The weighted mean of the above values is 0,1904 with a reduced -χ² of 1,14 ; the internal uncertainty is 0,0012; the external uncertainty 0,0013. This value is in good agreement with the theoretical adopted α_T (0,1918(19)).

The transition probability was deduced from the evaluated value (see below) of the emission intensity, using the adopted α_T.

Gamma Ray Emissions

- 159-keV gamma ray emission intensity is the weighted mean of :

83,65	0,50	(Chu – 1964Ch08)
83,48	0,38	(Hatch – 1966Ha03)
83,2	0,5	(Schötzig 1991 – [5])
83,9	0,6	(Coursey – 1992Co11)
83,81	0,32	(Janssen – 1992Ja15)
84,07	0,09	(Janssen – 1992Ja15)

The adopted value 83,99 is the weighted mean with an internal uncertainty of 0,08, and a reduced -χ² of 1,18.

[From the decay scheme and the α_T = 0,1918(19), the expected value is 83,90(14).]

- From α_T = 1099(33) and the decay scheme, the 88-keV gamma ray emission intensity is 0,0909(27). This value agrees with I_γ(88) = 0,0927(34), deduced from the ratio I_γ(159)/I_γ(88) = 906(33) measured by Raman (1972Ra07), using I_γ(159) = 83,99(8).

- The 247-keV gamma ray emission intensity of 0,000344(34) has been deduced from the ratio $I_{\gamma}(247)/I_{\gamma}(159) = 4,1(4) \cdot 10^{-6}$ measured by Raman (1973Ra32).

Conversion electrons

The conversion electron emission intensities have been calculated using conversion coefficients and gamma-ray emission intensities.

Atomic Data

The ω_K value is from Bambynek (1984) [6].

The ω_L value is from Schönfeld (1996Sc06).

The X-ray and Auger electron emission intensities have been calculated by using the program EMISSION (version 3.01) [4]

References

- [1] F. Lagoutine, N. Coursol, J. Legrand.
Table de Radionucléides, CEA/DIMRI, F-91191 Gif-sur-Yvette cedex, ISBN 2 7272 0078 1
- [2] V. M. Gorozhankin, N. Coursol, E. A. Yakushev. ICC99v3a: A computer program for interpolating internal conversion coefficients from Hager and Seltzer, Rosel et al., and from Band et al. (1999), BNM - CEA/LNHB
- [3] V. M. Gorozhankin, N. Coursol, E. A. Yakushev.
Appl. Rad. Isot. **56**, 189 (2002)
- [4] E. Schönfeld, H. Janssen. The program EMISSION Computer Code, PTB
Appl. Radiat. Isot. **52**, 595 (2000).
- [5] U. Schötzgig, H. Schrader, K. Debertin. Proc. Int. Conf. Nuclear Data Sci. Techn., Jülich, 13-17 May (1991)
- [6] W. Bambynek. X-84 Proc. X-Ray and Inner-Shell Processes in Atoms, Molecules and Solids, A. Meisel Ed., Leipzig Aug. 20-23 (1984)
- 1955Fa40 – L. W. Fagg, E. A. Woliki, R. O. Bondelid, K. L. Dunning, S. Snyder.
Phys. Rev. **100**, 1299 (1955).
- 1955Go21 – N. Goldberg, S. Frankel.
Phys. Rev. **100**, 1350 (1955).
- 1964Al28 – D. G. Alkhozov, V. D. Vasilev, G. M. Gusinskii, I. K. Lemberg, V. A. Nabichvrishvili.
Bull. Acad. Sci. USSR, Phys. Ser. **28**, 1575 (1965).
- 1964Ch18 – Y. Y. Chu, M.L. Perlman.
Phys. Rev. **135**, B319 (1964).
- 1964Ch08 – Y. Y. Chu, O. C. Kistner, A. C. Li, S. Monaro, M. L. Perlman.
Phys. Rev. **133**, B1361 (1964).
- 1966Gu02 – S. L. Gupta, M. M. Bajaj, N. K. Saha.
Nucl. Phys. **80**, 471 (1966).
- 1966Ha03 – E. N. Hatch, G. W. Eakins, G. C. Nelson, R. E. McAdams.
Proc. Intern. Conf. Internal Conversion Process, Nashville, Tenn. (1965), J.H. Hamilton, Ed., Academic Press, Inc., New York, p. 183 (1966).
- 1968Ka20 - R. A. Kalinauskas, K. V. Makaryunas, E. K. Makaryunene, R. I. Davidonis.
Bull. Acad. Sci. USSR, Phys. Ser. **32**, 187 (1969).
- 1969To02 – S. Tornkvist, S. Strom, L. Hasselgren.
Nucl. Phys. **A130**, 604 (1969).
- 1970EmZY – J. F. Emery, S. A. Reynolds, E. I. Wyatt.
ORNL-4466, p. 75 (1970).

- 1972Ra07 – S. Raman
Nucl. Instrum. Methods **103** (1972) 407
- 1973Ra32 – S. Raman, R. L. Auble, W. T. Milner.
Phys. Lett. **47B**, 19 (1973).
- 1974RaZO – S. Raman, R. L. Auble, W. T. Milner, T. A. Walkiewickz, R. Gunnink, B. Martin.
Report ORNL-4937, p. 144 (1974).
- 1977Kr13 – K. S. Krane.
At. Data Nucl. Data Tables **19**, 363 (1977).
- 1978Ro22 – F. Rosel, H. M. Fries, K. Alder, H. C. Pauli.
At. Data Nucl. Data Tables **21**, 92 (1978).
- 1992Ja15 – H. Janssen, E. Schönfeld, R. Klein.
Appl. Radiat. Isot. **43**, 1309 (1992).
- 1992Co11 – B. M. Coursey, D. B. Golas, D. H. Gray, D. D. Hoppes, F. J. Schima.
Nucl. Instrum. Methods Phys. Res. **A312**, 121 (1992).
- 1996Sc06 – E. Schönfeld, H. Janssen.
Nucl. Instrum. Methods Phys. Res. **A369**, 527 (1996).
- 2002Ba85 – I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr., P. O. Tikkanen, S. Raman.
At. Data Nucl. Data Tables **81**, 1 (2002).

¹²³I – Comments on evaluation of decay data by V. Chisté and M.M. Bé

1) Decay Scheme

There are 2 excited levels at 247 keV and 532 keV in ¹²³Te that have not been reported here. The 247 keV isomer ($T_{1/2} = 119,7$ d) is not populated in the electron capture decay of ¹²³I, and the expected electron capture population to the level 532 keV, if any, is very small.

2) Nuclear Data

The Q value is from Audi and Wapstra (1995Au04)

Level energies, spin and parities are from S. Ohya and T. Tamura (1993Oh07).

For level E= 687 keV, there are two possible spin values : 3/2+ and 5/2+. The 5/2+ value was suggested by Schoeters (1979Sc23) not after a measurement but by considering a proposal from Walters (1976Wa13). On the other hand, the 3/2+ value was measured by Sergolle ($\gamma\gamma$ coincidence (1969Se09) and Coulomb excitation (1970Se03)), Lien ((d,p) reaction (1975Li22)) and Andreev (Coulomb excitation (1975An16)). Then, the adopted value is 3/2+.

The half-life value, calculated by the Lweight program (version 3), is the weighted mean of :

$T_{1/2}$

Reference	Value (h)	Comments
Anderson (1964An03)	13,30 (5)	
Hupf (1968Hu01)	13,02 (4)	
Jonsson (1968Jo02)	13,4 (5)	
Karim (1973Ka01)	13,50 (11)	
Lagoutine (1982La13)	13,21 (2)	
Hoppes (1982Ho26)	13,219 (7)	Superseded 1992Un03
Unterweger (1992Un03)	13,2235 (19)	
Silva (2003Si04)	13,2228 (29)	
Schrader (2003Sc49)	13,232 (6)	

The original uncertainty given by Hupf (1968Hu01) (= 0,02) seems under estimated and has been multiplied by 2 by the evaluator. The uncertainty adopted by Lagoutine (1982La13) is the sum of the statistical uncertainty assessed at 3σ and the systematic uncertainty at 1σ ; consequently, the standard deviation cannot be obtained dividing the original uncertainty by 3 and we adopted the value 0,02. With this set of data, the reduced χ^2 is 4,7. The largest contribution comes from the value of Unterweger (1992Un03), amounting to 62%. The program Lweight 3 increases the uncertainty for the 1992Un03 value from 0,0019 to 0,00242 in order to reduce its relative weight from 62% to 50%.

The adopted value is the weighted mean : 13,2234 h, with the external uncertainty of 0,0037 h.

2.1) Electron Capture Transitions

The partial sub-shell capture probabilities are calculated with the program EC -Capture for the Allowed and 1st Forbidden transitions.

The electron capture probabilities and the related uncertainties have been deduced from the imbalance on each level of the decay scheme, assuming no EC transition to the ground state and to the 599 keV level. If this transition exists its intensity is of the order of a few per thousands.

2.3) Gamma Transitions

For the 159, 280, 346, 440 and 624 keV gamma transitions, the adopted δ (mixing of different multipolarities) are from the Krane evaluation (1977Kr06) of experimental measurements in which angular distribution and correlation data have been analyzed. For other transitions, the values of δ are from S. Ohya and T. Tamura (1993Oh07).

The internal conversion coefficients are calculated by ICC Computer Code (program Icc99v3a – GETICC dialog). The adopted values are interpolated from Rösler tables.

For the 159 keV gamma transition, many values of δ^2 have been found in the literature, as shown in the following table:

Reference	Value of d^2	Value of α_T
Goldberg et al – Phys. Rev. 100(1955)1350	0,013(1)	1,919 10^{-1}
Fagg et al – Phys. Rev. 100(1955)1299	0,0034(20)	1,905 10^{-1}
Chu et al – Phys. Rev. 133(1964)B1361	0,0067(11)	1,909 10^{-1}
Gupta et al – Nucl. Phys. 80(1966)471	0,011(8)	1,916 10^{-1}
Alkhozov et al – Phys. Serv. 28(1964)1575	0,004(5)	1,906 10^{-1}
Törnkvist et al – Nucl. Phys. A130(1969)604	0,0119(9)	1,917 10^{-1}
Krane et al - Atomic Data and Nuclear Data Tables 19(1977)19	0,01232 (47) (adopted value)	1,918 10^{-1}

It can be noted that even with values of δ^2 quite different the resulting α_T values are close with differences smaller than 1%; thus the adopted uncertainty is 1%.

For the 440 keV gamma transition, the following values of δ^2 have been found in the literature:

Reference	Value of d^2	Value of α_T
Sergolle et al – Nucl. Phys. A139(1969)554	0,149	0,0129912
Sergolle et al – Nucl. Phys. A145(1970)351	0,16	0,0129803
Roney et al – Nucl. Phys. A236(1974)165	4,41	0,0120886
Schoeters et al – Nucl. Phys. A323(1979)1	10,11	0,0119637
Krane - et al - Atomic Data and Nuclear Data Tables 19(1977)19	4,41 (adopted value)	0,0120886

In his articles (1969 and 1970), Sergolle deduced two values of δ for the 440 keV transition from 2 values of δ^2 for the 159 keV transition. The one reported here ($\delta^2(440)=0,149$) was calculated with $\delta^2(159) = 0,0119$ (Törnkvist). Nevertheless, this value is not close to the adopted one.

The 1% mixture of the 505 transition is from Sergolle (1969).

For the other transitions, measurements aren't precise, and only ranges of values are given for δ^2 .

Uncertainties calculations:

* For the 257 and 330 keV transitions (E2 pure), the α_T , α_K and α_L uncertainties are taken to be 3% from the calculated values with ICC Computer Code (program Icc99v3a).

* For the other transitions, the uncertainties calculations were made as follow : α_T was calculated for a pure M1 (or M3) transition and for a pure E2 transition. The difference between these values, normalized by α_T , is the uncertainty (%) of α_T . The same method is used for α_K and α_L uncertainties.

3) Atomic Data

Atomic values (ω_K , ω_L and n_{KL}) are from Schönfeld (1996Sc33).

The X-ray and Auger electron emission probabilities are calculated from the data set values by using the program EMISSION.

4) Radiation emissions

4.2) Gamma ray emissions

Gamma ray emission energies are from S. Ohya and T. Tamura (1993Oh07) and W. B. Walters (1976Wa13).

The measured emission intensities are given in table 1, they are relative to a value of 100 for the 159 keV gamma ray. Energy values are in keV.

Remarks to table 1 :

The original uncertainties given by Jacquemin (1987Ja10) for the 440, 528 and 538 lines have been multiplied by 2 by the evaluator to take into account some important factors:

- 1) During the measurement, there was a contamination that was not taken into account (Te-123m) by the author ;
- 2) As the value given is an absolute value, the uncertainty on the relative intensity given in table 1, has been estimated using the normalization factor and its uncertainty taking from the reference quoted by Jacquemin.

Two sets of values (R. C. Ragaine (1968Ra11) and E. H. Spejewski(1970Sp03)) were omitted in several cases from the analysis due to discrepancy with the other data.

For the 528 keV gamma line, the value given by R. K. Gupta (1960Gu14) was also omitted because it did not agree with the other values.

The normalization factor to convert the relative emission intensities to absolute intensities is calculated with the formula:

$$\text{Normalization} = \frac{100}{(\sum(1 + a_T)P_{rel})}$$

where the sum is to be done over all the gamma transitions to the ground state.

From the calculated α_T and the evaluated relative emission intensities (Table 1), the deduced normalization factor is **83,25 (2I)**. The uncertainties were calculated through their propagation on the above formula.

Absolute emission intensities are given on the last line in table 1.

4.2) Conversion electrons

The conversion electron emission intensities were deduced from the ICC values and from the gamma -ray emission probabilities. To our knowledge, there are no measured values for the conversion electron emission intensities.

Additional Reference

F. Lagoutine, Table de Radionucléides, CEA-LMRI(1984)

References

- 1960Gu14 - R.K. Gupta, Nucl. Phys. 14(1960)606 [E_γ , P_γ , I_γ]
 1964Ch08 - Y.Y. Chu , O.C. Kistner, A.C. Li, S. Monaro, M.L. Perlman, Phys. Rev. B133(1964)1316 [δ]
 1964Ch09 - Y. Y. Chu, M. L. Perlman, Phys. Rev. B135(1964)319 [α]
 1964An03 - G. Anderson, G. Rudstam, G. Sorensen, Ark. Fys. 28(1964)3 [$T_{1/2}$]
 1965Ha05 - E.N. Hatch, G.W. Eakins, G.C. Nelson, R.E. Mc Adams, Proc. Inter. Conf.: Internal Conversion Process (1965) [α]
 1966Gu01 – R. K. Gupta et al, Nucl. Phys. 80(1966)471 [δ]
 1967Se05 - H. Sergole, G. Albouy, J. Bouloume, J. M. Lagrange, L. Marcus, M.P. Pautrat, Le journal de Physique 28(1967)383 [E_γ , P_γ , I_γ]
 1968Hu01 - H.B. Hupf, J.S. Eldridge, J.E. Breaver, Int. J. Appl. Radiat. Isotop.. 19(1968)345 [$T_{1/2}$]
 1968Jo02 – G.G. Jonsson, B. Forkman, Nucl. Phys. A107(1968)52 [$T_{1/2}$]
 1968Ra11 - R.C. Ragaini , W.B. Walters, G.E. Gordon, P.A. Baedeker, Nucl. Phys. A115(1968)611 [E_γ , P_γ , I_γ]
 1968Ra02 - B.V.N. Rao, Swani Jnanananda, Phys. Rev.165(1968)1296 [E2 transitions]
 1968Se06 - H. Sergole, G. Albouy, M. Jourdain, J.M. Lagrange, N. Poffe, M. Pautrat, Le journal de Physique C1(1968)187 [E_γ , P_γ , I_γ]
 1969Se09 - H. Sergole , J. Vanhorenbeeck, Nucl. Phys. A139(1969)554 [E_γ , P_γ , δ]
 1969To02 - S. Törnkvist, S. Strom, L. Hasselgren, Nucl. Phys. A130(1969)604 [δ]
 1970Sc21 - H. Schrader, R. Stippler, F. Munich, Nucl. Phys. A151(1970)331 [E_γ , P_γ , I_γ]
 1970Se03 - H. Sergolle et al, Nucl. Phys. A145(1969)351 [E_γ , P_γ]
 1970Sp03 - E. H. Spejewski, P.K. Hopke, F. W. Loeser Jr., Nucl. Phys. A146(1970)182 [E_γ , I_γ]
 1971Ho02 - A. Hogleend, S.G. Malmstog, F. Munich, H. Schrader, Nucl. Phys. A165(1971)513 [M1 and E2 Transitions]
 1971Ch43 - H.C. Cheung, S.K. Mark, Nucl. Phys. A176(1971)489 [$T_{1/2}$]
 1971St08 - R. Stippler, D. Code, H. Schrader, F. Munich, Z. Phys. 242(1971)121 [E_γ , P_γ , I_γ , α_K]
 1972Ra07 - S. Raman, Nucl. Instrum. Methods 103(1972)407 [α]
 1973Ka01 - H.M.A. Karim, Radiochim. Acta 19(1973)1 [$T_{1/2}$]
 1973So04 - V.J. Sodd, J.W. Blue, K.L. Scholz, M.C. Oselka, Int. J. Appl. Radiot. Isotop. 24(1973)171 [E_γ , I_γ]
 1974Ro40 - W.N. Roney, D.W. Gebbie, R.R. Borchers, Nucl. Phys. A236(1974)165 [δ]
 1975Li22 - Lien et al, Nucl. Phys. A253(1975)165 [E_γ , I_γ , Spin]
 1975An16 - Andreev et al, Bull. Acad. Sci. USSR, Phys. Serv. 39(1975)55, n° 8 [E_γ , I_γ , Spin]
 1976Wa13 - W.B. Walters, R. A. Meyer, Phys. Rev. C14(1976)1925 [E_γ , I_γ]
 1977Fo02 - D.B. Fossan et al., Phys. Rev. C15(1977)1732 [Gamma-ray transitions, Spin]
 1977Ha09 - U. Hagemann, H-J. Keller, H-F Brinckman, Nucl. Phys. A289(1977)292 [E_γ , I_γ]
 1977Kr06 – K. S. Krane, At. Data Nucl. Data Tables 19(1977)19 [δ]
 1979Sc23 - E. Schoeters, J. Geenen, C. Nuytten, L. Vanneste, Nucl. Phys. A323(1979)1 [Spin, δ]
 1982Ho26 – D. D. Hoppes, J. M. R. Hutchinson, F. J. Schima, M. P. Unterweger, NBS -SP-626(1982)85 [Half-life]
 1982La13 – F. Lagoutine, J. Legrand, Int. J. Appl. Radiot. Isotop. 33(1982)711 [$T_{1/2}$]
 1982Sh08 - R.E. Shroy et al., Phys. Rev. C26(1982)1089 [E_γ , I_γ]
 1986Ag01 – V.A. Ageev – Report INIS-SU 392(1986)1 [E_γ , P_γ , I_γ]

Comments on evaluation

- 1987Ja10 - R. Jacquemin, Appl. Rad. Isotopes 38(1987)1087 [$T_{1/2}$, I_{γ}]
1992Un03 - M.P. Unterweger, D.D. Hoppes, F.J. Schima, Nucl. Instrum. Meth. A312(1992)349 [$T_{1/2}$]
1993Go01 - R. Goswani, B. Sethi, P. Barnerjee, P. K. Chattopadhyay, Phys. Rev. C47(1993)1013 [E_{γ}]
1993Oh07 - S. Ohya and T. Tamura, Nucl. Data Sheets 70(1993)531 [E_{γ} , I_{γ} , Spin]
1995Au04 - G. Audi, A.H. Wapstra, Nucl. Phys. A595(1995)409 [Q]
1996Sc33 - E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. A369(1996)527 [Atomic data]
2004Si04 - M. A. L. da Silva, M. C. M. de Almeida, C. J. da Silva, J. U. Delgado, Appl. Rad. Isotopes 60(2004)301 [Half-life].
2004Sc49 - H. Schrader, Appl. Rad. Isotopes 60(2004)317 [Half-life].

Table 1.I-123, gamma emission intensities, relative values to the 158 keV and, absolute values

04/12/01

Ref	174,2	182,61	192,17	197,26	198,25	206,82	207,82	242,32	247,96	257,51	278,36
60Gu14											
68Ra11		0,03(2)	0,03(2)						0,08(1)		
70Sp03		0,03(1)	0,03(2)						0,07(2)		
73So04		0,028(4)	0,025(4)		0,005(2)		0,0022(16)		0,068(6)		
76Wa13	0,0010(3)	0,0155(5)⌘	0,0238(8)	0,0004(2)	0,004(1)	0,004(1)	0,0013(4)	0,0004	0,0854(15)**	0,0018(5)	0,0027(5)
86Ag01									0,0864(31)	0,0026(12)	
87Ja10											
Adopted	0,0010(3)	0,022(6)	0,0239(8)	0,0004(2)	0,0042(9)	0,004(1)	0,00135(4)	0,0004	0,0838(27)	0,0019(5)	0,0027(5)
N	1	4	4	1	2	1	2	1	5	2	1
chi**2/N-1	0	2,07	0,09	0	0,2	0	0,3	0	2,16	0,38	0
Method		LWM, exp.unc	LWM, int. unc.		LWM, int. unc.		LWM, int. unc.		LWM, ext. unc.	LWM, int. unc.	
Abs. Value	0,00083(25)	0,0183(50)	0,0199(7)	0,00033(17)	0,0035(7)	0,0033(8)	0,00112(32)	0,0003330(8)	0,0698(23)	0,00160(22)	0,00225(42)

** = Input uncertainty multiplied by 1,75 in the program LWEIGHT

⌘ = Input uncertainty multiplied by 7,30 in the program LWEIGHT

exp.unc. = LWM expanded the uncertainty so range includes the most precise value.

int.unc. = internal uncertainty

ext.unc. = external uncertainty

Normalization factor = 83,25 (21)

Table 1.I-123, gamma emission intensities, relative values to the 158 keV and, absolute values

04/12/01

Ref	281,03	295,09	329,38	330,7	343,73	346,35	405,02	437,5	440,02	454,76	505,33
60Gu14	0,14(3) £			0,012(3)		0,16(3)			0,44(9)		0,280(6)
68Ra11	0,08(1)					0,12(2) (O)			0,42(2) (O)		0,31(5)
70Sp03	0,08(3)					0,11(3) (O)			0,42(8) (O)		0,32(8)
73So04	0,09(1)			0,017(6)		0,12(1)			0,46(2)	0,004(1)	0,27(3)
76Wa13	0,095(1)	0,0019	0,0031(7)	0,0139(5)	0,0051(5)	0,151(1)	0,0035(7)	0,0009(9)	0,514(6)	0,0047(6)	0,379(3)
86Ag01	0,095(44)			0,0142(7)	0,0055(5)	0,152(6)	0,0036(3)		0,524(21)	0,0051(3)	0,376(2)
87Ja10									0,450(29) ®		
Adopted	0,0948(1)	0,0019	0,0031(7)	0,01398(40)	0,00530(35)	0,151(1)	0,00358(28)	0,0009(9)	0,508(5)	0,00495(26)	0,32(5)
N	5	1	1	4	2	4	2	1	5	3	6
chi**2/N-1	0,68	0	0	0,27	0,32	3,22	0,02	0	2,98	0,66	3,8
Method	LWM, int. unc.			LWM, int. unc.	LWM, int. unc.	LWM, int. unc.	LWM, int. unc.		LWM, int. unc.	LWM, int. unc.	LWM, int. unc.
Abs. Value	0,0789(9)	0,0015818(40)	0,0026(6)	0,01164(33)	0,00441(29)	0,1257(9)	0,00298(23)	0,0007(7)	0,4229(43)	0,00412(22)	0,266(42)

® = Initial uncertainty multiplied by 2 by the evaluator

int.unc. = internal uncertainty

£ = Data rejection parameters for deviation from weighted average
(Chauvenet's criteria)

(O) = omitted value

Normalization factor = 83,25 (21)

Table 1.I-123, gamma emission intensities, relative values to the 158 keV and, absolute values

04/12/01

Ref	528,96	538,54	556,05	562,79	578,26	599,69	610,05	624,57	628,26	687,95	735,78
60Gu14	2,0(3) (O)										
68Ra11	1,27(11) (O)	0,32(2) (O)						0,08(1)		0,03(1)	0,04(1) (O)
70Sp03	1,26(24) (O)	0,31(6) (O)						0,07(2) (O)		0,04(2) £	0,05(2) (O)
73So04	1,40(5)	0,38(4)	0,0033(4)	0,0012(3)				0,085(5)		0,030(2)	0,06(3)
76Wa13	1,670(5)	0,458(5)	0,0037(5)	0,0013(5)	0,0018(5)	0,0031(11)	0,0013(4)	0,100(1)*	0,0019(3)	0,0321(15)	0,0739(14)
86Ag01	1,66(5)	0,460(21)		0,0014(1)	0,0015(1)	0,0032(2)		0,101(5)	0,0020(2)	0,0329(9)	0,0742(35)
87Ja10	1,41(6)®	0,379(31)®									
Adopted	1,58(10)	0,455(5)	0,00346(31)	0,00138(9)	0,00151(1)	0,0032(2)	0,0013(4)	0,0958(24)	0,00197(17)	0,0323(7)	0,074(1)
N	4	4	2	3	2	2	1	4	2	4	3
chi**2/N-1	8,34	3,3	0,39	0,21	0,35	0,01	0	3,28	0,08	0,5	0,11
Method	LWM, exp.unc.	LWM, int. unc.	LWM, int. unc.	LWM, int.unc.	LWM, int.unc.	LWM, int. unc.		LWM, ext. unc.	LWM, int. unc.	LWM, int. unc.	LWM, int. unc.
Abs. Value	1,32(8)	0,3788(43)	0,00288(26)	0,00115(7)	0,00126(8)	0,00266(17)	0,00108(33)	0,0798(20)	0,00164(14)	0,0269(6)	0,0616(8)

* = Input uncertainty multiplied by 3,33 in the program LWEIGHT

exp.unc. = LWM expanded the uncertainty so range includes the most precise value

® = Initial uncertainty multiplied by 2 by the evaluator

int.unc. = internal uncertainty

£ = Data rejection parameters for deviation from weighted average (Chauvenet's criteria)

ext.unc. = external uncertainty

(O) = omitted value

Normalization factor = 83,25 (21)

Table 1.I-123, gamma emission intensities, relative values to the 158 keV and, absolute values

04/12/01

Ref	783,59	837,1	877,52	894,8	909,12	1036,63	1068,12
60Gu14							
68Ra11	0,05(1) (O)						
70Sp03	0,05(2) (O)						
73So04	0,068(5)	0,0008(2)	0,0010(2)	0,0017(5)	0,0017(4)	0,0010(2)	0,0014(2)
76Wa13	0,0713(14)	0,0006(1)	0,0013(8)	0,0011(3)	0,0016(3)	0,0012(3)	0,0017(1)
86Ag01	0,0718(35)	0,00070(1)	0,0010(1)	0,0012(1)	0,0017(1)	0,0012(1)	0,0018(1)
87Ja10							
Adopted	0,0712(13)	0,000699(10)	0,00100(9)	0,00121(9)	0,00169(9)	0,00116(9)	0,00171(8)
N	3	3	3	3	3	3	3
chi**2/N-1	0,22	0,62	0,07	0,55	0,05	0,41	1,61
Method	LWM, int. unc.	LWM, int. unc.	LWM, int. unc.	LWM, int. unc.	LWM, int.unc.	LWM, int. unc.	LWM, ext.unc.
Abs. Value	0,0591(11)	0,000582(8)	0,00083(7)	0,00101(7)	0,00141(8)	0,00097(7)	0,00142(7)

(O) = omitted value

int.unc. = internal uncertainty

ext.unc. = external uncertainty

Normalization factor = 83,25 (21)

¹²⁴Sb - Comments on evaluation of decay data
by M.M. Bé and V. Chisté

This evaluation was completed in December 2008. The literature available by this date was included as well as the results obtained as a part of a specific exercise dedicated to the ¹²⁴Sb activity and γ -ray emission intensity measurements organized by the Euramet organisation (Project 907, full report to be published). In the following, the participants in the Euramet 907 project will be referred as E907- *n*, where *n* is a serial number.

1. Decay Scheme

This decay scheme is complete and is based on those proposed by Goswamy (1993Go10), Patil (2006Pa16) and the results obtained in the Euramet-907 project.

A good agreement was found between the effective Q value of 2906 (8) keV computed from the decay scheme data and the adopted Q value of 2904,3 (15) keV from the mass adjustment of Audi *et al.*

2. Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2003Au03).

Experimental half-life values (in days) are listed below:

Reference	T _{1/2}	Uc	Comments
Macklin (1957Ma50)	60,4	0,2	
C.H.Johnson (1958Jo01)	59,9	0,5	
J.P.Cali (1959Ca12)	60,1	0,3	
S.A.Reynolds (1968Re04)	60,3	0,2	
D.M.Fleming (1966Fl01)	60,20	0,03	calorimetry
I.A.Kharitonov (2000Kh04)	60,11	0,07	4 $\pi\beta$ - γ coincidence method
* E907- 8	60,212	0,011	Ionization chamber
Adopted	60,208	0,011	Reduced $\chi^2 = 1$; critical $\chi^2 = 4,6$

*Euramet 907 participant number 8

The adopted value is the weighted mean of the three most precise values with the external uncertainty.

2.1 Beta transitions

β^- transition energies have been energies are calculated from the Q value and the level energies.

The β^- transition probabilities were deduced from the γ transition probability balance at each level of the decay scheme. The adopted values are compared with the measured values in the following table:

	(0, 1) 2301 keV %	(0, 3) 1579 keV %	(0, 5) 946 keV %	(0, 10) 610 keV %	(0, 20) 210 keV %
Langer (1953La35)	21	7	9	49	14
Moreau (1954Mo83)	22	7	9	53	9
Azuma (1955Az29)	22	6	4	56	12
Hsue (1965Hs02)	23	5			
Zolotavin (1956Zo06)	28	10	4	49	9
Adopted	23,44 (28)	4,815 (29)	2,295 (7)	51,21 (19)	8,663 (27)
Nature	1 st S=q2+(1p2+16(2) (Hsue) S=k(1-0,25W- 0,06/W+0,041W2)(Hsue) S=q2+(1p2+7(2) (Canty) S=0,9q2+p2 (Johnson)				

The weak beta transition probabilities are based on the γ transition probability balance at each level of the decay scheme, especially in the upper part of the decay scheme (from level 2886-keV to level 2483-keV) where there are only gamma transitions depopulating these levels. In this evaluation, only the gamma rays observed in several independent experiments have been retained (see § 4.2) so the corresponding levels can be considered definitely established.

2.2 Gamma transitions and internal conversion coefficients

γ -ray measurements carried out by Doll *et al.* (2000Do11) confirmed the doublet structure of the 2039 level ; one with J^π assignment 2^+ and the second with 3^+ ; with a spacing of 129 eV.

The γ transitions with energy : 2039,4- ; 790,8- ; 1436,7- ; 713,9-keV start from level with $J^\pi = 2^+$ and, those with energy : 790,7- ; 1436,6- ; 713,8-keV from level with $J^\pi = 3^+$. They are shown as doublets in the following table.

Internal conversion coefficients

Multipolarity and multipole mixing ratio (δ) for some transitions were determined using the techniques of directional correlation and nuclear orientation measurements, these are summarized in Table 1 :

Table 1 :

Transition energy (keV)	multipole mixing ratio (d)	Multipolarity	Reference
444	0,57 (17) or 0,06 (8)		Robinson <i>et al.</i> 1983
646	0,013 (9) 0,000 (1)	E2, M3	Goswamy <i>et al.</i> 1993 Baker <i>et al.</i> 1972
709	- 0,8 (+3, -4) - 1 (+6, -8) - 0,18 (5) 0,04 (3, -5)	M1, E2	Goswamy <i>et al.</i> 1993 Goswamy <i>et al.</i> 1993 Robinson <i>et al.</i> 1983 Grabowski <i>et al.</i> 1971
714	- 0,65 (+38, -0,54) 1,15 (16, - 25) 1,5 (7)	M1, E2	Goswamy <i>et al.</i> 1993 Subrahmanyeswara <i>et al.</i> 1990 Robinson <i>et al.</i> 1983

Transition energy (keV)	multipole mixing ratio (d)	Multipolarity	Reference
	1,5 (6) 0,98 (19)		Baker <i>et al.</i> 1972 Grabowski <i>et al.</i> 1971
723	3,74 (12) - 3,8 (2) - 3,4 (3) - 3,3 (2) - 3,4 (1) - 7,5 (20) - 3,4 (6)	M1, E2	Goswamy <i>et al.</i> 1993 Subrahmanyeswara <i>et al.</i> 1990 Robinson <i>et al.</i> 1983 Baker <i>et al.</i> 1972 Grabowski <i>et al.</i> 1971 Sites <i>et al.</i> 1970 Stelson, 1967
791	- 0,15 (+5, -2) - 0,3 (+52, -14)	E2, M3	Goswamy <i>et al.</i> 1993 Goswamy <i>et al.</i> 1993
968	0,038 (3) - 0,35 (8) - 0,02 (2) - 0,03 (6, -5) - 0,02 (8)	E1, M2	Goswamy <i>et al.</i> 1993 Subrahmanyeswara <i>et al.</i> 1990 Robinson <i>et al.</i> 1983 Baker <i>et al.</i> 1972 Sites <i>et al.</i> 1970
1045	- 0,14 (+3, -4) - 0,03 (2) 0,041 (47, -41) - 0,1 (1)	E1, M2	Goswamy <i>et al.</i> 1993 Robinson <i>et al.</i> 1983 Baker <i>et al.</i> 1972 Sites <i>et al.</i> 1970
1356	- 0,32 (+25, -18)	E2, M1	Goswamy <i>et al.</i> 1993
1368	- 0,28 (6) - 0,02 (1) - 0,045 (90) - 0,01 (8)		Subrahmanyeswara <i>et al.</i> 1990 Robinson <i>et al.</i> 1983 Baker <i>et al.</i> 1972 Sites <i>et al.</i> 1970
1376	0,26 (11) < 0,29 - 0,01 (3)	E1, M2	Goswamy <i>et al.</i> 1993 Goswamy <i>et al.</i> 1993 Robinson <i>et al.</i> 1983
1437	0,51 (+13, -11) 1,5 (8) 3,7 (27, -20)	M1, E2	Goswamy <i>et al.</i> 1993 Robinson <i>et al.</i> 1983 Baker <i>et al.</i> 1972
1445	0,015 (80) 0,10 (9)	E1, M2	Goswamy <i>et al.</i> 1993 Robinson <i>et al.</i> 1983
1489	0,10 (23) - 3,4 (9, -15)		Robinson <i>et al.</i> 1983 Baker <i>et al.</i> 1972
1691	- 0,009 (22) - 0,06 (3) - 0,02 (1) 0,00 (3)	E1, M2	Goswamy <i>et al.</i> 1993 Subrahmanyeswara <i>et al.</i> 1990 Baker <i>et al.</i> 1972 Sites <i>et al.</i> 1970
2091	0,031 (6) 0,032 (32) 0,00 (2, -3) 0,07 (3)	E1, M2	Goswamy <i>et al.</i> 1993 Subrahmanyeswara <i>et al.</i> 1990 Baker <i>et al.</i> 1972 Sites <i>et al.</i> 1970

Moreover, two sets of measured values of the conversion electron intensities (I_{ce_i}) are also available: by Grigor'eev *et al.* (1968), and by Johnson (1974)Jo03). These values as well as their weighted means are summarized in where α_{K602} is the theoretical K conversion coefficient interpolated from Band's tables using the program BrIcc with the "frozen orbital approximation" (Kibédi *et al.* 2008Ki07) for an E2 transition ; I_{ce_i} are the conversion electron intensities, and I_{γ} , the relative gamma-ray emission probabilities as summarized in Table 3.

The experimental α_{K_i} conversion coefficients have been compared with the theoretical ICC, the deduced mixing ratios δ are in good agreement with those determined by directional correlation and nuclear orientation measurements summarized in Table 1.

Table 2. Then, the experimental K conversion coefficients α_{K_i} were deduced from the relation:

$$\alpha_{K_i} = \alpha_{K602} \times I_{ce_i} / I\gamma_i$$

where α_{K602} is the theoretical K conversion coefficient interpolated from Band's tables using the program BrIcc with the "frozen orbital approximation" (Kibédi *et al.* 2008Ki07) for an E2 transition ; I_{ce_i} are the conversion electron intensities, and $I\gamma_i$, the relative gamma-ray emission probabilities as summarized in Table 3.

The experimental α_{K_i} conversion coefficients have been compared with the theoretical ICC, the deduced mixing ratios δ are in good agreement with those determined by directional correlation and nuclear orientation measurements summarized in Table 1.

Table 2 :

Energy	Johnson		Grigor'eev		$\alpha_k (602)=$		0,00420 0,00006		$\alpha_k =$		Multipolarity	delta	%	α_k theo	α_T theo.
	Iec	Uc	Ice	Uc	Ice WM	Uc dopt.	Ig rel.	Uc Ig	Ice/Ig * ak602	uc α_k					
159	2,3	0,2			2,3	0,2	0,0050	0,0006	1,93	0,29					
254	0,10	0,08			0,1	0,08	0,0145	0,0009	0,0290	0,0232	E1 ?			0,01269 (18)	0,01465 (21)
336	0,12	0,08			0,12	0,08	0,0741	0,0009	0,0068	0,0045	E1			0,00611 (9)	0,00704 (10)
371	0,1	0,08			0,1	0,08	0,0292	0,0011	0,0144	0,0115					
400	0,45	0,08			0,45	0,08	0,128	0,0027	0,0148	0,0027	E2			0,01323 (2)	0,01566 (2)
444	0,35	0,15			0,35	0,15	0,192	0,009	0,0077	0,0033	M1+E2	0,06	26,5	0,01092 (16)	0,01261 (18)
469	< 0,14				< 0,14		0,0469	0,0027			E1			0,00268 (4)	0,00309 (5)
481	< 0,07				< 0,07		0,0237	0,0032							
525	0,14	0,08			0,14	0,08	0,1484	0,0036	0,0040	0,0023	M1+E2	1	50	0,0066 (3)	0,0077 (3)
602	100		100		100		100		0,00420	0,00006	E2			0,00420 (6)	0,00490 (7)
646	5,4	0,5	6,6	0,3	6,28	0,53	7,591	0,015	0,0035	0,0003	E2+M3	0,006	0,0036	0,00351 (5)	0,00409 (6)
709	1,4	0,5	1,2	0,1	1,21	0,10	1,3941	0,0046	0,0036	0,0003	M1+E2	-0,18	3,1	0,00349 (5)	0,00402 (6)
713	1,6	0,5	1,6	0,2	1,60	0,19	2,325	0,007	0,0029	0,0003	M1+E2	1	50	0,0031 (4)	0,0036 (4)
722	5,7	0,5	7,5	0,3	7,02	0,79	10,952	0,022	0,0027	0,0003	M1+E2	-3,4	92	0,00271 (4)	0,00314 (5)
735	0,04	0,02			0,04	0,02	0,1342	0,0016	0,0013	0,0006					
766	0,035	0,02	0,06	0,02	0,048	0,014	0,0105	0,0009	0,0190	0,0059	E0, M1			0,019 (6)	0,021 (7)
790	0,44	0,08	0,44	0,03	0,440	0,028	0,7584	0,0025	0,0024	0,0002	E2			0,00214 (6)	0,00248 (8)
968	0,24	0,08	0,33	0,03	0,319	0,030	1,93	0,01	0,0007	0,0001	E1(+M2)	-0,2	3,8	0,000569 (9)	0,000653 (11)
1045	0,18	0,08	0,25	0,03	0,241	0,028	1,894	0,014	0,0005	0,0001	E1(+M2)	-0,03	0,09	0,000494 (9)	0,000567 (10)
1325	0,35	0,1	0,30	0,03	0,304	0,029	1,623	0,007	0,0008	0,0001	E2			0,000693 (10)	0,000827 (12)
1355	0,17	0,1	0,20	0,02	0,199	0,020	1,0649	0,0039	0,0008	0,0001	E2(+M3)	-0,32	9,3	0,0009 (5)	0,0011 (5)
1368	0,14	0,05	0,22	0,03	0,199	0,035	2,680	0,008	0,0003	0,0001	E1(+M2)	-0,02	0,04	0,000303 (5)	0,000478 (7)
1376	0,035	0,03			0,035	0,03	0,5113	0,0044	0,0003	0,0002	E1(+M2)	-0,01	0,01	0,000300 (5)	0,000479 (7)
1418	0,25	0,1			0,25	0,1	0	0							
1436	0,28	0,1	0,17	0,03	0,18	0,03	1,262	0,008	0,0006	0,0001	M1+E2	1,5	69,23	0,00063 (5)	0,00078 (5)
1489	0,14	0,1	0,13	0,02	0,13	0,02	0,6924	0,0038	0,0008	0,0001	M1+E2	0,1	0,9901	0,000659 (14)	0,000829 (16)
1526	0,035	0,03	< 0,04		0,035	0,03	0,4232	0,0048	0,0003	0,0003	E1			0,000252 (6)	0,000535 (8)
1657	0,2	0,1			0,2	0,1	0,00	0,00							
1691	2,7	0,4	2,5	0,2	2,54	0,18	48,54	0,19	0,00022	0,00002	E1+M2	0,01	0,01	0,000213 (4)	0,000615 (9)
2090,9	0,24	0,06	0,20	0,04	0,212	0,033	5,618	0,025	0,00016	0,00002	E1(+M2)	0,03	0,1	0,0001522 (23)	0,000838 (12)

3. Atomic Data

The fluorescence yield data are from 1996Sc06 (Schönfeld and Janssen).

3.1 X Radiations

The relative K x-ray emission probabilities are from 1996Sc06.

3.2 Auger Electrons

The ratios $P(KLX)/P(KLL)$ and $P(KXY)/P(KLL)$ are from 1996Sc06.

4. Radiation Emissions

4.1 Electron Emissions

The β - emission energies and intensities were deduced from γ transition probabilities (§ 2.1).

The conversion electron emission intensities have been calculated from the γ -ray emission intensities in sect. 4.2, and the internal-conversion coefficients in sect. 2.2.

The Auger electron emission intensities were calculated by the EMISSION program from PTB using the γ -ray emission probabilities, the atomic data of sect. 3, and the internal-conversion coefficients of sect. 2.2.

4.2 Photon Emissions

The X-ray absolute emission intensities were calculated using the EMISSION program and the γ -ray emission intensities, the atomic data given in sect. 3, and the internal-conversion coefficients in sect. 2.2. They are compared with the three sets of absolute values measured by participants in the Euramet exercise. They are, in general, in good agreement.

Energy (keV)	E907- 2		E907- 3		E907- 8		Calculated	
	I %	Uc	I %	Uc	I %	Uc	I %	Uc
27,2 (K α 2)			0,128	0,002	0,130	0,003	0,1252	0,0018
27,5 (K α 1)			0,264	0,004	0,230	0,006	0,233	0,003
30,9 (K β '2)			0,068	0,001	0,063	0,002	0,0667	0,0012
31,7 (K β '1)			0,0170	0,0005	0,0136	0,0006	0,0145	0,0005
K α	0,35	0,07	0,392	0,0045	0,359	0,007	0,358	0,0035
K β	0,087	0,018	0,085	0,0011	0,076	0,0018	0,081	0,0013
K X Total	0,437	0,072	0,476	0,005	0,436	0,007	0,439	0,004

The X-ray relative emission intensities given by Euramet participants 2 and 3 are compared, in the following table, with the published values of Patil (2006) and Goswamy (1993).

Energy (keV)	E907- 2		E907- 3		Patil (2006)		Goswamy (1993)	
	Rel. Int.	Uc	Rel. Int.	Uc	Rel. Int.	Uc	Rel. Int.	Uc
K α : 27,3	0,361	0,076	0,4000	0,0046	0,3681	0,0066	0,366	0,017
K β : 30,9 – 31,8	0,089	0,018	0,0864	0,0014	0,0852	0,0017	0,084	0,050

g-ray energies

The γ -ray energies in the following table are from Helmer (2000He14). The other energies were deduced from the level energy differences.

E (keV)	Uc (keV)	E (keV)	Uc (keV)
602,7260	0,0023	1045,125	0,004
645,8520	0,0019	1325,504	0,004
713,776	0,004	1368,157	0,005
722,782	0,003	1436,554	0,007
790,706	0,007	1690,971	0,004
968,195	0,004	2090,930	0,007

g-ray emission intensities

The 6 participants in the Euramet project sent their γ -ray emission intensities in both relative and absolute scales, since they also carried out activity measurements of the solution.

Moreover, eight sets of measured values published in the literature are available. All of them are relative to the most intense 602-keV γ -ray line (Table 3).

Among the 111 γ rays mentioned before or in this exercise, some weak lines were observed once and not confirmed by other measurements, these are summarized below:

- Weak gamma rays of weak intensities observed by one Euramet participant often described being “barely visible” and then not adopted in the decay scheme:

2871-keV ; 2274-keV ; 2253-keV ; 2151-keV ; 1970-keV (just detection limits) ; 1950-keV ; 1657-keV ; 1557-keV ; 1428-keV ; 1269-keV ; 1202-keV ; 1198-keV ; 1180-keV ; 1163-keV ; 669-keV ;

- Weak gamma rays of weak intensities observed by Patil but by none of the Euramet participant and not adopted in the decay scheme:

2814-keV ; 2746-keV ; 2515-keV ; 2490-keV ; 2386-keV (just detection limits) ; 2373-keV ; 2256-keV ; 2232-keV ; 2145-keV ; 1418-keV ; 795-keV ; 743-keV ; 592-keV ; 186-keV.

A number of weak gamma rays were observed by some Euramet participants or by others:

- 2224-keV, 2204-keV the reported intensities are quite discrepant so they were omitted;

- 1453-keV could be between levels 2701,6 and 1248,5-keV, but the reported intensities are quite discrepant so this γ -ray was omitted ;

- 476-keV could be between levels 2701,6 and 2224,8-keV, but the reported intensities are quite discrepant so this ray has not been retained ;

- 1757-keV ; 1509-keV ; 1253-keV ; 1097-keV ; 1014-keV ; 937-keV ; 553-keV ; 498,4-keV ; 385-keV ; 346-keV ; do not correspond to levels differences, they have not been retained.

- 1235-keV ; 997-keV ; 159,8-keV were accepted but not placed in the decay scheme.

602-keV absolute g-ray emission intensity

1) A first attempt was made to determine the 602-keV line absolute emission intensity using the results of the absolute measurements carried out in the framework of the Euramet project:

Participant	I _{g602} in %	Uc
E907- 2	97,5	0,7
E907- 3	97,8	0,9
E907- 5	97,6	0,7
E907- 6	91	1
E907- 7	97,84	0,34
E907- 8	98,1	1,5

Chi2	0,1		$\chi^2 / (n-1)$
Chi2 crit:	3,3		Unweighted mean
UWM:	97,787		Weighted mean
WM:	97,769		Internal uncertainty
Uc (int):	0,26		External uncertainty
Uc (ext) :	0,07		
LWM :	97,77	0,26	Limited WM

The value of participant 6 was found to be an outlier based on Chauvenet’s criterion. Value of participant 7 contributes to 58 % to the weighted mean (WM). The set of the five remaining values is consistent, then the evaluated value (LWM) is the weighted mean with the internal uncertainty.

All absolute γ -ray emission intensities measured by the Euramet participants are summarized in Table 4.

2) A second attempt using all the available measurements was done. Since the Euramet participating laboratories sent their results as relative values also, these six sets of results were used as well as the previous measurements published in the literature. So, 14 sets of data were included in the evaluation (Table 3).

In the Euramet project, the participants sent their results as values relative to the reference line $I_{\gamma 602} = 100$; with its uncertainty included in the uncertainties of the other γ -ray lines.

In the other publications, when an author gave an uncertainty on this $I_{\gamma 602}$ reference line, then this uncertainty was included into each individual value using the relation : $Uc = \text{sqrt} (Uc_{rel} * Uc_{rel} + Uc_{I_{\gamma 602}} * Uc_{I_{\gamma 602}})$. So, all gamma rays have been treated with emission intensities relative to $I_{\gamma 602} = 100$ (with no uncertainty).

Since no beta transition populating the ground state level in Tellurium 124 is expected, the sum of the gamma transition probabilities with energy 2807-, 2693-, 2681-, 2455-, 2323-, 2294-, 2182-, 2039-, 1657-, 1325-, 602-keV which populate the ground state must be equal to 100. That is:

$$\sum_i I_{g_i} [1 + \alpha_{T_i}] = \frac{100}{N}$$

Where: I_{γ_i} is the relative emission probability of the gamma-ray, α_{T_i} is its total conversion coefficient, and N is a normalisation factor between the relative and absolute scales.

N, the normalization factor, is then deduced from the measured relative I_{γ_i} values:

$$N = \frac{100}{\sum_i I_{g_i} [1 + \alpha_{T_i}]} \quad \text{and} \quad dN^2 = \sum_i \left(\frac{\partial N}{\partial I_{g_i}} dI_{g_i} \right)^2 + \sum_i \left(\frac{\partial N}{\partial \alpha_{T_i}} d\alpha_{T_i} \right)^2$$

The α_{T_i} coefficients are theoretical values interpolated from Band’s tables (2002Ba85) using the program BrIcc with the “frozen orbital approximation” (Kibédi *et al.* 2008Ki07). All transitions with a measured multipolarity are E2.

This leads to $N = 0,977\ 75$ (20).

The absolute emission intensity of the 602-keV g-ray is then deduced to be: 97,775 (20) %.

This value is in full agreement with the above value of 97,77 (26) %. However, because of the normalization procedure used, its uncertainty is ten times smaller.

Having in mind that the energies of the involved transitions are relatively high and their respective multipolarities are E2, the conversion coefficient values deduced from theoretical calculations can be considered very reliable. Hence, this second absolute intensity value and its associated uncertainty were adopted here.

All the measured relative gamma emission intensities are summarized in Table 3, with the unweighted mean for each set of values given, as well as the weighted mean, the reduced χ^2 and the internal and external uncertainties, the adopted relative emission intensity value and its uncertainty and the deduced and adopted absolute values.

All the absolute gamma-ray emission intensities measured by the participants in the Euramet 907 project are summarized in Table 4. The most intense lines are compared to those obtained from relative values and conversion coefficients (Table 3) in the following table. The agreement is very good.

g-ray energy keV	From absolute measurements (Table 4)	From relative measurements and ICC (Table 3)
602	97,77 (26)	97,775 (20)
645	7,414 (21)	7,422 (15)
709	1,3635 (43)	1,363 (5)
713	2,269 (11)	2,273 (7)
722	10,712 (31)	10,708 (22)
968	1,880 (6)	1,887 (10)
1045	1,835 (6)	1,852 (14)
1325	1,583 (6)	1,587 (7)
1368	2,615 (9)	2,620 (8)
1690	47,39 (22)	47,46 (19)
2090	5,491 (26)	5,493 (24)

7. References

- 1953La35 L.M. LANGER, N.H.Lazar, R.J.D.Moffat. Phys. Rev. 91 (1953) 338. Beta emission probabilities
1954Mo83 J.Moreau. Comp. Rend. Acad. Sci. (Paris) 239 (1954) 800. Beta emission probabilities
1955Az29 T.Azuma. J. Phys. Soc. Jpn 10 (1955) 167. Beta emission probabilities
1956Zo06 A.V.Zolotavin, E.P.Grigoriev, M.A.Abrovian. Izvest.Akad.Nauk SSSR, Ser.Fiz.20 (1956) 289. Columbia Tech.Transl. 20, 271 (1957)
1957Ma50 R.L.Macklin. Nucl. Instrum. Methods 1 (1957) 335. Half-life
1956Zo06 A.V.Zolotavin, E.P.Grigoriev, M.A.Abrovian. Izvest.Akad.Nauk SSSR, Ser.Fiz. 20, 289 (1956) ; Columbia Tech.Transl.20 (1957) 271. Beta emission probabilities
1958Jo01 C.H.Johnson, A.Galonsky, J.P.Ulrich. Phys. Rev.109 (1958) 1243. Half-life
1959Ca12 J.P.Cali, L.F.Lowe. Nucleonics 17, 10 (1959) 86. Half-life
1965Hs02 S.T.HSUE, L.M.Langer, S.M.Tang, D.A.Zollman. Nucl. Phys. 73 (1965) 379. Beta emission probabilities
1966Fl01 D.M.FLEMING, I.T. MYERS. Int. J. Appl. Radiat. Isotop. 17 (1966) 251. Half-life
1967ST05 P.H. Stelson. Phys. Rev. 157 (1967) 1098. ICC
1968Gr24 E.P.GRIGORIEV, A.V. ZOLOTAVIN, V.O. SERGEEV, M.I. SOVTSOV. Izv. Akad. Nauk SSSR. Ser. Fiz. 32 (1968) 733. K-Conv. Elec. emission probabilities
1968Re04 S.A.Reynolds, J.F.Emery, E.I.Wyatt. Nucl. Sci. Eng. 32 (1968) 46. Half-life
1968Gr24 E.P.GRIGORIEV, A.V. ZOLOTAVIN, V.O. SERGEEV, M.I. SOVTSOV. Bull. Ac. Sci. USSR. Phys. Ser. 32 (1968) 711. K-Conv. Elec. emission probabilities
1969Ra31 R.C.RAGAINI, W.B. WALTERS, R.A. MEYER. Phys. Rev. 187 (1969) 1721. K ICC, Mixing Ratio
1969Me04 R.A. MEYER, W.B. WALTERS, R.C. RAGAINI. Nucl. Phys. A127 (1969) 595. Gamma-ray emission probabilities, Spin and Parity, Gamma-ray energies
1970Si17 J.R.SITES, W.A. STEYERT. Nucl. Phys. A156 (1970) 19. Mixing Ratio
1971GR14 Z.W.Grabowski, K.S.Krane, R.M.Steffen. Phys. Rev. C3 (1971) 1649. Mixing Ratio
1972BA38 K.R.Baker, J.H.Hamilton, A.V.Ramayya, G.Highland. Nucl. Phys. A186 (1972) 493. Mixing Ratio
1974Jo03 J.R.JOHNSON, K.C. MANN. Can. J. Phys. 52 (1974) 406. Gamma-ray emission probabilities, K-Conv. Elec. emission probabilities
1979Sh08 A.K.SHARMA, R.KAUR, H.R. VERNA, K.K. SURJ, P.N. TREHAN. J. Phys. Soc. Jap. 46 (1979) 1057. Gamma-ray emission probabilities
1983RO13 S.J.Robinson, W.D.Hamilton, D.M.Snelling. J. Phys. (London) G9 (1983) 921. Mixing ratio
1984MA13 G.Mardirosian, N.M.Stewart. Z.Phys. A315, 213 (1984) Gamma-ray emission probabilities
1984Iw03 Y.IWATA, M.YASUHARA, K.MAEDA, Y.YOSHIZAWA. Nucl. Instrum. Methods 219 (1984) 123. Gamma-ray emission probabilities

- 1988YO05 You Jianming, Liu Yunzuo, Hu Dailing. *Z. Physik A331* (1988) 391. Gamma-ray emission probabilities
- 1990ME15 R.A. Meyer. *Fizika 22 (Zagreb)* (1990) 153. Gamma-ray emission probabilities
- 1990Su10 S.Subrahmanyaswara Rao, K.Bhaskara Rao, V.Seshagiri Rao, H.C.Padhi. *Nuovo Cim. 103A* (1990) 803. ICC
- 1993Go10 J.Goswamy, B.Chand, D.Mehta, N.Singh, P.N.Treha. *Appl. Rad. Isotopes 44* (1993) 541. Gamma-ray emission probabilities
- 1996Sc06 E. Schönfeld, H. Janssen. *Nucl. Instrum. Meth. Phys. Res. A369* (1996) 527. Atomic data
- 2000Kh04 I.A.Kharitonov, T.E.Sazonova, S.V.Sepman, T.I.Shilnikova, A.V.Zanevsky. *Appl. Rad. Isotopes 52* (2000) 415. Half-life
- 2000He14 R.G. Helmer, C. van der Leun. *Nucl. Instrum. Methods Phys. Res. A450* (2000) 35. Gamma energy
- 2000Do11 C.Doll, H.Lehmann, H.G.Borner, T.von Egidy. *Nucl. Phys. A672* (2000) 3. Nuclear structure
- 2002Ba85 I.M.Band, M.B.Trzhaskovskaya. *At. Data. Nucl. Data Tables 88,1* (2002). Theoretical ICC
- 2003AU03 G. Audi, A.H.Wapstra, C. Thibault. *Nucl. Phys. A729* (2003) 337-676. Q value
- 2006Pa16 A.Patil, D.Santhosh, K.V.Sai, M.Sainath, K.Venkataramaniah. *Appl. Rad. Isotopes 64* (2006) 693. Gamma-ray emission probabilities
- 2008Ki07 T.Kibédi, T.W.Burrows, M.B.Trzhaskovskaya, P.M.Davidson, C.W.Nestor, Jr. *Nucl. Instrum. Methods Phys. Res. A589* (2008) 202. ICC

Table 3 : Relative gamma ray intensities and absolute values calculated with ⁶⁰Ig602 = 97,775 (20) %.
 (i, j) refers to initial and final levels, (-1, n) transition not placed in the decay scheme. DL = Detection Limit

	(14, 12) 148keV	(-1, 1) 159keV	(16, 12) 186keV	(14, 10) 189 keV	(20, 14) 209keV	(10, 6) 254keV	(23, 14) 291keV
	Value Uc	Value Uc	Value Uc	Value Uc	Value Uc	Value Uc	Value Uc
E907- 2	0,012 0,005	0,005 0,001		0,002 0,001	^(o) 0,0088 0,0012	0,009 0,001	0,0046 0,0008
E907- 3	DL=0,0031	DL=0,0032	DL=0,0041	DL=0,0042	DL=0,0043	0,014 0,002	DL=0,0053
E907- 5							
E907- 6							
E907- 7	0,0053 0,0012	0,0070 0,0014		0,010 0,006	0,0047 0,0024	0,0159 0,0015	0,0092 0,0010
E907- 8	0,0028 0,0008	0,0045 0,0007		0,0049 0,0005	0,0054 0,0010	0,0165 0,0014	0,0059 0,0012
Patil (2006Pa16)			0,0020 0,0036		^(o) 0,0147 0,0005	0,0137 0,0006	0,0070 0,0006
Goswamy (1993Go10)	0,0037 0,0007			0,0037 0,0007	0,0055 0,0010	0,0163 0,0008	0,0088 0,0008
Jianming (1988Yo05)	0,006 0,002			0,006 0,002	0,0062 0,0028	0,0214 0,0041	0,012 0,006
Mardirosian (1984Ma13)						^(o) 0,030 0,007	
Iwata (1984Iw03)							
Johnson (1974Jo03)							
Meyer (1990Me15)							
Sharma (1979Sh08)							
Chi2	1,4	1,2		1,6	0,1	4,3	4,0
Chi2 crit:	3,8	4,6		3,3	3,8	2,8	3,0
UWM:	0,00449	0,00552		0,00530	0,00546	0,01520	0,00795
WM:	0,00382	0,00504		0,00441	0,00543	0,01447	0,00713
Uc (int):	0,00047	0,00054		0,00039	0,00066	0,00041	0,00036
Uc (ext) :	0,00057	0,00060		0,00049	0,00015	0,00086	0,00073
LWM :	0,0038 0,0006	0,0050 0,0006		0,00441 0,00049	0,0054 0,0007	0,0145 0,0009	0,0071 0,0007
I Abs.*	0,0037 0,0006	0,0049 0,0006	omitted	0,0043 0,0005	0,0053 0,0007	0,0142 0,0009	0,0069 0,0007

^(o) Outlier

Table 3 (Cont'd) : Relative gamma ray intensities and absolute Values calculated with ^(*)Ig602 = 97,775 (20) %.
(i, j) refer to initial and final levels, (-1, n) transition not placed in the decay scheme. DL = Detection Limit

	(10, 5) 336 keV		346,5 keV		(20, 11) 371 keV		385 keV		(20, 10) 400 keV		(14, 6) 444 keV		(20, 9) 469 keV	
	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc
E907- 2	0,079	0,008	0,0034	0,0016	0,034	0,011	0,038	0,026	0,128	0,008	0,190	0,004	0,053	0,009
E907- 3	0,073	0,004	DL=0,0064		0,033	0,006	DL=0,0078		0,120	0,006	0,198	0,006	0,038	0,003
E907- 5	0,072	0,021			>0,0217	<0,0338			0,146	0,011	0,198	0,011	0,045	0,006
E907- 6									0,175	0,066	0,211	0,076		
E907- 7	0,0733	0,0016	0,0018	0,0018	0,0295	0,0027	DL=0,0024		0,130	0,007	0,1981	0,0024	0,0518	0,0028
E907- 8	0,0708	0,0026	0,0036	0,0025	0,0333	0,0022		0,1246	0,0037	0,1901	0,0047	0,0449	0,0021	
Patil (2006Pa16)	0,076	0,002			0,0257	0,0015		0,125	0,007	0,1830	0,0021	0,0364	0,0023	
Goswamy (1993Go10)	0,0750	0,0021	0,0060	0,0013	0,034	0,008		0,124	0,013	0,1920	0,0028	0,047	0,003	
Jianming (1988Yo05)	^(o) 0,086	0,006	0,013	0,005	0,036	0,006		0,155	0,013	0,204	0,010	0,053	0,003	
Mardirosian (1984Ma13)	0,078	0,007			0,024	0,006		0,168	0,012	0,226	0,015	^(o) 0,079	0,005	
Iwata (1984Iw03)					^(o) 0,051	0,009		0,129	0,016	0,205	0,010	0,058	0,008	
Johnson (1974Jo03)					0,03	0,01		0,132	0,015	0,173	0,015	0,031	0,010	
Meyer (1990Me15)								0,15	0,01	0,20	0,01			
Sharma (1979Sh08)					0,0315	0,0025		^(o) 0,215	0,006	0,221	0,006	0,064	0,003	
Chi2	0,5		1,3		1,4			2,2		4,7		7,4		
Chi2 crit:	2,6		3,8		2,4			2,2		2,1		2,3		
UWM:	0,07459		0,00369		0,03103			0,13890		0,19929		0,04749		
WM:	0,07407		0,00414		0,02925			0,12934		0,19237		0,04685		
Uc (int):	0,00094		0,00083		0,00097			0,00219		0,00116		0,00098		
Uc (ext) :	0,00069		0,00096		0,00115			0,00323		0,00252		0,00267		
LWM :	0,0741	0,0009	0,0041	0,0010	0,0292	0,0011		0,1293	0,0032	0,199	^(e) 0,016	0,0469	0,0027	
I Abs.*	0,0725	0,0009	omitted		0,0286	0,0011	omitted	0,1264	0,0031	0,195	0,016	0,0459	0,0026	

^(o) Outlier

^(e) expanded uncertainty so range to include the most precise Value

Table 3 (Cont'd) : Relative gamma ray intensities and absolute Values calculated with ^(*)Ig602 = 97,775 (20) %.
(i, j) refer to initial and final levels, (-1, n) transition not placed in the decay scheme. DL = Detection Limit

	?(21, 9) 476 keV		(23, 10) 481 keV		498 keV		(14, 5) 525 keV		(26, 12) 530 keV		553 keV		(26, 10) 572 keV	
	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc
E907- 2	0,046	0,017	0,024	0,007	0,038	0,014	0,1428	0,005	0,043	0,006	0,019	0,005	0,020	0,004
E907- 3	DL=0,0069		^(o) 0,015	0,006			0,140	0,005	0,022	0,003			0,013	0,004
E907- 5			>0,0197	<0,0298			0,182	0,009						
E907- 6			^(o) 0,163	0,055			^(o) 0,055	0,046						
E907- 7	DL=0,0018		0,0253	0,0014	DL=0,0018		0,140	0,005	0,0281	0,0012	0,0019	0,0008	0,0153	0,0017
E907- 8	0,0020	0,0009	0,0269	0,0014	0,0007	0,0005	0,1451	0,0034	0,0431	0,0015				
Patil (2006Pa16)			0,0205	0,0010			0,1429	0,0076	0,0421	0,0013			0,0184	0,0010
Goswamy (1993Go10)			0,024	0,0020			0,14	0,02	0,043	0,002			0,0193	0,0013
Jianming (1988Yo05)			0,029	0,0080			0,165	0,010	0,047	0,011			0,025	0,010
Mardirosian (1984Ma13)			0,030	0,005			0,178	0,012						
Iwata (1984Iw03)							0,117	0,012						
Johnson (1974Jo03)							0,132	0,010						
Meyer (1990Me15)							0,16	0,01						
Sharma (1979Sh08)							0,162	0,004						
Chi2	3,5		3,1				4,5		20,6				1,3	
Chi2 crit:	6,6		2,8				2,2		2,8				3,0	
UWM:	0,02402		0,02567				0,14975		0,03828				0,01843	
WM:	0,02402		0,02367				0,14837		0,03675				0,01799	
Uc (int):	0,01171		0,00065				0,00168		0,00068				0,00070	
Uc (ext) :	0,02198		0,00115				0,00357		0,00310				0,00080	
LWM :	0,024	0,022	0,0237	0,0032			0,1484	0,0036	0,037	^(e) 0,009			0,018	0,0008
I Abs.*	omitted		0,0232	0,0031	omitted		0,1451	0,0035	0,036	0,009	omitted		0,0176	0,0008

^(o) Outlier

^(e) expanded uncertainty so range to include the most precise Value

Table 3 (Cont'd) : Relative gamma ray intensities and absolute Values calculated with ^(*)Ig602 = 97,775 (20) %.
(i, j) refer to initial and final levels, (-1, n) transition not placed in the decay scheme. DL = Detection Limit

	592 keV		(1, 0)	602 keV		(5, 3)	632 keV		(2, 1)	646 keV		(21, 6)	662 keV		669 keV		(5, 2)	709 keV	
	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	
E907- 2			100		0,100	0,008			7,57	0,07		0,041	0,004					1,358	0,019
E907- 3			100		0,098	0,004			7,59	0,10		DL=0,0063						1,388	0,016
E907- 5			100		0,109	0,007			7,603	0,027		<0,0157						1,396	0,007
E907- 6			100						7,69	0,14								1,484	0,072
E907- 7			100		0,1073	0,0010			7,58	0,03		0,0139	0,0009					1,397	0,006
E907- 8			100		0,1053	0,0028			^(o) 7,35	0,16		0,0227	0,0012	0,180	0,004			1,36	0,03
Patil (2006Pa16)	0,014	0,002	100		0,0990	0,0013			7,69	0,09		0,0148	0,0010					1,39	0,02
Goswamy (1993Go10)			100		0,1070	0,0015			7,55	0,11		0,032	0,002					1,34	0,02
Jianming (1988Yo05)			100		0,101	0,006			7,55	0,13		0,035	0,011					1,38	0,04
Mardirosian (1984Ma13)			100		0,118	0,007			^(o) 7,82	0,22		0,043	0,005					1,49	0,07
Iwata (1984Iw03)			100		0,114	0,006			7,61	0,04		0,016	0,005					1,399	0,012
Johnson (1974Jo03)			100		0,12	0,03			7,53	0,16		0,015	0,003					1,38	0,09
Meyer (1990Me15)			100		0,10	0,01			7,55	0,05								1,38	0,02
Sharma (1979Sh08)			100		0,111	0,003			7,52	0,15		0,0148	0,0015					1,465	0,029
Chi2					3,4				0,3			18,7						1,6	
Chi2 crit:					2,2				2,2			2,4						2,1	
UWM:					0,10692				7,5861			0,02480						1,4008	
WM:					0,10524				7,5911			0,01790						1,3941	
Uc (int):					0,00064				0,0152			0,00050						0,0036	
Uc (ext) :					0,00118				0,0084			0,00217						0,0046	
LWM :					0,1052	^(e) 0,0021			7,591	0,015		^(u) 0,025	^(e) 0,011					1,394	0,005
I Abs.*	Omitted		97,775	0,020	0,1029	0,0021			7,422	0,015		0,024	0,011	omitted				1,363	0,005

^(o) Outlier

^(e) expanded uncertainty so range to include the most precise Value

^(u) unweighted mean

Table 3 (Cont'd) : Relative gamma ray intensities and absolute Values calculated with ^(*)Ig602 = 97,775 (20) %.
(i, j) refer to initial and final levels, (-1, n) transition not placed in the decay scheme. DL = Detection Limit

	(6, 3) Value	713 keV Uc	(3, 1) Value	722 keV Uc	(23, 6) Value	735 keV Uc	743 keV Value	Uc	(7, 3) Value	766 keV Uc	(25, 6) Value	775 keV Uc	(6, 2) Value	790 keV Uc
E907- 2	2,26	0,03	10,81	0,10	0,137	0,005			0,012	0,003	0,0104	0,0017	0,756	0,008
E907- 3	2,324	0,026	10,95	0,13	0,132	0,006			DL=0,0072		0,000		0,753	0,012
E907- 5	2,327	0,012	10,950	0,037	0,125	0,013			>0,0177	<0,0268			0,756	0,008
E907- 6	2,33	0,09	10,95	0,20	^(o) 0,22	0,06							^(o) 0,824	0,073
E907- 7	2,33	0,01	10,96	0,04	0,1338	0,0016			0,0080	0,0012	0,0097	0,0005	0,758	0,004
E907- 8	2,26	0,05	10,73	0,24	0,1245	0,0030			0,0089	0,0014	0,0100	0,0014	^(o) 0,733	0,016
Patil (2006Pa16)	2,29	0,03	10,88	0,16	0,1399	0,0024	0,0058	0,0011	^(o) 0,0039	0,0003	0,0119	0,0012	0,766	0,012
Goswamy (1993Go10)	2,27	0,04	10,77	0,18	0,129	0,002			0,0124	⁽ⁱ⁾ 0,0002	0,0093	0,0018	0,752	0,012
Jianming (1988Yo05)	2,29	0,05	10,99	0,19	0,145	0,021			0,0092	0,0041	0,0112	0,0041	0,753	0,013
Mardirosian (1984Ma13)	2,46	0,09	^(o) 11,46	0,16	0,142	0,005			0,009	0,005	0,0112	0,0041	0,766	0,008
Iwata (1984Iw03)	2,338	0,015	11,02	0,06	0,133	0,009							0,758	0,009
Johnson (1974Jo03)	2,43	0,10	11,16	0,20	0,14	0,03							0,763	0,015
Meyer (1990Me15)	2,32	0,03	11,0	0,2	0,14	0,01							0,76	0,01
Sharma (1979Sh08)	2,42	0,05	^(o) 11,31	0,22	0,146	0,004							0,734	0,016
Chi2	1,4		0,6		2,8				2,2		0,5		0,2	
Chi2 crit:	2,1		2,2		2,2				3,0		2,8		2,3	
UWM:	2,3317		10,9300		0,1359				0,00986		0,01052		0,75832	
WM:	2,3250		10,9525		0,1342				0,01053		0,01002		0,75842	
Uc (int):	0,0061		0,0224		0,0010				0,00059		0,00041		0,00247	
Uc (ext) :	0,0072		0,0176		0,0016				0,00089		0,00030		0,00120	
LWM :	2,325	0,007	10,952	0,022	0,1342	0,0016			0,0105	0,0009	0,01002	0,00041	0,7584	0,0025
I Abs.*	2,273	0,007	10,708	0,022	0,1312	0,0016	omitted		0,0103	0,0009	0,0098	0,0004	0,7415	0,0024

^(o) Outlier

⁽ⁱ⁾ This original uncertainty was increased in order to limit the relative weight to 50 %

Table 3 (Cont'd) : Relative gamma ray intensities and absolute Values calculated with ⁶⁰Ig602 = 97,775 (20) %
(i , j) refer to initial and final levels, (-1, n) transition not placed in the decay scheme. DL = Detection Limit

	795 keV		(23, 5)	817 keV	(8, 3)	856 keV	(9, 3)	899 keV	937 keV		(10, 3)	968 keV	(9, 2)	976 keV
	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc
E907- 2			0,081	0,007	0,0203	0,006	0,023	0,009	0,0206	0,005	1,907	0,055	0,084	0,005
E907- 3			0,074	0,007	0,017	0,006	0,026	0,016	DL=0,0085		1,921	0,024	^(o) 0,095	0,011
E907- 5			0,076	0,013	>0,0187	<0,0288	<0,0187				1,909	0,013	0,088	0,013
E907- 6											^(o) 2,857	0,118		
E907- 7			0,0735	0,0037	0,0228	0,0007	0,0175	0,0010	DL=0,0012		1,926	0,008	0,0862	0,0011
E907- 8			0,0745	0,0021	0,0243	0,0017	0,0176	0,0015	0,0032	0,0012	1,873	0,042	0,0833	0,0023
Patil (2006Pa16)	0,0368	0,0012			0,0216	0,0011	0,020	0,001			^(o) 2,105	0,031	0,0841	0,0013
Goswamy (1993Go10)			0,074	0,002	0,024	0,001	0,0175	0,0014			1,92	0,028	0,0845	0,0019
Jianming (1988Yo05)			0,074	0,007	0,032	0,006	0,020	0,006			1,945	0,030	0,088	0,005
Mardirosian (1984Ma13)			0,086	0,008	0,027	0,006					2,038	0,024	0,088	0,012
Iwata (1984Iw03)			0,079	0,006	0,029	0,007	0,016	0,009			1,919	0,015	0,088	0,008
Johnson (1974Jo03)			^(o) 0,065	0,006	0,022	0,006	0,011	0,004			2,03	0,04	^(o) 0,102	0,020
Meyer (1990Me15)											1,93	0,03	0,09	0,01
Sharma (1979Sh08)			0,083	0,003	0,029	0,003	0,028	0,004			2,03	0,04	^(o) 0,097	0,004
Chi2			1,1		1,2		1,4		5,3		3,5		0,4	
Chi2 crit:			2,4		2,3		2,4		6,6		2,2		2,4	
UWM:			0,0775		0,02447		0,01962		0,0119		1,9457		0,08639	
WM:			0,0761		0,02315		0,01825		0,0119		1,9304		0,08512	
Uc (int):			0,0012		0,00048		0,00059		0,0038		0,0053		0,00070	
Uc (ext) :			0,0012		0,00052		0,00071		0,0087		0,0099		0,00042	
LWM :			0,0761	0,0012	0,0232	0,0005	0,0183	0,0007	0,012	0,009	1,93	0,01	0,0851	0,0007
I Abs.*	omitted		0,0744	0,0012	0,0227	0,0005	0,0179	0,0007	omitted		1,887	0,010	0,0832	0,0007

^(o) Outlier

Table 3 (Cont'd) : Relative gamma ray intensities and absolute Values calculated with ^(*)Ig602 = 97,775 (20) %
(i, j) refer to initial and final levels, (-1, n) transition not placed in the decay scheme. DL = Detection Limit

	(-1, 2) Value	997 keV Uc	1014 keV Value	Uc	(10, 2) Value	1045 keV Uc	(4, 1) Value	1053 keV Uc	(12, 2) Value	1086 keV Uc	1097 keV Value	Uc	1163 keV Value	Uc
E907- 2	0,025	0,007			1,884	0,036			0,041	0,004	0,034	0,008		
E907- 3	DL=0,0091		DL=0,0093		1,867	0,024	DL=0,0097		0,042	0,009			DL=0,0108	
E907- 5					1,861	0,017			0,050	0,008				
E907- 6					2,00	0,11								
E907- 7	0,0014	0,0014 ⁽ⁱ⁾	0,0025	0,0025	1,880	0,008	0,0026	0,0026	0,0369	0,0012	DL=0,0019		DL=0,0019	
E907- 8	0,0046	0,0009	0,0046	0,0014	1,841	0,041	0,0036	0,0012	0,0368	0,0018	0,0026	0,0012	0,0033	
Patil (2006Pa16)					2,026	0,022			0,0358	0,0016				
Goswamy (1993Go10)					1,87	0,03	0,005	0,002	0,038	0,002				
Jianming (1988Yo05)					1,90	0,03			0,043	0,005				
Mardirosian (1984Ma13)					2,01	0,02	0,007	0,001 ^(o)	0,058	0,005				
Iwata (1984Iw03)					1,86	0,02			0,038	0,009				
Johnson (1974Jo03)					1,92	0,04			0,031	0,005				
Meyer (1990Me15)					1,88	0,04								
Sharma (1979Sh08)					1,97	0,04			0,046	0,004				
Chi2	5,8		0,5		6,2		1,9		1,2					
Chi2 crit:	4,6		6,6		2,1		3,8		2,3					
UWM:	0,01033		0,00354		1,9123		0,00457		0,03985					
WM:	0,00343		0,00408		1,8936		0,00538		0,03739					
Uc (int):	0,00099		0,00124		0,0053		0,00070		0,00074					
Uc (ext) :	0,00238		0,00088		0,0133		0,00097		0,00081					
LWM :	0,0034	0,0024	0,0041	0,0012	1,894	0,014	0,0054	0,0010	0,0374	0,0008				
I Abs.*	0,0033	0,0023	Omitted		1,852	0,014	0,0053	0,0010	0,0366	0,0008	omitted		omitted	

^(o) Outlier

⁽ⁱ⁾ This original uncertainty was increased in order to limit the relative weight to 50 %

Table 3 (Cont'd) : Relative gamma ray intensities and absolute Values calculated with ^(*)Ig602 = 97,775 (20) %
(i, j) refer to initial and final levels, (-1, n) transition not placed in the decay scheme. DL = Detection Limit

	1180 keV		1198 keV		1205 keV		(-1, 3) 1235 keV		1253 keV		(15, 2) 1263 keV		1269 keV	
	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc
E907- 2							0,028	0,006	0,042	0,009	0,043	0,004		
E907- 3			DL=0,0112		DL=0,012				DL=0,0117		0,030	0,010	DL=0,0118	
E907- 5											0,031	0,008		
E907- 6														
E907- 7			DL=0,002		DL=0,016		0,0047	⁽ⁱ⁾ 0,0010	DL=0,0019		0,0413	0,0015	DL=0,0019	
E907- 8	0,630	0,014	0,0031	0,0009	0,0314	0,0012	0,0094	0,0012		0,0382	0,0018		0,0037	0,0013
Patil (2006Pa16)										0,0482	0,0015			
Goswamy (1993Go10)										0,042	0,002			
Jianming (1988Yo05)										0,043	0,005			
Mardirosian (1984Ma13)										0,054	0,010			
Iwata (1984Iw03)										0,046	0,015			
Johnson (1974Jo03)										0,045	0,010			
Meyer (1990Me15)														
Sharma (1979Sh08)										0,057	0,005			
Chi2							9,8				3,1			
Chi2 crit:							4,6				2,2			
UWM:							0,0141				0,0432			
WM:							0,0075				0,0432			
Uc (int):							0,00086				0,0008			
Uc (ext) :							0,00269				0,0014			
LWM :							0,0075	0,0027			0,0432	^(e) 0,0019		
I Abs.*	Omitted		omitted		omitted		0,0073	0,0026		omitted	0,0422	0,0019		omitted

^(e) expanded uncertainty so range to include the most precise Value

⁽ⁱ⁾ This original uncertainty was increased in order to limit the relative weight to 50 %

Table 3 (Cont'd) : Relative gamma ray intensities and absolute Values calculated with ^(*)Ig602 = 97,775 (20) %.
(i, j) refer to initial and final levels, (-1, n) transition not placed in the decay scheme. DL = Detection Limit

	(17, 2)	1301 keV	(3, 0)	1325 keV	(5, 1)	1355 keV	(20, 3)	1368 keV	(21, 3)	1376 keV	(22, 3)	1385 keV	1418 keV	
	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc
E907- 2	0,032	0,004	1,637	0,033	1,066	0,031	2,650	0,045	0,521	0,021	0,072	0,006		
E907- 3	0,047	0,010	1,599	0,027	1,059	0,022	2,628	0,034	0,481	0,016	0,051	0,012		
E907- 5	0,037	0,009	1,603	0,016	1,055	0,022	2,686	0,017	0,505	0,009	0,064	0,008		
E907- 6			1,582	0,137	1,011	0,114	2,65	0,13	0,516	0,099	^(o) 0,20	0,08		
E907- 7	0,0339	0,0021	1,621	0,007	1,062	0,004	2,682	0,011	0,5130	⁽ⁱ⁾ 0,0034	0,070	0,002		
E907- 8	0,037	0,003	^(o) 1,768	0,040	1,070	0,024	2,633	0,061	0,493	0,011	0,060	0,002		
Patil (2006Pa16)	0,0256	0,0013	1,707	0,026	1,093	0,017	2,7	0,034	0,543	0,007	0,064	0,002	0,005	0,002
Goswamy (1993Go10)	0,035	0,001	1,61	0,03	1,05	0,015	2,64	0,04	0,493	0,008	0,062	0,003		
Jianming (1988Yo05)	0,039	0,005	1,645	0,028	1,103	0,021	2,696	0,041	0,496	0,011	0,071	0,006		
Mardirosian (1984Ma13)	^(o) 0,061	0,008	1,69	0,29	1,108	0,022	2,758	0,069	0,531	0,046	0,079	0,025		
Iwata (1984Iw03)	0,041	0,015	1,584	0,023	1,042	0,027	2,67	0,03	0,50	0,02	0,061	0,026		
Johnson (1974Jo03)			1,67	0,04	^(o) 1,14	0,04	2,76	0,06	0,54	0,03	^(o) 0,03	0,01		
Meyer (1990Me15)			1,66	0,04	1,06	0,04	2,68	0,05	0,51	0,04				
Sharma (1979Sh08)	0,045	0,004	1,71	0,04	1,17	0,02	^(o) 2,82	0,06	^(o) 0,572	0,012	0,053	0,003		
Chi2	5,5		2,0		1,1		0,7		2,9		3,5			
Chi2 crit:	2,4		2,2		2,2		2,2		2,2		2,3			
UWM:	0,0372		1,6399		1,06493		2,6794		0,51101		0,06434			
WM:	0,0327		1,6233		1,06491		2,6796		0,51128		0,06337			
Uc (int):	0,0007		0,0051		0,00363		0,0076		0,00258		0,00096			
Uc (ext) :	0,0017		0,0073		0,00387		0,0063		0,00438		0,00180			
LWM :	^(u) 0,0372	^(e) 0,0022	1,623	0,007	1,0649	0,0039	2,680	0,008	0,5113	0,0044	0,063	^(e) 0,006		
I Abs.*	0,0364	0,0022	1,587	0,007	1,0412	0,0038	2,620	0,008	0,4999	0,0043	0,062	0,006	omitted	

^(o) Outlier

^(e) expanded uncertainty so range to include the most precise Value

^(u) unweighted mean

Table 3 (Cont'd) : Relative gamma ray intensities and absolute Values calculated with ^(*)Ig602 = 97,775 (20) %.
(i, j) refer to initial and final levels, (-1, n) transition not placed in the decay scheme. DL = Detection Limit

	1428 keV		(6, 1)	1436 keV	(20, 2)	1445 keV	(21, 2) ?	1453 keV	(7, 1)	1489 keV	1509 keV		(23, 2)	1526 keV
	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc
E907- 2	0,049	0,007	1,253	0,026	0,336	0,008	0,032	0,007	0,686	0,018	0,052	0,016	0,421	0,019
E907- 3			1,266	0,022	0,309	0,014	DL=0,0163		0,684	0,016			0,404	0,012
E907- 5			1,244	0,031	0,350	0,012			0,693	0,015			0,443	0,016
E907- 6			1,19	0,10	0,38	0,09			0,71	0,09			0,43	0,08
E907- 7	DL=0,0026		1,257	⁽ⁱ⁾ 0,005	0,336	⁽ⁱ⁾ 0,002	DL=0,0027		0,700	0,007			0,4184	⁽ⁱ⁾ 0,0026
E907- 8	0,0276	⁽ⁱ⁾ 0,0017	1,313	0,030	0,384	0,009	0,080	0,002	0,667	0,015	0,0074	0,0025	0,398	0,010
Patil (2006Pa16)			1,27	0,017	0,335	0,032			0,692	0,009	0,008	0,001	0,451	0,006
Goswamy (1993Go10)			1,25	0,016	0,334	0,005			0,687	0,009			0,414	0,006
Jianming (1988Yo05)			1,236	0,021	0,346	0,011			0,71	0,05			0,434	0,010
Mardirosian (1984Ma13)			1,34	0,27	0,329	0,014			0,72	0,02			0,433	0,008
Iwata (1984Iw03)			1,225	0,024	0,358	0,017			0,68	0,02			0,41	0,02
Johnson (1974Jo03)			1,38	0,04	0,30	0,03			0,70	0,03			0,45	0,02
Meyer (1990Me15)			1,26	0,05	0,34	0,04			0,71	0,03			0,41	0,03
Sharma (1979Sh08)			1,37	0,03	0,41	0,01			^(o) 0,80	0,02			^(o) 0,49	0,01
Chi2	4,9		2,5		7,2		21,4		0,7		3,7		3,4	
Chi2 crit:	6,6		2,1		2,1		6,6		2,2		4,6		2,2	
UWM:	0,0383		1,2748		0,3460		0,0561		0,6959		0,0225		0,4241	
WM:	0,0383		1,2619		0,3423		0,0561		0,6924		0,0081		0,4232	
Uc (int):	0,0049		0,0051		0,0021		0,0052		0,0038		0,0009		0,0022	
Uc (ext) :	0,0107		0,0080		0,0057		0,0241		0,0031		0,0018		0,0040	
LWM :	0,038	0,011	1,262	0,008	0,342	^(e) 0,007	0,056	0,024	0,6924	0,0038	0,0081	0,0018	0,423	0,005
I Abs.*	omitted		1,234	0,008	0,334	0,007	omitted		0,677	0,0037	omitted		0,414	0,005

^(o) Outlier

^(e) expanded uncertainty so range to include the most precise Value

⁽ⁱ⁾ This original uncertainty was increased in order to limit the relative weight to 50 %

Table 3 (Cont'd) : Relative gamma ray intensities and absolute Values calculated with ^(*)Ig602 = 97,775 (20) %.
(i, j) refer to initial and final levels, (-1, n) transition not placed in the decay scheme. DL = Detection Limit

	1557 keV		(25, 2)	1565 keV		(8, 1)	1580 keV		(9, 1)	1622 keV		(4, 0)	1657 keV		(10, 1)	1691 keV		(11, 1)	1720 keV	
	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc
E907- 2					0,441	0,018	0,041	0,003					46,72	1,16			0,098	0,005		
E907- 3			DL=0,0105		0,422	0,015	0,043	0,008			DL=0,0089		48,08	0,57			^(o) 0,090	0,004		
E907- 5			>0,0197	<0,0298	0,414	0,008							48,28	0,21			0,100	0,007		
E907- 6									^(o) 0,22	0,05			49,12	0,94			^(o) 0,135	0,044		
E907- 7	DL=0,0017		0,012	0,001	^(r) 0,145	0,001	0,041	0,001			DL=0,0012		48,70	0,18			0,0967	0,0007		
E907- 8	0,014	0,007	0,006	⁽ⁱ⁾ 0,001	0,354	0,009	0,042	0,001			0,0086	0,0034	46,35	1,13			0,0963	0,0025		
Patil (2006Pa16)					0,460	0,006	0,0477	0,0013					46,63	0,65			0,097	0,0180		
Goswamy (1993Go10)			0,015	0,004	0,427	0,007	0,042	0,001					49,32	0,74			0,096	0,0022		
Jianming (1988Yo05)			0,013	0,004	0,42	0,04	0,040	0,004					48,73	0,78			0,102	0,0041		
Mardirosian (1984Ma13)					^(r) 0,238	0,007	0,047	0,004					50,88	0,88			0,101	0,005		
Iwata (1984Iw03)					^(r) 0,155	0,012	0,035	0,012					48,58	0,25			0,097	0,0070		
Johnson (1974Jo03)					^(r) 0,15	0,05	^(o) 0,03	0,01					51,3	1,0			0,096	0,007		
Meyer (1990Me15)					0,42	0,03							48,4	0,8						
Sharma (1979Sh08)					0,49	0,01	0,047	0,003					50,6	1,0			^(o) 0,104	0,003		
Chi2			2,9		0,4		3,0						3,0				0,3			
Chi2 crit:			3,8		3,3		2,4						2,1				2,4			
UWM:			0,0114		0,4203		0,0425						48,692				0,09794			
WM:			0,0111		0,4217		0,0425						48,545				0,09684			
Uc (int):			0,0007		0,0047		0,0005						0,108				0,00063			
Uc (ext) :			0,0012		0,030		0,0009						0,186				0,00035			
LWM :			0,0111	0,0012	0,422	0,005	0,0425	^(e) 0,0019					48,54	0,19			0,0968	0,0006		
I Abs.*	omitted		0,0109	0,0012	0,412	0,005	0,0416	0,0019					47,46	0,19			0,0946	0,0006		

⁽ⁱ⁾ This original uncertainty was increased in order to limit the relative weight to 50 %
^(r) Removed from analysis
^(o) Outlier
^(e) expanded uncertainty so range to include the most precise Value

Table 3 (Cont'd) : Relative gamma ray intensities and absolute Values calculated with ^(*)Ig602 = 97,775 (20) %.
(i, j) refer to initial and final levels, (-1, n) transition not placed in the decay scheme. DL = Detection Limit

	1757 keV		1852 keV		1918 keV		1950 keV		1970 keV		2016 keV		2039 keV	
	Value	Uc	(13, 1) Value	Uc	(16, 1) Value	Uc	Value	Uc	Value	Uc	(18, 1) Value	Uc	(6, 0) Value	Uc
E907- 2					0,056	0,005					0,013	0,002	0,0633	0,004
E907- 3			DL=0,0077		0,051	0,003			DL=0,016		0,008	0,002	0,064	0,003
E907- 5					0,054	0,008							0,064	0,006
E907- 6	0,007	0,021	^(o) 0,341	0,061	^(o) 0,077	0,038								
E907- 7	DL=0,0009		0,0054	0,0006	0,0537	0,0005	DL=0,0006				0,0092	⁽ⁱ⁾ 0,0003	0,0636	0,0006
E907- 8			0,0008	⁽ⁱ⁾ 0,0001	0,0529	0,0019	0,053	0,011			0,0098	0,0011	^(o) 0,0753	0,0020
Patil (2006Pa16)			0,0026	0,0001	0,058	0,016					0,0090	0,0009	0,0661	0,0020
Goswamy (1993Go10)	0,0049	0,0023	0,0062	0,0009	0,055	0,002					0,0112	0,0010	0,066	0,0021
Jianming (1988Yo05)			^(o) 0,0112	0,0031	0,06	0,03					0,0124	0,0007	0,068	0,0021
Mardirosian (1984Ma13)	0,0188	0,0035	0,0025	0,0025	0,055	0,003					0,0112	0,0025	0,068	0,003
Iwata (1984Iw03)					0,052	0,004					0,0093	0,0026	^(o) 0,0589	0,0029
Johnson (1974Jo03)					0,058	0,004					0,007	0,002	0,067	0,004
Meyer (1990Me15)					0,05	0,01							0,07	0,01
Sharma (1979Sh08)					0,059	0,002					0,012	0,001	0,067	0,003
Chi2	4,0		10,5		0,8						2,9		0,9	
Chi2 crit:	4,6		3,3		2,2						2,3		2,3	
UWM:	0,01032		0,00350		0,05494						0,01017		0,06611	
WM:	0,01170		0,00314		0,05405						0,00999		0,06446	
Uc (int):	0,0024		0,0003		0,00046						0,00026		0,00051	
Uc (ext) :	0,0049		0,0009		0,00042						0,00044		0,00049	
LWM :	0,0117	0,0049	0,0031	0,0009	0,0541	0,0005					0,0100	^(e) 0,0008	0,0645	0,0005
I Abs.*	omitted		0,0030	0,0009	0,0529	0,0005	omitted		omitted		0,0098	0,0008	0,0631	0,0005

^(o) Outlier

^(e) expanded uncertainty so range to include the most precise Value

⁽ⁱ⁾ This original uncertainty was increased in order to limit the relative weight to 50%

Table 3 (Cont'd) : Relative gamma ray intensities and absolute Values calculated with ^(*)Ig602 = 97,775 (20) %.
(i, j) refer to initial and final levels, (-1, n) transition not placed in the decay scheme. DL = Detection Limit

	(19, 1)	2079 keV	(20, 1)	2090,9 keV	(21, 1)	2099 keV	(22, 1)	2108 keV	2145 keV	2151 keV	(23, 1)	2172 keV	
	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	
E907- 2	0,0289	0,003	5,28	0,20	0,046	0,003	0,052	0,003					
E907- 3	0,024	0,001	5,56	0,08	0,058	0,001	0,048	0,001			0,0030	0,0003	
E907- 5	0,018	0,002	5,59	0,05	0,054	0,003	0,057	0,004					
E907- 6													
E907- 7	0,0206	0,0006	5,63	0,02	0,0448	⁽ⁱ⁾ 0,0004	0,0430	⁽ⁱ⁾ 0,0003		DL=0,0002	0,0014	⁽ⁱ⁾ 0,0001	
E907- 8	0,0213	0,0008	5,34	0,14	0,0532	0,0016	0,0457	0,0013		0,0010	0,0005	0,0057	0,0002
Patil (2006Pa16)	^(o) 0,0741	0,0019	5,40	0,07	0,0572	0,0013	0,0501	0,0009	0,00068	0,0000			
Goswamy (1993Go10)	0,0268	0,0014	5,74	0,09	0,047	0,001	0,045	0,002			0,0021	0,0005	
Jianming (1988Yo05)	0,0163	0,0025	5,69	0,11	0,046	0,002	0,044	0,002					
Mardirosian (1984Ma13)	0,037	0,009	5,92	0,1	0,037	0,005	0,035	0,005			0,0046	0,0010	
Iwata (1984Iw03)	0,0163	0,0025	5,59	0,03	0,045	0,006	0,0438	0,0027					
Johnson (1974Jo03)	^(r) 0,081		5,86	0,14	0,051	0,020	0,056	0,010					
Meyer (1990Me15)			5,7	0,1	0,04	0,01	0,04	0,01					
Sharma (1979Sh08)	0,0305	0,0010	5,75	0,12	0,04	0,01	0,047	0,002					
Chi2	12,2		2,8		12,0		6,1				74,3		
Chi2 crit:	2,4		2,2		2,2		2,2				3,3		
UWM:	0,02393		5,6195		0,04762		0,04667				0,00337		
WM:	0,02286		5,6176		0,04824		0,04540				0,00301		
Uc (int):	0,00036		0,0150		0,00042		0,00037				0,00011		
Uc (ext) :	0,0012		0,025		0,00146		0,00092				0,00094		
LWM :	0,0229	^(e) 0,0023	5,618	0,025	0,0482	^(e) 0,0034	0,0454	^(e) 0,0024			0,0030	^(e) 0,0016	
I Abs.*	0,0224	0,0022	5,493	0,024	0,0471	0,0033	0,0444	0,0023	omitted	omitted	0,0029	0,0016	

⁽ⁱ⁾ This original uncertainty was increased in order to limit the relative weight to 50 %

^(o) Outlier

^(e) expanded uncertainty so range to include the most precise Value

Table 3 (Cont'd) : Relative gamma ray intensities and absolute Values calculated with ^(*)Ig602 = 97,775 (20) %.
 (i, j) refer to initial and final levels, (-1, n) transition not placed in the decay scheme. DL = Detection Limit

	(8, 0) 2182 keV		? (24, 1) 2204 keV		? (9, 0) 2224 keV		2232 keV		2253 keV		2256 keV		2274 keV	
	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc
E907- 2	0,043	0,003	0,030	0,002	0,021	0,013								
E907- 3	0,041	0,001												
E907- 5	0,042	0,008												
E907- 6														
E907- 7	0,0422	0,0004	0,0004	0,0002	0,0002	0,0001			DL=0,00014		DL=0,00015			
E907- 8	0,0424	0,0011	0,0051	0,0002	0,0020	0,0003			0,0006	0,0001			0,0008	0,0003
Patil (2006Pa16)	^(o) 0,036	0,007	0,0310	0,0007			0,001	0,003			0,0006	0,0002		
Goswamy (1993Go10)	0,044	0,001												
Jianming (1988Yo05)	0,045	0,002												
Mardirosian (1984Ma13)	^(o) 0,048	0,002												
Iwata (1984Iw03)	0,0398	0,0019												
Johnson (1974Jo03)	0,041	0,003												
Meyer (1990Me15)	0,04	0,01												
Sharma (1979Sh08)	0,044	0,001												
Chi2	1,0		706,4		11,9									
Chi2 crit:	2,3		3,8		4,6									
UWM:	0,04217		0,01671		0,00773									
WM:	0,04241		0,00415		0,00109									
Uc (int):	0,00032		0,00015		0,00020									
Uc (ext) :	0,00031		0,00392		0,00068									
LWM :	0,04241	0,00032	0,017	0,016	0,0011	0,0009								
I Abs.*	0,04147	0,00031	Omitted		omitted		omitted		omitted		omitted		omitted	

^(o) Outlier

Table 3 (Cont'd) : Relative gamma ray intensities and absolute Values calculated with ^(*)Ig602 = 97,775 (20) %.
(i, j) refer to initial and final levels, (-1, n) transition not placed in the decay scheme. DL = Detection Limit

	(27, 1) 2283 keV		(10, 0) 2294 keV		(11, 0) 2323 keV		2373 keV		2386 keV		(13, 0) 2455 keV		2490 keV	
	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc
E907- 2	^(o) 0,024	0,015	^(r) 0,082	0,007	^(o) 0,0098	0,0044					0,0093	0,0034		
E907- 3	0,0051	0,0004	0,029	⁽ⁱ⁾ 0,001	DL=0,005		DL=0,0049		DL=0,004					
E907- 5			0,032	0,002										
E907- 6														
E907- 7	0,0046	0,0006	0,0342	0,0010	0,0020	⁽ⁱ⁾ 0,0001					0,0015	0,0002		
E907- 8	0,0064 ^(o)	0,0004	^(o) 0,413	0,011	0,0037	0,0003					0,0019	0,0003		
Patil (2006Pa16)	0,0422	0,0010	0,056	0,023	^(o) 0,0060	0,0003	0,0009	0,0003	0,00024	0,00002	^(r) 0,0092	0,0001	0,0020	0,0010
Goswamy (1993Go10)	0,0101	0,0008	^(r) 0,076	0,005	0,0027	0,0003					0,0018	0,0002		
Jianming (1988Yo05)	0,0076	0,0014	0,031	0,005	0,0025	0,0007					0,0016	0,0006		
Mardirosian (1984Ma13)	0,010	0,002	0,045	0,002	0,004	0,001					0,0010	0,0005		
Iwata (1984Iw03)	0,0041	0,0013	0,031	0,010										
Johnson (1974Jo03)	0,007	0,002	0,025	0,005										
Meyer (1990Me15)	0,008	0,001	0,031	0,001										
Sharma (1979Sh08)	0,0051	0,0006	0,059	0,002										
Chi2	5,8		43,2		5,7						0,8			
Chi2 crit:	2,4		2,4		3,3						3,3			
UWM:	0,00677		0,0374		0,00298						0,00156			
WM:	0,00596		0,03335		0,00260						0,00164			
Uc (int):	0,00020		0,00042		0,00014						0,00012			
Uc (ext) :	0,00048		0,0027		0,00034						0,00011			
LWM :	0,0060	0,0005	0,0334	^(e) 0,0042	0,0026	^(e) 0,0006					0,00164	0,00012		
I Abs.*	0,0059	0,0005	0,0327	0,0041	0,0025	0,0006	omitted		omitted		0,00160	0,00012	omitted	

^(o) Outlier

^(e) expanded uncertainty so range to include the most precise Value

^(r) removed from analysis

⁽ⁱ⁾ This original uncertainty was increased in order to limit the relative weight to 50 %

Table 3 (Cont'd) : Relative gamma ray intensities and absolute Values calculated with ^(*)Ig602 = 97,775 (20) %.
(i, j) refer to initial and final levels, (-1, n) transition not placed in the decay scheme. DL = Detection Limit

	2515 keV		2682 keV		(20, 0) 2693 keV		2746 keV		(24, 0) 2807 keV		2814 keV		2871 keV	
	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc	Value	Uc
E907- 2			^(o) 0,007	0,003	0,0048	0,0021			^(o) 0,0069	0,003				
E907- 3					0,0019	0,0001								
E907- 5					0,0025	0,0003								
E907- 6														
E907- 7			0,0017	0,0001	0,0033	0,0001			0,0007	0,0002			0,0002	0,0001
E907- 8			0,0019	0,0001	^(o) 0,0433	0,0012			0,0016	0,0002				
Patil (2006Pa16)	0,00049	0,00001			0,0003	0,0001	0,0010	0,0001			0,0035	0,0002		
Goswamy (1993Go10)			0,0020	0,0004	0,0047	0,0005			0,0015	0,0002				
Jianming (1988Yo05)			0,0018	0,0006	0,0026	0,0016			0,0020	0,0008				
Mardirosian (1984Ma13)			^(o) 0,0025	0,0010	0,0056	0,0010								
Iwata (1984Iw03)					0,0027	0,0019								
Johnson (1974Jo03)					0,0024	0,0005								
Meyer (1990Me15)					0,0026	0,0003								
Sharma (1979Sh08)					0,0066	0,0005								
Chi2			0,7		48,2				4,5					
Chi2 crit:			3,8		2,2				3,8					
UWM:			0,00187		0,00334				0,00145					
WM:			0,00180		0,00186				0,00121					
Uc (int):			0,00006		0,00005				0,00011					
Uc (ext) :			0,00005		0,00038				0,00024					
LWM :			0,00180	0,00006	^(u) 0,0033	^(e) 0,0014			0,0012	^(e) 0,0005				
I Abs.*	omitted		0,00176	0,00006	0,0032	0,0014	omitted		0,0012	0,0005	omitted		omitted	

^(o) Outlier

^(e) expanded uncertainty so range to include the most precise Value

^(u) unweighted mean

Table 4 : Absolute gamma ray intensity values measured by the participants in the Euramet project 907; in %.

	148 keV		158 keV		185 keV		189 keV		210 keV		254 keV		291 keV		
	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	
E907- 2	0,012	0,005	0,005	0,001	DL=0,0041		0,002	0,001	0,0086	0,0012	0,0089	0,0014	0,0045	0,0008	
E907- 3	DL=0,0031		DL=0,0032				DL=0,0042		DL=0,0043		0,013 0,002		DL=0,0053		
E907- 5															
E907- 6															
E907- 7	0,0052	0,0011	0,0069	0,0014			0,0096	0,0058	0,0046	0,0023	0,0155	0,0015	0,0090	0,0009	
E907- 8	0,0029	0,00084	0,0046	0,0007			0,0053	0,0005	0,0054	0,0010	0,0159	0,0014	0,0054	0,0011	
Chi2	3,0		1,1				2,5		2,5		5,2		6,9		
Chi2 crit:	4,6		4,6				4,6		4,6		3,8		4,6		
UWM:	0,00669		0,00549				0,0056		0,00621		0,01340		0,00631		
WM:	0,00385		0,00506				0,0050		0,00649		0,01345		0,00618		
Uc (int):	0,00067		0,00055				0,0005		0,00073		0,00076		0,00053		
Uc (ext) :	0,00115		0,00056				0,0008		0,00114		0,00172		0,00140		
LWM :	0,0038	0,0012	0,0051	0,0006			0,005	0,0008	0,0065	0,0011	0,0135 ^(e)	0,0025	0,0062 ^(e)	0,0017	

	335 keV		346 keV		370 keV		385 keV		400 keV		443 keV		468 keV	
	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc
E907- 2	0,077	0,007	0,0033	0,0016	0,033	0,011	0,037	0,025	0,124	0,008	^(o) 0,186	0,004	0,052	0,009
E907- 3	0,071	0,004	DL=0,0064		0,032	0,006	DL=0,0078		0,117	0,006	0,194	0,005	0,037	0,003
E907- 5	0,071	0,020			>0,0217	<0,0328			0,143	0,011	0,193	0,011	0,044	0,006
E907- 6									0,16	0,06	0,192	0,069		
E907- 7	0,0717	0,0015	0,0018	0,0018	0,0289	0,0026	DL=0,0023		0,13	0,01	0,1938	0,0023	0,0507	0,0027
E907- 8	0,0710	0,0024	0,0034	0,0023	0,0334	0,0021			0,125	0,003	0,1899	0,0037	0,0467	0,0021
Chi2	0,0		0,3		0,6				1,0		0,2		3,0	
Chi2 crit:	3,8		4,6		3,8				3,0		3,3		3,3	
UWM:	0,07116		0,00280		0,03186				0,13252		0,19259		0,04617	
WM:	0,07144		0,00276		0,03167				0,12446		0,19289		0,04577	
Uc (int):	0,00123		0,00105		0,00155				0,00236		0,00183		0,00139	
Uc (ext) :	0,00020		0,00053		0,00122				0,00232		0,00086		0,00239	
LWM :	0,0714	0,0012	0,0028	0,001	0,0317	0,0016			0,1245	0,0024	0,1929	0,0018	0,0458	0,0024

	476 keV		481 keV		498 keV		525 keV		530 keV		553 keV		571 keV	
	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc
E907- 2	0,045	0,016	0,023	0,007	0,037	0,014	0,1393	0,0045	0,042	0,005	0,019	0,005	0,020	0,004
E907- 3	DL=0,0069		0,014	0,006			0,1367	0,0044	0,022	0,003			0,012	0,004
E907- 5			>0,0187	<0,0288			^(o) 0,178	0,009						
E907- 6			0,148	0,050			^(o) 0,050	0,042						
E907- 7	DL=0,0018		0,0248	0,0013	DL=0,0018		0,1372	0,0050	0,0275	⁽ⁱ⁾ 0,0012	0,0019	0,0008	0,0149	0,0017
E907- 8	0,0019	0,0008	0,0250	0,0012	0,0007	⁽ⁱ⁾ 0,0005	0,1449	0,0026	0,0433	0,0014				
Chi2	3,5		1,0		3,5		1,3		31,2				0,9	
Chi2 crit:	6,6		3,8		6,6		3,8		3,8				4,6	
UWM:	0,02344		0,02175		0,01884		0,13952		0,03348				0,01575	
WM:	0,02344		0,02465		0,01884		0,14138		0,03338				0,01508	
Uc (int):	0,01146		0,00088		0,00968		0,00187		0,00086				0,00143	
Uc (ext) :	0,02156		0,00087		0,01816		0,00211		0,00480				0,00136	
LWM :	0,023	0,022	0,0247	0,0009	0,019	0,018	0,1414	0,0021	0,033	^(e) 0,006			0,0151	0,0014

	602 keV		632 keV		645 keV		662 keV		669 keV		709 keV		713 keV	
	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc
E907- 2	97,5	0,7	0,098	0,007	7,386	0,058	0,040	0,004			^(o) 1,325	0,019	2,205	0,029
E907- 3	97,8	0,9	0,096	0,004	7,42	0,07	DL=0,0063				1,358	0,009	2,273	0,015
E907- 5	97,6	0,7	0,106	0,007	7,420	0,053	<0,0157				1,362	0,011	2,270	0,018
E907- 6	^(o) 91	1			^(o) 7,00	0,11					1,350	0,065	^(o) 2,12	0,08
E907- 7	97,84	0,34	0,1050	0,0012	7,417	0,028	0,0136	0,0009			1,3671	0,0056	2,28	0,01
E907- 8	98,1	1,5	0,1052	0,0023	^(o) 7,33	0,11	0,0229	0,0011	0,1793	0,0029	1,350	0,021	2,21	0,04
Chi2	0,1		1,4		0,1		36,6				0,3		2,1	
Chi2 crit:	3,3		3,3		3,8		4,6				3,3		3,3	
UWM:	97,787		0,10209		7,4117		0,0254				1,35734		2,2475	
WM:	97,769		0,10440		7,4137		0,0190				1,36347		2,2694	
Uc (int):	0,260		0,00098		0,0214		0,0007				0,00426		0,0074	
Uc (ext) :	0,071		0,00115		0,0065		0,0045				0,00246		0,0107	
LWM :	97,77	0,26	0,1044	0,0011	7,414	0,021	0,019	0,005			1,3635	0,0043	2,269	0,011

	722 keV		735 keV		765 keV		775 keV		790 keV		816 keV		856 keV	
	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc
E907- 2	^(o) 10,538	0,084	0,134	0,005	0,0114	0,003	0,0101	0,002	0,737	0,007	^(o) 0,079	0,006	0,020	0,001
E907- 3	10,713	0,072	0,129	0,005	DL=0,0072		DL=0,0073		0,737	0,010	0,072	0,007	0,017	0,005
E907- 5	10,680	0,075	0,122	0,012	>0,0167	<0,0258			0,737	0,009	0,074	0,013	>0,0177	<0,0278
E907- 6	^(o) 9,96	0,16	^(o) 0,200	0,054					0,750	0,066				
E907- 7	10,72	0,04	0,1309	⁽ⁱ⁾ 0,0016	0,0078	0,0012	0,0095	0,0005	0,742	0,004	0,0719	0,0036	0,0223	0,0007
E907- 8	10,71	0,17	0,1173	0,0022	0,0085	0,0013	0,0093	0,0013	0,727	0,012	0,0745	0,0017	0,0230	0,0015
Chi2	0,1		6,4		0,6		0,1		0,3		0,2		3,1	
Chi2 crit:	3,8		3,3		4,6		4,6		3,0		3,8		3,8	
UWM:	10,7067		0,1266		0,00925		0,00962		0,73830		0,07313		0,02049	
WM:	10,7122		0,1260		0,00836		0,00951		0,73906		0,07398		0,02108	
Uc (int):	0,0310		0,0013		0,00084		0,00044		0,00293		0,00150		0,00044	
Uc (ext) :	0,0087		0,0033		0,00063		0,00013		0,00173		0,00060		0,00078	
LWM :	10,712	0,031	0,126	^(e) 0,005	0,0084	0,0008	0,00951	0,00044	0,7391	0,0029	0,0740	0,0015	0,0211	0,0008

	899 keV		937 keV		968 keV		976 keV		997 keV		1014 keV		1045 keV	
	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc
E907- 2	0,022	0,009	0,020	0,005	1,86	0,05	0,082	0,005	0,024	0,007	0		1,837	0,033
E907- 3	0,026	0,016	DL=0,0085		1,880	0,018	^(o) 0,093	0,011	DL=0,0091		DL=0,0093		1,826	0,019
E907- 5	<0,0187				1,863	0,017	0,086	0,013					1,816	0,020
E907- 6					⁽ⁱ⁾ 2,60	0,11							1,82	0,10
E907- 7	0,0171	0,0010	DL=0,0012		1,88	0,01	0,0843	0,0010	0,0014	0,0014	0,0025	0,0025	1,839	0,008
E907- 8	0,0171	0,0014	0,0030	⁽ⁱ⁾ 0,0012	1,87	0,03	0,0824	0,0019	0,0037	⁽ⁱ⁾ 0,0007	0,0068	0,0021	1,836	0,029
Chi2	0,2		5,3		0,4		0,4		5,4		1,8		0,3	
Chi2 crit:	3,8		6,6		3,3		3,8		4,6		6,6		3,0	
UWM:	0,02041		0,0116		1,8711		0,08360		0,00971		0,00463		1,8292	
WM:	0,01717		0,0116		1,8797		0,08379		0,00300		0,00497		1,8350	
Uc (int):	0,00079		0,0037		0,0064		0,00090		0,00097		0,00161		0,0063	
Uc (ext) :	0,00035		0,0085		0,0040		0,00054		0,00224		0,00215		0,0034	
LWM :	0,0172	0,0008	0,012	0,009	1,880	0,006	0,0838	0,0009	0,0030	0,0022	0,0050	0,0021	1,835	0,006

	1053 keV		1086 keV		1097 keV		1163 keV		1180 keV		1198 keV		1205 keV	
	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc
E907- 2	DL=0,0097		0,040	0,004	0,0335 0,008		DL=0,0108				DL=0,0112		DL=0,0115	
E907- 3			0,041	0,009										
E907- 5			^(o) 0,049	0,008										
E907- 6														
E907- 7	0,0026	0,0026	0,0361	0,0012	DL=0,0019		DL=0,0019				DL=0,002		DL=0,016	
E907- 8	0,0038	0,0013	0,0369	0,0017	0,0028	⁽ⁱ⁾ 0,0013	0,0016	0,0010	0,606	0,010	0,0030	0,0009	0,0142	0,0005
Chi2	0,2		0,4		7,3									
Chi2 crit:	6,6		3,8		6,6									
UWM:	0,00319		0,03850		0,01815									
WM:	0,00357		0,03660		0,01815									
Uc (int):	0,00116		0,00095		0,00569									
Uc (ext) :	0,00051		0,00057		0,01535									
LWM :	0,0036	0,0012	0,0366	0,0009	0,018	0,015								

	1235 keV		1253 keV		1263 keV		1269 keV		1301 keV		1325 keV		1355 keV	
	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc
E907- 2	0,027	0,006	0,041	0,009	0,042	0,004	DL=0,0118		0,031	0,004	1,597	0,030	1,039	0,029
E907- 3	DL=0,0117				0,029	0,010			0,046	0,010	1,565	0,024	1,036	0,020
E907- 5					0,030	0,008			0,036	0,009	1,564	0,018	1,029	0,022
E907- 6									^(o) 1,440	0,124	^(o) 0,92	0,10		
E907- 7	0,0046	⁽ⁱ⁾ 0,0009	DL=0,0018		0,0404	0,0014	DL=0,0019		0,0332	0,0021	1,586	0,006	1,0394	0,0043
E907- 8	0,0116	0,0015	0,0005		0,0016		0,0028	0,0010	0,0376	0,0030	^(o) 1,76	0,03	^(o) 1,06	0,02
Chi2	10,8		11,0						0,9		0,7		0,1	
Chi2 crit:	4,6		6,6						3,3		3,8		3,8	
UWM:	0,01456		0,0208						0,03669		1,57771		1,03606	
WM:	0,00867		0,0208						0,03442		1,58251		1,03894	
Uc (int):	0,00105		0,0061						0,00154		0,00581		0,00405	
Uc (ext) :	0,00346		0,0202						0,00144		0,00480		0,00112	
LWM :	0,0087	0,004	0,021		0,0014				0,0344	0,0015	1,583	0,006	1,0389	0,0040

	1368 keV		1376 keV		1385 keV		1428 keV		1436 keV		1445 keV		1453 keV	
	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc
E907- 2	2,585	0,041	0,508	0,020	0,070	0,006	0,048	0,007	1,222	0,024	0,328	0,007	0,031	0,007
E907- 3	2,571	0,025	0,471	0,015	0,050	0,011			1,238	0,019	0,303	0,013	DL=0,0163	
E907- 5	2,621	0,023	0,493	0,009	0,062	0,008			1,210	0,031	0,342	0,012		
E907- 6	^(o) 2,41	0,11	0,47	0,09	^(o) 0,18	0,07			^(o) 1,08	0,09	0,35	0,08		
E907- 7	2,624	0,011	0,5019	⁽ⁱ⁾ 0,0033	0,0682	0,0018	DL=0,0025		1,230	0,005	0,3286	⁽ⁱ⁾ 0,0023	DL=0,0026	
E907- 8	2,63	0,05	0,465	0,008	0,059	0,002	0,0262	0,0016	^(o) 1,31	0,02	0,367	0,006	0,077	⁽ⁱ⁾ 0,002
Chi2	1,1		3,4		3,4		5,3		0,2		6,8		20,6	
Chi2 crit:	3,3		3,0		3,3		6,6		3,8		3,0		6,6	
UWM:	2,6058		0,4849		0,06202		0,0371		1,2249		0,3364		0,0539	
WM:	2,6154		0,4904		0,06416		0,0371		1,2296		0,3363		0,0539	
Uc (int):	0,0088		0,0038		0,00125		0,0048		0,0048		0,0030		0,0050	
Uc (ext) :	0,0093		0,0070		0,00232		0,0109		0,0024		0,0078		0,0229	
LWM :	2,615	0,009	0,490	^(e) 0,012	0,0642	^(e) 0,0041	0,037	0,011	1,2296	0,0048	0,336	0,008	0,054	0,023

	1488 keV		1505 keV		1526 keV		1557 keV		1565 keV		1579 keV		1622 keV	
	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc
E907- 2	0,669	0,017	0,051	0,016	0,410	0,018					0,430	0,018	0,040	0,003
E907- 3	0,669	0,014			0,395	0,012			DL=0,0105		0,413	0,014	0,042	0,008
E907- 5	0,676	0,015			^(o) 0,432	0,016			>0,0187 <0,0298		0,404	0,008		
E907- 6	0,65	0,09			0,39	0,07							^(o) 0,200	0,046
E907- 7	0,685	0,007	DL=0,002		0,4094	0,0025	DL=0,0017		0,0114	⁽ⁱ⁾ 0,0007	^(o) 0,1420	0,0012	0,0397	0,0008
E907- 8	0,666	0,012	0,0084	0,0028	0,398	0,007	0,013	0,007	0,0053	0,0009	^(o) 0,353	0,007	0,0397	0,0012
Chi2	0,5		3,6		0,9				20,9		0,9		0,0	
Chi2 crit:	3,0		6,6		3,3				6,6		4,6		3,8	
UWM:	0,6692		0,02970		0,4005				0,00838		0,4155		0,04028	
WM:	0,6770		0,02970		0,4077				0,00838		0,4091		0,03971	
Uc (int):	0,0051		0,01118		0,0023				0,00066		0,0066		0,00066	
Uc (ext) :	0,0037		0,02130		0,0022				0,00304		0,0064		0,00012	
LWM :	0,677	0,005	0,030	0,021	0,4077	0,0023			0,0084	0,0030	0,409	0,007	0,0397	0,0007

	1657 keV		1690 keV		1720 keV		1757 keV		1851 keV		1918 keV		1950 keV	
	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc
E907- 2	DL=0,0089		45,56	1,09	0,095	0,005	DL=0,0077				0,055	0,004		
E907- 3			47,04	0,40	^(o) 0,088	0,004					0,049	0,003		
E907- 5			47,10	0,35	0,098	0,006					0,052	0,008		
E907- 6	DL=0,0012		^(o) 44,70	0,77	^(o) 0,123	0,041	0,007	0,019	^(r) 0,31	0,06	^(o) 0,070	0,035	DL=0,0006	
E907- 7			47,65	0,18	0,0946	0,0007	DL=0,0009	0,0053	0,0006	0,0526	0,0005			
E907- 8			0,009	0,003	46,03	0,87	0,0955	0,0020	0,0008	0,0001	0,0527	0,0017		
Chi2			2,2		0,2				28,9		0,3			
Chi2 crit:			3,3		3,8				6,6		3,3			
UWM:			46,68		0,09581				0,00304		0,05235			
WM:			47,39		0,09475				0,00304		0,05254			
Uc (int):			0,15		0,00065				0,00042		0,00049			
Uc (ext) :			0,22		0,00025				0,00225		0,00028			
LWM :			47,39	0,22	0,0947	0,0006			0,0030	0,0023	0,0525	0,0005		

	1970 keV		2015 keV		2039 keV		2078 keV		2090 keV		2099 keV		2108 keV	
	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc
E907- 2	DL=0,0016		0,013	0,002	0,062	0,004	0,028	0,003	5,15	0,19	0,045	0,003	0,051	0,003
E907- 3			0,008	0,001	0,063	0,003	0,023	0,001	5,44	0,08	0,056	0,001	0,047	0,001
E907- 5					0,062	0,006	0,017	0,002	5,45	0,06	0,052	0,003	0,056	0,004
E907- 6	DL=0,0012													
E907- 7			0,0090	0,0003	0,0622	0,0006	0,0201	0,0006	5,511	0,022	0,0439	⁽ⁱ⁾ 0,0004	0,0421	⁽ⁱ⁾ 0,0003
E907- 8			0,0092	0,0010	^(o) 0,0751	0,0016	0,0212	0,0007	5,33	0,11	0,0525	0,0013	0,0456	0,0011
Chi2			1,8		0,0		4,5		1,8		17,8		6,2	
Chi2 crit:			3,8		3,8		3,3		3,3		3,3		3,3	
UWM:			0,00968		0,06220		0,02198		5,3766		0,04998		0,0482	
WM:			0,00907		0,06221		0,02120		5,4909		0,04849		0,0444	
Uc (int):			0,00026		0,00058		0,00039		0,0193		0,00062		0,0006	
Uc (ext) :			0,00034		0,00010		0,00082		0,0256		0,00260		0,0014	
LWM :			0,00907	0,00034	0,0622	0,0006	0,0212	^(e) 0,0011	5,491	0,026	0,0485	^(e) 0,0046	0,048	^(e) 0,006

	2151 keV		2172 keV		2182 keV		2203 keV		2224 keV		2253 keV		2274 keV	
	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc
E907- 2					0,042	0,003	0,030	0,002	0,020	0,013				
E907- 3			0,0029	0,0003	0,040	0,001								
E907- 5					0,040	0,008								
E907- 6														
E907- 7	DL=0,0002		0,0014	0,0001	0,0413	0,0004	0,0004	⁽ⁱ⁾ 0,0002	0,0002	⁽ⁱ⁾ 0,0001	DL=0,0001		DL=0,0002	
E907- 8	0,0016	0,0008	0,0057	0,0002	0,0435	0,0010	0,0063	0,0003	0,0020	0,0003	0,0005	0,0001	0,0008	0,0003
Chi2			172,2		1,6		241,5		12,0					
Chi2 crit:			4,6		3,3		4,6		4,6					
UWM:			0,00335		0,04131		0,01210		0,00738					
WM:			0,00317		0,04145		0,00359		0,00107					
Uc (int):			0,00011		0,00035		0,00018		0,00019					
Uc (ext) :			0,00140		0,00045		0,00280		0,00067					
LWM :			0,0032	^(e) 0,0018	0,04145	0,00045	0,0036	^(e) 0,0032	0,0011	^(e) 0,0009				

	2283 keV		2293 keV		2323 keV		2454 keV		2682 keV		2693 keV		2808 keV	
	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc	I (%)	Uc
E907- 2	0,023	0,014	0,080	0,007	0,0096	0,0043	0,0091	0,0034	0,0071	0,0033	0,0047	0,0020	0,0067	0,0027
E907- 3	0,0049	0,0004	0,028	⁽ⁱ⁾ 0,001	DL=0,005		DL=0,0049		DL=0,004		0,0019	⁽ⁱ⁾ 0,0001	DL=0,0033	
E907- 5			0,032	0,002							0,0024	0,0003		
E907- 6														
E907- 7	0,0045	0,0006	0,0335	0,0010	0,0020	⁽ⁱ⁾ 0,0001	0,0015	0,0002	0,0017	0,0001	0,0032	0,0001	0,0007	0,0002
E907- 8	0,0062	0,0003	^(o) 0,414	0,009	0,0042	0,0003	0,0018	0,0003	0,0019	0,0001	^(o) 0,0434	0,0010	0,0009	0,0001
Chi2	3,5		20,3		12,0		2,9		2,8		22,6		2,9	
Chi2 crit:	3,8		3,8		4,6		4,6		4,6		3,8		4,6	
UWM:	0,00966		0,04337		0,00526		0,00413		0,00358		0,00305		0,00277	
WM:	0,00545		0,03123		0,00311		0,00159		0,00177		0,00251		0,00084	
Uc (int):	0,00023		0,00065		0,00024		0,00016		0,00006		0,00008		0,00010	
Uc (ext) :	0,00043		0,00295		0,00082		0,00028		0,00010		0,00038		0,00018	
LWM :	0,00545	0,00043	0,0312	0,0029	0,0031	^(e) 0,0011	0,00159	0,00028	0,00177	0,00010	0,0025	0,0006	0,00084	0,00018

	2871 keV	
	I (%)	Uc
E907- 2	0,0002	0,0001
E907- 3		
E907- 5		
E907- 6		
E907- 7		
E907- 8		
Chi2		
Chi2 crit:		
UWM:		
WM:		
Uc (int):		
Uc (ext) :		
LWM :		

⁽¹⁾ This original uncertainty was increased in order to limit the relative weight to 50 %

^(o) Outlier

^(e) expanded uncertainty so range to include the most precise I (%)

^(r) removed from analysis

¹²⁵Sb - Comments on evaluation
by R. G. Helmer and E. Browne

The initial ¹²⁵Sb decay data evaluation was done by R.G.Helmer in May 2004 . This current (revised) evaluation was carried out in November 2004. The literature available by November 2004 was included.

1. Decay Scheme

¹²⁵Sb decays by β^- emission to levels in ¹²⁵Te.

The γ ray at 109 keV depopulates the isomeric level at 144 keV (half-life of 57.4 days), so its intensity depends on any chemical separation and its grow-in time. It takes about 1 year for it to be in equilibrium with the other γ rays to within 1%. The level at 35 keV is primarily fed from higher lying levels, but 27% of the 35keV γ -ray intensity comes via the isomeric level when it is at equilibrium. So, for a chemically separated source, it needs about 8 months grow-in to be at equilibrium at the 1% level.

The (direct β^- , and indirect, through γ rays) population of the isomer is 22.9 (9) % calculated from this adopted decay scheme.

2. Nuclear Data

The decay energy of 766.7 (21) keV is from the 2003 mass evaluation (2003Au03).

For the adopted decay scheme, the total radiation energy per decay is calculated to be 767 (8) keV, which agrees well with the decay energy of 766.7 (21) keV and confirms the internal consistency of this decay scheme.

The population of several additional levels has been reported, especially by 1998Sa55, but these levels are uncertain; they are : 402-, 538-, 652- and 728- keV. Verification of the associated γ rays is needed. Thus, β and γ transitions to and from these levels have not been included here.

The adopted parent half-life is 1007.54 (9) days, or 2.75855 (25) years, from the following data:

2.7 y	1950Le09
2.6 (1) y	1960Kl04
2.78 (4) y	1961Wy01
2.71 (2) y	1965F102
2.81 (5) y	1966La13
1007.3 (3) d	1980Ho17
1008.1 (8) d	1983Wa26
1007.3 (3) d	1992Un01, superseded by 2002Un02
1007.56 (10) d	2002Un02
1007.54 (9) d	Weighted average

Adopted value is the weighted average of the three precise values (which are from after 1970) which are not superseded. The reduced- χ^2 value for this average is 0.58 and the value from 2002Un02 has 89% of the relative weight.

The values from other evaluations are 2.75856 (25) years from 1999Ka26, which did not have available the value from 2002Un02, and 1007.48 (21) days from 2004Wo02 where the relative weight of the value from 2002Un02 was presumably reduced to 50%.

The level half-lives are also taken from the evaluation 1999Ka26 and are as follows:

Energy (keV)	Half-life
0	Stable
35	1.48 (1) ns
144	57.40 (15) d
321	0.673 (13) ns
443	19.1 (6) ps
463	13.2 (5) ps
525	<160 ns
636	40 (20) ps
642	≤ 70 ps
671	1.26 (6) ps

The references that provide measured values of the level half-lives are: 1965An05, 1966In02, 1967Vo21, 1968Ho05, 1968Ko08, 1969Ho42, 1970Ba69, 1970Be47, 1970Be51, 1970Ma20, 1972Be21, 1972La21, 1972Sa08, 1972Sa33, 1988GeZS, and 1992De26. Half-lives for the levels at 443, 463, and 671 keV were calculated from B(E2) values from Coulomb excitation studies (1999Ka26).

2.1 β^- Transitions

The probabilities for the β^- transitions branches are computed from the intensity balances from the γ -ray transitions for the excited states above 150 keV. Upper limits for the β^- probabilities to the 0- and 35-keV levels can be computed from the $\log ft$ systematics (1998Si17); these values are 0.002% and 1.9%, respectively. In the adopted level scheme it is assumed that both of these values are 0. The resulting values are :

Level (keV)	P_{β^-} (%)	Character	$\log ft$
0	<0.002	unique 2 nd forb.	>13.9
35	≡0	2 nd forb.	>10.6
144	13.4(9)	unique 1 st forb.	9.77
321	7.54 (9)	1 st forb.	9.32
443	0.089 (10)	2 nd forb.	10.79
463	40.3 (4)	allowed	8.04
525	1.251 (12)	1 st forb.	9.23
636	18.07 (19)	allowed	7.23
642	5.82 (5)	allowed	7.66
671	13.58 (12)	allowed	6.93

For comparison, the measured values to the 144keV level are 13.6 (9)% by 1998Gr13, 13.4% by 1959Na06, and 13.7% by 1964Ma30.

2.2 γ Transitions

The γ -ray multiplicities and mixing ratios have been taken from 1999Ka26 and the internal α -conversion coefficients are interpolated from the tables of 1978Ro22, except the E5, which is from 1976Ba63. These values are as given in the following table. The uncertainties in the internal α -conversion coefficients are taken to be 3% of the value, unless otherwise given. The total theoretical conversion coefficient of the M4 109 keV γ ray, calculated from 1978Ro22, has been reduced by 2.5% as suggested by 1990Ne01.

Energy (keV)	Multi-polarity.	Δ	%E2 or M2	α	α_K
19	[M1]			11.3	0.0
35	M1+E2	0.029 (+3-2)	0.084 (18)	14.3	12.1
109	M4			354.6	182
117	E1			0.127	0.109
(144)	[E5]			265	39.8
172	M1(+E2)	-0.004 (8)	<0.014	0.151	0.129
176	M1+E2	-0.60 (2)	26.5 (18)	0.167	0.139
178	M1+E2			0.18 (4)	0.147 (26)
198	[E2]			0.154	0.123
204	M1+E2	+1.60 (3)	72 (3)	0.128	0.104
208	M1+E2	+0.105 (14)	1.1 (3)	0.092	0.0791
227	(M1+E2)			0.084 (13)	0.070 (11)
315	(E1)			0.00839	0.00726
321	E1			0.00798	0.0691
380	E2			0.0183	0.0154
408	M1+E2	+1.50 (7)	69 (6)	0.0152	0.0129
427	M1+E2	-0.538 (11)	22.4 (9)	0.0138	0.0119
443	M1+E2	-2.3 (1)	84 (7)	0.0118	0.0100
463	E2			0.0102	0.0086
497	[M2]			0.0318	0.0271
600	E2			0.00498	0.00421
606	E2			0.00485	0.00415
635	M1+E2	+0.332 (3)	9.9 (2)	0.00526	0.00455
672	E2			0.00373	0.00319

The references that provide data on the multiplicities and mixing ratios are: 1968An15 [from α_K], 1970Na12 [α_K , K/L], 1970Wy01 [$\gamma(\theta)$], 1971Kr11 [$\gamma(\theta)$ oriented nuclei], 1971Ro17 [$\gamma(\theta)$], 1971Sa24 [$\gamma(\theta)$], 1972Ba12 [$\gamma(\theta)$], 1972Br02 [L_i/L_j], 1975Ma32 [M_i/M_j], 1982Mu02 [α_K], 1982Si18 [$\gamma(\theta)$], 1983Si14 [$\gamma(\theta)$], 1997De38

$[\gamma(\theta)]$, 1998Ro20 $[\gamma(\theta)]$, 1998Sa36 $[\alpha_K, K/L]$, 1998Sa55 $[\alpha_K]$, and 1999Sa73 $[\alpha_K]$.

The γ -ray energies have been reported by 1969Ch09, 1970Na12, 1973Gu10, 1976Wa13, 1990He05, 1998Sa55, and 2000He14, with the last three references giving the more precise values. The calibration details are not given in 1998Sa55, so it is not possible to compare these values with the others. The values of 2000He14 are on the most recent energy scale on which the energy of the strong γ ray from the decay of ¹⁹⁸Au is 411.80205 (17) keV, while those from 1990He05 are on a scale for which this energy is 411.8044 (11) keV. No correction is made here for this difference. The energies are taken from 2000He14 if they are available there, from 1990He05 as a second choice, and as indicated otherwise. (Often these values are from use of energy combinations so they can not be averaged with direct measurements). These values are: from 2000He14: 176.314 (2), 204.138 (10), 208.077 (5), 427.874 (4), 443.555 (9), 463.365 (4), 600.597 (2), 606.713 (3), 635.950 (3), and 671.441 (6); from 1990He05: 35.489 (5), 172.719 (8), 178.842 (5), 198.654 (11), 227.891 (10), 380.452 (8), and 408.065 (10); 1976Wa13 and 1998Sa55: 19.981 (6), 110.86 (7), 314.96 (8), and 497.38 (9); 1973Gu10, 1976Wa13, and 1998Sa55: 109.27 (11), and 116.95 (7).

The recommended relative and absolute γ -ray emission probabilities are discussed in section 4.2.

3. Atomic Data

3.1 X rays and Auger electrons

The fluorescence yield data are from Schönfeld and Janßen (1996Sc06) and the EMISSION code; these values are ω_K , 0.875(4); mean ω_L , 0.086 (4); and η_{KL} , 0.917 (4).

The EMISSION code also supplies the Auger electron emission probabilities; these values are: KLL, 7.0 (4); KLX, 3.17 (17); and KXY, 0.359 (20).

4 Emissions

4.1 K x-rays

The relative K x-ray emission probabilities are from 1996Sc06 and the absolute probabilities have been computed from these relative probabilities, the above γ -ray emission probabilities, and internal -conversion coefficients by using the EMISSION code.

4.2 g rays

The measured relative γ -ray emission probabilities (or intensities) are given in the following table. The values for the 109-keV γ ray are for a source in equilibrium.

Part 1

Energy	68An15 ^a	68Se11 ^b	69Ch09	70Na12	73Gu10	74I102 ^c	76Wa13	77Ar10	77Ge12
19.9							0.068 (33)		
35.5				19.6 (20)		1.42 (9)			
58.3									
109.3		0.3	0.3 (1)	0.39 (4) ^f	0.18 (2)	0.36 (4)			
110.8		~0.05				0.170 (23)	0.0031 (3)		
117.0		0.75		1.13 (1) ^f	0.75 (4) ^f	0.96 (7)	0.866 (14)	0.89 (4)	0.910 (29)
172.6		0.8	0.9 (1)	0.90 (10)	0.65 (4)	0.47 (3)	0.618 (10)	0.65 (5)	
176.3		20.5	21.2 (11)	24.9 (20)	23.9 (8)	23.2 (13)	23.06 (7) ^g	22.9 (7)	23.9 (7)
178.7		~0.1			0.08 (1)	0.05 (1)	0.092 (14)	0.10 (2)	
198.6		~0.04			0.04 (1)		0.044 (10)	0.055 (10)	
204.1		0.9	1.0 (1)	1.15 (10)	1.21 (5)	1.10 (8)	1.097 (14)	0.99 (5)	1.15 (4)
208.1		0.7	0.8 (1)	0.85 (8)	0.90 (4)	0.83 (5)	0.802 (14)	0.79 (4)	0.829 (25)
227.9	0.4 (1)	0.4		0.44 (4)	0.47 (2)	0.64 (4)	0.448 (14)	0.45 (2)	
315.0							0.0143 (14)	0.020 (4)	
321.0	1.4 (2)	1.25	1.4 (1)	1.41 (10)	1.42 (5)	1.6 (1)	1.393 (14)	1.41 (7)	1.422 (16)
380.4	5 (1)	5	5.0 (4)	5.27 (40)	5.22 (17)	5.43 (32)	5.16 (3)	5.15 (20)	5.10 (5)
408.1	0.9 (4)	0.6		0.62 (6)	0.59 (3)	0.50 (3)	0.62 (2)	0.59 (3)	

Energy	68An15 ^a	68Se11 ^b	69Ch09	70Na12	73Gu10	74I102 ^c	76Wa13	77Ar10	77Ge12
427.9	100.	100.	100.	100.	100.	100.	100.0 (3)	100.	100.0 (10)
443.4	0.5 (3)	1		1.03 (10)	1.07 (4)	1.10 (7)	1.03 (2)	1.05 (5)	
463.4	33 (4)	35.5	35.3 (20)	35.4 (28)	35.3 (13)	35.2 (23)	35.50 (7)	35.2 (10)	35.26 (37)
497.0							0.0122(14)	0.011 (2)	
600.6		61	61.2 (34)	61.5 (49)	59.6 (18)	53.6 (32)	60.39 (10)	60.1 (18)	60.6 (6)
606.6		17	17.1 (12)	16.4 (12)	16.9 (6)	19.0 (11)	17.052 (34)	16.8 (5)	17.12 (17)
635.9	42 (2)	37	37.0 (22)	37.31 (30)	38.2 (12)	35.6 (23)	38.45 (7)	38.4 (11)	38.6 (4)
671.4	6.5 (5)	6	5.6 (5)	6.0 (5)	6.09 (20)	6.24 (38)	6.11 (14)	6.02 (24)	6.18 (6)

Part 2

Energy	79Pr08	80Ro22	83Si14	84Iw03	86Wa35	93Fa02	98Sa55	90He05
19.9			0.068 (2)			0.072 (6)	0.068 (3)	
35.5			14.53 (35)			14.79 (8) ^d	17.7 (2)	
58.3			0.091 (4)			0.093 (2)	0.0042 (20)	
109.3	0.26 (4)		0.232 (5)	0.241 (24)		0.235 (16)	0.232 (6)	
110.8	0.02 (1) ^h		0.0042 (3)				0.0039 (3)	
117.0	0.91 (5)	1.01 (12)	1.060(10) ^f	0.867 (25)		0.885 (5) ^j	0.945 (15)	0.867 (24)
172.6	0.74 (6)	0.89 (6)	0.86 (2) ^f	0.69 (4)		0.72 (4)	0.67 (4)	0.659 (11)
176.3	22.9 (6)	25.45 (60)	24.5 (8)	22.62 (21)	22.91 (41)	23.65 (34)	23.09 (20)	22.96 (24)

Energy	79Pr08	80Ro22	83Si14	84Iw03	86Wa35	93Fa02	98Sa55	90He05
178.7	0.11 (1)		0.130 (5)	0.11 (4)		0.099 (6)	0.121 (2) ^j	
198.6	0.06 (1)		0.081 (4) ^f	0.030 (11)		0.046 (9)	0.044 (3)	
204.1	1.12 (4)	1.19 (22)	1.14 (4)	1.08 (3)		1.19 (5)	1.014 (10)	1.080 (23)
208.1	0.80 (4)	0.96 (10)	0.82 (2)	0.788 (21)		0.89 (3)	0.860 (10)	0.825 (16)
227.9	0.42 (2)	0.42 (7)	0.44 (2)	0.433 (12)		0.465 (25)	0.442 (9)	0.443 (23)
315.0			0.013 (2)				0.0144 (15)	
321.0	1.48 (6)	1.46 (8)	1.30 (5)	1.391 (24)		1.45 (5)	1.43 (2)	1.41 (3)
380.4	5.18 (20)	5.26 (10)	6.02 (25) ^f	5.06 (4)	5.12 (15)	5.09 (3)	5.17 (4)	5.14 (5)
408.1	0.57 (4)	0.66 (8)	0.61 (3)	0.608 (21)		0.59 (2)	0.624 (7)	0.630 (19)
427.9	100.	100.	100.	100.0 (7)	100.	100.	100.	100.0 (8)
443.5	1.06 (2)	1.03 (8)	1.12 (5)	0.989 (23)		1.03 (1)	1.05 (11)	1.019 (29)
463.4	35.1 (8)	35.45 (84)	35.50 (7)	35.23 (14)	35.4 (9)	35.64 (10)	35.12 (18)	35.07 (28)
497.0			0.015 (3)	0.009 (8)		0.018 (3)	0.009 (1)	
600.6	60.4 (11)	59.3 (12)	60.50 (10)	59.54 (22)	60.95 (67)	59.70 (10)	59.22 (18)	59.09 (45)
606.6	16.6 (5)	16.25 (62)	17.2 (3)	16.94 (7)	16.97 (26)	16.98 (21)	16.92 (6)	16.70 (14)
635.9	38.7 (8)	37.7 (10)	39.1 (2)	37.87 (14)	37.47 (27)	38.78 (32)	38.32 (12)	37.52 (30) ^h
671.4	6.04 (16)	6.92 (14) ^f	5.9 (3)	6.039 (24)	5.65 (12)	5.97 (11)	6.03 (2)	6.05 (6)

Part 3 – Adopted relative and absolute values

Energy	Adopted	wtd. avg.	S_{int}	reduced- χ^2	σ_{ext}	σ_{LWM}	$P_{\gamma}(\%) \times$ 0.2955 (24)	90Lo03 eval.	1999Ka26 eval.
19.9	0.0683 (16)	0.0683	0.0016	0.14			0.0202 (5)	0.068 (2)	0.069 (3)
35.5	19.6 (6) ⁱ	16.0	0.13	43	0.9	1.7	5.79 (18)	14.53 (35)	15.2 (10)
58.3		^e						0.091 (4)	0.05 (4)
109.3	0.231 (4)	0.2310	0.0036	1.3	0.0041		0.0683 (12)	0.233 (5)	
110.8	0.0037 (3)	0.00373	0.00017	3.6	0.00033		0.00109 (9)	0.0036 (6)	0.0035 (4)
117	0.890 (9)	0.890	0.006	2.5	0.009		0.263 (4)	1.03 (4)	0.887 (9)
172.6	0.65 (3)	0.649	0.007	4.6	0.014	0.031	0.192 (9)	0.75 (5)	0.646 (24)
176.3	23.09 (15)	23.09	0.09	2.6	0.15		6.82 (7)	23.06 (14)	23.11 (5)
178.7	0.116 (5)	0.116	0.002	5.0	0.005		0.0343 (15)	0.110 (9)	0.114 (8)
198.6	0.0448 (24)	0.0448	0.0024	0.9			0.0132 (7)	0.054 (11)	0.0432 (20)
204.1	1.06 (5)	1.061	0.007	4.6	0.015	0.047	0.313 (15)	1.105 (11)	1.070 (21)
208.1	0.833 (27)	0.833	0.006	2.3	0.009	0.027	0.246 (8)	0.808 (9)	0.837 (14)
227.9	0.443 (9)	0.443	0.005	0.5			0.131 (3)	0.437 (12)	0.443 (6)
315	0.0144 (9)	0.0144	0.0009	0.8			0.0043 (3)	0.0138 (9)	0.0136 (16)
321	1.409 (8)	1.409	0.008	0.9			0.416 (4)	1.40 (2)	1.404 (9)
380.4	5.145 (13)	5.145	0.012	1.2	0.013		1.520 (15)	5.13 (4)	5.124 (19)
408.1	0.617 (5)	0.617	0.005	0.7			0.182 (2)	0.611 (12)	0.623 (6)
427.9							29.55 (24)	100	100

Energy	Adopted	wtd. avg.	S _{int}	reduced- χ^2	σ_{ext}	σ_{LWM}	P _γ (%) × 0.2955 (24)	90Lo03 eval.	1999Ka26 eval.
443.5	1.033 (7)	1.033	0.007	1.0			0.305 (4)	1.03 (2)	1.035 (6)
463.4	35.47 (4)	35.47	0.04	1.0			10.48 (9)	35.47 (5)	35.45 (10)
497	0.0109 (11)	0.0109	0.0007	2.4	0.0011		0.0032 (3)	0.013 (2)	0.014 (8)
600.6	60.1 (4)	60.07	0.05	6.0	0.13	0.43	17.76 (18)	60.36 (11)	59.62 (16)
606.6	16.997 (27)	19.997	0.027	1.0			5.02 (5)	17.03 (3)	16.83 (6)
635.9	38.31 (14)	38.31	0.05	4.7	0.11	0.14	11.32 (10)	38.36 (15)	37.9 (3)
671.4	6.036 (17)	6.036	0.014	1.5	0.017		1.783 (16)	6.06 (2)	6.049 (19)

^a All values from this reference omitted from analysis since 5 out of 8 were outliers in an initial averaging.

^b All values from this reference omitted from analysis since they do not have uncertainties.

^c All values from this reference omitted from analysis since 9 out of 19 were outliers in an initial averaging.

^d Uncertainty increased from 0.08 to 0.20 by evaluator.

^e No value adopted; data are very inconsistent, namely, 0.091, 0.093, and 0.004.

^f Omitted from average, outlier.

^g Uncertainty increased from 0.07 to 0.20 by evaluator.

^h Typographical error in reference.

ⁱ Equilibrium intensity deduced by evaluator from transition intensity balance.

^j Uncertainty increased in analysis to reduce relative weight to 50%.

Other γ rays have been reported in various papers, but have not been included in the scheme adopted here. For those from 1998Sa55 the energies and relative emission probabilities are listed here and for the other references only the energies are given. These lines are:

1968An15: 122.4, 489.8;

1968Se11: 105.8, 391.5;

1973Gu10: 81.8, 122.4;

1974II02: 81.8, 489.8;

1976Wa13: 146.1;

1979Pr08: 81.8, 122.1, 366.0, 402.0;

1983Si14: 642.1, 693.2, 729.8; and

1998Sa55: [I_γ]: 61.8 [0.0067 (27)]; 81.0 [0.017 (1)]; 132.8 [00029 (19)]; 209.3 [0.152 (9)]; 331.8 [0.0085 (8)]; 366.5 [0.027 (2)]; 401.9 [0.0221 (2)]; 489.7 [0.0046 (23)]; 491.2 [0.016 (8)]; 503.1 [0.013 (6)]; 538.6 [0.0047 (25)]; 617.4 [0.018 (2)]; and 652.8 [0.009 (3)].

The decay scheme normalization deduced here has assumed the sum of all the γ -ray transition probabilities (photons + conversion electrons) to the ground state and 35keV level (not including that of the 35 keV γ ray) to be equal to 100%. The relative equilibrium intensity (0.231 (4)) of the 109-keV γ ray has been reduced by 5.7% in the calculation because of its apparent increase due to the 57-day half-life of the 144-keV isomer from where it decays. Also, its total M4 theoretical conversion coefficient of 363.7 has been reduced by 2.5% to 354.6 as recommended in 1990Ne01. This reduction is usually applied to theoretical M4 conversion coefficients evaluated for the Evaluated Nuclear Structure Data File (ENSDF). This procedure has produced a decay scheme normalization factor of 0.2955 (24). The resulting γ -ray emission intensities are given in the third from the last column of the table given above. The last two columns give the relative probabilities from the evaluations of 1990Lo03 and 1999Ka26. The agreement is very good except for the line at 35 keV, where evaluators have preferred to use a value deduced from a γ -ray probability balance. The relative equilibrium intensity of 19.6 (6) for the 35-keV γ ray has been obtained from a transition probability balance at the 35keV level. Its absolute emission intensity is then 5.79 (18) %.

The γ ray at 109 keV depopulates the isomeric level at 144 keV (half-life of 58 days), so its intensity depends on any chemical separation and its grow-in time. It takes about 1 year for it to be in equilibrium with the other γ rays to within 1 %. The level at 35 keV is primarily fed from higher lying levels, but 27% of the 35-keV γ -ray intensity comes via the isomeric level when it is at equilibrium. So, for a chemically separated source, it needs about 8 months grow-in to be at equilibrium at the 1% level.

The population of the isomer was measured to be 24.3 (3) % (1998Gr13) compared to the 22.9 (9) % calculated from this adopted scheme.

4.3 Conversion electrons

From the adopted γ -ray intensities, and the conversion coefficients, one obtains the following conversion electron emission probabilities:

γ energy (keV)	shell	electron energy	emission prob. (%)
19.80	L	14.86	0.184 (7)
	M	18.79	0.0368 (14)
	N	19.63	0.0077 (3)
35.49	K	3.675	70 (3)
	L	30.55	9.5 (4)
	M	34.48	1.9 (1)

γ energy (keV)	shell	electron energy	emission prob. (%)
	N	35.35	0.46 (2)
109.28	K	77.46	12.4 (5)
	L	104.33	9.2 (5)
	M	108.27	2.1 (1)
	N	109.11	0.45 (2)
116.96	K	85.14	0.0287 (11)
	L	112.02	0.00371 (15)
172.72	K	140.90	0.0248 (10)
	L	167.78	0.0032 (1)
176.31	K	144.50	0.95 (4)
	L	171.37	0.150 (6)
	M	175.30	0.031 (1)
178.84	K	147.03	0.0050 (8)
	L	173.90	0.0009 (3)
198.65	K	166.84	0.00161 (10)
204.14	K	172.32	0.0322 (19)
	L	199.19	0.0059 (4)
	M	203.13	0.00120 (7)
208.08	K	176.26	0.0192 (8)
	L	203.13	0.00248 (10)
227.89	K	196.08	0.0090 (15)
	L	222.95	0.0014 (5)
321.04	K	289.23	0.00284 (11)
380.45	K	348.64	0.0231 (9)
	L	375.51	0.0035 (1)
408.06	K	376.25	0.00232 (9)
427.87	K	396.06	0.35 (2)
	L	422.94	0.0450 (18)
	M	426.87	0.0090 (3)
443.56	K	411.74	0.00302 (12)

γ energy (keV)	shell	electron energy	emission prob. (%)
463.36	K	431.55	0.090 (4)
	L	458.43	0.0128 (5)
	M	462.36	0.0026 (1)
600.60	K	568.78	0.074 (3)
	L	595.66	0.0101 (4)
	M	599.59	0.0020 (1)
606.72	K	574.90	0.0206 (8)
	L	601.77	0.0028 (1)
635.95	K	604.14	0.0509 (20)
	L	631.01	0.0063 (2)
671.44	K	639.62	0.00564 (22)
	L	666.50	0.0008

References

- 1950Le09 - G. R. Leader, W. H. Sullivan, *NNES* **9** (1950) 934 [$T_{1/2}$]
1959Na06 - R. S. Narcisi, Thesis, Harvard University (1959); AECU-4336 (1959) [I_{β}]
1960Kl04 - E. H. Klehr, A. F. Voigt, *J. Inorg. Nuclear Chem.* **16**(1960)8 [$T_{1/2}$]
1961Wy01 - E. I. Wyatt, S. A. Reynolds, T. H. Handley, W. S. Lyon, H. A. Parker, *Nucl. Sci. Eng.***11**(1961)74 [$T_{1/2}$]
1965An05 - G. Andersson, G. Rudstam, G. Sorensen, *Ark. Fys.* **28**(1965)37 [$T_{1/2}$ level]
1965Fl02 - K. F. Flynn, L. E. Glendenin, E. P. Steinberg, *Nucl. Sci. Eng.* **22**(1965)416 [$T_{1/2}$]
1966In02 - T. Inamura, T. Iwashita, S. Kageyama, *J. Phys. Soc. Japan* **21**(1966)2425 [$T_{1/2}$ level]
1966La13 - F. O. Lawrence, W. R. Daniels, D. C. Hoffman, *J. Inorg. Nucl. Chem.* **28**(1966)2477 [$T_{1/2}$]
1968An15 - D. S. Andreev, V. K. Bondarev, L. N. Laperin, A. Z. Ilyasov, I. K. Lemberg, *Bull. Acad. Sci. USSR, Phys. Ser.* **32**(1969)225 [I_{γ} , multipolarity]
1968Ho05 - C. Hohenemser, R. Rosner, *Nucl. Phys.* **A109**(1968)364 [$T_{1/2}$ level]
1968Ko08 - J. Kownacki, J. Ludziejewski, M. Moszynski, *Nucl. Phys.* **A113**(1968)561 [$T_{1/2}$ level]
1968Se11 - H. Sergolle, *Compt. Rend.* **267B**(1968)1042 [I_{γ}]
1969Ch09 - P. R. Christensen, A. Berinde, I. Neamu, N. Scintei, *Nucl. Phys.* **A129**(1969)337 [E_{γ} , I_{γ}]
1969Ho42 - R. R. Hosangdi, P. N. Tandon, S. H. Devare, *Indian J. Pure Appl. Phys.* **7**(1969)604 [$T_{1/2}$ level]
1970Ba69 - M. M. Bajaj, S. L. Gupta, N. K. Saha, *Proc. Nat. Inst. Sci. India* **36A**(1970)176 [$T_{1/2}$ level]
1970Be47 - E. E. Berlovich, V. V. Lukashevich, A. V. Popov, V. M. Romanov, *Sov. J. Nucl. Phys.***12**(1971)117 [$T_{1/2}$ level]
1970Be51 - B. Bengston, M. Moszynski, *Nucl. Instrum. Methods* **85**(1970)133 [$T_{1/2}$ level]
1970Ma20 - A. Marelus, J. Lindskog, Z. Awwad, K. G. Valivaara, S. E. Hagglund, J. Pihl, *Nucl. Phys.* **A148**(1970)433 [$T_{1/2}$ level]
1970Na12 - T. S. Nagpal, R. E. Gaucher, *Can. J. Phys.* **48**(1970)2978 [E_{γ} , I_{γ} , multipolarity]
1970Wy01 - L. D. Wyly, J. B. Salzberg, E. T. Patronis, Jr., N. S. Kendrick, C. H. Braden, *Phys. Rev.* **C1**(1970)2062 [Multipolarity]
1971Kr11 - K. S. Krane, J.R. Sites, W. A. Seyert, *Phys. Rev.* **C4**(1971)565 [Multipolarity]
1971Ro17 - M. Rots, R. Silverans, R. Coussement, *Nucl. Phys.* **A170**(1971)240 and private

- communication.(March 1972) [Multipolarity]
- 1971Sa24 - G. Satyanarayana, V. Lakshminarayana, Curr. Sci. (India) **40**(1971)458 [Multipolarity]
- 1972Ba12 - T. Badica, S. Dima, A. Gelberg, I. Popescu, Z. Phys. **249**(1972)321 [Multipolarity]
- 1972Be21 - B. Bengtson, M. Moszynski, Nucl. Instrum. Methods **100**(1972)293 [$T_{1/2}$ level]
- 1972Br02 - D. S. Brenner, M. L. Perlman, Nucl. Phys. **A181**(1972)207 [Multipolarity]
- 1972Sa08 - G. Satyanarayana, V. Lakshminarayana, D. S. Murty, Can. J. Phys. **50**(1972)600 [$T_{1/2}$ level]
- 1972Sa33 - G. Satyanarayana, V. V. Ramamurty, V. Lakshminarayana, J. Phys. (London)**A5**(1972)1243 [$T_{1/2}$ level]
- 1973Gu10 - J. B. Gupta, N. C. Singhal, J. H. Hamilton, Z. Phys. **261**(1973)137 [E_γ , I_γ]
- 1974II02 - P. Ila, K. Sudhakar, K. L. Narasimham, V. Lakshminarayana, Curr. Sci. (India) **43**(1974) 176 [I_γ]
- 1975Ma32 - B. Martin, D.Merkert, J.L. Campbell. Z. Phys. **A274**(1975)15 [Multipolarity]
- 1976Ba63 - I. M. Band, M. B. Trzhaskovskaya, M. A. Lis tengarten, Atomic Data Nucl. Data Tables **18**(1976)433 [α]
- 1976Wa13 - W.B. Walters, R.A. Meyer, Phys. Rev. **C14**(1976)1925 [E_γ , I_γ]
- 1977Ar10 - G. Ardisson, K. Abdmeziem, Radiochem. Radioanal. Lett. **29**(1977)1 [I_γ]
- 1977Ge12 - R. J. Gehrke, R.G. Helmer, R. C. Greenwood, Nucl. Instrum. Methods **147**(1977)405 [I_γ]
- 1978La21 - F. Lagoutine, J. Legrand, C. Bac, Int. J. Appl. Radiat. Isotop. **29**(1978)269 [$T_{1/2}$ level]
- 1978Ro22 - F. Rösel, H. M. Fries, K. Alder, H. C. Pauli, Atomic Data Nucl. Data Tables **21**(1978)92 [α]
- 1979Pr08 - R.Prasad, Czech. J. Phys. **B29**(1979)737 [I_γ]
- 1980Ho17 - H. Houtermans, O. Milosevic, F. Reichel, Int. J. Appl. Radiat. Isotop. **31**(1980)153 [$T_{1/2}$]
- 1980Ro22 - W. M. Roney, Jr., W. A. Seale, Nucl. Instrum. Methods **171**(1980)389 [I_γ]
- 1982Mu02 - P. Mukherjee, S. Bhattacharya, S. Sarkar, I. Mukherjee, B. K. Dasmahapatra, Phys. Rev. **C25**(1982)2120 [Multipolarity]
- 1982Si18 - K. Singh, H. S. Sahota, Indian J. Phys. **56A**(1982)291 [Multipolarity]
- 1983Si14 - K. Singh, H. S. Sahota, Indian J. Pure Appl. Phys. **21**(1983)19 [I_γ]
- 1983Wa26 - K. F. Walz, K. Debertin, H. Schrader, Int. J. Appl. Radiat. Isotop. **34** (1983)1191 [$T_{1/2}$]
- 1984Iw03 - Y. Iwata, M. Yasuhara, K. Maeda, Y. Yoshizawa. Nucl. Instrum. Methods **219**(1984)123 [I_γ]
- 1986Wa35 - Wang Xinlin, Li Xiaodi, Du Hongshan, Chin. J. Nucl. Phys. **8**(1986)371 [I_γ]
- 1988GeZS - A. M. Geidelman, Yu. S. Egorov, N. K. Kuzmenko, V. G. Nedovesov, V. P. Chechev, G. E. Shukin, Proc. Intern. Conf. Nuclear Data for Science and Technology, Mito, Japan, 1988, p.909 [$T_{1/2}$ level]
- 1990He05 - R. G. Helmer, Appl. Radiat. Isot. **41**(1990)75 [E_γ , I_γ]
- 1990Lo03 - L. Longoria -Gandara, M. U.Rajput, T. D. Mac Mahon, Nucl. Instrum. Methods Phys. Res. **A286**(1990)529 [I_γ evaluation]
- 1990Ne01 - Zs. Nemeth and A. Veres, Nucl. Instrum. Methods Phys. Res. **A286**(1990)601. [Theoretical conversion coefficients for M4 transitions]
- 1992De26 - C. C. Dey, B. K. Sinha, R. Bhattacharya, Nuovo Cim. **105A**(1992)523 [$T_{1/2}$ level]
- 1993Fa02 - N. I. Fawwaz, N. M. Stewart, J. Phys. (London) **G19**(1993)113 [I_γ]
- 996Sc06 - E. Schönfeld, H. Janßen, Nucl. Instrum. Methods **A369**(1996)527 [ω]
- 1997De38 - C. C. Dey, B. K. Sinha, R. Bhattacharya, Can. J. Phys. **75**(1997)591 [Multipolarity]
- 1998Gr13 - A. Grau Carles, L. Rodriguez Barquero, A. Jimenez de Mingo, Appl. Radiat. Isot. **49** (1998)1377 [I_β]
- 1998Ro20 - M. Roteta, E. Garcia-Torano, Appl. Radiat. Isot. **49**(1998)1349 [Multipolarity]
- 1998Sa36 - M. Sainath, K. Venkataramaniah, Nuovo Cim. **111A**(1998)223 [Multipolarity]
- 1998Sa55 - M. Sainath, K. Venkataramaniah, P. C. Sood, Phys. Rev. **C58** (1998)3730 [E_γ , I_γ , multipolarity]
- 1998Si17- B. Singh, J. L. Rodriguez, S. S. M. Wong, J. K. Tuli, Nucl. Data Sheets **84**(1998)487 [log ft sys.]
- 1999Ka26 - J. Katakura, Nucl. Data Sheets **86**(1999)955 [evaluation]
- 1999Sa73 - M.Sainath, K.Venkataramaniah, P.C.Sood, Pramana 53(1999)289 [Multipolarity]
- 2000He14 - R.G.Helmer, C.van der Leun, Nucl. Instrum. Methods Phys. Res. **A450**(2000)35 [E_γ]
- 2002Un02 - M.P.Unterweger, Appl. Radiat. Isot. **56**(2002)125 [$T_{1/2}$]
- 2003Au03 - G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. **A729**(2003)337 [Q]
- 2004Wo02 - M.J. Woods, S.M. Collins, Appl. Radiat. Isot. **60**(2004)257 [$T_{1/2}$ evaluation]

¹²⁵I - Comments on evaluation of decay data by V. Chisté, E. Schönfeld and M.M. Bé

This evaluation was completed in July 2010. Literature by July 2010 was included.

1 Decay Scheme

Given the adopted Q_{EC} value of 185.77 keV, there are three levels in the daughter nuclide ¹²⁵Te available for the EC decay of the ¹²⁵I ground state. The level at 144.8 keV ($J\pi=11/2^-$) would require a 3rd forbidden transition ($\Delta I = 3$ and parity change) and one may expect $\log ft > 15$ from systematics (Raman, 1973Ra10), which corresponds to a transition probability of $< 1 \cdot 10^{-8}$ per disintegration.

A direct decay to the ground state of ¹²⁵Te ($\Delta I = 2$, no parity change, non-unique 2nd forbidden) was not observed. Smith and Lewis (1966Sm05) have found that the transition probability of such a transition would be smaller than 0.01. From systematics (Raman, 1973Ra10), one may expect $\log ft > 11.0$ which corresponds to a transition probability of $1 \cdot 10^{-6}$ per disintegration.

The adopted decay scheme of ¹²⁵I presented in this evaluation is complete. Good agreement is found between the effective Q value (185.66 (42) keV) calculated from the decay scheme data and that recommended from the atomic mass evaluation of Audi (2003Au03).

2 Nuclear Data

The Q_{EC} value of 185.77 (6) keV is taken from Audi (2003Au03), while the spins, parities and the lifetime of the excited 35-keV level (1.48 (1) ns) are from the ENSDF evaluation of J. Katakura (1999Ka26).

Experimental ¹²⁵I half-life values (in days) are given in Table 1:

Table 1: Experimental values of ¹²⁵I half-life.

Reference	Experimental value (days)	Comments
A. F. Reid (1946Re**)	56	Not used: no uncertainty.
G. Friedlander (1951Fr21)	60.0 (5)	
M. Ia. Kuznetsova (1958Ku**)	60	Not used: no uncertainty.
C. M. E. Matthews (1960Ma36)	57.4 (2)	
G. I. Gleason (1963Ge**)	58.8 (2)	Private communication. Cited by 1965An07
H. Leutz (1964Le05)	60.25 (6)	
S. C. Anspach (1965An07)	59.83 (11)	Superseded by 2002Un02.
C. R. Richmond (1966Ri14)	58.76 (13)	
F. Lagoutine (1968La10)	59.89 (18)	Superseded by 1995Ra32 (2).
J. F. Emery (1972Em01)	60.18 (17)	
W. Künding (1979Kü**)	59.666 (16)	
H. Houtermans (1980Ho17)	59.156 (20)	
D. D. Hoppes (1982HoZJ)	59.47 (21)	Superseded by 2002Un02.
H. Kubo (1983Ku**)	59.56 (17)	
H. Schrader (1987Sc20)	59.39 (2)	
B. R. S. Simpson (1989Si19)	59.40 (5)	
P. de Felice (1990De09)	59.38 (3)	
M. J. Woods (1990Wo03)	59.416 (10)	
T. Altzitzoglou (1991Al05)	59.37 (6)	

G. Ratel (1995Ra32) - 1	59.29 (7)	SIR: result of AECL.
G. Ratel (1995Ra32) - 2	59.90 (11)	SIR: result of LNHB.
G. Ratel (1995Ra32) - 3	59.26 (3)	SIR: result of NCR.
M. P. Unterweger (2002Un02)	59.49 (13)	
Recommended value	59.388 (28)	$\chi^2 = 3.3$

The first twelve values are only cited for reasons of completeness (1946Re**, 1951Fr21, 1958Ku**, 1960Ma36, 1963Ge**, 1964Le05, 1965An07, 1966Ri14, 1968La10, 1972Em01, 1979Kü**, 1980Ho17). The values of 1982HoZJ, 1983Ku**, 1987Sc20, 1989Si19, 1990De09, 1990Wo03, 1991Al05, 1995Ra32 (1, 2 and 3) and 2002Un02 are in good agreement and this indicates that the true value is very close to 59,4 d. Taking this into account, the evaluators can classify the older values (1946Re** until 1980Ho17) (when looking at the uncertainty given by the authors) into values which are too low (1960Ma36 by 10 σ , 1966Ri14 by 4 σ , 1980Ho17 by 12 σ) or too high (1964Le05 by 14 σ , 1968La10 by 10 σ , 1972Em01 by 5 σ , 1979Kü** by 17 σ). These values have a large spread. The evaluators consider it to be sensible to calculate an average of these values only after enlarging their uncertainties by reasonable (but more or less arbitrarily chosen) factors. Values of 1946Re** and 1958Ku** are excluded because no uncertainty is given. Value of 1951Fr21 is a good value but because of its large uncertainty it does not contribute very much to the average.

As there are enough new accurate values the evaluators do not include all these old values into the averaging procedure. A weighted average of the eleven remaining values, of 1982HoZJ until 2002Un02, was calculated using the LWEIGHT computer code (version 3). The largest contribution to the weighted average comes from the value of M. J. Woods (1990Wo03), with a statistical weight of 63 %. The LWEIGHT computer code increases the uncertainty for the 1990Wo03 value from 0.010 to 0.013 in order to reduce its relative weight from 63 % to 50 %.

The adopted half-life value is 59.388 days with a final uncertainty of 0.028 days, expanded to include the most precise value of M. J. Woods (1990Wo03). The reduced- χ^2 value is 3.3.

2.1 Electron Capture Transition.

The energy of the electron capture transition has been obtained from the Q(EC) value (2003Au03) and the level energy given by J. Katakura (1999Ka26).

The adopted electron-capture transition probability and the associated uncertainty were deduced from the γ -ray transition probability balance at 35-keV level of the decay scheme.

The adopted P_K , P_L and P_M values were calculated from the table of Schönfeld (1995ScZY) using the adopted Q_{EC} value (program EC-Capture). The adopted P_K value of 0.8011 (17) can be compared with some experimental determination (Table 2):

Der Mateosian (1953De26) found $P_L/P_K = 0.23$ (3). Leutz and Ziegler (1964Le05) found $(P_L + P_M + P_N)/P_K = 0.2547$ (33) and 0.2539 (21) using two different extrapolation methods. The mean value 0.2543 (27) corresponds to $P_K = 0.797$ (3). Smith and Lewis (1966Sm05) found for the above ratio 0.253 (5). Karttunen et al. (1969Ka08) measured $P_K \omega_K = 0.685$ (18) (2 σ) whereas Plch and Zderadicka (1974Pl03) found 0.685 (12) and Tolea et al. (1974To04) 0.699 (30).

Table 2: Adopted and measured values of P_K .

1	0.797 (3)	Leutz and Ziegler (1964Le05) measured value.
2	0.798 (3)	Smith and Lewis (1966Sm05) measured value.
3	0.783 (11)	Karttunen et al. (1969Ka08) measured, recalculated and uncertainty related to 1 σ .
4	0.783 (15)	Plch and Zderadicka (1974Pl03) measured, recalculated.
5	0.799 (34)	Tolea et al. (1974To04) measured, recalculated.
6	0.801	Tolea et al. (1974To04); calculated from theory.
7	0.825 (35)	Kalyani et al. (1996Ka48) measured value.
8	0.8011 (17)	calculated from theory and adopted in the present evaluation.

Values 3, 4 and 5 are recalculated using the present adopted value of ω_K .

2.2 γ Transitions

The γ -ray transition probability for the 35-keV gamma-ray was calculated using the γ -ray emission intensity and the relevant internal conversion coefficient (see **5.2 Gamma Emissions**).

Multipolarity of the 35-keV γ -ray transition is M1 + E2. The mixing ratio (δ) was deduced by comparison between the experimental and theoretical total internal coefficients, the later calculated using the BrIcc computer code (2008Ki07).

The total coefficient of 35-keV γ -ray can be deduced from:

$$\alpha_T = \frac{P_{(\gamma+ce)35}}{I_{\gamma35}} - 1$$

where:

- $P_{(\gamma+ce)35}$ (100 %) is the transition probability of 35-keV γ -ray.
- $I_{\gamma35}$ (= 6.63 (6) %) is the weighted average of the experimental values of absolute emission intensity shown in Table 5 (see **5.2 Gamma Emissions**).

Table 3 shows the final results of experimental α_T , as well as the mixing ratio δ deduced from the comparison between experimental α_T value (column 2) and theoretical values of 13.63 (M1) and 77.3 (E2) given by the BrIcc computer code.

Table 3: Adopted conversion coefficient and mixing ratio.

E_γ (keV)	α_T experimental (given by equation above)	δ (mixing ratio)
35.4922 (5)	14.08 (14)	0.085 (13)

Then the internal conversion coefficients (ICC) and the associated uncertainties have been obtained using the BrIcc computer program with “the frozen orbital approximation” (2008Ki07).

These α_T and mixing ratio values can be compared with the experimental results:

- Geiger et al. (1965Ge04) compared their measured conversion electron results $L_1/L_2/L_3 = 1/0.089(4)/0.024(2)$ with the theoretical ratios derived from the table of Sliv and Band and found: 99.965 (20) % M1 + 0.035 (20) % E2.
- Mazets et al. (1966Ma49) found in the ^{125}Sb decay $L_1/L_2/L_3 = 10.7/1.0/0.2$ corresponding to 99.92 (3) % M1 + 0.08 (3) % E2.
- Karttunen et al. (1969Ka08) deduced from a comparison of experimental results with those of the Hager-Seltzer theory that a possible E2 admixture is smaller than 0.4 %.

- Casey et al. (1969Ca01) measured $L_1/L_2/L_3 = 1 / 0.106 (22) / 0.041 (2)$.
- Coursol (1979CoZG) measured more precisely $L_1/L_2/L_3 = 1 / 0.0820 (13) / 0.0190 (10)$ and derived from analysis an E2 admixture of 0.03 (2) %.
- Brabec et al. (1982Ba16) measured $L_1/L_2/L_3 = 1.00 (1) / 0.0954 (18) / 0.0229 (49)$ and derived from analysis of these and other data $\delta = 0.029 (+ 3 - 2)$ and 99.916 (+ 18 - 11) % M1 and 0.084 (+ 18 - 11) % E2.

The experimental conversion coefficient values of the 35.5 keV transition are compiled in the following table (Table 4).

	1 (1952Bo16)	2 (1969Ka08)	3 (1970Ma51)	4 (1979CoZG) ^μ	Adopted values (given by BrIcc)
α_K	11.7 (25)	11.78 (11)	11.8 (3)	11.90 (31)	11.70 (17)
α_L	1.6 (5)	1.62			1.91 (8)
α_M	0.3 (1)	0.25			0.386 (16)
α_N		0.044			0.075 (3)
α_O					0.00766 (23)
α_T	13.6 (26) [*]	13.65 (28) [£]		14.25 (64)	14.08 (22)

* $\alpha_T = \alpha_K + \alpha_L + \alpha_M$.

£ Value from the article.

μ The original uncertainties, given in this paper, are 3σ .

- 1 - Bowe and Axel (1952Bo16): α_L and α_M calculated from α_K and the measured ratio $K/L/M = 0.80 (5) / 0.11 (2) / 0.020 (4)$. ($\alpha_T = \alpha_K + \alpha_L + \alpha_M$)
- 2 - Karttunen et al. (1969Ka08): α_K recalculated from $\alpha_K\omega_K$ with $\omega_K (0.875 (4))$ as adopted here. The originally published value is $\alpha_K = 12.01 (18)$ (2σ uncertainty). The values of α_L , α_M , α_N are calculated from α_K using the ratio $K/L/M/N = 80/11/1.7/0.3$ as measured by Narcisi (1959Na06).
- 3 - Marelus et al. (1970Ma51).
- 4 - Coursol (1979CoZG, see also IAEA TecDoc-619 (1991)): measured values and the original uncertainties are 3σ (include in the Table 4).

3 Atomic Data

Atomic values, ω_K , ω_L , ω_M and n_{KL} and X-ray and Auger electron relative probabilities are from Schönfeld and Janßen (1996Sc06)

4 Electron emissions

The conversion electron emission probabilities have been deduced from the ICC and γ -ray emission probability values.

5 Emissions

5.1 K x-rays

The X-ray absolute intensities were deduced from the decay data using the EMISSION computer code.

For the total K X ray emission intensities some experimental values (per 100 % disintegrations) have been found:

1	137.9 (27)	Karttunen et al. (1969Ka08)
2	139.3 (25)	Tolea et al. (1974To04)
3	137.9 (23)	Plch and Zderadicka (1974PI03)
4	138.3 (20)	Konstantinov et al. (1989Ko**), recalculated from $P(KX + \gamma) = 1.45$ (2)
5	138.3 (12)	Weighted mean of four experimental values (1-4). $\chi^2 = 0.07$
6	137.9 (10)	Adopted value.

5.2 Photon emissions

The energy of the 35-keV γ -ray given in section 5.2 is from J. Katakura (1999Ka26).

The experimental absolute 35-keV γ -ray emission intensities from the decay of ^{125}I are given in the table 5.

Table 5: Absolute experimental γ -ray emission intensities for the 35-keV transition.

Reference	Absolute γ -ray intensity (%)	Comments
J. C. Bowe (1952Bo16)	7 (2)	
E. Karttunen (1969Ka08)	6.83 (14)	Original uncertainty = 2σ .
N. F. Coursol (1979CoZG) \ddagger	6.56 (9)	Original uncertainty = 3σ .
W. B. Mann (1985Ma**)	6.67 (22)	Not used: evaluated value.
A. Iwahara (1990Iw04)	6.68 (14)	Superseded by 2006Da20
U. Schötzig (1992ScZZ)	6.55 (13)	
M. A. L. da Silva (2006Da20)	6.67 (14)	
Recommended value	6.63 (6), $\chi^2 = 0.78$	

\ddagger see also IAEA TecDoc-619 (1991).

The adopted value is the weighted average of 6.63 % with an external uncertainty of 0.06 %. The reduced- χ^2 value is 0.78.

6 References

- 1946Re** A. F. Reid, A. S. Keston, Phys. Rev. 70(1946)987 [Half-life].
- 1951Fr21 G. Friedlander, W. C. Orr, Phys. Rev. 84(1951)484 [Half-life].
- 1952Bo16 J. C. Bowe, P. Axel, Phys. Rev. 85(1952)858 [α].
- 1953De26 E. der Mateosian, Phys. Rev. 92(1953)938 [P_{LM}/P_K].
- 1958Ku** M. Ia. Kuznetsova, V. N. Mekhedov, V. A. Khalkin, Sov. Phys. TEP 34(1958)759 [Half-life].
- 1959Na06 R. S. Narcisi, AECU-4336(1959) [α].
- 1960Ma36 C. M. E. Matthews, Phys. Med. Biol. 5(1960)45 [Half-life].
- 1963Gl** G. I. Gleason, Priv. Comm. cited by 1965An07 [Half-life].
- 1964Le05 H. Leutz, K. Ziegler, Nucl. Phys. 50(1964)648 [Half-life, P_K].
- 1965An07 S. C. Anspach, L. M. Carvalho, S. B. Garfinkel, J. M. R. Hutchinson, C. N. Smith, N.P. 15663 (260/9) [Half-life].
- 1965Ge04 J. S. Geiger, R. L. Graham, I. Bergstrom, F. Brown, Nucl. Phys. 68(1965)352 [δ].
- 1966Ri14 C. R. Richmond, J. S. Findlay, Health Phys. 12(1966)865 [Half-life].
- 1966Sm05 K. M. Smith, G. M. Lewis, Nucl. Phys. 89(1966)561 [P_{LMN}/P_K].

- 1966Ma49 E. P. Mazets, Y. V. Sergeenkov, *Izv. Akad. Nauk. SSSR, Ser. Fiz.* 30(1966)1185 / *Bull. Acad. Sci. USSR, Phys. Ser.* 30(1967)1237 [δ].
- 1968Go25 K. P. Gopinathan, W. Rubinson, *Bull. Am. Phys. Soc.* 13(1968)1452 [Q].
- 1968La10 F. Lagoutine, Y. Le Gallic, J. Legrand, *Int. J. Appl. Radiat. Isotop.* 19(1968)475 [Half-life].
- 1969Ca01 W. R. Casey, R. G. Albridge, *Z. Physik* 219(1969)216 [δ].
- 1969Ka08 E. Karttunen, H. U. Freund, R. W. Fink, *Nucl. Phys. A*131(1969)343 [P_γ].
- 1970Ma51 A. Marelius, K. G. Valivaara, Z. Awwad, J. Lindskog, J. Phil, S. -E. Hagglund, *Phys. Scr.* 1(1970)91 [α].
- 1972Em01 J. F. Emery, S. A. Reynolds, E. I. Wyatt, *Nucl. Sci. Eng.* 48(1972)319[Half-life].
- 1973Ra10 S. Raman, H. J. Kim, T. A. Walkiewicz, M. J. Martin, *Phys. Lett.* 44B(1973)255 [lg ft systematics].
- 1974Pl03 J. Plch, J. Zderadicka, *Czech. J. Phys.* B24(1974)1311 [P_{EC} , P_{XK}].
- 1974To04 F. Tolea, K. R. Baker, W. C. Smidth-Ott, R. W. Fink, *Z. Physik.* 268(1974)289 [P_{EC} , P_{XK}].
- 1977Kr13 K. S. Krane, *At. Data Nucl. Data Tables* 19(1977)363 [mixing ratio].
- 1979Kü** W. Künding, P. E. Müller, *Helv. Phys. Acta* 52(1979)555 [Half-life].
- 1979CoZG N. F. Coursol, Report CEA-R-5052 (1980) [α_T].
- 1980Ho17 H. Houtermans, O. Milosevic, F. Reichel, *Int. J. Appl. Radiat. Isotop.* 31(1980)153 [Half-life].
- 1982Ba16 V. Brabec, M. Rysavy, O. Dragoun, M. Fiser, A. Kovalik, Cs. Ujhelyi, D. Berenyi, *Z. Physik.* A306(1982)347 [δ].
- 1982HoZJ D. D. Hoppes, *NBS – 626*(1982)85 [Half-life].
- 1983Ku** H. Kubo, *Med. Phys.* 10(1983)889 [Half-life].
- 1983De11 K. Debertin, W. Pessara, *Int. J. Appl. Radiat. Isotop.* 34(1983)515 [P_γ].
- 1985Ma** W. B. Mann (Chairman), NCRP Report 58(1985)368 [P_γ].
- 1986Bo46 M. J. G. Borge, A. de Rujula, P. G. Hansen, B. Jonson, G. Nyman, H. L. Ravn, K. Riisager, *Phys. Scr.* 34(1986)591 [Q].
- 1987Sc20 H. Schrader, K. F. Walz, *Appl. Rad. Isotop.* 38(1987)763 [Half-life].
- 1989Ko** A. A. Konstantinov, T. E. Sazonova, S. V. Sepman, A. V. Zanevsky, Program and Thesis, Proc. 39th Ann. Conf. Nucl. Spectrosc. Struct. At. Nuclei, Leningrad, (1989) [P_γ , P_{XK}].
- 1989Si19 B. R. S. Simpson, B. R. Meyer, *Appl. Rad. Isotop.* 40(1989)819 [Half-life].
- 1990De09 P. de Felice, P. Ientile, C. Zicari, *Nucl. Instrum. Meth. Phys. Res.* A286(1990)514 [Half-life].
- 1990Iw04 A. Iwahara, M. H. H. Marechal, C. J. da Silva, R. Poledna, *Nucl. Instrum. Meth. Phys. Res.* A286(1990)370 [P_γ]
- 1990Li14 Sr. Little Flower, B. R. S. Babu, P. Venkataramaniah, H. Sanjeevia, *Nuovo Cim.* 130A(1990)553
- 1990Wo03 M. J. Woods, S. E. M. Lucas, *Nucl. Instrum. Meth. Phys. Res.* A286(1990)517 [Half-life].
- 1991Al05 T. Altzitzoglou, *Appl. Rad. Isotop.* 42(1991)493 [Half-life].
- 1992ScZZ U. Schötzgig, H. Schrader, K. Debertin, Proc. Conf. on Nuclear Data for Science and Technology, Jülich (1992)562 [P_γ].
- 1994Hi04 M. M. Hindi, R. L. Kozub, S. J. Robinson, *Phys. Rev.* C49(1994)3289 [Q].
- 1995Ra32 G. Ratel, *Nucl. Instrum. Meth. Phys. Res.* A366(1995)183 [Half-life].
- 1995ScZY E. Schönfeld, PTB-6 33-95-2 (1995) [Electron capture probabilities].
- 1996Ka48 V. D. M. L. Kalyani, G. S. K. Murty, N. V. S. V. Prasad, M. V. S. Chandrasekhar Rao, G. Satyanarayana, D. L. Sastry, *Nuovo Cim.* 109A(1996)1129 [P_{EC}].
- 1996Sc06 E. Schönfeld, H. Janßen, *Nucl. Instrum. Meth. Phys. Res.* A369(1996)527 [Atomic data].
- 1999Ka26 J. Katakura, *Nucl. Data Sheets* 86(1999)955 [Spin, parity, level energy].
- 2002Un02 M. P. Unterweger, *Appl. Rad. Isotop.* 56(2002)125 [Half-life].
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, *Nucl. Phys.* A729(2003)129 [Q].
- 2006Da20 M. A. L. da Silva, R. Poledna, A. Iwahara, C. J. da Silva, J. U. Delgado, R. T. Lopes, *Appl. Rad. Isotop.* 64(2006)1440 [P_γ].
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr., *Nucl. Instrum. Meth. Phys. Res.* A589(2008)202 [Theoretical ICC].

¹²⁷Sb - Comments on evaluation of decay data
by A. L. Nichols

Evaluated: February - March 2012

Evaluation Procedure

Limitation of Relative Statistical Weight Method (LWM) was applied to average the measured decay data when appropriate.

Decay Scheme

A reasonably complex decay scheme was constructed from the gamma-ray studies of 1967Ra13 and 1967Ta05. An earlier study involved the use of low-resolution NaI(Tl) detectors (1962Uh01), and these data have been set aside from consideration in this particular evaluation. The gamma-ray emission probabilities were expressed in terms of the emission probability of the 685.09-keV gamma ray (100 %), and weighted mean data were derived as appropriate. ¹²⁷Sb undergoes beta decay to both ^{127m}Te (defined as level 2, with a half-life of 106.1 days) and the ground state of ¹²⁷Te (half-life of 9.35 hours) – under these circumstances, the recommended ¹²⁷Sb decay scheme is rightly defined as ending effectively with the population of these two particular nuclear levels, while the subsequent IT and beta decay of ^{127m}Te are not included in the data file.

Nuclear Data

¹²⁷Sb undergoes beta decay to various nuclear levels of ¹²⁷Te, populating both ^{127m}Te (beta branch of 16.8 (6) %) and the ¹²⁷Te ground state (beta branch of 83.2 (6) %), latter more specifically by gamma decay only. These two daughter states possess significant half-lives of 106.1 days (^{127m}Te) and 9.35 hours (¹²⁷Te), and therefore their recommended decay data have been assembled as separate files.

Half-life (¹²⁷Sb)

The recommended half-life has been determined from the measurements of Sleight and Sullivan (1950S117), Bosch and Munczek (1957Bo96), Dropesky and Orth (1962Dr01), Uhler *et al.* (1962Uh01), Hagebø (1967Ha27), Takemoto *et al.* (1967Ta05), and Panontin and Sugarman (1972Pa13). A value of 3.85 days was derived in terms of LWM, with the uncertainty increased to the lowest experimental value of ± 0.07 days.

Half-life measurements (¹²⁷Sb).

Reference	Half-life (days)
1939Ab02	3.3* (80 h)
1946Gr06	4.0* (95 h)
1950S117	3.9 ± 0.1 (93 ± 3 h)
1957Bo96	3.7 ± 0.1 (88 ± 2 h)
1962Dr01	3.89 ± 0.07 (93.4 ± 1.7 h)
1962Uh01	3.9 ± 0.1 (94 ± 2 h)
1967Ha27	3.80 ± 0.08 [†] (91.2 ± 0.3 h)
1967Ta05	3.75 ± 0.10
1972Pa13	3.91 ± 0.07 (93.8 ± 1.6 h)
Recommended value	3.85 ± 0.07 [‡]

* no uncertainty quoted – not included in LWM analysis.

[†] possible systematic error proposed by 1967Ha27 – author suggests that the measured half-life may be too long by as much as 2 %; other measurements indicate this possibility may not be the case, although the uncertainty has been adjusted from ± 0.013 to ± 0.08 days as a sensible precaution.

[‡] uncertainty increased from ± 0.03 to the lowest experimental value of ± 0.07 days.

Q values

Q^- to the ¹²⁷Te ground state of 1582 (5) keV was adopted from the evaluated tabulations of 2011AuZZ, which compares with a value of 1581 (5) keV from 2003Au03. A value of 88.23 (7) keV was adopted for the energy of the ^{127m}Te nuclear level (2011Ha31), and was used to derive Q^- to ^{127m}Te of 1494 (5) keV.

Gamma RaysEnergies

Although gamma-ray energies have been measured to good accuracy by 1967Ra13 and 1967Ta05, the determination of the nuclear-level energies of ¹²⁷Te from a combination of decay data and the emissions from seven nuclear reactions are judged to be more robust (2011Ha31). Therefore, gamma transition energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies of 2011Ha31 were adopted, and used to determine the energies of the gamma-ray transitions between the depopulating-populating levels.

Adopted energies, spins and parities of the nuclear levels of ¹²⁷Te.

Nuclear level number	Nuclear level energy (keV)	Spin and parity
0	0.0	3/2 +
1	61.161 ± 0.019	1/2 +
2	88.23 ± 0.07 (^{127m} Te)	11/2 -
3	340.87 ± 0.06	(9/2 -)
4	473.26 ± 0.04	5/2 +
5	501.928 ± 0.010	3/2 +
6	631.40 ± 0.06	7/2 -
7	685.09 ± 0.07	7/2 +
8	762.64 ± 0.05	3/2 +
9	782.62 ± 0.03	5/2 +
10	786.13 ± 0.06	7/2 -
11	924.02 ± 0.18	7/2 +
12	1077.13 ± 0.17	5/2, 7/2, 9/2
13	1140.20 ± 0.07	5/2 +
14	1154.70 ± 0.09	5/2 +
15	1206 ± 5*	3/2 +, 5/2 +
16	1289.79 ± 0.08	5/2 +
17	1309.25 ± 0.07	3/2 +, 5/2+
18	1323.4 ± 0.8	
19	1378.58 ± 0.07	5/2 +

* Value of (1206.3 ± 0.7) keV adopted from the energy of the proposed depopulating γ transition (423.7 (7) keV) to the 782.62 (3)-keV nuclear level.

A number of gamma rays with very low emission probabilities have been observed only by 1967Ta05, and four of these particular transitions cannot be placed in the proposed decay scheme: 74.6-, 405.0-, 675.6- and 789.5-keV gamma rays. After inspection of the spectra (1967Ta05), these particular gamma rays have not been included in the recommended set of decay data.

Gamma-ray energies.

transition	E_{γ} (keV)		
	1967Ra13	1967Ta05	Recommended
$\gamma_{1.0}$ (Te)	61.0 (3)	61.1 (1)	61.16 (2)
–	–	74.6 (7)	unplaced/rejected
$\gamma_{10.6}$ (Te)	154.3 (5)	154.4 (7)	154.7 (1)
$\gamma_{3.2}$ (Te)	252.4 (3)	252.5 (5)	252.64 (9)
$\gamma_{9.5}$ (Te)	280.4 (5)	280.5 (10)	280.7 (1)
$\gamma_{6.3}$ (Te)	290.8 (5)	290.3 (15)	290.5 (1)
$\gamma_{11.6}$ (Te)	293.3 (9)	–	292.6 (2)
$\gamma_{9.4}$ (Te)	310.0 (7)	309.0 (10)	309.4 (1)
$\gamma_{12.7}$ (Te)	391.8 (5)	391.5 (7)	392.0 (2)
–	–	405.0 (10)	unplaced/rejected
$\gamma_{4.1}$ (Te)	412.1 (5)	411.6 (2)	412.10 (5)
$\gamma_{15.9}$ (Te)	–	423.7 (7)	423.7 (7)
$\gamma_{5.1}$ (Te)	441.0 (9)	440.7 (7)	440.77 (2)
$\gamma_{10.3}$ (Te)	445.1 (5)	444.9 (3)	445.3 (1)
$\gamma_{11.4}$ (Te)	451.0 (7)	451 (1)	450.8 (2)
$\gamma_{13.7}$ (Te)	–	456 (1)	455.1 (1)
$\gamma_{4.0}$ (Te)	473.0 (4)	473.0 (2)	473.26 (4)
$\gamma_{5.0}$ (Te)	502.8 (6)	501.5 (15)	501.93 (1)
$\gamma_{6.2}$ (Te)	543.3 (5)	543.0 (2)	543.2 (1)
$\gamma_{11.3}$ (Te)	584.2 (11)	–	583.2 (2)
$\gamma_{12.4}$ (Te)	603.5 (5)	603.6 (2)	603.9 (1)
$\gamma_{17.7}$ (Te)	–	624.0 (10)	624.2 (1)
$\gamma_{13.5}$ (Te)	637.8 (5)	638.5 (7)	638.3 (1)
$\gamma_{14.5}$ (Te)	652.3 (9)	653.5 (7)	652.8 (1)
$\gamma_{13.4}$ (Te)	667.5 (9)	666.9 (3)	666.9 (1)
–	–	675.6 (5)	unplaced/rejected
$\gamma_{14.4}$ (Te)	682.3 (10)	–	681.4 (1)
$\gamma_{7.0}$ (Te)	685.7 (5)	685.2 (3)	685.09 (7)
$\gamma_{10.2}$ (Te)	698.5 (5)	698.5 (3)	697.9 (1)
$\gamma_{9.1}$ (Te)	722.2 (5)	723.4 (7)	721.5 (1)
$\gamma_{19.6}$ (Te)	745.9 (5)	745.4(15)	747.2 (1)*
$\gamma_{8.0}$ (Te)	763.7 (8)	–	762.7 (1)
$\gamma_{9.0}$ (Te)	783.7 (5)	783.8 (3)	782.6 (1)
–	–	789.5 (15)	unplaced/rejected
$\gamma_{16.4}$ (Te)	817.0 (6)	817.3 (5)	816.5 (1)
$\gamma_{18.5}$ (Te)	820.6 (6)	820.1 (3)	821.5 (8)
$\gamma_{11.0}$ (Te)	924.4 (9)	923.5 (7)	924.0 (2)
$\gamma_{13.0}$ (Te)	1141.6 (8)	1141.2 (7)	1140.2 (1)
$\gamma_{14.0}$ (Te)	–	1155.2 (10)	1154.7 (1)
$\gamma_{16.0}$ (Te)	1290.3 (8)	1291.5 (15)	1289.8 (1)
$\gamma_{19.0}$ (Te)	1377.9 (9)	–	1378.6 (1)

* Significant adjustment has been made from 745.5 to 747.2 keV to give a recommended energy that can be satisfactorily placed in the proposed decay scheme.

Emission Probabilities

Although judged to be a rather limited data set, a reasonably consistent decay scheme was derived from the relative gamma-ray emission probabilities measured by Ragaini *et al.* (1967Ra13) and Takemoto *et al.* (1967Ta05). These relative emission probabilities were normalised to the 685.09-keV gamma ray (100 %).

Although the 61.16-keV gamma ray has been quantified by both 1967Ra13 and 1967Ta05, the assignment of this gamma transition in the decay scheme permits an accurate relative emission probability to be calculated from the gamma population-depopulation balance of the 61.161-keV nuclear level (with no populating beta transition). However, the observed gamma depopulation of the 762.64-keV nuclear level ($3/2^+$) is problematic, since no β^- or γ feeding of this level has been proposed – this unsatisfactory situation needs to be addressed in future experimental studies.

Nuclear-level studies by means of the ¹²⁶Te(n, γ) and (d,p) reactions have provided additional insight into the γ depopulation of many of the proposed ¹²⁷Te nuclear levels (2005Ho15). Thus, there is strong evidence for the depopulation of the 685.09-keV nuclear level by 212.2- and 183.7-keV gamma rays along with the main 685.09-keV gamma emission, but no detection of an equivalent 623.9-keV M3 gamma transition. The measurements of 2005Ho15 have been used to support the placing of gamma transitions and introduction of nuclear-level assignments throughout the proposed decay scheme.

Relative gamma-ray emission probabilities.

transition	E_γ (keV)	P_γ^{rel}		
		1967Ra13	1967Ta05	Recommended*
$\gamma_{1.0}$ (Te)	61.16 (2)	3.9 (3)	3.22 (4)	3.22 (4) [†]
–	74.6 (7)	–	0.11 (7)	unplaced/rejected
$\gamma_{10.6}$ (Te)	154.7 (1)	0.4 (2)	0.32 (7)	0.33 (7)
$\gamma_{3.2}$ (Te)	252.64 (9)	23.1 (9)	23.5 (4)	23.4 (4)
$\gamma_{9.5}$ (Te)	280.7 (1)	1.8 (4)	1.5 (1)	1.5 (1)
$\gamma_{6.3}$ (Te)	290.5 (1)	5.5 (3)	5.1 (2)	5.2 (2)
$\gamma_{11.6}$ (Te)	292.6 (2)	0.8 (4)	–	0.8 (4)
$\gamma_{9.4}$ (Te)	309.4 (1)	0.7 (3)	0.57 (10)	0.58 (10)
$\gamma_{12.7}$ (Te)	392.0 (2)	2.6 (2)	2.7 (3)	2.6 (2)
–	405.0 (10)	–	0.32 (5)	unplaced/rejected
$\gamma_{4.1}$ (Te)	412.10 (5)	10.4 (11)	9.6 (5)	9.7 (5)
$\gamma_{15.9}$ (Te)	423.7 (7)	–	0.28 (10)	0.28 (10)
$\gamma_{5.1}$ (Te)	440.77 (2)	1.9 (9)	0.7 (3)	1.9 (9) [‡]
$\gamma_{10.3}$ (Te)	445.3 (1)	11.8 (3)	11.8 (5)	11.8 (3)
$\gamma_{11.4}$ (Te)	450.8 (2)	0.5 (2)	1.1 (6)	0.6 (2)
$\gamma_{13.7}$ (Te)	455.1 (1)	–	0.3 (2)	0.3 (2)
$\gamma_{4.0}$ (Te)	473.26 (4)	70.1 (19)	70.1 (32)	70.1 (19)
$\gamma_{5.0}$ (Te)	501.93 (1)	2.1 (7)	1.7 (3)	1.8 (3)
$\gamma_{6.2}$ (Te)	543.2 (1)	8.0 (12)	7.4 (3)	7.4 (3)
$\gamma_{11.3}$ (Te)	583.2 (2)	0.9 (5)	–	0.9 (5)
$\gamma_{12.4}$ (Te)	603.9 (2)	12.1 (3)	11.7 (3)	11.9 (3)
$\gamma_{17.7}$ (Te)	624.2 (1)	–	0.18 (6)	0.18 (6)
$\gamma_{13.5}$ (Te)	638.3 (1)	1.2 (4)	1.0 (1)	1.0 (1)
$\gamma_{14.5}$ (Te)	652.8 (1)	1.0 (2)	0.7 (1)	0.8 (1)
$\gamma_{13.4}$ (Te)	666.9 (1)	2.0 (2)	1.0 (1)	1.5 (5)

transition	E_γ (keV)	P_γ^{rel}		
		1967Ra13	1967Ta05	Recommended*
–	675.6 (5)	–	0.18 (9)	unplaced/rejected
$\gamma_{14.4}$ (Te)	681.4 (1)	1.5 (7)	–	1.5 (7)
$\gamma_{7.0}$ (Te)	685.09 (7)	100	100.0	100
$\gamma_{10.2}$ (Te)	697.9 (1)	9.9 (2)	9.0 (2)	9.5 (5)
$\gamma_{9.1}$ (Te)	721.5 (1)	5.1 (3)	4.9 (2)	5.0 (2)
$\gamma_{19.6}$ (Te)	747.2 (1)	0.4 (2)	0.3 (1)	0.3 (1)
$\gamma_{8.0}$ (Te)	762.7 (1)	0.2 (1)	–	0.2 (1)
$\gamma_{9.0}$ (Te)	782.6 (1)	41.1 (9)	42.4 (12)	41.6 (9)
–	789.5 (15)	–	0.23 (4)	unplaced/rejected
$\gamma_{16.4}$ (Te)	816.5 (1)	1.1 (5)	0.75 (8)	0.76 (8)
$\gamma_{18.5}$ (Te)	821.5 (8)	0.6 (3)	0.32 (6)	0.33 (6)
$\gamma_{11.0}$ (Te)	924.0 (2)	1.4 (2)	1.29 (7)	1.30 (7)
$\gamma_{13.0}$ (Te)	1140.2 (1)	1.0 (2)	1.1 (3)	1.0 (2)
$\gamma_{14.0}$ (Te)	1154.7 (1)	–	0.11 (6)	0.11 (6)
$\gamma_{16.0}$ (Te)	1289.8 (1)	1.0 (3)	0.97 (9)	0.97 (9)
$\gamma_{19.0}$ (Te)	1378.6 (1)	0.2 (1)	–	0.2 (1)

* LWM of the measurements of 1967Ra13 and 1967Ta05, with the uncertainty increased to the lowest measured value when necessary.

† adopted on the basis of the γ population-depopulation balance of the 61.161-keV nuclear level, with no β^- transition.

‡ adopted from 1967Ra13, and in agreement with γ population-depopulation balance of the 501.93-keV nuclear level, with no β^- transition.

Multipolarities, Internal Conversion Coefficients and Internal-Pair Formation Coefficients

The nuclear level scheme specified by Hashizume (2011Ha31) has been used to define the multipolarities of the gamma transitions on the basis of known spins and parities. A significant number of important gamma-ray transitions possess (M1 + E2) multipolarity, and somewhat disparate studies have been undertaken to determine many of their mixing ratios (1972Kr15, 1974So03, 1985De04). Furthermore, assessments have been made of a more limited number of these mixing ratios by Krane (1977Kr13). These various data have been assessed by the evaluator, and specific selections have been made as follows:

61.16-keV gamma ray, 80.6%M1 + 19.4%E2, as derived from consideration of γ population-depopulation of the 61.161-keV nuclear level; 154.7-keV gamma ray, 92%M1 + 8%E2; 252.64-keV gamma ray, 18%M1 + 82%E2; 280.7-keV gamma ray, 99.2%M1 + 0.8%E2; 290.5-keV gamma ray, 86%M1 + 14%E2; 292.6-keV gamma ray, 98.5%E1 + 1.5%M2; 309.4-keV gamma ray, 99%M1 + 1%E2; 392.0-keV gamma ray, 97.8%M1 + 2.2%E2; 440.77-keV gamma ray, 80%M1 + 20%E2; 445.3-keV gamma ray, 50%M1 + 50%E2; 450.8-keV gamma ray, 67%M1 + 33%E2; 473.26-keV gamma ray, 96%M1 + 4%E2; 501.93-keV gamma ray, 89.6%M1 + 10.4%E2; 603.9-keV gamma ray, 98%M1 + 2%E2; 638.3-keV gamma ray, 85%M1 + 15%E2; 652.8-keV gamma ray, 94.6%M1 + 5.4%E2; 782.6-keV gamma ray, 95.8%M1 + 4.2%E2; 1140.2-keV gamma ray, 98%M1 + 2%E2; and 1289.8-keV gamma ray, 99.96%M1 + 0.04%E2.

Additional (M1 + E2) gamma transitions were arbitrarily assigned a mixing ratio of 1.0 ± 0.5 (50%M1 + 50%E2) in this reasonably comprehensive exercise (423.7, 455.1, 624.2, 666.9, 681.4, 762.7 and 816.5 keV). The 583.2- and 747.2-keV gamma rays were identified as E1 transitions, while the 412.10-, 543.2-, 685.09-, 697.9-, 721.5- and 924.0-keV gamma rays were defined as E2 transitions. These data were used to determine recommended internal conversion coefficients from the frozen orbital approximation of Kibédi *et al.* (2008Ki07), based on the theoretical tabulations of Band *et al.* (2002Ba25, 2002Ra45). Internal-pair formation coefficients were calculated by means of the methodology described by Kibédi *et al.* (2008Ki07).

Gamma-ray emissions: measured mixing ratios of (M1 + E2) transitions and (E1 + M2) transition.

E_γ (keV)	δ				
	1972Kr15	1974So03	1977Kr13	1985De04	Recommended
M1 + E2 transitions					
154.7 (1)	–	–	–	0.34 ± 0.21 or $-2.30^{+0.81}_{-2.02}$	0.3 ± 0.2
252.64 (9)	-0.56 ± 0.10 or -1.53 ± 0.24	-1.61 ± 0.39	-1.55 ± 0.20	-0.31 ± 0.03 or -2.55 ± 0.20	$-2.1 \pm 0.5^*$
280.7 (1)	–	–	–	-0.09 ± 0.02 or 7.80 ± 1.20	-0.09 ± 0.02
290.5 (1)	$0.27^{+0.21}_{-0.13}$ or 6^{+68}_{-3}	1.87 ± 0.51	1.9 ± 0.5	0.40 ± 0.03	0.40 ± 0.03
309.4 (1)	–	–	–	0.10 ± 0.03 or -2.13 ± 0.30	0.10 ± 0.03
392.0 (2)	$0.55^{+0.51}_{-0.19}$ or $2.8^{+2.5}_{-1.5}$ if $J^\pi(1077) = 5/2^+$ -0.29 ± 0.14 or -2.1 ± 0.7 if $J^\pi(1077) = 9/2^+$	–	–	0.15 ± 0.02 -0.31 ± 0.02	0.15 ± 0.02
440.77 (2)	–	–	–	$0.51^{+0.38}_{-0.22}$ or $-18.30^{+14.30}_{-60}$	0.5 ± 0.3
445.3 (1)	0.4 ± 0.2	-3.14 ± 0.76	-1.0 ± 0.3	-1.16 ± 0.30	-1.0 ± 0.5
450.8 (2)	–	–	–	$0.65^{+0.76}_{-0.12}$ or $1.16^{+0.22}_{-0.63}$	0.7 ± 0.5
473.26 (4)	-0.29 ± 0.06 or -1.56 ± 0.19	–	-0.29 ± 0.06	-0.10 ± 0.01 or -2.50 ± 0.05	$-0.20 \pm 0.10^\dagger$
501.93 (1)	–	–	–	0.34 ± 0.08 or 1.50 ± 0.22 $0.34^{+0.90}_{-0.24}$ or $2.13^{+0.38}_{-0.90}$	0.34 ± 0.08
603.9 (1)	0.00 ± 0.07 or 1.65 ± 0.25 if $J^\pi(1077) = 5/2^+$ pure E2 if $J^\pi(1077) = 9/2^+$	–	–	0.14 ± 0.08 or -2.32 ± 0.50 if $J^\pi(1077) = 5/2^+$ 0.05 ± 0.08	0.14 ± 0.08
638.3 (1)	–	–	–	-0.42 ± 0.03 or -5.50 ± 0.84	-0.42 ± 0.03
652.8 (1)	–	–	–	0.24 ± 0.07 or $2.08^{+0.26}_{-0.43}$	0.24 ± 0.07
697.9 (1)	-0.21 ± 0.03 or $-3.3^{+0.7}_{-0.3}$	–	–	–	defined as 100% E2
782.6 (1)	0.21 ± 0.01 or -11.7 ± 0.9	–	0.21 ± 0.01	–	0.21 ± 0.01
1140.2 (1)	$-0.14^{+0.14}_{-0.11}$ or $-2.2^{+0.9}_{-0.6}$	–	–	–	-0.14 ± 0.12
1289.8 (1)	$0.02^{+0.07}_{-0.09}$ or $-3.6^{+1.6}_{-0.9}$	–	–	–	$0.02^{+0.07}_{-0.09}$
E1(+M2) transition					
292.6 (2)	–	–	–	0.12 ± 0.13	0.12 ± 0.13

* LWM of 1972Kr15, 1974So03 and 1985De04 measurements.

† LWM of 1972Kr15 and 1985De04 measurements.

Gamma-ray emissions: recommended energies, multiplicities, theoretical internal conversion coefficients (frozen orbital approximation), and internal-pair formation coefficients.

E_γ (keV)	Multiplicity	α_K	α_L	α_{M+}	α_{IPF}	α_{tot}
61.16 (2)	80.6%M1+19.4% E2 $\delta = 0.49$ (6) [†]	2.93 (12)	0.99 (14)	0.28 (4)	–	4.2 (2)
154.7 (1)	92%M1 + 8%E2 $\delta = 0.3$ (2)	0.182 (14)	0.026 (5)	0.006	–	0.214 (20)
252.64 (9)	(18%M1 + 82% E2) $\delta = -2.1$ (5)	0.0541 (12)	0.0090 (4)	0.0021	–	0.0652 (17)
280.7 (1)	99.2%M1 + 0.8%E2 $\delta = -0.09$ (2)	0.0351 (5)	0.00445 (7)	0.00115	–	0.0407 (6)
290.5 (1)	(86%M1 + 14%E2) $\delta = 0.40$ (3)	0.0326 (5)	0.00430 (7)	0.0010	–	0.0379 (6)
292.6 (2)	98.5%E1(+ 1.5%M2) $\delta = 0.12$ (13)	0.0103 (60)	0.00136 (16)	0.000326	–	0.012 (7)
309.4 (1)	99%M1 + 1%E2 $\delta = 0.10$ (3)	0.0273 (4)	0.00345 (5)	0.00085	–	0.0316 (5)
392.0 (2)	(97.8%M1 + 2.2%E2) $\delta = 0.15$ (2)	0.01490 (21)	0.00187 (3)	0.00045	–	0.01722 (25)
412.10 (5)	E2	0.01210 (17)	0.001775 (25)	0.000435	–	0.01431 (20)
423.7 (7)	50%M1 + 50%E2 $\delta = 1.0$ (5)	0.0117 (4)	0.00158 (4)	0.00042	–	0.0137 (4)
440.77 (2)	80%M1 + 20%E2 $\delta = 0.5$ (3)	0.0109 (3)	0.001395 (22)	0.000305	–	0.0126 (3)
445.3 (1)	50%M1 + 50%E2 $\delta = -1.0$ (5)	0.0102(4)	0.001369 (23)	0.000431	–	0.0120 (4)
450.8 (2)	67%M1 + 33%E2 $\delta = 0.7$ (5)	0.0101 (4)	0.001318 (20)	0.000382	–	0.0118 (4)
455.1 (1)	50%M1 + 50%E2 $\delta = 1.0$ (5)	0.0097 (4)	0.001287 (19)	0.000313	–	0.0113 (4)
473.26 (4)	96%M1 + 4%E2 $\delta = -0.20$ (10)	0.00928 (14)	0.001159 (17)	0.000281	–	0.01072 (16)
501.93 (1)	89.6%M1 + 10.4%E2 $\delta = 0.34$ (8)	0.00795 (13)	0.000997 (14)	0.000243	–	0.00919 (14)
543.2 (1)	E2	0.00553 (8)	0.000761 (11)	0.000189	–	0.00648 (9)
583.2 (2)	E1	0.001622 (23)	0.000196 (3)	0.000052	–	0.00187 (3)
603.9 (1)	(98%M1 + 2%E2) $\delta = 0.14$ (8)	0.00513 (8)	0.000634 (10)	0.000156	–	0.00592 (9)
624.2 (1)	(50%M1 + 50%E2) $\delta = 1.0$ (5)	0.0043 (3)	0.000550 (24)	0.000150	–	0.0050 (4)
638.3 (1)	85%M1 + 15%E2 $\delta = -0.42$ (3)	0.00438 (7)	0.000544 (8)	0.000136	–	0.00506 (8)
652.8 (1)	94.6%M1 + 5.4%E2 $\delta = 0.24$ (7)	0.00423 (7)	0.000522 (8)	0.000128	–	0.00488 (8)
666.9 (1)	50%M1 + 50%E2 $\delta = 1.0$ (5)	0.0037 (3)	0.000464 (23)	0.000036	–	0.0042 (3)
681.4 (1)	50%M1 + 50%E2 $\delta = 1.0$ (5)	0.00347 (25)	0.000440 (22)	0.00009	–	0.0040 (3)

E_γ (keV)	Multipolarity	α_K	α_L	α_{M+}	α_{IPF}	α_{tot}
685.09 (7)	E2	0.00303 (5)	0.000399 (6)	0.000091	–	0.00352 (5)
697.9 (1)	E2	0.00289 (4)	0.000380 (6)	0.000090	–	0.00336 (5)
721.5 (1)	E2	0.00266 (4)	0.000348 (5)	0.000082	–	0.00309 (5)
747.2 (1)	E1	0.000951 (14)	0.0001142 (16)	0.0000278	–	0.001093 (16)
762.7 (1)	50%M1 + 50%E2 $\delta = 1.0$ (5)	0.00265 (20)	0.000332 (19)	0.000078	–	0.00306 (22)
782.6 (1)	95.8%M1 + 4.2%E2 $\delta = 0.21$ (1)	0.00277 (4)	0.000339 (5)	0.000081	–	0.00319 (5)
816.5 (1)	50%M1 + 50%E2 $\delta = 1.0$ (5)	0.00225 (17)	0.000282 (17)	0.000068	–	0.00260 (19)
821.5 (8)	–	–	–	–	–	–
924.0 (2)	E2	0.001491 (21)	0.000189 (3)	0.000045	–	0.001725 (25)
1140.2 (1)	98%M1 + 2%E2 $\delta = -0.14$ (12)	0.001179 (20)	0.0001427 (23)	0.0000348	0.00000150 (2)	0.001358 (23)
1154.7 (1)	M1 + E2	–	–	–	–	–
1289.8 (1)	99.96%M1 + 0.04%E2 $\delta = 0.02^{+0.07}_{-0.09}$	0.000901 (13)	0.0001087 (16)	0.0000265	0.0000188 (3)	0.001055 (15)
1378.6 (1)	M1 + E2	–	–	–	–	–

† adopted on the basis of the required transition probability in order to achieve γ population-depopulation balance for the 61.161-keV nuclear level.

A normalisation factor of 0.354 (4) was calculated from the internal conversion coefficients and relative emission probabilities of the gamma-ray transitions populating the 88.23-keV metastable (^{127m}Te) and ground (^{127g}Te) states, summed in conjunction with direct beta feeding to the 88.23-keV metastable state of $(2.0 \pm 0.5) \%$, as measured by 1967Ta05 (direct beta feeding to the ^{127}Te ground state was assumed to be zero on the basis of spin and parity considerations):

$$\left[\sum_{88.23 \text{ keV}} P_{\gamma+ce}^{rel} + \sum_{0.0 \text{ keV}} P_{\gamma+ce}^{rel} \right] x F + \beta_{0,2} = 100 \%$$

$$[41.8 (7) + 235.25 (238)] F + 2.0 (5) \% = 100 \%$$

$$F = 98.0 (5) / 277.05 (248) = 0.354 \pm 0.004$$

Beta-particle Emissions

Energies and emission probabilities

Beta-particle energies were calculated from the structural detail of the proposed decay scheme. Nuclear-level energies adopted from Hashizume (2011Ha31) and a Q_{β^-} value of 1582 (5) keV from the evaluated tabulations of 2011AuZZ were used to determine the energies and uncertainties of the beta-particle transitions.

The emission probability of the highest-energy beta-particle was measured by 1967Ta05 to be $(2.0 \pm 0.5) \%$, and this value was adopted to calculate the normalization factor of the relative gamma-ray emission probabilities. Direct beta population of the 762.64-, 501.93-, 61.161- and 0.0-keV nuclear levels of ^{127}Te were defined as zero on the basis of spin and parity

considerations. All other beta-particle emission probabilities were calculated on the basis of achieving population-depopulation balances with the relevant relative gamma transition probabilities, as derived from the relative gamma-ray emission probabilities, internal conversion coefficients, and normalization factor of 0.354 ± 0.004 .

Beta-particle emission probabilities per 100 disintegrations of ^{127}Sb .

Transition	E_{β} (keV)	P_{β}	Transition type	$\log ft$
$\beta_{0,19}^-$	203 ± 5	0.18 ± 0.04	allowed	7.42 ± 0.11
$\beta_{0,18}^-$	259 ± 5	0.12 ± 0.02	[allowed]	7.93 ± 0.08
$\beta_{0,17}^-$	273 ± 5	0.06 ± 0.02	(allowed)	8.30 ± 0.15
$\beta_{0,16}^-$	292 ± 5	0.61 ± 0.04	allowed	7.39 ± 0.04
$\beta_{0,15}^-$	376 ± 5	0.10 ± 0.04	(allowed)	8.53 ± 0.18
$\beta_{0,14}^-$	427 ± 5	0.85 ± 0.25	allowed	7.79 ± 0.13
$\beta_{0,13}^-$	442 ± 5	1.35 ± 0.21	allowed	7.64 ± 0.07
$\beta_{0,12}^-$	505 ± 5	5.17 ± 0.14	(allowed)	7.251 ± 0.021
$\beta_{0,11}^-$	658 ± 5	1.27 ± 0.25	allowed	8.26 ± 0.09
$\beta_{0,10}^-$	796 ± 5	7.72 ± 0.21	1 st forbidden non-unique	7.766 ± 0.018
$\beta_{0,9}^-$	799 ± 5	17.2 ± 0.3	allowed	7.425 ± 0.015
$\beta_{0,7}^-$	897 ± 5	34.4 ± 0.4	allowed	7.304 ± 0.013
$\beta_{0,6}^-$	951 ± 5	4.00 ± 0.21	1 st forbidden non-unique	8.33 ± 0.03
$\beta_{0,4}^-$	1109 ± 5	22.6 ± 0.8	allowed	7.826 ± 0.019
$\beta_{0,3}^-$	1241 ± 5	2.4 ± 0.3	(1 st forbidden non-unique)	8.98 ± 0.06
$\beta_{0,2}^-$	1494 ± 5	2.0 ± 0.5	1 st forbidden unique	10.21 ± 0.11

Σ 100.03

The proposed decay scheme is heavily dependent upon the absolute emission probability of the highest-energy β^- decay to the 88.23-keV $^{127\text{m}}\text{Te}$ nuclear level, as measured by Takemoto et al. (1967Ta05) to be $(2.0 \pm 0.5) \%$. There are also a number of gaps and uncertainties concerning some of the gamma-ray emissions. Under such unsatisfactory circumstances, spectroscopic measurements of the absolute γ -ray emission probabilities would assist greatly in addressing these specific issues, and so provide the means of deriving an evaluated decay scheme with much greater confidence.

Branching Fractions

$^{127}\text{Sb}(\beta^-)^{127\text{m}}\text{Te}$: summation of the γ and $\beta_{0,2}^-$ transitions populating the 88.23-keV metastable state.

$$\text{BF}(^{127}\text{Sb}(\beta^-)^{127\text{m}}\text{Te}) =$$

$$\sum_{i=1}^M [P_{\gamma}^{\text{rel}}(697.9 \text{ keV})(1 + \alpha) + P_{\gamma}^{\text{rel}}(543.2 \text{ keV})(1 + \alpha) + P_{\gamma}^{\text{rel}}(252.64 \text{ keV})(1 + \alpha)] \times F + \beta_{0,2}$$

$$= \{41.8 (7) \times 0.354 (4)\} + 2.0 (5) \% = 14.8 (3) \% + 2.0 (5) \% = 16.8 (6) \% \quad [0.168 (6)]$$

$^{127}\text{Sb}(\beta^-)^{127}\text{Te}$: summation of the γ transitions populating the 0.0-keV ground state (no direct population of the ground state by β^- decay).

$$\text{BF}(^{127}\text{Sb}(\beta^-)^{127}\text{Te}) =$$

$$\sum_{i=1}^M [P_{\gamma}^{\text{rel}}(1 + \alpha)] \times F$$

$$= 235.25 (238) \times 0.354 (4) = 83.3 (13) \% [0.833 (13)]$$

But lower value of 83.2 % [0.832] and uncertainty of $\pm 0.6 \% [\pm 0.006]$ adopted to achieve a precise balance with the equivalent BF for $^{127}\text{Sb}(\beta^-)^{127\text{m}}\text{Te}$.
 $\text{BF}(^{127}\text{Sb}(\beta^-)^{127}\text{Te}) = 83.2 (6) \% [0.832 (6)]$.

Atomic Data

The X-ray and Auger electron data have been calculated using evaluated X-ray data (1999ScZX, 2003De44), gamma-ray data, and atomic data from 1977La19, 1996Sc06 and 1998ScZM. Both the X-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.01, 28 January 2003). This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

K and L X-ray emission probabilities per 100 disintegrations of ¹²⁷Sb.

			Energy (keV)	Photons per 100 disint.
XL		(Te)	3.335 – 4.829	0.462 (23)
	XL ₁	(Te)	3.335	0.0089 (4)
	XL _α	(Te)	3.759 – 3.770	0.235 (10)
	XL _η	(Te)	3.605	0.00355 (19)
	XL _β	(Te)	4.030 – 4.302	0.184 (7)
	XL _γ	(Te)	4.572 – 4.829	0.0248 (10)
XK _α	XK _{α2}	(Te)	27.2020 (2)	1.11 (4)
	XK _{α1}	(Te)	27.4726 (2)	2.06 (7)
XK' _{β1}	XK _{β3}	(Te)	30.9446 (3))
	XK _{β1}	(Te)	30.9960 (4)) 0.591 (21)
	XK _{β5}	(Te)	31.236)
XK' _{β2}	XK _{β2}	(Te)	31.7008 (5))
	XK _{β4}	(Te)	31.774) 0.128 (6)
	XKO _{2,3}	(Te)	31.812)

Electron energies were determined from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

Data Consistency

Q_{β^-} values of 1582 (5) and 1494 (5) keV have been adopted for the β^- decay of ¹²⁷Sb to the ground and metastable states of ¹²⁷Te, respectively, based on the atomic mass evaluation of Audi and Wang (2011AuZZ) and the nuclear-level energy of ^{127m}Te (2011Ha31). An effective Q-value derived from these data has been compared with the Q-value calculated by summing the contributions of the individual emissions to the ¹²⁷Sb beta-decay process (i.e. β^- , electron, γ , etc.):

$$\text{effective Q-value} = \sum (Q_i \times BF_i) = 1567 (13) \text{ keV}$$

$$\text{calculated Q-value} = \sum (E_i \times P_i) = 1560 (15) \text{ keV}$$

The percentage deviation from the effective Q-value is $(0.5 \pm 1.3) \%$, which supports the derivation of a consistent decay scheme.

References

- 1939Ab02 P. ABELSON, An investigation of the products of the disintegration of uranium by neutrons, *Phys. Rev.* 56 (1939) 1-9. [half-life]
- 1946Gr06 W.E. GRUMMITT, G. WILKINSON, Fission products of U^{235} , *Nature* 158 (1946) 163. [half-life]
- 1950S117 N.R. SLEIGHT, W.H. SULLIVAN, Characteristics of 93 h Sb^{127} , *Radiochemical studies: the fission products*, C.D. CORYELL, N. SUGARMAN (editors), NNS 9 (1950) 928-930, McGraw-Hill, New York, USA. [half-life, Q_{β}]
- 1957Bo96 H. BOSCH, H. MUNCZEK, New half-life in the family of antimony isotopes, *Phys. Rev.* 106 (1957) 983-985. [half-life, E_{γ} , P_{γ}]
- 1962Dr01 B.J. DROPECKY, C.J. ORTH, A summary of the decay of some fission product tin and antimony isotopes, *J. Inorg. Nucl. Chem.* 24 (1962) 1301-1316. [half-life, E_{γ}]
- 1962Uh01 J. UHLER, G.H. NEUMANN, O. MELIN, T. ALVÄGER, Mass number assignments and γ -ray spectra of some neutron rich Sn and Sb isotopes, *Ark. Fys.* 21 (1962) 35-48. [half-life, E_{γ} , P_{γ}]
- 1967Ha27 E. HAGEBØ, Yields and isomeric yield ratios of antimony isotopes from the interaction of 159 MeV to 18.2 GeV protons with uranium, *J. Inorg. Nucl. Chem.* 29 (1967) 2515-2532. [half-life]
- 1967Ra13 R.C. RAGAINI, G.E. GORDON, W.B. WALTERS, Decay scheme of 3.9 d ^{127}Sb , *Nucl. Phys.* A99 (1967) 547-576. [E_{γ} , P_{γ}]
- 1967Ta05 I. TAKEMOTO, T. IWASHITA, S. KAGEYAMA, The energy levels of Te^{127} , *J. Phys. Soc. Japan* 23 (1967) 153-157. [E_{γ} , P_{γ} , Q_{β} , $P_{\beta 0,2}$]
- 1972Kr15 K.S. KRANE, W.A. STEYERT, Nuclear orientation study of the decays of $^{126,127,128}Sb$, *Phys. Rev.* C6 (1972) 2268-2275. [spin, parity, δ]
- 1972Pa13 J.A. PANONTIN, N. SUGARMAN, Mass yield distribution and charge dispersion in 450 MeV proton fission of ^{238}U , *J. Inorg. Nucl. Chem.* 34 (1972) 1485-1502. [half-life]
- 1974So03 J.C. SOARES, P. HERZOG, H. HÜBEL, Lifetime and magnetic moment of the $9/2^{-}$ anomalous coupling state in ^{127}Te , *Nucl. Phys.* A224 (1974) 358-366. [δ]
- 1977Kr13 K.S. KRANE, E2-M1 multipole mixing ratios in odd-mass nuclei $59 \leq A \leq 149$, *At. Data Nucl. Data Tables* 19 (1977) 363-416. [δ]
- 1977La19 F.P. LARKINS, Semiempirical Auger-electron energies for elements $10 \leq Z \leq 100$, *At. Data Nucl. Data Tables* 20 (1977) 311-387 [Auger and conversion electron energies]
- 1985De04 M.O.M.D. DE SOUZA, R.N. SAXENA, Directional correlation measurements for gamma transitions in ^{127}Te , *Phys. Rev.* C31 (1985) 593-601. [δ]
- 1996Sc06 E. SCHÖNFELD, H. JANßEN, Evaluation of atomic shell data, *Nucl. Instrum. Meth. Phys. Res.* A369 (1996) 527-533. [X_K , X_L , Auger electrons]
- 1998ScZM E. SCHÖNFELD, G. RODLOFF, Tables of the energies of K-Auger electrons for elements with atomic numbers in the range from $Z = 11$ to $Z = 100$, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]

- 1999ScZX E. SCHÖNFELD, G. RODLOFF, Energies and relative emission probabilities of K X-rays for elements with atomic numbers in the range from $Z = 5$ to $Z = 100$, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2002Ba25 I.M. BAND, M.B. TRZHASKOVSKAYA, C.W. NESTOR, Jr., P.O. TIKKANEN, S. RAMAN, Dirac–Fock internal conversion coefficients, *At. Data Nucl. Data Tables* 81 (2002) 1-334. [ICC]
- 2002Ra45 S. RAMAN, C.W. NESTOR, Jr., A. ICHIHARA, M.B. TRZHASKOVSKAYA, How good are the internal conversion coefficients now? *Phys. Rev. C* 66 (2002) 044312, 1-23. [ICC]
- 2003Au03 G. AUDI, A.H. WAPSTRA, C. THIBAULT, The AME2003 atomic mass evaluation (II). Tables, graphs and references, *Nucl. Phys. A* 729 (2003) 337-676. [Q-value]
- 2003De44 R.D. DESLATTES, E.G. KESSLER, Jr., P. INDELICATO, L. DE BILLY, E. LINDROTH, J. ANTON, X-ray transition energies: new approach to a comprehensive evaluation, *Rev. Mod. Phys.* 75 (2003) 35-99. [E_X]
- 2005Ho15 J. HONZÁTKO, V. BONDARENKO, I. TOMANDL, T. VON EGIDY, H.-F. WIRTH, D. BUCURESCU, V. YU. PONOMAREV, N. MĂRGINEAN, R. HERTENBERGER, Y. EISERMANN, G. GRAW, L. RUBÁČEK, Nuclear structure of ¹²⁷Te studied with (n,γ) and (\vec{d} ,p) reactions and interpreted with IBFM and QPM, *Nucl. Phys. A* 756 (2005) 249-307. [nuclear levels, E_γ]
- 2008Ki07 T. KIBÉDI, T.W. BURROWS, M.B. TRZHASKOVSKAYA, P.M. DAVIDSON, C.W. NESTOR, Jr., Evaluation of theoretical conversion coefficients using BrIcc, *Nucl. Instrum. Methods Phys. Res. A* 589 (2008) 202-229. [ICC]
- 2011AuZZ G. AUDI, M. WANG, Atomic mass evaluation 2011, private communication, Centre de Spectrométrie Nucléaire et de Spectrométrie de Masse (CSNSM), Orsay, France, April 2011. [Q-value]
- 2011Ha31 A. HASHIZUME, Nuclear data sheets for $A = 127$, *Nucl. Data Sheets* 112 (2011) 1647-1831. [nuclear levels]

¹²⁷Te - Comments on evaluation of decay data

by A. L. Nichols

Evaluated: February - May 2012

Evaluation Procedure

Limitation of Relative Statistical Weight Method (LWM) and other analytical procedures were applied to average the measured decay data when appropriate.

Decay Scheme

A simple decay scheme was constructed primarily from the gamma-ray studies of 1970Ap02 and 1965Au01 in which Ge(Li) gamma-ray detectors were used. An earlier study involved the use of low-resolution NaI(Tl) detectors (1956Kn20), and these data have not been considered in this particular evaluation. The gamma-ray emission probabilities were expressed in terms of the emission probability of the 417.99-keV gamma ray (100 %), and weighted mean data were derived as appropriate.

Nuclear Data

¹²⁷Te undergoes beta decay to various nuclear levels of ¹²⁷I through five β^- and nine subsequent γ emissions.

Half-life (¹²⁷Te)

The recommended half-life has been determined from the measurements of Seaborg *et al.* (1940Se01), Knight *et al.* (1956Kn20), Majumdar and Chatterjee (1963Ma20), Qaim and Ejaz (1968Qa02), and Bormann *et al.* (1970Bo22). A value of 9.35 hours was derived in terms of LWM, with the uncertainty increased from ± 0.06 to the lowest measured value of ± 0.10 hour.

Half-life measurements (¹²⁷Te).

Reference	Half-life (hours)
1940Se01	9.3 \pm 0.5
1956Kn20	9.35 \pm 0.10
1963Ma20	9.36 \pm 0.20
1968Qa02	9.23 \pm 0.13
1970Bo22	9.48 \pm 0.13
Recommended value	9.35 \pm 0.10*

* uncertainty increased from ± 0.06 to the lowest measured value of ± 0.10 hour.

Q values

Q^- of 702 (4) keV was adopted from the evaluated tabulations of 2011AuZZ, which compares with an earlier value of 702 (3) keV from 2003Au03.

Gamma RaysEnergies

Gamma transition energies were deduced from the structural details of the proposed decay scheme. The ¹²⁷I nuclear-level energies of 2011Ha31 were adopted, and used to determine the energies of the gamma-ray transitions between the depopulating-populating levels. Many of the lower-energy nuclear levels recommended by 2011Ha31 are primarily based on accurate gamma-ray energy measurements of the equivalent EC decay of ¹²⁷Xe by 1977Ge10 (observed gamma-ray emissions at 57.61 (2), 145.252 (10), 172.132 (10), 202.860 (10) and 374.991 (12) keV).

Adopted energies, spins and parities for the nuclear levels of ¹²⁷I.

Nuclear level number	Nuclear level energy (keV)	Spin and parity	¹²⁷ Te radionuclidic decay
0	0.0	5/2 +	¹²⁷ Te and ^{127m} Te
1	57.608 ± 0.011	7/2 +	¹²⁷ Te and ^{127m} Te
2	202.860 ± 0.008	3/2 +	¹²⁷ Te
3	374.992 ± 0.009	1/2 +	¹²⁷ Te
4	417.99 ± 0.06	5/2 +	¹²⁷ Te
5	618.31 ± 0.13	3/2 +	¹²⁷ Te
6	628.69 ± 0.16	7/2 +	^{127m} Te
7	650.92 ± 0.08	9/2 (+)	^{127m} Te
8	716.50 ± 0.06	(11/2 +)	^{127m} Te

Gamma-ray energies identified with β⁻ decay of ¹²⁷Te.

Transition	E _γ (keV)			
	1956Kn20	1965Au01	1970Ap02	Recommended*
γ _{1,0} (I)	58.5 (1)	57.6 (5)	57.63 (8)	57.608 (11)
γ _{2,1} (I)	145 (2)	145 (5)	145.2 (1)	145.252 (14)
γ _{3,2} (I)	–	–	172.1 (5)	172.132 (12)
γ _{2,0} (I)	203 (3)	203 (1)	202.9 (1)	202.860 (8)
γ _{4,2} (I)	215 (4)	214 (1)	215.1 (1)	215.13 (6)
γ _{4,1} (I)	360 (4)	360.0 (5)	360.3 (1)	360.38 (6)
γ _{3,0} (I)	–	–	375.0 (4)	374.991 (9)
γ _{4,0} (I)	418 (2)	417.0 (5)	417.9 (1)	417.99 (6)
γ _{5,0} (I)	–	–	618.6 (3)	618.31 (13)

* nuclear level energies of 2011Ha31 were used to determine the recommended energies of the gamma-ray transitions – gamma recoil of negligible impact on these data.

Emission Probabilities

Although judged to be a rather limited data set, a reasonably consistent decay scheme was derived from the relative gamma-ray emission probabilities measured by Auble and Kelly (1965Au01) and Apt *et al.* (1970Ap02) for a mixture of ¹²⁷Te and ^{127m}Te in secular equilibrium. These relative emission probabilities were normalised to the 100 % value assigned to the 417.99-keV gamma ray.

The 57.608-keV gamma-ray emission is common to both ¹²⁷Te and ^{127m}Te and has only been quantified by 1965Au01 and 1970Ap02 in terms of ¹²⁷Te-^{127m}Te mixture in secular equilibrium. However, the assignment of this gamma transition in the decay scheme of ¹²⁷Te permits an accurate relative emission probability to be calculated from the gamma population-depopulation balance of the 57.608-keV nuclear level, assuming no direct beta population of this particular 7/2⁺ level (3/2⁺ → 7/2⁺ would constitute a second forbidden non-unique transition):

$$TP_{\gamma}(57.608 \text{ keV}) = TP_{\gamma}(145.252 \text{ keV}) + TP_{\gamma}(360.38 \text{ keV}) = 0.59 (9) F + 13.9 (2) F \\ = 14.49 (22) F,$$

where TP is the transition probability of the relevant gamma ray, and

F is the normalisation factor for the relative γ -ray emission probabilities.

Thus, the relative γ -ray emission probability $P_{\gamma}^{rel}(57.608 \text{ keV})$ can be expressed as follows:

$$P_{\gamma}^{rel}(57.608 \text{ keV}) = \frac{TP_{\gamma}^{rel}(57.608 \text{ keV})}{(1 + \alpha_{tot})} = \frac{14.49 (22)}{(1 + 3.72 (6))} = 3.07 (6)$$

Relative gamma-ray emission probabilities for ¹²⁷Te, as adopted from measurements of a mixture of ¹²⁷Te and ^{127m}Te in secular equilibrium.

Transition	E _γ (keV)	P _γ ^{rel}		
		1965Au01	1970Ap02	Recommended*
γ _{1,0} (I) [†]	57.608 (11)	61 (1)	56 (5)	3.07 (6) [‡]
γ _{2,1} (I)	145.252 (14)	0.51 (6)	0.33 (3)	0.40 (6)
γ _{3,2} (I)	172.132 (12)	–	0.03 (2)	0.03 (2)
γ _{2,0} (I)	202.860 (8)	5.4 (2)	5.86 (21)	5.6 (2)
γ _{4,2} (I)	215.13 (6)	3.9 (2)	3.91 (17)	3.9 (2)
γ _{4,1} (I)	360.38 (6)	14.8 (1)	13.6 (1)	13.6 (2)
γ _{3,0} (I)	374.991 (9)	–	0.03 (2)	0.03 (2)
γ _{4,0} (I)	417.99 (6)	100	100	100
γ _{5,0} (I)	618.31 (13)	–	0.013 (2)	0.013 (2)

* weighted mean of appropriate measurements of 1965Au01 and 1970Ap02, from which NRM values were adopted.

[†] gamma transition common to the β⁻ decay of both ¹²⁷Te and ^{127m}Te.

[‡] derived from γ population-depopulation balance of the 57.608-keV nuclear level, assuming no direct population by β⁻ decay on the basis of spin-parity considerations (3/2⁺ → 7/2⁺).

Two specific numerical procedures were used to analyse the limited and somewhat disparate data set of P_γ^{rel} measurements of mixtures of ¹²⁷Te and ^{127m}Te in secular equilibrium: limitation of relative statistical weight method (LWM), and normalised residual method (NRM)

E _γ (keV)	Analytical method	P _γ ^{rel}	χ ² /(N-1)	χ ² /(N-1) _{critical}
57.608 (11)	LWM	61 (1)	0.96	6.63
	NRM	61 (1)	0.96	3.84
145.252 (14)	LWM	0.42 (9)	4.50	6.63
	NRM	0.40 (6)	2.63	3.84
202.860 (8)	LWM	5.6 (2)	2.52	6.63
	NRM	5.6 (2)	2.52	3.84
215.13 (6)	LWM	3.9 (2)	0.00	6.63
	NRM	3.9 (2)	0.00	3.84
360.38 (6)	LWM	14.2 (6)	72	6.63
	NRM	13.6 (2)	3.85	3.84

Multipolarities and Internal Conversion Coefficients

The nuclear level scheme specified by Hashizume (2011Ha31) has been used to define the multipolarities of the gamma transitions on the basis of known spins and parities. Many of the gamma-ray transitions possess (M1 + E2) multipolarity, and assessments have been made of a significant number of these mixing ratios by Krane (1977Kr13, 1980Kr22). Various proposed mixing ratios have been assessed by the evaluator, and specific selections have been made as follows:

- 57.608-keV gamma ray, 99.3 % M1 + 0.7 % E2;
- 172.132-keV gamma ray, 99.3 % M1 + 0.7 % E2;
- 202.860-keV gamma ray, 79 % M1 + 21 % E2;
- 215.13-keV gamma ray, 96.0 % M1 + 4.0 % E2;
- 360.38-keV gamma ray, 96.4 % M1 + 3.6 % E2; and
- 417.99-keV gamma ray, 99.4 % M1 + 0.6 % E2.

Additionally, the 618.31-keV (M1 + E2) gamma transition was arbitrarily assigned a mixing ratio of 1.0 ± 0.5 (50 % M1 + 50 % E2) in this reasonably comprehensive exercise. Both the 145.252- and 374.991-keV gamma rays were defined as E2 transitions. These data were used to determine recommended internal conversion coefficients from the frozen orbital approximation of Kibédi *et al.* (2008Ki07), based on the theoretical tabulations of Band *et al.* (2002Ba25, 2002Ra45).

Gamma-ray emissions: mixing ratios of (M1 + E2) transitions.

E_γ (keV)	δ				
	1965Au01	1967Ge10	1977Kr13	1980Kr22	Recommended
57.608 (11)	–	–0.084 (6) (M1 + 0.7(1)%E2)	–0.084 (6) (M1 + 0.7(1)%E2)	–0.083 (5) (M1 + 0.7(1)%E2)	–0.083 ± 0.005* (M1 + 0.7(1)%E2)
172.132 (12)	–	–	–0.084 (7) (M1 + 0.7(1)%E2)	–0.085 (6) (M1 + 0.7(1)%E2)	–0.085 ± 0.006* (M1 + 0.7(1)%E2)
202.860 (8)	–	+0.52 (5) (M1 + 21(3)%E2)	+0.52 (5) (M1 + 21(3)%E2)	–	+0.52 ± 0.05† (M1 + 21(3)%E2)
215.13 (6)	–0.20 (2) or > 200	–	–0.203 (15)	–	–0.203 ± 0.015† (M1 + 4.0(5)%E2)
360.38 (6)	0.18 (8) or 2.29 (7)	–	+0.194 (15)	–	+0.194 ± 0.015† (M1 + 3.6(5)%E2)
417.99 (6)	–	–	–0.08 (3)	–	–0.08 ± 0.03† (M1 + 0.6(3)%E2)
618.31 (13)	–	–	–	–	1.0 ± 0.5 (50%M1 + 50%E2)

* adopted directly from 1980Kr22.

† adopted directly from 1977Kr13.

Gamma-ray emissions: recommended energies, multiplicities, and theoretical internal conversion coefficients (frozen orbital approximation).

E_γ (keV)	Multiplicity	α_K	α_L	α_{M+}	α_{tot}	
57.608 (11)	99.3%M1 + 0.7%E2 $\delta = -0.083$ (5)	3.16 (5)	0.449 (8)	0.111	3.72 (6)	β^-
145.252 (14)	E2	0.357 (5)	0.0907 (13)	0.0233	0.471 (7)	β^-
172.132 (12)	99.3%M1 + 0.7%E2 $\delta = -0.085$ (6)	0.1419 (20)	0.0185 (3)	0.0046	0.1650 (24)	β^-
202.860 (8)	79%M1 + 21%E2 $\delta = +0.52$ (5)	0.0965 (17)	0.0142 (5)	0.0036	0.1143 (22)	β^-
215.13 (7)	96.0%M1 + 4.0%E2 $\delta = -0.203$ (15)	0.0782 (11)	0.01031 (16)	0.00249	0.0910 (13)	β^-
360.38 (7)	96.4%M1 + 3.6%E2 $\delta = +0.194$ (15)	0.0201 (3)	0.00256 (4)	0.00054	0.0232 (4)	β^-
374.991 (9)	E2	0.01671 (24)	0.00257 (4)	0.00062	0.0199 (3)	β^-
417.99 (6)	99.4%M1 + 0.6%E2 $\delta = -0.08$ (3)	0.01381 (20)	0.001741 (25)	0.000429	0.01598 (23)	β^-
618.31 (13)	50%M1 + 50%E2 $\delta = 1.0$ (5)	0.0047 (4)	0.00061 (3)	0.00019	0.0055 (4)	β^-

A normalisation factor of 0.009 97 (11) was calculated from the internal conversion coefficients and relative emission probabilities of the gamma-ray transitions populating the ground states of ¹²⁷Te and ¹²⁷I. An important feature of these calculations is the measurement of the ratio of the 417.99-keV γ -ray emission probability of ¹²⁷Te to the total β^- emission probability of ¹²⁷Te and ^{127m}Te in secular equilibrium by Apt *et al.* (1970Ap02), which has been adopted in the evaluation:

$$\frac{P_\gamma(417.99 \text{ keV})}{\sum(^{127}\text{Te} + ^{127m}\text{Te})\beta^-} = \frac{100 F}{[(122.33(43) + X) + 273.80(586)] F} = 0.0097(1),$$

where F is the normalisation factor for the relative γ -ray emission probabilities, and X is the relative emission probability of the β^- decay of ¹²⁷Te directly to the ground state of ¹²⁷I.

$$100 = 0.0097 (1) [396.13 (588) + X]$$

$$X = \frac{100}{0.0097(1)} - 396.13 (588) = 10309 (106) - 396.13 (588) = 9913 (106)$$

Therefore, within the β^- decay of ¹²⁷Te:

$$\sum(^{127}\text{Te})\beta^- = 9913(106)F + 122.33(43)F = 100 \%$$

$$F = 100 / 10035 (106) = 0.009 97 \pm 0.000 11$$

Beta-particle Emissions

Energies and emission probabilities

Beta-particle energies were determined from the structural detail of the proposed decay scheme. Nuclear-level energies adopted from Hashizume (2011Ha31) and a Q_{β^-} value of 702 (4) keV from the evaluated tabulations of 2011AuZZ were used to deduce the energies and uncertainties of the beta-particle transitions.

Absolute beta-particle emission probabilities were derived from γ population-depopulation of the various nuclear levels of ¹²⁷I, based on the relative emission probabilities of the γ rays, their normalisation factor of 0.009 97 (11), and the theoretical internal conversion coefficients. The $\beta_{0,0}^-$ emission directly to the ground state of ¹²⁷I can be derived by two routes:

- (i) relative β^- emission probabilities determined from γ population-depopulation of the nuclear levels of ¹²⁷I

$$P_{\beta_{0,0}^-} = 100 - \sum^{all\ other\ \beta^-} P_{\beta^-}^{rel} \times F$$

$$\begin{aligned} P_{\beta_{0,0}^-} &= 100 - [122.33 (43) \times 0.00997 (11)] = 100 - 1.220 (14) \\ &= (98.780 \pm 0.014) \% \end{aligned}$$

- (ii) relative γ -ray emission probabilities populating the ground state of ¹²⁷I directly

$$P_{\beta_{0,0}^-} = 100 - \sum^{\gamma\ to\ ground\ state} P_{\gamma}^{rel} (1 + \alpha_{tot}) \times F$$

$$\begin{aligned} P_{\beta_{0,0}^-} &= 100 - [0.013 (2) + 101.598 (23) + 0.03 (2) + 6.2 (2) + 14.5 (3)] \times 0.00997 (11) \\ &= 100 - [122.341 (361) \times 0.00997 (11)] = 100 - 1.220 (14) \\ &= (98.780 \pm 0.014) \% \end{aligned}$$

Beta-particle emission probabilities per 100 disintegrations of ¹²⁷Te.

Transition	E_{β} (keV)	P_{β}	Transition type	$\log ft$
$\beta_{0,5}^-$	84 ± 4	0.00013 ± 0.00002	allowed	8.38 ± 0.10
$\beta_{0,4}^-$	284 ± 4	1.19 ± 0.02	allowed	6.086 ± 0.022
$\beta_{0,3}^-$	327 ± 4	0.0006 ± 0.0003	allowed	9.58 ± 0.22
$\beta_{0,2}^-$	499 ± 4	0.025 ± 0.003	allowed	8.57 ± 0.06
$\beta_{0,0}^-$	702 ± 4	98.780 ± 0.014	allowed	5.490 ± 0.010

$$\sum 99.996 (25)$$

The proposed decay scheme is heavily dependent on the γ -ray studies of Apt *et al.* (1970Ap02), particularly their measurement of 0.0097 (1) for the $P_\gamma(417.99 \text{ keV})/\sum \beta^-$ ratio, and an estimate of 18.8 for the (¹²⁷Te + ^{127m}Te in secular equilibrium / ¹²⁷Te) ratio as applied to the 57.608-keV gamma-ray emission probability – current evaluation generates latter ratio of 18.9 (58/3.07). There is a lack of γ -ray spectroscopy measurements of ¹²⁷Te (and ^{127m}Te) decay with HPGe detectors that would assist greatly in quantifying the absolute γ -ray emission probabilities with much greater confidence, and hence derive a more satisfactory decay scheme.

Atomic Data

The X-ray and Auger electron data have been calculated from the evaluated X-ray data (1999ScZX, 2003De44), gamma-ray data, and atomic data from 1977La19, 1996Sc06 and 1998ScZM. Both the X-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.01, 28 January 2003). This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

K and L X-ray emission probabilities per 100 disintegrations of ¹²⁷Te.

			Energy (keV)	Photons per 100 disint.
XL		(I)	3.485 – 5.060	0.011 9 (6)
	XL ₁	(I)	3.485	0.000 226 (8)
	XL _{α}	(I)	3.927 – 3.938	0.005 97 (18)
	XL _{η}	(I)	3.779	0.000 088 (3)
	XL _{β}	(I)	4.221 – 4.508	0.004 76 (11)
	XL _{γ}	(I)	4.801 – 5.060	0.000 678 (17)
XK _{α}	XK _{α2}	(I)	28.3175 (4)	0.030 9 (7)
	XK _{α1}	(I)	28.6123 (3)	0.057 4 (12)
XK' _{β1}	XK _{β3}	(I)	32.2397 (3))
	XK _{β1}	(I)	32.2951 (4)) 0.016 5 (4)
	XK _{β5}	(I)	32.544)
XK' _{β2}	XK _{β2}	(I)	33.042 (2))
	XK _{β4}	(I)	33.120) 0.003 74 (12)
	XKO _{2,3}	(I)	33.166)

Electron energies were obtained from the electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

Data Consistency

A Q _{β^-} value of 702(4) keV has been adopted from the atomic mass evaluation of Audi and Wang *et al.* (2011AuZZ). This value has been compared with the Q-value calculated by summing the contributions of the individual emissions to the ¹²⁷Te beta-decay process (i.e. β^- , electron, γ , etc.):

$$\text{calculated Q-value} = \sum (E_i \times P_i) = 702 (4) \text{ keV}$$

Percentage deviation from the Q-value of Audi and Wang is (0.0 \pm 0.9) %, which supports the derivation of a highly consistent decay scheme.

References

- 1940Se01 G.T. SEABORG, J.J. LIVINGOOD, J.W. KENNEDY, Radioactive isotopes of tellurium, *Phys. Rev.* 57 (1940) 363-370. [half-life]
- 1956Kn20 J.D. KNIGHT, J.P. MIZE, J.W. STARNER, J.W. BARNES, Radiations of Te¹²⁷ and Te^{127m}, *Phys. Rev.* 102 (1956) 1592-1597. [half-life, E_γ, P_γ]
- 1963Ma20 N.K. MAJUMDAR, A. CHATTERJEE, 14.8 MeV neutron activation cross-section measurements of a few tellurium isotopes, *Nucl. Phys.* 41 (1963) 192-201. [half-life]
- 1965Au01 R.L. AUBLE, W.H. KELLY, A study of the excited states of ¹²⁷I populated in the decay of ¹²⁷Te and ^{127m}Te, *Nucl. Phys.* 73 (1965) 25-32. [E_γ, P_γ]
- 1966Ne02 J.F. NEESON, J.P. ROALSVIG, R.G. ARNS, Directional correlations in ¹²⁷I, *Can. J. Phys.* 44 (1966) 1313-1320. [γ-γ]
- 1967Ge10 J.S. GEIGER, Decay-scheme studies of ¹²⁵Xe and ¹²⁷Xe, *Phys. Rev.* 158 (1967) 1094-1104. [multipolarity, ICC]
- 1968Qa02 S.M. QAIM, M. EJAZ, Half-lives and activation cross-sections of some radioisotopes of iodine, tellurium and antimony formed in the interactions of iodine with 14.7 MeV neutrons, *J. Inorg. Nucl. Chem.* 30 (1968) 2577-2581. [half-life]
- 1970Ap02 K.E. APT, W.B. WALTERS, G.E. GORDON, Decay schemes of 109 d ^{127m}Te and 9.4 h ^{127g}Te, *Nucl. Phys.* A152 (1970) 344-353. [E_γ, P_γ]
- 1970Bo22 M. BORMANN, H.H. BISSEM, E. MAGIERA, R. WARNEMÜNDE, Total cross sections and isomeric cross-section ratios for (n,2n) reactions in the energy region 12-18 MeV, *Nucl. Phys.* A157 (1970) 481-496. [half-life]
- 1977Ge10 R.J. GEHRKE, R.G. HELMER, Absolute γ-ray intensities from ¹²⁷Xe decay, *Int. J. Appl. Radiat. Isot.* 28 (1977) 744-746. [E_γ]
- 1977Kr13 K.S. KRANE, E2-M1 multipole mixing ratios in odd-mass nuclei 59 ≤ A ≤ 149, *At. Data Nucl. Data Tables* 19 (1977) 363-416. [δ]
- 1977La19 F.P. LARKINS, Semiempirical Auger-electron energies for elements 10 ≤ Z ≤ 100, *At. Data Nucl. Data Tables* 20 (1977) 311-387 [Auger and conversion electron energies]
- 1980Kr22 K.S. KRANE, E2, M1 multipole mixing ratios, Supplement and corrections through December 1979, *At. Data Nucl. Data Tables* 25 (1980) 29-89. [δ]
- 1996Sc06 E. SCHÖNFELD, H. JANßEN, Evaluation of atomic shell data, *Nucl. Instrum. Meth. Phys. Res.* A369 (1996) 527-533. [X_K, X_L, Auger electrons]
- 1998ScZM E. SCHÖNFELD, G. RODLOFF, Tables of the energies of K-Auger electrons for elements with atomic numbers in the range from Z = 11 to Z = 100, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX E. SCHÖNFELD, G. RODLOFF, Energies and relative emission probabilities of K X-rays for elements with atomic numbers in the range from Z = 5 to Z = 100, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2002Ba25 I.M. BAND, M.B. TRZHASKOVSKAYA, C.W. NESTOR, Jr., P.O. TIKKANEN, S. RAMAN, Dirac-Fock internal conversion coefficients, *At. Data Nucl. Data Tables* 81 (2002) 1-334. [ICC]

- 2002Ra45 S. RAMAN, C.W. NESTOR, Jr., A. ICHIHARA, M.B. TRZHASKOVSKAYA, How good are the internal conversion coefficients now? Phys. Rev. C66 (2002) 044312, 1-23. [ICC]
- 2003Au03 G. AUDI, A.H. WAPSTRA, C. THIBAUT, The AME2003 atomic mass evaluation (II). Tables, graphs and references, Nucl. Phys. A729 (2003) 337-676. [Q-value]
- 2003De44 R.D. DESLATTES, E.G. KESSLER, Jr., P. INDELICATO, L. DE BILLY, E. LINDROTH, J. ANTON, X-ray transition energies: new approach to a comprehensive evaluation, Rev. Mod. Phys. 75 (2003) 35-99. [E_x]
- 2008Ki07 T. KIBÉDI, T.W. BURROWS, M.B. TRZHASKOVSKAYA, P.M. DAVIDSON, C.W. NESTOR, Jr., Evaluation of theoretical conversion coefficients using Br_{lcc}, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202-229. [ICC]
- 2011AuZZ G. AUDI, M. WANG, Atomic mass evaluation 2011, private communication, Centre de Spectrométrie Nucléaire et de Spectrométrie de Masse (CSNSM), Orsay, France, April 2011. [Q-value]
- 2011Ha31 A. HASHIZUME, Nuclear data sheets for A = 127, Nucl. Data Sheets 112 (2011) 1647-1831. [nuclear levels]

^{127m}Te - Comments on evaluation of decay data

by A. L. Nichols

Evaluated: February – May 2012

Evaluation Procedure

Limitation of Relative Statistical Weight Method (LWM) and other analytical procedures were applied to average the measured decay data when appropriate.

Decay Scheme

A simple decay scheme was constructed primarily from the gamma-ray studies of 1965Au01 and 1970Ap02 in which Ge(Li) gamma-ray detectors were used. An earlier study involved the use of low-resolution NaI(Tl) detectors (1956Kn20), and these data have not been considered in this particular evaluation. The relative emission probabilities of gamma rays emitted from mixtures of ¹²⁷Te and ^{127m}Te in secular equilibrium were quantified by 1965Au01 and 1970Ap02 in terms of the emission probability of the 417.99-keV gamma ray (100 %) in the β⁻ decay of ¹²⁷Te, and weighted mean data were derived as appropriate.

Nuclear Data

^{127m}Te undergoes IT decay directly to the ground state of ¹²⁷Te, with a small β⁻ branch to a number of nuclear levels of ¹²⁷I defined in terms of four β⁻ and five subsequent γ emissions.

Half-life (^{127m}Te)

The recommended half-life has been determined from the measurements of Seaborg *et al.* (1940Se01), Knight *et al.* (1956Kn20), Andersson *et al.* (1965An05), and Eastman and Krane (2008Ea01). A value of (106.1 ± 0.7) days was preferred as recommended by the Rajeval technique, rather than adopt the value determined by the LWM.

Half-life measurements (^{127m}Te).

Reference	Half-life (days)
1940Se01	92 ± 2
1951Co34	~ 115*
1956Kn20	105 ± 2
1965An05	109 ± 2
2008Ea01	106.1 ± 0.7
Recommended value	106.1 ± 0.7 [†]

* no uncertainty quoted, and therefore not included in the averaging procedures.

[†] Rajeval analysis adopted, in alignment with the measurement of Eastman and Krane.

Various procedures were considered in the analysis of the disparate data set: limitation of relative statistical weight method (LWM), normalised residual method (NRM), Rajeval technique, bootstrap method, and Mandel-Paule approach:

Analytical method	Half-life (days)	$\chi^2/(N-1)$	$\chi^2/(N-1)_{\text{critical}}$
LWM	104 ± 4	19.56	3.78
NRM	106.2 ± 0.6	2.47	2.60
Rajeval	106.1 ± 0.7	0.82	–
Bootstrap	104 ± 5	22.23	–
Mandel-Paule	103 ± 8	25.58	–

Q values

The nuclear-level energy of ^{127m}Te was adopted as Q_{IT} (88.23 (7) keV from 2011Ha31). A Q⁻ value of 790 (4) keV was obtained by summing the evaluated Q⁻ for the ground-state β⁻ decay taken from the tabulations of 2011AuZZ (702 (4) keV) with the nuclear-level energy for the metastable state of 88.23 (7) keV.

Gamma Rays

Energies

Gamma-ray transition energies were deduced from the structural details of the proposed decay scheme. The ¹²⁷Te and ¹²⁷I nuclear-level energies of 2011Ha31 were adopted, and used to determine the energies of the gamma-ray transitions between the depopulating-populating levels.

Adopted energies, spins and parities for the nuclear levels of ¹²⁷Te and ¹²⁷I.

Nuclear level number	Nuclear level energy (keV)	Spin and parity	¹²⁷ Te radionuclidic decay
¹²⁷Te nuclear level:			
0	0.0	3/2 +	^{127m} Te
1	61.161 ± 0.019	1/2 +	–
2	88.23 ± 0.07	11/2 –	^{127m} Te
¹²⁷I nuclear level:			
0	0.0	5/2 +	¹²⁷ Te and ^{127m} Te
1	57.608 ± 0.011	7/2 +	¹²⁷ Te and ^{127m} Te
2	202.860 ± 0.008	3/2 +	¹²⁷ Te
3	374.992 ± 0.009	1/2 +	¹²⁷ Te
4	417.99 ± 0.06	5/2 +	¹²⁷ Te
5	618.31 ± 0.13	3/2 +	¹²⁷ Te
6	628.69 ± 0.16	7/2 +	^{127m} Te
7	650.92 ± 0.08	9/2 (+)	^{127m} Te
8	716.50 ± 0.06	(11/2 +)	^{127m} Te

Gamma-ray energies identified with the IT and β⁻ decay modes of ^{127m}Te.

Transition	E _γ (keV)			
	1956Kn20	1965Au01	1970Ap02	Recommended*
γ _{1,0} (I)	58.5 (1)	57.6 (5)	57.63 (8)	57.608 (11)
γ _{2,0} (Te)	–	87 (1)	88.26 (8)	88.23 (7)
γ _{7,1} (I)	–	591 (1)	593.3 (1)	593.31 (8)
γ _{6,0} (I)	–	–	628.6 (3)	628.69 (16)
γ _{7,0} (I)	–	–	651.0 (2)	650.92 (8)
γ _{8,1} (I)	–	657 (1)	658.9 (1)	658.89 (6)

* nuclear level energies of 2011Ha31 were used to determine the recommended energies of the gamma-ray transitions – gamma recoil of negligible impact on these data.

Emission Probabilities

Although judged to be a rather limited data set, a reasonably consistent decay scheme was derived from the relative gamma-ray emission probabilities measured by Auble and Kelly (1965Au01) and Apt *et al.* (1970Ap02) for a mixture of ¹²⁷Te and ^{127m}Te in secular equilibrium. These relative emission probabilities were normalised to the 100 % value assigned to the 417.99-keV gamma ray, which is identified exclusively with the β⁻ decay of ¹²⁷Te.

Relative gamma-ray emission probabilities for ^{127m}Te, as adopted from measurements of a mixture of ¹²⁷Te and ^{127m}Te in secular equilibrium.

Transition	E _γ (keV)	P _γ ^{rel}		
		1965Au01	1970Ap02	Recommended [*]
γ _{1.0} (I) [†]	57.608 (11)	61 (1)	56 (5)	58 (1) [‡]
γ _{2.0} (Te)	88.23 (7)	25 (1)	12 (1)	8.56 (16) [#]
[γ _{4.0} (I)]	417.99 (6)	[100]	[100]	¹²⁷ Te decay only [100]
γ _{7.1} (I)	593.31 (8)	0.22 (4)	0.24 (2)	0.24 (2)
γ _{6.0} (I)	628.69 (16)	–	0.009 (2)	0.009 (2)
γ _{7.0} (I)	650.92 (8)	–	0.03 (1)	0.03 (1)
γ _{8.1} (I)	658.89 (6)	1.43 (6)	1.30 (10)	1.40 (6)

^{*} weighted mean of appropriate measurements of 1965Au01 and 1970Ap02 (identical LWM and NRM values).

[†] gamma transition common to the β⁻ decay of both ¹²⁷Te and ^{127m}Te.

[‡] determined from a weighted mean value of 61 (1) and subtraction of 3.07 (6) contribution from ¹²⁷Te β⁻ decay.

[#] calculated from IT branching fraction of 0.9727 (7), a normalisation factor of 0.00997 (11) for the relative γ-ray emission probabilities, and theoretical internal conversion coefficients of M4 88.23-keV γ transition.

Two specific numerical procedures were used to analysis the limited and somewhat disparate data set of P_γ^{rel} measurements of mixtures of ¹²⁷Te and ^{127m}Te in secular equilibrium: limitation of relative statistical weight method (LWM), and normalised residual method (NRM).

E _γ (keV)	Analytical method	P _γ ^{rel}	χ ² /(N-1)	χ ² /(N-1) _{critical}
57.608 (11)	LWM	61 (1)	0.96	6.63
	NRM	61 (1)	0.96	3.84
593.31 (8)	LWM	0.24 (2)	0.20	6.63
	NRM	0.24 (2)	0.20	3.84
658.89 (6)	LWM	1.40 (6)	1.24	6.63
	NRM	1.40 (6)	1.24	3.84

The 57.608-keV gamma-ray emission is common to both ¹²⁷Te and ^{127m}Te and has only been quantified by 1965Au01 and 1970Ap02 in terms of ¹²⁷Te-^{127m}Te mixture in secular equilibrium, with a relative emission probability of 61 (1). Nevertheless, an accurate relative emission probability of 3.07 (6) can be determined from the gamma population-depopulation balance of the 57.608-keV nuclear level in the β⁻ decay of ¹²⁷Te, based on the assumption of no direct beta transition to this particular 7/2⁺ level (3/2⁺ → 7/2⁺ would represent a second forbidden non-unique transition). Hence, an equivalent relative emission probability of 58 (1) can be calculated for the 57.608-keV gamma ray in the β⁻ decay of ^{127m}Te.

Multipolarities and Internal Conversion Coefficients

The nuclear level scheme specified by Hashizume (2011Ha31) has been used to define the multipolarities of the gamma transitions on the basis of known spins and parities. Detailed internal-conversion coefficient studies have been carried out by Kalinauskas *et al.* (1972Ka31, 1972Ka61) and Soni *et al.* (1977So06) on the 88.23-keV gamma ray that arises from the IT decay mode. These measurements of various ICC ratios and quantification of α_K are given below, and have been compared with equivalent theoretical internal conversion coefficients obtained from the frozen orbital approximation for M4 gamma transition (M4 BrIccFO).

Comparison of measured internal-conversion coefficient data with BrIcc frozen orbital calculations for the 88.23-keV M4 gamma transition.

	Ratios				α_K
	K : L : M : N+O	L _I : L _{II} : L _{III}	M _I : M _{II+III} : M _{IV+V}	(N+O) : L	
1972Ka31	0.99 (5) : 1 : 0.248 (24) : 0.050 (4)	0.599 (19) : 0.144 (8) : 1	1 : 2.29 (14) : 0.093 (23)	–	–
M4 BrIccFO	0.960 (21) : 1 : 0.238 (5) : 0.050 (1)	0.596 (14) : 0.137 (3) : 1	1 : 1.98 (4) : 0.0694 (15)	–	–
1972Ka61	–	–	–	0.050 (4)	–
M4 BrIccFO	–	–	–	0.050 (1)	–
1977So06	–	–	–	–	484 (23)
M4 BrIccFO	–	–	–	–	486 (7)

Mixing ratios for the 57.608- and 593.31-keV gamma transitions with (M1 + E2) multipolarity have been determined by 1965Au01 and 1967Ge10, and the data assessed by Krane (1977Kr13, 1980Kr22). Specific mixing ratios have been selected to give the following multiplicities: (99.3 % M1 + 0.7 % E2) for the 57.608-keV gamma ray, and (95 % M1 + 5 % E2) for the 593.31-keV gamma ray. An additional (M1 + E2) gamma transition of 628.69 keV was arbitrarily assigned a mixing ratio of 1.0 ± 0.5 (50 % M1 + 50 % E2). The 650.92- and 658.89-keV gamma rays were defined as E2 transitions. These data were used to determine recommended internal conversion coefficients from the frozen orbital approximation of Kibédi *et al.* (2008Ki07), based on the theoretical tabulations of Band *et al.* (2002Ba25, 2002Ra45).

Gamma-ray emissions from β^- decay: multiplicities and mixing ratios of (M1 + E2) transitions.

E_γ (keV)	δ				
	1965Au01	1967Ge10	1977Kr13	1980Kr22	Recommended
57.608 (11)	–	–0.084 (6) (M1 + 0.7(1)%E2)	–0.084 (6) (M1 + 0.7(1)%E2)	–0.083 (5) (M1 + 0.7(1)%E2)	$-0.083 \pm 0.005^*$ (M1 + 0.7(1)%E2)
593.31 (8)	–0.24 (13) or –5.68 (9)	–	–0.23 (3) (M1 + 5(1)%E2)	–0.23 (3) (M1 + 5(1)%E2)	$-0.23 \pm 0.03^*$ (M1 + 5(1)%E2)
628.69 (16)	–	–	–	–	1.0 ± 0.5 (50%M1 + 50%E2)
650.92 (8)	–	–	–	–	E2
658.89 (6)	–	–	–	–	E2

* Adopted directly from 1980Kr22.

Gamma-ray emissions: recommended energies, multiplicities, and theoretical internal conversion coefficients (frozen orbital approximation).

Transition	E_γ (keV)	Multiplicity	α_K	α_L	α_{M+}	α_{tot}	
$\gamma_{1,0}$ (I)	57.608 (11)	99.3%M1 + 0.7%E2 $\delta = -0.083$ (5)	3.16 (5)	0.449 (8)	0.111	3.72 (6)	β^-
$\gamma_{2,0}$ (Te)	88.23 (7)	M4	486 (7)	506 (8)	146 (3)	1138 (17)	IT
$\gamma_{7,1}$ (I)	593.31 (8)	95%M1 + 5%E2 $\delta = -0.23$ (3)	0.005 78 (9)	0.000 722 (11)	0.000 178	0.006 68 (10)	β^-
$\gamma_{6,0}$ (I)	628.69 (16)	50%M1 + 50%E2 $\delta = 1.0$ (5)	0.004 5 (4)	0.000 58 (3)	0.000 12	0.005 2 (4)	β^-
$\gamma_{7,0}$ (I)	650.92 (8)	E2	0.003 62 (5)	0.000 488 (7)	0.000 122	0.004 23 (6)	β^-
$\gamma_{8,1}$ (I)	658.89 (6)	E2	0.003 51 (5)	0.000 472 (7)	0.000 118	0.004 10 (6)	β^-

A normalisation factor of 0.009 97 (11) was calculated from the internal conversion coefficients and relative emission probabilities of the gamma-ray transitions populating the ground states of ¹²⁷Te and ¹²⁷I. An important feature of these calculations is the measurement of the ratio of the 417.99-keV γ -ray emission probability of ¹²⁷Te to the total β^- emission probability of ¹²⁷Te and ^{127m}Te in secular equilibrium by Apt *et al.* (1970Ap02), which was adopted in the evaluation:

$$\frac{P_{\gamma}(417.99 \text{ keV})}{\sum(^{127}\text{Te} + ^{127m}\text{Te})\beta^{-}} = \frac{100 F}{[(122.33 (43)+X) + 273.80 (586)] F} = 0.0097 (1),$$

where F is the normalisation factor for the relative γ -ray emission probabilities, and X is the relative emission probability of the β^{-} decay of ¹²⁷Te directly to the ground state of ¹²⁷I.

$$100 = 0.0097 (1) [396.13 (588) + X]$$

$$X = \frac{100}{0.0097 (1)} - 396.13 (588) = 10309 (106) - 396.13 (588) = 9913 (106)$$

Therefore, within the β^{-} decay of ¹²⁷Te:

$$\sum(^{127}\text{Te})\beta^{-} = 9913 (106) F + 122.33 (43) F = 100 \%$$

$$F = 100 / 10035 (106) = 0.009 97 \pm 0.000 11$$

Beta-particle Emissions

Energies and emission probabilities

Beta-particle energies were determined from the structural detail of the proposed decay scheme. Nuclear-level energies were adopted from Hashizume (2011Ha31), along with Q_{IT} and $Q_{\beta^{-}}$ values of 88.23 (7) and 702 (4) keV, respectively, from 2011Ha31 and the evaluated tabulations of 2011AuZZ, and used to deduce the energies and uncertainties of the beta-particle transitions.

Emission probabilities were derived from γ population-depopulation of the various nuclear levels of ¹²⁷I, based on the relative γ -ray emission probabilities, their normalisation factor of 0.009 97 (11), and the theoretical internal conversion coefficients. Direct beta decay to the ground state of ¹²⁷I would constitute a third forbidden non-unique transition ($11/2^{-} \rightarrow 5/2^{+}$), and has been assumed to be zero.

Beta-particle emission probabilities per 100 disintegrations of ^{127m}Te^m.

Transition	E_{β} (keV)	P_{β}	Transition type	log ft
$\beta_{2,8}^{-}$	74 ± 4	$0.014 1 \pm 0.000 6$	1 st forbidden non-unique	8.61 ± 0.08
$\beta_{2,7}^{-}$	139 ± 4	$0.002 7 \pm 0.000 2$	1 st forbidden non-unique	10.18 ± 0.05
$\beta_{2,6}^{-}$	161 ± 4	$0.000 09 \pm 0.000 02$	1 st forbidden unique	11.30 ± 0.11
$\beta_{2,1}^{-}$	732 ± 4	2.71 ± 0.07	1 st forbidden unique	9.873 ± 0.017

$$\sum 2.73 (7)$$

The proposed decay scheme is heavily dependent on the γ -ray studies of Apt *et al.* (1970Ap02), particularly their measurement of 0.0097 (1) for the $P_{\gamma}(417.99 \text{ keV})/\sum\beta^{-}$ ratio, and an estimate of 18.8 for the (¹²⁷Te + ^{127m}Te in secular equilibrium / ¹²⁷Te) ratio as applied to the 57.608-keV gamma-ray emission probability – current evaluation generates a latter value of 18.9 (58/3.07). There is a lack of γ -ray spectroscopy measurements of ^{127m}Te (and ¹²⁷Te) decay with HPGe detectors that would assist greatly in quantifying the absolute γ -ray emission probabilities with much greater confidence, and hence derive a more satisfactory decay scheme.

Branching Fractions and P_{γ}^{rel} (88.23-keV IT decay)

^{127m}Te(β^{-})¹²⁷I: summation of the β^{-} emissions deemed to populate specific nuclear levels of ¹²⁷I, based on the population-depopulation of the observed γ -ray emissions, their relative emission probabilities and associated normalisation factor of 0.009 97 (11), and the theoretical internal conversion coefficients. Direct beta decay to the ground state of ¹²⁷I has been assumed to be zero (spin and parity changes of $11/2^{-} \rightarrow 5/2^{+}$ would constitute a third forbidden non-unique transition).

$$\sum(^{127m}\text{Te})\beta^{-} = 273.80 (586) F$$

where F is the normalisation factor for the relative emission probabilities of the γ rays.

Thus, $BF(^{127m}\text{Te}(\beta^-)^{127}\text{I}) = 273.80 (586) \times 0.009 97 (11) = 2.73 (7) \% [0.0273 (7)]$

^{127m}Te(IT)¹²⁷Te: derived directly from ^{127m}Te(β⁻)¹²⁷I branching fraction.

$BF(^{127m}\text{Te}(\text{IT})^{127}\text{Te}) = 100 - 2.73 (7) = 97.27 (7) \% [0.9727 (7)]$

***P_γ^{rel}*(88.23-keV IT decay):**

IT branch = 97.27 (7) = $TP_{\gamma}^{abs}(88.23 \text{ keV})$

where $TP_{\gamma}^{abs}(88.23 \text{ keV})$ is the absolute transition probability of the 88.23-keV gamma emission.

$P_{\gamma}^{rel}(88.23 \text{ keV}) = \frac{TP_{\gamma}^{abs}(88.23 \text{ keV})}{(1 + \alpha_{tot}) * F} = \frac{97.27 (7)}{1139 (17) * 0.00997 (11)} = 8.56 (16)$

Atomic Data

The X-ray and Auger electron data have been calculated using evaluated X-ray data (1999ScZX, 2003De44), gamma-ray data, and atomic data from 1977La19, 1996Sc06 and 1998ScZM. Both the X-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.01, 28 January 2003). This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

K and L X-ray emission probabilities per 100 disintegrations of ^{127m}Te.

			Energy (keV)	Photons per 100 disint.
XL	(Te)		3.335 – 4.829	7.0 (3)
	L ₁	(Te)	3.335	0.146 (5)
	L _α	(Te)	3.759 – 3.770	3.86 (11)
	L _η	(Te)	3.605	0.036 9 (13)
	L _β	(Te)	4.030 – 4.302	2.45 (5)
	L _γ	(Te)	4.572 – 4.829	0.333 (8)
XK _α	XK _{α2}	(Te)	27.2020 (2)	10.3 (3)
	XK _{α1}	(Te)	27.4726 (2)	19.3 (5)
XK _{β1} '	XK _{β3}	(Te)	30.9446 (3)	5.51 (15)
	XK _{β1}	(Te)	30.9960 (4)	
	XK _{β5}	(Te)	31.236	
XK _{β2} '	XK _{β2}	(Te)	31.7008 (5)	1.20 (5)
	XK _{β4}	(Te)	31.774	
	XK _{O23}	(Te)	31.182	
XL	(I)		3.485 – 5.060	0.177 (9)
	XL ₁	(I)	3.485	0.003 36 (11)
	XL _α	(I)	3.927 – 3.938	0.089 (3)
	XL _η	(I)	3.779	0.001 30 (5)
	XL _β	(I)	4.221 – 4.508	0.070 7 (17)
	XL _γ	(I)	4.801 – 5.060	0.010 1 (3)
XK _α	XK _{α2}	(I)	28.3175 (4)	0.459 (12)
	XK _{α1}	(I)	28.6123 (3)	0.852 (21)
XK _{β1} '	XK _{β3}	(I)	32.2397 (3))

	XK _{β1}	(I)	32.2951 (4))	0.245 (7)
	XK _{β5}	(I)	32.544)	
XK' _{β2}	XK _{β2}	(I)	33.042 (2))	
	XK _{β4}	(I)	33.120)	0.055 5 (19)
	XKO _{2,3}	(I)	33.166)	

Electron energies were obtained from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

Data Consistency

Q-values of 88.23 (7) and 790 (4) keV have been adopted for the IT and β⁻ decay, respectively, based on the atomic mass evaluation of Audi and Wang (2011AuZZ) and the nuclear-level energy of ^{127m}Te (2011Ha31). An effective Q-value derived from these data has been compared with the Q-value calculated by summing the contributions of the individual emissions to the ¹²⁷Sb beta-decay process (i.e. β⁻, electron, γ, etc.):

$$\text{effective Q-value} = \sum (Q_i \times BF_i) = 107.4 (6) \text{ keV}$$

$$\text{calculated Q-value} = \sum (E_i \times P_i) = 106.1 (9) \text{ keV}$$

The percentage deviation from the effective Q-value is $(1.2 \pm 1.0) \%$, which indicates the derivation of a reasonably consistent decay scheme.

References

- 1940Se01 G.T. SEABORG, J.J. LIVINGOOD, J.W. KENNEDY, Radioactive isotopes of tellurium, Phys. Rev. 57 (1940) 363-370. [half-life]
- 1951Co34 J.M. CORK, A.E. STODDARD, C.E. BRANYAN, W.J. CHILDS, D.W. MARTIN, J.M. LEBLANC, Additional data on the radioactive isotopes of tin and tellurium, Phys. Rev. 84 (1951) 596-597. [half-life]
- 1956Kn20 J.D. KNIGHT, J.P. MIZE, J.W. STARNER, J.W. BARNES, Radiations of Te¹²⁷ and Te^{127m}, Phys. Rev. 102 (1956) 1592-1597. [half-life, E_γ, P_γ]
- 1965Au01 R.L. AUBLE, W.H. KELLY, A study of the excited states of ¹²⁷I populated in the decay of ¹²⁷Te and ^{127m}Te, Nucl. Phys. 73 (1965) 25-32. [E_γ, P_γ]
- 1965An05 G. ANDERSSON, G. RUDSTAM, G. SÖRENSEN, Decay data on some Xe, I and Te isotopes, Ark. Fys. 28 (1965) 37-43. [half-life]
- 1966Ne02 J.F. NEESON, J.P. ROALSVIG, R.G. ARNS, Directional correlations in ¹²⁷I, Can. J. Phys. 44 (1966) 1313-1320. [γ-γ, ICC]
- 1967Ge10 J.S. GEIGER, Decay-scheme studies of ¹²⁵Xe and ¹²⁷Xe, Phys. Rev. 158 (1967) 1094-1104. [multipolarity, ICC]
- 1970Ap02 K.E. APT, W.B. WALTERS, G.E. GORDON, Decay schemes of 109 d ^{127m}Te and 9.4 h ^{127g}Te, Nucl. Phys. A152 (1970) 344-353. [E_γ, P_γ]
- 1972Ka31 R.A. KALINAUSKAS, K.V. MAKARYUNAS, R.I. DAVIDONIS, Ratios of the internal conversion coefficients for M4 transitions in the nuclei Te^{121,123,125,127,129}, Sov J. Nucl. Phys. 15 (1972) 350-352. [M4 transition, ICC ratios]

- 1972Ka61 R.A. KALINAUSKAS, K.V. MAKARYUNAS, R.I. DAVIDONIS, Internal conversion in the N-shell for M4 transitions in ^{121,123,125,127,129}Te, Bull. Acad. Sci. USSR, Phys. Ser. 36 (1973) 2188-2189. [M4 transition, α_N]
- 1977Kr13 K.S. KRANE, E2-M1 multipole mixing ratios in odd-mass nuclei $59 \leq A \leq 149$, At. Data Nucl. Data Tables 19 (1977) 363-416. [δ]
- 1977La19 F.P. LARKINS, Semiempirical Auger-electron energies for elements $10 \leq Z \leq 100$, At. Data Nucl. Data Tables 20 (1977) 311-387 [Auger and conversion electron energies]
- 1977So06 S.K. SONI, A. KUMAR, S.L. GUPTA, S.C. PANCHOLI, Internal conversion coefficients of M4 transitions in ^{125m,127m,129m}Te decay, Z. Phys. A282 (1977) 49-53. [M4 transition, ICCs]
- 1980Kr22 K.S. KRANE, E2, M1 multipole mixing ratios, Supplement and corrections through December 1979, At. Data Nucl. Data Tables 25 (1980) 29-89. [δ]
- 1996Sc06 E. SCHÖNFELD, H. JANßEN, Evaluation of atomic shell data, Nucl. Instrum. Meth. Phys. Res. A369 (1996) 527-533. [X_K, X_L , Auger electrons]
- 1998ScZM E. SCHÖNFELD, G. RODLOFF, Tables of the energies of K-Auger electrons for elements with atomic numbers in the range from $Z = 11$ to $Z = 100$, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX E. SCHÖNFELD, G. RODLOFF, Energies and relative emission probabilities of K X-rays for elements with atomic numbers in the range from $Z = 5$ to $Z = 100$, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2002Ba25 I.M. BAND, M.B. TRZHASKOVSKAYA, C.W. NESTOR, Jr., P.O. TIKKANEN, S. RAMAN, Dirac-Fock internal conversion coefficients, At. Data Nucl. Data Tables 81 (2002) 1-334. [ICC]
- 2002Ra45 S. RAMAN, C.W. NESTOR, Jr., A. ICHIHARA, M.B. TRZHASKOVSKAYA, How good are the internal conversion coefficients now? Phys. Rev. C66 (2002) 044312, 1-23. [ICC]
- 2003Au03 G. AUDI, A.H. WAPSTRA, C. THIBAULT, The AME2003 atomic mass evaluation (II). Tables, graphs and references, Nucl. Phys. A729 (2003) 337-676. [Q-value]
- 2003De44 R.D. DESLATTES, E.G. KESSLER, Jr., P. INDELICATO, L. DE BILLY, E. LINDROTH, J. ANTON, X-ray transition energies: new approach to a comprehensive evaluation, Rev. Mod. Phys. 75 (2003) 35-99. [E_X]
- 2008Ea01 M.C. EASTMAN, K.S. KRANE, Neutron capture cross sections of even-mass tellurium isotopes, Phys. Rev. C77 (2008) 024303, 1-8. [half-life]
- 2008Ki07 T. KIBÉDI, T.W. BURROWS, M.B. TRZHASKOVSKAYA, P.M. DAVIDSON, C.W. NESTOR, Jr., Evaluation of theoretical conversion coefficients using BrIcc, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202-229. [ICC]
- 2011AuZZ G. AUDI, M. WANG, Atomic mass evaluation 2011, private communication, Centre de Spectrométrie Nucléaire et de Spectrométrie de Masse (CSNSM), Orsay, France, April 2011. [Q-value]
- 2011Ha31 A. HASHIZUME, Nuclear data sheets for $A = 127$, Nucl. Data Sheets 112 (2011) 1647-1831. [nuclear levels]

¹²⁷Xe - Comments on evaluation of decay data by X. Mougeot

This evaluation was completed in 2013, taking into account the available literature by December 2013. This radionuclide was first identified by 1940CR06 by using a (*p,n*) reaction on ¹²⁷I. The identification was confirmed later by using an (*α,n*) reaction on ¹²⁴Te (1950AN05).

1 Decay Scheme

The decay scheme is complete for this electron-capture decay. The ground state of ¹²⁷Xe is 1/2⁺ (1979AL14). From Q⁺ = 662,3 (20) keV (2012WA38), eight levels of ¹²⁷I are reachable (1970AP02): ground state 5/2⁺, 57 keV 7/2⁺, 203 keV 3/2⁺, 375 keV 1/2⁺, 418 keV 5/2⁺, 618 keV 3/2⁺, 629 keV 7/2⁺, 651 keV 9/2⁺. Notice that the parity of the latter level is uncertain (2011HA31).

Only three transitions lead to allowed electron-capture decays, namely to the 203 keV, 375 keV and 618 keV states. Up-to-now, no other transition has been observed. The following comment can be found in 1977GE10: “The electron-capture to the ground state is second forbidden for which the minimum log *ft* value is ~ 11 and thus the maximum intensity of the feeding branch is ~ 0,003 %”. The same comment can be applied for the electron-capture to the first excited state of ¹²⁷I since this transition is second forbidden unique. Identically, the 418 keV, 629 keV and 651 keV levels are not expected to be populated.

Consistency of the decay scheme was checked in Section 4.2.2. The decay scheme is consistent up to 99,9 (14) % confidence level. The effective Q value resulting from total energy balance considerations is: Q_{eff} = 662 (7) keV, consistent with Q⁺ = 662,3 (20) keV from 2012WA38 adopted in this evaluation.

2 Nuclear Data

The Q value adopted in this evaluation, Q⁺ = 662,3 (20) keV, is from Wang *et al.* 2012 (2012WA38).

2.1 ¹²⁷Xe Half-life

A full list of the half-life measurements available by December 2013 is summarized in Table 1. The most precise value comes from 2013UN, which is a corrected value of the NIST measurement from 2002UN02, originally being 36,3446 (28) d, due to a source positioning difficulty.

The value from 1954BA71 is 36,406 (16) d. It was extracted from the evolution of the abundance ratios ¹²⁷Xe/¹²⁶Xe and ¹²⁷Xe/¹²⁸Xe, assuming a constant ratio ¹²⁶Xe/¹²⁸Xe. But this latter ratio remained constant only within 0,5 %. The final uncertainty has thus been increased by 0,25 %, leading to: 36,41 (9) d.

The value from 1974CO05 is 36,3 (3) d, but after an average of their measured value with previous values and assuming a 4σ uncertainty. The measured value was preferred in this evaluation, namely 36,25 (9) d.

The LWEIGHT program excluded the results from 1940CR06, 1950AN05 and 1958FO48, which were found to be outliers by Chauvenet’s criterion. A weighted average with its external uncertainty was adopted from the resulting consistent data set, with a reduced-χ² value of 1,0. The main contribution comes from 2013UN, amounting to 58 %. This quite large contribution is kept because the value from 2013UN is the only one that comes from a metrological laboratory. The recommended value for the ¹²⁷Xe half-life is then:

$$T_{1/2}(^{127}\text{Xe}) = 36,358 (31) \text{ d.}$$

Note that the value from 1958FO48 agrees with other measurements within its uncertainty. A weighted average has been performed by the evaluator including this value. The result using the internal uncertainty is consistent with the recommended value: 36,357 (30) d.

Table 1: ¹²⁷Xe half-life measurements. The excluded value is crossed out.

Reference	T _{1/2} (¹²⁷ I) in d	Comments
4940CR06	34 (2)	Excluded by Chauvenet's criterion
4950AN05	32 (2)	Excluded by Chauvenet's criterion
1954BA71	36,41 (9)	
4958FO48	36,0 (5)	Excluded by Chauvenet's criterion
1964BR26	36,4 (1)	
1965WI12	36,5 (1)	
1974CO05	36,25 (9)	
2013UN	36,34 (4)	Revised NIST value
Adopted	36,358 (31)	

2.2 Half-lives of ¹²⁷I 57 keV, 203 keV and 375 keV states

The measured half-life values of the first excited state of ¹²⁷I, at 57 keV, are given in Table 2. The LWHEIGHT program excluded the result from 1964JH02, which was found to be an outlier by Chauvenet's criterion. A weighted average with its internal uncertainty was adopted with a reduced- χ^2 value of 0,2. The main contribution comes from 1976SA20, amounting to 96 %. The recommended value for the half-life of the first excited state of ¹²⁷I is:

$$T_{1/2}(\text{¹²⁷I, 57 keV}) = 1,95 (1) \text{ ns.}$$

Table 2: T_{1/2} half-life measurements of the 57 keV level of ¹²⁷I.

Reference	T _{1/2} in ns	Comments
1962TH12	2,0 (2)	
1964JH02	1,25 (21)	Excluded by Chauvenet's criterion
1967GE10	1,86 (11)	Same authors as in 1965GE04
1967SV01	1,93 (7)	
1969BEZH	1,90 (15)	
1976SA20	1,95 (1)	
Adopted	1,95 (1)	Consistent with the value adopted in 2011HA31

The measured half-life values of the second excited state of ¹²⁷I, at 203 keV, are given in Table 3. The value from 1965LA01 is cited in 1966GE13 but is not explicitly given in 1965LA01.

The LWHEIGHT program excluded the result from 1967SV01, which was found to be an outlier by Chauvenet's criterion. In the resulting consistent data set, a weighted average was used and the uncertainty of the value from 1967GE10 was expanded in order to take into account this value. The reduced- χ^2 value is 0,8. The two main contributions come from 1967GE10 and 1968KO01, amounting to 66 % and 32 % respectively. The recommended value for the half-life of the second excited state of ¹²⁷I is then:

$$T_{1/2}(\text{¹²⁷I, 203 keV}) = 0,387 (6) \text{ ns.}$$

Table 3: T_{1/2} half-life measurements of the 203 keV level of ¹²⁷I.

Reference	T _{1/2} in ns	Comments
1965LA01	0,39 (6)	Cited in 1966GE13 but not explicitly given in 1965LA01
1967GE10	0,381 (7)	Same authors as in 1966GE13
1967SV01	0,326 (21)	Excluded by Chauvenet's criterion
1968KO01	0,40 (1)	
1975AN19	0,39 (6)	
Adopted	0,387 (6)	Consistent with the value adopted in 2011HA31

The value from 1969LA08 of the half-life value of the third excited state of ¹²⁷I, at 375 keV, is: T_{1/2}(¹²⁷I, 375 keV) = 22 (4) ps. The adopted half-life value of the third excited state of ¹²⁷I, at 375 keV, is from 2011HA31: T_{1/2}(¹²⁷I, 375 keV) = 31 (8) ps.

2.3 Electron Capture Transition

From the absolute emission probabilities given in Section 4.2.2, together with the comment regarding the nature of the transition highlighted in Section 1, only the electron-capture transitions to the 203 keV, 375 keV and 618 keV states have to be considered. These decays are all allowed transitions, which can be calculated by the LOGFT program. The EC-capture program provides the $P_{K,L,M,N}$. Other values are from LOGFT, and the ones which can be deduced from them are in Table 4. A minimum relative uncertainty of 1 % has been taken by the evaluator.

Table 4: Electron-capture probabilities given by LOGFT and EC-capture for these allowed transitions.

	E (keV)	P_{ε} (%)	$\log ft$	P_K	P_L	P_M	P_{L2}/P_{L1}	P_{L1}	P_{L2}
$\gamma_{0,2}$	459,4 (20)	52,7 (14)	6,61	0,842 (8)	0,125 (1)	0,0272 (5)	0,0276 (3)	0,122 (1)	0,003 (1)
$\gamma_{0,3}$	287,3 (20)	47,3 (7)	6,21	0,830 (8)	0,134 (1)	0,0294 (6)	0,0276 (3)	0,130 (1)	0,004 (1)
$\gamma_{0,4}$	43,9 (21)	0,0142 (9)	7,42	0,31 (6)	0,523 (44)	0,137 (12)	0,0281 (3)	0,509 (43)	0,014 (43)

2.4 Gamma Transitions and Internal Conversion Coefficients

The evaluated energies of gamma transitions are from 2011HA31 (see Section 4.2.1).

2.4.1 Mixing ratios and multiplicities

The 57 keV and 203 keV transitions are the only ones which require a careful evaluation. The measurements of their mixing ratios or multiplicities are shown in Table 5 for the 57 keV transition, and in Table 6 for the 203 keV transition. Data published in the various publications are in bold text.

Table 5: Mixing ratio or multiplicity measurements for the 57 keV transition. Bold text stands for the original data given in the publications. Otherwise, the data were calculated by the evaluator.

Classification	Reference	$ \delta $	δ^2	Multiplicity	Adopted
M1 plus a possible admixture of E2	1964JH02	0,08 (25)	0,006 (39)	M1 + 0,6 (6) % E2	$\delta = 0,083 (5)$ $\delta^2 = 0,0069 (8)$ M1 + 0,68 (8) % E2
	1965LA01	0,08 (1)	0,0064 (16)	M1 + 0,64 (16) % E2	
	1966LE09	0,084 (6)	0,0071 (10)	M1 + 0,70 (10) % E2	
	1967GE10	0,08 (4)	0,007 (6)	M1 + 0,7 (1) % E2	

For the 57 keV transition, no outlier was found by the LWEIGHT program in this consistent data set. A weighted average with its internal uncertainty was adopted with a reduced- χ^2 value of 0,04. The main contribution comes from 1966LE09, amounting to 72 %. The recommended value is given in Table 5.

Table 6: Mixing ratio or multiplicity measurements for the 203 keV transition. Bold text stands for the original data given in the publications. Otherwise, the data were calculated by the evaluator.

Classification	Reference	$ \delta $	δ^2	Multiplicity	Adopted
M1 plus a possible admixture of E2	1966LE09	0,52 (5)	0,27 (5)	M1 + 21,3 (32) % E2	$\delta = 0,517 (27)$ $\delta^2 = 0,267 (28)$ M1 + 21,1 (17) % E2
	1967GE10	0,516 (32)	0,266 (33)	M1 + 21 (3) % E2	

For the 203 keV transition, no outlier was found by the LWEIGHT program in this consistent data set. A weighted average with its external uncertainty was adopted with a reduced- χ^2 value of 0,005. The main contribution comes from 1967GE10, amounting to 71 %. The recommended value is given in Table 6.

Considering the initial and final spins and parities, the 172 keV transition is classified as M1 plus a possible admixture of E2. 1967GE10 gives only an indication: M1 + (< 1%) E2. 1976LE23 provides two consistent values of the mixing ratios: $\delta = 0,088 (11)$ from the measurement of the 172 – 203 keV cascade and $\delta = 0,076 (45)$ from the 172 – 145 keV cascade. The recommended multiplicity of **the 172 keV transition** is the one from 2011HA31: $\delta = -0,085 (6)$, namely **M1 + 0,72 (10) % E2**.

For the 145 keV and 375 keV transitions, there is only an indication of their multiplicities given in

1967GE10: E2 + (< 25 %) M1 for the 145 keV transition, and E2 + (< 30 %) M1 for the 375 keV transition. Considering the initial and final spins and parities, both transitions are classified as E2 plus a possible admixture of M3. As recommended in 1967GE10 and consistently with the classification, the recommended multipolarity is **pure E2 for both the 145 keV and 375 keV transitions**.

Considering the initial and final spins and parities, the 618 keV transition is classified as M1 plus a possible admixture of E2. The adopted multipolarity of **the 618 keV transition** is the one from 2011HA31 with symmetrized uncertainty according to the method described in 2012WA38: $\delta = 0,083 (+16,-19)$ which becomes $\delta = 0,081 (18)$, namely **M1 + 0,65 (29) % E2**.

2.4.2 Internal conversion coefficients

The internal conversion coefficients measurements of the gamma transitions of ¹²⁷I were calculated with the BrIcc program (2008KI07) using the mixing ratios evaluated previously.

3 Atomic Data

3.1 Fluorescence yields

The fluorescence yields are taken from 1996SC06.

3.2 X Radiations and Auger electrons

The X-ray and Auger electron data were computed using the EMISSION program with the atomic data of Schönfeld and Janßen (1996SC06).

4 Gamma Emissions

4.2.1 Gamma-ray energies

The adopted energies of the ¹²⁷I levels are from 2011HA31. Independent evaluation, described below, was carried out by the evaluator. Results agree very well. The ones from 2011HA31 were preferred because of the completeness of their evaluation.

The gamma-ray energy measurements are given in Table 7. No value was excluded by the LWEIGHT program during the statistical process. All the data sets are consistent and a weighted average was used each time with the internal uncertainty. The adopted values are in red, and the reduced- χ^2 are also mentioned.

The energies of the gamma transitions were then deduced by adding the recoil energy of the nucleus. The proton and neutron masses, m_p and m_n , come from 2008MO18. The mass excess Δ of ¹²⁷I comes from 2012WA38. The largest recoil energy is for the 618 keV transition, with a value of 1,6 eV.

Table 7: E_γ measurements of the levels of ¹²⁷I. Final adopted values in red are from 2011HA31.

Reference	E_γ (keV)	E_γ (keV)	E_γ (keV)	E_γ (keV)	E_γ (keV)	E_γ (keV)
1958FO48	58,0 (10)	146,5 (5)	173,0 (5)	204,5 (5)	377,5 (5)	
1967GE10	57,60 (2)	145,22 (3)	172,10 (3)	202,84 (3)	374,96 (5)	
1968SC14			172,5 (3)			Evidence
1969GUZV		145,22 (20)	172,10 (20)	202,84 (20)	374,96 (20)	
1970AP02	57,63 (8)	142,2 (1)	172,1 (5)	202,9 (1)	375,0 (4)	618,6 (3)
1974CO05	57,6 (2)	145,2 (1)	172,1 (1)	202,8 (1)	375,0 (1)	
1977GE10	57,61 (2)	145,252 (10)	172,132 (10)	202,860 (10)	374,991 (12)	
Weighted average	57,606 (14)	145,248 (9)	172,100 (28)	202,858 (9)	374,989 (12)	618,6 (3)
χ^2	0,07	0,4	$2 \cdot 10^{-9}$	0,2	0,1	
2011HA31	57,61 (2)	145,252 (10)	172,132 (10)	202,860 (10)	374,991 (12)	618,41 (14)

4.2.2 Gamma-ray emission intensities

The measurements used to evaluate the gamma-ray emission intensities are summarized in the following tables. The data given in Table 8 from 1958FO48 are relative to the 203 keV gamma-ray intensity. This is the reference gamma-ray throughout this evaluation. The data crossed out were excluded by LWEIGHT according the Chauvenet's criterion during the statistical process.

The data from 1967GE10 given in Table 9 were recording from electron measurements. The use of internal conversion coefficients from BrIcc (see Section 2.4.2) allows calculating the corresponding relative gamma-ray emission intensities.

Data from 1974CO05 in Table 10 were given as absolute intensity measurements. The efficiency calibration was performed using standard calibrated sources, but no activity measurement of the ¹²⁷Xe source was given in the publication. We cannot therefore use the absolute intensity of the 203 keV transition to evaluate the other absolute gamma-ray intensities.

The data from 1977GE10 and from 2013RO are absolute intensity measurements and are given in Table 11 and Table 12. The efficiency calibration was performed using standard calibrated sources and an activity measurement of the ¹²⁷Xe source was given in each publication.

The relative gamma-ray intensities were evaluated with the LWEIGHT program and are given in Table 13. All the data sets are consistent and a weighted average was used each time. The type of uncertainty used and the reduced- χ^2 are also mentioned in Table 13.

Table 8: Relative data from 1958FO48 and their normalization to the 203 keV gamma-ray. Bold text stands for the original data given in the publications. Otherwise, the data were calculated by the evaluator. Data crossed out were excluded by LWEIGHT.

E (keV)	Relative I _γ	Relative I _γ	Comments
K X-rays	1170 (70)	117 (8)	Not used
57	21 (2)	2,10 (22)	
145	64 (10)	6,4 (10)	
172	342 (17)	34,2 (22)	Excluded by Chauvenet's criterion
203	1000 (40)	100	
375	301 (12)	30,1 (17)	Excluded by Chauvenet's criterion

Table 9: Relative electron data from 1967GE10. Bold text stands for the original data given in the publications. Otherwise, the data were calculated by the evaluator.

E (keV)	I _{ceK}	Multipol.	δ	α _K BrIcc	I _γ	Relative I _γ
57	0,632 (25)	M1+E2	-0,083 (5)	3,16 (5)	0,200 (9)	1,91 (9)
145	0,226 (4)	Pure E2		0,357 (5)	0,633 (14)	6,06 (17)
172	0,554 (10)	M1+E2	-0,085 (6)	0,1419 (20)	3,90 (9)	37,4 (10)
203	1	M1+E2	0,517 (27)	0,0964 (15)	10,45 (16)	100,0
375	0,0457 (10)	Pure E2		0,01671 (24)	2,73 (7)	26,2 (8)

Table 10: Absolute intensity measurements from 1974CO05, normalized to the 203 keV gamma-ray. Bold text stands for the original data given in the publications. Otherwise, the data were calculated by the evaluator.

E (keV)	Absolute I _γ	Relative I _γ
57	1,31 (8)	1,92 (12)
145	4,24 (21)	6,23 (33)
172	24,7 (10)	36,3 (16)
203	68,1 (13)	100,0
375	17,4 (10)	25,6 (15)

Table 11: Absolute intensity measurements from 1977GE10, normalized to the 203 keV gamma-ray. Bold text stands for the original data given in the publications. Otherwise, the data were calculated by the evaluator.

E (keV)	Absolute I _γ	Relative I _γ
57	1,22 (7)	1,79 (10)
145	4,29 (13)	6,28 (19)
172	25,5 (8)	37,3 (12)
203	68,3 (4)	100,0
375	17,2 (5)	25,2 (7)
618	0,014 (1)	0,0205 (15)

Table 12: Absolute and relative intensities measurements from 2013RO. Bold text stands for the original data given in the publications. Otherwise, the data were calculated by the evaluator.

E (keV)	Absolute I_γ	Relative I_γ	Comments
X K_α		83,4 (17)	Not used
X K_β		20,35 (42)	Not used
57	1,23 (7)	1,76 (10)	
145	4,33 (7)	6,20 (13)	
172	26,23 (44)	37,6 (8)	
203	69,8 (12)	100,0	
375	17,32 (28)	24,8 (5)	
618	0,0147 (20)	0,0211 (29)	

Table 13: Adopted relative gamma-ray emission intensities, the corresponding absolute emission intensities and the corresponding transition probabilities.

E (keV)	Relative I_γ	Comments	$I_\gamma \times 100$	Multipol.	δ	α_T BrIcc	$P_{\gamma+ce} \times 100$
57	1,858 (49)	External, $\chi^2 = 0,8$	1,272 (35)	M1+E2	-0,083 (5)	3,72 (6)	6,00 (18)
145	6,18 (9)	Internal, $\chi^2 = 0,22$	4,23 (7)	Pure E2		0,471 (7)	6,22 (11)
172	37,3 (5)	Internal, $\chi^2 = 0,18$	25,53 (38)	M1+E2	-0,085 (6)	0,1649 (24)	29,74 (45)
203	100		68,45 (45)	M1+E2	0,517 (27)	0,1142 (18)	76,3 (5)
375	25,21 (36)	Internal, $\chi^2 = 0,73$	17,26 (27)	Pure E2		0,0199 (3)	17,60 (28)
618	0,0206 (13)	Internal, $\chi^2 = 0,03$	0,0141 (9)	M1+E2	0,081 (18)	0,00609 (9)	0,0142 (9)

Only 1977GE10 and 2013RO can be conserved to evaluate the absolute gamma intensity of the 203 keV emission. The data are consistent. A weighted average using its internal uncertainty, with a reduced χ^2 of 1,4, gives: $I_\gamma(203 \text{ keV}) = 68,45 (45) \%$. The gamma transition probabilities are calculated from this value with the corresponding total internal conversion coefficients calculated with the BrIcc program (see Section 2.4.2) and given in Table 13.

Consistency of the decay scheme is tested summing all the gamma-rays falling to the ground state of ¹²⁷I. This sum is equal to: $\Sigma P_{\gamma+ce} = 99,9 (14) \%$. On the other hand, this sum can be set to 100 in order to determine the absolute gamma intensity $I_\gamma(203 \text{ keV})$:

$$\Sigma (1 + \alpha_T) \cdot I_{\gamma,rel.} = 145,9 (19),$$

$$\text{and as } I_\gamma(203 \text{ keV}) \cdot \Sigma (1 + \alpha_T) \cdot I_{\gamma,rel.} = 100,$$

$$\text{we have: } I_\gamma(203 \text{ keV}) = 68,5 (9) \%.$$

The best uncertainty associated with a consistent decay scheme is therefore obtained using the first method, which is used in this evaluation.

Consistency of the decay scheme is also tested from total energy balance considerations. The resulting effective Q value, $Q_{eff} = 662 (7) \text{ keV}$, is consistent with $Q^+ = 662,3 (20) \text{ keV}$ from 2012WA38 adopted in this evaluation.

From these absolute emission intensities, together with the comment regarding the nature of the transitions highlighted in Section 1, the following probabilities of electron-capture transitions were found: $P(\epsilon_{0,2}) = 52,7 (14) \%$, $P(\epsilon_{0,3}) = 47,3 (7) \%$, $P(\epsilon_{0,4}) = 0,0142 (9) \%$.

5. Electron Emissions

The energies of the conversion electrons were calculated from the gamma-transition energies given in Section 4.2.1 and the electron binding energies.

The emission probabilities of the conversion electrons were calculated using the internal conversion coefficients (see Section 2.4.2). The emission probabilities of K-Auger electrons were calculated using the transition probabilities given in Sections 2.3 and 2.4, the atomic data given in Section 3, and the internal conversion coefficients.

6. References

- 1940CR06 E.C. Creutz, L.A. Delsasso, R.B. Sutton, M.G. White, W.H. Barkas, Phys. Rev. 58, 481 (1940) [First measurement, $T_{1/2}$]
- 1950AN05 D.L. Anderson, M.L. Pool, Phys. Rev. 77, 142 (1950) [Identification, $T_{1/2}$]
- 1954BA71 S.J. Balestrini, Phys. Rev. 95, 1502 (1954) [$T_{1/2}$]
- 1958FO48 R.N. Forrest, H.T. Easterday, Phys. Rev. 112, 950 (1958) [$T_{1/2}$, E_γ , P_γ]
- 1962TH12 P. Thieberger, Arkiv Fysik 22, 127 (1962) [$T_{1/2}$]
- 1964BR26 M. Bresesti, F. Cappellani, A.M. Del Turco, Nucl. Phys. 58, 491 (1964) [$T_{1/2}$]
- 1964JH02 S. Jha, R. Leonard, Phys. Rev. 136, B1585 (1964) [$T_{1/2}$, Mixing ratios]
- 1965GE04 J.S. Geiger, R.L. Graham, I. Bergstrom, F. Brown, Nucl. Phys. 68, 352 (1965) [$T_{1/2}$]
- 1965LA01 H. Langhoff, Nucl. Phys. 63, 425 (1965) [Mixing ratios]
- 1965WI12 G. Winter, K. Hohmuth, J. Schintlmeister, Nucl. Phys. 73, 91 (1965) [$T_{1/2}$]
- 1966GE13 J.S. Geiger, R.L. Graham, Nucl. Phys. 89, 81 (1966) [$T_{1/2}$]
- 1966LE09 H.J. Leisi, Nucl. Phys. 76, 308 (1966) [Mixing ratios]
- 1967GE10 J.S. Geiger, Phys. Rev. 158, 1094 (1967) [Mixing ratios, E_γ , P_{cc}]
- 1967SV01 A.G. Svensson, R.W. Sommerfeldt, L.-O. Norlin, P.N. Tandon, Nucl. Phys. A95, 653 (1967) [$T_{1/2}$]
- 1968KO01 J. Kownacki, J. Ludziejewski, M. Moszynski, Nucl. Phys. A107, 476 (1968) [$T_{1/2}$]
- 1968SC14 W.-D. Schmidt-Ott, W. Weirauch, F. Smend, H. Langhoff, D. Gfoller, Z. Physik 217, 282 (1968) [J^π , E_γ]
- 1969BEZH R.B. Begzhanov, K.T. Salikhbaev, D. Gaffarov, Program and Theses, Proc. 19th Ann. Conf. Nucl. Spectrosc. Struct. At. Nuclei, Erevan, p.77 (1969) [$T_{1/2}$]
- 1969GUZV R. Gunnink, J.B. Niday, R.P. Anderson, R.A. Meyer, UCID-15439 (1969) [E_γ]
- 1969LA08 H. Langhoff, D. Gfoller, Nucl. Phys. A127, 379 (1969) [$T_{1/2}$]
- 1970AP02 K.E. Apt, W.B. Walters, G.E. Gordon, Nucl. Phys. A152, 344 (1970) [J^π , E_γ]
- 1974CO05 R. Collé, R. Kishore, Phys. Rev. C9, 981 (1974) [$T_{1/2}$, E_γ , P_γ]
- 1975AN19 D.S. Andreev, G.M. Gusinskii, K.I. Erokhina, V.S. Zvonov, A.A. Pasternak, Izv. Akad. Nauk SSSR, Ser. Fiz. 39, 2020 (1975); Bull. Acad. Sci. USSR, Phys. Ser. 39, No.10, 10 (1975) [$T_{1/2}$]
- 1976LE23 T. v. Ledebur, Helv. Phys. Acta 49, 661 (1976) [Mixing ratios]
- 1976SA20 S. Sanyal, A. Kumar, S.K. Soni, S.L. Gupta, S.C. Pancholi, Phys. Rev. C 14, 1285 (1976) [$T_{1/2}$]
- 1977GE10 R.J. Gehrke, R.G. Helmer, Int. J. Appl. Radiat. Isotop. 28, 744 (1977) [E_γ , P_γ]
- 1979AL14 W.P. Alford, R.E. Anderson, P.A. Batay-Csorba, R.A. Emigh, D.A. Lind, P.A. Smith, C.D. Zafiratos, Nucl. Phys. A 323, 339 (1979) [J^π]
- 1996SC06 E. Schönfeld, H. Janßen, Nucl. Instr. Meth. A369 (1996) 527 [ω_K , K x ray ratios, Auger e-ratios, atomic data]
- 2002UN02 M.P. Unterweger, Appl. Radiat. Isot. 56, 125 (2002) [$T_{1/2}$]
- 2008KI07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr. Nucl. Instr. and Meth. A589 (2008) 202-229 [BrICC]
- 2008MO18 P.J. Mohr, B.N. Taylor, D.B. Newell, Rev. Mod. Phys. 80, 633 (2008) [m_p , m_n]
- 2011HA31 A. Hashizume, Nucl. Data Sheets 112, 1647 (2011) [E_γ]
- 2012WA38 M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer, Chin. Phys. C 36, 1603 (2012) [Q-value]
- 2013RO M. Rodrigues, M.-C. Lépy, P. Cassette, X. Mougeot, M.-M. Bé, ICRM 2013 Conference, Appl. Radiat. Isot., in press (2013) [P_γ]
- 2013UN M.P. Unterweger, R. Fitzgerald, App. Radiat. Isot., in press (2013) [$T_{1/2}$]

¹²⁹I - Comments on evaluation of decay data by V. P. Chechev and V. O. Sergeev

1- Decay Scheme

The 2nd unique forbidden β^- -transition to the $1/2^+$ ground state of ¹²⁹Xe was not observed. In 1954 Der Matiosian and Wu (1954De17) showed experimentally that this β^- -branch intensity did not exceed 1 %. This limit gives a $\log f_{2ut} = 14.9$ (or $\log f_{0t} = 15.8$), which is consistent with the $\log f_{2ut}$ values of 14.6 – 15.8 tallied in 1998Si17 for ten cases from A=22 to A=138, excluding ¹⁰Be, with 13.8, and ²⁰⁹Po, with 14.36. The highest value of 15.8 corresponds to 0.13% for the transition considered.

Therefore, we have adopted the probability of the 2nd unique forbidden β^- -transition to the $1/2^+$ ground state of ¹²⁹Xe $P(\beta^-_{0,0}) = 0.05(5)\%$ with the uncertainty which provides the limits from 0 to 1% according to 1954De17.

2- Nuclear Data

The Q value has been computed on the basis of the spectrometric measurement of the $\beta^-_{0,1}$ energy by N. Coursol (1979CoZG) and the evaluated gamma-ray energy. This measurement gives a more accurate Q value than 194(3) keV, presented in the atomic mass evaluation (1995Au04).

The following four experimental values for the ¹²⁹I half-life are available (in units of 10⁷ years).

1.72(9)	1951 Ka16
1.56(6)	1957Ru65
1.57(4)	1972Em01
1.97(14)	1973Ku17

Use of the LRSW method leads to a higher uncertainty (0.047) in 1972Em01. Our recommended value has been obtained as the weighted mean with the external uncertainty 0.06 expanded due to the Student's factor (or MBAYS uncertainty) : 1.61(7). Thus our recommended value for the ¹²⁹I half-life is $1.61(7) \times 10^7$ years.

2.1. β^- -Transitions

The energy of the $\beta^-_{0,1}$ transition has been adopted from 1979CoZG (Coursol). For the probabilities $P(\beta^-_{0,1})$ and $P(\beta^-_{0,0})$ see discussion in sect.1.Decay Scheme.

2.2. Gamma-ray Transitions and Internal Conversion Coefficients

The correction for recoil has not changed the γ -ray transition energy.

The emission probability of the γ -ray transition (photons + electrons) has been adopted as 99.5(5)%. (see discussion in sect.1).

The multipolarity of the γ -ray transition was measured in 1965Ge04 (M1) and 1974Ra26 (M1 + 0.073(27)% E2).

ICC's have been interpolated from theoretical values of 1978Ro22 for the adopted multipolarity of $M1 + 0.07(3)\% E2$. The uncertainties in the theoretical values are as follows: 1% for α_K and 3% for α_L , α_M , α_{NO} . The ratio α_{NO}/α_M has been taken from 1971Dr11. The ICC interpolated from other tables (1968Ha53, 1969Ha61, 1978Band) agree with the adopted values within the limits of the stated uncertainties.

The interpolated value $\alpha_K^{\text{theory}} = 10.59(11)$ can be compared with the following experimental values: 10.6 (1968ReZY), 9.8(9) (1970Gy01), 10.2(4) (1970SaZI), 10.2(5) (1977Ra23), and 10.6(4) (1985Ba73), which have an unweighted average of 10.3.

3. Atomic Data

3.1. Fluorescence yields

The fluorescence yields have been taken from 1996Sc06 (Schönfeld and Janßen).

3.2. X rays

X-ray energies are based on the wavelengths given in the compilation of 1967Be65 (Bearden).

The relative K x-ray emission probabilities have been taken from 1996Sc06 and 1999Schönfeld.

3.3. Auger Electrons

The energies of Auger electrons are from 1977La19 (Larkins) and 1998Schönfeld.

The ratios $P(KLX)/P(KLL)$ and $P(KXY)/P(KLL)$ have been taken from 1996Sc06.

4. Electron emissions

The energies of the conversion electrons have been calculated from the γ -ray transition energy given in sect. 2.2 and the electron binding energies. Their absolute emission probabilities have been calculated using the conversion coefficients given in 2.2 and the absolute γ -ray emission probability.

For the L-shell the ratios $L_1:L_2:L_3 = 100:8.9(4):3.13(14)$ obtained from theoretical conversion coefficients can be compared with the experimental $L_1:L_2:L_3 = 100:10.0(4):3.1(3)$ from $^{129}\text{Cs} \rightarrow ^{129}\text{Xe}$ decay (1965Ge04).

Values of the emission probabilities of K-Auger electrons have been calculated using our recommended $P(\text{ceK})$ and $P(\text{ceL})$ values and atomic data given in 3.1.

The maximum energy of β^- particles with energy of 151 keV has been taken from 1979CoZG(Coursol). The average energy of β^- particles calculated with the LOGFT program, which uses an allowed spectral shape, is 40.6(3) keV. The SPEBETA program gives a different value of 37 keV (2001 Be). In 2001Be the shape factor $C(W) = q^2 + (0.10 \pm 0.01)p^2$ was used that given by E. der Matiosian and C. S. Wu (1953DE10) (measurement with a magnetic spectrometer). The value of 37 keV is supported also by the calculation of Kolobachkin et.al. (See the book "Beta emissions of fission products", authors: V. M. Kolobachkin, P. M. Rubtsov, V. G. Alexankin and P. A. Ruzhanskiy. - Moscow, Atomizdat, 1978, p.189. In Russian). They found 36 keV for the average energy of β^- particles of ^{129}I . So we adopt 37(1) keV as the recommended value.

5. Photon Emissions

5.1 X-Ray Emissions

Our recommended value for the total K x-ray absolute emission intensity has been calculated as $P_{\text{XK}}^{\text{eval.}} = \omega_{\text{K}}\alpha_{\text{K}}P_{\gamma}(39.6) = 69.8(11)\%$, based on the adopted value of ω_{K} , a theoretical value of α_{K} , and our recommended value of $P_{\gamma}(39.6) = 7.42(8)\%$. This K x-ray emission probability agrees well with the result of the measurement $P_{\text{XK}}^{\text{exp.}} = 70.2(8)\%$ in 1985Ba73, relative to $P_{\gamma}(39.6) = 7.46\%$ (or 69.8(8)%, relative to $P_{\gamma}(39.6) = 7.42\%$), and it also agrees with the less accurate experimental result from 1977Ra23: 73(6)%.

The absolute emission probabilities of the K x-ray components have been deduced from the total P_{XK} using the relative probabilities from sect. 3.2.

The total absolute emission probability of L x-rays has been deduced using the adopted values of ω_{L} and n_{KL} and the recommended values of $P(\text{ce}_{\text{K}}) = 78.6(12)$ and $P(\text{ce}_{\text{L}}) = 10.8(4)\%$.

5.2. Gamma Emissions

A γ -ray energy of 39.578(4) keV has been adopted from 1985Ba73 from an accurate measurement made with a planar HPGe detector. The adopted value coincides with 39.578(2) keV for the energy of the first excited level in ^{129}Xe (1996Te01), deduced from the decay of ^{129}Cs .

Other less accurate experimental values of $E(\gamma_{1,0})$ are (in keV): 39.58(3) (1965Ge04), 39.6(2) (1966Re10), 39.4(3) (1967Gr05), 39.58(5) (1972Ta15), and 39.581(15) (1976Me16).

The absolute γ -ray emission probability (P_{γ}) has been computed as $P(\beta_{1,0})/(1+\alpha_{\text{T}})$. The uncertainty in P_{γ} includes the uncertainty of 0.5% in $P(\beta_{1,0})$, and 1% in α_{T} .

References

- 1951KA16 S. Katcoff, O. A. Schaeffer, J. M. Hastings, Phys. Rev. 82(1951)688. (Half-life)
 1953DE10 Der E. Mateosian, C. S. Wu, Phys. Rev. 91(1953)497A. (Beta spectrum shape factor)
 1954DE17 Der E. Mateosian, C. S. Wu, Phys. Rev. 95(1954)458. (Gamma ray energy)
 1957RU65 H. T. Russell, Report ORNL – 2293(1965). (Half-life)
 1965GE04 J. S. Geiger, R. L. Graham, I. Bergstrom, F. Brown, Nucl. Phys. 68(1965)352. (Gamma ray energy, multipolarity)
 1966RE10 I. Rezanka, A. Spalek, J. Frana, A. Mastalka, Nucl. Phys. 89(1966)609. (Gamma ray energy)
 1967BE65 J. A. Bearden, Revs. Modern Phys. 39(1967)78. (X-ray energies)
 1967GR05 G. Graeffe, W. B. Walters, Phys. Rev. 153(1967)1321. (Gamma ray energy, K ICC)
 1968HA53 R. S. Hager, E. C. Seltzer. Nucl. Data Tables A4(1968)1. (Theoretical ICC)
 1968REZY S. A. Reynolds, J. F. Emery, ORNL-4343 (1968) p 78. (K ICC)
 1969HA61 R. S. Hager, E. C. Seltzer, Nucl. Data Tables A6(1969)1. (Theoretical ICC)
 1970GY01 F. N. Gyax, R. F. Jenefsky, H. J. Leisi, Phys. Letters 32B(1970)359. (K ICC)
 1970SAZI K. S. R. Sastry, R. E. Wood, J. M. Palms, P. V. Rao, Bull. Am. Phys. Soc. 15(4), JE12(1970)623. (K ICC)
 1971DR11 O. Dragoun, Z. Plajner, F. Schmutzler, Nucl. Data Tables A9(1971)119. (Theoretical ICC)
 1972Em01 J. F. Emery, S. A. Reynolds, E. I. Wyatt, G. I. Gleason, Nucl. Sci. Eng. 48(1972)319. (Half-life)

- 1972TA15 H. W. Taylor, B. Singh, J. Phys. Soc. Japan 32(1972)1472. (Gamma ray energy)
- 1973KU17 J. G. Kuhry, G. Bontems, Radiochem. Radioanal. Letters 15(1973)29. (Half-life)
- 1974MA24 G. Marest, R. Haroutunian, I. Berkes, M. Meyer, M. Rots, J. De Raedt, H. Van de Voorde, Van de, H. Oonis, R. Coussement, Phys. Rev. C10(1974)402. (Multipolarity)
- 1974RA26 K. Venkata Ramaniah, T. Seshi Reddy, K. Venkata Reddy, Current Sci. (India) 43 (1974)406. (Multipolarity, K ICC)
- 1976ME16 R. A. Meyer, F. F. Momyer, J. H. Landrum, E. A. Henry, R. P. Yaffe, W. B. Walters, Phys. Rev. C14(1976)1152. (Gamma ray energy)
- 1977RA23 T. K. Ragimov, D. F. Rau, V. I. Timoshin, Bull. Akad. Sci. USSR, Phys. Ser. (1977), 1941(6), 97. Izv Akad Nauk SSSR. Ser. Fiz. 41(1977)1222 (K ICC)
- 1979COZG N. F. Coursol, Thesis, Univ de Paris (1979). (K ICC, M ICC, β^- emission energy)
- 1985BA73 G. Barci-Funel, M. C. Kouassi, G. Ardisson, Nuclear Instrum. Methods 24(1985)252 (K ICC, gamma ray energy, gamma ray and K X-ray emission probabilities)
- 1995AU04 G. Audi and A. H. Wapstra, Nucl.Phys. A595(1995)409. (Q value)
- 1996SC06 E. Schönfeld, H. Janßen, Nucl. Instrum. Methods Phys. Res. A369(1996)527. (Atomic data)
- 1996TE01 Y. Tendow, Nucl. Data Sheets 77(1996)631 (Level energies)
- 1998Schönfeld E. Schönfeld and G. Rodloff, report PTB-6,11-98-1, Braunschweig, October 1998. (Energies of Auger electrons)
- 1998Si17 B. Singh, J. L. Rodriguez, S. S. M. Wong, J. K. Tuli, Nucl. Data Sheets 84(1998)487. (Systematics of log ft values)
- 1999Schönfeld E. Schönfeld and G. Rodloff, report PTB-6.11-1999-1999-1, Braunschweig, February 1999. (KX ray energies and relative emission probabilities)
- 2001Be M. M. Bé, INDC(NDS)-422, IAEA, Vienna, P.112. (Average β^- energy)

¹³¹I – Comments on evaluation of decay data by V. Chisté and M. M. Bé

The first evaluation was done in 2002; it has been updated in December 2013. Literature available by this date was included.

1) Decay Scheme

¹³¹I disintegrates by β^- emission via excited levels of ¹³¹Xe, including the isomeric state ^{131m}Xe ($T_{1/2} = 11,962$ (20) d).

The state of ideal balance, where the activity of ¹³¹I is equal to the activity of ^{131m}Xe, is for $tm = 14,04$ (9) days:

$$tm = \frac{1,4427 \times T_{1/2}({}^{131}\text{I}) \times T_{1/2}({}^{131m}\text{Xe}) \times \ln\left(\frac{T_{1/2}({}^{131m}\text{Xe})}{T_{1/2}({}^{131}\text{I})}\right)}{T_{1/2}({}^{131m}\text{Xe}) - T_{1/2}({}^{131}\text{I})}$$

The decay of ^{131m}Xe interferes with the decay of ¹³¹I only through the 163,9 keV gamma line. For this line, the gamma emission intensity is given, and is only valid, at tm (see above).

2) Nuclear Data

The Q value is from Wang *et al.* (2012Wa38).

Level energies, half-lives, spins and parities are from Yu. Khazov *et al.* (2006Kh09).

The measured ¹³¹I half-life values are, in days:

	$T_{1/2}$	
Reference	Value (d)	Comments
Livingood (1938Li05)	8,0 (2)	
Sreb (1951Sreb)	8,1409 (62)	
Sinclair (1951Si25)	8,04 (4)	
Lockett (1953Lo09)	8,06 (2)	
Seliger (1953Se28)	8,075 (22)	
Bartholomew (1953Ba10)	8,05 (1)	
Burkinshaw (1958Bu14)	8,054 (10)	
Keene (1958Ke26)	8,067 (7)	
Yakovleva (1967Ya02)	8,05 (6)	
Kemeny (1968Ke05)	8,04 (4)	
Rupp (1970Ru**)	8,026 (7)	Mean of three values
Zoller (1971Zo02)	8,117 (12)	
Emery (1972Em01)	8,040 (1)	Other result: 8,048 (16)
Karsten (1974Ka37)	8,031 (4)	
Lagoutine (1978La21)	8,020 (3)	
Houtermans (1980Ho17)	8,0213 (9)	

Reference	Value (d)	Comments
Hoppes (1982HoZF)	8,020 (2)	Superseded by 1992Un01
Walz (1983Wa26)	8,0207 (1)	Superseded by 2004Sc04
Unterweger (1992Un01)	8,0197 (22)	Superseded by 2012Fi12
da Silva (2004Da05)	7,999 (9)	
Schrader (2004Sc04)	8,0252 (6)	
Fitzgerald (2012Fi12)	8,0196 (22)	Superseded by 2014Un**
Unterweger (2014Un**)	8,020 (9)	
Crit (χ^2)	3,3	
$\chi^2 / (n-1)$	4,6	
UWM	8,0234	
LWM	8,0233	
uc(WM)int. :	0,0006	
uc(WM)ext. :	0,0012	
Adopted:	8,0233 (19)	

The half-life weighted average was calculated by the Lweight program (version 3).

The set of the 16 values (not superseded) is discrepant with a $\chi^2 / (n-1) = 19,1$; the unweighted mean of 8,037 (12) d would be chosen.

Only the six most recent values (1974Ka37, 1978La21, 1980Ho17, 2004Da05, 2004Sc04, 2014Un**) were included in the statistical processing. The da Silva *et al.* (2004Da05) value is rejected by the Lweight program, based on the Chauvenet's criterion. The program Lweight 3 increased the uncertainty of the 2004Sc04 value from 0,0006 to 0,00084 in order to reduce its relative weight to 50 %.

The adopted value is the weighted mean: 8,0233 d , with an uncertainty of 0,0019 d , expanded so range includes the most precise value of Schrader (2004Sc04).

2.1) β^- Transitions

The β^- probabilities and the associated uncertainties were deduced from γ transition probability balance at each level of the decay scheme, assuming no β^- transition to the ground state. The values of $lg ft$ were calculated with the program LOGFT for the Allowed, 1st Forbidden and 1st Unique Forbidden transitions.

2.2) Gamma Transitions

Probabilities

The transition probabilities were derived from the gamma emission intensities and the internal conversion coefficients.

Early experiments showed evidence for a weak β branch populating the 163 keV level and, that the following on isomeric transition goes directly to the ground state with a probability in the range 0,5 % - 1 %, see Bergström (1952Be55) and references therein. Meyer *et al.* (1974Me21) carried out a specific experiment and determined the total population of the 163 keV level to be 1,086 (7) %.

Mixing ratios and internal conversion coefficients

The adopted ICC(s) are the theoretical values interpolated by the BrIcc program (2008Ki07) from the tables of Band *et al.* (2002Ba85), accepting the "frozen orbital (no hole)" approximation.

The 163 keV transition is of M4 character, see ^{131m}Xe Comments for details.

For the 177, 272, 318, 324, 325, 364, 404 and 722 keV gamma transitions, which are all of M1+E2 character, the adopted δ (mixing ratio) are from the Krane's evaluation (1977Kr13) of experimental values obtained from angular distribution and correlation data.

For the **364 keV** gamma transition, several values of δ were found in the literature, as shown in the following table with the corresponding theoretical α_T coefficient:

Reference	Value of δ	Theoretical α_T
Johnson <i>et al.</i> – Phys. Rev. 120(1960)1777	-6,7 (5)	0,0227 (7)
Daniel <i>et al.</i> – Z. Phys. 179(1964)62	-4,7 (+4 -2)	0,0228 (4)
Langhoff– Nucl. Phys. A158(1970)657	-3,4 (5)	0,0228 (4)
Krane <i>et al.</i> – Phys. Rev. C5(1972)1671	-3,3 (6)	0,0228 (4)
Koene and Postman – Nucl. Phys. A219(1974)563	4,53 (12)	0,0228 (4)
Irving <i>et al.</i> – J. Phys. G5(1979)1595	3,8 (3)	0,0228 (4)
Krane - Atomic Data and Nuclear Data Tables 19(1977)363	-4,53 (11) (adopted value)	0,0228 (4)

It can be shown that the value of δ has a weak influence on the α_T value.

The corresponding theoretical α_K value of 0,0190 (3) is consistent with the measured values of 0,021 (Haskins and Kurbatov, 1952Ha**) and 0,0174 (Wolfson *et al.*, 1962Wo09).

For the **325 keV** gamma transition, a value of δ (= -4,4 (16)) measured by Koene *et al.* (1975Ko15) is not close to the adopted one (δ = -0,8 (7)) from Krane's evaluation which was derived from Bargholtz *et al.* (1976Ba42), and the two resulting α_K values are 0,0267 (4) and 0,0277 (8), respectively. They are not consistent with the measured by Graeffe and Walters (1967Gr05) α_K = 0,0236 (28) and Hargrove (1963Ha08) = 0,03. The uncertainties on the theoretical ICCs (with δ = -0,8 (7)) are enlarged to 3 % in the final tables.

For the **404 keV** gamma transition, values of δ (= -8,1 (+7,1 ; -4,3)) from Irving (1979Ir01) and $0,2 \leq \delta \leq 2,0$ from Koene and Postman (1974Ko02) were found in the literature. The Krane's recommendation is: δ = +1,0 (+10, -8).

The interpolated theoretical α_T for this δ is 0,0177 (12) .

For the **722 keV** gamma transition, the following values of δ were found in the literature:

Reference	Value of δ	Theoretical α_T
Koene and Postman – Nucl. Phys. A219(1974)563	0,207 (5)	0,00451 (7)
Irving <i>et al.</i> – J. Phys. G5(1979)1595	-0,12 (+5, -2)	0,00454 (7)
Krane - Atomic Data and Nuclear Data Tables 19(1977)363	0,207 (5) (adopted value)	0,00451 (7)

The theoretical α_K = 0,00390 (6) value can be compared with the measured α_K values of 0,0034 (Haskins) and 0,0038 (1963Ha08). The value of 0,0028 (30) from Bell (1952Be95) is consistent within the uncertainty limits.

For the other transitions, measurements aren't precise, and only ranges of values are given for δ . But some values were published for the ICCs.

Comments on evaluation

For the **80 keV** gamma transition, the theoretical values of α_K and α_L interpolated for a M1 character are consistent with the experimental results. A small contribution of E2 character can exist. An uncertainty of 3 % has been adopted in the final tables.

Reference	ICC, α_K	Multipolarity	Theoretical ICC
Bell (1952Be95)	1,73 (20)	M1	$\alpha_K = 1,32$ (4)
Jungclausen (1963Ju02)	1,33 (16)		
Graeffe (1967Gr05)	1,2 (3)		
Suri (1970SuZQ)	1,31 (18)		
Hargrove (1963Hr08)	1		
Daniel (1964Da19)	$\alpha_L = 0,17$ (2)	M1	$\alpha_L = 0,175$ (5)

For the **503 keV** gamma transition, an E2 character is adopted.

Reference	ICC, α_K	Multipolarity	Theoretical α_K
Graeffe (1967Gr05)	0,0085 (13)	E2	0,00748 (11)
Hargrove (1963Hr08)	0,006		

The **284** and **636 keV** transitions were found of E2 character (1960Jo11, 1963Ha08 1970Be46, 1970SuZQ, 1979Ir01).

Measured α_K values for the 284 keV: 0,052 (1952Ha**) and 0,036 (1963Hr08) for the 284 keV, theoretical value $\alpha_K = 0,0408$ (6).

Several results were published for the 636 keV transition:

Reference	ICC, α_K	Multipolarity	Theoretical ICC
Bell (1952Be95)	0,0037 (25)	E2	$\alpha_K = 0,00401$ (6)
Haskins (1952Ha**)	0,0040		
Graeffe (1967Gr05)	0,0043 (4)		
Hargrove (1963Hr08)	0,0039		
Daniel (1964Da19)	0,0039 (3)		

3) Atomic Data

Atomic values (ω_K , ω_L and n_{KL} , $K\beta/K\alpha$, etc.) are from Schönfeld and Janßen (1996Sc06).

The X-ray and Auger electron emission intensities were calculated from the decay scheme data by using the program EMISSION.

L and K X-rays were measured by Chand *et al.* (1989Ch45) in relative values. The evaluators have converted them to absolute values adopting $I_\gamma(364) = 81,2$ (5) %. K X-rays values of Lépy *et al.* (2014Le**) were given directly in absolute values. Both are compared with the values derived from the decay scheme data.

	Chand	Lépy	Calculated
L ℓ	0,0114 (16)		0,01220 (40)
L α	0,294 (18)		0,321 (9)
L β	0,233 (15)		0,256 (6)
L γ	0,0292 (24)		0,0377 (9)
K α	4,18 (12)	4,13 (6)	4,33 (7)
K' β 1	0,741 (25)		0,816 (19)
K' β 2	0,1705 (50)		0,193 (6)
K β	0,912 (25)	0,970 (14)	1,009 (20)
K $\beta/K\alpha$	0,218 (9)	0,235 (5)	0,233 (6)

4) Radiation emissions

4.1) Gamma ray emissions

Gamma ray energies (in keV) are from level energies and R. A. Meyer (1990Me15).

For gamma ray emission intensities, two publications of Meyer *et al.* (1974Me21, 1990Me15) are available. In both, the measurement results are given in absolute values and are the same but the uncertainties have been re evaluated in the second publication. Only the latest one was taken into account. They are compared in the following table with the results of Lépy *et al.* (2014Le**) who took the opportunity to determine absolute intensities from the activity value of a ¹³¹I solution carefully determined by using 4πβγ coincidence method. Both results are, generally, in agreement within the uncertainty limits.

Absolute γ ray intensities:

keV	Meyer	Lépy
80	2,60 (3)	2,593 (28)
85,9	0,00009 (5)	0,0051 (7)
177	0,263 (2)	0,2868 (33)
232	0,0014 (18)	
272	0,056 (1)	0,0581 (17)
284	6,01 (6)	6,227 (40)
295	0,0007 (4)	
302	0,005 (1)	
318	0,079 (3)	0,0830 (19)
324	0,022 (4)	0,02786 (47)
325	0,249 (4)	0,2698 (24)
358	0,0091 (2)	0,0162 (20)
364	80,6 (16)	81,3 (5)
404	0,056 (2)	0,0565 (18)
449	0,007 (2)	
503	0,358 (7)	0,3529 (34)
636	7,21 (9)	7,132 (45)
642	0,218 (4)	0,2183 (19)
722	1,79 (4)	1,795 (12)

The measured emission intensities listed in Table 1 are given in values relative to the 364 keV line adopted equal to 100.

The oldest values from 1952Be95, 1963Ju02, 1963Ha08, 1964Da19, 1966Mo10, 1967Gr05 and 1967Yt02 were omitted from the analysis because they are less precise than those of 1972Si**, 1989Ch45, 1990Me15 and 2014Le** and because they exhibit a large range of values, *e.g.* from 6,0 (10) to 8,2 (8) for the 284 keV ray.

The emission intensity values from Meyer (1990Me15) were converted to the 364-keV ray relative intensity taken equal to 100.

For the 163 keV gamma transition probability, $P_{abs}(163 \text{ keV})$, an absolute value of 1,086 (7) %, determined by Meyer, has been accepted.

The normalization factor N to convert the relative emission intensities to absolute intensities can be calculated using the formula:

$$N = (100 - 1,086(7)) / \sum (1 + \alpha_T) P_{rel}$$

where the sum is done over all probabilities of the gamma transitions to the ground state, except for 163 keV transition.

From the adopted α_T and the evaluated relative emission intensities (Table 1), the normalization factor N deduced in such a way is 0,814 (6). The uncertainties were obtained through their propagation on the above formula.

From the two absolute intensity values measured by Meyer and Lépy for the reference line (see table above) a weighted mean of 81,24 % with an external uncertainty of 0,48 is obtained. This value agrees well with the previous one and is somewhat more precise.

Then, the adopted intensity of the 364 keV γ ray is **81,2 (5) %**. And, the adopted normalization factor is 0,812 (5).

4.2) Conversion electrons

The conversion electron emission intensities were deduced from the gamma-ray emission intensities and the theoretical ICC values.

Energy conservation

The total average energy of 969 (5) keV, for one disintegration, calculated from the current evaluated data corresponds well to the available energy of 970,8 (6) keV (Q^-) from the mass tables confirming the consistency of the decay scheme and the reliability of this evaluation.

References

- 1938Li05 J. J. Livingood, G. T. Seaborg, Phys. Rev. 54 (1938) 775 [T_{1/2}]
 1951Sreb J. H. Sreb, Phys. Rev. 81 (1951) 643 [T_{1/2}]
 1951Si25 W. K. Sinclair, A. F. Holloway, Nature (London) 167 (1951) 365 [T_{1/2}]
 1952Be55 I. Bergstrom. Arkiv Fysik 5 (1952) 191. *The Isomers of Krypton and Xenon*
 1952Be95 R. E. Bell, R. L. Graham, Phys. Rev. 86 (1952) 212 [E _{γ} , I _{γ}]
 1952Ha** J. R. Haskins, J. D. Kurbatov. Phys. Rev. 88 (1952) 884 [ICC] *The Disintegration of I131*
 1953Lo09 E. E. Lockett, R. H. Thomas, Nucleonics 11 (1953) 14 [T_{1/2}]
 1953Se28 H. H. Seliger, L. Cavallo, S. V. Culpepper, Phys. Rev. 90 (1953) 443 [T_{1/2}]
 1953Ba10 R. M. Bartholomew, F. Brown, R. C. Hawkings, W. F. Merritt, L. Yaffe, Can. J. Chem. 31 (1953) 120 [T_{1/2}]
 1958Bu14 L. Burkinshaw, Phys. in Med. Biol. 2 (1958) 255 [T_{1/2}]
 1958Ke26 J. P. Keene, L. A. Mackenzie, C. W. Gilbert, Phys. in Med. Biol. 2 (1958) 360 [T_{1/2}]
 1960Jo11 C. E. Johnson, J. F. Schooley, D. A. Shirley, Phys.Rev. 120, 1777 (1960), *Nuclear Orientation of Iodine by Electric hfs Alignment*
 1962Wo09 J. L. Wolfson, J. J. H. Park, L. Yaffe. Nucl.Phys. 39, 613 (1962) *Internal Conversion Spectrum of I¹³¹ (8.06 d)*
 1963Ju02 H. Jungclaussen, J. Schintlmeiter, H. Sodan, Nucl. Phys. 43 (1963) 650 [E _{γ} , I _{γ}]
 1963Ha08 C. K. Hargrove, K. W. Geiger, A. Chatterjee, Nucl. Phys. 40 (1963) 566 [E _{γ} , I _{γ}]
 1964Da19 H. Daniel, O. Mehling, P. Schmidlin, D. Schotte, E. Thummernicht, Z. Phys. 179 (1964) 62 [E _{γ} , P _{γ} , δ]
 1966Mo10 G. A. Moss, D. O. Wells, D. K. McDaniels, Nucl. Phys. 82 (1966) 289 [E _{γ} , I _{γ}]
 1967Gr05 G. Graeffe, W. B. Walters, Phys. Rev. 153 (1967) 1321 [E _{γ} , I _{γ}]
 1967Ya02 G.V. Yakovleva, Program and Theses, Proc. 17th Ann. Conf. Nucl. Spectrosc. Struct. At. Nuclei, p.45 (1967)
 1967Yt02 C. Ythier, G. Ardisson, C. R. Acad. Sc. Paris 264C (1967) 944 [E _{γ} , I _{γ}]
 1968Ke05 P. Kemeny, Rev. Roumaine Phys. 13 (1968) 485 [T_{1/2}]
 1970Be46 M. Berman, G. B. Beard. Phys.Rev. C2, 1506 (1970) *Nuclear Resonance Fluorescence in Xe131*

- 1970La19 H. Langhoff. Nucl.Phys. A158, 657 (1970) *Lifetime Measurements in ^{131}Xe and ^{129}Xe*
- 1970Ru** A. F. Rupp, report ORNL-TM-2876 (1970) 5 [$T_{1/2}$]
- 1970SuZQ K. K. Suri, P. N. Trehan. Proc.Nucl.Phys.Solid State Symp., Madurai, India, Vol.II, p.489 (1970) *The Decay of ^{131}I*
- 1971Zo02 W. H. Zoller, P. K. Hopke, J. L. Fasching, E. S. Macias, W. B. Walters, Phys. Rev. C3 (1971) 1699 [$T_{1/2}$]
- 1972Em01 J. F. Emery, S. A. Reynolds, E. I. Wyatt, G. I. Gleason, Nucl. Scien. Eng. 48 (1972) 319 [$T_{1/2}$]
- 1972Kr07 K. S. Krane, C. E. Olsen, W. A. Steyert. Phys.Rev. C5, 1671 (1972) *Nuclear Orientation Study of the Decay of ^{239}Np Polarized in ZrFe_2 : Parity Mixing in ^{239}Pu and Nuclear Structure of ^{239}Pu and Fission Products $^{131},^{132},^{133}\text{Xe}$*
- 1972Si** N. Singh, S. S. Bhati, R. L. Dhingra, P. N. Trehan, Nucl. Phys. and Solid State Phys. Symp., Chandigarh, India (1972) [E_γ , I_γ]
- 1974Ka37 J. H. M. Karsten, P. G. Marais, F. J. Haasbroek, C. J. Visser, Agrochemophysica 6 (1974) 25 [$T_{1/2}$]
- 1974Ko02 B. K. S. Koene, H. Postman, Nucl. Phys. A219 (1974) 563 [δ]
- 1974Me21 R. A. Meyer, F. Momyer, W. B. Walters, Z. Phys. 268 (1974) 387 [E_γ , I_γ]
- 1975Ko15 B. K. S. Koene, H. Postman, H. Ligthart, Nucl. Phys. A250 (1975) 38 [δ]
- 1976Ba42 C. Bargholtz, S. Beshai, L. Gidefeldt. Nucl.Phys. A270, 189 (1976), *Angular Correlation Measurements in ^{131}Xe*
- 1977Kr13 K. S. Krane. At. Data Nucl. Data Tables 19 (1977) 363 [δ]
- 1978La21 F. Lagoutine, J. Legrand, C. Bac, Int. J. Appl. Radia. Isot. 29 (1978) 269 [$T_{1/2}$]
- 1979Ir01 A. D. Irving, P. D. Forsyth, I. Hall, D. G. E. Martin, J. Phys. G5 (1979) 1595 [δ]
- 1980Ho17 H. Houtermans, O. Milosevic, F. Reichel, Int. J. Appl. Radia. Isot. 31 (1980) 153 [$T_{1/2}$]
- 1982HoZF D. D. Hoppes, NBS-SP 626 (1982) 117 [$T_{1/2}$]
- 1983Wa26 K. F. Walz, K. Debertain, H. Schrader, Int. J. Appl. Radia. Isot. 34 (1983) 1191 [$T_{1/2}$]
- 1989Ch45 B. Chand, J. Goswamy, D. Mehta, N. Singh, P. N. Trehan, Nucl. Inst. Meth. A284 (1989) 393 [E_γ , I_γ]
- 1990Me15 R. A. Meyer, Fisika (Zagreb) 22 (1990) 153 [E_γ , I_γ]
- 1992Un01 M. P. Unterweger, D. D. Hoppes, F. J. Schima, Nucl. Instrum. Meth. A312 (1992) 349 [$T_{1/2}$]
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. A369 (1996) 527 [Atomic data]
- 2002Ba85 I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor Jr., P. O. Tikkanen, S. Raman. At. Data Nucl. Data Tables 81 (2002) 1. [Theoretical ICC]
- 2004Da05 M. A. L. da Silva, M. C. M. de Almeida, C. J. da Silva, J. U. Delgado, Applied Rad. Isotop. 60(2004)301 [$T_{1/2}$].
- 2004Sc04 H. Schrader, Applied Rad. Isotop. 60(2004)317 [$T_{1/2}$]
- 2006Kh09 Yu. Khazov, I. Mitropolsky, A. Rodionov. Data Sheets 107,11 (2006) 2715 [E_γ , I_γ , Spin]
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr. Nucl. Instrum. Methods Phys. Res. A589 (2008) 202. ICC
- 2012Fi12 R. Fitzgerald, J. Research Nat. Inst. Standards and Technology, 117 (2012) 80 [$T_{1/2}$]
- 2012Wa38 M. Wang, G. Audi, A. H. Wapstra, F. G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer. Chinese Physics C36 (2012) 1603. [Q]
- 2014Un** M. P. Unterweger, R. Fitzgerald. Applied Radiation Isotopes (2014), [$T_{1/2}$]
<http://dx.doi.org/10.1016/j.apradiso.2013.11.017>
- 2014Lépy** M.-C. Lépy, L. Brondeau, to be submitted to Applied Radiation Isotopes [I_γ]

Table 1 – Gamma emission intensities, relative values

Ref	80,1853	85,918	177,214	232,175	272,501	284,3047	295,848	302,444	318,093
1952Be95	2,71 (19)					6,6 (25)			
1963Ju02	2,6 (4)					6,0 (10)			
1963Ha08	3,5 (8)		0,29 (6)			7,9 (8)			
1964Da19	3,1 (2)		0,27 (10)			6,6 (3)			
1966Mo10	3,10 (18)		0,313 (26)			7,4 (6)			
1967Gr05	2,72 (15)		0,36 (2)		0,08 (1)	7,05 (40)			0,110 (15)
1967Yt02	3,4 (4)	~ 0,1	0,38 (8)		~ 0,07	8,2 (8)			~ 0,05
1972Si**	3,21 (5)		0,30 (2)			7,49 (5)			0,110 (5)
1989Ch45	3,26 (7)		0,334 (6)	0,0039 (5)	0,0735 (18)	7,56 (8)	0,0022 (10)	0,0057 (8)	0,096 (2)
1990Me15	3,23 (6)	0,00011 (6)	0,326 (7)	0,0017 (10)	0,0695 (19)	7,46 (15)	0,00087 (50)	0,0056 (11)	0,0980 (42)
2014Le**	3,189 (38)	0,0063 (9)	0,3528 (44)*		0,0715 (21)	7,66 (6)			0,1022 (23)
χ^2 crit.	3,8		3,8		4,6	3,8			3,8
$\chi^2/(n-1)$	0,3		5,7		1,2	1,7			3,0
UWM :	3,2223		0,328	0,0028	0,072	7,543	0,0015	0,0056	0,102
WM :	3,2098		0,3410		0,0716	7,5558			0,0994
uc(WM)int. :	0,0258		0,0031		0,0011	0,0339			0,0014
uc(WM)ext. :	0,0141		0,0075		0,0012	0,0441			0,0024
LWM:	3,210 (26)		0,341 (12)		0,0716 (12)	7,556 (44)			0,0994 (24)
Adopted	3,210 (38)	0,0063 (9)	0,341 (8)	0,0028 (11)	0,0716 (18)	7,56 (5)	0,0015 (7)	0,0057 (8)	0,0994 (24)

^o Value omitted from the data set used for averaging.

* The original uncertainty (as given in the table) has been increased to limit its weight to 50 %.

Table 1 – Gamma emission intensities, relative values (Cont.)

Ref	324,6307	325,791	358,419	364,49	404,816	503,005	636,991	642,7237	722,909
1952Be95				100			11,6 (19)		3,5 (31)
1963Ju02							9,0 (10)		3,0 (4)
1963Ha08		0,35 (8)		100		0,52 (17)	8,8 (7)		2,05 (16)
1964Da19		0,26 (10)		100		0,54 (5)	8,3 (3)		1,9 (1)
1966Mo10		0,279 (25)		100		0,45 (6)	9,1 (11)		2,05 (26)
1967Gr05	0,04 (1)	0,45 (3)	0,020 (4)	100	0,080 (7)	0,36 (2)	8,0 (4)	0,180 (15)	2,10 (15)
1967Yt02		0,37 (5)		100	~ 0,06	0,37 (8)	8,2 (8)		1,8 (2)
1972Si**		0,32 (1)		100	0,022 (5) ^o	0,30 (5) ^o	7,79 (10) ^o	0,13 (1) ^o	1,79 (9) ^o
1989Ch45	0,025 (3)	0,361 (5)	0,0304 (11)	100	0,066 (2)	0,438 (5)	8,75 (9)	0,269 (5)	2,19 (2)
1990Me15	0,0273 (50)	0,309 (8)	0,01129 (33)	100 (2)	0,0695 (28)	0,444 (12)	8,95 (21)	0,270 (7)	2,22 (7)
2014Le**	0,0343 (6)*	0,3319 (34)*		100,0 (8)	0,0695 (23)	0,4341 (47)	8,77 (7)	0,2685 (27)	2,208 (19)
χ^2 crit.	4,6	3,8			4,6	4,6	4,6	4,6	4,6
$\chi^2/(n-1)$	2,9	13,3			0,9	0,4	0,4	0,02	0,3
UWM :	0,029	0,330	0,020845		0,068	0,439	8,823	0,269	2,206
WM :	0,0300	0,3371			0,0680	0,4365	8,7746	0,2688	2,2002
uc(WM)int. :	0,0018	0,0028			0,0013	0,0033	0,0534	0,0022	0,0135
uc(WM)ext. :	0,0031	0,0101			0,0012	0,0020	0,0333	0,0003	0,0068
LWM:	0,0300 (31)	0,337 (10)			0,0680 (13)	0,4365 (33)	8,77 (5)	0,2688 (22)	2,200 (14)
Adopted	0,0300 (31)	0,337 (10)	0,021 (10)	100 (8)	0,068 (2)	0,436 (5)	8,77 (7)	0,2688 (27)	2,200 (19)

^o Value omitted from the data set used for averaging.

* The original uncertainty (as given in the table) has been increased to limit its weight to 50%

^{131m}Xe – Comments on evaluation of decay data by V. Chisté and M. M. Bé

The first evaluation was done in 2002, it has been reviewed in December 2013. Literature available by this date was included.

1) Decay Scheme

^{131m}Xe decays by a strongly converted gamma transition.

2) Nuclear Data

Level energies, spins and parities are from Yu. Khazov *et al.* (2006Kh09).

The ^{131m}Xe measured half-life values are, in days:

Reference	T _{1/2} Value (d)	Comments
Bergström (1952Be55)	12,0 (3)	
Andersson (1964An08)	11,8 (1)	
Knauf (1966Kn09)	11,94 (4)	
Emery (1972Em09)	12,00 (2)	
Meyer (1974Me21)	11,770 (12)	Omitted
Hoffman (1975Ho12)	11,92 (3)	
Tam (1990Ta02)	11,9 (2)	
Unterweger (2014Un**)	11,93 (3)	Replace 1992Un01, 2002Un02 and 2012Fi12
Adopted	11,962 (20)	$\chi^2/(n-1) = 1,4$; ext. unc. = 0,016

The set of the eight values is quite discrepant with a $\chi^2/(n-1)$ of 15,3.

The value from Meyer *et al.* (1974Me21) was omitted from the analysis because it significantly disagrees with the other values, then the Andersson (1964An08) value was rejected by the Lweight program (version 3), based on Chauvenet's criterion. The adopted value is the weighted mean of the six remaining results: 11,962 d, with the lowest experimental uncertainty of 0,020.

2.1) Gamma Transitions

The only gamma transition was demonstrated being of M4 character (1954Be36, 1962Ge09, 1969Fr04). The various theoretical conversion coefficients for this transition (Band *et al.*, Hager *et al.*, Rösel *et al.*) differ by about 2 - 4 %. The adopted ICC(s) are the theoretical values interpolated by the BrIcc program (2008Ki07) from the tables of Band *et al.* (2002Ba85), accepting the "frozen orbital (no hole)" approximation.

Reference	Measured α_K	Theoretical ICC
Bergström (1954Be36)	29 (4)	$\alpha_K = 31,6$ (5)
Knauf (1966Kn03)	32,1 (4)	
Geiger (1966Ge01)	30,1 (6)	

3) Atomic Data

Atomic quantities (ω_K , ω_L and n_{KL}) are from Schönfeld and Janßen (1996Sc33).

The X-ray and Auger electron emission probabilities have been calculated from the decay data parameters data by using the program EMISSION.

4) Radiation emissions

4.1) Conversion electrons

The conversion electron emission intensities were deduced from the ICC values and the gamma transition probability.

4.2) Gamma-ray emissions

Gamma-ray emission energy is from the level energy and R. A. Meyer (1990Me15).

The gamma-ray emission intensity has been deduced from the transition probability using the theoretical α_T to be: **1,942 (26) %**.

We have not found measured values for this emission intensity, the ^{131m}Xe radioisotope alone.

Additional Reference

F. Lagoutine, Table de Radionucléides, CEA-LMRI (1984)

References

- 1952Be55 I. Bergstrom. Arkiv Fysik 5, 191 (1952) *The Isomers of Krypton and Xenon* [$T_{1/2}$]
 1954Be36 I. Bergstrom, S. Thulin, A. H. Wapstra, B. Astrom. Arkiv Fysik 7 (1954) 255.
K-Conversion Coefficients of γ -Transitions Occurring in the Decay of Kr^{85m} , Xe^{129m} , Xe^{131m} , Xe^{133} , and Xe^{133m} . [α_K]
 1962Ge09 J. S. Geiger, R. L. Graham, F. Brown, Can. J. Phys. 40 (1962) 1258
 [α_K]
 1964An08 G. Andersson, Arkiv for Fysik 28 (1964) 37
 [$T_{1/2}$]
 1966Ge01 J. S. Geiger. Proc.Intern.Conf.Internal Conversion Process, Nashville, Tenn. (1965), J. H. Hamilton, Ed., Academic Press, Inc., New York, p.379 (1966). [α_K]
The Internal Conversion of High Multipole Order γ -Transitions
 1966Kn09 K. Knauf, H. Sommer, H. Klewe-Nebenius, Z. Phys. 197 (1966) 101
 [$T_{1/2}$, α_K]
 1969Fr04 K. Fransson, P. Erman, Arkiv för Fysik 39 (1969) 7
 [Multipolarity]
 1972Em09 J. F. Emery, S. A. Reynolds, E. I. Wyatt, G. I. Gleason, Nucl. Sci. Eng. 48 (1972) 319
 [$T_{1/2}$]
 1973Be06 P. A. Benson, H. Y. Gee, M. W. Nathans, J. Inorg. Nucl. Chem. 35 (1973) 2614
 [Branching Ratio]
 1974Me21 R. A. Meyer, F. Momyer, W. B. Walters, Z. Phys. 268 (1974) 387
 [Total Branch, $T_{1/2}$]
 1975Ho12 D. C. Hoffman, J. W. Barnes, B. J. Dropesky, F. O. Lawrence, G. M. Kelley, M. A. Ott, J. Inorg. Nucl. Chem. 37 (1975) 2336
 [$T_{1/2}$]

Comments on evaluation

- 1976Au08 R. L. Aube, H. R. Hiddleston, C. P. Browne, Nucl. Data Sheets 17 (1976) 573
[I_γ , Spin, Parity]
- 1990Me15 R. A. Meyer, Fisika (Zagreb) 22 (1990) 153 [E_γ , I_γ]
- 1990Ta02 N. C. Tam, A. Veres, I. Pavlicsek, L. Lakosi, J. Phys. G16 (1990) 1215
[$T_{1/2}$]
- 1992Un01 M. P. Unterweger, D. D. Hoppes, F. J. Schima, Nucl. Instrum. Meth. A312 (1992) 349
[$T_{1/2}$]
- 1996Sc33 E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. A369 (1996) 527
[Atomic data]
- 2002Ba85 I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor Jr., P. O. Tikkanen, S. Raman. At. Data
Nucl. Data Tables 81 (2002) 1. [Theoretical ICC]
- 2006Kh09 Yu. Khazov, I. Mitropolsky, A. Rodionov. Data Sheets 107,11 (2006) 2715
[E_γ , I_γ , Spin]
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr. Nucl.
Instrum. Methods Phys. Res. A589 (2008) 202. [ICC]
- 2012Fi12 R. Fitzgerald, J. Research Nat. Inst. Standards and Technology, 117 (2012) 80 [$T_{1/2}$]
- 2012Wa38 M. Wang, G. Audi, A. H. Wapstra, F. G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer.
Chinese Physics C36 (2012) 1603. [Q]
- 2014Un** M. P. Unterweger, R. Fitzgerald. Applied Radiation Isotopes (2014), [$T_{1/2}$]
<http://dx.doi.org/10.1016/j.apradiso.2013.11.017>

¹³²Te -Comments on evaluation of decay data

by A. L. Nichols

Evaluated: December 2007/March 2009

Evaluation Procedure

Limitation of Relative Statistical Weight Method (LWM) was applied to average the measured decay data when appropriate (see below).

Decay Scheme

A simple decay scheme was constructed from the gamma-ray studies of 1966Fr02 and 1981Yo02. An earlier study involved the use of low-resolution NaI(Tl) detectors, and these data have been set aside from consideration in this particular evaluation [1958Ch28]. The gamma-ray emission probabilities were expressed in terms of the emission probability of the 228.327-keV gamma ray (100 %), and weighted mean data were derived as appropriate.

All 100 % of the beta decay goes directly to the 277.86-keV nuclear level of ¹³²I, and the resulting four gamma cascade dominates the decay scheme.

Nuclear Data

¹³²Te undergoes beta decay to the 277.86-keV nuclear level of ¹³²I that undergoes gamma decay to the ground state of ¹³²I predominantly through the 49.72- and 228.327-keV gamma transitions.

Half-life (¹³²Te)

The recommended half-life has been determined from the measurements of Cheever *et al.* (1958Ch28), Andersson *et al.* (1965An05), Baba *et al.* (1971BaZW) and Walz *et al.* (1983Wa26). A value of 3.230 (13) days was derived in terms of LWM, with the uncertainty increased to the lowest measured value of ± 0.013 .

Half-life measurements (¹³²Te).

Reference	Half-life (days)
1956F115	2.8 \pm 0.1*
1958Ch28	3.2 \pm 0.2
1965An05	3.26 \pm 0.03
1971BaZW	3.28 \pm 0.02
1983Wa26	3.204 \pm 0.013
Recommended value	3.230 \pm 0.013

* set aside from the LWM analysis as an outlier.

Gamma Rays

Energies

Gamma-ray transition energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies of 2005Kh07 were adopted, and used to determine the energies of the gamma-ray transitions between the populated-depopulated levels, apart from the 228.327-keV gamma ray which was taken from 1979Bo26.

Emission Probabilities

Although judged to be a rather limited data set, a reasonably consistent decay scheme was derived from the relative gamma-ray emission probabilities measured by Fransson and Bemis (1966Fr02) and Yousif *et al.* (1981Yo02). These relative emission probabilities were normalised to the 228.327-keV gamma ray (100 %). The 49.72-keV gamma ray has only been quantified by 1966Fr02, and therefore the relative emission probability of this low-energy gamma ray was calculated from the population-depopulation balance of the 49.72-keV nuclear level (with no populating beta transition). A value of 2.1 (2) % was adopted for the relative emission probability of the 111.80-keV gamma ray on the basis of the population-depopulation balance of the 161.52-keV nuclear level (with no populating beta transition) and the measurement of Fransson and Bemis. Finally, the possible existence of a low-intensity 161.5-keV gamma transition from the 161.52-keV nuclear level to the ground state of I-132 was discarded on consideration of the population-depopulation of the 161.52-keV nuclear level.

Relative gamma-ray emission probabilities (%).

transition	E_γ (keV)	P_γ^{rel}		
		1966Fr02	1981Yo02	Recommended
$\gamma_{2,0}$ (I)	49.72 (1)	16.3 (11)	17.02 (34)*	17.14 (4)*
$\gamma_{4,2}$ (I)	111.80 (8)	2.1 (2)	1.98 (5)	2.1 (2)†
$\gamma_{5,4}$ (I)	116.34 (13)	2.2 (2)	2.23 (6)	2.23 (6)
$\gamma_{5,2}$ (I)	228.327 (3)	100 (6)	100 (2)	100 (2)

* deduced from decay scheme and calculated branching ratio (not measured directly).

† adopted from 1966Fr02 and on the basis of the population-depopulation balance of the 161.52-keV nuclear level.

Gamma-ray emissions: recommended energies, relative emission probabilities, multiplicities and theoretical internal conversion coefficients (frozen orbital approximation).

E_γ (keV)	P_γ^{rel}	Multiplicity	α_K	α_L	α_{M+}	α_{tot}
49.72 (1)	17.14 (4)	M1	4.83 (7)	0.64 (1)	0.15 (1)	5.62 (8)
111.80 (8)	2.1 (2)	M1 + E2 $\delta = 0.58$ (6)	0.562 (17)	0.115 (9)	0.033 (4)	0.71 (3)
116.34 (13)	2.23 (6)	M1 + E2 $\delta = 0.53$ (5)	0.489 (13)	0.093 (6)	0.024 (1)	0.606 (20)
228.327 (3)	100 (2)	E2	0.0802 (12)	0.0151 (2)	0.0037 (1)	0.0990 (14)

Multiplicities and Internal Conversion Coefficients

The nuclear level scheme specified by Khazov *et al.* (2005Kh07) has been used to define the multiplicities of the gamma transitions on the basis of known spins and parities. Somewhat

disparate mixing ratios were obtained by Fransson and Bemis (1966Fr02) and Yousif *et al.* (1981Yo02). All of the multipolarities recommended by Yousif *et al.* were adopted with improved uncertainties introduced for the (M1 + E2) transitions. These data were used to determine the internal conversion coefficients of the 49.7-, 111.8-, 116.3- and 228.327-keV gamma rays from the theoretical tabulations of Band *et al.* (2002Ba85, 2002Ra45) by means of the methodology of Kibédi *et al.* (2008Ki07) in which the frozen orbital approximation was adopted.

A normalisation factor of 0.8812 (13) was calculated from the internal conversion coefficients and relative emission probabilities of the gamma-ray transitions depopulating the 277.86-keV nuclear level of ¹³²I, assuming that there is no direct beta feeding to other levels as implied from the various spins and parities:

$$\sum P_{\gamma+ce}^{rel} = 100\%$$

$$P_{\gamma}(116.34 \text{ keV}) + P_{\gamma}(228.327 \text{ keV}) F = 100$$

$$[3.58(10) + 109.90(14)] F = 100$$

$$F = 0.8812 \pm 0.0013$$

Beta-particle Emission

Energy and emission probability

The single beta-particle energy was calculated from the structural detail of the proposed decay scheme. A nuclear level energy of 277.86(6) keV adopted from Khazov *et al.* (2005Kh07) and a Q_{β^-} value of 518 ± 4 keV from Audi *et al.* (2003Au03) were used to determine the energy and uncertainty of this beta-particle transition.

Beta-particle Emission Probability per 100 Disintegrations of ¹³²Te.

Transition	E _β (keV)	P _β	Transition type	logft
β _{0,5} ⁻	240 ± 4	100	allowed	4.85

Atomic Data

The x-ray and Auger electron data have been calculated using the evaluated gamma-ray data, and the atomic data from 1977La19, 1996Sc06, 1998ScZM and 1999ScZX.

References

- 1956Fl15 W.H. FLEMING, H.G. THODE, The mass assignment of the chain 2 min. Sb → 77.7 hr. Te → 2.25 hr. I, *Can. J. Chem.* 34 (1956) 408-409. [half-life]
- 1958Ch28 G.D. CHEEVER, W.S. KOSKI, D.R. TILLEY, L. MADANSKY, Decay of tellurium-132, *Phys. Rev.* 110 (1958) 922-923. [half-life, E_γ, α_K]
- 1965An05 G. ANDERSSON, G. RUDSTAM, G. SÖRENSEN, Decay data on some Xe, I, and Te isotopes, *Ark. Fys.* 28 (1965) 37-43. [half-life]
- 1966Fr02 K. FRANSSON, C.E. BEMIS Jr., The decay of ¹³²Te and levels in odd ¹³²I, *Nucl. Phys.* 78 (1966) 207-224. [E_γ, P_γ, δ, ICC]

- 1971BaZWS. BABA, H. BABA, H. UMEZAWA, T. SUZUKI, T. SATO, H. NATSUME, Decay analysis of some fission product nuclides with medium half-lives, Japan Atomic Energy Research Institute report JAERI-1211 (1971). [half-life]
- 1977La19 F.P. LARKINS, Semi empirical Auger-electron energies for elements $10 \leq Z \leq 100$, At. Data Nucl. Data Tables 20 (1977) 311-387. [Auger and conversion electron energies]
- 1979Bo26 H.G. BÖRNER, W.F. DAVIDSON, J. ALMEIDA, J. BLACHOT, J.A. PINSTON, P.H.M. VAN ASSCHE, High precision gamma-ray energy measurements of fission products, Nucl. Instrum. Methods 164 (1979) 579-586. [precise E_γ]
- 1981Yo02 A.A. YOUSIF, W.D. HAMILTON, E. MICHELAKAKIS, Gamma-ray transition strengths and multipolarities in the doubly odd nucleus ¹³²I, J. Phys. G: Nucl. Phys. 7 (1981) 445-453. [E_γ , P_γ , δ , α_K]
- 1983Wa26 K.F. WALZ, K. DEBERTIN, H. SCHRADER, Half-life measurements at the PTB, Int. J. Appl. Radiat. Isot. 34 (1983) 1191-1199. [half-life]
- 1996Sc06 E. SCHÖNFELD, H. JANßEN, Evaluation of atomic shell data, Nucl. Instrum. Meth. Phys. Res. A369 (1996) 527-533. [X_K , X_L , Auger electrons]
- 1998ScZM E. SCHÖNFELD, G. RODLOFF, Tables of the energies of K-Auger electrons for elements with atomic numbers in the range from $Z = 11$ to $Z = 100$, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX E. SCHÖNFELD, G. RODLOFF, Energies and relative emission probabilities of K X-rays for elements with atomic numbers in the range from $Z = 5$ to $Z = 100$, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2002Ba85 I.M. BAND, M.B. TRZHASKOVSKAYA, C.W. NESTOR, Jr., P.O. TIKKANEN, S. RAMAN, Dirac-Fock internal conversion coefficients, At. Data Nucl. Data Tables 81 (2002) 1-334. [ICC]
- 2002Ra45 S. RAMAN, C.W. NESTOR, Jr., A. ICHIHARA, M.B. TRZHASKOVSKAYA, How good are the internal conversion coefficients now? Phys. Rev. C66 (2002) 044312, 1-23. [ICC]
- 2003Au03 G. AUDI, A.H. WAPSTRA, C. THIBAUT, The AME2003 atomic mass evaluation (II). Tables, graphs and references, Nucl. Phys. A729 (2003) 337-676. [Q-value]
- 2005Kh07 Yu. KHAZOV, A.A. RODIONOV, S. SAKHAROV, B. SINGH, Nuclear data sheets for $A = 132$, Nucl. Data Sheets 104 (2005) 497-790. [nuclear levels]
- 2008Ki07 T. KIBÉDI, T.W. BURROWS, M.B. TRZHASKOVSKAYA, P.M. DAVIDSON, C.W. NESTOR, Jr., Evaluation of theoretical conversion coefficients using BrIcc, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202-229. [ICC]

**¹³³I - Comments on evaluation of decay data
by M. Galán**

1) Decay Scheme

¹³³I disintegrates by β^- emission to excited levels in ¹³³Xe, included the isomeric state ¹³³Xe^m at 233 keV ($T_{1/2} = 2,198$ (13) d).

¹³³I ground state has $J^\pi = 7/2^+$ (1976FU06).

2) Nuclear Data

The Q value is from AME2003 (2003Au03): $Q(\beta^-) = 1757$ (4) keV.

Level energies have been obtained from a least-squares fit to γ -ray energies (GTOL computer code) from 1976ME16. The energy of the isomeric level is from the ¹³³Xe^m evaluation. Spin and parities are from 1995RA12 except for the 1386-keV level. For this level the adopted value is $J^\pi(1386) = 7/2^+$ as proposed by 1976ME16 based on M1+E2 to $5/2^+$ (deduced from $\delta(856) = +3,7$ (3) (1974KO26 and 1977KR13)). J^π for 743-, 875-, 911-, and 1236-keV levels are uncertain.

The measured ¹³³I half-life values, in hours, are:

Reference	Value (h)	Comments
1968RE04	20,9 (1)	
1966EI01	20,8 (2)	
1965AN05	20,3 (3)	Rejected by Chauvenet's criterion
1955WA35	20,9 (3)	
1953KA28	20,8 (2)	
LWeight for Excel Code		
Nb of input values	4	
Reduced χ^2	0,10	
Weighted Mean	20,86	
Internal uncertainty	0,09	
External uncertainty	0,03	
Ave Tool Code		
Nb of input values	4	
	Mean	Reduced χ^2
LWM	20,87 (8)	0,11
NRM	20,87 (8)	0,11
RT	20,87 (8)	

The half-life was calculated by the Lweight for Excel code (version 2004) and by AveTool code. In both codes the value of 1965AN05 was rejected based on the Chauvenet's criterion. Ave Tool was run again without the value from 1965AN05. The results of the three statistical methods LWM (Limitation of

Relative Statistical Weight), NRM (Normalised Residual Method) and RT (Rajeval Technique) given by AveTool are also shown in the table. The recommended value is 20,87 (8) h.

2.1) b Transitions

The energies of the β transitions were deduced from the Q value and the level energies in ^{133}Xe , the later deduced from γ -ray transition energies. Some experimental values (1966Ei01) with the adopted ones are compared in the table:

Beta Transition	Adopted (keV)	1966Ei01 (keV)
$\beta_{0,9}$	521 (4)	500 (30)
$\beta_{0,6}$	882 (4)	890 (30)
$\beta_{0,3}$	1227 (4)	1230 (30)
$\beta_{0,1}$	1524 (4)	1540 (30)

The β^- probabilities and associated uncertainties have been deduced from γ -ray transition intensity balance at each level of the decay scheme, assuming no β^- transition to the ground state. These values are compared to the β^- emission probabilities measured by 1966EI01, 1971SA09 and 1976ME16. The $\lg ft$ values were calculated using the program LOGFT for the Allowed, 1st Forbidden and 1st Unique Forbidden β^- transitions.

Beta Transition	Adopted (%)	1966EI01 (%)	1971SA09 (%)	1976ME16 (%)
$\beta_{0,13}$	0,414 (15)	0,5	0,5	0,42
$\beta_{0,12}$	1,25 (4)	3,5	1,1	1,26
$\beta_{0,11}$	0,397 (12)	0,4	0,3	0,4
$\beta_{0,10}$	3,75 (7)	3,7	2,9	3,68
$\beta_{0,9}$	3,12 (6)	3,3	3,2	3,16
$\beta_{0,8}$	0,58 (5)	0,5	0,5	0,62
$\beta_{0,7}$	0,026 (18)	-	-	-
$\beta_{0,6}$	4,16 (13)	2,3	3,5	4,1
$\beta_{0,5}$	1,81 (6)	-	2,3	1,81
$\beta_{0,3}$	83,44 (21)	85,4	83,2	83,5
$\beta_{0,1}$	1,07 (6)	1,4	1,4	1,07

A beta transition of about 1080 keV to the 680-keV level was observed by 1966EI01 with a β^- probability = 0,3 %. 1971SA09 reported 0,2 % β^- probability for this transition.

2.2) g-ray Transitions

Transition Probabilities

For the 233-keV gamma transition probability, the adopted value is 2,88 (2) % measured by 1976ME16. Other transition probabilities have been calculated from the γ -ray emission probabilities using the recommended internal conversion coefficients.

Mixing ratios and internal conversion coefficients

For the 233-keV γ -ray transitions the adopted δ (mixing ratio) is from $^{133}\text{Xe}^m$ evaluation. The adopted δ values for the 417 -, 422-, 529-, 680- and 1298-keV are from 1977KR13. The adopted values were

deduced from angular correlation data. For the 768 -, 820 and 856 γ -ray transitions the adopted δ values are from 1974KO26 obtained by directional distributions of γ -rays. For the 909 -keV line a $\delta(909) = +0,40(6)$ has been adopted, as was reported by 1974KO26 if the $J^\pi(1589) = 5/2^+$.

The internal conversion coefficients (ICC) were calculated using the BrIcc computer code, which interpolated ICC values from tables of Band *et al.* (2002BA85).

Only experimental measurements of α_K and K/L values were found for the internal transition of 233-keV (see $^{133}\text{Xe}^m$ evaluation).

3) Atomic Data

Atomic values (ω_K , ω_L and η_{KL}) are from 1996SC06.

ω_K	$0,888 \pm 0,005$
ω_L	$0,097 \pm 0,005$
η_{KL}	$0,902 \pm 0,004$

The X-ray and Auger electron emission probabilities have been deduced from γ -ray and conversion electron data by using the computer code EMISSION. Results were verified with RADLST computer code.

4) Electron Emissions

The conversion electron emission probabilities have been computed from γ -ray emission probabilities and theoretical ICC values.

5) Photon Emissions

Energies

γ -ray energies and uncertainties are from level scheme. The isomeric transition γ -ray energy is from 2000HE14 (see ^{133m}Xe evaluation).

g-ray emissions

The gamma emission intensities are from 1976ME16. A 2% was increased by the evaluator in the uncertainty to account for uncertainty calibration, as cited by 1976ME16. Other experimental measurements are shown in table 1. In table 1 the absolute intensity values reported by 1974KO26 are just compared to the absolute intensity values recommended in this evaluation. The evaluator has not used the values of 1974KO26 in the present evaluation because detailed information, such as the detector calibration and uncertainty, calculation procedure or experimental conditions under which the absolute gamma intensities were achieved, are absent.

The normalization factor has been deduced from the decay scheme using the formulas:

$$N = \frac{100 - P_{g+ce}(233\text{keV})}{\sum_i I_{g_i} [1 + a_{T_i}]} \quad \text{and} \quad dN^2 = \left(\frac{\partial N}{\partial P_{g+ce}} \right)^2 + \sum_i \left(\frac{\partial N}{\partial I_{g_i}} dI_{g_i} \right)^2 + \sum_i \left(\frac{\partial N}{\partial a_{T_i}} da_{T_i} \right)^2,$$

where the sum is over all γ -ray transitions to the ground state (g.s.), thus considering no direct β^- feeding to the g.s. For the 233 -keV γ transition probability, $P_{\gamma+ce}(233 \text{ keV})$, an absolute value of 2,88 (2) %, determined by 1976ME16, has been accepted. From the estimated α_T (BrIcc) and the evaluated relative γ

emission intensities (Table 1) the deduced normalization factor is 0,0863 (16). This result was checked with the value of 0,0863 (16) reported by GABS computer code.

In Table 5.2 Gamma Emissions. The absolute gamma emission intensity of 0,293 (4) % for the 233-keV line has been estimated by the evaluator from $P_{\gamma+ce} = 2,88$ (2) % and $\alpha_T = 8,84$ (12).

References

- 1953KA28 Katcoff, S.; Rubinson, W. Phys. Rev 91(1953) 1458
[$T_{1/2}$]
- 1955WA35 Wahl, A. Phys. Rev. 99 (1955) 730
[$T_{1/2}$]
- 1959HO97 Holm, G. Ryde, H. Ark. Fysik 15 (1959) 387
[I_γ]
- 1960EI01 Eichler, E. ; Chase, J.W.; Johnson, N.R.; O'Kelley, G.D. Bull. Am. Phys. Soc 5 (1960) 448
[E_β]
- 1965AN05 Anderson, G.; Rudstam, G.; Sörensen, G. Ark. Fysik. 28 (1965) 37
[$T_{1/2}$]
- 1966EI01 Eichler, E. ; Chase, J.W.; Johnson, N.R.; O'Kelley, G.D. Phys. Rev. 146 (1966) 899
[$T_{1/2}, I_\beta$]
- 1968RE04 Reynolds, S.A.; Emery, J.F.; Wyatt, E.I. Nucl. Sci. Eng. 32 (1968) 46
[$T_{1/2}$]
- 1971SA09 Saxena, R.N.; Sharma, H.D. Nucl. Phys. A171 (1971) 593
[I_β, I_γ]
- 1972AC02 Achterberg, E.; Iglesias, F.C.; Jech, A.E. Moragues, J.A.; Otero, D.; Pérez, M.L.; Proto, A.N.; Rossi, J.J.; Scheuer, W.; Suárez, J.F. Phys. Rev. C5 (1972) 1759
[α_K, E_γ , multipolarity]
- 1972BE90 Begzhanov, R.B.; Kobilov, R.B.; Sabirov, KH.S.; Salimov, S. SK.; Khudaibergenov, U.KH. Bull. Acad. Sci. USSR 36 (1972) 2190
[Multipolarities]
- 1972KR07 Krane, K.S.; Olsen, C.E.; Steyert, W.A. Phys. Rev. C5 (1972) 1671
[δ , mixing ratios]
- 1974KO26 Koene, B.K.S.; Lightart, H.; Postma, H. Nucl. Phys. A235 (1974) 267
[δ , mixing ratios]
- 1976FU06 Fuller, G.H. J.Phys.Chem.Ref.Data 5, 835 (1976)
[Nuclear spins and moments]
- 1976ME16 Meyer, R.A.; Momyer, F.F.; Henry, E.A.; Yaffe, R.P.; Walters, W.B. Phys. Rev. C14 (1976) 1152
[I_β, I_γ , decay scheme]
- 1977KR13 Krane, K.S. At. Data Nucl. Data Tables 19 (1977) 363
[Mixing ratios]
- 1983LO08 Lönnroth, T.; Kumpulainen, J.; Tuokko, C. Physica Scripta 27 (1983) 228
[J^π]
- 1989RA17 Raghavan, P. At. Data and Nucl. Data Tables 42 (1989) 189
[Nuclear moments]
- 1995RA12 Rab, S. Nucl. Data Sheets 75 (1995) 527
[Decay scheme]
- 1996SC06 Schönfeld, E., Janssen, H. Nucl. Instrum. Meth. A 369 (1996) 527
[Atomic data]
- 2002BA85 Band I.M., Trzhaskovskaya M.B., Nestor C.W. Jr. At. Data Nucl. Data Tables 81(2002) 1
[Theoretical ICC]
- 2003AU03 Audi, G.; Wapstra, A.H.; Thibault, C. Nucl. Phys. A 729 (2003) 337
[Q value]

Reference	g150,39	g176,97	g203,7	g245,95	g262,702	g267,173	g345,43	g361,09	g372,05	g381,59
1959HO97							-			
1966EI01					0,18 (5)					
1971SA09					5,0	1,5	3,0			
1976ME16	0,34 (7)	0,9 (2)	0,05	0,4 (1)	4,13 (7)	1,35 (6)	1,2 (2)	1,3 (4)	0,11 (6)	0,52 (5)
Recommended	0,34 (7)	0,9 (2)	0,05	0,4 (1)	4,13 (11)	1,35 (7)	1,2 (2)	1,3 (4)	0,11 (6)	0,52 (5)
1974KO26					0,35 (3)	0,10 (2)	0,06 (2)	0,16 (3)	0,16 (3)	
Absolute	0,029 (6)	0,078 (18)	0,00432 (8)	0,035 (9)	0,356 (12)	0,117 (7)	0,104 (18)	0,11 (4)	0,009 (6)	0,045 (5)

Reference	g386,85	g417,56	g422,901	g438,87	g510,530	g510,82	g522,40	g529,872	g537,73	g554,8
1959HO97								1000		
1966EI01			4,0 (10)		24,8 (37)			1000		
1971SA09		1,6	3,0		17 (4)					
1976ME16	0,68 (5)	1,77 (11)	3,58 (6)	0,46 (5)	21,0 (2)	< 0,1	< 1	1000 (4)	0,41 (8)	< 0,01
Recommended	0,68 (5)	1,77 (11)	3,58 (9)	0,46 (5)	21,0(5)	< 0,1	< 1	1000 (20)	0,41 (8)	< 0,01
1974KO26		0,12 (2)	0,26 (2)		1,85 (5)			87,7 (2)		
Absolute	0,059 (5)	0,153 (10)	0,309 (10)	0,040 (5)	1,81 (6)	0,004 (5)	0,04 (5)	86,3 (2)	0,035 (7)	0,0004 (5)

Reference	g556,17	g567,1	g617,974	g648,76	g670,10	g678,65	g680,247	g706,578	g768,382	g789,59
1959HO97								20		
1966EI01			3,0 (8)				10 (2)	17,3 (26)	5,9 (15)	
1971SA09			4,2				8,8	18	5,4	0,6
1976ME16	0,23 (3)	0,04 (3)	6,25 (6)	0,65 (15)	0,49 (6)	0,25 (8)	7,47 (9)	17,3 (2)	5,29 (9)	0,58 (4)
Recommended	0,23 (3)	0,04 (3)	6,25 (14)	0,65 (15)	0,49 (6)	0,25 (8)	7,47 (17)	17,3 (4)	5,29 (14)	0,58 (4)
1974KO26			0,53 (2)				0,61 (2)	1,47 (4)	0,43 (2)	0,04 (1)
Absolute	0,020 (3)	0,003 (3)	0,539 (15)	0,056 (13)	0,042 (6)	0,022 (7)	0,645 (19)	1,49 (4)	0,457 (15)	0,050 (4)

Reference	g _{820,506}	g _{856,278}	g _{875,329}	g _{909,67}	g _{911,49}	g _{1018,1}	g _{1035,58}	g _{1052,296}	g _{1060,07}	g _{1087,71}
1959HO97			90					10		
1966EI01	2,2 (6)	13,7 (21)	58 (5)	4 (1)				7,2 (18)	1,6 (4)	
1971SA09	2,0	14	52	4,4				5,7	1,0	
1976ME16	1,78 (6)	14,3 (4)	51,8 (2)	2,46 (7)	0,53 (7)	0,07 (3)	0,10 (2)	6,39 (7)	1,59 (6)	0,14 (2)
Recommended	1,78 (6)	14,3 (4)	51,8 (11)	2,46 (9)	0,53 (7)	0,07 (3)	0,10 (2)	6,39 (15)	1,59 (7)	0,14 (2)
1974KO26	0,15 (1)	1,18 (4)	4,42 (11)	0,25 (2)				0,54 (2)	0,14 (1)	
Absolute	0,154 (6)	1,23 (4)	4,47 (12)	0,212 (9)	0,046 (6)	0,006 (3)	0,0086 (18)	0,551 (16)	0,137 (7)	0,0121 (18)

Reference	g _{1236,441}	g _{1298,223}	g _{1327,2}	g _{1350,38}	g _{1386,15}	g _{1589,94}
1959HO97	20	40				
1966EI01	17,2 (26)	27,4 (41)		1,6 (4)		
1971SA09	18	25		1,8		0,5
1976ME16	17,3 (2)	27,0 (2)	< 0,005	1,72 (4)	0,10 (3)	0,034 (5)
Recommended	17,3 (4)	27,0 (6)		1,72 (5)	0,10 (3)	0,034 (5)
1974KO26	1,45 (4)	2,25 (6)		0,14 (1)		
Absolute	1,49 (4)	2,33 (7)	0,00022 (22)	0,148 (5)	0,0086 (26)	0,0029 (4)

The 1959HO97 values were reported to $I(529) = 100$. In the table they have been reported to 1000 for the $I(529)$.

1966EI01 did not observe the 744-keV level, so they reported $I(509,8) = 25\%$ for the γ -transition from the 1385- to the 875-keV levels instead for the 744-233 keV transition. The 1966EI01 values were reported to $I(529) = 100$. In the table they have been reported to 1000 for the $I(529)$. The uncertainty in the 1966EI01 values are estimated by the evaluator following the notes given by the authors: $\pm 8\%$ for relative intensities > 5 ; $\pm 15\%$ for relative intensities > 1 ; $\pm 25\%$ for relative intensities < 1 .

The 1971SA09 values were reported to $I(529) = 100$. In the table they have been reported to 1000 for the $I(529)$.

In 1974KO26 the absolute γ emission probabilities are given but the details of the measurements are absent.

For the relative γ intensities less than ($<$) a certain value, the adopted absolute value is the result given by GABS computer code.

**¹³³Xe - Comments on evaluation of decay data
by M. Galán**

1) Decay Scheme

¹³³Xe disintegrates by β^- emission to excited levels in ¹³³Cs.

¹³³Xe ground state has $J^\pi = 3/2^+$. The isomeric state is at 233 keV and has $J^\pi = 11/2^-$ (1989RA17).

2) Nuclear Data

The Q value is from AME2003 (2003AU03): $Q_{\beta^-} = 427,4$ (24) keV.

Level energies have been obtained from a least-squares fit to γ -ray energies (GTOL computer code). Spin and parities are from 1995RA12.

The half-life of the 81-keV level has been deduced (using the AveTool computer code) from the values reported in 1965GE14, 1963GO17, 1962TH12, 1959BO56, 1958AL98, 1955LE18 and 1953GR07. Half-lives for other levels are from 1995RA12.

The measured ¹³³Xe half-life values, in days, are:

Reference	Value (d)	Comments
2002UN02, 1992UN01	5,2475 (5)	
1975HO18	5,25 (2)	
1975WO10	5,250 (13)	
1974CA27	5,245 (6)	
1974FOZY	5,240 (6)	
1972EM01	5,29 (1)	Rejected by Chauvenet's criterion
1968AL16	5,312 (25)	Rejected by Chauvenet's criterion
1950MA15	5,270 (2)	Rejected by Chauvenet's criterion
	Mean	Reduced χ^2
LWM	5,2474 (5)	0,44
NRM	5,2474 (5)	0,44
RT	5,2474 (5)	

The AveTool computer code has been used with these seven input values. This code calculates averages using three statistical methods: LWM (Limitation of Relative Statistical Weight), NRM (Normalised Residual Method) and RT (Rajeval Technique).

The values in 1950MA15, 1968AL16, 1972EM01 were rejected based on the Chauvenet's criterion. For the remaining values, the largest contribution to the weighted average comes from the value of Unterweger (2002UN02). The LWM method increased the uncertainty of this value 3.895 times in order to reduce its relative weight to 50 %.

The recommended value is therefore the LWM mean, **5,2474 (5) d**. Its uncertainty has been expanded to 0,009 d, so the half-life range includes the most precise value of 5,2475 d (1992UN01, 2002UN02).

2.1) β Transitions

The energies of the β transitions have been deduced from the Q value and the level energies in ¹³³Cs, the later deduced from γ -ray transition energies. The adopted values have been verified against those produced by the computer code GTOL.

All beta transitions of ¹³³Xe are allowed. The β⁻ probabilities and associated uncertainties have been deduced from γ-ray transition intensity balance at each level of the decay scheme, assuming no β⁻ transition to the ground state.

$$\%b_{0,3} = P_{g+ce}(384) + P_{g+ce}(303) + P_{g+ce}(223) = 0,0029(4) + 0,0061(8) + 0,000187(69) = 0,0092(9)$$

$$\%b_{0,2} = P_{g+ce}(80) + P_{g+ce}(161) - P_{g+ce}(384) = 0,78(8) + 0,088(10) - 0,000187(69) = 0,87(8)$$

$$\%b_{0,1} = 100 - [\%b_{0,3} + \%b_{0,2}] = 100 - [0,0092(9) + 0,87(8)] = 99,12(8)$$

These values have been compared to the β⁻ emission probabilities measured by 1952BE55, 1961ER04 and 1986SC34. Also, the lg ft values have been calculated using the program LOGFT for allowed β⁻ transitions, and compared to values reported in these references.

Such a comparison is given in the following table:

Reference	%β _{0,1}	Lg ft	%β _{0,2}	Lg ft	%β _{0,3}	Lg ft
1959JH17	0,1	5,7	2	7	98	5,6
1961ER04	0,006	-	0,71	7,5	99,28	5,7
1986SC34	0,0073	-	0,79	-	99,2	-
Recommended	0,0092 (9)	6,84	0,87 (8)	7,31	99,12 (8)	5,62

2.2) g-ray Transitions

Transition Probabilities

The γ-ray transition probabilities have been calculated from the γ-ray emission probabilities using our recommended internal conversion coefficients.

Mixing ratios and internal conversion coefficients

For the 81, 223, 302 and 384 keV γ-ray transitions the adopted δ (mixing ratio) are from 1977KR13. The adopted values were deduced from angular correlation data. For the 80 and 161 γ-ray transitions the adopted δ values are from 1995RA12.

The internal conversion coefficients (ICC) have been calculated using the BrIcc computer code, which interpolated ICC values from tables of Band et al. (2002BA85). Associated uncertainties are 1,4 %.

3) Atomic Data

Atomic values (ω_K, ω_L and η_{KL}) are from 1996SC06.

ω _K	0,894 ± 0,004
ω _L	0,104 ± 0,005
η _{KL}	0,895 ± 0,004

The X-ray and Auger electron emission probabilities have been deduced from γ-ray and conversion electron data by using the computer code EMISSION. Results were verified with the RADLST computer code. Differences between these results were < 1 %.

4) Electron Emissions

The conversion electron emission probabilities have been computed from γ -ray emission probabilities and theoretical ICC values.

5) Photon Emissions

Energies

γ -ray energies and uncertainties are from 2000HE14. These values have been deduced on a revised energy scale.

g-ray emissions

The available experimental relative gamma emission intensities are:

Reference	g79,6	g81	g161	g223	g303	g384
1958PL55	-	-	-	-	0,010	0,005
1959JH17	-	100	1,4	-	0,084	0,043
1961ER04	0,8 (1)	100	0,109 (10)	0,0004 (⁺⁴ ₋₃)	0,0123 (12)	0,0062 (9)
1968AL16	100 1,6 (7)	98,2 (59)	0,174 (9)	0,000647 (613)	0,0135 (4)	0,00618 (19)
1992MA05	100		0,242 (25)	0,00044 (18)	0,0193 (7)	0,000901 (41)
Weighted average			0,182	0,00046	0,0155	
Reduced χ^2			6,55	0,1	24	
Internal uncertainty			0,008	0,00017	0,0004	
External uncertainty			0,022	0,00006	0,0021	
Recommended	0,76 (9)	99,24 (9)	0,182 (22)	0,00046 (17)	0,0155 (21)	0,0076 (10)

1968AL16 relative intensities were reported to the group $\gamma_{79,6} + \gamma_{81} = 1000$. In the table they have been reported to 100 for that of the group $\gamma_{79,6} + \gamma_{81}$.

1995MA02 relative intensities were reported to the group $\gamma_{80} + \gamma_{81}$ and multiplied 10^{-5} . In this table they have been reported to 100 for that of group $\gamma_{79,6} + \gamma_{81}$.

To evaluate all relative intensities, the group $\gamma_{79,6} + \gamma_{81}$ has been taken as the reference line as measured 1968AL16 and 1992MA05.

The 79.6 keV line has been deduced using the ratio $\gamma_{79,6}/\gamma_{161}$ from ¹³³Ba decay (Chechev and Kuzmenko, 2004).

$$g_{79,6} = 0,182(22) \times \frac{4,27(8)}{1,028(8)} \Big|_{^{133}\text{Ba}} = 0,76(9)$$

$$\text{Therefore, } g_{81} = 100 - 0,76(9) = 99,24(9)$$

The relative γ -ray emission intensities for the 384 keV γ -ray has been deduced from the 303 keV γ -ray emission probability and the averaged ratio $\gamma_{384}/\gamma_{303}$ measured by:

Reference	$\gamma_{384}/\gamma_{303}$
1958PL55	0,50 (11)
1959JH17	0,512 (13)
1961ER041	0,504 (88)
1968AL16	0,458 (20)
1992MA05	0,467 (27)
Weighted mean	0,492
Reduced χ^2	1,53
Internal uncertainty	0,010
External uncertainty	0,012
Recommended	0,492 (12)

So that, $g_{384} = g_{303} \times \frac{g_{384}}{g_{303}} \Big|_{w.m.} = 0,0155(21) \times 0,492(12) = 0,0076(10)$

The normalization factor has been deduced from the decay scheme using the formulas:

$$N = \frac{100}{\sum_i I_{g_i} [1 + a_{T_i}]} \quad \text{and} \quad dN^2 = \sum_i \left(\frac{\partial N}{\partial I_{g_i}} dI_{g_i} \right)^2 + \sum_i \left(\frac{\partial N}{\partial a_{T_i}} da_{T_i} \right)^2,$$

where the sum is over all γ -ray transitions to the ground state (g.s.), thus considering no direct β^- feeding to the g.s. Therefore:

$$N = \frac{100}{99,24(9) \times [1 + 1,698(24)] + 0,182(22) \times [1 + 0,294(5)] + 0,0076(10) \times [1 + 0,0202(3)]}$$

The deduced normalization factor is 0,373 (3).

Additional reference:

F. Lagoutine, Table des Radionucléides, CEA-LMRI (1984).

References

- 1940WU05 Wu, C.S. Phys. Rev. 58 (1940) 926
[Production modes]
- 1941CL02 Clancy, E.P. Phys. Rev. 60 (1941) 87
[Production modes]
- 1945WU05 Wu, C.S.; Segré, E. Phys. Rev. 67 (1945) 142
[Production modes]
- 1950MA01 Macnamara, J.; Collins, C.B.; Thode, H.G. Phys. Rev. 78 (1950) 129
[$T_{1/2}$]
- 1952BE55 Bergström, I. Ark. Fysik 5 (1952) 191
[I_γ , I_β]

- 1953GR07 Graham, R.L.; Bell, R.E. Canadian J. Phys. 31 (1953) 377
[T_{1/2} level, α_K, K/L ratio]
- 1954BE36 Bergström, I.; Thulin, S.; Wapstra, A.H.; Aström, B. Ark. Fysik 7 (1954) 255
[α_K, K/L ratio]
- 1955LE18 Lehmann, P.; Miller, J. Comp. Rend. 240 (1955) 1525
[T_{1/2} level]
- 1958AL98 Alväger, T.; Johansson, B.; Zuk, W. Ark. Fysik. 14 (1958) 373
[T_{1/2} level]
- 1959BO56 Bodenstedt, E.; Körner, H.J.; Matthias, E. Nucl. Phys. 11 (1959) 584
[T_{1/2} level]
- 1961ER04 Erman, P.; Sujkowsky, Z. Ark. Fysik, 20 (1961) 209
[I_γ]
- 1962TH12 Thieberger, P. Ark. Fysik. 22 (1962) 127
[T_{1/2} level]
- 1963GO17 Govil, I.M.; Khurana, C.S.; Hans, H.S. Nucl. Phys. 45 (1963) 60
[T_{1/2} level]
- 1965GE04 Geiger, J.S.; Graham, R.L.; Bergström, I.; Brown, F. Nucl. Phys. 68 (1965) 352
[T_{1/2} level]
- 1966TH09 Thun, J.E.; Töknkvist, S.; Nielsen, K.B.; Snellman, H.; Falk, F.; Mocoroa, A. Nucl. Phys. 88 (1966) 289
[δ, mixing ratios]
- 1968AL16 Alexander, P.; Lau, J.P. Nucl. Phys. A121 (1968) 612
[T_{1/2}, I_γ]
- 1972EM01 Emery, J.F.; Reynolds, S.A.; Wyatt, E.I. Nucl. Sci. Eng. 48 (1972) 319
[T_{1/2}]
- 1974CA27 Cavallo, L.M.; Schima, F.J.; Unterweger, M.P. Phys. Rev. C10 (1974) 2631
[T_{1/2}]
- 1974FOZY Fontanilla, J.; Prindle, A.L.; Landrum, J.H.; Meyer, R.A. Bull. Amer. Phys. Soc. 19 (1974) 501
[T_{1/2}]
- 1975HO18 Hoffman, D.C.; Barnes, J.W.; Dropesky, B.J.; Lawrence, F.O.; Kelly, G.M.; Ott, M.A. J. Inorg. Nucl. Chem. 37 (1975) 2336
[T_{1/2}]
- 1975WO10 Woods, M.J.; Goodier, I.W.; Lucas, Sylvia E.M. Int. J. Appl. Radiat. Isot. 26 (1975) 485
[T_{1/2}]
- 1977KR13 Krane, K.S. At. Data Nucl. Data Tables 19 (1977) 363
[Mixing ratios]
- 1989RA17 Raghavan, P. At. Data and Nucl. Data Tables 42 (1989) 189
[Nuclear moments]
- 1992MA05 Martin, R.H.; Keller, N.A. Int. J. Appl. Radiat. Isot. 43 (1992) 463
[I_γ]
- 1992UN01 Unterweger, M.P.; Hoppes, D.D.; Schima, F.J. Nucl. Inst. Meth. A312 (1992) 349
[T_{1/2}]
- 1995RA12 Rab, S. Nucl. Data Sheets 75 (1995) 491
[Decay scheme]
- 1996SC06 Scöfnfeld, E.; Janssen, H.. Nucl. Instrum. Meth. A 369 (1996) 527
[atomic data]
- 2000HE14 Helmer, R.G.; van der Leun, C. Appl. Radiat. Isot. 52 (2000) 601
[γ-ray energies]
- 2002BA85 Band, I.M.; Trzhaskovskaya, M.B. Nestor, C.W. Jr. At. Data Nucl. Data Tables 81 (2002) 1
[ICC]
- 2002UN02 Unterweger, M.P. Nucl. Inst. Meth. A56 (2002) 125
[T_{1/2}]
- 2003AU03 Audi, G.; Wapstra, A.H.; Thibault, C. Nucl. Phys. A 729 (2003) 337
[Q value]

**¹³³Xe^m - Comments on evaluation of decay data
by M. Galán**

1) Decay Scheme

¹³³Xe^m disintegrates by a strong converted γ -transition to the ground state of ¹³³Xe.

2) Nuclear Data

The 233-keV isomeric state has $J^\pi = 11/2^-$ (1989RA17).

The measured ¹³³Xe^m half-life values, are:

Reference	Value (days)
1975HO18	2,19 (5)
1974FOZY	2,188 (8)
1968AL16	2,191 (29)
1961ER04	2,26 (2)
1951BE11	2,30 (8)
Number of input values	5
Reduced χ^2	3,22
Weighted Mean	2,198
Internal uncertainty	0,007
External uncertainty	0,013
NRM	2,200 (11)
RT	2,191 (8)
Adopted value	2,198 (13)

The AveTool program has been used with these five input values. This program calculates averages using three statistical methods: LWM (Limitation of Relative Statistical Weight), NRM (Normalised Residual Method) and RT (Rajeval Technique).

The recommended value for the ¹³³Xe^m half-life is the LWM mean of 2,198 d with an external uncertainty of 0,013 d.

2.1) Gamma-ray Transitions

The evaluated γ -ray transition energy is the photon energy plus the nuclear recoil energy.

The 233-keV γ -ray has an M4 multipolarity. The various theoretical conversion coefficients for this transition (Band *et al.* Häger and Seltzer, Rösel *et al.*) differ about 2 % from each other. The ICCs (α_T , α_K , α_L) have been interpolated from the new Band *et al.* tables (2002BA85) using the BrIcc Computer Code. The uncertainties on these conversion coefficients are estimated to be 1,4 %.

Some experimental values together with the theoretical values are shown in the table:

Reference	α_K	K/L+M
Experimental		
1954BE55	4,4 (14)	2,32 (15)
1968AL16	7,68 (25)	2,04 (12)
1972AC02	7,4 (14)	2,54 (20)
Theoretical		
1968HA52	6,37 (9)	2,51 (5)
1978RO22	6,35 (9)	2,44 (4)
2002BA85	6,25 (9)	2,41 (3)

3) Atomic Data

Atomic values (ω_K , $\bar{\omega}_L$ and η_{KL}) are from 1996SC06.

ω_K	$0,888 \pm 0,005$
$\bar{\omega}_L$	$0,097 \pm 0,005$
η_{KL}	$0,902 \pm 0,004$

The X-ray and Auger electron emission probabilities have been calculated from γ -ray and conversion electron data using the programs RADLST and EMISSION. Differences between these results were $< 0,6 \%$.

4) Radiation emissions

4.1 Conversion electrons

The conversion electron emission probabilities have been deduced from the ICC values and from the γ -ray emission probability.

The total conversion electron emission probability has been deduced from:

$$P_{ce} = 100 - P_\gamma = 100 - 10,16 (13) = 89,84 (13)$$

4.2 g-Ray Emissions

Various measurements of the γ -ray energy have been found in the bibliography:

Reference	Value (keV)
1976ME16	233,221 (15)
1972AC02	233,2 (4)
1952BE55	232,8 (3)
1951BE11	232,8 (4)
Number of input values	4

Reduced χ^2	1,02
Weighted Mean	233,219
Internal uncertainty	0,015
External uncertainty	0,015
NRM	233,219 (15)
RT	233,11 (12)
Adopted value	233,219 (15)

The recommended value is the LWM mean of 233,219 keV with an external uncertainty of 0,015.

The γ -ray emission intensity is given by:

$$P\gamma = 100 / (1 + \alpha) = 100 / [1 + 8,84 (13)] = 10,16 (13) \%$$

Additional reference:

F. Lagoutine, Table des Radionucléides, CEA-LMRI (1984)

References

- 1951BE11 Bergström, I. Phys. Rev. 81 (1951) 638
[$T_{1/2}$, E_γ , K/L ratio]
- 1952BE55 Bergström, I. Ark. Fysik 5 (1952) 191
[$T_{1/2}$, E_γ]
- 1954BE36 Bergström, I.; Thulin, S.; Wapstra, A.H.; Aström, B. Ark. Fysik 7 (1954) 255
[α_K , K/L ratio]
- 1961ER04 Erman, P.; Sujkowsky, Z. Ark. Fysik 20 (1961) 209
[$T_{1/2}$]
- 1968AL16 Alexander, P.; Lau, J.P. Nucl. Phys. A121 (1968) 612
[$T_{1/2}$, α_K]
- 1968HA52 Hager, R.S., Seltzer, E.C. Nucl. Data A4 (1968) 1
[Theoretical ICC]
- 1969FR09 Fransson, K.J.; Erman, P. Ark. Fysik 39 (1969) 7
[M4]
- 1972AC02 Achterberg, E.; Iglesias, F.C.; Jech, A.E. Moragues, J.A.; Otero, D.; Pérez, M.L.; Proto, A.N.; Rossi, J.J.; Scheuer, W.; Suárez, J.F. Phys. Rev. C5 (1972) 1759
[α_K , E_γ , multipolarity]
- 1974FOZY Fontanilla, J.; Prindle, A.L.; Landrum, J.H.; Meyer, R.A. Bu ll. Amer. Phys. Soc. 19 (1974) 501
[$T_{1/2}$]
- 1975HO18 Hoffman, D.C.; Barnes, J.W.; Dropesky, B.J.; Lawrence, F.O.; Kelly, G.M.; Ott, M.A. J. Inorg. Nucl. Chem. 37 (1975) 2336
[$T_{1/2}$]
- 1976ME16 Meyer, R.A.; Momyer, F.F.; Henry, E.A.; Yaffe, R.P.; Walters, W.B. Phys. Rev. C14 (1976) 1152
[E_γ]
- 1978RO22 Rosel, F., Fries, H.M., Alder, K. At. Data Nucl. Data Tables 21 (1978) 91
[Theoretical ICC]

- 1989RA17 Raghavan, P. At. Data and Nucl. Data Tables 42 (1989) 189
[Nuclear moments]
- 1995RA12 Rab, S. Nucl. Data Sheets 75 (1995) 491.
[Decay scheme]
- 1996SC06 Schönfeld, E. ; Janssen, H. Nucl. Instrum. Meth. A 369 (1996) 527
[Atomic data]
- 2002BA85 Band, I.M.; Trzhaskovskaya, M.B. Nestor, C.W. Jr. At. Data Nucl. Data Tables 81
(2002) 1
[Theoretical ICC]

¹³³Ba - Comments on evaluation of decay data

by V.P. Chechev and N.K. Kuzmenko

The first DDEP evaluation of ¹³³Ba decay data was completed by these authors in January 2004 (2004BeZR). The current updated evaluation was completed in May 2015 with a literature cut-off by the same date. The main changes compared to the initial evaluation are mainly due to new publications: 2012Wa38 (Q-value), 2008Ki07 (theoretical internal conversion coefficients), 2014Manohar (relative gamma and conversion electron probabilities) and 2010Sc08, 2012Fi12, 2014Un01 (half-life measurements and corrections).

1. DECAY SCHEME

¹³³Ba decays primarily by allowed electron capture (ϵ) branches to the $1/2^+$ and $3/2^+$ ¹³³Cs levels at 437 and 383 keV. As to the intensities of the other possible ϵ branches to the ¹³³Cs ground state and levels at 81 and 161 keV they can be estimated from $\log ft$ systematics. From that of 1998Si17, one expects the $\log ft$ of the unique 2nd forbidden decay to the ground state to be greater than 13.9 which corresponds to a branch of less than 0.0005%. Similarly, the $\log ft$ of the 2nd forbidden decays to the 81- and 161-keV levels are expected to be greater than 10.6 which corresponds to branches of less than 0.7% and 0.3%, respectively. Our evaluations for each of these branches obtained from the intensity balance at the above levels agree very well with this expectation (see section 2.1). Therefore, all of these unobserved β branches can be assumed as upper limits from the $\log ft$ systematics.

For comparison see also the evaluations made by Yu. Khazov, A. Rodionov, and F.G. Kondev (2011Kh02), Shaheen Rab (1995Ra12) and A.L. Nichols (1993Nichols) as well as the analyses made by R.B. Firestone (1990Fi03) and F.E. Chukreev (1992Chukreev).

2. NUCLEAR DATA

Q⁻ value is from 2012 mass evaluation by Wang *et al.* (2012Wa38).

The recommended half-life of ¹³³Ba is based on the experimental results given in Table 1.

Table 1. Experimental values of the ¹³³Ba half-life (in days)

N	Author(s) and year	Reference	T_{1/2} (d)	Method and comments
1	Wyatt <i>et al.</i> (1961)	1961Wy01	3908 (73)	γ-ray scintillation spectroscopy and a beta proportional counter;
2	Lagoutine <i>et al.</i> (1968)	1968La10	2849 (37)	Ionization chamber, γ-ray scintillation spectroscopy and 4πβ- proportional counter; <i>omitted</i> because of the large deviation from the mean value
3	Reynolds <i>et al.</i> (1968)	1968Re04	3894 (44)	γ-ray scintillation spectroscopy and a beta proportional counter
4	Walz <i>et al.</i> (1970)	1970Wa19	3781 (15)	4πγ ionization chamber; <i>omitted</i> as superseded by 12
5	Emery <i>et al.</i> (1972)	1972Em01	3981 (37)	γ-ray scintillation spectroscopy and a beta proportional counter; <i>omitted</i> on the Chauvenet's criterion
6	Lloyd <i>et al.</i> (1973)	1973Ll01	4127 (260)	<i>omitted</i> on the Chauvenet's criterion
7	Hansen <i>et al.</i> (1979)	1979HaYC	3850 (55)	Ge(Li), NaI(Tl), Si(Li)
8	Rutledge and Mouchel (1980)	1980RuZY	3785 (27)	4πγ ionization chamber, 4πβ-γ coincidence
9	Houtermans <i>et al.</i> (1980)	1980Ho17	3848.0 (11)	4πγ pressurized ionization chamber
10	Hoppes <i>et al.</i> (1982)	1982HoZJ	3828 (11)	4πγ ionization chamber; <i>omitted</i> as superseded by 19
11	Kits <i>et al.</i> (1983)	1983Ki08	3885.9 (43)	4π ionization chamber
12	Walz <i>et al.</i> (1983)	1983Wa26	3842 (18)	4πγ ionization chamber
13	Unterweger <i>et al.</i> (1992)	1992Un01	3853.6 (36)	4πγ ionization chamber; <i>omitted</i> as superseded by 19
14	Martin <i>et al.</i> (1979)	1997Ma75	3848.9 (7)	4πγ ionization chamber
15	Unterweger <i>et al.</i> (2002)	2002Un02	3854.7 (28)	4πγ ionization chamber; <i>omitted</i> as superseded by 19
16	Schrader (2004)	2004Sc04	3840.5 (65)	Ionization chamber; <i>omitted</i> as superseded by 17
17	Schrader (2010)	2010Sc08	3840.3 (89)	Ionization-chamber
18	Fitzgerald (2012)	2012Fi12	3854.7 (28)	4πγ ionization chamber; <i>omitted</i> as superseded by 19
19	Unterweger and Fitzgerald (2014)	2014Un01	3830 (7)	4πγ ionization chamber
Recommended value (days)			3849.3 (23)	LWM
Recommended value (years)			10.539 (6)	

The values before 1961 were omitted due to their large uncertainties (more than 1 year).

The values of 1970Wa19, 1982HoZJ, 1992Un01, 2002Un02 and 2012Fi12 were omitted as they were replaced by later results from the same group.

Also the value of 1968La10 was omitted on statistical considerations because of a great contribution into the χ^2 value (27 σ from adopted value).

The uncertainty of 1997Ma75 was increased to 0.98 days to adjust weights according to the Limitation of Relative Statistical Weight method (LWM). In consequence the LWIGHT program chose the weighted average of 3849.3 days and external uncertainty of 2.3 days. The internal uncertainty is 0.74 d and the smallest experimental uncertainty is 0.7 d. The ratio of the reduced $\chi^2 / (\chi^2)_{\text{crit}}$ is 10.0/2.4.

It should be noted that in the weighted average of the two values of 1980Ho17 and 1997Ma75 have altogether 95% of the relative weight. Since these two values agree, any weighted average will be about 3849 days that differs slightly from an unweighted average of about 3853 days.

The recommended value of ¹³³Ba half-life is **3849.3 (23) days**. It can be compared with earlier evaluations of 3849.7 (22) days (2004BeZR) and 3853.7 (40) days (2011Kh02).

2.1. Electron Capture Transitions

The energies of the electron capture, ϵ , transitions have been calculated from the Q value and the level energies which have been deduced from gamma transition energies (see also 2011Kh02: Adopted Levels, Gammas). The level spins, half-lives and parities of ¹³³Cs (Table 2) are taken from 2011Kh02 (Adopted Levels).

Table 2. ¹³³Cs levels populated in ¹³³Ba ϵ -decay

Level	Energy (keV)	Spin and parity	Half-life	P ϵ (%) – obtained from intensity balance	P ϵ (%) - adopted
0	0	7/2+	stable	0.0000 (5)	< 0.0005
1	80.9979 (11)	5/2+	6.283 (14) ns	0.08 (55)	< 0.7
2	160.6121 (16)	5/2+	172 (4) ps	0.04 (59)	< 0.3
3	383.8491 (12)	3/2+	44 (11) ps	14.46 (51)	14.5 (5)
4	437.0113 (13)	1/2+	≤ 150 ps	85.41 (53)	85.4 (5)

The electron capture probabilities P $\epsilon_{0,4}$ and P $\epsilon_{0,3}$ were calculated from the intensity balance for the 437 level and the 384 level, respectively, using the evaluated P γ_{+ce} values.

Similarly, the electron capture probabilities $P_{\epsilon_{0,2}}$ and $P_{\epsilon_{0,1}}$ are obtained from the intensity balance for 161-keV and 81-keV levels, respectively, as 0.08 (55) and 0.04 (59) per 100 disintegrations. From here the upper limit of 0.63 per 100 disintegrations for each of these transitions agree with theoretical estimations, and the evaluators adopted for their probabilities upper limits of theoretical estimations: $P_{\epsilon_{0,2}} < 0.3\%$ and $P_{\epsilon_{0,1}} < 0.7\%$. Similarly, $P_{\epsilon_{0,0}} < 0.0005\%$.

The P_K , P_L and P_M values for $\epsilon_{0,4}$ and $\epsilon_{0,3}$ transitions have been computed using the code LOGFT (<http://www.nndc.bnl.gov/logft/>).

The available experimental P_K values are presented in **Table 3**.

Table 3. The experimental and adopted P_K values in ¹³³Ba ϵ -decay.

	$P_K(\epsilon_{0,4})$	$P_K(\epsilon_{0,3})$
1968Na16	0.68 (5)	
1972Sc08	0.72 (4)	0.80 (7)
1974Da09	0.76 (6)	0.87 (14)
1975Ni07	0.75 (10)	
1983Si17	0.75 (4)	0.80 (4)
1983Si22	0.71 (11)	0.79 (5)
1988BeYQ	0.78 (4)	
1990Da11	0.76 (4)	
1990Bh01	0.730 (12)	0.81 (3)
1992Sa28	0.65 (3)	0.74 (4)
adopted	0.671 (5)	0.7727 (9)

Most of these values were obtained in 1974-1990 using the method of the X-, gamma-ray sum peak measurements. The measurement results exceed the adopted theoretical P_K values for the allowed $\epsilon_{0,4}$, $\epsilon_{0,3}$ - transitions and depend on conversion coefficients α_K and fluorescence yields ω_K used in those works.

2.2. Gamma Transitions and Internal Conversion Coefficients

The evaluated energies of gamma transitions are the energies of gamma rays with adding the recoil energy.

The multipolarities and mixing ratios δ are taken from the adopted data in the ENSDF evaluation (2011Kh02). Gamma-ray transition probabilities $P(\gamma+ce)$ except for the γ -ray of 81 keV ($\gamma_{1,0}$) have been deduced from their gamma-ray emission probabilities and the total ICC(s). The adopted ICC(s) are obtained from the theoretical values by using the adopted mixing ratios δ and the BrIcc computer program (<http://bricc.anu.edu.au>), accepting the “frozen orbital (no hole)” approximation (2008Ki07).

$P(\gamma+ce)$ of the of 81 keV γ -ray transition ($\gamma_{1,0}$) was calculated in order to reach the ground state gamma-ray transition intensity balance:

$$P(\gamma+ce) (\gamma_{81}) = 100\% - P(\gamma+ce) (\gamma_{161}) - P(\gamma+ce) (\gamma_{383}) = 90.05 (6) \%$$

3. ATOMIC DATA

The atomic data (fluorescence yields, X-ray energies and relative probabilities, and Auger electron energies and relative probabilities) have been deduced using the SAISINUC software.

4. ELECTRON EMISSIONS

The energies of the conversion electrons were obtained from the gamma-ray transition energies and the electron binding energies. The absolute emission probabilities of the conversion electrons were deduced using recommended $P\gamma$ and ICC values. The absolute emission probabilities of K and L Auger electrons were calculated using the EMISSION computer program.

5. PHOTON EMISSIONS

5.1. X-ray Emissions

The X-ray energies are based on the wavelengths in the compilation of 1967Be65 (Bearden).

The absolute emission probabilities of Cs KX- and LX- rays were calculated using the EMISSION computer program.

The measured values of the total absolute emission probability of KX-rays ($P_{\text{XK}} \times 100$) are given below in comparison with the calculated (adopted) value:

1972Sc08	1977Sc31	1989Egorov	Adopted
123.1 (17)	117.4 (22)	119.7 (11)	118.9 (13)

5.2 Gamma-Ray Emissions

The γ -ray energies are taken from the evaluation 2000He14 where the values are deduced on the revised energy scale. For the γ -ray of 81 keV see also the measurement of 1991We08.

The γ -ray absolute emission probabilities have been computed using the evaluated γ -ray relative emission probabilities and the absolute emission probability of the 356-keV γ -ray of 0.6205 (19) measured in 1980Chauvenet, 1983Ch11. This experimental value for the most intensive γ -ray in the decay of ¹³³Ba was obtained as a result of the international intercomparison ICRM-S-6 (1980Chauvenet). It is more preferable for normalizing of gamma-ray absolute emission probabilities than having been obtained from the ground state intensity balance 0.613 (34) – because of big uncertainties in multipolarity admixtures (and thus in ICC) as well as possible ambiguity in determination of some spins (see 1992Chukreev).

At the same time the relative gamma ray emission probabilities from ICRM-S-6 measured at the fifteen laboratories are used below in Table 1 equally with other measurements for averaging all the available data. The measurements of ICRM-S-6 have been lettered CRP and deduced from absolute emission probabilities published in 1980Chauvenet excluding an activity uncertainties ~ 0.2%.

Table 5. The experimental and adopted relative gamma ray emission probabilities.

	γ_{53}	γ_{80}	γ_{81}	γ_{161}	γ_{223}	γ_{276}	γ_{303}	γ_{356}	γ_{383}
1967Bi15	3.8 (8)	3.8 (4)	53 (4)	1.1 (3)	0.7 (3)	11.0 (7)†	30 (2)	100	14.5 (1)
1968Al16	3.3 (5)	-	-	1.20 (6)†	0.74 (6)	12.0 (4)†	30.6 (9)†	100	14.2 (5)
1968Bo04	4.2 (2)†	4.0 (4)	58.2 (15)	1.07 (5)	0.78 (6)	11.8 (3)	29.8 (8)	100	14.3 (10)
1968Do10	3.2 (4)	5.5 (7)†	52 (7)	0.99 (10)	0.72 (8)	11.6 (8)	29.4 (2)	100	14.3 (10)
1968No01	3.78 (9)	4.9 (6)	60 (7)	1.21 (5)†	0.80 (3)†	11.61 (17)	29.75 (29)	100	14.18 (26)
1969Gu15	2.91 (5)	4.54 (7)	53.7 (17)	1.13 (15)	-	11.2 (3)	29.3 (5)	100	14.03 (26)†
1972Sc08	3.54 (5)	3.9 (2)	52.6 (10)	1.16 (5)	0.74 (4)	11.4 (3)	30.2 (6)	100	14.4 (3)
1973In06	-	-	-	0.98 (7)	0.76 (5)	11.6 (5)	29.6 (11)	100	14.9 (6)†
1973Legrand	-	3.7 (4)	56 (6)	1.4 (2)†	0.66 (2)†	11.35 (25)	29.4 (6)	100	14.3 (3)
1973Mc18	-	-	-	-	-	11.43 (23)	29.3 (6)	100	14.5 (3)
1977Ge12	3.0 (4)	5.6 (15)†	52 (4)	1.12 (8)	0.85 (7)†	11.7 (8)	29.87 (21)	100	14.4 (11)
1977Sc31	3.49 (8)	4.29 (12)	55.8 (16)	0.97 (3)	0.73 (3)	11.41 (16)	29.4 (3)	100	14.33 (21)
1978He21	3.54 (18)	3.1 (3)†	49.2 (26)	1.08 (4)	0.745 (25)	11.7 (4)	29.8 (4)	100	14.36 (20)
1978Vylov	3.57 (12)	4.16 (18)	54.6 (17)	0.98 (8)	0.71 (4)	11.4 (3)	28.8 (8)†	100	14.3 (5)
1980Ro22	-	-	-	1.03 (7)	0.72 (5)	11.69 (16)	29.9 (4)	100	14.79 (27)†
1983Yo03	-	-	-	1.035 (28)	0.756 (16)	11.57 (7)	29.55 (18)	100	14.36 (9)
1987Lakshmi	2.96 (9)	4.67 (14)	55.3 (16)	-	-	-	-	100	-
1989Da11	3.6 (5)	3.7 (5)	52.3 (7)	1.032 (10)	0.713 (8)	11.51 (8)	29.51 (23)	100	13.99 (9)†
1990Me15	3.48 (7)	3.77 (9)	51.2 (4)	1.05 (3)	0.71 (2)	11.3 (2)	29.2 (3)	100	14.5 (2)
1998Hw07	-	-	-	0.950 (18)	0.715 (10)	11.64 (13)	29.31 (40)	100	14.52 (17)
CRP-1	-	-	-	1.11 (9)	0.85 (5)†	11.7 (4)	29.9 (11)	100	14.5 (5)
CRP-2	3.56 (14)	-	53.1 (19)	0.99 (4)	0.729 (28)	11.7 (3)	30.1 (9)	100	14.4 (5)
CRP-3	3.53 (8)	4.20 (12)	54.8 (12)	1.031 (24)	0.69 (3)	11.51 (14)	29.5 (3)	100	14.37 (16)
CRP-4	3.53 (7)	4.18 (11)	54.6 (12)	1.037 (20)	0.730 (22)	11.48 (14)	29.5 (4)	100	14.41 (16)
CRP-5	3.9 (7)	4.00 (15)	51.5 (19)	1.020 (27)	0.728 (22)	11.5 (3)	29.5 (9)	100	14.2 (5)
CRP-6	3.45 (8)	4.73 (12)	57.6 (14)	1.020 (25)	0.728 (18)	11.68 (28)	29.7 (7)	100	14.5 (4)
CRP-7	3.56 (8)	4.73 (12)	58.9 (15)	1.070 (27)	0.738 (18)	11.50 (28)	29.6 (7)	100	14.3 (4)
CRP-8	-	-	-	-	-	11.22 (27)	29.3 (6)	100	14.53 (28)
CRP-9	-	-	-	-	-	11.22 (24)	29.3 (5)	100	14.26 (25)
CRP-10	-	-	-	-	-	11.48 (25)	29.3 (5)	100	14.20 (22)
CRP-11	-	-	-	-	-	11.57 (19)	29.4 (4)	100	14.34 (26)
CRP-12	3.69 (18)	4.37 (16)	55.3 (18)	1.050 (19)	0.741 (15)	11.53 (16)	29.5 (4)	100	14.36 (20)
CRP-13	2.92 (16)	-	-	-	0.75 (3)	11.9 (4)	30.2 (11)	100	14.6 (5)
CRP-14	3.53 (8)	4.39 (11)	55.9 (12)	1.015 (20)	0.735 (10)	11.61 (13)	29.6 (4)	100	14.34 (18)

	γ_{53}	γ_{80}	γ_{81}	γ_{161}	γ_{223}	γ_{276}	γ_{303}	γ_{356}	γ_{383}
CRP-15	3.36 (18)	-	-	1.05 (4)	0.758 (28)	11.7 (5)	29.6 (10)	100	14.3 (4)
CRP-16	3.26 (17)	-	-	1.05 (4)	0.764 (26)	11.7 (4)	29.7 (6)	100	14.3 (3)
CRP-19	3.53 (5)	-	-	1.063 (17)	0.725 (17)	11.61 (12)	29.7 (3)	100	14.53 (13)
CRP-20	3.53 (6)	4.05 (8)	55.1 (9)	1.05 (5)	0.72 (4)	11.49 (21)	29.4 (6)	100	14.51 (22)
CRP-21	3.62 (6)	4.15 (12)	55.8 (9)	1.039 (15)	0.705 (11)	11.57 (17)	29.5 (4)	100	14.40 (20)
2014Manohar ^{††}	3.505 (50)	4.05 (7)	52.44 (98)	0.94 (3)	0.72 (2)	11.21 (7)	29.17 (18)	100	14.39 (9)
Number of input values	28	21	25	30	29	37	37		35
Reduced χ^2	7.0/1.7	5.6/1.9	4.0/1.8	1.9/1.7	0.8/1.7	0.9/1.6	0.4/1.6		0.2/1.6
Weighted average	3.454	4.241	53.36	1.029	0.725	11.497	29.51		14.40
Internal uncertainty	0.016	0.027	0.23	0.0047	0.0035	0.028	0.061		0.034
External uncertainty	0.042	0.064	0.45	0.0066	0.0030	0.026	0.038		0.015
Adopted value (LWM)	3.45 (9)	4.24 (30)	53.7 (5)^{†††}	1.029 (10)	0.725 (8)	11.49 (7)	29.51 (18)	100	14.40 (9)

[†] Omitted as outliers

^{††} None quantitative estimates of uncertainty *components* (uncertainty budget) are given in 2014Manohar for measured relative gamma-ray intensities. These uncertainties seem to be too underestimated. The reviewers adopted for them the smallest uncertainties from all previous measurements for each corresponding gamma-ray.

^{†††} Adopted value has been changed slightly from the weighted average for a precise ground state intensity balance to get. Such a small change only for one gamma-ray supports the adopted experimental value of 62.05 (19) % for the 356 keV γ -ray absolute emission probability and confirms the decay scheme. The adopted uncertainty of 0.5 is external.

In addition, in 1996Mi26 a special precise measurements of the absolute emission probabilities only for the two weak 161 and 223 keV gamma-rays were made by using a $4\pi\beta(\text{ppc})-\gamma(\text{HPGe})$ coincidence system. The adopted relative emission probabilities of these gamma-rays (1.029 (10) and 0.725 (8)) can be compared with relative emission probabilities (1.028 (8) and 0.730 (5), respectively) calculated from the absolute values (0.6383 (38) % and 0.4530 (31) %, respectively) from 1996Mi26.

6. ENERGY CONSERVATION

The total average energy of 516.8 (36) keV, for one disintegration, calculated from the current evaluated data corresponds well to the available energy of 517.3 (10) keV (Q^+) from the mass tables (2012Wa38) confirming the consistency of the decay scheme and the reliability of this evaluation.

7. REFERENCES

- 1961Wy01** Wyatt E.I., Reynolds S.A., Handley T.H., Lyon W.S., Parker H.A. (1961): Half-Lives Radionuclides. II. Nucl. Sci. Eng. 11, 74.
[Half-life]
- 1967Be65** Bearden, J.A. (1967) X-Ray Wavelengths, Rev. Mod. Phys. 39, 78
[X-ray energies]
- 1967Bl15** Blasi P., Bocciolini M., Maurenzig P.R., Sona P., Taccetti N. (1967): The Decay of ¹³³Ba and Nuclear Transitions in ¹³³Cs. Nuovo Cimento 50B, 298.
[γ -ray emission probabilities]
- 1968Al16** Alexander P., Lau J.P. (1968): Nuclear Structure in ^{133,135}Xe and ^{133,135}Cs. , Nucl. Phys. A121, 612.
[γ -ray emission probabilities, ICC(s)]
- 1968BO04** Bosch H.E., Haverfield A.J., Szichman E., Abecasis S.M. (1968): High-Resolution Studies in the Decay of ¹³³Ba with Semiconductor Counters Nucl. Phys. A108, 209.
[α_K (81 keV)]
- 1968Do10** Donnelly D.P., Reidy J.J., Wiedenbeck M.L. (1968): High-Resolution Gamma-Ray Spectroscopic Study of the Decay ¹³³Ba \rightarrow ¹³³Cs. Phys. Rev. 173, 1192.
[γ -ray emission probabilities]
- 1968La10** Lagoutine F., Le Gallic Y., Legrand J. (1968): Détermination Précise de Quelques Périodes Radioactives. Intern. J. Appl. Radiat. Isotop. 19, 475.
[Half-life]
- 1968Na16** Narang V., Houtermans H. (1968): The P(L)/P(K) Capture in ¹³³Ba. In: Proc. Conf. Electron Capture and Higher Order Processes in Nucl. Decays, Debrecen, Hungary, D. Berenyi, Ed., Eotvos Lorand Phys Soc., Budapest, p.97.
[L/K-capture ratio]
- 1968No01** Notea A., Gurfinkel Y. (1968): Transitions in ¹³³Cs from the Decay of ¹³³Ba, Nucl. Phys. A107, 193.
[γ -ray emission probabilities]
- 1968Re04** Reynolds S.A., Emery J.F., Wyatt E.I. (1968): Half-Lives of Radionuclides - III. Nucl. Sci. Eng. 32, 46.
[Half-life]
- 1969Gu15** Gunnink R., Niday J.B., Anderson R.P., Meyer R.A. (1969): Gamma-Ray Energies and Intensities. In: UCID-15439, and Gunnink, R., Nethaway, D. – Priv. Comm. (Jan 1969)
[γ -ray emission probabilities]
- 1970Wa19** Walz K.F., Weiss H.M. (1970): Messung der Halbwertszeiten von ⁶⁰Co, ¹³⁷Cs and ¹³³Ba. Z. Naturforsch. 25a, 921.
[Half-life]
- 1972Em01** Emery J.F., Reynolds S.A., Wyatt E.I., Gleason G.I. (1972): Half-Lives of Radionuclides - IV. Nucl. Sci. Eng. 48, 319.
[Half-life]

- 1972Sc08** Schmidt-Ott W.-D., Fink R.W. (1972): The Determination by an Independent Method of P(K) Electron Capture Probabilities in ¹³³Ba and ¹³⁹Ce Decays. Gamma Decay of ¹³³Ba. Z. Phys. 249, 286.
[K-capture probability, absolute XK emission probability, γ -ray emission probabilities, ICC]
- 1973In06** Inoue H., Yoshizawa Y., Morii T. (1973): Gamma-Ray Energies and Relative Intensities of ⁷⁵Se, ^{108m}Ag, ¹¹³Sn, ¹³¹I and ¹³³Ba. J. Phys. Soc. Japan 34, 1437.
[γ -ray emission probabilities]
- 1973Legrand** Legrand J. (1973): Calibration of γ -spectrometers. Nucl. Instrum. Methods 112, 229
[γ -ray emission probabilities]
- 1973LI01** Lloyd R.D., Mays C.W. (1973): A Note on the Half-Period of ¹³³Ba. Intern. J. Appl. Radiat. Isotop. 24, 189.
[Half-life]
- 1973Mc18** McNelles L.A., Campbell J.L. (1973): Absolute Efficiency Calibration of Coaxial Ge(Li) Detectors for the Energy Range 160-1330 keV. Nucl. Instrum. Methods 109, 241.
[γ -ray emission probabilities]
- 1974Da09** Mahapatra B.K. Das, Mukherjee P. (1974): K-Capture Probability in the Decay of ¹³³Ba from X-Ray-Gamma-Ray Summing in Ge(Li) Detectors. J. Phys. (London) A7, 388.
[K-capture probability]
- 1975Ni07** Nicaise W.F., Waltner A.W. (1975): A Single Detector Method for the Determination of P(K-1) in ¹³³Ba. Nucl. Instrum. Methods 131, 477.
[K-capture probability]
- 1977Ge12** Gehrke R.J., Helmer R.G., Greenwood R.C. (1977): Precise Relative gamma-Ray Intensities for Calibration of Ge Semiconductor Detectors. Nucl. Instrum. Methods 147, 405.
[γ -ray emission probabilities]
- 1977Sc31** Schötzig U., Debertain K., Walz K.F. (1977): Standardization and Decay Data of ¹³³Ba. Intern. J. Appl. Radiat. Isotop. 28, 503.
[XK-ray and γ -ray emission probabilities]
- 1978He21** Helmer R.G., Greenwood R.C., Gehrke R.J. (1978): Reevaluation of Precise gamma-Ray Energies for Calibration of Ge(Li) Spectrometers. Nucl. Instrum. Methods 155, 189.
[γ -ray emission probabilities]
- 1978Vylov** Vylov C., Osipenko B.P., Chumin V.G. (1978) In: Elementarnie chastitsi and atomnie yadra (Particles & Nuclei) 9, issue 6, p.1350 (in Russian).
[γ -ray emission probabilities]
- 1979HaYC** Hansen H.H., Mouchel D. (1979): Studies on the Decay of ¹³³Ba. In: NEANDC(E) 202U, Vol III, p.28.
[Half-life]
- 1980Chauvenet** Chauvenet B., Morel J., Legrand J. (1980): Report ICRM-S-6 (December 1980).
[Absolute γ -ray emission probabilities]
- 1980Ho17** Houtermans H., Milosevic O., Reichel F. (1980): Half-lives of 35 Radionuclides. Intern. J. Appl. Radiat. Isotop. 31, 153.
[Half-life]
- 1980Ro22** Roney W.M., Jr, Seale W.A. (1980): Gamma Ray Intensity Standards for Calibrating Ge(Li) Detectors for the Energy Range 200-1700 keV. Nucl. Instrum. Methods 171, 389.
[γ -ray emission probabilities]

- 1980RuZY** Rutledge A.R., Smith L.V., Merritt J.S. (1980): Decay Data for Radionuclides used for the Calibration of X- and gamma-Ray Spectrometers. In: AECL 6692.
[Half-life]
- 1982HoZJ** Hoppes D.D., Hutchinson J.M.R., Schima F.J., Unterweger M.P. (1982): Nuclear Data for X- or Gamma-Ray Spectrometer Efficiency Calibrations. In: NBS-SP-626, p.85.
[Half-life]
- 1983Ch11** Chauvenet B., Morel J., Legrand J. (1983): An International Intercomparison of Photon Emission-rate Measurements of X- and gamma-Rays Emitted in the Decay of ¹³³Ba. Intern. J. Appl. Radiat. Isotop. 34, 479.
[Absolute γ -ray emission probabilities]
- 1983Ki08** Kits J., Latal F., Choc M. (1983): The Half-Life of ¹³³Ba. Intern. J. Appl. Radiat. Isotop. 34, 935.
[Half-life]
- 1983Si17** Singh K., Sahota H.S. (1983): K-Capture Probabilities in the Decay of ¹³³Ba. J. Phys. Soc. Japan 52, 2336.
[K-capture probability]
- 1983Si22** Singh K., Sahota H.S. (1983): A New Approach to K-Electron-Capture Probabilities to the 437 and 384 keV Levels in the Decay of ¹³³Ba. J. Phys. (London) G9, 1565
[K-capture probability]
- 1983Wa26** Walz K.F., Debertain K., Schrader H. (1983): Half-Life Measurements at the PTB. Intern. J. Appl. Radiat. Isotop. 34, 1191.
[Half-life]
- 1983Yo03** Yoshizawa Y., Iwata Y., Katoh T., Ruan J.-Z., Kawada Y. (1983): Precision Measurements of Gamma-Ray Intensities IV. Low Energy Region: ⁷⁵Se and ¹³³Ba. Nucl. Instrum. Methods 212, 249.
[γ -ray emission probabilities]
- 1987Lakshmi** Lakshmi A.S.V. Subba, Venkataranam S., Reddy, S.B., Reddy, K.V. (1987): Relative intensities of the low energy gamma transitions in the decay of Ba-133. Curr. Sci. 56, 407.
[γ -ray relative emission probabilities]
- 1988BeYQ** Begzhanov R.B., Azimov K.Sh., Magrupov R.D., Mirakhmedov Sh.A., Mukhammadiev A., Narzikulov M., Salimov S.Kh. (1988): K-Capture Probabilities in ¹³³Ba Decay. In: Program and Theses, Proc. 38th Ann. Conf. Nucl. Spectrosc. Struct. At Nuclei, Baku, p 93.
[K-capture probabilities]
- 1989DA11** Danilenko V.N., Konstantinov A.A., Kurenkov N.V., Kurchatova L.N., Malinin A.B., Mamelin A.V., Matveev S.V., Sazonova T.E., Stepano E.K., Sepman S.V., Toporov Yu.G. (1989): Methods of Producing Radionuclides for Spectrometric Gamma-Ray Sources and Their Standardization - 1. Barium-133. Appl. Radiat. Isot. 40, 707.
[γ -ray emission probabilities]
- 1989Egorov** Egorov A.G., Egorov Yu.S., Nedovesov V.G., Shchukin G.E., Yakovlev K.P. (1989): In: Program and Thesis, Proc. 39th Ann. Conf. Nucl. Spectrosc. Struct. At Nuclei, Leningrad, p.505.
[X K-ray emission probabilities]
- 1989Ma60** Martins M.C., Marques M.I., Parente F., Ferreira J.G. (1989): Some K X-Ray Relative Transition Probabilities for Z=47, 49, 52, 55 and 56. J. Phys. (London) B22, 3167.
[K β '₂/K β '₁ ratio]

- 1990Bh01** Rao K. Bhaskara, Rao S.S., Rao V.S., Padhi H.C. (1990): K-Capture Probabilities in the Decay of ¹³³Ba. *Nuovo Cimento* 103A, 683.
[K-capture probability]
- 1990Da11** Dasmahapatra B., Bhattacharya S., Sen S., Saha M., Goswami A. (1990): Accurate Measurement of P(K) for the 437 keV State of ¹³³Cs in the Decay of ¹³³Ba (10.5y). *J. Phys. (London)* G16, 1227.
[K-capture probability]
- 1990Fi03** Firestone R.B. (1990): Analysis of alpha-, beta-, and gamma-Ray Emission Probabilities. *Nucl. Instrum. Methods Phys. Res.* A286, 584.
[A fully constrained analysis of the emission probabilities in the decay of ¹³³Ba]
- 1990Me15** Meyer, R.A. (1990): Multigamma-Ray Calibration Standards. *Fizika (Zagreb)* 22, 153.
[γ-ray emission probabilities]
- 1991We08** Wesselborg C., Alburger D.E. (1991): Precision Energy Measurements of Gamma Rays from ⁴⁴Ti and ¹³³Ba. *Nucl. Instrum. Methods Phys. Res.* A302, 89.
[γ-ray energies]
- 1992Chukreev** Chukreev F.E. (1992): Regarding to Selection of Radioactive Sources Problem for Gamma-ray Spectrometer Calibration. In: *Voprosi Atomnoi Nauki i Tekhniki, Ser.: Yadernie konstanti 2*, p.92 (in Russian).
[Decay scheme]
- 1992Sa28** Sahota G.P.S., Singh H., Binarh H.S., Pallah B.S., Sahota H.S. (1992): Sum Peak Comparison Measurement of K-Capture Probabilities to the Levels of ¹³³Cs. *J. Phys. Soc. Japan* 61, 3518.
[K-capture probability]
- 1992Un01** Unterweger M.P., Hoppes D.D., Schima F.J. (1992): New and Revised Half-Life Measurements Results. *Nucl. Instrum. Methods Phys. Res.* A312, 349.
[Half-life]
- 1993Nichols** Nichols A.L. (1993): AEA Technology Report AEA- RS-5449.
[Decay Scheme]
- 1995Ra12** Rab, S. (1995): Nuclear Data Sheets Update for A = 133, *Nucl. Data Sheets* 75, 491
[Decay scheme]
- 1996Mi26** Miyahara H., Usami K., Mori C. (1996): Precise Measurement of the Emission Probabilities for the Weak 161 and 223 keV Gamma-Rays of ¹³³Ba. *Nucl. Instrum. Methods Phys. Res.* A374, 193.
[γ-ray emission probabilities]
- 1996Sc06** Schönfeld E., Janßen H. (1996): *Nucl. Instrum. Methods Phys. Res.* A369, 527.
[Atomic Data]
- 1997Ma75** Martin R.H., Burns K.I.W., Taylor J.G.V. (1997): A Measurement of the Half-Lives of ⁵⁴Mn, ⁵⁷Co, ⁵⁹Fe, ⁸⁸Y, ⁹⁵Nb, ¹⁰⁹Cd, ¹³³Ba, ¹³⁴Cs, ¹⁴⁴Ce, ¹⁵²Eu. *Nucl. Instrum. Methods Phys. Res.* A390, 267.
[Half-life]
- 1998Hw07** Hwang H.Y., Lee C.B., Park T.S. (1998): *Appl. Radiat. Isot.* 49, 1201.
[γ-ray emission probabilities]
- 2000He14** Helmer R.G., van der Leun C. (2000): *Nucl. Instrum. Methods Phys. Res.* A450, 35.
[γ-ray energies]
- 2002Un02** Unterweger M.P. (2002): Half-Life Measurements at the National Institute of Standards and Technology. *Appl. Radiat. Isot.* 56, 125.
[Half-life]
- 2004BeZQ** Bé M.M., Chisté V., Dulieu C., Browne E., Chechev V., Kuzmenko N, Helmer R., Nichols A., Schönfeld E., Dersch R. (2004): In: *Monographie BIPM-5, Vol. 1*, p.263. Bureau International des Poids et Mesures.
[Previous ¹³³Ba decay data evaluation]

- 2004Sc04** Schrader H. (2004): Half-life measurements with ionization chambers - A study of Systematic effects and results Appl. Radiat. Isot. 60, 17.
[Half-life]
- 2008Ki07** Kibédi T., Burrows T.W., Trzhaskovskaya M.B., Davidson P.M., Nestor C.W. Jr. (2008): Evaluation of theoretical conversion coefficients using BrIcc. Nucl. Instrum. Methods Phys. Res. A589, 202.
[BrIcc computer program]
- 2010Sc08** Schrader H. (2010): Half-life measurements of long-lived radionuclides - New data analysis and systematic effects Appl. Radiat. Isot. 68, 1583.
[Half-life]
- 2011Kh02** Khazov Yu., Rodionov A., Kondev F.G. (2011): Nuclear Data Sheets for A = 133, Nucl. Data Sheets 112, 855.
[Decay scheme]
- 2012Fi12** Fitzgerald R., (2012): J. Res. Natl. Inst. Stand. Technol. 117, 80.
[Half-life]
- 2012Wa38** Wang M., Audi G., Wapstra A.H., Kondev F.G., MacCormick M., Xu X., Pfeiffer B., (2012): The AME2012 atomic mass evaluation, Chin. Phys. C36, 1603.
[Q]
- 2014Un01** Unterweger M.P., Fitzgerald R., (2014): Update of NIST half-life results corrected for ionization chamber source-holder instability. Appl. Radiat. Isot. 87, 92.
[Half-life]
- 2014Manohar** Manohar G.V. Sai, Yadav K.S., Sai K.V, Gowrishankar R., Venkataramaniah K., Deepa S., Rao D., (2014): Gamma and conversion electron measurements in ¹³³Cs. In: Proceedings of the DAE Symp. on Nucl. Phys. 59, p.286.
[γ-ray and electron relative emission probabilities]

**¹³⁴Cs - Comments on evaluation of decay data
by M.-M. Bé**

This evaluation was completed in February 2012.

1 Decay Scheme

¹³⁴Cs decays by β^- emission (99.9997 %) to excited levels of ¹³⁴Ba. A very weak electron capture (EC) branch to the 847-keV level of ¹³⁴Xe (0.0003 %) has been pointed out by Van Hise *et al.* (1975Va12).

The overall consistency of the decay scheme was checked by calculating the total energy carried away by the various emissions as determined below, it was found to be 2059 (1) keV per disintegration when the available energy is 2058.97 (33) keV.

2 Nuclear Data

Q values are from Audi (2011AuZZ).

The spins and parities are from A. A. Sonzogni (2004So32).

The ¹³⁴Cs half-life values are summarized in the tables below.

The results were converted from $a \Leftrightarrow d$, with $1 a = 365.242 198 78 d$.

Published measured values not used in the evaluation because less precise or superseded (as a rule only one result per laboratory is considered in the statistical process):

Reference	$T_{1/2}$ (a)	Remarks
1938Alexeeva	≥ 1	
1951Glendenin	2.3 (3)	
1957Geiger	2.07 (2)	
1957Meritt	2.19 (2)	
1958Bayly	2.15 (+8, -4)	
1958Edwards	2.26 (5)	
1961Wyatt	2.07 (2)	
1963Dietz	2.046 (4)	Superseded by 1973Di01
1965Flynn	1.99 (2)	
1978Bulovic	2.04 (3)	
	$T_{1/2}$ (d)	
1982HoZJ	754.19 (15)	Superseded by 2002Un02
1992Un01	753.88 (11)	Superseded by 2002Un02

Measured values used in the evaluation:

Reference	$T_{1/2}$ (d)	Remarks
1972La14	751.7 (15)	2.058 (12) a ; 3 σ uncertainty
1973Di01	753.1 (44)	2.062(5) a ; 3 σ uncertainty divided by 3
1980RuZY	753.78 (30)	
1980Ho17	754.50 (7)	Unrealistic uncertainty
1997Ma75	754.52 (18)	
2002Un02	753.88 (15)	Uc questionable
χ^2 crit.	0.3	
$\chi^2 / (n-1)$	0.4	
UWM	753.96	
WM	754.29	
Adopted	754.0 (5)	Or 2.0644 (14) a

In this set of data it is difficult to assess an uncertainty.

In Lagoutine *et al.*, they usually determined the uncertainty as: $\sigma = 3 \times u_{\text{stat}} + u_{\text{sys}}$

Where u_{stat} is the statistical component and u_{sys} the systematic component, their uncertainty cannot be simply divided by 3.

The uncertainty claimed by Houtermans (1980Ho17) is manifestly unrealistic.

The uncertainty given by Unterweger (2002Un02) was recently questioned (2012Un**).

Hence, the adopted value is the simple mean with an uncertainty which covers the most precise value.

2.1 β^- and Electron Capture transitions

The energies of β^- transitions were deduced from the Q values and the level scheme of ¹³⁴Ba.

The β^- transition probabilities were calculated from the gamma transition probability balance at each level of the decay-scheme. They are compared below with experimental results.

Comparison of adopted and measured values, as published in the two latest publications, of the β^- intensities, in %.

Ref	89-keV	415-keV	658-keV	891-keV	1454-keV
1968Hs01	27 (2)	3.0 (5)	70 (2)	0.045 (15)	0.008 (4)
1963Va06	28	1	71	0.045	0.005
Adopted	27.27 (3)	2.498 (8)	70.19 (8)	~0	0.06 (6)

A weak electron capture branch with 0.0003 % probability was observed by Van Hise *et al.* (1975Va12) from the measurement of a gamma ray with energy 847 keV, however this ray was not confirmed in later works, especially in the very precise measurements carried out by Miyahara (2002Mi06).

2.2 γ -ray Transitions

The γ -ray transitions probabilities were deduced from the γ -ray emission intensities and the internal conversion coefficients calculated with the BrIcc program v2.3-2011 (2008Ki07) for the “frozen approximation”.

The multipolarities are from Chand *et al.* (1990Ch47) who measured the γ - γ directional correlations for seven cascades in ¹³⁴Ba.

Comparison between calculated and some measured K conversion coefficients, $\times 10^{-3}$:

Ref	242-keV	475-keV	563-keV	569-keV	604-keV	795-keV
1965Br02		9.4 (1)	5.6 (6)	8.2 (9)	4.85 (20)	2.5 (3)
1968Ab01			6	7.5		2.9
1968Na11		9.84 (208)	6.40 (73)	8.80 (99)	5.07 (50)	2.90 (33)
1990Ch47	71 (24)	9.34 (72)	6.05 (45)	7.93 (42)	5.03	2.71 (10)
1998Ga24						2.59 (5)
Adopted	72.2 (12)	9.6 (4)	6.03(9)	8.05 (12)	5.03 (7)	2.58 (4)

Ref	801-keV	1038-keV	1167-keV	1365-keV
1965Br02	2.6 (4)	1.62 (18)	1.05 (10)	0.72 (7)
1968Ab01	2.4			
1968Na11	2.78 (58)	1.68 (20)	1.05 (13)	0.79 (9)
1990Ch47	2.49 (12)	1.74 (9)	1.11 (6)	0.855 (32)
1990Ma29		1.515 (20)		0.842 (18)
1998Ga24	2.66 (8)			
Adopted	2.54 (4)	1.79 (6)	1.122 (16)	0.820 (12)

3 Atomic Data

Atomic values for ω_K , ω_L , and η_{KL} , are from Schönfeld and Janssen (1996Sc06).

The X-ray and Auger electron emission intensities were derived from the decay scheme data, they are compared with the measured values of Chand (1988Ch44).

	Chand **	Adopted
K α	0.722 (15)	0.676 (6)
K' β 1	0.1386 (39)	0.1289 (19)
K' β 2	0.0312 (39)	0.0325 (8)

** Values converted with $I_{\gamma 604} = 97.63$ (8) %

4 Radiation Emissions

4.1 γ -ray emissions

The measured values of the gamma-ray energies are listed below. No significant discrepancies were observed. The adopted gamma-ray transition energies are the weighted means calculated using the Lweight program (version 3), values published in 1967Le** have been omitted because often not consistent with the others.

Measured and adopted energies of gamma-ray emissions, in keV.

Ref	242	326	475	563	569	604
1965Br02			475.26 (10)	563.11 (12)	569.24 (12)	604.64 (12)
1967Ra10	242.694 (41)	326.51 (10)	475.355 (38)	563.325 (41)	569.371 (47)	604.744 (27)
1968Ab01			475.2 (5)	563.2 (5)	569.38 (56)	604.67 (50)
1968Na11			475.57 (42) ^o	563.1 (5)	569.30 (51)	604.83 (54) ^o
1975Va12	242.89 (5)	326.45 (10)	475.35 (5)	563.26 (5)	569.29 (3)	604.660 (20)
1976Gr11				563.227 (15)	569.315 (15)	604.699 (15)
1985GoZK			475.365 (2)	563.250 (3)	569.333 (3)	604.721 (2)
1987Wa28	242.738 (8) ⁱ	326.589 (13)	475.364 (3)	563.240 (4)	569.328 (3)	604.720 (3)
χ^2 crit	4.6	4.6	3	2.6	2.6	2.8
χ^2 /n-1	4.9	1.2	0.3	1.5	0.8	2.1
WM	242.755	326.585	475.3646	563.2462	569.3301	604.7201
Adopted	242.76 (5)	326.585 (14)	475.365 (2)	563.246 (3)	569.330 (2)	604.720 (3)

Ref	795	801	1038	1167	1365
1965Br02	795.80 (16)	801.80 (16)	1038.46 (20)	1167.65 (25)	1364.97 (28)
1967Ra10	795.806 (50)	801.86 (28)	1038.61 (49)	1167.99 (39)	1365.08 (32)
1968Ab01	795.68 (49) ^o	801.54 (50) ^o	1038.17 (60)	1167.42 (50)	1364.93 (50)
1968Na11	796.02 (71) ^o	802.00 (71)	1038.02 (92)	1168.4 (10)	1365.4 (12) ^o
1975Va12	795.760 (20)	801.84 (3)	1038.50 (5)	1167.86 (6)	1365.13 (10)
1976Gr11	795.845 (22)	801.932 (22)	1038.571 (26)	1167.938 (26)	1365.152 (32)
1985GoZK	795.867 (4) ⁱ	801.956 (4)		1167.968 (5)	1365.200 (5)
1987Wa28	795.859 (5)	801.948 (5)	1038.610 (7)	1167.968 (5)	1365.185 (7)
χ^2 crit	3	2.8	2.8	2.6	2.8
χ^2 /n-1	5.8	2.8	1.3	1.1	1
WM	795.860	801.950	1038.605	1167.967	1365.1941
Adopted	795.86 (1)	801.950 (6)	1038.605 (8)	1167.967 (4)	1365.194 (4)

ⁱ Increased uncertainty ; ^o Outlier ; ^u unweighted mean

The measured relative γ ray intensities used for the statistical process are listed below. The different sets of data are consistent, then the adopted relative gamma-ray emission intensities are the weighted means, except as noted, calculated with the Lweight program (version 3). The intensity of 847-keV gamma-ray in ¹³⁴Xe (0.0003 %) and an upper intensity limit for the 232 keV gamma-ray of ¹³⁴Ba (0.0012 %) are from Van Hise *et al.* (1975Va12).

The normalization factor has been deduced from the decay scheme using the formulas:

$$N = \frac{100}{\sum_i I_{\gamma_i} [1 + \alpha_{T_i}]} \quad \text{and} \quad dN^2 = \sum_i \left(\frac{\partial N}{\partial I_{\gamma_i}} dI_{\gamma_i} \right)^2 + \sum_i \left(\frac{\partial N}{\partial \alpha_{T_i}} d\alpha_{T_i} \right)^2,$$

where the sum is over all γ -ray transitions to the ¹³⁴Ba ground state, thus considering no direct β^- feeding to the ground state.

Te relative emission intensities involved in these formulas are: the 847-, 1168- and 604-keV gamma-ray transitions, and α_{847} , α_{1168} and α_{1168} their internal conversion coefficients.

The calculated normalization factor is 0.9763 (8).

Then, the absolute intensity of the 604-keV γ ray is 97.63 (8) %, it can be compared with the experimental result of 97.65 (13) % obtained by Miyahara *et al.* (2002Mi06).

Measured and adopted relative emission intensities of gamma-ray emissions. The values are in %.

Ref	242-keV (4,3)	326-keV (5,4)	475-keV (4,2)	563-keV (2,1)	569-keV (5,3)	604-keV (1,0)
1962Ha10				8.0 (12)	12.0 (15) ^o	100
1965Br02			1.54 (15)	8.5 (8)	14.6 (14) ^o	100 (5)
1967Le**			1.53 (31)	8.2 (7)	15.2 (7)	100
1967Ra10	0.020 (10)	0.020 (10)	1.54 (16)	9.1 (9)	16.1 (11)	100
1968Ab01			1.43 (20) ^o	8.9 (10)	15.3 (16)	100
1968Na11			1.67 (11) ^o	8.8 (5)	13.6 (7) ^o	100 (3)
1970Ho06	0.0224 (20)		1.60 (8) ^o	9.0 (5)	16.3 (10)	100 (6)
1975Va12	0.0215 (8) ⁱ	0.015 (6)	1.50 (4)	8.59 (5)	15.82 (11)	100
1976De**			1.55 (3)	8.55 (12)	15.76 (23)	100
1980Yo05				8.57 (3)	15.78 (6)	100.0 (4)
1987Wa28	0.0322 (20)	0.0180 (15)	1.520 (10)	8.53 (6)	15.71 (10)	100.0 (7)
1988CH44	0.0294 (20)	0.0170 (17)	1.520 (20)	8.54 (7)	15.75 (3)	100.0 (7)
2002Mi06			1.503 (11)	8.530 (18)	15.728 (23)	100.00 (8)
χ^2 crit	3.3	3.8	2.6	2.2	2.4	
χ^2 /n-1	7.2	0.14	0.4	0.4	0.3	
UWM	0.0251	0.0175	1.525	8.600	15.745	
WM	0.0247	0.0175	1.515	8.544	15.741	
adopted	0.0247 (32)	0.0175 (11)	1.515 (7)	8.544 (14)	15.741 (17)	100.00 (8)

Ref	795-keV (3,1)	801-keV (5,2)	1038-keV (4,1)	1167-keV (2,0)	1365-keV (5,1)
1962Ha10					
1967Le**	87.3 (10)	8.9 (6)	1.12 (20) ^o	2.25 (20) ^o	3.37 (31)
1965Br02	90 (9)	9.0 (15)	1.06 (10)	1.99 (17)	3.5 (3)
1967Ra10	90 (7)	9.1 (8) ^o	1.04 (8)	2.00 (22)	3.3 (3)
1968Ab01	90 (9)	9.4 (10) ^o	1.1 (6) ^o	1.94 (20)	3.4 (3)
1968Na11	89 (4)	8.1 (4) ^o	1.06 (6)	2.06 (14)	3.55 (19)
1970Ho06	88 (4)	8.9 (4)	1.01 (6)	1.90 (10)	3.29 (17)
1975Va12	87.6 (4)	8.95 (4)	1.025 (10)	1.850 (27)	3.11 (4)
1976De**	87.4 (9)	8.85 (12)	1.023 (13)	1.84 (2)	3.09 (3)
1980Yo05	87.5 (3)	8.89 (3)	1.008 (5)	1.827 (8)	3.074 (13)
1987Wa28	87.5 (6)	8.97 (8)	1.016 (7)	1.841 (13)	3.109 (20)
1988CH44					
2002Mi06	87.54 (6)	8.898 (20)	1.021 (8)	1.834 (7)	3.094 (10)
χ^2 crit	2.3	2.6	2.5	2.4	2.3
χ^2 /n-1	0.05	0.4	0.6	0.7	1.4
UWM	88.35	8.920	1.029	1.908	3.262
WM	87.539	8.905	1.0150	1.834	3.092
adopted	87.54 (6)	8.905 (15)	1.0150 (33)	1.834 (5)	3.092 (8)

¹ – Increased uncertainty to reduce its weight to 50 %.

^o - Outlier

Omitted data in the statistical process:

- Ewan (1964Ew04), superseded by Brown (1965Br02);
- van Wijngaarden (1963Va06) because they were not able to separate the 563-569 keV lines and 795-801 keV lines;
- Bashandy (1966Ba57) because they are significantly discrepant with other results;
- Stelson (1973St14) because there are no details in the publication, the values are only mentioned in the decay scheme;
- Verhaeghe (1954Ve09) and Yamanoto (1960Ya**) given without uncertainties;
- Meyer (1990Me15), same as Van Hise (1975Va12).

4.2 Electron emissions

The conversion electron emission intensities have been obtained from the γ -ray emission intensities and theoretical ICC values.

5. References

- 1954Ve09 J.Verhaeghe, J.Demuynck. Compt.Rend. Ac. sciences 239 (1954) 1374
1960Ya** Y.Yamanoto. Thesis, Osaka University (1960)
1962Ha10 A. K. Hankla, J. H. Hamilton, R. V. Stockendal, Bull. Am. Phys. Soc. 7(1962)566
[Gamma-ray emission intensities]
1963Va06 W. van Wijngaarden, R.D. Connor. Can. J. Physics 42 (1963) 504
[Gamma-ray energies]
1965Br02 R. A. Brown, G. T. Ewan, Bull. Am. Phys. Soc. 10(1965)82
[Gamma-ray energies, Gamma-ray emission intensities]
1966Ba57 E.Bashandy, A.Abd El-Haliem. Atomkernenergie 11 (1966) 316
1967Le** J.Legrand, J.P.Boulanger. C.R. Acad. Sc. Paris, B265 (1967) 782
[Gamma-ray energies, Gamma-ray emission intensities]

- 1967Ra10 D. E. Raeside, J. J. Reidy, M. L. Wiedenbeck, Nucl. Phys. A98(1967)54
[Gamma-ray energies, Gamma-ray emission intensities]
- 1968Ab01 A. Abdul-Malek, R. A. Naumann, Nucl. Phys. A106(1968)225
[Gamma-ray energies, Gamma-ray emission intensities]
- 1968Hs01 S.T. Hsue, *et al.* Nucl. Phys. A109 (1968) 423
[Gamma-ray energies]
- 1968Na11 T. S. Nagpal, Can. J. Phys. 46(1968)2579
[Gamma-ray energies, Gamma-ray emission intensities]
- 1970Ho06 S. Hofmann, H. K. Walter, A. Weitsch, Z. Physik 230(1970)37
[Gamma-ray emission intensities]
- 1972La14 F. Lagoutine, J. Legrand, C. Perrot, J. P. Brethon, J. Morel, Int. J. Appl. Radiat. Isotop. 23(1972) 219
[Half-life]
- 1973Di01 L. A. Dietz, C. F. Pachucki, J. Inorg. Nucl. Chem. 35(1973)1769
[Half-life]
- 1973St14 P. H. Stelson, S. Raman, J. A. McNabb, R. W. Lide, C. R. Bingham, Phys. Rev. C8(1973)368
[Gamma-ray emission intensities]
- 1975Va12 J. R. Van Hise, D. C. Camp, R. A. Meyer, Z. Physik A274(1975)383
[Gamma-ray energies, Gamma-ray emission intensities]
- 1976De** K. Debertin, U. Schötzig, K.F. Walz. PTB Jaresberich (1976) 160
[Gamma-ray emission intensities]
- 1976Gr11 R. C. Greenwood, C. W. Reich, R. G. Helmer, R. J. Gehrke, R. A. Anderl, Phys. Rev. C14(1976)1906
[Gamma-ray energies.]
- 1980Ho17 H. Houtermans, O. Milosevic, F. Reichel, Appl. Rad. Isotopes 31(1980)153
[Half-life]
- 1980RuZY A. R. Rutledge, J. S. Merritt, L. V. Smith, AECL-6788(1980)45
[Half-life]
- 1980Yo05 Y. Yoshizawa, Y. Iwata, T. Kaku, T. Katoh, J. -Z. Ruan, T. Kojima, Y. Kawada, Nucl. Instrum. Methods Phys. Res. 174(1980)109
[Gamma-ray emission intensities]
- 1982HoZJ D. D. Hoppes, J. M. R. Huchinson, F. J. Schima, M. P. Unterweger, NBS – SP 626(1982)85
[Half-life]
- 1985GoZK V. Gorozhankin, *et al.* Report JINR P6 85.268 (1985)
[Gamma-ray energies]
- 1987Wa28 G. Wang, D. E. Alburger, E. K. Warburton, Nucl. Instrum. Methods Phys. Res. A260(1987)413
[Gamma-ray energies, Gamma-ray emission intensities]
- 1988Ch44 B. Chand, J. Goswamy, D. Mehta, S. Singh, M. L. Garg, N. Singh, P. N. Trehan, Nucl. Instrum. Methods Phys. Res. A273(1988)310
[Gamma-ray emission intensities]
- 1990Ch47 B. Chand, *et al.* Can. J. Phys. 68 (1990) 1479
[Multipolarities]
- 1990Ma29 N.M. Marchilashvili *et al.* Sov. J. Nucl. Phys. 51,1 (1990) 13
[Gamma-ray energies]
- 1992Un01 M. P. Unterweger, D. D. Hoppes, F. J. Schima, Nucl. Instrum. Meth. Phys. Res. A312(1992)349
[Half-life]
- 1996Sc06 E. Schönfeld *et al.* Nucl. Instrum. Meth. Phys. Res. A 369 (1996) 527
[Atomic Data]
- 1998Ga24 A.Gammal, *et al.* Phys. Rev. C58, 3 (1998)1829
[ICC]
- 1997Ma75 R. H. Martin, K. I. W. Burns, J. G. V. Taylor, Nucl. Instrum. Methods Phys. Res. A390(1997)267
[Half-life]

Comments on evaluation

- 2002Mi06 H. Miyahara, N. Hayashi, K. Fujiki, N. Takeuchi, Y. Hino, Appl. Rad. Isotopes 56(2002)131
[Gamma-ray emission intensities]
- 2002Un02 M. P. Unterweger, Appl. Rad. Isotopes 56(2002)125
[Half-life]
- 2004So32 A. A. Sonzogni, Nucl. Data Sheets 103(2004)1
[Spins, parity, multipolarity]
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr., Nucl. Instrum. Meth. Phys. Res. A589(2008)202
[Theoretical ICC]
- 2011AuZZ G. Audi, W. Meng, private communication
[Q]
- 2012Un** M. P. Unterweger, R. Fitzgerald. Appl. Rad. Isotopes 70,9 (2012) 1892
[Q]

¹³⁵Xe^m - Comments on evaluation of decay data
M. Galán

1) Decay Scheme

¹³⁵Xe^m disintegrates by IT (99,996 (2) %) to the ground state of ¹³⁵Xe and by β⁻ (0,004 (2) %) to ¹³⁵Cs excited levels. β⁻ branching has been reported by several authors: < 0,25 % (1976FE04); 0,004 % (1974MEZV and 1982WA21). 1974FOZY reported a transition from the 526 keV -level in ¹³⁵Xe to the 786,9 keV- level in ¹³⁵Cs with a $lg ft = 8,7$.

The β-decay scheme is that proposed by 1974MEZV (see also 2008SI01).

The ¹³⁵Xe^m isomeric state is at 526 keV and has $J_{\pi} = 11/2^{-}$ (1989RA17, 2008SI01).

2) Nuclear Data

$Q^{-}({}^{135}\text{Xe}^{\text{m}}) = 1692 (4) \text{ keV}$ has been deduced using a value of $Q({}^{135}\text{Xe}) = 1165 (4) \text{ keV}$ from 2003Au03.

The measured ¹³⁵Xe^m half-life values are:

Reference	Value (min)
1960AL12	15,8 (4)
1960KO02	15,65 (10)
1968AL16	15,2 (7)
1968TO20	15,4 (9)
1971HA13	15,287 (22)
1975FU12	15,29 (5)
Number of input values	6
Reduced χ^2	2,84
Weighted Mean	15,303
Internal uncertainty	0,020
External uncertainty	0,034
Adopted value	15,30 (3)

None of the values has been rejected by Chauvenet's criterion. The largest contribution to the weighted average comes from the value of Hawkins (1971HA13).

The recommended value for the ¹³⁵Xe^m half-life is the LWM mean of 15,30 with an external uncertainty of 0,03 d.

DECAY OF ¹³⁵Xe^m to ¹³⁵Xe**2.1) Gamma-ray Transition***Transition Energy*

The evaluated γ -ray transition energy is equal to the photon energy plus the nuclear recoil energy.

Isomeric Transition Probability

The 526-keV γ -ray has M4 multipolarity. The ICCs have been interpolated from the recent tables of Band *et al.* (2002BA85) using the BrIcc Computer Code. The uncertainties on these theoretical conversion coefficients (average deviations from the experimental values) are estimated to be 1,4 %.

Some experimental values (1960AL12, 1972AC02) together with the theoretical values (Band *et al.* 2002; Häger and Seltzer, 1968) are shown in the following table:

Reference	α_K	K/L
1960AL12	0,21 (5)	5,8 (11)
1972AC02	0,198 (12)	
1968HA52	0,193	
2002BA85	0,1908 (27)	5,25 (10)

A beta branching has been estimated as 0,004 (2) % (see below- **DECAY OF ¹³⁵Xe^m to ¹³⁵Cs**). Thus the recommended value of P(IT) is 99,996 (2) %.

3) Atomic Data

Atomic fluorescence yields (ω_K , ω_L and n_{KL}) are from 1996SC06

The X-ray and Auger electron emission probabilities have been calculated from γ -ray and conversion electron data using the EMISSION code.

4) Radiation emissions**4.1) Conversion electrons**

The conversion electron emission probabilities have been deduced from the ICC values and from the γ -ray emission probability.

The total conversion electron emission probability is:

$$P_{ce} = P(IT) - P_\gamma = 19,16 (25) \%$$

4.2) g-Ray Emission

Various measurements of the γ -ray energy found in the bibliography are given below:

Reference	Value (keV)
1960AL12	527,4 (8)
1960KO02	528 (3)
1972AC02	526,5 (3)
1979BO26	526,579 (7)
1982WA21	526,561 (7)
Number of input values	5
Reduced χ^2	3,32
Weighted Mean	526,570
Internal uncertainty	0,0050
External uncertainty	0,0054

The recommended value is the LWM mean of 526,570 keV with an external uncertainty of 0,005.

The absolute γ -ray emission probability is given by:

$$P_\gamma = 100 / (1 + \alpha_T) = 80,84 (20) \%$$

b⁻ DECAY OF ¹³⁵Xe^m to ¹³⁵Cs

2.1) Gamma-ray Transition

Transition Energy

The γ -ray transition energies are from 1974MEZV.

Mixing ratios and internal conversion coefficients

Neither mixing ratios nor internal conversion coefficients have been measured for these γ -ray transitions.

2.2) Gamma-ray Emission

γ -Ray Emission Probabilities

Only Meyer (1974) reported γ -ray intensities associated with a possible ¹³⁵Xe^m β -decay. The γ -ray relative intensities measured by 1974MEZV are those given in the following table (“?” purports “uncertain γ ”):

Transition energy (keV)	I_γ	Photons per 100 disint.
786,91	44 (22)	0,003 6 (18)
1133	3?	0,000 24
1192	0,4?	0,000 032
1358	2?	0,000 16

In the second column relative intensities I_γ are relative to 10^6 photons of 526 keV $-\gamma_{1,0}(\text{Xe})$ as reported in 1974MEZV. A 50 % uncertainty in $I_\gamma(787)$ has been assumed.

For the absolute γ intensities the total conversion coefficient of 0,237 (3) for the 526 keV transition has been taken into account. Then the absolute γ intensities are estimated by multiplying the relative intensities by 100/123,7.

2.3) b Transitions

The energies of the β⁻ transitions have been deduced from the Q value and the level energies in ¹³⁵Cs (2008Si01). The adopted values have been verified against those produced from a least -squares fit to gamma-ray energies by the computer code GTOL.

As no direct β⁻ transition to the ground state was reported by Meyer, the normalization factor was deduced assuming no feeding to the g.s. by using the equation:

$$[I\gamma(526) (1 + \alpha(526)) + I\gamma(787) (1 + \alpha(787))] N = 100 \%$$

The β⁻ emission probabilities in Sec. 2.1 are from the absolute gamma-ray emission probabilities, as given in the following table:

Transition	Energy (keV)	P(β) %	Log ft
β _{1,1}	905,1	0,003 6 (18)	8,7
β _{1,2}	559	0,000 24	9,2
β _{1,3}	500	0,000 032	9,9
β _{1,4}	334	0,000 16	8,7

Lg ft's were calculated with the LOGFT computer code. The adopted beta branching ratio is 0,004 (2) %.

The possible 1692-keV β transition

If there exists a beta transition to the ground state this might be a 1st forbidden unique transition. The lg ft value is > 8,5. Using the lg f tables of Gove and Martin (1971) or the LOGFT code, we have:

$$\lg f_i / f_0 = 0,935 \text{ and } \lg f_i = 3,35.$$

Now, $\lg(f_1 t) = \lg(f_1) + \lg(t)$ and $t = \frac{T_{1/2}(s)}{B.R.}$, with these two expressions we can estimate the β branching ratio.

$$\text{So, } \lg(t) > 8,5 - 3,35 = 5,15 \quad \longrightarrow \quad t > 1,42 \times 10^5$$

Finally we get, $B.R. < \frac{920}{1,42 \times 10^5} = 0,0065$ or $B.R. < 0,65 \%$ for the upper limit of the beta branching. If

we consider this beta feeding to the ground state, then the normalization factor can be estimated as:

$$[I\gamma(526) (1 + \alpha(526)) + I\gamma(787) (1 + \alpha(787))] N = 100 \% - 0,65 \%$$

Then the values would be:

$$P(IT) = 99,346 (2) \%$$

$$\beta = 0,0035 (18) \%$$

$$P_\gamma = 80,31 (20) \%$$

References

- 1960AL12** T. Alvager, Arkiv Fysik 17, 521 (1960)
[Half-life, γ -ray energy, α_K , K/L ratio]
- 1960KO02** K. Kotajima and H. Morinaga, Nuclear Phys. 16, 231 (1960)
[Half-life, γ -ray energy]
- 1968AL16** P. Alexander and J.P. Lau, Nucl.Phys. A121, 612 (1968)
[Half-life]
- 1968HA52** R.S. Hager and E.C. Seltzer, Nucl. Data A4 (1968) 1
[Theoretical ICC]
- 1968TO20** K. Tomura and N. Miyaji. Radiochim, Acta 10, 173 (1968)
[Half-life]
- 1971GO40** N.B. Gove and M. Martin, At. Data. Nucl. Data Tables A10 (1971) 205
[log f_t]
- 1971HA13** R.C. Hawkings, W.J. Edwards and W.J. Olmstead, Can. J. Phys. 49, 785 (1971)
[Half-life]
- 1972AC02** E. Achterberg, F.C. Iglesias, A.E. Jech, J.A. Moragues, D. Otero, M.L. Perez, A.N. Proto, J.J. Rossi, W. Scheuer and J.F. Suarez, Phys.Rev. C5, 1759 (1972)
[γ -ray energy, α_K , multipolarity]
- 1974MEZV** R.A. Meyer (private communication)
[γ -ray energy, γ probabilities, β probabilities]
- 1974FOZY** J. Fontanilla, A.L. Prindle, J.h. Landrum and R.A. Meyer, Bull. Am. Phys. Soc. 19 (1974) 501
[log f_t]
- 1975FU12** T. Fukuda and S. Omori, J. At. Energy Soc.Jap. 17, 177 (1975)
[Half-life]
- 1976FE04** H. Feuerstein and J. Oschinski, Inorg.Nucl.Chem.Lett. 12, 243 (1976)
[β probabilities]
- 1979BO26** H.G. Börner, W.F. Davidson, J. Almeida, J. Blachot and J.A. Pinston, P.H.M. Van Assche, Nucl. Instr. Meth. 164, 579 (1979)
[γ -ray energy]
- 1982WA21** W.B. Walters, S.M. Lane, N.L. Smith, R.J. Nagle and R.A. Meyer, Phys. Rev. C26, 2273 (1982)
[γ -ray energy, β probabilities]
- 1989RA17** P. Raghavan, At. Data and Nucl. Data Tables 42 (1989) 189
[Nuclear moments]
- 1996SC06** E.Schönfeld and H. Janssen, Nucl. Instrum. Meth. A 369 (1996) 527
[Atomic data]
- 2002BA85** I.M. Band, M.B. Trzhaskovskaya and C.W. Nestor, Jr, At. Data Nucl. Data Tables 81 (2002) 1
[Theoretical ICC]
- 2003AU03** G. Audi, A.H. Wapstra and C. Thibault, Nucl. Phys. A729, 337 (2003)
[Q value]
- 2008SI01** B. Singh, A.A. Rodionov and Y.L. Khazov, Nuclear Data Sheets 109, 517 (2008)
[Decay scheme, level energies, gamma-ray multiplicities]

¹³⁷Cs - Comments on evaluation of decay data by R.G. Helmer and V.P. Chechev

This evaluation was completed by R.G. Helmer in September 1996 with minor editing done in February 1998. Updating ¹³⁷Cs half-life and editing were done by V.P. Chechev in February 2006. The literature available by February 2006 was included.

1 Decay Scheme

There are as many as 2 supposed excited levels in ¹³⁷Ba below the decay energy that have not been reported in the ¹³⁷Cs decay and observed only in ¹³⁶Ba(d, p)-reaction (1997Tu04 evaluation). Since the possible 907 and 1044 levels do not have J^π assignments, and the de-exciting γ rays have not been reported, arguments about their feeding can not be made.

The decay scheme is internally consistent and essentially complete since the total decay energy computed by RADLIST is 1174 (3) keV compared to the Q value of 1175.63 (17) keV, a difference of 1.8 (28) keV.

The J^π values and half-lives of the excited levels in ¹³⁷Ba are from the evaluation of 1997Tu04.

2 Nuclear Data

Q value is from 2003Au03.

The experimental ¹³⁷Cs half-life values available are, in days (values published in years have been converted to days):

12053 (1096)	1951FIAA,	omitted from analysis
10957 (146)	1955Br06,	omitted from analysis
9715 (146)	1955Wi21,	omitted from analysis
10446 (+73-37)	1958MoZY,	omitted from analysis
11103 (146)	1961Fa03	
10592 (365)	1961Gl08	
10994 (256)	1962Fl09	
10840 (18)	1963Go03	
10665 (110)	1963Ri02	
10738 (66)	1964Co35	
10921 (183)	1965Fl01	
11286 (256)	1965Fl01	
11220 (47)	1965Le25	
11030 (110)	1966Re13,	replaced by 1972Em01
11041 (58)	1968Re04,	replaced by 1972Em01
11191 (157)	1970Ha32	
10921 (16)	1970Wa19,	replaced by 1983Wa26
11023 (37)	1972Em01	
11034 (29)	1973Co39	
11020.8 (41)	1973Di01	
10906 (33)	1978Gr08	
11009 (11)	1980Ho17	
10449 (147)	1980RuZX,	replaced by 1990Ma15
10678 (140)	1980RuZY,	replaced by 1990Ma15

10678 (140)	1982RuZV,	replaced by 1990Ma15
11206 (7)	1982HoZJ,	replaced by 1992Un01
10921 (19)	1983Wa26	
10941 (7)	1989KoAA	
10967.8 (45)	1990Ma15	
10940.8 (69)	1992Go24	
11015 (20)	1992Un01,	replaced by 2002Un02
11018.3 (95)	2002Un02	
10970 (20)	2004Sc04	
10976 (30)	Adopted	

If the four values from before 1960 are omitted as well as replaced values, the data set for analysis includes 21 values. The large reduced- χ^2 value (16.3) indicates that these data are quite discrepant; therefore, the adopted value will depend on the method of analysis.

Since no value in this data set contributes more than 50% of the relative weight, the Limitation of Relative Statistical Weight (LRSW) method does not adjust any of the input uncertainties; however, it may expand the final uncertainty to include the more precise value. The Normalized Residual (NORM, 1994Ka08) and RAJEVAL (1992Ra08) methods adjust the input uncertainties for the more discrepant values.

In 1997-1998 R.G. Helmer chose the Normalized Residual (NR) analysis for obtaining the recommended half-life value of 10964(9). That choice was based on a desire for reducing a large relative weight of the value from 1973Di01 and its big contribution to χ^2 value and also to avoid an expansion of the final uncertainty by use of the LRSW analysis. It was stated that the low evaluation result met the tendency of the last measurements (by 1992) and evaluation results to be lower. (Details of Helmer's analysis can be found in the book of 1999BeAA).

The updated NIST value, obtained as a result of continued measurements of six sources (2002Un02), changes the situation. This high value with a small uncertainty (half of that in 1992Un01) has shown that the discrepancy among the most recent and accurate measurements is still kept. Therefore, a small uncertainty of the evaluation result seems to be unrealistic.

Thus, at present we can use the LRSW analysis as one of the methods for the evaluation of the ¹³⁷Cs half-life.

The weighted average of the twenty one values is 10981.8, with an internal uncertainty of 2.3, a reduced χ^2 of 16.3, and an external uncertainty of 9.5. The unweighted average is 10967(37). The LWEIGHT computer program using the LRSW analysis has chosen the weighted average and expanded the final uncertainty to 39 so range includes the most precise value of 11020.8. Hence, use of the LRSW analysis leads to the evaluation of 10982(39) days for the ¹³⁷Cs half-life.

This evaluation agrees well with the recent independent evaluations. Woods and Collins (2004Wo02) used 11 experimental values since 1968 and recommended the value of 10990(40) days by similar evaluation technique. Helene and Vanin (2002He06) presented in their paper a very promising statistical procedure (BOOTSTRAP method) to deduce a best value and its standard deviation for a discrepant set of data. They used 19 experimental ¹³⁷Cs half-life values and obtained the evaluation result as 10987(30) days.

The NORM and RAJEVAL statistical procedures lead to the evaluation results of 10962(7) and 10971(6) days, with the small uncertainties. The Bayesian procedures (BAYS and MBAYS, 1994Ka08) give the equal result of 10982(10) days. Thus, different methods of statistical analysis have led to discrepant results. In such a way the best (the less worst ?) choice is derived from the BOOTSTRAP method. It gives an intermediate result (calculation of Helene and Vanin, 2006) between the unadjusted weighted mean and the adjusted values from different procedures and its uncertainty encompasses all the statistical results.

The adopted value of the ¹³⁷Cs half-life is **10976(30) days, or 30.05(8) years.**

2.1 Beta - Transitions

The emission probability (in %) of the β transition to the ground state has been measured as follows:

4.8 (3)	1957Ri41,	σ increased to 0.6
7.6 (8)	1958Yo01	
6.5 (2)	1962Da05,	σ increased to 0.6
4.8 (10)	1965Me03	
6.0 (5)	1966Hs02	
5.4 (3)	1969Ha05	
6.4 (5)	1978Gr09	
5.57 (7)	1983Be18	
5.69 (19)	Value from LRSW analysis	
5.64 (28)	Adopted value from sect. 4.2	

The uncertainties for early values of 1957Ri41 and 1962Da05 were increased by the evaluator to 0.6 to make them comparable with those of the values measured in the 1966 - 1978 period.

The LRSW analysis gives an internal uncertainty of 0.14, a reduced- χ^2 value of 2.03, and an external uncertainty of 0.19. In this analysis the uncertainty of the 1983Be19 value was increased from 0.07 to 0.19 in order to reduce its relative weight from 78% to 50%.

The average β energies and log ft values have been calculated using the LOGFT computer program.

The shape of the β spectra has been measured by 1983Be18, 1978Ch22, 1978Gr09, 1969Sc23, and 1966Hs02, which is useful in the determination of the relative β branch intensities.

The very detailed treatment of the expression for the shape of the β spectrum for the 2nd forbidden transition to the ground state argues that the measurement of 1983Be18 should replace all of the previous values. If this were done the $P_{\beta}(0)$ would decrease by 0.12% and $P_{\beta}(662)$ would increase by this amount. The $P_{\gamma}(662)$ would then increase by about 0.08%. However, the value of 1983Be18 has only been allowed to contribute 50% of the relative weight, as is our common practice. It should also be noted that this paper has additional influence since its data are also used in determining the $\alpha_T(662)$ value that is used in the calculation of $P_{\gamma}(662)$.

The adopted value $P_{\beta}(662)$ has been computed from the final adopted $P_{\gamma}(662)$ value. [The uncertainty has increased due to the inclusion of the uncertainty in $\alpha_T(662)$ twice.]

2.2 Gamma Transitions

The adopted $\alpha_T(662)$ value of 0.1102 (19) is from a LRSW analysis of the 5 measured values recommended in the 1985HaZA evaluation, except that the value of 1983Be18 is used in place of value of 1978Ch22; these values are 0.1100 (11) (1965Me03), 0.1121 (5) (1969Ha05), 0.1105 (10) (1973LeZJ), 0.1100 (6) (1975Go28), and 0.1083 (5) (1983Be18, where the uncertainty has been increased to match the lowest other value). For this average, internal uncertainty = 0.0003, the reduced- χ^2 = 7.3, and the external uncertainty = 0.0008. The final uncertainty was increased by the LRSW analysis from 0.0008 to 0.0019 to include the 2 most precise values. Due to the large discrepancies among the 12 measured α values reported, 1985HaZA chose not to recommend any value.

The theoretical α_T value interpolated from the tables of 1978Ro21 is 0.1143 34; but 1990Ne01 has suggested that the α_T values for M4's from 1978Ro21 should be multiplied by 0.975 which gives 0.1114; this agrees with the adopted value to 1.1% which is much smaller than the uncertainty in either value. The theoretical total ICC value interpolated from the tables of 1993Ba60 $\alpha_T(662)=0.1116$.

Other measurements of α_T listed in 1985HaZA include 0.114 (2) (1957Ri41), 0.114 (30) (1962Da05), 0.109 (20) (1963Bo31), 0.1167 (15) (1965Pa17), 0.112 (11) (1965Ra12), 0.1092 (8) (1978Ch22), and 0.114 (3) (1978Gr09).

The adopted value $\alpha_K(662)$ of 0.0896 (15) is from the LRSW analysis of the 4 values recommended in the 1985HaZA evaluation, except for the value of 1983Be18 which is used in place of that from 1978Ch22; these values are 0.0894 (10) (1965Me03), 0.0916 (4) (1969Ha05), 0.0901 (9) (1973LeZJ), and 0.0881 (2) (1983Be18). The LRSW analysis increases the uncertainty of the 1983Be18 value from 0.0002 to 0.00034 to reduce its relative weight from 75% to 50%. For this average, the internal uncertainty = 0.0002, the reduced- $\chi^2 = 14.8$, and the external uncertainty = 0.0009. The final uncertainty was increased by the LRSW analysis from 0.0009 to 0.0015 to include the most precise value.

The theoretical value $\alpha_K(662)$ interpolated from the tables of 1978Ro21 is 0.0929 28; but 1990Ne01 has suggested that the α_K values for M4's from 1978Ro21 should be multiplied by 0.975 which gives 0.0906; this agrees with the adopted value to 1.1% which is much smaller than the uncertainty in either value. The theoretical $\alpha_K(662)$ value interpolated from the tables of 1993Ba60 $\alpha_K(662)=0.0907$.

Other measured values of α_K listed in 1985HaZA are 0.097 (3) (1951Wa19), 0.095 (5) (1952He33), 0.11 (1) (1953Do31), 0.096 (5) (1954AZ01), 0.095 (8) (1957Mc34), 0.093 (1957Ri41), 0.092 (6) (1959Wa17), 0.0976 (55) (1958Yo01), 0.093 (6) (1959Hu23), 0.093 (6) (1960De17), 0.095 (4) (1961Hu12), 0.093 (3) (1962Da05), 0.0957 (10) (1965Pa17), 0.092 (9) (1965Ra12), 0.093 (7) (1966Hs01), 0.094 (5) (1966Hu02), 0.093 (9) (1967Ba80), 0.0925 (27) (1967HaZX), 0.0922 (22) (1973Wi10), 0.0901 (10) (1971BrAA), 0.0888 (70) (1978Ch22), and 0.093 (3) (1978Gr09).

3 Atomic Data

The data are from Schönfeld and Janßen (1996Sc06).

3.1 X Radiations

The data are from Schönfeld and Janßen (1996Sc06).

3.2 Auger Electrons

The data are from Schönfeld and Janßen (1996Sc06).

4 Radiation Emissions

4.1 Electron Emission

The β^- data are from RADLIST or LOGFT. The Auger and conversion electron data are from Schönfeld (1996Sc06) calculations. For comparison, these emission probabilities and those from RADLIST (with the atomic data from Schönfeld) are:

Electrons per decay

	Schönfeld	RADLIST
L Auger	0.0728 (12)	0.0728 (22)
K Auger	0.0076 (4)	0.0076 (3)
K-662	0.07644	0.076 (3)
L-662	0.01387	0.0142 (6)

4.2 Photon Emissions

The 662-keV γ -ray energy is from 2000He14 and that for the 283-keV γ is from 1997WaZZ, but more precise values of 283.46 6 and 283.53 4 are available from (n,n' γ) studies.

The intensity of the 662-keV γ ray has been deduced in two ways, (1) the ratio of the measured γ emission

rate and the measured source decay rate and (2) from the probability of β - decay to the 662-keV level and α_T (662). These two values are independent as long as they involve independent measurements. Of the many papers that quote P_γ values, several are listed in section 2.1 as giving $P_{\beta}(0)$ values and are not included here. References 1965Me03 and 1978ChZZ have been replaced by 1978MeZM and 1983Be18, respectively. This leaves the following three values of $P_\gamma(662)$ to consider:

85.3 (10)	1973LeZJ
86.0 (9)	1975Go28
84.7 (7)	1978MeZM
85.2 (5)	Weighted average with reduced- $\chi^2 = 0.65$

[It should be noted that in the evaluation of 1991BaZS the value of 1973LeZJ is quoted as 0.8456 (8), which is the value from 1978Ch22. The evaluation of 1997Tu04 adopts the 1991BaZS result and repeats this error.]

The second value of $P_\gamma(662)$ comes from the average $P_{\beta}(0) = 5.69\%$ (19) in section 2.1 and the $\alpha_T(662) = 0.1102$ (19) in section 2.2, $P_{\beta}(662)/[1.0+\alpha(662)] = 84.95\%$ (22). Then, the adopted value is taken to be the weighted average of the values 84.95% (22) and 85.2% (5) which is 84.99% (20).

The decay of ¹³⁷Cs to the first excited level in ¹³⁷Ba at 283 keV was observed in 1996Bi23 and 1997WaZZ. The γ -ray intensity relative to that of the 662-keV γ ray is 0.00053 (14) (1996Bi23) and 0.00061 (10) (1997WaZZ) which gives an average of 0.00058 (8) and a corresponding transition intensity of 0.00061 (8).

The final P_{β} values are adjusted to be in agreement with this result and are $P_{\beta}(662) = 94.36\%$ (28) and $P_{\beta}(0) = 5.64\%$ (28). [The uncertainties here are overestimated because the contribution from $\alpha_T(662)$ has been included twice.]

The X-ray emission probabilities are from the γ -ray emission probability, the internal-conversion coefficients, and the atomic data of 1996Sc06. The difference between the Schönfeld values given and the RADLIST values are within the uncertainties:

	Photons per decay	
	Schönfeld	RADLIST
$K_{\alpha 2}$	0.0195 (4)	0.0195 (7)
$K_{\alpha 1}$	0.0358 (7)	0.0359 (13)
K_{β}	0.0132 (3)	0.0132 (5)
Total K	0.0685 (13)	0.0686 (16)

Double-decay processes which might occur in lieu of the 662-keV γ ray have been studied; two γ 's (1960Be20, 1992BaAA, 1993Ba46); a K shell electron plus a γ (1969Lj01, 1971Lj01); and two electrons (1971Lj02, 1971Po04). The paper of 1993Ba46 suggests an upper limit of the ratio of 2 γ emission to 1 γ emission of 5.10^{-7} .

5 References

- 1951Wa19 M.A.Waggoner, Phys.Rev. **82** (1951) 906 [α_K]
 1952He33 R.L. Heath, Phys.Rev. **87** (1952) 1132 [α_K]
 1953Do31 V.M.Dolishnyuk, G.M.Drabkin, V.I.Orlov, L.I.Rusinov, Doklady Akad.Nauk SSSR **92** (1953) 1141 [α_K]
 1954Az01 T.Azuma, J.Phys.Soc.Japan **9** (1954) 1 [α_K]
 1954Ve09 J.Verhaeghe, J.Demuyne, Compt.Rend. **239** (1954) 1374 [K/L]
 1955Br06 F.Brown, G.R.Hall, A.J.Walter, J.Inorg.Nuclear Chem. **1** (1955) 241 [$T_{1/2}$]
 1955Wi21 D.M.Wiles, R.H.Tomlinson, Phys.Rev. **99** (1955) 188 [$T_{1/2}$]
 1957Mc34 F.K.McGowan, P.H.Stelson, Phys.Rev. **107** (1957) 1674 [α_K]

- 1957Ri41 R.A.Ricci, Physica **23** (1957) 693 [α_T , α_K , P_{β^-}]
- 1958MoZY A. J. Moses, H. D. Cook, report TID-7568, part 2 (1958) 192 [$T_{1/2}$]
- 1958Yo01 Y.Yoshizawa, Nuclear Phys. **5** (1958) 122 [α_K , P_{β^-}]
- 1959Hu23 S.Hultberg, R.Stockendal, Arkiv Fysik **14** (1959) 565 [α_K]
- 1959Wa17 A. H. Wapstra, G. J. Nijgh, N. Salomons-Grobbe, L. T. M. Ornstein, Nucl. Phys. **9** (1959) 538 [α_K]
- 1960Be20 W.Beusch, Helv.Phys.Acta **33** (1960) 363 [double particle emission]
- 1960De17 C. de Vries, E. J. Bleeker, N. Salomons-Grobbe, Nucl. Phys. **18** (1960) 454 [α_K]
- 1961Fa03 H. Farrar, A. K. Dasgupta, R. H. Tomlinson, Can. J. Chem. **39** (1961) 681 [$T_{1/2}$]
- 1961Gl08 M. P. Glazunov, A. I. Grivkova, B. A. Zaitsev, V. A. Kiselev, Atomic Energy **10** (1961) 622 (page 615 in English) [$T_{1/2}$]
- 1961Hu12 S.Hultberg, D.J.Horen, J.M.Hollander, Nucl.Phys. **28** (1961) 471 [α_K]
- 1962Da05 H.Daniel, H.Schmitt, Z.Physik **168** (1962) 292 [α_T , α_K , P_{β^-}]
- 1962Fl09 D. G. Fleishman, I. V. Burovina, V. P. Nesterov, Atomic Energy **13** (1962) 1225 (page 1224 in English) [$T_{1/2}$]
- 1962Ge09 J.S.Geiger, R.L.Graham, F.Brown, Can.J.Phys. **40** (1962) 1258 [K/L]
- 1963Bo31 H.E.Bosch, T.Urstein, Nucl.Instr.Methods **24** (1963) 109 [α_T]
- 1963Go03 S. G. Gorbics, W. E. Kunz, A. E. Nash, Nucleonics **21**, No. 1 (1963) 63 [$T_{1/2}$]
- 1963Ri02 B. F. Rider, J. P. Peterson, Jr., C. P. Ruiz, Nucl. Sci. Eng. **15** (1963) 284 [$T_{1/2}$]
- 1964Ch18 Y.Y.Chu, M.L.Perlman, Phys. Rev. **135** (1964) B319 [K/L]
- 1964Co35 H. D. Cook, C. J. Rettger, W. J. Sewalk, report WAPD-BT-30 (1964) [$T_{1/2}$]
- 1964Ge05 H.U.Gersch, E.Hentschel, P.Gippner, W.Rudolph, Nucl. Instr. Methods **25** (1964) 314 [K/L]
- 1965Fl01 K. F. Flynn, L. E. Glendenin, A. L. Harkness, E. P. Steinberg, J. Inorg. Nucl. Chem. **27** (1965) 21 [$T_{1/2}$]
- 1965Le25 R. E. Lewis, R. E. McHenry, T. A. Butler, Trans. Amer. Nucl. Soc. **8** (1965) 79 [$T_{1/2}$]
- 1965Me03 J. Merritt, J. G. V. Taylor, An. Chem. **37** (1965) 351 [α_K , α_T , P_{γ} , P_{β^-}]
- 1965Pa17 D.Parsignault, Thesis, Univ.Paris (1965); CEA-R-2631(1965) [α_T , α_K]
- 1965Ra12 M.R.Rao, S.Jnanananda, Nucl.Instr.Methods **36** (1965) 261; Phys.Abst. **69** (1966) 666, Abstr.7047 [α_T , α_K]
- 1966Hs01 S.T.Hsue, L.M.Langer, E.H.Spejewski, S.M.Tang, Nucl.Phys. **80** (1966) 657 [α_K]
- 1966Hs02 S. T. Hsue, L. M. Langer, S. M. Tang, Nucl. Phys. **86** (1966) 47 [P_{β^-}]
- 1966Hu02 S.Hultberg, A.A.Bartlett; J. H. Hamilton, Ed., Academic Press, Inc., New York, (1966) 141 [α_K , β shape]
- 1966Re13 S.A.Reynolds, ORNL-3889 (1966) 57 [$T_{1/2}$]
- 1967Ba80 E. Baldinger, E. Haller, Helv. Phys. Acta **40** (1967) 800 [α_K]
- 1967HaZX H.H.Hansen, M.Delabaye, Proc.Symp.Standardization of Radionuclides,Vienna, Austria (1966), Intern.At.Energy Agency, Vienna (1967) 361; CONF-661012 (1967) [α_K]
- 1968Re04 S.A.Reynolds, J.F.Emery, E.I.Wyatt, Nucl.Sci.Eng. **322** (1968) 46 [$T_{1/2}$]
- 1968Wo02 J.L.Wolfson, A.J.Collier, Nucl.Phys. **A112** (1968) 156 [E_{β^-}]
- 1969Ha05 H. H. Hansen, G. Lowenthal, A. Spornol, W. van der Eijk, R. Vaninbrouckx, Z. Phys. **218** (1969) 25 [P_{β^-} , P_{γ} , α_K]
- 1969Lj01 A.Ljubicic, B.Hrastnik, K.Ilakovac, V.Knapp, B.Vojnovic, Phys.Rev. **187** (1969) 1512 [double particle emission]
- 1969Sc23 H.Schneuwly, L.Schellenberg, O.Huber, W.Lindt, Helv.Phys.Acta **42** (1969) 743 [β shape]
- 1970Ha32 G. Harbottle, Radiochim. Acta **13** (1970) 132 [$T_{1/2}$]
- 1970Wa19 K. F. Walz, H. M. Weiss, Z. Naturforsch. **25a** (1970) 921 [$T_{1/2}$]
- 1971BrAA P. Brethon, report CEA-R-4196 (1971) [α_K]
- 1971Ca08 J.L.Campbell, H.J.Smith, I.K.Mackenzie, Nucl.Instrum.Methods **92** (1971) 237 [P_X]
- 1971Lj01 A.Ljubicic, B.Hrastnik, K.Ilakovac, M.Jurcevic, I.Basar, Phys.Rev. **C3** (1971) 824 [double particle emission]
- 1971Lj02 A.Ljubicic, M.Jurcevic, K.Ilakovac, B.Hrastnik, Phys.Rev. **C3** (1971) 831 [double particle emission]
- 1971Po04 F.T.Porter, M.S.Freedman, F.Wagner Jr., Phys.Rev. **C3** (1971) 2246 [double particle emission]
- 1972Em01 J. F. Emery, S. A. Reynolds, E. I. Wyatt, G. I. Gleason, Nucl. Sci. Eng. **48** (1972) 319 [$T_{1/2}$]
- 1973C039 J. A. Corbett, Nucl. Eng. Int. **18** (1973) 715 [$T_{1/2}$]

- 1973Di01 L. A. Dietz, C. F. Pachuck1, J. Inorg. Nucl. Chem. **35** (1973) 1769 [T_{1/2}]
- 1973LeZJ J. Legrand, J. P. Brethon, F. Lagoutine, CEAR-4428 (1973) [α_K , α_T , P _{γ} , T_{1/2}, P _{β}]
- 1973Wi10 J.B.Willett, G.T.Emery, Ann.Phys.(New York) **78** (1973) 496 [α_K]
- 1975Go28 I. W. Goodier, J. L. Makepeace, L. E. Stuart, Int. J. Appl. Radiat. Isot. **26** (1975) 490 [α_T , P _{γ} , P _{β}]
- 1976Bo16 G.L.Borchert, Z.Naturforsch. **31a** (1976) 387 [E _{γ}]
- 1978Ch22 P.Christmas, P.Cross, Metrologia **14** (1978) 157 [α_T , α_K]
- 1978Gr08 W. H. Gries, J. Steyn, Nucl. Instr. and Meth. **152** (1978) 459 [T_{1/2}, P _{β}]
- 1978Gr09 K.Y.Gromov, T.Kretsu, V.V.Kuznetsov, G.Makarie, Izv.Akad.Nauk SSSR, Ser.Fiz. **42** (1978) 790; Bull.Acad.Sci.USSR, Phys.Ser. **42**, No.4 (1978) 85 [α_T , α_K , β shape, P _{γ} , P _{β} , T_{1/2}]
- 1978MeZM J. S. Merritt, F. H. Gibson, report AECL-6203 (1978) [P _{γ}]
- 1978Ro21 F. Rösel, H. M. Fries, K. Alder, H. C. Pauli, Atomic Data Nucl. Data Tables **21** (1978) 91 and 292 [Theoretical ICC]
- 1980Ho17 H. Houtermans, O. Milosevic, F. Reichel, Intern. J. Appl. Radiat. Isot. **31** (1980) 153 [T_{1/2}]
- 1980RuZX A.R.Rutledge, J.S.Merritt, L.V.Smith, AECL-6788 (1980) 45 [T_{1/2}]
- 1980RuZY A.R.Rutledge, L.V.Smith, J.S.Merritt, AECL-6692 (1980) [T_{1/2}]
- 1982HoZJ D.D.Hoppes, J.M.R.Hutchinson, F.J.Schima, M.P.Unterweger, NBS-SP-626 (1982) 85 [T_{1/2}]
- 1982RuZV A.R.Rutledge, L.V.Smith, J.S.Merritt, NBS-SP-626 (1982) 5 [T_{1/2}]
- 1983Be18 H. Behrens, P. Christmas, Nucl. Phys. **399** (1983) 131 [α_T , α_K , P _{γ}]
- 1983Wa26 K.F.Walz, K.Debertin, H.Schrader, Int.J.Appl.Radiat.Isotop. **34** (1983) 1191 [T_{1/2}]
- 1985HaZA H.H.Hansen, European App.Res.Rept.Nucl.Sci.Technol. **6**, No.4 (1985) 777 [α_T , α_K]
- 1987Me02 D.Mehta, S.Singh, H.R.Verma, N.Singh, P.N.Trehan, Nucl.Instrum.Methods Phys.Res. **A254** (1987) 578 [P_X]
- 1989KoAA A. E. Kochin, Schmidt-Ott, H.Behrens, Z.Phys. **A337** (1990) 169 [T_{1/2}]
- 1990Ma15 R. H. Martin, J. G. V. Taylor, Nucl. Instr. Meth. **A286** (1990) 507 [T_{1/2}]
- 1990Ne01 Zs. Nemeth, A. Veres, Nucl. Instr. Meth. **A286** (1990) 601 [α_T , α_K]
- 1990Wo04 M. J. Woods, Nucl. Instr. Meth. **A286** (1990) 576 [T_{1/2}]
- 1991BaZS W.Bambynek, T.Barta, R.Jedlovsky, P.Christmas, N.Coursol, K.Debertin, R.G.Helmer, A.L.Nichols, F.J.Schima, Y.Yoshizawa, IAEA-TECDOC-619 (1991) [¹³⁷Cs decay data evaluation]
- 1992BaAA V. K. Basenko, A. N. Berlizov, and G. A. Prokopets, Bull. Russian Acad. Sci. **56** (1992) 1, 94 [P _{γ}]
- 1992Go24 J.-J. Gostely, Appl. Radiat. Isot. **43** (1992) 949 [T_{1/2}]
- 1992Ra08 M. U. Rajput, T. D. MacMahon, Nucl. Instr. Meth. **A312** (1992) 289 [T_{1/2} analysis]
- 1992Un01 M. P. Unterweger, D. D. Hoppes, F. J. Schima, Nucl. Instr. Meth. **A312** (1992) 349 [T_{1/2}]
- 1993Ba46 V. K. Basenko, A. N. Berlizov, and G. A. Prokopets, Bull. Russian Acad. Sci. **57** (1993) 55 [P _{γ}]
- 1993Ba60 I.M.Band and M.B.Trzhaskovskaya, At. Data and Nucl. Data Tables **55** (1993) 43. [Theoretical ICC]
- 1994Ka08 S.I.Kafala, T.D.MacMahon, P.W.Gray, Nucl.Instrum.Methods Phys.Res. **A339** (1994) 151 [T_{1/2} analysis]
- 1996Bi23 I. Bikit, I. Anicin, J. Slivka, M. Krmar, J. Puzovic, Lj.Conkic, Phys. Rev. C **54** (1996) 3270 [P _{γ}]
- 1996Sc06 Schönfeld, H. Janßen, Nucl. Instr. Meth. **A369** (1996) 527 [ω_K , ω_L , relative K x-ray, emission probabilities]
- 1997WaZZ B. K. Wagner, P. E. Garrett, M. Yeh, S. W. Yates, private communication (1997) [P _{γ}]
- 1997Tu04 J. K. Tuli, Nuclear Data Sheets **81** (1997) 579 [Decay scheme]
- 1999BeAA M.-M. Bé, E. Browne, V. Chechev, R. Helmer and E. Schönfeld. Table de Radionucléides. Comments on Evaluations. // CEA-ISBN 2-7272-0211-3. 1999.
- 2000He14 R. G. Helmer and C. van der Leun, Nucl. Instr. Meth. **A450** (2000) 35 [E _{γ}]
- 2002He06 O.Helene and V.R.Vanin, Nucl.Instrum.Methods Phys.Res. **A481** (2002) 626 [T_{1/2} analysis]
- 2003Au03 G.Audi, A.H.Wapstra, and C.Thibault, Nucl.Phys. **A729** (2003) 337 [Q value]
- 2004Wo02 M.J.Woods and S.M.Collins, Appl.Radiat.Isot. **60** (2004) 257 [T_{1/2} analysis]
- 2004Sc04 H.Schrader, **60** (2004) 317 [T_{1/2} analysis]

¹³⁸La - Comments on evaluation of decay data

by X. Mougeot, M.M. Bé

This evaluation was completed in November 2013 with literature available at this date included. The updated evaluation was done in June 2016 to account for the recent publication 2016Qu01.

Decay Scheme

¹³⁸La decays by electron capture and β^- disintegrations to the first excited level of ¹³⁸Ba and of ¹³⁸Ce respectively.

Q_{EC} and Q_{β^-} values for ¹³⁸La decay were adopted from Wang *et al.* (2012Wa38).

It should be noted that for the EC decay there exists one estimate of the Q_{EC} value: 1590 (40) keV, cited as 1620 (15) in 2012Wa38, which is far from the adjusted value of 1740,0 (34) keV resulting from a least-square calculation.

For the β^- decay, a maximum energy of 205 (10) keV and 370 (40) keV were determined by 1957Gl20 and 1970ElZR leading to a Q_{β^-} of 1015 (14) keV and of 1158 (40) keV respectively. The adjusted value in 2012Wa38 is 1047 (10) keV and was deduced from two mass measurements for ¹³⁸La and ¹³⁸Ce respectively. A new endpoint energy of 264,0 (43) keV was determined in 2016Qu01, corresponding to a Q_{β^-} of 1052,7 (43) keV. Following a private communication with G. Audi, the final value adopted here is $Q_{\beta^-} = 1051,7$ (40) keV. This value will be included within the next atomic mass evaluation.

The spins, parities and level energies are based on the mass-chain evaluation of Sonzogni (2003So13).

Half-life

The experimental half-life values used to calculate the recommended value are listed in the following table:

References	10¹¹ a	uc (10¹¹ a)	Comments
Pringle (1951Pr**)	2,0		Omitted
Mulholland (1952Mu04)	0,66		Omitted
Turchinets (1956Tu17)	1,0	0,1	Omitted – comparison with ⁴⁰ K ($A_\gamma = 3,4$)
Glover (1957Gl20)	1,16	0,11	Omitted – comparison with ⁴⁰ K ($A_\gamma = 3,47$)
Ellis (1972El02)	1,56	0,3	Omitted – comparison with ⁴⁰ K ($A_\gamma = 3,25$)
Marsol (1972Ma31)	1,27	0,18	Omitted – comparison with ⁴⁰ K
Taylor (1979Ta21)	1,28	0,12	Omitted – comparison with ⁴⁰ K ($A_\gamma = 3,26$)
Sato (1981Sa42)	1,03	0,03	Omitted – comparison with ⁴⁰ K, uc = 3%
Norman (1983No02)	1,06	0,03	WM of 1,07 (3) from mixed source with ⁴⁰ K and 0,97 (12) direct measurement
De Ruyter (1966De04)	1,04	0,03	Original uc = 0,014 multiplied by 2
Cesana (1977Ce**)	1,29	0,12	uc = 10% to account for systematic component, outlier
Norman (1983No02)	0,97	0,12	Direct measurement
Dalmasso (1994Da**)	1,14	0,05	Direct measurement
Nir-El (1997Ni12)	1,01	0,023	Direct measurement, original uc = 0,01 increased to limit its weight to 50%

Bernabei (2005Be73)	1,07	0,06	Direct measurement
Crit (χ^2)	3,32		
$\chi^2 / (n-1)$	1,56		
UWM	1,046		
WM	1,036		
uc(WM)int. :	0,016		
uc(WM)ext. :	0,020		
Adopted	1,036 (20)		

The six first results listed in the above table were derived from the comparison of the specific activity in ^{138}La of a lanthanum crystal with the specific activity of ^{40}K . The resulting values depend on the ^{40}K decay data which are not always the same in the various publications, and, when they are, the results are no longer independent. For these reasons these six values have not been used in the evaluation.

The six remaining results were obtained from direct measurement of the ^{138}La activity and calculation of the number of ^{138}La atoms. In this set, the value of Cesana was found to be an outlier. The original uncertainty of De Ruyter (1966De04) has been multiplied by 2, to account for a systematic component. The adopted value for Norman (1983No02) is the one obtained by direct measurement. The uncertainty given by Nir-El (1997Ni12) was increased from 0,01 to 0,023 to limit its weight to 50%, the resulting uncertainty of about 2% seems reasonable for a measurement of such a long half-life.

This set of five data adopted for statistical processing is consistent. The adopted value is the weighted mean with the external uncertainty (the lowest experimental uncertainty is 0,01), that is:

$$T_{1/2} = 103,6 (20) 10^9 \text{ a.}$$

From this latter value and the branching ratios obtained below, a value of $2,33 (6) 10^{-12} \text{ a}^{-1}$ for the partial decay constant λ_{β} is derived. It can be compared with the value of $2,37 (10) 10^{-12} \text{ a}^{-1}$ determined by geophysical methods (2000Ta24).

Gamma Transition probabilities and Internal Conversion Coefficients

The γ transitions with energy 1435 and 788 keV respectively are of pure E2 character.

The adopted ICC are the theoretical values interpolated by the BrIcc program (2008Ki07) from the tables of Band *et al.* (2002Ba85).

Electron Capture transition

- The electron capture probability to the 1435 keV level is deduced from the decay scheme balance.

$$P_{\text{EC}} = I_{\gamma_{1435}} (1 + \alpha_{\text{T}}) = 65,2 (6) \%.$$

Reference	L/K	uc	Comments
Mandal (1984Ma46)	0,48	0,16	$P_{\text{K}} = 0,676 (10)$
Quarati (2012Qu02)	0,44	0,044	
Logft	0,43		
Quarati (2012Qu02) - theoretical	0,402		$P_{\text{K}} = 0,69$; $P_{\text{L}} = 0,277$; $P_{\text{M}} = 0,032$
Quarati (2016Qu01) – experimental	0,391	0,003	$M/\text{K} = 0,102 (3)$; $M/\text{L} = 0,261 (9)$
Quarati (2016Qu01) - theoretical	0,405	0,007	$M/\text{K} = 0,0915 (17)$; $M/\text{L} = 0,226 (7)$

- The K/L capture ratio was measured by Mandal (1984Ma46) and Quarati (2012Qu02, 2016Qu01). Their results are in good agreement with the theoretical values calculated, for this 2nd unique forbidden transition, by the Logft program and Quarati.

- The atomic sub shell electron capture probabilities and the $lg ft$ values are calculated with the LOGFT program.

β^- transition

The probability of the β^- transition to the 788 keV level is deduced from the decay scheme balance.

$$P_{\beta^-} = 100 - P_{EC} = 34,8 (6) \%$$

An experimental shape factor was extracted from Quarati's data as the deformation to be applied on an allowed shape (see Fig. 1):

$$C(W) = 1 - 55,435 \times W + 9,53 / W + 68,53 \times W^2 - 24,565 \times W^3$$

The maximum energy of the β^- spectrum was deduced from the new Q_{β^-} value to be 263,0 (40) keV. The mean β^- energy was calculated by using the experimental shape factor from 2016Qu01 to be 91,1 (21) keV.




Figure 1 – La-138 β^- spectrum: measurement from 2016Qu01 (black); usual 2nd forbidden unique calculation (green); allowed spectrum (red); allowed spectrum distorted by the extracted experimental shape factor (blue – hidden by experimental spectrum).

Gamma Rays

The gamma ray energies are deduced from level energies.

Absolute γ ray intensities

The measured gamma ray intensities are:

References	$I_{\gamma 788}$ (%)	uc (%)	$I_{\gamma 1435}$ (%)	uc (%)	Comments
Turchinetz (1956Tu17)	47	5	53	5	omitted
Glover (1957Gl20)	30		70		omitted
Ellis (1972El02)	33		67		omitted
Cesana (1977Ce**)	35,1	1,9	64,9	1,9	Derived from partial $T_{1/2}$ ratio
Sato (1981Sa42)	33,8	2,0	65,9	2,0	Derived from partial $T_{1/2}$ ratio
Norman (1983No02)	33,1	2,0	66,9	2,0	Derived from partial $T_{1/2}$ ratio
De Ruyter (1966De04)	37,0	1,2	63,0	1,2	Derived from I_{γ} ratio; original uc = 0,6 multiplied by 2
Marsol (1972Ma31)	32,1	1,8	67,9	1,8	Direct measurement
Nir-El (1997Ni12)	35,0	0,8	65,0	0,8	Derived from I_{γ} ratio; original uc = 0,4 multiplied by 2
Crit. χ^2	3,32		3,02		
$\chi^2/(n-1)$	1,65		1,29		
UWM	34,20		65,6		
WM	34,908		65,053		
uc(WM)int.	0,571		0,547		
uc(WM)ext.	0,734		0,622		
Adopted	34,7 (6)		65,1 (6)		

- 1435 keV gamma intensity

Six values of the 1435 keV gamma intensity have been used, as they were derived from ratios some parameters, such as $T_{1/2}({}^{40}\text{K})$, cancel out.

The original uncertainty of De Ruyter (1966De04) has been multiplied by 2, to account for a systematic component. The uncertainty given by Nir-El (1997Ni12) was increased to limit its weight to 50%.

This set of the six data adopted for the statistical processing is consistent, the weighted mean is 65,1 (6) %.

- 788 keV gamma intensity

From the statistical processing of the six values above the weighted mean is 34,9 (6) %. But with $P_{(\gamma+ce)} = P_{\beta^-} = 100 - P_{EC} = 34,8$ (6) % and the adopted internal conversion coefficients, the I_{γ} (788) intensity must be equal to 34,7 (6) %. This latter value is adopted.

Atomic Data

The X-ray and Auger electron emission intensities are derived from the decay scheme parameters by using the program Emission, the derived total K X-ray intensities is 41,6 (5) %.

A K X-ray intensity of 0,4 X-ray $s^{-1} g^{-1}$ of natural Lanthanum was observed in an old publication (Pringle *et al.* (1951Pr**)).

References

The program LOGFT, National Nuclear Data Center, September 3, 2002.

- 1951Pr**** Phys.Rev. 84, 1006 (1951), R.W.Pringle *et al.*; *A further study of the natural activity of Lanthanum*
- 1952Mu04** Phys.Rev. 85, 144 (1952), G.L.Mulholland, T.P.Kohman; *Possible Natural Radioactivity of Neodymium*
- 1956Tu17** Phys.Rev. 103, 1000 (1956), W.Turchinets, R.W.Pringle; *Decay Scheme of La¹³⁸*
- 1957Gl20** Phil.Mag. 2, 49 (1957), R.N.Glover, D.E.Watt; *The Natural Radioactivity of Lanthanum*
- 1966De04** Physica 32, 991 (1966), A.W.De Ruyter, A.H.W.Aten, Jr., A.Van Dulmen, C.Krol-Koning, E.Zuidema; *Specific Gamma Emission of Natural Potassium and Lanthanum*
- 1972El02** Nucl.Phys. A179, 540 (1972), J.L.Ellis, H.E.Hall, Jr.; *Decay of Naturally Occurring ¹³⁸La*
- 1972Ma31** C.R.Acad.Sci., Ser.B, 274, 904 (1972), C.Marsol, F.Armanet, G.Ardisson ; *Étude de la Désintégration du Nucléide Naturel ¹³⁸La (T_{1/2}=1, 27.10¹¹ années)*
- 1977Ce**** Analytical Chemistry, Vol 49, 8 (1977) 1156, A. Cesana, M. Terrani; *Gamma-ray activity determination in large volume samples with a Ge-Li detector*
- 1979Ta21** J.Phys.Soc.Jpn. 47, 1395 (1979), H.W.Taylor, R.J.Bauer; *A Note on the Decay of ¹³⁸La*
- 1981Sa42** Radiochem.Radioanal.Lett. 46, 145 (1981), J.Sato, T.Hirose; *Half-Life of ¹³⁸La*
- 1983No02** Phys.Rev. C27, 1321 (1983), E.B.Norman, M.A.Nelson; *Half-Life and Decay Scheme of ¹³⁸La*
- 1984Ma46** J.Phys.(London) G10, 1765 (1984), A.M.Mandal, A.P.Patro; *(L+M+...)/K Capture Ratio in the Decay of ¹³⁸La*
- 1994Da**** Applied Radiation Isotopes 45, 3, 388 (1994), J. Dalmasso, G. Barci-Funel, G. Ardisson; *Reinvestigation of the Decay of the Naturally Occurring Nuclide ¹³⁸La*
- 1997Ni12** Radiochim.Acta 77, 191 (1997), Y.Nir-El; *Determination of the Half-Life and Gamma-Ray Emission Probability of ¹³⁸La*
- 2002Ba85** At.Data Nucl.Data Tables 81, 1 (2002), I.M.Band, M.B.Trzhaskovskaya, C.W.Nestor, Jr., P.O.Tikkanen, S.Raman; *Dirac-Fock Internal Conversion Coefficients*
- 2003So13** Nucl.Data Sheets 98, 515 (2003), A.A.Sonzogni ; *Nuclear Data Sheets for A = 138*
- 2005Be73** Nucl.Instrum.Methods Phys.Res. A555, 270 (2005), R.Bernabei, P.Belli, F.Montecchia, F.Nozzoli, A.d'Angelo, F.Cappella, A.Incicchitti, D.Prosperi, S.Castellano, R.Cerulli, C.J.Dai, V.I.Tretyak; *Performances and potentialities of a LaCl₃:Ce scintillator*
- 2008Ki07** Nucl.Instrum.Methods Phys.Res. A589, 202 (2008), T.Kibédi, T.W.Burrows, M.B.Trzhaskovskaya, P.M.Davidson, C.W.Nestor, Jr.; *Evaluation of theoretical conversion coefficients using BrIcc*
- 2000Ta24** Physical Review C 62, 017601-1 (2000), Masaharu Tanimizu
- 2012Qu02** Nucl.Instrum.Methods Phys.Res. A683, 46 (2012), F.G.A.Quarati, I.V.Khodyuk, C.W.E.van Eijk, P.Quarati, P.Dorenbos; *Study of ¹³⁸La radioactive decays using LaBr₃ scintillators*
- 2012Wa38** Chin.Phys.C 36, 1603 (2012), M.Wang, G.Audi, A.H.Wapstra, F.G.Kondev, M.MacCormick, X.Xu, B.Pfeiffer; *The AME2012 atomic mass evaluation (II). Tables, graphs and references*
- 2016Qu01** Applied Radiation Isotopes 109, 172-176 (2016), F.G.A.Quarati, P.Dorenbos, X. Mougeot; *Experiments and theory of ¹³⁸La radioactive decay*
- 2016Au** Priv. Comm., G. Audi, *Atomic Mass Evaluation 2016* [Q]

¹³⁹Ce - Comments on evaluation of decay data by M.M. Bé, R. G. Helmer, E. Schönfeld

1 Decay Scheme

This evaluation was completed in September 1996 and reviewed in 2007. The literature available by December 2007 was included.

This decay scheme is complete since the only excited level below the ¹³⁹Ce decay energy is populated (1989Bu12).

2 Nuclear Data

A Q value of 270 (3) keV is deduced from P_K measurements (see §2.1). It can be compared with a Q value of 264.6 (20) keV from measurement of the internal bremsstrahlung spectrum of 1996Hi14.

The ¹³⁹Ce half-life values available are, in days:

140 (1)	1948Po01	# (Pool and Krisberg as quoted in 1965An07)
137.5 (3)	1965An07	(Anspach et al.)
137.2 (4)	1972Em01	(Emery et al.)
137.63 (10)	1973MeYE	# (Merritt), replaced by 1982RuZV
137.65 (7)	1976Me	# (Merritt), replaced by 1982RuZV
137.66 (4)	1976Va30	(Vaninbroukx and Grosse)
137.59 (4)	1978La21	(Lagoutine et al.), uncertainty quoted as 0.12 at 3σ level
137.65 (3)	1980RuZY	# (Rutledge et al.), replaced by 1982RuZV
137.74 (8)	1982HoZJ	# (Hoppes et al.), replaced by 1992Un01
137.65 (3)	1982RuZV	(Rutledge et al.)
137.8 (2)	1982RyZX	(Rytz) BIPM value in NBS-SP-626
137.73 (9)	1992Un01	(Unterweger et al.)
137.641 (20)	Weighted average & adopted	

The value of 1948Po01 was omitted due to its large uncertainty. Omitting this value and the several (#) that were replaced by latter values, one has seven values to consider. The weighted average of these seven values is 137.641 with an internal uncertainty of 0.020 and a reduced- χ^2 of 0.83. No adjustments were made in the Limitation of Relative Statistical Weight method since the largest relative weight is less than 50 %, namely 44 % for the 1982RuZV value; also the set is consistent.

2.1 Electron Capture Transitions

The energies of the electron-capture transitions (ϵ) are calculated from the Q value and the level energies. The ϵ branch to the ground state is 2nd forbidden. From the log $f\bar{t}$ systematics (1998Si17), the expected log $f\bar{t}$ value is > 10.6 and the corresponding limit is $P_\epsilon(0) < 0.008 \%$ compared to the measured limit of $P_\epsilon(0) < 1 \%$ (1956Ke23) and $P_\epsilon(0) < 0.000097 \%$ (1993Mi20). If asymmetric uncertainties are used, the evaluator suggests the other ϵ branch probability is 99.9973 +27-53. If only symmetric uncertainties are used, 99.9973 (27) is suggested.

The P_K value for transition to the 165-keV level was deduced from the 17 measured values.

The available measured P_K values are listed in the following table as given in the original papers:

Value (uc)		ω_K	Reference	
0.87	(4)	Independant	Outlier	1954Pr31 (Pruett)
0.73	(2)	Independent		1956Ke23 (Ketelle)
0.68	(2)	Independent		1967Ma07 (Marelius)
0.75	(1)	Independent		1968Ad08 (B.Adamowicz)
0.69	(2)	Independent		1968Va08 (E.Vatai)
0.705	(20)	0.92 (1)		1972Ca07 (Campbell)
0.78	(3)	Independent		1972Sc08 (Schmidt-Ott)
0.73	(3)	(Martin ?)		1975Da08 (Dasmahapatra)
0.726	(10)	Independent		1975Ha43 (Hansen)
0.705	(20)	0.906 (26)		1975Pl06 (Plch)
0.801	(34)	0.906 (26)	Outlier	1976Ha36 (Hartl)
0.76	(3)	0.906 (26)		1978Se ** (Sergienko)
0.710	(24)	0.926		1987BeYL (Begzhanov)
0.68	(2)	0.91 (3)		1988Ko** (Konstantinov)
0.74	(3)	0.905 (4)		1994Ku43 (Kumar)
0.704	(6)	0.907, $K\beta = 0.193$		1996Hi14 (Hindi)
0.714	(25)	0.906 (26)		1997Ka** (Kalyani)
Critical χ^2	2			
Reduced χ^2	2.4			
WM	0.716	External Unc.= 0.006 Expanded Unc. = 0.012		
Adopted	0.716	0.006		

Two values (1954Pr31 and 1976Ha36) were found outlier due to Chauvenet's criterion. The remaining set of 15 values is slightly discrepant with a reduce χ^2 of 2.4.

The most important contribution comes from the Hindi's value amounting for 40 %, this value was deduced from the measurement of the Q value.

From this P_K value of 0.716 (6), a Q value of 270 (3) keV is derived.

A value of $Q=279$ (7) was obtained in 2003Au03 using the same methodology but with a reduce set of 10 P_K values (from 1954Pr31 to 1976Ha36).

See 1988Ri08 (Ruisager) for possible effects on the capture rates of the finite widths of the atomic levels.

2.2 Gamma Transitions

The probability for the 165-keV γ - transition is equal to the probability of the preceding ϵ - transition.

The γ - ray is mostly M1 and the %E2 is taken to be 0.0. The reported $\delta(E2/M1)$ are: +0.034 (34) [1963Ha07 from (γ , θ , T) and polarization]; 0.045 (+26-45) (1965Ge04 from $L_1/L_2/L_3$); 0.029 (+18-29) with the nuclear penetration parameter $\lambda = 2.8$ (13) (1979Ha21 from analysis of published data); and < 0.0055 with $\lambda = 4.2$ (8) (1977Ry01 from analysis of published measured data and a new calculation of a values). Also, $\lambda = 3.1$ (7) with $\delta = 0.0$ (1975Pl06 from experimental α_K and other published α data) and $\lambda = 3.6$ (18) with $\delta = 0.0$ (1975Mo12). The weighted average of these four λ values is 3.5 (5) with a reduced- $\chi^2 = 0.46$. Since much of the data used to determine these λ values are common to the various calculations, the values are correlated. Therefore, the uncertainty is increased to the smallest of the four uncertainties, and the value 3.5 (7) is recommended.

The K-shell and total internal-conversion coefficients are from the 1985HaZA evaluation. This evaluation lists the following values :

Retained in 85HaZA analysis				
α_K	a	Reference	α_K	a
0.22		1954Mi56		
0.20 (4)		1954Nu12		
0.20 (5)		1954Pr31		
0.22 (1)		1956Ke23		
0.263		1962Be31		
0.2148 (12)	0.2514 (11)	1962Ta03	yes	yes
0.209 (27)		1967HaZX		
	0.254 (6)	1971Ar43		yes
	0.2446 (12)	1973Le29+1973LeYP		
0.207 (9)		1975Mo12	yes [as 0.214 (5)]	
0.214 (2)	0.251 (2)	1975Pl06	yes	yes
0.2152 (33)	0.2520 (50)	1976Ha11	yes	yes
	0.2519 (6)	1977Sc**		yes [as 0.2519(10)]
0.2146 (10)	0.2516 (7)	1985HaZA recommended and adopted here		
	0.261 (4)	2005KiZW		Theory for M1 "Frozen orbital"
	0.337 (5)	2005KiZW		Theory for E2 "Frozen orbital"
	0.267	1978Ro22		Theory for M1
	0.264	1968Ha52		Theory for M1
	0.339	1978Ro22		Theory for E2
	0.339	1968Ha52		Theory for E2

The theoretical values are for $\lambda = 0.0$. The α_L and α_M values were computed from the adopted α_K value and the K/L and K/M ratios from the M1 theoretical values interpolated from the table of Rösler (1978Ro22). Since this transition is hindered and the aspect of nuclear penetration effect discussed by various authors (1975Mo12, 1977Ry01, 1979Ha21, ...) the adopted α values are the experimental ones.

3 Atomic Data

The fluorescence yield data are from 1996Sc06 (Schönfeld and Janssen).

3.1 X Radiation

The x-ray energies are based on the wave lengths in the compilation of 1967Be65 (Bearden). The relative K x-ray emission probabilities are taken from 1996Sc06. The value for $P(X_L)/P(K_{\alpha 1})$ is derived from the emission probabilities (sect. 4.2).

3.2 Auger Electrons

The ratios $P(KLX)/P(KLL)$ and $P(KXY)/P(KLL)$ are taken from 1996Sc06. The value for $P(eAL)/P(KLL)$ is derived from the emission probabilities (sect. 4.1).

4 Radiation Emission

4.1 Electron Emission

The electron emission probabilities are calculated from the X and γ -ray emission probabilities in sects. 2.1 and 4.2, the atomic data of sect. 3, and the internal-conversion coefficients of sect. 2.2.

4.2 Photon Emissions

The γ -ray energy is from the evaluation 2000He14 where the values are on a scale on which the strong line from the decay of ¹⁹⁸Au is 411.80205 (17).

The γ -ray emission intensity is calculated as $I_c(165)/[1 + \alpha(165)] = 79.90$ (4) which agrees well with the

measured value of 79.95 (6) as quoted in 1982RuZV and those of 79.88 (8) given in 1975Wa**.

Measured relative values, to the 165-keV γ line, of the X-ray emission intensities can be compared with the value deduced from the decay scheme data:

X-ray	Dasmahapatra	Kumar	Campbell	Pich	Decay scheme
γ - 165,40	100	100			79,90 (4)
K α	80,6 (35)	79,39 (111)			
K β 1	16,10 (69)	14,30 (21)			
K β 2	4,35 (19)				
K X				79,4 (9)	80,3 (8)
K X/ γ			1,010 (25)	0,99 (1)	1,005 (10)

Detailed measured values of the X-ray emissions carried out by 2001Sc08 are also compared with the values deduced from the decay scheme data:

X-ray	E (keV)	Schönfeld (2001Sc08)	Decay scheme
Ll	4,124	0,40 (11)	0,222 (6)
L η + L α	4,52 – 4,65	5,86 (5)	5,78 (13)
L β 1 + L β 4 + L β 3	5,04 – 5,14	4,26 (15)	4,21 (9)
L β 6 + L β 2 + L β 5	5,21 – 5,45	1,07 (4)	1,066 (25)
L γ 5 + L γ 1 + L γ 6	5,62 – 5,88	0,538 (18)	0,565 (15)
L γ 2 + L γ 3 + L γ 4	6,06 – 6,25	0,335 (15)	0,340 (9)
Total L X		12,46 (20)	12,19 (18)
K α 2	33,03	23,05 (28)	22,80 (24)
K α 1	33,44	41,96 (50)	41,9 (4)
K β 1	37,72 – 38,07	12,46 (15)	12,47 (18)
K β 2	38,73 – 38,83	3,11 (4)	3,16 (8)
Total K X		80,6 (6)	80,3 (8)

All the X ray intensities are strongly dependant of the adopted P_K value, the comparisons made in the two tables above show a good agreement between the measured values and those deduced from the decay scheme data. This suggests that the adopted decay scheme is consistent.

6 References

- 1948Po01 - M. L. Pool, N. L. Krisberg, Phys. Rev. **73** (1948) 1035 [$T_{1/2}$]
 1954Mi56 - A. C. G. Mitchell, E. Hebb, Phys. Rev. **95** (1954) 727 [α_K]
 1954Nu12 - R. H. Nussbaum, R. van Leishout, Physica **20** (1954) 440 [α_K]
 1954Pr31 - C. H. Pruett, R. G. Wilkinson, Phys. Rev. **96** (1954) 1340 [α_K , P_K]
 1956Ke23 - B. H. Ketelle, H. Thomas, A. R. Brosi, Phys. Rev. **103** (1956) 190 [$I_{EC}(0)$, α_K , P_K]
 1962Be31 - E.Y. Berlovich, G.M. Bukat, Y.K. Guser, V. V. Nikitin, M. K. Nikitin, Phys. Lett. **2** (1962) 344 [α_K]
 1962Ta03 - J.G.V.Taylor, J.S.Merritt, Bull.Am.Phys.Soc. **7** (1962) 352, XA4 [α , α_K]
 1963Ha07 - J.N.Haag, D.A.Shirley, D.H.Templeton, Phys.Rev. **129** (1965) 1601 [δ]
 1965An07- S.C.Anspach, L.M.Cavallo, S.B.Garfinkel, J.M.R.Hutchinson, C.N.Smith, NP-15663 (1965) [$T_{1/2}$]
 1965Ge04 - J.S.Geiger, R.L.Graham, I.Bergstrom, F.Brown, Nucl.Phys. **68** (1965) 352 [δ]
 1967Be65 - J.A.Bearden, Rev.Mod.Phys. **39** (1967) 78 [E_X]

- 1967HaZX - H. H. Hansen, M. Delabaye, Proc. Symp. Standardization of Radionuclides, Vienna (1966), Intern. 19At. Energy Agency, Vienna (1967) 361 [α_K]
- 1967Ma07 - A. Marelus, P. Sparrman, S.-E. Hågglund, Nucl. Phys. **A95** (1967) 632 [P_K]
- 1968Ad08 - B.Adamowicz, Z.Moroz, Z.Preibisz, A.Zglinski. Acta Phys.Polon. **34**, 529 (1968) [P_K]
- 1968Va08 - E.Vatai, K.Hohmuth. ATOMKI Kozlemen. **10**, 27 (1968) [P_K]
- 198Ha52 - R.S.Hager, E.C.Seltzer, Nucl.Data **A4** (1968) 1 [α , α_K]
- 1971Ar43 - E.A.Aristov, V.A.Bazhenov, Meas.Tech.(USSR) **14** (1971) 1883 [α]
- 1972Ca07 - J. L. Campbell, L. A. McNelles, Nucl. Instr. Meth. **98** (1972) 433 [P_K]
- 1972Em01 - J. F. Emery, S. A. Reynolds, E. I. Wyatt, G. I. Gleason, Nucl. Sci. Eng. **48** (1972) 319 [$T_{1/2}$]
- 1972Sc08 - W.-D. Schmidt-Ott, R. W. Fink, Z. Phys. **249** (1972) 286 [P_K]
- 1973Le29 - J. Legrand, M. Blondel, P. Magnier, Nucl. Instr. Meth. **112** (1973) 101 [α]
- 1973LeZO - J. Legrand, M. Blondel, P. Magnier, C. Perrot, J.-P. Brethon, report CEA-R-4427 (1973)[α]
- 1973MeYE - J. S. Merritt, J. G. V. Taylor, report AECL-4657 (1973) 30 [$T_{1/2}$]
- 1973Ra10 - S. Raman, N. B. Gove, Phys. Rev. C **7** (1973) 1995 [log ft sys.]
- 1975Da08 - B.K.Dasmahapatra, Pramana **4**, 5 (1975) 218 [P_K]
- 1975Ha43 - H. H. Hansen, D. Mouchel, Z. Phys. **A274** (1975) 335 [P_K]
- 1975Mo12 - A.Morinaga, K.Hisatake, J.Phys.Soc.Jap. **38** (1975) 322 [α_K]
- 1975Pl06 - J. Plch, J. Zderadi...ka, O. Dragoun, Intern. J. Appl. Radiat. Isot. **26** (1975) 579 [P_K]
- 1975Wa** - K.F. Walz, E.Funck, H.M.Weiss. PTB-Jahresbericht (Annual Report) (1975) 242 [I γ]
- 1976Ha11 - H.H.Hansen, D.Mouchel, Z.Phys. **A276** (1976) 303 [α , α_K]
- 1976Ha36 - W. Hartl, J. W. Hammer, Z. Phys. **A278** (1976) 183 [P_K]
- 1976Me - J. S. Merritt, J. G. V. Taylor, F. H. Gibson, report AECL-5546 (1976) 32 [$T_{1/2}$]
- 1976Va30 - R.Vaninbroukx, G.Grosse, Int.J.Appl.Radiat.Isotop. **27** (1977) 727 [$T_{1/2}$]
- 1977Ry01 - M.Rysavy, O.Dragoun, M.Vinduska, Czech.J.Phys. **27B** (1977) 538 [α , α_K]
- 1977Sc** - E. Schönfeld, R. Brust, Isotopenpraxis **13** (1977) 311 [α]
- 1978La21 - F.Lagoutine, J.Legrand, C.Bac, Int.J.Appl.Radiat.Isotop. **29** (1978) 269 [$T_{1/2}$]
- 1978Ro22 - F.Rösel, H.M.Friess, K.Alder, H.C.Pauli, At.Data Nucl.Data Tables **21** (1978) 92 [α , α_K]
- 1978Se** - V.A.Sergienko *et al.* Conference on nuclear spectroscopy and atomic nuclear structure. Alma-Ata, USSR. 28 - 31 Mar 1978. (1978) 57 [P_K]
- 1979Ha21 - H. H. Hansen, Z. Phys. **A291** (1979) 43 [d]
- 1980RuZY - A. R. Rutledge, L. V. Smith, J. S. Merritt, report AECL-6692 (1980) [$T_{1/2}$]
- 1982HoZJ - D.D.Hoppes, J. M. R. Hutchinson, F. J. Schima, M. P. Unterweger, NBS-SP-626 (1982) 85 [$T_{1/2}$]
- 1982RuZV - A.R.Rutledge, L.V.Smith, J.S.Merritt, NBS-SP-626 (1982) 5 [I γ , $T_{1/2}$]
- 1982RyZX - A. Rytz, NBS Special Publication 626 (1982) 32 [$T_{1/2}$]
- 1985HaZA - H.H.Hansen, European App.Res.Rept.Nucl.Sci.Technol. 6, No.4 (1985) 777 [α , α_K]
- 1987BeYL - R. B. Begzhanov, K. Sh. Azimov, D. A. Gladyshev, R. D. Magrupov, Sh. A. Mirakhmedov, A. Mukhammadiev, M. Narzikulov, S. Kh. Salimov, Proc. 37th Ann. Conf. Nucl. Spectroscopy Struct. At. Nuclei, Yurmala (1987) 528 [P_K]
- 1988Ko** - A.A. Konstantinov *et al.* Conference on nuclear spectroscopy and atomic nuclear structure. Baku (USSR), 12 - 14 Apr 1988 (1988) 553 [P_K]
- 1988Ri08 - K. Riisager, J. Phys. (London) **G14** (1988) 1301 [P_e]
- 1989Bu12 - T.W.Burrows, Nucl.Data Sheets **57** (1989) 337 [J ^{π}]
- 1992Un01 - M. P. Unterweger, D. D. Hoppes, F. J. Schima, Nucl. Instr. Meth. **A312** (1992) 349 [$T_{1/2}$]
- 1995Au04 - G. Audi, A. H. Wapstra, Nucl. Phys. **A595** (1995) 409 [Q_b]
- 1993Mi20 - M. Minowa *et al.* Phys. Rev. Letters **71**, 25 (1993) 4120 [P_e(0)]
- 1994KU43 - V.Kumar, Kawaldeep, K.Singh. Appl.Radiat.Isot. 45 (1994) 875
- 1995ScZY - E. Schönfeld, PTB-6.33-95-2 (1995)
- 1996Hi14 - M. M. Hindi and R. L. Kozub, Phys. Rev. **C54** (1996) 2709 [Q_e]
- 1996Sc06 - E. Schönfeld, H. Janßen, Nucl. Instr. Meth. **A369** (1996) 527 [ω_K , ω_L , relative K x-ray emission probabilities]
- 1997Ka** - V.D.M.L.Kalyani *et al.* Indian J. Phys. 71A(4) (1997) 493 [P_K]
- 1998Si17 - Nuclear Data Sheets v. 84 (1998) 487 [Log ft , P_e]
- 2000He14 - R. G. Helmer and C. van der Leun, Nucl. Instrum Methods Phys. Res. A450 (2000) 35 [E γ]
- 2001Sc08 - E.Schönfeld, U.Schötzig. Appl. Radiat. Isot. 54 (2001) 785 [X-rays]
- 2005KiZW - T. Kibedi, T.W. Burrows, M.B. Trzhaskovskaya, C.W. Nestor. Proc. Intern. Conf. Nuclear Data for Science and Technology, Santa Fe, New Mexico, 26 September-1 Oct;769 (2005) 268.

**¹⁴⁰Ba - Comments on evaluation of decay data
by V.P. Chechev and N.K. Kuzmenko**

The first DDEP evaluation of ¹⁴⁰Ba decay data was done by R. Helmer in 2003 (2004BeZQ). The current evaluation has been completed in March 2015 with a literature cut-off by the same date. The main changes compared to the DDEP initial evaluation are due to new publications: 2014Un01 (half-life), 2012Wa38 (Q-value), 2008Ki07 (theoretical internal conversion coefficients).

1. DECAY SCHEME

The structure of the adopted decay scheme of ¹⁴⁰Ba is based on the ENSDF evaluation by N. Nica (2007Ni07). ¹⁴⁰Ba disintegrates by β^- emission via ¹⁴⁰La excited levels. The level scheme is incomplete. There are 34 reported levels in ¹⁴⁰La below the β^- decay energy, so some levels in addition to the six reported here may be weakly populated in this decay.

2. NUCLEAR DATA

Q⁻ value is from 2012 mass evaluation by Wang *et al.* (2012Wa38).

The recommended half-life of ¹⁴⁰Ba is based on the experimental results given in Table 1.

Table 1. Experimental values of the ¹⁴⁰Ba half-life (in days)

N	Author(s) and year	Reference	T _{1/2} (d)	Method and comments
1	Simonet <i>et al.</i> (1965)	1965Si17	12.80 (5)	NaI detector
2	Baba <i>et al.</i> (1971)	1971Ba28	12.789 (20)	β counting, proportional counter; original uncertainty is 0.006
3	Debertin <i>et al.</i> (1982)	1982DeYX	12.746 (10)	4 $\pi\gamma$ ionization chamber; <i>omitted</i> as superseded by 5
4	Hoppes <i>et al.</i> (1982)	1982HoZJ	12.753 (2)	4 $\pi\gamma$ ionization chamber; <i>omitted</i> as superseded by 7
5	Walz <i>et al.</i> (1983)	1983Wa26	12.739 (22)	Ge(Li) spectrometer
6	Walz <i>et al.</i> (1983)	1983Wa26	12.751 (5)	4 $\pi\gamma$ ionization chamber
7	Unterweger <i>et al.</i>	1992Un01, 2002Un02	12.7527 (23)	4 $\pi\gamma$ ionization chamber; <i>omitted</i> as superseded by 8
8	Fitzgerald (2012)	2012Fi12	12.7525 (23)	4 $\pi\gamma$ ionization chamber; <i>omitted</i> as superseded by 9
9	Unterweger and Fitzgerald (2014)	2014Un01	12.753 (12)	4 $\pi\gamma$ ionization chamber
Recommended value			12.753 (5)	LWM

The value of 1971Ba28 disagrees with all of the later values, so the evaluators increased its uncertainty from 0.006 to 0.020 (see, also 2004BeZR). The values from 1982DeYX, 1982HoZJ, 1992Un01 and 2002Un02, and 2012Fi12 were not used because they were replaced ultimately by later results of the same laboratory.

The unweighted average of the remained five values is 12.766 (12) d. The weighted average is 12.753 d. The LWRIGHT computer program has chosen the weighted average with the internal uncertainty of 0.0044 d. The external uncertainty is 0.0048 d. The ratio of the reduced $\chi^2 / (\chi^2)_{\text{crit}}$ is 1.17/3.30. The smallest experimental uncertainty is 0.005 d.

The recommended value of ¹⁴⁰Ba half-life is **12.753 (5) days**. It can be compared with earlier evaluations of 12.753 (4) (2004BeZR) and 12.7527 (23) (2007Ni07). The latter value has just been taken from 1992Un01 and 2002Un02, later superseded.

2.1. Beta Transitions

The energies of β^- transitions have been obtained using the Q^- value and the ¹⁴⁰La level energies. The probabilities of β^- -transitions P_{β^-} have been deduced from the $P(\gamma+ce)$ balance at each level of ¹⁴⁰La. The level energies, spins and parities of ¹⁴⁰La (Table 2) are taken from 2007Ni07 (Adopted Levels).

Table 2. ¹⁴⁰La levels populated in ¹⁴⁰Ba β^- -decay

Level	Energy (keV)	Spin and parity	Half-life	P_{β^-} (%)
0	0	3-	1.67858 (21) d	\leq 0.000 01
1	29.9641 (6)	2-	0.25 ns	25.6 (42)
2	43.884 (18)	1-	0.52 ns	35.6 (31)
3	63.1790 (7)	4-		< 0.000 000 01
4	162.6591 (19)	2-	< 0.01 ns	4.14 (31)
5	467.63 (3)	1-	< 7.7 ns	9.71 (12)
6	581.106 (18)	0-		24.94 (50)

The limits for the very weak β^- branches to the ground state and the excited level with energy 63 keV of ¹⁴⁰La are estimated from the $\log ft$ systematics. The β^- transition to the ground state is a non-unique 3rd forbidden transition for which the $\log ft$ systematics of 1998Si17 list only one non-unique 3rd forbidden β^- decay and it has a $\log ft$ of 17.5. If we assume that this class of decays all have $\log ft > 15$, the corresponding I_{β^-} is $< 1 \times 10^{-5}$ %. Similarly, the β^- transition to the excited level with energy 63 keV is unique 3rd forbidden for which 1973Ra10 lists $\log ft$ values of 18.1 and 20.9.

(The corresponding $\log ft$ values in 1998Si17 are 20.7 and 21.4). If we assume that this class has $\log ft > 18$, I_{β^-} is $< 10^{-8}$ %.

2.2. Gamma Transitions and Internal Conversion Coefficients

The multipolarities and mixing ratios δ are from the adopted data in the ENSDF evaluation (2007Ni07). Gamma-ray transition probabilities have been deduced from their gamma-ray emission probabilities and the total ICC(s). The adopted ICC(s) are obtained by using the theoretical values with the BrIcc computer program, accepting the “frozen orbital (no hole)” approximation (2008Ki07).

3. ATOMIC DATA

The atomic data (fluorescence yields, X-ray energies and relative probabilities, and Auger electrons energies and relative probabilities) have been deduced by using the SAISINUC software.

4. ELECTRON EMISSIONS

The energies of the conversion electrons have been obtained from the gamma-ray transition energies and the electron binding energies. The absolute emission probabilities of the conversion electrons have been deduced using recommended P_{γ} and ICC values. The absolute emission probabilities of K and L Auger electrons have been calculated using the EMISSION computer program.

β^- average energies have been calculated using the LOGFT computer program.

5. PHOTON EMISSIONS

5.1. X-ray Emissions

The absolute emission probabilities of La KX- and LX- rays have been calculated using the EMISSION computer program. These probabilities are compared below with the experimental values deduced from the intensities measured relative to the 537-keV gamma ray intensity.

	1991Ch05	1982Ad02	1969Ka33	Adopted
LX	13.3 (6)	8 (2)		13.7 (4)
K α	0.73 (3)	1.6 (1)	2.5 (5)	1.566 (25)
K β	0.36 (3)	0.39 (4)	< 0.49 (7)	0.378 (8)

5.2. Gamma ray emissions

The evaluated energies of gamma rays in ¹⁴⁰La are the energies of the gamma transitions minus the recoil energy.

The measured relative intensities and the adopted values obtained with the LWEIGHT program are listed in Table 4. Many values have been scaled from their original normalizations (2004BeZR). All the values of 1966Mo16 were omitted since they do not have uncertainties. For $\gamma_{29.9}$, $\gamma_{113.6}$, $\gamma_{132.7}$, $\gamma_{162.7}$, and $\gamma_{304.9}$, the smallest uncertainties have been used in the adopted values.

Table 4. Experimental and adopted relative gamma-ray intensities in decay of ¹⁴⁰Ba

γ -ray energy (keV)	1991Ch05	1990Me03	1982Ad02	1977Ge12	1977De34	1976Li06	1975Ha50	1970Ke09	1969Ka33	Adopted
13.9	4.69 (12)	5.0 (7)	4.9 (6)						7.2 (25)	4.71 (12)
29.9	58.4 (10)	61.0 (40)	60 (3)					55 (8)	72 (12)*	58.6 (10)
43.8	0.054 (7)		< 0.007					< 0.005		
63.1		0.000 12 (6)								0.000 12 (6)
99.4		0.000 08 (5)								0.000 08 (5)
113.4	0.072 (6)	0.066 (5)	0.077 (16)					0.074 (8)		0.070 (5)
118.7	0.25 (1)	0.250 (3)	0.27 (3)				1.56 (16)*	0.28 (3)	0.21 (2)	0.248 (7)
132.6	0.81 (2)	0.83 (2)	0.90 (8)				2.14 (31)*	0.84 (5)	0.83 (7)	0.82 (2)
162.6	25.3 (3)	25.45 (29)	28.0 (8)	26.4 (8)	25.5 (3)	25.9 (7)	27.6 (16)	25.1 (10)	28.4 (9)	26.4 (10)
304.9	17.54 (15)	17.6 (2)	17.8 (5)	17.67 (18)	17.63 (21)	18.5 (7)*	17.9 (19)	17.2 (7)	17.3 (7)	17.60 (15)
418.4		0.015 (1)	< 0.04							
423.7	12.65 (12)	12.7 (1)	12.8 (5)	12.73 (14)	12.92 (16)	13.0 (6)	14.8 (12)*	12.7 (5)	12.8 (6)	12.73 (12)
437.6	7.91 (8)	7.91 (4)	7.80 (25)	7.82 (9)	7.91 (16)	8.5 (5)*	8.9 (4)*	7.8 (3)	7.8 (4)	7.89 (4)
467.7	0.29 (3)	< 0.002	< 0.01							
537.2	100 (1)	100.0 (3)	100	100.0 (10)	100.0 (9)	100.0 (23)	100.0 (23)	100.0 (20)	100	100
551.1	0.028 (4)	0.0128 (8)	0.027 (9)							0.020 (8)
848.9			0.02							

* These values have been rejected by the LWEIGHT computer program based on the Chauvenet's criterion.

These relative emission probabilities have been scaled by **0.246 (5)**. This scaling factor has been obtained from the transition intensity balance to the ground state of ^{140}La assuming that the probability of the beta transition to the ground state is $< 1 \times 10^{-5} \%$ (see sect.2.1).

The following normalization factors have been experimentally found: 0.2439 (22) (1977De34), 0.257 (6) (1975Ha50) and 0.236 (5) (1976Li06). In 1977De34 the normalization factor is based on the measured γ -emission rates for five lines and the source activity by 1977De34. Two other factors were determined for the 1596 keV line from ^{140}La decay. The discrepancy between the latter two values is 9 % and may result from difficulties in determining the efficiency at 1596 keV line where there is a lack of efficiency calibration lines. If the three values are averaged, the weighted mean is dominated by the 1977De34 value and is 0.2441 (22) with the internal uncertainty of 0.0019 and the external uncertainty of 0.0036, $\chi^2 / (\chi^2)_{\text{crit}} = 3.6/4.6$.

6. ENERGY CONSERVATION

The total average energy of 1040 (60) keV, for one disintegration, calculated from the current evaluated data corresponds to the available energy of 1048 (8) keV (Q^-) from the mass tables (2012Wa38) confirming the consistency of the decay scheme and the reliability of this evaluation.

7. REFERENCES

- 1965Si17** P. Simonet, G. Boile, G. Simonet, report CEA-R-2461 (1965).
[$T_{1/2}$]
- 1966Mo16** G.A. Moss, D.K. McDaniels, Nucl. Phys. 85 (1966) 513.
[P_γ]
- 1969Ka33** V.G. Kalinnikov, H.L. Ravn, Bull. Acad. Sci. USSR, Phys. Ser. **33** (1970) 1283.
[P_γ]
- 1970Ke09** J. Kern, G. Mauron, Helv. Phys. Acta **43** (1970) 272.
[P_γ]
- 1971Ba28** S. Baba, H. Baba, H. Natsume, J. Inorg. Nucl. Chem. **33** (1971) 589.
[$T_{1/2}$]
- 1973Ra10** S. Raman, N.B. Gove, Phys. Rev. C7 (1973) 1995.
[log ft systematics]
- 1975Ha50** J.T. Harvey, J.L. Meason, J.C. Hogan, H.L. Wright, Nucl. Sci. Eng. **58**, (1975) 31.
[P_γ]
- 1976Li06** C.-C. Lin, J. Inorg. Nucl. Chem. **38** (1976) 1409.
[P_γ]
- 1977De34** K. Debertin, U. Schötzig, K.F. Walz, Nucl. Sci. Eng. **64** (1977) 784.
[P_γ]
- 1977Ge12** R.J. Gehrke, R.G. Helmer, R.C. Greenwood, Nucl. Instrum. Methods **147** (1977) 405.
[P_γ]

- 1982Ad02** I. Adam, N.M. Antoneva, V.B. Brudanin, M. Budzynski, Ts. Vylov, V.A. Dzhashi, A. Zhumamuratov, A.I. Ivanov, V.G. Kalinnikov, A. Kugler, V.V. Kuznetsov, Li Zon Sik, T.M. Muminov, A.F. Novgorodov, Yu.N. Podkopaev, Z.D. Shavgulidze, V.L. Chikhladze, *Izv. Akad. Nauk. SSSR, Ser. Fiz.* **46** (1982) 2.
[P_γ, P_{KX}]
- 1982DeYX** K. Debertain, U. Schötzig, K.F. Walz, report NBS-SP-626 (1982) 101.
[T_{1/2}]
- 1982HoZJ** D.D. Hoppes, J.M.R. Hutchinson, F.J. Schima, M.P. Unterweger, report NBS-SP-626 (1982) 85.
[T_{1/2}]
- 1983Wa26** K.F. Walz, K. Debertain, H. Schrader, *Int. J. Appl. Radiat. Isotop.* **34** (1983) 1191
[T_{1/2}]
- 1990Me03** R.A. Meyer, K.V. Marsh, H. Seyfarth, S. Brant, M. Bogdanovic, V. Paar, *Phys. Rev. C* **41** (1990) 1172.
[P_γ]
- 1991Ch05** B. Chand, J. Goswamy, D. Mehta, N. Singh, P.N. Trehan, *Can. J. Phys.* **69** (1991) 90.
[P_γ]
- 1992Un01** M.P. Unterweger, D.D. Hoppes, F.J. Schima, *Nucl. Instr. Methods* **A312** (1992) 349.
[T_{1/2}]
- 1998Si17** B. Singh, J.L. Rodriguez, S.S. M Wong, J.K. Tuli, *Nucl. Data Sheets* **84** (1998) 487.
[log *ft* systematics]
- 2002Ba85** I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor Jr., P.O. Tikkanen, S. Raman. *At. Data Nucl. Data Tables* 81 (2002) 1.
[Theoretical ICC]
- 2002Un02** M.P. Unterweger, *Appl. Radiat. Isot.* 56 (2002) 125.
[T_{1/2}]
- 2004BeZR** M.M. Bé, V. Chisté, C. Dulieu, E. Browne, V. Chechev, N. Kuzmenko, R. Helmer, A. Nichols, E. Schönfeld, R. Dersch, *Monographie BIPM-5, Vol.1, Bureau International des Poids et Mesures* (2004).
[Previous ¹⁴⁰Ba decay data evaluation]
- 2007Ni07** N. Nica, *Nucl. Data Sheets* 108, 1287 (2007).
[Decay Scheme, ¹⁴⁰La adopted levels and gammas, multipolarities, mixing ratios]
- 2008Ki07** T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor Jr. *Nucl. Instrum. Methods Phys. Res. A* 589 (2008) 202.
[BrIcc computer program]
- 2012Fi12** R. Fitzgerald, *J. Res. Natl. Inst. Stand. Technol.* 117 (2012) 80.
[T_{1/2}]
- 2012Wa38** M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer, *The AME2012 atomic mass evaluation*, *Chin. Phys.* C36 (2012) 1603.
[Q]
- 2014Un01** M.P. Unterweger, R. Fitzgerald, *Appl. Radiat. Isot.* 87 (2014) 92.
[T_{1/2}]

**¹⁴⁰La - Comments on evaluation of decay data
by V.P. Chechev and N.K. Kuzmenko**

The first DDEP evaluation of ¹⁴⁰La decay data was done by R. Helmer in 2003 (2004BeZR). The current evaluation has been completed in May 2015 with a literature cut-off by the same date. The latter evaluators have followed fully the Helmer's approach to the evaluation of gamma ray energies and emission probabilities updating the evaluated values of ICC and other decay characteristics due to new publications of 2004-2014.

1. DECAY SCHEME

The structure of the adopted decay scheme of ¹⁴⁰La is based on the ENSDF (NDS) evaluation by N. Nica (2007Ni07). ¹⁴⁰La disintegrates by β^- emission via ¹⁴⁰Ce excited levels. The level scheme is incomplete as there are many levels in ¹⁴⁰Ce below the β^- decay energy of 3761 keV that are not reported in these decay data, so some other levels may be weakly populated.

2. NUCLEAR DATA

Q^- value is from 2012 mass evaluation by Wang et al. (2012Wa38).

The recommended half-life of ¹⁴⁰La is based on the experimental results given in Table 1.

Table 1. Experimental values of the ¹⁴⁰La half-life (in hours)

N	Author(s) and year	Reference	T _{1/2} (h)	Method and comments
1	Kirby and Salutsky (1954)	1954Ki08	40.224 (20)	β counting, proportional counter
2	Yaffe <i>et al.</i> (1954)	1954Ya02	40.31 (6)	β counting, proportional counter
3	Peppard <i>et al.</i> (1957)	1957Pe09	40.27 (5)	β counting, proportional counter
4	Wille and Fink (1960)	1960Wi10	40 (2)	β counting, proportional counter; <i>rejected</i> on Chauvenet's criterion
5	Simonet <i>et al.</i> (1965)	1965Si17	40.23 (3)	Ionization chamber
6	Karlsson <i>et al.</i> (1967)	1967Ka12	40.2 (2)	Ionization chamber
7	Reynolds <i>et al.</i> (1968)	1968Re04	40.2 (2)	Ionization chamber
8	Debertin <i>et al.</i> (1977)	1977DeYO	40.272 (7)	4 $\pi\gamma$ ionization chamber; <i>omitted</i> as superseded by 13

N	Author(s) and year	Reference	T _{1/2} (h)	Method and comments
9	Davis <i>et al.</i> (1978)	1978Da21	40.232 (67)	4πγ ionization chamber
10	Houtermans <i>et al.</i> (1980)	1980Ho17	40.280 (6)	4πγ ionization chamber
11	Olomo and MacMahon (1980)	1980OI03	40.295 (5)	4πγ ionization chamber
12	Hoppes <i>et al.</i> (1982)	1982HoZJ	40.279 (17)	4πγ ionization chamber; <i>omitted</i> as superseded by 15
13	Walz <i>et al.</i> (1983)	1983Wa26	40.270 (29)	Proportional counter
14	Abzouzi <i>et al.</i> (1989)	1989Ab18	40.284 (5)	4πγ ionization chamber
15	Unterweger <i>et al.</i>	1992Un01, 2002Un02, 2012Fi12	40.293 (12)	4πγ ionization chamber; <i>omitted</i> as superseded by 17
16	Adam <i>et al.</i> (2002)	2002Ad04	40.34 (4)	Ge(Li) detector
17	Unterweger and Fitzgerald (2014)	2014Un01	40.294 (7)	4πγ ionization chamber
Recommended value			40.286 (5)	LWM

The values 8, 12 and 15 were not used because they were replaced ultimately by later results of the same laboratory. The value 4 was rejected on Chauvenet's criterion.

The weighted average of the 13 remaining values is 40.2864 h. The internal uncertainty is 0.0027 h. The external uncertainty is 0.0036 h. The ratio of the reduced $\chi^2 / \chi^2_{\text{crit}}$ is 1.7/2.1. The smallest experimental uncertainty is 0.005 h.

The recommended value of ¹⁴⁰La half-life is **40.286 (5) h, or 1.678 58 (21) d**. It can be compared with earlier evaluations of 1.678 50 (17) d (2004BeZR) and 1.678 55 (12) d (2007Ni07).

2.1. Beta Transitions

The energies of β^- transitions have been obtained using the Q^- value and the ¹⁴⁰Ce level energies (Table 2) adopted from Helmer's least-squares fit to the γ -ray energies (see section 5.2 and 2004BeZR). The level spins, parities and half-lives of ¹⁴⁰Ce are adopted from 2007Ni07.

The probabilities of β^- -transitions P_{β^-} have been deduced from the $P(\gamma+ce)$ balance at each level of ¹⁴⁰Ce.

Table 2. ¹⁴⁰Ce levels populated in ¹⁴⁰La β^- -decay

Level	Energy(keV)	Spin and parity	Half-life	P_{β^-} (%)
0	0	0+	Stable	-
1	1596.213 (13)	2+	0.0916 (19) ps	4.5 (6)
2	1903.29 (4)	0+	0.40 (3) ns	-
3	2083.236 (14)	4+	3.45 (3) ns	20.8 (6)
4	2107.83 (4)	6+	7.3 (15) μ s	-
5	2347.868 (14)	2+	≤ 0.2 ns	5.03 (12)
6	2349.789 (9)	5+	≤ 10 ns	0.262 (22)
7	2411.997 (14)	3+	55 (15) ps	44.8 (4)
8	2464.055 (21)	3-	0.10 (2) ps	5.60 (7)
9	2480.91 (6)	4+	3.2 (3) ns	1.14 (2)
10	2515.749 (16)	3+		5.80 (4)
11	2521.414 (14)	2+	≤ 0.1 ns	11.11 (9)
12	2547.205 (26)	1+		0.636 (7)
13	2899.56 (7)	2+		0.112 (6)
14	3000.88 (9)	2+		0.085 (9)
15	3118.53 (10)	2+		0.027 (1)
16	3319.56 (24)	2+		0.0039 (3)
17	3394.82 (9)	(4-)		0.020 (4)
18	3473.55 (18)	3-		0.052 (7)
19	3520.8 (2)	(4+)		0.011 (3)

As discussed in 2004BeZR the β^- branches to the levels at 0, 1903 and 2107 keV are non unique 3rd forbidden. The $\log ft$ systematics of 1998Si17 give only one value, 17.5, for this class of β^- decays. From the data of 1998Si17, it is reasonable to assume a lower limit of $\log ft > 15$ for this class. The corresponding P_{β^-} limits are then $< 1 \times 10^{-4}$ %; $< 1 \times 10^{-5}$ %; and

$< 1 \times 10^{-5}$ %, respectively. Although there have been many analyses of the β^- spectrum, only 1966Dz05 has reported a branch to the ground state. Their intensity of $5(2) \times 10^{-5}$ % is compatible with the limit from the $\log ft$ systematics; however, since others have not seen this branch, this value is assumed to be too large. In any case, the value is negligible in determining the normalization of the γ -ray emission probabilities. These three P_{β^-} are all set to zero in the adopted decay scheme.

The average β^- energies and the $\log ft$ are from the LOGFT program (<http://www.nndc.bnl.gov/logft/>).

2.2. Gamma Transitions and Internal Conversion Coefficients

The γ -ray transition energies have been obtained as given in sect. 5.2.

The gamma-ray transition probability for $\gamma_{4,3}$ (24.6 keV) has been deduced from the intensity balance for level at 2107.8 keV assuming that corresponding $P_{\beta^-} < 1 \times 10^{-5}$ % (see sect. 2.1). The deduced value of 0.480 (11) % can be compared with the value for $P_{\gamma+ce} = 0.32(4)$ %, reported by 1967Ka12, which is presumably the contribution of the emission probability of L - conversion electrons.

The probability of the 1903-keV E0 transition ($\gamma_{2,0}$) of 0.0146 (15) % has been adopted from measurements of conversion electrons in 1991Ch05. Considering with this value the intensity balance at the level 2 (1903 keV) it is important to be defined with the value of $P_{\gamma_{11,2}}$ (618 keV) as the measured values of the relative $\gamma_{11,2}$ (618 keV)-ray intensity are very discrepant (Table 5). When using the values from 1980Ka32 and 1991Ch05 (the average value $P_{\gamma_{11,2}}$ (618 keV) = 0.041 (3) %), we have the significant intensity imbalance at the level 2. If we use for $P_{\gamma_{11,2}}$ (618 keV) the lower value of 0.014 (3) % measured in 1982Ad02, the balance becomes exact. Therefore we adopt the value of 0.014 (3) % for $P_{\gamma_{11,2}}$ (618 keV).

The remaining gamma-ray transition probabilities have been obtained from their gamma-ray emission probabilities and the total ICC(s).

The adopted ICC(s) are obtained from the theoretical values by using the adopted mixing ratios δ and the BrIcc computer program (<http://bricc.anu.edu.au>), accepting the “frozen orbital (no hole)” approximation (2008Ki07). The multipolarities and mixing ratios δ are from 2007Ni07.

3. ATOMIC DATA

The atomic data (fluorescence yields, X-ray energies and relative probabilities, and Auger

electrons energies and relative probabilities) have been deduced by using the SAISINUC software (1996Be46).

4. ELECTRON EMISSIONS

The energies of the conversion electrons have been obtained from the gamma-ray transition energies and the electron binding energies. The absolute emission probabilities of the conversion electrons have been deduced using recommended $P\gamma$ and ICC values. The absolute emission probabilities of K and L Auger electrons have been calculated using the EMISSION computer program (2000Sc47).

The number of electron-positron pairs per 100 disintegrations (Pcp) have been obtained using the adopted α_{π} values.

β^- average energies have been calculated using the LOGFT computer program (<http://www.nndc.bnl.gov/logft/>).

5. PHOTON EMISSIONS

5.1. X-Ray Emissions

The absolute emission probabilities of Ce KX- and LX- rays have been calculated using the EMISSION computer program. In Table 3 these calculated absolute emission probabilities of Ce KX- rays are compared with values deduced from the experimental relative intensities and the normalization factor (0.9540) and scaling factor (1.027), see sect. 5.2.

Table 3. Experimental and recommended Ce KX - ray emission probabilities (%) in decay of ¹⁴⁰La

KX-ray	1970Ka18	1982Ad02	1991Ch05	Recommended (calculated)
K α	2.4 (7)	1.73 (6)	1.68 (4)	1.673 (15)
K β	0.35 (8)	0.44 (2)	0.387 (14)	0.409 (7)

5.2. Gamma Ray Emissions

The γ -ray energies were determined from the reported values in Table 4. (2004BeZR). All of these 187 energies were entered into a simultaneous least-squares fit to determine the energies of the 19 excited levels. The adopted γ -ray transition energies were then computed from the differences between the level energies, with the corrections for recoil (2004BeZR).

The gamma-ray emission probability for $P_{\gamma+ce}$ for $\gamma_{4,3}$ (24.6 keV) has been deduced from its gamma-ray transition probability (see section 2.2) and the total ICC α_T .

The other gamma-ray emission probabilities have been deduced from the relative γ -ray intensities which were determined from the data in Table 5. Several of these sets of data were published as emission probabilities and have been scaled by Helmer (2004BeZR) to obtain values relative to the 1596-keV γ -ray. The LWEIGHT program using the limitation of relative statistical weight method (LWM) was used to compute the average values (Browne, E. Limitation of Relative Statistical Weights, a method for evaluating discrepant data. In: INDC(NDS)-363, IAEA, March 1998. MacMahon, T.D., and Browne, E. LWEIGHT, a computer program to calculate averages, Version 1.3. March 2000). In this calculation, if a particular value contributes more than 50% of the relative weight and the initial fit has a reduced- χ^2 of more than the critical reduced- χ^2 for the number of input values, the uncertainty of the most precise value is increase to reduce its relative weight to 50%. The critical reduced- χ^2 values are: 6.6 for 2 input values; 4.6 for 3; 3.8 for 4; 3.3 for 5; 3.0 for 6; 2.5 for 9; 2.4 for 10; 2.3 for 11; and 2.2 for 12 or 13. Some values have been deleted from the averaging, as indicated in the table and the evaluator (Helmer, 2004BeZR) has arbitrarily increased a few input uncertainties.

At the time when many of these measurements were made, there was a lack of good Ge detector efficiency calibration standards in the region of 1596 keV. Therefore, the evaluator (Helmer, 2004BeZR) has introduced an energy-dependent scaling factor based on the emission probabilities from 1977De34 for thirteen lines from 266 to 2521 keV. This factor, which is shown in Table 5 and varies by 3%, corrects for this assumed systematic deviation of the Ge detector efficiencies.

The total γ -ray feeding of the ground state is set to 100%, with no direct β^- decay, to obtain a normalization factor of **0.9540 (5)** to convert these relative emission probabilities to absolute probabilities as given in the last column of Table 5.

Table 4. Experimental and adopted γ -ray energies

1964Re09	1967Ka12	1968Ba18	1968Gu05	1970Ka18	1970Ke06	1972GeZG	1978Ar28	1979Bo26	1980Ka32	1982Ad02	Adopted (keV)
	24.595 (4)										24.595 (4)
	64.130 (7)	64.135 (10)									64.129 (4)
	68.916 (6)	69.0 (3)									68.923 (5)
	109.417 (6)	109.418 (7)				109.47 (20)				109.422 (11)	109.417 (4)
	131.122 (8)	131.121 (8)				131.15 (20)			130.97 (20)	131.117 (8)	131.121 (4)
	173.550 (11)	173.536 (12)				173.50 (20)			173.49 (17)	173.543 (9)	173.546 (5)
241.97 (3)	241.961 (22)	241.966 (12)				241.90 (8)	241.88 (10)		242.06 (9)	241.933 (30)	241.959 (6)
266.52 (6)	266.547 (22)	266.551 (14)				266.61 (6)	266.58 (10)		266.67 (7)	266.543 (12)	266.554 (5)
		306.9 (2)				306.5 (4)			307.1 (2)	306.9 (2)	307.08 (4)
328.789 (15)		328.768 (12)	328.752 (30)		328.745 (15)	328.76 (5)	328.80 (10)	328.746 (25)	328.78 (5)	328.762 (8)	328.761 (4)
	397.8 (3)	397.79 (11)				397.66 (10)			397.8 (1)	397.52 (5)	397.674 (6)
432.55 (8)	432.62 (6)	432.530 (29)			432.490 (20)	432.52 (4)	432.51 (10)		432.66 (4)	432.493 (12)	432.513 (8)
				438.5 (4)					438 (1)	438.5 (5)	438.178 (6)
									445 (1)	445.5 (5)	444.57 (4)
487.027 (24)	487.042 (29)	487.029 (19)	487.032 (30)		486.995 (30)	487.009 (30)	487.09 (10)	487.15 (25)	486.99 (3)	487.021 (12)	487.022 (6)
		618.2 (7)				617.7 (3)			618.2 (1)	618.12 (5)	618.12 (4)
752.42 (33)	751.75 (8)	751.83 (8)				751.655 (35)	751.66 (10)		751.65 (4)	751.637 (18)	751.653 (7)
815.82 (10)	815.85 (7)	815.80 (9)			815.735 (40)	815.775 (30)	815.80 (10)		815.78 (4)	815.772 (19)	815.781 (6)
867.9 (5)	867.87 (15)	867.82 (14)				867.842 (35)	867.85 (10)		867.80 (4)	867.856 (20)	867.839 (16)
	919.63 (15)	919.5 (2)				919.54 (4)	919.63 (10)		919.48 (6)	919.550 (23)	919.533 (10)

1964Re09	1967Ka12	1968Ba18	1968Gu05	1970Ka18	1970Ke06	1972GeZG	1978Ar28	1979Bo26	1980Ka32	1982Ad02	Adopted (keV)
924.1 (6)	925.24 (9)	925.20 (17)				925.188 (35)	925.21 (10)		925.14 (6)	925.189 (21)	925.198 (7)
	950.9 (3)	951.1 (4)		951.4 (4)		951.00 (6)			950.95 (6)	950.987 (26)	950.988 (20)
										992.9 (5)	992.64 (18)
						1045.2 (3)			1045.0 (1)	1045.05 (24)	1045.02 (9)
						1097.2 (3)			1097.2 (2)	1097.20 (23)	1097.58 (9)
									1303.3 (1)	1303.5 (4)	1303.34 (7)
						1404.5 (2)			1404.9 (2)	1405.20 (17)	1404.66 (9)
1596.34 (25)	1596.49 (24)	1596/6 (2)	1596.20 (4)		1596.170 (25)	1596.17 (6)	1596.22 (10)		1596.17 (6)	1596.210 (35)	1596.203 (13)
										1877.29 (19)	1877.33 (18)
	1903.15 (30)								1903 (1)		1903.28 (4)
						1924.2 (3)			1924.4 (1)	1924.62 (13)	1924.5 (2)
									2082.9 (2)	2083.2 (5)	2083.219 (14)
	2348.1 (7)	2348.8 (6)				2347.80 (6)			2347.82 (6)	2347.88 (5)	2347.847 (14)
				2465.3 (8)					2464.0 (1)	2464.1 (5)	2464.031 (20)
2519.7 (34)	2521.7 (5)	2522.2 (4)				2521.32 (6)	2522.03 (10)		2521.36 (6)	2521.40 (5)	2521.390 (14)
	2547.1 (8)	2548.6 (8)		2547.5 (6)		2547.14 (6)			2547.19 (7)	2547.34 (11)	2547.180 (23)
	2900 (2)	2899.7 (5)		2899.7 (8)		2899.5 (2)			2899.5 (2)	2899.61 (16)	2899.53 (7)
	3119 (2)	3118.3 (7)		3119.0 (8)		3118.52 (15)			3118.4 (2)	3118.51 (16)	3118.49 (10)
	3322 (4)	3319.7 (25)		3319.6 (9)		3319.4 (6)			3319.3 (3)	3320.4 (6)	3319.52 (24)

Table 5. Experimental and adopted relative gamma-ray intensities in decay of ^{140}La . Part 1: references from 1962 to 1975

Energy (keV)	1962Ha14	1967Ka12	1968Ba18	1969KuZV	1970Ka18	1974HeYW	1975Ha50
64					~ 0.01		
68			0.065 (13)		0.064 (16)		
109		0.50 (20)	0.27 (4)	0.23 (2)	0.210 (15)	0.17 (4)	0.20 (4)
131		1.05 (15)	0.61 (9)	0.47 (3)	0.50 (3)	0.42 (5)	0.58 (4)
173			0.13 (5)		0.130 (20)	0.60 (20)	
241		0.83 (10)	0.45 (6)	0.58 (6)	0.410 (30)	0.51 (8)	0.66 (3)
266		0.83 (10)	0.56 (6)	0.53 (4)	0.490 (30) @	0.50 (5)	0.34 (3)
307			0.022 (11)		0.035 (17)		
328		25.4 (20)	21.4 (11)	22.4 (4)	19.4 (1) @	19.6 (13)	18.8 (5)
397			0.054 (25)		0.110 (35)	0.12 (3)	
432		3.5 (3)	3.11 (16)	3.06 (9)	2.85 (15)	2.94 (20)	3.0 (2)
438					0.021 (10)		
444					~ 0.25		
487		49.6 (32)	49.4 (25)	48.2 (5)	45.0 (2) @	44.7 (30)	39.7 (5)
618		0.4 (3)	0.044 (22)		~ 0.045		
751		4.5 (4)	4.40 (22)	4.66 (23)	4.40 (20)	4.5 (3)	4.9 (2)
815		23.5 (20)	24.1 (12)	24.9 (2)	23.5 (7)	24.2 (15)	26.8 (11)
867		5.6 (5)	5.64 (28)	5.91 (24)	5.60 (30)	5.7 (3)	6.5 (1)

919		2.5 (6)	2.73 (16)	2.59 (10)	2.64 (16)	2.89 (20)	3.4 (2)
925		6.8 (6)	7.24 (43)	6.94 (21)	7.10 (30)	7.2 (4)	7.9 (3)
950		0.8 (3)	0.56 (5)	0.62 (9)	0.550 (30)	0.56 (4)	
992							
1045							
1097							
1303							
1405							
1596	100.	100.	100.	100.	100.	100.	100.
1877						0.05 (2)	
1903							
1924						0.023 (5)	
2083							
2347	0.86 (17)	1.0 (2)	0.901 (45)	0.85 (6)	0.90 (6)	0.89 (6)	
2464					0.0018 (6) #		
2521	3.0 (6)	3.5 (2)	3.52 (18)	3.37 (10)	3.60 (18)	3.59 (18)	4.9 (4)
2547		0.11 (2)	0.122 (9)		0.110 (7)	0.110 (6)	
2899	0.082 (17)	0.060 (10)	0.070 (5)		0.065 (6)	0.073 (8)	
3118	0.035 (10)	0.030 (10)	0.027 (3)		0.027 (4)	0.028 (3)	
3320			0.008 (4)		0.0047 (15)	0.050 (3)	

Table 5. Experimental and adopted relative gamma-ray intensities in decay of ¹⁴⁰La. Part 2: references from 1976 to 1991

Energy (keV)	1976Li06	1977De34	1977Ge12	1978Ar28	1980Ka32	1982Ad02	1991Ch05	LWM.	Reduced χ^2	Scaling factor	Adopted	Emission probability (%)
64						0.011 (4)	0.015 (2)	0.0142 (18)		1.027	0.0146 (20)	0.014 (2)
68					0.070 (16)	0.080 (6)	0.079 (2)	0.0785 (19)		1.027	0.0806 (20)	0.077 (2)
109	0.20 (9)				0.170 (10) @	0.220 (10)	0.230 (4)	0.221 (6)	1.9	1.027	0.227 (6)	0.217 (6)
131	0.46 (9)				0.44 (1) @	0.48 (3)	0.49 (1) *	0.479 (15)	2.9	1.027	0.492 (15)	0.47 (1)
173					0.120 (10)	0.110 (10)	0.133 (4)	0.129 (5)	2.2	1.027	0.132 (5)	0.126 (5)
241	0.52 (18)	0.6 (1)		0.51 (9)	0.450 (10)	0.460 (30)	0.434 (8) *	0.445 (10)	2.7	1.027	0.457 (10)	0.436 (10)
266	0.53 (6)	0.7 (1)		0.50 (3)	0.520 (10)	0.500 (30)	0.488 (8)	0.502 (9)	2.3	1.027	0.516 (9)	0.492 (9)
307					0.022 (6)	0.020 (5)	0.026 (7)	0.022 (3)		1.027	0.023 (5)	0.022 (5)
328	21.2 (6)	22 (2)	21.46 (22)	21.5 (6)	21.5 (4)	21.7 (4)	21.1 (3)	21.2 (3)	5.0	1.027	21.8 (3)	20.8 (3)
397					0.078 (3)	0.070 (5)	0.077 (5)	0.0763(24)	1.15	1.027	0.078 (3)	0.075 (3)
432	3.0 (4)	3.5 (2)	3.08 (3)	2.96 (16)	3.05 (3)	2.97 (15)	3.04 (3)	3.056 (17)	1.01	1.027	3.14 (3)	3.00 (3)
438					0.006 (3) *	<0.0014	0.041 (10)	0.0 18 (10)	4.1	1.027	0.018 (10)	0.017 (10)
444					0.005 (3)	0.0036 (12)	0.003 (1)	0.0034(7)		1.027	0.004 (1)	0.003 (1)
487	46.2 (11)	47 (2)	47.7 (5)	47.3 (9)	46.6 (9)	46.4 (8)	47.7 (6)	47.0 (4)	2.6	1.027	48.3 (5)	46.1 (5)
618					0.049 (6)	0.014 (3)	0.039 (4)	0.028 (14)	18.75	1.015	0.014 (3) ^a	0.014 (3)
751	4.40 (17)	4.6 (1)	4.65 (5)	4.37 (22)	4.45 (5)	4.36 (16)	4.54 (4)	4.536 (25)	1.10	1.015	4.60 (5)	4.39 (5)

Energy (keV)	1976Li06	1977De34	1977Ge12	1978Ar28	1980Ka32	1982Ad02	1991Ch05	LWM.	Reduced χ^2	Scaling factor	Adopted	Emission probability (%)
815	23.8 (6)	24.2 (4)	24.85 (25)	24.1 (5)	24.0 (4)	23.5 (7)	24.4 (2)	24.49 (13)	1.43	1.015	24.86 (20)	23.72 (20)
867	6.0 (5)	5.8 (3)	5.90 (6)	5.69 (10)	5.69 (6)	5.56 (19)	5.77 (7)	5.77 (3)		1.015	5.85 (7)	5.58 (7)
919	3.1 (4)	2.6 (2)	2.91 (4)	2.57 (14)	2.83 (4)	2.80 (9)	2.79 (3)	2.812 (24)	1.65	1.015	2.86 (3)	2.73 (3)
925	7.3 (8)	7.2 (3)	7.42 (8)	7.25 (16)	7.26 (8)	7.10 (21)	7.23 (7)	7.27 (4)		1.015	7.38 (7)	7.04 (7)
950	0.63 (12)	0.67 (6)			0.553 (7)	0.56 (3)	0.544 (7)	0.549 (5)		1.015	0.557 (7)	0.531 (7)
992						0.009 (3)	0.014 (5)	0.0103 (26)		1.015	0.010 (3)	0.010 (3)
1045					0.024 (4)	0.016 (4)	0.026 (15)	0.0202 (29)	1.08	1.015	0.021 (4)	0.020 (4)
1097					0.024 (5)	0.022 (5)	0.024 (5)	0.0233 (29)		1.015	0.024 (5)	0.023 (5)
1303					0.046 (6)	0.050 (7)	0.044 (7)	0.047 (4)		1.000	0.047 (6)	0.045 (6)
1405					0.066 (9)	0.068 (8)	0.062 (7)	0.065 (5)		1.000	0.065 (8)	0.062 (8)
1596	100.0	100.0 (3)	100 (1)	100.0 (3)	100.0	100.	100.0 (15)	100.0		1.000	100.0	95.40 (5)
1877						0.042 (6)	0.043 (4)	0.043 (3)		1.000	0.043 (6)	0.041 (6)
1903											-	-
1924					0.014 (3)	0.006 (2)	0.014 (2)	0.0115 (28)	5.0	1.000	0.012 (3)	0.011 (3)
2083					0.045 (3)	0.007 (2) #	0.031 (2)	0.038 (7)	11	1.000	0.038 (7)	0.036 (7)
2347		0.90 (4)	0.891 (16)		0.89 (1)	0.89 (3)	0.89 (3)	0.890 (7)		0.996	0.886 (16)	0.845 (16)
2464					0.012 (1)	0.008 (1)	0.012 (2)	0.0102 (14)	4.4	0.996	0.0102 (14)	0.0097 (13)
2521		3.5 (2)	3.62 (7)	3.65 (18)	3.58 (5)	3.61 (9)	3.63 (4)	3.591 (25)		0.996	3.58 (5)	3.41 (5)
2547			0.109 (3)		0.105 (2)	0.109 (5)	0.106 (3)	0.1070 (13)		0.996	0.107 (2)	0.102 (2)

Energy (keV)	1976Li06	1977De34	1977Ge12	1978Ar28	1980Ka32	1982Ad02	1991Ch05	LWM.	Reduced χ^2	Scaling factor	Adopted	Emission probability (%)
2899			0.069 (1)		0.070 (1)	0.069 (3)	0.070 (2)	0.0695 (6)		0.996	0.069 (1)	0.066 (1)
3118			0.027 (1)		0.027 (1)	0.028 (2)	0.026 (1)	0.0269 (5)		0.996	0.027 (1)	0.026 (1)
3320					0.0040 (3)	0.0045 (4)	0.0040 (3)	0.00413 (19)		0.996	0.0041 (3)	0.0039 (3)

Comments on Table 5:

* Uncertainties were increased in LWM analysis to reduce relative weight to 50%; this change is only made if the reduced- χ^2 is greater than the associated critical value. These changes were: 131 keV, 1991Ch05 0.010 to 0.012; 241, 1991Ch05 0.008 to 0.0087; and 438 keV, 1980Ka32 0.003 to 0.007.

@ Uncertainties were increased by evaluator due to large deviation from average. These changes were: 109 keV, 1980Ka32 0.01 to 0.02; 131, 1980Ka32 0.01 to 0.02; 266, 1970Ka18 0.03 to 0.06; 328, 1970Ka18 0.1 to 0.3; and 487, 1970Ka18 0.2 to 0.5.

Deleted from calculation.

^a Adopted from 1982Ad02. See section 2.2.

6. ENERGY CONSERVATION

The total average energy of 3763 (10) keV, for one disintegration, calculated from the current evaluated data corresponds very well to the available energy of 3760.9 (18) keV (Q^-) from the mass tables (2012Wa38) confirming the consistency of the decay scheme and the reliability of this evaluation.

7. REFERENCES

- 1954Ki08** H. W. Kirby, M. L. Salutsky, Phys. Rev. **93**, (1954) 1051. [T_{1/2}]
- 1954Ya02** L. Yaffe, H. G. Thode, W. F. Merritt, R. C. Hawkings, F. Brown, R. M. Bartholomew, Can. J. Chem. **32** (1954) 1017. [T_{1/2}]
- 1957Pe09** D. F. Peppard, G. W. Mason, S. W. Moline, J. Inorg. Nuclear Chem. **5** (1957) 141. [T_{1/2}]
- 1960An05** S. F. Antonova, S. S. Vasilenko, M. G. Kaganskii, D. L. Kaminskii, Soviet Phys. JETP **11** (1960) 554. [P γ]
- 1960Wi10** R. G. Wille, R. W. Fink, Phys. Rev. **118** (1960) 242. [T_{1/2}]
- 1962Ha14** P. G. Hansen, K. Wilsky, Nucl. Phys. **30** (1962) 405. [P γ]
- 1964Re09** J. J. Reidy, report TID-21826 (1964). [E γ]
- 1965Si17** P. Simonet, G. Boile, G. Simonet, report CEA-R-2461 (1965). [T_{1/2}]
- 1966Ba36** H. W. Baer, J. J. Reidy, M. L. Wiedenbeck, Nucl. Phys. **86** (1966) 332. [E γ]
- 1966Dz09** B. S. Dzelepov, N. N. Zhukovskii, A. G. Maloyan, V. P. Prikhodtseva, Bull. Acad. Sci. USSR, Phys. Ser. **30** (1967) 410. [P γ]
- 1966Ha20** G. I. Harris, D. V. Breitenbecher, Phys. Rev. **145** (1966) 866. [P γ]
- 1967Ka12** S.-E. Karlsson, B. Svahn, H. Pettersson, G. Malmsten, E. Y. De Aisenberg, Nucl. Phys. A100 (1967) 113. [E γ , P γ]
- 1968Ba18** H. W. Baer, J. J. Reidy, M. L. Wiedenbeck, Nucl. Phys. A113 (1968) 33. [E γ , P γ]
- 1968Gu05** R. Gunnink, R. A. Meyer, J. B. Niday, R. P. Anderson, Nucl. Instr. Methods **65**(1968)26. [E γ]
- 1968Re04** S. A. Reynolds, J. F. Emery, E. I. Wyatt, Nucl. Sci. Eng. **32** (1968) 46. [T_{1/2}]
- 1969GuZV** R. Gunnink, J. B. Niday, R. P. Anderson, R. A. Meyer, report UCID-15439 (1969). [P γ]
- 1970Ka18** V. G. Kalinnikov, H. L. Ravn, H. G. Hansen, N. A. Lebedev, Bull. Acad. Sci. USSR, Phys. Ser. **34** (1971) 815. [E γ , P γ]
- 1970Ke06** J. Kern, Nucl. Instr. Methods **79** (1970) 233. [E γ]
- 1972GeZG** R. J. Gehrke, report ANCR-1088, (1972) 392. [E γ]
- 1974HeYW** R. L. Heath, report ANCR-1000-2 (1974). [P γ]
- 1975Ha50** J. T. Harvey, J. L. Meason, J. C. Hogan, H. L. Wright, Nucl. Sci. Eng. **58** (1975) 431. [P γ]

- 1976Li06** C.-C. Lin, J. Inorg. Nucl. Chem. 38 (1976) 1409. [P γ]
- 1977De34** K. Debertain, U. Schötzig, K. F. Walz, Nucl. Sci. Eng. 64 (1977) 784. [P γ]
- 1977DeYO** K. Debertain, U. Schötzig and K. F. Walz, INDC(Ger)-10/L+Special (1977) 83. [T_{1/2}]
- 1977Ge12** R. J. Gehrke, R. G. Helmer, R. C. Greenwood, Nucl. Instrum. Methods 147 (1977) 405. [P γ]
- 1978Ar28** G. Ardisson, Nucl. Instr. Methods 151 (1978) 505. [E γ , P γ]
- 1978Da21** M. C. Davis, W. C. Bowman, J. C. Robertson, Int. J. Appl. Radiat. Isotop. 29 (1978) 331. [T_{1/2}]
- 1979Bo26** H. G. Börner, W. F. Davidson, J. Almeida, J. Blachot, J. A. Pinston, P. H. M. Van Assche, Nucl. Instr. Methods 164 (1979) 579. [E γ]
- 1980Ho17** H. Houtermans, O. Milosevic, F. Reichel, Int. J. Appl. Radiat. Isotop. **31** (1980) 153. [T_{1/2}]
- 1980Ka32** R. Kaur, A. K. Sharma, S. S. Sook, P. N. Trehan, J. Phys. Soc. Japan. 49 (1980) 2122. [E γ , P γ]
- 1980O103** J. B. Olomo, T. D. MacMahon, J. Phys. (London) G6 (1980) 367. [T_{1/2}]
- 1982Ad02** I. Adam, N. M. Anton²eva, V. B. Brudanin, M. Budzynski, Ts. Vylov, V. A. Dzhashi, A. Zhumamuratov, A. I. Ivanov, V. G. Kalinnikov, A. Kugler, V. V. Kuznetsov, Li Zon Sik, T. M. Muminov, A. F. Novgorodov, Yu. N. Podkopaev, Z. D. Shavgulidze, V. L. Chikhladze, Izv. Akad. Nauk SSSR, Ser. Fiz. 46 (1982) 2. [E γ , P γ]
- 1982HoZJ** D. D. Hoppes, J. M. R. Hutchinson, F. J. Schima, M. P. Unterweger, report NBS-SP-626 (1982) 85. [T_{1/2}]
- 1983Wa26** K. F. Walz, K. Debertain, H. Schrader, Int. J. Appl. Radiat. Isotop. 34 (1983) 1191. [T_{1/2}]
- 1989Ab18** A. Abzouzi, M. S. Antony, V. B. Ndocko Ndongue, J. Radioanal. Nucl. Chem. **137** (1989) 381. [T_{1/2}]
- 1991Ch05** B. Chand, J. Goswamy, D. Mehta, N. Singh, P. N. Trehan, Can. J. Phys. 69 (1991) 90. [P γ]
- 1992Un01** M. P. Unterweger, D. D. Hoppes, F. J. Schima, Nucl. Instr. Methods A312 (1992) 349. [T_{1/2}]
- 1996Be46** M.M. Bé, B. Duchemin, J. Lame, Nucl. Instrum. Methods Phys. Res. A369, 523 (1996) [SAISINUC software]
- 1998Si17** B. Singh, J. L. Rodriguez, S. S. M. Wong, J. K. Tuli, Nucl. Data Sheets **84** (1998) 565. [log ft systematics]
- 1999ScZX** E. Schönfeld, H. Janßen, laboratory report PTB-6.11-1999-1 (Feb. 1999). [P_X]
- 2000Sc47** E. Schönfeld and H. Janßen, Appl. Radiat. Isot. 52 (2000) 595. [EMISSION software]
- 2002Un02** M. P. Unterweger, Appl. Radiat. Isot. 56 (2002) 125. [T_{1/2}]

- 2002Ad04** J.Adam, A.G.Belov, R.Brandt, P.Chaloun, M.Honusek, V.G.Kalinnikov, M.I.Krivopustov, B.A.Kulakov, E.-J.Langrock, V.S.Pronskikh, A.N.Sosnin, V.I.Stegailov, V.M.Tsupko-Sitnikov, J.-S.Wan, W.Westmeier, Nucl.Instrum.Methods Phys.Res. B187 (2002) 419. [T_{1/2}]
- 2004BeZR** M.M.Bé, V.Chisté, C.Dulieu, E.Browne, V.Chechev, N.Kuzmenko, R.Helmer, A.Nichols, E.Schönfeld, R.Dersch, Monographie BIPM-5, Vol.1, Bureau International des Poids et Mesures (2004)
[Previous ¹⁴⁰Ba decay data evaluation, ¹⁴⁰La adopted levels and gammas]
- 2007Ni07** N.Nica, Nucl.Data Sheets 108, 1287 (2007)
[Decay Scheme, multipolarities, mixing ratios]
- 2008Ki07** T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor Jr. Nucl. Instrum. Methods Phys. Res. A589 (2008) 202. [BrIcc computer program]
- 2012Fi12** R.Fitzgerald, J.Res.Natl.Inst.Stand.Technol. 117 (2012) 80. [T_{1/2}]
- 2012Wa38** M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer, The AME2012 atomic mass evaluation, Chin. Phys. C36 (2012) 1603. [Q]
- 2014Un01** M.P. Unterweger, R., Fitzgerald. Appl. Radiat. Isot. 87 (2014) 92. [T_{1/2}]

**¹⁴¹Ce - Comments on evaluation of decay data
by E. Schönfeld and V.P. Chechev**

This evaluation was completed in 1998; it has been updated in February 2012. The literature available by this latter date has been included.

1. Decay Scheme and Decay Energy

¹⁴¹Ce decay scheme is complete as there are no other excited levels of ¹⁴¹Pr below the decay energy Q^- , except for the 7/2+ single level with an energy of 145.443 keV (2001Tu02).

Q^- value has been taken from the atomic mass adjustment by Audi and Wang (2012Au06).

2. Half-Life

The following values of the ¹⁴¹Ce half-life presented in Table 1 were considered here:

Table 1. Results of ¹⁴¹Ce half-life measurements (in days)

Reference	Author(s)	Value	Comments
1949Wa23	Walker	32.11 (23)	Omitted; uncertainty strongly underestimated in an unknown amount
1950Fr58	Freedman and Engelkemeir	32.50 (20)	
1957Ke26	Ketelle and Brozi	32.51 (2)	Omitted; uncertainty strongly underestimated in an unknown amount
1965An07	Anspach <i>et al.</i>	32.550 (7)	Omitted; superseded in 1992Un01
1967Ob01	O'Brien and Eldridge	32.38 (2)	Omitted; uncertainty strongly underestimated in an unknown amount
1971Ba28	S. Baba and H. Baba	32.60 (20)	
1971De11	Debertin	32.51 (6)	Omitted; superseded in 1983Wa26
1972Em01	Emery <i>et al.</i>	32.45 (13)	
1973MeYE	Merritt and Taylor	32.51 (6)	Omitted; superseded in 1980RuZY
1976Va30	Vaninbroukx and Grosse	32.501 (13)	
1980RuZY	Rutledge <i>et al.</i>	32.50 (3)	
1983Wa26	Walz <i>et al.</i>	32.51 (10)	
1992Un01	Unterweger <i>et al.</i>	32.510 (24)	Omitted; superseded in 2002Un02
2002Un02	Unterweger	32.510 (24)	

From the seven values (in boldface) used in the data analysis, the LWEIGHT computer program has consistently identified two outliers (1971Ba28 and 1972Em01), and deduced a weighted mean (32.503) and

an internal uncertainty (0.011) with $\chi^2/\nu = 0.03$. This result suggests that the uncertainties had been overestimated.

The recommended value for the ¹⁴¹Ce half-life is **32.503 (11) days**.

3. β^- Transitions

The energy of the $\beta^-_{0,1}$ - transition has been deduced from the Q^- value and the 145 keV ¹⁴¹Pr level energy. The emission probability of the $\beta^-_{0,1}$ - transition is equal to $P_{\gamma+ce}$ for the 145 keV gamma-ray transition. The probability of feeding the ground state was deduced from the relation $1-P(\beta^-_{0,1})$.

4. Gamma-Ray Transition

The energy was taken from the recommended data by Helmer and van der Leun (2000He14). The emission probability $P_{\gamma+ce}$ was deduced using the relation $P_{\gamma+ce} = P_{\gamma} (1 + \alpha_T)$. (For P_{γ} see Section 7.2). The multipolarity (M1+E2) is based on the measurements of conversion electrons of 1961Co04, 1961Ne12, 1965Ge04, 1966Di02, 1966Pa09, 1968Ge02, 1972Ca07, 1975Le09, 1979Ha09, 1992Sc24. The E2/M1 mixing ratio $\delta = 0.068$ (5) is a weighted average of measurements from 1962Sc11 (0.068 (8)), 1963Ha07 (0.066 (22)) and 1979Ha21 (0.069 (7)). The internal conversion coefficients (ICC) $\alpha_T, \alpha_K, \alpha_L, \alpha_{L'}, \alpha_M, \alpha_N, \alpha_O, \alpha_P$ and their associated uncertainties were interpolated from theoretical values of Band *et al.* (2002Ba85) using the BrIcc computer program (2008Ki07) for the “frozen orbital” approximation, version 2.3S.

The values of the total conversion coefficient α_T , measured and deduced (1966 - 1992); are presented below. A value for the total conversion coefficient of the 145 keV gamma transition was obtained from special coincidence measurements by Hansen *et al.* (1979Ha09) and Schönfeld *et al.* (1992Sc24). Another useful quantity used by them was the measured ratio of the emission probabilities of KX rays and the 145-keV gamma ray.

Total conversion coefficient α_T

1966Di02	0.440 (11)	Dingus <i>et al.</i>	deduced from α_K
1966Pa09	0.441 (9)	Pancholi	deduced from α_K
1975Le09	0.421 (21)	Legrand <i>et al.</i>	measured
1979Ha09	0.439 (13)	Hansen <i>et al.</i>	measured
1979Ha09	0.448 (7)	Hansen <i>et al.</i>	deduced from X_K/γ ratio
1979Ha09	0.436 (17)	Hansen <i>et al.</i>	coinc. meas., extrapol. technique
1992Sc24	0.452 (8)	Schönfeld <i>et al.</i>	coinc. meas., special technique
1992Sc24	0.435 (7)	Schönfeld <i>et al.</i>	deduced from X_K/γ ratio
	0.449 (7)	Present evaluation (BrIcc)	

5. Atomic Data

The fluorescence yields, X-ray energies and relative emission probabilities, and Auger electron energies and relative emission probabilities based on data in 1996Sc06 and 1977La19 are from the SAISINUC computer program.

6. Electron Emissions

The energies of the conversion electrons were obtained from the gamma-ray transition energy and the atomic electron binding energies in 1977La19.

The emission probabilities of the conversion electrons were deduced using the evaluated $P(\gamma)$ and internal conversion coefficient values for the various atomic shells.

The total absolute emission probabilities of K and L Auger electrons were calculated using the EMISSION computer program (1996Sc06, 2000Sc47).

7. Photon Emissions

7.1 X - Ray emissions

The Pr KX- and LX- absolute emission probabilities given in the Tables Section (Table 5.1) were deduced using the computer program EMISSION. Measured values of P_{X_K}/P_γ are compared with a value of 0.350 (6), which was deduced using the computer program EMISSION.

0.338 (5)	Nemet (1961Ne12)
0.347 (12)	Nemet (1961Ne12)
0.342 (9)	Campbell <i>et al.</i> (1971Ca49)
0.334 (9)	Campbell and Mc Nelles (1972Ca07)
0.349 (5)	Hansen <i>et al.</i> (1979Ha09)
0.339 (5)	Schönfeld <i>et al.</i> (1992Sc24)
0.350 (6)	Present evaluation

The recommended value in the present evaluation is in good agreement with the experimental results, especially with the value from 1979Ha09.

7.2 Gamma-Ray Emission

The recommended 145 keV gamma-ray absolute emission probability is the weighted mean of 4 values (2, 3, 4, 6). The following values (based on absolute activity determinations) were considered:

1	0.493 (6)	Eldridge	1966E109
2	0.4844 (41)	Legrand <i>et al.</i>	1975Le09
3	0.482 (3)	Hansen <i>et al.</i>	1979Ha09
4	0.485 (4)	Rutledge <i>et al.</i>	1980RuZY
5	0.489 (4)	Schötzig <i>et al.</i>	1980Sc07
6	0.480 (5)	Schönfeld <i>et al.</i>	1992Sc24
	0.4829 (19)	LWM (2, 3, 4, 6) recommended value. $\chi^2/\nu = 0.28$.	

Value 1 was not used when calculating the average because the uncertainty seems to be underestimated by an unknown amount. Value 5 was also not used because it is considered to be superseded by value 6. The remaining 4 values were used to calculate a weighted mean. (The uncertainty of value 2 is stated to be 3σ but is has been assumed here to be 1σ as this seems to be more realistic and comparable to the other values).

References

- 1949Wa23 D. Walker, Proc. Phys. Soc.(London) 62A, 799 (1949) [Half-life]
- 1950Fr58 M. S. Freedman and D. W. Engelkemeir, Phys. Rev. 79, 897–899 (1950) [Half-life]
- 1957Ke26 B.H. Ketelle, A.R. Brosi, Priv. Comm. (October 1957). Quoted in Nuclear Data Tables 8, No. 1-2, p. 115, 153 [Half-life]
- 1961Co04 J.R. Cook, Proc. Phys. Soc. (London) 77, 346 (1961) [Conversion electrons, γ -ray multipolarity]
- 1961Ne12 L. Nemet, Izvest. Akad. Nauk SSSR, Ser. Fiz. 25, 68 (1961); Columbia Tech. Transl. 25, 68 (1962) [P(KX)/P(γ) ratio]
- 1962Sc11 J.F. Schooley, D.D. Hoppes, A.T. Hirshfeld, J. Res. Nat. Bur. Std. 66A, 317 (1962) [E2/M1 mixing ratio]
- 1963Ha07 J.N. Haag, D.A. Shirley, and David H. Templeton, Phys. Rev. 129, 1601–1613 (1963) [E2/M1 mixing ratio]
- 1965An07 S.C. Anspach, L.M. Cavallo, S.B. Garfinkel, J.M.R. Hutchinson, C.N. Smith, NP 15663 (1965) [Half-life]
- 1965Ge04 J.S. Geiger, R.L. Graham, I. Bergstrom, F. Brown, Nucl. Phys. 68, 352 (1965) [Conversion electrons, γ -ray multipolarity]
- 1966Di02 R.S. Dingus, W.L. Talbert, Jr., M.G. Stewart, Nucl. Phys. 83, 545 (1966) [Experimental ICCs]
- 1966El02 J.S. Eldridge, P. Crowther, W.S. Lyon, Nucleonics 24, No.3, 62 (1966) [γ -ray emission probability]
- 1966Pa09 S.C. Pancholi, Nucl. Phys. 81, 417 (1966) [Experimental ICCs]
- 1967Ob01 H.A. O'Brien, Jr., J.S. Eldridge, Nucleonics 25, No.2, 41 (1967) [Half-life]
- 1968Ge02 W. Gelletly, J.S. Geiger, R.L. Graham, Phys. Rev. 168, 1336 (1968) [Conversion electrons, γ -ray multipolarity]
- 1971Ba28 S. Baba, H. Baba, H. Natsume, J. Inorg. Nucl. Chem. 33, 589 (1971) [Half-life]
- 1971Ca49 J.C. Campbell, P. O'Brien, L.A. McNelles, Nucl. Instrum. Methods 92, 269 (1971) [P(KX)/P(γ) ratio]
- 1971De11 K. Debertin, Z. Naturforsch. 26a, 596 (1971) [Half-life]
- 1972Ca07 J.L. Campbell, L.A. McNelles, Nucl. Instrum. Methods 98, 433 (1972) [P(KX)/P(γ) ratio]
- 1972Em01 J.F. Emery, S.A. Reynolds, E.I. Wyatt, G.I. Gleason, Nucl. Sci. Eng. 48, 319 (1972) [Half-life]
- 1973MeYE J.S. Merritt, J.G.V. Taylor, AECL 4657, p.30 (1973) [Half-life]
- 1975Le09 J. Legrand, J.P. Perolat, C. Bac, J. Gorry, Int. J. Appl. Radiat. Isotop. 26, 179 (1975) [γ -ray emission probabilities, experimental ICCs]
- 1976Va30 R. Vaninbroukx, G. Grosse, Int. J. Appl. Radiat. Isotop. 27, 727 (1977) [Half-life]
- 1977La19 F.P. Larkins, At. Data Nucl. Data Tables 20, 311 (1977) [Atomic electron binding energies]
- 1979Ha09 H.H. Hansen, E. Celen, G. Grosse, D. Mouchel, A. Nylandsted Larsen, R. Vaninbroukx, Z. Phys. A290, 113 (1979) [KX- and γ -ray emission probabilities, experimental ICCs]
- 1979Ha21 H.H. Hansen, Z. Phys. A291, 43 (1979) [Experimental ICCs, E2/M1 mixing ratio]
- 1980RuZY A.R. Rutledge, L.V. Smith, J.S. Merritt, AECL 6692 (1980) [Half-life, γ -ray emission probability]
- 1980Sc07 U. Schotzig, K. Debertin, K.F. Walz, Nucl. Instrum. Methods 169, 43 (1980) [γ -ray emission probability]
- 1983Wa26 K.F. Walz, K. Debertin, H. Schrader, Int. J. Appl. Radiat. Isot., Vol. 34, 1983, P. 1191-1199 [Half-life]

Comments on evaluation

- 1992Sc24 E. Schönfeld, H. Janssen, U. Schotzig, Appl. Radiat. Isot. 43, 1071 (1992) [KX- and γ -ray emission probabilities, experimental ICCs]
- 1992Un01 M.P. Unterweger, D.D. Hoppes, F.J. Schima, Nucl. Instrum. Meth. Phys. Res., Vol. A312, 1992, 349-352 [Half-life]
- 1996Sc06 E. Schönfeld, H. Janssen, Nucl. Instr. Methods A369 (1996) 527 [Atomic data]
- 2000He14 R.G. Helmer and C. van der Leun, Nucl. Instrum. Meth. Phys. Res. A450 (2000) 35 [Gamma-ray energy]
- 2000Sc47 E. Schönfeld, H. Janssen, Appl. Radiat. Isot. 52, 595 (2000) [Calculation of emission probabilities of X-rays and Auger electrons]
- 2001Tu01 J. Tuli, Nucl. Phys. A682, 236c (2001) [¹⁴¹Ce decay scheme, ¹⁴¹Pr levels]
- 2002Ba85 I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr., P. O. Tikkanen, S. Raman, Atomic Data Nucl. Data Tables 81(2002)1 [Theoretical ICCs]
- 2002Un02 M.P. Unterweger, Appl. Radiat. Isot. 56, 125 (2002) [Half-life]
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr., Nucl. Instrum. Methods Phys. Res. A589, 202 (2008) [Band-Raman theoretical ICCs]
- 2012Au06 M. Wang, G. Audi, *et al.* Chinese Physics C36 (2012) 1603 [Q β -]

**¹⁴⁴Ce – Comments on evaluation of decay data
by A. L. Nichols**

Evaluated: May 2013 and March/April 2014

Evaluation Procedures

Limitation of Relative Statistical Weight Method (LWM) and other analytical techniques were applied to average numbers throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

¹⁴⁴Ce ($T_{1/2} = 284.89$ days) undergoes 100% β^- decay ($Q(\beta^-) = 318.6$ (8) keV) via various excited levels of ¹⁴⁴Pr to the metastable state of ^{144m}Pr (branching fraction of 0.0115 (20), and ^{144m}Pr half-life of 7.2 min) and to the ground state of ¹⁴⁴Pr (branching fraction of 0.9885 (20), and ¹⁴⁴Pr half-life of 17.29 min). A reasonably well-defined decay scheme was derived from the gamma-ray measurements of 1969Gu15, 1970An15, 1970Fa03, 1970Po09, 1975De17, 1976Ch33, 1977Ge12, 1981OI04, 1982Yu02, 1983El14 and 1984Da13, coupled with comprehensive studies of conversion-electron spectra and extensive efforts to determine unambiguously the spin and parity of the ground state and the half-lives of the excited states of ¹⁴⁴Pr (1957Pa51, 1958Hi76, 1959Fr54, 1959Se57, 1960Ge05, 1961Ge09, 1962B105, 1962Bu09, 1962Bu22, 1962Fo04, 1963Co18, 1963Fu16, 1963Iw02, 1963Si10, 1963Cr11, 1963Kn05, 1963Bh11, 1964Az02, 1964Be36, 1964Mc22, 1965Re13, 1965Co19, 1966Be11, 1966Da04, 1967Gu17, 1968Da12, 1969Ge01, 1969Ma24, 1970An15, 1971Sa20 and 1975Ba32). While ¹⁴⁴Ce(β^-) ^{144m}Pr amounts to only 1.15 (20) % of the total decay, daughter ^{144m}Pr undergoes 99.94 (2) % isomeric-transition decay to ¹⁴⁴Pr by emission of a single 59.03(3)-keV gamma ray, and 0.06 (2) % β^- decay to ¹⁴⁴Nd. As a consequence of the short-lived decay of ^{144m}Pr the 59.03-keV M3 gamma transition has been included in both the ¹⁴⁴Ce and ^{144m}Pr decay-data files. Thus, the proposed decay scheme for ¹⁴⁴Ce consists of three β^- transitions and seven gamma-ray emissions, of which one gamma transition originates from ^{144m}Pr IT decay. Weighted-mean relative emission probabilities were calculated for some of the gamma rays, while equivalent data for other gamma transitions were primarily adopted from the comprehensive measurements of 1984Da13. All of these relative emission probabilities were defined in terms of the 133.515-keV gamma ray (100%).

Nuclear Data

¹⁴⁴Ce has been adopted as a burn-up monitor for irradiated fuel, and as a brachytherapy source in specific forms of cancer treatment. The β^- decay of daughter ¹⁴⁴Pr is important in decay heat assessments, design of irradiated fuel storage facilities, and other facets of radwaste management.

Half-life

The measurements of 1956Sc87, 1957Me47, 1965Fl02, 1968La10, 1968Re04, 1980Ho17, 1983Wa26, 1986OI01, 1997Ma75 and 2014Un01 were analysed by means of the limitation of relative statistical weight method (LWM), normalised residual method (NRM), Rajeval technique, bootstrap method, and Mandel-Paule approach. $\chi^2/(N-1)$ obtained from the LWM, bootstrap and Mandel-Paule analyses are significantly higher than the values obtained from the equivalent data manipulation by means of the NRM and Rajeval techniques. While NRM resulted in adjustments to the half-life data measured by 1980Ho17 and 1997Ma75, the equivalent Rajeval analysis consisted of a small adjustment to the uncertainty of 1965Fl02, and more substantial changes to the uncertainties quoted by 1980Ho17, 1983Wa26 and 1997Ma75. Under these circumstances, a half-life value of (284.89 ± 0.06) days was preferred and recommended, as quantified by the Rajeval analytical procedure.

Reference	Half-life (d)
1956Sc87	284.5 ± 1.0
1957Me47	285 ± 2
1960Ea02	277 ± 4*
1965Fl02	283.8 ± 0.6
1968La10	284.8 ± 1.0
1968Re04	284.9 ± 0.8
1980Ho17	285.8 ± 0.1
1983Wa26	284.45 ± 0.14
1986Ol01	284.893 ± 0.008 [§]
1992Un01	284.558 ± 0.038 [‡]
1997Ma75	286.14 ± 0.09
2002Un02	284.534 ± 0.032 [‡]
2012Fi12	284.35 ± 0.08 [Ⓞ]
2014Un01	284.35 ± 0.40
Recommended value	284.89 ± 0.06[†]

* Defined as an outlier, and therefore not adopted in the analytical procedures.

[§] Uncertainty adjusted to ± 0.06 to reduce weighting to no more than 0.50.

[‡] Superseded by study of 2012Fi12 after the discovery of problems in maintaining constant detector-source geometry within the NIST ionization chamber.

[Ⓞ] Replaced by 2014Un01 – uncertainty increased from ± 0.08 to 0.40 in the most recent NIST paper.

[†] Recommended value adopted from Rajeval analysis, with uncertainty increased from ± 0.008 to 0.06 in alignment with the smallest uncertainty adopted to calculate the average value.

Analytical method	Half-life (d)	$\chi^2/(N-1)$	$\chi^2/(N-1)_{\text{critical}}$
UWM	284.86 ± 0.22	–	–
LWM – WM	284.91 ± 0.05	31.85	2.41
LWM – WM [#]	285.3 ± 0.4	23.22	2.41
NRM	284.64 ± 0.16	2.01	1.88
Rajeval	284.892 ± 0.008	1.56	–
Bootstrap	284.79 ± 0.21	57.74	–
Mandel-Paule	285.0 ± 0.6	46.43	–

[#] Uncertainty of measurement by 1986Ol01 increased from ± 0.008 to 0.06 to reduce weighting to no more than 0.50.

Q values

The 59.03-keV IT gamma decay of ^{144m}Pr has been handled as part of the decay scheme of ¹⁴⁴Ce, with 100% β⁻ decay to ¹⁴⁴Pr and adoption of a Q⁻-value of 318.6 (8) keV (2012Wa38).

Gamma-ray energies and emission probabilities

Energies

The well-defined nuclear level energies of 2001So16 were used to calculate the gamma transition energies and their uncertainties. Greater confidence was placed on this approach because of the more wide-ranging origins of the gamma transition data even though the energies of the gamma-ray emissions have been measured by 1960Ge05, 1960Sa22, 1970An15, 1976Ch33, 1976Ra22, 1982Yu02, 1983El14 and 1984Da13. Daughter ^{144m}Pr undergoes 99.94 (2) % isomeric-transition decay to ¹⁴⁴Pr by means of a single 59.03(3)-keV gamma ray, and 0.06 (2) % β⁻ decay to ¹⁴⁴Nd – the 59.03-keV M3 gamma transition has been included in both the ¹⁴⁴Ce and ^{144m}Pr decay-data files.

Adopted energies, spins and parities for the nuclear levels of ¹⁴⁴Pr (2001So16).

Nuclear level number	Nuclear level energy (keV)	Spin and parity
0	0.0	0 – 17.29 (4) min ¹⁴⁴ Pr
1	59.03 ± 0.03	3 – 7.2 (2) min ^{144m} Pr
2	80.120 ± 0.004	1 – 136 (11) ps*
3	99.952 ± 0.009	2 – 0.67 (6) ns [#]
4	133.5152 ± 0.0020	1 – 7 (4) ps [†]

* Weighted mean of 125 (14) ps (1962Bu22 (same value also reported by 1962Bu09)) and 143 (11) ps (1964Mc22); uncertainty increased from ± 9 to 11, in alignment with the smallest measured uncertainty of the values used to calculate the weighted-mean value.

[#] Weighted mean of 0.69 (14) ns (1962Bu22 (same value also reported by 1962Bu09)) and 0.66 (6) ns (1964Be36).

[†] Determined by 1962Bu22 (same value also reported by 1962Bu09).

Measured and recommended gamma-ray energies.

E _γ (keV)													
1957Pa51*	1959Fr54	1959Se57	1960Ge05†	1960Sa22	1970An15	1976Ch33	1976Ra22	1978Mo22	1979Bo26	1982Yu02	1983El14	1984Da13	Recommended#
–	–	–	–	–	–	–	–	–	–	–	–	5.012 (50)	Pr X L _{α1} + L _{α2}
–	–	–	–	–	–	–	–	–	–	–	–	5.486 (50)	Pr X L _{β1} + L _{β3} + L _{β4}
–	–	–	–	–	–	–	–	–	–	–	–	5.851 (50)	Pr X L _{β2,15}
–	–	–	–	–	–	–	–	–	–	–	–	6.297 (50)	Pr X L _{γ1}
–	–	–	–	–	–	–	–	–	–	–	–	6.594 (50)	Pr X L _{γ2,3}
33.7	35 (X-ray)	36 (2)	33.57 (3)	34 (3)	33.6 (1)	33.622 (10)	33.6	–	–	33.552 (20)	33.56	33.568 (10)	33.563 ± 0.009
–	–	–	–	–	35.9 (1)	35.554 (10)	–	–	–	35.558 (20)	–	35.547 (10)	Pr X K _{α2}
–	–	–	–	–	–	36.030 (10)	–	–	–	36.033 (20)	–	36.026 (10)	Pr X K _{α1}
–	–	–	–	–	40.7	40.676 (10)	–	–	–	–	40.72	40.739 (20)	Pr X K _{β1}
41.0	–	–	40.93 (3)	–	41	40.892 (50)	40.8 (4)	–	–	–	–	40.98 (10)	40.92 ± 0.03
–	–	–	–	–	41.7 (1)	41.759 (10)	–	–	–	41.756 (15)	–	41.778 (10)	Pr X K _{β2}
53.0	–	–	53.41 (5)	54 (4)*	53.4 (1)	53.432 (10)	53.4 (1)	–	–	53.436 (10)	53.40	53.395 (5)	53.395 ± 0.005
–	–	–	59.03 (3)	–	59.0 (3)	–	58.5 (2)	–	–	–	58.96	58.98‡	59.03 ± 0.03 ^{144m} Pr IT decay
81.0	80	81 (4)	80.12 (3)	80 (3)	80.1 (1)	80.106 (5)	80.09 (8)	–	–	80.119 (10)	79.95	80.120 (5)	80.120 ± 0.004
100.6	–	–	99.95 (5)	101 (5)*	100.0 (1)	99.963 (20)	99.85 (8)	–	–	99.922 (15)	99.80	99.961 (15)	99.952 ± 0.009
134.1	133	134 (4)	133.53 (3)	134 (2)	133.5 (1)	133.544 (5)	133.51 (10)	133.515 (5)	133.528 (13)	133.514 (9)	133.29	133.515 (4)	133.5152 ± 0.0020

Determined from the nuclear level energies of 2001So16.

* Gamma-ray energies derived from coincidence measurements.

† Gamma-ray energies determined from conversion-electron measurements.

‡ Inadequate data arising from insufficient statistics for $P_{\gamma}^{rel} \leq 0.01$ (normalized with respect to a relative emission probability of 100% for the 133.515-keV gamma ray).

Emission Probabilities

Relative gamma-ray emission probabilities have been partially or fully determined or derived from the measurements of 1970An15, 1970Po09, 1976Ch33, 1977Ge12, 1982Yu02, 1983El14 and 1984Da13, and limited and complete studies of the absolute gamma-ray emission probabilities have been performed by 1960Ge05, 1969Gu15, 1970Fa03, 1975De17, 1976Ra22 and 1981Ol04. While considerable emphasis was placed on the comprehensive and well-defined measurements of 1984Da13, data from the studies of 1960Ge05, 1970An15, 1970Fa03, 1970Po09, 1976Ch33, 1977Ge12, 1982Yu02 and 1983El14 were also used in the weighted-mean analyses of the 33.563-, 40.92-, 53.395-, 80.120- and 99.952-keV gamma-ray emissions. All of the relative emission probabilities were suitably refined and quantified in terms of the emission probability of the 133.515-keV gamma ray (100.0%). Considerable difficulty has been experienced in resolving the 40.92-keV gamma-ray emission from Pr K_β X-rays (40.65 – 41.97 keV); hence, a speculative $P_{\gamma}^{rel}(40.92 \text{ keV})$ weighted-mean value of 2.94 (50) was adopted to achieve a gamma population-depopulation balance in conjunction with an assumed zero β^{-} population of the 99.952-keV nuclear level of ¹⁴⁴Pr. A weighted-mean normalization factor (F) of 0.1083 ± 0.0012 was adopted for the relative gamma-ray emission probabilities, as determined from the conversion-electron and gamma spectroscopic studies of Geiger *et al.* (1960Ge05), Debertain *et al.* (1975De17) and Olomo and MacMahon (1981Ol04) by means of a well-characterised β spectrometer and Ge(Li) detector systems.

Absolute emission probability of 133.515-keV gamma ray (normalization factor).

Reference	$P_{\gamma}^{abs}(133.515 \text{ keV})$
1960Ge05	0.108 (4) [#]
1975De17	0.1109 (16)
1981Ol04	0.1069 (12)
Recommended	0.1083 (12)[†]

[#] Calculated from theoretical internal conversion coefficients, measured conversion-electron and γ intensities, and $(K(133.515 \text{ keV}) / ^{144}\text{Pr } \beta \text{ spectrum})$ ratio of 0.053 (2), as specified in Table 2 and Section 6 of 1960Ge05.

[†] Recommended uncertainty adjusted from ± 0.0009 to 0.0012 to align with the smallest uncertainty of the values used to calculate the weighted mean value.

Multipolarities, and Internal Conversion Coefficients

The nuclear level scheme specified by Sonzogni has been used to define the multipolarities of the gamma transitions on the basis of known spins and parities (2001So16). Extensive conversion-electron, e- γ , γ - γ coincidence and directional angular correlation studies throughout the 1960s/1970s focused on defining gamma multipolarities and spin-parity assignments for the nuclear levels of ¹⁴⁴Pr (particularly for the 0 – ground state). The detailed conversion-electron measurements of Geiger *et al.* are particularly noteworthy (1960Ge05, 1961Ge09, 1969Ge01), along with equivalent studies of 1959Se57, 1963Iw02, 1965Co19, 1966Be11, 1966Da04, 1969Ma24, 1970An15, 1971Sa20, 1975Ba32 and 1976Ch33. Adopted multipolarities are 100%M1 for the 33.563-, 40.92-, 53.395-, 80.120-, and 133.515-keV gamma rays, and E2 for the 99.952-keV gamma ray. More specifically, evidence of 100%M1 for the 40.92-keV gamma transition arises from measurements of the relative conversion-electron emission probabilities of 1960Ge05, and the α_L value of 1.8 (6) determined by 1970An15. Recommended internal conversion coefficients have been determined from the frozen orbital approximation of Kibédi *et al.* (2008Ki07), based on the theoretical model of Band *et al.* (2002Ba85, 2002Ra45).

The observation of a low-intensity, 59.03-keV gamma ray was identified with the subsequent IT decay of ^{144m}Pr (half-life of 7.2 (2) min. – defined as M3 transition).

Branching fractions for ¹⁴⁴Ce(β^{-})¹⁴⁴Pr and ¹⁴⁴Ce(β^{-})^{144m}Pr

¹⁴⁴Ce undergoes β^{-} decay predominantly to the ground state of ¹⁴⁴Pr via both β^{-} population of the ground state, 80.120- and 133.5152-keV nuclear levels of ¹⁴⁴Pr, and γ depopulation/population of the ground state, 80.120-, 99.952- and 133.5152-keV nuclear levels of ¹⁴⁴Pr. However, there is also evidence for the existence of a small branch that populates the 59.03-keV nuclear level of ^{144m}Pr by means of the 40.92-keV γ transition, which arises as part of the 33.563–40.92(–59.03) γ cascade embracing the 133.5152-, 99.952- and 59.03-keV nuclear levels.

Adopting a recommended relative γ -ray emission probability of 2.94 (50) % for the 40.92-keV γ ray, and a normalisation factor of 0.1083 (12):

$$\text{BF}(^{144}\text{Ce}(\beta^{-})^{144m}\text{Pr}) = [P_{\gamma}^{rel} \times F] \times (1 + \alpha_{total})$$

where P_{γ}^{rel} is the relative γ -ray emission probability, F is the normalisation factor, and α_{total} is the total internal conversion coefficient.

Therefore, $\text{BF}(^{144}\text{Ce}(\beta^{-})^{144m}\text{Pr}) = [2.94 (50) \times 0.1083 (12)] \times 3.61 (4) = 1.149 (195) \%$.

$\text{BF}(^{144}\text{Ce}(\beta^{-})^{144m}\text{Pr}) = 0.0115 (20)$ when rounded-up and expressed as a fraction [i.e. 1.15 (20) %];

$\text{BF}(^{144}\text{Ce}(\beta^{-})^{144}\text{Pr}) = 0.9885 (20)$ when expressed as a fraction [i.e. 98.85 (20) %].

Published gamma-ray emission probabilities.

E_γ (keV)	P_γ (%)										
	1954Kr40*	1959Fr54*	1959Se57*	1960Ge05[▲]	1960Sa22*	1969Ma24*	1969Gu15⁰	1970An15*	1970Fa03[†]	1970Po09*	1975De17⁰
33.563 (9)	–	includes X-rays	144 (10)	0.23 (1)	95	80 (8)	–	2.0 (5)	0.15 (5)	2.3 (2)	–
40.92 (3)	–	–	–	0.36 (1)	–	–	–	4.4 (15)	0.50 (25)	5 (4)	–
53.395 (5)	–	–	–	0.127 (5)	–	–	–	1.2 (2)	0.09 (4)	1.1 (1)	–
59.03 (3) ^{144m} Pr IT decay	–	–	–	0.0011	–	–	–	~ 0.01	not observed	–	–
80.120 (4)	28.7 (43)	24 (5)	24 (5)	1.6 (1)	35	22 (2)	1.51 (15)	14.8 (10)	1.54 (15)	16 (1)	–
99.952 (9)	–	–	–	0.042 (2)	–	10 (2)	–	0.36 (5)	0.038 (4)	0.39 (4)	–
133.5152 (20)	100	100 (10)	100 (6)	10.8 (4)	100	100	11.0 (3)	100 (5)	(10.8)	100 (6)	11.09 (16)

E_γ (keV)	P_γ (%)						
	1976Ch33*	1976Ra22[‡]	1977Ge12⁺	1981O104⁰	1982Yu02*	1983El14*	1984Da13*
33.563 (9)	2.62 (15)	includes X-rays	–	–	2.14 (17)	1.77 (26)	1.8 (2)
40.92 (3)	3.59 (40)	0.72 (4)	–	–	3.76 (65) ^Δ	3.22 (230) [#]	1.38 (20)
53.395 (5)	0.86 (5)	0.07 (1)	7.8 (4)	–	1.05 (11)	0.84 (8)	0.90 (7)
59.03 (3) ^{144m} Pr IT decay	not observed	0.003 (2)	–	–	not observed	≈ 0.01	≤ 0.01
80.120 (4)	10.20 (100)	1.50 (4)	112 (6)	–	13.6 (10)	12.58 (100)	12.25 (49)
99.952 (9)	0.35 (3)	0.06 (1)	–	–	0.36 (4)	0.50 (7)	0.36 (4)
133.5152 (20)	100 (4)	10.0 (1)	836 (25)	10.69 (12)	100	100	100.0 (10)

* Emission probabilities expressed relative to P_γ(133.515 keV) of 100%.

[▲] Absolute emission probabilities calculated from theoretical internal conversion coefficients, measured conversion-electron and γ intensities, and (K(133.515 keV) / ¹⁴⁴Pr β spectrum) ratio of 0.053 (2), as specified in Table 2 and Section 6 of 1960Ge05.

⁰ Measured absolute emission probability (and probabilities).

[†] Absolute emission probabilities expressed on the basis of an absolute emission probability of 10.8% for the 133.515-keV gamma ray.

[‡] Emission probabilities expressed relative to an absolute emission probability of 10.0 (1) % for the 133.515-keV gamma ray.

⁺ Emission probabilities expressed relative to P_γ(696.5 keV) of 100% which is identified with the decay of daughter ¹⁴⁴Pr/^{144m}Pr.

^Δ Authors state that relative emission probability obtained by subtracting theoretical relative emission probability of Pr X K_{α1} from measured composite peak.

[#] Authors state that relative emission probability obtained by subtracting theoretical relative emission probabilities of Pr X K_{β1} and K_{β2} from measured composite peak.

Measured and recommended gamma-ray emission probabilities relative to P_γ(133.515 keV) of 100%.

E _γ (keV)	P _γ ^{rel} (%)									
	1954Kr40	1959Fr54	1959Se57	1960Ge05*	1960Sa22	1969Ma24	1969Gu15*	1970An15	1970Fa03*	1970Po09
33.563 (9)	–	–	144 (10) [#]	2.1 (1)	95 [#]	80 (8) [#]	–	2.0 (5)	1.4 (5)	2.3 (2)
40.92 (3)	–	–	–	3.3 (1)	–	–	–	4.4 (15)	4.6 (23)	5 (4)
53.395 (5)	–	–	–	1.18 (5)	–	–	–	1.2 (2)	0.8 (4)	1.1 (1)
59.03 (3) ^{144m} Pr IT decay	–	–	–	0.01	–	–	–	~ 0.01	not observed	–
80.120 (4)	28.7 (43)	24 (5)	24 (5)	14 (1)	35	22 (2)	13.7 (14)	14.8 (10)	14.3 (14)	16 (1)
99.952 (9)	–	–	–	0.39 (2)	–	10 (2)	–	0.36 (5)	0.35 (4)	0.39 (4)
133.5152 (20)	100	100 (10)	100 (6)	100 (4)	100	100	100	100 (5)	(100)	100 (6)

E _γ (keV)	P _γ ^{rel} (%)							Recommended [†]
	1976Ch33	1976Ra22*	1977Ge12*	1982Yu02	1983El14	1984Da13		
33.563 (9)	2.62 (15)	–	–	2.14 (17)	1.77 (26)	1.8 (2)	2.08 (10) ⁺	
40.92 (3)	3.59 (40)	7.2 (4)	–	3.76 (65) [‡]	3.22 (230) ⁰	1.38 (20)	2.94 (50) [▲]	
53.395 (5)	0.86 (5)	0.7 (1)	0.93 (6)	1.05 (11)	0.84 (8)	0.90 (7)	0.93 (5) ^Δ	
59.03 (3) ^{144m} Pr IT decay	–	0.03 (2)	–	not observed	≈ 0.01	≤ 0.01	0.0087 (18) [Ⓟ]	
80.120 (4)	10.20 (100)	15.0 (4)	13.4 (8)	13.6 (10)	12.58 (100)	12.25 (49)	12.9 (5) [×]	
99.952 (9)	0.35 (3)	0.6 (1)	–	0.36 (4)	0.50 (7)	0.36 (4)	0.38 (2) [§]	
133.5152 (20)	100 (4)	(100 (1))	100	100	100	100.0 (10)	100	

* Emission probabilities adjusted in order to be expressed relative to P_γ(133.515 keV) of 100%.

Unable to resolve 33.563-keV gamma transition from X-ray emissions – data not included in subsequent evaluation.

† LWM weighted-mean values adopted when judged appropriate.

* Stated as obtained by subtracting the theoretical Pr K_{α1} X-ray emission probability (36.033 keV) from the experimentally-measured composite peak.

⁰ Stated as obtained by subtracting the theoretical Pr K_{β1} and K_{β2} X-ray emission probabilities (40.739 and 41.778 keV) from the experimentally-measured composite peak.

+ Rajeval value adopted, with the recommended uncertainty adjusted from ± 0.07 to 0.10 to align with smallest uncertainty of the data used to calculate this weighted-mean value.

▲ Considerable difficulty has been experienced in resolving the 40.92-keV gamma-ray emission from Pr K_β X-ray emissions (40.65 – 41.97 keV) – hence, a speculative P_γ^{rel}(40.92 keV) weighted-mean value of 2.94 (50) was adopted to achieve a gamma population-depopulation balance in conjunction with an assumed zero β[–] population of the 99.952-keV nuclear level of ¹⁴⁴Pr.

Δ Rajeval value adopted, with the recommended uncertainty adjusted from ± 0.03 to 0.05 to align with smallest uncertainty of the data used to calculate this weighted-mean value.

Ⓟ Calculated from the recommended relative transition probability of 10.61 (215) % for the 40.92-keV γ ray which populates the 59.03-keV nuclear level of ¹⁴⁴Pr, and M3 total internal conversion coefficient of 1221 (18).

× Recommended uncertainty adjusted from ± 0.3 to 0.5 to align with smallest uncertainty of the data used to calculate the weighted-mean value.

§ Recommended uncertainty adjusted from ± 0.01 to 0.02 to align with smallest uncertainty of the data used to calculate the weighted-mean value.

Gamma-ray emissions: multiplicities, and theoretical internal-conversion coefficients (frozen orbital approximation).

E_γ (keV)		Multipolarity	α_K	α_L	α_{L1}	α_{L2}	α_{L3}	α_M	α_N	α_{O+}	α_{total}
33.563 (9)	(Pr)	M1	–	3.70 (6)	3.35 (5)	0.289 (4)	0.0603 (9)	0.780 (11)	0.1744 (25)	0.031	4.69 (7)
40.92 (3)	(Pr)	M1	–	2.06 (3)	1.87 (3)	0.1585 (23)	0.0329 (5)	0.434 (7)	0.0971 (14)	0.0168	2.61 (4)
53.395 (5)	(Pr)	M1	6.75 (10)	0.942 (14)	0.856 (12)	0.0710 (10)	0.01464 (21)	0.199 (3)	0.0444 (7)	0.0076	7.94 (12)
59.03 (3)	(Pr)	M3	408 (6)	618 (9)	233 (4)	34.9 (5)	351 (5)	155.0 (23)	34.7 (5)	5.20 (8)	1221 (18)
80.120 (4)	(Pr)	M1	2.08 (3)	0.288 (4)	0.263 (4)	0.0209 (3)	0.00427 (6)	0.0608 (9)	0.01359 (19)	0.0030	2.45 (4)
99.952 (9)	(Pr)	E2	1.214 (17)	0.710 (10)	0.1085 (16)	0.288 (4)	0.313 (5)	0.1599 (23)	0.0346 (5)	0.0049	2.12 (3)
133.5152 (20)	(Pr)	M1	0.486 (7)	0.0668 (10)	0.0613 (9)	0.00456 (7)	0.000919 (13)	0.01408 (20)	0.00315 (5)	0.00055	0.571 (8)

Recommended gamma-ray energies, relative and absolute emission probabilities, and transition probabilities, and conversion-electron probabilities.

	E_γ (keV)	P_γ^{rel}	P_γ^{abs} (%)	Transition probability (%)	$P_{ce}^{abs} = (TP - P_\gamma^{abs})$ (%)
$\gamma_{4,3}$	33.563 (9)	2.08 (10)	0.225 (11)	1.28 (6)	1.05 (6)
$\gamma_{3,1}$	40.92 (3)	2.94 (50)	0.32 (5)	1.16 (18)	0.84 (19)
$\gamma_{4,2}$	53.395 (5)	0.93 (5)	0.101 (5)	0.90 (4)	0.80 (4)
$\gamma_{1,0}$	59.03 (3)	0.0087 (18)	0.00094 (19)	1.15 (23)	1.149 (230)
$\gamma_{2,0}$	80.120 (4)	12.9 (5)	1.40 (5)	4.83 (17)	3.43 (18)
$\gamma_{3,0}$	99.952 (9)	0.38 (2)	0.041 (2)	0.128 (6)	0.087 (6)
$\gamma_{4,0}$	133.5152 (20)	100	10.83 (12)	17.01 (19)	6.18 (22)

β⁻ energies and emission probabilities

Energies

All β⁻ energies were derived from the structural details of the proposed decay scheme. The nuclear level energies of 2001So16 and evaluated Q-value of 318.6 (8) keV (2012Wa38) were used to determine the recommended energies and uncertainties of the β⁻ emissions.

Emission Probabilities

Emission probabilities were derived for the population-depopulation imbalances of the relative emission probabilities of the gamma rays, their theoretical internal-conversion coefficients, and the normalization factor of (0.1083 ± 0.0012) for the gamma-ray emissions, as noted above. β⁻ decay directly to the 99.952-keV nuclear level of ¹⁴⁴Pr and the 59.03-keV metastable state (^{144m}Pr) were both assumed to be zero on the basis of spin and parity considerations that would have resulted in first forbidden unique and third forbidden non-unique transitions, respectively. Logft values and average E_{β⁻} energies were determined by means of the LOGFT code.

Recommended energies and emission probabilities for the β⁻ decay of ¹⁴⁴Ce.

	E _β (keV)*	Av. E _{β⁻} (keV)	P _β	¹⁴⁴ Ce	¹⁴⁴ Pr	transition type	log ft
β ⁻ _{0,4}	185.1 (8)	50.29 (24)	19.2 (1)	0 +	1 -	first forbidden non-unique	7.27
β ⁻ _{0,2}	238.5 (8)	66.24 (25)	3.9 (2)	0 +	1 -	first forbidden non-unique	8.31
β ⁻ _{0,0}	318.6 (8)	91.3 (3)	76.9 (3)	0 +	0 -	first forbidden non-unique	7.424
			Σ 100				

* Determined from the nuclear level energies of 2001So16 and Q-value of 318.6(8) keV (2012Wa38).

Energies and emission probabilities of noteworthy internal-conversion electrons.

		Energy (keV)	Electrons per 100 disint.
ec _{4,3} T	(Pr)	26.728 – 33.563	1.05 (6)
ec _{4,3} L	(Pr)	26.728 – 27.599	0.83 (5)
ec _{4,3} M	(Pr)	32.052 – 32.632	0.175 (10)
ec _{4,3} N	(Pr)	32.259 – 33.561	0.039 (2)
ec _{4,3} O+	(Pr)	33.526 – 33.563	0.0069 (4)
ec _{3,1} T	(Pr)	34.09 – 40.90	0.84 (19)
ec _{3,1} L	(Pr)	34.09 – 34.96	0.66 (15)
ec _{3,1} M+	(Pr)	39.41 – 40.90	0.18 (4)
ec _{4,2} T	(Pr)	11.404 – 53.393	0.80 (4)
ec _{4,2} K	(Pr)	11.404 (5)	0.68 (3)
ec _{4,2} L	(Pr)	46.560 – 47.431	0.095 (5)
ec _{4,2} M+	(Pr)	51.884 – 53.393	0.025 (1)
ec _{1,0} T	(Pr)	17.04 – 59.03	1.15 (23)
ec _{1,0} K	(Pr)	17.04 (3)	0.38 (8)
ec _{1,0} L	(Pr)	52.20 – 53.07	0.58 (12)
ec _{1,0} M+	(Pr)	57.52 – 59.03	0.18 (4)
ec _{2,0} T	(Pr)	38.129 – 80.12	3.43 (18)
ec _{2,0} K	(Pr)	38.129 (4)	2.91 (15)
ec _{2,0} L	(Pr)	73.285 – 74.156	0.403 (21)
ec _{2,0} M	(Pr)	78.609 – 79.189	0.085 (4)
ec _{2,0} N	(Pr)	79.816 – 80.118	0.0190 (10)
ec _{2,0} O+	(Pr)	80.083 – 80.12	0.0042 (2)
ec _{4,0} T	(Pr)	91.524 – 133.515	6.18 (22)
ec _{4,0} K	(Pr)	91.524 (2)	5.26 (19)
ec _{4,0} L	(Pr)	126.680 – 127.551	0.723 (25)
ec _{4,0} M	(Pr)	132.004 – 132.584	0.152 (5)
ec _{4,0} N	(Pr)	133.211 – 133.513	0.0341 (12)
ec _{4,0} O+	(Pr)	133.478 – 133.515	0.00595 (21)

A consistent decay scheme was derived that contains three β^- transitions and seven gamma-ray emissions, of which one gamma transition originates from the IT decay of ^{144m}Pr. While the 59.03-keV gamma transition constitutes the major IT-decay mode of ^{144m}Pr, this emission has been adopted as part of the β^- decay scheme of ¹⁴⁴Ce, and can be found in the decay-data files of both ¹⁴⁴Ce and ^{144m}Pr.

Atomic Data

The X-ray and Auger-electron data have been calculated using the evaluated gamma-ray data, and atomic data from 1996Sc06, 1998ScZM and 1999ScZX. Both the X-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.02, 28 February 2012), as described in 2000Sc47. This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

K and L X-ray energies and emission probabilities of ¹⁴⁴Ce.

			Energy (keV)	Photons per 100 disint.	Relative probability
XL	(Pr)		4.453 – 6.617	1.54 (4)	35.0
		XL ₁	4.453	0.0282 (9)	
		XL _{α}	5.013 – 5.033	0.711 (21)	
		XL _{η}	4.929	0.0107 (4)	
		XL _{β}	5.489 – 5.851	0.673 (15)	
		XL _{γ}	6.327 – 6.617	0.116 (3)	
XK _{α}		XK _{α2}	35.5506 (2)	2.41 (5)	54.8
		XK _{α1}	36.0267 (2)	4.40 (9)	
XK' _{β1}		XK _{β3}	40.6533 (7))	30.5
		XK _{β1}	40.7487 (5)) 1.34 (3)	
		XK _{β5}	41.050)	
XK' _{β2}		XK _{β2}	41.774 (2))	7.80
		XK _{β4}	41.877) 0.343 (10)	
		XKO _{2,3}	41.968)	

Auger-electron energies and emission probabilities of ¹⁴⁴Ce.

		Energy (keV)	Electrons per 100 disint.	Relative probability
e _{AK}	(Pr)		0.80 (4)	
	KLL	28.162 – 29.890	0.514 (26)	100
	KLX	33.576 – 36.004	0.254 (13)	49.4
	KXY	38.97 – 41.95	0.0313 (17)	6.1
e _{AL}	(Pr)	2.90 – 4.91	9.88 (10)	1922

Pr: $\omega_K = 0.914$ (4); $\omega_L = 0.132$ (5); $n_{KL} = 0.871$ (4) were taken from 1996Sc06.

Electron energies were determined from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

Data Consistency

An effective Q_{β^-} -value of 318.6 (8) keV has been adopted from Wang *et al.* (2012Wa38), and compared with the Q -value calculated by summing the contributions of the individual emissions to the ¹⁴⁴Ce β^- -decay process (i.e. β^- , conversion electrons, γ , etc.):

$$\text{calculated } Q\text{-value} = \sum (E_i \times P_i) = 318.5 \text{ (13) keV}$$

The percentage deviation of the calculated Q -value from the effective Q -value is $(0.0 \pm 0.5) \%$, which supports the derivation of an extremely consistent decay scheme with a relatively modest variant.

References

- 1954Co60 J.M. Cork, M.K. Brick, L.C. Schmid, Energies of the Radiations from Ce¹⁴⁴ and Pr¹⁴⁴, Phys. Rev. 96 (1954) 1295-1297. [E_γ, E_e]
- 1954Em09 W.S. Emmerich, W.J. Auth, J.D. Kurbatov, The Nuclear Spectra of Ce¹⁴⁴ and Pr¹⁴⁴, Phys. Rev. 94 (1954) 110-112. [E_β, E_γ]
- 1954Kr40 W.E. Kreger, C.S. Cook, Gamma Radiation from Ce¹⁴⁴ and Pr¹⁴⁴, Phys. Rev. 96 (1954) 1276-1278. [P_γ]
- 1956Pu24 I. Pullman, P. Axel, Decay of Ce¹⁴⁴, Phys. Rev. 102 (1956) 1366-1372. [E_β, E_γ, E_e]
- 1956Sc87 R.P. Schuman, M.E. Jones, A.C. Mewherter, Half-lives of Ce¹⁴⁴, Co⁵⁸, Cr⁵¹, Fe⁵⁵, Mn⁵⁴, Pm¹⁴⁷, Ru¹⁰⁶ and Sc⁴⁶, J. Inorg. Nucl. Chem. 3 (1956) 160-163. [Half-life]
- 1957Me47 W.F. Merritt, P.J. Champion, R.C. Hawkings, Half-life Determination of Some Radionuclides, Can. J. Phys. 35 (1957) 16-20. [Half-life]
- 1957Pa51 V. Parfenova, N.V. Forafontov, V.S. Shpinel, Investigation of the Ce¹⁴⁴ Decay Using the β-γ Coincidence Method, Izvest. Akad. Nauk SSSR, Ser. Fiz. 21 (1957) 1601-1606. [E_β, P_β, E_γ]
- 1958Hi76 R.L. Hickok, W.A. McKinley, S.C. Fultz, Decay of Ce¹⁴⁴ and Pr¹⁴⁴, Phys. Rev. 109 (1958) 113-116. [E_β, P_β, α_K]
- 1959Fr54 N.J. Freeman, The Decay of ¹⁴⁴Ce, Proc. Phys. Soc. 74 (1959) 449-456. [E_β, P_β, transition type, E_γ, α_K, α_L, E_{ce}]
- 1959Se57 A.K. Sengupta, R. Bhattacharyya, J. Lahiri, P.N. Mukherjee, Level Scheme of Pr¹⁴⁴, Indian J. Phys. 33 (1959) 388-394. [E_β, P_β, E_γ, P_γ, α_K/α_(L+M)]
- 1960Ea02 H.T. Easterday, R.L. Smith, Balanced Ion Chamber Measurement of Half-lives of Several Radioisotopes, Nucl. Phys. 20 (1960) 155-158. [Half-life]
- 1960Ge05 J.S. Geiger, R.L. Graham, G.T. Ewan, Conversion Electron Spectrum from Ce¹⁴⁴ Decay, Nucl. Phys. 16 (1960) 1-26. [E_{ce}, P_{ce}, multipolarity]
- 1960Sa22 H.J. Sathoff, T. Azuma, Decay of Cerium-144, Ohio State University Research Report R-35051, TID-6080, Appendix 8 (1960). [E_γ, P_γ]
- 1961Ge09 J.S. Geiger, R.L. Graham, G.T. Ewan, The Decay of Ce¹⁴⁴, Nucl. Phys. 28 (1961) 387-406. [E_{ce}, P_{ce}, e-γ, γ-γ coincidence]
- 1962BI05 B. Blake, R. Bobone, H. Frauenfelder, H.J. Lipkin, Transverse Polarization of Conversion Electrons Following β-decay, Nuovo Cimento 25 (1962) 942-945. [133.5-keV conversion-electron process]
- 1962Bu09 J. Burde, M. Rakavy, G. Engler, Lifetime Determination of Three Excited Levels in the Odd-odd Nucleus Pr¹⁴⁴, Phys. Lett. 1 (1962) 147-149. [Nuclear-level lifetime]
- 1962Bu22 J. Burde, M. Rakavy, G. Engler, Absolute Transition Probabilities in the Odd-odd Nucleus Pr¹⁴⁴, Phys. Rev. 128 (1962) 325-332. [Nuclear-level lifetime]
- 1962Fo04 N.V. Forafontov V.S. Shpinel, Ts. Vasilev, Study of Ce¹⁴⁴ Decay by Coincidence Method, Nucl. Phys. 35 (1962) 260-272. [E_{ce}, E_{Ae}, P_{ce} summed subshell ratios]
- 1963Bh11 R. Bhattacharyya, S. Shastry, E2-M1 Mixing in the 100 keV Transition of Pr¹⁴⁴, Indian J. Phys. 37 (1963) 357-358. [γ-γ directional angular correlation, multipolarity]

- 1963Co18 W. Collin, H. Daniel, S. Margulies, O. Mehling, P. Schmidlin, H. Schmitt, K.S. Sibudhi, The Spins of Pr¹⁴⁴ from β - γ Correlation Measurements, Phys. Lett. 5 (1963) 329-331. [Spin]
- 1963Cr11 E. Creutz, J. de Raedt, J.P. Deutsch, L. Grenacs, D. Siddique, Spin of the Pr¹⁴⁴ Ground State Beta-decay Measurements on Ce¹⁴⁴ and Pr¹⁴⁴, Phys. Lett. 6 (1963) 329-331. [Ground-state spin]
- 1963Fu16 M. Fujishiro, T. Azuma, Beta-spectra of Ce-144, Annu. Report Radiation Center of Osaka Prefect. 4 (1963) 86-89. [E $_{\beta}$, P $_{\beta}$, E $_{\gamma}$, P $_{\gamma}$]
- 1963Iw02 T. Iwashita, T. Inamura, Y. Ikemoto, S. Kageyama, Decay of ¹⁴⁴Ce, J. Phys. Soc. Japan 18 (1963) 1358-1365. [γ - γ coincidence, γ - γ directional angular correlation, level structure, spin]
- 1963Kn05 U. Kneissel, H. Schneider, Bestimmung der Spins des Grundzustandes und des 134 keV Niveaus von ¹⁴⁴Pr durch Messung der β - γ -Zirkularpolarisation am β -Zerfall des ¹⁴⁴₅₈Ce und ¹⁴⁴₅₉Pr, Physik Verhandl 14 (1963) 125, 13. [Spin]
- 1963Si10 R.M. Singru, R.S. Raghavan, R.M. Steffen, Unique Spin Assignment to the Pr¹⁴⁴ Ground State, Phys. Lett. 6 (1963) 319-320. [Ground state spin, multipolarity]
- 1964Az02 T. Azuma, Y. Sato, β - γ Directional Angular Correlation of Pr-144, Annu. Report Radiation Center of Osaka Prefect. 5 (1964) 43-47. [γ - γ directional angular correlation, β transition type]
- 1964Be36 E.E. Berlovich, Yu.K. Gusev, D.M. Khai, I. Shenaikh, Lifetimes of Excited States of W¹⁸², Pr¹⁴⁴ and Eu¹⁵¹, Bull. Acad. Sci. USSR, Phys. Ser. 28 (1965) 77-84. [Nuclear-level lifetime]
- 1964Mc22 R.E. McAdams, E.N. Hatch, Some Lifetime Measurements of Isomeric States in the Rare Earth Region with Magnetic Coincidence Beta-ray Spectrometers, USAEC Research and Development Report IS-1071, TID-4500 (1964) 127-135. [Nuclear-level lifetime]
- 1965Co19 W. Collin, H. Daniel, B. Martin, P. Schmidlin, H. Schmitt, β - γ -Richtungskorrelationen und β - γ -Zirkularpolarisations-Korrelationen beim Einfach Verbotenen β -Zerfall von Ce¹⁴⁴, Pr¹⁴⁴, Tm¹⁷⁰ und Hg²⁰³, Kolloquium über Beta-Zerfall und Schwache Wechselwirkungen, Heidelberg (1965) 213-220. [β - γ correlation, nuclear-level structure]
- 1965Fl02 K.F. Flynn, L.E. Glendenin, E.P. Steinberg, Half-life Determinations by Direct Decay, Nucl. Sci. Eng. 22 (1965) 416-419. [Half-life]
- 1965Re13 W. Reiser, Spinbestimmung bei Praseodym 144 durch Winkelkorrelationsmessungen, Atomkernenergie 10 (1965) 307-310. [Spin]
- 1966Be11 H. Beer, H. Schneider, (54 – 80) keV- γ - γ -Winkelkorrelation von Pr¹⁴⁴, Z. Naturforsch. 21A (1966) 174-175. [Ground-state spin, δ (53.4 keV)]
- 1966Da04 H. Daniel, G.Th. Kaschl, Spektralformen von (0 \rightarrow 0)- β -Übergängen mit Paritätswechsel: ¹⁴⁴Ce, ¹⁴⁴Pr und ¹⁶⁶Ho, Nucl. Phys. 76 (1966) 97-117. [E $_{\beta}$, P $_{\beta}$]
- 1967Gu17 S.L. Gupta, N.K. Saha, Gamma-Gamma Directional Correlations in Pr¹⁴⁴, Indian J. Phys. 41 (1967) 48-54. [γ - γ coincidence, γ - γ directional correlation, ground-state spin]
- 1968Da12 H. Daniel, W. Collin, M. Kuntze, S. Margulies, B. Martin, O. Mehling, P. Schmidlin, H. Schmitt, Kernspektroskopische Untersuchungen an ⁸Li, ⁴²K, ⁸⁶Rb, ¹⁴¹Ce, ¹⁴⁴Ce, ¹⁴⁴Pr und ²⁰³Hg, Nucl. Phys. A118 (1968) 689-697. [β - γ directional correlation, ground-state spin]

- 1968La10 F. Lagoutine, Y. le Gallic, J. Legrand, Determination Précise de Quelques Périodes Radioactives, Int. J. Appl. Radiat. Isot. 19 (1968) 475-482. [Half-life]
- 1968Re04 S.A. Reynolds, J.F. Emery, E.I. Wyatt, Half-lives of Radionuclides III, Nucl. Sci. Eng. 32 (1968) 46-48. [Half-life]
- 1969Ge01 W. Gelletly, J.S. Geiger, L-subshell Conversion Line Intensity Ratios for Pure M1(1 ↔ 0) Transitions, Nucl. Phys. A123 (1969) 369-385. [L-subshell ratios]
- 1969Gu15 R. Gunnink, J.B. Niday, R.P. Anderson, R.A. Meyer, Gamma-ray Energies and Intensities, Lawrence Radiation Laboratory Report UCID-15439 (1969). [E_γ, P_γ]
- 1969Ma24 P.C. Mangal, P.N. Trehan, The Decay of ¹⁴⁴Ce, J. Phys. Soc. Japan 27 (1969) 1-7. [γ-γ coincidence, γ-γ angular correlation, nuclear structure, P_γ]
- 1970An15 A. Anttila, M. Piiparinen, The Levels of ¹⁴⁴Pr from the Decay of ¹⁴⁴Ce, Z. Phys. 237 (1970) 126-130. [E_γ, P_γ, α_K, α_L]
- 1970Fa03 J.L. Fasching, W.B. Walters, C.D. Coryell, Decay of 284-day ¹⁴⁴Ce, 17.3-min ¹⁴⁴Pr, and a New Isomer 7.2-min ^{144m}Pr, Phys. Rev. C1 (1970) 1126-1131. [E_γ, P_γ]
- 1970Po09 V.R. Potnis, G.P. Agin, C.E. Mandeville, The Decay of ¹⁴⁴Ce, J. Phys. Soc. Japan 29 (1970) 539-542. [P_γ]
- 1971Sa20 H.S. Sahota, Experimental Measurement of the K Conversion Coefficient of 80-keV Transition in ¹⁴⁴Pr, Curr. Sci., India 40 (1971) 289-290. [80-keV α_K]
- 1975Ba32 Chr. Bargholtz, S. Beshai, L. Eriksson, L.-E. Fröberg, L. Gidefeidt, The g-factor of the 80.1-keV Level in ¹⁴⁴Pr, Physica Scripta 11 (1975) 363-365. [δ(53 keV)]
- 1975De17 K. Debertain, U. Schötzig, K.F. Walz, H.M. Weiss, Gamma-ray Emission Probabilities of the Fission Products ¹⁴⁴Ce-¹⁴⁴Pr, ¹⁰⁶Rh, ⁹⁵Zr, Ann. Nucl. Energy 2 (1975) 37-43. [134-keV P_γ]
- 1976Ch33 J.M. Chatterjee-Das, R.K. Chattopadhyay, P. Bhattacharya, B. Sethi, S.K. Mukherjee, The Decay Scheme of the 284.3 d ¹⁴⁴Ce and Energy Levels and Transitions in ¹⁴⁴Pr, Radiochem. Radioanal. Lett. 27 (1976) 119-128. [E_γ, P_γ]
- 1976Ra22 B.V.N. Rao, G.N. Rao, Decay of ¹⁴⁴Ce to Levels in ¹⁴⁴Pr and ¹⁴⁴Nd, J. Phys. Soc. Japan 40 (1976) 1-5. [E_γ, P_γ]
- 1977Ge12 R.J. Gehrke, R.G. Helmer, R.C. Greenwood, Precise Relative γ-ray Intensities for Calibration of Ge Semiconductor Detectors, Nucl. Instrum. Methods 147 (1977) 405-423. [P_γ]
- 1977La19 F.P. Larkins, Semiempirical Auger-electron Energies for Elements 10 ≤ Z ≤ 100, At. Data Nucl. Data Tables 20 (1977) 311-387. [Auger-electron energies]
- 1978Mo22 T. Morii, Precision Energy Measurement of Gamma Rays of ⁵¹Cr, ¹⁰³Ru, ¹⁰⁹Cd, ¹⁴⁴Ce and ²⁰³Hg, Nucl. Instrum. Methods 151 (1978) 489-491. [133.5-keV E_γ]
- 1979Bo26 H.G. Borner, W.F. Davidson, J. Almeida, J. Blachot, J.A. Pinston, P.H.M. Van Assche, High Precision Gamma-ray Energy Measurements of Fission Products, Nucl. Instrum. Methods 164 (1979) 579-586. [133.5-keV E_γ]
- 1980Ho17 H. Houtermans, O. Milosevic, F. Reichel, Half-lives of 35 Radionuclides, Int. J. Appl. Radiat. Isot. 31 (1980) 153-154. [Half-life]

- 1981O104 J.B. Olomo, T.D. MacMahon, Gamma-ray Emission Probabilities in the Decay of ¹⁴⁴Ce and ¹⁴⁴Pr, Nucl. Energy 20 (1981) 237-240. [P_γ]
- 1982Yu02 B. Yu, F. Liu, X. Lu, S. Li, C. Yang, The Decay of ¹⁴⁴Ce, Radiochem. Radioanal. Lett. 53 (1982) 351-358. [E_γ, P_γ, P_X, P_X]
- 1983E114 M.R. El-Aasser, A. Abdel-Haliem, P. Asfour, M.N.H. Comsan, On the Decay of ¹⁴⁴Ce, Arab J. Nucl. Sci. Appl. 16-2 (1983) 283-292. [E_γ, P_γ]
- 1983Wa26 K.F. Walz, K. Debertin, H. Schrader, Half-life Measurements at the PTB, Int. J. Appl. Radiat. Isot. 34 (1983) 1191-1199. [Half-life]
- 1984Da13 J. Dalmaso, H. Maria, A. Hachem, G. Ardisson, Energie et Intensité des Photons X et de ¹⁴⁴Ce, Nucl. Instrum. Methods Phys. Res. 221 (1984) 564-565. [E_γ, P_γ]
- 1986O101 J.B. Olomo, Half-life of Radiotherapeutic ¹⁴⁴Ce Used as Beta-ray Applicator, Radiation Effects 94 (1986) 109-112. [Half-life]
- 1992Un01 M.P. Unterweger, D.D. Hoppes, F.J. Schima, New and Revised Half-life Measurements Results, Nucl. Instrum. Methods Phys. Res. A312 (1992) 349-352. [Half-life]
- 1996Sc06 E. Schönfeld, H. Janßen, Evaluation of Atomic Shell Data, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527-533. [ω_K, ω_L, K_β/K_α, K_{α2}/K_{α1}, KLX/KLL, KXY/KLL]
- 1997Ma75 R.H. Martin, K.I.W. Burns, J.G.V. Taylor, A Measurement of the Half-lives of ⁵⁴Mn, ⁵⁷Co, ⁵⁹Fe, ⁸⁸Y, ⁹⁵Nb, ¹⁰⁹Cd, ¹³³Ba, ¹³⁴Cs, ¹⁴⁴Ce, ¹⁵²Eu, Nucl. Instrum. Methods Phys. Res. A390 (1997) 267-273. [Half-life]
- 1998ScZM E. Schönfeld, G. Rodloff, Tables of the Energies of K-Auger Electrons for Elements with Atomic Numbers in the Range from Z = 11 to Z = 100, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX E. Schönfeld, G. Rodloff, Energies and Relative Emission Probabilities of K X-rays for Elements with Atomic Numbers in the Range from Z = 5 to Z = 100, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2000Sc47 E. Schönfeld, H. Janßen, Calculation of Emission Probabilities of X-rays and Auger Electrons Emitted in Radioactive Disintegration Processes, Appl. Radiat. Isot. 52 (2000) 595-600. [P_X, P_{Ae}]
- 2001So16 A.A. Sonzogni, Nuclear Data Sheets for A = 144, Nucl. Data Sheets 93 (2001) 599-762. [Nuclear levels]
- 2002Ba85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, Dirac-Fock Internal Conversion Coefficients, At. Data Nucl. Data Tables 81 (2002) 1-334. [ICC]
- 2002Ra45 S. Raman, C.W. Nestor, Jr., A. Ichihara, M.B. Trzhaskovskaya, How Good are the Internal Conversion Coefficients Now? Phys. Rev. C66 (2002) 044312, 1-23. [ICC]
- 2002Un02 M.P. Unterweger, Half-life Measurements at the National Institute of Standards and Technology, Appl. Radiat. Isot. 56 (2002) 125-130. [Half-life]
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr., Evaluation of Theoretical Conversion Coefficients using BrIcc, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202-229. [ICC]

- 2012Fi12 R. Fitzgerald, NIST Ionization Chamber "A" Sample-Height Corrections, J. Res. Natl. Inst. Stand. Technol. 117 (2012) 80-95. [Half-life]
- 2012Wa38 M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer, The AME2012 Atomic Mass Evaluation, (I). Tables, Graphs and References, Chin. Phys. C36 (2012) 1603-2014. [Q-value]
- 2014Un01 M.P. Unterweger, R. Fitzgerald, Update of NIST Half-life Results Corrected for Ionization Chamber Source-Holder Instability, Appl. Radiat. Isot. 87 (2014) 92-94. [Half-life]

**¹⁴⁴Pr – Comments on evaluation of decay data
by A. L. Nichols**

Evaluated: June/July 2013 and March/April 2014

Evaluation Procedures

Limitation of Relative Statistical Weight Method (LWM) and other analytical techniques were applied to average numbers throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average, unless stated otherwise.

Decay Scheme

¹⁴⁴Pr ($T_{1/2} = 17.29$ min) undergoes 100% β^- decay ($Q(\beta^-) = 2997.4$ (24) keV) via various excited levels to the ground state of ¹⁴⁴Nd. A reasonably well-defined decay scheme was derived from the studies of 1954Kr40, 1958Gr99, 1959Po77, 1968Ra01, 1968Sa05, 1969Gu15, 1969Ma24, 1970Fa03, 1974Be09, 1975De17, 1976Ra22, 1977Ge12, 1979Pr11, 1981Ol04, 1985Da16 and 2000He14, and can be defined in terms of ten β^- transitions and seventeen gamma-ray emissions.

Nuclear Data

¹⁴⁴Pr constitutes a major feature of the ¹⁴⁴Ce-¹⁴⁴Pr-¹⁴⁴Nd decay chain. The high-energy gamma rays generated in the β^- decay of ¹⁴⁴Pr are important in decay heat assessments and the design of short- and medium-term shielding requirements for irradiated fuel facilities.

Half-life

Half-life measurements of 1957Pe09, 1963Ho15, 1968Ra01 and 1970Fa03 were adopted to give a weighted-mean value of 17.29 ± 0.04 minutes, based on the LWM-NRM-Rajeval analysis techniques.

Reference	Half-life (min)
1957Pe09	17.27 ± 0.04
1963Ho15	17.30 ± 0.05
1968Ra01	17.3 ± 0.1
1970Fa03	17.3 ± 0.1
Recommended value	$17.29 \pm 0.04^*$

* Recommended uncertainty adjusted from ± 0.03 to 0.04, in alignment with the smallest uncertainty of the values used to calculate the weighted-mean value.

Analytical method	Half-life (d)	$\chi^2/(N-1)$	$\chi^2/(N-1)_{\text{critical}}$
LWM	17.29 ± 0.03	0.09	3.78
NRM	17.29 ± 0.03	0.09	2.60
Rajeval	17.29 ± 0.03	0.09	–
Bootstrap	17.295 ± 0.008	0.14	–
Mandel-Paule	null result	–	–

A half-life of (17.29 ± 0.04) minutes is recommended, as quantified by the LWM-NRM-Rajeval analytical procedures.

Q value

Q_{β^-} -value of 2997.4 (24) keV was adopted from Wang *et al.* (2012Wa38).

Gamma-ray energies and emission probabilities

Energies

The well-defined nuclear level energies of 2001So16 were used to calculate the gamma transition energies and their uncertainties, and these data were adjusted to account for gamma recoil in the formulation of recommended gamma-ray emission energies and uncertainties. Greater confidence was placed in this approach because of the more wide-ranging origins of the level energies compared with the measured gamma-ray decay data, even though the energies of a significant number of gamma-ray

emissions have been directly measured by 1968Ra01, 1970Fa03, 1974Be09, 1976Ra22, 1979Pr11 and 1985Da16. Nevertheless, the recommended gamma-ray energy standards proposed by 2000He14 have been adopted in the form of the 696.505(4)-, 1489.148(3)- and 2185.645(5)-keV gamma rays. The long half-life for the α decay of the ¹⁴⁴Nd ground state was determined from the measurements of 1954Wa05, 1956Po16, 1961Ma05, 1965Is01 and 1987A128, while other nuclear-level half-lives were adopted and calculated from the half-life and mean lifetime measurements of 1976CoZX, 1994Ro13 and 1998Hi09.

Adopted energies, spins and parities for the nuclear levels of ¹⁴⁴Nd (2001So16).

Nuclear level number*	Nuclear level energy (keV)	Spin and parity
0	0.0	0 + 2.3 (3) × 10 ¹⁵ a
1	696.561 ± 0.010	2 + –
2	1314.669 ± 0.013	4 + 7.4 (9) ps
3	1510.871 ± 0.021	3 – 0.56 (7) ps
4	1560.920 ± 0.013	2 + –
(5)	2072.91 ± 0.03	2 + 59 (10) fs
(6)	2084.68 ± 0.03	0 + 0.12 (5) ps
(7)	2185.75 ± 0.03	1 – 15 (2) fs
(8)	2368.82 ± 0.04	2 + 39 (14) fs
(9)	2582.32 ± 0.06	(3 +) –
(10)	2655.54 ± 0.03	1 + 9.9 (2) fs
(11)	2675.61 ± 0.08	0 + 0.2 (1) ps
(12)	2742.99 ± 0.07	0 + 64 (40) fs
(13)	2946.04 ± 0.10	(2, 3, 4) – –

* Nuclear level numbers in parentheses have been defined only on the basis of their population by the β^- decay of ¹⁴⁴Pr and ^{144m}Pr – many other nuclear levels of ¹⁴⁴Nd are known, but have not been considered in this assignment process.

Transition and gamma-ray energies for the β^- decay of ¹⁴⁴Pr (2001So16).

	Transition energy (keV)	Gamma-ray energy, E _γ (keV)
$\gamma_{(7,4)}$ (Nd)	624.83 (3)	624.83 ± 0.03
$\gamma_{(7,3)}$ (Nd)	674.88 (4)	674.88 ± 0.04
$\gamma_{1,0}$ (Nd)	696.561(10)	696.559 ± 0.010*
$\gamma_{3,1}$ (Nd)	814.310 (23)	814.308 ± 0.023
$\gamma_{4,1}$ (Nd)	864.359 (16)	864.356 ± 0.016
$\gamma_{(12,4)}$ (Nd)	1182.07 (7)	1182.06 ± 0.07
$\gamma_{(5,1)}$ (Nd)	1376.35 (3)	1376.34 ± 0.03
$\gamma_{(6,1)}$ (Nd)	1388.12 (4)	1388.11 ± 0.04
$\gamma_{(7,1)}$ (Nd)	1489.19 (3)	1489.18 ± 0.03*
$\gamma_{4,0}$ (Nd)	1560.920 (13)	1560.911 ± 0.013
$\gamma_{(8,1)}$ (Nd)	1672.26 (4)	1672.25 ± 0.04
$\gamma_{(11,1)}$ (Nd)	1979.05 (8)	1979.04 ± 0.08
$\gamma_{(12,1)}$ (Nd)	2046.43 (7)	2046.41 ± 0.07
$\gamma_{(5,0)}$ (Nd)	2072.91 (3)	2072.89 ± 0.03
$\gamma_{(7,0)}$ (Nd)	2185.75 (3)	2185.73 ± 0.03*
$\gamma_{(8,0)}$ (Nd)	2368.82 (4)	2368.80 ± 0.04
$\gamma_{(10,0)}$ (Nd)	2655.54 (3)	2655.51 ± 0.03

* Subsequently modified in line with the recommended energy standard of Helmer and van der Leun (2000He14).

Half-life of ¹⁴⁴Nd (α decay).

Reference	Half-life (years)
1954Wa05	(1.5 ± 0.8) × 10 ¹⁵ †
1956Po16	(2.2 ± 0.7) × 10 ¹⁵ ‡
1961Ma05	(2.4 ± 0.3) × 10 ¹⁵
1965Is01	(2.1 ± 0.4) × 10 ¹⁵
1987A128	(2.65 ± 0.37) × 10 ¹⁵
Recommended value	(2.3 ± 0.3) × 10 ¹⁵ *

† Estimated uncertainty – defined by 1954Wa05 as uncertain by a factor of at least 2.

‡ Estimated uncertainty – defined by 1956Po16 as possessing a margin of error of approximately 30%.

* Recommended half-life and uncertainty have been adjusted from (2.33 ± 0.19) × 10¹⁵ to (2.3 ± 0.3) × 10¹⁵ years, in alignment with the smallest uncertainty of the values used to calculate the weighted-mean value.

Measured and recommended gamma-ray energies.

E _γ (keV)									
1968Ra01	1968Sa05	1970Fa03	1974Be09	1976Ra22	1979Gr01	1979Pr11	1985Da16	2000He14	Recommended [#]
–	–	–	–	609 (1)	–	–	–	–	not adopted
–	–	–	–	–	–	618.2 (5)	617.8 (2)	–	^{144m} Pr
626 (3)	–	625.0 (10)	624.66 (20)	625.8 (4)	–	624.89 (20)	624.7 (7)	–	624.83 ± 0.03
675.3 (6)	–	675.0 (3)	675.02 (20)	674.9 (2)	–	675.02 (10)	674.95 (10)	–	674.88 ± 0.04
696.3 (4)	696.48 (9)	696.43 (6)	696.49 (2)	696.4 (1)	696.510 (6)	696.50 (2)	696.50 (5)	696.505 (4)	696.505 ± 0.004*
813.7 (6)	–	813.8 (3)	814.15 (15)	813.6 (2)	–	814.03 (10)	814.10 (10)	–	814.308 ± 0.023
863.6 (6)	–	863.9 (3)	864.53 (15)	863.5 (5)	–	864.44 (10)	864.45 (10)	–	864.356 ± 0.016
–	–	–	–	–	–	–	1182.0 (3)	–	1182.06 ± 0.07
–	–	–	–	–	–	1376.31 (30)	1376.27 (10)	–	1376.34 ± 0.03
1388.6 (6)	–	1388.0 (4)	1388.00 (15)	1388.2 (4)	–	1388.20 (10)	1388.02 (10)	–	1388.11 ± 0.04
–	–	–	–	1421.0 (5)	–	–	–	–	not adopted
1489.5 (4)	1489.14 (7)	1489.2 (2)	1489.15 (5)	1489.3 (2)	1489.160 (8)	1489.16 (5)	1489.17 (5)	1489.148 (3)	1489.148 ± 0.003*
–	–	–	1562 (2)	1558.4 (8)	–	1561.0	1560.97 (10)	–	1560.911 ± 0.013
–	–	–	–	–	–	–	1631.36 (10)	–	^{144m} Pr
–	–	–	–	–	–	–	–	–	1672.25 ± 0.04
–	–	–	–	–	–	–	1885.3 (2)	–	^{144m} Pr
–	–	–	–	–	–	1978.67 (10)	1978.82 (10)	–	1979.04 ± 0.08
–	–	–	–	–	–	2046.6 (5)	2046.3 (2)	–	2046.41 ± 0.07
–	–	–	–	–	–	2072.7 (7)	2072.9 (2)	–	2072.89 ± 0.03
2114.0 (20)	–	–	–	2114.0 (20)	–	–	–	–	not adopted
2185.6 (4)	2185.72 (5)	2185.6 (2)	2185.70 (6)	2185.5 (4)	2185.662 (11)	2185.60 (5)	2185.65 (5)	2185.645 (5)	2185.645 ± 0.005*
–	–	–	–	–	–	2368.4 (10)	2368.3 (3)	–	2368.80 ± 0.04
–	–	–	2654.6 (7)	2656.4 (4)	–	2654.1 (5)	2654.9 (2)	–	2655.51 ± 0.03
–	–	–	–	2802 (2)	–	–	–	–	not adopted
–	–	–	–	–	–	2842.9 (10)	–	–	not adopted

[#] Determined from the nuclear level energies of 2001So16 unless stated otherwise.

* Gamma-ray energy adopted from the recommended energy standards published by Helmer and van der Leun (2000He14).

Emission Probabilities

Relative gamma-ray emission probabilities have been partially or fully determined in the measurements of 1954Kr40, 1958Gr99, 1968Ra01, 1968Sa05, 1969Ma24, 1970Fa03, 1974Be09, 1975De17, 1976Ra22, 1977Ge12, 1979Pr11, 1981Ol04 and 1985Da16, and limited studies of the absolute gamma-ray emission probabilities have been performed by 1959Po77, 1969Gu15, 1975De17 and 1981Ol04.

While considerable emphasis was placed on the well-defined measurements of the absolute gamma-ray emission probabilities by 1975De17 and 1981Ol04, data from studies of both the absolute and relative gamma-ray emission probabilities as reported by 1954Kr40, 1958Gr99, 1959Po77, 1968Ra01, 1968Sa05, 1969Gu15, 1969Ma24, 1970Fa03, 1974Be09, 1976Ra22, 1977Ge12, 1979Pr11 and 1985Da16 were also used in weighted-mean analyses of the various gamma-ray emissions. All of the relative emission probabilities were suitably quantified in terms of the emission probability of the 696.505-keV gamma ray (100.0%). Although 696.505- and 814.308-keV gamma rays are emitted in the β^- decay of both ¹⁴⁴Pr and ^{144m}Pr, the total β^- branch of ^{144m}Pr is only 0.0006 (2) (i.e. (0.06 (2) %), and therefore this additional complication has no significant impact when determining the emission probabilities of these particular gamma rays in the β^- decay of the ¹⁴⁴Pr ground state.

Unfortunately, there are significant discrepancies between the gamma-ray emission probabilities measured by Debertain *et al.* (1975De17) and Olomo and MacMahon (1981Ol04), both undertaken with well calibrated Ge(Li) detectors. Thus, the absolute emission probability of the main 696.505-keV gamma ray differs by ~ 10% with measured values of 1.342 (13) % (1975De17) and 1.484 (12) % (1981Ol04). The LWM weighted-mean value of these data is (1.41 ± 0.07) %, and compares with a value of (1.49 ± 0.09) % measured at a significantly earlier date by means of a NaI(Tl) detector (1959Po77).

Difficulties have been experienced in determining the emission probability of the 1672.25-keV gamma ray, and a somewhat speculative P_{γ}^{rel} (1672.25 keV) value of 0.015 (4) was adopted from measurements of the relative gamma-ray emission probabilities of the 1672.25- and 2368.80-keV gamma transitions that depopulate the 2368.82-keV nuclear level following neutron capture by ¹⁴⁴Nd (1983Sn04, 1998Hi09):

Reference	E_{γ} (keV)	Relative intensity (I_{γ}^{rel})	Ratio [$I_{\gamma}^{rel}(1672.25 \text{ keV}) / I_{\gamma}^{rel}(2368.80 \text{ keV})$]
1983Sn04	1672.25	28.4 (2)	4.12 (7)
	2368.80	6.9 (1)	
1998Hi09	1672.25	84 (1)	5.25 (33)
	2368.80	16 (1)	
Mean			4.17 (23)
Evaluation	1672.25	0.015 (4) calculated	4.17
	2368.80	0.0036 (10)	

Despite the detection of a low-intensity 2114.0-keV gamma emission by Raman (1968Ra01) and 609-, 1421.0-, 2114.0- and 2802-keV gamma emissions by Rao and Rao (1976Ra22), the spectral observations are unconvincing, and therefore these particular gamma transitions have not be included in the evaluation of the decay scheme. Furthermore, Dalmaso *et al.* (1985Da16) searched unsuccessfully for the 2842.9-keV gamma emission observed by Pravikoff *et al.* (1979Pr11), and hence this high-energy gamma transition was also discarded.

Multipolarities, and Internal-Conversion and Internal-Pair Formation Coefficients

The nuclear level scheme specified by Sonzogni has been used to define the multipolarities of the gamma transitions on the basis of the known spins and parities (2001So16). Adopted multipolarities are 100%E1 for the 624.83-, 814.308-, 1489.148- and 2185.645-keV gamma rays, and 100%E2 for the 674.88-, 696.505-, 1182.06-, 1388.11-, 1560.911-, 1979.04-, 2046.41-, 2072.89- and 2368.80-keV gamma rays. Evidence for the mixing ratios and multipolarities of the 864.356-keV (51.5%M1 + 48.5%E2), 1376.34-keV (89.6%M1 + 10.4%E2) and 1672.25-keV (97.5%M1 + 2.5%E2) gamma transitions arise from γ - γ correlation studies by 1983Kr09, 1983Sn04 and 1998Hi09.

Extensive conversion-electron, e- γ , γ - γ coincidence and directional angular correlation studies throughout the 1960s focused on defining the spin-parity for the 0 – ground state of ¹⁴⁴Pr (1963Co18, 1963Cr11, 1963Iw02, 1963Kn05, 1963Si10, 1965Co19, 1965Re13, 1966Be11, 1967Gu17, 1968Da12, 1969Ge01 and 1969Ma24).

Recommended internal conversion coefficients have been determined from the frozen orbital approximation of Kibédi *et al.* (2008Ki07), based on the theoretical model of Band *et al.* (2002Ba85, 2002Ra45). A significant number of gamma transitions undergo decay via internal-pair formation, and the coefficient for this process has also been quantified from the tabulations of 2008Ki07.

Published gamma-ray emission probabilities - measured and recommended gamma-ray emission probabilities relative to P_γ(696.505 keV) of 100%.

E _γ (keV)	P _γ (%)									
	1954Kr40 ⁺	1958Gr99 ⁺	1959Po77 ⁰	1968Ra01 ⁺	1968Sa05 ⁺	1969Gu15 ⁰	1969Ma24 ⁺	1970Fa03 ⁺	1974Be09 ⁺	1975De17 ⁰
[609 (1)]	-	-	-	-	-	-	-	-	-	-
624.83 (3)	-	-	-	0.15 (4)	-	-	-	0.045 (20)	0.085 (15)	-
674.88 (4)	-	-	-	0.23 (6)	-	-	-	0.172 (17)	0.207 (15)	-
696.505 (4)	0.116 (9) → 100	100	1.49 (9) → 100	100	100	1.51 (15) → 100	100	100	100	1.342 (13) → 100
814.308 (23)	-	-	-	0.29 (7)	-	-	-	0.228 (22)	0.245 (13)	-
864.356 (16)	-	-	-	0.26 (7)	-	-	-	0.156 (15)	0.193 (12)	-
1182.06 (7)	-	-	-	-	-	-	-	-	-	-
1376.34 (3)	-	-	-	-	-	-	-	-	-	-
1388.11 (4)	-	-	-	0.45 (11)	-	-	-	0.420 (42)	0.445 (16)	-
[1421.0 (5)]	-	-	-	-	-	-	-	-	-	-
1489.148 (3)	0.0226 (18) → 19.5 (16)	17 (2)	0.29 (3) → 19.5 (20)	20.2 (20)	17.7 (5)	0.35 (7) → 23 (5)	22 (2)	19.5 (14)	20.3 (7)	0.279 (3) → 20.7 (3)
1560.911 (13)	-	-	-	-	-	-	-	-	~ 0.017	-
1672.25 (4)	-	-	-	-	-	-	-	-	-	-
1979.04 (8)	-	-	-	-	-	-	-	-	-	-
2046.41 (7)	-	-	-	-	-	-	-	-	-	-
2072.89 (3)	-	-	-	-	-	-	-	-	-	-
[2114.0 (20)]	-	-	-	0.03 (1)	-	-	-	< 0.016	-	-
2185.645 (5)	0.059 (5) → 51 (4)	50 (5)	0.68 (10) → 46 (7)	49.5 (50)	48 (9)	0.86 (17) → 57 (11)	64 (6)	49.5 (35)	52.2 (20)	0.700 (10) → 51.7 (10)
2368.80 (4)	-	-	-	-	-	-	-	-	-	-
2655.51 (3)	-	-	-	-	-	-	-	0.011 (3)	0.014 (2)	-
[2802 (2)]	-	-	-	-	-	-	-	-	-	-
[2842.9 (10)]	-	-	-	-	-	-	-	-	-	-

E _γ (keV)	P _γ (%)					
	1976Ra22*	1977Ge12*	1979Pr11*	1981O104#	1985Da16*	Recommended†
[609 (1)]	0.06 (2)	–	–	–	–	not adopted
624.83 (3)	0.09 (2)	–	0.092 (9)	–	0.084 (2)	0.084 (2)
674.88 (4)	0.23 (1)	–	0.195 (20)	–	0.22 (2)	0.212 (10)‡
696.505 (4)	100	100 (1)	100	1.484 (12) → 100 (1)	100	100
814.308 (23)	0.23 (1)	–	0.231 (25)	–	0.24 (2)	0.235 (10)‡
864.356 (16)	0.19 (1)	–	0.197 (20)	–	0.18 (2)	0.191 (10)‡
1182.06 (7)	–	–	–	–	0.004	0.004 (2)§
1376.34 (3)	–	–	0.029 (3)	–	0.029 (3)	0.029 (3)
1388.11 (4)	0.39 (3)	–	0.506 (20)	–	0.501 (4)	0.501 (4)
[1421.0 (5)]	0.03 (1)	–	–	–	–	not adopted
1489.148 (3)	20.0 (4)	20.27 (21)	21.8 (9)	0.277 (3) → 18.7 (3)	22.0 (12)	20.3 (2)¶
1560.911 (13)	0.11 (1)	–	0.016 (5)	–	0.015 (2)	0.015 (2)
1672.25 (4)	–	–	–	–	–	0.015 (4)Δ
1979.04 (8)	–	–	0.071 (7)	–	0.065 (6)	0.068 (6)‡
2046.41 (7)	–	–	0.021 (4)	–	0.020 (4)	0.021 (4)‡
2072.89 (3)	–	–	0.016 (4)	–	0.017 (2)	0.017 (2)‡
[2114.0 (20)]	0.03 (1)	–	–	–	–	not adopted
2185.645 (5)	49.9 (10)	52.3 (7)	55.7 (22)	0.768 (9) → 51.8 (7)	56.1 (25)	51.8 (7)‡
2368.80 (4)	–	–	0.0031 (10)	–	0.004 (1)	0.0036 (10)‡
2655.51 (3)	0.011 (2)	–	0.016 (2)	–	0.011 (2)	0.013 (2)‡
[2802 (2)]	0.004 (2)	–	–	–	–	not adopted
[2842.9 (10)]	–	–	0.008 (2)	–	–	not adopted

* Measured emission probabilities expressed relative to P_γ(133.515 keV) of 100% from β⁻ decay of parent ¹⁴⁴Ce; data have subsequently been adjusted to emission probabilities expressed relative to P_γ(696.505 keV) of 100% from β⁻ decay of ¹⁴⁴Pr.

* Emission probabilities expressed relative to P_γ(696.505 keV) of 100%.

° Measured absolute emission probabilities; data have subsequently been converted to relative emission probabilities (P_γ(696.505 keV) of 100%).

Measured absolute emission probabilities; authors also converted these data to relative emission probabilities (P_γ(696.505 keV) of 100%) – errors in Table 3 of 1981O104 have been noted (e-mail communications with T.D. MacMahon, 19/24 November 2013) and corrected.

† Weighted-mean values adopted when judged appropriate.

‡ Uncertainty adjusted upwards to align with smallest uncertainty of the values used to calculate the weighted-mean value.

§ Arbitrary uncertainty assigned of ± 0.002 (50%).

¶ Limitation of relative statistical weight method (LWM) produces a value of 20.07 (23) and χ²/(N-1) of 2.85, while the normalized residual method (NRM) defines a value of 20.31 (14) and χ²/(N-1) of 1.31; under these circumstances, a value of 20.3 (2) has been adopted, in good agreement with the well-defined measurement of 1975De17.

Δ Considerable difficulty has been experienced in resolving the 1672.25-keV gamma-ray emission, and therefore a speculative P_{γ^{rel}}(1672.25 keV) value of 0.015 (4) was adopted on the basis of the assumed relative probabilities of the 1672.25- and 2368.80-keV gamma transitions that depopulate the 2368.82-keV nuclear level of ¹⁴⁴Nd. A weighted mean for the P_{γ^{rel}}(1672.25 keV)/P_{γ^{rel}}(2368.80 keV) ratio of 4.17 (23) and a recommended P_{γ^{rel}}(2368.80 keV) of 0.0036(10) define P_{γ^{rel}}(1672.25 keV) of 0.015 (4).

Gamma-ray emissions: multiplicities, and theoretical internal-conversion (frozen orbital approximation) and internal-pair formation coefficients.

E_γ (keV)	Multipolarity	α_K	α_L	α_{L1}	α_{L2}	α_{L3}	α_{M+}	$\alpha_{totalICC}$	α_{IPF}	α_{total}
624.83 (3)	E1	0.002 07 (3)	0.000 267 (4)	0.000 242 (4)	0.000 011 55 (17)	0.000 0135 5 (19)	0.000 073	0.002 41 (4)	–	0.002 41 (4)
674.88 (4)	E2	0.004 60 (7)	0.000 686 (10)	0.000 547 (8)	0.000 088 3 (13)	0.000 051 1 (8)	0.000 184	0.005 47 (8)	–	0.005 47 (8)
696.505 (4)	E2	0.004 27 (6)	0.000 631 (9)	0.000 508 (8)	0.000 078 5 (11)	0.000 045 1 (7)	0.000 169	0.005 07 (7)	–	0.005 07 (7)
814.308 (23)	E1	0.001 198 (17)	0.000 152 8 (22)	0.000 140 7 (20)	0.000 005 46 (8)	0.000 006 56 (10)	0.000 040 2	0.001 391 (20)	–	0.001 391 (20)
864.356 (16)	51.5%M1 + 48.5%E2 $\delta = -0.97$ (15)	0.003 38 (14)	0.000 456 (16)	0.000 414 (18)	0.000 029 0 (12)	0.000 012 3 (13)	0.000 122	0.003 96 (16)	–	0.003 96 (16)
1182.06 (7)	(1983Kr09, 1983Sn04, 1998Hi09) E2	0.001 353 (19)	0.000 182 (3)	0.000 162 8 (23)	0.000 012 15 (17)	0.000 006 62 (10)	0.000 048	0.001 583 (19)	0.000 004 10 (6)	0.001 587 (23)
1376.34 (3)	89.6%M1 + 10.4%E2 $\delta = 0.34$ (10)	0.001 35 (3)	0.000 175 (4)	0.000 166 (4)	0.000 006 30 (12)	0.000 002 00 (14)	0.000 046	0.001 57 (3)	0.000 039 8 (6)	0.001 61 (4)
1388.11 (4)	(1983Sn04, 1998Hi09) E2	0.000 984 (14)	0.000 129 7 (19)	0.000 118 6 (17)	0.000 007 18 (10)	0.000 003 96 (6)	0.000 035	0.001 149 (14)	0.000 041 6 (6)	0.001 190 (17)
1489.148 (3)	E1	0.000 397 (6)	0.000 049 5 (7)	0.000 046 9 (7)	0.000 001 109 (16)	0.000 001 529 (22)	0.000 012 5	0.000 459 (6)	0.000 204 (3)	0.000 663 (10)
1560.911 (13)	E2	0.000 786 (11)	0.000 102 4 (15)	0.000 094 7 (14)	0.000 004 95 (7)	0.000 002 78 (4)	0.000 027 5	0.000 916 (11)	0.000 098 1 (14)	0.001 014 (15)
1672.25 (4)	97.5%M1 + 2.5%E2 $\delta = 0.16$ (7)	0.000 892 (14)	0.000 114 6 (18)	0.000 109 8 (17)	0.000 003 61 (6)	0.000 001 19 (4)	0.000 030 4	0.001 037 (14)	0.000 151 9 (22)	0.001 189 (18)
1799.04 (8)	(1983Sn04, 1998Hi09) E2	0.000 505 (7)	0.000 064 7 (9)	0.000 060 9 (9)	0.000 002 40 (4)	0.000 001 434 (20)	0.000 017 3	0.000 587 (8)	0.000 281 (4)	0.000 868 (13)
2046.41 (7)	E2	0.000 475 (7)	0.000 060 7 (9)	0.000 057 3 (8)	0.000 002 17 (3)	0.000 001 314 (19)	0.000 016 3	0.000 552 (7)	0.000 313 (5)	0.000 865 (13)
2072.89 (3)	E2	0.000 465 (7)	0.000 059 3 (9)	0.000 055 9 (8)	0.000 002 09 (3)	0.000 001 271 (18)	0.000 015 7	0.000 540 (7)	0.000 326 (5)	0.000 865 (13)
2185.645 (5)	E1	0.000 213 (3)	0.000 026 4 (4)	0.000 025 2 (4)	0.000 000 437 (7)	0.000 000 702 (10)	0.000 007 6	0.000 246 (3)	0.000 712 (10)	0.000 959 (14)
2368.80 (4)	E2	0.000 365 (6)	0.000 046 3 (7)	0.000 044 0 (7)	0.000 001 420 (20)	0.000 000 911 (13)	0.000 012 7	0.000 424 (6)	0.000 467 (7)	0.000 891 (13)
2655.51 (3)	M1+E2	–	–	–	–	–	–	–	–	–

The recommended normalization factor for the relative gamma-ray emission probabilities is based on the somewhat discrepant yet well-defined gamma-ray spectroscopy studies of 1975De17 and 1981OIo4. Measured absolute emission probabilities for the main 696.505-keV gamma ray differ by ~ 10%, with values of 1.342 (13) % (1975De17) and 1.484 (12) % (1981OIo4), to give (a) LWM weighted-mean value of (1.41 ± 0.07) % and $\chi^2/(N-1)$ of 59.7 and (b) NRM weighted-mean value of (1.41 ± 0.05) % and $\chi^2/(N-1)$ of 1.98 (NRM). Under these unsatisfactory circumstances, a normalization factor of (0.0141 ± 0.0007) has been adopted.

Recommended gamma-ray energies, transition probabilities, and relative and absolute emission probabilities.

E_γ (keV)	P_γ^{rel}	P_γ^{abs} (%)	Transition probability (%)
624.83 (3)	0.084 (2)	0.001 18 (3)	0.001 18 (3)
674.88 (4)	0.212 (10)	0.002 99 (14)	0.003 01 (14)
696.505 (4)	100	1.41 (7)	1.42 (7)
814.308 (23)	0.235 (10)	0.003 31 (14)	0.003 31 (14)
864.356 (16)	0.191 (10)	0.002 69 (14)	0.002 70 (14)
1182.06 (7)	0.004 (2)	0.000 06 (3)	0.000 06 (3)
1376.34 (3)	0.029 (3)	0.000 41 (4)	0.000 41 (4)
1388.11 (4)	0.501 (4)	0.007 06 (6)	0.007 07 (6)
1489.148 (3)	20.3 (2)	0.286 (3)	0.286 (3)
1560.911 (13)	0.015 (2)	0.000 21 (3)	0.000 21 (3)
1672.25 (4)	0.015 (4)	0.000 21 (6)	0.000 21 (6)
1979.04 (8)	0.068 (6)	0.000 96 (8)	0.000 96 (8)
2046.41 (7)	0.021 (4)	0.000 30 (6)	0.000 30 (6)
2072.89 (3)	0.017 (2)	0.000 24 (3)	0.000 24 (3)
2185.645 (5)	51.8 (7)	0.730 (10)	0.731 (10)
2368.80 (4)	0.003 6 (10)	0.000 051 (14)	0.000 051 (14)
2655.51 (3)	0.013 (2)	0.000 18 (3)	0.000 18 (3)

β^- energies and emission probabilities

Energies

All β^- energies were derived from the structural details of the proposed decay scheme. The nuclear level energies of 2001So16 and evaluated Q-value of 2997.4 (24) keV (2012Wa38) were used to determine the recommended energies and uncertainties of the β^- emissions.

Emission Probabilities

Emission probabilities were derived from the population-depopulation imbalances of the relative emission probabilities of the gamma rays, their theoretical internal-conversion coefficients, and a normalization factor of (0.0141 ± 0.0007) for the gamma-ray emissions, as noted above. The dominant first forbidden β^- transition directly to the ground state of ¹⁴⁴Nd ($0^- \rightarrow 0^+$) and resulting shape factor have been studied in detail to confirm the nature of the transition and parity of the ¹⁴⁴Pr ground state (1959Po77, 1971Na12, 1973Bo43). β^- decay to the 1510.871-keV nuclear level of ¹⁴⁴Nd was assumed to be zero on the basis of spin and parity considerations that would have resulted in a second forbidden unique transition. Logft values and average E_{β^-} energies were determined by means of the LOGFT code.

Recommended energies and emission probabilities for the β^- decay of ¹⁴⁴Pr.

E_β (keV)*	Av. E_{β^-} (keV)	P_β	¹⁴⁴ Pr	¹⁴⁴ Nd	transition type	log ft
$\beta^-_{(0,12)}$ 254.4 (24)	71.05 (8)	0.000 35 (6)	0 -	0 +	first forbidden non-unique	8.1
$\beta^-_{(0,11)}$ 321.8 (24)	92.21 (8)	0.000 96 (8)	0 -	0 +	first forbidden non-unique	8.0
$\beta^-_{(0,10)}$ 341.9 (24)	98.68 (8)	0.000 18 (3)	0 -	1 +	first forbidden non-unique	8.8
$\beta^-_{(0,8)}$ 628.6 (24)	213.04 (9)	0.000 27 (6)	0 -	2 +	first forbidden unique	9.7 ^{lu}
$\beta^-_{(0,7)}$ 811.7 (24)	267.12 (9)	1.021 (10)	0 -	1 -	allowed	6.32
$\beta^-_{(0,6)}$ 912.7 (24)	306.67 (10)	0.007 08 (6)	0 -	0 +	first forbidden non-unique	8.7
$\beta^-_{(0,5)}$ 924.5 (24)	322.77 (9)	0.000 65 (6)	0 -	2 +	first forbidden unique	10.2 ^{lu}
$\beta^-_{0,4}$ 1436.5 (24)	526.25 (10)	0.001 7 (3)	0 -	2 +	first forbidden unique	10.8 ^{lu}
$\beta^-_{0,1}$ 2300.8 (24)	894.90 (11)	1.116 (3)	0 -	2 +	first forbidden unique	9.17 ^{lu}
$\beta^-_{0,0}$ 2997.4 (24)	1221.990 (1)	97.852 (10)	0 -	0 +	first forbidden non-unique	6.530
		Σ 100.000 19				

* Determined from the nuclear level energies of 2001So16 and Q-value of 2997.4 (24) keV (2012Wa38).

A consistent decay scheme was derived that contains ten β^- transitions and seventeen gamma-ray emissions.

Atomic Data

The x-ray and Auger-electron data have been calculated using the evaluated gamma-ray data, and atomic data from 1996Sc06, 1998ScZM and 1999ScZX. Both the X-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.02, 28 February 2012), as described in 2000Sc47. This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

K and L X-ray energies and emission probabilities of ¹⁴⁴Pr.

			Energy (keV)	Photons per 100 disint.	Relative probability
XL	(Nd)		4.633 – 6.901	0.000 92 (3)	
	XL ₁	(Nd)	4.633	0.000 017 9 (9)	
	XL _α	(Nd)	5.208 – 5.230	0.000 449 (20)	
	XL _η	(Nd)	5.146	0.000 006 9 (4)	
	XL _β	(Nd)	5.722 – 6.090	0.000 384 (14)	
	XL _γ	(Nd)	6.604 – 6.901	0.000 061 5 (23)	
XK _α	XK _{α2}	(Nd)	36.8478 (3)	0.001 65 (9)	55
	XK _{α1}	(Nd)	37.3614 (2)	0.003 00 (15)	100
XK' _{β1}	XK _{β3}	(Nd)	42.1670 (4))	
	XK _{β1} '	(Nd)	42.2717 (3)) 0.000 92 (5)	30.7
	XK _{β5}	(Nd)	42.580)	
XK' _{β2}	XK _{β2}	(Nd)	43.335 (3))	
	XK _{β4}	(Nd)	43.451) 0.000 237 (13)	7.9
	XKO _{2,3}	(Nd)	43.548)	

Auger-electron energies and emission probabilities of ¹⁴⁴Pr.

		Energy (keV)	Electrons per 100 disint.	Relative probability
e _{AK}	(Nd)		0.000 52 (4)	
	KLL	29.154 – 30.978	0.000 333 (23)	100
	KLX	34.798 – 37.340	0.000 166 (12)	50
	KXY	40.42 – 43.53	0.000 020 7 (15)	6.2
e _{AL}	(Nd)	3.01 – 5.10	0.005 51 (18)	1655

Nd: ω_K = 0.918 (4); ω_L = 0.140 (6); η_{KL} = 0.866 (4) were taken from 1996Sc06.

Electron energies were determined from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

Data Consistency

Q_{β⁻}-value of 2997.4 (24) keV has been adopted from the atomic mass evaluation of Wang *et al.* (2012Wa38) while in the course of formulating the decay scheme of ¹⁴⁴Pr. This value has subsequently been compared with the Q-value calculated by summing the contributions of the individual emissions of the ¹⁴⁴Pr decay processes (i.e. β⁻, electron, γ, etc.):

$$\text{calculated Q-value} = \sum (E_i \times P_i) = 2997.3 (25) \text{ keV}$$

Percentage deviation from the Q-value of Wang *et al.* is (0.00 ± 0.12) %, which supports the derivation of an extremely consistent decay scheme with a modest variant.

References

- 1954Kr40 W.E. Kreger, C.S. Cook, Gamma Radiation from Ce¹⁴⁴ and Pr¹⁴⁴, Phys. Rev. 96 (1954) 1276-1278. [P_γ]
- 1954Wa05 E.C. Waldron, V.A. Schultz, T.P. Kohman, Natural Alpha Activity of Neodymium, Phys. Rev. 93 (1954) 254-255. [¹⁴⁴Nd half-life]
- 1956Po16 W. Porschen, W. Riezler, Sehr Langlebige Natürliche α-Strahler, Z. Naturforsch. 11a (1956) 143-151. [¹⁴⁴Nd half-life]
- 1957Pe09 D.F. Peppard, G.W. Mason, S.W. Moline, The Use of Dioctyl Phosphoric Acid Extraction in the Isolation of Carrier-free ⁹⁰Y, ¹⁴⁰La, ¹⁴⁴Ce, ¹⁴³Pr and ¹⁴⁴Pr, J. Inorg. Nucl. Chem. 5 (1957) 141-146. [Half-life]
- 1958Gr99 R.L. Graham, J.S. Geiger, T.A. Eastwood, Experimental Evidence for Axial Vector Interaction in the Disintegration of Pr¹⁴⁴, Can. J. Phys. 36 (1958) 1084-1111. [E_γ, P_γ, P_β, β-γ directional correlation, β spectral shape]
- 1959Po77 F.T. Porter, P.P. Day, 0⁻ to 0⁺ Beta Transition Pr¹⁴⁴ → Nd¹⁴⁴, Phys. Rev. 114 (1959) 1286-1296. [E_β, P_β, P_γ]
- 1961Ma05 R.D. MacFarlane, T.P. Kohman, Natural Alpha Radioactivity in Medium-Heavy Elements, Phys. Rev. 121 (1961) 1758-1769. [¹⁴⁴Nd half-life]
- 1963Co18 W. Collin, H. Daniel, S. Margulies, O. Mehling, P. Schmidlin, H. Schmitt, K.S. Sibudhi, The Spins of Pr¹⁴⁴ from β-γ Correlation Measurements, Phys. Lett. 5 (1963) 329-331. [Spin]
- 1963Cr11 E. Creutz, J. de Raedt, J.P. Deutsch, L. Grenacs, D. Siddique, Spin of the Pr¹⁴⁴ Ground State Beta-decay Measurements on Ce¹⁴⁴ and Pr¹⁴⁴, Phys. Lett. 6 (1963) 329-331. [Ground-state spin]
- 1963Ho15 D.C. Hoffman, Half-lives of Some Rare Earth Nuclides, J. Inorg. Nucl. Chem. 25 (1963) 1196-1198. [Half-life]
- 1963Iw02 T. Iwashita, T. Inamura, Y. Ikemoto, S. Kageyama, Decay of ¹⁴⁴Ce, J. Phys. Soc. Japan 18 (1963) 1358-1365. [γ-γ coincidence, γ-γ directional angular correlation, level structure, spin]
- 1963Kn05 U. Kneissel, H. Schneider, Bestimmung der Spins des Grundzustandes und des 134 keV Niveaus von ¹⁴⁴Pr durch Messung der β-γ-Zirkularpolarisation am β-Zerfall des ¹⁴⁴₅₈Ce und ¹⁴⁴₅₉Pr, Physik Verhandl 14 (1963) 125, 13. [Spin]
- 1963Si10 R.M. Singru, R.S. Raghavan, R.M. Steffen, Unique Spin Assignment to the Pr¹⁴⁴ Ground State, Phys. Lett. 6 (1963) 319-320. [Ground-state spin, multipolarity]
- 1965Co19 W. Collin, H. Daniel, B. Martin, P. Schmidlin, H. Schmitt, β-γ-Richtungskorrelationen und β-γ-Zirkularpolarisations-Korrelationen beim Einfach Verbotenen β-Zerfall von Ce¹⁴⁴, Pr¹⁴⁴, Tm¹⁷⁰ und Hg²⁰³, Kolloquium über Beta-Zerfall und Schwache Wechselwirkungen, Heidelberg (1965) 213-220. [β-γ correlation, nuclear-level structure]
- 1965Is01 A. Isola, N. Nurmia, Alpha Activity of Natural Neodymium, Z. Naturforsch. 20a (1965) 541-542. [¹⁴⁴Nd half-life]

- 1965Re13 W. Reiser, Spinbestimmung bei Praseodym 144 durch Winkelkorrelationsmessungen, Atomkernenergie 10 (1965) 307-310. [Spin]
- 1966Be11 H. Beer, H. Schneider, (54 – 80) keV- γ - γ -Winkelkorrelation von Pr¹⁴⁴, Z. Naturforsch. 21a (1966) 174-175. [Ground-state spin, δ (53.4 keV)]
- 1967Gu17 S.L. Gupta, N.K. Saha, Gamma-Gamma Directional Correlations in Pr¹⁴⁴, Indian J. Phys. 41 (1967) 48-54. [γ - γ coincidence, γ - γ directional correlation, ground-state spin]
- 1968Da12 H. Daniel, W. Collin, M. Kuntze, S. Margulies, B. Martin, O. Mehling, P. Schmidlin, H. Schmitt, Kernspektroskopische Untersuchungen an ⁸Li, ⁴²K, ⁸⁶Rb, ¹⁴¹Ce, ¹⁴⁴Ce, ¹⁴⁴Pr und ²⁰³Hg, Nucl. Phys. A118 (1968) 689-697. [β - γ directional correlation, ground-state spin]
- 1968Ra01 S. Raman, Decay of ¹⁴⁴Pr to Levels in ¹⁴⁴Nd, Nucl. Phys. A107 (1968) 402-410. [Half-life, E γ , P γ]
- 1968Sa05 A.R. Sayres, C.C. Trail, Precision Measurements of the Energies of the Gamma Rays from ¹⁴⁴Nd, Nucl. Phys. A113 (1968) 521-528. [E γ , P γ]
- 1969Ge01 W. Gelletly, J.S. Geiger, L-subshell Conversion Line Intensity Ratios for Pure M1(1 \leftrightarrow 0) Transitions, Nucl. Phys. A123 (1969) 369-385. [L-subshell ratios (59.03 keV), Ground-state spin]
- 1969Gu15 R. Gunnink, J.B. Niday, R.P. Anderson, R.A. Meyer, Gamma-ray Energies and Intensities, Lawrence Radiation Laboratory Report UCID-15439 (1969). [E γ , P γ]
- 1969Ma24 P.C. Mangal, P.N. Trehan, The Decay of ¹⁴⁴Ce, J. Phys. Soc. Japan 27 (1969) 1-7. [P γ]
- 1970Fa03 J.L. Fasching, W.B. Walters, C.D. Coryell, Decay of 284-day ¹⁴⁴Ce, 17.3-min ¹⁴⁴Pr, and a New Isomer 7.2-min ^{144m}Pr, Phys. Rev. C1 (1970) 1126-1131. [Half-life, E γ , P γ , P β]
- 1971Na12 T. Nagarajan, M. Ravindranath, K.V. Reddy, The 0⁻ \rightarrow 0⁺ Decay of ¹⁴⁴Pr and the Pseudoscalar Interaction, Nuovo Cimento 3A (1971) 699-708. [(0⁻ \rightarrow 0⁺) β^- shape factor]
- 1973Bo43 H.E. Bosch, M. Behar, M.C. Cambiaggio, G.G. Bermudez, L. Szybisz, Investigation of the 0⁻ \rightarrow 0⁺ and 0⁻ \rightarrow 2⁺ Beta Transitions in the Decay of ¹⁴⁴Pr, Can. J. Phys. 51 (1973) 2260-2269. [(0⁻ \rightarrow 0⁺) and (0⁻ \rightarrow 2⁺) β^- shape factors]
- 1974Be09 M. Behar, Z.W. Grabowski, S. Raman, Angular Correlation Studies in ¹⁴⁴Nd, Nucl. Phys. A219 (1974) 516-524. [E γ , P γ , δ , BF]
- 1975De17 K. Debertin, U. Schötzig, K.F. Walz, H.M. Weiss, Gamma-ray Emission Probabilities of the Fission Products ¹⁴⁴Ce-¹⁴⁴Pr, ¹⁰⁶Rh, ⁹⁵Zr, Ann. Nucl. Energy 2 (1975) 37-43. [P γ]
- 1976CoZX J.P. Collins, C.T. Wunker, R.L. Place, D.R. Ober, Nuclear Lifetime Measurement of the 1314-keV Level of Nd-144, Bull. Amer. Phys. Soc. 21, No. 2 (1976) 149, D1. [Nuclear-level half-life (1314.67 keV)]
- 1976Ra22 B.V.N. Rao, G.N. Rao, Decay of ¹⁴⁴Ce to Levels in ¹⁴⁴Pr and ¹⁴⁴Nd, J. Phys. Soc. Japan 40 (1976) 1-5. [E γ , P γ]
- 1977Ge12 R.J. Gehrke, R.G. Helmer, R.C. Greenwood, Precise Relative γ -ray Intensities for Calibration of Ge Semiconductor Detectors, Nucl. Instrum. Methods 147 (1977) 405-423. [P γ]
- 1977La19 F.P. Larkins, Semiempirical Auger-electron Energies for Elements 10 \leq Z \leq 100, At. Data Nucl. Data Tables 20 (1977) 311-387. [Auger-electron energies]

- 1979Gr01 R.C. Greenwood, R.G. Helmer, R.J. Gehrke, Precise γ -ray Energies for Calibration of Ge Semiconductor Spectrometers to 3.5 MeV, Nucl. Instrum. Methods 159 (1979) 465-482. [E $_{\gamma}$]
- 1979Pr11 M.S. Pravikoff, G. Barci-Funel, G. Ardisson, Sur la Décroissance de ^{144m+g}Pr et les Niveaux de ¹⁴⁴Nd, Radiochem. Radioanal. Lett. 40 (1979) 123-132. [E $_{\gamma}$, P $_{\gamma}$]
- 1981O104 J.B. Olomo, T.D. MacMahon, Gamma-ray Emission Probabilities in the Decay of ¹⁴⁴Ce and ¹⁴⁴Pr, Nucl. Energy 20 (1981) 237-240. [P $_{\gamma}$]
- 1983Kr09 K.S. Krane, S. Raman, F.K. McGowan, First-excited 0⁺ State in ¹⁴⁴Nd, Phys. Rev. C27 (1983) 2863-2868. [Spin, δ]
- 1983Sn04 D.M. Snelling, W.D. Hamilton, Gamma-gamma Directional Correlation Measurements in ¹⁴⁴Nd, J. Phys. G: Nucl. Phys. 9 (1983) 763-778. [Nuclear structure, δ]
- 1985Da16 J. Dalmaso, H. Forest, G. Ardisson, ¹⁴⁴Nd Levels Fed in β decay of ¹⁴⁴Pr Isomers, Phys. Rev. 32 (1985) 1006-1009. [E $_{\gamma}$, P $_{\gamma}$, BF]
- 1987A128 B. Al-Bataina, J. Jänecke, Half-lives of Long-lived Alpha Emitters, Radiochim. Acta 42 (1987) 159-164. [¹⁴⁴Nd half-life]
- 1994Ro13 S.L. Robinson, J. Jolie, H.G. Börner, P. Schillebeeckx, S. Ulbig, K.P. Lieb, E2 and E3 Transitions from Quadrupole-Octupole Coupled States in ¹⁴⁴Nd, Phys. Rev. Lett. 73 (1994) 412-415. [Nuclear-level lifetimes]
- 1996Sc06 E. Schönfeld, H. Janßen, Evaluation of Atomic Shell Data, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527-533. [ω_K , ω_L , K $_{\beta}$ /K $_{\alpha}$, K $_{\alpha 2}$ /K $_{\alpha 1}$, KLX/KLL, KXY/KLL]
- 1998Hi09 S.F. Hicks, C.M. Davoren, W.M. Faulkner, J.R. Vanhoy, Structural Characteristics of ¹⁴⁴Nd Through γ -ray Spectroscopy Following Inelastic Neutron Scattering, Phys. Rev. 57 (1998) 2264-2280. [Nuclear structure, δ]
- 1998ScZM E. Schönfeld, G. Rodloff, Tables of the Energies of K-Auger Electrons for Elements with Atomic Numbers in the Range from Z = 11 to Z = 100, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX E. Schönfeld, G. Rodloff, Energies and Relative Emission Probabilities of K X-rays for Elements with Atomic Numbers in the Range from Z = 5 to Z = 100, PTB Report PTB-6.11-1999-1, February 1999. [X $_K$]
- 2000He14 R.G. Helmer, C. van der Leun, Recommended Standards for γ -ray Energy Calibration, Nucl. Instrum. Methods Phys. Res. A450 (2000) 35-70. [E $_{\gamma}$]
- 2000Sc47 E. Schönfeld, H. Janßen, Calculation of Emission Probabilities of X-rays and Auger Electrons Emitted in Radioactive Disintegration Processes, Appl. Radiat. Isot. 52 (2000) 595-600. [P $_X$, P $_{Ae}$]
- 2001So16 A.A. Sonzogni, Nuclear Data Sheets for A = 144, Nucl. Data Sheets 93 (2001) 599-762. [Nuclear levels]
- 2002Ba85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, Dirac-Fock Internal Conversion Coefficients, At. Data Nucl. Data Tables 81 (2002) 1-334. [ICC]
- 2002Ra45 S. Raman, C.W. Nestor, Jr., A. Ichihara, M.B. Trzhaskovskaya, How Good are the Internal Conversion Coefficients Now? Phys. Rev. C66 (2002) 044312, 1-23. [ICC]

- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr., Evaluation of Theoretical Conversion Coefficients using BrIcc, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202-229. [ICC]
- 2012Wa38 M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer, The AME2012 Atomic Mass Evaluation, (I). Tables, Graphs and References, Chin. Phys. C36 (2012) 1603-2014. [Q-value]

^{144m}Pr – Comments on evaluation of decay data
by A. L. Nichols

Evaluated: May/June 2013 and March/April 2014

Evaluation Procedures

Limitation of Relative Statistical Weight Method (LWM) and other analytical techniques were applied to average numbers throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

^{144m}Pr ($T_{1/2} = 7.2$ min) undergoes 99.94% IT decay ($Q_{IT} = 59.03$ (3) keV) and 0.06% β^- decay ($Q(\beta^-) = 3056.4$ (24) keV) via various excited levels to the ground state of ¹⁴⁴Nd. The decay scheme is dominated by a simple IT decay mode consisting of a single 59.03-keV gamma transition, whereas the small β^- decay mode is poorly characterized. Despite the latter inadequacies, a reasonably well-defined decay scheme can be derived primarily from 1960Ge05 and 1985Da16 in terms of three low-intensity β^- transitions and six gamma transitions, in which the conversion-electron emissions identified with the 59.03-keV M3 gamma transition play a dominant role.

Nuclear Data

^{144m}Pr constitutes a minor branch in the ¹⁴⁴Ce-¹⁴⁴Pr-¹⁴⁴Nd decay chain. A limited number of reasonably high-energy gamma rays of low intensity are generated in the β^- decay mode of ^{144m}Pr.

Half-life

Only half-life measurements by Fasching *et al.* (1970Fa03) and Chatterjee-Das *et al.* (1976Ch33) have been published, of which the value reported by 1976Ch33 was described as being tentative.

Reference	Half-life (min)
1970Fa03	7.2 ± 0.2
1976Ch33	$6.6 \pm 1.0^*$
Recommended value	7.2 ± 0.2

* Half-life assigned tentatively to ^{144m}Pr.

Although such a small data set still permits analysis by both the limitation of relative statistical weight method (LWM) and the normalised residual method (NRM), weighted uncertainty adjustments have not been implemented because of the nature of the 1976Ch33 measurement. Thus, a half-life value of (7.2 ± 0.2) minutes is recommended, as determined by 1970Fa03.

Q values

Q_{IT} -value of 59.03 (3) keV and Q_{β^-} -value of 3056.4 (24) keV were adopted for the ^{144m}Pr decay modes (2012Wa38).

Gamma-ray energies and emission probabilities

Energies

The well-defined nuclear level energies of 2001So16 were used to calculate the gamma transition energies and their uncertainties. This approach was adopted because of the more wide-ranging origins of the level energies compared with the measured gamma-ray decay data. Nevertheless, the recommended gamma-ray energy standard proposed by 2000He14 has been adopted in the form of the 696.505(4)-keV gamma ray. Various nuclear-level half-lives were adopted and calculated from the half-life and mean lifetime measurements of 1976CoZX, 1994Ro13 and 1998Hi09.

Adopted energies, spins and parities for the nuclear levels of ¹⁴⁴Pr and ¹⁴⁴Nd (2001So16).

	Nuclear level number	Nuclear level energy (keV)	Spin and parity	
¹⁴⁴ Pr	0	0.0	0 –	17.29 (4) min
^{144m} Pr	1	59.03 ± 0.03	3 –	7.2 (2) min
¹⁴⁴ Nd	0	0.0	0 +	2.3 (3) × 10 ¹⁵ a
	1	696.561 ± 0.010	2 +	–
	2	1314.669 ± 0.013	4 +	7.4 (9) ps
	3	1510.871 ± 0.021	3 –	0.56 (7) ps
	4	1560.920 ± 0.013	2 +	–
	(5)	2072.91 ± 0.03	2 +	59 (10) fs
	(6)	2084.68 ± 0.03	0 +	0.12 (5) ps
	(7)	2185.75 ± 0.03	1 –	15 (2)
	(8)	2368.82 ± 0.04	2 +	39 (14) fs
	(9)	2582.32 ± 0.06	(3 +)	–
	(10)	2655.54 ± 0.03	1 +	9.9 (2) fs
	(11)	2675.61 ± 0.08	0 +	0.2 (1) ps
	(12)	2742.99 ± 0.07	0 +	64 (40) fs
(13)	2946.04 ± 0.10	(2, 3, 4) –	–	

Transition and gamma-ray energies of ^{144m}Pr (2001So16).

		Transition energy (keV)	Gamma-ray energy, E _γ (keV)	Decay mode
γ _{1,0}	(Pr)	59.03 (3)	59.03 ± 0.03	IT
γ _{2,1}	(Nd)	618.108 (16)	618.107 ± 0.016	β ⁻
γ _{1,0}	(Nd)	696.561 (10)	696.559 ± 0.010*	β ⁻
γ _{3,1}	(Nd)	814.310 (23)	814.308 ± 0.023	β ⁻
γ _(13,2)	(Nd)	1631.37 (10)	1631.36 ± 0.10	β ⁻
γ _(9,1)	(Nd)	1885.76 (6)	1885.75 ± 0.06	β ⁻

* Subsequently modified in line with the recommended energy standard of Helmer and van der Leun (2000He14).

Emission Probabilities

The gamma-ray emission probabilities were only partially obtained from the measurements of 1979Pr11 and 1985Da16. Thus, the emission probabilities of the 1631.36- and 1885.75-keV gamma rays were adopted in this manner, while the emission probability of the 618.107-keV gamma ray was arbitrarily adjusted from a measured value of 0.016 (7) to 0.030 (3) to achieve a population-depopulation balance of the 1314.67-keV nuclear level of ¹⁴⁴Nd. Equivalent emission probabilities for the 696.505- and 814.308-keV gamma rays were assigned to be 0.06 (2) and 0.02 (1), respectively, after due consideration of the β⁻-decay mode in which zero β⁻ feeding of the 1314.67-, 696.561-keV and ground states of ¹⁴⁴Nd were assumed on the basis of spin-parity considerations or balanced γ population-depopulation of these particular nuclear levels.

Studies of the β⁻ feeding of the 1510.87-keV nuclear level of ¹⁴⁴Nd have led to values of 0.05 (3) % (1970Fa03) and < 0.02 (1) % (1985Da16) – after careful consideration, a value of 0.02 (1) % has been recommended. An approximate normalization factor of (1.00 ± 0.10) was adopted in conjunction with these recommended gamma-ray emission probabilities. Under these somewhat unsatisfactory circumstances, a β⁻-decay mode with a branching fraction of 0.0006 (2) has been derived to give a matching IT branching fraction of 0.9994 (2). Finally, the absolute emission probability of the 59.03-keV M3 gamma transition was calculated from the IT branching fraction and the total internal conversion coefficient to be 0.0818 (12) %.

Multipolarities, Internal-Conversion and Internal-Pair Formation Coefficients

The nuclear level scheme specified by Sonzogni has been used to define the multipolarities of the gamma transitions on the basis of the known spins and parities (2001So16). Adopted multipolarities are 100%M3 for the 59.03-keV gamma ray constituting the IT decay mode, 100%E2 for the 618.107- and 696.505-keV gamma rays, and 100%E1 for the 814.308-keV gamma rays. Evidence for the mixing ratio and multipolarity of the 1885.75-keV (98.3%M1 + 1.7%E2) gamma transition arises from γ–γ correlation studies of 1998Hi09.

Measured and recommended gamma-ray energies.

E_γ (keV)										
1968Ra01	1968Sa05	1970An15	1970Fa03	1974Be09	1976Ra22	1979Gr01	1979Pr11	1985Da16	2000He14	Recommended[#]
–	–	59.0 (3)	–	–	58.5 (2)	–	–	–	–	59.03 ± 0.03
–	–	–	–	–	–	–	618.2 (5)	617.8 (2)	–	618.107 ± 0.016
696.3 (4)	696.48 (9)	–	696.43 (6)	696.49 (2)	696.4 (1)	696.510 (6)	696.50 (2)	696.50 (5)	696.505 (4)	696.505 ± 0.004 [*]
813.7 (6)	–	–	813.8 (3)	814.15 (15)	813.6 (2)	–	814.03 (10)	814.10 (10)	–	814.308 ± 0.023
–	–	–	–	–	–	–	–	1631.36 (10)	–	1631.36 ± 0.10
–	–	–	–	–	–	–	–	1885.3 (2)	–	1885.75 ± 0.06

[#] Determined from the nuclear level energies of 2001So16 unless stated otherwise.

^{*} Gamma-ray energy adopted from the recommended energy standards published by Helmer and van der Leun (2000He14).

Published relative gamma-ray emission probabilities.

E_γ (keV)	P_γ^{rel}			
	1970An15[†]	1976Ra22[‡]	1979Pr11[§]	1985Da16[§]
59.03 (3)	~ 0.01	0.003 (2)	–	–
618.107 (16)	–	–	0.015 (7)	0.016 (7)
696.505 (4)	–	–	–	–
814.308 (23)	–	–	–	–
1631.36 (10)	–	–	–	0.030 (3)
1885.75 (6)	–	–	–	0.010 (3)

[†] Emission probability expressed relative to P_γ(133.515 keV) of 100 % in ¹⁴⁴Ce β[–] decay.

[‡] Emission probability expressed relative to P_γ(133.515 keV) of 10.0 (1) % in ¹⁴⁴Ce β[–] decay.

[§] Emission probability expressed relative to P_γ(696.505 keV) of 100 % in ¹⁴⁴Pr β[–] decay.

Recommended absolute gamma-ray emission probabilities and transition probabilities.

E_γ (keV)	P_γ^{abs} (%)[*]	Transition probability (%)
59.03 ± 0.03	0.0818 ± 0.0012 [†]	99.94 ± 0.02
618.107 ± 0.016	0.030 ± 0.003 [‡]	0.030 ± 0.003
696.505 ± 0.004	0.06 ± 0.02 [#]	0.06 ± 0.02
814.308 ± 0.023	0.02 ± 0.01 [§]	0.02 ± 0.01
1631.36 ± 0.10	0.030 ± 0.003	0.030 ± 0.003
1885.75 ± 0.06	0.010 ± 0.003	0.010 ± 0.003

^{*} Absolute emission probabilities either adopted directly from 1979Pr11 and 1985Da16, or adjusted and derived on the basis of the proposed decay scheme, with an estimated normalization factor of (1.00 ± 0.10) applied to the relative gamma-ray emission probabilities as originally quoted.

[†] Determined from a calculated branching fraction for the IT-decay mode of 0.9994 (2) and total ICC of 1221 (18) for the 59.03-keV M3 gamma transition.

[‡] Arbitrarily assigned a value of 0.030 (3) to achieve a population-depopulation balance involving the 618.107- and 1631.36-keV gamma emissions identified with the 1314.67-keV nuclear level of ¹⁴⁴Nd to which β[–] feeding was assumed to be negligible.

[#] Arbitrarily assigned a value of 0.06 (2) to achieve a population-depopulation balance involving the populating 618.107-, 814.31- and 1885.75-keV gamma emissions and the depopulating 696.505-keV gamma emissions identified with the 696.56-keV nuclear level of ¹⁴⁴Nd to which β[–] feeding was assumed to be negligible.

[§] Arbitrarily assigned a value of 0.02 (1) based on β[–] feeding to the 1510.87-keV nuclear level of ¹⁴⁴Nd, as determined to be 0.05 (3) % by Fasching *et al.* (1970Fa03) and < 0.02 % by Dalmasso *et al.* (1985Da16).

Gamma-ray emissions: multipolarities, and theoretical internal-conversion (frozen orbital approximation) and internal-pair formation coefficients.

E_γ (keV)		Multipolarity	α_K	α_L	α_{L1}	α_{L2}	α_{L3}	α_{M+}	$\alpha_{ICtotal}$	α_{IPF}	α_{total}
59.03 (3)	(Pr)	M3	408 (6)	618 (9)	233 (4)	34.9 (5)	351 (5)	195	1221 (18)	–	1221 (18)
618.107 (16)	(Nd)	E2	0.005 68 (8)	0.000 869 (13)	0.000 673 (10)	0.000 123 1 (18)	0.000 072 7 (11)	0.000 241	0.006 79 (10)	–	0.006 79 (10)
696.505 (4)	(Nd)	E2	0.004 27 (6)	0.000 631 (9)	0.000 508 (8)	0.000 078 5 (11)	0.000 045 1 (7)	0.000 169	0.005 07 (7)	–	0.005 07 (7)
814.308 (23)	(Nd)	E1	0.001 198 (17)	0.000 152 8 (22)	0.000 140 7 (20)	0.000 005 46 (8)	0.000 006 56 (10)	0.000 040 2	0.001 391 (20)	–	0.001 391 (20)
1631.36 (10)	(Nd)	–	–	–	–	–	–	–	–	–	–
1885.75 (6)	(Nd)	(98.3%M1 + 1.7%E2) $\delta = 0.13$ (5) (1998Hi09)	0.000 686 (10)	0.000 087 8 (13)	0.000 084 3 (13)	0.000 002 59 (4)	0.000 000 926 (17)	0.000 023 3	0.000 797 (10)	0.000 255 (4)	0.001 052 (15)

The detailed conversion-electron measurements of Geiger *et al.* for the 59.03-keV gamma transition are particularly noteworthy (1960Ge05, 1969Ge01) – this M3 gamma transition constitutes the IT-decay mode of ^{144m}Pr.

Recommended internal conversion coefficients have been determined from the frozen orbital approximation of Kibédi *et al.* (2008Ki07), based on the theoretical model of Band *et al.* (2002Ba85, 2002Ra45). The (98.3%M1 + 1.7%E2) 1885.75-keV gamma transition also undergoes internal-pair formation, and the coefficient for this process has also been quantified from the tabulations of 2008Ki07.

β⁻ energies and emission probabilities

Energies

All β⁻ energies were derived from the structural details of the proposed decay scheme. The nuclear level energies of 2001So16 and evaluated Q-value of 3056.4 (24) keV (2012Wa38) were used to determine the recommended energies and uncertainties of the β⁻ emissions.

Emission Probabilities

Absolute emission probabilities were derived for the population-depopulation balances of the relative gamma-ray emission probabilities, their theoretical internal-conversion coefficients, and a normalization factor of (1.00 ± 0.10) for the gamma-ray emissions, as proposed above. β⁻ decay to the 1314.67-, 696.561-keV and ground-state levels of ¹⁴⁴Nd were assumed to be zero on the basis of the limited and rather questionable gamma-ray emission probabilities involved in the population-depopulation of these particular nuclear levels. Log ft values and average E_{β⁻} energies were determined by means of the LOGFT code.

Recommended energies and emission probabilities for the β⁻ decay of ^{144m}Pr.

	E _β (keV)*	Av. E _{β⁻} (keV)	P _β	¹⁴⁴ Pr	¹⁴⁴ Nd	transition type	log ft
β ⁻ _(1,13)	110.4 (24)	29.0 (7)	0.030 (3)	3 –	(2, 3, 4) –	(allowed)	4.65
β ⁻ _(1,9)	474.1 (24)	143.0 (8)	0.010 (3)	3 –	(3 +)	(first forbidden non-unique)	7.15
β ⁻ _{1,3}	1545.5 (24)	570.0 (11)	0.02 (1)	3 –	3 –	allowed	8.7
			Σ 0.06 (2)				

* Determined from the nuclear level energies of 2001So16 and Q-value of 3056.4 (24) keV (2012Wa38).

Energies and emission probabilities of noteworthy internal-conversion electrons.

	Energy (keV)	Electrons per 100 disint.
ec _{1,0T} (Pr)	17.04 – 59.03	99.86 (2)
ec _{1,0K} (Pr)	17.04 (3)	33.37 (1)
ec _{1,0L} (Pr)	52.20 – 53.07	50.54 (1)
ec _{1,0M} (Pr)	57.52 – 58.10	12.68 (1)
ec _{1,0N+} (Pr)	58.73 – 59.03	3.27 (1)

A consistent decay scheme was derived that contains three low-intensity β⁻ transitions and six gamma-ray emissions (one gamma transition constitutes the main IT-decay mode, and the other five gamma rays are associated with the small β⁻-decay mode).

Atomic Data

The X-ray and Auger-electron data have been calculated using the evaluated gamma-ray data, and atomic data from 1996Sc06, 1998ScZM and 1999ScZX. Both the X-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.02, 28 February 2012), as described in 2000Sc47. This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

K and L X-ray energies and emission probabilities of ^{144m}Pr.

			Energy (keV)	Photons per 100 disint.	Relative probability
XL	(Pr)		4.453 – 6.617	10.5 (5)	
		XL ₁	(Pr) 4.453	0.227 (7)	
		XL _α	(Pr) 5.013 – 5.033	5.74 (16)	
		XL _η	(Pr) 4.929	0.0498 (15)	
		XL _β	(Pr) 5.489 – 5.851	3.89 (7)	
		XL _γ	(Pr) 6.327 – 6.617	0.625 (13)	
XK _α		XK _{α2}	(Pr) 35.5506 (2)	8.66 (19)	54.8
		XK _{α1}	(Pr) 36.0267 (2)	15.8 (4)	100
XK' _{β1}		XK _{β3}	(Pr) 40.6533 (7))	
		XK _{β1} "	(Pr) 40.7487 (5)) 4.81 (12)	30.4
		XK _{β5}	(Pr) 41.050)	
XK' _{β2}		XK _{β2}	(Pr) 41.774 (2))	
		XK _{β4}	(Pr) 41.877) 1.23 (4)	7.78
		XKO _{2,3}	(Pr) 41.968)	
XL		(Nd)	4.633 – 6.901	0.000 067 (8)	
XK _α		XK _{α2}	(Nd) 36.8478 (3)	0.000 119 (23)	54.1
		XK _{α1}	(Nd) 37.3614 (2)	0.000 22 (5)	100
XK' _{β1}		XK _{β3}	(Nd) 42.1670 (4))	
		XK _{β1} "	(Nd) 42.2717 (3)) 0.000 067 (13)	30.5
		XK _{β5}	(Nd) 42.580)	
XK' _{β2}		XK _{β2}	(Nd) 43.335 (3))	
		XK _{β4}	(Nd) 43.451) 0.000 017 (4)	7.73
		XKO _{2,3}	(Nd) 43.548)	

Auger-electron energies and emission probabilities of ^{144m}Pr.

		Energy (keV)	Electrons per 100 disint.	Relative probability
e _{AK}	(Pr)		2.87 (15)	
	KLL	28.162 – 29.890	1.85 (10)	100
	KLX	33.576 – 36.004	0.91 (5)	49.2
	KXY	38.97 – 41.95	0.113 (7)	6.11
e _{AL}	(Pr)	2.90 – 4.91	69 (10)	3730
e _{AK}	(Nd)		0.000 038 (8)	
	KLL	29.154 – 30.978	0.000 024 (5)	100
	KLX	34.798 – 37.340	0.000 012 0 (24)	50
	KXY	40.42 – 43.53	0.000 001 5 (3)	6.25
e _{AL}	(Nd)	3.01 – 5.10	0.000 40 (5)	1667

Pr: $\omega_K = 0.914$ (4); $\omega_L = 0.132$ (5); $n_{KL} = 0.871$ (4), and Nd: $\omega_K = 0.918$ (4); $\omega_L = 0.140$ (6); $n_{KL} = 0.866$ (4) were taken from 1996Sc06.

Electron energies were determined from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

Data Consistency

An effective Q-value has been calculated from the Q_{IT} -value of 59.03 (3) keV and Q_{β^-} -value of 3056.4 (24) keV derived and adopted from the atomic mass evaluation of Wang *et al.* (2012Wa38), and their recommended branching fractions formulated during the evaluation of the decay scheme of ^{144m}Pr:

$$\begin{aligned} \text{effective Q-value} &= \sum (Q_i \times BF_i) = [59.03 (3) \times 0.9994 (2)] + [3056.4 (24) \times 0.0006 (2)] \\ &= 59.00 (12) + 1.8 (6) = 60.8 (6) \text{ keV} \end{aligned}$$

This value has been compared with the Q-value calculated by summing the contributions of the individual emissions to the ^{144m}Pr decay processes (i.e. β^- , conversion electrons, γ , etc.):

$$\text{calculated Q-value} = \sum (E_i \times P_i) = 60.9 (5) \text{ keV}$$

Percentage deviation from the effective Q-value is $-(0.2 \pm 1.3) \%$, which supports the derivation of a consistent decay scheme with a reasonably significant variant.

References

- 1960Ge05 J.S. Geiger, R.L. Graham, G.T. Ewan, Conversion Electron Spectrum from Ce¹⁴⁴ Decay, Nucl. Phys. 16 (1960) 1-26. [59.03-keV E_{ce} , P_{ce} , multipolarity]
- 1968Ra01 S. Raman, Decay of ¹⁴⁴Pr to Levels in ¹⁴⁴Nd, Nucl. Phys. A107 (1968) 402-410. [E_{γ}]
- 1968Sa05 A.R. Sayres, C.C. Trail, Precision Measurements of the Energies of the Gamma Rays from ¹⁴⁴Nd, Nucl. Phys. A113 (1968) 521-528. [E_{γ}]
- 1969Ge01 W. Gelletly, J.S Geiger, L-subshell Conversion Line Intensity Ratios for Pure M1(1 \leftrightarrow 0) Transitions, Nucl. Phys. A123 (1969) 369-385. [L-subshell ratios (59.03 keV)]
- 1970An15 A. Anttila, M. Piiparinen, The Levels of ¹⁴⁴Pr from the Decay of ¹⁴⁴Ce, Z. Phys. 237 (1970) 126-130. [59.03-keV E_{γ} , P_{γ} , α_K , α_L]
- 1970Fa03 J.L. Fasching, W.B. Walters, C.D. Coryell, Decay of 284-day ¹⁴⁴Ce, 17.3-min ¹⁴⁴Pr, and a New Isomer 7.2-min ^{144m}Pr, Phys. Rev. C1 (1970) 1126-1131. [Half-life, partial β^- branching fraction]
- 1974Be09 M. Behar, Z.W. Grabowski, S. Raman, Angular Correlation Studies in ¹⁴⁴Nd, Nucl. Phys. A219 (1974) 516-524. [E_{γ}]
- 1976Ch33 J.M. Chatterjee-Das, R.K. Chattopadhyay, P. Bhattacharya, B. Sethi, S.K. Mukherjee, The Decay Scheme of the 284.3 d ¹⁴⁴Ce and Energy Levels and Transitions in ¹⁴⁴Pr, Radiochem. Radioanal. Lett. 27 (1976) 119-128. [Half-life, E_{γ} , P_{γ}]
- 1976CoZX J.P. Collins, C.T. Wunker, R.L. Place, D.R. Ober, Bull. Amer. Phys. Soc. 21, No. 2 (1976) 149, D1. [Nuclear-level half-life (1314.67 keV)]
- 1976Ra22 B.V.N. Rao, G.N. Rao, Decay of ¹⁴⁴Ce to Levels in ¹⁴⁴Pr and ¹⁴⁴Nd, J. Phys. Soc. Japan 40 (1976) 1-5. [E_{γ} , P_{γ}]
- 1977La19 F.P. Larkins, Semiempirical Auger-electron Energies for Elements $10 \leq Z \leq 100$, At. Data Nucl. Data Tables 20 (1977) 311-387. [Auger-electron energies]
- 1979Gr01 R.C. Greenwood, R.G. Helmer, R.J. Gehrke, Precise γ -ray Energies for Calibration of Ge Semiconductor Spectrometers to 3.5 MeV, Nucl. Instrum. Methods 159 (1979) 465-482.[E_{γ}]
- 1979Pr11 M.S. Pravikoff, G. Barci-Funel, G. Ardisson, Sur la Décroissance de ^{144m+g}Pr et les Niveaux de ¹⁴⁴Nd, Radiochem. Radioanal. Lett. 40 (1979) 123-132. [E_{γ} , P_{γ}]

- 1985Da16 J. Dalmaso, H. Forest, G. Ardisson, ¹⁴⁴Nd Levels Fed in β decay of ¹⁴⁴Pr Isomers, Phys. Rev. 32 (1985) 1006-1009. [E _{γ} , P _{γ} , partial β^- branching fraction]
- 1994Ro13 S.L. Robinson, J. Jolie, H.G. Börner, P. Schillebeeckx, S. Ulbig, K.P. Lieb, E2 and E3 Transitions from Quadrupole-Octupole Coupled States in ¹⁴⁴Nd, Phys. Rev. Lett. 73 (1994) 412-415. [Nuclear-level lifetimes]
- 1996Sc06 E. Schönfeld, H. Janßen, Evaluation of Atomic Shell Data, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527-533. [ω_K , ω_L , K _{β} /K _{α} , K _{$\alpha 2$} /K _{$\alpha 1$} , KLX/KLL, KXY/KLL]
- 1998Hi09 S.F. Hicks, C.M. Davoren, W.M. Faulkner, J.R. Vanhoy, Structural Characteristics of ¹⁴⁴Nd Through γ -ray Spectroscopy Following Inelastic Neutron Scattering, Phys. Rev. 57 (1998) 2264-2280. [Nuclear structure, δ]
- 1998ScZM E. Schönfeld, G. Rodloff, Tables of the Energies of K-Auger Electrons for Elements with Atomic Numbers in the Range from Z = 11 to Z = 100, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX E. Schönfeld, G. Rodloff, Energies and Relative Emission Probabilities of K X-rays for Elements with Atomic Numbers in the Range from Z = 5 to Z = 100, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2000He14 R.G. Helmer, C. van der Leun, Recommended Standards for γ -ray Energy Calibration, Nucl. Instrum. Methods Phys. Res. A450 (2000) 35-70. [E _{γ}]
- 2000Sc47 E. Schönfeld, H. Janßen, Calculation of Emission Probabilities of X-rays and Auger Electrons Emitted in Radioactive Disintegration Processes, Appl. Radiat. Isot. 52 (2000) 595-600. [P_X, P_{Ae}]
- 2001So16 A.A. Sonzogni, Nuclear Data Sheets for A = 144, Nucl. Data Sheets 93 (2001) 599-762. [Nuclear levels]
- 2002Ba85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, Dirac-Fock Internal Conversion Coefficients, At. Data Nucl. Data Tables 81 (2002) 1-334. [ICC]
- 2002Ra45 S. Raman, C.W. Nestor, Jr., A. Ichihara, M.B. Trzhaskovskaya, How Good are the Internal Conversion Coefficients Now? Phys. Rev. C66 (2002) 044312, 1-23. [ICC]
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr., Evaluation of Theoretical Conversion Coefficients using BrIcc, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202-229. [ICC]
- 2012Wa38 M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer, The AME2012 Atomic Mass Evaluation, (I). Tables, Graphs and References, Chin. Phys. C36 (2012) 1603-2014. [Q-value]

¹⁴⁷Nd - Comments on evaluation of decay data by V. Chisté and M. M. Bé

This evaluation was completed in March 2011, including all publications by this date.

1 Decay Scheme

¹⁴⁷Nd disintegrates 100 % by beta minus emissions to excited levels of ¹⁴⁷Pm. If a transition to the ground state level exists, it is less than 0.15 % (1971Na11, 1966Be09).

A good agreement was found between the effective Q value (890 (60) keV) calculated from the decay scheme data and the adopted and recommended value from the mass adjustment of Audi (20012Au06), confirming the consistency of the adopted decay scheme.

2 Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2009AuZZ).

Experimental ¹⁴⁷Nd half-life values (in days) are given in Table 1:

Table 1: Experimental values of ¹⁴⁷Nd half-life.

Reference	Experimental value (d)	Comments
W. Bothe (1946Bo25)	11.1 (2)	
W. S. Emmerich (1951Em23)	11.1 (5)	
E. Kondaiah (1951Ko01)	11.6 (3)	
J. A. Marinsky (1951Ma**)	11.0 (3)	
W. C. Rutledge (1952Ru10)	11.9 (3)	Outlier
H. W. Wright (1957Wr37)	11.06 (4)	
R. G. Wille (1960Wi10)	11.5 (5)	
D. C. Hoffman (1963Ho15)	11.02 (5)	
S. Baba (1971Ba28)	10.98 (1)	
Recommended value	10.987 (11)	$\chi^2 = 1.4$

A weighted average has been calculated using LWEIGHT computer program (version 3). The Rutledge value (1952Ru10) has been shown to be outlier, based on the Chauvenet's criterion and thus was omitted in the final calculation. The largest contribution to the weighted average comes from the value of S. Baba (1971Ba28), with a statistical weight of 90 %.

The adopted value is the weighted average of 10.987 d with an external uncertainty of 0.011 d. The reduced- χ^2 value is 1.4.

2.1 β^- Transitions

The maximum energies of the β^- transitions in the decay of ¹⁴⁷Nd \rightarrow ¹⁴⁷Pm have been obtained from the Q value (2009AuZZ) and the level energies given in Table 2 from N. Nica (2009Ni02).

Table 2: ¹⁴⁷Pm levels populated in the decay of ¹⁴⁷Nd and the adopted β⁻ transition probabilities.

Level Number	Level energy, (keV) ^μ	Spin and Parity ^a	Half-life [*]	Adopted P _{β⁻} (%)
0	0	7/2 ⁺		0 (5)
1	91.1049 (20)	5/2 ⁺	2.50 (5) ns	81 (5)
2	408.54 (5)	9/2 ⁺		
3	410.512 (13)	3/2 ⁺	0.139 (14) ps	0.715 (34)
4	489.255 (16)	7/2 ⁺		0.781 (15)
5	531.012 (15)	5/2 ⁺	0.083 (15) ns	14.6 (9)
6	632.93 (7)	1/2 ⁺		0.0190 (27)
7	641.27 (8) ^μ			
8	649.03 (4)	11/2 ⁻	27 (3) ns	0.258 (19)
9	680.44 (4)	7/2 ⁺		0.0897 (28)
10	685.890 (15)	5/2 ⁺	0.25 (10) ns	2.184 (16)

* Given by N. Nica (2009Ni02).

^a Given by N. Coursol et al. (1987Table).

^μ Not used in this evaluation. No direct experimental evidences for this level, only speculative propositions: an unobservable weak β⁻ transition of ¹⁴⁷Nd decay (1997Sa53) or one 573-keV γ-ray transition (¹⁴⁸Nd(2p,γ)¹⁴⁷Pm), 1977Ko24) that populated it.

The adopted β⁻ transition probabilities and the associated uncertainties (Table 2) were deduced from the γ transition probability balance at each level of the decay scheme.

For the ground state level, the adopted β⁻ transition probability of 0 (5) % is in agreement with the experimental values of < 0.15 % (1966Be09, 1971Na11) and < 0.25 % (1962Sh02).

The values of log ft and average β⁻ energies have been calculated with the program LOGFT for all β⁻ transitions.

2.2 γ Transitions

The γ-ray transition probabilities were calculated using the γ-ray emission intensities and the relevant internal conversion coefficients (see 5.2 γ Emissions).

For all γ transitions, the internal conversion coefficients (ICC) and the associated uncertainties were interpolated from theoretical values of I. M. Band et al. (2002Ba85) using the BrIcc computer program (2008Ki07) for the “frozen orbital” approximation.

For multiplicities and mixing ratios, the evaluators used:

- 1) Multiplicities of γ-ray transitions listed in the Table 3 are from N. Nica (2009Ni02).

Table 3: Multiplicities of γ-ray transitions.

	Multipolarity	E _γ (keV)
¹⁴⁷ Pm	[M2]	31.3 (2)
	[E2]	53.1 (2), 541.83 (7)
	[E3]	36.75 (10)
	[M1,E2]	80.82 (27), 149.3 (2), 154.7 (2), 191.0 (3), 589.35 (4), 680.52 (15)
	M2	159.7 (2), 649.04 (8)
	E1	240.5 (2)
	E2	410.48 (3)
	E3	117.95 (8)

Multipolarity	E _γ (keV)
M1 + E2	196.64 (4), δ = 0.20 (8) (1977Al34) 271.87 (6), δ = 0.10 (3) (1979Se05) 408.52 (6), δ = 0.57 (3)

2) For other γ-ray transitions, the adopted mixing ratios (δ) are the weighted means of the δ values found in the literature (given by 1977Kr13) and shown in Table 4. A good agreement has been found between the experimental values of K and L internal coefficients and the calculated ones obtained by using the evaluated δ values and the BrIcc program.

Table 4: Experimental and recommended conversion coefficients and mixing ratios for the γ-ray transitions.

E _γ (keV)	δ experimental (mixing ratio)	α experimental	α theoretical (given by BrIcc)
91.105 (2)	0.10 (9) (1957Bi86)* 0.229 (143) (1961Ar09) 0.089 (11) (1961Ew02)* 0.13 (2) (1961We07) ^μ 0.18 (6) (1963Ph02) 0.28 (9) (1967Ba06) 0.67 (15) (1967Ba22) [@] 0.089 (5) (1969Ba32) 0.13 (2) (1970B112) 0.082 (10) (1977Al34)	α _K = 1.73 (6) (1997Sa53)	α _K = 1.714 (24)
Recommended value	 δ = 0.090 (5)	Reduced χ² = 1.7	
120.48 (5)	0.12 (3) (1963Ph02) 0.158 (15) (1970B112) 0.050 (21) (1977Al34)	α _K = 0.79 (3) α _L = 0.113 (6) (1997Sa53)	α _K = 0.772 (11) α _L = 0.112 (4)
Recommended value	 δ = 0.116 (42)	Reduced χ² = 8	
275.374 (15)	0.077 (14) (1960Bo17)* 0.13 (1) (1960Bo17)* 0.11 (11) (1961Ar09) 0.089 (11) (1961Ew02)* 0.14 (2) (1961We07) ^μ 0.05 (7) (1963Sp07)* 0.16 (4) (1966Go25)* 0.34 (12) (1967Ba06) [@] 0.112 (6) (1969Ba32) 0.58 (25) (1970B112) [@] 0.17 (4) (1974Bh02)* 0.16 (4) (1976Si08)* 0.14 (3) (1977Al34) 0.107 (7) (1979Se05)	α _K = 0.081 (3) α _L = 0.0109 (6) (1997Sa53)	α _K = 0.0792 (11) α _L = 0.01095 (16)
Recommended value	 δ = 0.112 (5)	Reduced χ² = 1.8	
319.411 (18)	0.40 (2) (1957Li40)* 0.38 (1) (1960Bo17)* 0.27 (1) (1960Ma03)* [@] 9.95 (11) (1961Ar09) [@] 0.36 (2) (1961We07) ^μ 0.38 (6) (1963Ph02)* 0.39 (4) (1963Sp07)* 0.34 (2) (1966Go25)* 0.31 (10) (1967Ba06) [@] 0.55 (5) (1969Ba32) [@] 0.35 (4) (1970B112)	α _K = 0.052 (2) α _L = 0.0079 (4) (1997Sa53)	α _K = 0.0514 (8) α _L = 0.00734 (11)

E_γ (keV)	$ \delta $ experimental (mixing ratio)	α experimental	α theoretical (given by BrIcc)
	0.011 (16) (1974Bh02) ^{*@} 0.38 (2) (1976Si08) [*] 0.41 (3) (1977Al34)		
Recommended value	$\delta = 0.378$ (9)	Reduced $\chi^2 = 0.9$	
398.155 (20)	0.31 (3) (1960Bo17) [*] 0.50 (7) (1966Go25) [*] 0.17 (7) (1970B112) 0.18 (6) (1974Bh02) [*] 0.30 (3) (1977Al34)	$\alpha_K = 0.0292$ (11) (1997Sa53)	$\alpha_K = 0.0293$ (5)
Recommended value	$\delta = 0.297$ (37)	Reduced $\chi^2 = 3.9$	
439.895 (22)	0.63 (5) (1960Bo17) [*] 0.70 (12) (1961Sa13) [*] 0.82 (65) (1961We07) ^μ 0.59 (7) (1963Sp07) [*] 0.56 (5) (1966Go25) [*] 0.62 (6) (1968Ra28) [*] 0.70 (9) (1969Ba32) 0.6 (1) (1970B112) 0.62 (7) (1974Bh02) [*] 0.59 (5) (1976Si08) [*] 0.77 (10) (1977Al34) [@]	$\alpha_K = 0.0212$ (9) $\alpha_L = 0.0028$ (2) (1997Sa53)	$\alpha_K = 0.0210$ (4) $\alpha_L = 0.00300$ (5)
Recommended value	$\delta = 0.609$ (21)	Reduced $\chi^2 = 0.4$	
489.24 (3)	0.79 (+23,-45) (1977Al34) 1.2 (+28,-8) (1961Sa13) [*]	$\alpha_K = 0.018$ (1) (1997Sa53)	$\alpha_K = 0.0152$ (16) $\alpha_K = 0.014$ (4)
Recommended value	$\delta = 0.79$ (+23,-45)		
531.016 (22)	0.75 (25) (1957Bi86) [*] 0.95 (30) (1961We07) ^μ 0.69 (32) (1969Ba32) 0.40 (3) (1977Al34)	$\alpha_K = 0.0133$ (3) (1997Sa53)	$\alpha_K = 0.01374$ (23)
Recommended value	$\delta = 0.407$ (35)	Reduced $\chi^2 = 1.4$	
594.80 (3)	0.66 (15) (1961Sa13) [*] 0.34 (16) (1963Sp07) [*] 0.66 (9) (1968Ra28) [*] 0.48 (8) (1974Bh02) [*]	$\alpha_K = 0.0071$ (5) (1997Sa53)	$\alpha_K = 0.00995$ (23)
Recommended value	$\delta = 0.55$ (6)	Reduced $\chi^2 = 1.5$	
685.90 (4)	0.95 (30) (1961We07) ^μ 0.87 (29) (1967Ba06) 1.05 (65) (1969Ba32) 0.95 (30) (1977Al34)	$\alpha_K = 0.0068$ (4) (1997Sa53)	$\alpha_K = 0.0063$ (4)
Recommended value	$\delta = 0.92$ (20)	Reduced $\chi^2 = 0.04$	

[@] Value has been shown to be outlier, based on the Chauvenet's criterion and thus was omitted in the final calculation.

^μ Not used: superseded by 1969Ba32.

^{*} Given by 1977Kr13.

3 Atomic Data

Atomic values, ω_K , ω_L , ω_M and n_{KL} are from Schönfeld and Janßen (1996Sc06).

The X-ray and Auger electron emission probabilities are calculated from the data set values using the program EMISSION.

4 Electrons Emissions

The conversion electron emission probabilities were deduced from the ICC values and the γ -ray emission intensities.

5 Photon emissions

5.1 K x-rays

The X-ray absolute intensities were deduced from the decay data using the EMISSION computer code and are compared in Table 5 with measured values found in the literature. The experimental and calculated values are in agreement within the uncertainty limits, supporting the overall consistency of the decay scheme data.

Table 5: Experimental and recommended (calculated) values of X-ray absolute intensities (%).

	J. Goswamy (1995Go**)	Recommended values
K α_2 x-ray	12.3 (5)	12.9 (9)
K α_1 x-ray	21.6 (9)	23.5 (15)
K β_1 x-ray	6.4 (3)	7.3 (5)
K β_2 x-ray	1.64 (6)	1.87 (13)

5.2 Gamma emissions

The energies of the γ -rays given in section 5.2 are from N. Nica (2009Ni02).

The experimental relative γ -ray emission intensities from ¹⁴⁷Nd have been obtained from all the available relative values. The normalization factor to convert relative γ -ray emission probabilities to absolute values is calculated with the formula:

$$\text{Normalization} = \frac{100 - P_{\beta^-}(\text{g.s.})}{\sum(1 + \alpha_T)P_{rel}} = 0.127 \text{ (9)}$$

where the sum is to be done over all the gamma transitions to the ground state, and $P_{\beta^-}(\text{g.s.}) = 0$ (5) %, deduced from the probability balance at the ground state (g.s.) level (see Table 2, **2.1 β^- Transitions**). From the theoretical α_T and the evaluated relative emission intensities (Table 6), the calculated normalization factor is 0.127 (9).

The experimental γ -ray emission probabilities relative to 100 for the 531-keV γ -ray are given in Table 6.

The adopted relative γ -ray intensity values are the weighted means calculated by the LWEIGHT program (version 3).

It should be noted that in the 50-150 keV region, only a few points of calibration exist to establish an efficiency curve for γ -ray detectors. Then the γ -ray intensity measurements in this region cannot lead to results with uncertainties better than 2-4 %. For this reason, the values of γ -ray intensities relative to the 91-keV γ -ray (Table 7) were omitted from averaging. The use of these values renormalized to the 531-keV γ -ray would introduce an increase of the uncertainties.

Our recommended relative and absolute γ -ray emission probabilities are given in Table 8.

Table 6: Experimental data sets of the relative γ -ray emission intensities (%).

Reference Energy (keV)	1966Ar16	1967Ca18	1967Do07	1967Hi04	1967Ja05	1974Ra30	1979Vo09	1997Sa53	1998Po**	Evaluated	Reduced χ^2
31.3 (2)											
36.75 (10)											
53.1 (2)											
80.82 (27)								0.0068 (9)		0.0068 (9)	
91.105 (2)	275 (50) ^μ	211 (42)	248 (13)	227 (35)	300 (100) ^μ	220 (14)	239 (5)	210.0 (43)		224 (14)	4.6
117.98 (5)								0.120 (10)		0.120 (10)	
120.48 (5)	2.6 (4)	2.5 (5)	2.1 (2)	3.3 (5)	8 (1) ^μ	3.3 (5)	3.05 (10)	2.810 (46)		2.84 (11)	3.5
149.3 (2)								0.0290 (30)		0.0290 (30)	
154.7 (2)					< 0.5			0.0310 (30)		0.0310 (30)	
159.7 (2)								0.0400 (30)		0.0400 (30)	
191.0 (3)								0.0280 (30)		0.0280 (30)	
196.64 (4)	1.3 (2)	1.30 (13)	1.0 (1) ^μ	1.5 (6)	2 (1) ^μ	1.4 (4)	1.38 (6)	1.420 (15)		1.416 (14)	0.3
230.77 (8)											
240.5 (2)								0.320 (20)		0.320 (20)	
271.87 (6)								0.099 (7)		0.099 (7)	
275.374 (15)	6.6 (7)	6.5 (6)	6.1 (5)	6.8 (14)	7 (2)	6.7 (7)	6.05 (10)	6.81 (8)	6 (1)	6.10 (9)	0.4
310							< 0.1				
319.411 (18)	15.0 (15)	14.2 (14)	15.8 (10)	16.3 (24)	15 (5)	16.5 (10)	15.0 (3)	15.91 (17)	15 (2)	15.68 (15)	1.2
398.155 (20)	7.0 (7)	6.4 (6)	6.7 (5)	6.8 (11)	5 (2) ^μ	6.5 (7)	6.59 (10)	6.82 (8)		6.73 (6)	0.6
408.52 (6)								0.140 (10)		0.140 (10)	
410.48 (3)	1.3 (1)	1.30 (13)	0.9 (2)	1.2 (5)	1.0 (6)	1.2 (3)	0.93 (5)	1.120 (13)		1.077 (47)	2.7
439.895 (22)	8.8 (9)	9.2 (9)	9.7 (6)	9.3 (11)	8 (2) ^μ	9.8 (2)	9.19 (14)	9.54 (10)		9.47 (9)	1.3
489.24 (3)	0.70 (8)	1.5 (8)	1.2 (3)	1.1 (5)	1.0 (5)	1.4 (4)	1.12 (6)	1.160 (14)		1.07 (9)	3.9
531.016 (22)	100	100	100	100	100	100	100	100	100	100	
541.83 (7)	0.20 (5)							0.140 (20)		0.148 (21)	1.2
589.35 (4)	0.40 (6) ^μ		0.26 (6)	0.31 (14)		0.29 (8)	0.30 (3)	0.290 (20)		0.291 (16)	0.09
594.80 (3)	2.2 (2)	2.20 (22)	1.6 (2) ^μ	1.9 (4)	2 (1)	2.0 (3)	1.92 (6)	2.120 (26)	2.0 (3)	2.089 (28)	1.5
649.04 (8)								0.0390 (30)		0.0390 (30)	
680.52 (15)			< 0.05	0.23 (16)		0.06	0.30 (5)	0.220 (10)		0.223 (11)	1.2
685.90 (4)	7.0 (7)	6.6 (7)	5.0 (4) ^μ	5.9 (10)	6 (1)	6.7 (6)	6.1 (2)	6.63 (7)		6.57 (7)	1.2

μ: the experimental value has been shown to be an outlier value by the Lweight program.

Table 7: Omitted experimental data sets of the relative γ -ray emission intensities (%).

Reference Energy (keV)	1963Ph02	1967Ba21	1971Si20	1974HeYW	1995Go**	2010Gh**
31.3 (2)						
36.75 (10)	106 (16)					
53.1 (2)	7.5 (10)					
80.82 (27)	8 (1)					
91.105 (2)	100	100	100	100	100	100
117.98 (5)						
120.48 (5)	2.0 (2)	1.4 (1)	1.42 (18)	1.42 (15)	1.64 (5)	1.540 (3)
149.3 (2)						
154.7 (2)					0.0250 (10)	< 0.034
159.7 (2)	1.5 (2)					
191.0 (3)						
196.64 (4)	1.6 (2)	0.72 (7)	0.73 (12)	0.73 (6)	0.610 (12)	1.012 (27)
230.77 (8)						
240.5 (2)						
271.87 (6)						
275.374 (15)	2.5 (2)	3.0 (2)	3.05 (20)	2.87 (18)	2.720 (40)	3.320 (5)
310	< 2	< 0.2	0.13 (5)			
319.411 (18)	7.0 (6)	6.8 (5)	7.60 (70)	7.0 (4)	6.80 (12)	8.010 (12)
398.155 (20)	< 2.5	3.1 (3)	3.35 (25)	3.12 (30)	3.050 (43)	3.680 (7)
408.52 (6)						
410.48 (3)	3.7 (3)	0.8 (1)	0.55 (15)	0.50 (3)	0.360 (20)	0.790 (2)
439.895 (22)	4.5 (4)	4.2 (3)	5.1 (3)	4.3 (3)	4.20 (9)	5.200 (7)
489.24 (3)	< 0.2	0.7 (1)	0.6 (1)	0.55 (3)	0.49 (11)	0.530 (7)
531.016 (22)	58 (2)	47 (3)	53.5 (15)	46.9 (26)	45.9 (10)	47.20 (24)
541.83 (7)						
589.35 (4)		0.13 (2)	0.20 (2)	0.164 (16)	0.1580 (25)	0.224 (6)
594.80 (3)	1.6 (1)	0.9 (1)	1.1 (1)	0.95 (6)	0.850 (13)	0.586 (14)
649.04 (8)						
680.52 (15)			0.17 (8)	0.070 (15)	0.0560 (31)	0.072 (5)
685.90 (4)	4.7 (1)	3.3 (2)	3.5 (2)	2.91 (18)	2.850 (41)	2.430 (5)

Table 8: Recommended relative and absolute γ -ray intensities (%).

E_{γ} (keV)	Relative γ -ray intensity (%)	Absolute γ -ray intensity (%)
31.3 (2)		
36.75 (10)		
53.1 (2)		
80.82 (27)	0.006 8 (9)	0.000 86 (11)
91.105 (2)	224 (14)	28.4 (18)
117.98 (5)	0.120 (10)	0.015 2 (13)
120.48 (5)	2.84 (11)	0.361 (14)
149.3 (2)	0.029 0 (30)	0.003 68 (38)
154.7 (2)	0.031 0 (30)	0.003 94 (38)
159.7 (2)	0.040 0 (30)	0.005 08 (38)
191.0 (3)	0.028 0 (30)	0.003 56 (38)
196.64 (4)	1.416 (14)	0.179 8 (18)
230.77 (8)		
240.5 (2)	0.320 (20)	0.040 6 (25)
271.87 (6)	0.099 (7)	0.012 6 (9)
275.374 (15)	6.10 (9)	0.775 (11)
310		
319.411 (18)	15.68 (15)	1.991 (19)
398.155 (20)	6.73 (6)	0.855 (8)
408.52 (6)	0.140 (10)	0.017 8 (13)
410.48 (3)	1.077 (47)	0.137 (6)
439.895 (22)	9.47 (9)	1.203 (11)
489.24 (3)	1.07 (9)	0.136 (11)
531.016 (22)	100	12.7 (9)
541.83 (7)	0.148 (21)	0.018 8 (27)
589.35 (4)	0.291 (16)	0.037 0 (20)
594.80 (3)	2.089 (28)	0.265 3 (36)
649.04 (8)	0.039 0 (30)	0.004 95 (38)
680.52 (15)	0.223 (11)	0.028 3 (14)
685.90 (4)	6.57 (7)	0.834 (9)

6 References

- 1946Bo25 W. Bothe, Z. Naturforsch. 1(1946)179 [Half-life].
1951Em23 W. S. Emmerich, J. D. Kurbatov, Phys. Rev. 83(1951)40 [Half-life].
1951Ko01 E. Kondaiah, Phys. Rev. 81(1951)1056 [Half-life].
1951Ma** J. A. Marinsky, National Nucl. Energ. Series 9(1951)1229 [Half-life].
1952Ru10 W. C. Rutledge, J. M. Cork, S. B. Burson, Phys. Rev. 86(1952)775 [Half-life].
1957Bi86 G. R. Bishop, M. A. Grace, C. E. Jonhson, H. R. Lemmer, J. Perez y Jorba, Phil. Mag 2(1957)534 [Mixing ratio - angular distribution].
1957Li40 T. Lindqvist, E. Karlsson, Ark. Fysik 12(1957)519 [Mixing ratio angular distribution].
1957Wr37 H. W. Wright, E. I. Wyatt, S. A. Reynolds, W. S. Lyon, T. H. Handley, Nucl. Sci. Eng. 2(1957)427 [Half-life].
1960Bo17 E. Bodenstedt, H. J. Korner, F. Frisius, D. Hovestadt, E. Gerdau, Z. Phys. 160(1960)33 [Mixing ratio - angular distribution].
1960Ma03 G. Manning, J. D. Rogers, Nucl. Phys. 15(1960)166 [Mixing ratio - angular distribution].
1960Wi10 R. G. Wille, R. W. Fink, Phys. Rev. 118(1960)242 [Half-life].
1961Ar09 A. P. Arya, Phys. Rev. 122(1961)1226 [δ].
1961Ew02 G. T. Ewan, R. L. Graham, J. S. Geiger, Bull. Am. Phys. Soc. 6(1961)238 [Mixing ratio - angular distribution].

- 1961Sa13 B. Saraf, R. Jambunathan, M. R. Gunye, Phys. Rev. 124(1961)178 [Mixing ratio - angular distribution].
- 1961We07 G. A. Westenbarger, D. A. Shirley, Phys. Rev. 123(1961)1812 [δ].
- 1962Sh08 R. P. Sharma, S. H. Devare, B. Saraf, Phys. Rev. 125(1962)2071 [P_β].
- 1963Ho15 D. C. Hoffman, J. Inorg. Nucl. Chem. 25(1963)1196 [Half-life].
- 1963Ph02 C. Philis, Thesis, Univ. Paris, CEA-2355 (1963) [I_γ , δ].
- 1963Sp07 E. Spring, Phys. Lett. 7(1963)218 [Mixing ratio - angular distribution].
- 1966Ar16 E. A. Arutyunyan, J. Vrzal, B. S. Dzhelepov, J. Liptak, Ya. Urbanets, Yu. V. Khol'nov, Bull. Acad. Sci. USSR, Phys. Ser. 30(1967)1317 [I_γ].
- 1966Be09 H. Beekhuis, P. Boskma, J. Van Klinken, H. de Waard, Nucl. Phys. 79(1966)220 [P_β].
- 1966Go25 K. P. Gopinathan, AEET-267(1966)44 [Mixing ratio - angular distribution].
- 1967Ba06 E. Bashandy, A. Abd. El-Haliem, Z. Naturforsch. 22a(1967)154 [δ].
- 1967Ba21 A. Bäcklin, S. G. Malmskog, Ark. Fysik 34(1967)459 [I_γ].
- 1967Ba22 A. Bäcklin, G. Malmskog, Ark. Fysik 34(1967)531 [δ].
- 1967Ca18 M. J. Canty, R. D. Connor, Nucl. Phys. A104(1967)35 [I_γ].
- 1967Do07 P. W. Dougan, B. Earlandsson, Z. Phys. 207(1967)105 [I_γ].
- 1967Hi04 J. C. Hill, M. L. Wiedenbeck, Nucl. Phys. A98(1967)599 [I_γ].
- 1967Ja05 E. Jacobs, K. Heyde, M. Dorinkens, J. Demuynck, L. Dorinkens-Vanpraet, Nucl. Phys. A99(1967)411 [I_γ].
- 1968Ra28 M. S. Rajput, M. L. Sehgal, Indian J. Phys. 42(1968)393 [Mixing ratio - angular distribution].
- 1969Ba32 P. H. Barrett, D. A. Shirley, Phys. Rev. 184(1969)1181 [δ].
- 1970B112 N. Blaskovich Jr., A. P. Arya, Phys. Rev. C2(1970)1881 [δ].
- 1971Ba28 S. Baba, H. Baba, H. Natsume, J. Inorg. Nucl. Chem. 33(1971)589 [Half-life].
- 1971Na11 T. Nagarajan, M. Ravindranath, K. V. Reddy, Nuovo Cimento 3A(1971)689 [P_β].
- 1971Si20 H. Singh, B. Sethi, S. K. Mukerjee, Nucl. Phys. A174(1971) 437 [I_γ]
- 1974Bh02 S. S. Bhati, N. Singh, P. C. Mangal, P. N. Trehan, J. Phys. Soc. Jpn. 36(1974)326 [Mixing ratio - angular distribution].
- 1974HeYW R. L. Heath, ANCR - 1000-2(1974) [I_γ].
- 1974Ra30 C. Rangacharyulu, S. N. Chaturvedi, G. K. Mehta, N. Nath, Aust. J. Phys. 27(1974)869 [I_γ].
- 1976Si08 B. K. Sinha, S. Sen, R. Bhattacharya, J. Phys. (London) G2(1976)159 [Mixing ratio - angular distribution].
- 1977Al34 T. Al-Janabi, W. D. Hamilton, D. D. Warner, J. Phys. (London) G3(1977)1415 [δ].
- 1977Ko24 M. Kortelahti, A. Pakkanen, M. Piiparinen, T. Kompa, R. Komu, Nucl. Phys. A288(1977)365 [Level energies].
- 1977Kr13 K. S. Krane, At. Data Nucl. Data Tables 19(1977)363 [δ].
- 1979Se05 T. Seo, T. Hayashi, Y. Miyatake, K. Aoki, Nucl. Phys. A321(1979)341 [δ].
- 1979Vo09 N. A. Voinova, A. A. Rodionov, Yu. V. Sergeenkov, P. A. Sushkov, M. A. Elizbarashvili, Bull. Acad. Sci. USSR, Phys. Ser. 43(1979)70 [I_γ].
- 1987Table F. Lagoutine, N. Coursol, J. Legrand, Table de Radionucléides. ISBN-2-7272-0078-1. LMRI, 1982-1987, BP 52, 91191 Gif-sur-Yvette Cedex, France [Spin and parity].
- 1995Go** J. Goswamy, B. Chand, D. Mehta, N. Singh, P. N. Trehan, Radiat. Phys. Chem. 45(1995)733 [I_γ , X-rays].
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
- 1997Sa53 M. Sainath, K. Venkataramaniah, P. C. Sood, Phys. Rev. C56(1997)2468 [I_γ , α].
- 1999Po** Yu. S. Popov, N. Yu. Nezhgorov, G. A. Timofeev, Radiochemistry 41(1999)25 [I_γ].
- 2002Ba85 I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr., P. O. Tikkanen, S. Raman, Atomic Data Nucl. Data Tables 81(2002)1 [Theoretical ICC].
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr., Nucl. Instrum. Meth. Phys. Res. A589(2008)202 [Theoretical ICC].
- 2009Ni02 N. Nica, Nucl. Data Sheets 110(2009)749 [E_γ , E_{level}].
- 2012Au06 M. Wang, G. Audi, *et al.* Chinese Physics C36 (2012) 1603 [$Q\beta^-$]
- 2010Gh** S. S. Ghumman, C. Singh, S. Singh, Asian J. Chem. 22(2010)3021 [I_γ].

¹⁴⁷Pm - Comments on evaluation of decay data by V. Chisté and M. M. Bé

This evaluation was completed in May 2011, including all publications by this date.

1 Decay Scheme

¹⁴⁷Pm disintegrates 100 % by beta minus emissions to the ¹⁴⁷Sm ground state mainly.

A good agreement was found between the effective Q value (224.5 (4) keV) calculated from the decay scheme data and the adopted and recommended value from the mass adjustment of Audi (2003Au03).

2 Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2003Au03).

Experimental ¹⁴⁷Pm half-life values (in years) are given in Table 1:

Table 1: Experimental values of ¹⁴⁷Pm half-life.

Reference	Experimental value (a)	Comments
E. A. Melaika (1955Me52)	2.52 (8)	Outlier.
R. P. Schuman (1956Sc87)	2.66 (2)	Outlier.
W. F. Merritt (1957Me47)	2.64 (2)	
J. P. Cali (1959Ca**)	2.7 (1)	Outlier.
E. I. Wyatt (1961Wy01)	2.50 (3)	Superseded by 1968Re04.
F. P. Roberts (1963Ro20)	2.67 (6)	Superseded by 1965Wh04.
S. C. Anspach (1965An07)	2.618 (7)	
J. F. Eichelberger (1965Ei04)	2.6226 (20)	Superseded by 1967Jo07.
K. F. Flynn (1965Fl02)	2.60 (2)	
E. J. Wheelwright (1965Wh04)	2.620 (5)	
K. C. Jordan (1967Jo07)	2.6234 (4)	
S. A. Reynolds (1968Re04)	2.62 (1)	
Recommended value	2.6234 (4)	$\chi^2 = 0.64$

A weighted average has been calculated using LWEIGHT computer program (version 3). The Melaika (1955Me52), Schuman (1956Sc87) and Cali (1959Ca**) values have been shown to be outlier, based on the Chauvenet's criterion and thus were omitted in the final calculation. The largest contribution to the weighted average comes from the value of K. C. Jordan (1967Jo07), with a statistical weight of 98 %.

The adopted value is the weighted average of 2.6234 a with an internal uncertainty of 0.0004 a. The reduced- χ^2 value is 0.64.

For ¹⁴⁷Sm, the experimental half-life values (in years) are given in Table 2:

Table 2: Experimental values of ¹⁴⁷Sm half-life.

Reference	Experimental value (10 ¹¹ a)	Comments
W. F. Libby (1934Li03)	0.92 (6)	Corrected for (Sm nat./Sm-147) = 0.1498 by R. D. MacFarlane (1961Ma05).
R. Hosemann (1936Ho**))	1.5 (1)	Corrected for (Sm nat./Sm-147) = 0.1498 by R. D. MacFarlane (1961Ma05).
P. Cuer (1946Cu**))	1.3 (1)	
E. Picciotto (1949Pi**))	1.00 (5)	Corrected for (Sm nat./Sm-147) = 0.1498 by R. D. MacFarlane (1961Ma05).
G. Beard (1954Be69)	1.25 (6)	Superseded by 1958Be78.
G. E. Leslie (1956Le55)	1.15 (6)	
G. Beard (1958Be78)	1.06 (4)	Corrected for wrong Sm content by P. M. Wright (1961Wr02). Original value: 1.28 (4). Superseded by 1987Al28.
M. Karras (1960Ka**))	1.14 (5)	Superseded by 1960Ka23.
M. Karras (1960Ka23)	1.17 (5)	
R. D. MacFarlane (1961Ma05)	1.15 (5)	Superseded by 1970Gu14.
P. M. Wright (1961Wr02)	1.05 (2)	
D. Donhoffer (1964Do01)	1.04 (3)	
K. Valli (1965Va16)	1.08 (2)	
M. C. Gupta (1970Gu14)	1.06 (2)	
B. Al-Batrina (1987Al28)	1.05 (4)	
J. B. Martins (1992Ma56)	1.06 (4)	Corrected for wrong Sm content by F. Begemann (2001Be81). Original value: 1.23 (4).
N. Kinoshita (2003Ki26)	1.17 (2)	Questioned by 2009Ko15.
K. Kossert (2009Ko15)	1.070 (9)	
Recommended value	1.079 (12)	$\chi^2 = 3.9$

The first 3 values (1934Li03, 1936Ho** and 1946Cu**) have been shown outliers, based on the Chauvenet's criterion and thus were omitted in the final calculation. With the eleven remaining values (1949Pi**, 1956Le55, 1960Ka23, 1961Wr02, 1964Do01, 1965Va16, 1970Gu14, 1987Al28, 1992Ma56, 2003Ki26 and 2009Ko15), a weighted average has been calculated using LWEIGHT computer program (version 3). The largest contribution to the weighted average comes from the value of K. Kossert (2009Ko15), with a statistical weight of 46 %.

The adopted value is the weighted average of $1.079 \cdot 10^{11}$ a with an external uncertainty of $0.012 \cdot 10^{11}$ a. The reduced- χ^2 value is 3.9.

2.1 β^- Transitions

The maximum energies of the β^- transitions in the decay of ¹⁴⁷Pm \rightarrow ¹⁴⁷Sm have been obtained from the Q⁻ value (2003Au03) and the level energies from N. Nica (2009Ni02) (Table 3).

Table 3: ¹⁴⁷Sm levels populated in the decay of ¹⁴⁷Pm and adopted β⁻ transition probabilities.

Level Number	Level energy, (keV) ^a	Spin and parity	Half-life [*]	Adopted P _{β⁻} (%)
0	0	7/2 ⁻	1.060 (11)·10 ⁻¹¹ s	99.994 56 (13)
1	121.223 (12)	5/2 ⁻	0.798 (17) ns	0.005 42 (13)
2	197.298 (11)	3/2 ⁻	1.25 (3) ns	0.000 000 40 (7)

^{*} Given by N. Nica (2009Ni02),

^a from least-squares fit to E_γ's.

The adopted β⁻ transition probabilities and the associated uncertainties (Table 3) were deduced from the γ transition probability balance at each level of the decay scheme.

The values of log ft and average β⁻ energies have been calculated with the program LOGFT for the unique 1st and 1st forbidden β⁻ transitions.

2.2 γ Transitions

The γ-ray transition probabilities were calculated using the γ-ray emission intensities and the relevant internal conversion coefficients (see 4.2 γ Emissions).

For all γ transitions, the internal conversion coefficients (ICC) and the associated uncertainties were interpolated from theoretical values of I. M. Band et al. (2002Ba85) using the BrIcc computer program (2008Ki07) for the “frozen orbital” approximation.

For multipolarity and mixing ratio of the γ-ray transitions, the evaluators used:

1) The multiplicities of the 76- and 197-keV γ-ray transitions are from N. Nica (2009Ni02):

76-keV γ-ray: M1 + E2, δ = 0.655 (34);

197-keV γ-ray: E2.

2) For the 121-keV γ-ray transition, the adopted mixing ratio (δ) is the weighted mean of the δ values found in the literature and shown in the Table 4.

Table 4: Experimental and recommended mixing ratio and ICC.

E _γ (keV)	δ experimental (mixing ratio)	Comments
121.220 (7)	0.25 (21) (1958An36) -0.06 (2) (1962Al19) ^a -0.33 (4) (1962Sc09) 0.34 (3) (1966Av02) -0.35 (4) (1966Go26) [‡] -0.38 (3) (1970Va38) -0.40 (+26,-15) (1971Be53) -0.278 (20) (1989Ad10)	Calculated [*] from K/L = 6.2 (6). Calculated by 1977Kr13 (γγ(θ)). Calculated [*] from L ₁ :L ₂ :L ₃ = 1.0 (2): 0.24 (4): 0.16 (2). Calculated [*] from K: L ₁₊₂ :L ₃ = 450 (40): 73 (7): 10 (1). Calculated by 1977Kr13 (γγ(θ)). Superseded by 1970Va38. Calculated by 1977Kr13 (γγ(θ)). Calculated by 1977Kr13 (γγ(θ)).
Recommended value	- 0.317 (19), χ ² = 1.7	α _K (BRICC – 121-keV γ-ray) = 0.815 (12).

^a Outlier value, based on the Chauvenet's criterion and thus was omitted in the final calculation.

[‡] Superseded by 1970Va38.

^{*} Using BrIccMixing program, version 2.2a (same package of BrIcc computer program).

3 Atomic Data

Atomic values, ω_K , ω_L , n_{KL} and the X-ray relative probabilities are from Schönfeld and Janßen (1996Sc06).

4 Photon Emissions

4.1 X-ray Emissions

The X-ray absolute intensities were deduced from the decay data using the EMISSION computer code.

4.2 Gamma emissions

The energies of the γ -rays given in Table 5 were derived from the ¹⁴⁷Sm adopted levels (Table 2).

The experimental absolute values of the 121-keV γ -ray emission intensities in the decay of ¹⁴⁷Pm are given in the table 5.

Table 5: Absolute experimental γ -ray emission intensities for the 121-keV transition.

Reference	Absolute γ -ray intensity (10^{-3} %)	Comments
H. Langevin-Joliot (1956La17)	3.0 (5)	
N. Starfelt (1957St05)	3.4 (5)	
R. S. Mowatt (1970Mo02)	2.73 (18)	
D. McConnon (1971Mc09)	2.93 (14)	
H. H. Hansen (1973HaHY)	3.0 (3)	
U. Schötzig (1990Sc08)	2.65 (6)	
Recommended value	2.72 (6)	$\chi^2 = 1.33$

The adopted value is the weighted average of $2.72 \cdot 10^{-3}$ % with an external uncertainty of $0.06 \cdot 10^{-3}$ %. The reduced- χ^2 value is 1.33.

For the 197-keV γ -ray emission, the adopted value of the γ -ray relative intensity ($1.2 (2) \cdot 10^{-4}$) comes from the unique measurement found in the literature given by P. H. Barrett (1969Ba33).

Our recommended γ -ray emission probabilities are given in Table 6.

Table 6: Recommended relative and absolute γ -ray intensities (%).

E_γ (keV)	Relative γ -ray intensity (%)	Absolute γ -ray intensity (%)
(76.073 (10))^a	$4.1 (7) \cdot 10^{-6}$	$1.1 (2) \cdot 10^{-8}$
121.220 (17)	100	$2.72 (6) \cdot 10^{-3}$
197.299 (12)	$1.2 (2) \cdot 10^{-4}$	$3.3 (5) \cdot 10^{-7}$

^a not observed in this decay scheme.

The 76-keV γ -ray transition has been observed in the ¹⁴⁷Eu electron capture decay, but not in the ¹⁴⁷Pm β^- decay. From the ¹⁴⁷Eu electron capture decay (1989Ad10):

$I_\gamma(76 \text{ keV})/I_\gamma(197 \text{ keV}) = 0.0344 (11)$ and

$I_\gamma(197 \text{ keV}) = 1.2 (2) \cdot 10^{-4}$. Then $I_\gamma(76 \text{ keV}) = 4.1 (7) \cdot 10^{-6}$.

This very weak transition was included in the decay scheme.

5 References

- 1934Li03 W. F. Libby, Phys. Rev. 46(1934)196 [¹⁴⁷Sm half-life].
- 1936Ho** R. Hosemann, Z. Physik 99(1936)405 [¹⁴⁷Sm half-life].
- 1946Cu** P. Cuer, C. M. G. Lattes, Nature 158(1946)197 [¹⁴⁷Sm half-life].
- 1949Pi** E. Picciotto, Compt. Rend. (Paris) 229(1949)117 [¹⁴⁷Sm half-life].
- 1954Be69 G. B. Beard, M. L. Wiedenbeck, Phys. Rev. 95(1954)1245 [¹⁴⁷Sm half-life].
- 1955Me52 E. A. Melaika, M. J. Parker, J. A. Petruska, R. H. Tomlinson, Can. J. Chem. 33(1955)830 [Half-life].
- 1956La17 H. Langevin-Joliot, M. Lederer, J. Phys. Radium 17(1956)497 [P_γ].
- 1956Le55 G. E. Leslie, Nucl. Sci. Abstr. 10(1956)1099 [¹⁴⁷Sm half-life].
- 1956Sc87 R. P. Schuman, M. E. Jones, A. C. Mewherter, J. Inorg. Nucl. Chem. 3(1956)160 [Half-life].
- 1957Me47 W. F. Merritt, P. J. Champion, R. C. Hawkings, Can. J. Phys. 35(1957)16 [Half-life].
- 1957St05 N. Starfelt, J. Cederlund, Phys. Rev. 105(1957)241 [P_γ].
- 1958An36 N. M. Anton'eva, A. A. Bashilov, B. S. Dzheleпов, B. K. Preobrashenskii, Bull. Acad. Sci. USSR, Phys. Ser. 22(1959)899 [121-keV mixing ratio].
- 1958Be78 G. B. Beard, W. H. Kelly, Nucl. Phys. 8(1958)207 [¹⁴⁷Sm half-life].
- 1959Ca** J. P. Cali, L. F. Lowe, Nucleonics 17(1959)86 [Half-life].
- 1960Ka** M. Karras, N. Nurmi, Nature 185(1960)601 [¹⁴⁷Sm half-life].
- 1960Ka23 M. Karras, Ann. Acad. Sci. Fenn. A VI(1960)65 [¹⁴⁷Sm half-life].
- 1961Ma05 R. D. MacFarlane, T. P. Kohman, Phys. Rev. 121(1961)1758 [¹⁴⁷Sm half-life].
- 1961Wy01 E. I. Wyatt, S. A. Reynolds, T. H. Handley, W. S. Lyon, H. A. Parker, Nucl. Sci. Eng. 11(1961)74 [Half-life].
- 1961Wr02 P. M. Wright, E. P. Steinberg, L. E. Glendenin, Phys. Rev. 123(1961)205 [¹⁴⁷Sm half-life].
- 1962Al19 Y. A. Aleksandrov, B. Bemmer, Bull. Acad. Sci. USSR, Phys. Ser. 26(1963)1171 [121-keV mixing ratio].
- 1962Sc09 C. F. Schwerdtfeger, H. J. Prask, J. W. Mihelich, Nucl. Phys. 35(1962)168 [121-keV mixing ratio].
- 1963Ro20 F. P. Roberts, E. J. Wheelwright, W. Y. Matsumoto, HW-77296 (1963)[Half-life].
- 1964Do01 D. Donhoffer, Nucl. Phys. 50(1964)489 [¹⁴⁷Sm half-life].
- 1965An07 S. C. Anspach, L. M. Cavallo, S. B. Garfinkel, J. M. R. Hutchinson, C. N. Smith, NP – 15663 (1965) [Half-life].
- 1965Ei04 J. F. Eichelberger, Report MLM – 1221(1965)5 [Half-life].
- 1965Fl02 K. F. Flynn, L. E. Glendenin, E. P. Steinberg, Nucl. Sci. Eng. 22(1965)416 [Half-life].
- 1965Va16 K. Valli, Ann. Acad. Sci. Fenn. A VI(1965)177 [¹⁴⁷Sm half-life].
- 1965Wh04 E. J. Wheelwright, D. M. Fleming, F. P. Roberts, J. Phys. Chem. 69(1965)1220 [Half-life].
- 1966Av02 M. P. Avotina, E. P. Grigorev, A. Z. Zolotavin, V. O. Sergeev, V. E. Tre-Nersesyants, J. Vrzal, N. A. Lebedev, J. Liptak, Y. Urbanets, Bull. Acad. Sci. USSR, Phys. Ser. 30(1967)1350 [121-keV mixing ratio].
- 1966Go26 T. Goworek, J. Wawryszczuk, Acta Phys. Pol. 29(1966)655 [121-keV mixing ratio].
- 1967Jo07 K. C. Jordan, MLM – 1399(1967)16 [Half-life].

- 1968Re04 S. A. Reynolds, J. F. Emery, E. I. Wyatt, Nucl. Sci. Eng. 32(1968)46 [Half-life].
- 1969Ba33 P. H. Barrett, D. A. Shirley, Phys. Rev. 184(1969)1185 [197-keV I_γ].
- 1970Gu14 M. C. Gupta, R. D. MacFarlane, J. Inorg. Nucl. Chem. 32(1970)3425 [¹⁴⁷Sm half-life].
- 1970Mo02 R. S. Mowatt, J. S. Merritt, Can. J. Phys. 48(1970)453 [P_γ].
- 1970Va38 Y. Vavryshchuk, T. Goworek, K. Klishchevska, Bull. Acad. Sci. USSR, Phys. Ser. 34(1971)1922 [121-keV mixing ratio].
- 1971Be53 R. A. Belt, E. G. Funk, J. W. Mihelich, Nucl. Phys. A175(1971)129 [121-keV mixing ratio].
- 1971Mc09 D. McConnon, Int. J. Appl. Radiat. Isotop. 22(1971)253[P_γ].
- 1973HaXY H. H. Hansen, Proc. Int. Conf. Inner Shell Ionization Phenomena and Future Applications, Atlanta 3(1972)2157 [P_γ].
- 1977Kr13 K. S. Krane, At. Data and Nucl. Data Tables 19(1977)363 [mixing ratio].
- 1987Al28 B. Al-Bataina, J. Jänecke, Radiochimica Acta 42(1987)159 [¹⁴⁷Sm half-life].
- 1989Ad10 I. Adam, Zh. T. Zhelev, D. Zakoutski, B. Kracik, I. Penev, Bull. Acad. Sci. USSR, Phys. Ser. 53(1989)6 [¹⁴⁷Eu ε decay to ¹⁴⁷Sm].
- 1990Sc08 U. Schötzg, Nucl. Instrum. Meth. Phys. Res. A286(1990)523 [P_γ].
- 1992Ma56 J. B. Martins, M. L. Terranova, M. M. Correa, Nuovo Cim. 105(1992)1621 [¹⁴⁷Sm half-life].
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
- 2001Be81 F. Begemann, K. R. Ludwig, G. W. Lugmair, K. Min, L. E. Nyquist, P. J. Patchett, P. R. Renne, C. –Y. Shih, I. M. Villa, R. J. Walker, Geochim. Cosmochim. Acta 65(2001)111 [Correction for ¹⁴⁷Sm half-life].
- 2002Ba85 I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr., P. O. Tikkanen, S. Raman, Atomic Data Nucl. Data Tables 81(2002)1 [Theoretical ICC].
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)337 [Q].
- 2003Ki26 N. Kinoshita, A. Yokoyama, T. Nakanishi, J. Nucl. Radiochem. Sci. 4(2003)5 [¹⁴⁷Sm half-life].
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr. , Nucl. Instrum. Meth. Phys. Res. A589(2008)202 [Theoretical ICC].
- 2009Ko15 K. Kossert, G. Jörg, O. Nähle, C. Lierse, V. Gostomski, Appl. Rad. Isotop. 67(2009)1702 [¹⁴⁷Sm half-life].
- 2009Ni02 N. Nica, Nucl. Data Sheets 110(2009)749 [E_γ, E_{level}, spin, parity].

¹⁴⁸Pm - Comments on evaluation of decay data

by M.A. Kellett

This evaluation was completed in February 2013 with the same literature cut-off date.

Limitation of Relative Statistical Weights Method (LWM) was applied to average the decay data when appropriate (unless otherwise stated) by use of the *LWEIGHT* program and/or the Excel implementation of same. All uncertainties are given as the combined uncertainty to one standard deviation.

The ¹⁴⁸Pm samples used in the various measurements were produced via the ¹⁴⁸Nd(p,n), (d,2n), ¹⁴⁷Pm(n,γ) or U(p,f) reactions.

Impurities are other promethium isotopes (^{149,150}Pm) produced from further capture reactions on the original target, or from an incomplete purification of the production source, leading to the presence of other neodymium isotopes (^{149,150}Nd). However, radiochemical separation or ion-exchange chromatography was performed before measurements in almost all of the published reports in order to reduce the level of impurities.

The most important complication in measuring the emissions intensities is the fact that the ground state is in equilibrium with the metastable state (with its significantly longer half-life of 41.29 (13) d), and both are produced in almost equal quantities via the ¹⁴⁷Pm(n,γ) reaction. All emission intensity measurements must be made at “short” and “long” times following the source production, so that the measured intensities can be associated with the decay of the relevant isomer. In general, multiple measurements over a period of weeks and months have been performed.

1 Decay Scheme

¹⁴⁸Pm decays 100 % via β⁻-transitions to nine excited levels and the ground state of ¹⁴⁸Sm.

The ¹⁴⁸Pm decay scheme is complete and consistent. Its overall consistency is verified by the comparison between $Q_{\text{calc}} = 2478$ (24) keV deduced from the evaluated average energies of all emissions, and Q_{β^-} (2471 (6) keV) from the 2012 atomic mass evaluation of M. Wang et al. (2012Wa38). There are two gamma rays observed by 1977Ka14, but only as upper limits, not placed in this decay scheme. The level energies, spins, and parities, along with the multipolarities and mixing ratios of the γ-rays presented in this evaluation, are taken from 1977Ka14, as also used in 2000Bh03. Further details are provided in section 2.2.

It should be noted that both sets of the *absolute* and *relative* intensity measurements for the 550.27 keV γ-ray are discrepant.

2 Nuclear Data

2.1. Half-life

The experimental half-life values of ¹⁴⁸Pm are given in Table 1, resulting in a recommended value of **5.370 (15) d**.

Table 1: Experimental values of ¹⁴⁸Pm half-life (in days)

Reference	T _{1/2} values (d)	Uncertainty
1943Ku**	5.3 ^c	
1947Pa**	5.3 ^c	
1951Fo**	5.3 ^c	
1952Lo01	5.3 ^c	
1955Fo18	5.3 ^c	
1959Bh95	4.2 ^d	0.2
1959Ei31	5.5 ^d	0.1
1961Ei02	5.4	0.1
1962Re03	5.39	0.06

Reference	T _{1/2} values (d)	Uncertainty
1962Sc04	5.0 ^d	0.5
1963Ba31 ^a	5 ^e	
1971Ca23 ^b	5.3675	0.0153
1971Mo04	5.37	0.04
Recommended^c	5.370	0.015

LWEIGHT reduced $\chi^2 = 0.076$ (critical $\chi^2 = 3.786$)

^a 1963Ba31 quotes the same value as 1963Ew01 and 1963Ba06 by the same authors (hence not listed)

^b Unweighted mean of four published values, two in each of 1970Ca09 and 1971Ca23, and using simple arithmetic mean of the four uncertainties

^c Not used – all six cases where no uncertainty given

^d Rejected – three values rejected by the LWEIGHT program due to the Chauvenet principle

^e Recommended value is the weighted mean of 1961EI02, 1962Re03, 1971Ca23 and 1971Mo04, where the value of 1971Ca23 amounts to 81 % of the total weight, due to its far lower uncertainty, which is also adopted as the uncertainty on the recommended value

The decay daughter ¹⁴⁸Sm is effectively stable, with a half-life of **7 (3) × 10¹⁵ a**.

2.2. Gamma Transitions

Energies

The measured γ -ray transition energies in ¹⁴⁸Sm following the ¹⁴⁸Pm decay are given in Table 2. These have been measured by a small number of authors, with only one complete set (1977Ka14). Since all published values are compatible, the values of 1977Ka14, with their significantly lower uncertainties, have been adopted.

Table 2: Measured energies (keV) of γ -rays in ¹⁴⁸Sm from the decay of ¹⁴⁸Pm

1959Ei31		1962Re03		1967CI05		1970GrYP		1970FoZZ		1971Mo04*		1977Ka14		1984LaZZ§			
560	5	550	5	550.1	0.2	550.4	0.5	550.22	0.06	550.1	0.1	303.59	0.03	550.22	0.06		
												362.8	0.2				
												393.80	0.03				
												550.27	0.03				
												592.83	0.03				
611.2	0.6	611.1	0.1	611.26	0.03	819.27	0.03	874.17	0.23	874.18	0.03	874.2	0.25				
														896.36	0.3		
														896.42	0.03		
														903.94	0.03		
														914.97	0.12		
910	10	910	5	914.9	0.2	914.1	0.1	914.85	0.03	914.85	0.03	915.0	0.1				
														1113.88	0.03		
														1152.5	0.2		
														1371.3	0.2		
														1454.21	0.03		
1470	15	1460	10	1465.1	0.2	1464.5	1.5	1465.19	0.1	1465.1	0.1	1465.12	0.03	1465.2	0.1		
																1507.68	0.03
																1664.15	0.03
																1734.12	0.03
																1763.7	0.2
																2284.39	0.03
																2314.0	0.2

* Uncertainty of 0.1 keV on energy assumed

§ The values in 1984LaZZ are from 1970FoZZ, with the uncertainties rationalised

Emission Intensities

A number of relative (see Table 3) and absolute (see Table 4) measurements have been published for the γ -ray intensities. The relative measurements of 1977Ka14 are the only complete set, with twenty-

two γ -rays identified, although two of these were only measured as intensity upper limits and have not been included in the decay scheme.

Table 3: Measured relative intensities of γ -rays in ¹⁴⁸Sm from the decay of ¹⁴⁸Pm

	Energy	1959Bh95*		1962Sc04	1971Mo04§		1977Ka14		1984LaZZ†	
$\gamma_{1,0}$	303.59						0.17	0.02		
$\gamma_{4,0}$	362.8						<0.01			
$\gamma_{5,0}$	393.80						0.07	0.01		
$\gamma_{6,0}$	550.27	75	15	82	105	3	99.1	0.7	127	9
$\gamma_{9,0}$	592.83						1.59	0.03		
$\gamma_{2,1}$	611.26				5.0	0.3	4.60	0.05		
$\gamma_{3,1}$	819.27						0.06	0.01		
$\gamma_{4,1}$	874.18						1.06	0.03	1.59	0.09
$\gamma_{5,1}$	896.42						4.42	0.04	10.0	0.2
$\gamma_{6,1}$	903.94						0.19	0.01		
$\gamma_{7,1}$	914.85	60	12	65	56.4	1.5	51.6	0.4	68.2	6.8
$\gamma_{8,1}$	1113.88						0.10	0.01		
$\gamma_{9,1}$	1152.5						0.013	0.006		
$\gamma_{10,1}$	1371.3						0.062	0.006		
$\gamma_{5,2}$	1454.21						0.23	0.01		
$\gamma_{8,2}$	1465.12	100	20	100	100	3	100		100	3
$\gamma_{10,2}$	1507.68						0.025	0.004		
$\gamma_{8,5}$	1664.15						0.051	0.005		
$\gamma_{9,5}$	1734.12						0.174	0.003		
$\gamma_{8,6}$	1763.7						0.028	0.003		
$\gamma_{1,0}$	2284.39						0.20	0.01		
$\gamma_{4,0}$	2314.0						<0.001			

* Original values of 60 (12), 40 (8) and 100 (20) were revised as quoted by 1961El02 due to presence of ¹⁴⁴Pm

§ 1971Mo04 quotes absolute intensities also, but these are based on assuming 50 % β -feeding to ¹⁴⁸Sm ground state

† Although 1984LaZZ is a private communication, it is a detailed report of 55 pages which was never formally published. I also believe the report relates to work carried out in the early 1970s (see 1970FoZZ, 1974DuZO and 1974LoZL) and I note the report's first author was a summer visitor at Vanderbilt University (probably in 1974). The report was only registered in NSR in 1984.

After careful consideration of the earlier “absolute” measurements, e.g. 1961El02, 1962Re03, 1963Ba31 and 1970FoZZ, it was decided to use the only true absolute intensity, measured by 1971Ca23, for the 1465.12 keV γ ray of **0.222 (5)** to normalise the relative measurements. Once this was achieved, it was clear that the earlier measurements using NaI detectors (and those unpublished in 1970FoZZ/1984LaZZ which used a GeLi detector) were incompatible with the measurements of 1971Mo04 and 1977Ka14. The evaluator has thus chosen to use the weighted means of the three γ ray intensity measurements common to both sets to calculate the recommended absolute intensities (see Table 4). In each case the two values are consistent and the external uncertainty has been quoted on the recommended value.

Table 4: Measured/recommended absolute intensities (per decay) of γ -rays in ¹⁴⁸Sm from the decay of ¹⁴⁸Pm

	Energy	1961El02*		1962Re03		1963Ba31		1970FoZZ		1971Ca23		1971Mo04		1977Ka14§		Recommended† (per decay)	
$\gamma_{1,0}$	303.59													0.377×10^{-3}	0.045×10^{-3}	0.377×10^{-3}	0.045×10^{-3}
$\gamma_{4,0}‡$	362.8													$< 2.2 \times 10^{-5}$			
$\gamma_{5,0}$	393.80													1.55×10^{-4}	0.22×10^{-4}	1.55×10^{-4}	0.22×10^{-4}
$\gamma_{6,0}$	550.27	0.25	0.02	0.31	0.02	0.27	0.05	0.28	0.02			0.233	0.007	0.220	0.005	0.225	0.006
$\gamma_{9,0}$	592.83													0.0035	0.0001	0.0035	0.0001
$\gamma_{2,1}$	611.26											0.0112	0.0007	0.0102	0.0003	0.0104	0.0004
$\gamma_{3,1}$	819.27													1.33×10^{-4}	0.22×10^{-4}	1.33×10^{-4}	0.22×10^{-4}
$\gamma_{4,1}$	874.18							0.0035	0.0002					0.0024	0.0001	0.0024	0.0001
$\gamma_{5,1}$	896.42							0.0220	0.0005					0.0098	0.0002	0.0098	0.0002
$\gamma_{6,1}$	903.94													4.2×10^{-4}	0.2×10^{-4}	4.2×10^{-4}	0.2×10^{-4}
$\gamma_{7,1}$	914.85	0.13	0.02	0.17	0.02	0.15	0.03	0.150	0.015			0.125	0.003	0.115	0.003	0.120	0.005
$\gamma_{8,1}$	1113.88													2.22×10^{-4}	0.23×10^{-4}	2.22×10^{-4}	0.23×10^{-4}
$\gamma_{9,1}$	1152.5													2.9×10^{-5}	1.3×10^{-5}	2.9×10^{-5}	1.3×10^{-5}
$\gamma_{10,1}$	1371.3													1.38×10^{-4}	0.14×10^{-4}	1.38×10^{-4}	0.14×10^{-4}
$\gamma_{5,2}$	1454.21													5.11×10^{-4}	0.25×10^{-4}	5.11×10^{-4}	0.25×10^{-4}
$\gamma_{8,2}$	1465.12	0.20	0.02	0.24	0.02	0.23	0.05	0.220	0.007	0.222	0.005	0.222	0.005	0.222	0.005	0.222	0.005
$\gamma_{10,2}$	1507.68													5.6×10^{-5}	0.9×10^{-5}	5.6×10^{-5}	0.9×10^{-5}
$\gamma_{8,5}$	1664.15													1.13×10^{-4}	0.11×10^{-4}	1.13×10^{-4}	0.11×10^{-4}
$\gamma_{9,5}$	1734.12													3.86×10^{-4}	0.11×10^{-4}	3.86×10^{-4}	0.11×10^{-4}
$\gamma_{8,6}$	1763.7													6.2×10^{-5}	0.7×10^{-5}	6.2×10^{-5}	0.7×10^{-5}
$\gamma_{1,0}$	2284.39													4.44×10^{-4}	0.24×10^{-4}	4.44×10^{-4}	0.24×10^{-4}
$\gamma_{4,0}‡$	2314.0													$< 2 \times 10^{-6}$			

* Uncertainty of 0.02 on intensity assumed – as no uncertainty given

§ 1977Ka14 values are calculated from relative measurements normalised to recommended absolute emission of 1465.12 keV γ -ray

† Recommended values are the same as the normalised 1977Ka14 values, except for the 550.27 and 914.85 keV γ -ray values which are weighted means of the five available absolute values, including that of 1977Ka14 after initial normalisation through the 1465 keV γ -ray

‡ Two γ -rays at 362.8 and 2314.0 keV not placed in scheme (measured as upper limit only)

Measurement conditions

The γ -ray intensities were measured with the detectors and under the timing conditions shown in Table 5.

Table 5: Conditions of the γ -ray intensity measurements

Reference	Detector	Timescale
1959Bh95	NaI	2 days to 3 months
1959Ei31	NaI	No timing details given
1961Ei02	NaI	Up to 250 days
1962Sc04	NaI	At 24 hrs and “equilibrium”
1962Re03	NaI	“Immediately after irradiation” to “several months”
1963Ba31	NaI	2 days to 140 days
1967Cl05	GeLi	No details given, but measured combined energy spectrum
1970FoZZ	GeLi	Immediately after irradiation, then two and twelve weeks later
1970GrYP	GeLi	No timing details given
1971Ca23	NaI	During first 34 – 50 hrs, and a further 40 days, following irradiation
1971Mo04	GeLi	3 to 75 days
1977Ka14	GeLi	Four sources produced periodically over 10 month measuring period
1984LaZZ	Same measurements as reported in 1970FoZZ	

Multipolarities and Internal Conversion Coefficients

The γ -ray multipolarities and the mixing ratios have been taken from the adopted gamma data within the evaluation of 2000Bh03 based on mainly on 1977Ka14 (γ -ray directional correlations), but also considering 1962Re03, 1968Wy02 (γ -ray directional correlations), 1962Sc04, 1963Ba31 (measured conversion electrons and γ -ray directional correlations), and 1970GrYP (measured conversion electrons).

All internal conversion coefficients have been calculated using the BrIcc code (2008Ki07), as implemented in the SAISINUC evaluation tool (2008DuZX). In Table 6 these values are compared with available measured data, where general agreement is seen.

Table 6: Comparison of BrIcc and measured K-shell internal conversion coefficients (α_K)

	Energy	1962Sc04*		1963Ba31		1970GrYP		BrIcc	
$\gamma_{6,0}$	550.27	0.0081	-			0.0079	0.0006	0.008 25	0.000 12
$\gamma_{2,1}$	611.26					0.0025	0.0008	0.002 39	0.000 05
$\gamma_{4,1}$	874.18	0.0050	0.0014					0.002 80	0.000 04
$\gamma_{7,1}$	914.85			0.000 68	0.000 19			0.001 050	0.000 015
$\gamma_{8,2}$	1465.12			0.000 47	0.000 14			0.000 449	0.000 007
$\gamma_{10,2}$	1507.68	0.0006	0.0002					0.000 428	0.000 006

* Obtained from ¹⁴⁸Eu EC decay for γ -rays in common

2.3. Beta Transitions*Energies and Emission Probabilities*

Beta-particle energies were determined from the difference between the adopted Q-value and the final level energies of daughter ¹⁴⁸Sm. Beta-particle emission probabilities were determined by balancing the proposed decay scheme through consideration of the $\beta\gamma$ -population and γ -depopulation of these same levels. Only the 2nd level has no beta-feeding based on published data. A comparison with

measured energies and absolute emission intensities is presented in Table 7. The adopted values are consistent, within the uncertainty limits, with the two sets of results having uncertainties.

Table 7: Measured/recommended absolute intensities (%) of β -emissions from the decay of ¹⁴⁸Pm

	1961EI02		1962Sc04		1962Re03		1963Ba31†		Recommended	
	E (keV)	I $_{\beta}$ (%)	E (keV)	I $_{\beta}$ (%)	E (keV)	I $_{\beta}$ (%)	E (keV)	I $_{\beta}$ (%)	E (keV)	I $_{\beta}$ (%)
$\beta_{0,0}$	2520	55	2460 (20)	22	2600 (200)	45 (9)	2480 (30)	50 (5)	2471 (6)	55.5 (7)
$\beta_{0,1}$	1970	12	1900 (50)	6	2000 (200)	14*	1930 (30)	10 (2)	1921 (6)	9.3 (6)
$\beta_{0,3}$									1047 (6)	0.236 (9)
$\beta_{0,4}$									1017 (6)	0.093 (3)
$\beta_{0,5}$	1060	33	1020 (15)	72	1100 (200)	41 (9)	1020 (30)	40 (5)	1006 (6)	33.3 (6)
$\beta_{0,6}$									807 (6)	0.018 (3)
$\beta_{0,7}$									549 (6)	0.0138 (14)
$\beta_{0,8}$									413 (6)	1.360 (22)
$\beta_{0,9}$									187 (6)	0.0965 (34)
$\beta_{0,10}$									157 (6)	0.0091 (15)

* 2000 keV intensity deduced by 1962Re03 from decay scheme, not directly measured

† Same data as published in 1963Ew01 and 1963Ba06 by the same authors

2.4. Conversion electrons

The energies of the internal conversion electrons have been deduced using γ -ray energies and electron binding energies from 1977La19 and 1996FiZX. Absolute conversion electron emission intensities have been calculated by using γ -ray internal conversion coefficients and absolute emission intensities.

3 Atomic Data

The recommended fluorescence yields shown in Table 8 were obtained from 1996Sc06.

Table 8: Recommended atomic data

ω_K	0.926 (4)
ω_L	0.158 (6)
η_{KL}	0.857 (4)

The energies of the K-Auger electrons were obtained from 1998ScZM, whereas the energies and the yield of Sm X-rays were obtained from 1999ScZX (data stored in SAISINUC). The X-ray and Auger electron emission intensities have been calculated from the decay scheme parameters using the computer program EMISSION (v3.10, 28-Jan-2003) described in 2000Sc47.

References

- 1943Ku** J.D.Kurbatov, M.L.Pool, Phys. Rev. 63 (1943) 463, 1
[Half-life, Beta transition energy]
- 1947Pa** G.W.Parker, P.M.Lantz, M.G.Inghram, D.C.Hess Jr, R.J.Hayden, Phys. Rev. 72 (1947) 85
[Half-life, Beta and gamma transition energies]
- 1948Se40 G.T.Seaborg, I.Perlman, Rev. Mod. Phys. 20 (1948) 585
[Half-life, Beta and gamma transition energies]
- 1951Fo** R.L.Folger, P.C.Stevenson, G.T.Seaborg, Report UCRL-1195, Univ. California (1951) 22
[Half-life, Beta and gamma transition energies]
- 1952Lo01 J.K.Long, M.L.Pool, Phys. Rev. 85 (1952) 137
[Half-life, Beta and gamma transition energies]
- 1955Fo18 R.L.Folger, P.C.Stevenson, G.T.Seaborg, Phys. Rev. 98 (1955) 107
[Half-life, Beta and gamma transition energies, IT branching fraction]
- 1955He64 N.P.Heydenburg, G.M.Temmer, Phys. Rev. 100 (1955) 150
[Sm-148 levels]
- 1955Ma77 H.Mark, G.T.Paulissen, Phys. Rev. 100 (1955) 813
[Sm-148 levels]
- 1959Bh95 S.K.Bhattacharjee, B.Sahai, C.V.K.Baba, Nucl. Phys. 12 (1959) 356
[Half-life, Beta and gamma transition energies and intensities, IT branching fraction, Sm-148 levels]
- 1959Ei31 J.A.Eisele, Diss. Abst. Int. 20 (1960) 3794
[Half-life, Beta and gamma transition energies, Sm-148 levels]
- 1959Ei31 J.A.Eisele, Thesis, Ohio State Univ. (1959)
[Half-life, Beta and gamma transition energies, Sm-148 levels]
- 1960Sc09 C.F.Schwerdtfeger, E.G.Funk, J.W.Mihelich, Bull. Am. Phys. Soc. 5 (1960) 425, P
[Half-life, Beta and gamma transition energies, Sm-148 levels]
- 1960Sc12 R.P.Schuman, J.R.Berreth, R.L.Heath, C.W.Reich, Bull. Am. Phys. Soc. 5 (1960) 494, C
[Beta and gamma transition energies and intensities, IT branching fraction, Sm-148 levels]
- 1961Br41 M.K.Brice, C.W.Reich, R.G.Helmer, Report IDO-16710 (1961)
[Half-life, Gamma transition energies and intensities, ICCs and multipolarities, IT branching fraction]
- 1961Ei02 J.S.Eldridge, W.S.Lyon, Nucl. Phys. 23 (1961) 131
[Half-life, Beta and gamma transition energies and intensities, IT branching fraction]
- 1962Gr34 R.W.Grant, D.A.Shirley, Report UCRL-10624, Univ. California (1962)
[Pm-148 and Sm-148 level spins]
- 1962Re03 C.W.Reich, R.P.Schuman, J.R.Berreth, M.K.Brice, R.L.Heath, Phys. Rev. 127 (1962) 192
[Half-life, Beta and gamma transition energies and intensities, IT branching fraction, Conversion electrons, Sm-148 levels]
- 1962Sc04 C.F.Schwerdtfeger, E.G.Funk Jr, J.W.Mihelich, Phys. Rev. 125 (1962) 1641
[Half-life, Beta and gamma transition energies and intensities, IT branching fraction, Conversion electrons, Sm-148 levels]

- 1963Al22 D.Ali, R.Marrus, Bull. Am. Phys. Soc. 8 (1963) 619, S
[Pm-148 level spin]
- 1963Ba06 C.V.K.Baba, G.T.Ewan, J.F.Suarez, Phys. Lett. 3 (1963) 232
[Pm-148 and Sm-148 level spins]
- 1963Ba31 C.V.K.Baba, G.T.Ewan, J.F.Suarez, Nucl. Phys. 43 (1963) 264
[Beta and gamma transition energies and intensities, IT branching fraction, Conversion electrons, Pm-148 and Sm-148 levels]
- 1963Ba32 C.V.K.Baba, G.T.Ewan, J.F.Suarez, Nucl. Phys. 43 (1963) 285
[Sm-148 levels]
- 1963Ew01 G.T.Ewan, C.V.K.Baba, J.F.Suarez, Bull. Am. Phys. Soc. 8 (1963) 73, VA
[Beta transition energies, intensities and shape, Sm-148 levels]
- 1963Gr10 R.W.Grant, D.A.Shirley, Phys. Rev. 130 (1963) 1100
[Pm-148 and Sm-148 level spins]
- 1964Ha17 J.W.Harpster, K.J.Casper, Nucl. Phys. 52 (1964) 497
[Gamma multiplicities, Sm-148 levels]
- 1964Ke03 R.A.Kenefick, R.K.Sheline, Phys. Rev. 133 (1964) B25
[Sm-148 level energies]
- 1965Al16 D.Ali, Nucl. Phys. 71 (1965) 441
[Pm-148 level spin]
- 1967Cl05 J.E.Cline, R.L.Heath, Report IDO-17222 (1967)
[Gamma transition energies and intensities]
- 1968Wy02 L.D.Wyly, E.T.Patronis Jr, C.H.Braden, Phys. Rev. 172 (1968) 1153
[Sm-148 level spins]
- 1970Ca09 M.J.Cabell, M.Wilkins, J. Inorg. Nucl. Chem. 32 (1970) 1409
[Half-life]
- 1970FoZZ J.W.Ford Jr, Thesis, Vanderbilt Univ. (1970)
[Gamma transition energies and absolute intensities, Multipolarities, Sm-148 levels and spins]
- 1970FoZZ J.W.Ford Jr, Diss. Abst. Int. 31B (1970) 3631
[Gamma transition energies and absolute intensities, Multipolarities, Sm-148 levels and spins]
- 1970GrYP E.P.Grigorev, A.V.Zolotavin, V.O.Segreev, M.I.Sovtsov, Program and Theses, Proc. 20th Ann. Conf. Nucl. Spectrosc. At. Nuclei, Leningrad (in Russian) (1970) 100
[Conversion electrons, Multipolarities, Pm-148 and Sm-148 levels]
- 1970Pe22 L.K.Peker, Izv. Akad. Nauk SSSR Ser. Fiz. 34 (1970) 2214; Bull. Acad. Sci. USSR Phys. Ser. (English translation) 34 (1971) 1975
[Pm-148 level spin]
- 1971Ca23 M.J.Cabell, M.Wilkins, J. Inorg. Nucl. Chem. 33 (1971) 1957
[Half-life, Beta and gamma transition energies and absolute intensities]
- 1971Mo04 R.S.Mowatt, W.H.Walker, Can. J. Phys. 49 (1971) 108
[Half-life, Beta and gamma transition energies and intensities]
- 1971Sh08 K.Shoda, A.Suzuki, M.Sugawara, T.Saito, H.Miyase, S.Oikawa, B.N.Sung, Phys. Rev. C3 (1971) 2006
[Pm-148 level spin]

- 1972AmZX R.A.Amadori, J.R.Horgan Jr, K.S.R.Sastry, Proc. Nucl. Phys. Solid State Phys. Symp., Bombay, India 14B (1972) 337
[Pm-148 level spin, Sm-148 level spins, beta transition, logft]
- 1974DuZO D.R.Dunn, A.V.Ramayya, J.W.Ford Jr, J.H.Hamilton, W.Lourens, J.J.Pinajian, Bull. Am. Phys. Soc. 19 (1974) 1124, FB8
[Gamma transition energies, Sm-148 levels]
- 1974LoZL W.Lourens, V.Lakshminarayana, J.H.Hamilton, A.V.Ramayya, D.R.Dunn, S.M.Brahmavar, J.J.Pinajian, Bull. Amer. Phys. Soc. 19 (1974) 1124, FB7
[Sm-148 levels]
- 1977Ka14 C.A.Kalfas, J. Phys. (London) G3 (1977) 929
[Gamma transition energies and intensities, Multipolarities, Mixing ratios, Sm-148 levels and spins]
- 1977La19 F.P.Larkins, At. Data Nucl. Data Tables 20 (1977) 311
[Electron Binding Energies]
- 1984LaZZ V.Lakshminarayana, B.Van Nooijen, W.Lourens, A.V.Ramayya, J.H.Hamilton, J.W.Ford Jr, D.R.Dunn, J.J.Pinajian, Priv. Comm. to NNDC (unpublished report) (1984)
[Gamma transition energies and intensities, Multipolarities, Sm-148 levels and spins]
- 1988No02 E.B.Norman, K.T.Lesko, A.E.Champagne, Phys. Rev. C37 (1988) 860
[Pm-148 level energies and spins]
- 1989Le01 K.T.Lesko, E.B.Norman, R.-M.Larimer, J.C.Bacelar, E.M.Beck, Phys. Rev. C39 (1989) 619
[Pm-148 level energies and spins]
- 1996FiZX R.B.Firestone, Table of Isotopes 8th Ed., John Wiley and Sons Inc. 2 (1996)
[Electron Binding energies]
- 1996Sc06 E.Schönfeld, H.Janssen, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527
[Atomic Data]
- 1998ScZM E.Schönfeld, G.Rodloff, PTB Report 6.11-98-1 6.11 (1998) 1
[Auger electrons]
- 1999ScZX E.Schönfeld, G.Rodloff, PTB Report 6.11-1999-1 6.11 (1999) 1
[K X-rays]
- 2000Bh03 M.R.Bhat, Nucl. Data Sheets 89 (2000) 797
[Sm-148 levels, Multipolarities, Mixing ratios]
- 2000Sc47 E.Schönfeld, H.Janssen, Appl. Radiat. Isot. 52 (2000) 595
[X-ray and Auger Electron emission probabilities]
- 2008DuZX C.Dulieu, M.M.Bé, V.Chisté, Proc. Int. Conf. on Nuclear Data for Science and Technology, 22-27 April 2007, Nice, France (2008) 97
[SAISINUC software]
- 2008Ki07 T.Kibédi, T.W.Burrows, M.B.Trzhaskovskaya, P.M.Davidson, C.W.Nestor Jr, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202
[Theoretical ICCs]
- 2012Wa38 M.Wang, G.Audi, A.H.Wapstra, F.G.Kondev, M.MacCormick, X.Xu, B.Pfeiffer, Chin. Phys. C36 (2012) 1603
[Q]

^{148m}Pm - Comments on evaluation of decay data

by M.A. Kellett

This evaluation was completed in March 2013 with the same literature cut-off date.

Limitation of Relative Statistical Weights Method (LWM) was applied to average the decay data when appropriate (unless otherwise stated) by use of the LWEIGHT program and/or the Excel implementation of same. All uncertainties are given as the combined uncertainty to one standard deviation.

The ^{148m}Pm samples used in the various measurements were produced via the ¹⁴⁸Nd(p,n), (d,2n), ¹⁴⁷Pm(n,γ) or U(p,f) reactions.

Impurities are other promethium isotopes (^{149,150}Pm) produced from further capture reactions on the original target, or from an incomplete purification of the production source, leading to the presence of other neodymium isotopes (^{149,150}Nd). However, radiochemical separation or ion-exchange chromatography was performed before measurements in almost all of the published reports in order to reduce the level of impurities.

The most important complication in measuring the emission intensities is the fact that the metastable state is in equilibrium with the ground state (with its significantly shorter half-life of 5.370 (15) d), and both are produced in almost equal quantities via the ¹⁴⁷Pm(n,γ) reaction. All emission intensity measurements must be made at “short” and “long” times following the source production, so that the measured intensities can be associated with the decay of the relevant isomer. In general, multiple measurements over a period of weeks and months have been performed.

1 Decay Scheme

^{148m}Pm decays 94.4 (5) % via β⁻-transitions to four excited levels of ¹⁴⁸Sm, and via an isomeric transition of 5.6 (5) %.

After a certain time (~ 50 d), the ratio of the activities A(^{148m}Pm)/A(¹⁴⁸Pm) becomes constant.

The ^{148m}Pm decay scheme is complete and consistent. Its overall consistency is verified by the comparison between Q_{calc} = 2612 (21) keV deduced from the evaluated average energies of all emissions, and Q_{eff} (2608 (6) keV) from the 2012 atomic mass evaluation of M. Wang et al. (2012Wa38). The level energies, spins, parities along with the multipolarities and mixing ratios of the γ-rays presented in this evaluation are taken from the evaluation of 2000Bh03. Further details are provided in section 2.3.

2 Nuclear Data

2.1. Half-life

The experimental half-life values of ^{148m}Pm are given in Table 1, resulting in a recommended value of **41.29 (13) d**.

Table 1: Experimental values of ^{148m}Pm half-life (in days)

Reference	T _{1/2} values (d)	Uncertainty
1951Fo**	43 ^c	
1952Ki25	42	1
1952Lo01	48 ^c	
1955Fo18	43 ^c	
1959Bh95	45.8 ^d	2.9
1959Ei31	50 ^b	1
1961El02	41.8	0.2
1961Ha23	46 ^c	
1962Sc04	45.5 ^d	0.5

Reference	T _{1/2} values (d)	Uncertainty
1962Re03	40.6	0.4
1963Ba31 ^a	41 ^c	
1971Mo04	40.9	0.2
1971BaZW	41.4	0.8
1971Wa05	41.3	0.1
Recommended*	41.29	0.13

LWEIGHT reduced $\chi^2 = 2.76$ (critical $\chi^2 = 3.02$)

^a 1963Ba31 quotes the same value as 1963Ew01 and 1963Ba06 by the same authors (hence not listed)

^b Rejected by the LWEIGHT program due to the Chauvenet principle

^c Not used – all five cases where no uncertainty given

^d Two values from 1959Bh95 and 1962Sc04 have been rejected by the evaluator, as they are not consistent with other measured values and since their reported values for ¹⁴⁸Pm were also rejected due to the Chauvenet principle, it is judged that their half-life values are not reliable

* Recommended value is the weighted mean of 1952Ki25, 1961El02, 1962Re03, 1971Mo04, 1971BaZW and 1971Wa05, where the value of 1971Wa05 amounts to 63 % of the total weight

The IT decay daughter ¹⁴⁸Pm has an evaluated half-life of **5.370 (15) d** and the β^- -decay daughter ¹⁴⁸Sm is effectively stable, with a half-life of **7 (3) × 10¹⁵ a**.

2.2. Branching fraction

^{148m}Pm decays via β^- (**94.4 (5) %**) to four excited levels of ¹⁴⁸Sm and via IT (**5.6 (5) %**). Table 2 shows the measured and recommended IT branching fraction.

Table 2: Experimental branching fractions for ^{148m}Pm IT decay

Reference	%IT	Comments	Method
1952Ki25	7.4	$I_{\beta(2700)}$ following IT decay ¹	(¹⁴⁸ Nd) ₂ O ₃ (p,n), E _p =8.9 MeV; Al, Be, Pb absorbers, 180° low resolution beta-ray spectrometer
1955Fo18	5–10	$I_{\beta(2700)}$ following IT decay ¹	U(p,f), E _p =340 MeV; Al absorber and GM tube
1959Bh95	~20		(¹⁴⁸ Nd) ₂ O ₃ (p,n), E _{p(max)} = 12 MeV; measured I_γ with NaI(Tl), I_β with anthracene scintillator, $\gamma\gamma^-$, $\gamma\beta^-$ -coin
1960Sc12 ^a	~5	No information on how this was deduced.	¹⁴⁷ Pm(n _{th} , γ) in Materials Testing Reactor
1961El02	5–10	Qualitative estimate only of $I_{\beta(2300)}$ following IT decay	(¹⁴⁸ Nd) ₂ O ₃ (p,n), E _p not given and ¹⁴⁷ Pm(n _{th} , γ) in Oak Ridge Research Reactor; measured I_γ with NaI(Tl), I_β with prop. counter and absorbers, $\gamma\gamma^-$ -coin
1961Br41 ^a	5.8 (15)	$I_{\gamma(1460)}$ of 1.5 (4) % following IT decay ²	No details of production method; measured I_{ce} with magnetic spectrograph
	5.5 (18)	$I_{\gamma(75)}$ – decay of intermediate 75 keV level ³	

Reference	%IT	Comments	Method
1962Sc04	8	By comparing the $I_{\gamma(1460)}$ to the $I_{\gamma(630)}$ at equilibrium	$(^{148}\text{Nd})_2\text{O}_3(\text{p},\text{n})$, E_p not given; measured I_{γ} with NaI(Tl), I_{ce} with magnetic spectrograph and intermediate-image spectrometer, I_{β} with Pilot B crystal, $\gamma\gamma$ -, $\gamma\beta$ -coin
1962Re03	~7	From $I_{\beta(2600)}$ following IT decay ¹	$^{147}\text{Pm}(\text{n}_{\text{th}},\gamma)$ in Materials Testing Reactor; measured I_{γ} with NaI(Tl), I_{ce} with magnetic spectrograph, I_{β} with prop. counter and anthracene scintillator
	6.5 (17)	From $I_{\gamma(1460)}$ of 1.6 (3) % following IT decay ⁴	
	5.1 (18)	From $I_{\gamma(75)}$ – decay of intermediate 75 keV level ⁵	
1963Ba31	6.6 (10)	From $I_{\gamma(76)}$ – decay of intermediate 76 keV level ⁶	$^{148}\text{Nd}(\text{d},2\text{n})$, $E_d=12$ MeV; Measured I_{γ} with NaI(Tl), I_{ce} and I_{β} with “Copenhagen” six-gap spectrometer
	6.1 (10)	From $I_{\gamma(62)}$ – decay of isomeric 137 keV level ⁷	
	7.6 (16)	From $I_{\beta(2600)}$ following IT decay ¹	
	10 (4)	From $I_{\gamma(1460)}$ of 1.6 (3) % following IT decay ⁴	
	6.9 (7)	Mean value of the 4 values quoted by the authors	
	6.6 (6)	Mean value of the 4 values calculated by evaluator	
1970GrYP	~7-7.5	From I_{β} 's deduced from I_{γ} and conversion electron measurements	Measured I_{γ} with Ge(Li) and I_{ce} with β -spectrometer
1971Mo04	4.6 (5)	From mean of three relative intensity ratios at long decay time, i.e. once equilibrium reached: $I_{\gamma(550)}/I_{\gamma(1465)}$, $(I_{\gamma(611)}+I_{\gamma(630)})/I_{\gamma(1465)}$ and $(I_{\gamma(1014)}+I_{\gamma(915)}+I_{\gamma(726)}+I_{\gamma(433)}+I_{\gamma(414)})/I_{\gamma(1465)}$	$^{147}\text{Pm}(\text{n}_{\text{th}},\gamma)$ in NRU reactor; measured I_{γ} with GeLi detector
Recommended	5.6 (5)	Using method described in 1971Mo04, with evaluated relative intensities measured at equilibrium (described below)	

^a Superseded by 1962Re03

¹ Measured absolute intensity of the high energy beta group (~2600–2700 keV) emitted by ^{148}Pm ground state, as fed from the IT decay at long times, thus indicates the IT branching fraction.

² Measured the relative intensity of the 1460 keV γ ray as 1.5 (4) % of the long-lived decays, then based on relative intensity calculated branching as 5.8 (15) %. Used ~ 26 % as absolute intensity per decay for the 1460 keV γ ray.

³ Calculated absolute transition probability of 5.5 (18) % for the 75 keV γ ray from measured intensity of 1.2 (4) % and calculated ICCs for M1 transition, with $\alpha_T=3.64$ (see Table VII of 1961Br41).

⁴ Measured the relative intensity of the 1460 keV γ ray as 1.6 (3) % (see Table II of 1962Re03, i.e. $0.017(3) \times 0.94 \times 100$) of the long-lived decays, then based on relative intensity calculated branching as 6.5 (17) %. Used ~ 24 % (see Table I of 1962Re03) as absolute intensity per decay for the 1460 keV γ ray.

⁵ Calculated absolute transition probability of 5.1 (18) % for the 75 keV γ ray from measured intensity of 1.0 (4) % (see Table II of 1962Re03, i.e. $0.011(4) \times 0.94 \times 100$) and calculated ICCs for M1 transition, with $\alpha_T=3.64$ (see Table III of 1962Re03).

⁶ Calculated absolute transition probability of 6.6 (10) % for the 76 keV γ ray from measured intensity of 1.6 (4) % (see Table 10 of 1963Ba31) assuming M1 transition.

⁷ Calculated absolute transition probability of 6.1 (10) % for the 62 keV γ ray from measured K conversion electron intensity of < 0.6 % (see Table 10 of 1963Ba31) assuming E4 transition.

The recommended IT branching fraction has been determined using the method described in 1971Mo04, which involves calculating the ratios of three different sets of relative γ -ray emission intensities at equilibrium. Each ratio yields a value for the branching fraction, and then the weighted mean of these three values is used to calculate the final recommended value, but due to the strong correlation in this method, the lowest experimental uncertainty is assigned to the value, rather than the calculated uncertainty of the weighted mean.

From 1971Mo04 the ratio of the decay rate of ^{148}Pm to $^{148\text{m}}\text{Pm}$ at equilibrium is:

$$R_i = \frac{\lambda_i}{\lambda_m} \left(1 - \frac{\lambda_m}{\lambda_g}\right)^{-1} \quad (1)$$

where λ_i is the IT partial decay constant, λ_g and λ_m are the ground and metastable state decay constants, then λ_i/λ_m is the IT transition probability. Using the recommended half-lives of 5.370 (15) d and 41.29 (13) d then:

$$R_i = 1.149 (5) \frac{\lambda_i}{\lambda_m} \quad (2)$$

Since all of the $^{148\text{m}}\text{Pm}$ β decays are to the 1594.31 keV level or above, and these feed γ transitions passing through either of the levels at 1161.53 keV or 1180.24 keV, then with reference to the decay scheme presented here, one finds:

$$1 - \frac{\lambda_i}{\lambda_m} = I_{550} = (I_{611} + I_{630}) = (I_{414} + I_{433} + I_{726} + I_{915} + I_{1014}) \quad (3)$$

where in each case I_x denotes the transition probability for the transition of energy x .

Due to the existence of the IT branch, all measured transition intensities include a minor contribution from the ^{148}Pm formed, which can be taken into account by including a correction term based on the absolute intensity evaluated for transition x following decay of the ground state.

For example, for the 550 keV transition is the sum of two components from $^{148\text{m}}\text{Pm}$ IT decay AND ^{148}Pm β decay, which contribute to the measured intensity, giving a combined intensity:

$$I_{550}^C = 1 - \frac{\lambda_i}{\lambda_m} + 0.227R_i = 1 - 0.739 \frac{\lambda_i}{\lambda_m} \quad (4)$$

since the absolute transition probability (γ -ray and conversion electrons) of the 550 keV γ -ray from ^{148}Pm is 0.227 (6) (from companion evaluation of ^{148}Pm) and by using Eqn. 2.

Equivalent equalities can be found for the $(I_{611} + I_{630})$ cascade:

$$I_{611}^C + I_{630} = 1 - \frac{\lambda_i}{\lambda_m} + 0.01043R_i = 1 - 0.988 \frac{\lambda_i}{\lambda_m} \quad (5)$$

and the $(I_{414} + I_{433} + I_{726} + I_{915} + I_{1014})$ cascade;

$$I_{414} + I_{433} + I_{726} + I_{915}^C + I_{1014} = 1 - \frac{\lambda_i}{\lambda_m} + 0.12R_i = 1 - 0.862 \frac{\lambda_i}{\lambda_m} \quad (6)$$

where the 611 keV and 915 keV γ -rays include the two components.

Finally, the 1465 keV γ -ray's metastable component can be shown to be:

$$I_{1465} = 0.222R_i = 0.255 \frac{\lambda_i}{\lambda_m} \quad (7)$$

By dividing each of Eqns. 4 – 6 by Eqn. 7, and using the relative intensities given in Table 4 we can determine three values for λ_i/λ_m of 0.056 (4), 0.056 (5) and 0.056 (5) respectively, giving a weighted mean of **0.056 (5)** and using the highest individual uncertainty.

This recommended IT branching fraction of **5.6 (5) %** is consistent with experimental values determined from direct measurements of the 75 keV γ ray of 1961Br41: 5.5 (18) %, 1962Re03: 5.1 (18) % and 1963Ba31: 6.6 (10) %.

2.3. Gamma Transitions

Energies

The measured γ -ray transition energies in ^{148}Sm following the $^{148\text{m}}\text{Pm}$ β^- decay and from the isomeric transition are given in Table 3. These have been measured by a number of authors.

The evaluated ^{148}Sm transition energies are deduced from the adopted level energy differences from 2000Bh03 and are consistent with the measured values given in 1977Ka14.

For the isomeric transition, the level energy is assumed to be the same as the sum of the energy of the two γ -rays inferred from the conversion electron measurements of 1970GrYP, i.e. 137.1 keV, not 137.9 keV as in 2000Bh03, where an inconsistency exists between the “adopted energy levels”, the actual energy level quoted for the isomer and the two γ -rays cascading from it. This decision is also supported by the $^{149}\text{Sm}(d,^3\text{He})$ reaction measurements reported in 1988No02 and 1989Le01 for example.

Seven unplaced γ -rays consist of five seen by 1970FoZZ (hence also 1984LaZZ), as well as one each observed by 1959Ei31 and 1967Cl05.

Table 3: Measured and recommended energies (keV) of γ -rays in ^{148}Sm from the decay of $^{148\text{m}}\text{Pm}$

	1959Ei31	1962Re03	1967Cl05	1970FoZZ	1970Gr09	1970GrYP	1971Mo04*	1977Ka14	1984LaZZ§	Recommended
^{148}Pm										
$\gamma_{2,1}$		60.1 0.5				61.3 0.05				61.30 0.05
$\gamma_{1,0}$		74.6 0.5	75.6 0.2	75.33 0.4	75.9 0.6	75.8 0.1			75.3 0.4	75.8 0.1
^{148}Sm										
$\gamma_{9,8}$		98 0.5	98.5 0.2	98.17 0.25	99.2 0.6	98.5 0.1	98.5 0.1	98.48 0.03	98.15 0.25	98.48 0.03
$\gamma_{8,7}$		188 1	189.5 0.3	189.61 0.06	190.2 0.6	189.7 0.2	189.5 0.1	189.63 0.03	189.61 0.06	189.63 0.03
$\gamma_{9,7}$	290 5	286 2	288 0.2	288.09 0.06	288.4 0.6	288.1 0.2	288 0.1	288.11 0.03	288.09 0.06	288.11 0.03
Unplaced				295.06 0.1						
$\gamma_{9,6}$				298.88 0.25				299.1 0.2		299.1 0.2
$\gamma_{7,4}$		309 2	311.7 0.3	311.62 0.06	312 0.6	311.5 0.4	311.7 0.1	311.63 0.03	311.62 0.06	311.63 0.03
$\gamma_{8,5}$				362.48 0.1				362.09 0.03	362.5 0.1	362.09 0.03
$\gamma_{4,3}$	420 5	412 2	414.1 0.3	414.02 0.06	414.1 0.3	413.8 0.4	414.1 0.1	414.07 0.03	414.02 0.06	414.07 0.03
$\gamma_{4,2}$		431 2	432.7 0.3	432.71 0.06	432.9 0.1	432.5 0.4	432.7 0.1	432.78 0.03	432.71 0.06	432.78 0.03
$\gamma_{9,5}$			443.1 0.4	460.5 0.1				460.57 0.03	460.5 0.1	460.57 0.03
$\gamma_{8,4}$		498 2	501.1 0.4	501.25 0.06	501.4 0.1	501.1 0.5	501.1 0.1	501.26 0.03	501.25 0.06	501.26 0.03
Unplaced				533.59 0.15					533.6 0.15	
$\gamma_{1,0}$		560 5	548 5	550.22 0.06	550.2 0.3	550.4 0.5	550.1 0.1	550.27 0.03	550.22 0.06	550.27 0.03
$\gamma_{5,3}$				553.26 0.19				553.24 0.03	553.3 0.2	553.24 0.03
$\gamma_{5,2}$				571.88 0.15				571.95 0.03	571.9 0.15	571.95 0.03
Unplaced			592.4 0.4							
$\gamma_{9,4}$		596 5	599.5 0.3	599.69 0.06	599.7 0.1	599.6 0.6	599.5 0.1	599.74 0.03	599.69 0.06	599.74 0.03
$\gamma_{2,1}$			611.1 0.3	611.23 0.06	611.4 0.1	611.2 0.6	611.1 0.1	611.26 0.03	611.23 0.06	611.26 0.03
$\gamma_{3,1}$	630 5	627 5	629.9 0.2	629.95 0.06	629.8 0.3	629.7 0.6	629.9 0.1	629.97 0.03	629.95 0.06	629.97 0.03
$\gamma_{6,3}$								714.7 0.2		714.7 0.2
$\gamma_{7,3}$	740 5	723 5	725.6 0.2	725.67 0.08	725.6 0.1	725.6 0.7	725.6 0.1	725.70 0.03	725.67 0.08	725.70 0.03
Unplaced				874.17 0.23					874.2 0.25	
Unplaced			896.2 0.3	896.36 0.3					896.4 0.3	
From g.s.				914.97 0.12		914			915 0.1	
$\gamma_{8,3}$	910 10	913 5	914.9 0.2	915.31 0.06	915.1 0.3	915.6 0.8	914.9 0.1	915.33 0.03	915.31 0.06	915.33 0.03
$\gamma_{9,3}$	1005 10	1011 10	1013.7 0.3	1013.8 0.08	1013.6 0.3	1013.6 0.9	1013.7 0.1	1013.81 0.03	1013.8 0.08	1013.81 0.03
Unplaced				1121.26 0.33					1121.3 0.3	
$\gamma_{6,1}$				1345.02 0.3				1344.6 0.2	1345 0.3	1344.6 0.2
From g.s.	1470 15	1460 10	1465.1 0.2	1465.19 0.1	1465.2 0.6	1464.5 1.5	1465.1 0.1		1465.2 0.1	1465.12 0.03
Unplaced	1610 15									

* Uncertainty of 0.1 keV on energy assumed on 1971Mo04 data

§ The values in 1984LaZZ are from 1970FoZZ, with the uncertainties rationalised

Emission Intensities

A number of relative (see Table 4) measurements have been published for the γ -ray intensities. No absolute measurements have been made. Details of the detectors used and the measurement timescale are given in Table 5.

Table 4: Measured relative intensities of γ -rays in ^{148}Sm from the decay of $^{148\text{m}}\text{Pm}$, the 629,97 keV γ ray being the reference line

	Energy	1959Bh95*	1961EI02	1962Sc04	1962Re03	1963Ba31	1969Gr32	1970FoZZ	1970Gr09	1970GrYP	1971Mo04	1977Ka14	Recommended
^{148}Pm													
$\gamma_{2,1}$	61.30												
$\gamma_{1,0}$	75.8	~4	2.4		1.1 0.4	1.7 0.5		7.3 0.8	1.1 0.2				
^{148}Sm													
$\gamma_{9,8}$	98.48	~4		9.1	3.2 0.4	3.4 0.8		3.3 1.7	3.5 0.5	6.4 0.7	4.29 0.12	2.78 0.05	3.33 0.29
$\gamma_{8,7}$	189.63	11	7.2	3.8	~1.0	1.5		1.2 0.1	1.1 0.2	1.6 0.1	1.39 0.05	1.24 0.03	1.31 0.07
$\gamma_{9,7}$	288.11				11 2		17	13.0 0.2	14 2	15 1	13.9 0.3	14.1 0.1	13.7 0.4
γ	295.06							0.08 0.01					
$\gamma_{9,6}$	299.1	20 5	16.9	14.0		16 3		0.20 0.02				0.1 0.02	0.15 0.05
$\gamma_{7,4}$	311.63				5 2		5	4.07 0.09	4.5 0.7	3.7 0.3	4.46 0.12	4.4 0.05	4.29 0.11
$\gamma_{8,5}$	362.09							0.19 0.02				0.2 0.02	0.195 0.014
$\gamma_{4,3}$	414.07	20 4	25.3	13.9	19 4	24 4	25	20.2 0.5	22 3	22.5 1.5	20.8 0.5	21.0 0.2	20.9 0.2
$\gamma_{4,2}$	432.78			13.9	8 3			5.7 0.1	5.8 0.9	6.2 0.7	6.35 0.20	6.0 0.1	5.9 0.1
$\gamma_{9,5}$	460.57							0.45 0.01				0.47 0.02	0.45 0.01
$\gamma_{8,4}$	501.26				8 3			7.5 0.02	8.1 1.2	7.9 0.8	7.74 0.19	7.6 0.1	7.51 0.02
γ	533.6					}105 15		0.32 0.02					
$\gamma_{1,0}$	550.27	100 20	120.5	100.0	99 5		120	106.2 3.2	109.5 16.4	112.6 4.6	106.1 1.9	106.6 0.8	106.5 0.7
$\gamma_{5,3}$	553.24							0.34 0.04				0.45 0.04	0.4 0.05
$\gamma_{5,2}$	571.95							0.24 0.01				0.24 0.01	0.24 0.007
$\gamma_{9,4}$	599.74						12	14.1 0.3	14 2	15.1 0.5	13.9 0.3	14.09 0.13	14.07 0.11
$\gamma_{2,1}$	611.26						6	6.6 0.1	7 1	7.0 0.3	6.29 0.22	6.16 0.1	6.4 0.2
$\gamma_{3,1}$	629.97	100 20	100	100.0	100	100	100	100.0 3.2	100 15	100	100 1.4	100 0	100
$\gamma_{6,3}$	714.7											0.051 0.006	0.051 0.006
$\gamma_{7,3}$	725.70	30 6	45.8	37.0	33 3	38 7	35	36.3 0.8	34.9 5.2	36.3 1.2	36.8 0.7	36.89 0.29	36.79 0.25
$\gamma_{8,3}$	915.33	23 5	25.3	20.8	20 2	22 4	23	20.4 0.8	22.1 3.3	22.0 1.7	22.1 0.6	19.29 0.2	20.4 0.5
$\gamma_{9,3}$	1013.81	25	25.3	23.1	21 2	21 4	21	21.4 0.5	23.2 3.5	19.5 1.8	22.9 0.6	22.79 0.19	22.6 0.23
$\gamma_{6,1}$	1344.6							0.06 0.01				0.066 0.005	0.0648 0.0045
γ	1465.12		2.4	4.9	1.7 0.3	2.5 1		1.5 0.1	1.7 0.3	2.1 0.2	1.43 0.16		1.59 0.1

* States intensities are "correct to 20%" above 400 keV, with larger uncertainties below 400 keV

Measurement conditions

The γ -ray intensities were measured with the detectors and under the timing conditions shown in Table 5.

Table 5: Conditions of the γ -ray intensity measurements

Reference	Detector	Timescale
1959Bh95	NaI	2 days to 3 months
1961El02	NaI	Up to 250 days
1962Sc04	NaI	At 24 hrs and “equilibrium”
1962Re03	NaI	“Immediately after irradiation” to “several months”
1963Ba31	NaI	2 days to 140 days
1969Gr32	GeLi	2-3 hours after irradiation
1970FoZZ	GeLi	Immediately after irradiation, then two and twelve weeks later
1970Gr09	GeLi	Made 24 measurements, but no timescale given
1970GrYP	GeLi	No timing details given
1971Mo04	GeLi	3 to 75 days
1977Ka14	GeLi	Four sources produced periodically over 10 month measuring period

The absolute intensities have been determined from placing the γ -rays into the decay scheme, and assuming the 550 keV γ -ray has a $P_{(\gamma+ce)}$ of 94.4 (5) %, i.e. it is emitted in every single β decay, so all higher levels decay through this level to the ground state, with no direct beta-feeding to the ground state. Assuming the transition is E2, then $\alpha_T = 0.009\ 98$ (14), gives $P_{(\gamma)} = 93.5$ (14) % for a relative intensity of 106.5 (7). Thus for the 629.95 keV γ -ray, with a relative intensity of 100, then $P_{(\gamma)} = 87.8$ (14) %, where the normalisation factor of 0.878 (14) is obtained from $93.5 \times 100/106.5$.

Multipolarities and Internal Conversion Coefficients

The γ -ray multipolarities and the mixing ratios have been taken from the adopted gamma data within the evaluation of 2000Bh03 based on 1961Br41, 1961Ha23 (measured conversion electrons and deduced multipolarities), 1962Re03, 1962Sc04, 1963Ba31 (measured conversion electrons and γ -ray directional correlations), 1968Wy02, 1977Ka14 (γ -ray directional correlations), and 1963Ku09, 1970GrYP (measured conversion electrons).

All internal conversion coefficients have been obtained using the BrIcc code (2008Ki07), as implemented in the SAISINUC evaluation tool (2008DuZX). In Table 6 these values are compared with available measured data.

Table 6: Comparison of BrIcc and measured K-shell internal conversion coefficients (α_K)

	Energy	1963Ba31		1970GrYP		BrIcc	
¹⁴⁸ Pm							
$\gamma_{1.0}$ ¹⁴⁸ Sm	75.8	2.9	0.4	2.5	0.8	2.9	0.4
$\gamma_{9.8}$	98.15	1.9	0.8	1.0	0.2	1.488	0.021
$\gamma_{8.7}$	189.61	0.17		0.11	0.04	0.176 9	0.025
$\gamma_{9.7}$	288.09	0.062	0.020	0.048	0.009	0.076 3	0.001 1
$\gamma_{7.4}$	311.62	0.012	0.004	0.011	0.002	0.011 41	0.000 16
$\gamma_{4.3}$	414.02	0.006 5	0.002 0			0.005 72	0.000 09
$\gamma_{4.2}$	432.71	0.017	0.005	0.017	0.003	0.015 44	0.000 22
$\gamma_{8.4}$	501.25	0.003 3	0.000 9	0.003 2	0.000 8	0.003 69	0.000 07
$\gamma_{1.0}$	550.22			0.007 9	0.000 6	0.008 25	0.000 12
$\gamma_{9.4}$	599.69			0.002 0	0.000 7	0.002 49	0.000 04
$\gamma_{2.1}$	611.23	0.002 4	0.000 8	0.002 5	0.000 8	0.002 37	0.000 04
$\gamma_{3.1}$	629.95	0.006 0	0.001 2			0.005 91	0.000 09
$\gamma_{7.3}$	725.67	0.003 8	0.000 7	0.004 6	0.000 4	0.004 24	0.000 06
$\gamma_{8.3}$	915.31	0.002 5	0.000 5	0.002 5	0.000 4	0.002 54	0.000 04
$\gamma_{9.3}$	1013.8	0.002 0	0.000 4	0.002 4	0.000 4	0.002 06	0.000 04

2.4. Beta Transitions

Energies and Emission Probabilities

Beta-particle energies were determined from the difference between the adopted Q-value and the final level energies of daughter ^{148}Sm . Beta-particle emission probabilities were determined by balancing the proposed decay scheme through consideration of the $\beta\gamma$ -population and γ -depopulation of these same levels. Only the 2nd level has no beta-feeding based on published data. A comparison with measured energies and emission intensities is presented in Table 7. Note that all experimental measurements were made relative and only appear absolute after consideration of the decay scheme.

Table 7: Measured intensities and recommended absolute intensities of β -emissions from the decay of ^{148m}Pm

	1962Sc04		1962Re03		1963Ba31†		Recommended	
	E (keV)	I_{β} (%)	E (keV)	I_{β} (%)	E (keV)	I_{β} (%)	E (keV)	I_{β} (%)
$\beta_{0,9}$	390 (30)	45			400 (30)	48 (10)	414 (6)	54.0 (9)
$\beta_{0,8}$	490 (30)	13	510	37	500 (30)	28 (10)	513 (6)	18.1 (9)
	560 (25)	16	610	19				
$\beta_{0,7}$	680 (20)	18	680	8	690 (30)	16 (5)	702 (6)	21.8 (7)
			800	24				
$\beta_{0,4}$							1014 (6)	0.93 (45)

† Same data as published in 1963Ew01 and 1963Ba06 by the same authors

2.5. Conversion electrons

The energies of the internal conversion electrons have been deduced using γ -ray energies and electron binding energies from 1977La19 and 1996FiZX. Absolute conversion electron emission intensities have been calculated by using γ -ray internal conversion coefficients and absolute emission intensities.

3 Atomic Data

The recommended fluorescence yields shown in Table 8 were obtained from 1996Sc06.

Table 8: Recommended atomic data

Sm		Pm	
ω_K	0.926 (4)	ω_K	0.922 (4)
ω_L	0.158 (6)	ω_L	0.148 (6)
η_{KL}	0.857 (4)	η_{KL}	0.861 (4)

The energies of the K-Auger electrons were obtained from 1998ScZM, whereas the energies and the yield of Sm X-rays were obtained from 1999ScZX (data stored in SAISINUC). The X-ray and Auger electron emission probabilities have been calculated using the computer program EMISSION (v3.10, 28-Jan-2003) described in 2000Sc47.

References

- 1951Fo** R.L.Folger, P.C.Stevenson, G.T.Seaborg, Report UCRL-1195, Univ. California (1951) 22
[Half-life, Beta and gamma transition energies]
- 1952Ki25 V.Kistiakowsky, Phys. Rev. 87 (1952) 859
[Half-life, Beta and gamma transition energies]
- 1952Lo01 J.K.Long, M.L.Pool, Phys. Rev. 85 (1952) 137
[Half-life, Beta and gamma transition energies]
- 1955Fo18 R.L.Folger, P.C.Stevenson, G.T.Seaborg, Phys. Rev. 98 (1955) 107
[Half-life, Beta and gamma transition energies, IT branching fraction]
- 1955He64 N.P.Heydenburg, G.M.Temmer, Phys. Rev. 100 (1955) 150
[Sm-148 levels]
- 1955Ma77 H.Mark, G.T.Paulissen, Phys. Rev. 100 (1955) 813
[Sm-148 levels]
- 1959Bh95 S.K.Bhattacharjee, B.Sahai, C.V.K.Baba, Nucl. Phys. 12 (1959) 356
[Half-life, Beta and gamma transition energies and intensities, IT branching fraction, Sm-148 levels]
- 1959Ei31 J.A.Eisele, Thesis, Ohio State Univ. (1959); Diss. Abst. Int. 20 (1960) 3794
[Half-life, Beta and gamma transition energies, Sm-148 levels]
- 1960Sc09 C.F.Schwerdtfeger, E.G.Funk, J.W.Mihelich, Bull. Am. Phys. Soc. 5 (1960) 425, P
[Half-life, Beta and gamma transition energies, Sm-148 levels]
- 1960Sc12 R.P.Schuman, J.R.Berreth, R.L.Heath, C.W.Reich, Bull. Am. Phys. Soc. 5 (1960) 494, C
[Beta and gamma transition energies and intensities, IT branching fraction, Sm-148 levels]
- 1961Br41 M.K.Brice, C.W.Reich, R.G.Helmer, Report IDO-16710 (1961)
[Half-life, Gamma transition energies and intensities, ICCs and multipolarities, IT branching fraction]
- 1961El02 J.S.Eldridge, W.S.Lyon, Nucl. Phys. 23 (1961) 131
[Half-life, Beta and gamma transition energies and intensities, IT branching fraction]
- 1961Ha23 B.Harmatz, T.H.Handley, J.W.Mihelich, Phys. Rev. 123 (1961) 1758
[Half-life, Gamma transition intensity, ICCs]
- 1962Gr34 R.W.Grant, D.A.Shirley, Report UCRL-10624, Univ. California (1962)
[Pm-148 and Sm-148 level spins]
- 1962Re03 C.W.Reich, R.P.Schuman, J.R.Berreth, M.K.Brice, R.L.Heath, Phys. Rev. 127 (1962) 192
[Half-life, Beta and gamma transition energies and intensities, IT branching fraction, Conversion electrons, Sm-148 levels]
- 1962Sc04 C.F.Schwerdtfeger, E.G.Funk Jr, J.W.Mihelich, Phys. Rev. 125 (1962) 1641
[Half-life, Beta and gamma transition energies and intensities, IT branching fraction, Conversion electrons, Sm-148 levels]
- 1963Ba06 C.V.K.Baba, G.T.Ewan, J.F.Suarez, Phys. Lett. 3 (1963) 232
[Pm-148 and Sm-148 level spins]
- 1963Ba31 C.V.K.Baba, G.T.Ewan, J.F.Suarez, Nucl. Phys. 43 (1963) 264
[Beta and gamma transition energies and intensities, IT branching fraction, Conversion electrons, Pm-148 and Sm-148 levels]

- 1963Ba32 C.V.K.Baba, G.T.Ewan, J.F.Suarez, Nucl. Phys. 43 (1963) 285
[Sm-148 levels]
- 1963Ew01 G.T.Ewan, C.V.K.Baba, J.F.Suarez, Bull. Am. Phys. Soc. 8 (1963) 73, VA
[Beta transition energies, intensities and shape, Sm-148 levels]
- 1963Gr10 R.W.Grant, D.A.Shirley, Phys. Rev. 130 (1963) 1100
[Pm-148 and Sm-148 level spins]
- 1963Ku09 T.J.Kurey Jr, R.R.Roy, Nucl. Phys. 44 (1963) 670
[Conversion electrons, Multipolarities, Sm-148 levels]
- 1967Cl05 J.E.Cline, R.L.Heath, Report IDO-17222 (1967)
[Gamma transition energies and intensities]
- 1968Wy02 L.D.Wyly, E.T.Patronis Jr, C.H.Braden, Phys. Rev. 172 (1968) 1153
[Sm-148 level spins]
- 1969Gr32 Z.G.Gritchenko, T.P.Makarova, Y.T.Oganesyan, Y.E.Penionzhkevich, A.V.Stepanov,
Yadern. Fiz. 10 (1969) 929; Soviet J. Nucl. Phys. (English translation) 10 (1970) 536
[Gamma transition energies and intensities]
- 1970FoZZ J.W.Ford Jr, Thesis, Vanderbilt Univ. (1970); Diss. Abst. Int. 31B (1970) 3631
[Gamma transition energies and absolute intensities, Multipolarities, Sm-148 levels
and spins]
- 1970Gr09 W.M.Greenberg, H.J.Fischbeck, Z. Phys.233 (1970) 391
[Gamma transition energies and intensities]
- 1970GrYP E.P.Grigorev, A.V.Zolotavin, V.O.Segreev, M.I.Sovtsov, Program and Theses, Proc.
20th Ann. Conf. Nucl. Spectrosc. At. Nuclei, Leningrad (in Russian) (1970) 100
[Conversion electrons, Multipolarities, Pm-148 and Sm-148 levels]
- 1971BaZW S.Baba, H.Baba, H.Umezawa, T.Suzuki, T.Sato, H.Natsume, Report JAERI-1211
(1971)
[Half-life]
- 1971Mo04 R.S.Mowatt, W.H.Walker, Can. J. Phys. 49 (1971) 108
[Half-life, Beta and gamma transition energies and intensities]
- 1971Wa05 F.W.Walker, T.A.Devito, F.M.Rourke, H.M.Eiland, J. Inorg. Nucl. Chem. 33(1971)
1208
[Half-life]
- 1974DuZO D.R.Dunn, A.V.Ramayya, J.W.Ford Jr, J.H.Hamilton, W.Lourens, J.J.Pinajian, Bull.
Am. Phys. Soc. 19 (1974) 1124, FB8
[Gamma transition energies, Sm-148 levels]
- 1977Ka14 C.A.Kalfas, J. Phys. (London) G3 (1977) 929
[Gamma transition energies and intensities, Multipolarities, Mixing ratios, Sm-148
levels and spins]
- 1977La19 F.P.Larkins, At. Data Nucl. Data Tables 20 (1977) 311
[Electron Binding Energies]
- 1979Dz07 B.S.Dzhelepov, S.A.Shestopalova, Izv. Akad. Nauk SSSR Ser. Fiz. 43 (1979) 2261;
Bull. Acad. Sci. USSR Phys. Ser. (English translation) 43 (1979) 16
[Gamma transition intensity ratio]
- 1984LaZZ V.Lakshminarayana, B.Van Nooijen, W.Lourens, A.V.Ramayya, J.H.Hamilton,
J.W.Ford Jr, D.R.Dunn, J.J.Pinajian, Priv. Comm. to NNDC (unpublished report)
(1984)
[Gamma transition energies and intensities, Multipolarities, Sm-148 levels and spins]

- 1988No02 E.B.Norman, K.T.Lesko, A.E.Champagne, Phys. Rev. C37 (1988) 860
[Pm-148 level energies and spins]
- 1989Le01 K.T.Lesko, E.B.Norman, R.-M.Larimer, J.C.Bacelar, E.M.Beck, Phys. Rev. C39
(1989) 619
[Pm-148 level energies and spins]
- 1996FiZX R.B.Firestone, Table of Isotopes 8th Ed., John Wiley and Sons Inc. 2 (1996)
[Electron Binding energies]
- 1996Sc06 E.Schönfeld, H.Janssen, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527
[Atomic Data]
- 1998ScZM E.Schönfeld, G.Rodloff, PTB Report 6.11-98-1 6.11 (1998) 1
[Auger electrons]
- 1999ScZX E.Schönfeld, G.Rodloff, PTB Report 6.11-1999-1 6.11 (1999) 1
[K X-rays]
- 2000Bh03 M.R.Bhat, Nucl. Data Sheets 89 (2000) 797
[Sm-148 levels, Multipolarities, Mixing ratios]
- 2000Sc47 E.Schönfeld, H.Janssen, Appl. Radiat. Isot. 52 (2000) 595
[X-ray and Auger Electron emission probabilities]
- 2008DuZX C.Dulieu, M.M.Bé, V.Chisté, Proc. Int. Conf. on Nuclear Data for Science and
Technology, 22-27 April 2007, Nice, France (2008) 97
[SAISINUC software]
- 2008Ki07 T.Kibédi, T.W.Burrows, M.B.Trzhaskovskaya, P.M.Davidson, C.W.Nestor Jr, Nucl.
Instrum. Methods Phys. Res. A589 (2008) 202
[Theoretical ICCs]
- 2012Wa38 M.Wang, G.Audi, A.H.Wapstra, F.G.Kondev, M.MacCormick, X.Xu, B.Pfeiffer,
Chin. Phys. C36 (2012) 1603
[Q]

¹⁵¹Sm - Comments on evaluation of decay data
by M.M. Bé

This evaluation was completed in November 2014 with literature available at this date included.

Decay Scheme

¹⁵¹Sm decays by β^- disintegrations to the first excited level and the ground state of ¹⁵¹Eu. Level energy, spin and parities were taken from 2009Si01.

Q^{β^-} values for ¹⁵¹Sm decay was adopted from Wang *et al.* (2012Wa38).

Half-life

The experimental half-life values used to calculate the recommended value are listed in the following table:

References	$T_{1/2}$ (a)	$T_{1/2}$ uc (a)	Comments
1950Inghram01	122		No uncertainty, omitted
1952Rutledge10	> 20		
1952Karraker26	73	33	+25 -14 ; omitted
1955Melaika52	93		No uncertainty, omitted
1965Flynn02	87	10	
1968Reynolds04	93	8,6	
2009Ming He22	96,6	2,4	
2015Bé**	94,6	0,6	
Crit (χ^2)	3,8		
$\chi^2 / (n-1)$	0,5		
UWM	92,80		
WM	94,68		
uc(WM)int. :	0,58		
uc(WM)ext. :	0,38		
Adopted	94,7	0,6	

Only the four latest values are used in the statistical process. The set of the four data is consistent; the weighted mean is adopted with the internal uncertainty.

β^- transitions

Two beta emissions were measured by Achor *et al.* (1959) with energy 75,9 keV and 54,2 keV and respective intensity of 98,3 (3) % and 1,7 (3) %. Both transitions were found either first forbidden non-unique shape, or allowed. This latter point was confirmed by a series of measurements described in 2015Be**.

The measured intensity values of the 54 keV β^- transition are:

Reference	I β (%)	uc	Comments
Achor (1959Ac28)	0,9	0,3	Re calculated by Freedman
Freedman (1975Fr01)	0,85	0,06	Mean of 0,86 and 0,84 ; uc from 1975Fr01
Simple mean	0,87	0,06	

The simple mean of the results of 1959Ac28 and 1975Fr01 is 0,87 (6) %, where the uncertainty is the lowest experimental uncertainty.

These results are in good agreement with the adopted intensity of 0,927 (41) % which was derived from the α_T coefficient and the γ intensity of the 21,5 keV emission (see the following paragraphs).

Then, the intensity of the transition which populates the ground state is 99,073 (41) %.

Both values were rounded to 0,93 (4) % and 99,07 (4) % respectively.

Gamma transition

Multipolarity and conversion coefficient of the 21,5 keV transition were determined by:

Reference		uc	Multipolarity
Achor (1959Ac28)	$\alpha_L = 20$	4	M1
Avotina (1966Av05 and 1968Gr25)	$\delta = 0,035$	0,007	M1 + 0,12 (3) % E2
Grigorev (1968Gr25)	$\alpha_L = 20$	5	
Antman (1970An17)	$\delta^2 = 0,00088$	0,00007	M1 + 0,088 (7) % E2
Ford (1970Fo02)	$\alpha_L = 24$	4	M1 + < 6 % E2
Freedman (1975Fr01)	$\delta = 0,022$		M1 \leq 0,05 % E2
Veluri(1977Ve01)	$\alpha_T = 27,5$	1,5	
Artamonova (1981Ar17)	$\alpha_L = 23,9$	5,0	M1 + 0,083 (5) % E2
Adopted	$\delta = 0,029$ (2)		M1 + 0,085 (5) % E2
BrIcc theory FO	$\alpha_T = 25,9$	0,4	M1
BrIcc theory FO	$\alpha_T = 27,6$	0,5	M1 + 0,085 (5) % E2
Adopted	$\alpha_T = 27,6$	0,5	M1 + 0,085 (5) % E2

The adopted multipolarity is the weighted mean of the values of Avotina (1968Gr25), Antman (1970An17) and Artamonova (1981Ar17) with the lowest experimental uncertainty.

The corresponding internal conversion coefficients are the theoretical values interpolated by the BrIcc program (2008Ki07) from the tables of Band *et al.* (2002Ba85), accepting the “frozen orbital (no hole)” approximation.

The theoretical and adopted α_T is in good agreement with the experimental result of Veluri (1977Ve01) and the α_L theoretical value of 21,7 (4) is in agreement, within the uncertainty limits, with the results of Achor (1959Ac28), Grigorev (1968Gr25), Ford (1970Fo02) and Artamonova (1981Ar17).

21,5 keV γ ray intensity

- From the β emission intensity to the ground state level (0,87 (6) %) and the adopted total conversion coefficient (27,6 (5)) the 21,5 keV γ ray intensity is 0,0304 (20) %.
- From the publication of Laird (1980La02) a value of 0,0323 (21) % can be deduced.
- A direct measurement of Shen (2011Sh37) leads to 0,0324 (13) %.

The adopted value is the direct measurement result of Shen (2011Sh37), *i.e.* $I_\gamma = 0,0324$ (13) %. This value is in good agreement with the two indirect results, within the uncertainty limits.

From this latter result and the α_T coefficient, the intensity of the beta transition to the 21,5 keV level is deduced to be 0,927 (41) %.

Atomic Data

Several studies were conducted in order to measure the internal ionisation probabilities in the K and L shells following the ¹⁵¹Sm beta decay. Various quantities were determined, they are summarized below.

The measured relative intensities of the X photons emitted in the ¹⁵¹Sm beta decay are listed in the table below and compared with the values derived from the decay scheme parameters with $I_\gamma = 0,0324$ (13) %.

x-rays	Unus (1981Un02)	Campbell (1971Ca44)	Deduced values
L ℓ	0,073 (4)		0,0434 (28)
L α , η	1,73 (9)		1,08 (6)
L β	2,65 (13)		2,15 (10)
L γ	0,636 (32)		0,475 (23)
γ 21,5 keV	1	1	1
K α	4,0 (2) 10^{-3}	5,8 (2) 10^{-3}	
K β	1,0 (2) 10^{-3}		

Intensity of L x-rays due to internal ionisation per 100 β decays

Reference	I Lx (%)
Unus (1981Un02)	4,7 (5) 10^{-2}
Freedman (1975Fr01)	5,4 (10) 10^{-2}

Internal Ionisation Probabilities

Reference	PK (%)	PL (%)
Unus (1981Un02)	1,59 (13) 10^{-4}	31 (3) 10^{-2}
Laird (1980La02)	2,4 (3) 10^{-4}	
Law (1975La20)	2,3 (3) 10^{-4}	30 (6) 10^{-2}
Freedman (1975Fr01)	2,4 (3) 10^{-4}	
Campbell (1971Ca44)	0,041 (6) 10^{-4}	

The internal ionisation probability in the K shell is 2,0 (2) 10^{-4} % and 31 (3) 10^{-2} % in the L shell.

References

- 1950In01** M.G.Inghram, R.J.Hayden, D.C.Hess. Phys.Rev. 79, 271 (1950) [T_{1/2}]
U²³⁵ Fission Yields in the Rare Earth Region
- 1952Ka26** D.G.Karraker, R.J.Hayden, M.G.Inghram. Phys.Rev. 87, 901 (1952) [T_{1/2}]
Half-Lives of Eu¹⁵², Eu¹⁵⁴ and Sm¹⁵¹
- 1952Ru10** W.C.Rutledge, J.M.Cork, S.B.Burson. Phys.Rev. 86, 775 (1952) [T_{1/2}]
Gamma-Rays Associated with Selected Neutron-Induced Radioactivities
- 1955Me52** E.A.Melaika, M.J.Parker, J.A.Petruska, R.H.Tomlinson. Can.J.Chem. 33, 830 (1955) [T_{1/2}]
The Relative Abundances of Neodymium and Samarium Isotopes in the Thermal Neutron Fission of U²³⁵ and U²³³
- 1959Ac28** W.T.Achor, W.E.Phillips, J.I.Hopkins, S.K.Haynes. Phys.Rev. 114, 137 (1959)
Low-Intensity Conversion Lines from Sm¹⁵¹ and Sn¹¹³
- 1965F102** K.F.Flynn, L.E.Glendenin, E.P.Steinberg. Nucl.Sci.Eng. 22, 416 (1965) [T_{1/2}]
Half-Life Determinations by Direct Decay
- 1966Av05** M.P.Avotina, E.P.Grigorev, A.V.Zolotavin, V.O.Sergeev, J.Vrzal, J.Liptak, N.A.Lebedev, Y.Urbanets. Izv.Akad.Nauk SSSR, Ser.Fiz. 30, 1304 (1966); Bull. Acad. Sci. USSR, Phys.Ser. 30, 1362 (1967)
Decay of Gd¹⁴⁹, Eu¹⁴⁹ and Gd¹⁵¹
- 1968Re04** S.A.Reynolds, J.F.Emery, E.I.Wyatt. Nucl.Sci.Eng. 32, 46 (1968) [T_{1/2}]
Half-Lives of Radionuclides - III
- 1968Gr25** E.P.Grigorev, A.V.Zolotavin, V.O.Sergeev, M.I.Sovtsov, J.Vrzal, N.A.Lebedev, J.Liptak, J.Urbanets, P.P.Dmitriev, N.N.Krasnov, Y.G.Sevastyanov. Izv.Akad.Nauk SSSR, Ser.Fiz. 32, 786 (1968); Bull. Acad. Sci.USSR, Phys.Ser. 32, 723 (1969)
Decay of ¹⁵¹Gd
- 1970An17** S.Antman, H.Pettersson, Z.Zehlev, I.Adam. Z.Phys. 237, 285 (1970)
High Resolution L- and M-Subshell Internal Conversion Studies of Low Energy Transitions in ¹⁴⁹Sm and ¹⁵¹Eu
- 1970Fo02** J.W.Ford, A.V.Ramayya, J.J.Pinajian. Nucl.Phys. A146 (1970) 397
The Decay of ¹⁵¹Gd to the Levels of ¹⁵¹Eu
- 1971Ca44** J.L.Campbell, L.A.McNelles, J.Law. Can.J.Phys. 49, 3142 (1971)
K-Shell Auto-Ionization in the Beta Decays of ¹⁵¹Sm, ¹⁶⁹Er, and ²¹⁰Bi
- 1975Fr01** M.S.Freedman, D.A.Beery. Phys.Rev.Lett. 34, 406 (1975)
K and L Shakeoff Probability in ¹⁵¹Sm
- 1975La20** J.Law, J.L.Campbell. Phys.Rev. C12, 984 (1975); Erratum Phys.Rev. C14, 2347 (1976)
Mechanisms Contributing to K Electron Ejection in Nuclear β Decay
- 1977Ve01** V.R.Veluri and P. Venugopala Rao. Z. Phys. A280 (1977) 317 [α_T]
- 1980La02** C.E.Laird, P.C.Hummel, H.-C.Liu, Phys.Rev. C21, 723 (1980)
K-Shell Autoionization Accompanying the Beta Decays of ⁹⁹Tc, ¹⁴⁷Pm, ¹⁵¹Sm, and ²⁰⁴Tl
- 1981Ar17** K.P.Artamonova, N.B.Grachev, E.P.Grigorev, A.V.Zolotavin, V.O.Sergeev. Bull. Acad. Sci. USSR, Phys. Ser. 45,1 (1981) 93
Penetration effect for γ-Transitions in ¹⁴⁹Sm and ¹⁵¹Eu
- 1981Un02** I.J.Unus, P.A.Indira, P.Venugopala Rao. J.Phys.(London) G7, 1683 (1981)
Internal Ionisation Probability in the Beta Decay of ¹⁵¹Sm
- 2002Ba85** I.M.Band, M.B.Trzhaskovskaya, C.W.Nestor, Jr, P.O.Tikkanen, S.Raman. At. Data Nucl. Data Tables 81, 1 (2002) [ICC]
Dirac-Fock Internal Conversion Coefficients

- 2008Ki07** T.Kibédi, T.W.Burrows, M.B. Trzhaskovskaya, P.M.Davidson, C.W.Nestor, Jr. Nucl. Instrum. Methods Phys.Res. A589 (2008) 202 [ICC]
Evaluation of theoretical conversion coefficients using BrIcc
- 2009He22** M.He, H.Shen, G.Shi, X.Yin, W.Tian, S.Jiang. Phys.Rev. C 80, 064305 (2009) [T_{1/2}]
Half-life of ¹⁵¹Sm remeasured
- 2009Si01** B.Singh. Nucl.Data Sheets 110, 1 (2009)
Nuclear Data Sheets for A = 151
- 2011Sh37** H.Shen, M.He, X.d.Ruan, K.Dong, S.Jiang. Phys.Rev. C 84, 054307 (2011)
Determination of the emission probability of the 21.54 keV γ ray in the decay of ¹⁵¹Sm
- 2012Wa38** M.Wang, G.Audi, A.H.Wapstra, F.G.Kondev, M.MacCormick, X.Xu, B.Pfeiffer. Chin.Phys. C 36, 1603 (2012) [Q]
The AME2012 atomic mass evaluation (II). Tables, graphs and references
- 2015Bé**** M.M. Bé *et al.* Accepted for publication in RadioChimica Acta
Determination of ¹⁵¹Sm half-life

¹⁵²Eu – Comments on evaluation of decay data

Vito R. Vanin and Ruy M. de Castro

Physics Institute, University of São Paulo, Brazil

Edgardo Browne

*Lawrence Berkeley National Laboratory, Berkeley, California***Evaluation Procedures**

We used the *Limitation of Relative Statistical Weights* (LWM) method (1985ZiZY, 1992Ra08) for averaging quantities throughout this evaluation. This method provides a uniform approach for the analysis for discrepant data.

Decay Scheme

¹⁵²Eu decays by electron capture (EC) to ¹⁵²Sm, and by β^- to ¹⁵²Gd. Only excited levels are populated in the daughter nuclei since decay to the respective ground states are highly hindered by spin selection rules. Therefore, we used the sum of the total γ -ray transition emission probabilities (photons + electrons) to the ground states of ¹⁵²Sm and ¹⁵²Gd to normalize the decay scheme of ¹⁵²Eu. We have deduced the following branchings: 72.1(3)% (EC), and 27.9(3)% (β^-). This normalization is virtually the same as that based on the measurement of the absolute γ -ray emission probabilities (See **Gamma Rays**).

Nuclear Data

We have considered the following measured values of the half-life of ¹⁵²Eu for deducing a recommended value.

1.	4934.1 (23) d	2004Sc04	Duration of measurement: about 26 years
2.	4936.6 (20) d	1998Si12	Duration of measurement: 20 years
3.	4948 (7) d	1997Ma75	Duration of measurement: about 2 years
4.	4945.5 (23) d	1992Un01	Duration of measurement: 13.5 years
5.	4943 (4) d	1986Wo05	
6.	4792(37) d	1983Ba29	
7.	4939 (6) d	1983Wa26	
8.	4892.3 (82) d	1980RuZX	
9.	4785 (19) d	1978La21	
10.	4821 (110) d	1972Em01	

Our recommended value of 4939 (6) d (or 13.522 (16) a) is a weighted average (LWM) ($\chi^2/\nu=12$) of the results from 2004Sc04, 1997Ma75, 1992Un01, 1986Wo05, and 1980RuZX. Values given by 1978La21, 1972Em01, and 1983Ba29 have not been included because they significantly disagree with most of the other results, suggesting that they may have been affected by systematic uncertainties. 1983Wa26 and 1998Si12 have been superseded by 2004Sc04 (same research groups, PTB).

Electron Capture, Positrons (β^+), and β^- Transitions

EC and positron transition energies to levels in ¹⁵²Sm have been deduced from $Q(\text{EC}) = 1874.3$ (7) keV (1995Au04) and the individual level energies. Transition probabilities (P_{EC}) are from γ -ray transition probability balance at each level. They are given as branchings ($P_{\text{EC}} \times 100$) in Sections 2.1 – 2.3. Fractional atomic sub-shell electron-capture probabilities (i.e., P_K, P_L, P_M, P_N) are theoretical values (1998Sc28) calculated with the computer program EC-CAPTURE [1].

Positrons are energetically possible and allowed by spin selection rules to the 121- and 366-keV levels only. Their transition probabilities, presented here as branchings ($P_{\beta^+} \times 100$), have been deduced from theoretical β^+/EC ratios (1957Zw01).

β^- endpoint energies for the decay to levels in ¹⁵²Gd have been deduced from $Q(\beta^-) = 1818.8$ (11) keV (1995Au04). Their transition probabilities, presented here as branchings ($P_{\beta^-} \times 100$), have been deduced from γ -ray transition probability balance at each level.

Gamma Rays

Energies. The precise energies of strong γ rays given here are from 2000He14. These values are based on a revised energy scale that uses the new fundamental constants and wave lengths deduced from an updated value of the lattice spacing in Si crystals (1987Co39). All other (less precise) energies are values adjusted to the new energy scale and recommended in 1996Ar09 evaluation.

Emission Probabilities. For a γ -ray transition, its absolute transition probability (photons + electrons) is given by $P_\gamma(1 + \alpha) \times 100$, where P_γ is the absolute γ -ray emission intensity, and α , its theoretical (1978Ro22, [4]) conversion coefficient. We have deduced the P_γ values used here as follows:

1. By averaging (LWM) the experimental relative emission intensities reported by 1970No06, 1970Ri19, 1971Ba63, 1972Ba05, 1977Ge12, 1980Sh15, 1984Iw03, 1986Me10, 1989Da12, 1990Me15, 1990St02, 1992Ya12, 1993Ka30, 1998Hw07, and from the fourteen measurements (ICRM01, ICRM02, ICRM08, ICRM10, ICRM12, ICRM15, ICRM16, ICRM17, ICRM18, ICRM20, ICRM25, ICRM27, ICRM28, and ICRM29) of the study participants [5] from the International Committee on Radioactivity Measurements (ICRM), which 1991BaZS considered reliable. These data are presented in Table 1 and Table 2.
2. By normalizing the above mentioned relative emission intensities to absolute values. We normalized these scales by using $P_\gamma(1408) = 0.2085$ (8), which was determined from an inter-comparison of measured absolute emission intensities produced by participants from various laboratories and coordinated by the ICRM [5]. This value agrees very well with $P_\gamma(1408) = 0.2086$ (21), deduced by evaluators from the sum of the relative γ -ray transition probabilities (photons + electrons) to the respective ground states of ¹⁵²Sm and ¹⁵²Gd. The larger uncertainty in the latter value is due mostly to that in the conversion coefficient of the 121-keV γ -ray (taken as 3%). We used 47.46 (20) for the relative intensity of the 1086-keV γ ray that de-excites the 1086-keV level in ¹⁵²Sm. We deduced this value from our recommended relative emission intensity of 48.63 (20) for the 1086-“doublet” (See Table 2) and subtracting 1.17 (4) for the contribution of the 1084-keV γ ray (1990Me15). The excellent agreement between these two normalizations confirms the completeness and self-consistency of the ¹⁵²Eu decay scheme and the good quality of our recommended data. We have preferred not to statistically combine these normalizations because of the correlations that exist between them. Absolute γ -ray emission intensities (P_γ) are given in Section 4.1.

Conversion Coefficients. Values given in Section 2.3 are the result of theoretical calculations (1978Ro22, [4]), interpolated for the recommended transition energies presented here, and for adopted multiplicities and mixing ratios from the 1996Ar09 evaluation, uncertainties have been taken being 3 %. For transitions with E0 multipolarity, the adopted values are derived from experiments.

Atomic Data

X-Rays. X-ray energies and relative emission probabilities are from Schönfeld and Rodloff [6]. Absolute X-ray emission probabilities have been calculated with the computer program EMISSION [2] using absolute γ -ray emission probabilities from Section 4.1, theoretical conversion coefficients (1978Ro22) from Section 2.3, and fluorescence yields from 1996Sc06. These calculated X-ray emission probabilities agree well with the experimental results shown in Table 2, and thus support the correctness of our recommended γ -ray data and the self-consistency of the ¹⁵²Eu decay scheme.

Electron Emission

Conversion-electron energies are from γ -ray energies given in Section 4.2 and the atomic binding energies reported by Larkins [7]. Absolute electron emission intensities are from γ -ray emission probabilities given in Section 4.1, and the theoretical (1978Ro22) conversion coefficients presented in Section 2.3.

Energies of K-Auger electrons are from Schönfeld and Rodloff [8]. Absolute emission intensities of Auger electrons are values calculated with the computer program EMISSION [2] using absolute γ -ray emission intensities from Section 4.2, theoretical conversion coefficients (1978Ro22) given in Section 2.3, and the electron-capture probabilities presented in Section 2.1. The same emission probabilities, but renormalized to a scale where $P_{KLL} = 1.0$, are given as relative emission probabilities in Section 3.2.

Total Average Radiation Energy

We show below the total average radiation energy released (by β^- , β^+ , neutrinos, γ rays, atomic electrons, and nuclear recoil) in the electron-capture and β^- decay of ¹⁵²Eu, as well as the total decay energies from mass differences, Q-values, and decay branchings (1995Au04).

	Total Average Radiation Energy* (keV)	Total Decay Energy ^{&} (Q x branching) (keV)
¹⁵² Eu EC decay	1345 (18)	1351 (6)
¹⁵² Eu β^- decay	508 (2)	507 (5)

* Calculated with the computer program RADLST [3], and using the recommended radiation data given in this evaluation.

[&] Q-values (Q(EC) and Q(β^-)) are from 1995Au04. Branchings are from this evaluation.

The agreement between these values confirms the quality, completeness, and self-consistency of the ¹⁵²Eu decay scheme presented in this evaluation.

References

- [1] E. Schönfeld, F.Y. Chu, E. Browne,
EC-CAPTURE, a computer program to calculate electron -capture probabilities to atomic
sub-shells, 1998.
[P_K, P_L, P_M, P_N]
- [2] H. Janßen and E. Schönfeld,
EMISSION, a computer program to calculate X-ray and Auger -electron emission
probabilities, 1998.
[X-ray, Auger-electron probabilities]
- [3] Thomas W. Burrows,
The Program RADLST, Report BNL-NCS-52142, February 1988.
[Average radiation energy]
- [4] E. Yakusev and N. Coursol,
ICC, a computer program to interpolate internal conversion coefficients, 1998.
[Theor. internal conversion coefficients]
- [5] K. Debertin, Nucl. Instrum. Methods **158**, 479 (1979);
K. Debertin, Report PTB-Ra-7 (and Report ICRM-S-3) (1978);
K. Debertin, Preliminary Summary communicated to ICRM study participants (1977);
K. Debertin, private communication (1987). Reference quoted in 91BaZS.
[Rel. P_γ]
- [6] E. Schönfeld, G. Rodloff,
Report PTB-6.11-1999-1, February 1999
[X-ray energies, P_{KX}]
- [7] F. B. Larkins,
At. Data and Nucl. Data Tables **20**, 313 (1977).
[Atom. electron binding energies]
- [8] E. Schönfeld, G. Rodloff,
Report PTB-6.11-98-1, October 1998.
[X rays, ω_K]
- 1957Zw01 - P. F. Zweifel, Phys. Rev. **107**, 329 (1957). [Theor. β⁺/EC ratios]
- 1970No06 - A. Notea, E. Elias, Nucl. Instrum. Methods **86**, 269 (1970). [Rel. P_γ]
- 1970Ri19 - L. L. Riedinger, N. R. Johnson, J. H. Hamilton, Phys. Rev. C **2**, 2358 (1970).
[Rel. P_γ]
- 1971Ba63 - J. Barrette, M. Barrette, A. Boutard, G. Lamoureux, S. Monaro, S. Markiza, Can. J.
Phys. **49**, 2462 (1971). [Rel. P_γ]
- 1972Ba05 - K. R. Baker, J. H. Hamilton, A. V. Ramayya, Z. Phys. **256**, 387 (1972). [Rel. P_γ]
- 1972Em01 - Emery J. F., Reynolds S. A., Wyatt E. I., Gleason G. I., Nucl. Sci. Eng. **48**, 319
(1972). [T_{1/2}]
- 1977Ge12 - R. J. Gehrke, R. G. Helmer, R. C. Greenwood, Nucl. Instrum. Methods **147**, 405
(1977) [Rel. P_γ]
- 1978La21 - Lagoutine F., Legrand J., Bac C., Int. J. Appl. Radiat. Isot. **29**, 269 (1978) [T_{1/2}]
- 1978Ro22 - F. Rösel, H. M. Friess, K. Alder, H. C. Pauli, At. Data Nucl. Data Tables **21**, 92
(1978). [Theor. ICC]
- 1980RuZX - Rutledge A. R., Smith L. V., Merritt J. S., NBS Special Publication 626, 5 (1982).
Compilation of work published in AECL Reports 3668 (1970), 4205 (1972), 5546 (1976), 5802
(1977), 6788 (1980). Quoted in 91BaZS. [T_{1/2}]

- 1980Sh15 - A. K. Sharma, R. Kaur, H. R. Verma, P. N. Trehan, J. Phys. Soc. Jpn. **48**, 1407 (1980). [Rel. P_γ]
- 1983Ba29 - Baba S., Ichikawa K., Gunji K., Sekine T., Baba H., Komori T., Int. J. Appl. Radiat. Isot. **34**, 891 (1983). [$T_{1/2}$]
- 1983Wa26 - Walz K., Debertain K., Schraeder H., Int. J. Appl. Radiat. Isot. **34**, 1191 (1983). [$T_{1/2}$]
- 1984Iw03 - Y. Iwata, M. Yasuhara, K. Maeda, Y. Yoshizawa, Nucl. Instrum. Methods **219**, 123 (1984). [Rel. P_γ]
- 1985ZiZY - W. L. Zijp, Report FYS/RASA -85/19 (1985) [Limitation of Relative Statistical Weights]
- 1986Me10 - D. Mehta, M. L. Garg, J. Singh, N. Singh, T. S. Cheema, P. N. Trehan, Nucl. Instrum. Methods Phys. Res. **A219**, 447 (1986). [Rel. P_γ]
- 1986Wo05 - M. J. Woods, S. E. M. Lucas, Int. J. Appl. Radiat. Isot. **37**, 1157 (1986). [$T_{1/2}$]
- 1987Co39 - E. R. Cohen and B. N. Taylor, Rev. Mod. Phys. **59**, 1121 (1987) [Fundamental Constants]
- 1989Da12 - V. N. Danilenko, N. P. Gromova, A. A. Konstantinov, N. V. Kurenkov, A. B. Malinin, S. V. Matveev, T. E. Sazonova, E. K. Stepanov, S. V. Sepman, I. N. Tronova, Appl. Radiat. Isot. **40**, 711 (1989). [Rel. P_γ]
- 1990Me15 - R. A. Meyer, Fizika (Zagreb) **22**, 153 (1990). [Rel. P_γ]
- 1990St02 - N. M. Stewart, E. Eid, M. S. S. El -Daghmah, J. K. Jabber, Z. Phys. **A335**, 13 (1990). [Rel. P_γ]
- 1991BaZS - W. Bambynek, T. Barta, R. Jedlovsky, P. Christmas, N. Coursol, K. Debertain, R. G. Helmer, A. L. Nichols, F. J. Schima, Y. Yoshizawa, Report IAEA -TECDOC-619 (1991) [Rel. P_γ]
- 1992Ra08 - M. U. Rajput, T. D. Mac Mahon, Nucl. Instrum. Methods Phys. Res. **A312**, 289 (1992). [Limitation of Relative Statistical Weights]
- 1992Un01 - M. P. Unterweger, D. D. Hoppes, F. J. Schima, Nucl. Instrum. Methods Phys. Res. **A312**, 349 (1992). [$T_{1/2}$]
- 1992Ya12 - Y. Yan, H. Sun, D. Hu, J. Huo, Y. Liu, Z. Phys. **A344**, 25 (1992). [Rel. P_γ]
- 1993Ka30 - Kawaldeep, V. Kumar, K. S. Dhillon, K. Singh, J. Phys. Soc. Jpn. **62**, 901 (1993). [Rel. P_γ]
- 1995Au04 - G. Audi and A. H. Wapstra, Nucl. Phys. **A595**, 409 (1995). [Q(EC), Q(β^-)]
- 1996Ar09 - Agda Artna-Cohen, Nucl. Data Sheets **79**, 1 (1996). [E_γ , Rel. P_γ , Multp.]
- 1996Sc06 - E. Schönfeld, H. Janssen, Nucl. Instrum. Methods Phys. Res. **A369**, 527 (1996) [K-Fluorescence Yield ω_K]
- 1997Ma75 - R. H. Martin, K. I. W. Burns, J. G. V. Taylor, Nucl. Instrum. Methods Phys. Res. **A390**, 267 (1997) [$T_{1/2}$]
- 1998Hw07 - H.Y. Hwang, T. S. Park, J. M. Lee, Appl. Radiat. Isot. **49**, 1201 (1998). [Rel. P_γ]
- 1998Sc28 - E. Schönfeld, Appl. Radiat. Isot. **49**, 1353 (1998). [P_K , P_L , P_M , P_N]
- 1998Si12 - H. Siegert, H. Schraeder, U. Schotzig, Appl. Radiat. Isot. **49**, 1397 (1998). [$T_{1/2}$]
- 2000He14 - R. G. Helmer, C. van der Leun, Nucl. Instrum. Methods in Phys. Res. **A450**, 35 (2000). [E_γ]

Table 1. Relative g-Ray Emission Probabilities Evaluated in this Revision (Uncertainty given below the value)

E(keV)	1970NO06	1970RI19	1971BA63	1972BA05	1977GE12	1980SH15	1984IW03	1986ME10	1989DA12	1990ME15	1990ST02	1992YA12	1993KA30	1998HW07*
121.8	145.0	138.5	132.9	144.6	141.0	140.6	136.9	136.7	139.0	136.2	136.6		133.5	136.9
	4.1	6.4	4.0	4.7	4.0	2.8	1.3	0.7	1.0	1.6	1.8		1.8	3.9
125.7										0.057	0.115			
										0.009	0.013			
148.0			0.077			0.154				0.190	0.218		0.231	
			0.026			0.013				0.040	0.026		0.026	
166.9											0.051		0.010	
											0.013		0.004	
173.1										0.002	0.038		0.081	
										0.001	0.013		0.003	
192.6										0.033	0.023	0.031	0.029	
										0.001	0.006	0.008	0.005	
202.6										0.018	0.028			
										0.009	0.006			
207.6			0.064	0.035		0.038				0.021	0.031	0.022	0.035	
			0.038	0.012		0.013				0.006	0.006	0.003	0.003	
209.4			0.077	0.038		0.026				0.021	0.038	0.027	0.026	
			0.038	0.026		0.013				0.006	0.013	0.003	0.013	
212.6		0.086	0.103	0.097		0.103				0.094	0.115		0.077	
		0.037	0.026	0.029		0.026				0.003	0.026		0.026	
237.3			0.051							0.045	0.064		0.012	
			0.026							0.004	0.026		0.004	
239.4				0.321							0.051		0.019	
				0.154							0.013		0.004	
244.7	39.4	36.2	35.8	36.4	36.6	35.8	36.2	36.5	36.5	35.9	38.0			36.8
	1.3	1.8	1.0	1.2	1.1	0.6	0.3	0.4	0.3	0.6	0.5			0.9
251.6		0.333	0.372	0.359		0.359				0.300	0.321		0.308	
		0.051	0.064	0.051		0.013				0.010	0.026		0.026	
269.9				0.015						0.039				
				0.006						0.004				

Comments on evaluation

¹⁵²Eu

Eg(keV)	1970NO06	1970RI19	1971BA63	1972BA05	1977GE12	1980SH15	1984IW03	1986ME10	1989DA12	1990ME15	1990ST02	1992YA12	1993KA30	1998HW07*
271.1#		0.359	0.359	0.374		0.410				0.389	0.372		0.436	
		0.051	0.064	0.038		0.026				0.011	0.026		0.013	
275.5		0.141	0.154	0.154		0.218				0.161	0.205		0.128	
		0.038	0.038	0.013		0.026				0.050	0.026		0.013	
286.0										0.053	0.064		0.044	
										0.005	0.026		0.004	
295.9	2.37	1.94	2.09	2.04		2.06	2.13	2.22	2.12	2.11	2.21		2.08	
	0.19	0.12	0.14	0.06		0.05	0.04	0.04	0.02	0.05	0.06		0.05	
315.2#		0.218	0.237	0.228		0.308				0.253	0.231		0.231	
		0.038	0.043	0.040		0.026				0.008	0.038		0.038	
316.2			0.045	0.023						0.010				
			0.019	0.012						0.006				
320.0										0.008				
										0.003				
324.8		0.333	0.385	0.346		0.359				0.360	0.346			
		0.038	0.064	0.051		0.026				0.010	0.013			
329.4		0.564	0.615	0.577		0.628	0.707			0.590	0.603		0.410	
		0.051	0.103	0.064		0.038	0.015			0.010	0.026		0.038	
330.5				0.029						0.360				
				0.008						0.050				
340.4			0.103	0.117						0.130	0.141		0.182	
			0.051	0.012						0.030	0.038		0.010	
344.3	128.2	128.2	128.2	128.2	127.2	128.2	127.1	126.9	128.2	127.5	128.2		128.2	128.2
	3.6	5.9	3.8	4.2	1.3	2.6	0.7	0.9	0.8	0.9	1.7		1.8	2.9
351.7			0.077	0.086		0.103				0.043	0.090		0.103	
			0.026	0.018		0.026				0.003	0.026		0.026	
357.3										0.023			0.013	
										0.003			0.004	
367.8	3.78	4.04	4.14	4.08	4.19	4.15	4.13	4.14	4.18	4.05	4.05		4.04	4.13
	0.32	0.23	0.15	0.14	0.04	0.09	0.04	0.07	0.04	0.08	0.06		0.08	0.10
379.4										0.004	0.051			
										0.001	0.013			

Comments on evaluation

¹⁵²Eu

Eg(keV)	1970NO06	1970RI19	1971BA63	1972BA05	1977GE12	1980SH15	1984IW03	1986ME10	1989DA12	1990ME15	1990ST02	1992YA12	1993KA30	1998HW07*
385.7				0.109						0.024	0.269		0.167	
				0.049						0.003	0.026		0.026	
387.9										0.014	0.017		0.018	
										0.001	0.006		0.005	
391.3										0.006				
										0.001				
395.0											0.038		0.026	
											0.013		0.013	
406.7										0.004				
										0.001				
411.0	10.14	10.32	10.77	10.59	10.71	10.55	10.84	10.73	10.80	10.70	10.82		10.72	10.70
	0.54	0.51	0.38	0.27	0.11	0.22	0.07	0.10	0.10	0.10	0.15		0.23	0.29
416.0		0.487	0.513	0.500		0.513				0.530	0.526		0.500	
		0.051	0.064	0.051		0.026				0.010	0.026		0.026	
423.5										0.013	0.027	0.022	0.013	
										0.003	0.006	0.010	0.005	
440.9										0.052			0.069	
										0.009			0.006	
444.0		13.2	13.5	13.6										
		0.8	0.5	0.8										
444.0		1.15	1.67	1.28										
		0.38	0.26	0.26										
444.0@	15.47	14.36	15.13	14.87	15.00	14.95	15.01	14.81	14.90	14.80	15.06		15.18	13.78
	0.33	0.86	0.57	0.81	0.15	0.13	0.11	0.13	0.20	0.20	0.22		0.22	0.39
482.3		0.141	0.115	0.128		0.167				0.130	0.154			
		0.026	0.026	0.026		0.013				0.010	0.026			
488.7		1.90	1.95	1.91	1.98	1.95	2.03		1.95	1.95	2.01		1.95	1.97
		0.12	0.13	0.06	0.02	0.03	0.02		0.04	0.02	0.04		0.05	0.05
493.5		0.115	0.154	0.218		0.179				0.190	0.179		0.103	
		0.051	0.038	0.038		0.026				0.010	0.026		0.026	
496.3				0.038		0.051				0.044	0.064		0.040	
				0.015		0.013				0.003	0.026		0.009	

Comments on evaluation

¹⁵²Eu

Eg(keV)	1970NO06	1970RI19	1971BA63	1972BA05	1977GE12	1980SH15	1984IW03	1986ME10	1989DA12	1990ME15	1990ST02	1992YA12	1993KA30	1998HW07*
503.5		0.705	0.718	0.705		0.718	0.768			0.730	0.782		0.474	
		0.038	0.077	0.038		0.026	0.018			0.010	0.051		0.256	
520.2		0.231	0.269	0.256		0.282				0.257	0.231			
		0.051	0.038	0.038		0.026				0.007	0.026			
523.1			0.051	0.031						0.071	0.103		0.096	
			0.026	0.010						0.004	0.038		0.123	
526.9			0.051	0.046		0.064				0.063	0.077		0.060	
			0.026	0.014		0.026				0.003	0.026		0.029	
534.4			0.179	0.179										
			0.051	0.051										
535.4#		0.205	0.218	0.205		0.231				0.206	0.192		0.167	
		0.051	0.053	0.052		0.026				0.005	0.038		0.026	
538.3										0.020				
										0.003				
556.6										0.091	0.077			
										0.005	0.013			
556.5#			0.115	0.090		0.051				0.110	0.128		0.090	
			0.026	0.026		0.026				0.006	0.018		0.013	
557.9										0.019	0.051			
										0.004	0.013			
561.2				0.013						0.005				
				0.006						0.001				
562.9				0.18										
				0.06										
564.0#		2.40	2.46	2.38		2.31	2.43		2.36	2.36	2.32			
		0.19	0.19	0.09		0.06	0.04		0.06	0.05	0.05			
566.4		0.526	0.564	0.577		0.679	0.640			0.620	0.551		0.697	
		0.128	0.128	0.051		0.038	0.060			0.010	0.026		0.022	
571.8										0.023			0.025	
										0.004			0.008	
586.3		2.08	2.28	2.22	2.24	2.27	2.19		2.22	2.20	2.24			2.14
		0.27	0.14	0.09	0.05	0.05	0.08		0.05	0.05	0.05			0.05

Comments on evaluation

¹⁵²Eu

Eg(keV)	1970NO06	1970RI19	1971BA63	1972BA05	1977GE12	1980SH15	1984IW03	1986ME10	1989DA12	1990ME15	1990ST02	1992YA12	1993KA30	1998HW07*
595.6											0.154		0.015	
											0.051		0.008	
616.1			0.064	0.049		0.038				0.043	0.051	0.038	0.064	
			0.026	0.015		0.026				0.004	0.013	0.013	0.026	
644.4			0.064	0.029		0.038				0.028	0.051	0.027	0.028	
			0.038	0.009		0.026				0.004	0.013	0.010	0.009	
656.5	0.590		0.744	0.679		0.654	0.710			0.690	0.718		0.692	
	0.064		0.090	0.051		0.038	0.050			0.010	0.038		0.026	
664.8			0.045	0.017		0.038				0.090	0.064		0.051	
			0.019	0.008		0.026				0.010	0.026		0.038	
671.3	0.059		0.090	0.109		0.064				0.110	0.077	0.091	0.051	
	0.027		0.051	0.038		0.026				0.010	0.038	0.009	0.026	
674.7	0.385		0.744	0.615										
	0.103		0.103	0.064										
675.0#	0.846		0.872	0.744		0.949	0.940			0.890	0.936		0.846	
	0.154		0.115	0.082		0.038	0.050			0.030	0.051		0.038	
678.6	2.06		2.31	2.19	2.30	2.31	2.28		2.21	2.21	2.41		2.24	2.22
	0.15		0.14	0.14	0.03	0.06	0.05		0.03	0.04	0.08		0.05	0.07
686.6			0.192	0.128						0.092				
			0.051	0.051						0.008				
688.7	3.88		4.15	4.14	4.12	4.08	4.20		4.12	4.09	4.06		4.17	4.06
	0.22		0.22	0.27	0.04	0.10	0.04		0.05	0.08	0.08		0.08	0.11
696.9											0.077		0.014	
											0.038		0.005	
703.3				0.073						0.025	0.103		0.013	
				0.022						0.004	0.038		0.009	
712.8	0.346		0.462	0.423		0.487				0.460	0.474			
	0.090		0.077	0.090		0.038				0.010	0.038			
719.3#	1.42		1.64	1.53		1.67	1.67		1.51	1.56	1.62		1.58	
	0.13		0.17	0.13		0.05	0.03		0.02	0.03	0.04		0.04	
719.3			0.283	0.282										
			0.077	0.038										

Comments on evaluation

¹⁵²Eu

Eg (keV)	1970NO06	1970RI19	1971BA63	1972BA05	1977GE12	1980SH15	1984IW03	1986ME10	1989DA12	1990ME15	1990ST02	1992YA12	1993KA30	1998HW07*
728.0				0.044		0.051				0.054	0.064	0.051	0.064	
				0.009		0.013				0.050	0.026	0.013	0.013	
735.4										0.028				
										0.005				
756.1										0.026			0.301	
										0.004			0.013	
764.9		0.821	0.910	0.885			0.950			0.840	0.962		0.936	
		0.141	0.103	0.115			0.050			0.040	0.051		0.038	
768.9		0.372	0.397	0.346		0.410				0.430	0.500		0.449	
		0.103	0.064	0.038		0.038				0.040	0.038		0.038	
778.9		59.7	62.6	59.9	62.6	62.5	62.16	62.1	62.2	61.9	62.1		62.5	63.7
		2.9	1.4	0.7	0.6	1.2	0.22	0.5	0.4	0.8	0.9		1.3	1.4
794.8		0.192	0.141	0.141		0.192				0.118	0.192		0.136	
		0.051	0.064	0.090		0.026				0.006	0.038		0.014	
805.7				0.077						0.061	0.090		0.050	
				0.026						0.005	0.026		0.009	
810.5		1.38	1.56	1.50		1.55	1.56		1.51	1.52	1.55		1.50	
		0.12	0.10	0.06		0.05	0.04		0.02	0.02	0.04		0.03	
839.4			0.077	0.079						0.079	0.064		0.077	
			0.038	0.045						0.005	0.013		0.013	
841.6			0.769	0.769						0.780	0.769		0.859	
			0.090	0.115						0.010	0.038		0.051	
867.4		19.23	20.09	19.31	20.54	20.29	20.33	20.36	20.40	19.90	20.33		20.45	20.92
		0.90	0.49	0.35	0.21	0.51	0.10	0.17	0.30	0.40	0.27		0.42	0.48
896.6											0.269		0.323	
											0.051		0.010	
901.2		0.295	0.385	0.359		0.346	0.400			0.440	0.397		0.449	
		0.090	0.064	0.077		0.038	0.050			0.030	0.038		0.038	
906.0										0.072			0.087	
										0.006			0.008	
919.3		1.88	2.06	1.91		2.14	2.08		2.09	2.09	2.04		2.05	2.05
		0.14	0.24	0.07		0.06	0.06		0.04	0.05	0.05		0.06	0.12

Comments on evaluation

¹⁵²Eu

Eg(keV)	1970NO06	1970RI19	1971BA63	1972BA05	1977GE12	1980SH15	1984IW03	1986ME10	1989DA12	1990ME15	1990ST02	1992YA12	1993KA30	1998HW07*
926.3		1.167	1.308	1.218		1.333	1.380		1.290	1.270	1.346		1.359	1.340
		0.103	0.128	0.115		0.051	0.060		0.040	0.040	0.641		0.051	0.058
930.6		0.308	0.333	0.346		0.359	0.370			0.350	0.385		0.308	
		0.077	0.064	0.051		0.038	0.060			0.010	0.038		0.038	
937.1				0.010						0.015	0.051			
				0.004						0.005	0.026			
958.6			0.064	0.077		0.064				0.110	0.103			
			0.038	0.038		0.026				0.010	0.038			
963.4			0.628	0.487										
			0.103	0.103										
964.1#		67.44	69.86	68.08	70.40	70.45	70.14	71.03	70.50	69.20	69.67		70.50	67.96
		3.33	1.79	1.79	0.70	1.41	0.23	0.40	0.60	0.90	0.95		1.49	1.93
974.1			0.045	0.051		0.064				0.069	0.090		0.065	
			0.019	0.013		0.013				0.005	0.026		0.009	
990.2		0.167	0.128	0.154		0.179				0.148	0.167		0.179	
		0.051	0.064	0.051		0.026				0.006	0.038		0.038	
1001.1										0.019			0.023	
										0.009			0.005	
1005.3		3.04	3.13	3.00	3.57	3.59	3.08		3.35	3.10	3.46		2.73	3.11
		0.31	0.32	0.21	0.07	0.13	0.02		0.04	0.07	0.13		0.12	0.13
1086.0		47.69	50.64	47.59	48.70	49.62	48.15	47.84	49.60	48.70	49.19		49.60	47.96
		2.82	1.54	0.86	0.50	1.28	0.16	0.31	0.40	0.80	0.67		0.94	1.06
1089.7		8.00	8.46	7.90	8.26	8.59	8.35	8.19		8.20	7.97		8.19	8.19
		0.64	0.77	0.37	0.09	0.26	0.04	0.10		0.10	0.51		0.17	0.19
1109.2			0.897	0.808			1.000			0.880				
			0.385	0.179			0.050			0.020				
1112.0#		63.59	65.77	63.99	65.00	65.64	65.67	65.45	65.90	65.80	65.23		62.47	
		3.21	1.85	0.87	0.70	1.28	0.22	0.78	0.50	0.90	0.99		1.12	
1112.0			64.87	63.18			64.67			64.90				
			1.79	0.86			0.21			0.90				
1139.0										0.006			0.006	
										0.002			0.002	

Comments on evaluation

¹⁵²Eu

Eg(keV)	1970NO06	1970RI19	1971BA63	1972BA05	1977GE12	1980SH15	1984IW03	1986ME10	1989DA12	1990ME15	1990ST02	1992YA12	1993KA30	1998HW07*
1170.9		0.167	0.167	0.167		0.256				0.171	0.231		0.141	
		0.038	0.038	0.038		0.026				0.006	0.038		0.038	
1206.1			0.064	0.038		0.038				0.072	0.064		0.051	
			0.038	0.013		0.026				0.005	0.026		0.013	
1212.9		6.55	7.05	6.74	6.67	6.72	6.85		6.83	6.70	6.97		6.85	6.70
		0.35	0.26	0.26	0.07	0.14	0.05		0.05	0.08	0.18		0.15	0.19
1249.9		0.795	0.885	0.833		0.962	0.875			0.880	0.923		0.859	0.921
		0.090	0.077	0.064		0.038	0.024			0.050	0.051		0.064	0.039
1261.3		0.154	0.167	0.167		0.192				0.157	0.192		0.162	
		0.038	0.038	0.038		0.026				0.006	0.026		0.060	
1292.8		0.487	0.474	0.474		0.500	0.460			0.490	0.641		0.654	
		0.090	0.077	0.077		0.026	0.030			0.030	0.064		0.077	
1299.1		7.71	8.23	7.88	7.76	7.97	7.80		7.88	7.80	7.94		8.08	
		0.40	0.41	0.44	0.08	0.19	0.05		0.06	0.10	0.19		0.36	
1314.7			0.019	0.018		0.038					0.038	0.024	0.026	
			0.009	0.006		0.013					0.013	0.005	0.013	
1348.1		0.058	0.090	0.081		0.090				0.081	0.090	0.078	0.115	
		0.023	0.013	0.010		0.013				0.006	0.013	0.008	0.013	
1363.8		0.108	0.128	0.126		0.141				0.117	0.128		0.132	
		0.031	0.013	0.015		0.013				0.005	0.013		0.012	
1390.4			0.026	0.019						0.023	0.031	0.024	0.015	
			0.013	0.006						0.006	0.010	0.005	0.010	
1408.0		99.5	103.6	97.7	100.0	99.9	100.0	100.0	100.0	100.0	99.2		102.6	
		5.0	2.7	2.8	1.0	1.9	0.3	0.6	0.5	0.3	1.1		1.4	
1457.6		2.45	2.46	2.40	2.52	2.46	2.39		2.35	2.36	2.38			
		0.13	0.19	0.13	0.09	0.05	0.03		0.03	0.05	0.10			
1486.0											0.027		0.014	
											0.012		0.005	
1528.1		1.67	1.28	1.46		1.27	1.35		1.38	1.27	1.26		1.47	
		0.09	0.08	0.09		0.04	0.01		0.02	0.03	0.10		0.05	
1537.4		0.007		0.010		0.012								
		0.003		0.003		0.004								

Comments on evaluation

¹⁵²Eu

Eg(keV)	1970NO06	1970RI19	1971BA63	1972BA05	1977GE12	1980SH15	1984IW03	1986ME10	1989DA12	1990ME15	1990ST02	1992YA12	1993KA30	1998HW07*
1605.6		0.035	0.038	0.037		0.051				0.036	0.038	0.044	0.041	
		0.008	0.008	0.008		0.013				0.003	0.013	0.004	0.009	
1608.4		0.029	0.023	0.027						0.024	0.027	0.029		
		0.006	0.008	0.006						0.002	0.006	0.004		
1635.2										0.0007				
										0.0002				
1643.6		0.024				0.005								0.009
		0.005				0.003								0.003
1647.4		0.033	0.028	0.031		0.038					0.041	0.024	0.031	
		0.006	0.006	0.006		0.013					0.006	0.004	0.003	
1674.3										0.029				
										0.004				
1769.0		0.042	0.041	0.042		0.038				0.042	0.038	0.049	0.046	
		0.004	0.006	0.005		0.013				0.003	0.013	0.003	0.006	

* Evaluators considered unwarranted the precision of the values given by 98Hw07. Their uncertainties have been doubled.

Value includes the contribution from the weakest component of the doublet.

@ Value is the sum of the components of the doublet.

Table 1. Relative g-Ray Emission Probabilities Evaluated in this Revision (Uncertainty given below the value), continuation

Eg (keV)	ICRM01	ICRM02	ICRM08	ICRM10	ICRM12	ICRM15	ICRM16	ICRM17	ICRM18	ICRM20	ICRM25	ICRM27	ICRM28	ICRM29
121.8	135.0	135.7	136.4	131.5	135.8		133.4		139.2	137.0		136.4	132.5	134.8
	1.9	0.8	0.5	4.3	0.9		1.4		2.9	1.0		3.0	2.9	2.0
244.7	35.5	35.5	36.3	36.2	35.9		36.3	36.7		35.7	35.7		36.3	36.4
	0.5	0.3	0.2	1.0	0.5		0.3	1.1		0.4	0.4		0.7	0.4
344.3	128.9	127.2	127.4	123.9	127.6	130.6	130.4	127.1		127.2	126.7	126.2	128.9	128.8
	1.5	0.8	0.6	2.8	0.4	2.9	1.2	1.1		1.0	1.1	3.4	2.4	1.3
411.0	10.46	10.67	10.80	10.27	10.75	10.77	10.90	10.71	10.90	10.72	10.90	10.62	10.72	10.86
	0.16	0.07	0.06	0.22	0.04	0.12	0.12	0.11	0.23	0.10	0.33	0.67	0.26	0.12
444.0@	14.68	14.84	14.96	14.35	15.07	15.25	15.33	14.88	15.3	14.95	14.73	14.64	15.15	15.22
	0.21	0.09	0.07	0.4	0.06	0.12	0.18	0.15	0.26	0.13	0.43	0.89	0.32	0.15
778.9	62.4	62.6	62.25		62.12	62.6	62.4	62.6	61.8	61.9	61.1	61.0	62.0	62.4
	0.8	0.4	0.19		0.23	0.4	1.2	0.6	1.2	0.4	0.9	1.0	1.0	0.5
964.1	69.62	69.82	70.10		70.41	70.40	69.80	70.30	69.90	70.30	70.90	69.30	68.40	70.10
	0.84	0.42	0.23		0.22	0.60	0.90	0.70	1.00	0.40	1.00	1.00	1.10	0.50
1086.0	48.89	48.61	49.13	47.43	48.83	49.10	47.90	48.70	48.90	48.40		48.50		48.59
	0.59	0.29	0.19	0.60	0.14	0.40	0.60	0.50	0.50	0.30		0.90		0.30
1112.0	64.28	64.45	65.25	64.00	65.26	65.70	64.70	64.30	66.70	64.90	67.20	64.50	65.50	65.30
	0.77	0.32	0.27	0.80	0.20	0.70	0.40	0.60	0.80	0.50	0.90	1.10	1.00	0.50
1408.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
	1.2	0.5	0.3	1.5	0.3	0.9	0.9	1.0	1.2	0.5	1.2	1.5	2.3	0.7

Eg (keV)	ICRM30	ICRM31	ICRM34	ICRM35
121.8	136.8	135.5	138.9	134.9
	4.1	2.0	4.3	1.2
244.7	37.9	35.6		36.4
	1.2	0.5		0.2
344.3	132.7	126.6	133.9	126.4
	4.0	1.3	5.5	0.9
411.0	11.21	10.52	11.18	10.57
	0.39	0.14	0.53	0.08
444.0		14.89	16.15	14.81
		0.19	0.73	0.16

Eg (keV)	ICRM30	ICRM31	ICRM34	ICRM35
778.9	61.2	61.3	64.2	62.0
	1.9	0.7	2.1	0.5
964.1	69.80	70.00	71.20	69.90
	2.20	0.80	2.30	0.50
1086.0	50.70	48.00	50.00	
	1.50	0.50	1.20	
1112.0	64.70	65.40	66.50	64.20
	2.00	0.80	1.50	0.70
1408.0	100.0	100.0	100.0	100.0
	3.0	1.0	2.9	1.2

Table 2. Recommended Relative g-Ray Emission Probabilities (Uncertainty given below the value).

E_g(keV)	Recommended	c2/n	Remarks	E_g(keV)	Recommended	c2/n	Remarks	E_g(keV)	Recommended	c2/n	Remarks
121.8	136.35	1.3		271.1	0.374	1.9	[2]	379.4	0.004		[5]
	0.25				0.014				0.001		
125.7	0.09	9.9		275.5	0.155	2.3		385.7	0.024		[6]
	0.03				0.008				0.003		
148.0	0.166	5.8		286.0	0.048	1.2		387.9	0.0142		[5]
	0.024				0.003				0.0010		
166.9			[18]	295.9	2.123	1.6		391.3	0.006		[13]
					0.013				0.001		
173.1			[18]	315.2	0.238	1.1	[3]	395.0			[18]
					0.008						
192.6	0.0326	1.1		316.2	0.015		[3]	406.7	0.004		[13]
	0.0010				0.005				0.001		
202.6			[18]	320.0	0.008		[13]	411.0	10.735	0.95	
					0.003				0.020		
207.6	0.0285	2.1		324.8	0.354	0.27		416.0	0.523	0.4	
	0.0019				0.007				0.008		
209.4	0.0266	0.60		329.4	0.62	11		423.5	0.0155	1.7	
	0.0025				0.03				0.0023		
212.6	0.094	0.23		330.5	0.029		[4]	440.9	0.064	2.5	
	0.003				0.008				0.005		
237.3	0.012		[1]	340.4	0.151	4.6		444.0	13.46		[7]
	0.004				0.016				0.09		
239.4	0.036	3.2		344.3	127.53	0.66		444.0	1.53		[7]
	0.016				0.20				0.09		
244.7	36.23	1.5		351.7	0.067	2.2		444.0	14.99	1.2	[7]
	0.08				0.011				0.03		
251.6	0.322	2.4		357.3	0.0194	4.0		482.3	0.141	1.3	
	0.007				0.0024				0.008		
269.9	0.029	8.0		367.8	4.136	0.77		488.7	1.985	1.8	
	0.012				0.018				0.008		

Eg(keV)	Recommended	c2/n	Remarks	Eg(keV)	Recommended	c2/n	Remarks	Eg(keV)	Recommended	c2/n	Remarks
493.5	0.178	2.1		571.8	0.023	0.10		719.3	1.29	0.33	[12]
	0.016				0.004				0.06		
496.3	0.044	0.31		586.3	2.215	0.57		719.3	0.282	0.0	[12]
	0.004				0.019				0.035		
503.5	0.735	1.0		595.6	0.015		[11]	728.0	0.051	0.37	
	0.008				0.008				0.006		
520.2	0.257	0.46		616.1	0.044	0.32		735.4	0.028		[13]
	0.006				0.003				0.005		
523.1	0.054	2.7		644.4	0.030	0.65		756.1	0.026		[13]
	0.010				0.003				0.004		
526.9	0.062	0.39		656.5	0.689	0.63		764.9	0.912	0.94	
	0.003				0.008				0.021		
534.4	0.176	0.56	[8]	664.8	0.046	6.6		768.9	0.424	1.5	
	0.009				0.014				0.016		
535.4	0.029		[8]	671.3	0.093	1.3		778.9	62.17	0.8	
	0.010				0.006				0.09		
538.3	0.020			674.7			[18]	794.8	0.126	2.3	
	0.003								0.005		
556.6			[18]	675.0	0.897	1.3		805.7	0.060	1.0	
					0.021				0.004		
556.5	0.085	1.7	[9]	678.6	2.256	1.3		810.5	1.519	0.57	
	0.005				0.015				0.011		
557.9	0.021	5.5	[9]	686.6	0.096	2.1		839.4	0.077	0.29	
	0.003				0.008				0.004		
561.2	0.0052	1.7		688.7	4.037	0.60		841.6	0.782	0.62	
	0.0010				0.021				0.009		
562.9	0.18		[4]	696.9	0.014		[11]	867.4	20.35	1.3	
	0.06				0.005				0.07		
564.0	2.19		[10]	703.3	0.025	3.6		896.6	0.321	1.0	
	0.06				0.004				0.010		
566.4	0.628	3.2		712.8	0.461	0.48		901.2	0.404	1.0	
	0.018				0.009				0.016		

Eg(keV)	Recommended	c2/n	Remarks	Eg(keV)	Recommended	c2/n	Remarks	Eg(keV)	Recommended	c2/n	Remarks
906.0	0.077	2.2		1112.0			[18]	1528.1	1.349	4.4	
	0.005								0.021		
919.3	2.06	1.1		1139	0.006		[13]	1537.4			[18]
	0.02				0.002						
926.3	1.309	0.73		1170.9	0.175	2.2		1605.6	0.0388	0.54	
	0.019				0.006				0.0020		
930.6	0.350	0.37		1206.1	0.065	1.7		1608.4	0.0255	0.38	
	0.009				0.004				0.0016		
937.1	0.013	1.4		1212.9	6.79	0.95		1635.2	0.0007		[13]
	0.003				0.03				0.0002		
958.6	0.101	1.10		1249.9	0.894	0.89		1643.6	0.0070	0.89	[16]
	0.009				0.015				0.0020		
963.4	0.644		[14]	1261.3	0.161	0.56		1647.4	0.0305	1.1	
	0.009				0.005				0.0019		
964.1	69.55	0.62	[14]	1292.8	0.499	1.6		1674.3	0.029		[13]
	0.10				0.015				0.004		
974.1	0.066	0.76		1299.1	7.83	0.48		1769.0	0.0441	0.63	
	0.004				0.03				0.0016		
990.2	0.151	0.39		1314.7	0.023	0.73					
	0.006				0.003						
1001.1	0.022	0.15		1348.1	0.084	1.2					
	0.005				0.004						
1005.3	3.19	9.6		1363.8	0.123	0.75					
	0.11				0.004						
1086.0	48.63	1.9	[17]	1390.4	0.023	0.36					
	0.20				0.003						
1089.7	8.30	0.78		1408.0	100.00	0.22					
	0.03				0.12						
1109.2	0.892		[15]	1457.6	2.388	0.82					
	0.018	1.7			0.017						
1112.0	64.30		[15]	1486.0			[18]				
	0.09										

REMARKS

- Evaluator's recommended relative γ -ray emission probabilities deduced using the *Limitation of Relative Statistical Weights* method, unless otherwise specified.
- For absolute intensity per 100 disintegrations, multiply by 0.2085 (8).

- [1]. From 1993Ka30.
- [2]. $I_\gamma =$ weighted average ($I_\gamma(271)$ doublet) - $I_\gamma(269) = 0.403$ (7) - 0.029 (12) = 0.374 (14). $\chi^2/\nu = 1.9$.
- [3]. $I_\gamma =$ weighted average ($I_\gamma(315)$ doublet) - $I_\gamma(316) = 0.253$ (7) - 0.015 (5) = 0.238 (8). $\chi^2/\nu = 1.1$.
- [4]. From 72Ba05.
- [5]. From 1990Me15. Value agrees with <0.006 (1990St02).
- [6]. From 1990Me15. Author removed double-escape contribution from 1408-keV γ ray.
- [7]. $I_\gamma =$ weighted average ($I_\gamma(444)$ doublet) - $I_\gamma(444, 810 \text{ level}) = 14.99$ (3) - 1.53 (9) = 13.46 (9).
 $\chi^2/\nu = 1.2$. $I_\gamma(444, 810 \text{ level})$ is from ¹⁵²Eu(9.3h) EC decay branching.
- [8] $I_\gamma =$ weighted average ($I_\gamma(535)$) - $I_\gamma(534) = 0.205$ (5) - 0.176(9)=0.029(10)
- [9]. $I_\gamma =$ weighted average ($I_\gamma(556.5)$ doublet) - weighted average $I_\gamma(557.8) = 0.106$ (5) - 0.021 (4) = 0.085 (6)
- [10]. $I_\gamma =$ weighted average ($I_\gamma(563.8)$ doublet) - $I_\gamma(562.9) = 2.37$ (2) - 0.18 (6) = 2.19 (6). $\chi^2/\nu = 0.64$.
 $I_\gamma(562.9) = 2.37$ (2) from transition intensity balance.
- [11]. From 1993Ka30, close to upper limit of ⁹²Yb12.
- [12]. $I_\gamma =$ weighted average ($I_\gamma(719)$ doublet, $\chi^2/\nu = 3.4$) - weighted average $I_\gamma(719.4) = 1.57$ (2) - 0.282 (35) = 1.29 (6).
- [13]. From 1990Me15.
- [14]. $I_\gamma =$ weighted average ($I_\gamma(964)$ doublet) - $I_\gamma(963) = 70.19$ (10) - 0.644 (9) = 69.55 (10).
 $I_\gamma(963) = 0.644$ (9) is from ¹⁵²Eu(9.3h) EC decay branching.
- [15]. $I_\gamma =$ weighted average ($I_\gamma(1112)$ doublet, $\chi^2/\nu = 1.5$) - weighted average $I_\gamma(1109, \chi^2/\nu = 1.7) = 65.19$ (9) - 0.895 (18) = 64.30 (9)
- [16]. Weighted average of values from 1980Sh15 and 1993Ka30.
- [17] $I_\gamma = I_\gamma(1084) + I_\gamma(1086) = 1.17$ (4) (1990Me15) + 47.46 (20) = 48.63 (20)
- [18]. Existence is uncertain.

Table 3. Absolute Emission Probabilities of KX Rays

P_{KX}*	70No06	Faerman[†]	72Da23	Bylov[‡]	79De36, 83De11	85Se18	86Me10	93Ka30	P_{KX} (Avg.)^{&}	P_{KX}(Cal.)[@]
Sm KA	0.492(35)	0.592(21)	0.501(16)	0.595(9)	0.591(12)	0.595(9)	0.589(9)	0.595(90)	0.584(11)	0.585(7)
Sm KB	0.122(9)	0.173(9)	0.122(8)	0.143(8)	0.149(3)	0.143(8)	0.144(2)	0.137(5)	0.144(3)	0.1482(24)
Gd KA			0.0068(2)	0.00636(14)	0.00648(22)	0.00636(14)	0.00459(11) [#]		0.00645(8)	0.00680(18)
Gd KB			0.00167(50)	0.00163(4)	0.00176(18)	0.00163(4)	0.00171(3)		0.00167(2)	0.00174(5)

* Absolute emission probabilities renormalized to Pg(121)=0.2841(13), Pg(344)=0.2658(12), or Pg(1408)=0.2084(9).

& Weighted average (LWM).

Outlier, not used for calculating the average.

† Faermann S, Notea A., Segal Y., Trans. Am. Nuc. Soc. 14, 500 (1971).

‡ Bylov T., Osipenko B.D., Chudin V.G., EchA Ya no. 9, 1350 (1978) (quoted by 85Se18).

@ Calculated by evaluators using recommended γ -ray data and K-fluorescence yields.

**¹⁵³Sm - Comments on evaluation of decay data
by M.M. Bé, R. G. Helmer and E. Schönfeld**

First evaluation was done in 2001 by R.G. Helmer and E. Schönfeld, it has been updated in June 2005, including new half-life and gamma intensity values.

1 Decay Scheme

There are many levels in ¹⁵³Eu below the decay energy, so other levels may be weakly populated in this decay.

2 Nuclear Data

The Q value is from Audi and Wapstra 2003 (2003Au03). Level energy, spin and parity data are from 1998He06.

The half-life values available are, in hours:

1942Ku03	47	1	as quoted in 1990Le13
1946Mi06	46		as quoted in 1990Le13
1952Ru10	46.5	1	as quoted in 1990Le13
1954Le08	47	0.3	as quoted in 1990Le13
1958Co76	47.1	0.1	
1958Gu09	46.7	1.6	
1960Wi10	45	8	outlier
1961Gr18	46.2	0.1	
1961Wy01	46.8	0.1	
1962Ca24	47.1	0.1	
1963Ho15	46.5	0.5	
1970Ch09	46.75	0.09	
1971Ba28	46.44	0.08	
1987Co04	46.27	0.01	superseded by 1992Un01
1989Ab05	46.70	0.05	
1989Po21	45.6	1.6	outlier
1992Un01	46.2853	0.0014	
1998Bo18	46.285	0.004	
1999Sc12	46.274	0.007	superseded by 2004Sc
2004Sc04	46.281	0.007	<i>Corrected value and uncertainty</i>
Adopted	46.2851	0.0013	or 1.92855 (5) d

A mistake appears in the value of the Sm -153 half-life published by 2004Sc04 in Applied Radiation Isotopes 60 (2004) 317 ; after discussion with the author the correct value is 1.92838 (29) d instead of 1.9284 (29) d.

Data are very discrepant, ranging from 46.281 (7) to two values of 47.1 (1), a difference of about 8 σ .

The Limitation of Relative Statistical Weight, LRSW, analysis (1985ZiZY, 1992Ra08), with the Lweight 3 program, shows that the values from 1960Wi10 and 1989Po21 are outlier due to Chauvenet's criterion, the reduced- χ^2 is 18.9 and the uncertainty of 1992Un01 value is increased to 0.0034 to reduce its weight to 50 %. The weighted mean is 46.2874 with α_{int} of 0.0024 and α_{ext} of 0.011. Then, the program recommends

the unweighted mean and expands the uncertainty to include the most precise value, this leads to a value of 46.64 (36) h.

The average of the measured values has decreased with time and the last three unreplaced values, which are from metrology laboratories, are among the lowest values and they are consistent. The weighted average of these three values is 46.2851 with a σ_{int} of 0.0013, a reduced χ^2 of 0.18, and a σ_{ext} of 0.0006. This weighted average and the internal uncertainty are adopted.

2.1 β^- Transitions

The probabilities for the β^- branches are primarily from the intensity balances from the γ -ray transition probabilities for all levels including the ground state. This is possible because one has measurements of the absolute emission probabilities for the 69- and 103-keV γ -rays (1987Co04, 1998Bo18, 1999Sc12, 2006Le).

The measured β^- probabilities (in %) from the decomposition of the β^- spectra are:

Level (keV)	Values (%)
0	15 (1952Ba49), 20 (1954Gr19), 21 (1954Le08), 20 (1955Ma62), 22 (1956Du31), 20 (1957Jo24), and 20 (1958Co76) compared to the adopted value of 19.5(15) %.
103	67 (1950Hi17), 35 (1952Ba49), 49 (1954Gr19), 70 (1954Le08), 35 (1955Ma62), 38 (1956Du31), 65 (1957Jo24), and 40 (1958Co76) which have an average of 50(14) compared to the adopted value of 49.2(17)% from the probability balance.
172	50 (1952Ba49), 30 (1954Gr19), 43 (1955Ma62), 40 (1956Du31), 15 (1957Jo24), and 40 (1958Co76) which have an average of 36(11) compared to the adopted value of 30.4(8)% from the probability balance.

2.2 Gamma Transitions

The energies and multiplicities are from the adopted gamma data in Nuclear Data Sheets (1998He06) and they are based on the internal-conversion electron data of 1961Mo07, 1962Su01, 1969Sm04, and 1970PaZI. Gamma transition probabilities are deduced from the gamma emission intensities and the conversion electron coefficients interpolated from the tables of Band *et al.* (2002Ba85).

The 19-keV gamma transition probability is deduced from the probability balance at the 83-keV level.

3 Atomic Data

The fluorescence yields and K x-ray relative intensities are from 1996Sc06.

4 Emissions

4.1 Electron Emission

Data were computed by EMISSION for the Auger electrons and with LOGFT for the average β^- energies.

4.2 Photon Emission

From the evaluation 2000He14, the curved-crystal spectrometer data for the decay of ¹⁵³Sm and ¹⁵³Gd give the energies for the γ -rays of 69, 75, 83, 89, 97, 103, and 172 keV on a scale on which the strong line from the decay of ¹⁹⁸Au is 411.80205(17). The γ -ray energies from the (n, γ) study of 1970Mu04 have been adjusted to this energy scale to provide values at 54, 68, 96, 118, 151, 166, and 172 keV. The values for 14 and 19 keV are from level energy differences.

The other γ -ray energies are from the data in the following table 1.

Table 1: Gamma-ray energies

1969Un03	1985Ab08	1969Pa03	Adopted	
412.05 (20)	412.26 (30)	411.9 (1)	412.05 (20)	doubly placed
424.38 (20)	424.79 (32)	424.2 (2)	424.4 (3)	
	431.65 (10)			
436.83 (20)	437.10 (30)	436.7 (2)	436.9 (3)	
	443.24 (45)		443.2 (5)	
		462.0 (3)	462.0 (3)	
463.67 (15)	463.93 (35)	463.4 (2)	463.6 (2)	
485.03 (20)	485.12 (40)	484.5 (2)	485.0 (2)	
	487.75 (23)		487.75 (23)	
509.11 (15)	510.36 (35)	509.0 (1)	509.15 (20)	
521.28 (15)	521.62 (26)	521.1 (1)	521.30 (25)	
		523.8 (6)		
531.38 (15)	531.43 (34)	531.6 (3)	531.40 (15)	
533.34 (15)	533.17 (25)	533.1 (1)	533.2 (2)	
539.03 (10)	539.10 (20)	539.2 (3)	539.1 (2)	
542.60 (20)	543.01 (45)	542.7 (6)	542.7 (2)	
545.75 (15)	545.68 (42)		545.75 (15)	
554.94 (10)	554.73 (37)	555.0 (1)	554.94 (10)	
	555.71 (15)			
574.01 (30)	574.32 (51)		574.1 (3)	
578.66 (15)	578.94 (30)	578.8 (1)	578.75 (20)	
584.49 (20)	584.67 (32)	584.8 (5)	584.55 (20)	
587.47 (20)	587.73 (22)	587.7 (6)	587.60 (25)	
	589.3			
590.96 (20)	591.03 (21)	590.7 (6)	590.96 (20)	
596.72 (15)	596.29 (30)	596.9 (2)	596.7 (2)	
598.4 (3)	598.13 (30)		598.3 (3)	doubly placed
603.39 (15)	604.04 (26)	603.5 (2)	603.6 (4)	doubly placed
609.22 (10)	610.21 (42)	609.4 (1)	609.5 (3)	doubly placed
		612 (1)		
615.41 (20)	616.28 (22)	615.5 (6)	615.8 (4)	doubly placed
617.71 (20)	618.07 (24)	618.0 (6)	617.9 (3)	
	623.73 (24)			
630.70 (30)	630.33 (26)	630 (1)	630.5 (4)	
634.61 (30)	634.92 (32)		634.8 (3)	
636.45 (25)	636.73 (30)	636.4 (2)	636.5 (2)	
657.55 (25)	657.68 (25)	657.4 (4)	657.55 (25)	doubly placed
		662.4 (6)	662.4 (6)	
676.9 (5)	677.09 (30)	676 (1)	677.0 (3)	
		682.0 (6)	682.0 (6)	
685.6 (3)	686.64 (21)	685.9 (3)	686.0 (4)	
694.4 (4)	694.02 (25)	694 (1)	694.1 (3)	
701.5 (4)	702.08 (24)	701.7 (10)	701.8 (4)	
706.2 (4)	707.29 (28)	706 (1)	706.8 (5)	
713.6 (3)	713.98 (22)	714.1 (6)	713.9 (3)	
718.5 (4)	719.26 (28)	719.1 (6)	719.0 (4)	
760.2 (3)	760.92 (38)	760.3 (6)	760.5 (4)	
	763.8	763.8 (6)	763.8 (6)	

For the relative γ -ray emission probabilities, the data listed in Table 2 were available. The values of 1969Un03 and 1985Ab08 were not listed since they do not have individual uncertainties and those of 1969Sm04 were not used because the ¹⁵³Sm was just a background in an (n, γ) study.

Some gamma emissions with weak intensities and reported by only one or two authors are not listed in Table 2, they are : 54.1 ; 68.2 ; 96.8 ; 118.1 ; 166.5 ; 487.7 ; 574.1 ; 630.5 ; 677.0 ; 682.0 ; 694.1 ; 701.8 ; 706.8 ; 719.0 ; 763.8 keV.

The emission intensities assigned to each of the components of the doublets at 598, 603, 609, 615 and 657 keV are equal, as there is no information on how to split the total intensity for the doublet.

For all cases with three or more values, the weighted average is computed by the Limitation of Relative Statistical Weight method. If the reduced- χ^2 is $>$ critical χ^2 and one value has a relative weight $>$ 50%, the uncertainty of this value is increased in order to reduce the relative weight to 50% and this is noted in the table. If the reduced χ^2 is \leq critical χ^2 , no such change is made, but if the relative weight is over 70% this is noted. For all weighted averages the internal uncertainty is given, and if the reduced χ^2 is $>$ 1.0 the external uncertainty is also given. In some cases the LRSW method expands the uncertainty to include the most precise value; this uncertainty is given as σ_{LRSW} . The adopted values are given in the last row.

The relative γ -ray emission probabilities adopted in Table 2 were normalized to γ 's per 100 decays by consideration of the absolute emission probabilities measured by 1987Co04, 1998Bo18, 1999Sc12 and 2006Le. Of the five γ rays that are given in all papers, the three strongest, at 69, 97, and 103 keV, were considered. Since the weighted average of the data for the 97-keV γ -ray gave a reduced- χ^2 value of 20, it was omitted.

For the 69-keV γ -ray, the weighted average of the four values is 4.668 γ 's per 100 decays with an internal uncertainty of 0.026, a reduced χ^2 of 3.1, and an external uncertainty of 0.047. The latter uncertainty was adopted.

For the 103-keV γ ray, the weighted average of the four values is 29.19 γ 's per 100 decays with an internal uncertainty of 0.12, a reduced χ^2 of 1.8, and an external uncertainty of 0.16. The value of 29.19 (16) was adopted and used to convert the relative values into absolute values as listed in the latest line in Table 2.

Table 3. Absolute emission intensities

	103.18 keV		69.6 keV		97.4 keV	
	I %	Uc	I %	Uc	I %	Uc
1987Co04	29.82	0.36	4.85	0.07	0.847	0.011
1998Bo18	28.5	0.5	4.67	0.05	0.794	0.017
1999Sc12	29.23	0.18	4.65	0.05	0.755	0.007
2006Le	29.07	0.2	4.59	0.05	0.738	0.013
chi2	1.8		3.1		19.7	
WM	29.19	0.16	4.668	0.047	0.778	0.024

X-ray emissions

The measured x-ray emission intensities (in %) are compared with the calculated values deduced from the decay scheme :

XK	K α 2	K α 1	K α	K β ' 1	K β ' 2	K β
1992Ch44			44.43 1.31	8.55 0.29	2.23 0.09	
1999Sc12	16.27 0.18	29.4 0.4	45.7 0.5	9.26 0.12	2.444 0.027	11.7 0.13
2006Le	16.03 0.27	28.53 0.20	44.56 0.3	9.03 0.07	2.37 0.06	11.4 0.12
LWM	16.20 0.15	28.70 0.35	44.85 0.35	9.07 0.10	2.417 0.041	11.54 0.15
Calculated	16.6 0.4	30.0 0.7	46.6 1.1	9.45 0.25	2.44 0.08	11.9 0.3

XL	LI	L α	L β	L γ
1992Ch44	0.190 0.018	4.90 0.26	4.20 0.26	0.651 0.044
1999Sc12	0.216 0.011	4.94 0.11	4.26 0.09	0.615 0.01
2006Le	0.245 0.012	5.06 0.15	4.33 0.13	0.0628 0.022
LWM	0.222 0.014	4.97 0.08	4.28 0.07	0.40 0.22
Calculated	0.213 0.007	5.20 0.15	4.63 0.10	0.755 0.017

6 References

- 1942Ku03 - J. D. Kurbatov, D. C. MacDonald, M. L. Pool, L. L. Quill, Phys. Rev. **61**(1942)106A [T_{1/2}]
1946Mi06 - L. C. Miller, L. F. Curtiss, Phys. Rev. **70**(1946)983 [T_{1/2}]
1950Hi17 - J. M. Hill, L. R. Shepherd, Proc. Phys. Soc. (London) **63A**(1950)126 [P _{β}]
1952Ba49 - R. C. Bannerman, Proc. Phys. Soc. (London) **65A**(1952)565 [P _{β}]
1952Ru10 - W. C. Rutledge, J. M. Cork, S. B. Burson, Phys. Rev. **86**(1952)775 [T_{1/2}]
1954Gr19 - R. L. Graham, J. Walker, Phys. Rev. **94**(1954)794A [P _{β}]
1954Le08 - M. E. Lee, R. Katz, Phys. Rev. **93**(1954)155 [T_{1/2}, P _{β}]
1955Ma62 - N. Marty, J. Phys. Radium **16**(1955)458 [P _{β}]
1956Du31 - V. S. Dubey, C. E. Mandeville, M. A. Rothman, Phys. Rev. **103**(1956)1430 [P _{β}]
1957Jo24 - M. C. Joshi, B. N. Subba Rao, B. V. Thosar, Proc. Indian Acad. Sci. **45A**(1957)390 [P _{β}]
1958Co76 - J. M. Cork, M. K. Brice, R. G. Helmer, R. M. Woods, Jr., Phys. Rev. **110**(1958)526 [T_{1/2}, P _{β}]
1958Gu09 - G. Gueben, J. Govaerts, Inst. Interuniv. Sci. Nucleaires (Bruxelles), Monographie No. 2 (1958) [T_{1/2}]
1960Su08 - R. E. Sund, M. L. Wiedenbeck, Phys. Rev. **120**(1960)1792 [I _{γ} , ce]
1960Wi10 - R. G. Wille, R. W. Fink, Phys. Rev. **118**(1960)242 [T_{1/2}]
1961Gr18 - R. E. Green, W. H. Walker, Can. J. Phys. **39**(1961)1216 [T_{1/2}]
1961Mo07 - E. Monnard, A. Moussa, Nuclear Phys. **25**(1961)292 [ce]
1961Ru01 - L. I. Rusinov, R. L. Aptekar, V. S. Gvozdev, S. L. Sakharov, Yu. L. Khazov, Soviet Phys. JETP **3**(1961)55 [I _{γ}]
1961Wy01 - E. I. Wyatt, S. A. Reynolds, T. H. Handley, W. S. Lyon, H. A. Parker, Nucl. Sci. Eng **11**(1961)74 [T_{1/2}]
1962Ca24 - M. J. Cabell, J. Inorg. Nuclear Chem. **24**(1962)749 [T_{1/2}]
1962Su01 - T. Suter, P. Reyes-Suter, S. Gustafsson, I. Marklund, Nuclear Phys. **29**(1962)33 [ce]
1963Ch25 - P. Chedin, A. Moussa, J. Phys. **24**(1963)930 [ce]
1963Ho15 - D. C. Hoffman, J. Inorg. Nucl. Chem. **25**(1963)1196 [T_{1/2}]
1964Al09 - P. Alexander, Phys. Rev. **134**(1964)B499 [I _{γ}]
1964No08 - T. Novakov, J. M. Hollander, Nucl. Phys. **60**(1964)593 [ce]
1966Bl06 - P. H. Blichert-Toft, E. G. Funk, J. W. Mihelich, Nucl. Phys. **79**(1966)12 [I _{γ}]
1966Ne06 - H. A. Neumann, Z. Naturforsch. **21a** (1966)1328 [ce]
1968Re04 - S. A. Reynolds, J. F. Emery, E. I. Wyatt, Nucl. Sci. Eng. **32**(1968)46 [T_{1/2}]
1969Pa03 - Y. Patin, Compt. Rend. **268B**(1969)574 [E _{γ} , I _{γ}]
1969Sm04 - R. K. Smither, E. Bieber, T. von Egidy, W. Kaiser, K. Wien, Phys. Rev. **187**(1969)1632 [I _{γ} , ce]
1969Un03 - J. Ungrin, M. W. Johns, Nucl. Phys. **A127**(1969)353 [E _{γ} , I _{γ}]
1970Ch09 - Y. Y. Chu, E. M. Franz, G. Friedlander, Phys. Rev. **C1**(1970)1826 [T_{1/2}]
1970Me26 - R. Y. Metskhvarishvili, M. A. Elizbarashvili, V. M. Gachechiladze, L. V. Bodokiya, Bull. Acad. Sci. USSR, Phys. Ser. **34**(1971)1993 [ce]
1970Mi15 - J. Milanovic, R. Stepic, D. Krpic, Fizika **2**(1970)109 [ce]
1970Mu04 - K. Muhlbauer, Z. Phys. **230**(1970)18 [E _{γ}]

- 1970PaZI - Y. Patin, Thesis, Paris Univ. (1970); NP-18835 (1970) [ce]
1971Ba28 - S. Baba, H. Baba, H. Natsume, J. Inorg. Nucl. Chem. **33**(1971)589 [$T_{1/2}$]
1974HeYW - R. L. Heath, report ANCR-1000-2 (1974) [I_{γ}]
1985Ab08 - S. Abdel-Malak, S. M. Darwish, M. Abou -Leila, N. Walley El -Din, A. M. Hassan, Z. Phys. **A322**(1985)163 [E_{γ} , I_{γ}]
1985ZiZY - W. L. Zijp, report ECN FYS/RASA-85/19 (1985) [averaging methods]
1987Co04 - B. M. Coursey, D. D. Hoppes, F. J. Schima, M. P. Unterweger, Appl. Radiat. Isot. **38**(1987)31 [$T_{1/2}$, P_{γ}]
1989Ab05 - A. Abzouzi, M. A. Antony, V. B. Ndocko Ndongue, J. Radioanal. Nucl. Chem.**131**(1989)1 [$T_{1/2}$]
1989Po21 - Yu. S. Popov, N. Yu. Nezhgorov, G. A. Timofeev, Sov. J. Radiochem. **31**(1989)1 [$T_{1/2}$]
1990Le13 - M. A. Lee, Nucl. Data Sheets **60**(1990)419 [$T_{1/2}$]
1992Ch44 - B. Chand, J. Goswamy, D. Mehta, N. Singh, P. N. Trehan, Appl. Radiat. Isot. **43**(1992)997 [I_{γ}]
1992Ra08 - M. U. Rajput and T. D. MacMahon, Nucl. Instr. Meth. **A312**(1992)289 [averaging methods]
1992Un01 - M. P. Unterweger, D. D. Hoppes, F. J. Schima, Nucl. Instr. Meth. **A312**(1992)349 [$T_{1/2}$]
1994Co02 - B. M. Coursey, J. M. Calhoun, J. Cessna, D. B. Golas, F. J. Schima, M. P. Unterweger, Nucl. Instr. Meth. **A339**(1994)26 [$T_{1/2}$]
1995Ch70 - V. P. Chechev, V. O. Sergeev, Bull. Rus. Acad. Sci. Phys. **59**(1995)900
1996Sc06 - E. Schönfeld, H. Janßen, Nucl. Instr. Meth. **A369**(1996)572 [P_x]
1998Bo18 - N. E. Bowles, S. A. Woods, D. H. Woods, S. M. Jerome, M. J. Woods, P. de Lavison, S. Lineham, J. Keightley, and I. Poupaki, Appl. Radiat. Isot. **49**(1998)1345 [P_{γ} , $T_{1/2}$]
1998He06 - R. G. Helmer, Nuclear Data Sheets **83**(1998)285 [multipolarities]
1999Sc12 - U. Schötzgig, E. Schönfeld, E. Günther, R. Klein, H. Schrader, Appl. Radiat. Isot.**51**(1999)169 [$T_{1/2}$, P_{γ}]
2000He14 - R. G. Helmer and C. van der Leun, Nucl. Instr. Meth. **A450**(2000)35 [E_{γ}]
2002Ba85 – I.M.Band, M.B.Trazhaskovskaya, C.W.Nestor, S.Raman. At. Data and Nucl. Data Tables 81, 1&2 (2002) 1 [ICC]
2003Au04 - G. Audi, A. H. Wapstra, C. Thibault. Nucl. Phys. **A729**(2003) 337 [Q]
2004Sc04 – H. Schrader. Appl. Rad. Isotopes **60** (2004) 317 [$T_{1/2}$]
2006Le – M.-C. Lépy, *et al.* Appl. Rad. Isotopes **64** (2006) 1428 [I_{γ} , IXK]

Table 2 : gamma relative and absolute emission intensities (1)

keV	69		75		83		89		97		151		172		412 ^(a)	
1964Al09	1730 ^(o)	100	61	4	75	4	58	3	263	13	3.2	0.5	21 ^(o)	2		
1966B106															0.64	0.2
1969Pa03											5.1 ^(o)	1.6	24	5	0.73	0.13
1974HeYW	1620	140	110 ^(o)	12	63	6	32	4	233	20	3	0.5	28	3	0.8	0.1
1987Co04	1626	21	117 ^(o)	5	68	4			284	4			27	0.4		
1992Ch44	1620	50	55	2	63	2	59	2	255	4	3.5	0.1	25	0.4	0.65	0.02
1998Bo18	1639	18	65	4	58	4			279	6			25.3	1.1		
1999Sc12	1591	17	80 ^(o)	7	72	4	53.4	2.4	258.3	2.4	3.93	0.21	24.5	0.24	0.65	0.04
2006Le	1579	17	61	7	69.8	3.4	37	8	253.9	4.5	3.47	0.21	25	0.6	0.38 ^(o)	0.05
Chi2	1.53		2.02		2.55		10.71		8.14		1.34		4.91		0.63	
Chi2 crit	3.02		3.79		2.80		3.32		2.80		3.32		2.80		3.32	
UWM:	1612.5		60.500		66.971		47.880		260.886		3.42		25.543		0.694	
WM:	1606.644		57.839		66.094		53.918		262.783		3.538		25.151		0.656	
Uc (int):	8.865		1.590		1.276		1.277		1.613		0.081		0.175		0.017	
Uc (ext):	10.967		2.258		2.038		4.181		4.603		0.093		0.388		0.014	
LWM :	1607	11	58	2.3	66.1	2	54	5	262.8	4.6	3.54	0.09	25.2	0.7	0.656	0.017
Abs	4.691	0.041	0.169	0.007	0.193	0.006	0.158	0.015	0.767	0.014	0.01033	0.00027	0.0736	0.0021	0.00191	0.00005

Table 2 : gamma relative and absolute emission intensities (2)

keV	424		436		443		462		463		485		509	
1964Al09														
1966B106	0.75	0.2	0.48	0.12					5.1	0.8	0.12	0.06	0.85	0.16
1969Pa03	0.73	0.13	0.5	0.1			0.5	0.1	4.7	0.4	0.12	0.06	0.61	0.2
1974HeYW	0.7	0.1	0.8 ^(o)	0.1					5.3	0.4			1	0.1
1987Co04														
1992Ch44	0.65	0.02	0.53	0.02	0.030	0.005	0.7	0.2	4.3	0.8	0.13	0.01	0.62	0.03 ^(U)
1998Bo18														
1999Sc12	0.62	0.04	0.57	0.03					4.34	0.06	0.12	0.03	0.63	0.06
2006Le	0.758	0.036	0.546	0.038	0.243	0.041			3.93	0.25			0.46	0.10
Chi2	1.80		0.42		13.49		0.80		2.03		0.05		3.61	
Chi2 crit	3.02		3.32		6.63		6.63		3.02		3.79		3.02	
UWM:	0.701		0.525		0.137		0.60		4.612		0.123		0.695	
WM:	0.669		0.541		0.137		0.540		4.349		0.129		0.651	
Uc (int):	0.016		0.015		0.029		0.089		0.057		0.009		0.030	
Uc (ext):	0.021		0.010		0.107		0.080		0.081		0.002		0.058	
LWM :	0.669	0.021	0.541	0.015	0.140	0.11	0.54	0.09	4.35	0.08	0.129	0.009	0.65	0.06
Abs	0.00195	0.00006	0.001579	0.000045	0.00041	0.00032	0.00158	0.00026	0.01270	0.00024	0.000377	0.000026	0.00190	0.00018

Table 2 : gamma relative and absolute emission intensities (3)

keV	521		531		533		539		542		545		554	
1964Al09														
1966B106	3.5 ^(o)	0.7	22.3	2	11.6	1	9.1	1.4					1.93	0.3
1969Pa03	2.5	0.9	23	3	8.8	2.5	8.2	2.5	0.6	0.5			1.6	0.13
1974HeYW	2.8 ^(o)	0.2	23.8	2	11.9	0.8	8.6	0.6	1.4 ^(o)	0.1	0.3	0.1	2	0.2
1987Co04														
1992Ch44	2.3	0.1	18.9	1.3	10.4	0.1	7.2	0.2	0.77	0.08	0.26	0.01 ^(U)	1.61	0.04
1998Bo18			19.3	2.1	9.8	2.1								
1999Sc12	2.31	0.04	18.37	0.21	10.02	0.09	7.04	0.09	0.75	0.06	0.41	0.17	1.62	0.03
2006Le	2.281	0.024	18.74	0.17	9.91	0.07	7.09	0.05	0.85	0.048	0.368	0.027	1.484	0.047
Chi2	0.15		2.38		4.04		1.84		0.69		2.91		2.35	
Chi2 crit	3.79		2.80		2.80		3.02		3.79		3.79		3.02	
UWM:	2.348		20.63		10.347		7.872		0.743		0.335		1.707	
WM:	2.289		18.646		10.066		7.094		0.803		0.312		1.595	
Uc (int):	0.020		0.13		0.048		0.043		0.034		0.018		0.021	
Uc (ext):	0.008		0.20		0.097		0.058		0.028		0.031		0.032	
LWM :	2.29	0.02	18.65	0.2	10.07	0.16	7.09	0.06	0.803	0.034	0.312	0.031	1.595	0.032
Abs	0.00668	0.00007	0.0544	0.0007	0.02939	0.00049	0.02070	0.00021	0.00234	0.00010	0.00091	0.00009	0.00466	0.00010

Table 2 : gamma relative and absolute emission intensities (4)

keV	578		584		587		590		596		598 ^(d)		603 ^(d)		
1964Al09															
1966B106	1.38	0.2	0.54 ^(o)	0.1					4.4 ^(o)	0.7				2	0.4
1969Pa03	1.15	0.23	0.45	0.15	0.1	0.1	0.45	0.15	4.2 ^(o)	0.6				1.8	0.3
1974HeYW	1.3	0.2	0.4	0.1	0.2	0.03	0.5	0.1	4.5 ^(o)	0.3	0.4	0.1		1.9	0.2
1987Co04															
1992Ch44	1.07	0.03	0.36	0.01	0.16	0.04	0.38	0.01	3.8	0.1	0.61	0.09		1.53	0.05
1998Bo18															
1999Sc12	1.17	0.03	0.352	0.027	0.161	0.027	0.421	0.027	3.56	0.1	0.70	0.03		1.49	0.03
2006Le	1	0.019	0.405	0.02	0.154	0.022	0.448	0.009 ^(U)	3.11	0.05 ^(U)	0.725	0.032		1.388	0.031
Chi2	5.19		1.20		0.52		6.38		17.69		3.50			3.26	
Chi2 crit	3.02		3.32		3.32		3.32		4.61		3.79			3.02	
UWM:	1.178		0.393		0.155		0.440		3.490		0.609			1.685	
WM:	1.063		0.368		0.165		0.417		3.395		0.693			1.462	
Uc (int):	0.015		0.008		0.014		0.007		0.050		0.021			0.020	
Uc (ext):	0.034		0.009		0.010		0.017		0.210		0.039			0.035	
LWM :	1.18	0.18	0.368	0.009	0.165	0.014	0.417	0.031	3.4	0.29	0.693	0.039		1.68	0.19
Abs	0.0034	0.0005	0.001074	0.000027	0.000482	0.000041	0.00122	0.00009	0.0099	0.0008	0.00202	0.00011		0.0049	0.0006

Table 2 : gamma relative and absolute emission intensities (5)

keV	609 ^(d)		615 ^(d)		618		634		636		657 ^(d)		662	
1964Al09														
1966B106	5.5	0.8	0.6 ^(o)	0.12					0.81	0.12	0.13	0.03		
1969Pa03	5.2	0.8	0.21	0.1	0.32	0.14			0.74	0.08	0.12	0.03	0.03	0.01
1974HeYW	5.1	0.4	0.3	0.1	0.3	0.1	0.20	0.03	0.7	0.1	0.1	0.03		
1987Co04														
1992Ch44	4.5	0.1	0.14	0.02	0.2	0.02	0.20	0.05	0.7	0.02	0.14	0.01	0.007	0.002
1998Bo18														
1999Sc12	4.04	0.14	0.233	0.024	0.304	0.027	0.15	0.03	0.595	0.027	0.14	0.024		
2006Le	4.59	0.20	0.159	0.020	0.213	0.022	0.168	0.011	0.65	0.06	0.112	0.009	0.197	0.040
Chi2	2.88		2.80		2.82		0.61		2.45		1.09		11.06	
Chi2 crit	3.02		3.32		3.32		3.79		3.02		3.02		4.61	
UWM:	4.822		0.208		0.267		0.180		0.699		0.124		0.078	
WM:	4.420		0.173		0.230		0.171		0.668		0.125		0.023	
Uc (int):	0.073		0.012		0.013		0.010		0.015		0.006		0.007	
Uc (ext):	0.125		0.020		0.022		0.007		0.023		0.006		0.023	
LWM :	4.42	0.12	0.173	0.020	0.230	0.022	0.171	0.01	0.668	0.023	0.125	0.006	0.023	0.023
Abs	0.01290	0.00036	0.00050	0.00006	0.00067	0.00006	0.000499	0.000029	0.00195	0.00007	0.000365	0.000018	0.00007	0.00007

Table 2 : gamma relative and absolute emission intensities (6)

keV	686		713		760	
1964Al09						
1966B106			0.11	0.03	0.013	0.004
1969Pa03	0.09	0.01	0.066	0.02	0.027	0.015
1974HeYW			0.1	0.03		
1987Co04						
1992Ch44	0.077	0.008	0.077	0.008	0.01	0.002
1998Bo18						
1999Sc12	0.072	0.021	0.09	0.04		
2006Le						
Chi2	0.62		0.53		0.81	
Chi2 crit	4.61		3.32		4.61	
UWM:	0.080		0.089		0.017	
WM:	0.081		0.079		0.011	
Uc (int):	0.006		0.007		0.002	
Uc (ext):	0.005		0.005		0.002	
LWM :	0.081	0.006	0.079	0.007	0.011	0.0018
Abs	0.000236	0.000018	0.000231	0.000020	0.000032	0.000005

^(u) Original uncertainty given, was increased in LRSW analysis to reduce the relative weight to 50%.

^(o) Omitted or outlier

^(a) γ is doubly placed, an undivided intensity is given

¹⁵³Gd - Comments on evaluation of decay data by R. G. Helmer and E. Schönfeld

1 Decay Scheme

In addition to the 5 levels populated in the daughter nucleus, there may be a few others with $J \leq 7/2$ in ¹⁵³Eu, so the completeness of the scheme depends on the failure to observe other γ -rays.

There are some serious discrepancies and ambiguities in the data for some of these five levels.

The recent mass evaluations give the decay energy as 484 keV. However, several measurements of the K-capture probability to the 172-keV level of ¹⁵³Eu (1962Bl11, 1964Cr08, 1967Bo11, 1980Se01, and 1985Si03) have been interpreted to indicate that the decay energy is 235 to 245 keV. In an attempt to resolve this conflict, 1981Gr19 looked for the 166-keV γ -ray which deexcites the 269-keV level and reported an emission probability of 0.0003(3) per 100 decays; so this result is not definitive since it allows 'no population' within the 1σ uncertainty. The problem with the K-capture probability measurements or their interpretation, if any, has not been resolved.

2 Nuclear Data

Q value is from Audi and Wapstra 1995 (1995Au04).

The half-life values available are, in days:

225	1949Ke01	as quoted in 1990Le13
236 (3)	1950He18	
200	1958An34	as quoted in 1990Le13
242 (1)	1963Ho15	
240.9 (6)	1970LyZZ	superseded by 1972Em01 2 nd value
241.6 (2)	1972Em01	
240.9 (6)	1972Em01	
239.63 (4)	1982HoZJ	superseded by 1992Un01 value
226.7 (21)	1989Po21	
239.47 (7)	1992Un01	
240.4 (10)	Adopted value, from LRSW weighted average	

The weighted average of the six remaining values with uncertainties is 239.71 with σ_{int} of 0.07, a reduced- χ^2 of 30.0, and σ_{ext} of 0.36. In the Limitation of Relative Statistical Weight (LRSW) method (1985ZiZY, 1992Ra09), the uncertainty for the 1992Un01 value is increased from 0.07 to 0.185 so that its relative weight is reduced from 88% to 50%. The weighted average is then 240.44 with σ_{int} of 0.13, a reduced- χ^2 of 21.8, and σ_{ext} of 0.61. This method then increases the final uncertainty from 0.61 to 1.0 to include the most precise value, namely, 239.47. In this LRSW analysis, the values of 1972Em01 and 1992Un01 provide 43% and 50% of the relative weight, respectively. The values of 1972Em01, 1989Po21, and 1992Un01 contribute 6.7, 8.6, and 5.5, respectively, to the reduced- χ^2 value.

The value from 1989Po21 differs from this average by about 6σ . The omission of this value would not make a significant difference; in the LRSW analysis without this value the weighted average

would only change to 240.49 with a reduced- χ^2 of 16.6. A more aggressive analysis would increase the uncertainties for the extreme values of 226.7(21) and 241.6(2) and thereby drive the result nearer the value of 1992Un01 and give a smaller final uncertainty. However, the evaluator feels that the larger uncertainty of 1.0 is justified by the large spread in the measured values. This large spread is illustrated by the fact that none of the 1σ ranges of the other five values overlap the value from 1992Un01.

2.1 Electron Capture Transitions

The probabilities for the ϵ branches are from the intensity balances from the γ -ray transition probabilities. It is possible to derive the ϵ intensities because one has a direct measurement of the 97-keV γ -ray emission probability (1987Co04). There is a question as to whether the 151-keV and 269-keV levels are fed in the ¹⁵³Gd decay; see the discussion in section 4.2. In the decay scheme adopted here, they are omitted.

2.2 Gamma Transitions

The multiplicities and mixing ratios are from the ¹⁵³Eu Adopted γ data in the Nuclear Data Sheets (1998He06).

3 Atomic Data

The atomic data are from 1996Sc06.

3.1 and 3.2

The relative K x-ray probabilities are from 1996Sc06.

The x-ray emission probabilities (in %) are:

	RADLST	EMISSION	Measured
K_a	97.2 (21)	96.6 (23)	94.2 (30)
K_b	24.8 (7)	24.6 (7)	24.0 (8)

The EMISSION values were adopted.

The K Auger electron intensities are from RADLST.

4.1 Electron Emission

Data were computed with RADLST for the conversion electrons and for the Auger electrons.

4.2 Photon Emission

From the Helmer and van der Leun evaluation (2000He14), the curved-crystal spectrometer data for the decay of ¹⁵³Sm and ¹⁵³Gd give the energies for the γ -rays of 69.6, 75.4, 83.3, 89.4, 97.4, 103.1, and 172.8 keV on a scale on which the strong line from the decay of ¹⁹⁸Au is 411.80205 (17) keV. In addition, the values from the ¹⁵²Eu(n, γ) study of 1970Mu04 have been adjusted to this energy scale and are used for the γ -rays at 54.1, 68.2, 96.8, 118.1, 151.6, 166.5, and 172.3 keV. The remaining two γ -ray energies, 14.0 and 19.8 keV, were computed from the deduced level energies.

The adopted values for the relative γ -ray emission probabilities were generally taken to be the

weighted averages of the data in the table below. The values for several γ -rays are very discrepant (e.g., χ_R^2 greater than 3.0) and are discussed below. The uncertainties have been chosen by the evaluator as shown in the table. The relative γ -ray emission probabilities given in 1990GeZZ have not been included since they are the same as those in 1992Ch16.

The 21.2-keV γ -ray has not been placed in the scheme.

The values for the 19-keV γ -ray form two groups, namely, the large values of 0.089 (9), 0.072 (11), and 0.06 (2) and the small values of < 0.03, 0.019 (3), and 0.006 (1); so the weighted average does not give a useful value. If one assumes that there is no electron capture feeding of the 83-keV level, a requirement of an intensity balance at this level gives the transition intensity of the 19-keV γ -ray as 1.55 (14) in the units of the table. Then, with $\alpha(19,E2) = 3290$, the γ intensity is $1.55/3291 = 0.00047$ (5). Also, from conversion electron data of 1963Gr09 (a private communication to the ENSDF system), $I_{ce}(LM) = 1.17$ (in the table units), which, with $\alpha(19,E2) = 3290$, gives the γ intensity of 0.0004. If these two independent values are correct, then none of the values in the table are correct, except the upper limit.

The measured intensities of the γ -ray which are proposed to depopulate the 151-keV level are not consistent with those from other modes of populating this level (see the 1998He06 for the other modes of population). These values are :

E_γ	Relative I_γ			
	¹⁵³ Sm β^-	(n, γ)	(d,3n γ)	¹⁵³ Gd ϵ
54	17.1 (18)	26 (4)	25 (3)	330 (130)
68	11 (3)	21.0 (21)	326 (47)	
151	100 (13)	100 (8)	100 (17)	100 (16)

If the ϵ feeding of the 151-keV level in the ¹⁵³Gd decay is simply computed from the intensities of the reported intensities of the 54- and 68-keV γ -rays, it is about 0.2%. On the other hand, the log ft systematics for 2nd forbidden transitions (1998Si17) give log $ft > 11.0$ which corresponds to an upper limit of branch intensity 0.02%. (Also, the intensity data in the table on the next page for the 54- and 151-keV lines are quite discrepant, with reduced- χ^2 values of 121 and 9.1, respectively.) Therefore, no adopted values are given for the 54- and 68-keV γ -rays. [A good new measurement of the intensities of the weak lines is desirable.]

As noted in section 1, it is not known if the level at 269 keV in ¹⁵³Eu is populated in this decay. If it is, the depopulating γ -rays are at 96.8, 118.1, 166.5, and 172.3 keV as shown from other modes of population. From the reported intensity of the 166-keV γ -ray (1981Gr19), this level would be fed in 0.008 (8) % of the decays. This level is omitted here.

The relative γ -ray intensities were normalized to γ 's per 100 decays based on the absolute intensity for the 97-keV line reported by 1990GeZZ; this gives a scaling factor of 0.290 (8), where the published 2σ uncertainty has been divided by 2.

The relative intensities of the K x-rays, on the scale of the table below, are $K_\alpha = 333$ (8) and $K_\beta = 84.8$ (24) as calculated from the decay scheme and 325 (5) and 82.6 (12), respectively, as adopted from the measured values in the table.

Relative Gamma emission Intensities

γ -ray energy (keV)	1974HeYW	1974Se08	1985Si03	1988Su13	1988Ve05	1992Ch16	1992Ch44	1993Eg05	1995Ku34	Weighted average ^e value	σ_{int}	χ_{R^2}	σ_{ext}	σ_{LRSW}	Adopted value
K α_2						114 (2) ^d		114 (4) ^d							
K α		321 (11)	150 (4) ^a	340 (4)	313 (8)		302 (8)		323 (8)	325 (2)		4.5	(5)	(15)	325 (5)
K α_1						204 (4) ^d		208 (8) ^d							
K β_1'						65.2 (14) ^d		65 (3) ^d	69.2 (19)						
K β		78 (11)	32.9 (5) ^a	84.9 (8)	78.9 (11)		76.4 (21)			82.6 (5)		5.3	(12)	(23)	82.6 (12)
K β_2'						17.5 (4) ^d		17.5 (7) ^d	16.84 (26)						
14.0			0.054 (9)	0.146 (15)	0.09 (1)		0.11 (3)	0.10 (3)	0.051 (5) ^g	0.068 (4)		9.2	(13)	(17)	0.068 (17)
19.8			0.089 (9)	0.072 (11)	0.006 (1) ^g		0.06 (2)	< 0.03	0.019 (3)	0.018 (2)		27.5	(10)	^f	0.0004 ⁱ
21.2				0.07 (2)				< 0.03	0.078(16)	0.075 (12)		0.10	(12)	(12)	0.075 (12) ^h
54.1		<0.01	0.091 (3)	0.058 (8)					0.027 (2) ^g	0.057 (2)		121	(22)	(30)	
68.2		0.04 (1)		0.071 (11)	0.035 (14)		0.064 (17)		0.071(11)	0.056 (5)		2.2	(8)	(16)	
69.6	7.8 (2)	8.4 (3)	8.35 (32)	8.60 (15)	8.31 (13)	8.41 (22)	7.97 (20)		8.20 (26)	8.28 (7)		1.9	(10)	(10)	8.28 (10)
75.4	0.30 (3)	0.26 (8)	0.26 (8)	0.278 (31)	0.27 (1) ^g		0.28 (2)		0.26 (2)	0.272 (8)		0.25	(8)	(8)	0.272 (8)
83.3	0.80 (8)	0.70 (7)	0.69 (7)	0.67 (4)	0.69 (3)		0.66 (2)		0.71 (4)	0.680 (14)		0.68	(14)	(14)	0.680 (14)
89.4	0.30 (3)	0.23 (7)	0.23 (6)	0.218 (26)	0.22 (2)		0.29 (2)		0.22 (2)	0.245 (10)		2.12	(14)	(45)	0.245 (14)
97.4	100 (5)	100.	100.	100.	100.0	100 (3)	100.0 (15)	100.	100.0	100					100
103.1	73.5 (10)	71.0 (15)	71.1 (15)	74.8 (7)	69.6 (10)	73.4 (17)	73.7 (12)		72.1 (14)	72.9 (4)		3.2	(7)	(19)	72.9 (7)
151.6	0.0130 (13)	<0.06	0.31 ^b	0.060 (15)	0.02 (1)		<0.010		0.021 (1)	0.0172 (9)		9.1	(27)	(38)	0.017 (4) ^h
172.8	0.130 (13)	0.10 (10)	0.28 ^c	0.144 (26)	0.10 (2)		0.13 (1)		0.12 (1)	0.125 (6)		0.56	(6)	(6)	0.125 (6)

^a Value is uniquely low, omitted from weighted average calculation.

^b Value is uniquely high, omitted from weighted average calculation.

^c No uncertainty, omitted from weighted average calculation.

^d Sum of K α_1 and K α_2 and sum of K β_1' and K β_2' used in weighted average calculation.

^e Limits are omitted from weighted average calculation.

^f LRSW method gives unweighted average of 0.049 (43).

^g LRSW method increased uncertainty in order to reduce relative weight to 50%.

^h Value is not consistent with one upper limit.

ⁱ Computed from γ -ray intensity balance at 83-keV level and $\alpha(19,E2)$ and from internal-conversion electron data and $\alpha(19,E2)$.

6 References

- 1949Ke01 - B. H. Kettle, ORNL-229(1949)34 [T_{1/2}]
 1950He18 - R. E. Hein, A. F. Voigt, Phys. Rev. **79**(1950)783 [T_{1/2}]
 1954Le08 - M. E. Lee, R. Katz, Phys. Rev. **93**(1954)155 [T_{1/2}]
 1958An34 - N. M. Antoneva, A. A. Bashilov, B. S. Dzhelepov, B. K. Preobrazhenskii, Columbia Tech. Transl. **22**(1959)134 [T_{1/2}]
 1958Co76 - J. M. Cork, M. K. Brice, R. G. Helmer, R. M. Woods, Jr., Phys. Rev. **110**(1958)526 [T_{1/2}]
 1961Ca24 - M. J. Cabell, J. Inorg. Nucl. Chem. **24**(1961)749 [T_{1/2}]
 1961Gr18 - R. E. Green, W. H. Walker, Can. J. Phys. **39**(1961)1216 [T_{1/2}]
 1961Wy01 - E. I. Wyatt, S. A. Reynolds, T. H. Handley, W. S. Lyon, H. A. Parker, Nucl. Sci. Eng. **11** (1961)74 [T_{1/2}]
 1962Bl11 - L. Blok, W. Goedbloed, E. Mastenbroek, J. Blok, Physica **28**(1961)993 [P_K]
 1963Ho15 - D. C. Hoffman, J. Inorg. Nucl. Chem. **25**(1963)1196 [T_{1/2}]
 1964Al09 - P. Alexander, Phys. Rev. **134** 1964)B499 [P_γ]
 1964Cr08 - T. Cretzu, K. Hohmuth, G. Winter, Nucl. Phys. **56**(1964)415 [P_K]
 1966Bl06 - P. H. Blichert-Toft, E. G. Funk, J. W. Mihelich, Nucl. Phys. **79**(1966)12 [P_γ]
 1967Bo11 - P. Boyer, P. Chedin, J. Oms, Nucl. Phys. **A99**(1967)213 [P_K]
 1969Pa03 - Y. Patin, Compt. Rend. **268B**(1969)574 [E_γ, P_γ]
 1969Sm04 - R. Smither, E. Bieber, T. von Egidy, W. Kaiser, K. Wien, Phys. Rev. **187**(1969)1632 [E_γ]
 1969Un03 - J. Ungrin, M. W. Johns, Nucl. Phys. **A127**(1969)353 [E_γ, P_γ]
 1970Ch09 - Y. Y. Chu, E. M. Franz, G. Friedlander, Phys. Rev. **C1**(1970)1826 [T_{1/2}]
 1970LyZZ - W. S. Lyon, H. H. Ross, L. C. Bate, F. F. Dyer, J. S. Eldridge, J. F. Emery, T. H. Handley, H. Kubota, S. B. Lupica, S. A. Reynolds, E. Ricci, J. E. Strain, H. E. Zittel, K. J. Northcutt, ORNL-4636 (1970)24 [T_{1/2}]
 1970Mu04 - K. Muhlbauer, Z. Phys. **230**(1970)18 [E_γ]
 1971Ba28 - S. Baba, H. Baba, H. Natsume, J. Inorg. Nucl. Chem. **33**(1971)589 [T_{1/2}]
 1972Em01 - J. F. Emery, S. A. Reynolds, E. I. Wyatt, G. I. Gleason, Nucl. Sci. Eng. **48**(1972)319 [T_{1/2}]
 1974HeYW - R. L. Heath, report ANCR-1000-2 (1974)[P_γ]
 1974Se08 - V. A. Sergienko, V. M. Lebedev, Bull. Acad. Sci. USSR, Phys. Ser. **38**, no. 4 (1974)122 [P_γ]
 1980Se01 - V. A. Sergienko, Ts. Vylov, S. M. Sergeev, S. L. Smolskii, Bull. Acad. Sci. USSR, Phys. Ser. **44**, No. 1 (1980)100 [P_K]
 1982HoZJ - D. D. Hoppes, J. M. R. Hutchinson, F. J. Schima, M. P. Unterweger, NBS-SP-626(1982)85 [T_{1/2}]
 1985Si03 - K. Singh, B. S. Grewal, H. S. Sahota, J. Phys. (London) **G11**(1985)399 [P_K]
 1985Ab08 - S. Abdel-Malak, S. M. Darwish, M. Abou-Leila, N. Walley El-Din, A. M. Hassan, Z. Phys. **A322**(1985)163 [E_γ, P_γ]
 1985ZiZy - W. L. Zijp, report ECN-179, Petten (1985) [analysis methodology]
 1986BrZQ - E. Browne, R. B. Firestone, Table of Radioactive Isotopes, Appendix C, Table 7a (John Wiley & Sons, New York, 1986) [P_x]
 1987Co04 - B. M. Coursey, D. D. Hoppes, F. J. Schima, M. P. Unterweger, Appl. Radiat. Isot. **38**(187)31 [P_γ]
 1988Su13 - S. Subrahmanyeswara Rao, K. Bhaskara Rao, V. Seshagiri Rao, H. C. Padhi, J. Phys. G: Nucl. Phys. **14**(1988)1259 [P_γ]
 1988Ve05 - N. Venkateswara Rao, G. S. Sri Krishna, S. Bhuloka Reddy, C. V. Raghavaiah, G. Satyanarayana, D. L. Sastry, Il Nuove Cimento **99A**(1988)303 [P_γ]
 1989Po21 - Yu. S. Popov, N. Yu. Nezgovorov, G. A. Timofeev, Sov. J. Radiochem. **31**(1989)1 [T_{1/2}]
 1990GeZZ - A. M. Geidelman, A. G. Egorov, Yu. S. Egorov, V. G. Nedovesov, G. E. Shchukin, Ann. Conf. Nucl. Spectrosc. Struct. Atomic Nuclei(1990)485 [P_γ]
 1990Le13 - M. A. Lee, Nucl. Data Sheets **60**(1990)419 [T_{1/2} data]
 1992CH16 - V. P. Chechev, A. G. Egorov, Nucl. Instr. Meth. **A312**(1992)378 [P_γ]
 1992Ch44 - B. Chand, J. Goswamy, D. Mehta, N. Singh, P. N. Trehan, Appl. Radiat. Isot. **43**(1992)997 [P_γ]
 1992Ra08 - M. U. Rajput, T. D. Mac Mahon, Nucl. Instr. Meth. **A 312**(1992)289 [analysis methodology]
 1992Un01 - M. P. Unterweger, D. D. Hoppes, F. J. Schima, Nucl. Instr. Meth. **A312**(1992)349 [T_{1/2}]
 1993Eg05 - A. G. Egorov, V. P. Chechev, G. E. Shchukin, Bull. Russ. Acad. Sci. Phys. **57**, no. 5 (1993) 911 [P_γ]
 1994Co02 - B. M. Coursey, J. M. Calhoun, J. Cessna, D. B. Golas, F. J. Schima, M. P. Unterweger, Nucl. Instr. Meth. **A339**(1994)26 [T_{1/2}]
 1995Au04 - G. Audi, A. H. Wapsrta, Nucl. Phys. **A595**(1995)409 [Q]
 1995Ku34 - V. Kumar, Kawaldeep, K. Singh, J. Radioanal. Nucl. Chem. **189**(1995)3 [P_γ]

Comments on evaluation

- 1995ScZY - E. Schönfeld, Appl. Radiat. Isot. **49**(1998)1353 [P_K,P_L]
1996Sc06 - E. Schönfeld, H. Janßen, Nucl. Instr. Meth. **A369**(1996)527 [P_x]
1998He06 - R. G. Helmer, Nuclear Data Sheets **83**(1998)285 [Multipolarities]
2000He14 - R. G. Helmer and C. van der Leun, Nucl. Instr. Meth **A450**(2000)35 [E_γ]

¹⁵⁴Eu – Comments on evaluation of decay data by V. P. Chechev and N. K. Kuzmenko

This evaluation was done in June 1999, and revised in January 2003. The literature available by 2003 was included.

1. Decay Scheme

The decay scheme is based on the evaluation of Reich (1998Re22).

The ¹⁵⁴Eu→¹⁵⁴Gd decay scheme has not been completed yet as there are a few unplaced ¹⁵⁴Gd gamma transitions. These transitions are weak, so they do not greatly influence the intensity balances.

The 3rd forbidden β⁻ transitions to the ground states of ¹⁵⁴Gd and ¹⁵⁴Sm have not been observed. From the log ft systematics (1998Si17), their log ft values should be greater than 17,6 and the corresponding upper limits of their intensities would be expected less than 5·10⁻⁵ % and less than 3·10⁻⁷ %, respectively.

In the “Adopted Levels” of 1998Re22, there are several ¹⁵⁴Gd levels with energies below Q⁻ that have not been observed in the ¹⁵⁴Eu β⁻ decay. Their energies are 1900,2; 1911,5; 1912,1; 1943,9; 1948,5 and 1963,8 keV. Their respective spins and parities are not known exactly except those for the 1911,5 keV, which is a 6⁺ level. The β⁻ transition to this 1911,5 keV level is 3rd forbidden and its intensity is expected to be less than 5·10⁻¹⁰ % (log ft > 17,6). On the assumption that the remaining levels can be populated by β⁻ transitions with an order of forbiddenness not lower than 2, their log ft values should be greater than 11 and their corresponding branch intensities expected to be less than 0,001%.

Likewise, the intensity of the 3rd forbidden electron-capture transition to the ¹⁵⁴Sm 543,7 keV 6⁺ level in the decay ¹⁵⁴Eu→¹⁵⁴Sm is expected to be less than 10⁻⁸ % (from log ft > 17,6).

Therefore, all of the above transitions can be neglected, and thus they are not shown in the ¹⁵⁴Eu decay scheme.

2. Nuclear Data

Q⁺, Q⁻ values are from 1995Au04.

The evaluated half-life of ¹⁵⁴Eu has been obtained by applying the evaluation procedure from 2000Ch01 (Chechev and Egorov). This value is based on the measured results given in Table .

Table 1. Set of experimental data for the evaluation of ¹⁵⁴Eu half-life (in days)

Reference	Author	Data set "1" $\chi^2=22,83$ $(\chi^2)_8^{0,05}=15,51$	Data set "2" $\chi^2=22,79$ $(\chi^2)_7^{0,05}=14,07$	Data set "3" $\chi^2=22,79$ $(\chi^2)_7^{0,05}=14,07$
2002Un02	Unterweger	3145,2(11) ^a	3145,2(11)	3145,2(11)
1998Si12	Siegert et.al	3138,1(16) ^b	3138,1(16)	3138,1(16)
1998Si12	Siegert et.al	3146(11) ^c	3146(11)	3146(11)
1983Th04	Thompson et.al	3170(55)	3170(55)	3170(55)
1992ScZZ	Schötzig et.al	3139,0(20)	3139,0(20)	3139,0(20)
1988RaZM	Rajput et.al	3143(59)	3143(59)	3143(59)
1986Wo05	Woods et.al	3138,0(20)	3138,0(20)	3138,0(20)
1983Wa26	Walz et.al	3136(4)	3136(4)	3136(4)
1972Em01	Emery et.al	3105(180)	Omitted ^d	-

^a Latest value from this laboratory. Previous measurements at NIST gave 3101(41) – 1982 HoZJ and 3138,2(61) – 1992Un01.

^b Measured with a pressured 4πγ ionization chamber.

^c Measured with semiconductor detectors.

^d Omitted on the basis of statistical considerations.

Data set "1" is the original data; set "2" has the discrepant values deleted, and set "3" would have the uncertainty increased for any value having more than 50% of the relative weight. There are none of the latter values, so set "3" is the same as set "2".

It should be noted that there are available the early half-life measurement results which have been omitted because of the very low accuracy: 5,4 years (without uncertainty) – 1949Ha04 and 16(4) years – 1952Ka26. There are also unpublished measurement results of 1978ScZO (7,45 - 10,5 years) and 1978GrZR (8,8(1) years) which have not been included in the set "1".

The weighted mean of data from the final set "3" is 3141,5(14) where the uncertainty has been obtained as an external uncertainty 1,35 multiplied by the Student's coefficient at the confidence level of 0,68 for 7 degrees of freedom (see 2000Ch01). The internal uncertainty is 0,75.

The adopted value of the ¹⁵⁴Eu half-life is 3141,5(14) days, or 8,601(4) years (converted to years with 365,24219 d/y).

2.1. β^- Transition and Electron Capture Transition

2.1.1. β^- Transitions

The energies of β^- transitions have been computed from the Q^- value and the level energies adopted from 1998Re22. The corrections to the level energies taking into account the evaluated values of gamma transition energies from section 2.2 are negligible.

The probabilities of β^- transitions have been obtained from the $P(\gamma+ce)$ balance for each level of ¹⁵⁴Gd based on the $P(\gamma)$ normalization factor of 0,3489(34) (see section 4.2.). Since 0,018 % (13) of the decays are *via* electron capture, the value of $P_{\beta_1}=10,3(5)$, to the first excited level in ¹⁵⁴Gd, has been obtained from $P_{\beta_1}=99,982(13) - \sum P_{\beta_i}, i>1$. From the $P(\gamma+ce)$ balance for this level $P_{\beta_1}=10,5(13)$. The more precise value has been adopted.

The more inaccurate experimental values from 1966Ha36 and 1968Ng01 obtained by direct measurements using magnetic beta-spectrometry and beta-gamma coincidences do not conflict with the calculated ones, as seen from Table 2 (except $\beta_{0,2}$).

Table 2. Comparison of the measured and evaluated (calculated) values of β^- transition probabilities.

	E_{β} , keV	P_{β} , % 1966Ha36	P_{β} , % 1968Ng01	Evaluated (calculated) values
$\beta_{0,26}$	248,8(11)		29,1(25)	28,32(22)
$\beta_{0,16}$	570,9(11)		37,8(35)	36,06(35)
$\beta_{0,8}$	840,6(11)		17,0(39)	17,33(18)
$\beta_{0,6}$	972,1(11)		4,6(38)	2,82(18)
$\beta_{0,5}$	1152,9(11)		0,67(49)	0,33(3)
$\beta_{0,2}$	1597,4(11)	0,19(5)		0,31(7)
$\beta_{0,1}$	1845,3(11)	9,2(15)	10,8(12)	10,3(5)

We are listing below the ¹⁵⁴Gd levels from the ¹⁵⁴Eu β⁻ decay (see 1998Re22).

Level number	Energy, keV	Spin and parity	Half-life	Probability of β ⁻ transition (× 100)
0	0,0	0 ⁺	Stable	
1	123,071	2 ⁺	1,18 ns	10,3(5)
2	371,00	4 ⁺	45 ps	0,31(7)
3	680,66	0 ⁺	4,0 ps	
4	717,7	6 ⁺	7,8 ps	
5	815,5	2 ⁺	6,4 ps	0,33(3)
6	996,26	2 ⁺	0,95 ps	2,82(18)
7	1047,6	4 ⁺		0,108(18)
8	1127,8	3 ⁺		17,33(18)
9	1136,0	1,2 ⁺		
10	1233,2			
11	1241,3	1 ⁻		
12	1251,6	3 ⁻		0,289(6)
13	1263,78	4 ⁺		0,707(7)
14	1277,0			
15	1294,2	(2) ⁺		
16	1397,5	2 ⁻		36,06(35)
17	1414,4	1 ⁻		
18	1418	2 ⁺		0,075(2)
19	1510,1	(1 ⁻)		0,021(2)
20	1531,3	2 ⁺		0,330(13)
21	1560,0	(4 ⁻)		0,100(4)
22	1617,1	3 ⁻		1,78(3)
23	1645,8	4 ⁺		0,148(4)
24	1660,9	3 ⁺		0,849(9)
25	1698,5	(4 ⁺)		0,0100(4)
26	1719,56	2 ⁻		28,32(22)
27	1770,2	5 ⁺		0,0022(4)
28	1790,2	(4 ⁺)		0,022(1)
29	1797,0	3 ⁻		0,060(6)
30	1838,6	2 ⁺		0,017(5)
31	1861,5	4 ⁻		0,034(3)
32	1878,5			0,0042(3)
33	1894,7	2 ⁺		0,0035(6)

2.1.2. Electron Capture Transitions

The energies of the electron capture, ε, transitions have been calculated from the Q⁺ value and the level energies from 1998Re22 (see below).

List of ¹⁵⁴Sm levels from the ¹⁵⁴Eu electron capture decay

Level number	Energy, keV	Spin and parity	Half-life	Probability of electron capture (× 100)
0	0,0	0 ⁺	Stable	
1	81,98	2 ⁺	3,02 ns	0,013(13)
2	266,79	4 ⁺	172 ps	0,0047(8)
3	543,73	6 ⁺	22,7 ps	

The transition probabilities have been obtained from the $P(\gamma+ce)$ balance for each ¹⁵⁴Sm level using a $P(\gamma)$ normalization factor of 0,3489(34).

Fractional electron capture probabilities P_K , P_L , P_M have been calculated from 1998Sc28 using the program EC-CAPTURE.

2.2. Gamma Transitions and Internal Conversion Coefficients

The evaluated energies of gamma-ray transitions include the recoil energy of $E_\gamma^2/2Mc^2$, where M is mass of the daughter nucleus (¹⁵⁴Gd or ¹⁵⁴Sm).

The gamma-ray transition probabilities have been deduced from their emission probabilities and total internal conversion coefficients (ICC).

The ICC are theoretical values from 1978Ro22 for the adopted energies and multiplicities. Other values have been taken from the evaluation 1998Re22, based on experimental data from 1957Ke08, 1962Lu03, 1966Za02, 1969An01, 1972Na21, 1977Ya04 and 1996Al31. Total ICC values for $\gamma_{1,0}(\text{Gd})$ have been obtained as weighted averages of measured values, 1,200(20) - 1962Lu03 and 1,194(19) - 1995Ma03, and taking into account the rule of "the smallest experimental uncertainty" (see 2000Ch01).

The relative uncertainties of α_K , α_L , α_M for pure multiplicities have been adopted 2%.

3. ATOMIC DATA

3.1. Fluorescence Yields

The fluorescence yield data are from 1996Sc06 (Schönfeld and Janßen).

3.2. X-Radiations

The X-ray energies are based on their wavelengths in the compilation of 1967Be65 (Bearden). The relative KX-ray emission probabilities have been taken from 1996Sc06 and 1999Schönfeld.

3.3. Auger Electrons

The energies of Auger electrons are from 1977La19 (Larkins) and 1987Lagoutine.

The ratios $P(\text{KLX})/P(\text{KLL})$, $P(\text{KXY})/P(\text{KLL})$ are taken from 1996Sc06.

4. PHOTON EMISSIONS

4.1. X-Ray Emissions

The total absolute emission probability of Gd KX -rays has been computed using the adopted value of $\omega_K(\text{Gd})$ and the evaluated total absolute emission probability of K conversion electrons in the decay ¹⁵⁴Eu→¹⁵⁴Gd, namely, $P_{ceK} = 27,3(6)\%$. The emission probability of Sm KX -rays has been computed using the adopted value of $\omega_K(\text{Sm})$, the evaluated probability of K electron capture to ¹⁵⁴Sm levels $P_{eK} = 0,015(11)\%$ and the evaluated emission probability of K conversion electrons in the decay ¹⁵⁴Eu→¹⁵⁴Sm, namely, $P_{ceK} = 0,007(4)\%$.

The absolute emission probabilities of the Gd KX-ray components have been computed using the relative probabilities from Section 3.2 and the total value of $P_{XK}(\text{Gd}) = 25,4(6)\%$.

4.2. Gamma-Ray Emissions

The energies of prominent gamma -rays $\gamma_{1,0}(123,1)$, $\gamma_{2,1}(247,9)$, $\gamma_{5,2}(444,5)$, $\gamma_{26,8}(591,7)$, $\gamma_{6,2}(625,2)$, $\gamma_{5,1}(692,4)$, $\gamma_{26,6}(723,3)$, $\gamma_{8,2}(756,8)$, $\gamma_{24,5}(845,4)$, $\gamma_{6,1}(873,2)$, $\gamma_{13,2}(892,8)$, $\gamma_{26,5}(904,1)$, $\gamma_{12,1}(1128,5)$, $\gamma_{13,1}(1140,7)$, $\gamma_{22,2}(1246,1)$, $\gamma_{16,1}(1274,4)$, $\gamma_{22,1}(1494,0)$, $\gamma_{26,1}(1596,5)$ have been taken from 2000He14 (Helmer and Van der Leun).

The energies of the gamma rays $\gamma_{26,20}(188,2)$, $\gamma_{16,6}(401,2)$, $\gamma_{26,12}(467,8)$, $\gamma_{26,11}(478,3)$, $\gamma_{3,1}(557,6)$, $\gamma_{16,5}(582,0)$, $\gamma_{7,2}(676,6)$, $\gamma_{20,5}(715,8)$, $\gamma_{5,0}(815,5)$, $\gamma_{20,3}(850,6)$, $\gamma_{12,2}(880,6)$, $\gamma_{7,1}(924,6)$, $\gamma_{6,0}(996,3)$,

$\gamma_{8,1}(1004,7)$, $\gamma_{11,1}(1118,5)$, $\gamma_{20,2}(1160,4)$, $\gamma_{21,2}(1188,1)$, $\gamma_{11,0}(1241,4)$, $\gamma_{24,2}(1290,5)$, $\gamma_{19,1}(1397,4)$, $\gamma_{24,1}(1537,8)$ have been evaluated using the experimental data of 1990He05, 1992Sm02, 1990Me15 along with taking into account a correction of the gamma -ray energetic scale in 2000He14 (lowering by 5,8 ppm) (Table 3).

Table 3. Measured and evaluated values of some gamma ray energies in the decay of ¹⁵⁴Eu (keV)

	1990He05	1990Me05	1992Sm02	Evaluated
$\gamma_{26,20}$	188,252(8)	188,22(4)	188,29(7)	188,24(2)
$\gamma_{16,6}$	401,258(14)	401,30(5)		401,259(14)
$\gamma_{26,12}$	467,84(5)			467,84(5)
$\gamma_{26,11}$		478,26(5)	478,29(7)	478,27(5)
$\gamma_{3,1}$		557,56(5)	557,61(7)	557,58(5)
$\gamma_{16,5}$		582,00(5)	582,03(7)	582,01(5)
$\gamma_{7,2}$	676,600(12)	676,60(5)		676,596(12)
$\gamma_{20,5}$	715,786(18)	715,77(5)	715,75(7)	715,77(3)
$\gamma_{5,0}$		815,57(5)	815,45(7)	815,53(5)
$\gamma_{20,3}$	850,643(12)	850,66(5)	850,61(7)	850,64(3)
$\gamma_{12,2}$	880,61(3)			880,60(3)
$\gamma_{7,1}$	924,64(5)			924,63(5)
$\gamma_{6,0}$	996,262(6)	996,35(4)	996,21(3)	996,25(5)
$\gamma_{8,1}$	1004,725(7)	1004,79(4)	1004,67(3)	1004,718(7)
$\gamma_{11,1}$		1118,53(6)		1118,52(6)
$\gamma_{20,2}$	1160,37(8)			1160,36(8)
$\gamma_{21,2}$	1188,10(4)	1188,60(10)		1188,34(17)
$\gamma_{11,0}$	1241,38(5)	1241,62(9)		1241,43(10)
$\gamma_{24,2}$	1290,51(10)			1290,50(10)
$\gamma_{19,1}$	1397,35(5)			1397,34(5)
$\gamma_{24,1}$	1537,80(4)	1537,84(5)		1537,81(4)

The energies of the gamma rays $\gamma_{15,8}(165,9)$, $\gamma_{22,17}(202,5)$, $\gamma_{14,7}(229,0)$, $\gamma_{22,5}(801,2)$ have been taken from 1992E111. The energy of the gamma ray $\gamma_{1,0}$ Sm (82,0) has been adopted from measurements of conversion electrons (1958Ch36). The unplaced gamma ray 197 keV has been reported in 1980Sh15 and 1989Ki10. The energy of the gamma ray $\gamma_{7,4}(329,9)$ has been adopted from 1974HeYW. The energy 533,1 keV (twice placed - $\gamma_{24,8}$ and $\gamma_{29,13}$) has been computed from the level energies. The energy and relative emission probability of the gamma ray $\gamma_{3,0}(680,7)$ has been taken from 1969An01. The energy of the unplaced gamma-ray γ 1316,4 keV has been adopted from 1970Ri19.

The energies of the remaining weak gamma rays have been taken from 1968Me18.

The measured and evaluated values of relative gamma ray emission probabilities are shown in Table 4.

Table 4. Measured and evaluated values of relative gamma ray emission probabilities in the decay of ¹⁵⁴Eu

keV	1968Me18	1969Va09	1970RiZY	1980Ro22	1980Sh15	1984Iw03	1986Wa35	1989Ki10	1989 Schima	1990Me15	1990He05	1992E111	1992Ha02	1992Sm02	1992Sa04	Evaluated value
58,4	0,0113(11)															0,0113(11)
80,4	0,008(4)															0,008(4)
82,0	0,009(6)															0,009(6)
123,1			116(6)		115,4(23)	118,5(13)	111,7(16)	122,1(36)	117,0(11)	114,1(20)	116,5(12)		115,6(15)	113,0(15)	115,4(7)	115,9(8)
125,4	0,0197(56)															0,020(6)
129,5	0,039(6)															0,039(6)
131,6	0,0310(14)				0,037			0,025					0,035(3)			0,0317(13)
134,8	0,0203(11)				0,03			0,024					0,027(6)			0,0205(11)
146,0	0,073(3)		0,085(27)		0,12(1)			0,078(28)					0,075(10)			0,074(3)
156,2	0,0282(12)				0,025			0,019					0,027(3)			0,0280(11)
159,9	<0,003												0,0030(15)			0,0030(15)
162,1	0,0028(14)												0,0035(17)			0,0031(11)
165,9	0,0065(14)				0,021			0,019					0,012(4)			0,0071(14)
180,7	0,0127(28)	0,0058(58)			0,015			<0,001					0,0116(17)			0,0115(17)
184,7	0,0113(28)				0,017			0,003					0,010(3)			0,011(3)
188,2		0,692(17)	0,61(12)		0,70(12)			0,88(10)		0,682(22)			0,658(27)	0,651(15)		0,684(15)
195,5	0,0056(28)															0,006(3)
197					0,005			0,004								0,0045(5)
202,5												0,08(2)				0,08(2)
209,4	0,0068(23)												0,0072(16)			0,0071(16)
219,4	0,0065(25)												0,0067(19)			0,0066(19)
229,0	0,0056(22)												0,0085(25)			0,0069(22)
232,0	0,0677(30)		0,079(43)		0,081(40)			0,059(22)					0,068(6)			0,068(3)
237,0	0,017(11)				0,026			0,024					0,019(9)			0,018(9)
247,9			20,1(10)	20,51(20)	19,34(37)	19,91(14)	19,615(98)	23,04(59)	19,82(16)	19,72(32)	19,8(2)		19,65(44)	19,5(2)	19,857(93)	19,76(9)
260,9	0,0056(25)							0,017					0,0066(20)			0,0062(20)
267,4	0,039(2)				0,023			<0,001					0,037(7)			0,039(2)
269,8	0,0197(28)				0,01			0,017					0,022(4)			0,0205(28)
274,0	0,0113(6)												0,0105(12)			0,0111(6)
279,9	0,0085(4)												0,0092(21)			0,0085(4)
290,0	0,0096(5)												0,010(2)			0,0096(5)
295,7	0,0068(4)												0,0073(15)			0,0068(4)
296,0	0,0039(25)															0,004(3)
301,3	0,0282(12)				0,032			0,03					0,032(2)			0,0292(12)
305,1	0,0496(22)	0,058(12)			0,07			0,078					0,055(7)			0,050(2)
308,2	≤0,005				0,01								0,0068(17)			0,0068(17)
312,3	0,0414(19)	0,055(12)			0,06			0,069					0,059(5)			0,053(4)
315,4	0,0130(7)	0,037(12)			0,03			0,027					0,027(6)			0,021(4)
320	0,0028(20)															0,0028(20)

keV	1968Me18	1969Va09	1970RiZY	1980Ro22	1980Sh15	1984Iw03	1986Wa35	1989Ki10	1989 Schima	1990Me15	1990He05	1992Ei11	1992Ha02	1992Sm02	1992Sa04	Evaluated value
322,0	0,189(9)	0,193(9)	0,16(4)		0,21(4)			0,168(22)					0,189(10)			0,189(9)
329,9	0,0259(4)		0,036(26)		0,032			0,023					0,031(10)			0,0260(14)
346,7	0,085(3)				0,067								0,075(6)			0,083(3)
368,2	0,0085(4)												0,0081(17)			0,0085(4)
370,7	0,015(4)				0,03			0,007					0,018(6)			0,016(4)
375,2	0,0051(28)												0,0059(23)			0,0056(23)
382,0	0,0285(12)				0,028			0,006					0,027(3)			0,0283(12)
397,1	0,085(3)	0,066(9)	0,12(5)		0,12(4)			0,070(16)					0,076(8)			0,082(3)
401,3		0,55(3)	0,58(10)		0,57(8)	0,49(4)		0,58(6)		0,56(3)	0,543(6)		0,54(3)			0,543(6)
403,5	0,076(3)		0,054(32)		0,042(40)								0,067(8)			0,075(3)
414,3	0,0141(18)												0,015(2)			0,0142(18)
419,4	0,011(6)												0,0094(41)			0,010(6)
422,1	≤0,0034												0,0062(24)			0,0062(24)
435,9	≤0,0073												0,011(3)			0,011(3)
444,5		1,64(4)	1,69(15)	1,53(6)	1,54(3)	1,63(3)	1,87(11)	2,11(6)		1,58(3)	1,600(15)		1,66(7)	1,628(17)	1,564(38)	1,606(15)
463,9	0,0121(7)												0,019(8)			0,0122(7)
467,8	0,161(7)	0,173(17)	0,20(9)		0,16(8)			0,18(3)					0,184(7)			0,173(7)
478,2		0,605(22)	0,69(15)		0,63(10)	0,626(27)		0,64(5)		0,68(3)	0,644(6)		0,63(3)	0,648(12)		0,643(6)
480,6	0,0138(8)															0,0138(8)
483,7	0,0141(8)				0,04			0,045					0,033(12)			0,0142(8)
484,6	0,0113(6)															0,0113(6)
488,3	0,020(9)												0,021(10)			0,020(9)
506,4	0,017(6)							0,017					0,018(4)			0,018(4)
510	0,103(5)		0,17(8)		0,14(8)			0,28(5)					0,19(3)			0,17(2)
512,0	≤0,17	0,092(20)														0,092(20)
518,0	0,132(6)	0,144(26)	0,16(9)		0,18(8)			0,17(5)					0,144(18)			0,135(6)
533,1 \$	0,031(6)				0,032			0,04					0,034(8)			0,032(6)
545,6	0,047(6)	0,035(29)											0,036(6)			0,041(6)
557,6		0,75(3)	0,74(10)		0,72(10)	0,758(24)		0,80(10)		0,73(3)	0,778(11)		0,75(3)	0,767(12)		0,767(11)
563,4												0,008(2)				0,008(2)
569,2	0,0282(12)				0,044			0,024					0,0410(64)			0,0286(23)
582,0		2,62(7)	2,53(23)	2,86(11)	2,45(5)	2,61(3)	2,45(5)	2,72(12)		2,51(3)	2,543(2)		2,53(3)	2,53(23)		2,54(2)
591,7		14,44(31)	14,8(8)	13,62(24)	13,57(26)	14,35(6)	14,05(14)	15,84(66)	14,19(11)	14,14(15)	14,21(11)		14,18(31)	14,0(14)	14,338(117)	14,18(7)
597,5	0,0158(9)															0,0158(9)
598,3	0,0172(10)				0,026								0,0280(54)			0,0176(21)
600,0	0,017(11)															0,017(11)
602,8					0,1			0,15					0,096(8)			0,096(4)
613,3	0,262(11)	0,288(20)	0,22(8)		0,25(8)			0,29(7)					0,265(19)			0,267(11)
620,5	0,0262(14)												0,023(6)			0,0260(14)
625,2		0,922(32)	0,89(12)		0,84(5)	0,927(21)	0,90(4)	0,92(9)		0,90(3)			0,91(2)	0,906(10)		0,909(10)
642,4	0,011(6)							0,040(28)					0,013(5)			0,013(5)

keV	1968Me18	1969Va09	1970RiZY	1980Ro22	1980Sh15	1984Iw03	1986Wa35	1989Ki10	1989 Schima	1990Me15	1990He05	1992Ei11	1992Ha02	1992Sm02	1992Sa04	Evaluated value
649,4	0,214(9)		0,28(11)		0,25(8)			0,30(10)					0,26(2)			0,223(9)
650,6	0,0282(12)															0,0282(12)
664,7	0,082(3)				0,072			0,03					0,088(15)			0,082(3)
668,9	0,034(8)				0,042			0,031					0,042(7)			0,038(7)
676,6		0,432(30)	0,43(11)		0,52(10)	0,47(5)	0,45(27)	0,53(11)		0,45(3)			0,46(5)			0,45(3)
692,4		5,07(13)	4,97(30)	4,86(8)	4,92(10)	5,182(29)	5,14(5)	5,75(15)		5,10(9)	5,09(4)		5,13(12)	5,04(5)	5,085(59)	5,12(3)
715,8		0,40(6)	0,32(13)		0,61(8)			0,27(12)		0,592(28)			0,52(2)	0,57(3)		0,54(3)
723,3		56,5(12)	60,1(31)	55,40(41)	55,33(106)	58,19(27)	57,23(46)	64,9(21)	57,6(4)	57,2(6)	57,3(4)		57,78(89)	56,9(6)	58,107(276)	57,46(27)
737,6	≤0,024												0,018(7)			0,018(7)
756,8		12,71(23)	12,9(6)	12,51(11)	12,62(24)	13,18(8)	12,89(13)	13,61(20)		12,99(15)	12,9(11)		13,02(24)	12,8(2)	13,035(127)	12,98(8)
774,4	0,028(14)												0,022(11)			0,024(11)
790,1	0,031(8)												0,029(9)			0,030(8)
800,2	0,092(14)							0,09					0,088(30)			0,091(14)
815,6		1,38(6)	1,38(18)	1,45(8)	1,47(10)	1,51(5)	1,48(3)	1,63(12)		1,44(3)	1,455(14)		1,52(4)	1,481(15)		1,467(14)
830,3	≤0,0141				0,02								0,023(8)			0,023(8)
845,4		1,614(62)	1,60(22)		1,58(10)	1,687(22)	1,64(10)	1,61(61)		1,66(3)	1,737(20)		1,69(3)	1,659(17)		1,68(2)
850,7		0,663(30)	0,60(13)		0,67(8)	0,692(23)		0,68(13)		0,68(3)			0,68(2)	0,699(14)		0,692(14)
873,2		33,72(75)	34,8(17)	33,6(25)	34,47(70)	35,18(16)	34,66(21)	35,7(13)	34,95(31)	34,65(30)	34,81(28)		35,01(44)	34,5(4)	34,342(266)	34,87(16)
880,6	0,231(10)	0,14(6)	0,20(8)		0,28(8)			0,22(11)					0,26(4)			0,231(10)
892,8		1,41(4)	1,31(10)	1,38(12)	1,43(3)	1,497(26)	1,55(3)	1,51(10)		1,49(3)			1,48(5)	1,416(16)		1,473(16)
898,4	0,0056(14)															0,0056(14)
904,1		2,45(7)	2,42(17)	2,47(8)	2,49(5)	2,62(3)	2,65(8)	2,74(13)		2,54(6)	2,537(22)		2,58(5)	2,54(3)		2,551(22)
906,1	0,0338(16)															0,0338(16)
919,2	0,0352(16)												0,025(11)			0,0350(16)
924,5	0,166(8)	0,173(29)	0,19(10)		0,18(10)			0,13(6)					0,189(8)			0,177(8)
928,4	≤0,0141												0,013(6)			0,013(6)
981,3	0,023(6)												0,025(5)			0,024(5)
984,5	0,018(11)												0,029(6)			0,027(6)
996,3		29,39(71)	29,4(15)	29,7(21)	30,30(65)	30,09(15)	30,87(12)	31,0(19)	29,9(3)	30,14(30)	29,78(23)		30,29(51)	29,9(3)	29,206(269)	30,1(1)
1004,7		50,4(11)	50,6(25)	50,93(32)	51,40(103)	52,04(25)	52,05(31)	54,84(225)	51,9(5)	51,8(6)	51,55(40)		52,07(89)	51,6(4)	51,233(276)	51,17(25)
1012,8	0,0082(34)															0,008(3)
1023	0,020(8)												0,019(7)			0,019(7)
1033,4	0,0338(16)												0,029(8)			0,0336(16)
1047,4	0,141(7)				0,23(10)			0,17(6)					0,16(5)			0,142(7)
1049,4	0,0493(22)															0,0493(22)
1072,2	≤0,0113												0,010(4)			0,010(4)
1110	0,008(6)															0,008(6)
1118,5		0,403(58)	0,30(8)		0,37(10)			0,04		0,296(25)			0,31(3)			0,31(4)
1124,2	0,0197(28)															0,020(3)
1128,5		0,89(6)	0,79(9)		0,94(8)	0,90(4)		0,88(6)		0,885(25)	0,952(15)		0,89(5)	0,892(10)		0,91(1)
1136,1	0,0211(28)							0,042								0,021(3)

keV	1968Me18	1969Va09	1970RiZY	1980Ro22	1980Sh15	1984Iw03	1986Wa35	1989Ki10	1989 Schima	1990Me15	1990He05	1992E111	1992Ha02	1992Sm02	1992Sa04	Evaluated value
1140,7		0,634(30)	0,69(10)		0,73(8)	0,671(14)		0,75(6)		0,65(3)	0,671(8)		0,68(4)	0,682(11)		0,673(8)
1153,1	0,039(11)												0,024(10)			0,031(10)
1160,3	0,124(6)		0,10(3)		0,13(10)			0,12(4)					0,131(12)			0,125(6)
1170,7	0,012(6)												0,010(3)			0,010(3)
1188,6		0,27(1)	0,23(5)		0,29(8)			0,25(4)		0,25(3)			0,265(20)			0,266(20)
1216,8	≤0,010												0,0096(28)			0,010(3)
1232,1	0,026(17)												0,021(14)			0,023(14)
1241,6		0,43(3)	0,30(7)		0,40(5)	0,38(5)		0,45		0,366(17)			0,38(4)			0,380(17)
1246,1		2,54(7)	2,40(22)	2,35(5)	2,48(10)	2,49(4)	2,52(5)	2,51(12)		2,48(3)	2,449(23)		2,45(8)	2,48(2)	2,403(48)	2,470(23)
1274,4	100	100	100	100	100	100	100	100	100	100	100		100	100	100	100
1290,1	0,0324(15)		0,068(26)		0,086(20)			0,064					0,077(9)			0,071(9)
1292,0	0,0369(17)												0,035(3)			0,0364(15)
1295,5	0,0254(29)				0,026(3)			0,061					0,027(3)			0,026(3)
1316,4			0,074(29)		0,053(10)			0,029(19)								0,050(10)
1387,0	0,056(6)	<0,029											0,055(5)			0,055(5)
1397,4	0,0084(28)							0,012					0,0093(22)			0,0090(22)
1408,5	0,059(8)				0,082(10)								0,063(8)			0,066(8)
1415,0	0,0113(6)				0,004			0,02					0,017(6)			0,0114(6)
1418,6	0,0208(12)		0,027(16)		0,039			0,041(11)					0,037(5)			0,031(5)
1419,0	0,0056(3)															0,0056(3)
1425,9	0,0037(22)												0,0031(19)			0,0034(19)
1489,6	0,0084(14)												0,0081(12)			0,0082(12)
1494,0			1,88(9)	2,10(4)	1,91(8)	2,058(17)	1,99(2)	1,72(8)		1,99(4)	1,979(16)		2,04(8)	2,00(3)		2,00(2)
1510,0	0,0141(28)	<0,012											0,013(4)			0,014(3)
1522	0,0017(8)															0,0017(8)
1531,4	0,0172(12)		0,009(5)		0,018(5)								0,018(2)			0,0171(12)
1537,9			0,15(2)		0,15(1)			0,12(1)		0,155(6)			0,160(13)			0,151(6)
1554	≤0,004												0,0032(15)			0,0032(15)
1596,5			5,15(26)	5,19(8)	4,81(10)	5,247(30)	5,237(84)	4,54(18)	5,08(5)	5,13(8)	5,078(40)		5,12(17)	5,08(5)	5,083(22)	5,11(3)
1667,3	0,0056(8)												0,0053(12)			0,0055(8)
1674,9	0,0039(11)				0,006(1)			0,004					0,0041(16)			0,0049(11)
1716,9	0,0017(11)							0,0017(9)					0,0017(9)			0,0017(9)
1773	0,0008(6)							0,0010(6)					0,0010(6)			0,0010(6)
1838,0	0,0023(6)							0,0027(11)					0,0027(11)			0,0024(6)
1895	0,0017(6)							0,0020(9)					0,0020(9)			0,0018(6)

§ This energy corresponds to the two gamma-rays: $\gamma_{24,8}$ and $\gamma_{29,13}$. The former one was added in 1998Re22 with a relative emission probability of 0,020(7). Considering the experimental intensity of 0,032(5) as a sum of intensities $\gamma_{24,8}$ and $\gamma_{29,13}$, it leads to the $\gamma_{29,13}$ relative emission probability of 0,012(8)-see section 4.2.

The gamma ray emission probabilities have been computed from their relative evaluated emission probabilities given in Table 3 using the normalization factor $K = 0,3489(34)$. This value has been obtained from the intensity balance for gamma transitions to the ground states of ¹⁵⁴Gd and ¹⁵⁴Sm assuming that the ground states are not populated directly by beta or electron capture decay. Then, $P_{\gamma+ce}(\gamma_{i,0} \text{ Sm}) + \sum P_{\gamma+ce}(\gamma_{i,0} \text{ Gd}) = 100\%$ where $i=1, 3, 5, 6, 9, 11, 17, 18, 19, 20, 30, 33$.

There are several measurements of the absolute emission probabilities (P_γ) of some prominent gamma rays in the decay ¹⁵⁴Eu → ¹⁵⁴Gd.

The evaluated (calculated) value of $P_{\gamma_{1,0}}$ (123,07 keV) = 40,4(5)% agrees well with the value of 40,6(7)% measured in 1991ZaZZ.

The evaluated value of $P_{\gamma_{16,1}}$ (1274,43 keV) = 34,9(3)% agrees well with the value of 34,8(2)% measured in 1994Co02, and it differs somewhat from the value of 35,32(12)% obtained in 1992Ha02.

The values of $P_{\gamma_{2,1}}$ (247,93 keV) = 6,96(8) % and $P_{\gamma_{6,0}}$ (996,26 keV) = 10,36(18)% measured in 1997Ka47 agree with the evaluated (calculated) values of 6,89(7)% and 10,50(10)%, respectively.

5. Electron Emissions

The energies of the conversion electrons have been calculated from the gamma transition energies given in 2.2 and the electron binding energies.

The emission probabilities of conversion electrons have been deduced from the evaluated P_γ and ICC values.

The absolute total emission probabilities of Gd and Sm K Auger electrons have been computed by using their corresponding evaluated total $P(\text{ce}_K)$ for Gd and Sm and their adopted ω_K from section 3.

The absolute total emission probabilities of Gd and Sm L Auger electrons have been computed using their corresponding evaluated total $P(\text{ce}_L)$ and $P(\text{ce}_{L'})$ for Gd and Sm and their adopted ω_L and n_{KL} from section 3.

Average energies of β^- spectrum components have been calculated using the LOGFT program.

6. References

- 1949Ha04 - R. J. Hayden, J. H. Reynolds, M. G. Inghram - Phys. Rev. 75(1949)1500 [Half-life].
 1952Ka26 - D. G. Karraker, R. J. Hayden, M. G. Inghram - Phys. Rev. 87(1952)901 [Half-life].
 1957Ke08 - V. M. Kelman, V. A. Romanov, R. Y. Metskhvarishvili and V. A. Kolyunov - Nucl. Phys. 2(1957)395 [Gamma multipolarities].
 1958Ch36 - E. L. Chupp, J. W. M. DuMond, F. J. Gordon, R. C. Jopson, H. Mark - Phys. Rev. 112(1958)518 [Gamma-ray energies].
 1962Lu03 - D.S. Lu and R. S. Dingus - Phys. Lett. 3(1962)44 [Gamma multipolarities, experimental internal conversion electrons].
 1966Ha36 - P.C. Hansen, H. L. Nielsen and K. Wilsky - Nucl. Phys. 89(1966)571 [Beta transitions probabilities].
 1966Ja02 - J. S. W. Jansen, J. H. Hamilton and E. F. Zganjar - Proc. Intern. Conf. Internal Conversion Process, Nashville, Tenn. (1965), J.H. Hamilton, Ed., Academic Press, Inc., New York, (1966)257 [Experimental ICC, gamma multipolarities].
 1967Be65 - J. A. Bearden - Rev. Mod. Phys. 39(1967)78 [X-ray energies].
 1968Me18 - R. A. Meyer - Phys. Rev. 174(1968)1478 [Gamma-ray energies and relative emission probabilities].
 1968Ng01 - L. K. Ng, K. S. Mann and T. G. Walton - Nucl. Phys. A116(1968)433 [Beta transition probabilities].
 1969An01 - G. I. Anderson and G. T. Evans - Nucl. Phys. A123 (1969)609 [Gamma multipolarities].
 1969Va09 - L. Varnell, J. D. Bowman and J. Trischuk - Nucl. Phys. A127(1969)270 [Gamma-ray relative emission probabilities].
 1970Ri19 - L. L. Riedinger, N. R. Johnson, J. H. Hamilton - Phys. Rev. C2(1970)2358 [Gamma-ray energies].
 1970RiZY - L. L. Riedinger, E. Eichler, J. Fuglsang G.B. Hagemann and B. Herskind - Proc. Intern. Conf. Nucl. Reactions Induced by Heavy Ions, Heidelberg, Germany (1969) R. Bock, W. R. Hering, Eds., North-Holland Publishing Co., Amsterdam. (1970) 442 [Gamma-ray relative emission probabilities].

- 1972Em01 - J. F. Emery, S. A. Reynolds, E. I. Wyatt and G. I. Gleason - Nucl. Sci. Eng. 48(1972)319 [Half-life].
- 1972Na21 - T. S. Nagpal and R. E. Caucher – Can. J. Phys. 50(1972)2688 [Gamma multipolarities].
- 1974HeYW - R. L. Heath, Gamma -Ray Spectrum Catalogue. In: ANCR 1000 2(1974) [Gamma -ray energies].
- 1977La19 - F. P. Larkins - Atomic Data and Nuclear Data Tables 20(1977)313 [Auger electron energies].
- 1977Ya04 - H. Yamada, H. Kawakami, M. Koike and K. Komura – J. Phys. Soc. Jap. 42(1977)1448 [Gamma multipolarities].
- 1978GrZR - G.-F. Grisham, (1978): No title. (Priv. Comm; June) [Half-life].
- 1978Ro22 - F. Rosel, H. M. Friess, K. Alder and H. C. Pauli - At. Data Nucl. Data Tables 21(1978)92. [Theoretical ICC].
- 1978ScZO - K. Schreckenbach, (1978): No title. (Priv. Comm; July) [Half-life].
- 1980Ro22 - W. M. Roney, Jr., and W. A. Seale – Nucl. Instr. Meth. 171(1980)389 [Gamma -ray relative emission probabilities].
- 1980Sh15 - A. K. Sharma, R. Kaur, H. R. Verma and P. N. Trehan – J. Phys. Soc. Jap. 48(1980)1407 [Gamma-ray relative emission probabilities].
- 1982HoZY - D. D. Hopes, J. M. R. Hutchinson, F. J. Schima and M. P. Unterweger - NBS-SP 626(1982)85 [Half-life]
- 1983Wa26 - K. F. Walz, K. Debertin and H. Schrader - Intern. J. Appl. Radiat. Isotop. 34(1983)1191 [Half-life].
- 1984Iw03 - J. Iwata, M. Yasuhara, K. Maeda and Y. Yoshizawa - Nucl. Instr. Meth. 219(1984)123 [Gamma-ray relative emission probabilities].
- 1986Wa35 - Wang Xinlin, Li Xiaodi and D. Hongshan - Chin. J. Nucl. Phys. 8(1986)371 [Gamma-ray relative emission probabilities].
- 1986Wo05 - M. J. Woods and S. F. M. Lucas - Intern. J. Appl. Rad. Isot. 37(1986)1157 [Half-life].
- 1987Lagoutine - F. Lagoutine, N. Coursol and J. Legrand. Table de Radionucléides, ISBN -2-7272-0078-1 (LMRI, 1982-1987) [Energy of Auger electrons].
- 1988RaZM - M. U. Rajput and T. D. Mac Mahon - ZFK-732. (1988)313 [Half-life].
- 1989Ki10 - L. L. Kiang, G. C. Kiang, P. K. Teng, G. C. Jon, T. H. Yuan and Y. M. Hsu - Z. Phys. A333(1989)19 [Gamma-ray relative emission probabilities].
- 1989Schima - F. J. Schima (1989), IAEA -CRP informal report GS/59; Cited in IAEA -TECDOC-619 (1991) [Gamma-ray relative emission probabilities].
- 1990He05 - R. G. Helmer - Appl. Rad. Isot. 41(1990)75 [Gamma -ray energies and relative emission probabilities].
- 1990Me15 - R. A. Meyer - Fizika (Zagreb) 22(1990)153 [Gamma -ray energies and relative emission probabilities].
- 1991ZaZZ - A. V. Zanevsky, G. A. Isaakyan, N. V. Kurenkov, A. B. Malinin, T. E. Sazonova and S. V. Sepman- Program and Thesis, Proc. 41st Ann. Conf. Nuclear Spectrosc. Struct. At. Nuclei, Minsk. (1991)477 [123 keV gamma-ray absolute emission probabilities].
- 1992El11 - S. U. El-Kameesy, M. S. Abdel-Wahab, L. Al-Houty and H. Abou-Leila - Acta Phys. Acad. Sci. Hung. 71(1992)161 [Gamma-ray relative emission probabilities].
- 1992Ha02 - M. A. Hammed, I. M. Lowles and T. D. Mac Mahon - Nucl. Instr. Meth. Phys. Res. A312(1992)308 [Gamma-ray relative emission probabilities and 1274 keV gamma-ray absolute emission probability].
- 1992Sa04 - T. E. Sazonova, G. A. Isaakyan, N. I. Karmalitsyn, S. V. Sepman and A. V. Zanevsky - Nucl. Instrum. Methods Phys. Res. A312(1992)372 [Gamma-ray relative emission probabilities].
- 1992ScZZ - U. Schötzig, H. Schrader and K. Debertin - Proc. Intern. Conf. Nuclear Data for Science and Technology, Julich, Germany. (1992)562 [Half-life].
- 1992Sm02 - D. Smith, D. H. Woods, S. A. Woods, J. L. Makepeace, R. A. Mercer and C. W. A. Downey - Nucl. Instrum. Methods Phys. Res. A312(1992)353 [Gamma-ray energies and emission probabilities].
- 1992Un01 - M. P. Unterweger, D. D. Hoppes and F. J. Schima - Nucl. Instrum. Methods Phys. Res. A312(1992)349 [Half-life]
- 1993Th04 - J. L. Thompson and A. R. Cartwright - Appl. Radiat. Isot. 44(1993)707 [Half-life].
- 1994Co02 - B. M. Coursey, J. M. Calhoun, J. Cessna, D. B. Golas, F. J. Schima and M. P. Unterweger - Nucl. Instrum. Methods Phys. Res. A339(1994)26 [1274 keV gamma-ray absolute emission probability].
- 1995Au04 - G. Audi and A. H. Wapstra - Nucl. Phys. A595(1995)409 [Q-value].

- 1995Ma03 - N. Mach and B. Foggerberg - Phys. Rev. C51 (1995)509 [Experimental internal conversion electrons].
- 1996AL31 - I. Alfter, E. Bodenstedt, W. Knichel, J. Schuth - Z. Phys. A355(1996)277 [Experimental gamma multiplicities].
- 1996Sc06 - E. Schönfeld and H. Janßen - Nucl. Instrum. Methods Phys. Res. A369(1996)527 [Atomic data].
- 1997Ka47 - S. I. Kafala, T. D. Mac Mahon and S. V. Borzakov - J. Radioanal. Nucl. Chem. 215(1997)193 [248 keV and 996 keV gamma-ray absolute emission probability].
- 1998Si12 - H. Siegert, H. Schrader and U. Schotzig - Appl. Radiat. Isot. 49(1998)1397 [Half-life].
- 1998Sc28 - E. Schönfeld - Appl. Radiat. Isot. 49(1998)1353 [Fractional electron capture probabilities P_K , P_L , P_M].
- 1998Re22 - C. W. Reich, R. G. Helmer - Nuclear Data Sheets 85(1998)171 [Level scheme, multiplicities, total ICC].
- 1999Schönfeld - E. Schönfeld and G. Rodloff - PTB-6.11-1999-1999-1, Braunschweig, Februar 1999 [KX-ray energies and relative emission probabilities].
- 2000Ch01 - V. P. Chechev and A. G. Egorov - Appl. Rad. Isot. 52(2000)601[Evaluation technique].
- 2000He14 - R. G. Helmer and C. van der Leun - Nucl. Instrum. Methods Phys. Res. A450(2000)35 [Gamma-ray energies].
- 2002Un02 - M. P. Unterwiesing - Appl. Rad. Isot. 56(2002)125 [Half-life].

¹⁵⁵Eu – Comments on evaluation of decay data

by V. P. Chechev and V. O. Sergeev

1. DECAY SCHEME

The ¹⁵⁵Eu decay scheme is complete. The most intense allowed β^- -transitions occur to the excited levels with energy of 105.31 keV (46.1%) and 86.55 keV (25.5%).

The 1st forbidden β^- -transitions populate the 60.01 keV (9,2%) and 146.07 keV (1.9%) levels.

The ground state in ¹⁵⁵Gd is populated by the intense allowed β^- -transition (16.6%).

The 2nd forbidden β^- -transition to the excited level of 107.58 keV was not observed. From the log ft systematics its log ft should be more than 11.1 and the upper limit on this β^- branch intensity is expected less than 0.01%.

2. NUCLEAR DATA

Q value is from 1995Au04 .

The evaluated value of the ¹⁵⁵Eu half-life has been taken from 2000Ch01 (Chechev and Egorov). It is based on the measurement results given in Table 1.

Table 1. Set of experimental data for the evaluation of ¹⁵⁵Eu half-life (in days)

Reference	Author	Data set "1"	Data set "2"	Data set "3"
		$\chi^2 = 334.9$ $(\chi^2)_6^{0.05} = 14.1$	$\chi^2 = 6.14$ $(\chi^2)_5^{0.05} = 12.6$	$\chi^2 = 5.68$ $(\chi^2)_5^{0.05} = 12.6$
1998Si12	Siegert <i>et al.</i>	1739(8)	1739(8)	1739(8)
1993Th04	Thompson <i>et al.</i>	1735(22)	1735(22)	1735(22)
1992Un01	Unterweger <i>et al.</i>	1739.0(5)	1739.0(5)	1739(7) ^b
1983Wa26	Walz <i>et al.</i>	1737(23)	1737(23)	1737(23)
1974Da24	Daniels <i>et al.</i>	1708(18)	1708(18)	1708(18)
1972Em01	Emery <i>et al.</i>	1812(4)	Omitted ^a	-
1972Su09	Subba Rao	1653(51)	1653(51)	1653(51)
1970Mo23	Mowatt <i>et al.</i>	1698(74)	1698(74)	1698(74)

^a The value from 1972Em01 has been omitted on the basis of statistical considerations.

^b The rule of "50% weight"(LRSW) leads to a significant increase of the 1992Un01 uncertainty.

In 2002Un02 the new NIST measurement result was published for the ¹⁵⁵Eu half-life: $T_{1/2} = 1739.06(45)$ d. It does not differ practically from 1992Un01 and its use instead of 1992Un01 does not change this evaluation.

The weighted mean of the experimental values from the final data “set 3” is 1736(5) days where the uncertainty is internal. The adopted value of the ¹⁵⁵Eu half-life is 1736(5) days, or 4.753(14) years.

2.1. β^- -Transitions

The energies of the β^- transitions have been computed from the Q value and the level energies adopted from 1986Sc25, where the reaction ¹⁵⁴Gd(n, γ)¹⁵⁵Gd was studied. For the level energies see also the evaluation in Nuclear Data Sheets (1994Re10).

The probabilities of the β^- transitions have been obtained from the $P_{\gamma+ce}$ balance for each level based on the P_{γ} normalization factor of 0.307(3) (see sect.4.2.3). The calculated $P(\beta_{0,0})$ agrees with the unweighted mean of 18(4)% of the five measurement results of 1949Ma58, 1954Le08, 1956Du31, 1959Am16, 1960Su04.

2.2. Gamma Transitions and Internal Conversion Coefficients

The evaluated energies of gamma transitions are energies of gamma rays (E_{γ}) with adding the recoil energy of $E_{\gamma}^2 / 2Mc^2$ where M – mass of the ¹⁵⁵Gd nucleus. The latter changes the energy only for $\gamma_{6,0}$.

The gamma transition probabilities have been calculated from the gamma emission probabilities and the internal conversion coefficients (ICC).

For gamma transitions with energies more than 25 keV the ICC have been evaluated using theoretical values from 1978Ro22 for the adopted multiplicities. For these transitions the following uncertainties for theoretical values have been adopted 1% for α_K and 3% for α_L , α_M , α_{NO} . The ICC interpolated from other tables (1968Ha53, 1978Band) do not differ from the evaluated values within limits of adopted uncertainties.

For low-energy gamma transitions $\gamma_{5,4}$, $\gamma_{3,2}$, $\gamma_{4,2}$ the ICC have been evaluated using theoretical values from 1993Ba60. The ICC values in 1968Ha53 and 1978Ro22 for these energies differ considerably or are absent.

The adopted E2 admixtures for (M1+E2)-transitions $\gamma_{5,4}$, $\gamma_{3,2}$, $\gamma_{5,2}$, $\gamma_{1,0}$ and $\gamma_{2,0}$ have been evaluated using measurement results from 1959De29, 1961Su13, 1962Ha24, 1966As02, 1967Fo11, 1967Ko12, 1975Ch04, 1975Kr04, 1986Sc25 and 1990GoZS. In these works the intensity ratios $L_1:L_2:L_3$ were measured for conversion electrons in decays of ¹⁵⁵Eu and ¹⁵⁵Tb and also in the ¹⁵⁴Gd(n, γ) reaction. Also $\gamma\gamma(\theta)$ -correlations were studied in ¹⁵⁵Tb decay and in Coulomb excitation of the ¹⁵⁵Gd levels - ¹⁵⁵Gd (p, p γ) (see Table 2).

Table 2. Measured and evaluated E2 admixtures for the (M1+E2) multiplicities of gamma transitions in the decay of ¹⁵⁵Eu

E_{γ} , keV	Measurement result, % E2	NSR code	Method	Evaluated (adopted) value, % E2
10.418	0.11(5) 0.4(3)	1975Ch04 1967Fo11	$L_1; L_2; L_3, ^{155}\text{Tb}$ $L_1; L_2; L_3, ^{155}\text{Eu}$	0.11(5)
18.763	7.4(6) 6.3(8) 7.1(4) 5.6(12) 6.3(14)	1990GoZS 1967Fo11 1975Ch04 1962Ha24 1975Kr04	$L_1; L_2; L_3, ^{155}\text{Eu}$ $L_1; L_2; L_3, ^{155}\text{Eu}$ $L_1; L_2; L_3, ^{155}\text{Tb}$ $L_1; L_2; L_3, ^{155}\text{Tb}$ $\gamma\gamma, ^{155}\text{Eu}$	7.1(4) WM
31.444	17(5)	1986Sc25	$L_1; L_2; L_3, ^{154}\text{Gd}(n,\gamma)$	17(5)

60.009	4.0(4) 3.3(10) 4.4(4) 3.7(10) 3.5(9) 3.8(10) 4.9(24)	1967Fo11 1967Ko12 1986Sc25 1962Ha24 1975Kr04 1961Su13 1966As62	L ₁ ; L ₂ ; L ₃ , ¹⁵⁵ Eu L ₁ ; L ₂ ; L ₃ , ¹⁵⁵ Tb L ₁ ; L ₂ ; L ₃ , ¹⁵⁴ Gd(n,γ) L ₁ ; L ₂ ; L ₃ , ¹⁵⁵ Tb γγ, ¹⁵⁵ Eu γγ, ¹⁵⁵ Eu ¹⁵⁵ Gd (p, p' γ)	4.1(4) WM
86.059	2.5(6) 3.5(10) 4.9(15) 3.5(16)	1986Sc25 1975Kr04 1966As02 1959De29	L ₁ ; L ₂ ; L ₃ , ¹⁵⁴ Gd(n,γ) γγ, ¹⁵⁵ Eu ¹⁵⁵ Gd (p, p' γ) ¹⁵⁵ Gd (p, p' γ)	3.0(6) WM

3. ATOMIC DATA

3.1. Fluorescence yields

The fluorescence yields are taken from 1996Sc06 (Schönfeld and Janßen).

3.2. X Radiations

The X-ray energies are based on the wavelengths in the compilation of 1967Be65 (Bearden). The relative KX-ray emission probabilities are taken from 1996Sc06, 1999Schönfeld and 1974Sa28.

3.3. Auger Electrons

The energies of Auger electrons are from 1977La19 (Larkins) and 1987Table. The ratios P(KLX)/P(KLL) and P(KLY)/P(KLL) are taken from 1996Sc06.

4. PHOTON EMISSIONS

4.1 X-Ray Emissions

The total absolute emission probability of KX -rays (P_{XK}) has been computed using the adopted value of ω_{K} and the evaluated total absolute emission probability of K conversion electrons $P_{\text{ce}_{\text{K}}} = 25.17(46)$ per 100 disintegrations. The absolute emission probabilities of the KX -ray components have been computed from P_{XK} using the relative probabilities from Sect.3.2.

The measured values of the total absolute emission probability of KX -rays given below can be compared to the calculated (adopted) value of $P_{\text{XK}}^{\text{eval}} = 23.6(5)$ per 100 disintegrations:

1967Fo11	1967Bl11	1968Om01	1969Me09	1971Ge11	1994Eg01	WM
22.9(10)	25.2(25)	21.3(23)	21.1(6)	22.5(12)	23.50(19)	23.3(2) ^a

^a Weighted mean of all 6 values. The value of 1969Me09 gives the 80% contribution to χ^2 . With omitting this value the weighted mean of 5 values is 23.5(2).

The total absolute emission probability of LX -rays has been computed using the adopted values of ω_{L} and n_{KL} and the evaluated values of $P(\text{ce}_{\text{K}}) = 25.17(46)$ and $P(\text{ce}_{\text{L}}) = 21.2(24)\%$.

4.2. Gamma-Ray Emissions

4.2.1. Gamma-Ray Energies

The measured and evaluated values of gamma ray energies are given in Table 3.

The evaluated values of E_γ have been obtained as weighted means omitting outliers contradicting to the energies of excited levels measured in 1986Sc25. The values of 1969Me09 have been omitted as the author in 1990Me15 replaces them.

4.2.2. Gamma-Ray Relative Emission Probability

The measured and evaluated values of relative gamma ray emission probabilities ($P'(\gamma)$) are shown in Table 4.

The evaluated values of $P'(\gamma)$ have been obtained as weighted means apart from $P'(\gamma_{5,4})$ and $P'(\gamma_{4,2})$. The $P'(\gamma_{5,4})$ has been evaluated from the intensity balance for the 107.58 keV - level. The $P'(\gamma_{4,2})$ has been calculated from data on conversion electrons (1967Fo11) and the adopted ICC using the measured in 1967Fo11 ratio $P(\text{ce}_{4,2} \text{L3})/P(\text{ce}_{3,0} \text{K}) = 0,115(6)$ and the adopted values of $\alpha_{L3}(\gamma_{4,2})$ and $\alpha_K(\gamma_{3,0})$.

The values of 1969Me09 have been omitted as the author in 1990Me15 replaces them. Other values have been omitted due to absence of uncertainties or as statistical outliers.

Our evaluated value $P'(\gamma_{3,0}) = 68.8(14)$ for the intense gamma ray with energy of 105.31 keV is supported by the results of measurements of the intensity ratio $P(\text{ce}_{3,0} \text{K})/P(\text{ce}_{2,0} \text{K}) = 0.408(8)$ in 1967Fo11 (see Table 5) which leads to the value $P'(\gamma_{3,0}) = 68.7(17)$ if the adopted α_K in sect.2.2 is used.

4.2.3. Gamma-Ray Absolute Emission Probabilities

Two absolute measurements of the emission probability are available for the 86,55 keV gamma ray: 31.1(4)% in 1994Co02 and 30.5(3)% in 1994Eg01. The weighted mean of these values has been adopted as the evaluated $P(\gamma_{2,0}) = 30.7(3)\%$. Here the uncertainty is the external one of WM.

The absolute emission probabilities of other gamma rays have been computed from the evaluated emission probabilities (P') given in Table 4 and the evaluated absolute emission probability of $\gamma_{2,0}$ (86.55 keV).

It should be noted that the absolute emission probability of $\gamma_{3,0}$ (105.31 keV) was measured in 1992Sa04: $P(\gamma_{3,0}) = 20.39(13)\%$. This value is considerably less than the evaluated one and measured in 1994Eg01 and 1996Ch27. If it is adopted without changing of the evaluated $P(\gamma_{2,0}) = 30.7(3)\%$ the relative emission probability of $\gamma_{3,0}$ will be 66.4(9), essentially less than the average of the eight measurement results (Table 4 and comment in sect.4.2.2.). On other hand, if the value of 1992Sa04 is adopted together with the evaluated $P'(\gamma_{3,0}) = 68.8(14)$, the $P(\gamma_{2,0})$ will be obtained as 29.6(6)%, less than both results of direct measurement of the absolute emission probability of this gamma ray (1994Co02 and 1994Eg01).

Therefore we consider the value of 1999Sa04 as too small and do not take it into account.

Table 3. Measured and evaluated values of gamma ray energies in the decay of ¹⁵⁵Eu

	1959Ha07	1967Fo11	1969Me09	1970Re08	1970Ra37	1975Ch04 ^a	1975Kr04	1986Sc25 ^b	1990Me15	1990GoZS	Evaluated (adopted) value
$\gamma_{5.4}$		10.40(2)*				10.40(2)*		10.4183(13)			10.4183(13)
$\gamma_{3.2}$		18.776(35)*	18.776(35)*			18.749(19)*	18.73(3)*	18.760(4)	18.784(35)*	18.764(2)	18.763(2) ^c
$\gamma_{4.2}$		21.02(2)				21.02(2)		21.030(10)		21.036(4)	21.035(4)
$\gamma_{2.1}$			26.513(21)*				26.49(5)	26.530(23)	26.532(21)		26.531(21)
$\gamma_{5.2}$			31.40(10)*	31.55(12)				31.444(7)	31.40(10)		31.444(7)
$\gamma_{3.1}$	45.29(1)	45.3(2)*	45.299(13)*	45.299(2)	45.2972(13)		45.27(5)*	45.3000(10)	45.295(13)		45.2990(10)
$\gamma_{5.1}$			57.983(30)*	57.970(26)	57.9805(20)		57.99(4)	57.989(1)	57.986(30)		57.989(1)
$\gamma_{1.0}$	60.00(2)		60.019(15)*	60.006(4)	60.0100(18)		60.01(4)	60.008(2)	60.022(15)	60.0086(10)	60.0086(10) ^c
$\gamma_{6.1}$		86.01(20)	86.0(5)	86.062(23)	86.062(5)		86.03(7)	86.0590(10)			86.05910(10)
$\gamma_{2.0}$	86.56(1)	86.82(20)	86.539(15)*	86.541(3)	86.5452(33)		86.53(3)	86.5470(10)	86.554(15)		86.5479(10)
$\gamma_{3.0}$	105.32(3)	105.28(20)	105.315(15)*	105.302(4)	105.308(3)		105.30(3)	105.3090(10)	105.338(15)		105.3083(10)
$\gamma_{3.0}$			146.05(2)*	146.061(5)			146.04(10)	146.0710(10)	146.090(90)		146.0710(10)

^a Decay of ¹⁵⁵Tb^b Reaction ¹⁵⁴Gd(n, γ)¹⁵⁵Gd^c The data of 1976Me10 (decay of ¹⁵⁵Tb) have been taken into consideration additionally: E($\gamma_{3.2}$)=18.769(15) keV and E($\gamma_{1.0}$)=60.012(3) keV.

* Omitted from averaging. Values of 1969Me09 are superseded by those of 1990Me15.

Table 4. Measured and evaluated values of relative gamma ray emission probabilities in the decay of ¹⁵⁵Eu.

	E _γ , keV	1959Ha07	1967Be11	1968Al01	1969Me09	1970Re08	1971Ge11	1975Kr04	1990Me15	1994Eg01	1996Ch27	Evaluated value
γ _{5,4}	10.418											0.0115(13) ^a
γ _{3,2}	18.763	≈0,1*			0.16(4)*		0.17(3)	0.13(3)	0.16(4)			0.16(2) ^{b,c}
γ _{4,2}	21.035											1.5(3)·10 ⁻³ ^d
γ _{2,1}	26.531	≈4*		≈1*	1.03(6)*		1.00(10)	1.10(13)	1.03(6)			1.03(6) ^c
γ _{5,2}	31.444				0.023(5)*	0.03(2)			0.023(5)			0.023(5) ^c
γ _{3,1}	45.299	2.3*		2.8(7)*	4.18(17)*	3.6(7)	4.1(3)	3.95(40)	4.21(20)	4.36(12)	4.3(10)	4.27(12) ^c
γ _{5,1}	51.989			0.20(3)	0.217(18)*	0.22(5)		0.23(3)	0.221(18)	0.213(30)		0.217(18) ^c
γ _{1,0}	60.009	4,0*	5.1(20)*	3.8(2)	3.60(10)*	4.3(3)	3.9(9)	3.8(4)	3.60(10)	3.99(12)	3.9(9)	3.96(12) ^c
γ _{6,1}	86.059			0.50(5)		0.49(5)		0.54(11)				0.50(5) ^c
γ _{2,0}	86.548	100	100	100	100	100	100	100	100	100	100	100
γ _{3,0}	105.308	64*	65.7(65)	67.9(35)	66.8(27)*	68.3(27)	68(4)	69.9(35)	66.8(27)	68.5(14)	69.5(16)	68.8(14) ^{c,e}
γ _{6,0}	146.071		0.16(5)		0.167(10)*	0.19(2)		0.14(2)	0.167(10)			0.166(10) ^c

^a Evaluated from the intensity balance for the 107.58 keV level

^b In addition the value of 0.16(2) from 1974HeYW has been taken into account

^c Weighted mean

^d Evaluated from the conversion electron intensity and ICC

^e In addition the value of 69.1(9) from 1982Co05 has been taken into account

* Omitted from averaging. Values of 1969Me09 are superseded by those of 1990Me15.

5. ELECTRON EMISSIONS

The energies of the conversion electrons have been calculated from the gamma β^- -transition energies given in 2.2 and the electron binding energies.

The emission probabilities of the conversion electrons have been calculated using the evaluated P_{γ} and ICC. In Table 5 the relative intensities of conversion electrons $P'_{ce}(exp.)$ measured in 1967Fo11 are compared to the relative intensity values $P'_{ce}(calc.)$ calculated from the evaluated absolute emission probabilities (in units $P'_{ce}(K) = 1000$).

Table 5. Comparison of experimental and calculated values of relative intensity of conversion electrons in the ^{155}Eu decay.

	Energy, keV	$P'_{ce}(exp)$	$P'_{ce}(calc.)$
ec _{5,4} L	2.043-3.175	305(27)	206(30)
ec _{1,0} K	9.770(3)	1870(100)	2000(130)
ec _{3,2} L	10.387-11.520	2730(110)	3080(400)
ec _{4,2} L	12.659-13.792	212(8)	218(30)
ec _{6,1} K	35.820(3)	66(5)	91(12)
ec _{2,0} K	36.309(3)	2450(50)	2440(50)
ec _{3,1} L	36.923-38.053	90(5)	100(5)
ec _{1,0} L	51.633-52.766	420(10)	418(16)
ec _{3,0} K	55.069(3)	1000	1000
ec _{2,0} L	78.172-79.305	380(9)	382(13)
ec _{3,0} L	96.933-98.066	152(6)	152(8)

As seen from Table 5 the experimental and calculated values agree well with the exception of ec_{5,4} L and ec_{6,1} K. The disagreement for ec_{5,4} L can be connected with experimental difficulties of measurement of the 2-3 keV conversion electrons on the background of intense L Auger electrons, and for ec_{6,1} K – of measurement on the background of intense conversion line of ec_{2,0} K.

The total absolute emission probability of K Auger electrons has been computed using the total $P(ce_K) = 25.17(46)\%$ and the adopted ω_K in sect.3.

The total absolute emission probability of L Auger electrons has been computed using the evaluated total $P(ce_K)$ and $P(ce_L) = 21.2(24)\%$ and the adopted ω_L and n_{KL} in sect.3.

The values of β^- average energies have been calculated using the LOGFT program.

References

- 1949MA58 - J. A. Marinsky, L. E. Glendenin, F. Metzger, In: MIT -LNS Progr Rept. p.67 (1949) NP -1272. (Beta emission energies and probabilities)
- 1954LE08 - M. R. Lee, R. Katz, Phys. Rev. 93(1954)155. (Beta emission energies and probabilities)
- 1956DU31 - V. S. Dubey, C. E. Mandeville, M. A. Rothman, Phys. Rev. 103(1956)1430 (Beta emission energies and probabilities)
- 1959AM16 - M. C. Joshi, B. V. Thosar, Proc. Indian Acad. Sci. 50A(1959)342. (Beta emission energies and probabilities)
- 1959DE29 - J. de Boer, M. Martin, P. Marmier, Helv. Phys. Acta. 32(1959)658.(Multipolarity mixing ratios)
- 1959HA07 - E. N. Hatch, F. Boehm, Z. Phys. 155(1959)609.(Gamma ray energies and relative emission probabilities)
- 1960SU04 - B. N. Subba Rao, Nuovo Cimento 16(1960)283. (Beta emission energies and probabilities)
- 1961SU13 - B. N. Subba Rao, Nucl. Phys. 28(1961)503.(Multipolarity mixing ratios)
- 1962HA24 - B. Harmatz, T. H. Handley, J. W. Mihelich, Phys. Rev. 128(1962)1186. (Multipolarity mixing ratios)
- 1966AS02 - D. Ashery, A. E. Blaugrund, R. Kalish, Nucl. Phys. 76(1966)336.(Multipolarity mixing ratios)
- 1967BE11 - R. E. Berg, J. L. Snelgrove and E. Kashy, Phys. Rev. 153(1967)1165. (Gamma ray relative emission probabilities)
- 1967BL11 - P. H. Blichert -Toft, E. G. Funk, J. W. Mihelich, Nucl. Phys. A96(1967)190. (KX ray emission probabilities)
- 1967FO11 - C. Foin, J. Oms, J. -L. Barat, J. Phys. (Paris) 28(1967)861.(Gamma ray energies, KX β^- -ray emission probabilities, conversion electron emission probabilities, multipolarity mixing ratios)

- 1967KO12 - J. Kormicki, H. Ni ewodniczanski, Z. Stachura, K. Zuber, A. Budziak, Nucl. Phys. A102(1967)253. (Multipolarity mixing ratios)
- 1968AL01 - P. Alexander, Nucl. Phys. A108(1968)145. (Gamma ray relative emission probabilities)
- 1968HA53 - R. S. Hager, E. C. Seltzer, Nucl. Data Tables A4(1968)1. (Theoretical ICC)
- 1968OM01 - J. Oms, C. Foin, A. Baudry, Compt. Rend. 266B(1968)1292. (KX ray emission probabilities)
- 1969ME09 - R. A. Meyer, J. W. T. Meadows, Nucl. Phys. A132(1969)177. (Gamma and KX ray energies and emission probabilities)
- 1970MO23 - R. S. Mowatt, Can. J. Phys. 48(1970)1933. (Half-life)
- 1970RA37 - D. E. Raeside, Nuclear Instrum. Methods 87(1970)7. (Gamma ray energies)
- 1970RE08 - J. D. Reiersen, G. C. Nelson, E. N. Hatch, Nucl. Phys. A153(1970)109. (Gamma ray energies and relative emission probabilities)
- 1971GE11 - R. J. Gehrke, R. A. Lokken, Nuclear Instrum. Methods 97(1971)219. (Gamma ray relative emission probabilities)
- 1972EM01 - J. F. Emery, S. A. Reynolds, E. I. Wyatt, G. I. Gleason, Nucl. Sci. Eng. 48(1972)319 (Half-life)
- 1972SU09 - B. N. Subba Rao, Current Sci. (India) 41(1972)692. (Half-life)
- 1974DA24 - W. R. Daniels, D. W. Barr, G. F. Grisham, F. O. Lawrence, J. Inorg. Nucl. Chem. 36(1974) 3874. (Half-life)
- 1974HEYW - R. L. Heath, Report ANCR 1000-2 (1974). (Gamma ray relative emission probabilities)
- 1974SA28 - S. I. Salem, S. L. Panossian, R. A. Krause, At. Data Nucl. Data Tables 14(1974)91. (KX ray relative emission probabilities)
- 1975CH04 - P. Christmas, P. Cross, J. Phys. (London) G1(1975)113. (Gamma ray energies, multipolarity mixing ratios)
- 1975KR04 - H. J. Krell, S. Hofmann, Z. Phys. A272(1975)257. (Gamma ray energies and relative emission probabilities, multipolarity mixing ratios)
- 1976ME10 - R. A. Meyer, R. Gunnink, C. M. Lederer, E. Browne, Phys. Rev. C13(1976)2466. (Gamma ray energies)
- 1977LA19 - F. P. Larkins, Atomic Data and Nuclear Data Tables 20(1977)313 (Auger electron energies)
- 1978BAND - I. M. Band, M. B. Trzhaskovskaya, Special report of Leningrad nuclear physics institute, 1978. (Theoretical ICC)
- 1978RO22 - F. Rösel, H. M. Friess, K. Alder, H. C. Pauli, At. Data Nucl. Data Tables 21(1978)92. (Theoretical ICC)
- 1982CO05 - B. M. Coursey, D. D. Hoppes, F. J. Schima, Nuclear Instrum. Methods 193(1982)1. (Gamma ray relative emission probabilities)
- 1983WA26 - K. F. Walz, K. Debertin, H. Schrader, Int. J. Appl. Radiat. Isotop. 34(1983)1191. (Half-life)
- 1986SC25 - H. H. Schmidt, W. Stoffl, T. Egidy, P. Hungerford, H. J. Scheerer, K. Schreckenbach, H. G. Börner, D. D. Warner, R. E. Chrien, R. C. Greenwood, C. W. Reich, J. Phys. (London) G12(1986)411. (Gamma ray and level energies, multipolarity mixing ratios)
- 1987Table - F. Lagoutine, N. Coursol, J. Legrand, Table de Radionucléides - ISBN-2-7272-0078-1 (LMRI, 1982-1987) (Energy of Auger electrons)
- 1990GOZS - V. M. Gorozhankin, A. Kovalik, L. E. Ir, M. A. Makhmud, Ya. Novak, A. F. Novgorodov. In: Program and Thesis, Proc. 40th Ann. Conf. Nucl. Spectrosc. Struct. At Nuclei, Leningrad, (1990) p 97. (Gamma ray energies, multipolarity mixing ratios)
- 1990ME15 - R. A. Meyer, Fizika (Zagreb) 22(1990)153. (Gamma ray energies and relative emission probabilities)
- 1992SA04 - T. E. Sazonova, G. A. Isaakyan, N. I. Karmalitsyn, S. V. Sepman, A. V. Zanevsky, Nucl. Instrum. Methods Phys. Res. A312(1992)372. (105,31 keV gamma ray absolute emission probability)
- 1992UN01 - M. P. Unterweger, D. D. Hoppes, F. J. Schima, Nucl. Instrum. Methods Phys. Res. A312(1992)349. (Half-life)
- 1993BA60 - I. M. Band, M. B. Trzhaskovskaya, At. Data Nucl. Data Tables 55(1993)43(Theoretical ICC)
- 1993TH04 - J. L. Thompson, A. R. Cartwright, Appl. Radiat. Isot. 44(1993)707. (Half-life)
- 1994CO02 - B. M. Coursey, J. M. Calhoun, J. Cessna, D. B. Golas, F. J. Schima, M. P. Unterweger, Nucl. Instrum. Methods Phys. Res. A339(1994)26. (86,55 keV gamma ray absolute emission probability)
- 1994EG01 - A. G. Egorov, V. P. Chechev, Nucl. Instrum. Methods Phys. Res. A339(1994)248. (Gamma and KX ray absolute emission probabilities)
- 1994RE10 - C. W. Reich, Nucl. Data Sheets 71(1994)709. (Decay scheme and level energies)
- 1995AU04 - G. Audi, A. H. Wapstra, Nucl. Phys. A595(1995)409. (Q value)
- 1996Ch27 - V. P. Chechev, A. G. Egorov, G. E. Shchukin, Appl. Radiat. Isot. 47(1996)329 (Gamma ray relative emission probabilities)
- 1996Sc06 - E. Schönfeld, H. Janßen, Nucl. Instrum. Methods Phys. Res. A369(1996)527. (Atomic data)
- 1998Si12 - H. Siegert, H. Schrader, U. Schötzg, Appl. Radiat. Isot. 49(1998)1397 (Half-life)
- 1999Schönfeld E. Schönfeld, G. Rodloff, PTB -6.11-1999-1, Braunschweig, February 1999 (KX ray energies and relative emission probabilities)
- 2000Ch01 - V. P. Chechev, A. G. Egorov, Appl. Radiat. Isot. 52(2000)601. (Evaluation of half-life)
- 2002Un02 - M. P. Unterweger, Appl. Radiat. Isot. 56(2002)125. (Half-life)

¹⁵⁹Gd - Comments on evaluation of decay data by R. G. Helmer

This evaluation was completed in 2004. The literature available by March 2005 was included.

1 Decay Scheme

¹⁵⁹Gd decays by β^- emission to levels in ¹⁵⁹Tb.

2 Nuclear Data

Q value is 970.5(7) from Audi et al. 2003 mass evaluation (2003Au03).

For the adopted decay scheme, the total radiation energy per decay is calculated to be 970(12) keV which agrees well with the decay energy of 970.5(7) keV from the 2003 mass evaluation (2003Au03) which confirms the internal consistency of this scheme.

The half-life values available are, in hours:

18.0	1948Kr03
18.0(2)	1949Bu01
18.0(3)	1960Wi10
18.56(8)	1966Da19
18.479(4)	1989Ab05

18.479(7) Adopted value

The weighted average of the last four values in the Limitation of Relative Statistical method, as implemented in the LWEIGHT code, is completely dominated by the value of 1989Ab05 which has 99.7% of the relative weight. The data of 1949Bu01 and 1960Wi10 contribute 2.8 to the reduced χ^2 value of 3.1, but since this value is less than the critical reduced χ^2 value of 3.8 used in LWEIGHT for four input values, the relative weight of the dominate input value is not reduced. The internal uncertainty for this average is 0.004 and the external uncertainty is 0.007, which is adopted.

2.1 β^- Transitions

The probabilities for the β^- branches are from the probability balances from the γ -ray transitions for the excited levels and from the measurement of 1975BaXG for the ground state. These values are:

Level (keV)	Value (%)
0	57.8(12)
58	29.6(12)
137	0.012(9)
348	0.315(4)
363	12.19(6)
580	0.0626(8)
617	0.0300(9)
674	0.00388(10)
854	0.0162(5)
891	0.0009(4)

The other measured values from 1975BaXG are 24(4) for the level at 58 keV and 13(2) for the levels at 348 and 363 keV.

2.2 g Transitions

The multiplicities are from the Adopted data in the Nuclear Data Sheets (2003He11). See sect. 4.2 for comments on the γ -ray and level energies and the normalization of relative photon emission probabilities to absolute values. The multiplicities are as follows:

() indicates a tentative assignment, based on experimental data;

[] indicates an assignment based on the spins and parities of the associated levels:

Levels and $J\pi$'s	γ energy (keV)	multiplicity	mixing ratio	%E2
58 5/2+ 0 3/2+	58	M1+E2	+0.119(2)	1.40(6)
137 7/2+ 58 5/2+	79	M1+E2	+0.126(8)	1.56(20)
0 3/2+	137	[E2]		
348 5/2+ 137 7/2+	210	[M1,E2]		
58 5/2+	290	[M1,E2]		
0 3/2+	348	M1+E2	+0.43(+10, -9)	16(6)
363 5/2- 137 7/2+	226	E1		
58 5/2+	305	E1		
0 3/2+	363	E1		
580 1/2+ 0 3/2+	580	[M1,E2]		
617 3/2+ 58 5/2+	559	M1+E2	0.67(+58, -1)	31(+30, -1)
0 3/2+	617	(M1)		
674 5/2+ 137 7/2+	536	(M1)		
58 5/2+	616	(M1)		
0 3/2+	674	(M1)		
854 (1/2-) 617 3/2+	237	[E1]		
580 1/2+	274	[E1]		
0 3/2+	854	[E1]		
891 (5/2-) 617 3/2+	273	[E1]		
137 7/2+	753	[E1]		

See section 4.2 for the γ -ray energies and emission probabilities.

3 Atomic Data

3.1 X rays and Auger electrons

The fluorescence yield data are from Schönfeld and Janßen (1996Sc06) and the EMISSION code. These give $\omega_K = 0.935(4)$, the average $\omega_L = 0.186(8)$, and $\eta_{KL} = 0.847(4)$.

The Auger electron emission intensities are from the EMISSION code and based on the adopted γ -ray emission probabilities and conversion coefficients. These values are KLL 0.94(7)%, KLX 0.49(4)%, and KXY 0.063(5)%.

4 Emissions

4.1 K x-rays

The K x-ray electron emission probabilities are from the EMISSIONS code and based on the adopted γ -ray emission probabilities and conversion coefficients.

4.2 Photon Emission

Values for the γ -ray energies are available from 1968Hi03, 1969Br05, and 1995Mo08. Any weighted average would be dominated by the values of 1995Mo08, so the values from the latter reference are adopted.

The γ -ray energies from these references are:

1968Hi03	1969Br05	1995Mo08
58.00(1)	58.00(5)	58.0000(22)
79.45(2)	79.52(2)	79.5132(27)
137.7(3)	137.4(2)	137.515(5)
210.8(3)	210.9(5)	210.783(3)
226.00(4)	226.2(2)	226.0406(18)
236.9(4)	237.5(2)	237.341(5)
		273.62(12)
274.2(6)	274.2(2)	274.163(19)
290.2(3)	290.3(2)	290.2865(25)
305.6(2)	305.5(2)	305.5492(20)
348.17(8)	348.1(2)	348.2807(18)
363.56(3)	363.3(2)	363.5430(18)
		479.84(6)
536.7(4)	536.8(2)	536.730(12)
559.9(3)	559.56(15)	559.623(6)
581.1(3)	580.84(15)	580.808(6)
	616.5(3)	616.233(18)
617.7(3)	617.7(2)	617.615(18)
	674.3(5)	674.26(5)
		753.74(6)
854.5(4)	854.9(2)	854.947(20)

For the relative γ -ray emission probabilities, the following data were used. All the values of 1965Fu14 are omitted since the normalization value of 100 has a 30% uncertainty.

g ray (keV)	1964Pe07	1965Fu14	1968Hi03	1969Br05	1985Da31	1994St05	1995Mo08	2001Ma01	Adopted	Reduced c ²
58			18.0(30)	21(2)	19.1(8)	22.7(4)	18.9(9)	20.7(3) ^{ac}	21.1(6)	6.2
79		0.44(8)	0.38(7)	0.38(4)	0.37(6)	0.36(2)	0.417(11)	0.388(14)	0.397(9)	1.52
137		0.10(3)	0.042(26)	0.06(1)	0.05(1)	0.05(1)	0.0550(13) ^f		0.0549(13)	0.25
210			0.090(35)	0.165(25)	0.16(3)	0.192(23)	0.178(4) ^c		0.170(12)	1.66
226			1.8(1)	1.96(10)	1.80(4)	1.92(10)	1.89(4)	1.83(1)	1.842(18)	0.99
237			0.055(36)	0.072(11)	0.059(12)	0.064(12)	0.0652(14) ^f		0.0653(14)	0.33
246					0.012(7)		< 0.0008			
269					0.013(9)		< 0.0004			
273		0.065(3) ^b	0.065(40) ^b	0.054(13) ^b	0.056(11) ^b	0.055(12) ^b	0.0065(25)		0.006(3)	
274							0.0478(25)		0.048(3)	
290			0.24(3)	0.28	0.23(5)	0.27(3)	0.275(5)	0.274(8)	0.274(4)	0.43
305			0.54(4)	0.55(4)	0.51(2)	0.52(2)	0.527(10)	0.527(9)	0.526(6)	0.25
348	2.0(3)		2.0(1)	2.00(15)	1.99(8)	2.04(10)	2.05(4)	1.99(1) ^{cc}	2.031(21)	1.86
363	≡ 100(5)	100(30)	100	100(5)	100	100	100	100	100	
371					0.006(4)		< 0.0003			
429					0.005(4)		< 0.0003			
536	0.07 (4)	< 0.02	0.018(12)	0.010(3)	0.018(9)	0.013(3)	0.0137(4) ^f		0.0136(4)	0.48
559	0.25(10)	0.23(4)	0.17(3)	0.20(2)	0.19(2)	0.19(1)	0.187(6)		0.188(5)	0.20
581	0.70(15)	0.5(2)	0.55(4)	0.57(4)	0.60(4)	0.57(2)	0.578(19)	0.581(5) ^f	0.588(5)	0.24
616	0.20(8) ^d	0.02(1)	0.009(6)	0.020(5)	0.016(6)	0.026(8)	0.0159(7) ^f		0.0160(7)	0.90
617		0.15(5)	0.13(4)	0.13(2)	0.15(3)	0.14(1)	0.134(5) ^f		0.135(4)	0.15
674				0.0034(10)	< 0.008	0.0034(13)	0.00263(20) ^f		0.00268(19)	0.044
753							0.00153(17)		0.00153(17)	
854		0.015(7)	0.014(8)	0.021(3)	0.020(6)	0.021(2)	0.0212(18)		0.0209(12)	0.20

^a Authors also give value of 20.1(8). The most precise value is adopted.

^b Value is for sum of 273 and 274 lines.

^c Authors also give value of 2.11(3), both values are included in the calculation of the average.

^d Value is for sum of 616 and 617 lines.

^e This uncertainty was increased in the averaging process to reduce the relative weight to 50%.

^f Value contributes over 70% of the relative weight in the calculation of the average, but since the input values are consistent this weight is not reduced.

These relative γ -ray emission probabilities have been scaled by 0.1178(5) to obtain absolute values based on the measured emission probability of 11.78(5)% from 2001Ma01.

5. Electron emissions

The internal-conversion electron emission probabilities are from the adopted γ -ray emission probabilities and the associated conversion coefficients. These values for the stronger lines are:

g-ray energy (keV)	shell, energy	emission probability (%)
58	K, 6.004	22.8(9)
	L, 49.292	3.86(16)
	M, 56.032	0.85(4)
	N+, 57.602	0.235(10)
79	K, 27.518	0.17(7)
	L, 70.805	0.0273(11)
	M, 77.546	0.00604(23)
	N+, 70.115	0.00167(5)
137	K, 85.519	0.00307(12)
	L, 128.807	0.00179(7)
210	K, 158.787	0.0036(11)
226	K, 174.045	0.00629(19)
290	K, 238.291	0.0024(8)
348	K, 296.285	0.0134(5)
	L, 339.573	0.00201(6)
363	K, 311.547	0.104(3)
	L, 354.835	0.0145(4)
	M, 361.576	0.00313(9)
581	K, 528.812	0.00084(25)

6 References

- 1948Kr03 - N. L. Krisberg, M. L. Pool, C. T. Hibdon, Phys. Rev. 74(1948)1249 [T_{1/2}].
 1949Bu01 - F. D. S. Butement, Phys. Rev. 75(1948)1276 [T_{1/2}].
 1960Wi10 - R. G. Wille, R. W. Fink, Phys. Rev. 118(1960)242 [T_{1/2}].
 1964Pe07 - L. Persson, Arkiv Fysik 25(1964)307 [I _{γ}].
 1965Fu14 - L. Funke, H. Graber, K.-H. Kaun, H. Sodan, L. Werner, Nucl. Phys. 70(1965)353 [I _{γ}].
 1966Da19 - W. R. Daniels, D. C. Hoffman, J. Inorg. Nucl. Chem 28(1966)2424 [T_{1/2}].
 1968Hi03 - J. C. Hill, M.L. Wiedenbeck, Nucl. Phys. A111(1968)457 [E _{γ}].
 1969Br05 - R. A. Brown, K. I. Roulston, G. T. Ewan, G. I. Andersson, Can. J. Phys. 47(1969)1017 [E _{γ} , P _{γ}].
 1975BaXG - N. B. Badalov, S. O. Omanov, Proc. 25th Ann. Conf. Nucl. Spectrosc. and Structure At. Nuclei, Leningrad (1975)120 [P _{β}].
 1985Da31 - S. M. Darwish, S. Abdel -Malak, M. Abou-Leila, S. M. El -Bahi, A. M. Hassan, Nucl. Sci. J. (Taiwan) 22(1985)83 [I _{γ}].
 1994St05 - N. M. Stewart, N. I. Fawwaz, F. S. Radhi, Z. Phys. A348(1994)9 [I _{γ}].
 1995Mo08 - M. Moralles, P. R. Pascholati, V. R. Vanin, O. Helene, Appl. Radiat. Isot. 46(1995) 133 [E _{β} , I _{γ}].
 1996Sc06 - E. Schönfeld, H. Janßen, Nucl. Instr. Meth. Phys. Res. A369(1996)527 [ω].
 1996ScZX - E. Schönfeld, H. Janßen, report PTB-6.11-1999-1 (Feb. 1999) [I_x].
 2001Ma01 - N. Marnada, H. Miyahara, N. Ueda, N. Hayashi, K. Ikeda, ApplRadiat. Isot. 54(2001)695 [I _{γ} , P _{γ}].
 2003Au03 - G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)337 [Q].

¹⁶⁶Ho - Comments on evaluation of decay data by E. Schönfeld and R. Dersch

1 Decay Scheme

Below the Q value of 1854,5 keV there are several other excited levels of ¹⁶⁶Er which are populated in the disintegration of ¹⁶⁶Ho^m ($T_{1/2} = 1200$ a) and ¹⁶⁶Tm ($T_{1/2} = 7,70$ h). Beta transitions from ¹⁶⁶Ho to these levels, if existing, would have high degrees of forbiddenness so that they are not populated in the ¹⁶⁶Ho decay (or with extremely low transition probabilities). Thus, the decay scheme, given on page 1, can be considered as complete. Spins, parities and half-lives of the excited levels, and $lg\ f_t$ were taken from Ignatovich et al. (1987).

2 Nuclear Data

Following half-life measurements have been taken into account ($T_{1/2}$ in h):

1	27,5	Inghram and Hayden	1947
2	26,8(4)	Grant and Hill	1949
3	26,9(1)	Cork et al.	1958
4	26,8(2)	Funke et al.	1963
5	26,74(5)	Daniel and Kaschl	1966
6	27,00(4)	Venkata Ramaniah et al.	1976
7	26,827(5)	Abzouzi et al.	1989
8	26,78(1)	Calhoun et al.	1991
9	26,7663(44)	Unterweger et al.	1992
10	26,795(29)	adopted value	1999

Value 1 is only of historical interest. Value 8 is replaced by value 9, value 6 is considered as outlier (or its accuracy is overestimated). The adopted value is the LWM of values 2-5, 7 (with doubled uncertainty to take account for systematical errors) and 9. LWM has used weighted average and expanded the uncertainty so range includes the most precise value 9. The rather large uncertainty reflects the discrepancy between the values 7 and 9.

2.1 β^- Transitions

The maximum beta energy of the transition to the ground state of ¹⁶⁶Er and the transition probability of this transition have been determined as follows:

1	1840	25 %	Sunyar 1954
2	1854(5)	51,6 %	Graham et al. 1955
3	1839(5)	47 %	Cork et al. 1958
4	1844	52 %	Marklund et al. 1960
5	1840	46 %	Cline et al. 1962
6	1859(3)	48,8 %	Funke et al. 1963
7	1857(3)	48,8 %	Daniel and Kaschl 1966
8	1854,7(15)	51,2 %	Grigoriev et al. 1974
9	1845(2)	52 %	Venkata Ramaniah et al. 1976
10	1854,8(17)		weighted average of values 2, 6 - 9 (see text below)
11	1854,5(9)		Audi and Wapstra 1995. Here adopted too

For the calculation of the average value 10, the originally given uncertainty of value 9 has been doubled before inserting it in the averaging procedure because the uncertainty seems to be overestimated. The unweighted average for the transition probability to the ground state (including values 2 to 9) is 49,6 %. This value agrees satisfactorily with the adopted value 48,2(15) % which was derived in the balancing procedure from the gamma transition probabilities.

2.2 Gamma Transitions

The energies of the gamma transitions are calculated from the gamma ray energies (section 4.2) taking the recoil energies into account which can be neglected in most cases. The probabilities $P_{\gamma+ce}$ are calculated from the gamma ray emission probabilities and the total conversion coefficients.

The conversion coefficients are interpolated from the tables of Rösels et al. (1978). Very much work has been spent for the study of the conversion of the 80,57 keV gamma transition. The K conversion coefficient of this transition was found to be

1	1,69(9)	Ramaswamy and Brahmavar	1963
2	1,63(5)	Falkstroem et al.	1968
3	1,72(6)	Nelson and Hatch	1969
4	1,69(6)	Campbell et al.	1971
5	1,66(6)	Campbell et al.	1972
6	1,65(5)	interpolated from Rösels et al.	1978; adopted value

For the K/L ratio the following values were found:

1	0,390(18)	Bogdanovic et al.	1968
2	0,426(11)	Nilsson et al.	1968
3	0,414(13)	Kartashov et al.	1977
4	0,411(12)	interpolated from Rösels et al.	1978; adopted value

Kartashov et al. (1977) have also determined the ratios M/L, N/M and O/N. From their measurements the following set can be derived:

$$\alpha_K = 1,650(33)$$

$$\alpha_L = 3,983(170)$$

$$\alpha_M = 0,990(50)$$

$$\alpha_N = 0,200(12)$$

$$\alpha_{OP} = 0,048(3)$$

$$\alpha_t = 6,87(18)$$

The total conversion coefficient of this transition was determined by Brandtley et al. (1966) to be $\alpha = 6,94(48)$. Several other authors have determined L subshell ratios (Hermann et al. (1966), Gelletly et al. (1966, 1967), Karlsson et al. (1966), Zylicz et al. (1966), Arnoux and Gizon (1967), Bogdanovic et al. (1968)). Also M and N subshell ratios were determined (Hoegberg et al. (1968), Dragoun et al. (1972), Bulgakov et al. (1981)).

The conversion coefficients contained in table 2.2 are interpolated from the tables of Rösels et al. (1978).

3 Atomic Data

The atomic data are taken from Schönfeld and Janßen (1996).

3.1 X Radiation

The energies are based on the X ray wave lengths compiled by Bearden (1967). The relative probabilities are calculated using the ratios $P(K_{\beta_2})/P(K_{\alpha_1})$ and $P(K_{\beta_1})/P(K_{\alpha_1})$ as given by Schönfeld and Janßen (1996). The relative probability of X_L radiation is calculated from the absolute value putting $P(K_{\alpha_1}) = 100$.

3.2 Auger Electrons

The energies are taken mainly from the report of Larkins (1977). The relative probabilities are calculated using the ratios $P(KLX)/P(KLL)$ and $P(KXY)/P(KLL)$ as given in the cited report of Schönfeld and Janßen (1995). The relative probability of e_{AL} electrons is calculated from the absolute value putting $P(KLL) = 100$.

4 Radiation Emission

4.1 Electron Emission

The numbers of Auger electrons per disintegration are calculated using the program EMISSION and the atomic data as given in Section 3. The numbers of conversion electrons per disintegration are calculated using the conversion coefficients and the probabilities $P_{\gamma+ce}$ as given in 2.2. Spectra of the conversion electrons from the 80,6 keV

transition, the 1379,4 keV transition and the $0^+ \rightarrow 0^+$ 1460 keV E0 transition were measured by Grigoriev et al. (1974). The data for the emission of β particles are those already given in 2.1.

4.2 Photon Emission

Most of the gamma-ray energies were taken from Ardisson et al. (1992) ($\gamma_{1,0}$, $\gamma_{4,3}$, $\gamma_{3,1}$, $\gamma_{3,0}$, $\gamma_{4,1}$, $\gamma_{5,0}$, $\gamma_{6,1}$, $\gamma_{6,0}$, $\gamma_{8,1}$, $\gamma_{7,0}$, $\gamma_{8,0}$).

The following measurements of relative photon emission probabilities have been taken into account (the relative emission probability of the 1379,4 keV line was arbitrarily set to 1):

E in keV	1	2	3	4	5	6	7
80,6	6,67(43)	-	7,04(30)	6,72(70)	7,22(8)	6,56(40)	7,02(14)
184,4	-	0,0022(5)	-	0,0013(3)	0,0023(1)	0,0010(1)	0,0016(7)
521,0	-	-	-	0,00032(11)	0,0005(2)	0,00038(1)	0,00038(2)
674,2	0,032(2)	0,022(2)	0,034(2)	0,0176(9)	0,023(1)	0,0201(4)**	0,0212(18)
705,4	0,020(3)	0,016(2)	0,023(1)	0,0137(7)	0,0170(10)	0,0144(3)**	0,0156(13)
785,9	0,016(3)	0,014(2)	0,012(5)	0,0125(7)	0,0140(10)	0,0128(3)**	0,01288(27)
1263,0	-	-	-	0,0015(2)	0,0017(1)	0,0016(3)	0,00166(9)
1379,4	1	1	1	1	1	1	1
1447,5	-	-	-	0,00105(10)	0,0012(1)	0,0014(5)	0,00113(10)
1528,2	-	-	-	0,0002	-	0,00010(1)	0,00015(5)
1581,8	0,206(10)	0,195(10)	0,215(10)*	0,197(7)	0,199(5)	0,197(5)	0,1994(28)
1662,4	0,129(7)	0,125(6)	0,099(5)*	0,130(5)	0,127(4)	0,130(2)**	0,126(5)
1731,5	-	-	-	-	-	0,00005(2)	0,00005(2)
1749,8	0,033(1)*	0,027(2)	0,030(17)	0,028(2)	0,028(1)	0,0285(6)**	0,0292(9)
1812,8	-	-	-	-	-	0,00006(2)	0,00006(2)
1830,5	0,0100(8)*	0,0086(11)	0,0081(5)	0,0089(5)	0,0085(2)	0,0089(3)	0,0087(2)

1 Burson et al. 1967

2 Reich and Cline 1970

3 Venkata Ramaniah et al. 1976

4 Allab et al. 1977

5 Chand et al. 1989

6 Ardisson et al. 1992

7 values adopted in this evaluation (LWM)

* classified as outlier (appearing only in values of references 1 and 3)

** input uncertainty slightly increased (only for some values of reference 6 and one value of reference 5)

Earlier results of Marklund et al. (1960), Hansen et al. (1961), Cline et al. (1962), Funke et al. (1963) and Neumann (1966) were not taken into account because they are less accurate, incomplete and given without uncertainties.

The absolute emission probability for the gamma rays from the transition $\gamma_{1,0}$ (80,6 keV) has been determined as follows (gamma rays per 100 disintegrations):

1	6,55(30)	Venkata Ramaniah et al. 1976
2	6,25(60)	Allab et al. 1977
3	6,60(40)	Sekine and Baba 1981
4	6,55(8)	Calhoun et al. 1991; Coursey et al. 1994

In the present evaluation value 4 is adopted. Combining it with the relative emission probability of the 80,6 keV transition, the normalization factor 0,933(16) is obtained.

5 Main Production Modes

Taken from Firestone (1995).

6 References

- M. G. Inghram and R. J. Hayden, *Phys. Rev.* 71 (1947) 130
[$T_{1/2}$]
- P. J. Grant and J. M. Hill, *Nature* 163 (1949) 524
[$T_{1/2}$]
- M. Antoneva, A. A. Bashilov, B. S. Dzheleпов, *Izv. Acad. Nauk SSSR* 14 (1950) 299
[$T_{1/2}$]
- J. M. Cork, M. K. Brice, R. G. Helmer, R. M. Woods Jr., *Phys. Rev.* (2) 110 (1958) 526
[$T_{1/2}$]
- D. C. Hoffman, *J. Inorg. Nucl. Chem.* 25 (1963) 1196
[$T_{1/2}$]
- M. K. Ramaswamy and S. M. Brahmavar, *Curr. Sci.* (India) 32 (1963) 451
[α_K]
- W. H. Brantley, A. R. Polderman and W. H. G. Lewin, *Bull. Amer. Phys. Soc.* 11 (1966) 824
[α_4]
- H. A. Neumann, *Z. Naturforschg.* 21a (1966) 1328
[P_γ]
- P. Erman, G. T. Emery and M. L. Perlman, *Phys. Rev.* 147 (1966) 858
[$L_1/L_2/L_3$]
- W. Gelletly, J. S. Geiger and R. L. Graham, *Bull. Amer. Phys. Soc.* 11 (1966) 352
[$L_1/L_2/L_3$]
- S. E. Karlsson, I. Andersson, O. Nilsson, G. Malmsten, C. Nordling and K. Siegbahn, *Nucl. Phys.* 89(1966) 513
[$L_1/L_2/L_3$]
- J. Zylicz, M. H. Jorgensen, O. B. Nielsen and O. Skilbreid, *Nucl. Phys.* 81 (1966) 88
[K/L/M]
- M. Arnoux and A. Gizon, *Compt. Rend. Acad. Sci.* (Paris) 264 (1967) 1518
[L/M]
- W. Gelletly, J. S. Geiger and R. L. Graham, *Phys. Rev.* 157 (1967) 1043
[L/M]
- J. A. Bearden, *Rev. Mod. Phys.* 39 (1967) 78
[E(KX)]
- M. Bogdanovic, M. Mladjenovic and R. Stepic, *Z. Phys.* 216 (1968) 267
[K/L]
- M. Bogdanovic, M. Mladjenovic and R. Stepic, *Nucl. Phys.* A106 (1968) 209
[K/L]
- E. Falkstroem, S. Nilsson and S. Boreving, *Ark. Phys.* 39 (1968) 1
[α_K]
- S. Hoegberg, J. E. Bergmark, G. Malmsten and O. Nilsson, *Nucl. Phys.* A120 (1968) 569
[$M_1/M_2/M_3/M_4/M_5$]
- O. Nilsson, I. Thoren, G. Malmsten and S. Hoegberg, *Nucl. Phys.* A120 (1968) 561
[K/L]
- G. C. Nelson and E. N. Hatch, *Nucl. Phys.* A127 (1969) 560
[α_K]
- J. L. Campbell, J. J. Smith and I. K. MacKenzie, *Nucl. Instr. Meth.* 92 (1971) 237
[α_K]
- J. L. Campbell, R. J. Goble and J. J. Smith, *Radioactivity in Nuc. Spec.* J. H. Amilton and J. C. Manthuruthil (eds.) Gordon and Breach, New York (1972) p. 1387
[α_K]
- O. Dragoun, B. Martin, D. Merkert and M. Vinduska, *Nucl. Phys.* A183 (1972) 390
[$N_1/N_2/N_3/N_4/N_5$]
- F. P. Larkins *Atomic Data and Nuclear Data Tables* 20 (1977) 313
[E(KLL), E(KLX)]
- V. V. Bulgakov, V. I. Gavriilyuk, A. A. Klyuchnikov, A. P. Lasko, P. N. Muzalev, N. V. Strelchuk, A. I. Feokistov and Y. E. Frantsev, *Summaries of the 30. Conf. Nucl. Spectr. and Struct.*, Samarkand, (1981) p. 293
[L/M/N]
- R. B. Firestone and V. S. Shirley, *Table of Isotopes*, Wiley, New York, 1996
[Production modes]

Other references can be found in the Tables Part.

¹⁶⁶Ho^m - Comments on evaluation of decay data by E. Schönfeld, R. Dersch

1 Decay Scheme

The decay scheme was taken from Ardisson *et al.* 1992. It contains 54 gamma transitions between 17 excited levels of ¹⁶⁶Er or to the ground state of this nuclide. This decay scheme is not complete. 12 additional gamma rays have been reported, six of them from branching in Tm-166 EC decay (see 2.2).

The half-lives of the excited level in ¹⁶⁶Er indicated in the decay scheme are taken from Shursikow and Timofeeva (1992).

2 Nuclear Data

The half-life was determined by Faler (1965) to be 1200 a. The uncertainty was estimated to be 180 a. New measurements are desirable. The Q-value is 6,0 keV above Q(¹⁶⁶Ho). This is the energy difference between the isomer level and the ground state of ¹⁶⁶Ho. The Q-value of ¹⁶⁶Ho was derived from β -ray endpoint energies to be 1854,5(9) keV. Thus, the Q-value of ¹⁶⁶Ho^m is 1860,5(9) keV.

2.1 β^- Transitions

There are seven β transitions to excited levels of ¹⁶⁶Er. The most important transitions are the allowed transitions to levels no. 17 and 16 (17,2(4) % and 74,8(12) %). Weak transitions are feeding the levels 11, 10, 9, 6 and 3. Transitions to the levels 15, 14, 13, 12, 8, 7, 5, 4, 2, 1 and the ground state ($\Delta J_0 = 7$) have not been observed. All these transitions are at least second forbidden except a transition to level 8 which is unique first forbidden.

The energies of these transitions were calculated by subtracting the level energy from the Q-value. The transition probabilities P_β were calculated from the transition probabilities $P_{\gamma+ec}$ using the relations which correspond to the decay scheme.

2.2 Gamma transitions

The level differences are equal to the gamma-ray energies as the recoil energies are small compared with the uncertainties of the latter. The gamma-ray energy of the 80,6 keV emission has been determined as follows (energy in keV):

1	80,573	Reich and Cline 1970
2	80,589(5)	Morii et al. 1975 .
3	80,572(15)	Souch et al. 1982
4	80,585(15)	Adam et al. 1988
5	80,574(8)	Hardell and Nilsson 1962; cryst.-spektr.
6	80,5725(13)	Helmer and van der Leun 2000; here also adopted

The energies of gamma transitions between the levels 0, 1, 2, 3, 5, 6, 7, 8, 9, 10 and the transitions $\gamma_{16,5}$ and $\gamma_{17,3}$ are taken from Helmer and van der Leun (2000). The energies of all other transitions are either taken from Ardisson *et al.* (1992) or based on values given by these authors.

The probabilities $P_{\gamma+ce}$ were calculated from the gamma -ray emission probabilities P_{γ} using the values for the total conversion coefficients α_t . The conversion coefficients α_K , α_L and α_t were interpolated from the tables of Röseler *et al.* (1978). The normalization factor which is necessary to convert relative emission probabilities (related to 100 for the 184 keV gamma rays) can be calculated from balancing conditions using cuts between the levels 0 and 1, 1 and 2, 2 and 3. This is possible because the levels 2, 1 and 0 (the ground state) are not populated by β transitions. The cut between the levels 0 and 1 contains the emission probability of the 80,6 keV gamma transition. The conversion coefficient of this transition has a relatively large uncertainty, the calculation of the normalization factor from the cuts 1-2 and 2-3 is therefore preferred here. Moreover, the normalization factor was determined using absolute activity measurements:

1	0,732(37)	Reich and Cline, 1970
2	0,699(14)	Danilenko et al., 1989
3	0,7258(22)	Miyahara et al., 1994
4	0,7021(35)	Morel et al., 1996
5	0,7235(67)	Hino et al., preliminary value, 1999
6	0,7214(72)	from cut between levels 1 and 2, this evaluation 1999
7	0,7298(75)	from cut between levels 2 and 3, this evaluation 1999
8	0,725(3)	adopted value

The value 8 is the LWM between values 1, 3, 5, 6 and 7 where the uncertainty of value 3 has been doubled in order to contribute less than 50 % to the mean. Values 2 and 4 are considered to be significantly too low by the evaluator and were not included in the averaging procedure. The reduced χ^2 of the LWM is 0,2. The adopted value of the normalization factor is in excellent agreement with the value 0,726(9) evaluated by Shursikow and Timofeeva (1992).

The K-conversion coefficients were calculated using the tables of Röseler *et al.* (1978). The multiplicities of the transitions were determined from the spin and parity assignments as made by Ardisson *et al.* (1992) and Shursikow and Timofeeva (1992). There is reasonable agreement between measured and calculated conversion coefficient for the 80,6 keV transition:

1	1,76(15)	Marklund et al. 1960
2	1,72(6)	Nelson and Hatch 1969
3	1,69(6)	Campbell et al. 1971
4	1,65(3)	E2 Theory, Röseler et al. 1978

The following gamma rays are not included in the decay scheme and in the tables 2.2 and 4.2:

E_{γ} in keV	P_{rel} (related to 100 for the 184,4 keV line)	
96,85(5)	0,00307	*
170,31(3)	0,0184(11)	*
255,20(12)	0,0059(13)	
410,80(5)	0,0231(7)	*
520,945(15)	0,00039(7)	*
617,0(5)	0,031(9)	
712,89(13)	0,41(12)	*
736,02(8)	0,19(2)	
1446,72(13)	< 0,01	
1521,99(4)	0,018(5)	
1562,57(4)	0,0040(11)	

* Deduced from branching in Tm-166 EC decay where also the 73 keV transition, contained in Table 2, occurs. These data are taken from Shursikow and Timofeeva (1992), see also Adam *et al.* (1979).

For several transitions, mixing ratios were determined from γ - γ angular correlation measurements. Most of them are compiled in the following table:

E2-M1 mixing ratios for γ -transitions in ¹⁶⁶Er following the decay of ¹⁶⁶Ho^m

E_r in keV	d	d (adopted)	% M1
119,0	$\pm 1,79(12)[1]$ $1,75(12)[2]$	1,79(12)	24(2)
140,7	$\pm 1,43(10)[1]$ $1,67(11)[2]$	1,43(10)	33(3)
160,1	$1,45(11)[1]$	1,45(11)	32(4)
464,8	$-(3,1+1,5-0,9)[3]$ $-80<\delta<+30[4]$ $-(32+98-14)[5]$ $-(63+19-12)[6]$	-50(20)	(0,04+0,07-0,02)
529,8	$-(85+8-45)[7]$ $-25(3)[4]$ $-5,0(25)[3]$ $-(25+5-4)[5]$ $-(62+40-17)[8]$ $-(60+45-19)[8]$	-30(20)	(0,11+0,9-0,07)
594,1	$-(9+319-5)[4]$ $(9+8-5)[5]$ $-(12+29-5)[8]$ $-(8+15-3)[8]$ $-(59+74-21)[2]$	-10(5)	(1+3-0,5)
644,5	$ \delta >2[4]$ $+1,6+1,0-0,55[3]$ $-0,75(20)[3]$ $<-1 \text{ or } >+4[8]$ $-(13,4+3,3-2,2)[2]$	3-2+3	(10+40-7)
670,5	$6,3+8-2,9[3]$ $-(1,15+0,80-0,35)[3]$ $-(20+90-9)[4, 5]$ $(10,0+1,6-1,2)[8]$ $9,4+2,9-1,6[8]$ $(19+5-3)[2]$	12(5)	(0,69+1,31-0,35)
691,3	$3,3+3,0-1,2[9]$ $-(10+27-4)[4]$ $-(16+27-4)[5]$ $-(28+7-5)[2]$ $-(16+8-9)[8]$ $-(16+8-10)[8]$	-16(8)	(0,39-0,22+1,15)
705,2	$ \delta =25[10]$ $38+8-24[9]$ $19+38-9[9]$ $-(55+13-9)[2]$	50(10)	(0,04+0,02-0,01)
778,8	$-(20+8-13)[3]$ $-(18+8-9)[4]$ $-(19+8-10)[5]$ $-(20+4-2)[8]$ $-(18+8-5)[8]$ $-(109+26-17)[2]$	18(6)	(0,31+0,35-0,14)
810,3	$37+10-7[7]$ $-16,4+3,2-2,3[11]$ $-20(4)[4]$ $-(84+8-57)[3]$ $-(20+4-3)[5]$ $-(36+11-7)[6]$ $-21(2)[8]$ $-15(1)[8]$	25(5)	(0,16+0,09-0,05)
830,6	$70+260-30[7]$ $-(42+25-13)[11]$ $-(22+7-5)[4,5]$ $-(37+8-17)[3]$ $-(18+3-2)[6]$ $-23(4)[8]$ $-(16,6+1,8-1,5)[8]$ $-(15,3+2,3-1,7)[2]$	-18(3-2)	0,31(8)

- [1] Wagner 1992, measured
- [2] Wagner 1992, calculated
- [3] West et al. 1976
- [4] Baker et al. 1975
- [5] Lange et al. 1981
- [6] Alzner et al. 1985
- [7] Reich and Cline 1965
- [8] Krane and Moses 1981
- [9] Domingos et al. 1972
- [10] McGowan et al. 1978
- [11] Miyokawa et al. 1972 as cited in the paper of Krane and Moses 1981

Some of the measurements are discrepant. However, the influence of the results on the conversion coefficients is in most cases small. Gerda et al. (1963) determined some mixing ratios from γ - γ angular correlations. Some of them deviate from the results of later publications (411 keV 95 % E1 + 5 % M2; 712 keV 99,6 % E1 + 0,4 % M2; 810 keV 99,1 % E2 + 0,9 % M1; 831 keV 96,1 % E2 + 3,9 % M1).

If two multiplicities are mentioned in Table 2.2, then the mixing ratio was taken into account when calculating the conversion coefficients. If a second multiplicity is given in brackets, then the conversion coefficients are calculated for the first multiplicity but an admixture of the second multiplicity is not ruled out.

3 Atomic Data

The atomic data are taken from Schönfeld and Janßen (1996).

3.1 X Radiations

The energies are based on the wavelengths of Bearden (1967). The relative probabilities are taken from Schönfeld and Janßen (1996). The relative probability of the L X rays is calculated from the absolute value (Table 4) setting $P(K_{\alpha 1}) = 1$.

3.2 Auger electrons

The energies are taken mainly from Larkins (1977). The relative probabilities are taken from Schönfeld and Janßen (1996). The relative probability of the L Auger electrons is calculated from the absolute value (Table 4) setting $P(KLL) = 1$.

4 Radiation Emissions

4.1 Electron Emissions

The energies of the Auger electrons are the same as in 3.2. The energies of the conversion electrons are calculated from the transition energy (2.2) and the binding energies. The emission probabilities of the Auger electrons are calculated from P_{γ} 's and conversion coefficients using the program EMISSION (PTB, 1997).

The emission probabilities of the conversion electrons are calculated using the conversion coefficients given in Table 2.2, the atomic data given in Section 3, and the emission probabilities of the gamma rays given in Table 4.2.

4.2 Photon Emissions

The energies of the X rays are the same as in Table 3.1. Measured K X-ray emission probabilities (Chand *et al.* 1988, Morel *et al.* 1996) are in good agreement with the calculated values. If the measured values are related to the here adopted emission probability of the 184-keV gamma rays, the following values are obtained (quanta per 100 disintegrations):

	E in keV	P_X (Chand)	P_X (Morel)	P_X (calc)
Er $K_{\alpha 2}$	48,221	10,95(23)	10,63(8)	10,81(21)
Er $K_{\alpha 1}$	49,128	18,4(3)	19,17(13)	19,2(4)
Er $K'_{\beta 1}$	55,624	5,70(9)	6,03(5)	6,24(14)
Er $K'_{\beta 2}$	57,239	1,41(3)	1,594(20)	1,62(5)

The calculated emission probabilities of the X-rays (calculated from P_{γ} 's and conversion coefficients using the program EMISSION (PTB, 1997)) are compiled in the last column.

The energies of the gamma rays are taken either from Helmer and van der Leun (2000) or from Ardissone *et al.* (1992) (see Sect. 2.2). Their uncertainties are to be considered as standard uncertainties.

The relative emission probabilities of gamma rays (related to 100 for the emission probabilities of the 184,4 keV transition $\gamma_{2,1}$) as measured by 17 authors are compiled in the following table. The last column in this table contains the LWM except of $\gamma_{1,0}$ where balance conditions are taken into account. The transition probability of the transition $\gamma_{1,0}$ is very well known as there is only one other transition to the ground state which is very weak ($\gamma_{4,0}$):

$$f_N [P_{\text{rel}}(\gamma_{1,0}) (1 + \alpha_t) + P_{\text{rel}}(\gamma_{4,0}) (1 + \alpha_t)] = 100$$

The conversion coefficient is, of course, to put for the assigned gamma transition. This yields for the transition $\gamma_{1,0}$

$$P_{g+ce} = 100 - f_N P_{\text{rel}}(\gamma_{4,0}) (1 + \alpha_t)$$

$$\text{With } f_N = 0,725(3), P_{\text{rel}}(\gamma_{4,0}) = 0,026(5), \alpha_t(\gamma_{4,0}) = 0,00566(12)$$

we obtain :

$P_{g+ce}(\gamma_{1,0}) = 99,981(4)$ per 100 disintegrations. With the conversion coefficient of the transition $\gamma_{1,0}$ this yields:

$$P_\gamma(\gamma_{1,0}) = 12,66(23) \text{ per 100 disintegrations, in relative units: } 17,46(31).$$

Gamma relative emission intensities, references 1 to 6 :

$g_{i,f}$	E_g (keV)	1	2	3	4	5	6
$\gamma_{1,0}$	80,577(7)	14,5(29)	14,55(47)	17,1(9)	14,48(48)	16,83(42)	16,7(10)
$\gamma_{16,15}$	94,679(9)	0,16(3) ¹⁾	-	0,19(1)	0,3	0,21(3)	-
$\gamma_{8,7}$	119,035(10)	-	-	0,24(3)	-	0,23(3)	-
$\gamma_{16,14}$	121,175(10)	0,7(5) ¹⁾	-	0,36(5)	0,78(18) ¹⁾	0,54(5) ¹⁾	-
$\gamma_{17,15}$	135,257(14)	0,1(1)	-	0,14(2)	-	-	-
$\gamma_{9,8}$	140,702(20)	-	-	0,059(14)	-	-	-
$\gamma_{10,9}$	160,077(20)	0,35(10)	-	0,134(16)	0,36(15) ¹⁾	0,16(3)	-
$\gamma_{17,14}$	161,707(14)	-	-	0,15(2)	-	0,16(3)	-
$\gamma_{3,2}$	184,404(7)	100	100	100	100	100	100
$\gamma_{16,13}$	190,747(16)	-	-	0,30(3)	-	0,31(4)	-
$\gamma_{16,12}$	214,79(3)	-	-	0,75(10) ¹⁾	-	-	-
$\gamma_{8,5}$	215,871(10)	3,8(4)	4,15(7)	3,6(4)	3,94(9)	3,96(8)	4,1(2) ²⁾
$\gamma_{17,12}$	231,32(4)	0,3(2)	0,32(5)	0,33(4)	0,36(3) ¹⁾	0,31(4)	-
$\gamma_{9,7}$	259,70(3)	1,8(5) ¹⁾	1,42(10)	1,50(11)	1,77(12) ¹⁾	1,52(5)	-
$\gamma_{9,2}$	280,468(7)	39,5(28)	43,6(6) ¹⁾	40,7(29)	38,61(46)	39,63(126)	40,2(18)
$\gamma_{10,9}$	300,731(9)	4,8(4)	5,45(8)	5,12(37)	4,77(9)	4,92(12)	4,97(22)
$\gamma_{9,6}$	305,03(5)	-	-	-	-	-	-
$\gamma_{11,9}$	339,75(5)	-	-	0,23(3)	-	0,23(4)	-
$\gamma_{6,3}$	365,736(9)	2,9(3) ¹⁾	3,72(8)	3,44(25)	2,93(6)	3,25(10)	3,30(11)
$\gamma_{16,10}$	410,950(8)	15,8(12)	16,8(3) ¹⁾	15,8(12)	15,50(19)	14,77(30)	15,27(50)
$\gamma_{17,10}$	451,528(9)	3,5(7)	4,30(9)	4,18(30)	3,48(7) ¹⁾	3,84(13)	3,99(13)
$\gamma_{10,6}$	464,819(12)	2,0(4)	1,66(8)	1,68(14)	2,00(7)	1,50(8)	-
$\gamma_{15,9}$	476,38(6)	0,4(2) ¹⁾	-	-	-	-	-
$\gamma_{12,8}$	496,86(4)	-	-	-	-	-	-
$\gamma_{4,2}$	520,85(5)	-	-	-	-	-	-
$\gamma_{8,3}$	529,811(10)	10,3(10) ¹⁾	13,00(42)	13,9(10)	10,16(32) ¹⁾	12,36(25)	12,78(42)
$\gamma_{16,9}$	570,940(10)	6,8(7)	7,08(16)	7,86(56)	6,77(14)	7,04(14)	7,45(24)
$\gamma_{5,2}$	594,536(24)	1,2(4) ¹⁾	0,74(10)	0,96(8) ¹⁾	1,28(18) ¹⁾	0,70(5)	-
$\gamma_{17,9}$	611,620(17)	1,4(10)	1,59(32)	1,90(15)	1,48(27)	1,67(9)	-
$\gamma_{11,7}$	615,84(9)	-	-	-	-	-	-
$\gamma_{13,7}$	639,97(9)	-	-	0,22(7) ¹⁾	-	-	-
$\gamma_{11,6}$	644,689(15)	0,27(15)	0,31(105) ¹⁾	0,25(3)	-	-	-
$\gamma_{9,3}$	670,565(12)	7,0(7)	7,35(30)	7,88(56)	7,01(25)	6,98(16)	7,37(24)
$\gamma_{7,2}$	691,304(12)	1,9(4)	1,62(8)	2,09(15) ¹⁾	1,85(9)	1,60(10) ¹⁾	1,800(59)
$\gamma_{4,1}$	705,09(7)	-	-	-	-	-	-
$\gamma_{16,8}$	711,680(8)	72,5(60)	71,5(10)	80,2(57) ¹⁾	71,65(68)	71,10(142)	74,5(25)
$\gamma_{13,5}$	736,70(7)	0,45(15)	0,50(5)	0,14(5) ¹⁾	0,46(4)	0,45(5)	-
$\gamma_{17,8}$	752,332(10)	16,1(12)	15,20(34) ¹⁾	17,9(13)	16,06(40)	15,98(32)	16,57(54)
$\gamma_{8,1}$	778,862(12)	3,8(3)	3,88(7)	4,51(33)	3,72(7)	4,16(12)	4,13(13)
$\gamma_{4,0}$	785,81(7)	-	-	-	-	-	-
$\gamma_{8,2}$	810,325(10)	76(8)	76,40(110)	85,7(61) ¹⁾	76,38(82)	75,71(151)	78,1(28)
$\gamma_{10,3}$	830,601(15)	12,5(10)	12,90(32)	14,5(11)	12,07(28)	12,83(26)	13,26(44)
$\gamma_{7,1}$	875,63(5)	1,15(15)	0,91(4)	1,08(10)	1,14(7)	1,00(9)	0,979(32)
$\gamma_{9,2}$	950,963(10)	3,6(6)	3,16(13) ¹⁾	4,15(30) ¹⁾	3,50(14) ¹⁾	3,74(16)	3,68(12)
$\gamma_{11,3}$	1010,27(6)	-	0,11(340)	0,12(2)	-	-	-
$\gamma_{14,3}$	1120,35(5)	-	0,26(2)	0,31(3)	0,30	-	-
$\gamma_{15,3}$	1146,81(9)	0,38(6) ¹⁾	0,26(2)	0,30(3)	0,38(5) ¹⁾	-	0,274(9)
$\gamma_{16,3}$	1241,52(2)	1,25(25)	1,06(4)	1,37(10) ¹⁾	1,22(5)	1,17(12)	1,098(37)
$\gamma_{17,3}$	1282,06(6)	0,80(15) ¹⁾	0,22(2)	0,31(3)	0,38(4) ¹⁾	0,24(5)	0,241(8)
$\gamma_{12,2}$	1306,60(15)	-	-	-	-	-	-
$\gamma_{13,2}$	1331,04(13)	-	-	-	-	-	-
$\gamma_{14,2}$	1400,79(2)	0,93(9) ¹⁾	0,72(2)	0,75(6)	0,86(5) ¹⁾	-	0,670(22)
$\gamma_{15,2}$	1427,24(2)	0,69(7)	0,69(2)	0,81(6) ¹⁾	0,65(3)	-	0,665(23)
-	1446,7(2)	-	-	-	-	-	-

Gamma relative emission intensities, references 7 to 12 :

$g_{i,f}$	E_g (keV)	7	8	9	10	11	12
$\gamma_{1,0}$	80,577(7)	17,51(61)	16,56(8)	17,8(4)	16,97(13)	17,2(8)	16,59(39)
$\gamma_{16,15}$	94,679(9)	0,221(12)	-	0,22(1)	0,20(1)	0,190(26)	-
$\gamma_{8,7}$	119,035(10)	0,222(12)	-	0,27(2) ¹⁾	0,24(1)	0,243(13)	-
$\gamma_{16,14}$	121,175(10)	0,337(15)	-	0,45(2) ¹⁾	0,35(2)	0,346(14)	-
$\gamma_{17,15}$	135,257(14)	0,126(10)	-	0,14(1)	0,14(1)	0,128(6)	-
$\gamma_{9,8}$	140,702(20)	0,059(9)	-	0,06(1)	0,07(1)	0,060(4)	-
$\gamma_{10,9}$	160,077(20)	0,109(8)	-	0,14(1)	0,14(2)	0,124(4)	-
$\gamma_{17,14}$	161,707(14)	0,135(8)	-	0,15(1)	0,15(2)	0,140(7)	-
$\gamma_{3,2}$	184,404(7)	100	100	100	100	100	100
$\gamma_{16,13}$	190,747(16)	0,304(15)	-	0,31(1)	0,33(2)	0,291(10)	-
$\gamma_{16,12}$	214,79(3)	0,586(23)	-	0,61(2)	0,61(2)	-	0,60(5)
$\gamma_{8,5}$	215,871(10)	3,54(13)	4,04(4)	3,67(9)	3,60(13)	4,14(17) ²⁾	3,61(13)
$\gamma_{17,12}$	231,32(4)	0,284(15)	-	0,30(1)	0,33(3)	0,289(11)	0,263(20)
$\gamma_{9,7}$	259,70(3)	1,446(52)	-	1,53(3)	1,52(3)	1,47(5)	1,50(5)
$\gamma_{9,2}$	280,468(7)	40,79(141)	41,26(28)	41,0(5)	40,6(5)	40,4(15)	40,9(8)
$\gamma_{10,9}$	300,731(9)	5,12(18)	5,22(4)	5,17(8)	5,11(8)	5,04(19)	5,13(10)
$\gamma_{9,6}$	305,03(5)	-	-	-	0,023(3)	0,030(3)	-
$\gamma_{11,9}$	339,75(5)	0,234(16)	-	0,21(1)	0,21(3)	0,222(8)	-
$\gamma_{6,3}$	365,736(9)	3,327(117)	3,30(3)	3,49(6)	3,46(6)	3,33(12)	3,44(7)
$\gamma_{16,10}$	410,950(8)	15,25(53)	15,65(10)	15,9(2)	15,5(4)	15,3(5)	15,93(28)
$\gamma_{17,10}$	451,528(9)	4,02(15)	3,85(5)	4,17(5)	4,04(11)	4,00(14)	4,12(9)
$\gamma_{10,6}$	464,819(12)	1,651(61)	-	1,67(3)	1,73(7)	1,59(5)	1,69(6)
$\gamma_{15,9}$	476,38(6)	-	-	-	0,052(6)	0,050(3)	-
$\gamma_{12,8}$	496,86(4)	-	-	0,18(3)	0,17(1)	0,170(6)	-
$\gamma_{4,2}$	520,85(5)	-	-	0,22(3)	0,21(1)	0,20(3)	0,240(24) ¹⁾
$\gamma_{8,3}$	529,811(10)	13,10(45)	12,48(10)	13,3(2)	13,18(34)	12,83(39)	13,46(26)
$\gamma_{16,9}$	570,940(10)	7,53(27)	7,22(6)	7,65(9)	7,64(20)	7,42(24)	7,81(15)
$\gamma_{5,2}$	594,536(24)	0,773(34)	-	0,77(2)	0,80(9)	0,769(24)	0,80(4)
$\gamma_{17,9}$	611,620(17)	1,951(72)	-	1,86(4)	1,86(12)	1,85(7)	1,95(11)
$\gamma_{11,7}$	615,84(9)	-	-	-	0,044(13)	0,163(8)	-
$\gamma_{13,7}$	639,97(9)	0,122(16)	-	0,12(1)	0,11(1)	0,124(6)	-
$\gamma_{11,6}$	644,689(15)	0,213(19)	-	0,19(1)	0,23(6)	0,186(6)	-
$\gamma_{9,3}$	670,565(12)	7,37(26)	7,28(6)	7,53(9)	7,16(20)	7,32(22)	7,60(14)
$\gamma_{7,2}$	691,304(12)	1,871(69)	-	1,87(4)	1,86(9)	1,79(6)	1,84(5)
$\gamma_{4,1}$	705,09(7)	-	-	-	0,011(1)	0,025(15)	-
$\gamma_{16,8}$	711,680(8)	74,48(258)	72,37(39)	75,7(8)	75,33(177)	73,8(32)	76,4(14)
$\gamma_{13,5}$	736,70(7)	0,506(26)	-	0,51(2)	0,50(4)	0,530(18)	0,547(23)
$\gamma_{17,8}$	752,332(10)	16,57(56)	16,26(12)	17,0(2)	17,08(43)	16,5(5)	16,98(33)
$\gamma_{8,1}$	778,862(12)	4,17(15)	4,00(3)	4,25(6)	4,22(14)	4,13(13)	4,27(8)
$\gamma_{4,0}$	785,81(7)	-	-	-	0,019(4)	0,023(3)	-
$\gamma_{8,2}$	810,325(10)	78,66(273)	76,94(44)	80,1(8)	79,31(177)	78,2(26)	80,3(12)
$\gamma_{10,3}$	830,601(15)	13,34(47)	12,99(10)	13,5(2)	13,51(35)	13,3(4)	13,62(26)
$\gamma_{7,1}$	875,63(5)	0,993(35)	-	0,99(4)	1,00(5)	0,987(31)	1,002(25)
$\gamma_{9,2}$	950,963(10)	3,71(14)	3,65(4)	3,89(6)	3,87(12)	3,74(12)	3,85(8)
$\gamma_{11,3}$	1010,27(6)	0,096(8) ¹⁾	-	0,11(1)	0,13(3) ¹⁾	0,107(4)	-
$\gamma_{14,3}$	1120,35(5)	0,327(15) ¹⁾	-	0,35(1) ¹⁾	0,28(5)	0,268(8)	-
$\gamma_{15,3}$	1146,81(9)	0,271(14)	-	0,30(1)	0,29(4)	0,279(9)	0,281(26)
$\gamma_{16,3}$	1241,52(2)	1,142(41)	-	1,21(4)	1,21(6)	1,118(34)	1,12(4)
$\gamma_{17,3}$	1282,06(6)	0,246(13)	-	0,29(1)	0,28(4)	0,240(11)	0,271(19)
$\gamma_{12,2}$	1306,60(15)	-	-	-	0,010(2)	0,0044(4)	-
$\gamma_{13,2}$	1331,04(13)	-	-	-	0,010(1)	0,0051(6)	-
$\gamma_{14,2}$	1400,79(2)	0,686(25)	-	0,74(2)	0,76(4)	0,672(21)	0,720(27)
$\gamma_{15,2}$	1427,24(2)	0,667(25)	-	0,72(2)	0,77(4) ¹⁾	0,673(22)	0,708(21)
-	1446,7(2)	-	-	-	<0,01	<0,0006	-

Gamma relative emission intensities, references 13 to 17 :

$g_{i,f}$	$E_g(\text{keV})$	13	14	15	16	17	18
$\gamma_{1,0}$	80,577(7)	17,00(22)	16,7(5)	17;6(4)	16,050(120)	17,18(15)	17,46(31)
$\gamma_{16,15}$	94,679(9)	0,208(10)	0,198(5)	0,23(3)	-	0,1977(50)	0,202(5)
$\gamma_{8,7}$	119,035(10)	-	0,236(7)	0,23(3)	-	0,2384(72)	0,238(4)
$\gamma_{16,14}$	121,175(10)	0,307(11)	0,326(9)	0,38(3)	-	0,343(9)	0,333(9)
$\gamma_{17,15}$	135,257(14)	-	0,1358(35)	0,15(3)	-	0,142(9)	0,1350(25)
$\gamma_{9,8}$	140,702(20)	-	0,0584(19)	0,07(1)	-	0,051(7)	0,059(3)
$\gamma_{10,9}$	160,077(20)	0,153(7)	0,139(3)	0,14(3)	-	0,140(11)	0,134(5)
$\gamma_{17,14}$	161,707(14)	-	0,160(5)	0,15(3)	-	0,1580(80)	0,151(5)
$\gamma_{3,2}$	184,404(7)	100	100	100	100	100	100
$\gamma_{16,13}$	190,747(16)	-	0,273(8) ¹⁾	0,31(3)	-	0,3010(62)	0,296(6)
$\gamma_{16,12}$	214,79(3)	-	0,671(17)	0,61(4)	-	0,600(10)	0,614(14)
$\gamma_{8,5}$	215,871(10)	3,594(37)	3,60(9)	3,49(14)	3,447(26)	3,566(85)	3,67(24)
$\gamma_{17,12}$	231,32(4)	0,283(6)	0,260(7)	0,30(4)	-	0,2933(55)	0,302(8)
$\gamma_{9,7}$	259,70(3)	1,529(34)	1,507(34)	1,45(5)	1,434(25)	1,480(12)	1,487(9)
$\gamma_{9,2}$	280,468(7)	41,41(51)	41,8(9)	39,8(9)	40,634(167)	40,66(29)	40,75(21)
$\gamma_{10,9}$	300,731(9)	5,339(58)	5,29(12)	4,98(13)	5,079(39)	5,118(36)	5,15(4)
$\gamma_{9,6}$	305,03(5)	-	0,020(10)	0,023(3)	-	0,026(6)	0,0252(16)
$\gamma_{11,9}$	339,75(5)	-	0,221(6)	0,22(3)	-	0,2250(36)	0,2229(27)
$\gamma_{6,3}$	365,736(9)	3,589(45)	3,51(9)	3,34(9)	3,439(47)	3,404(24)	3,39(4)
$\gamma_{16,10}$	410,950(8)	16,49(19)	16,02(36)	15,0(4)	15,424(74)	15,81(11)	15,65(22)
$\gamma_{17,10}$	451,528(9)	4,235(60)	4,11(10)	3,89(13)	4,023(30)	4,062(42)	4,02(5)
$\gamma_{10,6}$	464,819(12)	1,729(35)	1,73(4)	1,66(7)	2,027(31)	1,665(19)	1,73(6)
$\gamma_{15,9}$	476,38(6)	-	0,0494(26)	0,052(7)	-	-	0,0500(18)
$\gamma_{12,8}$	496,86(4)	-	0,175(4)	0,17(3)	-	0,174(16)	0,173(4)
$\gamma_{4,2}$	520,85(5)	-	0,276(14) ¹⁾	0,21(3)	-	0,212(13)	0,211(8)
$\gamma_{8,3}$	529,811(10)	13,19(15)	-	12,6(4)	13,380(126)	13,33(10)	13,0(6)
$\gamma_{16,9}$	570,940(10)	7,964(91)	-	7,27(23)	7,505(71)	7,71(6)	7,49(27)
$\gamma_{5,2}$	594,536(24)	0,761(22)	-	0,78(7)	-	0,880(20)	0,80(8)
$\gamma_{17,9}$	611,620(17)	2,097(26)	-	1,86(11)	1,952(60)	1,911(36)	1,81(29)
$\gamma_{11,7}$	615,84(9)	-	0,138(11)	0,044(13)	-	0,160(10)	0,13(4)
$\gamma_{13,7}$	639,97(9)	-	0,137(4)	0,11(2)	-	0,138(9)	0,130(4)
$\gamma_{11,6}$	644,689(15)	-	0,206(5)	0,21(4)	-	0,189(12)	0,198(5)
$\gamma_{9,3}$	670,565(12)	7,718(84)	-	6,98(22)	7,618(45)	7,56(6)	7,36(28)
$\gamma_{7,2}$	691,304(12)	1,872(40)	-	1,78(9)	1,914(17)	1,862(21)	1,82(10)
$\gamma_{4,1}$	705,09(7)	-	0,0272(7)	0,011(2)	-	-	0,019(9)
$\gamma_{16,8}$	711,680(8)	77,51(62)	-	72,0(19)	76,30(35)	76,3(6)	75,7(16)
$\gamma_{13,5}$	736,70(7)	0,510(12)	-	0,49(4)	-	0,524(16)	0,514(7)
$\gamma_{17,8}$	752,332(10)	17,16(14)	-	16,2(5)	16,973(84)	16,98(12)	16,8(4)
$\gamma_{8,1}$	778,862(12)	4,279(56)	-	4,04(14)	4,257(28)	4,242(33)	4,15(11)
$\gamma_{4,0}$	785,81(7)	-	0,0312(11)	0,019(4)	-	-	0,026(5)
$\gamma_{8,2}$	810,325(10)	80,81(59)	-	76,1(20)	80,52(38)	80,3(6)	79,1(14)
$\gamma_{10,3}$	830,601(15)	13,87(18)	-	12,9(4)	13,639(79)	13,64(10)	13,41(23)
$\gamma_{7,1}$	875,63(5)	1,003(21)	-	0,97(6)	-	0,501(9) ¹⁾	0,994(11)
$\gamma_{9,2}$	950,963(10)	3,898(48)	-	3,68(12)	3,789(25)	3,793(30)	3,785(21)
$\gamma_{11,3}$	1010,27(6)	-	0,1113(28)	0,11(2)	-	0,107(6)	0,1095(21)
$\gamma_{14,3}$	1120,35(5)	-	0,281(8)	0,28(3)	-	0,278(10)	0,275(5)
$\gamma_{15,3}$	1146,81(9)	0,290(6)	0,289(8)	0,27(3)	-	0,279(6)	0,284(3)
$\gamma_{16,3}$	1241,52(2)	1,211(10)	-	1,14(5)	-	1,121(14)	1,17(4)
$\gamma_{17,3}$	1282,06(6)	0,268(12)	0,263(7)	0,27(3)	-	0,2434(30)	0,252(9)
$\gamma_{12,2}$	1306,60(15)	-	0,00610(3)	0,010(2)	-	-	0,0076(15)
$\gamma_{13,2}$	1331,04(13)	-	0,0025(10)	0,010(2)	-	-	0,0059(16)
$\gamma_{14,2}$	1400,79(2)	0,707(17)	-	0,70(3)	-	0,689(7)	0,700(7)
$\gamma_{15,2}$	1427,24(2)	0,705(28)	-	0,68(3)	-	0,696(12)	0,687(7)
$\gamma_{15,2}$	1427,24(2)	-	-	-	<0,01	-	-

¹⁾Outlier

²⁾214, 8 + 215, 8 keV doublet

Upper limits for a possible 1446,7 keV transition have been determined by authors 10, 11, 16.

1	Burson et al. 1967
2	Gunther and Parsignault 1967
3	Reich and Cline 1970
4	Lavi 1973
5	Lingeman et al. 1974
6	Gehrke et al. 1977
7	Sampson 1978
8	Blagojevic and Wood 1982
9	Sooch et al. 1982
10	Ogandaga et al. 1986
11	Adam et al. 1988 (give also values for six additional very weak transitions)
12	Danilenko et al. 1989
13	Wang Xin Lin 1992
14	Wagner 1992 (gives additionally four weak transitions)
15	Ardisson 1992
16	Miyahara et al. 1994
17	Morel et al. 1996
18	Adopted value

The final values of Hino et al. (2000) were not available when this evaluation was carried out. The absolute emission probabilities (Table 4.2) are calculated by multiplying the relative values by the normalization factor $f_N = 0,725$ (3). The transition probabilities (Table 2.2) are calculated by multiplying the emission probabilities by $(1 + \alpha_t)$.

5 Main Production Mode

Taken from Firestone (1996).

6 References

References are given only in those cases where the reference is not already included in the list of references in the Tables Part.

- C. W. Reich and J. E. Cline, *Phys. Rev.* 137 (1965) B1424 [δ]
 J. M. Domingos, G. D. Symons and A. C. Douglas, *Nucl. Phys.* A180 (1972) 600 [δ]
 T. Miyokawa, I. Katayama, S. Morinobu and H. Ikegami, *Int. Conf. on Nuclear Moments and Nuclear Structure*, Osaka, Japan, 1972, edited by H. Horie and K. Sugimoto, *J. Phys. Soc. Jpn. Suppl.* 34 (1972) 247 [δ]
 K. R. Baker, J. H. Hamilton, J. Lange, A. V. Ramayya, L. Varnell, V. Maruhn-Rezwani, J. J. Pinajian and A. Maruhn, *Phys. Lett.* 57B (1975) 441 [δ]
 R. L. West, E. G. Funk, A. Visvanathan, J. P. Adams and J. W. Mihelich, *Nucl. Phys.* A270 (1976) 300 [δ]
 F. K. McGowan, W. T. Milner, R. L. Robinson, P. H. Stelson and Z. W. Grabowski, *Nucl. Phys.* A297 (1978) 51 [δ]
 J. Lange, K. R. Baker, J. H. Hamilton, A. V. Ramayya, L. Varnell, J. J. Pinajian, V. Maruhn-Rezwani, *Z. Phys.* A303 (1981) 31 [δ]
 K. S. Krane and J. D. Moses, *Phys. Rev.* C24 (1981) 654 [δ]
 A. Alzner, E. Bodenstedt, B. Gemünden, J. van den Hoff and H. Reif, *Z. Phys.* A322 (1985) 467 [δ]
 W. Wagner, *Bull. Russ. Ac. Sci.* 56 (1992) 675 [δ]

Comments on evaluation of decay data
by M.M. Bé, B.Q. Lee, T. Kibédi, A.E. Stuchbery

This evaluation was completed in June 2015 with a literature cut-off at the same date.

Decay Scheme

¹⁶⁹Er decays by beta minus emission to the first excited level and the ground state of ¹⁶⁹Tm and with a very small branch to the second excited level at 118 keV.

Level energies, spins and parities (J^π) were taken from the evaluation by Baglin (2008Ba31).

Nuclear Data

The Q^- value is from Wang *et al.* (2012Wa38).

Half-life

The recommended value was derived from measurements listed in Table 1.

Table 1. Measurement results and recommended value of ¹⁶⁹Er half-life.

Reference	$T_{1/2}$ (d)	Uncertainty (d)	Remarks
1948Ke11	9,4	0,2	
1956Bi30	9,0	0,2	
1958Pa16	9,5		Omitted, no uncertainty
1960Wi10	9,8	0,5	
1961Bj02	9,6	0,1	
1963Ra15	9,0	0,1	
1977My02	9,40	0,02	
2004Sc04	9,36	0,04	
$\chi^2/n-1 =$	4		
χ^2 crit. =	2,8		
UWM =	9,36		
LWM =	9,370	0,048	
Adopted	9,38	0,02	Simple mean of 1977My02 and 2004Sc04, with lowest uncertainty

The set of the seven values is discrepant with a $\chi^2/n-1$ of 4, the *Limitation of Relative Statistical Weight Method* recommends the increase of the uncertainty given in 1977My02 to 0,034 in order to limit its weight to 50%, then the weighted mean is 9,37 with the external uncertainty of 0,05.

However, the earliest values range from 9 to 9,8 d, where the two latest ones are in good agreement and more precise, thus the simple mean of 1977My02 and 2004Sc04 is adopted with the lowest experimental uncertainty, i.e. **9,38 (2) d**.

Characteristics of Electron Emissions

Two experimental results exist for the intensity of the β transition to the 8,4 keV level: 42 (1) % (1959Ch31) and 45 (5) % (1965Du02). The adopted value is the simple mean with the uncertainty taken from 1965Du02 (the uncertainty given by 1959Ch31 includes only the statistical part). Then $I_{\beta} = 43,5$ (50) % rounded to 44 (5) %. Consequently, the β transition intensity to the ground state is 56 (5) %.

The intensity of the first forbidden unique β transition to the 118 keV level was derived from the intensities of the 109,7 and 118 keV γ rays and the conversion coefficients (see following sections), that is : $P\beta \sim 0,016$ %.

The $\lg ft$ values were calculated using the Logft program and the mean energy using the BetaShape program (2014Mo20 and 2012Mo38).

Conversion electron energies and intensities were derived from the decay scheme parameters.

The full energy spectrum of the Auger electrons has been calculated using the procedures and programs described in Lee *et al.* (2012Le09). Based on the adopted values of the transition energies, absolute gamma-ray intensities, multiplicities and mixing ratios (see below), an ENSDF input file was created. As the ¹⁶⁹Er decays by β^- decay, the main source of the primary atomic vacancies is internal conversion, dominantly of the 8,4102 keV transition. The BrIcc program (2008Ki07) was used to calculate all relevant sub-shell conversion coefficients. For the main part of the calculations, the BrIccEmis code (2012Le09) was used. It evaluates the primary vacancy distribution, then it propagates the atomic vacancies using a full Monte Carlo approach. Atomic transition energies were calculated with the RAINE code (2002Ba85), taking into account the presence of all atomic vacancies during the recombination process. Atomic transition rates have been taken from the Evaluated Atomic Data Library, Perkins *et al.* (1991PeZZ). The Auger (and X-ray) data presented here have been obtained from a calculation based on 1 million radioactive decays.

Detailed Auger and Coster-Kronig electrons in the M sub shells.

M1-Shell					
M1 vacancies = 20,4					
	Auger M1-XY	CK M1M2X	CK M1M3X	CK M1M4X	CK M1M5X
(keV)	1,787	0,0374	0,112	0,482	0,528
%	2,1	5,1	9,97	1,34	1,89

M2-Shell				
M2 vacancies = 12,34				
	Auger M2-XY	CK M2M3X	CK M2M4X	CK M2M5X
(keV)	1,468	0,026	0,265	0,311
%	2,5	0,9	7,1	1,36

M3-Shell			
M3 vacancies = 18,2			
	Auger M3-XY	CK M3M4X	CK M4M5X
(keV)	1,263	0,190	0,324
%	5,41	1,54	11,1

Comments on evaluation

	M4-Shell		M5-Shell	
	M4 vacancies = 10,1		M5 vacancies = 14,4	
	Auger M4-XY	CK M4M5X	Auger M5-XY	
(keV)	0,89 – 0,95		0,85	
%	9,88	0,08	14,4	

Summary of the Auger electron emissions

Auger transition	Mean energy (keV)	Electrons per 100 decays
L _{total}	5,293	0,0103
MMX	0,274	40,3
MYX	1,211	34,3
M _{total}	0,704	74,6
NNN	0,092	75,8
NNX	0,092	34,4
NXY	0,225	0,59
N _{total}	0,099	116,1
OOX	0,0095	3,86
OXY	0,030	8,75
O _{total}	0,024	12,6
Total	0,317	203,3

Gamma-ray transitions

The 8,4 keV γ transition is strongly converted, several studies of the conversion in the sub-shells lead to attribute it a M1+E2 character. The measured ICC ratios and total ICC values are summarized below:

Reference	Shell	Icc(Unc)	Type
1958Sh64	M2/M1	0,42 (6)	Ratio
1958Sh64	M3/M1	0,48 (7)	Ratio
1958Sh64	N23/M1	0,25 (3)	Ratio, excluded
1958Sh64	M45/M1	0,043 (11)	Ratio
1958Sh64	O123/M1	0,029 (10)	Ratio, excluded
1959Ch31	MN	106 (6)	ICC, Excluded, discrepant > 20 σ
1964Ka02	T	325 (35)	Total ICC
1965Du02	M2/M1	0,33 (6)	Ratio
1965Du02	M3/M1	0,39 (6)	Ratio
1967Ma42	M2/M1	0,31 (2)	Ratio
1967Ma42	M3/M1	0,34 (2)	Ratio
1967Ma42	N2/N1	0,25 (2)	Ratio
1967Ma42	N3/N1	0,41 (4)	Ratio
1967Ma42	M/N	4,2 (3)	Ratio
1967Ma42	N/O	6,4 (6)	Ratio
1968Ca06	M2/M1	0,368 (14)	Ratio
1968Ca06	M3/M1	0,380 (14)	Ratio
1968Ca06	M4/M1	0,009 (2)	Ratio
1968Ca06	M5/M1	0,010 (2)	Ratio
1968Ca06	N1/M1	0,221 (6)	Ratio
1968Ca06	N2/M1	0,064 (3)	Ratio

1968Ca06	N3/M1	0,076 (2)	Ratio
1968Ca06	N45/M1	0,0020 (2)	Ratio, outlier
1968Ca06	O1/M1	0,037 (1)	Ratio
1968Ca06	O23/M1	0,0208 (6)	Ratio, outlier
1968Wa05	T	220 (50)	Total ICC
2001KoZQ	O2/O1	0,30 (4)	Ratio
2001KoZQ	O3/O1	0,24 (4)	Ratio
2001KoZQ	P1/O1	0,054 (12)	Ratio
2015In**	M2/M1	0,34 (4)	Ratio
2015In**	N3/N1	0,40 (6)	Ratio
Reduced χ^2		2,7	
Mixing Ratio		0,0306 (16)	Deduced from parabolic search

The adopted mixing ratio of 0,0306 (16) was deduced from the experimental values listed above by the program BrIccMixing. This program uses two procedures to determine the best parameter values and uncertainties:

- (a) Chi-squared fit, based on the CFIT [1980Ry04] program;
- (b) Searching the $\chi^2(\text{ICC}_{\text{Theor}}, \text{ICC}_{\text{Exp}})$ hypersurface.

The total conversion coefficient is then $\alpha_T = 251 (10)$. This value is consistent with the experimental ICC of 220 (50) determined by Wagner (1968Wa05), but not with the value of 325 (35) obtained by Kankeleit *et al*, (1964Ka02) or 106 (6) by Charpak and Suzor (1959Ch31), it is also not consistent with the value of 148 (26) deduced from the Yb-169 decay scheme. It is noteworthy that this latter value was derived from a series of measurements of the absolute γ ray intensities, including the 8,7 keV ray, carried out by several laboratories, thus it should be considered as confident.

The 109,7 keV γ transition is a mixture of M1 and E2, the adopted mixing ratio is 2,17 (4) % as determined in the ¹⁶⁹Yb evaluation (2004BeZQ).

With the conversion coefficients given by the BrIcc program, the transition probabilities are:

$$P_{\gamma_{109}} = 0,0152 (3) \%$$

$$P_{\gamma_{118}} \sim 0,0013 \%$$

Gamma-ray emissions

From the β intensity and the α_T coefficient as determined above, the intensity of the 8,4 keV γ ray is 0,174 (21) %,

Bisi (1956Bi30) reported that if a γ -ray of 110 keV occurs it would have an intensity lower than 10^{-4} per β decay. Later, Sharma *et al*, (1970Sh09) measured the gamma spectrum in the decay of ¹⁶⁹Er, observed two rays with energy 109,7 and 118,2 keV respectively, but gave only the intensity of the sum, *i.e.*, 0,005 (1) %, Looking at the spectrum shown in 1970Sh09, the intensity of the 109,7 keV ray seems to be about four times greater than the intensity of the 118,2 keV ray. But in the decay of ¹⁶⁹Yb, which has been extensively studied (2002Be05), the ratio $\gamma_{109}/\gamma_{118}$ is 9,28 (7); this leads to $I_{\gamma_{109}} = 0,0045 (9) \%$ and $I_{\gamma_{118}} \sim 0,0005 \%$,

The energy spectrum of the X-rays has been evaluated using the same procedure as for the Auger electrons. The mean energy and intensity of each major X-ray group is for 100 β decays.

X-ray transition	Mean energy (keV)	X rays (%)
M _{total}	1,496	0,529
N _{total}	0,208	0,043
O _{total}	0,037	1,44
Total	0,674	2,03

References

- 1948Ke11** B.H.Ketelle, W.C.Peacock. Phys. Rev. 73,10 (1948) 1269 (Minutes of the meeting at Chicago, Dec. 29-31, 1947). Half-life
- 1956Bi30** A.Bisi, S.Terrani, L.Zappa. Nuovo Cimento 4 (1956) 758. Half-life
- 1958Pa16** F.I.Pavlotskaia, A.K.Lavrukhina. Soviet Phys. JETP 7 (1958) 732. Half-life
- 1958Sh64** K.N.Shliagin, P.S.Samoilov. Soviet Phys. JETP 7 (1958) 20. Mixing ratio
- 1959Ch31** G.Charpak, F.Suzor. J. Phys. Radium 20 (1959) 513. Beta emission intensities, M ICC, N ICC
- 1960Wi10** R.G.Wille, R.W.Fink. Phys. Rev. 118 (1960) 242. Half-life
- 1961Bj02** S.Bjornholm, H.L.Nielsen, O.B.Nielsen, G.Sidenius, O.Skilbreid, A.Svanheden. J. Inorg. Nucl. Chem. 21 (1961) 193. Half-life
- 1962Gr16** Z. Grabowski, J.E Thun, B.Lindstrom. Z. Phys. 169 (1962) 303. K ICC, M ICC, Gamma-ray energies, Gamma-ray emission intensities
- 1963Mc13** R.E.McAdams, G.W.Eakins, E.N.Hatch. Phys. Lett. 6 (1963) 219. Half-life isomeric level
- 1963Ra15** G.V.S.Rayudu, L.Yaffe. Can. J. Chem. 41 (1963) 2544. Half-life
- 1964Ka02** E.Kankeleit, F.Boehm, R.Hager. Phys. Rev. 134B (1964) 747. T ICC
- 1965Du02** J.-C.Duperrin, A.Gizon-Juillard. C. R. Acad. Sci. (Paris) 261B (1965) 98. Beta emission intensities
- 1967Ma42** M.I.Marques, M.T.Ramos. C. R. Acad. Sci. (Paris) 265B (1967) 1209. M/N, N/O
- 1968Wa05** F.E.Wagner. Z. Physik 210 (1968) 361. ICC
- 1968Ca06** T.A.Carlson, P.Erman, K.Fransson. Nucl. Phys. A111 (1968) 371. Gamma-ray energies
- 1970Sh09** R.P.Sharma, H.L.Nielsen, P.G.Hansen. Nucl. Phys. A152 (1970) 225. Beta emission intensities
- 1971MaXG** M.I.Macias-Marques. Thesis, Univ. Orsay, FRNC-TH136 (1971) 36. M/N, N/O
- 1972Mc05** E.J.McGuire. Phys. Rev. A5 (1972) 1043. Auger electron emission intensities
- 1977My02** W.A.Myers. J. Inorg. Nucl. Chem. 39 (1977) 925. Half-life
- 1980Ry04** M. Ryšavý and O. Dragoun. Computer Phys. Comm. 19 (1980) 93
- 1991PeZZ** S.T.Perkins, D.E.Cullen, M.H.Chen, J.H.Hubbell, J.Rathkopf, J.Scofield, UCRL-50400 vol.30 (1991) Atomic Data
- 2001KoZQ** A.Kovalik, E.A.Yakushev, A.F.Novgorodov, V.M.Gorozhankin, M.Mahmoud Program and Thesis, Proc.51st Ann.Conf.Nucl.Spectrosc.Struct.At.Nuclei, Sarov (2001) 177. Mixing ratio
- 2002Ba85** I.M.Band, M.B.Trzhaskovskaya, C.W.Nestor Jr., P.O.Tikkanen, S.Raman. At. Data. Nucl. Data Tables 81 (2002) 1. Theoretical ICC
- 2002Be05** M.M.Bé, E.Schönfeld, J.Morel. Appl. Radiat. Isotopes 56 (2002) 181. Gamma-ray emission intensity ratio
- 2004Sc04** H.Schrader. Appl. Radiat. Isotopes 60 (2004) 317. Half-life
- 2004BeZQ** M.M.Bé, V.Chisté, C.Dulieu, E.Browne, V.Chechev, N.Kuzmenko, R.Helmer, A.Nichols, E.Schönfeld, R.Dersch. Monographie BIPM-5, Vol.2, Bureau International des Poids et Mesures (2004) Mixing ratio
- 2008Ba31** C.M.Baglin. Nucl. Data Sheets 109 (2008) 2033. Spin and Parity
- 2008Ki07** T.Kibédi, T.W.Burrows, M.B.Trzhaskovskaya, P.M.Davidson, C.W.Nestor Jr. Nucl. Instrum. Methods Phys. Res. A589 (2008) 202. Theoretical ICC

Comments on evaluation

- 2012Le09** B.Q.Lee, T.Kibédi, A.E.Stuchbery, K.A.Robertson, *Comput.Math.Meth.Med.* 2012, 651475 (2012) Auger electron intensities
- 2012Mo38** X.Mougeot, M.-M.Bé, C.Bisch, M.Loidl. *Phys. Rev. A* 86 (2012) 042506. Mean Beta energies
- 2012Wa38** M.Wang, G.Audi, A.H.Wapstra, F.G.Kondev, M.MacCormick, X.Xu, B.Pfeiffer. *Chin. Phys. C* 36 (2012) 1603. Q
- 2014Mo20** X.Mougeot, M.-M.Bé, C.Bisch, M.Loidl. *Nucl. Data Sheets* 120 (2014) 129. Mean Beta energies
- 2015In**** A.Kh.Inoyatov, A. Kovalik, D.V. Filosofov, M. Rysavy, L.L.Perevoshchikov and Yu.B. Gurov. *Eur. Phys. J. A* 51, 6 (2015) 65. Mixing Ratio

¹⁶⁹Yb - Comments on evaluation of decay data by M. M. Bé and E. Schönfeld

1. Decay Scheme

The decay scheme tries to be complete : the confirmed gamma rays (even the weakest, are placed), the questionable gamma transitions are mentioned but not placed.

The J^π values and the level half-lives are taken from NDS 64,2 (1991).

2. Nuclear Data

- To determine the half-life of ¹⁶⁹Yb the following values have been taken in account ($T_{1/2}$ in d):

1	31,83(21)	Walker 1949 (49Wa23)
2	31,97(5)	Lagoutine et al. 1975 (75La16)
3	32,022(8)	Houtermans et al. 1980 (80Ho17)
4	32,015(9)	Rutledge et al. 1980 (80RuZY)
5	32,032(20)	Funck et al. 1983 (83Fu12)
6	32,07(8)	Kits et al. 1988 (88Ki12)
7	31,88(12)	Parker 1990 (90Pa08)
8	32,0147(93)	Unterweger et al. 1992 (92Un01)
9	32,001(34)	Iwahara et al. 1999
10	32,018(5)	weighted mean, adopted value

Value 1 was measured with a Geiger counter, value 2 with a proportional counter, value 7 with a Ge(Li) detector. For all the other measurements an ionisation chamber was used.

This set is a consistent one with a reduced $-\chi^2$ of 0,59. The largest weights are those of values 3 (36 %), 4 (28 %) and 8 (27 %).

Several others values with greater or without uncertainty can be found: 33,0(15) d (Bothe 1946); 32,4 d (Cork 1954), 33,0(15) d (Don Martin 1951), 32 d (Michel, 1954), 30,6(2) d (Cork 1956).

- The Q value is from Audi and Wapstra (1995).

2.1 Electron Capture Transitions

The probabilities and uncertainties are deduced from the gamma transition probability balance on each level.

The balance on level 13 (570 keV) introduces the possible existence of a second forbidden transition to populate this level. This solution is preferred to those of a possible gamma transition from level 19 (878 keV) with energy 307,5 keV, this gamma line being not mentioned in any publication. The existence of gamma rays from levels 14, 16, 17 has not been pointed out in any process.

From spin and parity it follows that a transition to the ground state ($\Delta J^\pi = 3^+$) would be unique second forbidden and an EC transition to the 8,4 keV ($\Delta J^\pi = 2^+$) level would be non-unique second forbidden. If these transitions exist, the limits of their probabilities, which are based on $\lg ft$ systematics, are 0,001% and 0,1% respectively.

EC transitions to the 118 keV ($J^\pi = 5/2^+$) and 139 keV ($J^\pi = 7/2^+$) levels of the rotational band ($K^\pi = 1/2^+$) could also be possible and would both be allowed. Nevertheless the projection of the angular momentum J on the rotational symmetry axis K, is $1/2$, this involves a transfer of 3 units of angular momentum rather than the 0 or 1 unit indicated by the J value. Due to the fact that this nucleus is a deformed nucleus and from $\lg ft > 9$, it results that the intensities of the EC transitions, if exist, are very low.

In the proposed decay scheme the sum of the electron capture transition probabilities is 100,0
(19)

From experimental emission probabilities and balancing conditions, and taking into account the uncertainties of the gamma transitions feeding and leaving these levels, it seems not necessary to introduce the EC transitions mentioned to the 118 keV and 139 keV levels.

The fractional capture probabilities given in section 2.1 have been calculated on the basis of the table of Schönfeld (1998) and the Q value of Audi and Wapstra (1995). Sahota *et al.* (1982) have determined experimental values of P_K with a relative uncertainty of 3 to 5 % [$P_K(472) = 0,812(29)$; $P_K(379) = 0,823(34)$; $P_K(316) = 0,825(43)$]; their values agree within the uncertainties with the more accurate theoretical values.

The $\lg ft$ values were calculated from the half-life, the evaluated EC transition probabilities and the transition energies using the $\log-f$ tables for beta decay of Gove and Martin (1971).

2.2 Gamma Transitions

Precise γ -ray energies of the main γ -rays have been determined by Borchert *et al.* 1975 and Kessler *et al.* 1979. The values of nine lines (i. e., 63, 93, 109, 118, 130, 177, 197, 261, and 307) given in the table in Section 4.2 are taken from Helmer (2000He14). They are based on a value of 411,80205(17) keV for the 412 keV line following the ¹⁹⁸Au decay. The energies of the weaker γ -rays are taken from Vagner (1990). The remaining energies (316, 328, 425, 614 keV) were computed from these energies and the relationships in the decay scheme. In order to calculate the level differences which are given in section 2.2 the recoil energies have been taken in account. The γ -ray energies can be found in section 4.2.

The transition probabilities $P_{\gamma+ce}$ were calculated from the measured relative γ -ray emission probabilities (see section 4.2), the total conversion coefficients and from the absolute intensity value of the 198 keV line 35,93(12) which was derived from statistical treatment of measured values (see section 4.2).

The conversion coefficients were interpolated from the table of Rösel *et al.* 1978. Mixing ratios are taken from angular correlation measurements and from $L_1/L_2/L_3$ ratios respectively $M_1/M_2/M_3/M_4/M_5$ ratios (Günther *et al.* 1969, Agnihotry *et al.* 1972, Krane *et al.* 1972, Akhmetov *et al.* 1985, Davaa *et al.* 1987, Kracikova *et al.* 1987, Wagner *et al.* 1990). The mixing ratios were derived by comparing the subshell ratios from theory and experiment.

The uncertainties of the conversion coefficients are assumed to be 1,5 % for the three well studied transitions 2,1; 4,3; 4,2; 10 % for the less accurate measured transitions 6,3; 7,3; 7,4 and those above 330 keV, and 3 % for all other transitions.

Recently Dey *et al.* (1997) found from angular correlation measurements evidence for a pure M1 character of the 94 keV transition, almost pure E2 character for the 198 keV transition and only 4 % E2 admixture in the 177 keV transition. The corresponding change in $\alpha_i(94)$ from 3,89 to 3,88 is negligible, the change in $\alpha_i(177)$ from 0,59 to 0,62 is small, but $\alpha_i(198)$ would become markedly lower and lead to disagreement when determining the normalisation factor from different cuts through the decay scheme. Also, considering the recent measurements carried out by Baratova *et al.* (1993) who found a E2 admixture of : 3,4 % in the 94 keV; 16 % in the 177 keV and, 11 % in the 198 keV transition these results being in agreement with the other experiments; the values of Dey *et al.* (1997) were not used for the present evaluation.

Comparison between measured α_k and theoretical value from Rösel and from new tables of Band *et al.*(1993) for some important lines which are M1+E2 or E2 :

Eg	93,6	109,8	130	177,2	198	307,8
Adopted admixture %E2	3,25 (25)	2,17 (4)	100	15,8 (3)	9,0 (6)	100
Grabowski (1962)	3,3 (3)	2,15 (20)		0,52 (4)	0,41 (3)	0,048 (5)
Agnihotry (1972)				0,445 (35)	0,30 (2)	0,049
Zheltonozhsky (1995)		2,04 (2)	0,545 (5)	0,515 (5)	0,388 (4)	
α_K theoretical Rösler	3,18 (10)	2,03 (3)	0,538 (17)	0,484 (7)	0,370 (6)	0,0482 (15)
α_K theoretical Band	3,06 (10)	1,95 (3)	0,529 (16)	0,467 (6)	0,358 (5)	0,0477 (14)

3. Atomic data

- The values of ω_K , ω_L , n_{KL} are taken from Schönfeld and Janßen 1996.
- The energies of the X rays are based on the wavelengths given by Bearden (1967).

4.1 X-ray emissions

The emission intensities of the L- and K- X-rays are calculated with the EMISSION program (version 102) from the data set evaluated in this study : electron capture transition probabilities, gamma emission probabilities and from the internal conversion coefficients (α_K , α_{L1} , α_{L2} , α_{L3}) from Rösler *et al.* and the partial capture coefficients P_K , P_L taken from the PTB EC-CAPTURE program with the ratio $P_{L2} / P_{L1} = 0,0527$.

These values are compared with experimental values (see table enclosed), they are generally in good agreement within the uncertainty limits. The measurements were performed with a Si-Li detector for Reference 1-E, an HP-Ge for References 7-E, 10-E1, 10-E2 and 3, a Si-Li and HP-Ge for References 1 and 2 and a low energy photon spectrometer for Reference 4.

4.2 Gamma Emissions

The gamma emission probabilities taken in consideration are from the EUROMET exercise 410 (Morel *et al.*) and from several other authors.

List of laboratories which took part in the EUROMET exercise (all details can be found in the report- 1999MoZV) :

- Institute for Physics and Nuclear Engineering (Romania)
- Institut de Radiophysique Appliquée (Switzerland)
- Institute for Reference Materials and Measurements (Belgium)
- V.G. Khlopin Radium Institute (Russia)
- Laboratorio Nacional de Metrologia das Radiações Ionizantes (Brazil- Iwahara *et al.*)
- Laboratoire Primaire des Rayonnements Ionisants (France)
- National Physical Laboratory (U.K.)
- National Office of Measures (Hungary)
- Radioisotope Centre POLATOM (Poland)
- Physikalisch-Technische Bundesanstalt (Germany – Schönfeld *et al.*)
- D.I.Mendeleyev Institute for Metrology (Russia – Sazonova *et al.*)

An arbitrary code number was assigned to each participant. The same code number is used here to reference the results.

The recent references : Schönfeld *et al.* (1999), Sazonova *et al.*(2000), Iwahara *et al.* (2000) have not been included as independent reference because they were participants in the EUROMET exercise and then, their results are *de facto* included.

In the EUROMET exercise 410, references 1-E to 11-E, the values were given in absolute value, they have been converted relatively to the 198 keV line.

The other references used are :

1: Artomonova *et al.* 1976 (below 308 keV) and Balalaev *et al.* 1972 (above 308 keV), in this reference the values are given relatively to the 307 keV gamma-ray. As described, from V.S Aleksandrov the absolute intensity for this ray was taken as 10,1(5) % and those of the 198 keV gamma-ray is 34,34 (264). For this study the values given by Balalaev were converted relatively to the 198 keV ray taken as 100, with respect to the above absolute values used in the quoted paper.

2: Gehrke *et al.* 1977

3: Funck *et al.* 1983 (below 308 keV), Georgieva and Tumbev 1976 (above 308 keV)

4: Mehta *et al.* 1986 (uncertainties above 130 keV multiplied by a factor 2 to be compatible with the results of other authors)

5: Vagner *et al.* 1990, this work is supposed to be the continuation of the work of I. Adam, V. Vagner *et al.* (1986).

6: Bhattacharya *et al.* 1996

7: Miyahara 1998

The less accurate values of the following references were not taken into account for the present evaluation:

Alexander and Boehm 1963

Brown and Hatch 1967

Sen *et al.* 1972

Agnihotry *et al.* 1972

Potnis *et al.* 1972

Lavy *et al.* 1973

Aleksandrov *et al.* 1973

Verma *et al.* 1976

• Other remarks :

- The gamma given at the 205,99 energy by Vagner and at the 206,2 energy by Mehta are processed together in the same line.

- The intensity of the 51 keV is from the imbalance of level 7.

- Some weak gamma transitions were seen in only one spectrum :

105,2 ; 193,1 ; 213,9 ; 226,3 ; 291,2 ; 294,5 ; 316,2 ; 328,0 ; 356,7 ; 425,0 ; 500,3 ; 507,8 ; 546,1 ; 614,1 ; 633,3 ; 693,5 ; 710,3 ; 739,4 ; 760,2 and 781,6 lines.

The 616,2 and the 614,1 lines can not be placed in the decay scheme.

- Four EUROMET participants and Funck made the measurement of the resulting gamma emission of the 8,4 keV transition with the $L_{\beta 2}$ and $L_{\beta 15}$ X-rays emission. The LWEIGHT program running on these 5 values gives for this line $(\gamma_{8,4} + L_{\beta 2,15}) = 4,68(14)\%$

On the other hand, we obtain with the EMISSION program : $L_{\beta 2,15} = 3,93(10)\%$ for the X-ray emission.

The gamma emission absolute intensity can be deduced : $4,68 - 3,93 = 0,75(17)\%$

From the balance on the levels 1 and 0 of the decay scheme, a probability of 95,1 % for the 8,4 keV transition is deduced. As the decay scheme is quite consistent in every part, this value is certainly good.

The consequence is that the deduced ICC total is : 125(16)

This is not consistent with the theoretical ICC obtained from the Rosel table for a M1+0,108%E2 transition which is = 273(13)

It can be noted that with a pure M1 transition the Rosel ICC is 177(8)

The E2 admixture to the M1 multipolarity is deduced from the M1/M2/M3/M4/M5 ratio measured by T.A. Carlsson, *et al.* They compared their measured ratio with those from the Tables of Hager and Seltzer. Their calculations, taking the Rösler *et al.* conversion coefficients, were repeated and confirmed their result of 0,108(5) % E2 admixture. There are also some older less accurate values giving 0,10(2) %.

It also exists an old measurement of $\alpha_{MN} = 106(6)$ from G. Charpak and F. Suzor (1959).

Without other confirmation of this value, we will stay with the theoretical ICC for a M1+0,108%E2 transition calculated from Rösler *et al.*

This leads to the **adopted absolute value of 0,347(17)%** for the emission intensity.

This approach was also followed by Artomonova who gave a value of 0,33(4)% for the 8,4 keV gamma line emission intensity.

- Determination of the absolute emission intensity of the 198 keV line

During the EUROMET exercise the absolute activity measurement of Yb -169 sources was carried out by several methods and the absolute intensity of the 198 gamma -ray line deduced. This gives 8 measurements made by independent laboratories (references from 1-E to 11 -E), moreover 3 others absolute measurements are available (references 3, 7, 8). In these conditions a statistical treatment by using the program LWEIGHT has been done to determine the absolute emission intensity of the 198 keV line.

Absolute values of the 198 keV line from EUROMET exercise and others :

1-E	(36,26 ± 0,18)	EUROMET, 1999
3-E	(37,3 ± 0,5)	EUROMET, 1999
4-E	(35,7 ± 0,6)	EUROMET, 1999
7-E	(36,3 ± 1,1)	EUROMET, 1999
8-E	(35,9 ± 0,8)	EUROMET, 1999
9-E	(35,49 ± 0,39)	EUROMET, 1999
10-E1	(36,06 ± 0,15)	EUROMET, 1999
11-E	(35,9 ± 0,5)	EUROMET, 1999
3	(36,0 ± 0,5)	Funck et al. 1983
7	(35,14 ± 0,28)	Miyahara et al. 1999
8	(35,5 ± 0,4)	Coursey et al. 1994

The reference 3-E is rejected due to deviation from the weighted average (Chauvenet criteria), this leads to process 10 values. No value contributes more than 50%, the reduce $-\chi^2$ is 1,64 ; the weighted mean and external uncertainty is chosen. Then **the adopted value is 35,93(12)%**.

This value is quite close to those obtained by Schönfeld et al. (35,91(13)) by considering the balance of the decay scheme.

5. Electron Emissions

Auger Electrons

The energies of the KLL Auger electrons are taken from Larkins (1977), the others are calculated from the binding energies using approximations. The probabilities of L - and K-Auger electrons are calculated with the PTB program Emission (version 102).

Conversion Electron Emissions

The energies were calculated from the gamma transition energies and from the binding electron energies on the electronic shells.

The emission probabilities were calculated using the adopted gamma emission probabilities and conversion coefficients.

The comparison between measured internal conversion electron intensities and calculated values gives a good agreement which confirms the consistency of the evaluated data set.

E gamma	Agnihotry (1972)	Artamonova (1976)	Calculated
8,4 keV - Ie M		71 (7)	76 (5)
20,7 keV - Ie L		7,5 (4)	8,6 (3)
Ie M		1,7 (1)	1,93 (7)
63 keV - Ie K		36 (7)	39,6 (12)
- Ie L		7,16 (15)	7,2 (3)
93 keV - Ie K		7,5 (7)	8,18 (27)
- Ie L		1,5 (1)	1,4 (5)
109 keV - Ie K		34,9 (11)	35,2 (6)
- Ie L		5,7 (1)	5,68 (9)
118 keV - Ie K		1,28 (6)	1,30 (4)
- Ie L			1,37 (4)
130 keV - Ie K		6,2 (3)	6,1 (2)
- Ie L		5,4 (2)	5,3 (2)
177 keV - Ie K	10,1 (5)	10,7 (7)	10,8 (2)
- Ie L		2,1 (1)	1,94 (3)
198 keV - Ie K	10,8 (5)	13,5 (5)	13,29 (22)
- Ie L		2,16 (5)	2,17 (3)
240 keV - Ie K	0,0043 (4)	0,0045 (5)	0,0042 (5)
- Ie L		0,0010 (5)	0,00075 (8)
261 keV - Ie K	0,047 (7)	0,040 (4)	0,040 (1)
- Ie L			0,0060 (2)
307 keV - Ie K	0,53	0,50 (2)	0,484 (15)
- Ie L		0,15 (2)	0,142 (4)

6. Main Production Modes

From Firestone (1996) and Shirley (1991)

References of the programs used

LWEIGHT : A computer program to calculate averages, D. MacMahon, E. Browne

EC-CAPTURE : Calculation of electron capture probabilities. PTB

EMISSION-102 : Calculation of X-rays and Auger electrons emission probabilities. PTB

ICC Database : ICC computer code, CEA-BNM/LNHB technical note LPRI/98/002

References not used

Von W. Bothe. Z. Naturforschg. 1 (1946) 173

J. M. Cork *et al.* Phys. Rev. 78,2 (1950) 95

Don S. Martin *et al.* Phys. Rev. 82,5 (1951) 579

M. C. Michel *et al.* Phys. Rev. 93 (1954) 1422

J. M. Cork *et al.* Phys. Rev. 101,3 (1956) 1042

E. N. Hatch *et al.* Phys. Rev. 104,3 (1956) 745

Other quoted references can be found in the Tables Part.

**¹⁷⁰Tm - Comments on evaluation of decay data
by V. P. Chechev and N. K. Kuzmenko**

1. Decay Scheme

Since ¹⁷⁰Tm has spin and parity 1^- , it decays with detectable probability to the 0^+ ground states and 2^+ first excited levels in both ¹⁷⁰Yb and ¹⁷⁰Er. The only other levels below the decay energies are at 277 keV (4^+) and 573 keV (6^+) in ¹⁷⁰Yb and 260 keV in ¹⁷⁰Er. From the log ft systematics of 1998Si17, one expects the log ft 's of the 3^{rd} forbidden decays to the 4^+ levels to be greater than 16, which corresponds to a β branch of less than 0,000 002% to the 4^+ level of ¹⁷⁰Yb and weaker branch to 4^+ in ¹⁷⁰Er. Since the branch to the 6^+ level will be a 5^{th} forbidden decay, it will be even much weaker. Therefore, all of these unobserved β branches will be negligible.

For decay scheme see also Baglin (1996Ba01).

2. Nuclear Data

Q value is from Audi and Wapstra (1995Au04).

The ¹⁷⁰Tm half-life values are available, in days

125 (2)	1962Bo12	
134,2 (8)	1965F102	Omitted as outlier
128 (1)	1967Ke13	
128,6 (3)	1968Re04	
127,1 (3)	1969La34	(the original value of uncertainty is $3\sigma = 0,9$)
127,8 (6)	Average	

The outlier value of 1965F102 was omitted on the statistical considerations of its large deviation from the mean.

For statistical processing one third of the total 3σ -uncertainty, 0,9 days, stated in 1969La34, was used. Then, the weighted average is 127,8 d with an internal uncertainty of 0,21 d, a reduced- χ^2 of 4,85 and an external uncertainty of 0,45 d. In this case the different statistical procedures using the weighted average give the following values for a final uncertainty, in days: UINF - 0,45; PINF - 0,45; BAYS - 0,79; MBAYS - 0,56; LWM - 0,77; tS- 0,54. The LWEIGHT program using the LWM method has expanded the uncertainty to 0,77 d to include the accurate value of the 1968Re04. The EV1NEW program chooses the tS or MBAYS procedure for this case and gives 0,6 d. The latter value was adopted for the final uncertainty of the average.

It should be noted that without rejecting 1965F102 the Normalised Residuals technique leads almost to the same average of 127,9(6) days. It inflates the uncertainty of the 1965F102 value to 2,7 days and of each of the 1968Re04 and 1969La34 to 0,5 days.

A considerable discrepancy of few available experimental data on the ¹⁷⁰Tm half-life, all obtained before 1970, requires new ¹⁷⁰Tm half-life measurements.

2.1 β^- - Transitions

The β^- -decay probabilities have been computed from the $P_{\gamma+\text{ce}}(\text{Yb})$ of section 2.3 and balance correlations.

2.2. Electron Capture Transitions

The values of the electron capture probabilities to the ¹⁷⁰Er ground state and the level of 78,6 keV have been obtained from the balance correlations including the X K - and gamma emission probabilities. Indeed, we can write:

$$P_{XK}(Yb) = \omega_K(Yb) \alpha_K(84) P\gamma(84)$$

$$P_{XK}(Er) = \omega_K(Er) [P_K^{0,0} P(\epsilon_{0,0}) + P_K^{0,1} P(\epsilon_{0,1}) + \alpha_K(79) P\gamma(79)]$$

From here:

$$S \equiv \frac{P_{XK}(Er)}{P_{XK}(Yb)} = \frac{w_K(Er)}{w_K(Yb)} \cdot \frac{1}{a_K(84) \cdot P_g(84)} [P_K^{0,0} \cdot P(\epsilon_{0,0}) + P_K^{0,1} \cdot P(\epsilon_{0,1}) + a_K(79) \cdot P_g(79)]$$

Finally, for $P(\epsilon_{0,0})$ and $P(\epsilon_{0,1})$ the following expressions are obtained (see also 1988Kuzmenko):

$$P(\epsilon_{0,0}) = \frac{P_g(84)}{P_K^{0,0}} \left\{ a_K(84) \cdot S \cdot \frac{w_K(Yb)}{w_K(Er)} - \frac{P_g(79)}{P_g(84)} [a_K(79) + P_K^{0,1} (1 + a_T(79))] \right\}$$

$$P(\epsilon_{0,1}) = P_g(79) \cdot (1 + a_T(79))$$

In this calculation, the adopted values of ICC, P_K , ω_K , P_γ and the ratio of $S = 0,035(1)$ measured in 1986Ve05 were used.

The fractional electron capture probabilities to the specific atomic shells (P_K , P_L , P_M ...) have been deduced from the tables of Schönfeld (1998Sc28).

2.3. Gamma Transitions and Internal Conversion Coefficients

The energies of gamma transitions are the energies of gamma rays with the recoil energy added. The probabilities of gamma transitions $P_{\gamma+ce}$ have been computed using the gamma -ray emission probabilities and the total internal conversion coefficients (ICC).

The theoretical values of ICC from Rosel et al. (1978Ro21) have been adopted for the gamma transitions which have the same multipolarity E2. The evaluated α_{NO} values have been computed from $\alpha_M(\text{theoretical})$ using the ratio $\alpha_M / \alpha_{NO} = 3,77(9)$ (1968Ni06).

The weighted mean of the eight measurement results for $\alpha_K(\gamma84)$ [1,48(5) (1966Di02), 1,41(4) (1969Ne02), 1,37(4) (1970Mo07), 1,41(5) (1971Ca08), 1,46(7) (1973Pl08), 1,39(3) (1985Me18), 1,41(3) (1986Ve01), and 1,43(4) (1990Ke01)] is 1,41 with an internal uncertainty of 0,014 ; a reduced χ^2 of 0,6 and an external uncertainty of 0,011. Taking into account that a systematic error of the measurement method can contribute mainly to the measurement uncertainties, the smallest of the input uncertainties has been chosen as a final uncertainty of the weighted mean. The average value of $\alpha_K(\gamma84)$ (experimental), equal 1,41(3), agrees well with the theoretical value of 1,39(2). The relative uncertainty of the theoretical ICC has been adopted of 1,5%. This value of uncertainty provides overlapping $\alpha_K(\gamma84)$ (theoretical) and $\alpha_K(\gamma84)$ (experimental).

3. Atomic Data

The fluorescence yields are taken from 1996Sc06 (Schönfeld and Janßen). The X-ray energies are based on the wavelengths in the compilation of 1967Be65 (Bearden). The relative KX-ray emission $K\beta/K\alpha$, $K\alpha_2/K\alpha_1$, $K'\beta_2/K'\beta_1$ probabilities and the ratios $P(KLX)/P(KLL)$, $P(KLY)/P(KLL)$ are taken from 1996Sc06. The energies of Auger electrons are from 1977La19 (Larkins).

4. Photon Emissions

4.1. X-Ray Emissions

The absolute XK(Er), XK(Yb), XL(Yb) emission probabilities have been computed on the basis of the relative intensities P_X/P_γ (84) measured in 1985Me18 and 1986Ve05. The absolute measurement results of 1989Egorov for XK(Yb) [$K\alpha_2 = 1,00(2)$, $K\alpha_1 = 1,69(4)$, $K'\beta_1 = 0,54(2)$, $K'\beta_2 = 0,14(1)$] agree well with our evaluated values. The total absolute XK(Er) emission probability of 0,089(5) measured in 1990EgZY disagrees with the evaluated value of section "X Radiations".

The weighted mean of the two measurement results for the Yb $K\alpha_1$ -ray, 0,675(17), was adopted as the evaluated value and the values on $K\alpha_2$, $K'\beta_1$, $K'\beta_2$ were computed from the relative probabilities from 1996Sc06. The analogous procedure was made for the Er with the $K\alpha_1$ value from the measurements of 1986Ve05 and the other values from the relative probabilities from 1996Sc06.

P_{XK}/P_γ (84) for Er

Er	1985Me18	1986Ve05	adopted
$K\alpha_2$	} 0,0248 (6)	0,0133 (4)	0,0134 (4)
$K\alpha_1$	}	0,0238 (4)	0,0238 (4)
$K'\beta_1$	6,3 (2)·10 ⁻³	7,7 (3)·10 ⁻³	0,0077 (3)
$K'\beta_2$	1,45 (6)·10 ⁻³	2,2 (1)·10 ⁻³	0,0020 (1)

P_{XK}/P_γ (84) for Yb

Yb	1985Me18	1986Ve05	average (EV1NEW)	adopted
$K\alpha_2$	0,377 (9)	0,381 (11)	0,379 (9)	0,383 (9)
$K\alpha_1$	0,680 (17)	0,668 (20)	0,675 (17)	0,675 (17)
$K'\beta_1$	0,2145 (32)	0,228 (7)	0,221 (12)	0,222 (7)
$K'\beta_2$	0,0533 (9)	0,0604 (19)	0,057(1)	0,058 (2)

P_{XL}/P_γ (84) for Yb

Yb	adopted (1985Me18)
Ll	0,0238 (8)
$L\alpha+L\eta$	0,573 (18)
$L\beta$	0,603 (19)
$L\gamma$	0,0974 (31)
ΣXL	1,297 (27)

The total absolute Er LX emission probability has been computed using the adopted values of ω_K , ω_L , n_{KL} , the evaluated total KX absolute emission probability and the evaluated total absolute emission probabilities of L conversion electrons and electron capture.

It should be noticed that the absolute XK - emission probabilities of $P_{XK}(Er)=0,113(6)$ and $P_{XK}(Yb)=3,27(12)$ per 100 disintegrations, calculated from the adopted values of ω_K , the evaluated total absolute emission probabilities of K conversion electrons (P_{ceK}) and the electron capture ($P_{\epsilon K}$), agree well with the evaluated, 0,116 (3) and 3,31 (8), respectively.

For $P_{XL}(Yb)$ such a calculation gives 2,93 (15) per 100 disintegrations - in comparison with the value of 3,22 (13), adopted from experimental data on P_{XL}/P_γ (84).

The evaluated values of $P_{XK}(Er) = 0,116$ (3)%, $P_{XK}(Yb) = 3,31$ (8)% and $P_{XL}(Yb) = 3,22$ (13)% have been obtained directly from relative measurements of the intensity of peaks in the ¹⁷⁰Tm photon spectrum ($P_X/P_\gamma(84)$) with use of the $P_\gamma(84)$ value evaluated from independent experimental data. Unlike that the calculated value of $P_{XK}(Er) = 0,113$ (6) has been founded on the adopted semiempirical and theoretical values ω_K , $P_K(\epsilon_{0,1})$, and $\alpha_K(\gamma79)$ as well as the evaluated $P_\gamma(79)$. In the calculation of $P_{XK}(Yb) = 3,27$ (12)% the same value of $P_\gamma(84)$ is used as in the evaluation of 3,31 (8)%. However, the adopted $\omega_K(Yb)$ and theoretical value of $\alpha_K(\gamma84)$ have been used instead of the experimental relative intensity $P_{XK}/P_\gamma(84)$.

Above agreement of the evaluated and calculated values shows a concordance of the obtained decay characteristics for ¹⁷⁰Tm.

4.2. Gamma Emissions

The energy of 78,6 keV γ -ray has been obtained as the weighted mean of the following three measurements results: 78,59 (2) keV (1958Ch36), 78,7 (5) keV (1969Ha20) and 78,6 (4) keV (1970Mo07).

The 84,25 keV γ -ray energy has been adopted from 2000He14.

The absolute emission probability for the γ -ray of 84,25 keV (per 100 disintegrations) has been obtained with use of the weighted mean of the three measurement results: 2,54 (6) (1973Pl08), 2,56 (4) (1987GeZU, 1988GeZS) and 2,37 (4) (1990Ke01). This weighted average is 2,48 with an internal uncertainty of 0,03, a reduced $-\chi^2$ of 6,3 and an external uncertainty of 0,06. In this case the different statistical procedures using the weighted average give the following values for a final uncertainty: UINF - 0,064; PINF - 0,064; BAYS - 0,091; MBAYS - 0,091; LWM - 0,109; tS - 0,084. The EVINEW program has chosen MBAYS for this case and hence the uncertainty of 0,09. This value was adopted as the uncertainty of the evaluated $P_\gamma(84)$. It should be noted that the Rajeval technique leads to the same result of 2,48(9). The normalised Residuals technique gives only slightly greater value of 2,51(4).

The absolute emission probability for the γ -ray of 78,6 keV has been obtained with use of the weighted mean of the results of measurements of the ratio of $P_\gamma(79)/P_\gamma(84)$: 0,00122 (24) (1970Mo07), 0,0015 (2)(1985Me18) and 0,00140 (8) (1986Ve01). The LRSW method has expanded the uncertainty of the 1986Ve01 from 0,00008 to 0,00015 in order to reduce its relative weight from 79% to 50%. Then, the weighted mean is 0,00139 with an internal uncertainty of 0,00011, a reduced $-\chi^2$ value of 0,4 and an external uncertainty of 0,00007. The adopted value of $P_\gamma(79)/P_\gamma(84)$ is 0,00139 (11).

5. Electron Emissions

The energies of the conversion electrons have been calculated from the gamma \rightarrow transition energies given in 2.3 and the electron binding energies. The energies of the Auger electrons are taken from 1977La19 (Larkins).

The emission probabilities of the conversion electrons have been calculated using the conversion coefficients given in 2.3. The values of the emission probabilities of K-Auger electrons have been calculated using the transition probabilities given in 2.1 and 2.2, the atomic data given in 3. and the conversion coefficients given in 2.3.

6. References

- 1958Ch36 E. L. Chupp *et al.*, Phys. Rev., 1958, V.112. P.518. [Energy of $\gamma79$]
 1962Bo12 Bonner *et al.*, Phys. Rev., 1962, V. 127. P. 217. [Half-life]
 1965Fl02 K. F. Flynn *et al.*, Nucl. Sci. Eng., 1965, V. 22. P. 416. [Half-life]
 1966Di02 R. S. Dingus *et al.*, Nucl. Phys., 1966, V. 83. P.545. [$\alpha_K(\gamma84)$]
 1967Be65 J. A. Bearden, Rev. Mod. Phys., 1967, V.39. P.78. [X-ray energies]
 1967Ke13 W. I. Kerrigan, J. Inorg. Nucl. Chem., 1967, V. 29. P. 2657. [Half-life]
 1968Ni06 O. Nilsson *et al.*, Nucl. Phys., 1968, V. A120. P.561. [α_M/α_{NO}]

- 1968Re04 S. A. Reynolds *et al.*, Nucl. Sci. Eng., 1968, V.32. P.46. [Half-life]
- 1969Ha20 H. H. Hansen, S. Hellström, Z. Phys., 1969, Bd.223. S.139. [Energy of γ 79]
- 1969La34 F. Lagoutine *et al.*, Int. J. Appl. Rad. Isot., 1969, V. 20. P. 868. [Half-life]
- 1969Ne02 G. C. Nelson, E. N. Hatch, Nucl. Phys., 1969, V. A127. P.560. [$\alpha_K(\gamma$ 84)]
- 1970Mo07 S. Mohan, Phys. Rev., 1970, V. C1, N1. P. 254. [$\alpha_K(\gamma$ 84), energy and relative probability of γ 79]
- 1971Ca08 J. L. Campbell *et al.*, Nucl. Instrum. Meth., 1971, V.92. P.237. [$\alpha_K(\gamma$ 84)]
- 1973Pl08 J. Plch, J. Zderadicka, L. Kokta, Czech. J. Phys., 1973, V. 23, N 4. P. 1181. [$\alpha_K(\gamma$ 84), absolute γ 84 emission probability]
- 1977La19 F. P. Larkins, Atomic Data and Nuclear Data Tables, 1977, V.20. P.313. [Auger electron energies]
- 1978Ro21 F. Rösel, H. M. Fries, K. Alder, H. C. Pauli, Atomic Data and Nuclear Data Tables, 1978, V.21. P. 293. [Internal conversion coefficients]
- 1985Me18 D. Mehta *et al.*, Nucl. Instrum. Meth., 1985, V. A242, N1. P.149. [$\alpha_K(\gamma$ 84), $P_X/P_\gamma(84)$, $P_\gamma(79)/P_\gamma(84)$]
- 1986Ve01 Rao N. Venkateswara *et al.*, J. Phys.(London), 1986, V. G12. P.45. [$\alpha_K(\gamma$ 84), $P_\gamma(79)/P_\gamma(84)$]
- 1986Ve05 Rao N. Venkateswara *et al.*, Indian J. Phys., 1986, v.60A. P.162. [$P_X/P_\gamma(84)$]
- 1987GeZU A. M. Geidelman *et al.*, Abstracts of 37th Conference on Nuclear Spectroscopy and Atomic Nuclear Structure, Yurmala, 14-17 April, 1987, LO Nauka, Leningrad, 1987.P. 133. [Absolute γ 84 emission probability]
- 1988GeZS A. M. Geidelman, Yu. S. Egorov, N. K. Kuzmenko, V. G. Nedovesov, V. P. Chechev, G. E. Shukin, Proc. Intern. Conf. Nuclear Data for Science and Technology, Mito, Japan, 1988, p 909. [Absolute γ 84 emission probability]
- 1988Kuzmenko N. K. Kuzmenko, V. G. Nedovesov, V. P. Chechev, Izmeritel'nay Tekhnika, 1988, N 9, P. 47. [1988 decay data evaluation, absolute γ 84 emission probability]
- 1989Egorov A. G. Egorov, Yu. S. Egorov, V. G. Nedovesov, G. E. Shchukin, K. P. Yakovlev, Program and Thesis, Proc. 39th Ann. Conf. Nucl. Spectrosc. Struct. At Nuclei, Leningrad, 1989, p 505. [Yb XK-ray emission probabilities]
- 1990EgZY A. G. Egorov, V. G. Nedovesov, G. E. Shchukin, K. P. Yakovlev, Program and Thesis, Proc. 40th Ann. Conf. Nucl. Spectrosc. Struct. At Nuclei, Leningrad, 1990, p 486. [Er X K-ray emission probability]
- 1990Ke01 T. Kempisty *et al.*, Nucl. Instrum. Meth. Phys. Res., 1990, V. A286, P. 535. [Absolute γ 84 emission probability, $\alpha_K(\gamma$ 84)]
- 1995Au04 G. Audi and A. H. Wapstra, Nucl. Phys. , 1995, V. A595, P. 409.[Q values]
- 1996Ba01 C. M. Baglin, Nuclear Data Sheets, 1996, V.77, N1. P.125.[Decay scheme]
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instr. Meth. Phys. Res., 1996, V.A369. P.527.[Atomic data]
- 1998Sc28 E. Schönfeld, Appl. Rad. Isotopes, 1998, V.49. P.1353. [Fractional electron capture probabilities]
- 1998Si17 B. Singh, J. L. Rodriguez, S. S. Wong, J. K. Tuli, Nucl. Data Sheets, 1998, 84, 487. [lg f_t]
- 2000He14 R. G. Helmer and C. van der Leun, Nucl. Instrum. Meth., 2000, V. A450, P. 35. [γ -ray energies]

¹⁷⁷Lu - Comments on Evaluation of Decay Data for β^- Decay

F. G. Kondev

Evaluation Procedures

The *Limitation of Relative Statistical Weight* (LWM) [1] method for averaging numbers has been applied throughout this evaluation.

1. Decay Scheme

The decay scheme for ¹⁷⁷Lu is taken from the recent evaluations of Kondev (2002KoXX) and Browne (1993Br06). The ground state has been assigned $J^\pi = 7/2^+$ and the $7/2^+[404]$ ($g_{7/2}$) Nilsson configuration. It decays via β^- emission ($P_{\beta^-} = 100\%$) to levels of the stable ¹⁷⁷Hf daughter isotope. While the decay branches to the ¹⁷⁷Hf ground state ($J^\pi = 7/2^-$) and to the 112.9499 keV ($J^\pi = 9/2^-$), and 321.3162 keV ($J^\pi = 9/2^+$) levels are well established, there is some ambiguity in the literature regarding the direct β^- -decay feeding into the $J^\pi = 11/2^-$ level at 249.6744 keV.

2. Nuclear Data

Half-life

The half-life of the ¹⁷⁷Lu ground state has been measured by several authors and the results are summarized in Table 1. In all cases the source was prepared using the ¹⁷⁶Lu(n,γ) reaction, where a three-quasiparticle isomer ($K^\pi = 23/2^-$ and excitation energy of 970 keV), with a half-life that is significantly longer ($T_{1/2} = 160.44(6)$ d), when compared to that for the ¹⁷⁷Lu ground state, is also produced. Since the isomer de-excites partially via gamma emission ($P_\gamma = 21.4\%(8)$), its half-life and relative population should be taken into account when determining the $T_{1/2}$ for the ground state. The recommended value for the ¹⁷⁷Lu ground state half-life is $T_{1/2} = 6.647(4)$ d. It is the weighted average of the 6.645(30) d (1982La25), 6.65(1) d (2001Zi01) and 6.646(5) d (2001Sc23) values. The half-lives reported by 1958Be41, 1960Sc19, 1972Em01 and 1990Ab02 were excluded from this analysis since authors did not consider the effect of the ¹⁷⁷Lu^m isomer ($T_{1/2} = 160.44$ d) was not taken into account. Although the relative statistical weight of the 2001Sc23 value was 78.3%, its uncertainty was not increased since the set is consistent. It should be noted that there may be a systematic uncertainty in the recommended $T_{1/2}$ value for the ¹⁷⁷Lu ground state, due to possible differences in the half-life values of ¹⁷⁷Lu^m and its population intensity that were used in 1982La25, 2001Zi01 and 2001Sc23.

Table 1 Measured and recommended values for the ¹⁷⁷Lu ground state half-life.

Reference	$T_{1/2}$, d	Comment
1958Be41	6.75 (5) #	
1960Sc19	6.74 (4) #	
1972Em01	6.71 (1) #	
1990Ab02	6.7479 (7) #	
1982La25	6.645 (30)	$T_{1/2}({}^{177m}\text{Lu}) = 159.5$ d (7) was used in the fitting procedure.
2001Zi01	6.65 (1)	Corrections for $T_{1/2}({}^{177m}\text{Lu})$ have been applied, but the value has not been reported.
2001Sc23	6.646 (5)	$T_{1/2}({}^{177m}\text{Lu}) = 160.4$ d was used in the fitting procedure.
Adopted	6.647 (4)	$c2/(N-1) = 0.07$

Contributions from the decay of the ¹⁷⁷Lu^m (T_{1/2} = 160.44 d) isomer have not been taken into account. The value is not used in the analysis.

Q value

The Q(β⁻) = 498.3(8) keV is from 1995Au04. It is in agreement with that of 496.8(17) keV (1962El02), deduced from the β⁻-decay endpoint energy to the ¹⁷⁷Hf ground state. The total average decay energy released in the β⁻-decay of the ¹⁷⁷Lu ground state is calculated using RADLST [2] as 497.4(25) keV. It agrees very well with the Q(β⁻) value that is reported by Audi (1995Au04), thus suggesting that the decay scheme is complete.

2.1 b- Decay Transitions

The β⁻ transition endpoint energies were determined from Q(β⁻) = 498.3(8) keV (1995Au04) and the individual level energies. The latter were deduced from a least-squares fit to the adopted gamma-ray energies that are given in Table 3. The β⁻ transition endpoint energies are in agreement with values measured by 1962El02 and 1955Ma12. The adopted values for the β⁻ transition probabilities per 100 disintegrations were determined from the total (photons + conversion electrons) transition probability balances at each level. In general, values deduced in the present evaluation are consistent with those from 2001Sc23, 1975El07 and 1993Br06, albeit in 2001Sc23 there is no report on a direct β⁻-decay feeding into the J^π = 11/2⁻ level.

Table 2 Measured and adopted values for the ¹⁷⁷Lu b⁻-decay transition probabilities

Reference	P _{β⁻} to J ^π = 7/2 ⁻	P _{β⁻} to J ^π = 9/2 ⁻	P _{β⁻} to J ^π = 11/2 ⁻	P _{β⁻} to J ^π = 9/2 ⁺
2001Sc23	79.3 (5)	9.1 (5)		11.58 (12)
1975El07	78.6 (10)	9.1 (10)	0.05 (2)	12.2 (7)
1993Br06				
1967Ha09	87.2 (11)	6.0 (8)	0.07 (2)	6.7 (3)
1964Al04	86.3 (13)	7 (1)	0.03 (3)	6.7 (3)
1962El02	90 (4)	2.95 (3)	0.31 (6)	6.72 (25)
1956Wi39	96	1.3	0.2	2.6
1955Ma12	90	3		7
1949Do05	65	17		
Adopted	79.3 (5)	9.1 (5)	0.012 (8)	11.64 (10)

There are, however, significant differences with the 1967Ha09, 1964Al04, 1962El02, 1956Wi39, 1955Ma12 and 1949Do05 work, as summarized in Table 2. The log *ft* values were calculated using the program LOGFT [3] using the adopted β⁻ transition probabilities.

2.2 Gamma Transitions and Internal Conversion Coefficients

The measured values for gamma-ray transition energies that follow the decay of the ¹⁷⁷Lu ground state are presented in Table 3. The gamma-ray energies reported by Matsui et al. (1989Ma56) were adopted in the present evaluation. These were measured with a high precision using a germanium spectrometer. The total (photon + conversion electrons) transition probabilities were deduced by multiplying the adopted values for the relative gamma-ray intensities (Table 10) by a normalization factor that was deduced from the values for the absolute intensity per 100 disintegrations of the 208.3662 keV gamma ray (Table 11). The total electron conversion coefficients were interpolated from the tables of Rösel (1978Ro22). Transition multiplicities are taken from 2002KoXX and 1996Br06. They are based on comparisons

between the measured electron conversion coefficients with theoretical values (1978Ro22), as well as on available angular correlation data.

Table 3 Measured and adopted values for gamma ray transition energies following β^- decay of ¹⁷⁷Lu

Reference	$\gamma_{1,0}$	$\gamma_{2,1}$	$\gamma_{2,0}$	$\gamma_{3,2}$	$\gamma_{3,1}$	$\gamma_{3,0}$
1989Ma56	112.9498 (4)	136.7245 (5)	249.6742 (6)	71.6418 (6)	208.3662 (4)	321.3159 (6)
1981Hn03	112.95 (2)	136.72 (2)	249.7 (5)	71.646	208.35 (2)	321.27 (5)
1967Ha09	112.95 (2)	136.72 (5)	249.65 (6)	71.66 (6)	208.34 (6)	321.32 (12)
1965Ma18	112.952 (2)	136.730 (6)	249.868 (25)	71.646 (2)	208.359 (10)	321.330 (40)
1964Al04	112.97 (2)	136.68 (2)	249.69 (10)	71.64 (2)	208.36 (6)	321.36 (20)
1961We11	112.97 (2)	136.70 (10)	249.70 (10)	71.60 (10)	208.38 (2)	321.34 (3)
1955Ma12	112.965 (20)		250.0 (5)	71.644 (20)	208.362 (20)	321.36 (10)
Adopted	112.9498 (4)	136.7245 (5)	249.6742 (6)	71.6418 (6)	208.3662 (4)	321.3159 (6)

Details about the mixing ratios values for E1+M2 and M1+E2 transitions are given below. The electron conversion coefficients are interpolated values from the tables of Rösler (1978Ro22). The quoted uncertainties reflect the corresponding uncertainties in the mixing ratios values. Adopted α_K , α_{L1} , α_{L2} , α_{L3} , and α_M values were also used as an input for the RADLST [2] and EMISSION (2001Sc08) programs.

2.2.1 112.9498 keV ($g_{1,0}$)

Values used in the analysis of the mixing ratios are summarized in Table 4. The unweighted average value is adopted, but its uncertainty was increased to 0.4, so that the range includes the most precise value of $\delta(\gamma_{1,0}) = -4.85(5)$ (1992De53). During the analysis, the uncertainty of the 1992De53 value was also increased to 0.056, so that its relative statistical weight is scaled down from 55.8% to 50%.

Table 4 Measured and adopted mixing ratios values for the 112.9498 keV transition

Reference	$\delta(\gamma_{1,0})$	Comment
1974Kr12	-4.7 (2)	From $\gamma(\theta)$ in ^{177m} Lu ($T_{1/2} = 160.44$ d) decay.
1974Ag01	-3.99 (25)	From $\gamma\gamma(\theta)$ in ¹⁷⁷ Lu ($T_{1/2} = 6.647$ d) β^- decay.
1970Hr01	-3.7 (3)	From $\gamma\gamma(\theta)$ in ¹⁷⁷ Lu ($T_{1/2} = 6.647$ d) β^- decay.
1961We11	-4.0 (2)	From $\gamma\gamma(\theta)$ in ¹⁷⁷ Lu ($T_{1/2} = 6.647$ d) β^- decay.
1972Ho54	-4.75 (7)	From $\gamma\gamma(\theta)$ in ¹⁷⁷ Lu ($T_{1/2} = 6.647$ d) β^- decay.
1972Ho39	-4.5 (3)	From ICC ratios in ¹⁷⁷ Lu ($T_{1/2} = 6.647$ d) β^- decay.
1977Ke12	-4.8 (2)	From $\gamma(\theta)$ in ¹⁷⁷ Lu ($T_{1/2} = 6.647$ d) β^- decay.
1992De53	-4.85 (5)	From $\gamma\gamma(\theta)$ in ¹⁷⁷ Lu ($T_{1/2} = 6.647$ d) β^- decay.
Adopted	-4.4 (4)	$c2/(N-1) = 5.61$

2.2.2 136.7245 keV ($g_{2,1}$)

The adopted mixing ratios values of $\delta(\gamma_{2,1}) = -3.0 (7)$ is from 1974Kr12.

2.2.3 321.3159 keV ($g_{3,0}$)

Values used in the analysis of the mixing ratios are summarized in Table 5. The unweighted average value is adopted, but the uncertainty was expanded so that the range includes the most precise value of $\delta(\gamma_{1,0}) = +0.17(1)$ (1974Kr12). The sign of $\delta(\gamma_{3,0})$ was determined to be positive by 1974Kr12.

Table 5 Measured and adopted mixing ratios values for the 321.3159 keV transition

Reference	$ \delta(\gamma_{3,0}) $	Comment
1974Kr12	0.17 (1)	From $\gamma(\theta)$ in ^{177m} Lu ($T_{1/2} = 160.44$ d) decay
	0.42 (1)	From comparison between experimental $\alpha_K = 0.087(3)$, weighted average from values reported by 1972Gr35, 1974Ag01, 1974Je02 and 1961We11, and theoretical $\alpha_K(E1)$, and $\alpha_K(M2)$ values from 1978Ro22.
	0.42 (1)	From comparison between experimental $\alpha_L = 0.0169(8)$, weighted average from values reported by 1972Gr35, 1974Ag01, 1974Je02 and 1961We11, and theoretical $\alpha_L(E1)$, and $\alpha_L(M2)$ values from 1978Ro22.
Adopted	0.34 (17)	$c^2/(N-1) = 208.33$

2.2.4 208.3662 keV ($g_{3,1}$)

Values used in the analysis of the mixing ratios are given in Table 6. The weighted average and the internal uncertainty were adopted. The sign of $\delta(\gamma_{3,1})$ is uncertain. It has been reported to be positive by 1974Kr12, but negative by 1977Ke12 and 1961We11.

Table 6 Measured and adopted mixing ratios values for the 208.3662 keV transition

Reference	$ \delta(\gamma_{3,1}) $	Comment
1974Kr12	0.07 (2)	From $\gamma(\theta)$ in ^{177m} Lu ($T_{1/2} = 160.44$ d) decay
1977Ke12	0.08 (2)	From $\gamma(\theta)$ in ¹⁷⁷ Lu ($T_{1/2} = 6.647$ d) β^- decay
1961We11	0.07 (3)	From $\gamma\gamma(\theta)$ in ¹⁷⁷ Lu ($T_{1/2} = 6.647$ d) β^- decay
Adopted	0.074 (13)	$c^2/(N-1) = 0.07$

2.2.5 71.6418 keV ($g_{3,2}$)

Values used in the analysis of the mixing ratios are shown in Table 7. None of them has a relative statistical weight greater than 50%, and hence the weighted average value was adopted. The sign of $\delta(\gamma_{3,2})$ is negative as determined by 1974Kr12 and 1970Hr01.

Table 7 Measured and adopted mixing ratios values for the 71.6418 keV transition

Reference	$ \delta(\gamma_{3,2}) $	Comment
1974Kr12	0.051(37)	From $\gamma(\theta)$ in ^{177m} Lu ($T_{1/2} = 160.44$ d) decay.
1974Ag01	0.049 (15)	From comparison between experimental $\alpha_K = 0.90(11)$ from 1974Ag01 and theoretical $\alpha_K(E1)$, and $\alpha_K(M2)$ values from 1978Ro22.
1970Hr01	0.017 (7)	From $\gamma\gamma(\theta)$.
	0.016 (6)	From comparison between experimental $\alpha_{L1} = 0.076(5)$, weighted average from values reported by 1972Gr35 and 1974Ag01, and theoretical $\alpha_{L1}(E1)$, and $\alpha_{L1}(M2)$ values from 1978Ro22.
	0.034 (14)	From comparison between experimental $\alpha_{L2} = 0.029(3)$, weighted average from values reported by 1972Gr35 and 1974Ag01, and theoretical $\alpha_{L2}(E1)$, and $\alpha_{L2}(M2)$ values from 1978Ro22.
Adopted	0.018 (4)	$c^2/(N-1) = 0.37$

3. Atomic Data

3.1 Hf

The data are from Schönfeld and Janssen (1996Sc06).

3.1.1 X Radiation

While the energies for $XK\alpha_2$ (Hf) and $XK\alpha_1$ (Hf) are from Schönfeld and Rodloff (1999ScZX), the $XK\beta$ and XL energies are from Firestone (1996FiZX). Relative emission probabilities were calculated using the program EMISSION (2001Sc08).

3.1.2 Auger Electrons

The energies for KLL (Hf), KLX (Hf) and KXY (Hf) are from Schönfeld and Rodloff (1998ScZM). Relative emission probabilities were calculated using the program EMISSION (2001Sc08).

4. Photon Emission

4.1 X-Ray Emission

While the energies for $XK\alpha_2$ (Hf) and $XK\alpha_1$ (Hf) are from Schönfeld and Rodloff (1999 ScZX), the $XK\beta$ and XL energies are from Firestone (1996FiZX). The adopted absolute intensities per 100 disintegrations were calculated using the program EMISSION (2001Sc08). Comparisons between calculated values and the experimental data in 2001Sc23 and 1987Me17, as well as values calculated using the program RADLST [2], are presented in Table 8. In general the agreement between various entries is fairly good, thus suggesting that the ¹⁷⁷Lu ground state decay scheme is complete.

Table 8 comparison between various X-ray emission intensities per 100 disintegration

	Energy KeV	2001Sc23	1987Me17	RADLST	EMISSION
XLI (Hf)	6.960	0.0735 (25)	0.087 (5)		0.0613 (16)
XL α_2 (Hf)	7.844	}	}		0.137 (4)
XL α_1 (Hf)	7.899	}	1.51 (3)	1.59 (6)	1.21 (3)
XL η (Hf)	8.139	}	}		0.0313 (9)
XL β_4 (Hf)	8.905	}	}		0.0335 (12)
XL β_1 (Hf)	9.023	}	1.34 (3)	}	1.15 (4)
XL β_6 (Hf)	9.023	}	}	1.76 (7)	0.0147 (4)
XL β_3 (Hf)	9.163	}	}		0.0435 (15)
XL $\beta_{2,15}$ (Hf)	9.342	0.274 (7)	}		0.248 (7)
XL γ_1 (Hf)	10.516	}	0.231 (6)	}	0.222 (6)
XL γ_6 (Hf)	10.733	}	}	}	0
				0.292 (12)	
XL γ_2 (Hf)	10.834	}	0.0223 (14)	}	0.00835 (19)
XL γ_3 (Hf)	10.890	}	}	}	0.0115 (4)
XL				3.08 (7)	3.18 (6)
XK α_2 (Hf)	54.6120 (7)	1.55 (3)	1.65 (3)	1.59 (5)	1.59 (3)
XK α_1 (Hf)	55.7909 (8)	2.73 (6)	2.84 (5)	2.78 (9)	2.78 (6)
XK β_1 (Hf)	62.985-63.662	0.885 (15)	0.919 (16)		0.917 (23)
XK β_2 (Hf)	64.942-65.316	0.238 (5)	0.252 (5)		0.245 (8)
XK β (Hf)				1.16 (4)	1.16 (3)

4.2 Gamma Emission

The measured relative intensities for transitions following the β^- decay of ¹⁷⁷Lu and their adopted values are presented in Table 9. The original values were normalized to $I_\gamma = 100.0$ for the 208.3662 keV ($\gamma_{3,1}$) gamma ray. The uncertainty in I_γ for the 321.3159 keV ($\gamma_{3,0}$) gamma ray was increased 1.86 times so that its statistical weight was lowered from 77.6% to 50%.

The measured absolute intensities for the 208.3662 keV ($\gamma_{3,1}$) gamma ray and its corresponding adopted value are presented in Table 10. The latter was used to normalize the relative intensities (Table 9) to absolute values per 100 disintegrations.

Table 9 - Relative gamma-ray intensities for transitions following β^- decay of ¹⁷⁷Lu

	$\gamma_{1,0}$	$\gamma_{2,1}$	$\gamma_{2,0}$	$\gamma_{3,2}$	$\gamma_{3,1}$	$\gamma_{3,0}$
2001Sc23	59.6 (6)	0.448 (8)	1.918 (17)	1.674 (21)	100.0	2.002 (19) *
1987Me17	59.6 (11)	0.457 (8)	2.00 (3)	1.71 (5)	100.0	2.17 (4)
1974Ag01	60 (5)	0.52 (5)	1.90 (20)	1.50 (10)	100.0	2.00 (20)
1964Al04	58 (4)	0.43 (3)	1.93 (14)	1.40 (10)	100.0	1.99 (14)
1961We11	62 (2)	0.47 (15)	2.00 (20)	0.30 (10) #	100.0	2.28 (10)
1955Ma12	45.5 #		1.36 #	0.91 #	100.0	1.45 (29) #
Adopted	59.7 (5)	0.453 (6)	1.938 (15)	1.663 (19)	100.0	2.08 (8)
c²/(N-1)	0.38	0.76	1.45	3.58		3.62

* The uncertainty was increased 1.86 times in order to reduce its statistical weight from 77.6% to 50%.

Value not used in the analysis.

Table 10 - Absolute emission probabilities per 100 disintegrations for the 208.3662 keV gamma ray

	Absolute Intensity for $\gamma_{3,1}$ per 100 disintegrations, %
2001Sc23	10.36 (7)
1964Cr02	10.7 (5)
1961We11	11.4 (6)
Adopted	10.38 (7)
c²/(N-1)	1.69

5. Electron Emission

The electron energies and emission probabilities were calculated using the RADLST [2] program. The average β^- energies were calculated using the LOGFT [3] program. The β^- transition endpoint energies were determined using $Q(\beta^-) = 498.3(8)$ keV (1995Au04) and the individual level energies that were deduced from a least-squares fit to the recommended gamma-ray energies. The adopted values for the β^- transition emission probabilities were determined from the total (photons + electrons) gamma-ray emission probability balances at each level.

References

[1] M. J. Woods and A. S. Munster, National Physics Laboratory, Teddington, UK, Rep. RS(EXT) 95, (1988)
 [2] The program RADLST, T. W. Burrows, report BNL-NCS-52142, February 29, 1988
 [3] The program LOGFT, NNDC
 1949Do05 - D. G. Douglas, Phys. Rev. 75, 1960 (1949) [P_{β^-}]
 1955Ma12 - P. Marmier, F. Boehm, Phys. Rev. 97, 103 (1955) [P_{β^-} , E_γ , I_γ]
 1956Wi39 - T. Wiedling, Thesis, Univ. Stockholm (1956) [P_{β^-}]
 1958Be41 - R. H. Betts, O. F. Dahlinger, D. M. Munro, Can. J. Phys. 36, 73 (1958) [$T_{1/2}$]

- 1960Sc19 - L. C. Schmid, W. P. Stinson, Nucl. Sci. Eng. 7, 477 (1960) [$T_{1/2}$]
 1961We11 - H. I. West, Jr., L. G. Mann, R. J. Nagle, Phys. Rev. 124, 527 (1961) [E_γ , I_γ , δ , ICC]
 1962El02 - M. S. El-Nesr, E. Bashandy, Nucl. Phys. 31, 128 (1962) [P_{β^-}]
 1964Al04 - P. Alexander, F. Boehm, E. Kankeleit, Phys. Rev. 133, B284 (1964) [P_{β^-} , E_γ , I_γ]
 1964Cr02 - D. F. Crouch, L. D. McIsaac, IDO-16932, p.26 (1964) [I_γ]
 1965Ma18 - B. P. K. Maier, Z. Physik 184, 153 (1965) [E_γ]
 1967Ha09 - A. J. Haverfield, F. M. Bernthal, J. M. Hollander, Nucl. Phys. A94, 337 (1967) [P_{β^-} , E_γ , I_γ]
 1970Hr01 - B. Hrastnik, I. Basar, M. Diksic, K. Ilakovac, V. Kos, A. Ljubicic, Z. Phys. 239, 25 (1970) [δ]
 1972Em01 - J. F. Emery, S. A. Reynolds, E. I. Wyatt, G. I. Gleason, Nucl. Sci. Eng. 48, 319 (1972) [$T_{1/2}$]
 1972Gr35 - V. N. Grigorev, D. M. Kaminker, Y. V. Sergeenkov, Izv. Akad. Nauk SSSR, Ser. Fiz. 36, 842 (1972); Bull. Acad. Sci. USSR, Phys. Ser. 36, 762 (1973) [ICC]
 1972Ho39 - S. Hogberg, R. Jadrny, S. E. Karlsson, G. Malmsten, O. Nilsson, Z. Phys. 254, 89 (1972) [δ]
 1972Ho54 - L. Holmberg, V. Stefansson, J. Becker, C. Bargholtz, L. Gidefeldt, Phys. Scr. 6, 177 (1972) [δ]
 1974Ag01 - A. P. Agnihotry, K. P. Gopinathan, H. C. Jain, Phys. Rev. C9, 336 (1974) [I_γ , ICC, δ]
 1974Je02 - B. D. Jeltama, F. M. Bernthal, Phys. Rev. C10, 778 (1974) [ICC]
 1974Kr12 - K. S. Krane, C. E. Olsen, W. A. Steyert, Phys. Rev. C10, 825 (1974) [$\gamma(\theta)$]
 1975El07 - Y. A. Ellis, B. Harmatz, Nucl. Data Sheets 16, 135 (1975) [nuclear data]
 1977Ke12 - H. E. Keus, W. J. Huiskamp, Physica 85B, 137 (1977) [δ]
 1978Ro22 - F. Rosel, H. M. Friess, K. Alder, H. C. Pauli, At. Data Nucl. Data Tables 21, 92 (1978) [ICC]
 1981Hn03 - V. Hnatowicz, Czech. J. Phys. B31, 260 (1981) [E_γ , I_γ]
 1982La25 - F. Lagoutine, J. Legrand, Int. J. Appl. Radiat. Isotop. 33, 711 (1982) [$T_{1/2}$]
 1987Me17 - D. Mehta, B. Chand, S. Singh, M. L. Garg, N. Singh, T. S. Cheema, P. N. Trehan, Nucl. Instrum. Methods Phys. Res. A260, 157 (1987) [I_γ]
 1989Ma56 - S. Matsui, H. Inoue, Y. Yoshizawa, Nucl. Instrum. Methods Phys. Res. A281, 568 (1989) [E_γ]
 1990Ab02 - A. Abzouzi, M. S. Antony, A. Hachem, V. B. Ndocko Ndongue, J. Radioanal. Nucl. Chem. 144, 359 (1990) [$T_{1/2}$]
 1992De53 - C. C. Dey, B. K. Sinha, R. Bhattacharya, Nuovo Cim. 105A, 1307 (1992) [δ]
 1993Br06 - E. Browne, Nucl. Data Sheets 68, 747 (1993) [nuclear data]
 1995Au04 - G. Audi, A. H. Wapstra, Nucl. Phys. A595, 409 (1995) [atomic masses]
 1996FiZX - R. B. Firestone, Table of Isotopes, 8th Ed., V. S. Shirley, C. M. Baglin, S. Y. F. Chu, J. Zipkin Eds., John Wiley and Sons, Inc., New York, Vol.2 (1996) [nuclear data]
 1996Sc06 - E. Schönfeld and H. Janssen, Nucl. Instrum. and Methods A369, 527 (1996) [atomic data]
 1998ScZM - E. Schönfeld, G. Rodloff, PTB-6.11-98-1 (1998) [KLL, KLX, KXY energies]
 1999ScZX - E. Schönfeld, G. Rodloff, PTB-6.11-1999-1 (1999) [X-ray energies]
 2001Sc08 - E. Schönfeld, U. Schötzig, Appl. Radiat. Isot. 54, 785 (2001) [Emission probabilities]
 2001Sc23 - U. Schötzig, H. Schrader, E. Schönfeld, E. Gunther, R. Klein, Appl. Radiat. Isot. 55, 89 (2001) [P_{β^-} , $T_{1/2}$, I_γ]
 2001Zi01 - B. E. Zimmerman, M. P. Unterweger, J. W. Brodack, Appl. Radiat. Isot. 54, 623 (2001) [$T_{1/2}$]
 2002KoXX - F. G. Kondev, Nuclear Data Sheets, 98, 801 (2003) [nuclear data]

¹⁸²Ta - Comments on evaluation of decay data by V. Chisté and M. M. Bé

This evaluation was completed in September 2010, including all publications by this date.

1 Decay Scheme

¹⁸²Ta disintegrates 100 % by beta minus emissions to excited levels of ¹⁸²W.

A good agreement was found between the effective Q value (1821 (19) keV) calculated from the decay scheme data and the adopted and recommended value from the mass adjustment of Audi (2003Au03).

2 Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2003Au03).


Experimental ¹⁸²Ta half-life values (in days) are given in Table 1:

Table 1: Experimental values of ¹⁸²Ta half-life

Reference	Experimental value (d)	Comments
R. V. Zumstein (1943Zu**)	117 (3)	
L. Seren (1947Se33)	117 (3)	
L. Meitner (1948Me**)	117.5 (18)	
W. K. Sinclair (1951Si25)	111 (1)	
G. G. Eichholz(1952Ei02)	111.2 (5)	
H. W. Wright (1957Wr37)	115.05 (25)	Superseded by 1972Em01
J. P. Keene (1958Ke26)	114.80 (12)	
A. Specke (1958Sp17)	118.2 (5)	
D. A. Walker (1967Wa29)	117.3 (10)	
J. F. Emery (1972Em01)	115.0 (2)	
C. J. Visser (1973Vi13)	114.740 (24)	Original uncertainty is 3σ (0.08)
U. Schötzig (1980Sc07)	114.43 (4)	
Recommended value	114.61 (13)	$\chi^2 = 16$

For the data of L. Meitner (1948Me**), the evaluators have chosen to use the average value of 117.5 (18), calculated from two experimental values given in the paper to produce a single DDEP value for each laboratory. A weighted average has been calculated using LWEIGHT computer program (version 3). Originally, the largest contributions to the weighted average come from the values of U. Schötzig (1980Sc07) and C. J. Visser (1973Vi13), amounting to 43 % and 70 %, respectively. LWEIGHT increased the uncertainty of 1973Vi13 value from 0.024 to 0.037 in order to reduce its relative weight from 70 % to 50 %.

The adopted value is the weighted average of 114.61 d with an uncertainty of 0.13 d (expanded so range to include the most precise value of C. J. Visser (1973Vi13)). The reduced- χ^2 value is 16, that reflects the high discrepancy of the set of data. See Graphic 1.

Graphic 1: ¹⁸²Ta half-life values

2.1 β^- Transitions

The maximum energies of the β^- transitions in the decay of ¹⁸²Ta \rightarrow ¹⁸²W have been obtained from the Q value (2003Au03) and the level energies given in Table 2 from B. Singh (2010Si13).

Table 2: ¹⁸²W levels populated in the decay of ¹⁸²Ta and the adopted β^- transition probabilities

Level Number	Level energy, (keV)	Spin and parity	Half-life	Adopted P _{β^-} (%)
0	0	0 ⁺		
1	100.10598 (7)	2 ⁺	1.40 (2) ns	~ 0*
2	329.4268 (6)	4 ⁺	62 (3) ps	~ 0*
5	1221.4001 (10)	2 ⁺	0.434 (11) ps	1.6 (22)
6	1257.4121 (11)	2 ⁺	1.71 (13) ps	0.22 (21)
7	1289.1498 (10)	2 ⁻	1.12 (4) ns	45.1 (23)
8	1331.1153 (10)	3 ⁺	< 0.6 ns	2.39 (15)
9	1373.8301 (10)	3 ⁻	78 (10) ps	19.9 (7)
10	1442.836 (9)	4 ⁺	0.32 (3) ps	0.563 (10)
11	1487.5018 (10)	4 ⁻	< 49 ps	1.5 (7)
12	1510.25 (7)	4 ⁺		0.1414 (39)
13	1553.2240 (10)	4 ⁻	1.27 (4) ns	29.0 (7)

* Measured values by 1967Ba01 are 0.058 (6) and 0.096 (10), respectively. These values are inconsistent with intensity balance at each level of the decay scheme.

The adopted β^- transition probabilities and the associated uncertainties (Table 2) were deduced from the γ transition probability balance at each level of the decay scheme.

The values of log ft and average β^- energies have been calculated with the program LOGFT for the allowed and 1st forbidden β^- transitions.

2.2 γ Transitions

The γ -ray transition probabilities were calculated using the γ -ray emission intensities and the relevant internal conversion coefficients (see **5.2 γ Emissions**).

For all γ transitions, the internal conversion coefficients (ICC) and the associated uncertainties were interpolated from the theoretical values of I. M. Band et al. (2002Ba85) using the BrIcc computer program (2008Ki07) for the “frozen orbital” approximation.

For multipolarities and mixing ratios, the evaluators used:

1) Multipolarities and mixing ratios of the γ -ray transitions listed in the Table 3 are from B. Singh (2010Si13).

Table 3: Multipolarities of γ -ray transitions

	Multipolarity	E_γ (keV)
¹⁸² W	[E2]	(121.50 (14)), 829.80 (12)
	[E1]	110.388 (9)
	E3	1373.824 (3)
	E2	100.10595 (7), 198.35187 (29), 229.3207 (6), 264.0740 (3), 351.06 (6), 891.9710 (12), 927.9828 (13), 1035.80 (14), 1221.395 (3), 1257.407 (3), 1410.14 (7)
	E1	31.7377 (15), 42.7148 (14), 116.4179 (6), 152.42991 (26), 156.3864 (3), 222.1085 (3), 1158.0711 (12),
	E0	1135.91 (14)
	M1 + E2	1231.004 (3) $\delta = -33$ (+6, -9) 1180.82 (7) $\delta = -2.8$ (10) 1001.6856 (12) $\delta = -8.9$ (+18, -21)
¹⁸² W	M2	1289.145 (3)
	M2 + E3	1387.390 (3) $\delta = 2.6$ (3)
	E1 + M2 + E3	1273.719 (3) $\delta(M2/E1) = 0.36$ (10), $\delta(E3/E1) = -0.28$ (12)

2) For the 84-, 113- and 179-keV γ -ray transitions (M1 + E2), the adopted mixing ratios (δ) are the weighted means of the δ values found in the literature and shown in Table 4. A good agreement has been found between the experimental values of K and L internal coefficients and the calculated ones obtained by using the evaluated δ values and the BrIcc program.

Table 4: Experimental and recommended conversion coefficients and mixing ratios for 84-, 113- and 179-keV γ -ray transitions

E_γ (keV)	δ experimental (mixing ratio)	α experimental	α theoretical (given by BRICC)
84.68024 (26)	0.30 (9) (1964Ba12) 0.33 (17) (1966Gr21) 0.35 (7) (1967Ni03) 0.352 (45) (1972He10) 0.30 (2) (1972Kr05) 0.31 (5) (1975Qu01) 0.30 (2) (1980Sp01) 0.32 (3) (1983Ri05)	$\alpha_K = 5.2$ (10) $\alpha_L = 1.44$ (10) (1963Ni07)	$\alpha_K = 5.88$ (9) $\alpha_L = 1.36$ (4)
Recommended value	0.309 (12)	$\chi^2 = 0.26$	

E_γ (keV)	δ experimental (mixing ratio)	α experimental	α theoretical (given by BRICC)
113.67170 (22)	0.21 (6) (1964Ba12) 0.32 (17) (1966Gr21) 0.30 (14) (1967Ni03) 0.36 (8) (1972He10) 0.31 (2) (1972Kr05) 0.31 (5) (1975Qu01) 0.36 (3) (1980Sp01) 0.36 (2) (1983Ri05)	$\alpha_K = 2.4$ (10) $\alpha_L = 0.50$ (5) (1963Ni07)	$\alpha_K = 2.52$ (4) $\alpha_L = 0.519$ (10)
Recommended value	0.338 (12)	$\chi^2 = 0.70$	
179.39381 (25)	2.8 (8) (1964Ba12) 0.72 (26) (1966Gr21) 0.84 (32) (1967Ni03) 0.92 (10) (1972Kr05) 1.26 (15) (1981Ka22) 2.1 (3) (1983Ri05) 2.2 (2) (1992Ch26)	$\alpha_K = 0.49$ (5) $\alpha_L = 0.15$ (2) (1963Ni07)	$\alpha_K = 0.44$ (8) $\alpha_L = 0.148$ (7)
Recommended value	1.21 (29)	$\chi^2 = 7$	

3) For the eleven remaining γ -ray transitions, the mixing ratios (δ) were deduced by comparison between the weighted mean of the experimental values of internal coefficients and the theoretical ICC calculated using the BrIcc computer code (2008Ki07), shown in the Table 5.

Table 5: Experimental and recommended conversion coefficients and mixing ratios

E_γ (keV)	α experimental	δ (mixing ratio)	α theoretical (given by BRICC)	Multipolarities
65.72215 (15)	$\alpha_L = 2.5$ (1) (1963Ni07)	0.094 (43)	$\alpha_L = 2.3$ (1)	M1 + E2
67.74970 (10)	$\alpha_L = 0.18$ (2) (1963Ni07)	0.018 (9)	$\alpha_L = 0.17$ (2)	E1 + M2
959.7203 (12)	$\alpha_K = 7$ (5) 10^{-3} (1961Gr21) $\alpha_K = 9.2$ (24) 10^{-3} (1966Dz01) $\alpha_K = 9.08$ (20) 10^{-3} (1976He18) α_K (LWM) = 9.08 (20) 10^{-3}	- 5.48 (44)	$\alpha_K = 9.01$ (15) 10^{-3}	M2 + E3
1044.4001 (12)	$\alpha_K = 2.4$ (6) 10^{-3} (1966Dz01) $\alpha_K = 4.4$ (20) 10^{-3} (1969Ga23) $\alpha_K = 4.35$ (10) 10^{-3} (1976He18) α_K (LWM) = 4.36 (10) 10^{-3}	0.48 (1)	$\alpha_K = 4.44$ (12) 10^{-3}	E1 + M2
1113.406 (9)	$\alpha_K = 4.8$ (8) 10^{-3} (1972Ga23) $\alpha_K = 3.59$ (13) 10^{-3} (1975We22) $\alpha_K = 3.02$ (6) 10^{-3} (1976He18) α_K (LWM) = 3.32 (30) 10^{-3}	5.6 (+13,-10) (from 1983Ri05)	$\alpha_K = 3.11$ (8) 10^{-3}	M1 + E2
1121.290 (3)	$\alpha_K = 3.9$ (2) 10^{-3} (1960Gr**) $\alpha_K = 3.2$ (2) 10^{-3} (1964Da15) $\alpha_K = 2.9$ (3) 10^{-3} (1966Dz01) $\alpha_K = 3.28$ (15) 10^{-3} (1966Ko12) $\alpha_K = 3.15$ (19) 10^{-3} (1972Ga23) $\alpha_K = 3.16$ (10) 10^{-3} (1975We22) $\alpha_K = 2.99$ (4) 10^{-3} (1976He18) α_K (LWM) = 3.036 (40) 10^{-3}	30 (+6,-4) (from 1983Ri05)	$\alpha_K = 2.97$ (5) 10^{-3}	M1 + E2

E_γ (keV)	α experimental	δ (mixing ratio)	α theoretical (given by BRICC)	Multipolarities
1157.3022 (11)	$\alpha_K = 2.7$ (3) 10^{-3} (1960Gr**) $\alpha_K = 3.5$ (5) 10^{-3} (1964Da15) $\alpha_K = 3.5$ (8) 10^{-3} (1966Dz01) $\alpha_K = 6.3$ (4) 10^{-3} (1966Ko12) $\alpha_K = 6.8$ (7) 10^{-3} (1972Ga23) α_K (LWM) = 4.1 (14) 10^{-3}	1.3 (7)	$\alpha_K = 3.9$ (11) 10^{-3}	M1 + E2
1189.040 (3)	$\alpha_K = 4.3$ (2) 10^{-3} (1960Gr**) $\alpha_K = 4.6$ (3) 10^{-3} (1964Da15) $\alpha_K = 3.6$ (4) 10^{-3} (1966Dz01) $\alpha_K = 4.22$ (40) 10^{-3} (1966Ko12) $\alpha_K = 4.10$ (21) 10^{-3} (1972Ga23) $\alpha_K = 4.18$ (14) 10^{-3} (1975We22) $\alpha_K = 3.88$ (4) 10^{-3} (1976He18) α_K (LWM) = 3.93 (6) 10^{-3}	δ (M2/E1) = 0.470 (17), δ (E3/E1) = - 0.662 (32)	$\alpha_K = 3.732$ (33) 10^{-3}	E1 + M2 + E3
1223.7928 (12)	$\alpha_K = 2.4$ (4) 10^{-3} (1976He18)	0.38 (7)	$\alpha_K = 2.4$ (5) 10^{-3}	E1 + M2
1342.72(5)	$\alpha_K = 1.9$ (10) 10^{-3} (1966Dz01) $\alpha_K = 2.20$ (85) 10^{-3} (1966Ko12) $\alpha_K = 2.28$ (6) 10^{-3} (1976He18) α_K (LWM) = 2.28 (6) 10^{-3}	- 0.11 (11)	$\alpha_K = 2.3$ (5) 10^{-3}	E2 + M3
1453.1118 (10)	$\alpha_K = 3.7$ (6) 10^{-3} (1966Dz01) $\alpha_K = 2.4$ (16) 10^{-3} (1966Ko12) $\alpha_K = 4.38$ (30) 10^{-3} (1976He18) α_K (LWM) = 4.28 (29) 10^{-3}	2.1 (4)	$\alpha_K = 4.3$ (3) 10^{-3}	M2 + E3

For the 1113-, 1121- and 1157-keV γ -ray transitions, from comparison between measured and calculated ICCs, the introduction of a third E0 component appears not to be necessary. This was proposed for:

1. 1113-keV by 1975We22, but not by 1972Ga23 and 1976He18;
2. 1121-keV by 1966Ko12 and 1975We22, but not by 1960Gr**, 1964Da15, 1966Dz01 and 1972Ga23;
3. 1157-keV by 1966Ko12 and 1972Ga23, but not by 1960Gr**, 1964Da15 and 1966Dz01.

3 Atomic Data

Atomic values, ω_K , ω_L , ω_M , n_{KL} and the X-ray and Auger electron relative probabilities are from Schönfeld and Janßen (1996Sc06).

4 Electron emissions

The conversion electron emission probabilities were deduced from the ICC values and the γ -ray emission intensities.

5 Photon emissions

5.1 X-ray emissions

The X-ray absolute intensities were deduced from the decay data using the EMISSION computer code and are compared in Table 6 with measured values found in the literature. A good agreement has been found between the experimental and calculated values, supporting the overall consistency of the decay scheme data.

Table 6: Experimental and recommended (calculated) values of X-ray absolute intensities (%)

	U. Schötzig (1980Sc07)	B. Chand (1992Ch26) [@]	Recommended values
K α x-ray	28.02 (52)	27.82 (39)	27.54 (34)
K β_1 x-ray		6.01 (12)	5.79 (13)
K β_2 x-ray		1.51 (5)	1.59 (5)
L x-ray		21.84 (44)	24.4 (4)

[@]Using a normalization factor of 0.3517 (33) (see **5.2 Gamma Emissions**)

5.2 Photon emissions

The energies of the γ -rays in Table 7 are from R. G. Helmer (2000He14). For other γ -rays, the energy values come from B. Singh (2010Si13).

Table 7: γ -ray energies given by R. G. Helmer (2000He14).

E$_{\gamma}$ (keV)	65.722 15 (15), 67.749 70 (10), 84.680 24 (26), 100.105 95 (7), 113.671 70 (22), 116.417 9 (6), 152.429 91 (26), 156.386 4 (3), 179.393 81 (25), 198.351 87 (29), 222.108 5 (3), 229.320 7 (6), 264.074 0 (3), 1121.290 (3), 1189.040 (3), 1221.395 (3), 1231.004 (3), 1257.407 (3), 1273.719 (3), 1289.145 (3), 1373.824 (3), 1387.390 (3)
--------------------------------------	---

The experimental relative γ -ray emission intensities from ¹⁸²Ta were obtained from all the available relative values (Table 8).

The normalization factor to convert the relative emission intensities to absolute emission intensities was calculated using the formula:

$$N = \left(\frac{100}{\sum (1 + \alpha_T) P_{rel}} \right) = 0.3517 (33),$$

where the sum is over all the γ transitions to the ground state (100-, 1135-, 1221-, 1257-, 1289- and 1373-keV) and α_T is the relevant coefficient. The uncertainty was calculated through its propagation on the formula given above.

The experimental γ -ray emission probabilities relative to 100 for the 1121-keV γ -ray are given in Table 8, except for the Edwards's values (1965Ed01) who measured only the low energy γ -rays until 264-keV relatively to the 100 keV line.

Our recommended relative and absolute γ -ray emission probabilities are given in Table 9.

The adopted values are the weighted means calculated by the LWEIGHT program (version 3).

Were omitted from analysis:

* : N. A. Voinova (1959Vo27), V. D. Vitman (1961Vi07) and N. A. Voinova (1961Vo05), because these values come from the same laboratory that B. S. Dzhelepov (1966Dz01);

\$: Idem for 1976He18 superseded by 1977Ge12 only for high energy γ -rays (1121-keV to 1453-keV), where both measured the same energies. For other energies (891-keV to 1113-keV), the evaluators used the values given by 1976He18;

£ : Idem to 1983EI02, superseded by 1990Ja02;

§ : Idem to 1989Ka20, superseded by 1992Ch26;

⌘ : 1969Wh03 - superseded by 1970Wh03;

μ : the set of value from H. Daniel (1964Da15), because of a lack of information on the γ -ray reference line more generally, on γ spectrometry part of the experiment;

& : the set of value from W. F. Edwards (1965Ed01), because of a lack of information in the article about the experimental measurements carried out and, therefore on the results.

Table 8: Experimental data sets of the relative γ -ray emission intensities (%) (cont'd. next pages)

Energy (keV) Reference	31	42 65		67	84	100	110	113	116	152
1959Vo27*										
1961Ry03						67.2 (47)£				33.2 (29)£
1961Vi07*										
1961Vo05*						15 (7)				18 (4)
1964Da15 μ			8.6 (5)	128 (6)	9.0 (6)	43.2 (25)		5.6 (3)	1.41 (12)	19.4 (8)
1965Ed01&		1.73 (9)	20.0 (10)	293 (15)	18.8 (9)	100		13.6 (7)	3.16 (19)	51.0 (20)
1965He07										
1966Dz01										
1966Ko12										
1969Sa25					7.0 (7)	40.7 (41)		5.2 (5)	1.2 (2)	19.5 (20)
1969Wh03 \square										
1970Wh03										21.3 (10)
1971Ja21						40.2 (10)				20.5 (5)
197MI01					6.70 (22)*	38.0 (23)		4.90 (29)	1.00 (8)*	19.7 (12)
1972Ga23					7.6 (8)	40.3 (40)		5.28 (40)	1.27 (13)	19.3 (14)
1974La15				121.0 (52)	7.82 (16)	37.43 (80)*		6.15 (14)*		18.70 (60)
1976He18 $\$$										
1977Ge12				122 (7)	7.80 (41)	40.8 (35)		5.43 (18)	1.260 (42)	20.5 (6)
1978MeZK/1990Me15	2.75 (6)	0.86 (7)	8.75 (17)	130 (10)	7.19 (14)	40.4 (5)	0.330 (20)	5.34 (5)	1.260 (20)	19.95 (19)
1980Ro22						40.6 (26)		4.95 (40)	1.18 (18)	19.59 (80)
1980Sc07	2.53 (6)	0.750 (21)	161.7 (36)		7.45 (17)	40.3 (10)		5.29 (19)	1.26 (5)	19.69 (28)
1981Is08	1.18 (12)*	0.82 (10)	13.5 (14)*		7.6 (6)	41.6 (14)*	0.25 (4)	5.87 (40)*	1.14 (8)*	23.15 (50)*
1983Ji01						40.3 (6)	0.25 (6)	5.36 (7)	1.260 (30)	19.94 (19)
1983El02£	1.40 (2)	0.80 (1)	8.45 (14)	118.3 (20)	6.81 (26)	40.50 (64)	0.25 (1)	5.47 (8)	1.22 (2)	19.52 (32)
1986Wa35					7.30 (22)	39.03 (64)*		4.44 (16)*		21.19 (39)
1989Ka20 $\$$					7.87 (14)	41.50 (51)		5.47 (8)	1.24 (4)	20.30 (25)
1990Ja02	2.21 (2)	0.82 (3)	8.55 (7)	120.0 (11)	7.31 (5)	40.45 (51)	0.30 (4)	5.31 (8)	1.28 (6)	19.86 (17)
1992Su09	1.80 (6)*	0.827 (24)	7.61 (16)	126.2 (24)	7.80 (16)	42.6 (9)*	0.37 (3)	5.64 (11)	1.33 (3)	20.94 (24)
1992Ch26	2.44 (7)	0.710 (21)	8.40 (21)	131.8 (24)	7.65 (10)	41.4 (5)*	0.300 (10)	5.27 (10)	1.230 (22)	20.40 (26)
1992Ke02	2.46 (5)	0.754 (18)	9.02 (22)		7.43 (12)	40.5 (5)	0.320 (20)	5.34 (6)	1.270 (21)	19.81 (22)
1998Mi17					7.58 (7)	38.50 (23)*		5.21 (5)		19.60 (11)
Evaluated	2.38 (17)	0.765 (18)	8.45 (20)	124.0 (40)	7.45 (14)	40.42(24)	0.305 (9)	5.315 (29)	1.264 (10)	19.93 (33)
χ^2	20	3.3	9.7	3.8	2.6	0.025	1.4	1.5	0.8	3.2

* Outliers values, based on the Chauvenet's criterion and thus were omitted in the final calculation.

£ Data rejection parameters for deviation from weighted mean (3σ).

Energy (keV)	156	179	198	222	229	264	351	829	891	927	959
Reference											
1959V _{o27} *									< 1.4	3 (2)	2.5 (15)
1961R _{y03}	13.7 (35)£	19.5 (39)£	7.7 (13)£	29.2 (18)£	14.2 (11)£	13.4 (41)£					
1961Vi ₀₇ *									< 0.5		
1961V _{o05} *	18 (3)	3 (1)		19 (3)		8 (1)					
1964Da _{15μ}	7.3 (4)	10.0 (5)	4.5 (3)	23.4 (7)	10.3 (5)	10.6 (7)					
1965Ed _{01&}	20.0 (9)	22.9 (10)	10.7 (6)	56.1 (22)	27.7 (12)	26.9 (12)					
1965He ₀₇									1	1	2
1966Dz ₀₁									~ 0.3	1.74 (26)	0.94 (24)
1966Ko ₁₂											
1969Sa ₂₅	7.5 (8)	8.7 (9)	4.3 (4)	21.2 (21)	10.5 (11)	10.3 (10)			0.20 (7)	1.6 (2)*	1.3 (2)£
1969Wh _{03□}									0.15 (2)	1.79 (9)	1.02 (6)
1970Wh ₀₃	8.07 (40)*	9.57 (5)£	4.40 (25)*	22.6 (12)*	10.9 (5)*	10.6 (4)			0.15 (2)	1.79 (9)	1.02 (6)
1971Ja ₂₁	7.6 (2)	8.8 (3)		21.30 (55)	10.3 (3)	10.1 (3)					
1971Mi ₀₁	7.5 (7)	8.70 (48)	4.20 (24)	21.5 (12)	10.3 (7)	10.0 (6)				1.50 (30)*	1.00 (10)
1972Ga ₂₃	7.13 (48)*	8.7 (6)	4.15 (28)	21.5 (15)	10.3 (7)	10.4 (7)				1.75 (20)	0.95 (11)
1974La ₁₅	7.78 (20)	8.57 (25)	3.75 (12)£	21.26 (62)	9.24 (26)£	9.46 (29)*				2.10 (8)*	1.12 (6)*
1976He _{18\$}									0.164 (19)	1.779 (27)	0.998 (18)
1977Ge ₁₂	7.77 (24)	9.10 (29)	4.31 (14)	21.9 (7)	10.60 (32)	10.50 (32)					
1978MeZK/1990Me ₁₅	7.59 (10)	8.82 (10)	4.19 (9)	21.60 (31)	10.39 (18)	10.26 (18)	0.034 (8)			1.730 (30)	0.980 (30)
1980Ro ₂₂	7.43 (40)	8.88 (66)	4.13 (28)	21.75 (56)	10.39 (34)	10.36 (52)				1.53 (45)*	0.92 (47)
1980Sc ₀₇	7.46 (12)	8.75 (11)	4.09 (6)	21.27 (28)	10.32 (13)	10.26 (16)					
1981Is ₀₈	7.6 (8)	9.1 (7)	4.2 (2)	21.3 (8)	10.2 (6)	9.98 (50)			0.21 (5)	1.64 (10)*	0.87 (8)*
1983Ji ₀₁	7.60 (7)	8.84 (9)	4.22 (6)	21.61 (20)	10.49 (10)	10.37 (7)	0.033 (19)	0.038 (15)	0.16 (5)	1.760 (40)	0.98 (5)
1983Ei _{02£}	7.26 (11)	8.38 (9)	3.91 (4)	21.12 (22)	10.33 (13)	9.81 (12)	0.24 (1)		0.14 (4)	1.85 (3)	1.06 (3)
1986Wa ₃₅	7.26 (18)*	8.85 (24)	4.14 (7)	21.62 (51)	10.24 (22)	9.9 (8)					
1989Ka _{20\$}	7.51 (15)	8.92 (10)	4.23 (4)	21.90 (24)	10.59 (13)	10.32 (12)				1.71 (4)	0.93 (2)
1990Ja ₀₂	7.59 (12)	8.83 (8)	4.12 (5)	21.80 (20)	10.38 (11)	10.14 (9)			0.15 (4)	1.77 (6)	1.01 (3)
1992Su ₀₉	7.89 (16)*	9.04 (18)	4.21 (10)	21.6 (5)	10.5 (2)	10.26 (22)	0.03 (1)	0.05 (2)	0.20 (6)	1.72 (8)	0.99 (8)
1992Ch ₂₆	7.54 (10)	8.93 (12)	4.19 (5)	21.90 (27)	10.43 (13)	10.37 (15)	0.0330 (10)	0.039 (8)	0.160 (10)	1.720 (24)	0.970 (21)
1992Ke ₀₂	7.51 (9)	8.81 (9)	4.12 (6)	21.43 (24)	10.43 (12)	10.43 (14)	0.028 (6)		0.174 (22)	1.75 (7)	0.99 (8)
1998Mi ₁₇	7.570 (46)	8.770 (48)	4.150 (24)	21.17 (12)	10.20 (6)	10.13 (6)					
Evaluated	7.570 (29)	8.811 (29)	4.155 (16)	21.45 (7)	10.334 (36)	10.243 (35)	0.0329 (10)	0.040 (7)	0.162 (7)	1.746 (13)	0.989 (11)
χ^2	0.3	0.4	0.4	0.9	0.7	0.8	0.2	0.1	0.3	0.4	0.2

* Outliers values, based on the Chauvenet's criterion and thus were omitted in the final calculation.

£ Data rejection parameters for deviation from weighted mean (3σ).

Energy (keV)	1001	1035	1044	1113	1121	1135	1157	1158 1180		1189	1221
Reference											
1959Vo27*	9 (3)				100			< 4		45 (8)	84 (8)
1961Ry03					100			4.2 (9)		47.5 (27)	81 (6)*
1961Vi07*	5 (2)		0.9 (8)		100			3.6 (10)		44 (3)	80 (6)
1961Vo05*					100					43	118
1964Da15μ											
1965Ed01&											
1965He07	6 (2)		2 (1)£		100			3		44	72
1966Dz01	5.4 (3)		1.2 (2)*		100			4.1 (12)		44.3 (15)*	77 (6)
1966Ko12	7.9 (26)£		< 1		100		2.67 (15)			48.1 (20)	85.1 (31)£
1969Sa25	5.6 (6)		0.8 (1)	1.2 (2)	100		2.0 (3)	0.76 (16)	0.25 (4)	46.3 (32)	77.3 (54)
1969Wh03□	5.98 (30)		0.69 (8)	1.13 (10)	100		1.84 (35)	0.99 (28)		47.7 (7)	79.3 (12)
1970Wh03	5.98 (30)		0.69 (8)	1.13 (10)	100		1.84 (35)	0.99 (28)		47.7 (7)	79.3 (12)*
1971Ja21					100					46.5 (7)	77.3 (12)
1971Mi01	5.4 (10)		0.60 (10)		100		2.60 (21)			47.2 (21)	78.0 (34)
1972Ga23	5.66 (40)		0.69 (10)	1.44 (20)	100		2.90 (20)		0.28 (4)	46.7 (23)	80.3 (41)*
1974La15	6.43 (11)*			1.11 (7)	100		2.96 (9)			46.1 (15)	78.4 (12)
1976He18\$	5.90 (8)		0.678 (14)	1.276 (19)	100		2.838 (39)		0.249 (15)	46.64 (46)	77.3 (6)
1977Ge12					100		2.850 (49)			46.5 (7)	77.0 (11)
1978MeZK/1990Me15	5.87 (6)			1.320 (30)	100		2.920 (31)			47.1 (8)	77.80 (38)
1980Ro22	5.99 (35)			1.18 (7)	100					47.61 (53)	78.1 (9)
1980Sc07					100					46.59 (46)	77.0 (8)
1981Is08	5.36 (11)		0.58 (10)	2.21 (20)£	100			2.65 (20)	0.56 (7)*	48.8 (17)*	77.9 (27)
1983Ji01	5.85 (10)		0.72 (7)	1.30 (3)	100		1.66 (24)	1.22 (21)	0.210 (40)	46.40 (20)	76.8 (6)
1983El02£	5.99 (6)		0.68 (3)	0.95 (4)	100		2.72 (6)		0.10 (3)	46.90 (45)	78.31 (79)
1986Wa35	5.89 (34)				100		2.92 (34)			47.02 (48)	77.3 (13)
1989Ka20§	5.92 (7)		0.66 (2)	1.15 (3)	100					47.18 (67)	78.38 (81)
1990Ja02	6.01 (5)		0.70 (8)	1.35 (15)	100		2.71 (20)		0.23 (9)	47.37 (9)	77.48 (34)
1992Su09	5.87 (13)	0.017 (6)	0.68 (5)	1.08 (5)	100		2.01 (7)	0.82 (5)	0.23 (4)	46.3 (19)	76.2 (15)
1992Ch26	5.86 (10)		0.660 (21)	1.240 (22)	100		2.830 (46)		0.250 (10)	46.6 (8)	
1992Ke02	5.89 (13)		0.72 (5)	1.19 (7)	100		2.87 (5)		0.22 (6)	47.0 (6)	78.0 (10) ^a
1998Mi17					100		2.930 (22)			46.70 (24)	
Evaluated	5.88 (13)	0.017 (6)	0.677 (10)	1.257 (19)	100		2.37 (36)	0.84 (5)	0.248 (8)	47.13 (9)	77.53 (20)
χ^2	2.2		0.5	2.7			12	1.3	0.3	1.7	0.3

* Outliers values, based on the Chauvenet's criterion and thus were omitted in the final calculation.

£ Data rejection parameters for deviation from weighted mean (3σ).

a Doublet with 1223 keV gamma-ray.

Energy (keV)	1223	1231	1257	1273	1289	1342 1373	1387		1410	1453
Reference										
1959Vo27*		35 (10)	6 (2)	3 (2)	5 (2)		< 1.4			
1961Ry03		29.1 (22)*	5.1 (9)£		< 2.9					
1961Vi07*		25 (5)	4 (1)							
1961Vo05*		118		~ 4	~ 4			~ 2		~ 0.4
1964Da15µ										
1965Ed01&										
1965He07		36	4.5	2	4	0.80 (16)	0.70 (14)			
1966Dz01		26 (5)£	3.8 (3)*	1.5 (3)*	3.7 (2)	0.60 (9)*	0.52 (9)	0.25 (6)	0.115 (17)	0.094 (12)
1966Ko12		28.6 (10)*	3.91 (15)£	1.64 (15)	3.67 (15)	0.79 (5)	0.70 (14)	0.184 (41)	0.13 (6)	0.14 (6)
1969Sa25	0.6 (1)	32.7 (23)	4.3 (3)	1.8 (1)	3.8 (3)	0.7 (1)	0.6 (1)	0.18 (2)	0.11 (2)	0.12 (2)
1969Wh03□		33.4 (5)	4.33 (7)	1.90 (4)	4.05 (7)	0.75 (2)	0.66 (2)	0.217 (10)	0.117 (8)	0.123 (10)
1970Wh03		33.4 (5)	4.33 (7)	1.90 (4)	4.05 (7)	0.75 (2)	0.66 (2)	0.217 (10)	0.117 (8)	0.123 (10)
1971Ja21		32.8 (5)								
1971Mi01		32.3 (14)	4.27 (19)	1.92 (10)	4.06 (19)	0.750 (38)	0.690 (36)	0.240 (21)	0.140 (30)	0.120 (30)
1972Ga23		34.5 (25)*	4.46 (45)*	1.96 (19)	4.10 (40)	0.80 (9)	0.70 (8)	0.225 (23)	0.130 (25)	0.10 (2)
1974La15		32.60 (52)	4.31 (8)	1.83 (5)	3.96 (8)	0.74 (3)	0.65 (3)	0.21 (1)		
1976He18§		32.92 (30)	4.269 (46)	1.864 (28)	3.87 (5)	0.720 (11)	0.628 (9)	0.2019 (39)	0.1152 (46)	0.0804 (32)
1977Ge12	0.778 (11)	32.96 (47)	4.26 (6)	1.860 (27)	3.86 (6)	0.718 (12)	0.628 (11)	0.202 (5)	0.112 (6)	0.0790 (31)
1978MeZK/1990Me15	0.30 (10)*	33.1 (5)	4.36 (8)	1.950 (31)	4.29 (8)	0.740 (10)	0.680 (10)	0.270 (10)*	0.1170 (40)	0.123 (8)
1980Ro22		32.32 (56)	4.33 (15)	1.66 (18)	4.06 (22)					
1980Sc07		32.81 (33)	4.250 (36)		3.860 (34)					
1981Is08		32.3 (11)	4.07 (30)*	1.67 (15)	3.65 (20)	0.66 (8)	0.58 (8)	0.20 (5)	0.10 (4)	0.10 (4)
1983Ji01	0.53 (24)	32.72 (14)	4.276 (24)	1.871 (13)	3.800 (32)	0.723 (7)	0.626 (6)	0.2040 (40)	0.111 (5)	0.0872 (24)
1983El02£		33.20 (22)	4.27 (2)	1.87 (1)	3.89 (4)	0.72 (1)	0.60 (1)	0.20 (1)	0.10 (1)	0.09
1986Wa35		33.42 (31)	4.36 (16)	1.73 (12)	4.17 (30)					
1989Ka20§		33.28 (45)	4.40 (6)	1.92 (3)	4.01 (5)	0.75 (1)	0.68 (2)			
1990Ja02	0.55 (12)	33.85 (22)	4.35 (6)	1.90 (4)	3.90 (5)	0.76 (4)	0.65 (2)	0.24 (3)	0.14 (2)	0.11 (2)
1992Su09	0.58 (10)	32.2 (7)	4.22 (9)	1.84 (5)	3.80 (8)	0.69 (3)	0.55 (2)	0.19 (2)	0.083 (10)*	0.11 (1)
1992Ch26		32.80 (48)	4.31 (7)	1.850 (33)	3.91 (6)	0.720 (12)	0.610 (11)	0.205 (6)	0.1090 (41)	0.0830 (31)
1992Ke02	78.0 (10) ^a	33.17 (37)	4.34 (7)	1.860 (31)	4.03 (7)	0.748 (21)	0.628 (13)	0.220 (11)	0.117 (8)	0.097 (7)
1998Mi17		33.18 (19)	4.320 (24)		3.940 (23)					
Evaluated	0.58 (6)	33.04 (10)	4.296 (13)	1.872 (9)	3.907 (33)	0.7284 (43)	0.633 (7)	0.2060 (24)	0.1136 (21)	0.106 (19)
χ^2	0.05	1.7	0.5	1.1	3.0	0.8	3.2	0.9	0.5	4.3

* Outliers values, based on the Chauvenet's criterion and thus were omitted in the final calculation.

£ Data rejection parameters for deviation from weighted mean (3σ).

a Doublet with 1221 keV gamma-ray.

Table 9: Recommended relative and absolute γ -ray intensities (%)

E_{γ} (keV)	Relative γ -ray intensity (%)	Absolute γ -ray intensity (%)	E_{γ} (keV)	Relative γ -ray intensity (%)	Absolute γ -ray intensity (%)	E_{γ} (keV)	Relative γ -ray intensity (%)	Absolute γ -ray intensity (%)
31	2.38 (17)	0.84 (6)	229	10.334 (36)	3.634 (36)	1180	0.247 (8)	0.0869 (29)
42	0.765 (18)	0.269 (7)	264	10.243 (35)	3.602 (36)	1189	47.13 (9)	16.58 (16)
65	8.45 (20)	2.97 (8)	351	0.0329 (10)	0.01157 (37)	1221	77.53 (20)	27.27 (27)
67	124.0 (40)	43.6 (15)	829	0.040 (7)	0.0141 (25)	1223	0.58 (6)	0.204 (21)
84	7.45 (14)	2.62 (6)	891	0.162 (7)	0.0570 (25)	1231	33.04 (11)	11.62 (12)
100	40.42 (24)	14.22 (16)	927	1.746 (13)	0.614 (7)	1257	4.296 (13)	1.511 (15)
110	0.305 (9)	0.1073 (33)	959	0.989 (11)	0.348 (5)	1273	1.872 (9)	0.658 (7)
113	5.315 (29)	1.869 (20)	1001	5.88 (13)	2.07 (5)	1289	3.906 (33)	1.374 (17)
116	1.264 (10)	0.445 (5)	1035	0.017 (6)	0.0060 (21)	1342	0.7284 (43)	0.2562 (28)
(121)*		0.0021 (7)	1044	0.677 (10)	0.2381 (42)	1373	0.633 (7)	0.2226 (32)
152	19.93 (33)	7.01 (13)	1113	1.257 (19)	0.442 (8)	1387	0.2060 (24)	0.0725 (11)
156	7.570 (29)	2.662 (27)	1121	100	35.17 (33)	1410	0.1136 (21)	0.0400 (8)
179	8.811 (29)	3.099 (31)	1135			1453	0.106 (19)	0.037 (7)
198	4.155 (16)	1.461 (15)	1157	2.37 (36)	0.83 (13)			
222	21.45 (7)	7.54 (7)	1158	0.84 (5)	0.295 (18)			

*Deduced from gamma-ray probability imbalance at level 4 (1135 keV) of the decay scheme.

6 References

- 1943Zu** R. V. Zumstein, J. D. Kurbatov, M. L. Pool, Phys. Res. 63(1943)59 [Half-life].
- 1947Se33 L. Seren, H. N. Friedlander, S. H. Turkel, Phys. Rev. 72(1947)888 [Half-life].
- 1948Me** L. Meitner, Ann. Phys. 6(1948)113 [Half-life].
- 1951Si25 W. K. Sinclair, A. F. Holloway, Nature 167(1951)365 [Half-life].
- 1952Ei12 G. G. Eichholz, L. A. Ficko, Phys. Rev. 86(1952)794 [Half-life].
- 1957Wr37 H. W. Wright, E. I. Wyatt, S. A. Reynolds, W. S. Lyon, T. H. Handley, Nucl. Sci. Eng. 2(1957)427 [Half-life].
- 1958Ke26 J. P. Keene, L. A. Mackenzie, C. W. Gilbert, Phys. Med. Biol. 2(1958)360 [Half-life].
- 1958Sp17 A. Speecke, J. Hoste, Bull. Soc. Chim. Belges 67(1958)131 [Half-life].
- 1959Vo27 N. A. Voinova, B. S. Dzhelepov, N. N. Zhukovskii, Bull. Acad. Sci. USSR, Phys. Ser. 23(1959)822 [I_γ].
- 1960Gr** V. S. Grodzev, L. I. Rusinov, Y. L. Khazov, Bull. Acad. Sci. USSR, Phys. Ser. 24(1960)1439 [α_K].
- 1961Ry03 H. Ryde, Z. Sujkowski, Ark. Fys. 20(1960)289 [I_γ].
- 1961Vi07 V. D. Vitman, N. A. Voinova, B. S. Dzhelepov, A. A. Karan, Bull. Acad. Sci. USSR, Phys. Ser. 25(1962)192 [I_γ].
- 1961Vo05 N. A. Voinova, B. S. Dzhelepov, Yu. V. Khol'nov, Bull. Acad. Sci. USSR, Phys. Ser. 25(1962)223 [I_γ].
- 1964Da15 H. Daniel, J. Huefner, Th. Lorenz, O. W. B. Schult, U. Gruber, Nucl. Phys. 56(1964)147 [I_γ].
- 1964Ba12 E. Bashandy, A. H. El-Farrash, M. S. El-Nesr, Nucl. Phys. 52(1964)61 [δ].
- 1964Ba47 V. A. Balalaev, N. A. Voinova, B. S. Dzhelepov, A. Meshter, S. A. Shestopalova, Bull. Acad. Sci. USSR, Phys. Ser. 28(1964)1596 [I_γ , I_{CE}].
- 1965He07 R. Henck, L. Stab, P. Siffert, A. Coche, Compt. Rend. Acad. Sci. (Paris) 260(1965)4991 [I_γ].
- 1965Ed01 W. F. Edwards, F. Boehm, J. Rogers, E. J. Seppi, Nucl. Phys. 63(1965)97 [I_γ].
- 1966Dz01 B. S. Dzhelepov, V. D. Vitman, Nucl. Phys. 75(1966)371 [I_γ].
- 1966Gr21 E. P. Grigorev, A. V. Zolotavin, V. O. Sergeev, V. S. Bekrenev, Yadern. Fiz. 4(1966)9; Soviet J. Nucl. Phys. 4(1967)5 [δ].
- 1966Ko12 K. Korkman, A. Bäcklin, Nucl. Phys. 82(1966)561 [I_γ].
- 1967Ba01 V. A. Balalaev, B. S. Dzhelepov, L. N. Moskvina, S. A. Shestopalova, N. A. Voinova, Nucl. Phys. A91(1967)465 [P_β].
- 1967Ni03 O. Nilsson, S. Hogberg, S. -E. Karlsson, G. M. El-Sayad, Nucl. Phys. A100(1967)351 [δ , α].
- 1967Wa29 D. A. Walker, Nucl. Instrum. Meth. 48(1967)277 [Half-life].
- 1969Ga23 P. Galan, T. Galanova, Z. Malek, N. Voinova, Z. Preibisz, K. Stryczniewicz, Nucl. Phys. A136(1969)673 [α].
- 1969Wh03 D. H. White, R. E. Birkett, Nucl. Phys. A136(1969)657 [I_γ].
- 1969Sa15 J. J. Sapyta, E. G. Funk, J. W. Mihelich, Nucl. Phys. A139(1969)161 [I_γ].
- 1970Wh03 D. H. White, R. E. Birkett, T. Thomson, Nucl. Instrum. Meth. 77(1970)261 [I_γ].
- 1971Ja21 L. J. Jardine, Nucl. Instrum. Meth. 96(1971)259 [I_γ].
- 1971Ml01 M. Mladenovic, M. Ninkovic, M. Stojanovic, Fizika 3(1971)01 [I_γ].
- 1972Em01 J. F. Emery, E. I. Wyatt, S. A. Reynolds, Nucl. Sci. Eng. 48(1972)319 [Half-life].
- 1972Ga23 P. Galan, M. Vejs, Fyz. Cas. 22(1972)60 [I_γ].
- 1972He10 P. Herzog, M. J. Canty, K. D. Killig, Nucl. Phys. A187(1972)49 [δ].
- 1972Kr05 K. S. Krane, J. R. Sites, W. A. Steyert, Phys. Rev. C5(1972)1104 [δ].
- 1973Vi13 C. J. Visser, J. H. M. Karsten, F. J. Haasbroek, P. G. Marais, Agrochemophysica 5(1973)15 [Half-life].
- 1974La15 N. Lavi, Nucl. Instrum. Meth. 116(1974)457 [I_γ].
- 1975Qu01 L. M. Quinones, Z. W. Grabowski, Nucl. Phys. A242(1975)243 [δ].
- 1975We22 L. Westerberg, L. O. Edvardson, G. C. Madueme, Nucl. Phys. A255(1975)427 [α_K].
- 1976He18 R. G. Helmer, Nucl. Phys. A272(1976)269 [I_γ].
- 1977Ge12 R. J. Gehrke, R. G. Helmer, R. C. Greenwood, Nucl. Instrum. Meth. 147(1977)405 [I_γ].
- 1978MeZK R. A. Meyer, LLNL M-100(1978) [I_γ].
- 1980Sc07 U. Schötzig, K. Debertain, K. F. Walz, Nucl. Instrum. Meth. 169(1980)43 [Half-life, I_γ].

- 1980Sp01 R. Spanhoff, M. J. Canty, H. Postma, G. Mennenga, Phys. Rev. C21(1980)361 [δ].
- 1980Ro22 W. N. Roney, W. A. Seale, Nucl. Instrum. Meth. 171(1980)389 [I_γ].
- 1981Is08 H. A. Ismail, M. Morsy, H. Hanafi, S. Abdle-Malak, H. El-Samman, Rev. Roum. Phys. 26(1981)455 [I_γ].
- 1981Ka22 R. Kaur, A. K. Sharma, S. S. Sooch, H. R. Verma, P. N. Trehan, Indian J. Pure Appl. Phys. 19(1981)133 [δ].
- 1983Ji01 J. Jin, J. Takada, Y. Iwata, Y. Yoshizawa, Nucl. Instrum. Meth. 212(1983)259 [I_γ].
- 1983El02 M. S. S. El-Daghmah, N. M. Stewart, Z. Phys. A309(1983)219 [I_γ].
- 1983Ri05 J. Rikovska, D. Novakova, J. Ferencei, M. Finger, Z. Phys. A311(1983)185 [δ].
- 1986Wa35 W. Xinlin, Li Xiaodi, Du Hongshan, Chin. J. Nucl. Phys. 8(1986)371 [I_γ].
- 1988Fi05 R. B. Firestone, Nuclear Data Sheets 54(1988)307 [Spin, parity, energy levels, multipolarity].
- 1989Ka20 R. Kaur, P. N. Trehan, Appl. Rad. Isotopes 40(1989)727 [I_γ].
- 1990Ja02 J. K. Jabber, N. M. Stewart, J. Phys. (London) G16(1990)271 [I_γ].
- 1990Me15 R. A. Meyer, Fizika (Zagreb) 22(1990)153 [I_γ].
- 1992Su09 Sun Huibin, Liu Yunzou, Zhou Jiewen, Wu Yaodong, Z. Phys. A342(1992)141 [I_γ].
- 1992Ch26 B. Chand, J. Goswamy, D. Mehta, N. Singh, P. N. Trehan, Can. J. Phys. 70(1992)242 [I_γ].
- 1992Ke02 T. Kempisty, K. Pochwalski, Nucl. Instrum. Meth. Phys. Res. A312(1992)390 [I_γ].
- 1996Sc06 E. Schönfeld, H. Janssen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
- 1998Mi17 H. Miyahara, H. Nagata, T. Furusawa, N. Murakami, C. Mori, N. Takeuchi, T. Genka, Appl. Rad. Isotopes 49(1998)1383 [I_γ].
- 2000He14 R. G. Helmer, C. van der Leun, Nucl. Instrum. Meth. Phys. Res. A450(2000)35 [E_γ].
- 2002Ba85 I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr., P. O. Tikkanen, S. Raman, Atomic Data Nucl. Data Tables 81(2002)1 [Theoretical ICC].
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)1 [Q].
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr. , Nucl. Instrum. Meth. Phys. Res. A589(2008)202 [Theoretical ICC].
- 2010Si13 B. Singh, J. C. Roediger, Nucl. Data Sheets 111(2010)2081 [Spin, parity, energy levels, multipolarity].

¹⁸⁶Re - Comments on evaluation of decay data by E. Schönfeld and R. Dersch

This evaluation was completed in November 1998 and the half-life value has been updated in May 2004.

1 Decay Scheme

The decay scheme is taken from Baglin (1997). It is based mainly on the work of Fogelberg (1972), Seegmiller et al. (1972) and Maly et al. (1964). The latter two authors did not only study gammas, but also conversion electrons. There are EC branches to the 122 keV level and the ground state of ¹⁸⁶W (together 7,53 %) and beta branches to the ground state (70,9 %) and the excited states (21,5 %) in ¹⁸⁶Os. Spins and parities of the levels are taken from Baglin (1997), also the half-lives of the excited levels in ¹⁸⁶Os. The splitting into the EC and the beta part was calculated from the measured total W K-X ray emission probability. Beside the four excited levels of ¹⁸⁶Os given in the decay scheme, there is a level at 868,94(4) keV (6+). A direct beta transition to this level would be fifth forbidden and, therefore, would be too weak to be observed. The next higher level in ¹⁸⁶Os is at 1070,5 keV which is already above the adopted Q_{β} -value if the latter is correct.

¹⁸⁶W has below the Q_{EC} value a further level at 396,26 keV (4+; 36 ps). An EC transition to this level would be third forbidden, so this branch will be very weak, thus the decay scheme given on page 1 can be considered to be complete.

2 Nuclear Data

The following values of the half-life have been considered ($T_{1/2}$ in d):

1	3,750	Sinma et al. (1939); Fajans et al. (1940); Chu (1950)
2	3,792	Cork et al. (1940); Grant <i>et al.</i> (1945); Dybvig <i>et al.</i> (1950)
3	3,867(8)	Yamasaki et al. (1940)
4	3,867(8)	Goodman and Pool (1947)
5	3,704(8)	Porter et al. (1956)
6	3,775(13)	Gueben and Govaerts (1958)
7	3,777(4)	Michel and Herpers (1971)
8	3,775(1)	Abzouzi et al. (1989)
9	3,7187(29)	Unterweger et al. (1992)
10	3,7183(11)	Schönfeld et al. (1994) ; superseded by 11
11	3,7186(5)	Schrader (2004)
12	3,7186(17)	by the present evaluator adopted value

The adopted value is mainly based on values 9 and 11. The values 1 to 4 are considered to be only of historical interest. The remaining six values are discrepant: there is a group of three low values (5, 9, 11) and three high values (6, 7, 8). If values 6, 7 and 8 would be included in an averaging procedure, the mean value would be larger than value 12 and also its uncertainty. The present evaluator has not included values 6, 7 and 8 into the averaging procedure because of the well agreeing values 9, 10 and 11 which were measured in well equipped national institutes by experienced scientists whereas the consideration of radioactive impurities and other systematical uncertainties is not convincing in the papers 7 and 8. The value 10 is superseded by value 11 and was then not used for the mean.

Both Q values are taken from Audi and Wapstra (1995).

2.1 β^- Transitions

The maximum beta energy of the transition to the 137 keV level have been measured to be (values in keV)

1	934,3(13)	Porter et al. (1956)
2	927(2)	Johns et al. (1956)
3	937(14)	Bashandi and El Nesr (1963)
4	939(3)	Maly et al. (1964)
5	927(3)	Andre and Liaut (1968)
6	945(5)	Trudel et al. (1970)
7	932,8(21)	weighted mean

By adding the level energy of 137,1 keV to the weighted mean we obtain 1069,9 keV for the Q value which is in good agreement with the value given for Q_{β^-} by Audi and Wapstra: 1069,5(9) keV.

The energy of the $\beta_{0,1}$ transition in table 2.1 is deduced from the adopted Q_{β} value and the gamma ray energy. The spectra of the β transitions to the ground state and to the 137 keV level which are both non - unique first forbidden were found to have an almost allowed shapes. The total beta emission probability is calculated by subtracting the total EC probability (Section 2.2) from 1.

2.2 Electron Capture Transitions

The fractional capture probabilities of the transitions $\epsilon_{0,1}$ and $\epsilon_{0,0}$ were calculated using the data of Schönfeld (1998). The energies are derived from the Q values and the level energies. From the emission probability of the 122 keV γ ray (which was found to be 0,00603(6); original value of Schönfeld et al., 1994) and the conversion coefficient of this transition, the transition probability $P_{\gamma+ce}$ (which is also the transition probability of the electron capture branch to the 122 keV level) is obtained to be $P_{\gamma+ce} = P_{EC}(0,1) = 0,0169(3)$.

The transition probability of the electron capture transition feeding the ground state of ^{186}W can be calculated from the total emission probability of W KX rays. This emission probability is given by

$$P(\text{W KX}) = \left\{ P_{EC}(0,1) \left[P_K(0,1) + a_K / (1 + a_t) \right] + P_{EC}(0,0) P_K(0,0) \right\} w_K.$$

Using the known values for P_K (Table 2.2), the conversion coefficients a_K and a_t (Table 2.3), and the fluorescence yield w_K for tungsten, the transition probability $P_{EC}(0,0)$ can be extracted from the above expression. Using $P(\text{W KX}) = 0,0602(8)$ as determined by Schönfeld et al. (1994), one obtains $P_{EC}(0,0) = 0,0584(12)$.

Thus, the total electron capture probability amounts to $P_{EC}(0,1) + P_{EC}(0,0) = 0,0169(3) + 0,0584(12) = 0,0753(12)$.

2.3 Gamma Transitions

Concerning the energies see Sect. 4.2. The transition probabilities $P_{\gamma+ce}$ are calculated from the emission probabilities (Sect.4.2) and the total conversion coefficients. The conversion coefficients were interpolated from the tables of Rösel et al. (1978). Maly et al. have determined the K conversion coefficients as follows: $\alpha_K(122 \text{ keV}) = 0,53(5)$, $\alpha_K(137 \text{ keV}) = 0,44(2)$. Both are pure E2 transitions.

These values are in agreement with the theoretical ones. Maly et al. have also determined the ratios K/L/M/N for these two transitions. Mixing ratios for the transitions $\gamma_{4,3}$, $\gamma_{4,2}$, $\gamma_{3,1}$ and $\gamma_{4,1}$ were taken from Baglin (1997).

3 Atomic Data

The atomic data are taken from Schönfeld and Janßen (1996).

3.1 X Radiation

The energy values are calculated from the wave lengths in Å* as given by Bearden (1967). The relative emission probabilities of K X rays are taken from Schönfeld and Janßen (1996). The relative emission probabilities of L X rays is calculated from the absolute emission probability given in Table 4.2 setting $P(K_{a_1}) = 1$.

3.2 Auger Electrons

The energy values are taken from Larkins (1977) (KLL) and the Table de Radionucl éides (LMRI 1982) (KLX, KXY). The relative emission probabilities of K Auger electrons are taken from Schönfeld and Janßen (1996). The relative emission probabilities of the L Auger electrons is calculate d from the value in the table 4.1 putting $P(KLL) = 1$.

4 Radiation Emission

4.1 Electron Emission

The energies of the Auger electrons are the same as in 3.2. The energies of the conversion electrons are calculated from the transition energy (2.2) and the binding energies.

The emission probabilities of the conversion electrons are calculated using the conversion coefficients given in 2.2. The values of the emission probabilities of the Auger electrons are calculated using the transition probabilities given in 2.1 and 2.2, the atomic data given in 3, and the conversion coefficients given in 2.2 using the Programm EMISSION.

4.2 Photon Emission

The energy of the X rays are from 3.1. The energy of the 137 keV gamma rays was determined by Marklund and Lindström (1963) using a curved-crystal spectrometer. The energies of the other γ rays are taken from Baglin (1997) who took into account also coulomb excitation and n, γ reactions.

The emission probability (photons per disintegration) of the 137 keV γ rays in ¹⁸⁶Os has been determined to be 0,0945(16) by Coursey et al. (1991) and 0,0939(9) by Schönfeld et al. (1994). Together with Baglin (1997) we take the unweighted mean 0,0942(6) as adopted value in the present evaluation in order to compare the results of different authors who carried out relative measurements. Then we have (normalized to this value) the following emission probabilities:

	1	2	3	4	5
W L X	0,0308(?)	-	-	0,0192(2)	0,0166(4)
W K _{a2}	0,0178(4)	-	0,0172(5)	0,0176(4)	0,01736(30)
W K _{a1}	0,0312(4)	-	0,0297(8)	0,0303(6)	0,0302(5)
W K _a	0,0490(6)	0,0445(13)	0,0469(10)	0,0479(8)	0,0475(8)
W K' _{b1}	0,0109(2)	-	0,0099(4)	0,00989(20)	0,01000(23)
W K' _{b2}	0,0034(2)	-	0,0026(2)	0,00269(6)	0,00274(8)
W K _b	0,0143(3)	-	0,0125(4)	0,01258(21)	0,1273(29)
W K X	0,0633(7)	-	0,0594(11)	0,0605(8)	0,0603(10)
Os L X	0,0300(3)	-	-	0,0306(34)	0,0299(7)
Os K _{a2}	0,0114(2)	-	0,0113(4)	0,0112(3)	0,01128(26)
Os K _{a1}	0,0199(4)	-	0,0193(6)	0,0196(4)	0,0194(5)
Os K _a	0,0313(5)	0,0286(6)	0,0306(7)	0,0308(5)	0,0307(7)
Os K' _{b1}	0,0067(2)	-	0,0066(3)	0,00635(14)	0,00650(18)
Os K' _{b2}	0,00198(20)	-	0,00170(6)	0,00186(4)	0,00182(6)
Os K _b	0,0087(2)	-	0,0083(3)	0,00821(15)	0,00833(23)
Os K X	0,0400(6)	-	0,0389(7)	0,0390(5)	0,0390(9)
W γ 122	0,00603(20)	0,00598(10)	0,00604(23)	0,00605(6)	0,00603(6)
Os γ 137	≅0,0942(6)	≅0,0942(6)	≅0,0942(6)	≅0,0942(6)	≅0,0942(6)
Os γ 630	-	0,00032(3)	0,000292(6)	0,000294(6)	0,000293(6)
Os γ 767	-	0,00037(4)	0,000324(7)	0,000328(6)	0,000327(6)

- 1 Seegmiller et al. (1972)
 2 Coursey et al. (1991)
 3 Goswamy et al. (1991)
 4 Schönfeld et al. (1994)
 5 calculated with EMISSION (X rays); values adopted by the present evaluator (gammas)

In all cases there is excellent agreement. Relative values for the emission probabilities of the gamma rays were also determined by Johns et al. (1956), Maly et al. (1964) and Rao et al. (1969). These values are less accurate and were not taken into account in the present evaluation. The emission probabilities and the energies of the gamma rays of the very weak gamma transitions in ¹⁸⁶Os (not contained in the above table) were determined by Fogelberg (1972) which is the only one to report these values.

Multiplying the adopted value for $P_{\gamma}(122)$ by $1 + a_{\gamma}(122)$ we obtain, in agreement with table 2.2, $P_{EC}(122) = 0,0169$.

Values, recently measured by Miyahara *et al.* (2000) and Woods *et al.* (2000) are also in good agreement with the here adopted values.

5 Main Production Modes

Taken from the „Table des Radionucléides“, LMRI, 1982.

6 References

- K. Sinma and H. Yamasaki, *Phys. Rev.* 55 (1939) 320
 $[T_{1/2}]$
- K. Fajans and W. H. Sullivan, *Phys. Rev.* 58 (1940) 276
 $[T_{1/2}]$
- J. M. Cork, R. G. Shreffler and C. M. Fowler, *Phys. Rev.* 74 (1940) 1657
 $[T_{1/2}]$
- H. Yamasaki and K. Sinma, *Sci. Pap. Inst. Phys. Chem. Res. Tokyo* 37 (1940) 10
 $[T_{1/2}]$
- P. J. Grant and R. Richmond, *Proc. Phys. Soc.* 62A (1945) 573
 $[T_{1/2}]$
- L. J. Goodman and M. L. Pool, *Phys. Rev.* 71 (1947) 288
 $[T_{1/2}]$
- A. T. Dybvig and M. L. Pool, *Phys. Rev.* 80 (1950) 126
 $[T_{1/2}]$
- T. C. Chu, *Phys. Rev.* 79 (1950) 582
 $[T_{1/2}]$
- D. Guss, L. Killion, F. T. Porter, *Phys. Rev.* 95 (1954) 627
 $[E, \lg ft]$
- M. W. Johns, C. C. Mac Mullen, I. R. Williams, S. V. Nablo, *Can. J. Phys.* 34 (1956) 69
 $[P_{\beta}]$
- F. T. Porter, M. S. Freedman, T. B. Novey, F. Wagner jr., *Phys. Rev.* 103 (1956) 921
 $[E_{\gamma}, P_{\beta}]$
- T. B. Novey, M. S. Freedman, F. T. Porter, F. Wagner, Jr., *Phys. Rev.* 103 (1956) 942
 $[E_{\beta}, E_{\gamma}, \text{angular correlation}]$
- E. L. Chupp, A. F. Clark, J. W. M. Dumond, F. J. Gordon, H. Mark, *Phys. Rev.* 107 (1957) 745
 $[E_{\gamma}]$
- G. Gueben and J. Govaerts, *Monographie* Nr. 2, Inst. Interuniv. Sciences Nucleares, Bruxelles (1958)
 $[T_{1/2}]$
- G. T. Emery, W. R. Kane, M. Mac Keown, M. L. Perlman, G. Scharff-Goldhaber,
Phys. Rev. 129 (1963) 2597
 $[E_{\gamma}, P_{\gamma}]$
- B. Harmatz, T. H. Handley, *Nucl. Phys.* 56 (1964) 1
 $[E_{\gamma}]$
- R. Michel, U. Herpers, *Radiochim. Acta* 16 (1971) 115
 $[T_{1/2}]$
- A. Abzouzi, M S. Antony, V. B. Ndocko Ndongue, *J. Radioanal. Cl. Chem. Letters* 137 (1989) 381
 $[T_{1/2}]$

For additional references see also § References in the Tables Part.

¹⁹⁵Au - Comments on evaluation of decay data by V. Chisté and M. M. Bé

This evaluation was completed in June 2012, including all publications by this date.

1 Decay Scheme

¹⁹⁵Au disintegrates 100 % by electron-capture transitions to the ground state level and excited levels of ¹⁹⁵Pt. Good agreement is found between the effective Q^+ value (227 (5) keV) calculated from the decay scheme data and that recommended (226.8 (10) keV) from the atomic mass evaluation of Audi and Meng (2012Au06).

2 Nuclear Data

The Q^+ value is from the atomic mass evaluation of Audi *et al.* (2011AuZZ).

The recommended ¹⁹⁵Au half-life has been deduced from the experimental values (in days) given in Table 1:

Table 1: Experimental values of ¹⁹⁵Au half-life.

Reference	Experimental value (d)	Comments
R. M. Steffen (1949St17)	180 (15)	
G. Wilkinson (1949Wi08)	185 (3)	
A. Bisi (1959Bi07)	192 (5)	Omitted, outlier.
M. Bresesti (1960Br11)	199 (3)	Omitted, outlier.
N. A. Bonner (1962Bo12)	185 (1)	
G. Harbottle (1963Ha17)	182.9 (5)	
D. D. Hoppes (1982HoZJ)	186.09 (4)	Omitted, superseded by 2002Un02.
M. P. Unterweger (2002Un02)	186.098 (47)	
Recommended value	184.7 (14)	$\chi^2 = 6$

A weighted average was calculated by using LWEIGHT computer program (version 3). The Bisi (1959Bi07) and Bresesti (1960Br11) values were showed to be outliers, based on the Chauvenet's criterion, and thus were omitted in the final calculation. The largest contribution to the weighted average comes from the value of Unterweger (2002Un02), with a relative statistical weight of 99 %. The LWEIGHT program increased the uncertainty of the 2002Un02 value from 0.047 to 0.44 in order to reduce its relative statistical weight to 50 %.

The recommended value of ¹⁹⁵Au half-life is the weighted average of 184.7 d with a final uncertainty of 1.4 d, expanded to include the most precise value of M. P. Unterweger. The reduced- χ^2 value is 6.

2.1 Electron capture transition

The energies of the electron-capture transitions in the decay of ¹⁹⁵Au \rightarrow ¹⁹⁵Pt have been obtained from the Q^+ value (2011AuZZ) and the level energies given in Table 2 from C. Zhou (1999Zh11).

Table 2: ^{195}Pt levels populated in the decay of ^{195}Au and the evaluated electron-capture transition probabilities.

Level Number	Level energy, (keV)	Spin and Parity ^a	Evaluated P_{ec} (%)
0	0	1/2	9.5 (4)
1	98.882 (4)	3/2 ⁻	57.6 (35)
2	129.777 (5)	5/2 ⁻	32.8 (30)
3	199.526 (12)	3/2 ⁻	0.0149 (14)
4	211.398 (6)	3/2 ⁻	0.0210 (18)

^a Given by C. Zhou (1999Zh11).

For the ^{195}Pt ground state, the adopted electron-capture transition probability of 9.5 (4) % is from S. C. Govere (1973Go05).

The electron-capture transition probabilities to the ^{195}Pt excited levels and the associated uncertainties (Table 2) were deduced from the γ transition probability balance at each level of the decay scheme.

The partial electron-capture transition probabilities P_K , P_L , P_{MNO} and log ft values were calculated for the 1st forbidden and 1st forbidden unique electron-capture transitions using the LOGFT computer code.

2.2 γ Transitions

The γ transition probabilities were obtained using the γ -ray emission intensities and the relevant internal conversion coefficients (see **5.2 Gamma Emissions**).

For all γ transitions, the internal conversion coefficients (ICC) and the associated uncertainties were interpolated from theoretical values of I. M. Band *et al.* (2002Ba85) using the BrIcc computer program (2008Ki07) for the “frozen orbital” approximation.

For multipolarity and mixing ratio of the γ -ray transitions, the evaluators used:

1) The multipolarities of the 129-, 199- and 211-keV γ transitions are from C. Zhou (1999Zh11):

129-keV γ -ray: E2;

199-keV γ -ray: M1 + E2, $|\delta| = 1.2$ (2);

211-keV γ -ray: M1 + E2, $|\delta| = 0.38$ (3).

2) For the 30- and 98-keV γ transitions (M1 + E2), the mixing ratios (δ) were calculated from experimental ICC's (α), using BrIccMixing program, version 2.2a (the same package of BrIcc computer program, <http://bricc.anu.edu.au/index.php>) and the adopted values of δ are shown in the table 3.

Table 3: Experimental ICC's (α) and adopted mixing ratios (δ).

E_γ (keV)	Experimental α	Adopted mixing ratio (δ)
30.895 (7)	$\alpha_L = 30.2$ (39); $\alpha_M = 6.9$ (9) (1969Fi08) $\alpha_{L1} = 17.9$ (46); $\alpha_{L2} = 1.40$ (64); $\alpha_{L3} = 0.25$ (8) (1970To19) $\alpha_{L1} = 23.0$ (28); $\alpha_{L2} = 2.50$ (30); $\alpha_{L3} = 0.43$ (5) (1970Ah05)	- 0.013 (7), $\chi^2 = 1.7$
98.882 (4)	$\alpha_K = 8.4$ (5) (1959Bi07) $\alpha_K = 5.8$ (15) (1959Mc69) $\alpha_K = 6.01$ (15) (1964Go19) $\alpha_K = 5.8$ (5); $\alpha_L = 0.82$ (7); $\alpha_M = 0.186$ (15) (1969Fi08) $\alpha_K = 6.9$ (15); $\alpha_{L1} = 0.92$ (20); $\alpha_{L2} = 0.088$ (17); $\alpha_{L3} = 0.027$ (6) (1970To19) $\alpha_K = 5.6$ (7); $\alpha_{L1} = 0.870$ (36); $\alpha_{L2} = 0.119$ (8); $\alpha_{L3} = 0.033$ (3) (1970Ah05)	-0.122 (+14,-13), $\chi^2 = 3.3$

3 Atomic Data

Atomic values, ω_K , ω_L and n_{KL} are from Schönfeld and Janßen (1996Sc06).

The X-ray and Auger electron emission probabilities were calculated from the data set values using the computer program EMISSION.

4 Electron emissions

The conversion electron emission probabilities were deduced from the ICC values and the γ -ray emission intensities.

5 Photon Emissions

5.1 X-rays

The X-ray absolute intensities were deduced from the decay data using the EMISSION computer code and are compared in Table 4 with measured values found in the literature. A reasonable agreement has been found between the experimental and calculated values.

Table 4: Experimental and recommended (calculated) values of the total K X-ray absolute intensities.

	1964Go19 *	1967Sc18 *	1968Ja11 [*]	1970Ah05 *	1972Ha21	Recommended
K X-rays	92.5	99 (13)	87.2	98 (7)	99 (5)	94.6 (35)

*Using normalization factor of 0.1121 (15) (see **5.2 Gamma Emissions**)

5.2 Gamma emissions

The γ -ray energies given in section 5.2 were deduced from the decay scheme using the ¹⁹⁵Pt level energies adopted by C. Zhou (1999Zh11).

The experimental relative γ -ray emission probabilities in ¹⁹⁵Au decay were obtained by averaging all the available measured values. The normalization factor to convert relative γ -ray emission probabilities to absolute values was calculated with the formula:

$$\text{Normalization} = \frac{100 - P_{ec}(g.s.)}{\sum(1 + \alpha_T)P_{rel}} = 0.1121 (15)$$

where the sum is to be done over all the gamma transitions populated the ground state, P_{rel} is a relative γ -ray emission probability and $P_{ec}(g.s.) = 9.5 (4) \%$, given by S. C. Govere (1973Go05). From the theoretical total ICC α_T and the evaluated relative γ -ray emission probabilities (Table 5), the calculated normalization factor is 0.1121 (15).

The experimental γ -ray emission probabilities relative to the 98-keV γ -ray taken equal to 100 are given in Table 5.

The evaluated relative γ -ray emission probability values are the weighted means calculated with the LWEIGHT computer program (version 3).

Our recommended relative and absolute γ -ray emission probabilities are given in Table 6.

Table 5: Experimental and evaluated relative γ -ray emission probabilities (%).

Reference	1965Ha13	1967Sc18	1970Ah05	1972Ha2 1	1974HeYW	Evaluated	Reduced χ^2
Energy (keV)							
30.895 (7)	12.3 (18)		6.8 (5)	7.08 (41)		7.1 (7)	4.3
98.882 (4)	100	100	100	100	100	100	
129.777 (5)	7.7 (8)	7.2 (8)	7.4 (5)	7.64 (44)	8.0 (6)	7.62 (26)	0.2
199.526 (12)		0.093 (10)			0.078 (9)	0.083 (7)	0.9
211.398 (6)	0.25 (3)*	0.119 (11)			0.102 (11)	0.108 (9)	0.8

* the experimental value has been shown to be an outlier value according to Lweight computer program.

Table 6: Recommended relative and absolute γ -ray emission probabilities (%).

E_γ (keV)	Relative γ -ray emission probability (%)	Absolute γ -ray emission probability (%)
30.895 (7)	7.1 (7)	0.80 (8)
98.882 (4)	100	11.21 (15)
129.777 (5)	7.62 (26)	0.854 (29)
199.526 (12)	0.083 (7)	0.0093 (8)
211.398 (6)	0.108 (9)	0.0121 (10)

6 References

- 1949St17 R. M. Steffen, O. Huber, F. Humbel, *Helv. Phys. Acta* 22(1949)167 [Half-life].
1949Wi08 G. Wilkinson, *Phys. Rev.* 75(1949)1019 [Half-life].
1959Bi07 A. Bisi, E. Germagnoli, L. Zappa, *Nuovo Cim.* 11(1959)843 [Half-life, Experimental α 's].
1959Mc69 F. K. McGowan, P. H. Stelson, *Phys. Rev.* 116(1959)154 [Experimental α 's].
1960Br11 M. Bresesti, J. C. Roy, *Can. J. Chem.* 38(1960)197 [Half-life].
1962Bo12 N. A. Bonner, W. Goishi, W. H. Hutchin, G. M. Iddings, H. A. Tewes, *Phys. Rev.* 127(1962)217 [Half-life].
1963Ha17 G. Harbottle, *Nucl. Phys.* 41(1963)604 [Half-life].
1964Go19 W. Goedbloed, E. Mastenbroek, A. Kemper, J. Blok, *Physica* 30(1964) [Experimental α 's, I_γ].
1965Ha13 J. R. Harris, G. M. Rothberg, N. Benczer-Koller, *Phys. Rev.* 138(1965)B554 [I_γ].
1967Sc18 R. Schöneberg, D. Gföller, A. Flammersfeld, *Z. Phys.* 203(1967)453 [I_γ].
1968Ja11 A. Jasinski, C. J. Herrlander, *Arkiv Fysik* 37(1968)585 [K X-rays].
1969Fi08 T. Fink, N. Benczer-Koller, *Nucl. Phys.* A138(1969)337 [Experimental α 's].
1970Ah05 B. Ahlesten, A. Bäcklin, *Nucl. Phys.* A154(1970)303 [Experimental α 's, I_γ].
1970To19 L. H. Toburen, R. G. Albridge, *Z. Phys.* 240(1970)185 [Experimental α 's].
1972Ha21 J. S. Hansen, J. C. McGeorge, R. W. Fink, R. E. Wood, P. V. Rao, J. M. Palms, *Z. Phys.* 249(1972)373 [I_γ , K X-rays].
1973Go05 S. C. Goverse, J. van Pelt, J. van den Berg, J. C. Klein, J. Blok, *Nucl. Phys.* A201(1973)326 [$P_{ec}(g.s.)$].
1974HeYW R. L. Heath, ANCR - 1000 - 2(1974) [I_γ].
1982HoZJ D. D. Hoppes, J. M. R. Hutchinson, F. J. Schima, M. P. Unterweger, NBS - SP - 626(1982)85 [Half-life].
1996Sc06 E. Schönfeld, H. Janßen, *Nucl. Instrum. Meth. Phys. Res.* A369(1996)527 [Atomic data].
1999Zh11 C. Zhou, *Nucl. Data Sheets* 86(1999)676 [Spin, level energies].
2002Un02 M. P. Unterweger, *Appl. Rad. Isotopes* 56(2002)125 [Half-life].
2002Ba85 I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr., P. O. Tikkanen, S. Raman, *Atomic Data Nucl. Data Tables* 81(2002)1 [Theoretical ICC].
2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr., *Nucl. Instrum. Meth. Phys. Res.* A589(2008)202 [Theoretical ICC].
2012Au06 M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer, *Chinese Physics C* 36 (2012) 1603 [Q].

**¹⁹⁸Au - Comments on evaluation of decay data
by V.P. Chechev and N.K. Kuzmenko**

The first DDEP evaluation of ¹⁹⁸Au decay data was completed by E. Schönfeld and R. Dersch in 1998 with a minor update in 2004 (2004BeZQ). The current evaluation was completed in January 2014 with a literature cut-off by the same date.

1. DECAY SCHEME

The structure of the adopted scheme of ¹⁹⁸Au is based on the ENSDF evaluation by Huang Xiaolong (2009Hu11). ¹⁹⁸Au disintegrates by β⁻ emission via ¹⁹⁸Hg excited levels of 411.8 keV (2⁺) and 1087.7 keV (2⁺). In addition to these levels, ¹⁹⁸Hg has an excited level at 1048.5 keV (4⁺, half-life of 7.2 (3) ps) which is below the Q⁻ value. A β⁻ transition from the ¹⁹⁸Au ground state (2⁻) to this level (ΔJ = 2 and the parity changes, E_β = 324.3 keV) would be unique 1st forbidden and was not observed. In 2004BeZQ an upper limit of 0.004 for the transition probability to this level was derived from *lg ft* systematics (*lg ft* ≥ 8.5).

Iwata and Yoshizawa (1980Iw03) estimated the probability of a possible EC decay of ¹⁹⁸Au to the ground state of ¹⁹⁸Pt (unique first forbidden EC transition) to be less than 0.0017 % from *lg ft* systematics, i.e. negligible for most purposes.

2. NUCLEAR DATA

The Q⁻ value is from the 2012 mass evaluation by Wang *et al.* (2012Wa38).

The recommended half-life of ¹⁹⁸Au is based on the experimental results given in Table 1.

Table 1. Experimental values of the ¹⁹⁸Au half-life (in days)

N	Author(s) and year	Reference	T _{1/2}	Method ^c and comments
1	Amaldi <i>et al.</i> (1935)	1935Am01	2.7	β counting; <i>omitted</i>
2	McMillan <i>et al.</i> (1937)	1937Mc04	2.7	β counting; <i>omitted</i>
3	Pool <i>et al.</i> (1937)	1937Po04	2.5	β counting; <i>omitted</i>
4	Sherr <i>et al.</i> (1941)	1941Sh08	2.7	β counting; <i>omitted</i>
5	Diemer and Groendijk (1946)	1946Di**	2.73 (2)	Ionization chamber; <i>omitted</i>
6	Seren <i>et al.</i> (1947)	1947Se33	2.7	β counting, Geiger counter; <i>omitted</i>
7	Saxon (1948)	1948Sa36	2.66 (1)	β counting, magnetic spectrometer; <i>omitted</i>
8	Saxon and Heller (1949)	1949Sa18	2.69 (2)	β counting, magnetic spectrometer; <i>omitted</i>
9	Steffen <i>et al.</i> (1949)	1949St17	2.7	β and γ counting; <i>omitted</i>
10	Cavanagh <i>et al.</i> (1951)	1951Ca06	2.66 (1)	β counting, Geiger counter; <i>omitted</i>
11	Silver (1951)	1951Si91	2.69 (1)	Ionization chamber, electrometer; <i>omitted</i>
12	Sinclair and Holloway (1951)	1951Si25	2.73 (1)	β counting, liquid counter, electroscopes; <i>omitted</i>

13	Lockett and Thomas (1953)	1953Lo09	2.697 (3)	Ionization chamber
14	Bell and Yaffe (1954)	1954Be61	2.699 (3)	Ionization chamber
15	Tobailem (1955)	1955To07	2.686 (5)	γ counting, ionization chambers
16	Johansson (1956)	1956Jo24	2.697 (5)	γ timing, NaI(Tl) detector
17	Sastre and Price (1956)	1956Sa75	2.694 (6)	Geiger counter
18	Keene (1958)	1958Ke26	2.704 (4)	Ionization chamber
19	Robert (1960)	1960Ro22	2.699 (4)	Calorimetry on gold samples
20	Starodubtsev <i>et al.</i> (1963)	1963St20	2.687 (5)	β counting, β spectrometer
21	Anspach <i>et al.</i> (1965)	1965An07	2.694 (4)	4π , 2π ionization chambers, proportional counter
22	Goodier (1968)	1968Go22	2.695 (7)	4π ionization chamber; <i>omitted</i> as superseded by 29
23	Lagoutine <i>et al.</i> (1968)	1968La10	2.697 (5)	Ionization chambers, 4π β counters, γ counter
24	Reynolds <i>et al.</i> (1968)	1968Re04	2.693 (5)	End window Geiger counter
25	Cabell and Wilkins (1969)	1969Ca23	2.6946 (10)	γ counting
26	Vuorinen and Kaloinen (1969)	1969Vu04	2.695 (2)	4π β γ coincidence counter
27	Costa Paiva and Martinho (1970)	1970Co14	2.696 (4)	γ counting, NaI(Tl) detector
28	Debertin (1971)	1971De**	2.693 (3)	γ counting, Ge(Li) detector
29	Goodier (1971)	1971GoYM	2.695 (3)	4π ionization chamber
30	Hoppes <i>et al.</i> (1982)	1982HoZJ	2.695 (2)	4π ionization chamber, γ counting; <i>omitted</i> as superseded by 33
31	Rutledge <i>et al.</i> (1982)	1982RuZV	2.6935 (4)	γ counting, GeLi detector
32	Abzouzi <i>et al.</i> (1990)	1990Ab02	2.6966 (7)	γ counting, GeLi detector
33	Unterweger <i>et al.</i> (1992)	1992Un01	2.69517 (21)	4π ionization chamber; <i>omitted</i> as superseded by 35
34	Mignonsin (1994)	1994Mi03	2.6837 (50)	γ counting, GeLi detector
35	Unterweger <i>et al.</i> (2002)	2002Un02	2.69516 (21)	4π ionization chamber; <i>omitted</i> as superseded by 36
36	Unterweger and Lindstrom (2004)	2004Un01	2.69555 (30)	4π ionization chamber; <i>omitted</i> as superseded by 48
37	Lindstrom <i>et al.</i> (2005)	2005Li66	2.6924 (11)	γ counting, Ge detector, 4π ionization chamber
38	Novkovic <i>et al.</i> (2006)	2006No10	2.6947 (6)	γ counting, Ge detector
39	Goodwin <i>et al.</i> (2007)	2007Go39	2.6951 (8) ^a	γ counting, Ge detector; <i>omitted</i> as superseded by 44
40	Spillane <i>et al.</i> (2007)	2007Sp01	2.726 (22) ^a	γ counting, Ge detector; <i>omitted</i> on the Chauvenet's criterion
41	Kumar <i>et al.</i> (2008)	2008Ku09	2.6973 (20) ^a	γ counting, Ge detector;
42	Ruprecht <i>et al.</i> (2008)	2008Ru05	2.6937 (4) ^a	γ counting, 2 Ge detectors
43	Fortak <i>et al.</i> (2010)	2010Fo13	2.6847 (51) ^a	γ counting, Ge detector
44	Goodwin <i>et al.</i> (2010)	2010Go25	2.6948 (9)	γ counting, Ge detector; <i>omitted</i> as superseded by 46

45	Lindstrom <i>et al.</i> (2011)	2011Li52	2.6782 (3) ^b	γ counting, Ge detector; <i>omitted</i> on the Chauvenet's criterion
46	Hardy <i>et al.</i> (2012)	2012Ha23	2.69445 (32)	γ counting, Ge detector; stat. unc.: (20), syst. unc.: (25)
47	Fitzgerald (2012)	2012Fi12	2.69516 (21)	4 π ionization chamber; <i>omitted</i> as superseded by 48
48	Unterweger and Fitzgerald (2014)	2014Un**	2.6934 (37)	4 π ionization chamber
Recommended value			2.6943 (3)	LWM

^a Average of measurement results at room and low (10-12 K) temperatures; measurements 39-43 showed temperature independence of ¹⁹⁸Au half-life.

^b Average of all reported data for two sources (sphere, wire); measurements 2010Li48 and 2011Li52 showed independence of ¹⁹⁸Au half-life on the source shape.

^c For measurements 1 to 44, the method is as listed in the compilation of 2011Ch22.

Values 1 to 12 from very early measurements (1935-1951) have been omitted as they are much less accurate. Values 22, 30, 33, 35, 36, 39, 44, 47 were not used because they were replaced ultimately by later results of the same laboratory.

Values 40 and 45 have been rejected by the LWEIGHT computer program based on Chauvenet's criterion. An unweighted average of the remaining twenty six values is 2.6940 (9) d. A weighted average is 2.69427 d. The LWEIGHT program, using the limitation of relative statistical weights method (LWM), has chosen the weighted average with the internal uncertainty of 0.00021 d. The external uncertainty is 0.00024 d. The largest contributions to the weighted average are the values 38 (12 %), 42 (26 %) and 46 (47 %). The ratio of the reduced $\chi^2 / (\chi^2)_{\text{crit}}$ is 1.35/1.80. The smallest experimental uncertainty is 0.00032 d.

The recommended value of ¹⁹⁸Au half-life is thus **2.6943 (3) days**. It can be compared with earlier evaluations of 2.6944 (8) (2004BeZQ), 2.6947 (3) (2009Hu11) and 2.6948 (12) (2011Ch22).

2.1. Beta Transitions

The energies of the β^- transitions have been obtained using the Q^- value and the ¹⁹⁸Hg level energies adopted from 2009Hu11 (Table 2). The probabilities of β^- -transitions P_{β^-} have been deduced from the $P(\gamma+ce)$ balance at each level of ¹⁹⁸Hg.

Table 2. ¹⁹⁸Hg levels populated in ¹⁹⁸Au β^- -decay

Level	Energy (keV)	Multipolarity	Half-life	P_{β^-} (%)
0	0	0 ⁺	Stable	0.025 (5)
1	411.80250 (17)	2 ⁺	23.16 (12) ps	98.99 (6)
2	1087.6874 (5)	2 ⁺	2.5 (2) ps	0.985 (5)

The adopted energies of the β^- transitions can be compared with the experimental values. $E_{\beta_{0,2}}$ (keV): 285.1 (5) – adopted and 290 (15) - measured (1951Br52).

$E_{\beta_{0,0}}$ (keV): 1372.8 (5) – adopted and 1371 (4) measured (1955E111).

For $E_{\beta_{0,1}}$, the adopted value of 961.0 (5) keV can be compared with the averaged value of 960.4 (10) keV evaluated by Schönfeld and Dersch (2004BeZQ) which came from many measurement results of 1954-1965.

2.2. Gamma Transitions and Internal Conversion Coefficients

Gamma-ray transition probabilities have been deduced from their gamma-ray emission probabilities and the total ICC(s). The adopted ICC(s) are the theoretical values interpolated by the BrIcc computer program (2008Ki07) from the tables of Band et al. (2002Ba85), accepting the “frozen orbital (no hole)” approximation. The multipolarities and mixing ratio δ have been taken from 2009Hu11.

For the 411.8 keV γ -ray transition:

- The adopted α_K of 0.0300 (5) can be compared with the experimental values: 0.0301 (5) – 1963Le11 ; 0.0302 (4) – 1964Be33 ; 0.0299 (2) – 1965Ke04 ; 0.0308 (9) – 1965Pe05 ; 0.0299 (2) – 1965Pa08 ; 0.0302 (4) – 1967BoZZ ; 0.0301 (3) – 1972Na22 ; 0.03035 (45) – 1973El10 and 0.0300 (3) – 1976Re**.

- The adopted α_T of 0.0439 (7) can be compared with the experimental values of 0.0444 (5) – 1965Ke04 ; 0.0447 (6) – 1967BoZZ ; 0.043 (4) – 1976Re** and 0.0445 (9) – 1980Iw03.

For γ 676 keV: the adopted value of α_K is 0.0216 (17) and the measured values are 0.0224 (19) – 1954El04 ; 0.019 (5) – 1956Vo20 and 0.03 (1) – 1967BoZZ.

For γ 1088 keV, the adopted value of α_K is 0.00414 (6) and the measured values are 0.00450 (31) – 1954El04 ; 0.0046 (6) – 1956Vo20, 0.0046 (6) – 1967BoZZ.

The adopted mixing ratio δ of 1.07 (14) for the (M1+E2) 676 keV γ -ray transition is based on the results of angular correlation measurements of 1971Pa06, 1971Be09 and 1969BeZR for 676 keV - 412 keV gamma-gamma cascade in the decay of 5.3 h ¹⁹⁸Tl, corrected for the current values of constants including the mean lifetime and g-factor of the 412 keV level in ¹⁹⁸Hg (2009Hu11). The most careful investigation was made in 1971Pa06 using Ge(Li), NaI(Tl), plastic detectors and a Ge(Li)-Ge(Li) setting. Earlier measurements of δ (before 1968) did not include two Ge(Li) and could contain systematic errors. Their uncorrected results are 0.96 (10) – 1953Sc19; 1.22 (22) – 1953Sc23; 1.43 (14) – 1964Sa11; 1.26 (8) – 1966Uh01; 1.34 (9) – 1967Ko13.

The $\delta = 1.07$ (14) corresponds to the adopted M1 contribution of 46.6 (65) %.

The latest angular correlation measurements, after 1971, gave the results of 54 (2) % (1972Ve03) and 39.4 (25) % (1974Ka18), divergent with each other, but compatible to the adopted value within the uncertainty limits.

3. ATOMIC DATA

The atomic data (fluorescence yields, X-ray energies and relative probabilities, and Auger electrons energies and relative probabilities) have been deduced by using the SAISINUC software.

4. ELECTRON EMISSIONS

The energies of the conversion electrons have been obtained from the gamma-ray transition energies and the electron binding energies.

The absolute emission probabilities of the conversion electrons have been deduced using recommended P_γ and ICC values.

The absolute emission probabilities of K and L Auger electrons have been calculated using the EMISSION computer program.

β^- average energies have been calculated using the LOGFT computer program.

5. PHOTON EMISSIONS

5.1. X-ray Emissions

The absolute emission probabilities of Hg KX- and LX-rays have been calculated using the EMISSION computer program. In Table 3 the calculated values are compared to experimental data.

Table 3. Experimental and recommended Hg X-ray emission probabilities in decay of ¹⁹⁸Au

X-ray	1989Ch45	2010Mo06	Recommended (calculated)
L1	0.027 (3)		0.0270 (8)
L α	0.592 (17)		0.527 (13)
L η	0.0105 (15)		0.0105 (3)
L β	0.643 (19)		0.536 (12)
L γ	0.124 (5)		0.1024 (23)
K α 2	0.816 (2)	0.813 (10)	0.807 (15)
K α 1	1.41 (4)	1.374 (17)	1.369 (24)
K' β 1	0.485 (12)	0.460 (7)	0.465 (11)
K' β 2	0.137 (7)	0.135 (3)	0.136 (4)

5.2. Gamma ray emissions

The energies of the gamma rays in ¹⁹⁸Hg have been adopted from 2009Hu11

The measured relative intensities and the adopted values obtained with the LWEIGHT program are listed in Table 4. Absolute gamma ray emission probabilities measured by Hammed *et al.* (1992Ha02) and Moreira *et al.* (2010Mo06) have been converted into relative intensities.

Table 4. Experimental and adopted relative gamma-ray intensities in decay of ¹⁹⁸Au

Reference	411.8 keV	675.9 keV	1087.7 keV
1951Ca06	100	1.5	0.4
1951Hu18	100	1.4 (1)	0.25 (5)
1951Br52	100	1	0.2
1954Ma19	100	1.3	0.25
1954E104	100	0.842 (56)	0.170 (12)
1955Dz41	100	1.11 (5)	0.26 (2)
1965Ke04	100	1.0	0.28
1967BoZZ	100	0.75	0.15
1980Iw03	100	0.841 (5)	0.1664 (22)
1989Ch45	100	0.846 (11)	0.165 (4)
1992Ha02	100	0.842 (11)	0.1669 (29)
2010Mo06	100	0.840 (8)	-
Recommended	100	0.841 (5)	0.1664 (22)

The normalization factor (0.956 17 (64) %) to convert the adopted relative gamma ray intensities to absolute emission probabilities has been deduced from the gamma ray transition intensity balance for the ground state assuming that the recommended probability of the beta transition to the ground state is equal to the measured value of 0.025 (5) % (1955E111).

6. ENERGY CONSERVATION

The total average energy of 1372.8 (9) keV, for one disintegration, calculated from the current evaluated data, corresponds very well to the available energy of 1372.8 (5) keV (Q) from the mass tables (2012Wa38) confirming the consistency of the decay scheme and the reliability of this evaluation.

7. REFERENCES

- 1935Am01** E. Amaldi, O.D. Agostino, E. Fermi, B. Pontecorvo, F. Rasetti, E. Segre. Proc. Roy. Soc. (London) 149A (1935) 522. [T_{1/2}]
- 1937Mc04** E. Mc Millan, M. Kamen, S. Ruben. Phys. Rev. 52 (1937) 531. [T_{1/2}]
- 1937Po04** M.L. Pool, J.M. Cork, R.L. Thornton. Phys. Rev. 52 (1937) 239. [T_{1/2}]
- 1941Sh08** R. Sherr, K.T. Bainbridge, H.H. Anderson. Phys. Rev. 60 (1941) 473. [T_{1/2}]
- 1946Di**** C. Diemer, H. Groendijk. Physika 11 (1946) 396. [T_{1/2}]
- 1947Se33** L. Seren, H.N. Friedlander, S.H. Turkel. Phys.Rev. 72 (1947) 888. [T_{1/2}]
- 1948Sa36** D. Saxon, Phys. Rev. 73 (1948) 811. [T_{1/2}]
- 1949Sa18** D. Saxon, R. Heller. Phys. Rev. 75 (1949) 909. [T_{1/2}]
- 1949St17** R.M. Steffen, O. Huber, F. Humbel. Helv. Phys. Acta 22 (1949) 167. [T_{1/2}]
- 1951Br52** A. Brosi, B. Ketelle, H. Zeldes, E. Fairstein. Phys. Rev. 84 (1951) 586. [I_γ]
- 1951Ca06** P.E. Cavanagh, J.F. Turner, D.V. Booker, H.J. Dunster. Proc. Phys. Soc. (London) 64A (1951) 13. [T_{1/2}, I_γ]
- 1951Hu18** M. Hubert, Compt. Rend. 232 (1951) 2201 [I_γ]
- 1951Si91** L.M. Silver, Can. J. Phys. 29 (1950) 59; Phys. Rev. 76 (1949) 589. [T_{1/2}]
- 1951Si25** W.K. Sinclair, A.F. Holloway. Nature 167 (1951) 365. [T_{1/2}]
- 1953Lo09** E.E. Lockett, R.H. Thomas. Nucleonics 11 (1953) 14. [T_{1/2}]
- 1953Sc19** C.D. Schrader. Phys. Rev. 92 (1953) 928. [Multipolarities 676 keV]
- 1953Sc23** D. Schiff, F.R. Metzger. Phys. Rev. 90 (1953) 849. [Multipolarities 676 keV]
- 1954Be61** R.E. Bell, L. Yaffe. Can. J. Phys. 32 (1954) 416. [T_{1/2}]
- 1954El04** L.G. Elliott, M.A. Preston, J.L. Wolfson. Can. J. Phys. 32 (1954) 153. [I_γ, Beta emission probabilities, Conv. Elec. emission probabilities]
- 1954Ma19** D. Maeder, R. Mueller, V. Wintersteiger, Helv. Phys. Acta 27 (1954) 3 [I_γ]
- 1955Dz41** B. S. Dzhelepov, N. N. Zhukovski, V. P. Prikhodtseva and Iu. V. Kholnov, Bull. Acad. Sci.(USSR) 19 (1955) 247 [I_γ]
- 1955El11** L.G. Elliott, M.A. Preston, J.L. Wolfson. Can. J. Phys. 33 (1955) 607. [Beta emission energies and probabilities]
- 1955To07** J. Tobailem, J. Phys. Rad. 16 (1955) 48. [T_{1/2}]
- 1956Jo24** K.-E. Johansson, Arkiv Fysik 10 (1956) 247. [T_{1/2}]
- 1956Sa75** C. Sastre, G. Price, Nucl. Sci. Eng. 1 (1956) 325. [T_{1/2}]
- 1956Vo20** J. Volpe, G. Hinmann. Phys. Rev. 104 (1956) 753. [Multipolarities 676 keV]
- 1958Ke26** J.P. Keene, L.A. Mackenzie, C.W. Gilbert. Phys. Med. Biol. 2 (1958) 360. [T_{1/2}]
- 1960Ro22** J. Robert, J. Phys. Rad. 21 (1960) 808. [T_{1/2}]
- 1963Le11** W.H.G. Lewin, B. van Nooyen, C.W.E. van Eijk, A.H. Wapstra. Nucl. Phys. 48 (1963) 159. [Beta emission probabilities]

- 1963St20** S.V. Starodubtsev, R.B. Begzhanov, S.L. Rakovitskii. *Izv. Akad. Nauk UzSSR, Ser. Fiz.-Mat. Nauk* 7, No.2 (1963) 44; *Nucl. Sci. Abstr.* 18 (1964) 9348. [T_{1/2}]
- 1964Be33** K.-E. Bergkvist, S. Hultberg. *Arkiv Fysik* 26 (1964) 239. [Experimental ICC]
- 1964Sa11** M. Sakai, M. Nozawa, H.I. Kegami, T. Yamazaki. *Nucl. Phys.* 53 (1964) 529. [Multipolarities]
- 1965An07** S.C. Anspach, L.M. Cavallo, S.B. Garfinkel, J.M.R. Hutchinson. *NBS Misc. Publ.* 260-9 (1965); NP-15663. [T_{1/2}]
- 1965Be07** K.-E. Bergkvist, S. Hultberg. *Arkiv Fysik* 27 (1965) 321. [Experimental ICC]
- 1965Ke04** W.J. Keeler, R.D. Connor. *Nucl. Phys.* 61 (1965) 513. [Conv. Elec. emission probabilities, Beta emission probabilities, I_γ]
- 1965Pa08** H. Paul. *Nucl. Phys.* 72 (1965) 326. [Beta emission probabilities]
- 1965Pe05** B.-G. Pettersson, L. Holmberg, T.R. Gerholm. *Nucl. Phys.* 65 (1965) 454. [Experimental ICC]
- 1966Uh01** M. Uhl, H. Warhanek. *Oesterr. Akad. Wiss. Sitzber. Math-Nat.Kl., Abt.II* 175 (1966) 77. [Multipolarities 676 keV]
- 1967BoZZ** H.E. Bosch, E. Szichman. *Bull. Am. Phys. Soc.* 12 (1967) 598. [Conv. Elec. emission probabilities, I_γ]
- 1967Ko13** J. Koch, F. Munnich, U. Schotzig. *Nucl. Phys. A103* (1967) 300. [Multipolarities 676 keV]
- 1968Go22** I.W. Goodier, *Int. J. Appl. Rad. Isotopes* 19 (1968) 823. [T_{1/2}]
- 1968La10** F. Lagoutine, Y. Le Gallic, J. Legrand. *Int. J. Appl. Rad. Isotopes* 19 (1968) 475. [T_{1/2}]
- 1968Re04** S.A. Reynolds, J.F. Emery, E.I. Wyatt. *Nucl. Sci. Eng.* 32 (1968) 46. [T_{1/2}]
- 1969BeZR** R. Beraud, I. Berkes, J. Daniere, R. Haroutunian, M. Levy, G. Marest, R. Rougny. *Phys. Rev.* 188 (1969) 1958. [Multipolarities 676 keV]
- 1969Ca23** M.J. Cabell, M. Wilkins. *J. Inorg. Nucl. Chem.* 31 (1969) 1229. [T_{1/2}]
- 1969Vu04** A. Vuorinen, E. Kaloinen. *Ann. Acad. Sci. Fennicae, Ser.A VI* (1969) No.310. [T_{1/2}]
- 1970Co14** M.M. Costa Paiva, E. Martinho. *Int. J. Appl. Radiat. Isot.* 21 (1970) 40. [T_{1/2}]
- 1971Be09** R. Beraud, I. Berkes, R. Haroutunian, G. Marest, M. Meyer-Levy, R. Rougny, A. Troncy, A. Baudry, V. Lopac. *Phys.Rev. C4* (1971) 1829. [Multipolarities 676 keV]
- 1971De**** V.K. Debertin, *Atomkernergie* 17 (1971) 97. [T_{1/2}]
- 1971GoYM** I.W. Goodier, M.J. Woods, A. Williams. *Proc. Int. Conf. Chemical Nucl. Data* (1971), Canterbury, M.L. Hurrell, Ed., p.175. [T_{1/2}]
- 1971Pa06** A. Pakkanen. *Nucl. Phys. A172* (1971) 193. [Multipolarities 676 keV]
- 1972Na22** T. Nagarajan, M. Ravindranath, K. Venkata Reddy. *J. Phys. A (Gen.Phys.)* 5 (1972) 1395. [Conv. Elec. emission probabilities, Beta emission probabilities]
- 1972Ve03** K. Venkata Ramana Rao, V. Lakshminarayana. *Nuovo Cimento* 8A (1972) 298. [Multipolarities 676 keV]
- 1973El10** M.S. El-Nesr, M.G. Mousa. *Atomkernenergie* 21 (1973) 207. [Experimental ICC]
- 1974Ka18** M. Kawamura, T. Tomiyama. *J. Phys. Soc. Japan* 36 (1974) 27. [Multipolarities 676 keV]
- 1976Re**** T.S. Reddy. Thesis Andhra Univ., Waltair, India (1976). [Conv. Elec. emission probabilities]
- 1980Iw03** Y. Iwata, Y. Yoshizawa. *Nucl. Instrum. Methods* 175(1980) 525. [P(EC), I_γ]
- 1982HoZJ** D.D. Hoppes, J.M.R. Hutchinson, F.J. Schima, M.P. Unterweger. *NBS-SP-626* (1982) 85. [T_{1/2}]

- 1982RuZV** A.R. Rutledge, L.V. Smith, J.S. Merritt. NBS-SP-626 (1982) 5. [T_{1/2}]
- 1989Ch45** B. Chand, J. Goswamy, D. Mehta, N. Singh, P.N. Trehan, Nucl. Instr. Meth. A284 (1989) 393. [P_γ, P_X]
- 1990Ab02** A. Abzouzi, M.S. Antony, A. Hachem, V.B. Ndocko Ndongue, J. Radioanal. Nucl. Chem. 144 (1990) 359. [T_{1/2}]
- 1992Ha02** M.A. Hammed, I.M. Lowles, T.D. Mac Mahon. Nucl. Instrum. Methods Phys. Res. A312 (1992) 308. [P_γ]
- 1992Un01** M.P. Unterweger, D.D. Hoppes, F.J. Schima. Nucl. Instrum. Methods Phys. Res. A312 (1992) 349. [T_{1/2}]
- 1994Mi03** E.P. Mignonsin, Appl. Radiat. Isotopes 45 (1994) 17. [T_{1/2}]
- 2002Ba85** I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor Jr., P.O. Tikkanen, S. Raman. At. Data Nucl. Data Tables 81 (2002) 1. [Theoretical ICC]
- 2002Un02** M.P. Unterweger, Appl. Radiat. Isot. 56, 125 (2002). [T_{1/2}]
- 2004BeZQ** M.-M. Bé, V. Chisté, C. Dullieu, E. Browne, V. Chechev, N. Kuzmenko, R. Helmer, A. Nichols, E. Schönfeld, and R. Dersch. *Table of Radionuclides (Vol. 2 – A = 151 to 242)*. Bureau International des Poids et Mesures, 2004. ¹⁹⁸Au. P. 121-128. [Previous ¹⁹⁸Au decay data evaluation]
- 2004Un01** M.P. Unterweger, R.M. Lindstrom. Appl. Radiat. Isot. 60 (2004) 325. [T_{1/2}]
- 2005Li66** R.M. Lindstrom, M. Blaauw, M.P. Unterweger. J. Radioanal. Nucl. Chem. 263 (2005) 311. [T_{1/2}]
- 2006No10** D. Novkovic, L. Nadder, A. Kandic, I. Vukanac, M. Durasevic, D. Jordanov. Nucl. Instrum. Methods Phys. Res. A566 (2006) 477. [T_{1/2}]
- 2007Go39** J.R. Goodwin, V.V. Golovko, V.E. Iacob, J.C. Hardy. Eur. Phys. J. A34 (2007) 271. [T_{1/2}]
- 2007Sp01** T. Spillane, F. Raiola, F. Zeng, H.W. Becker, L. Gialanella, R. Kunze, C. Rolfs, M. Romano, D. Schurmann, F. Streider. Eur. Phys. J. A31 (2007) 203. [T_{1/2}]
- 2008Ki07** T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor Jr. Nucl. Instrum. Methods Phys. Res. A589 (2008) 202. [BrIcc computer program]
- 2008Ku09** V. Kumar, M. Hass, Y. Nir-El, G. Haquin, Z. Yungriess. Phys. Rev. C 77 (2008) 051304. [T_{1/2}]
- 2008Ru05** G. Ruprecht, C. Vockenhuber, L. Buchmann, R. Woods, C. Ruiz, S. Lapi, D. Bemmerer. Phys. Rev. C 77, 065502 (2008). [T_{1/2}]
- 2009Hu11** Huang Xiaolong, Nuclear Data Sheets 110 (2009) 2533. [Decay scheme, ¹⁹⁸Hg adopted levels and gammas, multipolarities, mixing ratio for 676 keV gamma-ray].
- 2010Fo13** K. Fortak, R. Kunz, L. Gialanella, H.-W. Becker, J. Meijer, F. Strieder. Eur. Phys. J. A46 (2010) 161. [T_{1/2}]
- 2010Go25** J.R. Goodwin, N. Nica, V.E. Iacob, A. Dibidad, J.C. Hardy. Phys.Rev. C 82, 044320 (2010). [T_{1/2}]
- 2010Li48** R.M. Lindstrom, E. Fischbach, J.B. Buncher, G.L. Greene, J.H. Jenkins, D.E. Krause, J.J. Mattes, A. Yue. Nucl. Instrum. Methods Phys. Res. A622 (2010) 93. [Independence of T_{1/2} on the source shape]
- 2010Mo06** D.S. Moreira, M.F. Koskinas, M.S. Dias, I.M. Yamazaki. Appl. Radiat. Isot. 68 (2010) 1566. [P_γ]
- 2011Ch22** J. Chen, S.D. Geraedts, C. Ouellet, B. Singh. Appl. Radiat. Isot. 69 (2011) 1064. [Evaluation of ¹⁹⁸Au half-life by different statistical procedures]
- 2011Li52** R. M. Lindstrom, E. Fischbach, J.B. Buncher, J. H. Jenkins, A. Yue. Nucl. Instrum. Methods Phys. Res. A 659 (2011) 269. [T_{1/2}]

- 2012Ha23** J.C. Hardy, J.R. Goodwin, V.E. Jacob. Appl. Radiat. Isot. 70 (2012) 1931. [T_{1/2}]
- 2012Wa38** M. Wang, G. Audi, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer. Chin. Phys. C36 (2012) 1603. [Q]
- 2012Fi12** R. Fitzgerald, J. Res. Natl. Inst. Stand. Technol. 117 (2012) 80 [T_{1/2}]
- 2014Un**** M.P. Unterweger, R. Fitzgerald. Appl. Radiat. Isot. (2014). Article in press:
<http://dx.doi.org/10.1016/j.apradiso.2013.11.017>. [T_{1/2}]

²⁰¹Tl - Comments on evaluation of decay data by E. Schönfeld

This evaluation was completed in May 1997 and the half life value has been updated in May 2004.

1 Decay Scheme

Above the 167 keV level and below available energy there are three levels of ²⁰¹Hg: 384,601(18) keV (5/2-), 414,522(17) keV (7/2-); 21,3 ps, and 464,41(3) keV (5/2 -); 2,6 ps. EC transitions to these levels would be (in the above order) unique first forbidden / nonunique third forbidden and unique first forbidden. But, these transitions have not been observed in the decay of ²⁰¹Tl. If these transitions do not exist, then the decay scheme on page 1 is complete.

2 Nuclear Data

The following values of the half-life have been considered ($T_{1/2}$ in d):

1	3,00(13)	Neumann and Perlman (1950)
2	3,063(33)	Herrlander et al. (1960)
3	3,0380(7)	Debertin et al. (1979) ; superseded by 6
4	3,0408(14)	Lagoutine and Legrand (1982); originally given $3\sigma = 0,0040$ d
5	3,0447(9)	Hoppes et al. (1982) ; superseded by 7
6	3,043(3)	Schrader (1989) ; superseded by 10
7	3,0456(15)	Unterweger et al. (1992)
8	3,0400(28)	Simpson and Meyer (1994)
9	3,038(17)	de Souza (2004)
10	3,0486(30)	Schrader (2004)
11	3,0421(17)	adopted value with external uncertainty, present evaluation

Values 1 and 2 are only of historical interest. Value 5 is superseded by value 7 and value 3 by value 6 and then by value 10. The LWM of values 4, 7, 8, 9 and 10 is given as value 11, the reduced χ^2 is 4,3.

The Q_{EC} value 483(15) keV is taken from Audi and Wapstra (1995).

2.1 Electron Capture Transitions

The adopted values P_K , P_L , P_M , P_N were calculated from the table of Schönfeld (1995) using the Q_{EC} value of Audi and Wapstra (1995) and the binding energies of Hg. These values are:

ΔE keV	P_K	P_L	P_M	P_{NO}
316(15)	0,724(7)	0,206(7)	0,054(2)	0,016(2)
451(15)	0,758(3)	0,181(3)	0,0461(12)	0,025(2)
483(15)	0,763(3)	0,178(3)	0,0451(12)	0,014(2)

The above values are in excellent agreement with the values calculated by Funck and Nylandstedt Larsen (1983) although the latter have no assigned uncertainties:

to level keV	P_K	P_L	P_M
167	0,7230	0,2016	0,0549
32	0,7567	0,1813	0,0474
1,6 and 0	0,7613	0,1779	0,0464

They are also in agreement with the values given by Lagoutine in the Table des Radionucléides (1984). It has to be mentioned that Lagoutine used different transition energies. His values are:

ΔE keV	P_K	P_L	P_{MN}
321(15)	0,730(5)	0,206(3)	0,064(2)
456(15)	0,762(5)	0,182(3)	0,056(2)
488(15)	0,767(5)	0,178(3)	0,055(2)

The transition probabilities of the EC transitions were calculated by

$$P_{e_{0,4}} = P_{g+ce_{4,0}} + P_{g+ce_{4,1}} + P_{g+ce_{4,2}} + P_{g+ce_{4,3}}$$

$$P_{e_{0,3}} = P_{g+ce_{3,0}} + P_{g+ce_{3,1}} + P_{g+ce_{3,2}} - P_{g+ce_{4,3}}$$

$$P_{e_{0,1}} + P_{e_{0,0}} = 1 - (P_{e_{0,4}} + P_{e_{0,3}})$$

2.2 Gamma Transitions

The energies of the main transitions are measured by Herrlander et al. (1960) via the conversion energies. The present values are taken from S. Rab (1994).

Herrlander et al. (1960) have measured the $L_1/L_2/L_3$ ratios of the 30,6 keV, 32,19 keV, 135,34 keV and 167,43 keV. By comparing the experimental values with theoretical ones the multipolarity of all this transitions were proved to be M1. For the 165,88 keV an E 2 mixture of up to 7 % could not be excluded. The present multiplicities and conversion coefficients are taken from Rab (1994). The transition probabilities are calculated from the gamma-ray emission probabilities (4.2) and the total conversion coefficients.

3 Atomic data

The atomic data are taken from Schönfeld and Janßen (1996).

3.1 X Radiation

The energy values are calculated from the wavelengths in Å* as given by Bearden (1967).

The relative emission probabilities of K X rays are taken from Schönfeld and Janßen (1996).

3.2 Auger Electrons

The energy values are taken from Larkins (1977) (KLL) and the Table de Radionucléides (LMRI 1982) (KLX, KXY). The relative emission probabilities of K Auger electrons are taken from Schönfeld and Janßen (1996). The relative emission probabilities of the L Auger electrons is calculated from the value in the table 4.1 putting $P(KLL) = 1$.

4 Radiation Emission

4.1 Electron Emission

The energies of the Auger are the same as in 3.2. The energies of the conversion electrons are calculated from the transition energy (2.2) and the binding energies.

The emission probabilities of the conversion electrons are calculated using the conversion coefficients given in 2.2. The values of the emission probabilities of the Auger electrons are calculated using the transition probabilities given in 2.1 and 2.2, the atomic data given in 3 and the conversion coefficients given in 2.2.

4.2 Photon Emission

The energy of the X rays are the same as in 3.1. For the relative K X ray emission probabilities and the relative γ ray emission probabilities it has been found

E_γ in keV	1	2	3	4	5	6	7	8	9
30,60	2,2(2)	3,10(13)	2,35(25)	2,57(6)	2,60(8)	2,60(8)	2,53(5)	2,58(5)	-
32,19	2,2(2)	2,85(12)	2,69(34)	2,60(9)	2,60(7)	2,72(6)	2,58(5)	2,63(5)	-
68,90 K_{a_2}		274(9)	243(15)	261(7)		270(4)		268(4)	273(5)
70,82 K_{a_1}		466(14)	412(25)	446(12)		442(6)		446(6)	464(7)
K_a		740(23)	655(29)	707(14)	722(13)	712(7)		715(7)	737(11)
80,2 K_{b_1}				153(4)				153(4)	157(4)
82,5 K_{b_2}				45,9(15)				45,9(15)	46,1(13)
K_b		205(7)	182(11)	199(16)	205(4)	195(5)		202(5)	203(5)
135,34	26,5(13)	26,5(10)	31(4)	26,4(3)	26,5(4)	27,2(5)	25,65(18)	26,04(22)	-
165,88	1,6(1)	1,80(20)	1,6(3)	1,5(2)	1,46(20)	1,45(2)	1,55(5)	1,47(2)	-
167,43	100	100,0(17)	100(8)	100,0(11)	100,0(10)	100,0(12)	100	100,0(10)	-

1: Hofmann and Walcher (1975)

2: Nass (1977)

3: Martin (1976)

4: Debertain et al. (1978)

5: Funck et al. (1983)

6: Kawada et al. (1990)

7: Coursey et al. (1990)

8: LWM (without 3)

9: Calculated from atomic data, EC data and conversion coefficients. Adopted and recommended values for the X rays.

The values in column 8 are the LWM from 1, 2, 4 - 7 (the values 3 are less reliable). The uncertainties were taken not smaller than the minimum of a single value. Between values 8 and 9 there is not in all cases 1σ overlapping. The transformation from relative emission probabilities to absolute emission probabilities was made using the absolute transition probability for the 167 keV transition $P_\gamma(167) = 0,1000(10)$ as determined by Coursey et al. (1990) from absolute activity measurements..

5 Main Production Modes

Taken from the "Table de Radionucléides", LMRI, 1982.

6 References

R. K. Gupta, *Arkiv Fysik* 17 (1960) 337

C. J. Herrlander, R. Stockendal, R. K. Gupta, *Arkiv Fysik* 17 (1960) 315

[KLL, KXL, KXY]

D. Reyes-Suter, T. Suter, *Arkiv Fysik* 20 (1961) 415

[$T_{1/2}$ 32 keV level in ^{201}Hg]

S. Rab, *Nuclear Data Sheets* 71 (1994) 421

[Multipolarities]

M. Neumann, I. Perlman, *Phys. Rev.* 78 (1950) 191

[$T_{1/2}$]

And also see the Tables Part.

²⁰³Hg – Comments on evaluation of decay data by A.L. Nichols

Evaluated: April 2001

Re-evaluated: January 2004

Evaluation Procedures

Limitation of Relative Statistical Weight Method (LWM) was applied to average numbers throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

The simple and consistent decay scheme is dominated by beta decay to the first excited state of ²⁰³Tl, followed by a single gamma transition to the ground state.

Nuclear Data

The single well-characterised gamma ray at 279.1952(10) keV and the 46.6 -day half-life of ²⁰³Hg make this radionuclide of some value as a standard in the calibration of γ -ray detectors.

Half-life

Half-life adopted from the evaluation of Woods et al (2004) for the IAEA -CRP: Update of X- and Gamma-ray Decay Data Standards for Detector Calibration. The measurements of 1968La10, 1972Em01, 1980Ho17, 1980RuZY, 1983Wa26 and 1992Un01 were considered.

Reference	Half-life (days)
1968La10	47.000(30)*
1972Em01	46.760(80)*
1980Ho17	46.582(2)#
1980RuZY	46.600(10)
1983Wa26	46.612(19)
1992Un01	46.619(27)
Recommended value	46.593(7)

* Removed from evaluated data set due to large deviation from mean.

Uncertainty adjusted to ± 0.008 to reduce weighting below 0.5.

Woods evaluation for IAEA-CRP (2004WoZZ): recommended half-life of 46.594(12) days (using above dataset).

Gamma Rays

Energy

The gamma-ray energy and uncertainty recommended by 2000He14 were adopted. This energy is in good agreement with the nuclear level energy of the first excited state of ²⁰³Tl as specified by 1985Sc23 and 1993Ra11.

Emission Probability

The 279.1952 keV gamma transition is of mixed M1 + E2 multipolarity, and α_{tot} of 0.2271(12) and α_{K} of 0.1640(10) have been adopted from the evaluation of 1985HaZA, in good agreement with various measurements (1962Ta06, 1964He19, 1974Ha29, 2000Sc05). A small uncertainty was assigned to these two parameters because of the high degree of confidence in the data. The gamma transition probability of 0.9999(1) was deduced as described below, and used in conjunction with α_{tot} to calculate an absolute emission probability of 0.8148(8).

Multipolarity and Internal Conversion Coefficients of 279.1952 keV Gamma Ray

The comprehensive assessment of 1985HaZA provides accurate estimates for α_{tot} of 0.2271(12) and α_{K} of 0.1640(10), and a multipolarity of close to 25%M1 + 75%E2. These values have been adopted, and used to calculate the other α components in terms of the recommended value of α_{tot} . The selected data set used by 1985HaZA to determine α_{tot} and α_{K} is included in the table below (see footnotes); not all measurements are listed (see 1985HaZA for further details).

Internal conversion coefficients for 279.1952 keV gamma ray – selected measurements

	1956Wa30	1958Ni28	1960Pe22	1961Su10	1962Ta06*	1963Bu09*
α_{tot}	-	-	0.227(8)	-	0.2273(24) [#]	-
α_{K}	0.164(5) [#]	0.163(3) [#]	0.164(6) [#]	0.164(4) [#]	0.1642(21) [#]	0.165(9) [#]
α_{L}	0.049(2)	0.0487(12)	-	-	-	-
$\alpha_{\text{M+}}$	-	-	-	-	-	-

	1963Cr14	1964He19	1972Sa34	1972WaYL*	1974Ha29	2000Sc05
α_{tot}	-	-	0.149(9) 0.156(9)	0.2267(16) [#]	0.2279(24) [#]	0.2250(12)
α_{K}	0.162(3) [#]	0.163(3) [#]	-	-	0.1653(17) [#]	-
α_{L}	-	0.0484(6)	-	-	0.0475(13)	-
$\alpha_{\text{M+}}$	-	0.0153(4)	-	-	-	-

* Data adjusted by 1985HaZA from the published values.

Values adopted in an evaluation by 1985HaZA.

Internal conversion coefficients of 279.1952 keV gamma ray – theoretical values and 1985HaZA evaluation

	1978Ro22*	1985HaZA [‡]	Recommended Values
α_{tot}	0.231(7)	0.2271(12)	0.2271(12)
α_{K}	0.161(5)	0.1640(10)	0.1640(10)
α_{L}	0.053(2)	-	0.0476(2)
$\alpha_{\text{M+}}$	0.017(5)	-	0.0155(2)

* Interpolated values for 25%M1 + 75%E2, with 3% uncertainty.

[‡] Hansen used three α_{tot} and nine α_{K} values (see previous table) to derive recommended values, which were originally selected from six α_{tot} and twenty-eight α_{K} values respectively.

Beta-particle Emissions

Energies

The beta-particle energies were calculated from the proposed decay scheme. The nuclear level energies of 1993Ra11 and the Q-value were used to determine the energies and uncertainties of the beta -particle transitions to the first excited state (dominant) and ground level.

Emission Probabilities

The beta-particle emission probabilities were calculated from the limits set on the beta transition to the ground state by 1955Ma40 and 1956Wo09. Beta-decay branch to $\frac{1}{2}^+$ Ground State of ²⁰³Tl:

	1955Ma40	1956Wo09	Recommended Values
$P_{\beta} (5/2^- \rightarrow 1/2^+)$	<0.00004	<0.0003	0.0001(1)
$\log f^{nu}_t$	-	>11.3	11.6(4)

A value of 0.0001(1) was recommended from these studies. Hence, the beta-particle emission probability was defined as 0.9999(1) for the transition to the first excited state of ²⁰³Tl ($5/2^- \rightarrow 3/2^+$).

Beta-particle Emission Probabilities

$E_b(\text{keV})$	P_b
	Recommended Values*
212.6(12)	0.9999(1)
491.8(12)	0.0001(1)

* Recommended emission probabilities derived from the postulated limit of the beta branch to the ²⁰³Tl ground state.

Atomic Data

The X-ray data have been calculated using the evaluated gamma -ray data, and the atomic data from 1996Sc06, 1998ScZM and 1999ScZX.

References

- 1955Ma40 - N. Marty, Sur la Désintégration de ²⁰³Hg, C. R. Acad. Sc. Paris, 240B(1955)291-294. [P_β]
- 1956Wa30 - A. H. Wapstra and G. J. Nijgh, Indications for a Strong Influence of the Nuclear Size on Magnetic Dipole Conversion Coefficients, Nucl. Phys. 1(1956)245-258. [ICC]
- 1956Wo09 - J. L. Wolfson, High-energy Forbidden β-ray Transitions from Cs¹³⁴ (2.3 y), Co⁶⁰ (5.3 y), Sc⁴⁶ (84 d) and Hg²⁰³ (47.9 d), Can. J. Phys. 34(1956)256-264. [P_β, transition type]
- 1958Ni28 - G. J. Nijgh, A. H. Wapstra, L. Th. M. Ornstein, N. Salomons-Grobbe, J. R. Huizenga and O. Almén, Conversion Coefficients of Gamma Transitions in ²⁰³Tl, Nucl. Phys. 9(1958/59)528-537. [ICC]
- 1960Pe22 - R. W. Peelle, Determination of the Internal Conversion Coefficient of the 279-keV Gamma Ray in Tl²⁰³ by Absolute Coincidence Techniques, Oak Ridge National Laboratory Report ORNL 3016(1960)116-125. [ICC]
- 1961Su10 - Z. Sujkowski, Beta Spectrometric Study on the Decay of Pb²⁰³ and the K Auger Spectrum of Tl, Ark. Fys. 20, No. 16 (1961) 243-267. [ICC]
- 1962Ta06 - J. G. V. Taylor, The Total Internal Conversion Coefficient of the 279-keV Transition Following the Decay of Hg²⁰³ as Measured by a New Coincidence Method, Can. J. Phys. 40(1962)383-392. [ICC]

- 1963Bu09 - R. Burmeister, H. Graber, J. Schintlmeister and R. Weibrecht, Bestimmung des K Konversionskoeffizienten des 279 keV-Übergangs in Tl²⁰³, Nucl. Phys. 42(1963)56-61. [ICC]
- 1963Cr14 - W. L. Croft, B. G. Pettersson and J. H. Hamilton, The K-conversion Coefficient of the 279 keV Transition in Tl²⁰³ by a Coincidence Technique and Establishment as a Standard, Nucl. Phys. 48(1963)267-272. [ICC]
- 1964He19 - C. J. Herrlander and R. L. Graham, Penetration Effects in the K and L Internal Conversion Coefficients of the 279 keV Transition in Tl²⁰³, Nucl. Phys. 58(1964)544-560. [ICC]
- 1968La10 - F. Lagoutine, Y. Le Gallic and J. Legrand, Determination Precise de Quelques Perodes Radioactives, Int. J. Appl. Radiat. Isot. 19(1968)475-482. [Half-life]
- 1972Em01 - J. F. Emery, S. A. Reynolds, E. I. Wyatt and G. I. Gleason, Half -Lives of Radionuclides – IV, Nucl. Sci. Eng. 48(1972)319-323. [Half-life]
- 1972Sa34 - H. S. Sahota, On the Measurement of K-shell Internal Conversion Coefficients with an X-ray Scintillation Spectrometer, Indian J. Phys. 46(1972)86-92. [ICC]
- 1972WaYL - K. F. Walz, H. M. Wei ß and E. Funck, Bestimmung des Koeffizienten der inneren Konversion beim 279 keV-Übergang des ²⁰³Hg, PTB Jahresbericht 1971 (1972)150-151. [ICC]
- 1974Ha29 - H. H. Hansen and D. Mouchel, Internal Conversion Coefficients and Penetration Effect for the 279 keV Transition in ²⁰³Tl, Z. Phys. 267(1974)371-377. [ICC]
- 1978Ro22 - F. Rösler, H. M. Fries, K. Alder and H. C. Pauli, Internal Conversion Coefficients for all Atomic Shells, ICC Values for Z = 68-104, At. Data Nucl. Data Tables 21(1978)291-514. [ICC]
- 1980Ho17 - H. Houtermans, O. Milosevic and F. Reichel, Half -lives of 35 Radionuclides, Int. J. Appl. Radiat. Isot. 31(1980)153. [Half-life]
- 1980RuZY - A. R. Rutledge, L. V. Smith and J. S. Merritt, Decay Data for Radionuclides Used for the Calibration of X- and Gamma-Ray Spectrometers, AECL-6692(1980). [Half-life]
- 1983Wa26 - K. F. Walz, K. Debertin and H. Schrader, Half -Life Measurements at the PTB, Int. J. Appl. Radiat. Isot. 34(1983)1191. [Half-life]
- 1985HaZA - H. H. Hansen, Evaluation of K-shell and Total Internal Conversion Coefficients for Some Selected Nuclear Transitions, European Appl. Res. Rept. - Nucl. Sci. Technol. 6, No.4(1985)777-816; EUR 9478 EN. [ICC]
- 1985Sc23 - M. R. Schmorak, Nuclear Data Sheets for A = 203, Nucl. Data Sheets 46(1985)287. [Nuclear structure, Energies]
- 1992Un01 - M. P. Unterweger, D. D. Hoppes and F. J. Schima, New and Revised Half -Life Measurements Results, Nucl. Instrum. Meth. Phys. Res. A312(1992)349-352. [Half-life]
- 1993Ra11 - S. Rab, Nuclear Data Sheets Update for A = 203, Nucl. Data Sheets 70(1993)173. [Nuclear structure, Energies]
- 1995Au04 - G. Audi and A. H. Wapstra, The 1995 Update to the Atomic Mass Evaluation, Nucl. Phys. A595(1995)409. [Q value]
- 1996Sc06 - E. Schönfeld and H. Janßen, Evaluation of Atomic Shell Data, Nucl. Instrum. Meth. Phys. Res. A369(1996)527-533. [X_K, X_L, Auger electrons]
- 1998ScZM - E. Schönfeld and G. Rodloff, Tables of the Energies of K-Auger Electrons for Elements with Atomic Numbers in the Range from Z = 11 to Z = 100, PTB Report PTB -6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX - E. Schönfeld and G. Rodloff, Energies and Relative Emission Probabilities of K X-rays for Elements with Atomic Numbers in the Range from Z = 5 to Z = 100, PTB Report PTB -6.11-1999-1, February 1999. [X_K]
- 2000He14 - R. G. Helmer and C. van der Leun, Recommended Standards for γ-ray Energy Calibration (1999), Nucl. Instrum. Meth. Phys. Res. A450(2000)35-70. [E_γ]
- 2000Sc05 - E. Schönfeld, H. Janßen and R. Klein, Redetermination of the Total Internal Conversion Coefficient of the 279 keV Transition Following the Decay of ²⁰³Hg, Appl. Radiat. Isot. 52(2000)955-956. [ICC]
- 2004WoZZ - M. J. Woods, Half -life Evaluations for IAEA -CRP on “Update of X-ray and Gamma-ray Decay Data Standards for Detector Calibration and Other Applications” (2004). [Half-life evaluation]

²⁰³Pb - Comments on evaluation of decay data by V. Chisté and M. M. Bé

1 Decay Scheme

²⁰³Pb disintegrates by electron capture to ²⁰³Tl via excited levels. Spin and half-life of the 680-keV level are from the mass-chain evaluation of F. G. Kondev (2005Ko20).

2 Nuclear Data

The Q(EC) value is from the atomic mass adjustment of Audi et al. (2003Au03).

Experimental ²⁰³Pb half-life values (in hours) are given in Table 1:

Table 1: Experimental values of ²⁰³Pb half-life.

Reference	Experimental value (h)	Comments
K. Fajans (1941Fa04)	52,0 (5)	
J. R. Prescott (1954Pr04)	52 (1)	
A. A. Barlett (1958Bart)	52,1 (2)	
L. Persson (1961Pe12)	52,1 (2)	
G. A. Chackett (1971Ch54)	52,02 (10)	Original uncertainty increased (x 2) for missing details (systematic uncertainty).
H. Houtermans (1979Ho17)	51,88 (2)	
D. D. Hoppes (1982HoZJ)	51,92 (4)	Superseded by 2002Un02.
K. Lindenberg (2001Li17)	51,94 (1)	
M. P. Unterweger (2002Un02)	51,923 (37)	
Recommended value	51,929 (10)	$\chi^2 = 1,37$

The evaluators have chosen to take into account the eight values with associated uncertainty for the calculation. The original uncertainty given by Chackett (1971Ch54) has been multiplied by 2, in order to take into account the systematic uncertainties not considered by 1971Ch54. Then a weighted average of the eight values above has been calculated using LWEIGHT computer program (version 3). The largest contribution comes from the value of Lindenberg (2001Li17), amounting to 75 %.

The recommended value is the weighted average of 51,929 h, with an external uncertainty of 0,010 and a reduced χ^2 of 1,37.

Experimental 279-keV level half-life values (in ps) are given in Table 2.

Table 2: Experimental 279-keV level half-life.

Reference	Experimental value (ps)
R.E. Azuma (1955Az33)	300 (100)
E. E. Berlovich (1957Be57)	290 (30)
E. Bashandy (1960Ba16)	290 (20)
S. Gorodetzky (1960Go15)	283 (17)
B. Johansson (1960Jo15)	220 (30)
E.C. Pederson (1960Pe16)	282 (8)
A. Schwarzschild (1961Sc04)	281 (6)
J. de Boer (1962De14)	340 (3)
R. Rougny (1964Ro19)	283 (7)
J.C. Palathingal (1967Pa09)	280 (40)
Recommended value	282,3 (37)

The half-life weighted average has been calculated by the LWEIGHT program (version 3).

The evaluators have chosen to take into account for the calculation the ten experimental values shown in Table 2. The Azuma (1955Az33), Johansson (1960Jo15) and de Boer (1962De14) values were rejected by the LWEIGHT program, based on the Chauvenet's criterion, thus they were not used for averaging.

The recommended value is the weighted average of 282,3 ps, with an internal uncertainty of 3,7 and a reduced χ^2 of 0,05.

2.1 Electron Capture Transitions

The electron capture probabilities have been deduced from gamma -ray transition intensity imbalance for each level of the decay scheme.

P_K , P_L , P_M values have been calculated for 1st forbidden and 1st forbidden unique electron-capture transitions in the decay of ²⁰³Pb to the excited states in ²⁰³Tl using the LOGFT computer program.

2.2 g Transitions

Probabilities

The absolute transition probabilities have been deduced from the relative γ -ray emission intensities (see 5.2 Gamma ray emission), the internal conversion coefficients and the normalization of the decay scheme to an absolute radiation intensity scale.

Multipolarity and internal conversion coefficients

Multipolarities of γ -ray transitions in decay of ²⁰³Tl are from 2005Ko20:

279-keV γ -ray : M1 + E2, with $\delta = +1,17$ (6)

401-keV γ -ray : M1 + E2, with $\delta = 0,030$ (3) (1965Ka02)

680-keV γ -ray : E2

The internal conversion coefficients (ICC's) for these γ -ray transitions have been calculated using the BRICC computer program, which interpolates the new values in 2006Ra03.

For the 279-keV γ -ray, the evaluators have chosen to follow the recommendations of H. H. Hansen (1985HaZA). The 279-keV γ -ray transition is M1(l -forbidden) + E2. It takes place between the $d_{3/2}$ and $s_{1/2}$ shell model proton configurations. Thus nuclear penetration is significant (see 1979Ha21). The forbidness applies only to the M1 component. Therefore, the evaluators have chosen to use experimental values for α . The experimental data set given by 1985HaZA to determine α_T and α_K are included in Tables 3 and 4, respectively.

Table 3: Experimental values of α_T used by 1985HaZA.

Reference	Original value	Revised by Hansen (1985HaZA) and used value.	Comments
1960Pe22	0,227 (8)		Not used.
1962Ta06	0,2262 (19)	0,2273 (24)	The authors revised their values.
1965Ra12	0,210 (30)		Not used.
1965Wa13	0,222 (15)		Not used.
1971WaYL	0,2267 (7) 0,2240 (9)	0,2267 (16)	The author gives 2 results without explaining the reason of the discrepancy. Hansen has chosen the higher one, with the sum of their uncertainties quoted for both results.
1974Ha29	0,2279 (24)	0,2279 (24)	
2000Sc05	0,2250 (12)	0,2250 (12)	
Recommended value		0,2261 (8)	$\chi^2 = 0,60$.

Hansen's study provides, together with three experimental values, an α_T average of 0,2271 (12). The evaluators have included the most recent measurement of 2000Sc05 (0,2250 (12)) in their evaluation and, with four experimental values (1962Ta06, 1972WaYL, 1974Ha29, 2000Sc05), a weighted average has been calculated using the LWEIGHT computer program (version 3). The recommended value is the weighted average of 0,2261, with an internal uncertainty of 0,0008 and a reduced χ^2 of 0,60.

Table 4: Experimental values of α_K and α_L .

Reference	Original value of α_K	Revised by Hansen (1985HaZA) and used value.	Original value of α_L (10^{-2})	Comments
1952He18	0,23 (10)			Not used.
1954Th17	0,154 (15)			Not used.
1954Wa12	0,15 (1) 0,141 (15)			Not used.
1955Do12	0,147 (2)			Not used.
1955Ma40	0,205 (20)			Not used.
1956No26	0,159 (4)			Not used.
1956Of03	0,150 (10)		4,8 (3)	Not used.
1956Wa30	0,164 (5)	0,164 (5)	4,90 (17)	
1956Wo09	0,130 (10)			Not used.
1958Ni28	0,163 (3)	0,163 (3)	4,87 (12)	
1960Pe22	0,163 (6)	0,163 (6)		
1960Ra04	0,195 (14)			Not used.
1960St21	0,160 (15)			Not used.
1961Hu15	0,1750 (36)			Not used.
1961Su10	0,164 (4)	0,164 (4)	4,49 (34)	
1962Ta06	0,1633 (17)	0,1642 (21)		The authors revised their values.
1963Bu09	0,168 (8)	0,165 (9)		Result had to be corrected for ω_K .
1963Cr14	0,162 (3)	0,162 (3)		
1964He19	0,163 (3)	0,163 (3)	4,84 (6)	
1965Ra12	0,158 (24)			Not used.
1967Bo47	0,14 (3)			Not used.
1968Ra26	0,179 (13)			Not used.
1968Sa22	0,156 (7)			Not used.
1974Ha29	0,1653 (17)	0,1653 (17)	4,75 (13)	
Recommended values		0,1640 (10)	4,837 (48)	$\chi^2 = 0,16$; $\chi^2 = 0,22$

For the α_K recommended value, the evaluators, following the recommendations of H. H. Hansen (1985HaZA), used only nine experimental values with their associated uncertainties in the weighted average calculation, using the LWEIGHT computer program (version 3). A recommended value of 0,1640 for α_K (279-keV γ -ray) is a weighted average, with an internal uncertainty of 0,0010 and a reduced χ^2 of 0,16.

Evaluators' recommended α_L is 4,837 (48) 10^{-2} (reduced $\chi^2 = 0,22$), weighted average of values from: A. H. Wapstra (1956Wa30), G. J. Nijgh (1958Ni28), Z. Sujkowski (1961Su10), C. J. Herrlander (1964He19) and H. H. Hansen (1974Ha29).

3 Atomic Data

Atomic values, ω_K , ω_L and n_{KL} , are from Schönfeld and Janssen (1996Sc06).

3.1 X rays and Auger electrons

The X-ray and Auger electrons relative probabilities have been calculated from γ -ray data by using the EMISSION computer program.

4 Electron Emissions

The Auger electrons emission probabilities have been calculated from γ -ray data using the EMISSION computer program.

5 Photon emissions

5.1 K x-rays

X-ray emissions probabilities have been calculated from γ -ray data using the EMISSION computer program.

5.2 Gamma-ray emissions

The measured energies of γ -ray emissions are given in Table 6.

Table 6 : The measured energies of γ -ray emissions, in keV.

γ -ray	1954Pr04	1954Wa12	1958Ni28	1964He19	1969Cl11	1978He21	2000He14 (evaluated)	Recommended values (keV)
$\gamma_{1,0}$	280 (5)	279 (1)	279,12 (5)	279,16 (2)	279,16 (2)	279,1967 (12)	279,1952 (10)	279,1952 (10)
$\gamma_{2,1}$	400 (7)	400 (2)	403,8 (3)	401,27 (5)	401,28 (40)	401,325 (10)	401,320 (3)	401,320 (3)
$\gamma_{2,0}$	685 (10)	678 (3)			680,7 (6)	680,514 (10)	680,515 (3)	680,515 (3)

The evaluators have adopted the recommended values of R. G. Helmer (2000He14).

The measured relative emission intensities listed in Table 7 are given in values relative to 100 for the 279-keV γ ray.

Table 7: Measured relative γ emission intensity in %.

Energie (keV)	1954Pr04	1954Wa12	1989Ne05	Recommended value
279	100	100	100	100
401	4,7 (3)	4,30 (8)	4,14 (8)	4,24 (8)
680	0,87 (10)	0,80 (1)	0,932 (22)	0,932 (22)

For the 401-keV γ -ray, the recommended value is a weighted average (with an external uncertainty) calculated using the LWEIGHT computer program with these three experimental values. For the 680 -keV γ -ray, the calculation using the LWEIGHT computer program showed that the data are discrepant, so the evaluators have chosen to use the most recent and precise result of Zs. Németh (1989Ne05).

The normalization factor to convert the relative emission intensities to absolute emission intensities is calculated using the formula:

$$N = \left(\frac{100}{(\sum (1 + \alpha_T) P_{rel})} \right) \times 100$$

where the sum is over all the γ transitions to the ground state and α_T is the relevant coefficient. In this case, the contributions are from the 279 - and 680 -keV γ transitions. The uncertainty was calculated through the propagation on the formula given above.

From the recommended α_T (Table 5) and the evaluated relative emission intensities (Table 7), the deduced normalization factor is **80,94 (5)**.

The evaluated relative and absolute γ -ray emission intensities are given in Table 8.

Table 8 : Evaluated relative and absolute γ -ray emission intensities, in %.

Energy (keV)	Relative emission intensity	Absolute emission intensity
279	100	80,94 (5)
401	4,24 (8)	3,43 (6)
680	0,932 (22)	0,754 (18)

6 References

- 1941Fa04 – K. Fajans, A. F. Voigt, Phys. Rev. 60(1941)619 [$T_{1/2}$ (Pb-203)].
- 1952He18 – R. L. Heath, P. R. Bell, Phys. Rev. 87(1952)176A [$\alpha_K(279\text{-keV } \gamma\text{-ray})$].
- 1954Pr04 – J. R. Prescott, Proc. Phys. Soc. (London) A67(1954) 254 [$T_{1/2}$ (Pb-203), E_γ , I_γ].
- 1954Th17 – S. Thulin, K. Nybö, Arkiv för Fysik 7(1954)289 [$\alpha_K(279\text{-keV } \gamma\text{-ray})$].
- 1954Wa12 – A. H. Wapstra, D. Maeder, G. J. Nijgh, L. Th. M. Ornstein, Physica 20(1954)169 [$\alpha_K(279\text{-keV } \gamma\text{-ray})$, E. C. branching, E_γ , I_γ].
- 1955Az33 – R. E. Azuma, G. M. Lewis, Phil. Mag. 46(1955)1034 [$T_{1/2}$ (279-keV level, Tl-203)].
- 1955Do12 – R. K. Doerner, A. H. Weber, Phys. Rev. 99(1955)672A [$\alpha_K(279\text{-keV } \gamma\text{-ray})$].
- 1955Ma40 – N. Marty, Comptes Rendu 240(1955)291 [$\alpha_K(279\text{-keV } \gamma\text{-ray})$].
- 1956No26 – C. Nordling, K. Siegbahn, E. Sokolowski, Nucl. Phys. 1(1956)326 [$\alpha_{K,L}(279\text{-keV } \gamma\text{-ray})$].
- 1956OF03 – Z. O’Friel, A. H. Weber, Phys. Rev. 101(1956)1076 [$\alpha_K(279\text{-keV } \gamma\text{-ray})$].
- 1956Wa30 – A. H. Wapstra, G. J. Nijgh, Nucl. Phys. 1(1956)245 [$\alpha_{K,L}(279\text{-keV } \gamma\text{-ray})$].
- 1956Wo09 – J. L. Wolfson, Can. J. Phys. 34(1956)256 [$\alpha_K(279\text{-keV } \gamma\text{-ray})$].
- 1957Be57 – E. E. Berlovich, G. V. Dubinkin, Soviet Phys. JETP 5(1957)164 [$T_{1/2}$ (279-keV level, Tl-203)].
- 1958Bart – A. A. Bartlett, G. Rebka, Bull. Am. Phys. Soc. 3(2)(1958)64 [$T_{1/2}$ (Pb-203)].
- 1958Ni28 – G. J. Nijgh, A. H. Wapstra, L. Th. M. Ornstein, N. SalomonsGrobbe, J. R. Huizenga, Nucl. Phys. 9(1958/1959)528 [$\alpha_{K,L}(279\text{-keV } \gamma\text{-ray})$, E_γ].
- 1960Ba16 – E. Bashandy, T. R. Gerholm, J. Lindskog, Arkiv för Fysik 17(1960)421 [$T_{1/2}$ (279-keV level, Tl-203)].
- 1960De04 – B. I. Deutch, N. Goldberg, Phys. Rev. 117(1960)818 [$\delta(401\text{-keV } \gamma\text{-ray})$].
- 1960Go15 – S. Gorodetzky, R. Manquenouille, R. Richert, A. Knipper, Comptes Rendu 251(1960)65 [$T_{1/2}$ (279-keV level, Tl-203)].
- 1960Jo15 – B. Johansson, T. Alväger, Arkiv för Fysik 17(1960)163 [$T_{1/2}$ (279-keV level, Tl-203)].
- 1960Pe16 – E. C. B. Pederson, R. E. Bell, Nucl. Phys. 21(1960)393/ Corrigendum: Nucl. Phys. 29(1962) 694 [$T_{1/2}$ (279-keV level, Tl-203)].
- 1960Pe22 – R. W. Peelle, ORNL– 3016 (1960)116 / Nucl. Sci. Abstr. 15(1961)872, abstr. 6771 [$\alpha_{T,K}(279\text{-keV } \gamma\text{-ray})$].
- 1960Ra04 – M. K. Ramaswamy, P. S. Jastram, Nucl. Phys. 15(1960)510 [$\alpha_K(279\text{-keV } \gamma\text{-ray})$].
- 1960St21 – R. Stockendal, Arkiv för Fysik 17(1960)579 [$\alpha_K(279\text{-keV } \gamma\text{-ray})$].
- 1961Ge01 – T. R. Gerholm, B. G. Pettersson, B. van Nooijen, Z. Grabowski Nucl. Phys. 24(1961)177 [$\delta(401\text{-keV } \gamma\text{-ray})$].
- 1961Hu15 – J. P. Hurley, J. M. Ferguson, Nucl. Phys. 27(1961)75/ Addendum Nucl. Phys. 31(1962)690 [$\alpha_K(279\text{-keV } \gamma\text{-ray})$].
- 1961Pe12 – L. Persson, Z. Sujkowski, Arkiv för Fysik 19(1961)309 [$T_{1/2}$ (Pb-203)].
- 1961Sc04 – A. Schwarzschild, J. V. Kane, Phys. Rev. 122(1961)854 [$T_{1/2}$ (279-keV level, Tl-203)].
- 1961Su10 – Z. Sujkowski, Arkiv för Fysik 20(1961)243 [$\alpha_K(279\text{-keV } \gamma\text{-ray})$].
- 1962De14 – Th. J. de Boer, H. Voorthuis, J. Blok, Physica 28(1962)417 [$T_{1/2}$ (279-keV level, Tl-203)].
- 1962Ta06 – J. G. V. Taylor, Can. J. Phys. 40(1962)383 [$\alpha_T(279\text{-keV } \gamma\text{-ray})$].
- 1963Bu09 – R. Burmeister, H. Graber, J. Schintlmeister, R. Weibrecht, Nucl. Phys. 42(1963)56 [$\alpha_K(279\text{-keV } \gamma\text{-ray})$].
- 1963Cr14 – W. L. Croft, B.-G. Pettersson, J. H. Hamilton, Nucl. Phys. 48(1963)267 [$\alpha_K(279\text{-keV } \gamma\text{-ray})$].

- 1964He19 – C. J. Herrlander, R. L. Graham, Nucl. Phys. 58(1964)544 [δ (401-keV γ -ray), $\alpha_{K,L}$ (279-keV γ -ray), E_γ].
- 1964Ro19 – R. Rougny, J. J. Samueli, A. Sarazin, J. de Physique 25(1964)989 [$T_{1/2}$ (279-keV level, Tl-203)].
- 1965Ka02 – E. Karlsson, E. Matthias, S. Gustafsson, K. Johansson, Å. G. Svensson, S. Ogaza, P. da Rocha Andrade, Nucl. Phys. 61(1965)582 [δ (401-keV γ -ray)].
- 1965Ra12 – M. Raja Rao, S. Jnanananda, Nucl. Instrum. Meth. 35(1965)261 [$\alpha_{T,K}$ (279-keV γ -ray)].
- 1965Wa13 – A. Walthert, E. Baumgartner, P. Huber, Helv. Phys. Acta 38(1965)514 [α_T (279-keV γ -ray)].
- 1967Bo47 – H. E. Bosch, E. Szichman, A. Baseggio, R. Dolinkue, Nucl. Instrum. Meth. 52(1967)289 [α_K (279-keV γ -ray)].
- 1967Pa09 – J. C. Palathingal, M. L. Wiedenbeck, Nucl. Phys. A101(1967)193 [$T_{1/2}$ (279-keV level, Tl-203)].
- 1968Ra26 – M. S. Rajput, Current Sci. (India) 37(1968)639 [α_K (279-keV γ -ray)].
- 1968Sa22 – H. S. Sahota, B. S. Ghumman, B. S. Sood, Current Sci. (India) 37(1968)42 [α_K (279-keV γ -ray)].
- 1969Cl11 – J. E. Cline, IN – 1130(1969)39 [E_γ].
- 1971Ch54 – G. A. Chackett, K. F. Chackett, J. B. Welborn, Int. J. Appl. Rad. Isot. 22(1971)715 [I_γ (Pb 203)].
- 1971WaYL – K.F. Walz, H. M. Weiss, E. Funck, PTB Jahresbericht (1971)150 [α_T (279-keV γ -ray)].
- 1974Ha29 – H. H. Hansen, D. Mouchel, Z. Phys. 267(1974)371 [$\alpha_{T,L}$ (279-keV γ -ray)].
- 1978He21 – R. G. Helmer, R. C. Greenwood, R. J. Gehrke, Nucl. Instrum. Meth. 155(1978)189 [E_γ].
- 1979Ha21 – H.H. Hansen. Z.Phys. A291 (1979) 43 [α_T (279-keV γ -ray)].
- 1980Ho17 – H. Houtermans, O. Milosevic, F. Reichel, Int. J. Appl. Rad. Isot. 31(1980)153 [$T_{1/2}$ (Pb-203)].
- 1982HoZJ – D. D. Hoppes, NBS – SP – 626(1982)85 [$T_{1/2}$ (Pb-203)].
- 1985HaZA – H. H. Hansen, European Appl. Res. Rept.–Nucl. Sci. Technol. 6(1985)777 [$\alpha_{T,K}$ (279-keV γ -ray)].
- 1989Ne05 – Zs. Németh, T. Sekine, Y. Yoshihara, Appl. Radiat. Isot. 40(1989)519 [E. C. branching, I_γ].
- 1996Sc06 – E. Schönfeld, H. Janssen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
- 2000He14 – R. G. Helmer, C. van der Leun, Nucl. Instrum. Meth. Phys. Res. A450(2000)35 [E_γ].
- 2000Sc05 – E. Schönfeld, H. Janssen, R. Klein, Appl. Radiat. Isot. 52(2000)955 [α_T (279-keV γ -ray)].
- 2001Li17 – K. Lindenberg, F. Neumann, D. Galaviz, T. Hartmann, P. Mohr, K. Vogt, S. Volz, A. Zilges, Phys. Rev. C63(2001)047307 [$T_{1/2}$ (Pb-203)].
- 2002Un02 – M. P. Unterweger, Appl. Radiat. Isot. 56(2002)125 [$T_{1/2}$ (Pb-203)].
- 2003Au03 – G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].
- 2005Ko20 – F. G. Kondev, Nucl. Data Sheets 105(2005)1 [Spin, multipolarity, level energy].
- 2006Ra03 - S. Raman, M. Ertugrul, C. W. Nestor, Jr., M. B. Trzhaskovskaya, At. Data Nucl. Data Tables 92(2006)207 [Theoretical ICC].

²⁰⁴Tl – Comments on evaluation
by M. M. Bé and V. Chisté

The electron capture transition to the Hg -204 ground state is first forbidden unique, so the P_K/P_L ratio strongly depends on the decay energy. In this evaluation the Q^+ value from Audi and Wapstra has been adopted. However, if this value changes, P_K and P_L , as well as the decay branching ratios, must be reevaluated.

Nuclear Data

Spin and parity assignments are from Schmorak (1994Sc24).

Experimental Q^+ values

The following experimental values have been noted from publications :

Reference	Value in keV	Uc	
Biavati(1962Bi04)	310	10	393 quoted in Klein
Leutz (1962Le05)	410	+30 – 23	As quoted by Christmas
Christmas (1964Ch17)	313	+17 – 14	
Klein (1966Kl02)	324	+21 – 16	
Lancman (1973La17)	385	20	
Zide (1979Zi02)	357	15	
Audi (1995Au04)	347,5	15	
Audi (2002)	345,0	13	Adopted

In the 1995Au04 publication, Audi recommended 347,5(15) keV for the Q^+ energy, but a new mass determination of Hg-204 (2002Be) leads to the value of 345,0(13) keV (Audi on the AMDC web site) from the atomic mass differences. As these mass measurements were performed with Penning trap facility, the resulting Q value is considered to be more reliable than the other values quoted in the above table.

Adopted Q values

Q^- value is from Audi and Wapstra (1995Au04)

$Q^- = 763,72 (18) \text{ keV}$

$Q^+ = 345,0 (13) \text{ keV}$

Half-life

Reference	Value (years)	Uc	Comments
Anspach (1965An07)	3,754	0,004	
Horroks (1968Ho07)	3,825	0,003	
Bortels (1969Bo24)	3,774	0,008	Uc for 1 σ
Jordan (1969Jo02)	3,7730	0,0028	Uc for 1 $\sigma \times 1,5$
Harbottle (1970Ha32)	3,793	0,005	
Adopted	3,788	0,015	

The uncertainty for one standard deviation given by Jordan has been multiply by 1,5. The set of five values quoted above is quite discrepant with a reduced $-\chi^2$ of 64,3. The Lweight program has calculated a weighted average of 3,788 years with an external uncertainty of 0,013, which was increased to 0,015 to include the most precise value.

Electron capture sub shell probabilities

The adopted values have been calculated with the LOGFT program for a unique 1st forbidden transition and $Q = 345,0 (13) \text{ keV}$.

$$P_K = 0,5843(14) ; P_L = 0,3024(10) ; P_{M^+} = 0,1133(5)$$

Several measurements of the P_L/P_K ratio were carried out :

Reference	P_L/P_K	P_K/P_{b^-}	Branching ratio %
Christmas (1964Ch17)	0,600 (55)	0,01590 (36)	2,54 (12)
Joshi (1961JO12)	0,42 (5)	0,0155 (10)	
Leutz (1962Le05)	0,41 (3)		
Klein (1966Kl02)	0,55 (5)	0,0153 (5)	2,15 (6)
Weighted mean	0,47 (3)		
Adopted values	0,518 (2)		2,92 (13)

Branching ratios

From the Xk emissions intensities measured by Schötzig (1990Sc08), $I_{XK} = 1,64(7)$, and using $P_K = 0,5843(14)$ and $\omega_K = 0,962(4)$, the electron capture branching ratio $P\epsilon$ becomes:

$$P\epsilon = I_{XK} / (P_K \times \omega_K) = 2,92(13) \%$$

$$\text{And then } P\beta^- = 97,08(13) \%$$

Atomic data

All the atomic data : $\omega_K = 0,962(4)$ etc. and ratio K_β/K_α etc. are from Schönfeld (1996Sc06).

Photons emissions*X-ray emissions*

The X_K emission intensities are those measured by Schötzig.

Reference		I(%)	Uc
Schotzig (1990Sc08)	Hg- $K_{\alpha 2}$	0,474	0,020
	Hg- $K_{\alpha 1}$	0,812	0,034
	Hg- $K_{\beta 1}$	0,273	0,010
	Hg- $K_{\beta 2}$	0,081	0,003
	Pb- $K_{\alpha 2}$	$4,4 \cdot 10^{-3}$	0,3
	Pb- $K_{\alpha 1}$	$6,1 \cdot 10^{-3}$	0,3
	Pb- $K_{\beta 1}$	$2,7 \cdot 10^{-3}$	0,2
	Pb- $K_{\beta 2}$	$7,3 \cdot 10^{-4}$	0,2

The X_L emission intensities have been calculated by using the Emission program after addition of the PL1, etc. values.

The ratio $K\text{-Auger} / \beta^- = 6,7(8) \cdot 10^{-4}$, deduced from the evaluated data, can be compared with the measured value, $K\text{-Auger} / \beta^- = 4,9(28) \cdot 10^{-4}$ given by Park and Christmas (1967Pa08).

Internal bremsstrahlung

Internal bremsstrahlung accompanying capture of orbital electrons is about (3×10^{-5}) photons per K capture.

References

- 1962Bi04 M. H. Biavati, S. J. Nassiff, C. S. Wu, Phys. Rev. 125,4 (1961) 1364 ; Q
 1961Jo12 B. R. JOSHI, Proc. Phys. Soc. 77 (1961) 1205 ; PL/PK
 1962Le05 H. LEUTZ, K. ZIEGLER, Z. Phys. 166 (1962) 582 ; Q, P_L / P_K
 1964Ch17 P. CHRISTMAS, Nucl. Phys. 55 (1964) 577 ; Q
 1965An07 S. C. Anspach et al., Report NBS 260-9 (1965) ; $T_{1/2}$
 1966KI02 H. KLEIN, H. LEUTZ, Nucl. Phys. 79 (1966) 27 ; Q
 1967Pa08 J. J. H. Park, P. Christmas, Can. J. Phys. 45 (1967) 2621 ; $K\text{-Auger} / \beta^-$
 1968Ho07 D. L. HORROCKS, Nucl. Phys. A110 (1968) 238 ; $T_{1/2}$
 1969Bo24 G. BORTELS, Int. J. Appl. Radiat. Isotop. 20 (1969) 613 ; $T_{1/2}$
 1969Jo02 K. C. JORDAN, J. H. BIRDEN, B. C. BLANKE, J. Inorg. Nucl. Chem. 31 (1969) 2641 ; $T_{1/2}$
 1970Ha32 G. HARBOTTLE, Radiochim. Acta 13 (1970) 132 ; $T_{1/2}$
 1973La17 H. Lancman, A. Bond, Phys. Rev. C 7, 6 (1973) 2600 ; Q
 1979Zi02 A. ZIDE, H. LANCMAN, Phys. Rev. C19 (1979) 1053 ; Q
 1990Sc08 U. Schötzig, Nucl. Instrum. Methods A286 (1990) 523 ; KX emission intensities
 1994Sc24 M. R. Schmorak, Nucl. Data Sheets 72,3 (1994) 409 ; spin and parity
 1995Au04 G. Audi, A. H. Wapstra, Nucl. Phys. A 595 (1995) 409 ; Q
 1996Sc06 E. Schönfeld, H. Janssen, Nucl. Instrum. Methods A369 (1996) 527 ; atomic data
 2002Be I. Bergström et al., Nucl. Instrum. Methods A487 (2002) 618 ; Q^+

²⁰⁶Hg - Comments on evaluation of decay data by F. G. Kondev

This evaluation was completed in May 2011 with a literature cut off by the same date. The Saisinuc software (2008DuZX) and associated supporting programs were used in assembling the data following the established protocol within the DDEP collaboration.

1 Decay Scheme

The nuclide ²⁰⁶Hg disintegrates 100 % by β⁻ emissions. The strongest β⁻-decay branch of 62 (7) % populates the J^π = 0⁻ ground state of the daughter nuclide ²⁰⁶Tl. The level schemes of ²⁰⁶Hg and ²⁰⁶Tl are based on the ENSDF evaluations of Browne (1999Br39) and Kondev (2008Ko21).

2 Nuclear Data

Q(β⁻) value is taken from the evaluation of Audi *et al.* (2003Au03).

The experimental half-life data for the ²⁰⁶Hg ground state are presented in Table 1. These data were evaluated using different techniques (see for example 1992Ra08, 1994Ka08 and 2004Mb11 and references therein) and the results are presented in Table 2. The LRSW value of T_{1/2} = 8.32 (7) min is recommended here with χ²_v = 3.22 (χ²_v = χ²/N-1) which is smaller than the critical value of χ²_{v,crit} = 4.61 (99 % confidence level). The lifetimes assigned to the excited states of the daughter nuclide ²⁰⁶Tl are taken from the ENSDF evaluation of Browne (1999Br39).

Table 1. Experimental data for the half-life of ²⁰⁶Hg.

Author	T _{1/2} (min)	Used in the evaluation
1961Nu01	7.5 (10)	No
1962Ka27	8.5 (1)	Yes
1964Wo05	8.1 (4)	Yes
1968Wo08	8.15 (10)	Yes

Table 2. Evaluated values for the half-life of ²⁰⁶Hg.

Method/Author ^{a)}	Evaluated T _{1/2} (min)	χ ² /N-1
UWM	8.25 (13)	3.70
WM	8.32 (7)	3.22
LRSW	8.32 (7)	3.22
NRM	8.27 (8)	2.30
RM	8.18 (9)	0.38
1999Br39	8.15 (10)	

^{a)} UWM – Unweighted Mean; WM – Weighted Mean; LRSW – Limitation of Relative Statistical Weight; NRM – Normalized Residual; RM – Rajeval.

2.1 β^- Transitions

Information of level and maximum β^- -decay energies, $E_{\beta \text{ max}}$, and β^- -decay transition probabilities, P_{β} , and $\log ft$ values is presented in Table 3. The $E_{\beta \text{ max}}$ values for the $\beta_{0,2}$ and $\beta_{0,3}$ transitions were determined from $Q(\beta^-)$ (2003Au03) and the excitation energies for the 1^- states, deduced from the corresponding γ -ray transition energies (see section 2.2 and Table 4 for details). The $P_{\beta_{0,2}}$ and $P_{\beta_{0,3}}$ values were deduced from the decay scheme and the corresponding absolute γ -ray transition probabilities, as detailed in section 2.2 and Table 4. It was assumed that no direct β^- -decay feeding takes places to the first excited state at 265.8 keV ($J^\pi = 2^-$), since such a transition is a second-fold forbidden non-unique, and hence, the $\beta_{0,0}$ transition probability was determined as:

$$P_{\beta_{0,0}} = 100 - P_{\beta_{0,2}} - P_{\beta_{0,3}} \quad (1)$$

The $\log ft$ values were calculated using the LOGFT program from the ENSDF evaluation package.

Table 3. Level energies, $E_{\beta \text{ max}}$, P_{β} and $\log ft$ values in decay of ²⁰⁶Hg.

	Level energy (keV)	$E_{\beta \text{ max}}$ (keV)	P_{β} (%)	Nature	$\log ft$
$\beta_{0,0}$	0.0	1308 (20)	62 (7)	First forbidden non-unique	5.67 (10)
$\beta_{0,2}$	304.896 (6)	1003 (20)	35 (7)	First forbidden non-unique	5.24 (10)
$\beta_{0,3}$	649.42 (5)	659 (20)	3.0 (4)	First forbidden non-unique	5.41 (6)

2.2 γ Transitions

The γ -ray transition energies, multiplicities, absolute transition probabilities and electron internal conversion coefficients are presented in Table 4.

Table 4. Energies, multiplicities, absolute transition probabilities and electron internal conversion coefficients for γ -ray transitions following β^- -decay of ²⁰⁶Hg.

	Energy (keV)	$P_{\gamma+ce}$ (%)	Multi- polarity	α_K	α_L	α_M	α_T
$\gamma_{1,0}$	265.832 (5)	0.014 (7)	E2	0.0855 (12)	0.0561 (8)	0.01440 (21)	0.1603 (23)
$\gamma_{2,0}$	304.896 (6)	36 (7)	M1	0.308 (5)	0.0519 (8)	0.01211 (17)	0.375 (6)
$\gamma_{3,0}$	649.42 (5)	2.3 (3)	M1	0.0412 (6)	0.00681 (10)	0.001585 (23)	0.0501 (7)
$\gamma_{3,2}$	344.52 (17)	0.70 (14)	M1	0.221 (4)	0.0371 (6)	0.00866 (13)	0.269 (4)
$\gamma_{3,1}$	383.59 (6)	0.014 (7)	M1(+E2)	0.10 (7)	0.021 (7)	0.0050 (15)	0.13 (8)

The γ -ray transition energies and multiplicities are taken from the ENSDF evaluation of Browne (1999Br39). The $\gamma(3,1)$ energy is deduced from the adopted level energies difference. The electron internal conversion coefficients were calculated using a program supplied by the Saisinuc software (2008DuZX) which uses interpolated values of Band et al (2002Ba85) with the hole being taken into account. These are consistent with values given by the BrIcc program (2008Ki07). The $P_{\beta_{0,2}}$ value was deduced from the reported in 1968Wo08 absolute γ -ray transition probabilities for the 304.9 keV transition of $P_{\gamma_{2,0+ce}(304.9\gamma)} = 36 (7) \%$ and by taking into account a small feeding from the 1^- level at 649.4 keV via the 344.5 keV γ -ray transition. The $P_{\gamma+ce}$ values for the $\gamma(1,0)$ and $\gamma(2,1)$ transitions were determined from the absolute γ -ray emission

probabilities, P_γ , shown in Table 5, and the total electron internal conversion coefficients as:

Table 5

E level (keV)	Relative Intensity				Abs. Total Int. (%)
	E_γ (keV)	1970As05	1969La18	1976TuZY	1968Wo08
265.832	265.832 (5)				
304.896	304.896 (6)	100 (1)			36 (7)
649.42	344.52 (17)	2.4 (1)	1.4	1.4	
	383.59 (6)			0.011	
	649.42 (5)	8.4 (17)	5.6	7.7	5 (2)

3 Atomic Data

The atomic data (fluorescence yields, X-ray energies and relative probabilities, and Auger electrons energies and relative probabilities) were provided by the Saisinuc software (2008DuZX). Details regarding the origin of these data can be found in 1996Sc06, 1998ScZM, 1999ScZX, 2000Sc47 and 2003De44.

4 Emissions

4.1 Photon emissions

The number of γ rays per 100 disintegrations was evaluated from the available experimental data, as described in section 2.2 (see also Table 5).

5 Electron emissions

The energies of the conversion electrons were calculated from the γ -ray transition energies presented in Table 4 and the corresponding electron shell binding energies (1977La19). The number of conversion electrons of type $x=T,K$ and L where T stands for total, K and L for K - and L -shell electrons, per 100 disintegrations was calculated from the recommended in the present evaluation (see Table 5) numbers of photons per 100 disintegrations, $P_{\gamma 1,0}$, and the corresponding electron internal conversion coefficients (see Table 4), $\alpha_{x1,0}$: $ec_{1,0x} = P_{\gamma 1,0} \times \alpha_{x1,0}$.

The number of K and L Auger electrons per 100 disintegrations, $P(e_{AK(L)})$ was calculated from the number of vacancies in the K and L shells and the corresponding $P_{XK(L)}$ yield: $P(e_{AK}) = N_K - P_{XK}$ and $P(e_{AL}) = N_L - P_{XL}$.

6 References

- 1961Nu01 M. Nurmia, P. Kauranen, M. Karras, A. Siivola, A. Isola, G. Graeffe, A. Lyyjen, Nature 190(1961)427 (Half-life)
- 1968Wo08 G. K. Wolf, Nucl. Phys. A116(1968)387 (Half-life, gamma-ray energy and emission probabilities)
- 1969La18 R. C. Lange, G. R. Hagee, A. R. Campbell, Nucl. Phys. A133(1969)273 (Gamma-ray energy and emission probabilities)

- 1969La18 R. C. Lange, G. R. Hagee, A. R. Campbell, Priv. Comm. (1971) (Gamma-ray energy and emission probabilities)
- 1970As05 G. Astner, G. K. Wolf, , Nucl. Phys. A147(1970)481 (Gamma-ray energy and emission probabilities)
- 1976TuZY D. G. Tuggle, Report LBL – 4460, PhD Thesis Univ. California (1976) (Gamma-ray energy and emission probabilities)
- 1977La19 F. P. Larkins, At. Data Nucl. Data Tables 20(1977)311 (K Auger electron energies)
- 1992Ru08 M. U. Rajput, T. D. MacMahon, Nucl. Instrum. Meth. Phys. Res. A312(1992)289 (Evaluation techniques)
- 1994Ka08 S. I. Kafala, T. D. MacMahon, P. W. Gray, Nucl. Instrum. Meth. Phys. Res. A339(1994)151 (Evaluation techniques)
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 (K-shell fluorescence yields)
- 1998ScZM E. Schönfeld, G. Rodloff, Report PTB-6.11-98-1 Braunschweig (1998) (K Auger electron energies)
- 1999ScZM E. Schönfeld, G. Rodloff, Report PTB-6.11-1999-1 Braunschweig (1999) (K X-ray energies and relative emission probabilities)
- 1999Br39 E. Browne, Nucl. Data Sheets 88(1999)29 (Nuclear levels)
- 2000Sc47 E. Schönfeld, H. Janßen, Appl. Radiat. Isot. 52(2000)595 (EMISSION program and X-ray and Auger electron emission probabilities and energies)
- 2002Ba85 I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr., P. O. Tikkanen, S. Raman, Atomic Data Nucl. Data Tables 81(2002)1 (ICC's)
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)1 (Q value)
- 2003De44 R. D. Deslattes, E. G. Kessler Jr., P. Indelicato, L. de Billy, E. Lindroth, J. Anton, Rev. Mod. Phys. 75(2003)35 (K and L X-ray energies)
- 2004Mb11 T. D. MacMahon, A. Pearce, P. Harris, Appl. Radiat. Isot. 60(2004)275 (Evaluation techniques)
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr. , Nucl. Instrum. Meth. Phys. Res. A589(2008)202 (Theoretical ICCs)
- 2008Ko21 F. G. Kondev, Nucl. Data Sheets 109(2008)1527 (Nuclear levels)
- 2008DuZX C. Dulieu, M. M. Bé, V. Chisté, Proc. Int. Conf. On Nuclear Data for Science and Technology, Nice, France, 22-27 April 2007(2008)97 (SAISINUC software)

²⁰⁶Tl - Comments on Decay Data Evaluation

by F.G. Kondev

This evaluation was completed in September 2006 with a literature cut off by the same date. The Saisinuc software (2002BeXX) and associated supporting programs were used in assembling the data following the established protocol within the DDEP collaboration.

1. Decay Scheme

The nuclide ²⁰⁶Tl ($J^\pi=0^-$) disintegrates 100 % by β^- emissions. The strongest β^- -decay branch of 99.885 (14) % populates the $J^\pi=0^+$ ground state of the daughter nuclide ²⁰⁶Pb. The level schemes of ²⁰⁶Tl and ²⁰⁶Pb are based on the ENSDF evaluation of Browne (1999Br39).

2. Nuclear Data

$Q(\beta^-)$ value is taken from the evaluation of Audi *et al.* (2003Au03).

The experimental half-life data for the ²⁰⁶Tl ground state are presented in Table 1. These data were evaluated using different techniques (see for example 1992Ra08, 1994Ka08 and 2004MaXX and references therein) and the results are presented in Table 2. The value of 1961Nu01 was excluded from the data analysis, since no uncertainty was quoted in the original publication. The LRSW value of $T_{1/2}=4.202$ (11) min is recommended here with $\chi^2_v = 1.54$ ($\chi^2_v = \chi^2/N-1$) which is smaller than the critical value of $\chi^2_{v, \text{crit}} = 2.64$ (99 % confidence level). The lifetimes of the excited states of the daughter nuclide ²⁰⁶Pb are taken from the ENSDF evaluation of Browne (1999Br39).

Table 1. Experimental data for the half-life of ²⁰⁶Tl

Author	$T_{1/2}$, min	Used in the evaluation
1941Fa04	4.23 (3)	Yes
1953Sa11	4.19 (2)	Yes
1959Po64	4.29 (5)	Yes
1961Nu01	4.2	No
1970Fl12	4.27 (5)	Yes
1971Pe03	4.183 (17)	Yes
1972CoYX	4.14 (5)	Yes
1972Gr01	4.2 (2)	Yes
1972Wi18	4.27 (5)	Yes

Table 2. Evaluated values for the half-life of ²⁰⁶Tl

Method/Author ^{a)}	Evaluated T _{1/2} , min	c ² /N-1
UWM	4.222 (19)	2.02
WM	4.202 (11)	1.54
LRSW	4.202 (11)	1.54
NRM	4.202 (11)	1.54
RM	4.202 (11)	1.41
1999Br39	4.200 (17)	

^{a)} UWM – Unweighted Mean; WM – Weighted Mean; LRSW – Limitation of Relative Statistical Weight; NRM – Normalized Residual; RM – Rajeval.

2.1. b⁻ Transitions

The experimental data for the maximum $\beta_{0,0}$ energy, $E_{\beta_{0,0} \max}$, are presented in Table 3. The LRSW value of 1527 (3) keV ($\chi^2_{\nu} = 1.48$ is smaller than $\chi^2_{\nu \text{ crit}} = 4.61$ (99 % confidence level)) is comparable with $Q(\beta^-) = 1532.4$ (6) keV (2003Au03). The $E_{\beta \max}$ values for the $\beta_{0,1}$ and $\beta_{0,2}$ transitions were determined from $Q(\beta^-)$ (2003Au03) and the 2^+ and 0^+ level energies that were deduced from the corresponding transition energies (see section 2.2 and Table 4 for details). The $\beta_{0,1}$ and $\beta_{0,2}$ transition probabilities, P_{β} , were deduced from the decay scheme and the corresponding absolute γ -ray transition probabilities, $P_{\gamma+ce}$, as detailed in section 2.2 and Table 5. The P_{β} value for the $\beta_{0,1}$ transition is an upper limit, since the possible feeding from the 1166.4 keV level ($J^{\pi}=0^+$) via the yet unobserved 363.3 keV γ -ray transition ($\gamma_{2,1}$) was not taken into account. It should be noted that only a limit for $P_{\gamma_{2,1}}$ is reported in the literature (see section 2.2 for details). The $\beta_{0,0}$ transition probability was determined as:

$$P_{\beta_{0,0}} = 100 - P_{\beta_{0,1}} - P_{\beta_{0,2}}.$$

The $\lg ft$ values were calculated using the LOGFT program from the ENSDF evaluation package. The $\lg f$ values are based on the work of Gove and Martin (1971Go40). For the first forbidden $\beta_{0,0}$ transition ($0^- \rightarrow 0^+$) the shape factor was measured by several authors, as shown in Table 3. The fit to the experimental data using the expression $S(W) = 1 + aW + b/W$, where W is the electron energy, yields the shape factor coefficients, a and b , which are also presented in Table 3. The value of $a = -0.020$ (2) (with $b = 0.000$) (1972Wi18) is recommended in the present evaluation. It should be noted that using this parameterization of the shape factor, a $\lg f$ value of 2.85 for the $\beta_{0,0}$ transition ($0^- \rightarrow 0^+$) can be obtained. It is in a good agreement with $\lg f = 2.78$, deduced using the LOGFT program (1971Go40).

Table 3. Measured $E_{\beta_{0,0} \max}$ values and shape factor parameters a and b ($S(W)=1+aW+b/W$) for the first forbidden $0^- \rightarrow 0^+$ decay of ^{206}Tl

Author	a	b	$E_{\beta_{0,0} \max}$, keV	Used in the evaluation
1951Al14			1510 (10)	No
1961Ho17	-0.154	-0.484	1571 (10)	No
1970Fl12	-0.017 (5)	0.030 (9)	1523 (4)	Yes
1971Pe03	0.00 (1)	0.00	1534 (5)	Yes
1972Wi18	-0.020 (2)	0.000	1527 (4)	Yes
Adopted	-0.020 (2)	0.000	1532.4 (6)	

Table 4. Level energies, $E_{\beta \max}$, P_{β} and $\log ft$ values in decay of ^{206}Tl

	Level energy, keV	$E_{\beta \max}$, keV	$P_{\beta} \times 100$	Nature	$\log ft$
$\beta_{0,0}$	0.0	1532.4 (6)	99.885 (14)	First forbidden	5.1775 (13)
$\beta_{0,1}$	803.06 (3)	729.3 (6)	0.0051 (3)	First forbidden Unique	8.60 ^{1U} (3)
$\beta_{0,2}$	1166.4 (5)	366.0 (8)	0.110 (14)	First forbidden	5.99 (6)

2.2 Gamma Transitions and Electron Internal Conversion Coefficients

The γ -ray transition energies, multipolarities, absolute transition probabilities and electron internal conversion coefficients are presented in Table 5.

The γ -ray transition multipolarities are taken from the ENSDF evaluation of Browne (1999Br39). The recommended $\gamma_{1,0}$ transition energy of 803.06 (3) keV is determined as the weighted mean of 803.10 (5) keV (1972Ma63) and 803.04 (3) keV (1996Ra16), the two most precise values reported in the literature. The $\gamma_{2,0}$ transition between the excited 0^+ level and the 0^+ ground state is a pure E0, and hence, there is no γ -ray component associated with the decay of the former level. The transition energy is taken from the work of Draper *et al.* (1977Dr08) where the K-shell conversion electron energy was measured with a Si(Li) detector. The $\gamma_{2,1}$ transition was not observed and its energy is inferred from the energy difference between the excited 0^+ and 2^+ levels. The electron internal conversion coefficients were calculated using a program supplied by the Saisinuc software (2002BeXX) which uses interpolated values of Band *et al.* (2002Ba85) with the hole being taken into account. The $P_{\gamma+ce}$ values for the $\gamma_{1,0}$ and $\gamma_{2,1}$ transitions were determined from the absolute γ -ray emission probabilities, P_{γ} , shown in Table 6, and the total electron internal conversion coefficients as: $P_{g+ce} = P_g \times (1 + a_T)$.

Experimental and evaluated P_{γ} values are shown in Table 6. The LRSW value of $P_{\gamma_{1,0}} = 0.0050$ (3) % ($\chi^2_{\nu} = 2.40$ is smaller than $\chi^2_{\nu \text{ cryt}} = 4.61$ (99 % confidence level)) is recommended for the $\gamma_{1,0}$ transition. As stated above, the $\gamma_{2,1}$ transition was not observed experimentally and only a limit for its absolute

emission probability was given in 1972CoYX and 1972Gr01. The value of $P_{\gamma_{2,1}} < 0.00026 \%$ (1972CoYX) is adopted in the present evaluation. The $\gamma_{2,0}$ transition is a pure E0 ($0^+ \rightarrow 0^+$) and hence $P_{\gamma_{2,0}}$ is zero. The recommended $P_{\gamma+ce}(\gamma_{2,0})$ value here is deduced from the measured absolute KX-ray yield, $P_{KX}(\gamma_{2,0})$, the corresponding fluorescence yield, ω_K , and the K/T conversion electrons ratio. The value of $P_{KX}(\gamma_{2,0}) = 0.09 (1) \%$, deduced as a weighted mean of $0.08 (2) \%$ (1972CoYX) and $0.10 (2) \%$ (1972Gr01) (see Table 6), is adopted in the present work. It should be noted that an electron shake-off component of 0.02% has been taken into account in these values. The K-shell to total conversion electrons ratio of $K/T = 0.85 (6)$ was deduced from $K/L = 5.7 (4)$, a weighted mean of the measured $K/L = 5.61 (38)$ and $6 (1)$ in 1990Tr01 and 1977Dr08, respectively. This value is in very good agreement with that of $K/T = 0.855$, calculated using the electronic factors of $\Omega_K(E0)$ and $\Omega_L(E0)$ that are given by the BRICC program (2005KiZW). Using a K-fluorescence yield value of $\omega_K = 0.963 (4)$ (1996Sc06) one then obtains:

$$P_{g+ce}(g_{2,0}) = P_{ce}(g_{2,0}) = (P_{KX}(g_{2,0}) / \omega_K) / (K/T) = 0.110 (14) \%$$

Table 5. Energies, multipolarities, absolute transition probabilities and electron internal conversion coefficients for γ -ray transitions following β^- -decay of ²⁰⁶Tl

	Energy, keV	$P_{\gamma+ce} \times 100$	Multipolarity	α_K	α_L	α_M	α_N	α_T
$\gamma_{1,0}$	803.06 (3)	0.0051 (3)	E2	0.00801 (24)	0.00174 (5)	$4.19 (13)10^{-4}$	$1.06 (3)10^{-4}$	0.0103 (3)
$\gamma_{2,1}$	363.3 (5)	0.00015 (15)	(E2)	0.0414 (12)	0.0187 (6)	0.00476 (14)	0.00120 (4)	0.066 (2)
$\gamma_{2,0}$	1166.4 (5)	0.110 (14)	E0					

Table 6 Experimental and evaluated γ -ray emission probabilities.

Authors	$P_{g_{1,0}}, \%$	$P_{KX}(g_{2,0}) \%^{a)}$	$P_{g_{2,1}}, \%$	Comment ^{b)}
1968Zo02	0.0055 (5)			Not used
1970Zo02	0.0055 (4)			Expt.
1972CoYX	0.0041 (6)	0.08 (2)	<0.00026	Expt.
1972Gr01	0.004 (1)	0.10 (2)	<0.001	Expt.
Adopted	0.0050 (3)	0.09 (1)	<0.00026	Evaluated

^{a)} Absolute KX-ray yield

^{b)} Expt. – experimental value used in the present evaluation. The 1968Zo02 value is superseded by 1970Zo02

3. Atomic Data

The Atomic data (Fluorescence yields, X-Ray energies and Relative probabilities, and Auger electrons energies and Relative probabilities) were provided by the Saisinuc software (2002BeXX). Details regarding the origin of these data can be found in 1996Sc06, 1998ScZM, 1999ScZX, 2000ScXX and 2003DeXX.

4. Photon Emissions

4.1 X-Ray Emissions

The X-ray yield in β^- decay of ²⁰⁶Tl is produced entirely in the decay of the 1166.4 keV (E0, $0^+ \rightarrow 0^+$) transition. Contributions from the much weaker 803.06 and 363.3 keV transitions can be neglected, since their X-ray yields are several orders of magnitude smaller than that of the 1166.4 keV transition.

For the 1166.4 keV E0 ($0^+ \rightarrow 0^+$) transition, the number of vacancies in the K-shell per 100 disintegrations was determined as:

$$N_K = P_{ceK} = P_{XK} / w_K = 0.090 (10) / 0.963 (4) = 0.093 (11).$$

The corresponding number of vacancies in the L shell per 100 disintegrations was then determined as:

$$N_L = P_{ceL} + n_{KL} \times N_K = 0.0163 (22) + 0.811 (5) \times 0.093 (11) = 0.092 (11) \%$$

where $P_{ceL} = P_{ceK} / (K/L) = 0.0163 (22) \%$ with $K/L = 5.7 (4)$, a weighted mean of 5.61 (1990Tr01) and 6 (1) (1977Dr08). The number of X-rays per 100 disintegrations was then calculated as:

$$P_{XK} = w_K \times N_K \text{ and } P_{XL} = \bar{w}_L \times N_L$$

4.2 Gamma Emissions

The number of γ rays per 100 disintegrations was evaluated from the available experimental data, as described in section 2.2 (see also Table 6).

5. Electron Emissions

The energies of the conversion electrons were calculated from the γ -ray transition energies presented in Table 5 and the corresponding electron shell binding energies (1977La19). For the $\gamma_{1,0}$ transition, the number of conversion electrons of type $x = T, L, M, N$ and O , where T stands for total, L for L-shell electrons, etc., per 100 disintegrations was calculated from the absolute photon intensity ($P_{\gamma_{1,0}}$ per 100 disintegrations) recommended in the present evaluation (see Table 6), and the corresponding electron internal conversion coefficients (see Table 5), $\alpha_{x,1,0}: ec_{1,0,x} = P_{g_{1,0}} \times a_{x,1,0}$. For the $\gamma_{2,0}$ transition, the number of K and L conversion electrons per 100 disintegrations was determined from the measured P_{XK} yield, w_K value and the K/L sub-shell ratio, as detailed in section 4.1.

The number of K and L Auger electrons per 100 disintegrations, $P(e_{AK(L)})$ was calculated from the number of vacancies in the K and L shells and the corresponding $P_{XK(L)}$ yield: $P(e_{AK}) = N_K - P_{XK}$ and $P(e_{AL}) = N_L - P_{XL}$.

REFERENCES

- 1941Fa04 K. Fajans and A.F. Voigt, Phys. Rev. 60 (1941) 619.
(Half-life)
- 1951Al14 D.E. Alburger and G. Friedlander, Phys. Rev. 82 (1951) 977.
(Maximum β^- -decay energy)
- 1953Sa11 B.W. Sargent, L. Yaffe and A.P. Gray, Can. J. Phys. 31 (1953) 235
(Half-life)
- 1959Po64 A. Poularikas and R.W. Fink, Phys. Rev. 115 (1959) 989.
(Half-life)
- 1961Ho17 D.A. Howe and L.M. Langer, Phys. Rev. 124 (1961) 519.
(Maximum β^- -decay energy)
- 1961Nu01 M. Nurmi, P. Kauranen, M. Karras, A. Siivola, A. Isola, G. Graeffe and A. Lyyjy, Nature 190 (1961) 427.
(Half-life)
- 1968Zo02 W.H. Zoller, C. Botteron and W.B. Walters, MIT-905-133 (1968) p.4.
(γ -ray emission probability)
- 1970Zo02 W.H. Zoller and W.B. Walters, J. Inorg. Nucl. Chem. 32 (1970) 2465.
(γ -ray emission probability)
- 1970Fl12 D. Flothmann, R. Lohken, W. Wiesner and H. Rebel, Phys. Rev. Lett. 25 (1970) 1719.
(Half-life, Maximum β^- -decay energy)
- 1971Go40 N.B. Gove and N.J. Martin, Nuclear Data Tables 10 (1971) 205
(log *ft* values)
- 1971Pe03 B.I. Persson, I. Plessner and J.W. Sunier, Nucl. Phys. A167 (1971) 470.
(Half-life, Maximum β^- -decay energy)
- 1972CoYX L.L. Collins, G.D. O'Kelley and E. Eichler, ORNL-4791 (1972) p.14.
(Half-life)
- 1972Gr01 H.C. Griffin and A.M. Donne, Phys. Rev. Lett. 28 (1972) 107.
(Half-life)
- 1972Ma63 J.C. Manthuruthil, D.C. Camp, A.V. Ramayya, J.H. Hamilton, J.J. Pinajian and J.W. Doornebos, Phys. Rev. C6 (1972) 1870.
(Gamma-ray transition energies)
- 1972Wi18 W. Wiesner, D. Flothmann, H.J. Gils, R. Lohken, H. Rebel, Nucl. Phys. A191 (1972) 166.
(Half-life, Maximum β^- -decay energy)
- 1977Dr08 J.E. Draper, R.J. McDonald and N.S.P. King, Phys. Rev. C16 (1977) 1594. (Transition energies, K/L conversion electrons subshell ratio)
- 1977La19 F.P. Larkins, Atomic Data and Nuclear Data Tables. 20 (1977) 313.
(Electron shells binding energies)
- 1990Tr01 W.H. Trzaska, J. Kantele, R. Julin, J. Kumpulainen, P. Van Duppen, M. Huyse and J. Wauters, Z. Phys. A335 (1990) 475
(K/L conversion electrons subshell ratio)
- 1992Ra08 M.U. Rajput and T.D. MacMahon, Nucl. Instrum. Methods Phys. Res. A312 (1992) 289.
(Evaluation techniques)
- 1994Ka08 S.I. Kafala, T.D. MacMahon and P.W. Gray, Nucl. Instrum. Methods Phys. Res. A339 (1994) 151.
(Evaluation techniques)
- 1996Ra16 S. Raman, J.B. McGrory, E.T. Journey and J.W. Starner, Phys. Rev. C53 (1996) 2732.
(Gamma-ray transition energies)
- 1996Sc06 E. Schönfeld and H. Janßen, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527.
(Fluorescence yields)
- 1998ScZM E. Schönfeld and G. Rodloff, PTB-6.11-98-1, Braunschweig, October 1998
(K Auger electron energies)
- 1999Br39 E. Browne, Nucl. Data Sheets 88 (1999) 29.
(²⁰⁶Tl and ²⁰⁶Pb level schemes)
- 1999ScZX E. Schönfeld and G. Rodloff, PTB-6.11-1999-1, Braunschweig, February 1999
(KX-ray energies and relative emission probabilities)

- 2000ScXX E. Schönfeld and H. Janßen, Appl. Rad. Isot. 52 (2000) 595
(Program Emission)
- 2002Ba85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, P.O. Tikkanen and S. Raman, At. Data
Nucl. Data Tables. 91 (2002) 1.
(ICC)
- 2002BeXX M.M. Bé, R. Helmer and V. Chisté, J. Nucl. Scien. and Techn., suppl. 2 (2002) 481.
(Saisinuc & supporting software)
- 2003Au03 G. Audi and A.H. Wapstra, Nucl. Phys. A729 (2003) 337.
(Decay Q values)
- 2003DeXX R.D. Deslattes, E.G. Kessler, P. Indelicato, L. De Billy, E. Lindroth and J. Anton, Rev.
Mod. Phys. 77 (2003) 35.
(K and L X-ray energies)
- 2004MaXX D. MacMahon, A. Pearce and P. Harris, Appl. Rad. Isot. 60 (2004) 275
(Evaluation techniques)
- 2005KiZW T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya and C.W. Nestor, Jr., Proc. Intern. Conf.
Nuclear Data for Science and Technology, Santa Fe, New Mexico, 26 September-1
October, 2004, R.C. Haight, M.B. Chadwick, T. Kawano, P. Talou, Eds., AIP Conf. Proc.
769 (2005) 268.
(ICC)

²⁰⁷Tl - Comments on evaluation of decay data by F. G. Kondev

This evaluation was completed in September 2010, with a literature cut off by the same date, as a part of ANL commitment to the IAEA-CRP on “Updated Decay Data Library for Actinides”.

1 Decay Scheme

The nuclide ²⁰⁷Tl ($J^\pi = 1/2^+$) disintegrates 100 % by β^- emission. The strongest β^- -decay branch of 99.729 (10) % populates the $J^\pi = 1/2^-$ ground state of the daughter nuclide ²⁰⁷Pb. The level schemes of ²⁰⁷Tl and ²⁰⁷Pb, including level energies and J^π values, are based on the ENSDF evaluation of Kondev and Lalkovski (2011Ko04).

2 Nuclear Data

Adopted $Q(\beta^-)$ value of 1418 (5) keV is taken from the evaluation of Audi *et al.* (2003Au03).

The experimental half-life data for the ²⁰⁷Tl ground state are listed in Table 1. The LRSW value of $T_{1/2} = 4.774$ (12) min was adopted ($\chi^2_{\nu} = 0.38$, which is smaller than the critical value of $\chi^2_{\nu \text{ crit}} = 3.78$ (99 % confidence level)).

Table 1. Experimental data for the half-life of ²⁰⁷Tl.

Author	$T_{1/2}$ (min)	Used in evaluation
1931Cu01	4.76 (2)	Yes
1940Fa04	4.77 (5)	Yes
1953Sa11	4.79 (2)	Yes
1967Tr01	4.77 (3)	Yes

2.1 β^- Transitions

The values for the maximum β^- -decay energies, $E_{\beta^-, \text{max}}$, presented in Table 2, were deduced from $Q(\beta^-) = 1418$ (5) keV (2003Au03) and the adopted level energies of ENSDF (2011Ko04). The β^- -decay transition probabilities (P_β) were deduced from the decay scheme and the corresponding absolute γ transition probabilities. Comparisons with other measured values are given in Table 3. The $\log ft$ values were calculated using the LOGFT program from the ENSDF evaluation package, based on the work of Gove and Martin (1971Go40).

Table 2. Level energies, quantum numbers, $E_{\beta_{0,i} \text{ max}}$, P_β and $\log ft$ values in decay of ²⁰⁷Tl.

	Level energy (keV)	J^π	$E_{\beta^-, \text{max}}$ (keV)	P_β (%)	Nature	$\log ft$
$\beta_{0,2}$	897.698 (17)	3/2-	520 (5)	0.271 (10)	First forbidden non-unique	6.15
$\beta_{0,1}$	569.6982 (20)	5/2-	848 (5)	$< 8 \times 10^{-5}$	First forbidden unique	$> 10.8^{1U}$
$\beta_{0,0}$	0.0	1/2-	1418 (5)	99.729 (10)	First forbidden non-unique	5.11

Table 3. Beta-decay transition probabilities (P_β) in decay of ^{207}Tl .

	Present work (%)	1967Da10 (%)	1963Ch09 (%)	1988Hi14 (%)
$\beta_{0,2}$	0.271 (10)	0.24	0.155 (20)	
$\beta_{0,1}$	$< 8 \times 10^{-5}$	$< 1 \times 10^{-2}$		$< 8 \times 10^{-5}$
$\beta_{0,0}$	99.729 (10)	99.76	99.845 (20)	

2.2 γ Transitions

The γ -ray energies, multiplicities, absolute transition probabilities and electron internal conversion coefficients are listed in Table 4. γ transition multiplicities are taken from the ENSDF evaluation of Kondev and Lalkovski (2011Ko04), while the electron conversion coefficients were calculated using the BrIcc code (2008Ki07).

The $P_\gamma(897.77 \gamma)$ value of 0.263 (9) % was deduced from the intensity ratio of $I_\gamma(898\gamma)/I_\gamma(351\gamma) = 0.0202$ (7) (1988Hi14) and $P_\gamma(351\gamma)$ in ^{211}Bi α decay) = 13.02 (12) %. A $P_\gamma(328.10\gamma)$ value of 0.00142 (14) % was deduced from the intensity ratio of $I_\gamma(328.10\gamma)/I_\gamma(898\gamma) = 0.0054$ (5) (1988Hi14) and $P_\gamma(897.77\gamma) = 0.263$ (9) %, as described above. The absolute emission probability for the 569.698 γ of 0.00185 (19) % was deduced from the intensity balance at the 569-keV level and by neglecting the small β^- decay feeding contribution of $< 8 \times 10^{-5}$ reported in 1988Hi14. The mixing ratio of +0.091 (9) for the 569.698-keV transition was deduced as a weighted average of +0.096 (11) (1970K103), +0.075 (25) (1972Ha59), +0.075 (25) (1976Av01), and +0.11 (6) (1973Ba38).

Table 4. Energies, multiplicities, mixing ratios, absolute emission probabilities and electron internal conversion coefficients for γ transitions following the β^- -decay of ^{207}Tl .

	Energy (keV)	Multiplicity	δ	P_γ (%)	$\alpha_K \times 10^{-2}$	$\alpha_L \times 10^{-3}$	$\alpha_M \times 10^{-4}$	$\alpha_{N+} \times 10^{-4}$	α_T
$\gamma_{2,0}$	897.77 (12)	M1+E2	+0.091 (9)	0.263 (9)	1.92 (3)	3.18 (5)	7.41 (11)	2.30 (4)	0.0233 (4)
$\gamma_{2,1}$	328.10 (12)	[M1]		0.00142 (14)	27.3 (4)	46.6 (7)	109.0 (16)	33.8 (5)	0.334 (5)
$\gamma_{1,0}$	569.698 (2)	E2		0.00185 (19)	1.584 (23)	4.39 (7)	10.81 (16)	3.30 (5)	0.0216 (3)

3 Atomic Data

The atomic data (fluorescence yields, X-ray energies and relative probabilities, and Auger electron energies and relative probabilities) were provided by the SAISINUC software (2008DuZX). Details regarding the origin of these data can be found in 1996Sc06, 1998ScZM, 1999ScZX, 2000Sc47 and 2003De44.

4 Emissions

4.1 K x-rays

The X-ray data have been calculated using the evaluated gamma-ray data, and the atomic data from 1996Sc06, 1998ScZM and 1999ScZX. Both the X-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program. This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

5 Electron emissions

Electron energies were determined from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

6 References

- 1931Cu01 M. Curie, A. Debierne, A. S. Eve, H. Geiger, O. Hahn, S. C. Lind, S. Meyer, E. Rutherford, E. Schweidler. *Rev. Mod. Phys.* 3(1931)427 (Half-life)
- 1940Fa04 K. Fajans, A. F. Voigt. *Phys. Rev.* 58(1940)177 (Half-life)
- 1950Ev03 H. D. Evans. *Proc. Phys. Soc. (London)* 63A(1950)575 (Measured energies and probabilities of beta-transitions)
- 1953Sa11 B. W. Sargent, L. Yaffe, A. P. Gray. *Can. J. Phys.* 31(1953)235 (Half-life)
- 1961Cu05 S. Cuperman. *Nucl. Phys.* 28(1961)84 (Measured energies and probabilities of beta-transitions)
- 1963Ch09 P. R. Christensen, O. B. Nielsen, H. Nordby. *Phys. Letters* 4(1963)318 (Measured energies and probabilities of beta-transitions)
- 1967Da10 W. F. Davidson, C. R. Cothorn, R. D. Connor. *Can. J. Phys.* 45(1967)2295 (Measured energies and probabilities of beta-transitions)
- 1967Tr01 J. M. Trischuk, E. Kankeleit. *Nucl. Phys.* A90(1967)33 (Half-life, measured energies and probabilities of beta-transitions)
- 1968Br17 C. Briancon, C.F. Leang, R. Walen. *Compt. Rend.* 266B(1968)1533 (Measured energies and probabilities of gamma-transitions)
- 1970Kl03 H.V. Klapdor, P. von Brentano, E. Grosse, K. Haberkant. *Nucl. Phys.* A152 (1970) 263 (Measured energies and probabilities of gamma- and beta-transitions and ICCs)
- 1971Go40 N. B. Gove, M. J. Martin. *Nucl. Data Tables* A10(1971)205 (Log ft values)
- 1972Ha59 O. Hausser, F. C. Khanna, D. Ward. *Nucl. Phys.* A194(1972)113 (Multipolarity and mixing ratio)
- 1973Ba38 C. Bargholtz, L. Eriksson, L. Gidefeldt. *Phys. Scr.* 7(1973)254 (Multipolarity and mixing ratio)
- 1974Ha34 O. Hausser, D. B. Fossan, A. Olin, D. Ward, W. Witthuhn, R. E. Warner. *Nucl. Phys.* A225(1974) 425 (Measured energies and probabilities of gamma-transitions)
- 1976Av01 F. T. Avignone III, T. A. Girard. *Phys. Rev. C* 13(1976)2067 (Multipolarity and mixing ratio)
- 1977La19 F. P. Larkins. *At. Data Nucl. Data Tables* 20(1977)311 (Auger electron energies)
- 1988Hi14 M. M. Hindi, E. G. Adelberger, S. E. Kellogg, T. Murakami. *Phys. Rev. C* 38(1988)1370 (Measured energies and probabilities of gamma- and beta-transitions, B(M1), experimental ICCs)
- 1991Ar04 A. Artna-Cohen. *Nucl. Data Sheets* 63(1991)79 (Evaluation and gamma-ray normalisation factor)
- 1996Sc06 E. Schnfeld, H. Janssen. *Nucl. Instrum. Methods Phys. Res.* A369(1996)527 (Evaluation of K-shell fluorescence yields and X-ray emission probabilities)
- 1998ScZM E. Schnfeld, G. Rodloff. Report PTB-6 .11-98-1, Braunschweig (1998) (Auger electron energies)
- 1999ScZX E. Schnfeld, G. Rodloff. Report PTB-611-1999-1, Braunschweig (1999) (X-ray energies and emission probabilities)
- 2000Sc47 E. Schnfeld, H. Jansen. *Appl. Radiat. Isot.* 52(2000)595 (X-ray and Auger electron emission probabilities)
- 2000He14 R. G. Helmer, C. van der Leun. *Nucl. Instrum. Methods Phys. Res.* A450 (2000) 35 (Evaluated energies and probabilities of gamma-transitions)
- 2003De44 R. D. Deslattes, E. G. Kessler Jr., P. Indelicato, L. de Billy, E. Lindroth, J. Anton. *Rev. Mod. Phys.* 75(2003)35 (Evaluated X-ray transition energies)
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault. *Nucl. Phys.* A729(2003)337 (Q value)
- 2008Ki07 T. Kibedi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor Jr.. *Nucl. Instrum. Methods Phys. Res.* A589(2008)202 (Theoretical ICC)
- 2008DuZX C. Dulieu, M.M. B, V. Chist. *Proc. Int. Conf. on Nuclear Data for Science and Technology, Nice, France, 22-27 April 2007*(2008)97 (SAISINUC Software and atomic data)
- 2011Ko04 F. G. Kondev, S. Lalkovski. *Nucl. Data Sheets* 112(2011)707 (Nuclear levels)

**²⁰⁷Bi - Comments on evaluation of decay data
by M.M. Bé and V. Chisté**

This evaluation was completed in November 1997 and reviewed in December 2009.

1) Decay scheme

The J^π values are from NDS 70,2 (1993).

The level energies are deduced from the γ -ray energies.

2) Nuclear Data

- The Q value is from Audi *et al.* 2003
- The measured half-life values are, in years:

28 (3)	J. Sosniak et al., Can. J. Phys 37,1 (1959) 1
30,2 (5)	G. Harbottle, J. Inorg. Nucl. Chem. 12 (1959) 6
38 (3)	E.H. Appelman, Phys. Rev. 121,1 (1961) --- uncertainty divided
38 (4)	T. Rupnik, Phys. Rev. C6,4 (1972) 1433
33,4 (8)	M. Yanokura et al., Nuclear Physics A299 (1978) 92 --- omitted from analysis
	Hoppes et al (1982)
34,9 (4)	D.E. Alburger et al., Phys. Rev. C 41,5 (1990) 2320 --- uncertainty divided
32,7 (8)	W.J. Lin et al., J. Radioanal. Nucl. Chem. Letters 153,1 (1991) 51
31,55 (5)	M.P. Unterwegger et al., NIM A312 (1992) 349 --- replaces [82HoZJ], Hoppes et al.
31,549 (41)	M.P. Unterwegger, Applied Rad. Isot. 56 (2002) 125 – replaces the previous one

1. The value from M. Yanokura et al. has been omitted because it is dependent on:

- the EC/ α branching ratio of At-211,
- the probability for the 6868 keV α -transition from Po-211 to the 569,7 keV level in Pb-207,
- the half-life of At-211,
- the decay probability of Bi-207 feeding the 569,7 keV level in Pb-207.

All these data were updated since 1978, it should be necessary to re-calculate the value taking into account all these parameters. So, this value is not included in the data set.

2. The uncertainty on the Appelman's value is given for 3σ , it has been divided by 3 to give 38 (1).

3. The uncertainty on the Alburger's value is given for 2σ , it has been divided by 2 to give 34,9 (2).

Conclusion: The adopted value of 32,9 (14) a is from the LRSW analysis of the seven accepted values. The uncertainty on the Unterwegger (2002) value has been increased to (0,177) in order to reduce its relative weight to 50 %. Then $\sigma_{\text{int}} = 0,13$; $\sigma_{\text{ext}} = 0,75$ and, reduced- $\chi^2 = 36,2$. The final value is the weighted average and the uncertainty is expanded to include the most precise value. New measurements would be desirable.

2.1) Electron-Capture Transitions

- The EC transition energies are deduced from $Q(\text{EC}) = 2397,2 (21) \text{ keV}$ and from the individual level energies.
- The transition probabilities are deduced from the total gamma-ray transition probability balance at each level.
- The electron-capture sub shell ratios were calculated by using the LOGFT program.

LOGFT calculated values

level	P_K	P_L	P_{M+}	$(L+M+...)/K$
570	0,797(8)	0,150 (3)	0,049 (1)	0,25
1633	0,733(7)	0,199 (4)	0,069 (1)	0,365
2340		0,651 (6)	0,349 (6)	

Experimental values from **Mandal et al.** for the transitions to the 570-keV and 1633-keV levels:

level	P_K	$(L+M+...)/K$
570	0,59 (6)	0,68 (16)
1633	0,73 (6)	0,37 (12)

The P_K value for the transition to the 570 keV level is the weighted average of two values (0,62 (8) and 0,59 (6)) obtained by two different coincidence method measurements. These values are dependant on: ω_K , $K\alpha/(K\alpha + K\beta)$, α_K , α_T and on the EC branching ratios. By using, for these parameters, the values evaluated in this work the re-calculated value for the first P_K is = 0,73 instead of 0,62, in agreement, within the uncertainty limits, with the theoretical values from the LOGFT program.

Experimental values from **A De Beer et al.** and **M. Tan et al.** for the transition to the 2340-keV level:

A. De Beer, $P_L = 0,663 (14)$, this measurement does not depend on any other data.

M. Tan, $P_L = 0,57 (3)$, this measurement depends on α_K and α_T for the 570 keV γ -transition. In this case $\alpha_K = 0,0159$ and $\alpha_T = 0,0218$.

2.2) β^+ transitions

A weak β^+ transition to the 570-keV level was reported by **Rupnik** (1972) to be $(1,2 (2)) 10^{-2} \%$.

2.3) Gamma transitions

- Internal Conversion coefficients

The adopted values are from the LRSW analysis of all the values published after 1963. An earlier value from **R.A. Ricci** (1957) was not used due to its large uncertainty. The values from **E. Baldinger et al.** (1967) have been replaced by those of **E. Baldinger et al.** (1969). Two set of values were published by **Sen and Rizvi**, [1967Ri00, 1967Se15], one in a B.A.P.S. abstract (June), the other one in N.I.M. (July); only the last one was used because it gave a detailed description.

Internal Conversion coefficients measured values (All values are multiplied by 10²) :

- 570 keV gamma transition

	$\alpha_K - 570 \text{ keV}$	u_c	
1967KL02	1,56	0,07	1969Ba53 and 1974Mu16 are rejected due to the Chauvenet criterion Internal uncertainty = 0,009; external uncertainty = 0,24 Reduced- $\chi^2 = 0,15$ No value has a relative weight greater than 50 %. LRSW has used the weighted average and the external uncertainty. The evaluated value is = 1,574 (24)
1967VA25	1,59	0,06	
1967SE15	1,60	0,10	
1968AN04	1,56	0,05	
1969HE19	1,55	0,05	
1969AnZU	1,60	0,05	
1969BA53	1,50	0,15	
1974MU16	2,30	0,03	

	$\alpha_L - 570 \text{ keV}$	u_c	
1967SE15	0,49	0,03	Reduced- $\chi^2 = 1,1$ Internal uncertainty = 0,0060 External uncertainty = 0,0064 1988Fu05 amounts for 74 % LRSW has used the weighted average and the external uncertainty The evaluated value is = 0,452 (6)
1968AN04	0,452	0,047	
1969HE19	0,444	0,021	
1969BA53	0,50	0,10	
1974AV03	0,483	0,018	
1988FU05	0,446	0,007	

	$\alpha_M - 570 \text{ keV}$	u_c	
1967SE15	0,10	0,05	Reduced- $\chi^2 = 3$ Internal uncertainty = 0,003; external uncertainty = 0,005 weighted average and external uncertainty = 0,114 (5)
1974AV03	0,138	0,010	
1988FU05	0,112	0,003	

	$\alpha_{NOP} - 570 \text{ keV}$	u_c	
1974AV03	0,0288	0,0032	
1988FU05	0,0341	0,0017	

	$\alpha_{M+} - 570 \text{ keV}$	u_c	
1968AN04	0,172	0,047	Reduced- $\chi^2 = 1,5$ Internal uncertainty = 0,003; external uncertainty = 0,004 evaluated: weighted average = 0,1485 (39)
1969BA53	0,168	0,035	
1974AV03	0,167	0,010	
1988FU05	0,1461	0,0034	

- 897 keV gamma transition

	$\alpha_K - 897 \text{ keV}$	u_c	
1970AhZX	1,90	0,30	No value has a relative weight greater than 50 %.
1974AV03	1,81	0,25	Internal uncertainty = 0,13; external uncertainty = 0,07 reduced- $\chi^2 = 0,24$
1975JA04	1,60	0,30	LRSW has used the weighted average.
1988FU05	1,90	0,23	The evaluated value is = 1,82 (13)

- 1064 keV gamma transition

	$\alpha_K - 1064 \text{ keV}$	u_c	
1967SE15	8,5	0,5	1967Se15, 1967Kl02 and 1988Fu05 are rejected due to Chauvenet criterion Reduced- $\chi^2 = 0,03$ Internal uncertainty = 0,23; external uncertainty = 0,04 Adopted: weighted average = 9,53 (23)
1967KL02	9,0	0,9	
1969ANZU	9,4	0,9	
1969HE19	9,6	0,3	
1969AN00	9,4	0,9	
1969BA53	9,5	1,3	
1974AV03	9,43	0,47	
1974MU16	9,5	1,1	
1988FU05	9,86	0,35	

	$\alpha_L - 1064 \text{ keV}$	u_c	
1967SE15	2,33	0,15	Reduced- $\chi^2 = 1,3$ Internal uncertainty = 0,06; external uncertainty = 0,07 Adopted: weighted average = 2,47 (7)
1969BA53	2,97	0,46	
1974AV03	2,23	0,16	
1988FU05	2,51	0,10	

	$\alpha_M - 1064 \text{ keV}$	u_c	
1967SE15	0,44	0,09	Reduced- $\chi^2 = 2,2$; Critical- $\chi^2 = 4,6$ Internal uncertainty = 0,022; external uncertainty = 0,033 Adopted: weighted average = 0,591 (33)
1974AV03	0,55	0,05	
1988FU05	0,615	0,026	
	$\alpha_{M+} - 1064 \text{ keV}$	u_c	
1969BA53	1,05	0,17	

	$\alpha_{NOP} - 1064 \text{ keV}$		
1974AV03	0,17	0,03	Internal uncertainty = 0,012; external uncertainty = 0,010 Adopted: weighted average = 0,194 (12)
1988FU05	0,198	0,013	

- 1442 keV gamma transition

	$\alpha_K - 1442 \text{ keV}$	u_c	
1974AV03	0,27	0,04	
	$\alpha_L - 1442 \text{ keV}$		
1974AV03	0,042	0,008	

- 1770 keV gamma transition

	$\alpha_K - 1770 \text{ keV}$	u_c	
1971Al03	0,34	0,03	Reduced- $\chi^2 = 0,65$ Internal uncertainty = 0,018; external uncertainty = 0,014 Uncertainty increased to 0,025 to reduce weight to 50 % evaluated: weighted average = 0,346 (18)
1974AV03	0,30	0,05	
1988FU05	0,362	0,019	

	$\alpha_L - 1770 \text{ keV}$	u_c	
1974AV03	0,041	0,009	Mean = 0,0049 (8), WM = 0,053 Internal uncertainty = 0,04; external uncertainty = 0,07 evaluated: simple mean = 0,049 (8)
1988FU05	0,0569	0,0048	

	$\alpha_{M+} - 1770 \text{ keV}$		
1974AV03	0,0095	0,0024	Mean = 0,0126 (31), WM = 0,0136 Internal uncertainty = 0,0029; external uncertainty = 0,0017 evaluated: simple mean = 0,0126 (31)
1988FU05	0,0157	0,0017	

Comparison of experimental results and theoretical values : ($\alpha \times 10^2$)

Theoretical ICC values were derived from the Band *et al.* tables with the program BrIcc for the “frozen orbital” approximation (Kibédi *et al.*).

Multipolarities and mixing ratios were deduced from comparison between measured and theoretical ICC values and by comparison with δ values obtained by angular correlation measurements.

	α_K	α_L	α_M	α_{M+}	α_T	δ	Multipolarity
570 keV							
Exper.	1,574 (24)	0,452 (6)	0,114 (5)	0,1485 (39)	2,174 (9)		
BrIccFO	1,583 (23)	0,439 (7)	0,1081 (16)		2,16 (3)		E2
Adopted	1,583 (23)	0,439 (7)	0,1081 (16)		2,16 (3)		E2
897 keV	α_K						
Exper.	1,82 (13)						
BrIccFO	1,82 (8)					0,3 (3)	M1+8,3%E2
Adopted	1,82 (8)	0,304 (12)	0,071 (13)		2,22 (9)		
1064 keV	α_K	α_L	α_M	α_{NOP}	α_T		
Exper.	9,53 (23)	2,47 (7)	0,591 (3)	0,194 (12)	12,78 (24)		
BrIccFO	9,43 (14)	2,38 (4)	0,589 (9)	0,1833 (25)	12,57 (18)	0,01 (1)	M4+0,01%E5
Adopted	9,53 (23)	2,47 (7)	0,591 (33)	0,194 (12)	12,78 (24)		
1442 keV	α_K	α_L					
Exper.	0,27 (4)	0,042 (8)					
BrIccFO	0,271 (4)	0,0468 (7)					E2
Adopted	0,271 (4)	0,0468 (7)					
1770 keV	α_K	α_L	α_M	α_{M+}	α_T		
Exper.	0,346 (18)	0,049 (8)		0,0126 (31)	0,408 (20)		
BrIccFO	0,342 (5)	0,0555 (8)			0,442 (7)	0,05 (5)	M1+0,0025%E2
Adopted	0,342 (5)	0,0556 (8)			0,442 (7)	0,05 (5)	
Measured internal-pair formation coefficient, $\alpha_\pi = 0,025 (5) 10^{-2}$ (Allan 1971)							

- Gamma transition probabilities

The transition probabilities were calculated from the adopted values of the ICC and the absolute emission intensities.

4.1) X-ray emissions

- ω_K is from **Bambynek**, ω_L η_{KL} η_{LM} from **Schönfeld et al.**, ω_M from **Hubbell et al.**
A value of $\omega_K = 0,972$ (8) was measured by **Hansen et al.** (1972) and is in good agreement.
- X-ray energy: the wavelengths are from **Bearden** and converted into energy with $1 \text{ \AA} = 1,000\ 014\ 81$ (92) 10^{-10} m.
- The emission intensities are calculated with the EMISSION program from PTB.
The ratios used are in good agreement with the measured values from **Dasmahapatra et al.**

	EMISSION	Measured
$K\alpha_2/K\alpha_1 =$	0,5950 (25)	0,5984 (42)
$K\beta/K\alpha =$	0,279 (4)	0,283 (9)
$K\beta_2'/K\beta_1' =$	0,302 (5)	0,302 (30)

- Some others measurements were made by **Campbell et al.:**
 $K\beta_1/K\alpha_1 = 0,2215$ (30)
 $K\beta_2/K\alpha_1 = 0,083$ (1)

4.2) Gamma emissions

The γ -ray energies are from **Helmer et al.** for those of 569, 1063 and 1770 keV. Those at 897 and 1442 keV are from **Jardine** and 368 keV is from level energies.

All the experimental emission intensities were done relatively to that of the 570 keV gamma-ray, except **Lin et al.** where the absolute intensity is assumed to be 97,75.

The adopted values are from the LRSW analysis of all the known values, except Aubin et al. because no uncertainties were given.

897 keV	I_{rel}	u_c	
1969Ra13	0,150	0,015	Reduced- $\chi^2 = 1,23$; critical- $\chi^2 = 3,3$ Internal uncertainty = 0,0043; external uncertainty = 0,0048 LRSW has used the weighted average and the external uncertainty. The adopted value is = 0,1313 (48)
1975Ja04	0,14	0,02	
1980Yo05	0,122	0,013	
1989Sc**	0,1274	0,0052	
1991Li10	0,153	0,015	

1064 keV	I_{rel}	u_c	
1967Do09	78,4	2,40	Reduced- $\chi^2 = 2$; critical- $\chi^2 = 2,3$ Internal uncertainty = 0,15; external uncertainty = 0,22 LRSW has used the weighted average. The adopted value is = 76,29 (22)
1969Ra13	78,7	4,00	
1972Ro03	75,6	0,50	
1968He00	74,0	2,00	
1975JA04	75,5	2,3	
1973Wi10	77,7	0,45	
1980Yo05	75,79	0,25	
1989De**	76,5	0,50	
1989Sc**	76,584	0,367	
1990He16	76,4	0,50	
1991Li10	77,7	1,4	

1442 keV	I_{rel}	u_c	
1969Ra13	0,150	0,015	Internal uncertainty = 0,0025; external uncertainty = 0,0018 Reduced- $\chi^2 = 0,65$; critical- $\chi^2 = 3$ LRSW has used the weighted average and the internal uncertainty. The adopted value is = 0,1345 (23)
1975JA04	0,15	0,02	
1980Yo05	0,132	0,005	
1979Si17	0,144	0,024	
1989Sc**	0,1337	0,0027	
1991Li10	0,147	0,012	

1770 keV	I_{rel}	u_c	
1967Do09	7,07	0,35	
1969Ra13	7,5	0,4	<--- This value is rejected due to the Chauvenet criterion
1975JA04	6,95	0,20	Reduced- $\chi^2 = 0,14$; critical - $\chi^2 = 3,3$ Internal uncertainty = 0,026; external uncertainty = 0,01 The adopted value is = 7,028 (26)
1980Yo05	7,026	0,029	
1989Sc**	7,023	0,068	
1991Li10	7,11	0,13	

Gamma - 328 keV

A weak gamma emission was reported by **Schima**, with a relative intensity of 0,0045 (36).

Gamma - 1460 keV

A transition with $E\gamma = 1460$ keV was reported by **Singh et al.**, nevertheless in spite of its relatively great intensity (= 1,65 (6)), it has never been confirmed by other authors.

Absolute emission intensities:

Considering the decay scheme, the absolute emission intensity of the 570 keV gamma ray is calculated by:

$$\Sigma P(\gamma + ce)(570 + 897) = 100$$

The α_T coefficients are those determined above.

$E\gamma$	Absolute γ -ray intensity
328	0,0044 (35)
570	97,76 (3)
897	0,1284 (47)
1064	74,58 (22)
1442	0,1315 (22)
1770	6,871 (26)

5) Electron emissions

- The intensities of Auger electrons emitted were deduced from the decay scheme data by using the EMISSION program.
- The intensities of conversion electrons were calculated from the conversion coefficients and the gamma emission intensities.

6) Main production modes

From CEA/LMRI

7) References

- R.A.Ricci. *Physica* 23 (1957) 693. ICC, not used
- G.Harbottle. *J. Inorg. Nucl. Chem.* 12 (1959) 6. Half-life
- J.Sosniak, R.E.Bell. *Can. J. Phys.* 37,1 (1959) 1. Half-life
- E.H.Appelman. *Phys. Rev.* 121,1 (1961). Half-life
- A De Beer, H.P.Blok, J.Blok. *Physica* 30 (1964) 1938. Electron Capture Coefficients
- P.Kleinheinz, R.Vukanovic, L.Samuelsson, D.Krmpotic, H.Lindström, K.Siegbahn. *Nucl. Phys.* A93 (1967) 63. ICC
- D.P.Donnelly, H.W.Baer, J.J.Reidy, M.L.Wiedenbeck. *Nucl. Instrum. Methods* 57 (1967) 219. Gamma emission probabilities
- S.K.Sen, S.I.H.Rizvi. *Nucl. Instrum. Methods* 57 (1967) 227. ICC
- B.Van Nooijen, H.Van Krugten. *Phys. Lett.* 25 B,8 (1967) 510. ICC
- E.Baldinger, E.Haller. *Helv. Phys. Acta* 40 (1967) 800. ICC
- J.A.Bearden. *Rev. Mod. Phys.* 39,1 (1967) 78. X-Rays energies
- S.I.H.Rizvi, S.K.Sen. *B.A.P.S.* 12 (1967) 715. ICC, not used
- V.Andersen, C.J.Christensen. *Nucl. Phys.* A113 (1968) 81. ICC
- V.Andersen. *Riso Report* 195 (1969). ICC
- G.Hedin, A.Bäcklin. *Ark. Fysik* 38 (1969) 593. ICC, Gamma emission probabilities
- E.Baldinger, E.Haller. *Helv. Phys. Acta* 42 (1969) 949. ICC
- P.Venugopala Rao, R.E.Wood, J.M.Palms, R.W.Fink. *Phys. Rev* 178,4 (1969) 1997. Gamma emission probabilities
- G.Aubin, J.Barrette, M.Barrette, S.Monaro. *Nucl. Instrum. Methods* 76 (1969) 93. Gamma emission probabilities, not used
- B.Ahlesten, A.Backlin. *Report NP-18288(LF-26)* (1970). K ICC (897 keV)
- C.J.Allan. *Can. J. Phys.* 49,2 (1971) 157. ICC
- J.S.Hansen, J.C.McGeorge, R.W.Fink, R.E.Wood, P.Venugopala Rao, J.M.Palms. *Z. Phys.* 249 (1972) 373. K fluorescence yield, not used
- D.C.Robinson, J.M.Freeman. *Nucl. Phys.* A181 (1972) 645. Gamma emission probabilities
- T.Rupnik. *Phys. Rev.* C6,4 (1972) 1433. Half-life, Beta plus emission probability
- D.W.Nix, J.C.McGeorge, R.W.Fink. *Phys. Lett.* 46A,3 (1973) 205. X-Ray emissions, not used
- J.B.Willett, G.T.Emery. *Ann. Phys.* 78 (1973) 496. Gamma emission probabilities
- F.T.Avignone. *Nucl. Instrum. Methods* 116 (1974) 521. ICC
- P.Mukherjee, B.K.Dasmahapatra. *J. Phys.* A7,16 (1974) 2008. ICC
- L.J.Jardine. *Phys. Rev. C* 11,4 (1975) 1385. Gamma emission probabilities
- M.Yanokura, H.Kudo, H.Nakahara, K.Miyano, S.Ohya, O.Nitoh. *Nucl. Phys.* A299 (1978) 92. Half-life
- G.P.Singh, R.K.Mishra, A.K.Singh, A.Kumar. *Czech. J. Phys.* B29 (1979) 870. Gamma emission probabilities
- Y.Yoshikawa, Y.Iwata, T.Kaku, T.Katoh, J.Z.Ruan, T.Kojima, Y.Kawada. *Nucl. Instrum. Methods* 174 (1980) 109. Gamma emission probabilities
- M.Tan, R.A.Braga, R.W.Fink. *Nucl. Phys.* A388 (1982) 498. Electron Capture Coefficients
- W.Bambynek. *X-84 Proc. X-Ray and Inner-Shell Processes in Atoms, Molecules and Solids*, A. Meisel Ed., Leipzig Aug. 20-23 (1984). K fluorescence yield
- A.M.Mandal, A.P.Patro. *J. Phys.* G11 (1985) 1025. Electron Capture Coefficients, not used
- Y.Fujita, M.Imamura, K.Omata, Y.Isozumi, S.Ohya. *Nucl. Phys.* A484 (1988) 77. ICC
- F.J.Schima. *IAEA-CRP GS/59* (1989). Gamma emission probabilities
- K.Debertin, U.Schötzig. *IAEA-CRP GS/55* (1989). Gamma emission probabilities
- D.E.Alburger, G.Harbottle. *Phys. Rev. C* 41,5 (1990) 2320. Half-life
- R.G.Helmer. *Int. J. Appl. Radiat. Isotop.* 41 (1990) 791. Gamma emission probabilities
- TECDOC-619. *IAEA. A-1400 Vienna* (1991). X-Ray emission probabilities, not used
- W.J.Lin, G.Harbottle. *J. Radioanal. Nucl. Chem. Letters* 153,1 (1991) 51. Half-life, Gamma emission probabilities
- M.P.Unterwegger, D.D.Hoppes, F.J.Schima. *Nucl. Instrum. Methods Phys. Res.* A312 (1992) 349. Half-life
- LOGFT program, ENSDF. BNL (1993). lg ft

Comments on evaluation

- J.H.Hubbell, Trehan P.N., Nirmal Singh, Chand B., Mehta D., Garg M.L., Garg R.R., Surinder Singh, Puri S.J. Phys. Chem. Ref. Data 23-2 (1994) 339. M fluorescence yield
- B.Dasmahapatra, A.Mukherjee. Phys. Rev. A51,5 (1995) 3546. X-Ray emission probabilities, not used
- E.Schönfeld, H.Janssen. Nucl. Instrum. Methods Phys. Res. A369 (1996) 527. L fluorescence yield
- E.Schönfeld. EMISSION program, PTB (1997). Auger and X-ray emission probabilities
- R.G.Helmer, C.Van der Leun. Nucl. Instrum. Methods Phys. Res. A450 (2000) 35. Gamma energies
- I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, P.O. Tikkanen, S. Raman. Atom. Data and Nucl. Data Tables 91 (2002) 1. Theoretical internal conversion coefficients
- M. P. Unterweger. Appl. Rad. Isotopes 56 (2002) 125. Half-life
- G.Audi, A.H.Wapstra, C.Thibault. Nucl. Phys. A729 (2003) 129. Q
- T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor Jr. Nucl. Instrum. Methods Phys. Res. A589 (2008) 202. Theoretical ICC

**²⁰⁸Tl – Comments on evaluation of decay data
by A. L. Nichols**

Evaluated: July/August 2001

Re-evaluated: January 2004 and July 2010

Evaluation Procedures

Limitation of Relative Statistical Weight Method (LWM) was applied to average numbers throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

The ground state of ²⁰⁸Tl ($J^\pi = 5^+$) decays by beta minus emission to various excited levels of ²⁰⁸Pb. A consistent decay scheme has been derived, assuming no direct beta decay to both the 2614.55-keV nuclear level and ground state of ²⁰⁸Pb (based on spin-parity considerations). This decay scheme is primarily based on the gamma-ray measurements of 1960Em01, 1960Sc07, 1961Si11, 1969Au10, 1969La23, 1969Pa02, 1972DaZA/1973Da38, 1972Ja25, 1975Ko02, 1977Ge12, 1978Av01, 1982Sa36, 1983Sc13, 1983Va22, 1984Ge07, 1992Li05 and 1993El08.

Nuclear Data

²²⁸Th decay chain is important in quantifying the environmental impact of the decay of naturally-occurring ²³²Th. Specific radionuclides in this decay chain are noteworthy because of their decay characteristics (²²⁴Ra alpha decay to ²²⁰Rn; ²¹²Bi and ²⁰⁸Tl gamma-ray emissions).

Half-life

The half-life is the weighted mean of the measurements of 1957Ba05, 1967La20, 1970Mu21 and 1971Ac02, with the uncertainty increased artificially to encompass the most precise study.

Reference	Half-life (min)
1957Ba05	3.100 (15)
1967La20	3.055 (6)
1970Mu21	3.17 (5)
1971Ac02	3.0527 (33)*
Recommended value	3.058 (6)#

* Uncertainty adjusted to ± 0.0055 to reduce weighting to no more than 0.50.

Weighted mean adopted, with uncertainty increased to include most precise value.

Gamma Rays

Energies

Both the 583.187 (2)- and 2614.511 (10)-keV gamma-ray energies were taken from 2000He14. All other gamma-ray transition energies were calculated from the structural details of the proposed decay scheme; the nuclear level energies of 2007Ma45 were adopted, and used to determine the energies and associated uncertainties of the gamma-ray transitions between the various populated-depopulated levels.

Adopted nuclear levels of ²⁰⁸Pb: Energy, J^π and origins (2007Ma45).

Nuclear level	Nuclear level energy (keV)*	J ^π	Origins
0	0.0	0 +	²⁰⁸ Bi EC decay, ²⁰⁸ Tl β ⁻ decay, ²¹² Po α decay, ²⁰⁶ Pb(t,x), ²⁰⁷ Pb(n,γ), ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(γ, x), ²⁰⁸ Pb(e,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁸ Pb(α,x), ²¹⁰ Pb(p,x), ²⁰⁹ Bi(d,x), ²⁰⁹ Bi(t,x), Coulomb excitation, etc.
1	2614.552 ± 0.010	3 -	²⁰⁸ Bi EC decay, ²⁰⁸ Tl β ⁻ decay, ²¹² Po α decay, ²⁰⁶ Pb(t,x), ²⁰⁷ Pb(n,γ), ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(e,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁸ Pb(α,x), ²¹⁰ Pb(p,x), ²⁰⁹ Bi(d,x), ²⁰⁹ Bi(t,x), Coulomb excitation, etc.
2	3197.711 ± 0.010	5 -	²⁰⁸ Tl β ⁻ decay, ²¹² Po α decay, ²⁰⁶ Pb(t,x), ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(e,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁸ Pb(α,x), ²¹⁰ Pb(p,x), ²⁰⁹ Bi(d,x), ²⁰⁹ Bi(t,x), Coulomb excitation, etc.
3	3475.078 ± 0.011	4 -	²⁰⁸ Tl β ⁻ decay, ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁹ Bi(d,x), etc.
4	3708.451 ± 0.012	5 -	²⁰⁸ Tl β ⁻ decay, ²⁰⁶ Pb(t,x), ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(e,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁸ Pb(α,x), ²⁰⁹ Bi(d,x), ²⁰⁹ Bi(t,x), etc.
5	3919.966 ± 0.013	6 -	²⁰⁸ Tl β ⁻ decay, ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), etc.
6	3946.578 ± 0.014	4 -	²⁰⁸ Tl β ⁻ decay, ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁹ Bi(d,x), etc.
7	3961.162 ± 0.013	5 -	²⁰⁸ Tl β ⁻ decay, ²⁰⁶ Pb(t,x), ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(e,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁸ Pb(α,x), ²¹⁰ Pb(p,x), ²⁰⁹ Bi(d,x), ²⁰⁹ Bi(t,x), etc.
8	3995.438 ± 0.013	4 -	²⁰⁸ Tl β ⁻ decay, ²⁰⁷ Pb(n,γ), ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁹ Bi(d,x), etc.
9	4037.443 ± 0.014	7 -	²⁰⁶ Pb(t,x), ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(e,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁸ Pb(α,x), ²¹⁰ Pb(p,x), etc.
10	4051.134 ± 0.013	3 -	²⁰⁷ Pb(n,γ), ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁸ Pb(α,x), ²⁰⁹ Bi(d,x), etc.
11	4085.52 ± 0.04	2 +	²⁰⁶ Pb(t,x), ²⁰⁷ Pb(n,γ), ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(γ, x), ²⁰⁸ Pb(e,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁸ Pb(α,x), ²¹⁰ Pb(p,x), ²⁰⁹ Bi(d,x), Coulomb excitation, etc.
12	4125.347 ± 0.012	5 -	²⁰⁸ Tl β ⁻ decay, ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(e,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁸ Pb(α,x), ²⁰⁹ Bi(d,x), ²⁰⁹ Bi(t,x), etc.
13	4180.414 ± 0.014	5 -	²⁰⁸ Tl β ⁻ decay, ²⁰⁶ Pb(t,x), ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(e,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁸ Pb(α,x), ²¹⁰ Pb(p,x), ²⁰⁹ Bi(d,x), etc.
14	4206.277 ± 0.004	6 -	²⁰⁷ Pb(d,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁹ Bi(d,x), etc.
15	4229.590 ± 0.017	2 -	²⁰⁷ Pb(n,γ), ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x)
16	4254.795 ± 0.017	3 -	²⁰⁶ Pb(t,x), ²⁰⁷ Pb(n,γ), ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(e,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁸ Pb(α,x), ²¹⁰ Pb(p,x), ²⁰⁹ Bi(d,x)
17	4261.871 ± 0.013	4 -	²⁰⁸ Tl β ⁻ decay, ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁹ Bi(d,x), etc.
18	4296.560 ± 0.013	5 -	²⁰⁸ Tl β ⁻ decay, ²⁰⁶ Pb(t,x), ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁸ Pb(α,x), ²¹⁰ Pb(p,x), ²⁰⁹ Bi(d,x), etc.
19	4323.946 ± 0.014	4 +	²⁰⁸ Tl β ⁻ decay, ²⁰⁶ Pb(t,x), ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(e,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁸ Pb(α,x), ²¹⁰ Pb(p,x), ²⁰⁹ Bi(d,x), etc.
20	4358.670 ± 0.013	4 -	²⁰⁸ Tl β ⁻ decay, ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁹ Bi(d,x), etc.
21	4383.285 ± 0.017	6 -	²⁰⁸ Tl β ⁻ decay, ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(e,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁹ Bi(d,x), etc.
22	4423.647 ± 0.015	6 +	²⁰⁷ Pb(d,x), ²⁰⁸ Pb(e,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁸ Pb(α,x), ²¹⁰ Pb(p,x), ²⁰⁹ Bi(d,x), etc.
23	4480.746 ± 0.016	6 -	²⁰⁸ Tl β ⁻ decay, ²⁰⁷ Pb(d,x), ²⁰⁸ Pb(e,x), ²⁰⁸ Pb(n,n'γ), ²⁰⁸ Pb(p,x), ²⁰⁸ Pb(d,x), ²⁰⁸ Pb(α,x), ²⁰⁹ Bi(d,x), etc.

* Nuclear levels at 4144 (5) and 4447 (5) keV not included, although proposed in studies of the ²⁰⁹Bi(d, ³He) reaction.

Placements of gamma-ray transitions (2007Ma45).

	Adopted E _γ [*] (keV)	Proposed location in decay scheme (²⁰⁸ Pb nuclear levels)		Adopted E _γ [*] (keV)	Proposed location in decay scheme (²⁰⁸ Pb nuclear levels)
γ _{5,4}	211.52 (2)	3919.966 (13) – 3708.451 (12)	–	835.90 (11)	not placed in decay scheme
γ _{4,3}	233.37 (2)	3708.451 (12) – 3475.078 (11)	γ _{3,1}	860.53 (2)	3475.078 (11) – 2614.552 (10)
γ _{7,4}	252.71 (2)	3961.162 (13) – 3708.451 (12)	γ _{20,3}	883.59 (2)	4358.670 (13) – 3475.078 (11)
γ _{3,2}	277.37 (2)	3475.078 (11) – 3197.711 (10)	γ _{12,2}	927.64 (2)	4125.347 (12) – 3197.711 (10)
γ _{7,3}	486.08 (2)	3961.162 (13) – 3475.078 (11)	γ _{13,2}	982.70 (2)	4180.414 (14) – 3197.711 (10)
γ _{4,2}	510.74 (2)	3708.451 (12) – 3197.711 (10)	γ _{4,1}	1093.90 (2)	3708.451 (12) – 2614.552 (10)
γ _{2,1}	583.187 (2)	3197.711 (10) – 2614.552 (10)	γ _{19,2}	1126.24 (2)	4323.946 (14) – 3197.711 (10)
γ _{18,4}	588.11 (2)	4296.560 (13) – 3708.451 (12)	γ _{20,2}	1160.96 (2)	4358.670 (13) – 3197.711 (10)
γ _{12,3}	650.27 (2)	4125.347 (12) – 3475.078 (11)	γ _{21,2}	1185.57 (2)	4383.285 (17) – 3197.711 (10)
γ _{13,3}	705.34 (2)	4180.414 (14) – 3475.078 (11)	γ _{23,2}	1283.04 (2)	4480.746 (16) – 3197.711 (10)
γ _{5,2}	722.26 (2)	3919.966 (13) – 3197.711 (10)	γ _{8,1}	1380.89 (2)	3995.438 (13) – 2614.552 (10)
γ _{6,2}	748.87 (2)	3946.578 (14) – 3197.711 (10)	γ _{17,1}	1647.32 (2)	4261.871 (13) – 2614.552 (10)
γ _{7,2}	763.45 (2)	3961.162 (13) – 3197.711 (10)	γ _{20,12}	1744.12 (2)	4358.670 (13) – 2614.552 (10)
–	808.32 (13)	not placed in decay scheme	γ _{1,0}	2614.511 (10)	2614.552 (10) – 0
γ _{18,3}	821.48 (2)	4296.560 (13) – 3475.078 (11)			

* Values derived from the adopted energies of the ²⁰⁸Pb nuclear levels as specified in columns 3 and 6, with the uncertainties rounded upwards on the basis of the recommended uncertainties of the nuclear level energies (2007Ma45).

Emission Probabilities

A consistent decay scheme has been constructed from the gamma-ray measurements of 1960Em01, 1960Sc07, 1961Si11, 1969Au10, 1969Pa02, 1969La23, 1972Ja25, 1972DaZA/1973Da38, 1975Ko02, 1977Ge12, 1978Av01, 1982Sa36, 1983Sc13, 1983Va22, 1984Ge07, 1992Li05 and 1993El08. The study of 1975Ko02 is particularly comprehensive, along with the gamma-ray measurements of 1993El08 below 1000 keV. Gamma-ray emission probabilities have been expressed relative to the 2614.511-keV transition, and specific sets of data were adjusted accordingly (some of the original measurements were quantified relative to the 583.187-keV gamma ray or as absolute emission probabilities, while minor modifications were made to the relevant emission probabilities for the partially resolved 277.37-, 510.74- and 583.187-keV gamma rays as reported by 1983Sc13). 1993El08 observed additional gamma rays (808.32 and 835.90 keV) that have not been successfully placed in the proposed decay scheme – all nuclear levels of ²⁰⁸Pb below an energy of 4611 keV have been assessed in terms of shell-model calculations and particle-gamma coincidence measurements by 1997Sc21, arguing against the possible existence of additional nuclear levels below this energy that might accommodate either of these two gamma transitions.

Experimental studies have been made of weak crossover gamma transitions by Vasil’ev et al. (2006Va23) to provide upper limits for the emission probabilities of three such emissions:

E _γ (keV)	P _γ (%), expressed per 100 decays of ²⁰⁸ Tl
3197.7	≤ 0.0007
3475.0	≤ 0.0003
3708.5	≤ 0.0004

Other high-energy gamma emissions were identified as summation peaks. These crossover gamma transitions have not been included in the proposed decay scheme because of their somewhat ill-defined, low emission probabilities and tentative nature.

Published gamma-ray emission probabilities.

E_γ (keV)	P_γ						
	1960Em01	1960Sc07	1961Si11		1969Au10*	1969La23	1969Pa02
211.52 (2)	-	-	-	-	-	0.20 (5)	0.17 (8)
233.37 (2)	-	0.3	-	-	-	0.30 (5)	0.33 (17)
252.71 (2)	1.5 (7)	1.1	-	-	-	0.8 (1)	0.70 (11)
277.37 (2)	6.9 (8)	8.6	-	7.2 (7)	-	6.9 (5)	6.5 (4)
486.08 (2)	-	0.1 (1)	-	-	-	0.07 (4)	0.05 (2)
510.74 (2)	23 (2)	25.3 (12)	24 (3)	22.5 (25)	-	23 (1)	22.5 (12)
583.187 (2)	86.4 (56)	85.1 (40)	81 (5)	84 (5)	100	85 (4)	86 (4)
588.11 (2)	-	-	-	-	-	-	-
650.27 (2)	-	-	-	-	-	-	-
705.34 (2)	-	-	-	-	-	-	-
722.26 (2)	-	-)	-	-	0.3 (1)	0.27 (8)
748.87 (2)	-	-) 22.5 (20)	-	-	-	-
763.45 (2)	1.9 (5)	3.4 (2))	3.6 (7)	-	2.0 (2)	1.68 (8)
808.32 (13)	-	-	-	-	-	-	-
821.48 (2)	-	-	-	-	-	-	0.09 (4)
835.90 (11)	-	-	-	-	-	-	-
860.53 (2)	11.4 (12)	14.2 (6)	15.3 (20)	15.2 (15)	-	13 (1)	12.0 (8)
883.59 (2)	-	-	-	-	-	-	-
927.64 (2)	-	-	-	-	-	0.15 (5)	0.13 (3)
982.70 (2)	-	-	-	-	-	0.20 (5)	0.20 (3)
1004 (2)	-	-	-	-	-	-	~ 0.01
1093.90 (2)	-	0.7 (1)	~ 2	-	-	0.5 (1)	0.38 (5)
1126.24 (2)	-	-	-	-	-	-	-
1160.96 (2)	-	-	-	-	-	-	-
1185.57 (2)	-	-	-	-	-	-	-
1283.04 (2)	-	-	-	-	-	-	0.05 (2)
1380.89 (2)	-	-	-	-	-	-	0.02 (1)
1647.32 (2)	-	-	~ 3	-	-	-	~ 0.01
1744.12 (2)	-	-	-	-	-	-	-
2614.511 (10)	100	(100)	100	100	116.7 (24)	100	100

Published gamma-ray emission probabilities (cont.).

E_γ (keV)	P_γ (cont.)					
	1973Da38	1972Ja25	1975Ko02	1977Ge12*	1978Av01	1982Sa36†
211.52 (2)	0.16 (4)	-	0.17 (2)	-	-	-
233.37 (2)	~ 0.2	-	0.31 (3)	-	-	-
252.71 (2)	0.8 (2)	-	0.80 (5)	-	0.62 (4)	0.28 (3)
277.37 (2)	6.6 (13)	6.2 (7)	6.8 (3)	-	6.1 (2)	2.4 (1)
486.08 (2)	0.04 (1)	-	0.050 (5)	-	-	-
510.74 (2)	22.9 (23)	21.9 (7)	21.6 (9)	-	22.8 (7)	7.8 (4)
583.187 (2)	85.0 (85)	86.0 (4)	86 (3)	100	85	30.0 (14)
588.11 (2)	~ 0.04	-	0.04 (2)	-	-	-
650.27 (2)	-	-	0.036 (5)	-	-	-
705.34 (2)	~ 0.02	-	0.022 (4)	-	-	-
722.26 (2)	0.21 (6)	-	0.203 (14)	-	0.27 (2)	-
748.87 (2)	0.05 (1)	-	0.043 (4)	-	-	-
763.45 (2)	1.7 (3)	-	1.64 (9)	-	1.82 (9)	0.7 (1)
808.32 (13)	-	-	-	-	-	-
821.48 (2)	0.04 (1)	-	0.040 (4)	-	-	-
835.90 (11)	-	-	-	-	-	-
860.53 (2)	11.8 (12)	11.5 (10)	12.0 (4)	14.79 (15)	13.9 (6)	4.2 (2)
883.59 (2)	~ 0.025	-	0.031 (3)	-	-	-
927.64 (2)	0.13 (4)	-	0.125 (11)	-	-	-
982.70 (2)	0.20 (6)	-	0.197 (15)	-	-	-
1004 (2)	-	-	< 0.005	-	-	-
1093.90 (2)	0.37 (7)	-	0.37 (4)	-	-	-
1126.24 (2)	-	-	0.005 (2)	-	-	-
1160.96 (2)	-	-	0.011 (3)	-	-	-
1185.57 (2)	-	-	0.017 (5)	-	-	-
1283.04 (2)	~ 0.05	-	0.052 (5)	-	-	-
1380.89 (2)	-	-	0.007 (3)	-	-	-
1647.32 (2)	-	-	0.002 (1)	-	-	-
1744.12 (2)	-	-	0.002 (1)	-	-	-
2614.511 (10)	100	(100)	100	118.5 (16)	(100)	-

Published gamma-ray emission probabilities (cont.).

E_γ (keV)	P_γ (cont.)				
	1983Sc13 [‡]	1983Va22 [#]	1984Ge07 [*]	1992Li05	1993El08 [¶]
211.52 (2)	-	-	0.228 (20)	-	0.18 (1)
233.37 (2)	-	-	0.31 (4)	-	0.30 (1)
252.71 (2)	-	-	0.955 (13)	-	0.77 (2)
277.37 (2)	2.33 (7)	2.29 (4)	7.55 (6)	2.54 (7) [§]	6.88 (12)
486.08 (2)	-	-	-	-	0.055 (11)
510.74 (2)	7.90 (23)	8.31 (14)	26.9 (9)	-	22 (1)
583.187 (2)	30.7 (8)	30.8 (6)	100.0 (6)	29.4 (7) [§]	86 (3)
588.11 (2)	-	-	-	-	0.07 (1)
650.27 (2)	-	-	-	-	0.065 (11)
705.34 (2)	-	-	-	-	-
722.26 (2)	-	-	0.31 (6)	-	0.27 (2)
748.87 (2)	-	-	-	-	0.054 (9)
763.45 (2)	0.73 (5)	-	2.15 (2)	0.651 (40)	1.72 (8)
808.32 (13)	-	-	-	-	0.029 (7)
821.48 (2)	-	-	-	-	0.041 (17)
835.90 (11)	-	-	-	-	0.075 (11)
860.53 (2)	4.55 (12)	-	14.78 (9)	4.32 (15)	12.6 (7)
883.59 (2)	-	-	-	-	-
927.64 (2)	-	-	-	-	0.13 (1)
982.70 (2)	-	-	-	-	0.21 (1)
1004 (2)	-	-	-	-	-
1093.90 (2)	-	-	0.525 (8)	-	0.47 (4)
1126.24 (2)	-	-	-	-	-
1160.96 (2)	-	-	-	-	-
1185.57 (2)	-	-	-	-	-
1283.04 (2)	-	-	-	-	0.049 (13)
1380.89 (2)	-	-	-	-	-
1647.32 (2)	-	-	-	-	-
1744.12 (2)	-	-	-	-	-
2614.511 (10)	35.6 (11)	-	119.1 (21)	-	98.1 (13)

* Emission probabilities relative to $P_\gamma(583.187 \text{ keV})$ of 100.

† Emission probabilities relative to $P_\gamma(583.187 \text{ keV})$ of 30.0.

‡ Emission probabilities relative to $P_\gamma(583.187 \text{ keV})$ of 30.7.

Emission probabilities relative to $P_\gamma(583.187 \text{ keV})$ of 30.8.

¶ Absolute emission probabilities.

§ Unresolved overlap with another gamma-ray emission.

Equivalent measurements of specific emission probabilities deviate significantly between laboratories:

- 252.71-keV gamma ray: 1960Em01 and 1978Av01;
- 486.08-keV gamma ray: 1960Sc07;
- 510.74-keV gamma ray: 1960Sc07;
- 583.187-keV gamma ray: 1961Si11;
- 763.45-keV gamma ray: 1960Sc07 and 1961Si11;
- 860.53-keV gamma ray: 1960Sc07, 1961Si11 and 1978Av01;
- 927.64-keV gamma ray: 1969La23;
- 1093.90-keV gamma ray: 1960Sc07.

These particular values were judged to be outliers, and were not included in the weighted-mean analyses. Other gamma-ray emission probabilities were not reported with uncertainties within 1960Sc07, along with the 583.187-keV gamma-ray emission in 1978Av01; these data were also not included in the weighted-mean analyses. 1982Sa36, 1983Va22 and 1992Li05 reported measurements that did not include the main 2614.511-keV gamma-ray transition: the evaluated relative emission probability of the 583.187-keV gamma ray was adopted to create data sets comparable with the other studies, and therefore the assumed $P_\gamma(583.187 \text{ keV})$ in these particular calculations were not included in the subsequent analysis.

While an uncertainty of 0.8 % can be derived for the relative emission probability of the 2614.511-keV gamma ray from the emission probabilities and uncertainties determined experimentally by 1969Au10, 1977Ge12, 1983Sc13, 1984Ge07 and 1993El08, the precise nature of this transition in such a well-

defined area of the decay scheme permits the establishment of a recommended value of 100 % with no assigned uncertainty:

Reference	$P_{\gamma}(2614.511 \text{ keV})$
1969Au10	100 (2)
1977Ge12	100.0 (14)
1983Sc13	100 (3)
1984Ge07	100 (2)
1993El08	100.0 (13)
Weighted-mean value (LRSW)	100.0 (8) \rightarrow 100 (1)
Recommended value	100

Gamma-ray emission probabilities: Relative to $P_{\gamma}(2614.511 \text{ keV})$ of 100 %.

E_{γ} (keV)	P_{γ}^{rel}						
	1960Em01	1960Sc07	1961Si11		1969Au10	1969La23	1969Pa02
211.52 (2)	-	-	-	-	-	0.20 (5)	0.17 (8)
233.37 (2)	-	0.3 [§]	-	-	-	0.30 (5)	0.33 (17)
252.71 (2)	1.5 (7) [†]	1.1 [§]	-	-	-	0.8 (1)	0.70 (11)
277.37 (2)	6.9 (8)	8.6 [§]	-	7.2 (7)	-	6.9 (5)	6.5 (4)
486.08 (2)	-	0.1 (1) [†]	-	-	-	0.07 (4)	0.05 (2)
510.74 (2)	23 (2)	25.3 (12) [†]	24 (3)	22.5 (25)	-	23 (1)	22.5 (12)
583.187 (2)	86.4 (56)	85.1 (40)	81 (5) [†]	84 (5)	85.7 (18)	85 (4)	86 (4)
588.11 (2)	-	-	-	-	-	-	-
650.27 (2)	-	-	-	-	-	-	-
705.34 (2)	-	-	-	-	-	-	-
722.26 (2)	-	-)	-	-	0.3 (1)	0.27 (8)
748.87 (2)	-	-) 22.5 (20) [‡]	-	-	-	-
763.45 (2)	1.9 (5)	3.4 (2) [†])	3.6 (7) [†]	-	2.0 (2)	1.68 (8)
808.32 (13)	-	-	-	-	-	-	-
821.48 (2)	-	-	-	-	-	-	0.09 (4)
835.90 (11)	-	-	-	-	-	-	-
860.53 (2)	11.4 (12)	14.2 (6) [†]	15.3 (20) [†]	15.2 (15) [†]	-	13 (1)	12.0 (8)
883.59 (2)	-	-	-	-	-	-	-
927.64 (2)	-	-	-	-	-	0.15 (5) [†]	0.13 (3)
982.70 (2)	-	-	-	-	-	0.20 (5)	0.20 (3)
1004 (2)	-	-	-	-	-	-	~ 0.01
1093.90 (2)	-	0.7 (1) [†]	~ 2	-	-	0.5 (1)	0.38 (5)
1126.24 (2)	-	-	-	-	-	-	-
1160.96 (2)	-	-	-	-	-	-	-
1185.57 (2)	-	-	-	-	-	-	-
1283.04 (2)	-	-	-	-	-	-	0.05 (2)
1380.89 (2)	-	-	-	-	-	-	0.02 (1)
1647.32 (2)	-	-	~ 3	-	-	-	~ 0.01
1744.12 (2)	-	-	-	-	-	-	-
2614.511 (10)	100	(100)	100	100	100 (2)	100	100

Gamma-ray emission probabilities: Relative to $P_\gamma(2614.511 \text{ keV})$ of 100 % (cont.).

E_γ (keV)	P_γ^{rel} (cont.)					
	1973Da38	1972Ja25	1975Ko02	1977Ge12	1978Av01	1982Sa36
211.52 (2)	0.16 (4)	-	0.17 (2)	-	-	-
233.37 (2)	~ 0.2	-	0.31 (3)	-	-	-
252.71 (2)	0.8 (2)	-	0.80 (5)	-	0.62 (4) [†]	0.80 (9)
277.37 (2)	6.6 (13)	6.2 (7)	6.8 (3)	-	6.1 (2)	6.8 (3)
486.08 (2)	0.04 (1)	-	0.050 (5)	-	-	-
510.74 (2)	22.9 (23)	21.9 (7)	21.6 (9)	-	22.8 (7)	22.2 (11)
583.187 (2)	85.0 (85)	86.0 (4)	86 (3)	84.4 (11)	85 [§]	[85.2 (3)] [#]
588.11 (2)	~ 0.04	-	0.04 (2)	-	-	-
650.27 (2)	-	-	0.036 (5)	-	-	-
705.34 (2)	~ 0.02	-	0.022 (4)	-	-	-
722.26 (2)	0.21 (6)	-	0.203 (14)	-	0.27 (2)	-
748.87 (2)	0.05 (1)	-	0.043 (4)	-	-	-
763.45 (2)	1.7 (3)	-	1.64 (9)	-	1.82 (9)	2.0 (3)
808.32 (13)	-	-	-	-	-	-
821.48 (2)	0.04 (1)	-	0.040 (4)	-	-	-
835.90 (11)	-	-	-	-	-	-
860.53 (2)	11.8 (12)	11.5 (10)	12.0 (4)	12.48 (13)	13.9 (6) [†]	11.9 (6)
883.59 (2)	~ 0.025	-	0.031 (3)	-	-	-
927.64 (2)	0.13 (4)	-	0.125 (11)	-	-	-
982.70 (2)	0.20 (6)	-	0.197 (15)	-	-	-
1004 (2)	-	-	< 0.005	-	-	-
1093.90 (2)	0.37 (7)	-	0.37 (4)	-	-	-
1126.24 (2)	-	-	0.005 (2)	-	-	-
1160.96 (2)	-	-	0.011 (3)	-	-	-
1185.57 (2)	-	-	0.017 (5)	-	-	-
1283.04 (2)	~ 0.05	-	0.052 (5)	-	-	-
1380.89 (2)	-	-	0.007 (3)	-	-	-
1647.32 (2)	-	-	0.002 (1)	-	-	-
1744.12 (2)	-	-	0.002 (1)	-	-	-
2614.511 (10)	100	(100)	100	100.0 (14)	(100)	-

Gamma-ray emission probabilities: Relative to P_γ(2614.511 keV) of 100 % (cont.).

E _γ (keV)	P _γ ^{rel} (cont.)					
	1983Sc13	1983Va22	1984Ge07	1992Li05	1993El08	Recommended value*
211.52 (2)	-	-	0.19 (2)	-	0.18 (1)	0.18 (1)
233.37 (2)	-	-	0.26 (3)	-	0.31 (1)	0.31 (1)
252.71 (2)	-	-	0.80 (1)	-	0.78 (2)	0.78 (2)
277.37 (2)	6.5 (2)	6.3 (1)	6.34 (5)	7.36 (20) ^ψ	7.01 (12)	6.6 (3)
486.08 (2)	-	-	-	-	0.056 (11)	0.049 (4)
510.74 (2)	22.2 (6)	23.0 (4)	22.6 (8)	-	22 (1)	22.6 (2)
583.187 (2)	85.8 (22)	[85.2 (3)] [#]	84.0 (5)	[85.2 (3)] ^ψ	88 (3)	85.2 (3)
588.11 (2)	-	-	-	-	0.07 (1)	0.06 (1)
650.27 (2)	-	-	-	-	0.066 (11)	0.041 (5)
705.34 (2)	-	-	-	-	-	0.022 (4)
722.26 (2)	-	-	0.26 (5)	-	0.28 (2)	0.24 (4)
748.87 (2)	-	-	-	-	0.055 (9)	0.046 (3)
763.45 (2)	2.05 (14)	-	1.81 (2)	1.89 (12)	1.75 (8)	1.80 (2)
808.32 (13)	-	-	-	-	0.030 (7)	0.030 (7)
821.48 (2)	-	-	-	-	0.042 (17)	0.041 (4)
835.90 (11)	-	-	-	-	0.076 (11)	0.076 (11)
860.53 (2)	12.8 (3)	-	12.41 (8)	12.5 (4)	12.8 (7)	12.4 (1)
883.59 (2)	-	-	-	-	-	0.031 (3)
927.64 (2)	-	-	-	-	0.13 (1)	0.128 (7)
982.70 (2)	-	-	-	-	0.21 (1)	0.205 (8)
1004 (2)	-	-	-	-	-	-
1093.90 (2)	-	-	0.441 (7)	-	0.48 (4)	0.44 (1)
1126.24 (2)	-	-	-	-	-	0.005 (2)
1160.96 (2)	-	-	-	-	-	0.011 (3)
1185.57 (2)	-	-	-	-	-	0.017 (5)
1283.04 (2)	-	-	-	-	0.050 (13)	0.052 (5)
1380.89 (2)	-	-	-	-	-	0.007 (3)
1647.32 (2)	-	-	-	-	-	0.002 (1)
1744.12 (2)	-	-	-	-	-	0.002 (1)
2614.511 (10)	100 (3)	-	100 (2)	-	100.0 (13)	100

* Weighted mean values adopted when appropriate; remainder derived from proposed decay scheme; normalisation factor of 0.997 55 (4) calculated from total theoretical internal conversion coefficient of 2614.511-keV (0.002 46 (4)) gamma transition and transition probability of 100 % (1.00), with no direct β⁻ decay to the 2614.552-keV nuclear level and ground state of ²⁰⁸Pb.

† Rejected as outlier, and not included in weighted-mean analysis.

§ No uncertainty quoted; data not included in the weighted-mean analysis.

‡ Unresolved data not included in the weighted-mean analysis.

Measurements did not include determination of the 2614.511-keV gamma ray; therefore, relative emission probability of 85.2 (3) for the 583.187-keV gamma ray was used to convert all other data in this study to comparable relative values – under these circumstances, P_γ(583.187 keV) was not included in the weighted-mean analysis.

ψ unresolved overlap with another gamma-ray emission, and measurement did not include 2614.511-keV γ ray; therefore relative emission probability of 85.2 (3) was used for the 583.187-keV γ ray to convert other data in this study to comparable relative values – under these circumstances, P_γ(277.37 keV) and P_γ(583.187 keV) were not included in the weighted-mean analyses.

Multipolarities and Internal Conversion Coefficients

The major 583.187- and 2614.511-keV gamma rays were identified as E2 and E3 transitions, respectively. Many other gamma rays have mixed M1 + E2 multipolarities; these transitions were generally assumed to be 100 % M1, although estimated mixing ratios derived from the studies of 1954El07, 1957Kr56, 1957Vo22, 1963Da11, 1972Ja25, 1976Av03, 1978Av01 and 1990Go33 were used to determine specific multipolarities and theoretical internal conversion coefficients: ((97 % M1 + 3 % E2) for the 211.52-keV gamma transition, (67 % M1 + 33 % E2) for the 233.37-keV gamma transition, (86 % M1 + 14 % E2) for the 252.71-keV gamma transition, (99.96 % M1 + 0.04 % E2) for the 277.37-keV gamma transition, (99.75 % M1 + 0.25 % E2) for the 510.74-keV gamma transition, (91.2 % M1 + 8.8 % E2) for the 722.26-keV gamma transition, (99.0 % M1 + 1.0 % E2) for the 763.45-keV gamma transition, and (99.98 % M1 + 0.02 % E2) for the 860.53-keV gamma transition). The assigned multipolarity of the 860.53-keV gamma ray is particularly important in achieving the desired population-depopulation balance for the 2614.552-keV nuclear level. Recommended internal conversion coefficients have been determined from the frozen orbital approximation of Kibédi *et al.* (2008Ki07), based on the theoretical model of Band *et al.* (2002Ba85, 2002Ra45). Ion-pair formation coefficients were calculated by means of the methodology described by Kibédi *et al.* (2008Ki07). Uncertainties of ± 1.5 % were adopted for the E1 and E2 gamma transitions.

A normalisation factor (NF) of 0.997 55 (4) was calculated for the relative emission probabilities of the gamma rays, assuming no direct beta decay to the ground state of ²⁰⁸Pb:

absolute transition probability of 2614.511-keV gamma ray = 100 %,
 relative emission probability of 2614.511-keV gamma ray = 100 %, and
 total theoretical internal conversion coefficient (2614.511-keV E3 transition) = 0.002 46 (4) (2002Ba85, 2002Ra45, 2008Ki07) →

$$P_V^{abs} = P_V^{rel} \times NF = \frac{TP_V^{abs}}{[1 + \alpha_{total}]}$$

and, therefore:

$$NF = \frac{TP_V^{abs}}{[1 + \alpha_{total}] \times P_V^{rel}} = \frac{100}{[1 + 0.00246(4)] \times 100}$$

$$NF = 0.99755(4)$$

Gamma-ray emissions: multiplicities, theoretical internal conversion coefficients (frozen orbital approximation) and ion-pair formation coefficients.

E_γ (keV)	Multipolarity	α_K	α_L	α_{M+}	α_{IPF}	α_{total}
211.52 (2)	97%M1 + 3%E2 δ = 0.18(2) 1957Kr56, 1957Vo22	0.890 (14)	0.1570 (22)	0.049	–	1.096 (17)
233.37 (2)	67%M1 + 33%E2 δ = 0.70(7) 1957Kr56, 1957Vo22	0.51 (3)	0.1136 (18)	0.0364	–	0.66 (3)
252.71 (2)	86%M1 + 14%E2 δ = -0.40(4) 1957Vo22, 1963Da11, 1978Av01	0.495 (14)	0.0926 (14)	0.0284	–	0.616 (15)
277.37 (2)	99.96%M1 + 0.04%E2 δ = 0.02(1) 1957Vo22, 1963Da11, 1976Av03, 1978Av01, 1990Go33	0.432 (6)	0.0739 (11)	0.0231	–	0.529 (8)
486.08 (2)	[M1] 1957Vo22	0.0954 (14)	0.01608 (23)	0.00492	–	0.1164 (17)
510.74 (2)	99.75%M1 + 0.25%E2 δ = -0.05(5) 1957Vo22, 1963Da11, 1976Av03, 1978Av01	0.0835 (13)	0.01406 (21)	0.00434	–	0.1019 (16)
583.187 (2)	E2 1954El07, 1957Kr56, 1963Da11, 1972Ja25, 1978Av01	0.01509 (22)	0.00410 (6)	0.00131	–	0.0205 (3)
588.11 (2)	[M1]	0.0577 (8)	0.00968 (14)	0.00302	–	0.0704 (10)
650.27 (2)	[M1]	0.0444 (7)	0.00742 (11)	0.00228	–	0.0541 (8)
705.34 (2)	[M1]	0.0360 (5)	0.00599 (9)	0.00181	–	0.0438 (7)
722.26 (2)	91.2%M1 + 8.8%E2 δ = 0.31(3) 1976Av03, 1978Av01	0.0317 (6)	0.00534 (10)	0.00166	–	0.0387 (7)
748.87 (2)	[M1]	0.0308 (5)	0.00512 (8)	0.00158	–	0.0375 (6)
763.45 (2)	99.0%M1 + 1.0%E2 δ = -0.10(1) 1957Vo22, 1963Da11, 1978Av01, 1990Go33	0.0291 (4)	0.00484 (7)	0.00146	–	0.0354 (5)
808.32 (13)	–	–	–	–	–	–
821.48 (2)	[M1]	0.0242 (4)	0.00402 (6)	0.00128	–	0.0295(5)
835.90 (11)	–	–	–	–	–	–
860.53 (2)	99.98%M1 + 0.02%E2 δ = 0.015 1957Vo22, 1963Da11, 1972Ja25, 1976Av03, 1978Av01, 1990Go33	0.0215 (3)	0.00356 (5)	0.00114	–	0.0262 (4)
883.59 (2)	[M1]	0.0201 (3)	0.00333 (5)	0.00097	–	0.0244 (4)
927.64 (2)	[M1]	0.01774 (25)	0.00293 (5)	0.00093	–	0.0216 (3)
982.70 (2)	[M1]	0.01530 (22)	0.00253 (4)	0.00077	–	0.0186 (3)
1093.90 (2)	E2	0.00449 (7)	0.000844 (12)	0.000266	–	0.00560 (8)
1126.24 (2)	E1	0.001691 (24)	0.000256 (4)	0.000081	0.00000206 (3)	0.00203 (3)
1160.96 (2)	[M1]	0.01000 (14)	0.001641 (23)	0.000496	0.00000259 (4)	0.01214 (17)
1185.57 (2)	[M1]	0.00947 (14)	0.001555 (22)	0.000480	0.00000501 (7)	0.01151 (17)
1283.04 (2)	[M1]	0.00775 (11)	0.001269 (18)	0.000388	0.0000232 (4)	0.00943 (14)
1380.89 (2)	[M1]	0.00643 (9)	0.001050 (15)	0.000315	0.0000546 (8)	0.00785 (11)
1647.32 (2)	[M1]	0.00411 (6)	0.000669 (10)	0.000207	0.000194 (3)	0.00518 (8)
1744.12 (2)	[M1]	0.00356 (5)	0.000578 (8)	0.000177	0.000255 (4)	0.00457 (7)
2614.511 (10)	E3	0.001708 (24)	0.000292 (4)	0.000089	0.000371 (6)	0.00246 (4)

Beta Particles

Energies

All beta-particle energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies of 2007Ma45 and the Q_{β^-} value of 4999.0 (17) keV adopted from 2003Au03 were used to determine the energies and uncertainties of the beta-particle transitions to the various levels.

Emission Probabilities

The beta-particle emission probabilities were calculated from gamma-ray probability balances, through the recommended gamma-ray emission intensities, and adopted multiplicities and theoretical internal conversion coefficients. A significant majority of the beta-particle transitions were defined as first forbidden non-unique.

Beta-particle emission probabilities per 100 disintegrations of ²⁰⁸Tl.

E_{β} (keV)	P_{β}				Transition type	log ft
	1960Em01	1960Sc07	1967Os01	Recommended value*		
$\beta_{0,23}$ 518.3 (17)	—	—	—	0.052 (5)	1 st forbidden non-unique	6.67
$\beta_{0,21}$ 615.7 (17)	—	—	—	0.017 (5)	1 st forbidden non-unique	7.41
$\beta_{0,20}$ 640.3 (17)	—	—	4.15 (15)	0.045 (4)	1 st forbidden non-unique	7.04
$\beta_{0,19}$ 675.1 (17)	—	—	—	0.005 (2)	allowed	8.1
$\beta_{0,18}$ 702.4 (17)	—	—	—	0.102 (11)	1 st forbidden non-unique	6.82
$\beta_{0,17}$ 737.1(17)	—	—	—	0.002 (1)	1 st forbidden non-unique	8.6
$\beta_{0,13}$ 818.6 (17)	—	—	—	0.231 (9)	1 st forbidden non-unique	6.70
$\beta_{0,12}$ 873.7 (17)	—	—	—	0.174 (9)	1 st forbidden non-unique	6.92
$\beta_{0,8}$ 1003.6 (17)	—	—	—	0.007 (3)	1 st forbidden non-unique	8.5
$\beta_{0,7}$ 1037.8 (17)	3.6	4.6 (2)	< 0.6	3.17 (4)	1 st forbidden non-unique	5.92
$\beta_{0,6}$ 1052.4 (17)	—	—	—	0.048 (3)	1 st forbidden non-unique	7.76
$\beta_{0,5}$ 1079.0 (17)	—	—	—	0.63 (4)	1 st forbidden non-unique	6.68
$\beta_{0,4}$ 1290.5 (17)	24.3	23.9 (8)	21 (2)	24.1 (2)	1 st forbidden non-unique	5.38
$\beta_{0,3}$ 1523.9 (17)	20.6	22.7 (7)	22 (2)	22.1 (5)	1 st forbidden non-unique	5.69
$\beta_{0,2}$ 1801.3 (17)	51.3	48.8 (27)	52 (1)	49.2 (6)	1 st forbidden non-unique	5.61
				Σ 99.9 (8)		

*Recommended emission probabilities derived from evaluated gamma-ray emission probabilities and theoretical internal conversion coefficients.

Atomic Data

The x-ray and Auger-electron data have been calculated using the evaluated gamma-ray data, and atomic data from 1996Sc06, 1998ScZM and 1999ScZX. Both the x-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.01, 28 January 2003). This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

K and L X-ray emission probabilities per 100 disintegrations of ²⁰⁸Tl.

			Energy (keV)	Photons per 100 disint.
XL		(Pb)	9.184 – 15.216	2.75 (12)
	XL ₁	(Pb)	9.184	0.0671 (19)
	XL _α	(Pb)	10.450 – 10.551	1.27 (4)
	XL _η	(Pb)	11.349	0.0209 (7)
	XL _β	(Pb)	12.142 – 13.015	1.155 (25)
	XL _γ	(Pb)	14.765 – 15.216	0.220 (5)
	XK _α	XK _{α2}	(Pb)	72.8049 (8)
XK _{α1}		(Pb)	74.9700 (9)	3.42 (7)
XK' _{β1}	XK _{β3}	(Pb)	84.451)
	XK _{β1} "	(Pb)	84.937) 1.17 (3)
	XK _{β5}	(Pb)	85.470)
XK' _{β2}	XK _{β2}	(Pb)	87.238)
	XK _{β4}	(Pb)	87.580) 0.353 (11)
	XKO _{2,3}	(Pb)	87.911)

Electron energies were determined from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

Data Consistency

A Q_β-value of 4999.0 (17) keV has been adopted from the atomic mass evaluation of Audi *et al.* (2003Au03) while in the course of formulating the decay scheme of ²⁰⁸Tl. This value has subsequently been compared with the Q-value calculated by summing the contributions of the individual emissions to the ²⁰⁸Tl beta-decay process (i.e. β⁻, conversion electrons, γ, etc.):

$$\text{calculated Q-value} = \sum (E_i \times P_i) = 4989 (14) \text{ keV}$$

Percentage deviation from the Q-value of Audi *et al.* is (0.2 ± 0.3) %, which supports the derivation of a highly consistent decay scheme.

Acknowledgement

ALN would like to thank M.J. Martin (ORNL, USA) for an exchange of information in November 2006 with respect to an earlier DDEP evaluation of the β⁻ decay of ²⁰⁸Tl undertaken in January 2004. This further re-evaluation of July 2010 to May 2011 includes a number of adjustments and corrections based on this welcome communication.

References

- 1954El07 L.G. Elliott, R.L. Graham, J. Walker, J.I. Wolfson, Spins and Parities of Energy Levels in Pb^{208} , Phys. Rev. 93 (1954) 356. [α_K]
- 1957Ba05 D.L. Baulch, H.A. David, J.F. Duncan, The Half-life of Thorium C", Australian J. Chem. 10 (1957) 85-87. [Half-life]
- 1957Kr56 E.M. Krisiuk, A.G. Sergeev, G.D. Latyshev. K.I. Il'in, V.I. Fadeev, The Decay Scheme of Tl^{208} , Sov. Phys. JETP 6 (1958) 880-882. [Multipolarity]
- 1957Vo22 V.D. Vorob'ev, K.I. Il'in, T.I. Kol'chinskaia, G.D. Latyshev, A.G. Sergeev, Iu.N. Trofimov, V.I. Fadeev, Internal Conversion Electron Spectrum of Radiothorium, Part III. $H_p = 1380-2700$ and $3500-9000$ Gauss-cm Regions, Izvest. Akad. Nauk SSSR, Ser. Fiz. 21 (1957) 954; Columbia Technical Translation 21 (1958) 956-963. [P_{ce}]
- 1960Em01 G.T. Emery, W.R. Kane, Gamma-ray Intensities in the Thorium Active Deposit, Phys. Rev. 118 (1960) 755-762. [P_β, P_γ]
- 1960Sc07 G. Schupp, H. Daniel, G.W. Eakins, E.N. Jensen, Transition Intensities in the Tl^{208} Beta Decay, the $Bi^{212} \rightarrow Po^{212}$ Decay Scheme, and the Bi^{212} Branching Ratio, Phys. Rev. 120 (1960) 189-198. [P_β, P_γ]
- 1961Si11 L. Simons, M. Brenner, L. Kald, K-E. Nysten, E. Spring, Angular Correlations of Gamma-Gamma Cascades in Pb^{208} , Soc. Sci. Fenn. Comm. Phys. Math. 26 (1961) part 6. [P_γ]
- 1963Da11 H. Daniel, G. Luhns, Aufbau eines automatisch arbeitenden $\pi/2$ -Spektrometers und Untersuchung des Konversionslinienspektrums von In^{114m} und ThB mit Folgeprodukten, Z. Phys. 176 (1963) 30-44. [$P_{ce}, \alpha_K, multipolarity$]
- 1967La20 N.O. Lassen, N. Hornstrup, Half-life of ^{208}Tl (ThC"), Kgl. Danske Videnskab. Selskab., Mat.-Fys. Medd. 36, No. 4 (1967). [Half-life]
- 1967Os01 H. Ostertag, K.H. Lauterjung, Der β -Zerfall des ^{208}Tl , Z. Phys. 199 (1967) 25-40. [P_β]
- 1969Au10 G. Aubin, J. Barrette, G. Lamoureux, S. Monaro, Calculated Relative Efficiency for Coaxial and Planar Ge(Li) Detectors, Nucl. Instrum. Methods 76 (1969) 85-92. [P_γ]
- 1969La23 J.S. Larsen, B.C. Jorgensen, The Decay of ^{208}Tl : Gamma-ray Measurement, Z. Phys. 227 (1969) 65-70. [P_γ]
- 1969Pa02 A. Pakkanen, J. Kantele, P. Suominen, Levels in ^{208}Pb Populated in the Decay of ^{208}Tl (ThC"), Z. Phys. 218 (1969) 273-281. [P_γ]
- 1970Mu21 V.H. Mundschenk, Uber ein Verfahren zur Abtrennung kurzlebiger Radionuklide unter Ausnutzung des Ruckstoeffektes, Radiochim. Acta 14 (1970) 72-77. [Half-life]
- 1971Ac02 R. Ackerhalt, P. Ellerbe, G. Harbottle, The Half-life of $^{208}Tl/ThC$ ", Radiochem. Radioanal. Lett. 8 (1971) 75-77. [Half-life]
- 1972DaZA J. Dalmaso, Recherches sur le Rayonnement Gamma de Quelques Radioelements Naturels Appartenant  la Famille du Thorium, PhD thesis, University of Nice (1972). [P_γ]
- 1972Ja25 P. Jagam, D.S. Murty, Multipole Mixing Ratios of γ -transitions in ^{208}Pb , Nucl. Phys. A197 (1972) 540-552. [P_γ, δ]
- 1973Da38 J. Dalmaso, H. Maria, C. Ythier, tude du Rayonnement γ du Thorium 228 et de ses Derives, et plus Particulierement du Thallium 208 (ThC"), C. R. Acad. Sci. Paris 277B (1973) 467-470. [P_γ]
- 1975Ko02 M. Kortelahti, A. Pakkanen, J. Kantele, Electromagnetic Transition Rates in ^{208}Pb , Nucl. Phys. A240 (1975) 87-97. [P_γ, δ]
- 1976Av03 F.T. Avignone, S.M. Blankenship, W.W. True, Experimental and Theoretical Multipole Mixing Ratios in Transitions of ^{208}Pb , Phys. Rev. C14 (1976) 267-274. [δ]
- 1977Ge12 R.J. Gehrke, R.G. Helmer, R.C. Greenwood, Precise Relative γ -ray Intensities for Calibration of Ge Semiconductor Detectors, Nucl. Instrum. Methods 147 (1977) 405-423. [P_γ]
- 1977La19 F.P. Larkins, Semiempirical Auger-electron Energies for Elements $10 \leq Z \leq 100$, At. Data Nucl. Data Tables 20 (1977) 311-387. [Auger-electron energies]
- 1978Av01 F.T. Avignone, A.G. Schmidt, γ -ray and Internal-conversion Intensity Studies of Transitions in the Decay of ^{228}Th , Phys. Rev. C17 (1978) 380-384. [P_γ, δ]

- 1982Sa36 S. Sadasivan, V.M. Raghunath, Intensities of Gamma Rays in the ^{232}Th Decay Chain, Nucl. Instrum. Methods 196 (1982) 561-563. [P $_{\gamma}$]
- 1983Sc13 U. Schötzig, K. Debertin, Photon Emission Probabilities per Decay of ^{226}Ra and ^{232}Th in Equilibrium with their Daughter Products, Int. J. Appl. Radiat. Isot. 34 (1983) 533-538. [P $_{\gamma}$]
- 1983Va22 R. Vaninbrouckx, H.H. Hansen, Determination of γ -ray Emission Probabilities in the Decay of ^{228}Th and its Daughters, Int. J. Appl. Radiat. Isot. 34 (1983) 1395-1397. [P $_{\gamma}$]
- 1984Ge07 R.J. Gehrke, V.J. Novick, J.D. Baker, γ -ray Emission Probabilities for the ^{232}U Decay Chain, Int. J. Appl. Radiat. Isot. 35 (1984) 581-589. [P $_{\gamma}$]
- 1990Go33 L.I. Govor, A.M. Demidov, V.A. Kurkin, Spectrum and Angular Distribution of γ rays from the Reaction $^{208}\text{Pb}(n,n'\gamma)$, Bull. Acad. Sci. USSR, Phys. Ser. 54 (1990) 147-155. [δ]
- 1992Li05 W.-J. Lin, G. Harbottle, Gamma-ray Emission Intensities of the ^{232}Th Chain in Secular Equilibrium, of ^{235}U and the Progeny of ^{238}U , J. Radioanal. Nucl. Chem. 157 (1992) 367-372. [P $_{\gamma}$]
- 1993El08 O. El Samad, J. Dalmasso, G. Barci-Funel, G. Ardisson, Fast Radiochemical Separation and γ Spectroscopy of Short-lived Thallium Isotopes, Radiochim. Acta 62 (1993) 65-69. [P $_{\gamma}$]
- 1996Sc06 E. Schönfeld, H. Janßen, Evaluation of Atomic Shell Data, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527-533. [X $_K$, X $_L$, Auger electrons]
- 1997Sc21 M. Schramm, K.H. Maier, M. Rejmund, L.D. Wood, N. Roy, A. Kuhnert, A. Aprahamian, J. Becker, M. Brinkman, D.J. Decman, E.A. Henry, R. Hoff, D. Manatt, L.G. Mann, R.A. Meyer, W. Stoeffl, G.L. Struble, T.-F. Wang, Study of Excited States in ^{208}Pb by Particle- γ Coincidences with the $^{207}\text{Pb}(d,p)^{208}\text{Pb}$ and $^{209}\text{Bi}(t,\alpha)^{208}\text{Pb}$ Reactions, Phys. Rev. C56 (1997) 1320-1337. [Nuclear levels, spin, parity]
- 1998ScZM E. Schönfeld, G. Rodloff, Tables of the Energies of K-Auger Electrons for Elements with Atomic Numbers in the Range from $Z = 11$ to $Z = 100$, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX E. Schönfeld, G. Rodloff, Energies and Relative Emission Probabilities of K X-rays for Elements with Atomic Numbers in the Range from $Z = 5$ to $Z = 100$, PTB Report PTB-6.11-1999-1, February 1999. [X $_K$]
- 2000He14 R. G. Helmer, C. van der Leun, Recommended Standards for γ -ray Energy Calibration (1999), Nucl. Instrum. Methods Phys. Res. A450 (2000) 35-70. [E $_{\gamma}$]
- 2002Ba85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, Dirac-Fock Internal Conversion Coefficients, At. Data Nucl. Data Tables 81 (2002) 1-334. [ICC]
- 2002Ra45 S. Raman, C.W. Nestor, Jr., A. Ichihara, M.B. Trzhaskovskaya, How Good are the Internal Conversion Coefficients Now? Phys. Rev. C66 (2002) 044312, 1-23. [ICC]
- 2003Au03 G. Audi, A.H. Wapstra, C. Thibault, The AME2003 Atomic Mass Evaluation (II). Tables, Graphs and References, Nucl. Phys. A729 (2003) 337-676. [Q-value]
- 2006Va23 S.I. Vasil'ev, K.Ya. Gromov, A.A. Klimenko, Zh.K. Samatov, A.A. Smol'nikov, V.I. Fominykh, V.G. Chumin, Coincidence Summing in γ -ray Spectra and Determination of the Intensity of Weak Crossover γ Transitions, Instruments and Experimental Techniques 49 (2006) 34-40. [Crossover γ]
- 2007Ma45 M.J. Martin, Nuclear Data Sheets for $A = 208$, Nucl. Data Sheets 108 (2007) 1583-1806. [Nuclear structure, level energies]
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr., Evaluation of Theoretical Conversion Coefficients using BrIcc, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202-229. [ICC]

²⁰⁹Tl- Comments on Evaluation of Decay Data

By F.G. Kondev

This evaluation was completed in May 2011, with a literature cut off by the same date, as a part of ANL commitment to the IAEA-CRP on “Updated Decay Data Library for Actinides”.

1. Decay Scheme

The nuclide ²⁰⁹Tl ($J^\pi=1/2^+$) disintegrates 100 % by β^- emissions. The strongest β^- -decay branch of 97.70 (15) % populates the $J^\pi=1/2^-$ excited state at 2149.29 keV of the daughter nuclide ²⁰⁹Pb. The decay scheme of ²⁰⁹Tl was constructed by the evaluator, based on the work of Gromov (2000Gr35) and Ardisson (1998Ar03). The ENSDF evaluation of Martin (1991Ma16) was consulted for J^π and multipolarity assignments to levels in ²⁰⁹Pb.

2. Nuclear Data

Adopted $Q(\beta^-)$ value of 3976 (8) keV is taken from the evaluation of Audi *et al.* (2003Au03).

The experimental data for the half-life of the ²⁰⁹Tl ground state are very scarce. The value of 2.161 (7) min (1998Ar03) is adopted in the present evaluation. It is in agreement with the other known, but less precise, value of 2.20 (17) min (1950Ha64).

2.1. β^- Transitions

The values for the maximum β^- -decay energies, $E_{\beta^- \text{max}}$, presented in Table 1, were deduced from $Q(\beta^-) = 3976$ (8) keV (2003Au03) and the level energies deduced in the present evaluation, as detailed in section 2.2. The β^- -decay transition probabilities, P_β , were deduced from the decay scheme and the corresponding absolute γ -ray transition probabilities. The sum of the β^- intensities to levels above 2149 keV is 2.30 (15) %. Then the β^- feeding to the 2149-keV level is $(100 - 2.30 (15)) \% = 97.70 (15) \%$. The $\log ft$ values were calculated using the LOGFT program from the ENSDF evaluation package. The $\log f$ values are based on the work of Gove and Martin (1971Go40).

2.2. Gamma Transitions and Electron Internal Conversion Coefficients

The γ -ray transition energy data are presented in Table 2. Statistical analysis using the LWEIGHT program has been performed and the corresponding gamma-ray energies were deduced (the last column of Table 2). With those energies, the level scheme was fitted using the *gtol* program from the ENSDF analysis package and new level energies (shown in Table 1) were obtained.

The γ -ray transition multiplicities were taken from the ENSDF evaluation of Martin (1991Ma16) and the recent work of Gromov (2000Gr35) and Ardisson (1998Ar03). The electron conversion coefficients were calculated using the BrIcc code (2008Ki07).

Table 1. Level energies, quantum numbers, $E_{\beta_{0,1 \text{max}}}$, P_β and $\log ft$ values in decay of ²⁰⁹Tl ($J^\pi=1/2^+$)

Level energy (keV)	J^π	$E_{\beta^- \text{max}}$ (keV)	P_β (%)	Nature	$\log ft$
3388.96 (13)	(1/2,3/2)	587 (8)	0.420 (22)		
3361.36 (17)	(1/2,3/2)	615 (8)	0.10 (3)		
3069.72 (13)	3/2-	906 (8)	0.645 (16)	first forbidden	6.3
2905.14 (25)	3/2-	1071 (8)	0.70 (9)	first forbidden	6.5
2524.14 (25)	(1/2,3/2)+	1451 (8)	0.070 (15)	allowed	8.0
2460.8 (3)	(5/2)-	1515 (8)	0.031 (16)	first forbidden unique	9.2

2315.68 (13)	(3/2)-	1660 (8)	0.32 (11)	first forbidden	7.5
2149.29 (6)	1/2-	1827 (8)	97.70 (15)	first forbidden	5.2
2032.07 (6)	1/2+	1944 (8)	<0.1	allowed	> 8.3
1566.94 (5)	5/2+				
0.0	9/2+				

The gamma-ray emission probability data are presented in Table 3. The unplaced gamma rays and their emission probabilities are presented in Table 4. Future work is merited to obtain a more complete decay scheme of ²⁰⁹Tl.

3. Atomic Data

The Atomic data (Fluorescence yields, X-Ray energies and Relative probabilities, and Auger electrons energies and Relative probabilities) were provided by the Saisinuc software (2008DuZX). Details regarding the origin of these data can be found in 1996Sc06, 1998ScZM, 1999ScZX, 2000Sc47 and 2003De44.

Table 2. Measured, deduced and adopted gamma-ray energies in β^- -decay of ²⁰⁹Tl

2003ChZV	2000Gr35	1999GrZT	1998Ar03	1993El08	1989Ko26	1986He06	1981Di14	1977Vy02	adopted
	117.18 (10)	117.1 (3)	117.24 (5)	117.24 (1)	117.21 (1)		117.25 (5)	117.211 (21)	117.224 (7)
			284.04 (23)	284.04 (25)					284.04 (23)
			311.5 (3)	311.5 (3)					311.5 (3)
			375.5 (2)	375.5 (2)					375.5 (2)
465.2	465.21 (4)	465.0 (4)	465.10 (5)	465.10 (1)	465.14 (1)	465.4 (1)	465.1 (2)	465.065 (25)	465.128 (24)
582.4	582.4 (2)								582.4 (2)
	748.5 (3)		748.0 (3)	748.0 (3)					748.3 (2)
	755.6 (3)								755.6(3)
	873.5 (4)								873.5 (4)
920.2	920.8 (1)	919.9 (3)	920.34 (9)	920.34 (7)	920.2 (3)				920.43 (11)
1239.8	1239.7 (2)	1239.2 (3)	1239.76 (15)	1239.76 (15)					1239.66 (11)
1329.2	1329.3 (3)	1329.6 (3)	1329.3 (3)	1329.3 (3)	1239.5 (5)				1329.29 (16)
1566.9	1566.9 (3)	1566.7 (3)	1566.96 (5)	1566.96 (1)	1567.11 (2)		1566.9 (2)	1566.95 (6)	1566.93 (5)
	2149.0 (10)								2149.0 (10)
2315.9	2315.9 (3)	2315.7 (3)							2315.80 (21)

Table 3. Measured, deduced and adopted γ -ray emission probabilities for γ -ray transitions in β^- -decay of ²⁰⁹Tl

Eg, keV	2003ChZV	2000Gr35	1999GrZT	1998Ar03	1993EI08	1989Ko26	1981Di14	1977Vy02	adopted
117.24 (1)		78 (4)	74 (2)	73 (4)	73 (1)	85 (4) *	85.6 (59) *	84 (2) *	77.22 (27) ^{a)}
284.04 (25)				0.14 (7)	0.14 (7)				0.14 (7)
311.5 (3)				0.028 (14)	0.028 (14)				0.028 (14)
375.5 (2)				0.070 (15)	0.070 (15)				0.070 (15)
465.10 (1)	80.4 (21) *	97 (5)	93.2 (16) *	95 (5)	95 (5)	96 (4)	99.1 (64)	100 (3) *	96.62 (5) ^{a)}
582.4	0.33 (3)	0.28 (4)							0.312 (24)
748.5 (3)		0.07 (3)		0.09 (3)	0.086 (30)				0.080 (21)
755.6(3)		0.11 (2)							0.11 (2)
873.5 (4)		0.59 (8)							0.59 (8)
920.8 (1)	0.62 (3)	0.63 (5)	0.63 (2)	0.70 (7)	0.70 (7)	0.63 (6)			0.631 (15)
1239.8	0.45 (4)	0.42 (7)	0.41 (3)	0.31 (12)	0.31 (12)				0.420 (22)
1329.2	0.14	0.10 (3)	0.21 (3)	0.026 (5)	0.026 (5)	0.42 (4)			0.10 (3)
1566.9	100 (1)	100 (5)	100.0 (8)	100 (5)	100 (5)	100 (4)	100.6 (64)	93 (3) *	99.707 (5) ^{a)}
2149.0 (10)	<0.0006	0.015 (5)							0.015 (5)
2315.9 (3)	0.0284 (24)	0.03 (1)	0.030 (5)						0.0288 (21)

* not included in the statistical analysis

^{a)} deduced from $100/(1+\alpha_T)$ due to cascading.

Table 4. Gamma-ray energies and emission probabilities for transitions in β^- -decay of ²⁰⁹Tl, which were not placed in the decay scheme

2003ChZV		2000Gr35		1999GrZT		1998Ar03	
		469.9	0.12 (3)			469.7 (3)	0.03 (2)
		860.5 (3)	0.26 (4)				
		890.0 (4)	0.12 (3)				
970.3	0.054 (15)	902.8 (4)	0.10 (2)				
		1661.1 (5)	0.10 (2)				
		1673.2 (4)	0.48 (4)				
		1781.7 (5)	0.04 (2)				
				2005.3 (2)	0.020 (5)		
2032.1	<0.019	2032.1 (5)	0.001				
2548.2	0.015 (6)						

4. References

- 1950Ha64 F. Hagemann. Phys. Rev. 79 (1950) 534
(Half-life)
- 1971Go40 N.B. Gove, M.J. Martin. Nucl. Data Tables A10 (1971) 205
(Log ft values)
- 1977Vy02 T. Vylvov, N.A. Golovkov, B.S. Dzhelepov, R.B. Ivanov, M.A. Mikhailova, Y.V. Norseev, V.G. Chumin. Bull. Acad. Sci. USSR, Phys.Ser. 41 (1977) 85
(Gamma-ray emission energies and probabilities)
- 1981Di14 J.K. Dickens, J.W. McConnell. Radiochem. Radioanal. Lett. 47 (1981) 331
(Gamma-ray emission energies and probabilities)
- 1986He06 R.G. Helmer, C.W. Reich, M.A. Lee, I. Ahmad. Appl. Radiat. Isot. 37 (1986) 139
(Gamma-ray emission energies and probabilities)
- 1989Ko26 M.C. Kouassi, A. Hachem, C. Ardisson, G. Ardisson. Nucl. Instrum. Methods Phys. Res. A280 (1989) 424
(Gamma-ray emission energies and probabilities)
- 1991Ma16 M.J. Martin. Nucl. Data Sheets 63 (1991) 723
(Nuclear levels, multipolarities)
- 1993El08 O. El Samad, J. Dalmaso, G. Barci-Funel, G. Ardisson. Radiochim. Acta 62 (1993) 65
(Gamma-ray emission energies and probabilities)
- 1996Sc06 E. Schönfeld, H. Janssen. Nucl. Instrum. Methods Phys. Res. A369 (1996) 527
(K-shell fluorescence yields)
- 1998ScZM E. Schönfeld, G. Rodloff. Report PTB-6.11-98-1 Braunschweig (1998)
(K Auger electron energies)
- 1988Ar03 G. Ardisson, V. Barci, O. El Samad. Phys. Rev. C57 (1998) 612
(Gamma-ray emission energies and probabilities)
- 1999GrZT K.Ya. Gromov, Sh.R. Malikov, T.M. Muminov, Zh.K. Samatov, Zh. Sehrehehtehr, V.I. Fominykh, V.V. Tsupko-Sitnikov, V.G. Chumin. Proc. 49th Ann. Conf. Nucl. Spectrosc. Struct. At. Nuclei, Dubna (1999) 117
(Gamma-ray emission energies and probabilities)
- 1999Sc47 E. Schönfeld, G. Rodloff. Report PTB-6.11-1999-1 Braunschweig (1999)
(K X-ray energies and relative emission probabilities)
- 2000Sc47 E. Schönfeld, H. Janssen. Appl. Radiat. Isot. 52 (2000) 595
(EMISSION program and X-ray and Auger electron emission probabilities)
- 2000Gr35 K.Ya. Gromov, S.A. Kudrya, Sh.R. Malikov, T.M. Muminov, Zh.K. Samatov, Zh. Sereeter, V.I. Fominykh, V.G. Chumin. Bull. Rus. Acad. Sci. Phys. 64 (2000) 1770
(Gamma-ray emission energies and probabilities)
- 2003Au03 G. Audi, A.H. Wapstra, C. Thibault. Nucl. Phys. A729 (2003) 337
(Q value)
- 2003De44 R.D. Deslattes, E.G. Kessler Jr., P. Indelicato, L. de Billy, E. Lindroth, J. Anton. Rev. Mod. Phys. 75 (2003) 35
(K and L X-ray energies)
- 2003ChZV V.G. Chumin, V.I. Fominykh, K.Ya. Gromov, A.A. Klimenko, S.A. Kudrya, A.A. Smolnikov, S.I. Vasiliev. Proc. 53rd Ann. Conf. Nucl. Spectrosc. Struct. At. Nuclei, Moscow (2003) 105
(Gamma-ray emission energies and probabilities)
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor Jr.. Nucl. Instrum. Methods Phys. Res. A589 (2008) 202
(Theoretical ICCs)
- 2008DuZX C. Dulieu, M.M. Bé, V. Chisté. Proc. Int. Conf. on Nuclear Data for Science and Technology, Nice, France, 22-27 April 2007 (2008) 97
(SAISINUC software)

²⁰⁹Pb - Comments on evaluation of decay data by F. G. Kondev

This evaluation was completed in February 2011, with the same literature cut off date, as a part of ANL commitment to the IAEA-CRP on “Updated Decay Data Library for Actinides”.

1 Decay Scheme

The nuclide ²⁰⁹Pb ($J^\pi = 9/2^+$) disintegrates 100 % by β^- emissions with a single β^- -decay branch to the ground state ($J^\pi = 9/2^-$) of the daughter nuclide ²⁰⁹Bi. The level schemes of ²⁰⁹Pb and ²⁰⁹Bi, including level energies and J^π values, are based on the ENSDF evaluation of Martin (1991Ma16).

2 Nuclear Data

Adopted $Q(\beta^-)$ value of 644.0 (12) keV is taken from the evaluation of Audi *et al.* (2003Au03).

The experimental half-life data for the ²⁰⁹Pb ground state are listed in Table 1. The LRSW value of $T_{1/2} = 3.277$ (15) h was adopted ($\chi^2_v = 1.87$, which is smaller than the critical value of $\chi^2_{v, \text{crit}} = 3.32$ (99 % confidence level)).

Table 1. Experimental data for the half-life of ²⁰⁹Pb.

Author	$T_{1/2}$ (h)	Used in evaluation
1972Be44	3.253 (14)	Yes
1971Pe03	3.31 (3)	Yes
1959Po64	3.31 (3)	Yes
1942Ma03	3.3 (1)	Yes
1941Fa04	3.32(3)	Yes
1940Kr08	2.75 (5)	No

2.1 β^- Transitions

The decay of ²⁰⁹Pb proceeds with a single β^- transition directly to the ²⁰⁹Bi ground state. The maximum β^- -decay energy recommended in Table 2 was deduced from $Q(\beta^-) = 644.0$ (12) keV (2003Au03). The $\log ft$ value was calculated using the LOGFT program from the ENSDF evaluation package, which is based on the work of Gove and Martin (1971Go40).

Table 2. Level energy, quantum number, $E_{\beta_{0,1} \text{max}}$, P_β and $\log ft$ value in decay of ²⁰⁹Pb.

	Level energy (keV)	J^π	$E_{\beta\text{-max}}$ (keV)	P_β (%)	Nature	$\log ft$
$\beta_{0,0}$	0.0	9/2-	644.0 (12)	100	First forbidden non-unique	5.536 (4)

3 References

- 1940Kr08 R. S. Krishnan, E. A. Nahum. Proc. Cambridge Phil. Soc. 36(1940)490 (Half-life)
1941Fa04 K. Fajans, A. F. Voigt. Phys. Rev. 60(1941) 619 (Half-life)
1942Ma03 W. Maurer, W. Ramm. Z. Phys. 119 (1942)602 (Half-life)
1959Po64 A. Poularikas, R.W. Fink. Phys. Rev. 115(1959)989 (Half-life)
1971Go40 N. B. Gove, M. J. Martin. Nucl. Data Tables A10(1971)205 (log ft values)
1971Pe03 B. I. Persson, I. Plesser, J.W. Sunier. Nucl. Phys. A167(1971)470 (Half-life)
1972Be44 H. Behrens, M. Kobelt, W. G. Thies, H. Appel. Z. Phys. 252(1972)349 (Half-life)
1991Ma16 M. J. Martin. Nucl. Data Sheets 63(1991)723 (Nuclear levels)
2003Au03 G. Audi, A. H. Wapstra, C. Thibault. Nucl. Phys. A729(2003)337 (Q value)

²⁰⁹Po - Comments on evaluation of decay data by V. Chisté and M. M. Bé

This evaluation was completed in December 2009. The literature available by September 2009 was included.

1 Decay Scheme

²⁰⁹Po disintegrates by alpha emission to excited levels and to the ground state level of ²⁰⁵Pb and by electron capture to the 896-keV excited level of ²⁰⁹Bi.

A good agreement was found between the adopted Q_α and Q_{EC} values from Audi (2003Au03) and the effective Q values, Q_α (4957 (2) keV) and Q_{EC} (1891 (20) keV), calculated from the decay scheme data (and branching ratio).

2 Nuclear Data

The Q values are from the atomic mass evaluation of Audi *et al.* (2003Au03).

Experimental ²⁰⁹Po half-life values (in years) are given in Table 1.

Table 1: Experimental values of ²⁰⁹Po half-life.

Reference	Experimental value (a)	Comments
C. G. Andre (1956An05)	102 (5)	From ²⁰⁹ Po/ ²⁰⁸ Po mass and activity ratios and $T_{1/2}(\text{208Po}) = 2.898 (2) \text{ a}$ (see 1991Ma16).
R. Collé (2007Co07)	128 (7)	Decay data from two separate primary standardizations of a ²⁰⁹ Po solution standard, carried out ~ 12 years apart.
Recommended value	115 (13)	$\chi^2 = 6.9$

The value from 2007Co07 is not a direct measurement of the ²⁰⁹Po half-life. R. Collé said in a private communication: "My paper which stated the value **128 a** was not a new determination... The whole point was to show that there was evidence to suggest and support that the extant **102 a** value is very wrong, perhaps by 25 %".

However, to take into account all scarce information available, the evaluators have chosen to adopt the simple mean of the two existing values (1956An05 and 2007Co07) with an uncertainty covering them. Then, the recommended value is 115 (13) a.

2.1 α Transitions and Emissions

The energies of the α -particle transitions given in Section 2.1 have been obtained from the Q_α (2003Au03) and the ²⁰⁵Pb level energies from F. G. Kondev (2004Ko28), given in Table 2.

Table 2: ²⁰⁵Pb levels populated in the ²⁰⁹Po α -decay.

Level number	Energy (keV)	Spin and parity	Half-life	Probability of α transition (%)
0	0.0	5/2 ⁻	17.3 (7) 10 ⁶ a	19.8 (32)
1	2.329 (7)	1/2 ⁻	24.2 (4) μ s	79.2 (32)
2	262.833 (25)	3/2 ⁻		0.548 (7)

The energies of the $\alpha_{0,0}$, $\alpha_{0,1}$ and $\alpha_{0,2}$ emissions given in Section 4 are from A. M. Mandal (1989Ma05). In 1989Ma05, two weak alpha transitions of 4310 and 4110 keV respectively, were reported. These alpha transitions were not reported here by the evaluators, because other authors have doubts about their existence (1996Sc24, 2004Ko28).

The transition intensity of the $\alpha_{0,2}$ transition has been deduced from the $P(\gamma + ce)$ decay scheme balance. For the $(\alpha_{0,0} + \alpha_{0,1})$, an unresolved doublet, the evaluators decided to follow the explanation given by V. Chechev (private communication and based on the Schmorak article (1980Sc26)):

“Schmorak found, for odd-A nuclides, that all known cases of hindrance factors (HF) for $p_{1/2}$ to $p_{1/2}$ favoured alpha transitions are grouped around $HF = 1.3$. For the Po-209 alpha transition to the 2.3 keV excited level (1/2-) in Pb-205, this allows us to adopt $HF=1.3$. From $HF=1.3$, alpha intensity is derived being approximately 80 %.

Then, for $p_{1/2}$ to $f_{5/2}$ alpha transitions, which interest us, i.e. the Po-209 alpha transition to the 5/2- ground state of Pb-205, the $HF (= 6)$ is deduced from known (measured) HF for such alpha transitions (Rn-211 to Po-207: 6.75; Ra-213 to Rn-209: 6.39; Th-215 to Ra-211: 6.5) and then, an alpha intensity of approximately 20 % is derived.

Therefore, a ratio of 80/20 (1980Sc26) can be derived from HF systematic and analysis of level characteristics.”

Then the total alpha transition intensity was deduced from the $P(\alpha_{0,2}) (= 0.548 (7) \%)$ and $P_{EC} (= 0.454 (7) \%)$:

$$P(\alpha_{0,0} + \alpha_{0,1}) = 100 - P_{EC} - P(\alpha_{0,2}) = 98.998 (10) \%$$

With this value of $P(\alpha_{0,0} + \alpha_{0,1})$, using the estimation $P(\alpha_{0,1})/ P(\alpha_{0,0}) = 80/20$ and accepting a relative uncertainty equal to approximately 20 %, the evaluators have obtained the individual values:

$$P(\alpha_{0,1}) = 79.2 (32) \%$$

$$P(\alpha_{0,0}) = 19.8 (32) \%$$

2.2 Electron Capture Transition

The energy of the electron capture transition has been obtained from the $Q(EC)$ value (2003Au03) and the ²⁰⁹Bi level energy of 896.29 (5) keV given by M. J. Martin (1991Ma16).

The electron capture probability ($P_{EC} = 0.454 (7) \%$) has been deduced from gamma-ray transition intensity imbalance at the 896-keV level.

P_K, P_L, P_M values have been calculated for 2nd forbidden unique electron-capture transition in the decay of ²⁰⁹Po to the excited state in ²⁰⁹Bi using the LOGFT computer program.

2.3 γ Transitions

The gamma-ray transition with energy of 2.328 (7) keV has not been observed directly in the ²⁰⁹Po decay but it was studied in the decay of ²⁰⁹Bi to ²⁰⁵Pb (1971Jo06). The transition probability for this gamma-ray has been obtained from the intensity balance at the ²⁰⁵Pb 2.3-keV level.

The transitions probabilities values for the remaining gamma-rays have been deduced using the γ -ray emission probabilities and the relevant internal conversion coefficients (see **4.2 Gamma Emissions**).

For the three (M1 + E2) γ transitions (²⁰⁵Pb: 260- and 262-keV; ²⁰⁹Bi: 896-keV), the mixing ratios (δ) were deduced by comparison between experimental values of K internal conversion coefficients and the theoretical K ICC calculated using the BrIcc computer code (2008Ki07).

Table 3 shows the experimental and evaluated values of α_K , as well as the deduced mixing ratios.

Table 3: Experimental and evaluated internal conversion coefficients and mixing ratios.

Energy (keV) Reference	260	262	896
G. R. Hagee (1966Ha29)	0.495 (10)	0.495 (10)	0.0170 (5)
A. M. Mandal (1989Ma05)	0.49 (5)	0.49 (5)	
F. Schima (1996Sc24)	0.538 (20)	0.524 (20)	
Evaluated α_K	0.503 (12)	0.500 (9)	0.0170 (5)
χ^2	1.9	0.9	-
δ (mixing ratio)	0.16 (6)	0.05 (7)	-0.62 (4)
α_K theoretical (BRICC)	0.503 (12)	0.500 (9)	0.0170 (5)
α_T theoretical (BRICC)	0.617 (13)	0.612 (10)	0.0208 (6)

The theoretical internal conversion coefficients (ICCs) and the associated uncertainties given in Table 3 have been obtained using the BrIcc computer code with “the frozen orbital approximation” (2008Ki07).

3 Atomic Data

Atomic values, ω_K , ω_L and n_{KL} and X-ray and Auger electron relative probabilities are from Schönfeld and Janßen (1996Sc06).

4 Emissions

4.1 K x-rays

The X-ray absolute emission probabilities have been calculated from the decay scheme data using the EMISSION computer program and compared in Table 4 with the measured values. These values are in a slight agreement; it is difficult to draw a definite conclusion because there is an unresolved doublet, the Bi $K_{\alpha 2}$ – Pb $K_{\alpha 1}$ at 74.89 keV, which is difficult to separate in the spectrum analysis.

Table 4: Experimental and recommended (calculated) values of K_{α} X-ray absolute emission probabilities.

	F. Schima (1996Sc24)	Recommended values
Bi K_{α} X-ray (74.82 – 77.11 keV)	0.202 (5)	0.248 (3)
Pb K_{α} X-ray (72.80 – 74.97 keV)	0.136 (5)	0.128 (2)

4.2 Gamma emissions

The energies of the γ -rays given in section 6.2 are from F. G. Kondev (2004Ko28).

The experimental and recommended values of γ -ray emission probabilities are given in Table 5.

Table 5: Experimental and recommended values of γ -ray emission probabilities.

E_{γ} (keV)	G. R. Hagee (1966Ha29) ^a	A. M. Mandal (1989Ma05) ^b	F. Schima (1996Sc24)	Recommended values.
260.5 (1)	0.391	100	0.254 (3)	0.254 (3)
262.8 (1)	0.391	33.3 (16)	0.085 (2)	0.085 (2)
896.6 (1)	0.263	108.1 (75)	0.445 (7)	0.445 (7)

^a G. R. Hagee was unable to resolve the 260.5- and 262.8-keV γ -ray doublet and the values are given without uncertainties. Not used.

^b A. M. Mandal quoted relative intensities normalized to the 260.5-keV γ -ray.

The Mandal values (1989Ma05) were omitted from analysis because of a lack of information in the article

about the experimental measurements carried out and, therefore on the results. Then, the adopted values of the absolute emission intensities are the most recent values of F. J. Schima (1996Sc24).

5 Electron emissions

The conversion electron emission probabilities have been deduced from ICC values and γ -ray emission probabilities.

6 References

- 1956An05 C. G. Andre, J. R. Huizenga, J. F. Mech, W. J. Ramler, E. G. Rauh, S. R. Rocklin, Phys. Rev. 101(1956)645 [Half-life].
- 1966Ha29 G. R. Hagee, R. C. Lange, J. T. McCarthy, Nucl. Phys. 84(1966)62 [Gamma-ray emission probabilities].
- 1971Jo06 W. C. Johnston, W. H. Kelly, S. K. Haynes, K. L. Kosanke, W. C. McHarris, Phys. Rev. Lett. 26(1971)1043 [2.3 keV gamma-ray transition energy].
- 1980Sc26 M. R. Schmorak, Nucl. Data Sheets 31(1980)283 [$P_{\alpha}(0-, 2.3\text{-keV})$].
- 1989Ma05 A. M. Mandal, S. K. Saha, S. M. Sahakundu, A. P. Patro, J. Phys. (London) G15(1989)173 [Gamma-ray emission probabilities].
- 1991Ma16 M. J. Martin, Nucl. Data Sheets 63(1991)808 [^{209}Bi level energy, spin and parity].
- 1996Sc24 F. J. Schima, R. Collé, Nucl. Instrum. Meth. Phys. Res. A369(1996)498 [Gamma-ray emission probabilities].
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].
- 2004Ko28 F. G. Kondev, Nucl. Data Sheets 101(2004)586 [Level energies, spins and parities].
- 2007Co07 R. Collé, L. Laureano-Perez, I. Outola, Appl. Rad. Isotopes 65(2007)728 [Half-life].
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr., Nucl. Instrum. Meth. Phys. Res. A589(2008)202 [Theoretical ICCs].

**²¹⁰Tl - Comments on evaluation of decay data
by V. Chisté and M. M. Bé**

This evaluation was completed in 2007. Literature available by August 2007 has been included.

1 Decay Scheme

²¹⁰Tl disintegrates by beta minus emission to excited levels of ²¹⁰Pb. A weak delayed neutron emission was reported (1961St20 and 1957Ko42). Level energies, spins and parities are from the mass-chain evaluation of E. Browne (2003Br13) and B. Harmatz (1981Ha54).

This decay scheme is mainly based on the measurements of P. Weinzierl (1964We06). Several inconsistencies appeared :

- β^- branching to levels : 3879-, 3458-, and 3069-keV were deduced from γ -ray transition intensity imbalance. β^- feedings to the 1096 - and 1192-keV levels are uncertain. There is no experimental evidence for β^- transitions with energy > 3 MeV to these levels. β^- feedings the 1869-, 2208- and 2412-keV levels, suggested by γ -ray transition intensity imbalances ($< 10\%$, $< 9\%$ and $< 12\%$, respectively), are uncertain.
- An 83-keV γ -ray is not placed in the present decay scheme as suggested by B. Harmatz (1981Ha54) (transition between 1275-keV level and 1192-keV level), because there is no experimental evidence that the 1275-keV level in ²¹⁰Pb was populated in the β^- decay of ²¹⁰Tl.

These discrepancies cannot be resolved without new experimental results. New measurements are strongly suggested.

Some agreement was found between the adopted $Q(\beta^-)$ value of Audi and the effective $Q(\beta^-)$ value of 5470 (1000) keV calculated from decay scheme data, which indicates a consistency and correctness of the decay scheme.

2 Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2003Au03).

Experimental ²¹⁰Tl half-life values (in minutes) are given in Table 1:

Table 1: Experimental values of ²¹⁰Tl half-life.

Reference	Experimental value (min)	Comments
M. Curie (1931Cu01)	1.32	Not used. No uncertainty.
A.V. Kogan (1957Ko42)	1.50 (25)	
P. Weinzierl (1964We06)	1.30 (3)	
Recommended value	1.30 (3)	$\chi^2 = 0.63$

A weighted average has been calculated using Lweight computer program (version 3)The largest contribution to the weighted average comes from P. Weinzierl (1964We06), amounting to a statistical weight of 98 %.

The recommended value of ²¹⁰Tl half-life is the weighted average of 1.30 minutes with an internal uncertainty of 0.03 minutes. The reduced- χ^2 value is 0.63.

2.1 β^- Transitions and Emissions.

The end-point energies of the β^- transitions in the decay of $^{210}\text{Tl} \rightarrow ^{210}\text{Pb}$ have been obtained from the $Q(\beta^-)$ value (2003Au03) and the level energies given by E. Browne (2003Br13).

The adopted β^- transition probabilities were deduced from the $P(\gamma + ce)$ balance at each level of the decay scheme. Table 2 shows the adopted β^- transition probabilities compared with the only three β^- transitions reported by P. Weinzierl (1964We06). No β^- transitions with $E_{\beta^-} > 3\text{MeV}$ were observed by these authors.

Table 2: Experimental and recommended (calculated) values of β^- transition probabilities.

Level	Energy (keV)	P. Weinzierl (1964We06)	Adopted values
11	1380 (12)	25 %	2 %
10	1603 (12)		7 %
9	1860 (12)	56 %	24 %
8	2024 (12)		10 %
7	2413 (12)	19 %	10 %
3	4290 (12)		31 %
2	4386 (12)		13 %

The sum of the adopted β^- transition probabilities is equal to 97 %. The 3 % missing cannot be placed in the decay scheme without more information about the β^- decay of ^{210}Tl .

The values of $\lg ft$ and the average β^- energies have been calculated using the computer program LOGFT for β^- transitions.

2.2 g Transitions.

The transition probabilities were deduced from the absolute γ -ray emission intensities and the relevant internal conversion coefficients. (see 5.2 g Emissions).

Multipolarities of the γ -ray transitions were deduced from conversion electron measurements and K/L ratios of 1964We06:

83-keV γ -ray: [E2]	97-keV γ -ray: M1 + E2	296-keV γ -ray: E2
356-keV γ -ray: [M1]	356-keV γ -ray: [M1]	799-keV γ -ray: E2
1070-keV γ -ray: [E1]		

The internal conversion coefficients (ICC's) for these γ -ray transitions were calculated using the BrIcc computer program (calculation for 'frozen orbital approximation'), which interpolates from theoretical values of I. M. Band *et al.* (2002Ba85).

Due to the large uncertainty on the 83- and 97-keV transition energy, only estimated ICC values are given.

3 Atomic Data.

Atomic values, ω_K , ω_L and n_{KL} and the X-ray relative probabilities are from Schönfeld and Jäpfen (1996Sc06).

4 Electron Emissions.

The conversion electrons emission probabilities have been deduced using the γ -ray emission intensities and ICC's.

5 Photon Emissions.

5.1 X-ray Emissions.

The X-ray absolute intensities have been calculated from γ -ray data and ICC using the EMISSION computer program. The KX-ray intensity is compared in Table 3 to the measured value of P. Weinzierl (1964We06).

Table 3: Experimental and recommended (calculated) values of X-ray absolute intensities.

	P. Weinzierl (1964We04)	Recommended value
K x-ray	20 (4) %	23 (11) %

5.2 γ Emissions.

The energies of the γ -ray emissions given in Section 5 are from E. Browne (2003Br13).

The experimental relative γ -ray emission intensities measured by P. Weinzierl (1964We06) (single experimental data set found in the literature) given in Table 4 are relative to that of the 799-keV γ -ray. Only one set of measured data (1964We06) is available.

Table 4: The experimental data set of the relative γ -ray emission intensities.

Energy (keV)	Relative γ -ray Emission intensity (%) (1964We06)
83 ^(a)	2.0
97	4 (2)
296	80 (10)
356 ^(a)	4 (2)
382 ^(a)	3 (2)
480	2 (1)
670 ^(a)	2 (1)
799	100
860	7 (2)
910 ^(a)	3 (2)
1070	12 (5)
1110	7 (2)
1210	17 (4)
1316	21 (5)
1410	5 (2)
1490 ^(a)	2 (1)
1540 ^(a)	2 (1)
1590	2 (1)
1650 ^(a)	2 (1)
2010	7 (2)
2090 ^(a)	5 (2)
2270	3 (2)
2360	8 (3)
2430	9 (3)

(a) γ -ray not placed in level scheme as explained in Weinzierl (1964We06).

The normalization factor of **98.969 (30)** to convert the relative γ -ray emission intensities to absolute intensities was obtained using the formula of :

$$N = \left(\frac{100}{(1 + a_T)P_{rel}(799g)} \right)$$

The uncertainties were calculated through their propagation on the above formula.

The evaluated relative and absolute γ -ray emission intensities are given in Table 5.

Table 5: Evaluated relative and absolute γ -ray emission intensities.

Energy (keV)	Relative γ -ray Emission intensity (%)	Absolute γ -ray emission intensity (%)
83 ^(a)	2.0	1.98 (40)
97	4 (2)	4 (2)
296	80 (10)	79 (10)
356 ^(a)	4 (2)	4 (2)
382 ^(a)	3 (2)	3 (2)
480	2 (1)	2 (1)
670 ^(a)	2 (1)	2 (1)
799	100	98.969 (30)
860	7 (2)	6.9 (20)
910 ^(a)	3 (2)	3 (2)
1070	12 (5)	11.9 (49)
1110	7 (2)	6.9 (20)
1210	17 (4)	16.8 (40)
1316	21 (5)	20.8 (50)
1410	5 (2)	4.9 (20)
1490 ^(a)	2 (1)	2 (1)
1540 ^(a)	2 (1)	2 (1)
1590	2 (1)	2 (1)
1650 ^(a)	2 (1)	2 (1)
2010	7 (2)	6.9 (20)
2090 ^(a)	5 (2)	4.9 (20)
2270	3 (2)	3 (2)
2360	8 (3)	7.9 (30)
2430	9 (3)	8.9 (30)

(a) γ -ray not placed in level scheme as explained in Weinzierl (1964We06).

6 References

- 1931C01 M. Curie, A. Debierne, A. S. Eve, H. Geiger, O. Hahn, S. C. Lind, St. Meyer, E. Rutherford, E. Schweidler, Rev. Mod. Phys. 3(1931)427
[Half-life]
- 1957Ko42 A. V. Kogan, L. I. Rusinov, Sov. Phys. JETP 5(1957)365
[Half-life, neutron emission probability]
- 1961St20 G. Stetter, TID – 14880 (1961)
[Neutron emission probability]
- 1964We06 P. Weinzierl, E. Ujlaki, G. Preinreich, G. Eder, Phys. Rev. 134(1964)B257
[Half-life, E_{β} , I_{β} , E_{γ} , I_{γ}]
- 1981Ha54 B. Harmatz, Nucl. Data Sheets 34(1981)735
[Spin, parity, energy level, I_{β} , I_{γ}]
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
2002Ba85 – I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr., P. O. Tikkanen, S. Raman,
At. Data Nucl. Data Tables 81(2002)1
[Theoretical ICC]
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129
[Q]
- 2003Br13 E. Browne, Nucl. Data Sheets 99(2003)483
[Spin, parity, energy level, I_{β} , I_{γ}]

**²¹⁰Pb - Comments on evaluation of decay data
by V. Chisté and M. M. Bé**

This evaluation was completed in 2007. Literature available by October 2007 has been included.

1 Decay Scheme

²¹⁰Pb disintegrates by beta minus emission to an excited level and to the ground state level of ²¹⁰Pb. A weak alpha transition to the ²⁰⁶Hg ground state has been observed (1.9 (4) 10⁻⁶ %). Spins and parities are from the ENSDF mass-chain evaluations by E. Browne (2003Br13 for A = 210) and R. G. Helmer (1990He18 for A = 206).

The good agreement found between the adopted Q(β⁻) value of Audi and the effective Q(β⁻) value of 63.9 (11) keV calculated from decay scheme data indicates the completeness and correctness of the decay scheme.

2 Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2003Au03).

Experimental ²¹⁰Pb half-life values (in years) are given in Table 1:

Table 1: Experimental values of ²¹⁰Pb half-life.

Reference	Experimental value (a)	Comments
G. N. Antonoff (1910An**)	16.5	Not used. No uncertainty. ZnS counting.
I. Curie (1929Cu**)	23	Not used. No uncertainty. α counting.
M. Curie (1931Cu01)	19.5	Not used. No uncertainty.
F. Wagner (1950Wa**)	25.4 (15)	Ion Chamber.
R. J. Toboilem (1955To14)	19.40 (35)	Ion Chamber.
W. F. Merritt (1957Me47)	22.4 (4)	4π proportional counter.
G. Harbottle (1959Ha20)	20.4 (3)	Ion Chamber.
B. D. Pate (1959Pa03)	23.3 (5)	4π proportional counter.
W. R. Eckelmann (1960Ec01)	21.4 (5)	Geological.
L. Imre (1963Im02)	22.85 (70)	β counting.
H. Ramthun (1964Ra12)	21.96 (51)	Calorimetry.
H. R. von Gunten (1967Vo04)	22.2 (10)	Proportional counter.
A. Höndorf (1969Ho06)	22.26 (11)	α spectrometry.
G. A. Rech (2002Re18)	21.8 (3)	γ spectrometry.
Adopted value	22.23 (12)	χ ² = 1.53

The weighted average has been calculated using LWEIGHT computer program (version 3).

The evaluators have chosen to take into account the eleven experimental values with reported uncertainties found in the literature and given in Table 1. The values of Wagner (1950Wa**), Toboilem (1955To14) and Harbottle (1959Ha20) are rejected by the LWEIGHT program, because they are outliers, based on the Chauvenet's criterion. The largest contribution (71 %) to the weighted average comes from the value of Höndorf (1969Ho06).

The adopted value of ²¹⁰Pb half-life is a weighted average of **22.23 a** and the external uncertainty of **0.12 a**. The reduced-χ² value is 1.53.

2.1 a Transitions and Emissions

The transition energy of the α -particles group to the ground of ²⁰⁶Hg given in Section 2.1 is from Q_α (2003Au03).

For the probability of the α transition to the ground state of ²⁰⁶Hg, the available published data are given in Table 2.

Table 2: Experimental and adopted values of the α transition probability to the ground state of ²⁰⁶Hg.

Reference	Experimental value (10^{-6} %)	Comments
M. Nurmia (1961Nu01)	1.8 (5)	Superseded by 1962Ka27
P. Kauranen (1962Ka27)	1.7 (3)	
G. K. Wolf (1964Wo05)	2.7 (6)	
Adopted value	1.9 (4)	$\chi^2 = 2.22$

The adopted value of α transition to the ground state of ²⁰⁶Hg is the weighted average, calculated using LWEIGHT computer program, of **$1.9 \cdot 10^{-6}$ %** with the external uncertainty of **$0.4 \cdot 10^{-6}$ %**. The reduced- χ^2 value is 2.22.

2.2 b⁻ Transitions and Emissions

The end-point energies of the β^- transitions in the decay of ²¹⁰Pb \rightarrow ²¹⁰Bi have been obtained from the Q_{β^-} (2003Au03) value and the level energies of R. G. Helmer (1990He18), given in Table 3.

Table 3: ²¹⁰Bi level populated in the decay of ²¹⁰Pb.

Level Number	Level energy, (keV)	Spin and parity.
0	0	1^-
1	46.539 (1)	0^-

For these two levels, the adopted β^- transition probabilities and the associated uncertainties were deduced from the γ transition probability balance at each level of the decay scheme, taking into account, also the α transition probability to the ground state of ²⁰⁶Hg. In the table 4, our adopted values of β^- transitions probabilities are compared with the experimental results found in the literature: C. S. Wu (1953Wu28), J. Tousset (1957To16 and 1958To10), W. Stanners (1956St99) and I. M. Rogachev (1963Ro31). Except to C. S. Wu (1953Wu28), a fair agreement has been found, within the uncertainty limits, between the experimental results and the recommended values for the 17-keV and 63.5-keV β^- transitions.

Table 4: Adopted and experimental values of β^- transition probabilities.

	17-keV β^- transition	63.5-keV β^- transition
C. S. Wu (1953Wu28)	92 (5) %	8 (5) %
J. Tousset (1957To16)		19 (4) %
J. Tousset (1958To10)	81 (14) %	19 (4) %
W. Stanners (1956St99)	84.5 (35) %	15.5 (35) %
I. M. Rogachev (1963Ro31)		≤ 19 (2) %
Adopted value	80.2 (13) %	19.8 (13) %

The values of $\lg ft$ and average β^- energies have been calculated with the program LOGFT for the β^- forbidden transitions.

2.3 g Transitions

The 46.5-keV γ -ray transition probability was calculated using the γ -ray emission intensity (see **5.2 g Emissions**) and the relevant internal conversion coefficient. Multipolarity of this γ -ray transition is M1 (from E. Browne (2003Br13)).

The internal conversion coefficients (ICC) and their associated uncertainties for 46.5-keV γ -ray transition have been calculated using the BrIcc computer program (calculation for 'hole'), which interpolated from theoretical values of I. M. Band (2002Ba85). The α_T value is then 17.86 (25) compared to the previous value of 19.0 (6) from Rösels tables.

3 Atomic Data

Atomic values, ω_K , ω_L and n_{KL} and the X-ray relative probabilities are from Schönfeld and Jaßén (1996Sc06).

4 Electron Emissions

The conversion electrons emission probabilities have been deduced using the γ -ray emission intensities and ICC's. The calculated total conversion electrons intensity of 75.2 (10) % is in fair agreement with the measured value of 81 (4) % from W. Stanners (1956St99).

5 Photon Emissions

5.1 X-ray Emissions

The X-ray absolute intensities have been calculated from γ -ray data and ICC using the EMISSION computer program and compared in Table 5 with the measured values found in the literature. For L_I , L_α and L_η x-rays, a good agreement was found between the experimental results given by 1987Me17 and 1990Sc08 and the recommended values deduced from decay scheme balance.

Table 5: Experimental and recommended (calculated) values of L X-ray absolute intensities.

	R. W. Fink (1957Fi06)	R. J. Gehrke (1971Ge11)	D. Metha (1987Me17) ^a	U. Schötzig (1990Sc08)	Recommended Values
L_I			0.584 (18)	0.55 (3)	0.552 (17)
L_α			10.27 (32)	9.48 (17)	10.3 (3)
L_η			0.074 (4)	0.075 (4)	0.075 (2)
L_β			11.6 (4)	10.9 (4)	9.05 (13)
L_γ			2.64 (8)	2.36 (5)	1.97 (3)
L total	23.8 (20)	22.8 (15)	25.2 (3)	23.4 (4)	22.0 (5)

^a Normalized with $I_\gamma(46.5\text{-keV}) = 4.252(40)\%$ (see 5.2 γ Emissions.)

5.2 γ Emissions

The energy of the γ -ray emission given in Section 5 is from R. G. Helmer (1981He15 and 2000He14).

For the 46.5-keV γ -ray from ²¹⁰Bi, the experimental data set of absolute γ -ray emission intensity and adopted value in this evaluation are given in Table 6.

Table 6: The experimental data set of the relative γ -ray emission intensity.

Reference	Experimental values (%)	Comments
D. K. Butt (1951Bu37)	3.5 (4)	Not used by the evaluators.
C. S. Wu (1953Wu28)	2.8 (6)	Not used by the evaluators.
P. E. Damon (1954Da23)	3.8 (6)	Not used by the evaluators.
R. W. Fink (1957Fi06)	4.5 (4)	
I. Y. Krause (1958Kr71)	4.05 (8)	Not used by the evaluators.
K. Ya. Gromov (1969Gr33)	4.8 (6)	
K. Debertin (1983De11)	4.18 (9)	Superseded by 1990Sc08.
Y. Hino (1990Hi03)	4.26 (7)	
U. Schötzig (1990Sc08)	4.24 (5)	
Adopted value	4.252 (40)	$\chi^2 = 0.42$

The sets of values from D. K. Butt (1951Bu37), C. S. Wu (1953Wu28) and P. E. Damon (1954Da23) were omitted from analysis due to discrepancy with the other data and a lack of information in the articles about experimental measurements carried out and, therefore on the results.

The original uncertainty given by I. Y. Krause (1958Kr71) (= 0.08) seems under-estimated for the measurement method (NaI spectrometry) then it was decided to omit this value from the analysis.

The adopted value for 46.5-keV γ -ray emission intensity is the weighted average, calculated using LWEIGHT computer program, of **4.252 %** with the internal uncertainty of **0.040 %**. The reduced- χ^2 value is 0.42.

The evaluated absolute 46.5-keV γ -ray emission and transition probabilities are given in Table 7.

Table 7: Recommended absolute 46.5-keV γ -ray emission and transition probabilities.

Energy (keV)	Absolute γ -ray emission probability (%)	Absolute γ -ray transition probability (%)
46.539 (1)	4.252 (40)	80.2 (13)

6 References

- 1929Cu** P. Curie, I. Curie, J. Phys. Radium 10(1929)385 [Half-life].
 1929Cu** I. Curie, J. Phys. Radium 10(1929)388 [Half-life].
 1931Cu01 M. Curie, A. Debierne, A. S. Eve, H. Geiger, O. Hahn, S. C. Lind, St. Meyer, E. Rutherford, E. Schweidler, Rev. Mod. Phys. 3(1931)427 [Half-life].
 1950Wa** F. Wagner, ANL – 4490(1950)5 [Half-life].
 1951Bu37 D.K. Butt, W. D. Brodie, Proc. Phys. Soc. (London) 64A(1951)791 [I_γ].
 1953Wu28 C.S. Wu, F. Boehm, E. Nagel, Phys. Rev. 91(1953)319 [I_γ].
 1954Da23 P. E. Damon, R. R. Edwards, Phys. Rev. 95(1954)1698 [I_γ].
 1955To11 J. Toboilem, J. Phys. Radium 16(1955)235 [Half-life].
 1956St99 W. Stanners, M.A.S. Ross, Proc. Phys. Soc. 69A(1956)836 [I_β].
 1957To16 J. Tousset, Compt. Rendu 245(1957)1617 [I_β].
 1957Me47 W. F. Merritt, P. J. Champion, R. C. Hawkings, Can. J. Phys. 35(1957)16 [Half-life].
 1957Fi06 R. W. Fink, Phys. Rev. 106(1957)266 [I_γ].
 1958Kr71 I.Y. Krause, Z. Phys. 152(1958)586 [I_γ].
 1958To10 J. Tousset, J. Phys. Radium 19(1958)39 [I_β].
 1959Pa03 B.D. Pate, D. C. Santry, L. Yaffe, Can. J. Phys. 37(1959)1000 [Half-life].
 1959Ha20 G. Harbottle, J. Inorg. Nucl. Chem. 12(1959)6 [Half-life].
 1960Ec01 W.R. Eckelmann, W. S. Broecker, J. L. Kulp, Phys. Rev. 118(1960)698 [Half-life].
 1961Nu01 M. Nurmi, P. Kauranen, M. Karras, A. Siivola, A. Isola, G. Graeffe, A. Lyyjynen, Nature 190(1961)427 [Alpha branching ratio].
 1962Ka27 P. Kauranen, Ann. Acad. Sci. Fennicae, Ser. A VI 96(1962)7 [Alpha branching ratio].
 1963Im02 L. Imre, G. Fabry, I. Dezi, Nucl. Sci. Abstr. 17(1963)4186, Abstr. 31442 [Half-life].
 1963Ro31 I.M. Rogachev, Nucl. Sci. Abstr. 18(1964)2284, abstr. 17094 [I_β].
 1964Ra12 H. Ramthun, Z. Naturforsch. 19a(1964)1064 [Half-life].
 1964Wo05 G.K. Wolf, F. Lux, H. J. Born, Radiochim. Acta 3(1964)206 [Alpha branching ratio].
 1967Vo04 H.R. von Gunten, A. Wyttenbach, H. Dulakas, J. InorgNucl. Chem. 29(1967)2826 [Half-life].
 1969Ho06 A. Höhdorf, Z. Naturforsch. 24a(1969)612 [Half-life].
 1969Gr33 K. Ya. Gromov, B. M. Sabirov, J. J. Urbanets, Bull. Acad. Sci. USSR, Phys. Ser. 33(1970)1510, [II]
 1971Ge11 R.J. Gehrke, R.A. Lokken, Nucl. Instrum. Meth. 97(1971)219 [L x-rays].
 1981He15 R.G. Helmer, A.J. Caffrey, R.J. Gehrke, R.C. Greenwood, Nucl. Instrum. Meth. 188(1981)671 [I_γ].
 1983De11 K. Debertin, W. Pessara, Int. J. Appl. Radiat. Isotop. 34(1983)515 [I_γ].
 1987Me17 D. Metha, B. Chand, S. Singh, M.L. Garg, N. Singh, T.S. Chema, P.N. Trehan, Nucl. Instrum. Meth. Phys. Res. A260(1987)157 [L x-rays].

Comments on evaluation

- 1990He18 R.G. Helmer, M. A. Lee, Nucl. Data Sheets 61(1990)93 [Spin, parity, level energy].
1990Hi03 Y. Hino, Y. Kamada, Nucl. Instrum. Meth. Phys. Res. A286(1990)543 [I_γ].
1990Sc08 U. Schötzig, Nucl. Instrum. Meth. Phys. Res. A286(1990)523 [I_γ].
1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
1999Br39 E. Browne, Nucl. Data Sheets 88(1999)29 [Spin, parity, level energy].
2000He14 R.G. Helmer, C. van der Leun, Nucl. Instrum. Meth. Phys. Res. A450(2000)35 [I_γ].
2002Ba85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman,
Atomic Data Nucl. Data Tables 81(2002)1 [Theoretical ICC].
2002Re18 G.A. Rech, E. Browne, I. D. Goldman, F. J. Schima, E. B. Norman, Phys. Rev.
C65(2002)057302 [Half-life].
2003Au03 G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].
2003Br13 E. Browne, Nucl. Data Sheets 99(2003)483 [Spin, parity, level energy].

**²¹⁰Bi - Comments on evaluation of decay data
by V. Chisté and M. M. Bé**

This evaluation was completed in 2008. Literature available by January 2008 was included.

1 Decay Scheme

²¹⁰Bi disintegrates by beta minus emission to the ground state level of ²¹⁰Po. Weak alpha transitions to excited levels of ²⁰⁶Tl have been observed (1.40 (15) 10⁻⁴ %). Spins and parities are from the ENSDF mass-chain evaluations E. Browne (2003Br13 for A = 210). For ²⁰⁶Tl, spins and parities are from L. I. Rusinov measurements (1961Ru02).

2 Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2003Au03).

Experimental ²¹⁰Bi half-life values (in days) are given in Table 1:

Table 1: Experimental values of ²¹⁰Bi half-life.

Reference	Experimental value (d)	Comments
A. Pompéi (1935Po01)	5.02 (1)	Ionization chamber.
N. Hole (1944Ho**)	5.15 (10)	GM counter.
F. Begemann (1952Be22)	5.02 (2)	GM counter.
E. E. Lockett (1953Lo09)	4.989 (13)	Ionization chamber.
J. Robert (1956Ro18)	5.013 (5)	Ionization chamber. Superseded by 1959Ro51
J. Robert (1959Ro51)	5.013 (5)	Ionization chamber.
Recommended value	5.012 (5)	$\chi^2 = 1.32$

The weighted average has been calculated using the LWEIGHT computer program (version 3).

The evaluators have chosen to use just five experimental values with uncertainties given in Table 1. The value of Hole (1944Ho**) has been rejected by the LWEIGHT program because it is an outlier, based on the Chauvenet's criterion. With this data set, the largest contribution to the weighted average comes from the value of Robert (1959Ro51) amounting to 68 % of the total statistical weight.

The recommended value of ²¹⁰Bi half-life is the weighted average of **5.012 d** with an external uncertainty of **0.005 d**. The reduced- χ^2 value is 1.32.

2.1 a Transitions and Emissions

The recommended values of emission energies of the α -particles are given by A. Rytz (1991Ry01).

Table 2: Experimental values of emission energies of the α -particles.

Reference	$\alpha_{0,1}$ (keV)	$\alpha_{0,2}$ (keV)	Comments
R. J. Walen (1960Wa14)	4686 (2)	4649 (2)	Uncertainty given by Rytz.
P. Kauranen (1962Ka27)	4700	4660	Not used: no uncertainty.
R. C. Lange (1969La18)	4697 (5)	4660 (5)	Uncertainty given by Rytz.
Recommended value (1991Ry01)	4687 (4)	4650 (4)	$\chi^2 = 4.2$. External uncertainty.

Several experimental values of the α branching to ²⁰⁶Tl are given in Table 3.

Table 3: Experimental and recommended values of total α branching for $^{210}\text{Bi} \rightarrow ^{206}\text{Tl}$.

Reference	Experimental value (10^{-4} %)	Comments
E. Broda (1947Br36)	0.5	Not used: no uncertainty.
R. J. Walen(1959Wa05)	1.25	Not used: no uncertainty.
R. W. Fink (1956Fi09)	1.7 (2)	
M. Nurmia (1961Nu01)	1.9 (4)	Superseded by 1962Ka27
P. Kauranen (1962Ka27)	1.32 (10)	
Recommended value	1.40 (15)	$\chi^2 = 2.9$

The weighted average has been calculated using the LWEIGHT computer program (version 3).

The value given by M. Nurmia (1961Nu01) is from the same laboratory as 1962Ka27, thus, it was not included in the averaging procedure. Then, the recommended alpha transition branching is the average of the values given by R. W. Fink (1956Fi09) and P. Kauranen (1962Ka27).

The recommended value of α transitions to the excited levels of ^{206}Tl is the weighted average of **$1.40 \cdot 10^{-4}$ %** with an external uncertainty of **$0.15 \cdot 10^{-4}$ %**. The reduced- χ^2 value is 2.9.

The individual α particle probabilities to the 265 -keV and 304-keV levels are (1959Wa05, 1960 Wa14) $0.56 (6) \cdot 10^{-4}$ % and $0.84 (9) \cdot 10^{-4}$ %, respectively.

2.2 β^- Transitions and Emissions

The end-point energy of the β^- transition in the decay of $^{210}\text{Bi} \rightarrow ^{210}\text{Po}$ is from the Q_{β^-} (2003Au03). The recommended and experimental values are shown in Table 4.

Table 4: Experimental and recommended values of the end-point energy of the β^- transition.

Reference	E_{β^-} (keV)
A. Flammersfeld (1939F102)	1170
G. J. Neary (1940Ne04)	1170
E. A. Plassmann (1954Pl30)	1155 (5)
H. Daniel (1962Da03)	1160.5 (5)
S. T. Hsue(1967Hs01)	1161.5 (15)
D. Flothmann (1969F102)	1153
Recommended value (2003Au03)	1162.1 (8)

For the $\beta_{0,0}$ transition probability and associated uncertainty, the following relation was applied:

$$P_{\beta_{0,0}} = 100 \% - P_{\alpha},$$

where $P_{\alpha} = 1.40 (15) \cdot 10^{-4}$ % (see 2.2 α Transitions and Emissions). Then: $P_{\beta_{0,0}} = 99.99986 (2) \%$.

The $\lg ft$ value and the average β^- energy have been calculated with the program LOGFT for a 1st forbidden transition.

2.3 γ Transitions and Emissions

Multipolarity of γ -ray transitions are from L. I. Rusinov (1961Ru02):

265-keV γ -ray: E2

304-keV γ -ray: M1

The γ -ray transition probabilities following the α -decay of $^{210}\text{Bi} \rightarrow ^{206}\text{Tl}$ were deduced from the decayscheme balance using the recommended α -particle intensity values given in section 2.1 α Transitions and Emissions, shown in Table 5.

Table 5: Adopted values of α transition and γ -ray emission probabilities.

γ -ray energy (keV)*	α probability (%)	γ -ray absolute transition probability (%)	γ -ray absolute emission probability (%)
265.832 (5)	0.000 056 (6)	0.000 056 (6)	0.000 048 (5)
304.896 (6)	0.000 084 (9)	0.000 084 (9)	0.000 061 (7)

*From 1999Br39

The γ -ray emission intensities were obtained using the γ -ray transition probabilities (given in Table 6) and the relevant internal conversion coefficients, calculated using the BrIcc computer code (calculation for 'hole'), which interpolated from theoretical values of I. M. Band (2002Ba85).

3 Atomic Data

Atomic values, ω_K , ω_L and n_{KL} are from Schönfeld and Janßen (1996Sc06).

4 References

- 1935Po01 – A. Pompéi, J. Phys. Radium 6(1935)471 [Half-life].
 1939F102 – A. Flammersfeld, Z. Phys. 112(1939)727 [E_{β^-}].
 1940Ne04 – G. J. Neary, Proc. Roy. Soc. (London) 175A(1940)71 [E_{β^-}].
 1944Ho** – N. Hole, Ark. Mat., Astron. Fys. 31B(1944)1 [Half-life].
 1947Br36 – E. Broda, N. Feather, Proc. Roy. Soc. (London) 191A(1947)20 [α - branching].
 1952Be22 – F. Begemann, F. G. Houtermans, Z. Naturforsch. 7a(1952)143 [Half-life].
 1953Lo09 – E. E. Lockett, R. H. Thomas, Nucleonics 11(1953)14 [Half-life].
 1954P130 – E. A. Plassmann, L. M. Langer, Phys. Rev. 96(1954)1593 [E_{β^-}].
 1956Ro18 – J. Robert, J. Tobaillem, J. Phys. Radium 17(1956)440 [Half-life].
 1956Fi09 – R. W. Fink, G. W. Warren, B. L. Robinson, R. R. Edwards, Bull. Am. Phys. Sod.(1956)171 [α -branching].
 1959Ro51 – J. Robert, Ann. Phys. (Paris) 4(1959)440 [Half-life].
 1959Wa05 – R. J. Walen, G. Bastin-Scoffier, J. Phys. Radium 20(1959)589 [α - branching].
 1960Wa14 – R. J. Walen, G. Bastin-Scoffier, Nucl. Phys. 16(1960)246 [α - branching, E_{α}].
 1961Nu01 – M. Nurmia, P. Kauranen, M. Karras, A. Siivola, A. Isola, G. Graeffe, A. Lyyjynen, Nature 190(1961)427 [α - branching].
 1961Ru02 – L. I. Rusinov, Yu. N. Andreev, S. V. Golenetskii, M. I. Kislov, Yu. I. Filimonov, Sov. Phys. JETP 13(1961)707 [Spin, parity and multipolarity].
 1962Da03 – H. Daniel, Nucl. Phys. 31(1962)293 [E_{β^-}].
 1962Ka27 – P. Kauranen, Ann. Acad. Sci. Fennicae, Ser. A VI, 96(1962) [α - branching, E_{α}].
 1967Hs01 – S. T. Hsue, M. U. Kim, S. M. Tang, Nucl. Phys. A94(1967)146 [E_{β^-}].
 1969F102 – D. Flothmann, W. Wiesner, R. Löhken, H. Rebel, Z. Phys. 225(1969)164 [E_{β^-}].
 1969La18 – R. C. Lange, G. R. Hagee, A. R. Campbell, Nucl. Phys. A133(1969)273 [E_{α}].
 1991Ry01 – A. Rytz, At. Data Nucl. Data Tables 47(1991)205 [E_{α}].
 1996Sc06 – E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
 1999Br39 – E. Browne, Nucl. Data Sheets 88(1999)29 [level energy].
 2002Ba85 – I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr., P. O. Tikkanen, S. Raman, Atomic Data Nucl. Data Tables 81(2002)1 [Theoretical ICC].
 2003Au03 – G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].
 2003Br13 – E. Browne, Nucl. Data Sheets 99(2003)483 [Spin, parity, level energy].

**²¹⁰Po - Comments on evaluation of decay data
by V. Chisté and M. M. Bé**

This evaluation was completed in 2008. Literature available by February 2008 was included.

1 Decay Scheme

²¹⁰Po disintegrates by alpha emission to the 803 -keV excited level and ground state level of ²⁰⁶Pb. Energy levels, spins and parities are from the ENSDF mass -chain evaluations R.G. Helmer (1990He18) and E. Browne (1999Br39).

2 Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2003Au03).

Experimental ²¹⁰Po half-life values (in days) are given in Table 1:

Table 1: Experimental values of ²¹⁰Po half-life.

Reference	Experimental value (d)	Comments
E. V. Schweidler (1912Sc**)	136.5	Not used: no uncertainty.
M. Curie (1920Cu**)	140.0	Not used: no uncertainty.
A. Dorabialska (1931Do**)	137.6 (6)	Calorimetry.
A. S. Sanielevici (1936Sa**)	139.6 (14)	Calorimetry.
W. H. Beamer (1949Be54)	138.30 (14)	Calorimetry.
D. C. Ginnings (1953Gi10)	138.39 (14)	Calorimetry.
M. L. Curtis (1953Cu46)	138.374 (32)	α counting.
J. F. Eichelberger (1954Ei20)	138.400 (6)	Calorimetry. Not used. Superseded by 1964EiZZ.
J. F. Eichelberger (1964EiZZ)	138.3763 (17)	Calorimetry
Recommended value	138.3763 (17)	χ ² = 0.10

The weighted average has been calculated using LWEIGHT computer program (version 3).

The evaluators have chosen to use six experimental values with uncertainty found in the literature and given in Table 1. The values of A. Dorabialska (1931Do**) and A. S. Sanielevici (1936Sa**) have been rejected by the LWEIGHT program, they are statistical outliers, based on the Chauvenet's criterion. With this data set, the largest contribution (99 %) to weighted average comes from the value of J. F. Eichelberger (1964EiZZ).

The recommended value of ²¹⁰Po half-life is the weighted average of **138.3763 d** with an internal uncertainty of **0.0017 d**. The reduced-χ² value is 0.10.

2.1 a Transitions and Emissions

The recommended value of α_{0,0} emission energy is given by A. Rytz (1991Ry01), based on a measurement by D. J. Gorman (1973Go39). The experimental and recommended values of α_{0,0} emission energy are shown in Table 2.

Table 2: Experimental and recommended (calculated) values of $\alpha_{0,0}$ emission energy.

Reference	$\alpha_{0,0}$ emission energy (keV)	Comments
S. Rosenblum (1933Ro03)	5298 (6)	
W. B. Lewis (1934Le01)	5298 (21)	
E. R. Collins (1953Co64)	5304.3 (29)	
G. H. Briggs (1954Br07)	5300.6 (26)	Evaluated value reported by author.
I. I. Agapkin (1957Ag15)	5297.8 (15)	
F. A. White (1958Wh09)	5305.4 (10)	
C. P. Browne (1960Br20)	5308.6 (30)	
E. H. Beckner (1961Be13)	5302.5 (15)	
A. Rytz (1961Ry05)	5304.9 (6)	
D. J. Gorman (1973Go39)	5304.51 (7)	
Recommended value (1991Ry01)	5304.33 (7)	

For $\alpha_{0,1}$, the emission energy has been obtained from $Q_{\alpha}(2003Au03) = 5407.46 (7) \text{ keV}$ and the level energy given in Table 3 from R. G. Helmer (1990He18).

Table 3: ²⁰⁶Pb excited level populated in the decay of ²¹⁰Po.

Level Number	Level energy, (keV)	Spin and parity.
1	803.10 (5)	2 ⁺

The emission intensities of the α -particles have been deduced from the $P(\gamma + ce)$ decay scheme balance at each level and shown in Table 4.

Table 4: Emission intensities of the α -particles.

α emission energy (keV)	Emission Intensities (%)
4516.66 (9)	0.00124 (4)
5304.33 (7)	99.99876 (4)

The ratio $I_{\alpha}(4516)/I_{\alpha}(5304)$, with the recommended values (Table 4), is $1.24 (4) 10^{-5}$, which can be compared with the measured value of $1.07 (2) 10^{-5}$ (1958Ba45).

2.2 g Transitions

The transition probability was calculated using the experimental 803 -keV γ -ray emission intensity and the relevant internal conversion coefficient (see **4.2 g Emissions**).

Multipolarity of the 803-keV γ -ray transition (E2) is given by S. de Benedetti (1952De08).

The internal conversion coefficient (ICC) for the the 803 -keV γ -ray transition has been interpolated from theoretical values of I. M. Band (2002Ba85) using the BRICC computer program (calculation for 'hole').

3 Atomic Data

Atomic values, ω_K , ω_L and n_{KL} and the X-ray relative probabilities are from Schönfeld and Jaßßen (1996Sc06).

4 Photon Emissions

4.1 X-rays

The X-ray absolute intensities have been calculated from γ -ray data and ICC using the EMISSION computer program.

4.2 g Emissions

The energies of the γ -ray emission given in section 5.2 is from R. G. Helmer (1990He18).

For the 803-keV γ -ray, the experimental data set of γ -ray emission intensity is given in Table 5.

Table 5: The experimental data set of the γ -ray emission intensity.

Reference	Experimental values (10^{-3} %)	Comments
M. A. Grace (1951Gr15)	1.80 (14)	
M. Riou (1952Ri04)	1.6 (2)	
W. C. Barber (1952Ba20)	1.5 (4)	
O. Rojo (1955Ro30)	1.20 (12)	
R. W. Hayward (1955Ha09)	1.21 (6)	
A. Ascoli (1956As46)	1.21 (8)	
N. S. Shimanskaia (1956Sh24)	1.2 (2)	
V. V. Ovechkin (1957Ov09)	1.22 (9)	
Recommended value	1.23 (4)	$\chi^2 = 0.69$

The weighted average has been calculated using LWEIGHT computer program (version 3).

The evaluators have used the eight experimental values given with uncertainties in the literature and shown in Table 5. The value of M.A. Grace (1951Gr15) has been rejected by the LWEIGHT program, as statistical outlier, based on the Chauvenet's criterion. In the data set of seven values, the largest contribution (41%) to the weighted average comes from the value of R. W. Hayward (1955Ha09).

The recommended value of the relative γ -ray emission intensity is the weighted average of **1.23 10^{-3} %** with the internal uncertainty of **0.04 10^{-3} %**, and a reduced- χ^2 value of 0.69.

5 References

- 1912Sc** – E. V. Schweidler, Verh. Deutsch Phys. Ges. 14(1912)539 [Half-life].
 1920Cu** – M. Curie, J. Phys. Radium 1(1920)12 [Half-life].
 1931Do** – A. Dorabialska, Roczniki Chem. (Poland) 11(1931)475 [Half-life].
 1933Ro03 – S. Rosenblum, G. Dupouy, J. Phys. Radium 4(1933)262 [E_α].
 1934Le01 – W. B. Lewis, B. V. Bowden, Proc. Roy. Soc. (London) A145(1934)235 [E_α].
 1936Sa** – A. S. Sanielevici, J. Chim. Phys. 33(1936)759 [Half-life].
 1949Be54 – W. H. Beamer, C. R. Maxwell, J. Chem. Phys. 17(1949)1293 [Half-life].
 1951Gr15 – M. A. Grace, R. A. Allen, D. West, H. Halban, Proc. Roy. Soc. (London) A64(1951)493 [I_γ].
 1952Ba20 – W. C. Barber, R. H. Helm, Phys. Rev. 86(1952)275 [I_γ].
 1952Ri14 – M. Riou, J. Phys. Radium 13(1952)244 [I_γ].
 1952De08 – S. de Benedetti, G. H. Minton, Phys. Rev. 85(1952)944 [Multipolarity].
 1953Gi10 – D. C. Ginnings, A. F. Ball, D. T. Vier, J. Res. NBS 50(1953)75 [Half-life].
 1953Cu46 – M. L. Curtis, Phys. Rev. 92(1953)1489 [Half-life].
 1953Co64 – E. R. Collins, C. D. McKenzie, C. A. Ramm, Proc. Roy. Soc. (London) 216A(1953)219 [E_α].
 1954Br07 – G. H. Briggs, Rev. Mod. Phys. 26(1954)1 [E_α].
 1954Ei20 – J. F. Eichelberger, K. C. Jordan, S. R. Orr, J. R. Parks, Phys. Rev. 96(1954)719 [Half-life].
 1955Ha09 – R. W. Hayward, D. D. Hoppes, W. B. Mann, J. Res. Nat. Bur. Stand. 54(1955)47 [I_γ].
 1955Ro30 – O. Rojo, M. A. Hakeem, M. Goodrich, Phys. Rev. 99(1955)1629 [I_γ].
 1956As46 – A. Ascoli, M. Asdente, E. Germagnoli, Nuovo Cim. 4(1956)946 [I_γ].
 1956Sh24 – N. S. Shimanskaia, Soviet Phys. JETP 4(1957)165 [I_γ].
 1957Ag15 – I. I. Agapkin, L. L. Gol'din, Bull. Acad. Sci. USSR 21(1958)911 [E_α].
 1957Ov09 – V. V. Ovechkin, Bull. Acad. Sci. USSR 21(1958)1627 [I_γ].
 1958Ba45 – G. Bastin -Scoffier, R. J. Walen, Compt. Rend. Acad. Sci. (Paris) 247(1958)2333 [$I_\alpha(4516)/I_\alpha(5304)$].

- 1958Wh09 – F. A. White, F. M. Rourke, J. C. Sheffield, R. P. Schuman, J. R. Huizenga, Phys. Rev. 109(1958)437 [E_{α}].
- 1960Br20 – C. P. Browne, J. A. Galey, J. R. Erskine, K. L. Warsh, Phys. Rev. 120(1960)905 [E_{α}].
- 1961Ry05 – A. Rytz, H. H. Staub, H. Winkler, Helv. Phys. Acta 34(1961)960 [E_{α}].
- 1961Be13 – E. H. Beckner, R. L. Bramblett, G. C. Phillips, T. A. Eastwood, Phys. Rev. 123(1961)2100 [E_{α}].
- 1962Br22 – C. P. Browne, Phys. Rev. 126(1962)1139 [E_{α}].
- 1964EiZZ – J. F. Eichelberger, G. R. Grove, L. V. Jones, MLM – 1209(1964)11 [Half-life].
- 1973Go39 – D. J. Gorman, A. Rytz, Compt. Rend. Acad. Sci. (Paris), Ser. B277(1973)29 [E_{α}].
- 1990He18 – R. G. Helmer, Nucl. Data Sheets 61(1990)93 [Energy level, spin, parity].
- 1991Ry01 – A. Rytz, At. Data Nucl. Data Tables 47(1991)205 [E_{α}].
- 1996Sc06 – E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
- 1999Br39 – E. Browne, Nucl. Data Sheets 88(1999)29 [Spin, parity].
- 2002Ba85 – I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr., P. O. Tikkanen, S. Raman, Atomic Data Nucl. Data Tables 81(2002)1 [Theoretical ICC].
- 2003Au03 – G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].

²¹¹Pb - Comments on evaluation of Decay Data

By F.G. Kondev

This evaluation was completed in March 2011, with a literature cut off by the same date, as a part of ANL commitment to the IAEA-CRP on “Updated Decay Data Library for Actinides”.

1. Decay Scheme

The nuclide ²¹¹Pb ($J^\pi=9/2^+$) disintegrates 100 % by β^- emissions. The strongest β^- -decay branch of 91.28 (12) % populates the $J^\pi=9/2^-$ ground state of the daughter nuclide ²¹¹Bi. The level schemes of ²¹¹Pb and ²¹¹Bi, including J^π values, are based on the ENSDF evaluation of Browne (2004Br45).

2. Nuclear Data

Adopted $Q(\beta^-)$ value of 1367 (6) keV is taken from the evaluation of Audi *et al.* (2003Au03).

The experimental data for the half-life of the ²¹¹Pb ground state are very scarce. The value of 36.1 (2) min that is included in ENSDF (2004Br45) originates from the work of Sargent (1939Sa11) and Nurmia (1965Nu03). This value is adopted in the present evaluation, but new measurements are certainly required to confirm this value.

2.1. β^- Transitions

The values for the maximum β^- -decay energies ($E_{\beta^-,max}$, presented in Table 1) were derived from $Q(\beta^-) = 1367$ (6) keV (2003Au03) and the level energies deduced in the present evaluation, as detailed in section 2.2. The β^- -decay transition probabilities (P_β) were deduced from the decay scheme and the corresponding absolute γ -ray transition probabilities. Log ft values were calculated using the *LOGFT* program from the ENSDF evaluation package, based on the work of Gove and Martin (1971Go40).

2.2. Gamma Transitions and Electron Internal Conversion Coefficients

The γ -ray transition energy data are presented in Table 2. Statistical analysis using the *LWEIGHT* program has been performed, and the resulting gamma-ray energies are listed in column 13 of Table 2. With those energies, the level scheme was fitted using the *GTOL* program from the ENSDF analysis package, and new level energies were obtained (shown in Table 1). Then adopted gamma-ray energies were determined from the corresponding level energies.

The γ -ray transition multiplicities and mixing ratios were taken from the ENSDF evaluation of Browne (2004Br45). The electron conversion coefficients were calculated using the *BrIcc* code (2008Ki07).

Table 1. Level energies, quantum numbers, $E_{\beta_{0,1,max}}$, P_β and log ft values in the β^- decay of ²¹¹Pb ($J^\pi=9/2^+$).

Level energy (keV)	J^π	$E_{\beta^-,max}$ (keV)	P_β (%)	Nature	log ft
1270.75 (6)	(7/2,9/2,11/ 2)-	96 (6)	0.0172 (15)	first forbidden non- unique	5.93
1234.3 (4)		133 (6)	0.0009 (3)	-	-
1196.33 (5)		171 (6)	0.019 (4)	-	-
1109.509 (23)	9/2-	257 (6)	1.06 (4)	first forbidden non- unique	5.58
1103.52 (20)		263 (6)	0.0047 (7)	-	-
1080.64 (4)		286 (6)	0.0570 (24)	-	-
1014.38 (4)	(7/2,9/2,11/ 2)-	-	-	-	-
831.984 (12)	9/2-	535 (6)	6.32 (9)	first forbidden non- unique	5.73

Level energy (keV)	J ^π	E _{β- max} (keV)	P _β (%)	Nature	log ft
766.680 (13)	(9/2,11/2)-	600 (6)	< 0.09	first forbidden non-unique	> 7.7
404.834 (9)	7/2-	962 (6)	1.57 (9)	first forbidden non-unique	7.21
0.0	9/2-	1367 (6)	91.28 (12)	first forbidden non-unique	5.99

The gamma-ray emission probability data are presented in Table 3. The reported values were determined relative to $I_\gamma(351.07_\gamma) = 100\%$, where the 351.07 keV transition depopulates the first $3/2^+$ level of the ²⁰⁷Tl nuclide fed in the α decay of ²¹¹Bi. The statistical analysis using the *LWEIGHT* program has been performed and *deduced* intensities were obtained (column 14 of Table 3). Using the absolute emission probability of $I_\gamma(351.07_\gamma) = 13.06 (12)\%$ (2011Ko04) and $\% \alpha(^{211}\text{Bi}) = 99.724 (4)\%$ (2004Br45), a normalization factor of 0.1302 (12) was obtained. This value was used to determine the *adopted* gamma-ray emission probabilities, which are shown in the last column of Table 3.

A number of weak transitions, summarized in Table 4, have been assigned to the β^- decay of ²¹¹Pb by 1988Hi14 (five gamma rays), 1971Da34 (nine gamma rays), 1968Ha21 (three gamma rays) and 1968Br17 (one gamma ray). However, the experimental information presented in those articles is insufficient to assign these gamma rays unambiguously to the decay of ²¹¹Pb, and hence they were not placed in the proposed decay scheme. None of the above publications reported the same unplaced gamma rays, which facilitated the conclusion made in this evaluation to exclude them from the proposed decay scheme. Further work, including gamma-ray coincidence studies, is merited to obtain a more complete decay scheme for ²¹¹Pb.

3. Atomic Data

The atomic data (fluorescence yields, X-ray energies and relative probabilities, and Auger electrons energies and relative probabilities) were provided by the SAISINUC software (2008DuZX). Details regarding the origin of these data can be found in 1996Sc06, 1998ScZM, 1999ScZX, 2000Sc47 and 2003De44.

4. Data Consistency

The adopted Q_β-value of 1367 (6) keV (2003Au03) has subsequently been compared with the Q-value calculated by summing the contributions of the individual emissions to the ²¹¹Pb beta-decay process (i.e. β^- , conversion electrons, γ , etc.):

$$Q_{\beta\text{-(calc)}} = \sum (E_i \times P_i) = 1368 (6) \text{ keV}$$

Percentage deviation from the Q-value of Audi *et al.* is 0.0731 (5) %, which supports the derivation of a highly consistent decay scheme.

Table 2. Measured, deduced and adopted gamma-ray energies in β^- -decay of ²¹¹Pb.

1988Hi14	1976B113	1971Da34	1968Br17	1968Go15	1968Ha21	1967Da20	1967Da10	1965Co06	1965Me07	1963Va05	1962Gi03	deduced, keV	adopted, keV
	65.420 (14)		65.5 (2)	65.4 (2)		65.7 (10)	65.5 (5)		65.502 (8) g)			65.420 (14)	65.304 (18)
		95.0 (2)		95.0 (6)		94.5 (5)						95.0 (2)	95.13 (5)
313.64 (12)	313.58 (9)	313.8 (2)	313.6 (3)	313.6 (5)		313.5 (10)	313 (1)	290 (10)	310 (3)			313.59 (9)	313.96 (4)
342.02 (12)	342.90 (4)*	342.7 (2)		342.7 (3)			342.5 (10)		340 (3)			342.91 (4)	342.829 (26)
	362.062 (17)*		362.9 (5) ?									362.072 (17)	361.846 (16)
404.89 (12)	404.843 (10)	404.84 (4)	404.8 (1)	404.8 (1)	405	404.8 (5)	404.7 (3)	400 (7)	404.84 (4) g)	404	405 (5)	404.853 (10)	404.834 (9)
427.14 (12)	427.078 (10)	426.99 (4)	427.0 (1)	426.9 (1)	427	427.1 (5)	427.0 (3)	430 (7)	426.99 (4) g)	426	425 (5)	427.088 (10)	427.150 (15)
				429.1 (5)								429.1 (5)	429.65 (6)
504.12 (12)		503.3 (4)		503.6 (7)								504.12 (12)	504.07 (6)
	609.33 (4)*	609.5 (2)	609.1 (2)	609.3 (5)		610 (2)			612 (5) f)		615 (5)	609.38 (4)	609.55 (4)
	676.65 (7)*		676.6 (3)	675.2 (3)				650 (10)				676.69 (7)	675.81 (4)
704.66 (12)	704.59 (3)	704.5 (1)	704.5 (2)	704.3 (2)	702	703.3 (8)	703.8 (3)	706 (7)	702 (3)	700	700 (5)	704.64 (3)	704.675 (25)
766.45 (12)	766.47 (3)	766.34 (7)	766.3 (2)	766.4 (1)	766	766.2 (8)	766.2 (3)	758 (7)	766.34 (7) g)		755 (10)	766.51 (3)	766.680 (13)
832.02 (12)	831.96 (3)	831.83 (4)	831.8 (1)	831.8 (1)	832	831.8 (5)	831.7 (5)	830 (7)	831.83 (4) g)	830	830 (2)	832.01 (3)	831.984 (12)
865.87 (24)	865.88 (14)	865.6 (3)	865.5 (3)	865.2 (2)		866 (2)	864 (1)		860 (10)			865.93 (14)	865.92 (6)
1014.71 (12)	1014.59 (5)	1014.7 (2)	1014.4 (3)	1014.1 (5)		1014.8 (10)	1014 (1)		1020 (3)		1020 (10)	1014.64 (5)	1014.38 (4)
1080.10 (13)	1080.11 (6)	1080.2 (1)	1080.0 (3)	1080.0 (5)	1076	1080.9 (10)	1079 (1)	1060 (15)	1076 (3)			1080.16 (6)	1080.64 (4)
1103.52 (20)	1103.7 (8)	1103.4 (4)	1103.0 (6) ?									1103.52 (20)	1103.52 (20)
1109.48 (13)	1109.43 (5)	1109.5 (2)	1109.8 (3)	1109.1 (1)	1106	1109.6 (8)	1108.5 (10)	1100 (15)	1104 (2)	1104	1100 (5)	1109.48 (5)	1109.509 (23)
1196.15 (14)	1196.28 (5)	1196.6 (2)	1196.1 (3)	1195.5 (5)		1196.6 (10)	1194 (1)		1188 (2)			1196.33 (5)	1196.33 (5)
		1234.3 (4)	1234.6 (10)									1234.3 (4)	1234.3 (4)
1270.79 (18)	1270.66 (8)	1270.8 (2)	1270.3 (3)	1270.0 (5)	1265	1271.2 (10)	1269 (1)		1265 (2)			1270.71 (8)	1270.75 (6)

*) value omitted from the statistical data analysis.

g) value reported in 1965Me07, but measured by A. Green, PhD thesis, University of California at Davis with Ge detector (unpublished).

Table 3. Measured, deduced and adopted gamma-ray emissions probabilities for gamma-transitions in β^- -decay of ²¹¹Pb.

Eg, keV	1988Hi14	1976BI13	1971Da34	1968Br17	1968Go15	1968Ha21	1967Da20	1967Da10	1965Co06	1965Me07	1963Va05	1962Gi03	deduced, rel	adopted, %
65		0.57 (6)	0.60 (4)	0.58 (11)			0.10 (5) *	~0.35 *		0.5 (2)			0.59 (3)	0.077 (4)
95			0.14 (2)					-0.10					0.14 (2)	0.018 (3)
314		0.20 (3)	0.24 (3)	0.19 (4)	0.10 (5)		0.26 (5)	0.21 (7)	0.90 (21) *	~0.2 *			0.206 (16)	0.0268 (21)
343		1.63 (13) *	0.27 (4)		0.15 (5)					~0.3 *			0.22 (3)	0.029 (4)
362		0.326 (24)		0.30 (8)									0.324 (23)	0.042 (3)
405	29.3 (9)	30.2 (14)	30.0 (9)	30.8 (15)	29.6 (20)	28.6 (11)	28.0 (28)	29.9 (35)	30.6 (28)	27.4 (12)	26 (5)	34(4)	29.4 (4)	3.83 (6)
427	13.9 (4)	14.2 (7)	13.5 (6)	14.3 (8)	13.7 (10)	11.6 (7)	14.0 (14)	13.9 (17)	21.5 (21) *	14.5 (14)	12.5 (25)	22 (3) *	13.9 (3)	1.81 (4)
429					0.065 (25)								0.065 (25)	0.008 (3)
504	0.045 (6)		0.12 (2) *		~ 0.006 *								0.045 (6)	0.0059 (8)
610		0.407 (24) *	0.18 (3)	0.38 (4)	0.25 (5)		0.30 (6)	0.21 (7)		0.9 (2) *		0.76 (13) *	0.25 (7)	0.033 (9)
676		0.130 (8)		0.173 (15)	0.10 (5)				1.3 (3) *				0.139 (7)	0.0181 (9)
705	3.6 (1)	3.6 (3)	3.77 (19)	3.68 (23)	3.7 (3)	2.9 (1) *	3.0 (3)	3.3 (4)	5.3 (6) *	3.7 (2)	3.8 (11)	5.5 (4) *	3.61 (7)	0.47 (1)
767	4.94 (16)	5.1 (4)	5.55 (28)	5.04 (30)	4.9 (3)	4.5 (1)	4.0 (4)	5.1 (6)	6.3 (6)	5.2 (2)		6.1 (4)	4.8 (3)	0.62 (4)
832	26.7 (8)	25.4 (20)	29.8 (7) *	25.6 (23)	24.1 (17)	27.4 (4)	23.0 (23)	26.4 (35)	26.4 (28)	27.4 (12)	24.8 (25)	34.2 (13) *	26.9 (3)	3.50 (5)
866	0.042 (6)	0.033 (4)	0.050 (8)	0.053 (15)	0.07 (2)		0.03 (1)	0.0347 (14)		0.04 (2)			0.0354 (13)	0.0046 (2)
1014	0.129 (8)	0.122 (8)	0.14 (1)	0.128 (15)	0.15 (1)		0.13 (2)	0.125 (21)		0.14 (2)		0.38 (19) *	0.133 (4)	0.0173 (5)
1081	0.095 (6)	0.090 (7)	0.120 (12)	0.083 (10)	0.08 (1)	0.0025 (1) *	0.08 (2)	0.104 (14)	0.49 (7) *	0.13 (12) *			0.093 (4)	0.0121 (5)
1104	0.033 (4)	0.049 (5)	0.040 (6)	0.023 (5)									0.036 (5)	0.0047 (7)
1110	0.90 (3)	0.82 (6)	1.15 (8)	0.79 (8)	0.81 (6)	0.0105 (7) *	0.70 (15)	0.87 (10)	0.83 (14)	1.03 (10)	1.07 (16)	1.46 (19) *	0.891 (21)	0.116 (3)
1196	0.072 (5)	0.081 (6)	0.10 (1)	0.079 (15)	0.08 (1)		0.08 (2)	0.076 (14)		0.11 (3)			0.079 (3)	0.0103 (4)
1234			0.010 (2)	0.0053 (15)									0.0070 (23)	0.0009 (3)
1271	0.043 (4)	0.057 (5)	0.070 (7)	0.048 (8)	0.08 (1)	0.0006 (1) *	0.05 (1)	0.042 (7)		0.06 (2)			0.052 (9)	0.0068 (12)

*) value omitted in the statistical data analysis.

Table 4. Gamma-ray energies and emission probabilities (relative to $I_\gamma(351.07\gamma) = 100$) for transitions that were not placed in the proposed decay scheme of ²¹¹Pb.

E_γ , keV	1988Hi14	1971Da34	1968Ha21	1968Br17
81.0 (2)		0.35 (9)		
83.8 (1)		0.45 (7)		
88.2 (2)		0.13 (3)		
94.3 (3)		0.09 (2)		
97.3 (2)		0.09 (1)		
244			0.003 (1)	
275			0.004 (1)	
478.0 (4)		0.10 (2)		
479.6 (2)	0.04 (1)			
481.1 (4)		0.20 (4)		
481.92 (12)	0.08 (1)			
491.82 (12)	0.032 (6)			
494.2 (3)	0.013 (5)			
500.4 (5)		0.09 (2)		
502.0 (2)	0.028 (6)			
951			0.0017 (1)	
1090.5 (5)		0.020 (5)		
1120 (1)				0.0019 (11)

5. References

- 1939Sa11 B.W. Sargent. Can. J. Res. 17A (1939) 103.
(Half-life)
- 1962Gi03 M. Giannini, D. Prosperi, S. Sciuti. Nuovo Cimento 25 (1962) 1227.
(Gamma-ray emission energies and probabilities)
- 1963Va05 S.E. Vandenbosch, C.V.K. Baba, P.R. Christensen, O.B. Nielsen, H. Nordby. Nucl. Phys. 41 (1963) 482.
(Gamma-ray emission energies and probabilities)
- 1965Co06 C.R. Cothorn, R.D. Connor. Can. J. Phys. 43 (1965) 383.
(Gamma-ray emission energies and probabilities)
- 1965Me07 R.O. Mead, J.E. Draper. Phys. Rev. 139 (1965) B9.
(Gamma-ray emission energies and probabilities)
- 1965Nu03 M. Nurmi, D. Giessing, W. Sievers, L. Varga. Ann. Acad. Sci. Fennicae Ser. A VI (1965) No. 167.
(Half-life)
- 1967Da10 W.F. Davidson, C.R. Cothorn, R.D. Connor. Can. J. Phys. 45 (1967) 2295.
(Gamma-ray emission energies and probabilities)
- 1967Da20 J. Dalmaso, H. Maria. C.R. Acad. Sci. (Paris) 265B (1967) 822.
(Gamma-ray emission energies and probabilities)
- 1968Br17 C. Briançon, C.F. Leang, R. Walen. C.R. Acad. Sci. (Paris) 266B (1968) 1533.
(Gamma-ray emission energies and probabilities)
- 1968G015 S. Gorodetzky, F.A. Beck, T. Byrski, A. Knipper. Nucl. Phys. A117 (1968) 208.
(Gamma-ray emission energies and probabilities)
- 1968Ha21 W.D. Hamilton, K.E. Davies. Nucl. Phys. A114 (1968) 577.
(Gamma-ray emission energies and probabilities)

- 1971Da34 E.F. da Silveira, A.G. de Pinho, C.V. de Barros. *An. Acad. Brasil. Cienc.* 43 (1971) 97.
(Gamma-ray emission energies and probabilities)
- 1971Go40 N.B. Gove, M.J. Martin. *Nucl. Data Tables A10* (1971) 205.
(Log ft)
- 1976B113 K. Blaton-Albicka, B. Kotlinska-Filipek, M. Matul, K. Stryczniewicz, M. Nowicki, E. Ruchowska-Lukasiak. *Nukleonika* 21 (1976) 935.
(Gamma-ray emission energies and probabilities)
- 1988Hi14 M.M. Hindi, E.G. Adelberger, S.E. Kellogg, T. Murakami. *Phys. Rev. C* 38 (1988) 1370.
(Gamma-ray emission energies and probabilities)
- 1996Sc06 E. Schönfeld, H. Janssen. *Nucl. Instrum. Methods Phys. Res. A* 369 (1996) 527.
(K-shell fluorescence yields)
- 1998ScZM E. Schönfeld, G. Rodloff. Report PTB-6.11-98-1 Braunschweig (1998).
(K Auger electron energies)
- 1999ScZX E. Schönfeld, G. Rodloff. Report PTB-6.11-1999-1 Braunschweig (1999).
(K X-ray energies and relative emission probabilities)
- 2000Sc47 E. Schönfeld, H. Janssen. *Appl. Radiat. Isot.* 52 (2000) 595.
(X-ray and Auger electron emission probabilities and energies)
- 2003Au03 G. Audi, A.H. Wapstra, C. Thibault. *Nucl. Phys. A* 729 (2003) 337.
(Q value)
- 2003De44 R.D. Deslattes, E.G. Kessler Jr., P. Indelicato, L. de Billy, E. Lindroth, J. Anton. *Rev. Mod. Phys.* 75 (2003) 35.
(K and L X-ray energies)
- 2004Br45 E. Browne. *Nucl. Data Sheets* 103 (2004) 183
(Nuclear levels, multipolarities and mixing ratios)
- 2008DuZX C. Dulieu, M.M. Bé, V. Chisté. Proc. Int. Conf. on Nuclear Data for Science and Technology, Nice, France, 22-27 April 2007 (2008) 97.
(SAISINUC software)
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor Jr. *Nucl. Instrum. Methods Phys. Res. A* 589 (2008) 202.
(Theoretical ICCs)
- 2011Ko04 F.G. Kondev, S. Lalkovski. *Nucl. Data Sheets* 112 (2011) 707.
(Nuclear levels)

²¹¹Bi – Comments on Evaluation of Decay Data by A. Luca

This evaluation was completed in July 2009. The literature available by December 31st, 2008 was included.

1. Evaluation Procedures

The Limitation of Relative Statistical Weight (LWM) method was applied for averaging numbers throughout this evaluation; this method was implemented by using the computer code LWEIGHT, ver. 4 (designed for Excel, MS Office). The uncertainty assigned to an average value in this evaluation is never lower than the lowest uncertainty of any of the experimental input values.

2. Decay Scheme

²¹¹Bi decays 99.724 (4) % by alpha particle emissions, populating the ²⁰⁷Tl ground state (83.56 (23) %) and the 351.03 keV excited state (16.16 (23) %). ²¹¹Bi has also a weak beta minus decay branch (0.276 (4) %) to the ground state of ²¹¹Po; although these β⁻ particles were not observed experimentally (the low intensity beta-particle emission is obscured by the intense β⁻ particles emission from the ²¹¹Pb sources used for measurements), the existence of the beta minus decay and the adopted value of the corresponding branching ratio are based on the alpha-particle spectrometry measurements of the emission probabilities ratio, $I_{\alpha}({}^{211}\text{Po})/(I_{\alpha}({}^{211}\text{Po})+I_{\alpha}({}^{211}\text{Bi}))$, performed by several scientists (see references from Table 1). The adopted value represents the weighted mean of the experimental results published in the literature (see also Table 1, below); an earlier value, 0.32 % (without a quoted uncertainty), determined by Rutherford et al. (1931), was not taken into account. Another important study of the ²¹¹Bi decay scheme is presented in the reference 1966Go13. The most recent evaluations of the ²¹¹Bi nuclear structure, alpha and beta minus decay data, published in Nuclear Data Sheets, were made by M. J. Martin (1993) and E. Browne (2004). In the present evaluation, the spin and parity of the levels have been adopted from the above mentioned A = 207 and A = 211 ENSDF mass-chain evaluations (1993Ma73 and 2004Br45, respectively).

Table 1: Beta minus branching ratio for the ²¹¹Bi decay

Beta minus branching ratio (experimental), %	Reference
0.274 (4)	1967Da10
0.274 (10)	1965Nu03
0.29 (1)	1962Gi04
Recommended value: 0.276 (4) %	

3. Nuclear Data

The adopted alpha decay energy value $Q(\alpha) = 6750.33$ (46) keV, is from 2003Au03. This value is in very good agreement with the effective $Q(\alpha)$ value of 6750.63 keV (with an uncertainty of 0.21 keV), calculated from the decay scheme data, by using the SAISINUC software, version 2008 April. The adopted beta minus decay energy value $Q(\beta) = 574$ (5) keV is also from 2003Au03.

3.1. Half-life

In the literature, four measured ²¹¹Bi half-life ($T_{1/2}$) values are reported. All these measurements are old (the most recent is from 1970), so new half-life measurements are needed to improve the quality of the evaluation. The half-life values and their uncertainties are presented in Table 2.

The value recommended by Curie et al. (1931), with an estimated uncertainty added by the evaluator, was included, too. The uncertainty of other two results (1954Sp32 and 1965Nu03) was also estimated by the evaluator. The set of data is consistent and the recommended value, 2.15 minutes, with an uncertainty of 0.02 minutes, is the weighted average (LWM, $\chi^2_{\nu}=3.7$) of the four input values.

Table 2 : ²¹¹Bi Half-life values

T _{1/2} (minutes)	Uncertainty of T _{1/2} (minutes)	Reference
2.16	0.08	1931Cu01
2.15	0.02	1954Sp32
2.13	0.03	1965Nu03
2.22	0.06	1970Mu21

3.2. Alpha and Beta transitions and emissions

In the literature, the most important reference that studies measurements of alpha-particle energies and emission intensities for ²¹¹Bi alpha transitions is 1991Ry01.

For this evaluation, the two adopted alpha-particle emission energies were calculated as weighted means of the experimental values presented in Table 3 (both data sets are consistent):

Table 3: Energy of the alpha-particles emitted in the ²¹¹Bi decay

Alpha-particle group	Energy of the alpha particles (experimental), keV	Reference
$\alpha_{0,1}$	6300 (10)	1989It01
	6278.2 (7)	1991Ry01
	6279 (1)	1992Sc26
	Recommended energy value: 6278.5 (9) keV	
$\alpha_{0,0}$	6622.9 (6)	1971Gr17 and 1991Ry01
	6620 (10)	1989It01
	6621.33 (69)	1991Ry01
	6623 (1)	1992Sc26
	Recommended energy value: 6622.4 (6) keV	

The ratio of the 6278.5 keV to the sum of 6278.5 keV and 6622.4 keV alpha-particle emission probabilities was determined in a similar way, as the weighted mean of four experimental values reported in the literature and presented below, in Table 4. This data set is discrepant and, consequently, the uncertainty was expanded to include in its range the most precise relative value (16.43 (4) from 1967Da10); the adopted value is 16.20 (23). Considering both the experimental results and the normalization condition (modified to take into account the beta minus decay, see section 2), i.e. the sum of the two absolute alpha-particle emission probabilities must be 100 % - 0.276 (4) % = 99.724 (4) %, the computed absolute emission probability of the 6278.5 keV alpha-particles is 16.16 (23) %. The 6622.4 keV alpha-particles absolute emission probability is then 83.56 (23) %.

The beta minus transition is of the first order forbidden type (non-unique) and populates the ground state of ²¹¹Po. The beta particles must have a maximum energy of 574 keV (corresponding to the Q(β) value) and an absolute emission probability of 0.276 (4) %. The adopted values of the average beta minus energy (172.9 (18) keV) and log ft (5.99) were obtained by using the LOGFT computer program.

Table 4: Experimental values of the relative alpha-particles emission probability ratio (6278.5 keV) / (6278.5 keV + 6622.4 keV)

Alpha-particle emission probability ratio (6278.5 keV) / (6278.5 keV + 6622.4 keV) x 100	Reference
15.8 (1)	1962Gi04
15.9 (3)	1962Wa18
16.02 (5)	1966Go13
16.43 (4)	1967Da10

3.3. γ - transitions: γ rays and internal conversion electrons

There is a single gamma-ray transition following the ²¹¹Bi decay. Both its energy and emission probability were studied by many scientists. Table 5 summarizes the experimental results published in the literature. The adopted energy of this gamma-ray transition is the weighted mean of the 6 values from Table 5 (consistent data set): 351.03 (4) keV.

The absolute emission probability of this gamma-ray was determined from the alpha feeding of 16.16 (23) % to the ²⁰⁷Tl excited state: 16.16 (23) / 1.243 (4) = 13.00 (19) %, where 0.243 (4) is the total

internal conversion coefficient (total ICC), which is in good agreement with the experimental values reported in references 1976BI13 and 1982Mo30 (see Table 5).

All the internal conversion coefficients (ICCs) adopted in this evaluation were computed with the program BrIcc, version 2.2 /2008, using the "Frozen Orbitals" approximation (2008Ki07). The energy range of the internal conversion electrons corresponding to the gamma-ray transition is from 265.5 keV to 351.02 keV, whereas the total number of conversion electrons emitted per 100 disintegrations is 3.17 (7) (i.e.3.17 (7) %)

Table 5: ²¹¹Bi γ -ray Energy and Absolute Emission Probability (experimental values)

E_γ (keV)	Uncertainty E_γ (keV)	Absolute Emission Probability (%)	Uncertainty of absolute emission probability (%)	Reference
351.0	0.1	10.70	0.30	1968Br17
351.0	0.3			1973UrZX
351.01	0.04			1975VaYT
351.07	0.05	12.27	1.4	1976BI13
351.89	0.20	13.3	1.3	1982Mo30
351.06	0.12			1988Hi14

4. Atomic data

The K-shell fluorescence yield (ω_K), the mean L-shell fluorescence yield ($\bar{\omega}_L$) and the mean number of vacancies in the L-shell produced by one vacancy in the K-shell (η_{KL}) were determined using the computer program EMISSION v.3.10, 28-Jan-2003: 0.963 (4), 0.367 (15) and 0.812 (5) respectively.

4.1. Auger electrons and X-rays

The relative probability values of the K Auger electron emissions (KLL, KLX, KXY) normalized to the KLL value, were computed using the same EMISSION computer program. The total numbers of K and L Auger electrons emitted per disintegration were also calculated (in %): 0.096 (11) and 1.620 (21), respectively. The energy ranges for K and L Auger electrons were filled-in by the SAISINUC program, version 2008 April.

The relative probability (normalized to $K_{\alpha 1}$ X-rays emission) and the absolute emission probability values of the different groups of K and L X-rays were determined using the same EMISSION program. The adopted values (in %) of the total absolute emission probability of the KX-rays and LX-rays were 2.50 (6) and 0.931 (19), respectively. The energy range values of the K and L X-rays are from the tables linked to SAISINUC.

Only one reference reporting the measurement of the ²⁰⁷Tl KX-rays energies and emission probabilities was found in the literature (1976BI13). A comparison between these experimental values and the results of this evaluation is presented in Table 6.

For the two K_α X-rays the results are in very good agreement for energy and unsatisfactory for the absolute emission probability values. The Tl- $K_{\alpha 2}$ and Tl- $K_{\alpha 1}$ x-ray absolute emission probabilities reported in 1976BI13 are about 30 % lower than expected (See Table 6). The cause of this serious disagreement is unknown.

For the two K_β X-rays, the energy values are in good agreement, whereas the absolute emission probabilities values again are in clear disagreement. There are at least two possible causes of this disagreement:

- the evaluated values refer to a sum of three components, not only to $K\beta_1$, respectively $K\beta_2$ (see the Note below Table 6);
- the measurements reported in the article 1976BI13 include also the Rn $K_{\alpha 1}$ X-rays with an energy of 83.788 keV, situated just between the two components of interest; the presence of this additional peak makes the spectral analysis of this region more difficult, considering the software tools available in 1976 (a higher uncertainty than reported for the experimental results is possible).

This second assumption is supported by the very good agreement between the sum of Tl- $K\beta_1$ and Tl- $K\beta_2$ absolute emission probabilities (in %), according to Table 6: 0.542 (12) (evaluated) and 0.55 (6) (experimental).

Neither measurements of ²⁰⁷Tl LX-rays energies nor of emission probabilities were found in the literature, in order to compare them with the results of this evaluation.

Table 6: Comparison of the evaluated TI KX-rays energy and absolute emission probability values with experimental results from 1976BI13

X-rays identification	Evaluated energy (keV)	Evaluated Absolute Emission Probability (in %)	Experimental energy (keV)	Experimental absolute emission probability (in %) (1976BI13)
Tl-K α_2	70.832	0.726 (16)	70.839 (13)	0.51 (8)
Tl-K α_1	72.872	1.225 (27)	72.857 (10)	0.82 (12)
Tl-K β_1	82.577	0.417 (11)	83.019 (80)	0.24 (4)
Tl-K β_2	84.838	0.124 (4)	84.720 (50)	0.31 (4)

* Note: the evaluated absolute emission probabilities of the two K β X-rays include not only the contributions of the K β_1 and K β_2 components, but also K β_3 , K β_5 , K β_4 and K $O_{2,3}$.

5. Main production mode

The main production mode of ²¹¹Bi is by beta minus decay of the ²¹¹Pb nuclei (both nuclides are members of the Actinium-Uranium natural radioactive series). ²¹¹Bi can be produced also by the alpha decay of ²¹⁵At (a process of very low probability in the above mentioned radioactive series, because ²¹⁵At is produced by the weak beta minus decay branch of ²¹⁵Po, which is about 2.3·10⁻⁴ %).

6. References

- 1931Cu01 M. Curie, A. Debierne, A.S. Eve, H. Geiger, O. Hahn, S.C. Lind, St. Meyer, E. Rutherford and E. Schweidler, "The Radioactive constants as of 1930", Rev. Mod. Phys. 3, 427 (1931), citing: St. Meyer and F. Paneth, Mitt. Ra. Inst. 104, Wien. Ber. Ila, 127, 147 (1918).
- 1931Ru02 E. Rutherford, C.E. Wynn-Williams and W.B. Lewis, Proc. Roy. Soc. (London), Ser. A, 133, 351 (1931).
- 1954Sp32 F.N. Spiess, "Alpha-Emitting Isomer: Polonium-211", Phys. Rev. 94, 1292 (1954)
- 1962Gi04 M. Giannini, D. Prospero and S. Sciuti, "Intensity Measurements of Alpha Groups from ²¹¹Bi, ²¹¹Po, ²¹⁹Rn and ²²³Ra by Means of Solid State Counter Techniques", Nuovo Cimento 25, 1314 (1962)
- 1962Wa18 R.J. Walen, V. Nedovesov and G. Bastin-Scoffier, "Spectrographie α de ²²³Ra (AcX) et Ses Derivés", Nuclear Phys. 35, 232 (1962)
- 1965Nu03 M. Nurmia, D. Giessing, W. Sievers and L. Varga, "Studies of the Natural Actinium Radioactive Series", Ann. Acad. Sci. Fennicae, Ser.A VI, No.167 (1965)
- 1966Go13 S. Gorodetzky, F. Beck and A. Knipper, "Melange M1+E2 et Effets de Penetration dans la Conversion Interne des Rayons γ de 350 keV du ²⁰⁷Tl", Nucl. Phys. 82, 275 (1966)
- 1967Da10 W.F. Davidson, C.R. Cothorn and R.D. Connor, "Studies in the Decay of the Active Deposit of Actinium. III. Levels in ²¹¹Bi and its Daughter Products", Can. J. Phys. 45, 2295 (1967)
- 1968Br17 C. Briancon, C.F. Leang and R. Walen, "Etude du Spectre γ Emis par le Radium-223 et Ses Derives", Compt. Rend. 266B, 1533 (1968)
- 1970Mu21 Von H. Mundschenk, "Uber ein Verfahren zur Abtrennung kurzlebiger Radionuklide unter Ausnutzung des Ruckstosseffektes", Radiochim. Acta 14, 72 (1970)
- 1971Gr17 B. Grennberg and A. Rytz, "Absolute Measurements of α -Ray Energies", Metrologia 7, 65 (1971)
- 1971Ko37 G.A. Korolev, A.A. Vorobyov and Y.K.Zalite, "A Microwave Method for Lifetime Measurements of Nuclear States Excited By α -Particle Decay", Nucl. Instrum. Methods 97, 323 (1971)
- 1973UrZX D.F. Urquhart, "The Gamma Ray Spectra of Uranium and Thorium Ores by High Resolution Ge(Li) Spectrometry", Report AAEC / TM 634 (1973)
- 1975VaYT V.M. Vakhtel, T. Vylov, V.M. Gorozhankin, N.A. Galovkov, B.S. Dzhelepov, R.B. Ivanov, M.A. Mikhailova, Yu.V. Norseev and V.G. Chumin, Conf. Dubna, 149 (1975)
- 1976BI13 K. Blaton-Albicka, B. Kottinska-Filipek, M. Matul, K. Stryczniewicz, M. Nowicki and E. Ruchowska-Lukasiak, "Precision Gamma-Ray Spectroscopy of the Decay of ²²³Ra and its Daughter Products", Nukleonika 21, 935 (1976)
- 1982Mo30 M.H. Momeni, "Analyses of Uranium and Actinium Gamma Spectra: An Application to Measurements of Environmental Contamination", Nucl. Instrum. Methods 193, 185 (1982)

- 1988Hi14 M.M. Hindi, E.G. Adelberger, S.E. Kellogg and T. Murakami, "Search for the I-Forbidden Beta Decay $^{207}\text{Tl} \rightarrow ^{207}\text{Pb}^*(570 \text{ keV})$ ", Phys. Rev. C 38, 1370 (1988)
- 1989It01 J.T. Iturbe, "Alpha-Particle Spectrum of ^{219}Rn and its Daughters from Pitchblende Samples using Silicon Surface-Barrier Detectors", Nucl. Instrum. Methods Phys. Res. A 274, 404 (1989)
- 1991Ry01 A. Rytz, "Recommended Energy and Intensity Values of Alpha Particles from Radioactive Decay", At. Data Nucl. Data Tables 47, 205 (1991)
- 1992Sc26 P. Schuurmans, J. Wouters, P. De Moor, N. Severijns, W. Vanderpoorten, J. Vanhaverbeke and L. Vanneste, "Anisotropic Alpha-Emission in the ^{223}Ra Decay Chain", Hyperfine Interactions 75, 423 (1992)
- 1993Ma73 M.J. Martin, "Nuclear Data Sheets Update for A = 207", Nucl. Data Sheets 70, 315 (1993)
- 2003Au03 G. Audi, A.H. Wapstra and C. Thibault, "The AME2003 atomic mass Evaluation (II). Tables, graphs, and references", Nucl. Phys. A 729, 337 (2003).
- 2004Br45 E. Browne, "Nuclear Data Sheets for A = 211", Nucl. Data Sheets 103, 183 (2004)
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr., Nucl. Instrum. Meth. Phys. Res. A589, 202 (2008) (Theoretical ICC).

²¹¹Po – Comments on Evaluation of Decay Data by A. Luca

This evaluation was completed in August 2009. The literature available by December 31st, 2008 was included.

1. Evaluation Procedures

The Limitation of Relative Statistical Weight (LWM) method was applied for averaging numbers throughout this evaluation; this method was implemented by using the computer code LWEIGHT, ver. 4 (designed for Excel, MS Office). The uncertainty assigned to an average value in this evaluation is never lower than the lowest uncertainty of any of the experimental input values.

2. Decay Scheme

²¹¹Po decays 100 % by alpha particle emissions, mainly to the ground state of ²⁰⁷Pb. The most recent evaluations of the ²¹¹Po nuclear structure and decay data, published in Nuclear Data Sheets, were done by E. Browne (2004) and M.J. Martin (1993). In the present evaluation, the spin, parity and energy of the levels, together with the multipolarities and mixing ratios of the γ -ray transitions, have been adopted from the A=207 ENSDF mass-chain evaluation 1993Ma73. This data evaluation refers only to the decay of the ²¹¹Po ground state, and not to the decay of the ²¹¹Po metastable state at 1462 keV (with a half-life of 25.2 s).

3. Nuclear Data

The adopted alpha decay energy value $Q(\alpha)=7594.48(51)$ keV, is from 2003Au03. This value is in very good agreement with the effective $Q(\alpha)$ value of 7594.2 (20) keV, deduced from average radiation energies from the decay scheme data, by using the SAISINUC software, version 2008 April.

3.1. Half-life

In the literature, five measured ²¹¹Po half-life ($T_{1/2}$) values are reported. The value from 1931Cu01 is unrealistically low (in strong disagreement with all the other values), and was excluded from the data set, according to the Chauvenet's criterion implemented by the LWEIGHT computer code. The half-life values and their uncertainties are presented in Table 1. The value recommended by Curie et al. (1931), with an estimated uncertainty added by the evaluator, has been also included. The set of data, excluding the value given in 1931Cu01 is consistent and the recommended half-life value, 0.516 (3) s, is the weighted average (LWM, $\chi^2_{\nu}=3.7$) of the four input values. The reference *Nuclear Science References* (NSR) keynumbers are:

Table 1 : ²¹¹Po Half-life values

$T_{1/2}$ (seconds)	Uncertainty of $T_{1/2}$ (seconds)	Reference
0.005	0.005	1931Cu01
0.52	0.02	1954Sp32
0.5	0.1	1954Wi26
0.56	0.04	1958To25
0.516	0.003	1974Ba29

3.2. Alpha transitions and emissions

The most important reference in the literature that studies measurements of alpha-particle energies and emission intensities for ²¹¹Po alpha transitions is 1991Ry01.

For this evaluation, three adopted alpha-particle energies were deduced as weighted means of the experimental values presented in Table 2; the complete data set of the main alpha-particle group (7450.2 keV) is consistent, while from the other two data sets, the values from 1953AsZZ were rejected from the weighted mean computations, according to Chauvenet's criterion. The hindrance factors were determined by using the ALPHAD version 2.0a (2006) computer program (developed at BNL/NNDC, USA).

Table 2: Energy of the alpha-particles emitted in the ²¹¹Po decay

Alpha-particle group	Energy of the alpha particles (experimental), keV	Reference
$\alpha_{0,2}$	6570 (10)	Leininger et al. (1951)
	6570 (40)	1951Ne02
	6560 (10)	1953AsZZ
	6569 (20)	1953Ho49
	6571 (4)	1968GuZX
	6570.0 (25)	1969Go23
	6568 (1)	1978Ya04
	Recommended energy value: 6568.4 (10) keV	
$\alpha_{0,1}$	6900 (10)	Leininger et al. (1951)
	6900 (40)	1951Ne02
	6880 (10)	1953AsZZ
	6895 (20)	1953Ho49
	6890.7 (25)	1962Wa18
	6891 (4)	1968GuZX
	6892.5 (25)	1969Go23
	6891 (1)	1978Ya04
Recommended energy value: 6891.2 (10) keV		
$\alpha_{0,0}$	7430 (40)	1951Ne02
	7430 (20)	1953Ho49
	7442 (15)	1954Br07
	7450.3 (2)	1962Wa18, updated in 1991Ry01
	7440 (30)	1963Jo09
	7448 (4)	1968GuZX
	7449.8 (30)	1969Go23, updated in 1991Ry01
	7460 (20)	1969Ha32
	7448 (10)	1970Va13
	7443.3 (20)	1982Bo04, updated in 1991Ry01
	7456.2 (30)	1985La17, updated in 1991Ry01
	Recommended energy value: 7450.2 (3) keV	

The reference 1951Ne02 is the only one that reports the detection of another group of alpha particles emitted in the decay of ²¹¹Po, with an energy of 6340 (60) keV and an emission probability of $7 \cdot 10^{-4}$. In the report UCRL-2325 (1953), R.W. Hoff doesn't confirm the detection of this alpha-particle decay branch, but establishes that there are no alpha-particle groups with emission probabilities higher than $2 \cdot 10^{-4}$, in the energy range from 6.26 MeV to 6.57 MeV. In a similar study presented in 1969Go23, a maximum limit of $2 \cdot 10^{-5}$ is given for the emission probability of any alpha-particle group in the energy range (5.88 to 6.43) MeV. As there is no other experimental data to confirm the existence of this branch, the evaluator adopted a decay scheme with only the three alpha particle groups given in Table 2.

The recommended emission probabilities of the alpha-particle emissions of 6568.4 keV and 6891.2 keV are the weighted means of the published experimental values, presented in

Table 3. From the first data set in this table, the values of 1978Ya04 (0.58 (1) %) and 1951Ne02 (0.48 (5) %) were rejected by the Chauvenet's criterion. A similar procedure, applied to the second data set from Table 3, lead to the rejection of the value published in the reference 1962Wa18 (0.70 (14) %).

The adopted emission probability of the main alpha-particle emission, 7450.2 keV, was computed from the normalization condition (the sum of the three alpha-particle emission probabilities is 100 %): 98.936 (19) %.

Table 3: Emission probabilities of the alpha-particles emitted in the ²¹¹Po decay

Alpha-particles energy (keV)	Experimental emission probability (%)	Reference
6568.4	0.5 (1)	Leininger et al. (1951)
	0.48 (5)	1951Ne02
	0.53 (1)	1953AsZZ
	0.53 (5)	1953Ho49
	0.53 (3)	1968GuZX
	0.537 (19)	1975Ja04
	0.58 (1)	1978Ya04
	0.513 (9)	1985La17
Recommended emission probability: 0.523 (9) %; HF=17.9		
6891.2	0.6 (1)	Leininger et al. (1951)
	0.57 (5)	1951Ne02
	0.50 (1)	1953AsZZ
	0.50 (5)	1953Ho49
	0.70 (14)	1962Wa18
	0.57 (3)	1968GuZX
	0.546 (19)	1975Ja04
	0.60 (1)	1978Ya04
	0.524 (9)	1985La17
Recommended emission probability: 0.541 (17) %; HF=272		
7450.2	Recommended emission probability: 98.936 (19) %; HF=112	

3.3. γ - transitions: γ rays and internal conversion electrons

There are only few papers that report measurements of the γ -ray energies and emission probabilities following the ²¹¹Po decay: 1954Mi70 and 1975Ja04 (energy values), 1968Br17 and 1985La17 (absolute emission probabilities), respectively. 1975Ja04 and 1972As11 report relative emission probabilities.

The adopted gamma-ray energy values are the weighted means of the experimental values published in 1954Mi70 and 1975Ja04, as presented below in Table 4 (for the 328 keV photons just one measurement was made, and published in 1975Ja04):

Table 4: Gamma-rays energy values in the decay of ²¹¹Po

Experimental energy values (keV)	Reference
562 (5)	1954Mi70
569.65 (10)	1975Ja04
Recommended energy value: 569.65 (15) keV	
880 (8)	1954Mi70
897.8 (1)	1975Ja04
Recommended energy value: 897.8 (2) keV	
328.2 (2)	1975Ja04
Recommended energy value: 328.2 (2) keV	

Using the measured 328.2 keV gamma-ray relative photon intensity of 0.6 (2) (the intensity of the 569.65 keV photons is considered as 100, see reference 1975Ja04), the internal

conversion coefficients and the intensity balance for each of the two excited states of ²⁰⁷Pb, the corresponding absolute gamma-ray emission probabilities and their uncertainties were computed for all the three γ rays; these data are given below in Table 5. A comparison between the evaluated data and the experimental values (included in the same table, with the corresponding references) shows a good agreement, with the exception of the relative emission probability of 897.8 keV reported by 1985La17.

The internal conversion coefficients were computed with the program BrIcc, version 2.2b/20-Jan-2009, using the “Frozen Orbitals” approximation. In the article of L.J. Jardine (1975), an experimental value of 0.016 (3) was determined for the K-conversion coefficient associated to both gamma-ray transitions of 569.65 keV and 897.8 keV; this value is in good agreement with the theoretical ICC's, computed with BrIcc: 0.01583 (23), respectively 0.0192 (3).

Table 5: γ -rays absolute and relative emission probabilities in the decay of ²¹¹Po

E_γ (keV)	Recommended Absolute Emission Probability (%)	Experimental Absolute Emission Probability (%)	Evaluated relative emission probabilities	Experimental relative emission probabilities	Total ICC (α_T)
328.2	0.0032 (11)		0.6 (2)*	0.6 (2) ^c	0.334 (5)
569.65	0.534 (17)	0.534 (19) ^a 0.512 (36) ^b	100	100.0 (14) ^b 100 ^{c,d}	0.0216 (3) E2
897.8	0.507 (9)	0.535 (40) ^b	94.9 (35)	104.4 (20) ^b 97 (5) ^c 83 (11) ^d	0.0233 (4) M1

Note: a – reference 1968Br17; b – reference 1985La17; c – reference 1975Ja04;
d – reference 1972As11 (renormalized); * - value adopted from reference 1975Ja04.

4. Atomic data

The K-shell fluorescence yield (ω_K), the mean L-shell fluorescence yield ($\overline{\omega}_L$) and the mean number of vacancies in the L-shell produced by one vacancy in the K-shell (η_{KL}) were determined using the computer program EMISSION v.3.10, 28-Jan-2003: 0.963 (4), 0.379 (15) and 0.811 (5) respectively.

4.1. Auger electrons and X-rays

The relative probability values of the K Auger electron emissions (KLL, KLX, KXY) normalized to the KLL value, were computed using the EMISSION computer program. The total numbers of K and L Auger electrons emitted per 100 disintegrations were also calculated as 0.00071 (8) and 0.01216 (17), respectively. The energy ranges for K and L Auger electrons were filled-in by the SAISINUC program, version 2008 April.

The relative probability (normalized to $K_{\alpha 1}$ X-rays emission) and the absolute emission probability values of the different groups of K and L X-rays were deduced using the EMISSION program. The adopted values of the total absolute emission probability of the KX-rays and LX-rays were 0.0184 (5) % and 0.00740 (16) %, respectively. The energy range values of the K and L X-rays are from the tables linked to SAISINUC.

Neither measurement of ²⁰⁷Pb KX-rays and LX-rays energies nor of emission probabilities was found in the literature in order to compare it with the results of this evaluation.

5. Main production mode

The main production mode of ²¹¹Po is by β^- decay of the ²¹¹Bi nuclei (in the Actinium-Uranium natural radioactive series).

6. References

- 1931Cu01 M. Curie, A. Debierne, A.S. Eve, H. Geiger, O. Hahn, S.C. Lind, St. Meyer, E. Rutherford and E. Schweidler, "The Radioactive constants as of 1930", *Rev. Mod. Phys.* 3, 427 (1931).
- R.F. Leininger, E. Segrè, F.N. Spiess, "The Half-Life of Ac C' ", *Phys. Rev.* 82, A334, 1951.
- 1951Ne02 H.M. Neumann, I. Perlman, "Long-Lived Bi207 and Energy Levels of Pb207", *Phys. Rev.* 81, 958 (1951).
- 1953AsZZ F. Asaro, "The Complex Alpha Spectra of the Heavy Elements", Thesis, Univ. California (1953); UCRL-2180 (1953).
- 1953Ho49 R.W. Hoff, "Orbital Electron Capture in the Heaviest Elements", Thesis, Univ. California (1953); UCRL-2325 (1953).
- 1954Br07 G.H. Briggs, "The Energies of Natural Alpha Particles", *Rev. Mod. Phys.* 26, 1 (1954).
- 1954Mi70 J.W. Mihelich, A.W. Schardt, E. Segrè, "Energy Levels in Po210", *Phys. Rev.* 95, 1508 (1954).
- 1954Sp32 F.N. Spiess, "Alpha-Emitting Isomer: Polonium-211", *Phys. Rev.* 94, 1292 (1954).
- 1954Wi26 M.M. Winn, "Short-Lived Alpha Emitters Produced by ³He and Heavy Ion Bombardments", *Proc. Phys. Soc. (London)* 67A, 949 (1954).
- 1958To25 P.A. Tove, "Alpha-Emitters with Short Half-Life Induced by Protons on Heavy Elements", *Arkiv Fysik* 13, 549 (1958).
- 1962Wa18 R.J. Walen, V. Nedovesov, G. Bastin-Scoffier, "Spectrographie α de ²²³Ra (AcX) et Ses Derives", *Nuclear Phys.* 35, 232 (1962).
- 1963Jo09 W.B. Jones, "New Isomers of Astatine-212", *Phys. Rev.* 130, 2042 (1963).
- 1968Br17 C. Briancon, C.F. Leang, R. Walen, "Etude du Spectre γ Emis par le Radium-223 et Ses Derives", *Compt. Rend.* 266B, 1533 (1968).
- 1968GuZX L. Gueth, S. Gueth, E. Daroczy, B.S. Dzhelepov, Y.V. Narseev, V.A. Khalkin, "Investigation of the ²¹¹At, ²¹¹Po, ²¹⁰At, ²⁰⁹At, and ²⁰⁷At Alpha Spectra with Semiconductor Alpha Spectrometer", Report JINR-P6-4079 (1968).
- 1969Go23 N.A. Golovkov, S. Guetkh, B.S. Dzhelepov, Y.V. Narseev, V.A. Khalkin, V.G. Chumin, "Fine Structure of the Alpha-Particle Spectra of ²⁰⁹At, ²¹⁰At, ²¹¹At and ²¹¹Po", *Izv. Akad. Nauk SSSR, Ser. Fiz.* 33, 1622 (1969); *Bull. Acad. Sci. USSR, Phys. Ser.* 33, 1489 (1970).
- 1969Ha32 R.L. Hahn, M.F. Roche, K.S. Toth, "Alpha Decay of ²²⁷U", *Phys. Rev.* 182, 1329 (1969).
- 1970Va13 K. Valli, E.K. Hyde, J. Borggreen, "Production and Decay Properties of Thorium Isotopes of Mass 221-224 Formed in Heavy-Ion Reactions", *Phys. Rev. C*1, 2115 (1970).
- 1972As11 G. Astner, "Properties of ²¹¹At as observed in the decay of ²¹¹Rn", *Phys. Scr.* 5, 31 (1972).
- 1974Ba29 A.R. Barnett, J.S. Lilley, "Interaction of Alpha Particles in the Lead Region Near the Coulomb Barrier", *Phys. Rev. C*9, 2010 (1974).
- 1975Ja04 L.J. Jardine, "Decays of ²¹¹At, ²¹¹Po, and ²⁰⁷Bi", *Phys. Rev. C*11, 1385 (1975).
- 1978Ya04 M. Yanokura, H. Kudo, H. Nakahara, K. Miyano, S. Ohya, O. Nitoh, "The Half-Life of ²⁰⁷Bi and Decays of ²¹¹At and ²¹¹Po", *Nucl. Phys.* A299, 92 (1978).

- 1982Bo04 J.D. Bowman, R.E. Eppley, E.K. Hyde, "Alpha Spectroscopy of Nuclides Produced in the Interaction of 5 GeV Protons with Heavy Element Targets", Phys. Rev. C25, 941 (1982).
- 1985La17 R.M. Lambrecht, S. Mirzadeh, "Cyclotron Isotopes and Radiopharmaceuticals - XXXV. Astatine-211", Int. J. Appl. Radiat. Isotop. 36, 443 (1985).
- 1991Ry01 A. Rytz, "Recommended Energy and Intensity Values of Alpha Particles from Radioactive Decay", At. Data Nucl. Data Tables 47, 205 (1991).
- 1993Ma73 M.J. Martin, "Nuclear Data Sheets Update for A = 207", Nucl. Data Sheets 70, 315 (1993).
- 1996Sc06 E. Schonfeld, H. Janssen, "Evaluation of Atomic Shell Data", Nucl. Instrum. Methods Phys. Res. A369, 527 (1996).
- 2003Au03 G. Audi, A.H. Wapstra and C. Thibault, "The AME2003 atomic mass Evaluation (II). Tables, graphs, and references", Nucl. Phys. A 729, 337 (2003).
- 2004Br45 E. Browne, "Nuclear Data Sheets for A = 211", Nucl. Data Sheets 103, 183 (2004).
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr., "Evaluation of theoretical conversion coefficients using Brlcc", Nucl. Instrum. Methods Phys. Res. A589, 202 (2008).

²¹¹At - Comments on evaluation of decay data

by A. L. Nichols

Evaluated: August 2010**Evaluation Procedure**

Limitation of Relative Statistical Weight Method (LWM) was applied to average the decay data when appropriate.

Decay Scheme

A reasonably simple decay scheme was constructed from the α -particle and γ -ray measurements of Hoff (1953Ho49), Gray (1956Gr11), Golovkov *et al.* (1969Go23), Jardine (1975Ja04), and Chumin *et al.* (2001Ch66), and studies of the α branching fraction by Neumann and Perlman (1951Ne02), Golovkov *et al.* (1969Go23), Afanasiev *et al.* (1970AfZZ), Jardine (1975Ja04), Yanokura *et al.* (1978Ya04), and Lambrecht and Mirzadeh (1985La17).

Nuclear Data

²¹¹At is an important α -emitting radionuclide in therapeutic nuclear medicine, along with daughter ²¹¹Po.

Half-life

The recommended half-life of 7.216 (7) hours has been adopted from five sets of measurements (1956Gr11, 1959Ra08, 1961Ap01, 1962Th08, 1978Ya04).

Half-life measurements

Reference	Half-life (hours)
1956Gr11	7.20 ± 0.05
1959Ra08	7.23 ± 0.04
1961Ap01	7.214 ± 0.007
1962Th08	7.17 ± 0.09
1978Ya04	7.23 ± 0.02
Recommended value	7.216 ± 0.007

A half-life of 0.516 (3) second was adopted for ²¹¹Po from the DDEP evaluation of Luca (July-November 2009), while the ²⁰⁷Bi half-life of 32.9 (14) years was taken from the DDEP evaluation of Bé and Chisté (December 2009). More recently, a further re-evaluation of the half-life of ²⁰⁷Bi by Kondev and Lalkovski resulted in a recommended value of 31.55 (4) years (2011Ko04).

Branching fractions

Neumann and Perlman (1951Ne02), Golovkov *et al.* (1969Go23), Afanasiev *et al.* (1970AfZZ), Jardine (1975Ja04), Yanokura *et al.* (1978Ya04), and Lambrecht and Mirzadeh (1985La17) have determined the α branching fraction for ²¹¹At. These data were used to derive an alpha branch of 41.78 (8) %, along with a matching electron-capture branch of 58.22 (8) %.

Reference	BF _{α}
1951Ne02	0.409 ± 0.005
1969Go23	0.418 ± 0.002
1970AfZZ	0.413 ± 0.013
1975Ja04	0.419 ± 0.005
1978Ya04	0.4174 ± 0.0010
1985La17	0.4194 ± 0.0016
Recommended value	0.4178 ± 0.0008
α branch	(41.78 ± 0.08) %

Q values

Q_{EC} of 785.4 (25) keV and Q _{α} of 5982.4 (13) keV were adopted from the evaluated tabulations of Audi *et al.* (2003Au03).

Alpha Particles

Alpha-particle measurements reveal a relatively simple α -decay mode (1969Go23, 1975Ja04, 1991Ry01, 2001Ch66). The Q _{α} of 5982.4 (13) keV (2003Au03) and nuclear level energies as defined by Kondev and Lalkovski (2011Ko04) were used to calculate the alpha-particle energies, while the alpha-particle emission probabilities were primarily adopted from the measurements of Golovkov *et al.* (1969Go23), Afanasiev *et al.* (1970AfZZ), Jardine (1975Ja04) and Chumin *et al.* (2001Ch66).

Alpha-particle emission probabilities per 100 disintegrations of ²¹¹At, and hindrance factors.

E _{α} (keV)	P _{α}					HF
	1969Go23*	1970AfZZ [†]	1975Ja04 [#]	2001Ch66 [#]	Recommended value	
4895.4 (13)	–	–	–	< 0.000 04	< 0.000 04	> 9.6
4993.4 (13)	–	~ 0.000 4 ?	–	–	~ 0.000 4	~ 3.8
5140.3 (13)	0.001 7 (9)	0.001 5	0.001 0 (3)	0.001 1 (2)	0.001 1 (2)	10.1
5211.9 (13)	0.005 4 (8)	0.006 7	0.003 6 (8)	0.003 9 (3)	0.003 9 (3)	7.3
5869.0 (13)	41.8 (2)	[41.78]	41.93	41.80	41.78 (8)	1.59

* Calculated from measurements of the relative alpha-particle emission probabilities.

[†] Calculated from measurements of the relative alpha-particle emission probabilities, but no uncertainties listed; absolute emission probability of 41.78 % was adopted for the 5869.0-keV α particle to convert other data in this study to comparable absolute values.

[#] Calculated from measurements of the relative gamma-ray emission probabilities.

An unweighted mean value of 1.422 (13) was adopted for the radius parameter r₀(²⁰⁷Bi) as derived from the equivalent data for neighboring nuclei (1998Ak04), and used in the calculation of α -hindrance factors (HF):

$$\begin{aligned}
 r_0(^{207}\text{Bi}) &= [r_0(^{206}\text{Pb}) + r_0(^{208}\text{Po})] / 2 \\
 &= [1.40882(10) + 1.4343(34)] / 2 \\
 &= 1.422 (13)
 \end{aligned}$$

Gamma Rays

Energies

All gamma-ray transition energies and uncertainties were calculated from the structural details of the proposed decay scheme. Nuclear level energies were adopted from Browne for ²¹¹Po and from Kondev and Lalkovski for ²⁰⁷Bi (2004Br45, 2011Ko04).

Emission Probabilities

The absolute emission probabilities of the 149.72-, 222.69-, 669.77-, 742.74- and 892.46-keV gamma rays from the α -decay branch were derived from a combination of the alpha-particle emission probabilities populating the ground state and 669.77-, 742.74-, 892.46- and 992.43-keV nuclear levels of ²⁰⁷Bi (1969Go23, 1970AfZZ), relevant relative emission probabilities for these gamma rays (1975Ja04, 1985La17), theoretical internal conversion coefficients of Band *et al.* (2002Ba85, 2008Ki07), and depopulating ratios of the 149.72-, 222.69- and 892.46-keV gamma transitions as quantified by Kondev and Lalkovski (2011Ko04). A weighted mean value of 0.245 (12) was adopted for the absolute emission probability of the 687.7-keV gamma ray from the EC-decay branch, based on the gamma-ray spectroscopy studies of Jardine (1975Ja04) and Lambrecht and Mirzadeh (1985La17).

Gamma-ray emission probabilities relative to 100 % for the 569.7-keV gamma ray of daughter ²¹¹Po.

E_γ (keV)	P_γ^{rel}		
	1975Ja04	1985La17	Recommended value
[569.70]	100	100.0 (14)	–
669.77 (7)	1.1 (2)	–	–
687.2 (7)	79 (4)	83.0 (20)	82 (2)
742.74 (7)	0.3 (1)	–	–

Absolute gamma-ray emission probabilities per 100 disintegrations of ²¹¹At.

E_γ (keV)	P_γ^{abs}		
	1975Ja04*	1985La17	Recommended value
$\gamma_{3,2}$ (Bi) 149.72 (10)	–	–	~ 0.000 05
$\gamma_{3,1}$ (Bi) 222.69 (10)	–	–	~ 0.000 04
$\gamma_{1,0}$ (Bi) 669.77 (7)	0.003 4 (6)	–	0.003 8 (3)
$\gamma_{1,0}$ (Po) 687.2 (7)	0.245 (12)	0.247 (26)	0.245 (12)
$\gamma_{2,0}$ (Bi) 742.74 (7)	0.000 9 (3)	–	0.001 25 (19)
$\gamma_{3,0}$ (Bi) 892.46 (7)	–	–	~ 0.000 14

* Derived from an absolute emission probability of 0.31 (2) per 100 decay of ²¹¹At for the 569.70-keV gamma transition within the α decay of daughter ²¹¹Po.

Multipolarities and Internal Conversion Coefficients

The nuclear level schemes specified by Browne for ²¹¹Po and Kondev and Lalkovski for ²⁰⁷Bi have been used to define the multipolarities of the gamma transitions on the basis of known spins and parities (2004Br45, 2011Ko04). All known gammas are (M1 + E2) transitions, and their mixing ratios have been derived on the basis of the studies of Astner and Alpsten (1970As07), Schmidt-Ott and Dincklage (1978Sc12), and Herzog *et al.* (1983He09). Recommended internal conversion coefficients have been determined from the frozen orbital approximation of Kibédi *et al.* (2008Ki07), based on the theoretical model of Band *et al.* (2002Ba85, 2002Ra45).

Gamma-ray emissions: recommended energies, emission probabilities, multiplicities and theoretical internal conversion coefficients (frozen orbital approximation).

	E_γ (keV)	P_γ^{abs}	Multiplicity	α_K	α_L	α_{M+}	α_{tot}	
$\gamma_{3,2}$ (Bi)	149.72 (10)	$\sim 0.000\ 05$	86.2 % M1 + 13.8 % E2 $\delta = 0.40$ (20)	2.3 (3)	0.50 (4)	0.2	3.0 (3)	α
$\gamma_{3,1}$ (Bi)	222.69 (10)	$\sim 0.000\ 04$	86.2 % M1 + 13.8 % E2 $\delta = 0.40$ (10)	0.76 (5)	0.147 (2)	0.043	0.95 (5)	α
$\gamma_{1,0}$ (Bi)	669.77 (7)	0.003 8 (3)	94.1 % M1 + 5.9 % E2 $\delta = 0.25$ (3)	0.042 6 (8)	0.007 25 (12)	0.002 15	0.052 0 (9)	α
$\gamma_{1,0}$ (Po)	687.2 (7)	0.245 (12)	96.15 % M1 + 3.85 % E2 $\delta = -0.20$ (2)	0.043 7 (7)	0.007 52 (12)	0.002 38	0.053 6 (9)	E C
$\gamma_{2,0}$ (Bi)	742.74 (7)	0.001 25 (19)	91.7 % M1 + 8.3 % E2 $\delta = 0.30$ (3)	0.032 0 (6)	0.005 44 (10)	0.001 66	0.039 1 (7)	α
$\gamma_{3,0}$ (Bi)	892.46 (7)	$\sim 0.000\ 14$	33.8 % M1 + 66.2 % E2 $\delta = 1.4$ (2)	0.011 7 (11)	0.002 15 (16)	0.000 65	0.014 5 (13)	α

Electron-capture Transitions

Energies

Electron-capture energies were calculated from the nuclear level energies of Browne (2004Br45) and a Q_{EC} value of 785.4 ± 2.5 keV taken from Audi *et al.* (2003Au03).

Transition probabilities

The EC transition probabilities were calculated from BF_{EC} of 0.5822 (8) and the absolute emission probability and theoretical internal conversion coefficients of the 687.2-keV gamma ray.

EC transition probabilities per 100 disintegrations of ²¹¹At.

	E_{EC} (keV)	P_{EC}	Transition type	$\log ft$	P_K	P_L	P_M
EC _{0,1}	98.2 ± 2.6	0.258 ± 0.013	1 st forbidden non-unique	5.77	0.015 (17)	0.684 (10)	0.301 (7)
EC _{0,0}	785.4 ± 2.5	57.96 ± 0.08	1 st forbidden non-unique	5.97	0.773 1 (2)	0.169 3 (1)	0.057 58 (4)

Atomic Data

The x-ray and Auger-electron data have been calculated using the evaluated gamma-ray data, and atomic data from 1996Sc06, 1998ScZM and 1999ScZX. Both the x-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.01, 28 January 2003). This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

K and L X-ray emission probabilities per 100 disintegrations of ²¹¹At.

			Energy (keV)	Photons per 100 disint.
XL		(Bi)	9.420 – 15.709	0.000 136 (14)
	XL ₁	(Bi)	9.420	0.000 003 3 (4)
	XL _α	(Bi)	10.731 – 10.839	0.000 063 (7)
	XL _η	(Bi)	11.712	0.000 001 03 (15)
	XL _β	(Bi)	12.480 – 13.393	0.000 057 (6)
	XL _γ	(Bi)	15.248 – 15.709	0.000 011 0 (12)
XK _α	XK _{α2}	(Bi)	74.8157 (9)	0.000 098 (15)
	XK _{α1}	(Bi)	77.1088 (10)	0.000 164 (25)
XK' _{β1}	XK _{β3}	(Bi)	86.835)
	XK _{β1} "	(Bi)	87.344) 0.000 056 (9)
	XK _{β5}	(Bi)	87.862)
XK' _{β2}	XK _{β2}	(Bi)	89.732)
	XK _{β4}	(Bi)	90.074) 0.000 017 (3)
	XKO _{2,3}	(Bi)	90.421)
XL		(Po)	9.658 – 16.213	18.6 (8)
	XL ₁	(Po)	9.658	0.465 (12)
	XL _α	(Po)	11.016 – 11.130	8.53 (20)
	XL _η	(Po)	12.085	0.134 (4)
	XL _β	(Po)	12.823 – 13.778	7.76 (14)
	XL _γ	(Po)	15.742 – 16.213	1.53 (3)
XK _α	XK _{α2}	(Po)	76.864 (4)	12.66 (9)
	XK _{α1}	(Po)	79.293 (5)	21.08 (12)
XK' _{β1}	XK _{β3}	(Po)	89.256)
	XK _{β1} "	(Po)	89.807) 7.26 (12)
	XK _{β5}	(Po)	90.363)
XK' _{β2}	XK _{β2}	(Po)	92.263)
	XK _{β4}	(Po)	92.618) 2.26 (5)
	XKO _{2,3}	(Po)	92.983)

Electron energies were determined from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

Data Consistency

An effective Q-value of 2956.7 (16) keV has been adopted from the atomic mass evaluation of Audi *et al.* (2003Au03) while in the course of formulating the decay scheme of ²¹¹At. This value has subsequently been compared with the Q-value calculated by summing the

contributions of the individual emissions to the ²¹¹At alpha- and EC-decay processes (i.e. α , γ , conversion electrons, etc.):

$$\text{calculated Q-value} = \sum (E_i \times P_i) = 2957 (5) \text{ keV}$$

Percentage deviation from the effective Q-value of Audi *et al.* is $-(0.01 \pm 0.17) \%$, which supports the derivation of a highly consistent decay scheme.

References

- 1951Ne02 H.M. NEUMANN, I. PERLMAN, Long-lived Bi²⁰⁷ and energy levels of Pb²⁰⁷, Phys. Rev. 81 (1951) 958-962. [BF α]
- 1953Ho49 R.W. HOFF, Orbital electron capture in the heaviest elements, PhD thesis, University of California, Berkeley, September 1953. [α , Auger-electron and γ spectra]
- 1956Gr11 P.R. GRAY, Auger effect in the heaviest elements, Phys. Rev. 101 (1956) 1306-1314. [half-life, α and Auger-electron spectra]
- 1959Ra08 W.J. RAMLER, J. WING, D.J. HENDERSON, J.R. HUAZENGA, Excitation functions of bismuth and lead, Phys. Rev. 114 (1959) 154-162. [half-life]
- 1961Ap01 E.H. APPELMAN, Half-lives of At²¹¹ and Bi²⁰⁷, Phys. Rev. 121 (1961) 253-255. [half-life]
- 1962Th08 T.D. THOMAS, G.E. GORDON, R.M. LATIMER, G.T. SEABORG, Spallation-fission competition in astatine compound nuclei formed by heavy-ion bombardment, Phys. Rev. 126 (1962) 1805-1810. [half-life]
- 1969Go23 N.A. GOLOVKOV, Sh. GUETKH, B.S. DZHELEPOV, Yu.V. NORSEEV, V.A. KHALKIN, V.G. CHUMIN, Fine structure of the alpha-particle spectra of ²⁰⁹At, ²¹⁰At, ²¹¹At and ²¹¹Po, Bull. Acad. Sci. USSR, Phys. Ser. 33 (1970) 1489-1496. [E α , P α , BF α]
- 1970As07 G. ASTNER, M. ALPSTEN, Gamma-ray transitions in ²⁰⁷Bi following the decay of ²⁰⁷Po, Nucl. Phys. A140 (1970) 643-657. [multipolarity]
- 1970AfZZ V.P. AFANASIEV, M. BOCHVAROVA, N.A. GOLOVKOV, I.I. GROMOVA, R.B. IVANOV, V.I. KUZIN, Y.V. NORSEEV, V.G. CHUMIN, α -decay of ²¹¹Rn and ²¹²Rn, JINR-P6-4972 (1970). [E α , P α , BF α]
- 1975Ja04 L.J. JARDINE, Decays of ²¹¹At, ²¹¹Po, and ²⁰⁷Bi, Phys. Rev. C11 (1975) 1385-1391. [E α , P α , E γ , P γ , BF α]
- 1977La19 F.P. LARKINS, Semiempirical Auger-electron energies for elements $10 \leq Z \leq 100$, At. Data Nucl. Data Tables 20 (1977) 311-387. [Auger-electron energies]
- 1978Ya04 M. YANOKURA, H. KUDO, H. NAKAHARA, K. MIYANO, S. OHYA, O. NITOH, The half-life of ²⁰⁷Bi and decays of ²¹¹At and ²¹¹Po, Nucl. Phys. A299 (1978) 92-98. [half-life, BF α]
- 1978Sc12 W.-D. SCHMIDT-OTT, R.-D.V. DINCKLAGE, Spin of ^{207m}Po (2.79 s); conversion measurements in ^{207g}Po decay, Z. Phys. A286 (1978) 301-305. [multipolarity]

- 1983He09 P. HERZOG, H. WALITZKI, K. FREITAG, H. HILDEBRAND, K. SCHLÖSSER, Nuclear orientation and NMR/ON of ^{205,207}Po, Z. Phys. A – Atoms and Nuclei 311 (1983) 351-362. [multipolarity]
- 1985La17 R.M. LAMBRECHT, S. MIRZADEH, Cyclotron isotopes and radiopharmaceuticals – XXXV. Astatine-211, Int. J. Appl. Radiat. Isot. 36 (1985) 443-450. [E_α, P_α, E_γ, P_γ, E_x, P_x]
- 1991Ry01 A. RYTZ, Recommended energy and intensity values of alpha particles from radioactive decay, At. Data Nucl. Data Tables 47 (1991) 205-239. [E_α, P_α]
- 1996Sc06 E. SCHÖNFELD, H. JANßEN, Evaluation of atomic shell data, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527-533. [X_K, X_L, Auger electrons]
- 1998Ak04 Y.A. AKOVALI, Review of alpha-decay data from doubly-even nuclei, Nucl. Data Sheets 84 (1998) 1-114. [alpha decay, r₀ parameters]
- 1998ScZM E. SCHÖNFELD, G. RODLOFF, Tables of the energies of K-Auger electrons for elements with atomic numbers in the range from Z = 11 to Z = 100, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX E. SCHÖNFELD, G. RODLOFF, Energies and relative emission probabilities of K X-rays for elements with atomic numbers in the range from Z = 5 to Z = 100, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2001Ch66 V.G. CHUMIN, K.Ya. GROMOV, Sh.R. MALIKOV, Yu.V. NORSEEV, Zh.K. SAMATOV, V.I. FOMINYKH, A.P. CHEREVATENKO, L.V. YURKOVA, ²¹¹Po α-decay to the ²⁰⁷Pb 1633-keV level, Bull. Russian Acad. Sci., Physics 65 (2001) 27-30. [E_α, P_α]
- 2002Ba85 I.M. BAND, M.B. TRZHASKOVSKAYA, C.W. NESTOR Jr., P.O. TIKKANEN, S. RAMAN, Dirac–Fock internal conversion coefficients, At. Data Nucl. Data Tables 81 (2002) 1-334. [ICC]
- 2002Ra45 S. RAMAN, C.W. NESTOR Jr., A. ICHIHARA, M.B. TRZHASKOVSKAYA, How good are the internal conversion coefficients now? Phys. Rev. C66 (2002) 044312, 1-23. [ICC]
- 2003Au03 G. AUDI, A.H. WAPSTRA, C. THIBAUT, The AME2003 atomic mass evaluation (II). Tables, graphs and references, Nucl. Phys. A729 (2003) 337-676. [Q-value]
- 2004Br45 E. BROWNE, Nuclear data sheets for A = 211, Nucl. Data Sheets 103 (2004) 183-268. [Nuclear structure, level energies]
- 2008Ki07 T. KIBÉDI, T.W. BURROWS, M.B. TRZHASKOVSKAYA, P.M. DAVIDSON, C.W. NESTOR Jr., Evaluation of theoretical conversion coefficients using BrIcc, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202-229. [ICC]
- 2011Ko04 F.G. KONDEV, S. LALKOVSKI, Nuclear data sheets for A = 207, Nucl. Data Sheets, 112 (2011) 707-853. [Nuclear structure, level energies]

**²¹²Pb – Comments on evaluation of decay data
by A. L. Nichols**

Evaluated: July/August 2001

Re-evaluated: January 2004

Evaluation Procedures

Limitation of Relative Statistical Weight Method (LWM) was applied to average numbers throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

A reasonably simple and consistent decay scheme has been constructed from the gamma-ray measurements of 1960Ro16, 1961Gi02, 1972DaZA, 1978Av01, 1982Sa36, 1983Sc13, 1983Va22, 1984Ge07 and 1992Li05. Only five distinct gamma-ray emissions were identified with ²¹²Pb decay in all of these studies. A further gamma ray has been added in the evolution of the decay scheme (energy of 123.45 keV) to achieve the necessary population-depopulation balance of the 115.183 keV nuclear level of ²¹²Bi.

Low-energy gamma transitions have been postulated to exist in the decay scheme of ²¹²Pb (with energies between 40 and 60 keV). However, this possibility was rejected on the basis of insufficient experimental evidence in the open literature. Further studies are required to resolve this issue, and confirm the correctness of the proposed decay scheme.

Nuclear Data

²²⁸Th decay chain is important in quantifying the environmental impact of the decay of naturally occurring ²³²Th. Specific radionuclides in this decay chain are noteworthy because of their decay characteristics (²²⁴Ra alpha decay to ²²⁰Rn; ²¹²Bi and ²⁰⁸Tl gamma-ray emissions).

Half-life

The recommended half-life is the weighted mean of three elderly measurements (1952Bu72, 1953Ma26 and 1955To11). Further studies are merited to determine this value with greater confidence.

Reference	Half-life (h)
1952Bu72	10.67(5)
1953Ma26	10.64(3)
1955To11	10.643(12)
Recommended Value	10.64(1)

Gamma Rays

Energies

All gamma-ray transition energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies of 1992Ar05 were adopted, and used to determine the energies and associated uncertainties of the gamma-ray transitions between the various populated-depopulated levels.

Emission Probabilities

Weighted mean relative emission probabilities were determined for the 115.183, 176.64, 238.632 and 300.09 keV gamma rays, using the relevant data from the measurements of 1960Ro16, 1961Gi02, 1972DaZA, 1978Av01, 1982Sa36, 1983Sc13, 1983Va22, 1984Ge07 and 1992Li05. The relative emission probability of the 415.27 keV gamma ray was adopted from the studies of 1961Gi02, while a further gamma ray has been added in the evolution of the decay scheme (energy of 123.45 keV) to achieve the necessary population-depopulation balance of the 115.183 keV nuclear level of ²¹²Bi.

Gamma-ray Emission Probabilities: Relative to P_g(238.632 keV) of 100

E _g (keV)	P _g ^{rel}				
	1960Ro16	1961Gi02	1972DaZA	1978Av01	1982Sa36
115.183(5)	[observed]	1.4(3)	1.3(3)	1.4(1)	1.65(12)
123.45(1)	-	-	-	-	-
176.64(1)	~ 0.5	0.50(10)	0.10(3)	-	-
238.632(2)	100	100	100	100(3)	100(5)
300.09(1)	7.7(4)	6.9(4)	7.7(15)	6.3(2)	6.7(5)
415.27(1)	~ 0.3	0.33(5)	-	-	-

E _g (keV)	P _g ^{rel} (cont.)				
	1983Sc13	1983Va22	1984Ge07	1992Li05	Recommended Values*
115.183(5)	-	-	1.37(2)	-	1.43(5)
123.45(1)	-	-	-	-	0.22(1)
176.64(1)	-	-	0.12(1)	-	0.12(1)
238.632(2)	100(3)	100(1)	100(1)	100(2)	100(1)
300.09(1)	7.5(2)	7.3(1)	7.6(1)	7.6(3)	7.3(3)
415.27(1)	-	-	-	-	0.33(5)

* Weighted mean values adopted when appropriate using LWEIGHT; remainder derived from proposed decay scheme.

A weighted mean normalisation factor of 0.436(3) was calculated for the emission probabilities from the measurements of 1982Sa36, 1983Sc13, 1983Va22, 1984Ge07 and 1992Li05.

Absolute Gamma-ray Emission Probabilities: Normalisation Factor

E _g (keV)	P _g ^{abs}					Recommended Value*
	1982Sa36	1983Sc13	1983Va22	1984Ge07	1992Li05	
238.632(2)	0.430(20)	0.435(12)	0.440(6)	0.433(4)	0.441(10)	0.436(3)

* Weighted mean value adopted from LWEIGHT.

Multipolarities and Internal Conversion Coefficients

The nuclear level scheme specified by 1992Ar05 has been used to define the multipolarities of the gamma transitions on the basis of known spins and parities. Limited studies of the internal conversion coefficients support the proposed transition types: 100%M1 for the 115.183, 238.632 and 300.09 keV gamma rays (1957Ni11, 1957Kr49, 1959Se59, 1960Ro16, 1963Da11, 1969Kr06 and 1978Av01); the 176.64 and 415.27 keV gamma rays were also assigned 100%M1 multipolarity, while the 123.45 keV gamma transition was defined as E2.

Multipolarity Assignments

Reference	E _g (keV)	Multipolarity
1957Ni11	115.183(5)	M1 [K/L = 5(1)]
1957Kr49	115.183(5)	M1
	176.64(1)	E0 [K/L = 1 : 0.18(2)]
	238.632(2)	M1
	300.09(1)	M1

1959Se59	115.183(5)	M1 [L _I :L _{II} :L _{III} → 100 : 10.4(3) : 0.88(10)]
	238.632(2)	M1 [L _I :L _{II} :L _{III} → 100 : 10.4(2) : 0.74(5)]
1960Ro16	115.183(5)	M1 [α _K = 5.8(9)]
	238.632(2)	M1 [α _K = 0.74(7)]
1963Da11	238.632(2)	M1
	415.27(1)	M1 [α _K ~ 0.35]
1969Kr06	238.632(2)	M1
1978Av01	115.183(5)	E2
	238.632(2)	M1 (+ E2)
	300.09(1)	M1 + E2

Beta-particle Emissions

Energies

All beta-particle energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies of 1992Ar05 and the Q-value were used to determine the energies and uncertainties of the beta-particle transitions to the various levels.

Emission Probabilities

The beta-particle emission probabilities were calculated from gamma -ray transition probability balances, using the recommended gamma -ray emission probabilities and the theoretical internal conversion coefficients of 1978Ro22:

415.272 keV nuclear level:

[Σ P_{γ_i} (1 + α_i) depopulating 415.27 keV level]NF was calculated to be 11.65(47)NF; since NF = 0.436(3), beta-particle emission probability is calculated to be 5.1(2)% (0.051(2));

238.632 keV nuclear level:

{ [Σ P_{γ_i} (1 + α_i) depopulating 238.63 keV level] - P_γ(176.64 keV)(1 + α(176.64 keV)) }NF was calculated to be 192.7(34)NF; since NF = 0.436(3), beta -particle emission probability is calculated to be 84.0(14)% (0.840(14));

115.183 keV nuclear level:

spin and parity considerations support zero beta decay to this level;

population/depopulation by gamma transitions require balance of the form

Σ P_{γ_i} (1 + α_i) populating 115.18 keV level should equal P_γ(115.18 keV)(1 + α(115.18 keV));

hence, derivation of transition probability P_γ(123.45 keV) = 0.85(4)NF

ground state (0.0 keV):

(i) through population of ground state: [Σ P_{γ_i} (1 + α_i) populating ground state]NF + P_{b_{0,0}} = 100

and NF = 0.436(3) to give P_{b_{0,0}} = 10.9(14)% (0.109(14))

(ii) through summation of beta decay and NF = 0.436(3)

$$P_{b_{0,0}} = 10.9(14)\% (0.109(14))$$

Beta-particle Emission Probabilities per 100 Disintegrations of ²¹²Pb

E _b (keV)	P _b	
	1948Ma30	Recommended Values*
159(2)	-	5.1(2)
335(2)	-	84.0(14)
574(2)	12(2)	10.9(14)

* Recommended emission probabilities derived from evaluated gamma-ray emission probabilities and theoretical internal conversion coefficients.

Atomic Data

The x-ray data have been calculated using the evaluated gamma -ray data, and the atomic data from 1996Sc06, 1998ScZM and 1999ScZX.

References

- 1948Ma30 - D. G. E. Martin and H. O. W. Richardson, The Nuclear β -spectra of Thorium B \rightarrow C and C \rightarrow C', and the Intensities of Some β -ray Lines of Thorium (B+C+C''), Proc. Phys. Soc. (London) 195A(1948)287. [P_β]
- 1952Bu72 - H. V. Buttlar, Neubestimmung der Halbwertszeit des ThB (²¹²Pb), Naturwissenschaften 39(1952)575. [Half-life]
- 1953Ma26 - P. Marin, G. R. Bishop and H. Halban, The Absolute Standardization of the 2.615 MeV Gamma -Rays of ThC'' and the Cross Section for the Photodisintegration of the Deuteron at this Energy, Proc. Phys. Soc. (London) 66A(1953)608. [Half-life]
- 1955To11 - J. Tobailem and J. Robert, Mesure de la Periode du ThB (²¹²Pb), J. Phys. Radium 16(1955)115. [Half-life]
- 1957Ni11 - K. O. Nielsen, O. B. Nielsen and M. A. Waggoner, Internal Conversion Coefficients for M1 Transitions in ⁸³Bi²¹⁴, ⁸⁴Po²¹⁴ and ⁸³Bi²¹², Nucl. Phys. 2(1956/57)476. [P_{ce} , multipolarity]
- 1957Kr49 - E. M. Krisyouk, A. G. Sergeev, G. D. Latyshev and V. D. Vorobyov, Decay Scheme of Pb ²¹², Nucl. Phys. 4(1957)579. [P_{ce} , multipolarity]
- 1959Se59 - A. G. Sergeev, V. D. Vorobyev, A. S. Remenny, T. I. Kolchinskaya, G. D. Latyshev and Yu. S. Yegorov, Influence of the Finite Dimensions of the Nucleus on the Relative Conversion Coefficients in the L-subshells, Nucl. Phys. 9(1958/59)498. [P_{ce} , multipolarity]
- 1960Ro16 - P. G. Roetling, W. P. Ganley and G. S. Klaiber, The Decay of Pb ²¹², Nucl. Phys. 20(1960)347. [P_γ , multipolarity]
- 1961Gi02 - M. Giannini, D. Prosperi and S. Sciuti, Decay Scheme of ²¹²Pb, Nuovo Cimento 21(1961)430. [P_γ]
- 1963Da11 - H. Daniel and G. Lührs, Aufbau eines Automatisch Arbeitenden $\pi\sqrt{2}$ -Spektrometers und Untersuchung des Konversionslinienspektrums von In^{114m} und ThB mit Folgeprodukten, Z. Phys. 176(1963)30. [P_{ce} , multipolarity]
- 1969Kr06 - D. Krpic, R. Stepic, M. Bogdanovic and M. Mladenovic, K/L ₃, M/L and (N + O + ...) / M Ratios for 239 keV M1 Transition in ²¹²Bi, Fizika 1(1969)171. [P_{ce} , multipolarity]
- 1972DaZA - J. Dalmaso, Recherches sur le Rayonnement Gamma de Quelques Radioéléments Naturels Appartenant à la Famille du Thorium, PhD thesis, University of Nice (1972); J. Dalmaso, H. Maria and C. Ythier, Étude du Rayonnement γ du Thorium 228 et de ses Dérivés, et plus Particulièrement du Thallium 208 (ThC), C. R. Acad. Sci. Paris 277B(1973)467. [P_γ]
- 1978Av01 - F. T. Avignone and A. G. Schmidt, γ -ray and Internal-conversion Intensity Studies of Transitions in the Decay of ²²⁸Th, Phys. Rev. C17(1978)380. [P_γ , multipolarity]
- 1978Ro22 - F. Rösel, H. M. Fries, K. Alder and H. C. Pauli, Internal Conversion Coefficients for all Atomic Shells, ICC Values for Z = 68-104, At. Data Nucl. Data Tables 21(1978)291-514. [ICC]
- 1982Sa36 - S. Sadasivan and V. M. Raghunath, Intensities of Gamma Rays in the ²³²Th Decay Chain, Nucl. Instrum. Meth. 196(1982)561. [P_γ]
- 1983Sc13 - U. Schötzgig and K. Debertin, Photon Emission Probabilities per Decay of ²²⁶Ra and ²³²Th in Equilibrium with their Daughter Products, Int. J. Appl. Radiat. Isot. 34(1983)533. [P_γ]
- 1983Va22 - R. Vaninbrouckx and H. H. Hansen, Determination of γ -ray Emission Probabilities in the Decay of ²²⁸Th and its Daughters, Int. J. Appl. Radiat. Isot., 34(1983)1395. [P_γ]
- 1984Ge07 - R. J. Gehrke, V. J. Novick and J. D. Baker, γ -ray Emission Probabilities for the ²³²U Decay Chain, Int. J. Appl. Radiat. Isot. 35(1984)581. [P_γ]
- 1992Ar05 - A. Artna-Cohen, Nuclear Data Sheets for A = 212, Nucl. Data Sheets 66(1992)171. [Nuclear structure, energies]
- 1992Li05 - W-J. Lin and G. Harbottle, Gamma -ray Emission Intensities of the ²³²Th Chain in Secular Equilibrium of ²³⁵U and the Progeny of ²³⁸U, J. Radioanal. Nucl. Chem. 157(1992)367. [P_γ]
- 1995Au04 - G. Audi and A. H. Wapstra, The 1995 Update to the Atomic Mass Evaluation, Nucl. Phys. A595(1995)409. [Q value]
- 1996Sc06 - E. Schönfeld and H. Janßen, Evaluation of Atomic Shell Data, Nucl. Instrum. Meth. Phys. Res. A369(1996)527. [X_K , X_L , Auger electrons]
- 1998ScZM - E. Schönfeld and G. Rodloff, Tables of the Energies of K -Auger Electrons for Elements with Atomic Numbers in the Range from Z = 11 to Z = 100, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX - E. Schönfeld and G. Rodloff, Energies and Relative Emission Probabilities of K X-rays for Elements with Atomic Numbers in the Range from Z = 5 to Z = 100, PTB Report PTB-6.11-1999-1, February 1999. [X_K]

**²¹²Bi – Comments on evaluation of decay data
by A. L. Nichols**

Evaluated: July/August 2001

Re-evaluated: January 2004

Evaluation Procedures

Limitation of Relative Statistical Weight Method (LWM) was applied to average numbers throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

²¹²Bi undergoes beta decay to ²¹²Po (BF = 64.07(7)%), and alpha decay to ²⁰⁸Tl (BF = 35.93(7)%). The alpha branching fraction was calculated as the weighted mean of the measurements of 1960Sc07, 1962Be09, 1962Fl03 and 1965Wa09, with the uncertainty increased to include the most precise value of 36.00(3)%.

Reference	α-decay Branching Fraction (BF) %
1960Sc07	35.96(6)
1962Be09	35.81(4)
1962Fl03	36(1)
1965Wa09	36.00(3)*
Recommended Value	35.93(7)

*Uncertainty increased slightly so that weighting does not exceed 0.5.

A reasonably consistent decay scheme has been constructed from a combination of alpha-particle studies by 1951Ry17 (two main emissions modified), 1960Wa14, and 1962Be09, and the gamma-ray measurements of 1960Sc07, 1962Be09, 1962Fl03, 1967Be19, 1968Yt02, 1972DaZA, 1978Av01, 1982Sa36, 1983Sc13, 1983Va22, 1984Ge07 and 1992Li05.

Nuclear Data

²²⁸Th decay chain is important in quantifying the environmental impact of the decay of naturally occurring ²³²Th. Specific radionuclides in this decay chain are noteworthy because of their decay characteristics (²²⁴Ra alpha decay to ²²⁰Rn; ²¹²Bi and ²⁰⁸Tl gamma-ray emissions).

Half-life

The recommended half-life is the unweighted mean of two somewhat elderly measurements (1914Le01 and 1961Ap03). Further studies are merited to determine this value with greater confidence.

Reference	Half-life (min)
1914Le01	60.480(52)
1961Ap03	60.600(43)
Recommended Value	60.54(6)

Gamma Rays

Energies

All gamma-ray transition energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies of 1986Ma 17 were adopted, and used to determine the energies and associated uncertainties of the gamma-ray transitions between the various populated-depopulated levels.

Emission Probabilities

The gamma-ray measurements of 1960Sc07, 1962Be09, 1962Fl03, 1967Be19, 1968Yt02, 1972DaZA, 1978Av01, 1982Sa36, 1983Sc13, 1983Va22, 1984Ge07 and 1992Li05 were used to determine the emission probabilities of the major gamma rays. These data have been measured relative to widely differing decay parameters: beta⁻-decay mode, alpha⁻-decay mode, per decay of ²¹²Bi (ie., absolute emission probabilities), and relative to the 583.19 and 2614.51 keV gamma rays of ²⁰⁸Tl. All of these measured data were adjusted to absolute emission probabilities when appropriate, and weighted mean values determined.

Absolute emission probabilities were estimated for the 180.2 and 1800.9 keV gamma rays in the beta⁻-decay mode, and the 433.7, 492.7, 580.5, 620.4, 759 and 807 keV gamma rays in the alpha⁻-decay mode. The latter values were derived from measurements of the low-intensity alpha-particle emission probabilities by 1960Wa14, and involved the introduction of uncertainty estimates that varied between 10% and 50% (depending on the number of significant figures quoted in the measurement of the relevant alpha emission probability).

Published Gamma-ray Emission Probabilities

E _g (keV)	P _g							
	1960Sc07	1962Be09	1962Fl03	1967Be19	1968Yt02	1972DaZA	1978Av01	1982Sa36
	*		‡	#	s	s	Δ	¶
39.858(4) [α]	-	-	-	-	-	-	-	0.9(1)
180.2(2) [β ⁻]	-	-	-	-	-	-	-	-
288.08(6) [α]	-	0.775(40) [#]	-	0.82(2)	-	0.9(2)	0.97(5)	0.32(3)
327.94(6) [α]	-	0.299(23) [#]	-	0.33(1)	-	0.36(7)	-	-
433.7(2) [α]	-		-	0.04(1)	-	~ 0.025	-	-
452.8(1) [α]	-		-	0.84(2)	-	0.88(17)	1.10(6)	0.42(5)
		1.18(5) [#]						
473.6(2) [α]	-		-	0.122(8)	-	0.10(3)	-	-
492.7(1) [α]	-		-	< 0.008	-	-	-	-
580.5(3) [α]	-	-	-	-	-	-	-	-
620.4(3) [α]	-	-	-	-	-	-	-	-
727.33(1) [β ⁻]	11.1(7)		11.8(24)	-	-	17.6(17)	21.0(8)	6.9(4)
759(1) [α]	-	100 [†]	-	-	-	-	-	-
785.37(9) [β ⁻]	1.70(26)		-	-	-	2.8(6)	3.26(16)	1.01(7)
807(1) [α]	-	-	-	-	-	-	-	-
893.41(2) [β ⁻]	0.66(7)	4.9(3) [†]	0.5(1)	-	-	0.94(19)	-	0.49(8)
952.12(2) [β ⁻]	0.16(4)	-	-	-	-	0.46(9)	-	-
1073.6(2) [β ⁻]			-	-	-	~ 0.03	-	-
	0.99(8)	10.1(4) [†]						
1078.63(11) [β ⁻]			0.7(1)	-	-	1.4(2)	-	-
1512.70(8) [β ⁻]	0.49(5)	3.4(3) [†]	-	-	0.99(15)	0.8(1)	-	-
1620.74(1) [β ⁻]	2.81(20)	20.0(6) [†]	3.0(6)	-	4.85(50)	3.9(4)	-	-
1679.45(1) [β ⁻]	-	-	-	-	0.230(7)	0.16(3)	-	-
1800.9(2) [β ⁻]				-	-	-	-	-
	0.17(3)	1.4(2) [†]	0.5(1)					
1805.96(10) [β ⁻]				-	0.41(10)	0.25(5)	-	-

Published Gamma-ray Emission Probabilities (cont.)

E _g (keV)	P _g (cont.)			
	1983Sc13 ^ψ	1983Va22 ^ψ	1984Ge07 ^Δ	1992Li05 ^ψ
39.858(4) [α]	-	-	3.49(28)	-
180.2(2) [β ⁻]	-	-	-	-
288.08(6) [α]	0.274(23)	-	1.106(10)	0.389(57)
327.94(6) [α]	0.120(4)	-	0.423(20)	3.23(12)
433.7(2) [α]	-	-	-	-
452.8(1) [α]	0.256(23)	-	1.191(11)	0.370(49)
473.6(2) [α]	-	-	-	-
492.7(1) [α]	-	-	-	-
580.5(3) [α]	-	-	-	-
620.4(3) [α]	-	-	-	-
727.33(1) [β ⁻]	6.56(15)	7.00(18)	21.63(13)	6.93(18)
759(1) [α]	-	-	-	-
785.37(9) [β ⁻]	1.07(5)	-	3.62(4)	1.05(5)
807(1) [α]	-	-	-	-
893.41(2) [β ⁻]	0.352(36)	-	1.25(6)	-
952.12(2) [β ⁻]	-	-	-	-
1073.6(2) [β ⁻]	-	-	-	-
1078.63(11) [β ⁻]	0.58(4)	-	1.85(6)	0.555(41)
1512.70(8) [β ⁻]	0.276(42)	-	-	-
1620.74(1) [β ⁻]	1.38(8)	-	4.88(10)	1.44(9)
1679.45(1) [β ⁻]	-	-	-	-
1800.9(2) [β ⁻]	-	-	-	-
1805.96(10) [β ⁻]	-	-	-	-

* Emission probabilities expressed in terms of ²¹²Bi β⁻ decay mode only.

† Emission probabilities expressed in terms of (727 + 785) keV gamma rays of ²¹²Bi.

‡ Emission probabilities relative to ²¹²Po α decay.

Emission probabilities expressed in terms of ²¹²Bi α decay mode only.

§ Emission probabilities relative to P_γ(2614.51 keV) of ²⁰⁸Tl.

Δ Emission probabilities relative to P_γ(583.19 keV) of ²⁰⁸Tl.

¶ Emission probabilities relative to P_γ(238.63 keV) of ²¹²Pb specified as 0.430(20), compared with recommended value of 0.435(4).

ψ Absolute emission probabilities.

Absolute Gamma-ray Emission Probabilities per 100 Disintegrations of ²¹²Bi

E _g (keV)	P _g ^{abs}							
	1960Sc07	1962Be09	1962Fl03	1967Be19	1968Yt02	1972DaZA	1978Av01	1982Sa36
39.858(4) [α]	-	-	-	-	-	-	-	0.9(1)
180.2(2) [β ⁻]	-	-	-	-	-	-	-	-
288.08(6) [α]	-	0.278(14)	-	0.29(1)	-	0.3(1)	0.35(2)	0.32(3)
327.94(6) [α]	-	0.107(8)	-	0.12(1)	-	0.13(3)	-	-
433.7(2) [α]	-		-	0.014(4)	-	~ 0.009	-	-
452.8(1) [α]	-		-	0.30(1)	-	0.32(6)	0.40(2)	0.42(5)
		0.424(18)						
473.6(2) [α]	-		-	0.044(3)	-	0.04(1)	-	-
492.7(1) [α]	-		-	< 0.003	-	-	-	-
580.5(3) [α]	-	-	-	-	-	-	-	-
620.4(3) [α]	-	-	-	-	-	-	-	-
727.33(1) [β ⁻]	7.11(45)		7.6(15)	-	-	6.3(6)	7.6(3)	7.0(4)
759(1) [α]	-	[7.85]	-	-	-	-	-	-
785.37(9) [β ⁻]	1.09(17)		-	-	-	1.0(2)	1.17(6)	1.02(7)
807(1) [α]	-	-	-	-	-	-	-	-
893.41(2) [β ⁻]	0.42(4)	0.38(2)	0.32(6)	-	-	0.34(7)	-	0.50(8) ^s
952.12(2) [β ⁻]	0.10(3)	-	-	-	-	0.17(3)	-	-
1073.6(2) [β ⁻]			-	-	-	~ 0.01		-
	0.63(5)	0.79(3)						
1078.63(11) [β ⁻]			0.45(6)	-	-	0.50(7)	-	-
1512.70(8) [β ⁻]	0.31(3)	0.27(2)	-	-	0.36(5)	0.29(4)	-	-
1620.74(1) [β ⁻]	1.80(13)	1.57(5)	1.9(4)	-	1.74(18)	1.4(1)	-	-
1679.45(1) [β ⁻]	-	-	-	-	0.083(3) [†]	0.06(1)	-	-
1800.9(2) [β ⁻]				-	-	-	-	-
	0.11(2)	0.11(2)	0.32(6)					
1805.96(10) [β ⁻]				-	0.15(4)	0.09(2) [†]	-	-

Absolute Gamma-ray Emission Probabilities per 100 Disintegrations of ²¹²Bi (cont.)

E _g (keV)	P _g ^{abs} (cont.)				Recommended Values*
	1983Sc13	1983Va22	1984Ge07	1992Li05	
39.858(4) [α]	-	-	1.07(9) [¶]	-	1.01(3) [†]
180.2(2) [β ⁻]	-	-	-	-	0.003(1)
288.08(6) [α]	0.274(23)	-	0.339(3) [¶]	0.389(57)	0.32(2)
327.94(6) [α]	0.120(4) [¶]	-	0.129(6)	3.23(12) ^ψ	0.121(3)
433.7(2) [α]	-	-	-	-	0.0095(20) [‡]
452.8(1) [α]	0.256(23)	-	0.365(3) [¶]	0.370(49)	0.34(3)
473.6(2) [α]	-	-	-	-	0.044(3)
492.7(1) [α]	-	-	-	-	0.04(1) [‡]
580.5(3) [α]	-	-	-	-	0.0010(2) [‡]
620.4(3) [α]	-	-	-	-	0.0038(6) [‡]
727.33(1) [β ⁻]	6.56(15)	7.00(18)	6.62(4) [¶]	6.93(18) ^ψ	6.74(12)
759(1) [α]	-	-	-	-	0.00036(18) [‡]
785.37(9) [β ⁻]	1.07(5)	-	1.11(1)	1.05(5)	1.11(1)
807(1) [α]	-	-	-	-	0.000039(4) [‡]
893.41(2) [β ⁻]	0.352(36)	-	0.383(18)	-	0.38(1)
952.12(2) [β ⁻]	-	-	-	-	0.14(4)
1073.6(2) [β ⁻]	-	-	-	-	0.015(5) [#]
1078.63(11) [β ⁻]	0.58(4)	-	0.566(18) [¶]	0.555(41)	0.55(2)
1512.70(8) [β ⁻]	0.276(42)	-	-	-	0.29(1)
1620.74(1) [β ⁻]	1.38(8)	-	1.49(3) [¶]	1.44(9)	1.51(3)
1679.45(1) [β ⁻]	-	-	-	-	0.07(1)
1800.9(2) [β ⁻]	-	-	-	-	0.004(2)
1805.96(10) [β ⁻]	-	-	-	-	0.12(3)

* Weighted mean values adopted when appropriate; remainder derived from proposed decay scheme (see other footnotes).

† Determined directly from proposed decay scheme (calculated transition probability and total theoretical internal conversion coefficient).

‡ Calculated from low-intensity alpha-particle emission probabilities of 1960Wa14.

Estimated from the approximate measurement of 1972DaZA, and used to define P_γ for 180.2 and 1800.9 keV gamma rays.

¶ Uncertainty increased so that weighting does not exceed 50%.

§ Datum rejected as outlier, and not included in weighted mean analysis.

ψ Unresolved overlap with other gamma-ray emission(s); data not included in the weighted-mean analysis.

Multipolarities and Internal Conversion Coefficients

Many of the M1 + E2 gamma transitions in the alpha -decay mode were assumed to be close to 100%M1, based on the studies of 1978Av01 and 1982Be09. Specific exceptions to this assumption include:

- 99.55 %M1 + 0.45 %E2 for 288.08keV,
- 99.2 %M1 + 0.8 %E2 for 785.37 keV,
- 99.8 %M1 + 0.2 %E2 for 893.41 keV,
- 70 %M1 + 30 %E2 for 952.12 keV,
- 98.2 %M1 + 1.8 % E2 for 1078.63 keV,
- 90 %M1 + 10 %E2 for 1620.74 keV gamma rays.

Multipolarity Assignments

Reference	E _g (keV)	Multipolarity
1978Av01	288.08(6) [α decay]	M1 + E2
	452.8(1) [α decay]	72%M1 + 28%E2
	727.33(1) [β^- decay]	E2
	785.37(9) [β^- decay]	98%M1 + 2%E2
1982Be09	785.37(9) [β^- decay]	99.2%M1 + 0.8%E2
	893.41(2) [β^- decay]	M1 (+ \leq 0.25%E2)
	952.12(2) [β^- decay]	70%M1 + 30%E2
	1078.63(11) [β^- decay]	98.2%M1 + 1.8%E2

Reasonable consistency was achieved from the proposed gamma γ -ray emission probabilities, internal conversion coefficients and alpha α -particle emission probabilities. The 39.858 keV gamma ray is particularly important in the alpha branch, and further measurements are required to determine the emission probability of this transition with greater confidence. A value of 1.01(3)% (0.0101(3)) was adopted on the basis of the relevant alpha α -particle emission probability, gamma-ray transition probability and a total internal conversion coefficient of 24.6(7).

Alpha-particle EmissionsEnergies

All alpha-particle energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies specified by 1986Ma17 and 1992Ar05, and Q-values were used to determine the energies and uncertainties of the alpha α -particle transitions to the various levels, while allowing for the significant recoil components.

Emission Probabilities

The main alpha α -particle emission probabilities emitted directly by ^{212}Bi were calculated from the evaluated gamma-ray emission probabilities (see above) and theoretical internal conversion coefficients, combined with an alpha branching fraction of 0.3593(7). These data are in excellent agreement with the measured emission probabilities of the two main alpha transitions (1951Ry17, 1960Wa14 and 1962Be09), but deviate considerable for the low γ -intensity transitions that are poorly resolved. Under such circumstances, the low γ -intensity alpha-particle data of 60Wa14 were adopted when appropriate, while others were derived from the gamma-ray studies.

Alpha-particle Emission Probabilities

E _a (keV)	P _a ^{rel}			P _a ^{abs}	
	1951Ry17	1960Wa14	1962Be09		Recommended Values*
5298(1)	0.016	0.00011(1)	-	-	5298(1)
5345(1)	0.147	0.001	-	-	5345(1)
5481.3(3)	-	0.014	~ 0.04	~ 0.02	5481.3(3)
5606.63(14)	1.08	1.19))	5606.63(14)
) 1.35(6)) 1.22(2)	
5625.4(2)	-	0.1625))	5625.4(2)
5768.27(10)	1.67	1.78	1.63(11)	1.67(2)	5768.27(10)
6050.92(4)	69.86 [#]	69.7	70.2(3)	70.2(2)	6050.92(4)
6090.02(4)	27.16 [#]	27.1	27.0(5)	26.8(2)	6090.02(4)
9498.79(12) [†]	-	-	-	-	9498.79(12) [†]
10432.95(12) [†]	-	-	-	-	10432.95(12) [†]
10552.1(3) [†]	-	-	-	-	10552.1(3) [†]

* Recommended emission probabilities derived from evaluated gamma-ray emission probabilities, theoretical internal conversion coefficients and alpha branching fraction of 0.3593(7), unless stated otherwise (expressed per 100 disintegrations of ^{212}Bi).

‡ Data reported by 1960Wa14 were adopted and adjusted for alpha branch; uncertainties were estimated when not quoted.

† Arises from $\beta^- \alpha$ decay (long-range alpha particles).

Data reported incorrectly; re-assigned by evaluator.

Alpha-particle emissions from the $\beta^- \alpha$ decay mode have been observed at energies greater than 9 MeV by 1951Ry17, 1962Be09 and 1965Le08. Some of the excited states of ^{212}Po populated by the beta of ^{212}Bi undergo subsequent alpha decay (in competition with the gamma γ -ray decay). These nuclear levels at 1800.9, 1679.45 and 727.33 keV emit high γ -energy alpha particles (energies of 10552.1, 10432.95 and 9498.79 keV, respectively). All measurements were expressed relative to 10^6 emission probability for the 8785.18 keV alpha particle of ^{212}Po , but with no quoted uncertainties. These long γ -range alpha particles constitute part of the ^{212}Bi decay; and their emission probabilities were determined from the measurements of 1951Ry17, 1962Be09 and 1965Le08:

Alpha-particle Emissions ($\beta^- \alpha$ Decay)

E_a (keV)	P_a^{rel}			
	1951Ry17	1962Be09	1965Le08	Mean Value
[8785.18(11)]*	10^6	10^6	10^6	10^6
9498.79(12)	35	45	34	38
10432.95(13)	20	17	10	16
10552.1(3)	170	167	160	166
Total α (of $\beta^- \alpha$)	225	229	204	219(15)

* ^{212}Po alpha decay.

Total α emissions from $\beta^- \alpha$ decay have an estimated mean value of 219 relative to 10^6 for the emission probability of the 8785.18 keV alpha particle of ^{212}Po , with an uncertainty of 15 to cover the range of measured data. Therefore, a mean value of 0.00014 was estimated for the $\beta^- \alpha$ branching fraction, combined with an uncertainty of approximately 7% ($\text{BF}(\beta^- \alpha) = 0.00014(1)$). Absolute alpha α -particle emission probabilities for this small branch were calculated from the mean values and $\text{BF}(\beta^- \alpha)$.

Beta-particle Emissions

Energies

All beta-particle energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies of 1992Ar05 and the Q-value were used to determine the energies and uncertainties of the beta-particle transitions to the various levels.

Emission Probabilities

The beta-particle emission probabilities were calculated from gamma γ -ray transition intensity balances, using the recommended gamma γ -ray emission probabilities and the theoretical internal conversion coefficients of 1978Ro22.

Beta-particle Emission Probabilities

E_b (keV)	P_b	
	1957Bu34	Recommended Values*
448(2)	8.5	0.68(5)
453(2)	-	0.029(1)
575(2)	-	0.21(5)
633(2)	6	1.90(4)
741(2)	-	1.45(2)
1527(2)	10	4.58(21)
2254(2)	63	55.23(21)

* Recommended emission probabilities derived from evaluated gamma-ray emission probabilities, theoretical internal conversion coefficients, beta branching fraction of 64.06(7) % and beta-alpha branching fraction of 0.00014(1) (expressed per 100 disintegrations of ^{212}Bi).

Atomic Data

The x-ray data have been calculated using the evaluated gamma γ -ray data, and the atomic data from 1996Sc06, 1998ScZM and 1999ScZX.

References

- 1914Le01 - F. V. Lerch, Verdampfungserscheinungen der ThB - und ThC -Verbindungen, Sitzber. Akad. Wiss., Wien, Math.-Naturw. Kl. Abt. Ila 123(1914)699. [Half-life]
- 1951Ry17 - A. Rytz, Nouvelles Expériences sur le Spectre Magnétique Alpha du Thorium C et des Longs Parcours du Thorium C', C. R. Acad. Sci. Paris 233(1951)790. [E_{α} , P_{α}]
- 1957Bu34 - J. Burde and B. Rozner, Beta Spectrum of Bi²¹² (ThC), Phys. Rev. 107(1957)531. [P_{β}]
- 1960Em01 - G. T. Emery and W. R. Kane, Gamma -ray Intensities in the Thorium Active Deposit, Phys. Rev. 118(1960)755. [P_{γ} , high-energy α]
- 1960Sc07 - G. Schupp, H. Daniel, G. W. Eakins and E. N. Jensen, Transition Intensities in the Tl²⁰⁸ Beta Decay, the Bi²¹² \rightarrow Po²¹² Decay Scheme, and the Bi²¹² Branching Ratio, Phys. Rev. 120(1960)189. [P_{γ} , BF]
- 1960Wa14 - R. J. Walen and G. Bastin-Scoffier, Spectre α du ²¹⁴Bi et Remarques sur Quelques Émetteurs α du Bi, Nucl. Phys. 16(1960)246. [P_{α}]
- 1961Ap03 - K. R. Applegate, E. M. Morimoto, M. Kahr and J. D. Knight, Redetermination of the Half -Life of Bismuth (ThC), J. Inorg. Nucl. Chem. 19(1961)375. [Half-life]
- 1962Be09 - G. Bertolini, F. Cappellani, G. Restelli and A. Rota, Excited States of Tl -208 and Po-212, Nucl. Phys. 30(1962)599. [P_{α} , P_{γ} , BF]
- 1962Fl03 - F. C. Flack and J. E. Johnson, The Gamma Radiation from ²¹²Po (ThC'), Proc. Phys. Soc. (London) 79(1962)10. [P_{γ} , BF]
- 1965Le08 - C-F. Leang, Spectres α de Long Parcours des Poloniums 214 (RaC') et 212 (ThC'), C. R. Acad. Sci. Paris 260(1965)3037. [high-energy α]
- 1965Wa09 - J. Walker and T. Salgir, The Branching Ratio of ²¹²Bi (ThC), Proc. Phys. Soc. (London) 86(1965)423. [BF]
- 1967Be19 - R. Benoit, G. Bertolini, F. Cappellani and G. Restelli, Decay of the Excited Levels of ²⁰⁸Tl, Nuovo Cimento 49B(1967)125. [P_{γ}]
- 1968Yt02 - C. Ythier, H. Forest, G. Ardisson and H. Maria, Sur le Rayonnement γ de Haute Énergie Accompagnant la Désintégration du Bismuth-212, C. R. Acad. Sci. Paris Series B 267(1968)1362. [P_{γ}]
- 1972DaZA - J. Dalmasso, Recherches sur le Rayonnement Gamma de Quelques Radioéléments Naturels Appartenant à la Famille du Thorium, PhD thesis, University of Nice (1972); J. Dalmasso, H. Maria and C. Ythier, Étude du Rayonnement γ du Thorium 228 et de ses Dérivés, et plus Particulièrement du Thallium 208 (ThC"), C. R. Acad. Sci. Paris 277B(1973)467. [P_{γ}]
- 1978Av01 - F. T. Avignone and A. G. Schmidt, γ -ray and Internal-conversion Intensity Studies of Transitions in the Decay of ²²⁸Th, Phys. Rev. C17(1978)380. [P_{γ} , multipolarity]
- 1978Ro22 - F. Rösel, H. M. Fries, K. Alder and H. C. Pauli, Internal Conversion Coefficients for all Atomic Shells, ICC Values for Z = 68-104, At. Data Nucl. Data Tables 21(1978)291-514. [ICC]
- 1982Be09 - B. Bengtson, H. L. Nielsen, N. Rud and K. Wilsky, Half-life of the α -emitting Excited 0⁺ State in ²¹²Po, Nucl. Phys. A378(1982)1. [Multipolarity]
- 1982Sa36 - S. Sadasivan and V. M. Raghunath, Intensities of Gamma Rays in the ²³²Th Decay Chain, Nucl. Instrum. Meth. 196(1982)561. [P_{γ}]
- 1983Sc13 - U. Schötzgig and K. Debertin, Photon Emission Probabilities per Decay of ²²⁶Ra and ²³²Th in Equilibrium with their Daughter Products, Int. J. Appl. Radiat. Isot. 34(1983)533. [P_{γ}]
- 1983Va22 - R. Vaninbrouckx and H. H. Hansen, Determination of γ -ray Emission Probabilities in the Decay of ²²⁸Th and its Daughters, Int. J. Appl. Radiat. Isot. 34(1983)1395. [P_{γ}]
- 1984Ge07 - R. J. Gehrke, V. J. Novick and J. D. Baker, γ -ray Emission Probabilities for the ²³²U Decay Chain, Int. J. Appl. Radiat. Isot. 35(1984)581. [P_{γ}]
- 1986Ma17 - M. J. Martin, Nuclear Data Sheets for A = 208, Nucl. Data Sheets 47(1986)797. [Nuclear structure, energies]
- 1992Ar05 - A. Artna-Cohen, Nuclear Data Sheets for A = 212, Nucl. Data Sheets 66(1992)171. [Nuclear structure, energies]
- 1992Li05 - W-J. Lin and G. Harbottle, Gamma -ray Emission Intensities of the ²³²Th Chain in Secular Equilibrium, of ²³⁵U and the Progeny of ²³⁸U, J. Radioanal. Nucl. Chem. 157(1992)367. [P_{γ}]
- 1995Au04 - G. Audi and A. H. Wapstra, The 1995 Update to the Atomic Mass Evaluation, Nucl. Phys. A595(1995)409. [Q values]
- 1996Sc06 - E. Schönfeld and H. Janßen, Evaluation of Atomic Shell Data, Nucl. Instrum. Meth. Phys. Res. A369(1996)527. [X_K , X_L , Auger electrons]
- 1998ScZM - E. Schönfeld and G. Rodloff, Tables of the Energies of K-Augur Electrons for Elements with Atomic Numbers in the Range from Z = 11 to Z = 100, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX - E. Schönfeld and G. Rodloff, Energies and Relative Emission Probabilities of K X-rays for Elements with Atomic Numbers in the Range from Z = 5 to Z = 100, PTB Report PTB-6.11-1999-1, February 1999. [X_K]

²¹²Po – Comments on evaluation of decay data by A. L. Nichols

Evaluated: July/August 2001

Re-evaluated: January 2004

Evaluation Procedures

Limitation of Relative Statistical Weight Method (LWM) was applied to average numbers throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

²¹²Po is an extremely short-lived radionuclide populated via the beta decay of ²¹²Bi and the alpha decay of ²¹⁶Rn. Alpha decay of ²¹²Po occurs directly to the ground state of ²⁰⁸Pb.

Nuclear Data

Half-life

Po-212 is an extremely short-lived radionuclide populated primarily via the alpha decay of Rn -216 and the beta decay of Bi-212. The recommended half-life of $3.00(2) \times 10^{-7}$ sec is based on the weighted mean of five sets of measurements (1949Bu09, 1962F103, 1963As02, 1972Mc29 and 1975Sa06).

Reference	Half-life (s)
1949Bu09	$3.04(4) \times 10^{-7}$
1962F103	$3.05(25) \times 10^{-7}$
1963As02	$3.05(5) \times 10^{-7}$
1972Mc29	$3.04(8) \times 10^{-7}$
	$3.00(8) \times 10^{-7}$
1975Sa06	$2.96(2) \times 10^{-7*}$
Recommended Value	$3.00(2) \times 10^{-7}$

* Uncertainty adjusted to $\pm 0.03 \times 10^{-7}$ to reduce weighting below 0.5.

Alpha-particle Emission

Energy

The Q-value of 1995Au04 was used to determine the energy and uncertainty of the single alpha-particle transition to the ground state of ²⁰⁸Pb, while allowing for the significant recoil component. Thus, an alpha-particle energy of 8785.18(11) keV was calculated.

Emission Probability

The emission probability of the single alpha particle was defined as 100% (1.00).

Alpha-particle Emission Probabilities per 100 Disintegrations of ²¹²Po

E_a(keV)	P_a
	Recommended Value*
8785.18(11)	100.0

* Only one α transition directly to the ground state of ²⁰⁸Pb.

References

- 1949Bu09 - D. E. Bunyan, A. Lundby and W. Walker, Experiments with the Delayed Coincidence Method, Including a Search for Short-Lived Nuclear Isomers, Proc. Phys. Soc. (London) 62A(1949)253. [Half-life]
- 1962Fl03 - F. C. Flack and J. E. Johnson, The Gamma Radiation from ²¹²Po (ThC), Proc. Phys. Soc. (London) 79(1962)10. [Half-life]
- 1963As02 - G. Astner, I. Bergstrom, L. Eriksson, U. Fagerquist, G. Holm and A. Persson, A Hindered E2 Ground State Transition in Po²⁰⁷, Nucl. Phys. 45(1963)49. [Half-life]
- 1972Mc29 - G. W. McBeth and R.A. Winyard, Isotope Identification and Radioassay by Time Interval Analysis, Int. J. Appl. Radiat. Isotop. 23(1972)527. [Half-life]
- 1975Sa06 - S. Sanyal, R. K. Garg, S. D. Chauhan, S. L. Gupta and S. C. Panchol i, Half-Life Measurement of the ²¹²Po Ground State, Phys. Rev. C12(1975)318. [Half-life]
- 1995Au04 - G. Audi and A. H. Wapstra, The 1995 Update to the Atomic Mass Evaluation, Nucl. Phys. A595(1995)409. [Q value]

²¹³Bi - Comments on evaluation of the decay data

by Huang Xiaolong, Wang Baosong

This evaluation was completed in 2006. Literature available by January 2006 was included.

1. Decay Scheme

²¹³Bi disintegrates 97.91 (3) % by β^- emission to levels in ²¹³Po and 2.09 (3) % through α decay to ²⁰⁹Tl. ²¹³Bi ground state has $J^\pi = 9/2^-$ (1992Ak01).

The ²¹³Bi β^- decay scheme was built from the γ - γ coincidence measurements of 1998Ar03 and 2000Gr35. The ²¹³Bi α decay scheme was built from the α - γ coincidence measurements of 1964Gr11, the singles α -particle measurements of 1997Ch53 and γ - γ coincidence measurements of 1998Ar03.

The decay branching ratios have been deduced by the evaluator using the absolute photon intensity (96.58 (10), 1991Ma16) adopted for the 465 keV γ -ray from ²⁰⁹Tl β^- decay and measured absolute intensity (2.022 (26), 1986He06) of the same γ -ray following the ²¹³Bi α decay. Our recommended α decay branching ratio is $I_\alpha = 2.09$ (3) %, thus $I_{\beta^-} = 97.91$ (3) %.

The three values of the ²¹³Bi α decay branching ratio found in the literature are presented in Table 1. The corresponding β^- branching ratios are: $I_{\beta^-} = 97.84$ (11) %, (deduced by 1964Gr11); $I_{\beta^-} = 97.91$ (3) %, (deduced from the measurements of 1986He06); $I_{\beta^-} = 97.80$ (3) %, measured in equilibrium with ²¹³Po by 1997Ch53.

Table 1: Measured and recommended branching ratio for ²¹³Bi α decay.

I_α (%)	References	Comments
2.16 (11)	1964Gr11	Deduced from measured I_α
2.09 (3)	1986He06	Deduced from the $P_\gamma(465 \text{ keV})$ from ²⁰⁹ Tl following ²¹³ Bi α decay and measured value by 1986He06
2.20 (3)	1997Ch53	Measured in equilibrium with ²¹³ Po
2.15 (4)		LWM
2.09 (3)		Recommended

The recommended $Q(\alpha)$ value of 5983 (6) keV and $Q(\beta^-)$ values of 1423 (5) keV in Audi(2003Au03) agrees with the $Q(\alpha)$ value of 5979 (2) keV and $Q(\beta^-)$ values of 1422 (6) keV, calculated by the evaluator (using program RADLST) from average radiation energies. This agreement supports the completeness and correctness of the decay scheme.

2. Nuclear Data

The Q values are from the 2003Au03 evaluation.

Level energies have been obtained from a least-squares fit to γ -ray energies (GTOL computer code). Spin and parities are from 1992Ak01, 1998Ar03 and 2000Gr35.

The measured and recommended ²¹³Bi half-life values are listed in Table 2.

Table 2: Measured half-life values of ²¹³Bi and recommended value

$T_{1/2}$ (min)	References	Measurement method
46	1947En03	
47 (1)	1950Ha52	Alpha pulse analyzer, 9 $T_{1/2}$
46 (1)	1964Gr11	
45.59 (6)	1973Po16	ZnS(Ag), weighted average of 2 sources, 8 $T_{1/2}$
46.2 (4)		Unweighted mean (except 1947En03)
45.60 (6)		LWM (except 1947En03), $\chi^2 = 1.07$
45.59 (6)	Recommended value	From 1973Po16

The half-life weighted average has been calculated by LWM program. The recommended value is taken from the measurement of 1973Po16.

2.1 β^- transitions

The maximum energies of the β^- transitions in the decay of ²¹³Bi have been deduced from the Q value (2003Au03) and the level energies.

The adopted β^- transition probabilities and the associated uncertainties were deduced from the γ -ray transition probability balance at each level of the decay scheme. Measured and adopted β^- transition probabilities are given in table 3.

Table 3: Measured and adopted probabilities (%) of β^- transitions

Level energy (keV)	1955Ma61	1968Va17	Adopted value
0		65 (3)	66.2 (4)
292.8			0.21 (9)
440.4	32	35 (3)	30.8 (4)
600.8			0.002 5 (19)
868			0.012 9 (6)
1003.6			0.064 8 (23)
1045.6			0.020 (4)
1100.2			0.595 (17)
1119.4			0.060 8 (20)
1328.2			0.0014 (2)

The values of $lg ft$ and average β^- energies have been calculated with the program LOGFT.

2.2 γ -ray Transitions

The γ -ray transition probabilities were calculated using the γ -ray emission intensities and the relevant internal conversion coefficients.

Multipolarities and mixing ratios of γ -ray transitions are from 2000Gr35, 1977Vy02 and 1969DzZZ.

The internal conversion coefficients (ICC) and their associated uncertainties for γ -ray transitions have been obtained using the BrIcc computer program, which uses the ‘‘Frozen Orbital’’ approximation (2002Ba85). Experimental and theoretical conversion coefficients are compared in Table 4.

Table 4: Comparison of the calculated and measured conversion coefficients

E_γ (keV)	Multipolarity	Mixing ratio	α (theory)	α (exp.) 2000Gr35
147.7	E2		$\alpha_T = 1.453, \alpha_K = 0.31, \alpha_L = 0.85$	$\alpha_K = 0.33$ (14)
292.8	M1+E2	1.2 (+11 -8)	$\alpha_T = 0.3, \alpha_K = 0.22, \alpha_L = 0.06$	$\alpha_K = 0.20$ (5)
323.7	E2+M1	1.26 (16)	$\alpha_T = 0.178, \alpha_K = 0.134, \alpha_L = 0.03$	$\alpha_K = 0.131$ (15)
440.44	M1		$\alpha_T = 0.179, \alpha_K = 0.146$	$\alpha_K = 0.13$ (2)

2.3 α Transitions

1997Ch53 measured the upper limit for intensity of the E_α (868 keV level) = 5018 keV, and the value is $< 10^{-4}$. But this measurement did not support the assumption that the 868 keV level is excited in ²⁰⁹Tl by the ²¹³Bi α decay. Thus the 868 keV level is not considered here.

Measured energies of alpha particles are listed in table 5. The measured values are in good agreement with the calculated results from Q_α (2003Au03) and the level energies. Our recommended values are from 1964Gr11.

Table 5: Measured and recommended values of α -particle energies (in keV) from ²¹³Bi α decay

1947En02	1964Gr11	1967Dz02	Deduced	Recommended
	5549 (10)		5553 (6)	5549 (10)
5860 (30)	5869 (10)	5870 (6)	5871 (6)	5869 (10)

Experimental and recommended α -particle relative intensities to 100 % α decay are listed in Table 6. Our recommended α -particle relative intensities are deduced from the calculated results of the γ transition probability balance. These calculated results are in good agreement with the measured relative intensities of 1964Gr11 and 1997Ch53.

Table 6: Experimental, recommended α -particle relative intensities to 100 % α decay

E_α (keV)	I_α (%)		
	1964Gr11	1997Ch53	Recommended
5549 (10)	7.4 (14)	6.8 (1)	8.9 (2)
5869 (10)	92.6 (14)	93.2 (14)	91.1 (14)

The recommended α -particle emission intensities are the relative intensities values recommended in table 6 multiplied by 0.0209 (3).

3. Atomic data

Atomic fluorescence yields ($\omega_K, \omega_L, \omega_M, \eta_{KL}$ and η_{LM}) are from Schönfeld (1996Sc06).

The X-ray and Auger electron emission intensities have been deduced from γ -ray and conversion electron data by using the computer code RADLST. Measured and calculated X-ray emission intensities are compared in Table 7.

Table 7: Comparison of the calculated and measured X-ray emission intensities

	1972Dz14	Adopted (deduced)
$K_{\alpha 1}$	1.6 (2)	1.6 (3)
$K_{\alpha 2}$	0.93 (12)	0.99 (15)

The deduced KX-ray emission intensities agree with the measured value of 1972Dz14, thus confirming the completeness of the decay scheme.

4. Electron Emissions.

The conversion electron emission probabilities have been deduced from γ -ray transition data.

5. Photon Emissions

5.1 γ -ray energy values

The measurements of the γ -ray energy value of ²¹³Bi are listed in Table 8 associated with their LWM average value. The recommended values are taken from the LWM of the measurements of 1977Vy02, 1981Di14, 1989Ko26, 1998Ar03 and 2000Gr35, or from 1998Ar03 in the case of only one energy measurement (402.8 keV, 884.6 keV, 886.66 keV, 897.0 keV and 1328.2 keV).

Table 8: Measured and recommended values of γ -ray energy (in keV) for ²¹³Bi

1977Vy02	1981Di14	1989Ko26	1998Ar03	2000Gr35	Recommended
		147.63 (8)	147.66 (5)	147.7 (1)	147.70 (4)
292.86 (10)	292.85 (2)	292.80 (1)	292.76 (5)	292.81 (1)	292.80 (1)
323.81 (5)	323.7 (2)	323.71 (3)	323.69 (5)	323.80 (4)	323.70 (2)
			402.8 (3)		402.8 (3)
440.42 (2)	440.4 (2)	440.46 (1)	440.43 (5)	440.44 (1)	440.44 (1)
			574.8 (3)	575.2 (5)	574.9 (3)
			600.7 (3)	601.0 (2)	600.9 (2)
			604.9 (3)	604.94 (21)	604.93 (17)
			646.03 (9)	646.0 (1)	646.0 (1)
659.81 (10)	659.7 (2)	659.8 (1)	659.77 (5)	659.74 (2)	659.75 (2)
		710.8 (1)	710.81 (21)	710.82 (3)	710.82 (3)
807.36 (4)	807.3 (2)	807.36 (1)	807.38 (5)	807.37 (1)	807.37 (1)
		826.8 (2)	826.47 (6)	826.59 (5)	826.55 (4)

1977Vy02	1981Di14	1989Ko26	1998Ar03	2000Gr35	Recommended
		868.0 (2)	867.98 (3)	867.93 (3)	867.96 (2)
			880.2 (3)	880.91 (1)	880.91 (1)
			884.6 (3)		884.6 (3)
			886.66 (14)		886.66 (14)
			897.0 (3)		897.0 (3)
		1003.57 (3)	1003.55 (5)	1003.59 (3)	1003.58 (2)
			1045.70 (9)	1045.10 (40)	1045.67 (8)
1100.14 (6)	1100.1 (2)	1100.16 (2)	1100.12 (5)	1100.18 (2)	1100.16 (1)
1119.60 (14)		1119.4 (1)	1119.29 (5)	1119.50 (4)	1119.42 (8)
			1328.2 (3)		1328.2 (3)

5.2 Relative values of the γ -ray intensities

The measurements of the relative γ -ray intensities of ²¹³Bi α decay and β^- decay are listed in table 9 and table 10, respectively.

For α decay, the recommended values are taken from the LWM average of the measurements of 1989Ko26, 1998Ar03 and 2000Gr35. For β^- decay, the recommended values are taken from the LWM average of the measurements of 1986He06, 1989Ko26, 1998Ar03 and 2000Gr35 (according to the availability of the reported data).

Table 9: Measured and recommended relative γ -ray intensities for ²¹³Bi α decay
(the intensity of the 440.44 keV γ -ray is considered 100)

E_γ (keV)	I_γ						
	1969ArZV	1977Vy02	1981Di14	1989Ko26	1998Ar03	2000Gr35	Recommended
323.70(2)	0.67 (10)	1.12 (8)	0.660 (15)	0.619 (37)	0.567 (46)	0.618 (32)	0.607 (2)

Table 10: Measured and recommended relative γ -ray intensities for ²¹³Bi γ -decay

E (keV)	I_γ							
	1969ArZV	1977Vy02	1981Di14	1986He06	1989Ko26	1998Ar03	2000Gr35	Recommended
147.70 (4)					0.0429 (43)	0.0567 (46)	0.087 (32)	0.049 (3)
292.80 (1)	1.81 (14)	2.65 (38)	1.555 (87)	1.644 (27)	1.571 (63)	1.594 (88)	1.58 (4)	1.613 (20)
402.8 (3)						0.00038 (12)		0.00038 (12)
440.44 (1)	100	100	100	100	100	100	100	100
574.9 (3)						0.00241 (65)	0.0099 (39)	0.0026 (6)
600.9 (2)						0.00268 (84)	0.0165 (32)	0.010 (7)
604.93 (17)						0.00192 (69)	0.0091 (24)	0.0055 (17)
646.0 (1) ^x						0.00885 (84)	0.0095 (39)	0.0089 (8)
659.75 (2)	0.19 (10)	0.53 (5)	0.185 (6)		0.1476 (89)	0.1383 (77)	0.173 (12)	0.165 (20)
710.82 (3)					0.0429 (43)	0.0391 (42)	0.0469 (39)	0.043 (2)
807.37 (1)	1.14 (14)	1.59 (5)	1.152 (26)	1.119 (46)	1.048 (42)	0.923 (57)	1.114 (71)	1.10 (5)
826.55 (4)					0.0271 (16)	0.0218 (19)	0.0303 (51)	0.0249 (14)
867.96 (2)					0.0476 (29)	0.0425 (42)	0.0484 (43)	0.0467 (21)
880.91 (1) ^x						0.0111 (38)	0.0165 (16)	0.015 (2)
884.6 (3) ^x						0.00111 (38)		0.0011 (4)
886.66 (14)						0.00391 (73)		0.0039 (7)
897.0 (3) ^x						0.00119 (35)		0.0012 (4)
1003.58 (2)					0.205 (12)	0.192 (19)	0.209 (12)	0.205 (8)
1045.67 (8)						0.069 (12)	0.134 (75)	0.071 (12)
1100.16 (1)	1.05 (14)	1.71 (8)	1.000 (24)		1.095 (44)	0.992 (61)	0.988 (67)	1.016 (19)
1119.42 (8)		0.214 (25)			0.238 (14)	0.192 (12)	0.201 (12)	0.208 (7)
1328.2 (3)						0.0015 (5)		0.0015 (5)

^x: not placed in level scheme.

5.3 Absolute values of the γ -ray emission probabilities

There is only one measurement of the absolute γ -ray emission probability of the 440.44 keV from ²¹³Bi β^- decay which was measured in equilibrium with ²²⁹Th by 1986He06 in 1986. This measurement can be adopted as the normalization factor N, that is, $N = 0.261$ (3).

The evaluated absolute γ -ray emission probabilities are the relative values evaluated in table 9 and table 10 multiplied by 0.261 (3).

6. References

- 1947En03 A.C.English, T.E.Cranshaw, P.Demers, J.A.Harvey, E.P.Hincks, J.V.Jelley, A.N.May, Phys.Rev. 72, 253 (1947) [$T_{1/2}$].
- 1950Ha52 F.Hagemann, L.I.Katzin, M.H.Studier, G.T.Seaborg, A.Ghiorso, Phys.Rev. 79, 435 (1950) [$T_{1/2}$].
- 1955Ma61 L.B.Magnusson, F.Wagner, Jr., D.W.Engelkemeir, M.S.Freedman, ANL-5386 (1955) [Multipolarity].
- 1964Gr11 G.Graeffe, K.Valli, J.Aaltonen, Ann.Acad.Sci.Fenn., Ser.A VI, No.145 (1964) [$E\alpha$, $I\alpha$].
- 1969ArZV R.Arlt, B.S.Dzhelepov, R.B.Ivanov, M.A.Mikhailova, L.N.Moskvin, V.O.Sergeev, L.G.Tsaritsyna, K.Shtrusnyi, B.S.Dzhelepov, Program and Theses, Proc. 19th Ann. Conf. Nucl. Spectroscopy and Struct. of At. Nuclei, Erevan, p.152 (1969) [$E\gamma$, $I\gamma$].
- 1969DzZZ B.S.Dzhelepov, A.V.Zolotavin, R.B.Ivanov, M.A.Mikhailova, V.O.Sergeev, M.I.Sovtsov, O.M.Shumilo, Program and Theses, Proc.19th Ann. Conf. Nucl. Spectroscopy and Struct. Of At. Nuclei, Erevan, p.153 (1969) [Multipolarity].
- 1972Dz14 B.S.Dzhelepov, R.B.Ivanov, Bull.Acad.Sci.USSR, Phys.Ser. 36, 1832 (1973) [X-ray intensities]
- 1973Po16 P.Polak, Radiochim.Acta 19, 148 (1973) [$T_{1/2}$].
- 1977Vy02 T.Vylov, N.A.Golovkov, B.S.Dzhelepov, R.B.Ivanov, M.A.Mikhailova, Y.V.Norseev, V.G.Chumin, Bull.Acad.Sci.USSR, Phys.Ser. 41, No.8, 85 (1977) [$E\gamma$, $I\gamma$].
- 1981Di14 J.K.Dickens, J.W.McConnell, Radiochem.Radioanal.Lett. 47, 331 (1981) [$E\gamma$, $I\gamma$].
- 1986He06 R.G.Helmer, C.W.Reich, M.A.Lee, I.Ahmad, Int.J.Appl.Radiat.Isotop. 37, 139(1986) [$E\gamma$, $I\gamma$, $P\gamma$].
- 1989Ko26 M.C.Kouassi, A.Hachem, C.Ardisson, G.Ardisson, Nucl.Instrum. Methods Phys.Res. A280, 424 (1989) [$E\gamma$, $I\gamma$].
- 1991Ma16 M.J.Martin, Nucl.Data Sheets 63, 723 (1991) [NDS].
- 1992Ak01 Y.A.Akovali, Nucl.Data Sheets 66, 237 (1992) [NDS].
- 1996Sc06 E.Schönfeld, H.Janssen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
- 1997Ch53 V.G.Chumin, J.K.Jabber, K.V.Kalyapkin, S.A.Kudrya, V.V.Tsupko-Sitnikov, K.Ya.Gromov, V.I.Fominykh, T.A.Furyaev, Bull.Rus.Acad.Sci.Phys. 61, 1606 (1997) [$E\alpha$, $I\alpha$].
- 1998AR03 G.Ardisson, V.Barci, O.El Samad, Phys.Rev. C57, 612 (1998) [$E\gamma$, $I\gamma$].
- 2000GR35 K.Ya.Gromov, S.A.Kudrya, Sh.R.Malikov, T.M.Muminov, Zh.K.Samatov, Zh.Sereeter, V.I.Fominykh, V.G.Chumin, Bull.Rus.Acad.Sci.Phys. 64, 1770 (2000) [$E\gamma$, $I\gamma$].
- 2002Ba85 I.M.Band, M.B.Trzhaskovskaya, C.W.Nestor, Jr., P.O.Tikkanen, S.Raman, At.Data Nucl.Data Tables 81, 1 (2002) [calculated ICC]
- 2003Au03 G.Audi, A.H.Wapstra, C.Thibault, Nucl. Phys. A729(2003)129 [Q].

**²¹³Po - Comments on evaluation of the decay data
by Huang Xiaolong, Wang Baosong**

This evaluation was completed in 2007. Literature available by December 2007 was included.

1 Decay Scheme

²¹³Po disintegrates 100 % by α emissions to levels in ²⁰⁹Pb. ²¹³Po ground state has $J^\pi = 9/2^+$ (2007Ba19).

2 Nuclear Data

The Q value is from the 2003Au03 evaluation.

The level energies, spin and parities are from 2007Ba19.

The measured and evaluated ²¹³Po half-life values are listed in Table 1.

Table 1 - Measured half-life values of ²¹³Po and evaluated value, in μ s.

$T_{1/2}$ (μ s)	References	measurement method
4.2 (8)	1948Je05	
3.74 (2)	1995WaZQ	Superseded by 1998Wa25
3.70 (3)	1997VaZV	Superseded by 1998Wa25
3.75 (4)	1997Wa27	Si(Au), delayed β - α coincidences
3.65 (4)	1998Wa25	Three-dimensional single-crystal scintillation time spectrometer
3.65	2002Mo46	HPGe and 4π autocorrelation single-crystal scintillation time spectrometer. No uncertainty given
3.70 (5)		Unweighted mean of 1997Wa27 and 1998Wa25
3.70 (5)		Weighted mean of 1997Wa27 and 1998Wa25, $\chi^2=3.1$
3.70 (5)	Recommended value	

Values given by 1995 WaZQ, 1997VaZV, 1997Wa27, and 1998 Wa25 have authors in common, thus, they may not be independent of each other. A recommended value of 3.70 (5) μ s has been estimated by the evaluator.

2.1 g Transitions

The γ -ray transition probability is calculated using the γ -ray emission intensity and the relevant internal conversion coefficient.

Multipolarity of 778.8 keV γ -ray is from level scheme (not measured).

The internal conversion coefficient (ICC) and their associated uncertainties for γ -ray transitions have been obtained using the BRICC computer program, which uses the "Frozen Orbital" approximation (2002Ba85)

2.2 a Transitions

Measured and recommended alpha particles energies are listed in table 2. The recommended values are from 1964Va20 and 1991Ry01.

Table 2 - Measured and recommended value of α -particle energy from ²¹³Po decay

1964Va20	1982Bo04 ^a	1991Ry01 ^b	Recommended value
7614 (10)			7614 (10)
8377 (5)	8376 (3)	8375.9 (25)	8375.9 (25)

^a: Original energies of 1982Bo04 have been increased by 2 keV due to changes in calibration energies (1991Ry01).

^b: evaluation.

The measured and recommended alpha particle emission probabilities are listed in table 3. The recommended alpha particle emission probabilities have been deduced from γ -ray transition intensity balance.

Table 3 - Measured and recommended α -particle emission probabilities from ²¹³Po decay

E_α (keV)	P_α			
	1964Va20	1969LeZW	1997Ch53	Recommended
7614 (10)	0.003 (1)	0.006 (2)	0.0031 (2)	0.0050 (5)
8375.9 (25)	100	100	99.997 (31)	99.9950 (5)

$P_\alpha = 0.0031$ (2) % in 1997Ch53 is from an α -particle spectrum. This very weak peak is at the low-energy tail of the intense 8376-keV α -particle group. Thus, the evaluator has considered its reported intensity to be quite inaccurate, despite the value reported in 1997Ch53.

3. Photon Emissions

There is only one γ -ray emitted from ²¹³Po α decay. Only 1989Ko26 measured the γ -ray energy: 778.8 (3) keV. The present recommended γ -ray energy has been taken from this measurement.

The recommended absolute γ -ray emission probability has been obtained as follows: 1989Ko26 measured the ratio: $I_\gamma(779 \text{ keV}) / I_\gamma(440 \text{ keV})$ (in ²¹³Bi β^- decay) = 0.000181 (18). Using $P_\gamma(440 \text{ keV}) = 26.1$ (3) % and $\% \beta^- = 0.9791$ (3) (2007HuXX) then $P_\gamma(778 \text{ keV}) = 0.0048$ (5) %.

4. References

- 1948Je05 J.V.Jelley, Can.J.Res. 26A, 255 (1948) [$T_{1/2}$].
 1964Va20 K.Valli, Ann.Acad.Sci.Fennicae, Ser.A VI, No.165 (1964) [E_α].
 1969LeZW C.-F.Leang, Thesis, Univ.Paris (1969) [E_α].
 1982Bo04 J.D.Bowman, R.E.Eppley, E.K.Hyde, Phys.Rev. C25, 941 (1982) [E_α].
 1989Ko26 M.C.Kouassi, A.Hachem, C.Ardisson, G.Ardisson, Nucl.Instrum.Methods Phys.Res. A280, 424 (1989) [E_γ , I_γ].
 1991Ry01 A.Rytz, At.Data Nucl.Data Tables 47, 205 (1991) [Evaluation]
 1995WaZQ J. Wawryszczuk, M.B.Yuldashev, K.Ya. Gromov, T.M. Mumino v, Program and Thesis, Proc.45th Ann. Conf. Nucl.Spectrosc.Struct.At.Nuclei, St.Petersburg, p.107 (1995) [$T_{1/2}$].
 1997Ch53 V.G. Chumin, J.K. Jabber, K. V. Kalyapkin, S.A.Kudrya, V.V. Tsupko-Sitnikov, K.Ya. Gromov, V.I. Fominykh, T.A. Furyaev, Bull. Rus. Acad. Sci.Phys. 61, 1606 (1997) [P_α].
 1997VaZV Ya. Vavryshchuk, K.Ya. Gromov, V.B. Zlokazov, V.G. Kalinnikov, V.A. Morozov, N.V. Morozova, V.I. Fominykh, V.V. Tsupko-Sitnikov, I.N. Churin, JINR -P6-97-180 (1997) [$T_{1/2}$].
 1997Wa27 J. Wawryszczuk, K.V. Kalyapkin, M.B. Yuldashev, K.Ya. Gromov, V.I. Fominykh, Bull. Rus. Acad. Sci. Phys. 61, 25 (1997) [$T_{1/2}$].
 1998Wa25 J. Wawryszczuk, K.Ya. Gromov, V.B. Zlokazov, V.G. Kalinnikov, V.A. Morozov,

Comments on evaluation

- N.V.Morozova, V.I. Fominikh, V.V. Tsupko-Sitnikov, I.N. Churin, Phys.Atomic Nuclei 61, 1322 (1998) [$T_{1/2}$].
- 2002Mo46 V.A.Morozov, N.V.Morozova, Yu.V.Norseev, Zh.Sereeter, V.B.Zlokazov, Nucl. Instrum. Methods Phys.Res. A484, 225 (2002) [$T_{1/2}$].
- 2003Au03 G.Audi, A.H.Wapstra, C.Thibault, Nucl. Phys. A729(2003)129 [Q].
- 2007Ba19 M.S.Basunia, Nucl.Data Sheets 108, 633 (2007) [NDS]
- 2007HuXX Huang Xiaolong, Wang Baosong, Nuclear Science and Techniques, Vol. 108, 261(2007) [Evaluation].

**²¹⁴Pb - Comments on evaluation of decay data
by V. Chisté and M. M. Bé**

This evaluation was completed in 2007. Literature available by January 2007 was included.

1 Decay Scheme

²¹⁴Pb disintegrates by beta minus emission to the excited levels and to the ground state of ²¹⁴Bi. Spins and parities are from the mass-chain evaluation of Y. A. Akovali (1988Ak01 and 1995E107 for A = 214).

A good agreement was found between the recommended Q value of Audi and the effective Q value (1024 (11) keV) calculated from the decay scheme data.

2 Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2003Au03).

The recommended value of ²¹⁴Pb half-life is 26.8 minutes with an uncertainty of 0.9 minutes from M. Curie (1931Cu01). No recent reference was found in the literature.

2.1 β⁻ Transitions and Emissions

The maximum energies of the β⁻ transitions in the decay of ²¹⁴Pb → ²¹⁴Bi were obtained from the Q⁻ value and the level energies given in Table 1 from Y. A. Akovali (1995E107).

Table 1: ²¹⁴Bi levels populated in the decay of ²¹⁴Pb.

Level number	Level energy, (keV)	Spin and parity	Half-life
0	0.0	1 ⁻	19.9 (4) min
4	295.224 (2)	1 ⁻	≤ 0.05 ns
5	351.932 (2)	0 ⁻ , 1 ⁻	≤ 0.10 ns
7	533.67 (2)	(1 ⁻)	
8	797.24 (9)		
9	839.00 (4)	1 ⁺	

The adopted β⁻ transition probabilities were deduced from the P(γ + ce) balance at each level of the decay scheme. In the Table 2, the recommended values of β⁻ transition probabilities are compared with the experimental results found in the literature: E. E. Berlovich (1952Be78) and S. Kageyama (1953Ka40) observed only two β⁻ transitions 672-keV and 729-keV and H. Daniel (1956Da28) and K. O. Nielsen (1957Ni11) observed the 1024-keV β⁻ transition. A fair agreement has been found between the results given by S. Kageyama and the recommended value for the 729-keV β⁻ transition.

Table 2: Recommended and experimental values of β⁻ transition probabilities.

	672-keV β ⁻ transition	729-keV β ⁻ transition	1024-keV β ⁻ transition
E. E. Berlovich (1952Be78)	25 %	75 %	
S. Kageyama (1953Ka40)	56 %	44 %	
H. Daniel (1956Da28)			6.3 (20) %
K. O. Nielsen (1957Ni11)			< 10 %
Recommended	46.52 (37) %	41.09 (39) %	9.2 (7) %

The values of $lg ft$ and average β^- energies have been calculated with the program LOGFT for the β^- transitions.

2.2 g Transitions

The γ -ray transition probabilities were deduced using the γ -ray emission intensities and the relevant internal conversion coefficients.

Multipolarities and δ (recommended by 1995El07) of these γ -ray transitions and the internal conversion coefficients (ICC's) are shown in Table 3. The internal conversion coefficients have been obtained using:

- A - the Icc99v3a computer program (GETICC dialog) which is based on the new tables of Band *et al.* (2002Ba85) (calculation for 'no hole') and Rösels (1978Ro22).
- B - the BrIcc computer program ("Frozen orbital approximation") which interpolated from theoretical values of Band *et al.* (2002Ba85).

Table 3: Multipolarities of γ -ray transitions.

E_γ (keV)	Multipolarity	α_T (Band) ^a	α_T (Rösels) ^a	α_T (BRICC) ^b
53.2275 (21)	M1 + E2, $\delta = 0.030$ (10)	1.212 (36) E+01	1.288 (39) E+01	1.214 (19) E+01
241.997 (3)	M1 (+E2), $\delta = 0.00$ (15)	8.37 (25) E-01	8.88 (27) E-01	8.38 (18) E-01
258.87 (3)	M1	6.95 (21) E-01	7.37 (22) E-01	6.96 (10) E-01
274.80 (5)	M1 + E2, $\delta = 1.0$	3.73 (11) E-01	3.92 (12) E-01	3.74 (6) E-01
295.224 (2)	M1 + E2, $\delta = 0.30$ (13)	4.54 (14) E-01	4.82 (14) E-01	4.6 (3) E-01
305.26 (3)	[E1]	2.91 (9) E-02	2.95 (9) E-02	2.92 (4) E-02
351.932 (2)	M1 (+E2), $\delta = 0.00$ (35)	3.00 (9) E-01	3.19 (10) E-01	3.00 (25) E-01
480.43 (2)	M1 (+E2), $\delta = 0.0$ (10)	1.302 (39) E-01	1.384 (42) E-01	1.3 (5) E-03
487.09 (7)	(E1)	1.046 (31) E-02	1.058 (32) E-02	1.047 (15) E-03
533.66 (2)	[M1,E2]	6.24 (19) E-02	6.57 (20) E-02	6 (4) E-02
543.81 (7)	[E1]	8.34 (25) E-03	8.43 (25) E-03	8.34 (12) E-03
580.13 (3)	(E1)	7.32 (22) E-03	7.40 (22) E-03	7.32 (11) E-03
785.96 (9)	E1	4.07 (12) E-03	4.10 (12) E-03	4.06 (6) E-03
839.00 (4)	(E1)	3.60 (11) E-03	3.63 (11) E-03	3.59 (5) E-03

a: A fractional uncertainty of 3 % was adopted for all conversion coefficients.

b: Associated uncertainties are calculated by BrIcc.

The evaluators have adopted the internal conversion coefficients interpolated from the Rösels tables, because these ICCs lead to a better decay scheme, where the sum of all the β^- transition probabilities is equal to 100.6 %. The others two ICC's set of values, Band and BrIcc, lead to an inconsistent decay scheme, where the sum of all β^- transitions probabilities would be of the order of 102– 103 %. Moreover, the effective Q value, of 1024 (11) keV, calculated from the decay scheme data with Rösels's Icc, is closer to the recommended value of 1019 (11) keV than the 1029 (15) keV with the "No hole" approximation.

3 Atomic Data

Atomic values, ω_K , ω_L , ω_M , n_{KL} and ω_{LM} and the X-ray and Auger electrons relative probabilities are from Schönfeld and Janßen (1996Sc06).

4 Electron Emissions

The conversion electron emission probabilities have been calculated from γ -ray transition data.

5 Photon Emissions

5.1 X-ray Emissions

The X-ray absolute intensities were calculated from γ -ray data and Rösel's ICC using the EMISSION computer program and compared in Table 4 with the measured values of U. Schötzig (1983Sc13) and E. W. A. Lingeman (1969Li11). A good agreement was found between the experimental results given by 1969Li11 and 1983Sc13 and the recommended values deduced from the decay scheme balance. For the $K\beta$ x-ray, a fair agreement was found between 1969Li11 and the recommended one.

Table 4: Experimental and recommended (calculated) values of X-ray.

	U. Schötzig (1983Sc13)	E. W. A. Lingeman (1969Li10)	Recommended values
$K\alpha$ x-ray (74.82 + 77.11 keV)	16.3 (4) %	17.3 (20) %	16.73 (23) %
$K\beta$ x-ray		4.3 (8) %	4.69 (10) %

5.2 γ Emissions

The γ -ray energy emissions given are from Y. A. Akovali (1995El07).

The experimental relative γ emission intensities in ²¹⁴Bi are based on all available relative and absolute measurements of γ -rays for the ²²⁶Ra decay chain. The normalization factor to convert the relative emission intensities to absolute intensities is the weighted average of the measured absolute γ -ray emission intensities (Table 5) of the most intense line in ²²⁶Ra decay chain, presents in the ²¹⁴Bi disintegration namely the 609.3 keV line.

Table 5: The experimental absolute 609.3 keV gamma-ray emission intensity.

References	Experimental values (%)	Comments
E. W. A. Lingeman (1969Li10)	42.8 (40)	
D. G. Olson (1983Ol01)	45.0 (7)	
U. Schötzig (1983Sc13)	44.6 (5)	
W. -J. Lin (1991Li11)	46.1 (5)	
J. Morel (1998Mo14)	44.8 (6)	Omitted (superseded in 2004Mo07)
J. Morel (2004Mo07)	45.57 (18)	
Recommended value	45.49 (19)	$\chi^2 = 1.45$

The recommended normalization factor is the weighted average of the five experimental values: 45.49 with an external uncertainty of 0.19.

The experimental relative γ emission intensities given in Table 6 are relative to the ²¹⁴Bi 609-keV γ -ray.

Table 6: The experimental data set of the relative γ emission intensities (next page)

Reference	53-keV γ -ray	107-keV γ -ray	137-keV γ -ray	141-keV γ -ray	170-keV γ -ray	196-keV γ -ray	205-keV γ -ray	216-keV γ -ray	241-keV γ -ray
1964Ew04									16.0 (16)
1969Li10									17.1 (18) ^b
1969Wa27									19.33 (30) ^a
1969Gr33	3.15 (34) ^a								16.2 (17) ^a
1970Mo28									16.10(21)
1975Ha31						0.16 (7) ^a			17.5 (17) ^a
1977Zo01									16.06 (19) ^a
1982Ak03						0.14 (2) ^a			16.1 (24) ^a
1982Fa10					0.020 (8) ^a				16.53 (31) ^a
1983Ol01									16.49 (29)
1983Sc13	2.44 (11)								15.65 (25)
1990Mouze		0.015 (3)			0.032 (6)	0.15 (2)	0.025 (6)	0.022 (5)	16.23 (10)
1991Li11									16.33 (25)
2000Sa32						0.16 (8)	0.026 (12)		16.1 (10)
2002De03	2.329 (23)		0.10 (4)	0.06 (3)					15.896 (48)
2002MoZP	2.329 (23)								15.98 (6)
2004Mo07	2.329 (23) ^a								15.880 (48) ^a
Recommended	2.331 (16)	0.015 (3)			0.032 (6)	0.151 (9)	0.025 (5)	0.022 (5)	15.977 (48)
χ^2	0.5					0.015	0.005		2.0
Reference	258-keV γ -ray	274-keV γ -ray	295-keV γ -ray	305-keV γ -ray	314-keV γ -ray	323-keV γ -ray	351-keV γ -ray	462-keV γ -ray	480-keV γ -ray
1964Ew04			40.46 (40)				77 (8)		
1969Li10	1.32 (22)	1.10 (22) ^b	42.6 (44) ^b		0.220 (44)	0.066 (22)	80 (9)	0.46 (11)	0.66 (15)
1969Wa27			47.87 (91) ^a				87.2 (19) ^a		
1969Gr33	1.16 (7) ^a	1.01 (10) ^a	40.2 (40) ^a		0.137 (23) ^a		79 (7) ^a	0.444 (46) ^a	
1970Mo28			41.45 (56) ^b				79.7 (11)		
1975Ha31	1.24 (12) ^a	0.71 (7) ^a	40.2 (40) ^a	0.050 (25) ^a	0.198 (50) ^a	0.062 (25) ^a	86 (9) ^a	0.446 (50) ^a	0.73 (7) ^a
1977Zo01			42.01 (53) ^a				80.42 (81) ^a		
1982Ak03	1.17 (15) ^a	0.86 (16) ^a	42.2 (54) ^a	0.075 (16) ^a	0.185 (28) ^a	0.072 (40) ^a	82 (11) ^a	0.44 (7) ^a	0.75 (10) ^a
1982Fa10	1.72 (4) ^a		42.52 (59) ^a				81.3 (8) ^a		0.68 (2) ^a
1983Ol01			40.8 (6)				78.7 (11)		
1983Sc13			40.0 (7)				77.2 (9)		
1990Mouze	1.23 (6)	0.84 (6)	41.85 (26) ^a	0.068 (10)	0.17 (2)	0.06 (1)	81.48 (48) ^a	0.40 (4)	0.71 (5)
1991Li11	1.152 (25)	1.042 (25) ^b	42.43 (47) ^a				82.7 (9) ^a	0.486 (20)	0.703 (24)
2000Sa32	1.15 (4)	0.83 (8)	40.8 (12)	0.080 (15)	0.158 (20)	0.084 (20)	78.5 (24)	0.470 (14)	0.74 (3)
2002De03	1.171(9)	0.787 (23)	40.36 (12)				78.16 (23)		0.749 (10)
2002MoZP			40.61 (13)				78.34 (23)		
2004Mo07	1.171(9) ^a	0.760(27) ^a	40.32 (12) ^a				78.10 (23) ^a		0.75 (1) ^a
Recommended	1.169 (8)	0.796 (21)	40.48 (31)	0.0692 (47)	0.169 (13)	0.063 (7)	78.26 (16)	0.469 (12)	0.741 (9)
χ^2	0.56	0.43	0.57	0.56	0.82	0.65	0.52	1.24	0.95

Reference	487-keV γ -ray	533-keV γ -ray	538-keV γ -ray	543-keV γ -ray	580-keV γ -ray	765-keV γ -ray	785-keV γ -ray	839-keV γ -ray
1964Ew04								
1969Li10	0.77 (18)	0.37 (9)			0.70 (13)		2.31 (33)	1.30 (18)
1969Wa27								
1969Gr33	0.91 (23) ^a	0.501 (46) ^a			0.89 (9) ^a		2.41 (23) ^a	1.41 (14) ^a
1970Mo28								
1975Ha31	0.88 (10) ^a	0.408 (50) ^a		0.050 (16) ^a	0.80 (7) ^a		2.48 (25) ^a	1.42 (14) ^a
1977Zo01								
1982Ak03	0.88 (11) ^a	0.42 (5) ^a		0.14 (2) ^a	0.79 (11) ^a		2.32 (32) ^a	1.33 (19) ^a
1982Fa10	0.83 (3) ^a							1.30 (3) ^a
1983O101								
1983Sc13							2.286 (45)	
1990Mouze	0.83 (7)	0.39 (3)	0.044 (6)	0.15 (2)	0.76 (6)	0.17 (3)	2.33 (17)	1.29 (10)
1991Li11	0.928 (35)	0.409 (20)			0.774 (31)		2.396 (45)	1.290 (20)
2000Sa32	0.90 (5)	0.39 (3)	0.037 (20)	0.100 (10)	0.74 (4)	0.11 (1)	2.33 (7)	1.29 (4)
2002De03	0.961 (12)				0.823 (11)			
2002MoZP								
2004Mo07	0.961 (12) ^a				0.824 (10) ^a			
Recommended	0.951 (14)	0.399 (14)	0.043 (6)	0.11 (2)	0.811 (13)	0.116 (18)	2.339 (28)	1.290(18)
χ^2	1.54	0.18	0.11	5	1.80	3.6	0.75	0.001

a: Not used by the evaluators (see below).

b: the experimental value has been shown to be outlier value by the Lweight program.

There were omitted from analysis:

a) four sets of values, A. Hachem (1975Ha31), G. Mouze (1981Mo28), H. Akcay (1982Ak03), G. Mouze (1990Mo08) and O. Diallo (1993Di09), because these values comes from the same laboratory of G. Mouze (1990Mo**).

b) the sets of values from K. Ya. Gromov (1969Gr33), G. Wallace (1969Wa27) and M. A. Farouk (1982Fa10), because a lack of information in the articles describing their experimental measurements.

c) the set of values from V. Zobel(1977Zo01), because these values changedthe consistency of the data set when introduced in the preliminary calculation with Lweight program, and produced inconsistent set of data for gamma emission intensities. Therefore, in the case of 295keV and 351-keV γ -rays, the values of G. Mouze (1990Mouze) and W. –J. Lin (1991Li11), consistent with Zobel values, were not used by the evaluators for the weighted mean calculations.

d) the relative γ emission intensity values given by 2004Mo07, because they are those measured by J. U. Delgado (2002De03). In 2004Mo07 article, the author measured the 609.3 keV absolute emission probability (Table 2) and normalized the 2002De03 data set with this value of 45.57 (18).

The adopted values are the weighted means calculated by the Lweight program (version 3).

The evaluated relative and absolute γ -ray emission intensities are given in Table 7.

Table 7: Evaluated relative and absolute γ -ray emission intensities

Energy (keV)	Relative emission intensity	Absolute emission intensity (%)
53.2275 (21)	2.331 (16)	1.060 (9)
107.22 (9)	0.015 (3)	0.0068 (14)
137.45 (30)	0.10 (4)	0.045 (18)
141.3 (6)	0.06 (3)	0.027 (14)
170.07 (6)	0.032 (6)	0.0146 (27)
196.20 (5)	0.151 (9)	0.069 (9)
205.68 (9)	0.025 (5)	0.0114 (23)
216.47 (7)	0.022 (5)	0.0100 (23)
241.997 (3)	15.977 (48)	7.268 (22)
258.87 (3)	1.169 (8)	0.5318 (43)
274.80 (5)	0.796 (21)	0.362 (10)
295.224 (2)	40.48 (8)	18.414 (36)
305.26 (3)	0.0692 (47)	0.0315 (21)

Energy (keV)	Relative emission intensity	Absolute emission intensity (%)
314.32 (7)	0.169 (13)	0.077 (6)
323.83 (4)	0.063 (7)	0.0287 (32)
351.932 (2)	78.26 (16)	35.60 (7)
462.00 (7)	0.469 (12)	0.213 (6)
480.43 (2)	0.741 (9)	0.3371 (43)
487.09 (7)	0.951 (14)	0.433 (7)
533.66 (2)	0.399 (14)	0.182 (6)
538.41 (8)	0.043 (6)	0.0196 (27)
543.81 (7)	0.11 (2)	0.050 (9)
580.13 (3)	0.811 (13)	0.369 (6)
765.96 (9)	0.116 (18)	0.053(8)
785.96 (9)	2.339 (28)	1.064 (13)
839.04 (9)	1.290 (18)	0.587 (8)

6 References

- 1931C01 – M. Curie, A. Debierne, A. S. Eve, H. Geiger, O. Hahn, S. C. Lind, St. Meyer, E. Rutherford, E. Schweidler, *Rev. Mod. Phys.* 3(1931)427 [Half-life].
- 1952Be78 – E. E. Berlovich, *Izvest. Acad. Nauk.SSSR, Ser. Fiz.* 16(1952)314, *Bull.Acad. Sci. USSR, Phys. Ser.* 16(1952)314 [I_{β}].
- 1953Sa40 – K. Sageyama, *J. Phys. Soc. (Japan)* 8(1953)689 [I_{β}].
- 1956Da06 – H. Daniel, *Z. Naturforsch.* 11a(1956)212 [Half-life].
- 1956Da28 – H. Daniel, *Z. Naturforsch.* 11a(1956)759 [I_{β}].
- 1957Ni11 – K. O. Nielsen, O. B. Nielsen, M. A. Waggoner, *Nucl. Phys.* 2(1957)476 [I_{β}].
- 1964Ew04 – G. T. Ewan, J. Tavendale, *Can. J. Phys.* 42(1964)2286 [I_{γ}].
- 1969Gr33 – K. Ya. Gromov, B. M. Sabirov, J. J. Urbanets, *Bull. Acad. Sci. USSR, Phys. Ser.* 33(1970)1510 [I_{γ}].
- 1969Li10 – E. W. A. Lingeman, J. Konijn, P. Polak, A. H. Wapstra, *Nucl. Phys.* A133(1969)630 [I_{γ}].
- 1969Wa27 – G. Wallace, G. E. Coote, *Nucl. Instrum. Meth.* 74(1969)353 [I_{γ}].
- 1970Mo28 – R. S. Mowatt, *Can. J. Phys.* 48(1970)2606 [I_{γ}].
- 1975Ha31 – A. Hachem, *Compt. Rend. (Paris)* 281B(1975)45 [I_{γ}].
- 1977Zo01 – V. Zobel, J. Eberth, U. Eberth, E. Eube, *Nucl. Instrum. Meth.* 141(1977)329 [I_{γ}].
- 1978Ro22 – F. Rösel et al., *At. Data Nucl. Data Tables* 21(1978)91 [Theoretical ICC].
- 1981Mo28 – G. Mouze, *Compt. Rend. (Paris)* 292(1981)1243 [I_{γ}].
- 1982Ak03 – H. Akcay, G. Mouze, D. Maillard, Ch. Ythier, *Radiochem. Radioanal. Lett.* 51(1982)1 [I_{γ}].
- 1982Fa10 – M. A. Farouk, A. M. Al-Soraya, *Nucl. Instrum. Meth.* 200(1982)593 [I_{γ}].
- 1983Ol01 – D. G. Olson, *Nucl. Instrum. Meth.* 206(1983)313 [I_{γ}].
- 1983Sc13 – U. Schötzig, K. Debertin, *Int. J. Appl. Radiat. Isot.* 34(1983)533 [I_{γ}].
- 1984Pe13 – I. Penev, W. Andretscheff, Ch. Prochostow, *Zh. Fiz. i Khim. At. Energ.* 31(1984)213 [Half-life (E_{γ} = 53 keV)].
- 1988Ak01 – Y. A. Akovali, *Nucl. Data Sheets* 55(1988)665 [Energy level, spin, parity and multipolarity].
- 1990Mo08 – G. Mouze, O. Diallo, P. Bechlich, C. Ythier, J. F. Comanducci, *Radiochimica Acta* 49(1990)13 [I_{γ}].
- 1990Mo** – G. Mouze, C. Ythier, J. F. Comanducci, *Rev. Roumaine Phys.* 35(1990)337 [I_{γ}].
- 1991Li11 – W. –J. Lin, G. Harbottle, *J. Radioanal. Nucl. Chem. Lett.* 153(1991)137 [I_{γ}].
- 1993Di09 – O. Diallo, G. Mouze, C. Ythier, J. F. Comanducci, *Nuovo Cimento* 106A(1993)1321 [I_{γ}].
- 1995El07 – Y. A. Akovali, *Nucl. Data Sheets* 76(1995)127 [Energy level, spin, parity and multipolarity].
- 1996Sc06 – E. Schönfeld, H. Janßen, *Nucl. Instrum. Meth. Phys. Res.* A369(1996)527 [Atomic data].
- 1998Mo14 – J. Morel, M. Etcheverry, J. L. Picolo, *Appl. Radiat. Isot.* 49(1998)1387 [I_{γ}].
- 2000Sa32 – D. Sardari, T. D. MacMahon, *J. Radioanal. Nucl. Chem.* 244(2000)463 [I_{γ}].
- 2002Ba85 – I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr., P. O. Tikkanen, S. Raman, *At. Data Nucl. Data Tables* 81(2002)1 [Theoretical ICC].

Comments on evaluation

- 2002De03 – J. U. Delgado, J. Morel, M. Etcheverry, Appl. Radiat. Isot. 56(2002)137 [I_γ].
2002MoZP - G. L. Molnar, Z. S. Révay, T. Belgya, 11th Int. Symp. on capture gamma-ray spectroscopy, 2-6 Sep. 2002, Pruhonice (2003)522 [I_γ].
2003Au03 – G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].
2004He** – R. G. Helmer, IAEA – CRP Report to be published (2004) [I_γ].
2004Mo07 – J. Morel, S. Speman, M. Rasko, E. Terechtchenko, J. U. Delgado, Appl. Radiat. Isot. 60(2004)341 [I_γ].

**²¹⁴Bi - Comments on evaluation of decay data
by V. Chisté and M. M. Bé**

This evaluation was completed in 2007. Literature available by January 2007 was included.

1 Decay Scheme

²¹⁴Bi disintegrates by beta minus emissions to excited levels and to the ground state of ²¹⁴Po (99.979 (13) %) and by alpha emission to excited levels of ²¹⁰Tl (0.0210 (13) % (1960Wa14)), some alpha emissions of long range from excited levels in ²¹⁴Po to excited levels in ²¹⁰Pb have been observed. Spins and parities are from the mass-chain evaluation of Y. A. Akovali (1988Ak01 and 1995El07 for A = 214) and E. Browne (2003Br13 for A = 210).

A good agreement was found between the adopted $Q(\beta^-)$ value of Audi and the effective $Q(\beta^-)$ value of 3261 (10) keV calculated from decay scheme data.

2 Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2003Au03).

The recommended value of ²¹⁴Bi half-life is 19.9 minutes with an uncertainty of 0.4 minutes from H. Daniel (1956Da06). No recent references were found in the literature.

2.1 β^- Transitions and Emissions

The maximum energies of the β^- transitions in the decay of ²¹⁴Bi \rightarrow ²¹⁴Po have been obtained from the Q^- value (2003Au03) and the level energies given in Table 1 from Y. A. Akovali (1995El07).

Table 1: ²¹⁴Po levels populated in the decay of ²¹⁴Bi.

Level Number	Level energy, (keV)	Spin and parity	Level Number	Level energy, (keV)	Spin and parity
0	0	0 ⁺	24	2293.34 (5)	1 ⁽⁺⁾ , 2 ⁺
1	609.316 (7)	2 ⁺	25	2348.3 (9)	1 ⁻ , 1 ⁺ , 2 ⁺
4	1377.675 (12)	2 ⁺	26	2360.8 (4)	1 ⁻ , 1 ⁺ , 2 ⁺
5	1415.489 (19)	0 ⁺	27	2423.19 (15)	1 ⁺ , 2 ⁻ , 2 ⁺
6	1543.375 (14)	2 ⁺	28	2447.70 (6)	1 ⁻
7	1661.28 (3)	2 ⁺	29	2482.46 (4)	(2) ⁺
8	1712.93 (20)	(3) ⁺	30	2505.21 (15)	1 ⁽⁻⁾ , 2 ⁺
9	1729.611 (13)	2 ⁺	31	2508.2 (2)	
10	1742.98 (3)	0 ⁺	32	2544.9 (3)	
11	1764.498 (14)	1 ⁺	34	2562.4 (3)	
12	1847.431 (14)	2 ⁺	35	2604.66 (14)	(2) ⁺
13	1890.287 (21)	2 ⁺	36	2630.85 (17)	1 ⁻ , 1 ⁺ , 2 ⁺
14	1994.63 (3)	(2) ⁻	37	2662.29 (12)	(2) ⁺
15	2010.81 (4)	2 ⁺	38	2694.6 (2)	1 ⁽⁻⁾ , 2 ⁺
16	2017.3 (5)	0 ⁺	39	2698.8 (3)	1 ⁽⁻⁾ , 2 ⁺
17	2088.41 (12)	1 ⁻ , 1 ⁺ , 2 ⁺	40	2699.2 (2)	1 ⁽⁻⁾ , 2 ⁺
18	2118.552 (17)	1 ⁺	41	2719.22 (9)	1 ⁻ , 1 ⁺ , 2 ⁺
19	2147.78 (6)	1 ⁽⁻⁾ , 2 ⁺	42	2728.59 (4)	(1,2) ⁺
20	2192.56 (4)	2 ⁺	43	2769.9 (2)	1 ⁻ , 1 ⁺ , 2 ⁺
21	2204.13 (9)	1 ⁺	44	2785.9 (2)	1 ⁻ , 1 ⁺ , 2 ⁺
23	2266.39 (18)	1 ⁽⁻⁾ , 2 ⁺	47	2827.0 (2)	1 ⁻ , 1 ⁺ , 2 ⁺

Level Number	Level energy, (keV)	Spin and parity	Level Number	Level energy, (keV)	Spin and parity
48	2861.1 (3)	1 ⁻ , 1 ⁺ , 2 ⁺	61	2986.2 (2)	(1 ⁻ , 2 ⁻ , 2 ⁺)
49	2869.6 (2)		62	3000.0 (2)	1 ⁽⁻⁾ , 2 ⁺
50	2880.3 (2)	1 ⁻ , 1 ⁺ , 2 ⁺	65	3014.1 (3)	1 ⁻ , 1 ⁺ , 2 ⁺
51	2893.6 (2)	1 ⁻ , 1 ⁺ , 2 ⁺	69	3053.9 (2)	1 ⁻ , 1 ⁺ , 2 ⁺
52	2897.0 (3)		70	3068.3 (8)	
53	2919.5 (3)		72	3081.7 (3)	1 ⁻ , 1 ⁺ , 2 ⁺
54	2921.8 (4)	1 ⁻ , 1 ⁺ , 2 ⁺	73	3094.0 (4)	(1 ⁻ , 2 ⁺)
55	2928.6 (3)	1 ⁻ , 1 ⁺ , 2 ⁺	75	3142.6 (4)	1 ⁻ , 1 ⁺ , 2 ⁺
56	2934.5 (3)	1 ⁻ , 1 ⁺ , 2 ⁺	76	3149.2 (5)	1 ⁻ , 1 ⁺ , 2 ⁺
57	2940.6 (2)	1 ⁽⁻⁾ , 2 ⁻ , 2 ⁺	77	3160.4 (6)	1 ⁻ , 1 ⁺ , 2 ⁺
58	2962.8 (7)		79	3173.3 (6)	
60	2978.8 (2)	1 ⁻ , 1 ⁺ , 2 ⁺	80	3183.6 (4)	1 ⁻ , 1 ⁺ , 2 ⁺

The adopted β^- transition probabilities and the associated uncertainties were deduced from the γ transition probability balance at each level of the decay scheme.

The values of $\log ft$ and average β^- energies have been calculated with the program LOGFT for the allowed and 1st forbidden β^- transitions.

2.2 g Transitions

The γ -ray transition probabilities were calculated using the γ -ray emission intensities and the relevant internal conversion coefficients (see 4.2 g Emissions).

Multipolarities of γ -ray transitions are from Y. A. Akovali (1995El07 for A = 214) and E. Browne (2003Br13 for A = 210) and shown in Table 2.

Table 2: Multipolarities of γ -ray transitions.

	Multipolarity	E_γ (keV)
²¹⁰ Tl	(M1)	62.5 (10)
²¹⁴ Po	[M1,E2]	221 (1), 386.823 (18), 452.92 (10), 469.756 (18), 474.52 (5), 543.0 (2), 595.32 (7), 633.14 (10), 634.72 (21), 649.19 (7), 661.1 (2), 697.88 (20), 740.73 (18), 752.84 (3), 814.885 (10), 878.03 (12), 915.74 (15), 939.6 (5), 991.49 (19), 1051.964 (31), 1103.61 (20), 1104.79 (19)
	[M1]	252.80 (6), 349.009 (24), 388.941 (50), 461.15 (20), 703.11 (4), 788.6 (5), 1594.81 (8)
	[E1]	268.614 (26), 333.35 (6), 454.850 (26), 487.95 (13), 572.76 (7), 615.53 (10), 617.0 (2), 683.22 (6), 704.9 (3), 786.1 (4), 904.29 (10), 917.8 (3), 1032.37 (8), 1069.96 (8), 1207.70 (3), 1385.314 (31)
	[E2]	405.74 (4), 528 (1), 639.62 (10), 832.38 (11), 1133.664 (31), 1172.98 (10), 1543.375 (14)
	(E2)	1407.98 (4)
	(M1 + E2)	1401.494 (41) $\delta = 1.6$ (5)
	E2	609.316 (7), 719.869 (37), 806.173 (20), 1377.675 (12), 1661.28 (6), 1729.611 (13),
	E1	665.445 (23), 2447.86 (10)
	M1 + E2	768.359 (14) $\delta = 2.8$ (7)
		934.059 (16) $\delta = -0.3$ (1)
		1120.295 (15) $\delta = 0.18$ (2)
		1155.182 (16) $\delta = 0.33$ (6)
		1238.115 (12) $\delta = -0.03$ (3)
		1509.236 (15) $\delta = -0.053$ (35)
		1583.244 (40) $\delta = -0.20$ (10)
	M1	821.18 (3), 826.46 (20), 1280.97 (2), 1764.498 (14), 2118.552 (30), 2204.21 (4)

The internal conversion coefficients (ICC) and the associated uncertainties for these γ -ray transitions have been obtained using the BRICC computer program (calculation for 'hole'), which interpolated the new values from theoretical values of I. M. Band (2002Ba85).

2.3 a Transitions and Emissions

The energies of the α -particle transitions given in Section 2.3 have been obtained from Q $_{\alpha}$ (2003Au03) and the level energies given by E. Browne (2003Br13).

The adopted $\alpha_{0,0}$, $\alpha_{0,2}$ and $\alpha_{0,3}$ emission energies are the recommended values of A. Rytz (1991Ry01) and the other α emission energies are from E. Browne (2003Br13).

The recommended α emission probabilities come from the measured values of R. J. Walen (1960Wa14).

For the α of long range, the energy and emission probabilities are from the measurements of C. -F. Leang (1965Le08).

3 Atomic Data

Atomic values, ω_K , ω_L , ω_M , n_{KL} and \bar{n}_{LM} and the X-ray and Auger electron relative probabilities are from Schönfeld and Janßen (1996Sc06).

4 Electron Emissions

The conversion electron emission probabilities have been deduced from γ -ray transition data.

5 Photon Emissions

5.1 X-ray Emissions

The X-ray absolute intensities have been calculated from γ -ray data and ICC using the EMISSION computer program and compared in Table 3 with the measured values of U. Schötzig (1983Sc13). These values are not consistent, it is difficult to draw a conclusion since, as said in 1983Sc03, "the x-ray spectrum is rather complex, as the Po and Bi x-ray peaks overlap, a deconvolution is difficult".

Table 3: Experimental and recommended (calculated) values of X-ray absolute intensities.

	U. Schötzig (1983Sc13)	Recommended values
K α x-ray	1.77 (5) %	1.135 (25) %
K β x-ray	4.94 (12)	0.320 (9) %

5.2 γ Emissions

The γ -ray energies are from Y. A. Akovali (1995El07 for A = 214) and E. Browne (2003Br13 for A = 210).

For the ^{210}Tl γ -rays, the absolute γ -ray emission intensities have been deduced from the α emission intensities measured by R. J. Walen (1960Wa14).

The experimental relative γ -ray emission intensities in ^{214}Po are based on all available relative and absolute measurements of γ -rays for the ^{226}Ra decay chain. The normalization factor to convert the relative γ -ray emission intensities to absolute intensities is the weighted average of the measured values of the 609.3 keV γ -ray absolute intensity (Table 4).

Table 4: The experimental values of 609.3-keV γ -ray absolute intensity.

References	Experimental values (%)	Comments
E. W. A. Lingeman (1969Li10)	42.8 (40)	
D. G. Olson (1983Ol01)	45.0 (7)	
U. Schötzig (1983Sc13)	44.6 (5)	
W. –J. Lin (1991Li11)	46.1 (5)	
J. Morel (1998Mo14)	44.8 (6)	Omitted: superseded by 2004Mo07
J. Morel (2004Mo07)	45.57 (18)	
Recommended value	45.49 (19)	$\chi^2 = 1.45$

Evaluators' recommended normalization factor is the weighted average of the five experimental values: 45.49 with an external uncertainty of 0.19.

The experimental relative γ -ray emission intensities are given in Table 5 relatively to the ²¹⁴Bi 609-keV γ -ray intensity.

The evaluated relative and absolute γ -ray intensities are given in Table 6.

The adopted values are the weighted means calculated by the Lweight program (version 3).

Table 5: The experimental data set of the relative γ -ray emission intensities. (see next pages)

Energy (keV)	1969Li10	1696Wa27*	1969Gr33*	1975Ha31*	1977Zn01	1982AK03*	1982Fa10*	1983OI01	1983Sc13	1990Mouze	1991Li11	2000Sa32	2002De03	2002MoZP	2004Mo07*	Evaluated	χ^2
221						0.012 (7)						0.130 (13)				0.130 (13)	
230												0.0063 (21)				0.0063 (21)	
252						0.033 (7)				0.028 (4)		0.019 (7)				0.0258 (39)	1.3
268						0.031 (8)				0.035 (4)		0.059 (28)				0.0355 (40)	0.72
273	0.18 (9)			0.384 (50)		0.25 (5)				0.27 (3)		0.29 (10)	0.265 (23)		0.278 (17)	0.264 (18)	0.33
280	0.132 (22)			0.136 (50)		0.13 (2)				0.13 (2)		0.17 (4)				0.136 (14)	0.42
304	0.18 (9)			0.074 (25)		0.069 (15)				0.055 (5)		0.065 (20)				0.056 (5)	1.1
333			0.148 (23)	0.15 (7)		0.16 (3)				0.14 (1)		0.13 (3)				0.139 (9)	0.1
334	0.132 (44)			0.074 (37)		0.072 (14)				0.066 (8)		0.090 (17)				0.072 (10)	1.8
348										0.34 (5)		0.20 (5)				0.27 (7)	3.9
386	0.68 (26)		1.41 (18)	0.64 (7)		0.64 (10)				0.63 (5)		0.70 (15)	0.651 (12)		0.647 (11)	0.650 (12)	0.10
388	0.81 (26)			0.83 (7)		0.87 (12)				0.85 (1)	0.92 (6)	0.86 (4)	0.888 (14)		0.89 (13)	0.864 (10)	1.5
394				0.019 (9)		0.033 (4)				0.032 (3)		0.024 (3)				0.0280 (40)	3.6
396				0.050 (25)		0.060 (9)				0.059 (7)		0.053 (10)				0.057 (4)	0.24
405	0.33 (9)		0.341 (34)	0.40 (7)		0.38 (5)				0.37 (2)		0.39 (3)				0.375 (16)	0.28
452						0.068 (11)				0.067 (8)						0.067 (8)	
454	0.62 (11)		0.64 (7)	0.64 (7)		0.67 (8)	0.63 (2)			0.64 (3)		0.59 (3)	0.640 (12)		0.642 (12)	0.634 (10)	0.82
461						0.078 (13)				0.14 (2)		0.10 (3)				0.128 (18)	1.2
469						0.30 (5)				0.27 (2)		0.34 (3)				0.292 (32)	3.8
474	0.15 (7)			0.24 (7)		0.23 (4)				0.22 (2)		0.190 (20)				0.203 (14)	0.86
485						0.052 (11)				0.048 (9)		0.035 (20)				0.046 (8)	0.35
487										0.061 (20)						0.061 (20)	
494						0.031 (5)				0.031 (4)		0.019 (3)				0.023 (6)	5.8
496						0.015 (4)				0.015 (4)						0.015 (4)	
501				0.038 (7)		0.041 (7)				0.040 (5)		0.035 (19)				0.0397 (48)	0.06
519				0.0124 (50)		0.035 (6)				0.036 (4)		0.039 (11)				0.0364 (38)	0.07
524				0.033 (12)		0.038 (6)				0.037 (4)		0.039 (13)				0.0372 (38)	0.02
528						0.025 (5)				0.024 (3)		0.022 (11)				0.0239 (29)	0.03
536				0.124 (50)		0.14 (2)				0.14 (2)		0.12 (3)				0.134 (17)	0.31
543	0.22 (7)		0.296 (34)	0.20 (7)		0.14 (2)				0.13 (2)		0.27 (4)				0.194 (46)	3.4
547	0.066 (22)			0.071 (14)		0.08 (1)				0.08 (1)		0.074 (7)				0.075 (6)	0.22
551										0.012 (3)						0.012 (3)	
572	0.132 (44)		0.159 (23)	0.161 (25)		0.17 (2)				0.16 (2)		0.16 (4)				0.156 (17)	0.17

Energy (keV)	1969Li10	1696Wa27*	1969Gr33*	1975Ha31*	1977Zn01	1982Ak03*	1982Fa10*	1983OI01	1983Sc13	1990Mouze	1991Li11	2000Sa32	2002De03	2002MoZP	2004Mo07*	Evaluated	χ^2
595						0.035 (7)				0.038 (4)		0.039 (6)				0.0383 (33)	0.02
600										0.018 (8)						0.018 (8)	
609	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	
615	0.20 (7)			0.099 (25)		0.13 (5)				0.12 (2)		0.11 (3)				0.121 (16)	0.71
617				0.074 (25)		0.066 (44)				0.053 (6)		0.077 (11)				0.059 (10)	3.7
626						0.036 (6)						0.009 (3)				0.009 (3)	
630			0.228 (34)	0.037 (12)		0.039 (6)				0.035 (4)		0.039 (5)				0.0366 (31)	0.39
633	0.110 (44)			0.124 (12)		0.12 (2)				0.11 (1)		0.130 (10)				0.120 (7)	1.0
634						0.014 (5)				0.014 (5)						0.014 (5)	
639				0.074 (25)		0.061 (11)				0.065 (10)		0.085 (10)				0.075 (10)	2.0
649	0.110 (44)			0.124 (12)		0.114 (15)				0.13 (2)		0.10 (3)				0.119 (16)	0.37
658						0.037 (8)				0.046 (8)		0.030 (8)				0.038 (8)	2.0
661				0.094 (37)		0.077 (13)				0.11 (2)		0.120 (10)				0.118 (9)	0.2
665	3.08 (44)		3.49 (30)	3.59 (37)		3.36 (37)	2.87 (6)			3.51 (20)	3.21 (7)	3.33 (10)	3.359 (17)	3.386 (21)	3.364 (17)	3.364 (15)	1.4
677						0.012 (5)				0.012 (5)						0.012 (5)	
683	0.176 (44)		0.296 (46)	0.186 (25)		0.18 (3)				0.18 (2)		0.190 (20)				0.184 (13)	0.08
687				0.012 (6)		0.016 (5)				0.015 (4)		0.014 (5)				0.0146 (31)	0.02
693				0.012 (6)		0.012 (5)				0.015 (6)		0.012 (4)				0.0129 (33)	0.17
697	0.154 (44)		0.501 (46)	0.100 (50)		0.14 (2)				0.14 (2)		0.150 (10)				0.148 (9)	0.11
699						0.044 (9)				0.035 (10)						0.035 (10)	
703	1.03 (13)		1.55 (16)	1.14 (12)		1.08 (15)	0.82 (3)			1.11 (7)	1.038 (27)	1.12 (8)				1.053(24)	0.57
704										0.11 (3)		0.113 (29)				0.112 (21)	0.006
708						0.031 (9)				0.042 (11)		0.025 (3)				0.0262 (43)	2.2
710	0.13 (7)		0.364 (34)	0.161 (50)		0.16 (2)				0.16 (2)		0.170 (9)				0.168 (8)	0.25
719	0.84 (11)		1.22 (13)	0.94 (12)		0.90 (13)				0.91 (8)	0.833 (24)	0.91 (3)				0.865 (22)	1.5
722				0.099 (50)		0.075 (11)				0.073 (9)		0.107 (15)				0.082 (15)	3.8
733	0.066 (22)			0.087 (25)		0.086 (12)				0.085 (8)		0.092 (17)				0.084 (7)	0.45
740						0.11 (2)				0.088 (13)		0.095 (5)				0.0941 (47)	0.25
752	0.24 (7)			0.31 (7)		0.30 (4)				0.28 (2)		0.28 (4)				0.278 (17)	0.15
768		9.90 (21)	10.6 (10)	11.4 (12)	10.90 (15)	11.9 (17)	10.64 (20)		10.46 (16)	10.91 (8)	10.86 (14)	10.39 (31)	10.66 (5)	10.77 (3)	10.68 (5)	10.755 (36)	2.3
786	0.64 (18)											0.70 (10)				0.69 (10)	0.09
788										0.041 (8)		0.020 (10)				0.033 (10)	2.7
806	2.42 (44) μ		2.68 (25)	2.97 (37)		2.92 (43)	2.49 (6)			2.90 (22)	2.682 (45)	2.76 (11)	2.788 (22)	2.777 (14)	2.791 (20)	2.774 (13)	1.2
815	0.088 (44)			0.050 (25)		0.087 (13)				0.081 (8)		0.110 (20)				0.085 (7)	0.91
821	0.35 (9)			0.40 (7)		0.37 (6)				0.36 (3)		0.37 (3)				0.364 (21)	0.04

Energy (keV)	1969Li10	1696Wa27*	1969Gr33*	1975Ha31*	1977Zn01	1982Ak03*	1982Fa10*	1983OI01	1983Sc13	1990Mouze	1991Li11	2000Sa32	2002De03	2002MoZP	2004Mo07*	Evaluated	χ^2
826	0.29 (13)			0.21 (7)		0.29 (4)				0.28 (3)		0.29 (4)				0.284 (24)	0.02
832	0.066 (22)			0.062 (25)		0.064 (10)				0.062 (6)		0.080 (3)				0.076 (5)	3.7
847				0.037 (12)		0.052 (12)				0.057 (7)		0.053 (15)				0.035 (13)	0.06
873						0.032 (10)				0.042 (9)		0.040 (10)				0.041 (7)	0.02
878						0.022 (7)				0.026 (6)						0.026 (6)	
904	0.15 (7)			0.198 (50)		0.15 (2)				0.14 (2)		0.16 (4)				0.144 (17)	0.1
915				0.050 (12)		0.070 (14)				0.065 (8)		0.043 (6)				0.051 (11)	4.8
917						0.010 (7)				0.010 (7)						0.010 (7)	
930						0.058 (13)				0.10 (2)		0.08 (3)				0.094 (17)	0.31
934	6.8 (7)	6.26 (18)	7.0 (7)	7.3 (7)	6.93 (10)	7.0 (9)	6.54 (13)		6.75 (9)	6.88 (5)	6.66 (9)	6.70 (20)	6.783 (34)	6.83 (4)	6.788 (34)	6.814 (22)	1.05
939						0.030 (8)				0.028 (8)		0.045 (9)				0.036 (8)	2.0
943			0.205 (34)	0.037 (12)		0.034 (8)				0.037 (6)		0.050 (26)				0.038 (6)	0.24
949						0.009 (6)				0.012 (5)						0.012 (5)	
952										0.013 (5)						0.013 (5)	
961						0.046 (12)				0.03 (2)		0.022 (3)				0.0222 (30)	0.16
964	0.81 (11)		0.78 (8)	0.85 (9)		0.82 (10)				0.80 (5)	0.796 (38)	0.80 (7)				0.799 (27)	0.01
976				0.050 (25)		0.029 (8)				0.033 (5)		0.035 (13)				0.0333 (47)	0.02
991				0.0031 (15)		0.009 (6)				0.022 (5)		0.050 (22)				0.023 (6)	1.5
1013				0.022 (11)						0.018 (3)		0.034 (11)				0.0191 (41)	1.9
1021				0.025 (12)						0.034 (6)		0.036 (15)				0.034 (6)	0.02
1032	0.154 (44)			0.161 (50)						0.13 (1)		0.17 (3)				0.135 (9)	0.9
1038				0.025 (12)						0.018 (3)		0.030 (10)				0.0190 (33)	1.3
1045				0.062 (12)						0.051 (6)		0.037 (20)				0.050 (6)	0.45
1051	0.73 (9)		0.71 (7)	0.68 (7)			0.76 (3)			0.66 (5)	0.692 (24)	0.72 (4)				0.713 (17)	1.1
1067				0.062 (25)						0.055 (20)		0.051 (24)				0.053 (15)	0.02
1069	0.57 (9)		0.73 (14)	0.62 (7)						0.56 (4)	0.605 (33)	0.65 (6)				0.595 (23)	0.59
1103	0.35 (7)			0.21 (10)						0.21 (3)		0.24 (7)				0.233 (33)	1.7
1104			0.250 (34)	0.17 (7)						0.16 (3)						0.16 (3)	
1118										0.015 (8)		0.034 (11)				0.022 (9)	1.9
1120	33.0 (33)	31.90 (73)	29.4 (28)	34.0 (35)	32.72 (39)		33.52 (42)	32.73 (48)	32.31 (46)	33.13 (22)	33.19 (46)	32.3 (10)	32.71 (10)	32.77 (12)	32.74 (10)	32.77 (7)	0.64
1130				0.099 (25)						0.078 (9)		0.080 (11)				0.079 (7)	0.02
1133	0.55 (11)		0.478 (46)	0.62 (6)						0.56 (3)	0.545 (29)	0.57 (3)				0.558 (17)	0.12
1155	3.74 (44)		3.72 (34)	3.96 (50)			3.65 (7)			3.5 (4)	3.583 (46)	3.4 (7)	3.594 (36)	3.595 (17)	3.597 (32)	3.594 (15)	0.06
1167				0.021 (17)						0.027 (4)		0.028 (10)				0.0271 (37)	0.01
1172	0.066 (22) μ			0.113 (41)						0.098 (12)		0.132 (9)				0.120 (16)	5.1

Energy (keV)	1969Li10	1696Wa27*	1969Gr33*	1975Ha31*	1977Zn01	1982Ak03*	1982Fa10*	1983OI01	1983Sc13	1990Mouze	1991Li11	2000Sa32	2002De03	2002MoZP	2004Mo07*	Evaluated	χ^2
1207	1.03 (13)		0.89 (9)	1.10 (11)						0.98 (6)	0.991 (35)	1.04 (7)				0.998 (27)	0.18
1226				0.058 (19)						0.028 (11)		0.074 (20)				0.039 (18)	4.1
1230										0.015 (6)		0.08 (4)				0.016 (10)	2.6
1238	13.4 (13) μ	12.77 (12)	12.8 (11)	14.9 (15)	12.94 (17)		13.25 (22)	13.01 (18)	12.71 (16)	12.87 (9)	12.73 (18)	12.7 (4)	12.83 (6)	12.80 (4)	12.85 (5)	12.819 (29)	0.43
1280	3.30 (44) μ		2.92 (28)	3.59 (50)			3.22 (6)			3.17 (17)	3.144 (46)	3.15 (11)	3.147 (28)	3.159 (16)	3.151 (28)	3.155 (13)	0.05
1284										0.052 (12)		0.020 (7)				0.028 (14)	5.3
1303	0.24 (7)		0.284 (34)	0.25 (6)						0.21 (2)	0.246 (15)	0.20 (5)				0.231 (12)	0.83
1316	0.154 (44)			0.198 (50)						0.16 (2)		0.20 (3)				0.170 (16)	0.69
1330				0.024 (11)						0.026 (3)		0.039 (17)				0.0264 (30)	0.57
1341				0.050 (25)						0.046 (6)		0.059 (29)				0.047 (6)	0.19
1351			0.205 (34)							0.008 (2)		0.014 (4)				0.0092 (24)	1.8
1353				0.0099 (25)						0.008 (2)						0.008 (2)	
1377	9.5 (11) μ	8.70 (48)	9.0 (9)	9.9 (11)	8.87 (15)		8.66 (16)			8.82 (12)	8.79 (14)	8.52 (25) μ	8.689 (19)	8.79 (3)	8.720 (44)	8.722 (25)	2.5
1385	1.76 (33)	1.29 (30)	1.66 (17)	2.04 (20)						1.81 (8) μ	1.664 (40) μ	1.76 (5)	1.744 (17)	1.755 (16)	1.750 (19)	1.750 (11)	1.8
1392				0.041 (19)						0.018 (4)		0.035 (15)				0.0191 (42)	1.2
1401	3.30 (44) μ		3.03 (28)	3.47 (37)						2.91 (16)	2.792 (45)	3.0 (4)	2.924 (20)	2.934 (13)	2.927 (20)	2.923 (16)	2.3
1407	5.7 (7)		5.9 (6)	6.2 (7)						5.37 (6)	4.73 (13) μ	5.5 (5)	5.233 (26)	5.250 (19)	5.245 (42)	5.252 (17)	1.3
1419				0.0111 (25)						0.011 (3)		0.013 (3)				0.0120 (21)	0.22
1470										0.020 (3)		0.035 (15)				0.0206 (29)	0.96
1479	0.110 (44)			0.124 (50)						0.11 (1)		0.14 (3)				0.113 (9)	0.45
1509	4.84 (44)		4.77 (46)	5.45 (50)	4.78 (9)		4.77 (9)		4.57 (11)	4.76 (5)	4.64 (9)	4.63 (15)	4.61 (6)	4.67 (3)	4.64 (6)	4.679 (21)	0.95
1515										0.015 (2)		0.039 (10)				0.0159 (46)	5.5
1538	1.17 (13) μ		0.72 (7)	1.14 (12)						0.95 (6)	0.827 (31)	0.98 (5)				0.882 (49)	4.1
1543	0.75 (18)			0.74 (7)						0.68 (4)	0.44 (11)	0.67 (3)				0.664 (29)	1.5
1583	1.60 (15)		1.47 (5)	1.86 (19)			1.57 (3)			1.58 (8)	1.517 (34)	1.64 (17)		1.556 (13)		1.555 (11)	0.39
1594	0.66 (20)		0.51 (6)	0.69 (6)						0.61 (4)	0.55 (8)	0.63 (10)				0.603 (33)	0.21
1599	0.75 (20)		0.66 (7)	0.85 (11)						0.72 (4)	0.51 (12)	0.73 (7)				0.707 (33)	0.98
1636				0.040 (12)						0.024 (3)		0.06 (3)				0.0244 (36)	1.4
1657				0.16 (7)						0.10 (1)		0.14 (3)				0.104 (12)	1.6
1661	2.55 (26)		2.49 (20)	2.72 (25)			2.55 (5)			2.33 (12)	2.53 (7)	2.37 (22)	2.271 (34)	2.299 (14)	2.284 (34)	2.304(20)	2.5
1665										0.018 (3)		0.046 (9)				0.032 (14)	4.8
1683	0.53 (9)		0.52 (6)	0.56 (6)						0.49 (3)	0.475 (13)	0.43 (4)		0.481 (9)		0.478 (7)	0.52
1711												0.050 (10)				0.050 (10)	
1729	7.03 (9) μ	6.94 (20)	6.6 (6)	7.5 (7)	6.29 (10)		6.56 (12)			6.60 (4) μ	6.42 (9)	6.33 (15)	6.226 (31)	6.25 (3)	6.229 (31)	6.251 (22)	1.2
1751										0.002 (1)						0.002 (1)	

Energy (keV)	¹⁹⁶⁹ Li10	¹⁶⁹⁶ Wa27*	¹⁹⁶⁹ Gr33*	¹⁹⁷⁵ Ha31*	¹⁹⁷⁷ Zo01	¹⁹⁸² Ak03*	¹⁹⁸² Fa10*	¹⁹⁸³ OI01	¹⁹⁸³ Sc13	¹⁹⁹⁰ Mouze	¹⁹⁹¹ Li11	²⁰⁰⁰ Sa32	²⁰⁰² De03	²⁰⁰² MoZP	²⁰⁰⁴ Mo07*	Evaluated	χ^2
1764	36.7 (35) μ	35.34 (10)	34.4 (34)	40.0 (40)	34.23 (44)		34.91 (41)		33.2(7)	34.48 (25)	33.85 (46)	33.3 (10)	33.54 (10)	33.63 (9)	33.56 (10)	33.66 (10)	2.5
1813				0.026 (10)						0.024 (2)		0.020 (10)				0.0238 (20)	0.15
1838	0.81 (11)		0.88 (8)	0.89 (10)						0.74 (3)		0.77 (4)				0.753(23)	0.32
1847	5.1 (7) μ		4.76 (46)	5.32 (50)	4.52 (9)		4.59 (9)			4.57 (6)		4.35 (13)	4.448 (36)	4.42 (3)	4.457 (31)	4.451 (26)	1.6
1873	0.48 (11)		0.478 (46)	0.557 (50)						0.46 (2)		0.51 (5)				0.467 (18)	0.44
1890	0.22 (7)		0.205 (46)	0.21 (7)						0.17 (1)		0.17 (3)				0.171 (9)	0.25
1895	0.40 (9)		0.432 (46)	0.37 (6)						0.31 (2)		0.35 (4)				0.321 (18)	0.8
1898				0.136 (50)						0.11 (2)		0.10 (3)				0.107 (17)	0.08
1935			0.432 (46)	0.111 (50)						0.067 (7)		0.16 (4)				0.070 (16)	5.2
1994										0.005 (1)		0.010 (5)				0.0052 (10)	0.96
2010	0.081 (13)			0.111 (12)						0.100 (5)		0.093 (5)				0.0954 (37)	1.1
2021				0.074 (12)						0.045 (5)		0.057 (11)				0.0471 (46)	0.99
2052	0.154 (44)		0.250 (34)	0.173 (25)						0.15 (1)		0.16 (3)				0.151 (9)	0.52
2085	0.022 (7)			0.0198 (50)						0.018 (1)						0.0181 (10)	0.32
2089	0.110 (22)		0.137 (46)	0.124 (12)						0.096 (5)		0.12 (3)				0.0973 (48)	0.49
2109	0.220 (44)		0.20 (6)	0.235 (50)					0.180 (9)	0.19 (1)		0.17 (3)				0.185 (6)	0.48
2118	2.86 (33) μ	2.76 (13)	2.61 (23)	3.03 (31)	2.53 (5)		2.51 (5)		2.57 (7)	2.56 (3)		2.65 (25) μ	2.536 (20)	2.548 (21)	2.537 (20)	2.545 (12)	0.17
2147	0.026 (7)			0.036 (10)						0.029 (2)		0.050 (10)				0.0295 (28)	2.3
2160										0.004 (1)		0.026 (1)				0.015 (11)	
2176										0.007 (1)		0.015 (6)				0.0072 (13)	1.7
2192	0.154 (44)			0.161 (50)					0.070 (13)	0.073 (6)		0.093 (5)				0.084 (7)	3.4
2204	11.7 (11) μ	11.22 (47)	11.37 (24)	12.38 (27)	10.77 (20)		10.66 (20)		10.95 (70)	11.02 (9)		11.1 (3)	10.74 (5)	10.75 (9)	10.76 (5)	10.80 (5)	1.8
2251				0.015 (7)						0.012 (1)						0.012 (1)	
2260			0.057 (23)	0.0149 (50)						0.019 (1)		0.020 (3)				0.0191 (9)	0.1
2266	0.033 (7)			0.045 (12)						0.037 (2)		0.034 (4)				0.0362 (17)	0.34
2270				0.0111 (50)						0.0029 (5)		0.010 (5)				0.0030 (7)	2.0
2284										0.011 (1)		0.011 (3)				0.0110 (9)	
2287										0.010 (1)						0.010 (1)	
2293	0.73 (9)		0.67 (7)	0.83 (9)			0.67 (2)		0.662 (20)	0.67 (3)		0.72 (6)	0.665 (17)	0.677 (10)	0.665 (17)	0.673 (8)	0.57
2310										0.003 (2)						0.003 (2)	
2312	0.020 (7)			0.0235 (50)						0.019 (2)		0.018 (5)				0.0189 (18)	0.029
2319										0.0009 (3)		0.0050 (10)				0.0030 (20)	8.4
2325				0.0040 (20)						0.0037 (4)		0.009 (3)				0.0038 (7)	3.1
2331	0.046 (9)		0.034 (11)	0.0557 (50)						0.048 (3)		0.076 (7)				0.056 (9)	5.7
2348										0.0003 (2)						0.0003 (2)	

Energy (keV)	1969Li10	1696Wa27*	1969Gr33*	1975Ha31*	1977Zn01	1982Ak03*	1982Fa10*	1983OI01	1983Sc13	1990Mouze	1991Li11	2000Sa32	2002De03	2002MoZP	2004Mo07*	Evaluated	χ^2
2353										0.0008 (3)						0.0008 (3)	
2361				0.0040 (12)						0.0033 (3)		0.0060 (10)				0.0046 (14)	3.6
2369										0.006 (1)		0.008 (3)				0.0062 (9)	0.4
2376	0.0132 (44)		0.057 (23)	0.022 (7)						0.019 (1)		0.034 (7)				0.0190 (17)	3.2
2390				0.0042 (10)						0.0034 (3)		0.006 (3)				0.00343 (30)	0.74
2405										0.0009 (3)		0.0040 (10)				0.0024 (16)	4.8
2423	0.0132 (44)			0.0115 (40)						0.010 (1)		0.018 (4)				0.0106 (14)	2.1
2444										0.018 (9)						0.018 (9)	
2447	3.63 (40) μ	3.32 (6)	3.79 (28)	3.96 (37)	3.32 (8)		3.28 (6)			3.42 (3)		3.30 (10)	3.402 (24)	3.41 (4)	3.408 (24)	3.403 (16)	0.50
2459										0.0031 (5)						0.0031 (5)	
2482				0.0046 (19)						0.0021 (4)						0.0021 (4)	6.1
2505	0.0154 (44)			0.0149 (37)						0.012 (1)		0.025 (7)				0.0124 (13)	1.9
2550										0.0007 (2)						0.0007 (2)	
2562										0.0004 (3)						0.0004 (3)	
2564										0.0003(2)						0.0003(2)	
2604				0.00099 (25)						0.0008 (2)						0.0008 (2)	
2630				0.0020 (10)						0.0018 (3)		0.0050 (17)				0.0019 (5)	3.4
2662										0.0006 (2)		0.0004 (1)				0.00044 (9)	0.8
2694	0.068 (9)		0.072 (34)	0.079 (7)			0.078 (2)			0.066 (3)		0.062 (4)				0.072 (6)	4.5
2699	0.0110 (44)			0.0050 (19)						0.0061 (5)						0.0062 (5)	1.2
2719	0.0033 (11)			0.0040 (12)						0.0038 (4)						0.00374 (38)	0.18
2769	0.057 (9)		0.057 (23)	0.062 (7)			0.047 (2)			0.053 (3)		0.048 (15)				0.0494 (17)	1.2
2785	0.0110 (22)			0.0149 (25)						0.012 (1)		0.030 (11)				0.0120 (11)	1.4
2826	0.0046 (11)			0.0062 (12)						0.0048 (4)		0.011 (6)				0.00480 (38)	0.55
2861				0.00074 (37)						0.0009 (2)		0.008 (5)				0.00091 (28)	2.01
2880	0.0176 (33)		0.019 (6)	0.024 (7)						0.020 (2)		0.030 (3)				0.0222 (35)	4.8
2893	0.0132 (33)		0.016 (7)	0.0149 (37)						0.012 (1)		0.017 (3)				0.0126 (10)	1.3
2921	0.035 (7)		0.032 (11)	0.037 (6)						0.029 (1)		0.035 (4)				0.0295 (11)	1.4
2928				0.0026 (10)						0.0024 (2)						0.0024 (2)	
2934				0.00124 (50)						0.0010 (2)		0.005 (3)				0.00102 (27)	1.8
2978	0.031 (7)		0.038 (23)	0.037 (6)			0.029 (2)			0.030 (1)		0.034 (7)				0.0302 (9)	0.85
2999	0.0220 (44)		0.015 (7)	0.024 (6)						0.019 (1)		0.030 (5)				0.0195 (15)	2.5
3053	0.046 (7)		0.046 (23)	0.053 (7)						0.041 (2)		0.057 (3)				0.048 (7)	1.8
3081	0.0110 (44)			0.0124 (37)						0.011 (1)		0.020 (4)				0.0115 (15)	2.4
3093				0.00111 (37)						0.0008 (1)		0.0010 (3)				0.00082 (9)	0.4

Energy (keV)	1969Li10	1696Wa27*	1969Gr33*	1975Ha31*	1977Zn01	1982Ak03*	1982Fa10*	1983OI01	1983Sc13	1990Mouze	1991Li11	2000Sa32	2002De03	2002MoZP	2004Mo07*	Evaluated	χ^2
3142	0.0022 (9)			0.0035 (12)						0.0026 (2)		0.0060 (28)				0.00260 (19)	0.84
3149										0.00019						0.00019	
3160	0.00110 (44)			0.00111 (50)						0.0010 (2)		0.0030 (17)				0.00104 (18)	0.7
3183	0.00110 (44)			0.0032 (10)						0.0028 (2)		0.0060 (10)				0.0023 (10)	1.3

*: Not used by the evaluators (see below).

μ : the experimental value has been shown to be outlier value by the Lweight program.

There were omitted from analysis:

- four sets of values, A. Hachem (1975Ha31), G. Mouze (1981Mo28), H. Akcay (1982Ak03), G. Mouze (1990Mo08) and O. Diallo (1993Di09), because these values come from the same laboratory of G. Mouze (1990Mo**).
- the sets of values from K. Ya. Gromov (1969Gr33), G. Wallace (1969Wa27) and M. A. Farouk (1982Fa10), because of a lack of information in the articles about the experimental measurements carried out and, therefore on the results.
- the relative γ -ray intensity values given in 2004Mo07, because they are those measured by J. U. Delgado (2002De03). In 2004Mo07, the author measured the absolute 609.3-keV γ -ray emission probability (Table 5) and normalized the 2002De03 data set with their value of 45.57 (18).

Table 6: Evaluated relative and absolute γ -ray intensities.

Energy (keV)	Relative γ -ray intensity (%)	Absolute γ -ray intensity (%)	Energy (keV)	Relative γ -ray intensity (%)	Absolute γ -ray intensity (%)	Energy (keV)	Relative γ -ray intensity (%)	Absolute γ -ray intensity (%)	Energy (keV)	Relative emission intensity (%)	Absolute γ -ray intensity (%)
221	0.130 (13)	0.059 (6)	703	1.053(24)	0.479 (11)	1238	12.819 (29)	5.831 (14)	2204	10.80 (5)	4.913 (23)
230	0.0063 (21)	0.0029 (10)	704	0.112 (21)	0.051 (10)	1280	3.155 (13)	1.435 (6)	2251	0.012 (1)	0.0055 (5)
252	0.0258 (39)	0.0117 (18)	708	0.0262 (43)	0.0119 (20)	1284	0.028 (14)	0.013 (6)	2260	0.0191 (9)	0.0087 (4)
268	0.0355 (40)	0.0161 (18)	710	0.168 (8)	0.076 (4)	1303	0.231 (12)	0.105 (5)	2266	0.0362 (17)	0.0165 (8)
273	0.264 (18)	0.120 (8)	719	0.865 (22)	0.393 (10)	1316	0.170 (16)	0.077 (7)	2270	0.0030 (7)	0.0014 (3)
280	0.136 (14)	0.062 (6)	722	0.082 (15)	0.037 (7)	1330	0.0264 (30)	0.0120 (14)	2284	0.0110 (9)	0.0050(4)
304	0.056 (5)	0.0255 (23)	733	0.084 (7)	0.038 (3)	1341	0.047 (6)	0.0214 (27)	2287	0.010 (1)	0.0046 (5)
333	0.139 (9)	0.063 (4)	740	0.0941 (47)	0.0428 (21)	1351	0.0092 (24)	0.0042 (11)	2293	0.673 (8)	0.306 (4)
334	0.072 (10)	0.033 (5)	752	0.278 (17)	0.126 (8)	1353	0.008 (2)	0.0036 (9)	2310	0.003 (2)	0.0014 (9)
348	0.27 (7)	0.123 (32)	768	10.755 (36)	4.892 (16)	1377	8.722 (25)	3.968 (11)	2312	0.0189 (18)	0.0086 (8)
386	0.650 (12)	0.296 (5)	786	0.69 (10)	0.31 (5)	1385	1.750 (11)	0.796 (5)	2319	0.0030 (20)	0.0014 (9)
388	0.864 (10)	0.394 (5)	788	0.033 (10)	0.015 (5)	1392	0.0191 (42)	0.0087 (19)	2325	0.0038 (7)	0.0017 (3)
394	0.0280 (40)	0.0127 (18)	806	2.774 (13)	1.262 (6)	1401	2.923 (16)	1.330 (7)	2331	0.056 (9)	0.026 (4)
396	0.057 (4)	0.0259 (18)	815	0.085 (7)	0.039 (3)	1407	5.252 (17)	2.389 (8)	2348	0.003 (2)	0.0014 (9)
405	0.375 (16)	0.171 (7)	821	0.364 (21)	0.166 (10)	1419	0.0120 (21)	0.0055 (10)	2353	0.0008 (3)	0.00036 (14)
452	0.067 (8)	0.031 (4)	826	0.284 (24)	0.129 (11)	1470	0.0206 (29)	0.0094 (13)	2361	0.0046 (14)	0.0021 (6)
454	0.634 (10)	0.288 (5)	832	0.076 (5)	0.035 (2)	1479	0.113 (9)	0.051 (4)	2369	0.0062 (9)	0.0028 (4)
461	0.128 (18)	0.058 (8)	847	0.035 (13)	0.016 (6)	1509	4.679 (21)	2.128 (10)	2376	0.0190 (17)	0.0086 (8)
469	0.292 (32)	0.133 (15)	873	0.041 (7)	0.019 (3)	1515	0.0159 (46)	0.0072 (21)	2390	0.00343 (30)	0.00156 (14)
474	0.203 (14)	0.092 (6)	878	0.026 (6)	0.0118 (27)	1538	0.882 (49)	0.401 (22)	2405	0.0024 (16)	0.0011 (7)
485	0.046 (8)	0.021 (4)	904	0.144 (17)	0.066 (8)	1543	0.664 (29)	0.302 (13)	2423	0.0106 (14)	0.0048 (6)
487	0.061 (20)	0.028 (9)	915	0.051 (11)	0.023 (5)	1583	1.555 (11)	0.707 (5)	2444	0.018 (9)	0.008 (4)
494	0.023 (6)	0.011 (3)	917	0.010 (7)	0.005 (3)	1594	0.603 (33)	0.274 (15)	2447	3.403 (16)	1.548 (7)
496	0.015 (4)	0.0068 (18)	930	0.094 (17)	0.043 (8)	1599	0.707 (33)	0.322 (15)	2459	0.0031 (5)	0.00141 (23)
501	0.0397 (48)	0.0181 (22)	934	6.814 (22)	3.100 (10)	1636	0.0244 (36)	0.0111 (16)	2482	0.0021 (4)	0.00096 (18)
519	0.0364 (38)	0.0166 (17)	939	0.036 (8)	0.016 (4)	1657	0.104 (12)	0.047 (5)	2505	0.0124 (13)	0.0056 (6)

Energy (keV)	Relative γ -ray intensity (%)	Absolute γ -ray intensity (%)	Energy (keV)	Relative γ -ray intensity (%)	Absolute γ -ray intensity (%)	Energy (keV)	Relative γ -ray intensity (%)	Absolute γ -ray intensity (%)	Energy (keV)	Relative emission intensity (%)	Absolute γ -ray intensity (%)
524	0.0372 (38)	0.0169 (17)	943	0.038 (6)	0.017 (3)	1661	2.304(20)	1.048 (9)	2550	0.0007 (2)	0.00032 (9)
528	0.0239 (29)	0.0109 (13)	949	0.012 (5)	0.0055 (23)	1665	0.032 (14)	0.015 (6)	2562	0.0004 (2)	0.00018 (9)
536	0.134 (17)	0.061 (8)	952	0.013 (5)	0.0059 (23)	1683	0.478 (7)	0.217 (3)	2564	0.0003(2)	0.00014 (9)
543	0.194 (46)	0.088 (21)	961	0.0222 (30)	0.0101 (14)	1711	0.050 (10)	0.023 (5)	2604	0.0008 (2)	0.00036 (9)
547	0.075 (6)	0.034 (3)	964	0.799 (27)	0.363 (12)	1729	6.251 (22)	2.844 (10)	2630	0.0019 (5)	0.00086 (23)
551	0.012 (3)	0.0055 (14)	976	0.0333 (47)	0.0151 (21)	1751	0.002 (1)	0.0009 (5)	2662	0.00044 (9)	0.00020 (4)
572	0.156 (17)	0.071 (8)	991	0.023 (6)	0.011 (3)	1764	33.66 (10)	15.31 (5)	2694	0.072 (6)	0.033 (3)
595	0.0383 (33)	0.0174 (15)	1013	0.0191 (41)	0.0087 (19)	1813	0.0238 (20)	0.0108 (9)	2699	0.0062 (5)	0.00282 (23)
600	0.018 (8)	0.008 (4)	1021	0.034 (6)	0.016 (3)	1838	0.753(23)	0.343 (10)	2719	0.00374 (38)	0.00170 (17)
609	100	45.49 (19)	1032	0.135 (9)	0.061 (4)	1847	4.451 (26)	2.025 (12)	2769	0.0494 (17)	0.0225 (8)
615	0.121 (16)	0.055 (7)	1038	0.0190 (33)	0.0086 (15)	1873	0.467 (18)	0.212 (8)	2785	0.0120 (11)	0.0055 (5)
617	0.059 (10)	0.027 (5)	1045	0.050 (6)	0.023(3)	1890	0.171 (9)	0.078 (4)	2826	0.00480 (38)	0.00218 (17)
626	0.009 (3)	0.0041 (14)	1051	0.713 (17)	0.324 (8)	1895	0.321 (18)	0.146 (8)	2861	0.00091 (28)	0.00041 (13)
630	0.0366 (31)	0.0166 (14)	1067	0.053 (15)	0.024 (7)	1898	0.107 (17)	0.049 (8)	2880	0.0222 (35)	0.0101 (16)
633	0.120 (7)	0.055 (3)	1069	0.595 (23)	0.271 (10)	1935	0.070 (16)	0.032 (7)	2893	0.0126 (10)	0.0057 (5)
634	0.014 (5)	0.0064 (23)	1103	0.233 (33)	0.106 (15)	1994	0.0052 (10)	0.0024 (5)	2921	0.0295 (11)	0.0134 (5)
639	0.075 (10)	0.034 (5)	1104	0.16 (3)	0.073 (14)	2010	0.0954 (37)	0.0434 (17)	2928	0.0024 (2)	0.00109 (9)
649	0.119 (16)	0.054 (7)	1118	0.022 (9)	0.010 (4)	2021	0.0471 (46)	0.0214 (21)	2934	0.00102 (27)	0.00046 (12)
658	0.038 (8)	0.017 (4)	1120	32.77 (7)	14.91 (3)	2052	0.151 (9)	0.069 (4)	2978	0.0302 (9)	0.0137 (4)
661	0.118 (9)	0.054 (4)	1130	0.079 (7)	0.036 (3)	2085	0.0181 (10)	0.0082 (5)	2999	0.0195 (15)	0.0089 (7)
665	3.364 (15)	1.530 (7)	1133	0.558 (17)	0.254 (8)	2089	0.0973 (48)	0.0443 (22)	3053	0.048 (7)	0.022 (3)
677	0.012 (5)	0.0055 (23)	1155	3.594 (15)	1.635 (7)	2109	0.185 (6)	0.084 (3)	3081	0.0115 (15)	0.0052 (7)
683	0.184 (13)	0.084 (6)	1167	0.0271 (37)	0.0123 (17)	2118	2.545 (12)	1.158 (5)	3093	0.00082 (9)	0.00037 (4)
687	0.0146 (31)	0.0066 (14)	1172	0.120 (16)	0.055 (7)	2147	0.0295 (28)	0.0134 (13)	3142	0.00260 (19)	0.00118 (9)
693	0.0129 (33)	0.0059 (15)	1207	0.998 (27)	0.454 (12)	2160	0.015 (11)	0.007 (5)	3149	0.00019	0.00019
697	0.148 (9)	0.067 (4)	1226	0.039 (18)	0.018 (8)	2176	0.0072 (13)	0.0033 (6)	3160	0.00104 (18)	0.00047 (8)
699	0.035 (10)	0.016(5)	1230	0.016 (10)	0.007 (5)	2192	0.084 (7)	0.038 (3)	3183	0.0023 (10)	0.0011 (5)

6 References

- 1956Da06 H. Daniel, Z. Naturforsch. 11a(1956)212 [Half-life].
- 1960Wa14 R. J. Walen, G. Bastin-Scoffier, Nucl. Phys. 16(1960)246 [α branching ratio, I_{α}].
- 1965Le08 C.-F. Leang, Compt. Rend. Acad. Sci. (Paris) 260(1965)3037 [E_{α} , I_{α}].
- 1969Gr33 K. Ya. Gromov, B. M. Sabirov, J. J. Urbanets, Bull. Acad. Sci. USSR, Phys. Ser. 33(1970)1510 [I].
- 1969Li10 E. W. A. Lingeman, J. Konijn, P. Polak, A. H. Wapstra, Nucl. Phys. A133(1969)630 [I _{γ}].
- 1969Wa27 G. Wallace, G. E. Coote, Nucl. Instrum. Meth. 74(1969)353 [I _{γ}].
- 1975Ha31 A. Hachem, Compt. Rend. (Paris) 281B(1975)45 [I _{γ}].
- 1977Zo01 V. Zobel, J. Eberth, U. Eberth, E. Eube, Nucl. Instrum. Meth. 141(1977)329 [I _{γ}].
- 1981Mo28 G. Mouze, Compt. Rend. (Paris) 292(1981)1243 [I _{γ}].
- 1982Ak03 H. Akcay, G. Mouze, D. Maillard, Ch. Ythier, Radiochem. Radioanal. Lett. 51(1982)1 [I _{γ}].
- 1982Fa10 M. A. Farouk, A. M. Al-Soraya, Nucl. Instrum. Meth. 200(1982)593 [I _{γ}].
- 1983Ol01 D. G. Olson, Nucl. Instrum. Meth. 206(1983)313 [I _{γ}].
- 1983Sc13 U. Schötzig, K. Debertain, Int. J. Appl. Radiat. Isot. 34(1983)533 [I _{γ}].
- 1988Ak01 Y. A. Akovali, Nucl. Data Sheets 55(1988)665 [Energy level, spin, parity and multipolarity].
- 1990Mo08 G. Mouze, O. Diallo, P. Bechlich, C. Ythier, J. F. Comanducci, Radiochimica Acta 49(1990)13 [I].
- 1990Mo** G. Mouze, C. Ythier, J. F. Comanducci, Rev. Roumaine Phys. 35(1990)337 [I _{γ}].
- 1991Li11 W. -J. Lin, G. Harbottle, J. Radioanal. Nucl. Chem. Lett. 153(1991)137 [I _{γ}].
- 1993Di09 O. Diallo, G. Mouze, C. Ythier, J. F. Comanducci, Nuovo Cimento 106A(1993)1321 [I _{γ}].
- 1995El07 Y. A. Akovali, Nucl. Data Sheets 76(1995)127 [Energy level, spin, parity and multipolarity].
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
- 1998Mo14 J. Morel, M. Etcheverry, J. L. Picolo, Appl. Radiat. Isot. 49(1998)1387 [I _{γ}].
- 2000Sa32 D. Sardari, T. D. MacMahon, J. Radioanal. Nucl. Chem. 244(2000)463 [I _{γ}].
- 2002Ba85 I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr., P. O. Tikkanen, S. Raman, At. Data Nucl. Data Tables 81(2002)1 [Theoretical ICC].
- 2002De03 J. U. Delgado, J. Morel, M. Etcheverry, Appl. Radiat. Isot. 56(2002)137 [I _{γ}].
- 2002MoZP G. L. Molnar, Z. S. Révay, T. Belgya, 11th Int. Symp. on capture gamma-ray spectroscopy, 2-6 Sep. 2002, Pruhonice (2003)522 [I _{γ}].
- 2002Ba85 I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr., P. O. Tikkanen, S. Raman, Atomic Data Nucl. Data Tables 81(2002)1 [Theoretical ICC].
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].
- 2003Br13 E. Browne, Nucl. Data Sheets 99(2003)483 [Energy level, spin, parity and multipolarity].
- 2004He** R. G. Helmer, IAEA – CRP Report to be published (2004) [I _{γ}].
- 2004Mo07 J. Morel, S. Speman, M. Rasko, E. Terechtchenko, J. U. Delgado, Appl. Radiat. Isot. 60(2004)341 [I _{γ}].

**²¹⁴Po - Comments on evaluation of decay data
by V. Chisté and M. M. Bé**

This evaluation was completed in 2007. Literature available by January 2007 was included.

1 Decay Scheme

²¹⁴Po disintegrates by alpha emissions mainly to the ground state level of ²¹⁰Pb. Spins and parities are from the mass-chain evaluation of Y. A. Akovali (1995El07 for A = 214) and E. Browne (1992Br01 and 2003Br13 for A = 210).

A good agreement was found between the recommended Q value of Audi and the effective Q value (7833.24 (10) keV) calculated from decay scheme data.

2 Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2003Au03).

Experimental ²¹⁴Po half-life values (in μ s) are given in Table 1:

Table 1: Experimental values of ²¹⁴Po half-life.

Reference	Experimental value (μ s)	Comments
J. V. Dunworth (1939Du**)	150 (20)	Not used.
J. Rotblat (1941Ro**)	145 (5)	Not used.
A. G. Ward (1942Wa04)	148 (6)	Not used.
J. C. Jacobsen (1943Ja**)	155 (5)	Not used.
G. von Dardel (1950Vo02)	163.7 (18)	Original uncertainty increased
R. Ballini (1953Ba60)	158 (2)	
K. W. Ogilvie (1960Og01)	159.5 (30)	
T. Dobrowolski (1961Do02)	164.3 (18)	
A. Erlik (1971Er02)	165 (3)	
J. W. Zhou (1993Zh30)	160 (12)	
Recommended value	162.3 (12)	$\chi^2 = 1.6$

The first four and less precise historical values were omitted from analysis. The G. von Dardel uncertainty value (1950Vo02) of 0.2, which seems not realistic, was increased to 1.8 the smallest of the other experimental values obtained with the same method.

Using the LWEIGHT computer program (version 3) with the remaining set of 6 data the weighted average is **162.3 ms** with an external uncertainty of **1.2 ms**. The reduced- χ^2 value is 1.82.

The largest contribution to weighted average comes from the value of G. von Dardel (1950Vo02) and T. Dobrowolski (1961Do02), each of them amounting per 28 %.

2.1 a Transitions and Emissions

The energies of the α -particle transitions given in Section 2.1 were obtained from Q_α (2003Au03) and level energies.

The energy of $\alpha_{0,0}$ emission given in section 4 is the weighted average of the measured values of A. Rytz (1961Ry02) and B. Grennberg (1971Gr17), with the recommendations given by A. Rytz (1991Ry01) where the original energies given by 1961Ry02 and 1971Gr17 have been readjusted due to changes in calibration

energies. For the $\alpha_{0,1}$ and $\alpha_{0,2}$, the emission energies were deduced from Q_α (2003Au03), level energy and taking the nucleus recoil into account.

The α emission probabilities have been deduced from the value of the γ -ray transition probability decay-scheme balances for the corresponding levels. (see **2.2 Gamma Transitions**).

2.2 g Transitions

The γ -ray transition probabilities were obtained using the γ -ray emission intensities, measured by 1976Ku08, and the relevant internal conversion coefficients (see **4.2 g Emissions**).

Multipolarities of the γ -ray transitions (E2) are from 1992Br01 and 2003Br13.

The internal conversion coefficients (ICC) for the γ -ray transitions have been deduced using the BrIcc computer program (calculation for 'hole'), which interpolated the new values from 2006Ra03.

3 Atomic Data

Atomic values, ω_K , $\overline{\omega}_L$ and n_{KL} and the X-ray relative probabilities are from Schönfeld and Jaßien (1996Sc06).

4 Photon Emissions

4.1 X-ray Emissions

The X-ray absolute intensities were calculated from γ -ray data and ICC using the EMISSION computer program.

4.2 g Emissions

The γ -ray energies given in section 5.2 are from W. Kurcewicz (1976Ku08).

The absolute γ -ray emission intensities have been deduced from the relative γ -ray emission intensities measured by W. Kurcewicz (1976Ku08) in relative value and normalized with the 324.2-keV γ -ray in ²²²Ra decay, as measured by A. Peghaire (1969Pe17) to be 2.77 (8) %. In the table 2, the relative emission intensities and the recommended values of absolute emission intensities are shown.

Table 2: Recommended (deduced) values of γ -ray absolute emission intensities

Energy (keV)	Relative Emission Intensity (%)	Recommended value
298 (1)	0.06 (2)	0.000052 (18) %
799.7 (1)	11.9 (5)	0.0104 (6) %

5 References

- 1939Du** – J. V. Dunworth, Nature (London) 144(1939)152 [Half-life].
 1941Ro** – J. Rotblat, Proc. Roy. Soc. (London) A177(1941)260 [Half-life].
 1942Wa04 – A. G. Ward, Proc. Roy. Soc. (London) A181(1942)183 [Half-life].
 1943Jo** – J. C. Jacobsen, T. Sigurgeirsson, Kgl. D. Vid. Selsk. Medd. 20(1943)11 [Half-life].
 1950Vo02 – G. von Dardel, Phys. Rev. 79(1950)734 [Half-life].
 1953Ba60 – R. Ballini, Ann. Phys. (Paris) 8(1953)441 [Half-life].
 1960Og01 – K. W. Ogilvie, Proc. Phys. Soc. (London) 76(1960)299 [Half-life].

- 1961Do02 – T. Dobrowolski, J. Young, Proc. Phys. Soc. (London) 77(1961)1219 [Half-life].
- 1961Ry02 – A. Rytz, Helv. Phys. Acta 34(1961)240 [E_{α}].
- 1969Pe17 – A. Peghaire, Nucl. Instrum. Meth. 75(1969)66 [I_{γ}].
- 1971Er02 – A. Erlik, J. Felsteiner, H. Lindeman, M. Tatcher, Nucl. Instrum. Meth. 92(1971)45 [Half-life].
- 1971Gr17 – B. Grennberg, A. Rytz, Metrologia 7(1971)65 [E_{α}].
- 1976Ku08 – W. Kurcewicz, N. Kaffrell, N. Trautmann, A. Plochocki, J. Kylicz, K. Stryczniewicz, I. Yutlandov, Nucl. Phys. A270(1976)175 [I_{γ} , E_{γ} , I_{α}].
- 1991Ry01 – A. Rytz, At. Data and Nucl. Data Tables 47(1991)205 [E_{α} , I_{α}].
- 1992Br01 – E. Browne, Nucl. Data Sheets 65(1992)209 [Energy level, spin, parity and multipolarity].
- 1993Zh30 – J. W. Zhou, P. de Marcillac, N. Coron, S. Wang H. H. Stroke, O. Redi, J. Leblanc, G. Dambier, M. Barthelemy, J. P. Torre, O. Testard, G. Beyer, H. Ravn, J. Mangin, Nucl. Instrum. Meth. Phys. Res. A335(1993)443 [Half-life].
- 1995E107 – Y. A. Akovali, Nucl. Data Sheets 76(1995)127 [I_{α} , E_{α} , I_{γ} , E_{γ} , spin and parity].
- 1996Sc06 – E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
- 1998Ak04 – Y. A. Akovali, Nucl. Data Sheets 84(1998)1 [I_{α} , E_{α}].
- 2002Ba85 – I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr, P. O. Tikkanen, S. Raman, At. Data Nucl. Data Tables 81(2002)1 [Theoretical ICC].
- 2003Au03 – G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].
- 2003Br13 – E. Browne, Nucl. Data Sheets 99(2003)483 [Energy level, spin, parity and multipolarity].
- 2006Ra03 – S. Raman, M. Ertugrul, C. W. Nestor, Jr., M. B. Trzhaskovskaya. At. Data Nucl. Data Tables 92, 207 (2006) *Ratios of internal conversion coefficients*

**²¹⁵Bi – Comments on evaluation of decay data
by A. L. Nichols and F. G. Kondev**

Evaluated: June 2011

Evaluation Procedure

Limitation of Relative Statistical Weight Method (LWM) was applied to average numbers throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

The ²¹⁵Bi ground state ($J^\pi = (9/2^-)$) decays 100 % by β^- emission to various excited levels and the ground state of ²¹⁵Po. A reasonably complex but inadequate decay scheme has been constructed primarily from the gamma-ray measurements of Kurpeta *et al.* (2003Ku26) in which 19 distinct gamma-ray emissions were identified with the β^- decay of ²¹⁵Bi. Although these authors assessed that there is no direct beta decay to the ground state of ²¹⁵Po, their reported absolute emission probabilities for the gamma rays populating the ground state are in conflict with this proposal.

Direct β^- feeding to the ground state of daughter ²¹⁵Po has not been satisfactorily determined. Therefore, the evaluators resorted to comparisons with the β^- decay of other odd-even Bi radionuclides (²¹³Bi) and β^- -decay theory in order to define the β^- and γ emission probabilities in absolute terms. Further studies are required to clarify and define more clearly the ²¹⁵Bi decay scheme, particularly with respect to the absolute gamma-ray emission probabilities and quantification of direct β^- feeding to the ground state of daughter ²¹⁵Po.

Nuclear Data

²¹⁵Bi is part of the (4n + 3) naturally-occurring decay chain, and of relevance in quantifying the environmental impact of ²³⁵U and decay-chain products. Specific radionuclides in this decay chain are noteworthy because of their decay characteristics (²¹⁵Po, ²¹¹Bi and ²¹¹Po alpha decay).

Half-life

²¹⁵Bi was first observed by 1953Hy83, and assigned a half-life of (8 ± 2) min. However, the recommended half-life is the weighted mean of three more recent measurements (1965Nu03, 1989Bu09 and 1990Ru02).

Reference	Half-life (min)
1965Nu03	7.4 (6)
1989Bu09	7.5 (4)
1990Ru02	7.7 (2)
Recommended value	7.6 (2)

²¹⁵Po half-life of 1.781 (4) millisecond was adopted from the evaluation of Browne (2001Br31).

Q value

Q^- of 2189 (15) keV was adopted from the evaluated tabulations of Audi *et al.* (2003Au03).

Beta particlesEnergies

All beta-particle energies were calculated from the structural details of the proposed decay scheme. A combination of nuclear level energies recommended by 2001Br31 and derived from 2003Ku26, and a Q-value of 2189 (15) keV (2003Au03) were used to determine the energies and uncertainties of the beta-particle emissions to the various levels.

Adopted nuclear levels of ²¹⁵Po: Energy, J^π and origins (2001Br31, 2003Ku26).

Nuclear level	Nuclear level energy (keV)	J ^π	Origins
0	0.0	9/2 +	²¹⁵ Bi β ⁻ decay, ²¹⁹ Rn α decay
1	271.228 ± 0.010	7/2 +	²¹⁵ Bi β ⁻ decay, ²¹⁹ Rn α decay
2	293.56 ± 0.04	11/2 +	²¹⁵ Bi β ⁻ decay, ²¹⁹ Rn α decay
3	401.812 ± 0.010	5/2 +	²¹⁵ Bi β ⁻ decay, ²¹⁹ Rn α decay
4	517.60 ± 0.06	7/2 +, 9/2 +	²¹⁵ Bi β ⁻ decay, ²¹⁹ Rn α decay
5	608.30 ± 0.07	(11/2 +, 13/2 +)	²¹⁵ Bi β ⁻ decay, ²¹⁹ Rn α decay
6	676.66 ± 0.07		²¹⁵ Bi β ⁻ decay, ²¹⁹ Rn α decay
7	708.1 ± 0.5		²¹⁹ Rn α decay
8	732.7 ± 0.4		²¹⁹ Rn α decay
9	835.32 ± 0.22		²¹⁵ Bi β ⁻ decay, ²¹⁹ Rn α decay
10	877.2 ± 0.6		²¹⁹ Rn α decay
11	891.1 ± 0.3		²¹⁹ Rn α decay
12	930 ± 1		²¹⁹ Rn α decay
13	1073.7 ± 0.4	(5/2 +)	²¹⁹ Rn α decay
14	1077.6 ± 2.0*		²¹⁵ Bi β ⁻ decay
15	1094.2 ± 1.0		²¹⁹ Rn α decay
16	1176.2 ± 2.0*		²¹⁵ Bi β ⁻ decay
17	1294.5 ± 0.1*		²¹⁵ Bi β ⁻ decay
18	1398.8 ± 0.4*		²¹⁵ Bi β ⁻ decay

* Calculated from the energies of the depopulating gamma rays (2003Ku26), and the lower-energy nuclear levels that they populate.

Emission Probabilities

Direct beta-particle feeding to the ground state of ²¹⁵Po has not been unambiguously defined from the various γ-ray measurements. Under these circumstances, a systematic assessment of the appropriate properties of odd-even Bi nuclides in the vicinity of ²¹⁵Bi has been undertaken to explore whether a reasonable approximation can be made of beta decay directly to the ground state of ²¹⁵Po (1991Ma16, 2001Br31, 2003Ak06, 2004Br45, 2007Ba19).

(a) Spin and parity of ²¹⁵Bi

Nuclide	²⁰⁹ Bi	²¹¹ Bi	²¹³ Bi	²¹⁵ Bi	²¹⁷ Bi
β ⁻ decay	stable	0.28 %	97.91 %	100 %	100 %
Direct β ⁻ decay to ground state	–	0.28 %	65.9 %	?	?
α decay	stable	99.72 %	2.09 %	–	–
Spin and parity	9/2 ⁻	9/2 ⁻	9/2 ⁻	(9/2 ⁻)	?
Spin and parity of Po ground state	1/2 ⁻	9/2 ⁺	9/2 ⁺	9/2 ⁺	(11/2 ⁺)

Spins and parities of 9/2⁻ are well defined for ^{209,211,213}Bi, and can be similarly assigned with reasonable confidence as (9/2⁻) for ²¹⁵Bi.

(b) Direct beta-particle feeding of ²¹⁵Bi to the ground state of ²¹⁵Po

Population-depopulation balances have been calculated on the basis of the relative emission probabilities of the gamma rays (see below) in order to derive relative beta-particle emission probabilities to all of the excited nuclear levels of ²¹⁵Po.

The β⁻ decay of ²¹⁵Bi was assumed to occur primarily via first forbidden non-unique transitions to the ground state (9/2⁺) and 293.56-keV nuclear level (11/2⁺) of ²¹⁵Po. The preparation of recommended decay-data files for DDEP necessitates the formulation of decay schemes that are based on absolute emission and transition probabilities that frequently encompass well-defined normalization factors in conjunction with accurate relative emission probabilities and various other nuclear parameters (e.g. internal conversion coefficients). This ideal situation cannot be achieved for ²¹⁵Bi because of existing inadequacies in the measured data. Therefore, the main β⁻ branches populate the 293.56-keV nuclear level and ground state of ²¹⁵Po, and their important emission probabilities have been derived somewhat unusually through application of the fifth-power law of β⁻ decay (1933Sa01, 1955Ev23, 1963KaZZ).

A general approximation has been formulated for the ratio of allowed beta-particle emission probabilities, based on the observation that the mean life (τ) for partial β⁻ decay is inversely proportional to the fifth power of the β⁻ end-point energy (1955Ev23, 1963KaZZ):

$$\frac{1}{\tau_{\beta}} \propto [(M(Z) - M(Z \pm 1)c^2)]^5$$

where

$$\tau_{\beta} = \frac{\tau_{exp}}{P_{\beta}} \quad \text{and} \quad \tau_{exp} \text{ is the lifetime of the parent nuclide.}$$

Therefore

$$\frac{1}{\tau_{\beta}} \sim (E_{\beta})^5$$

This approximation has been applied to the major first-forbidden non-unique beta-particle emissions of ²¹⁵Bi directly to the ground state of ²¹⁵Po ((9/2⁻) → 9/2⁺)

$$\frac{1}{\tau_{0,0}} \sim (E_{\beta_{0,0}})^5 \tag{1}$$

and to the 293.56-keV nuclear level of ²¹⁵Po ((9/2⁻) → 11/2⁺)

$$\frac{1}{\tau_{0,2}} \sim (E_{\beta_{0,2}})^5 \tag{2}$$

Combining equations (1) and (2):

$$\frac{\tau_{0,2}}{\tau_{0,0}} = \frac{P_{\beta_{0,0}}}{P_{\beta_{0,2}}} \sim \left(\frac{E_{\beta_{0,0}}}{E_{\beta_{0,2}}}\right)^5 = \left[\frac{2189(15)}{1895(15)}\right]^5 = 1.155^5 \sim 2.055$$

where $P_{\beta_{0,0}}$ and $P_{\beta_{0,2}}$ are the β-particle emission probabilities to the ground state and 293.56-keV nuclear level, respectively.

The proposed decay scheme, recommended relative emission probabilities of the gamma rays and α_{total} have been used to determine a $P_{\beta_{0,2}}^{rel}$ value of 125 (7) by the appropriate summation of the measured gamma population/depopulation of the 293.56-keV nuclear level. Therefore:

$$P_{\beta_{0,0}}^{rel} \sim 2.055 \times 125 (7) = 257 (14)$$

with an uncertainty assigned in a somewhat arbitrary manner on the basis of the uncertainty derived for $P_{\beta_{0,2}}^{rel}$.

The normalization factor (NF) for the relative emission probabilities of both the β^- particles and γ rays has been determined from the total $\beta\gamma$ transitions populating the ground state of ²¹⁵Po directly:

$$P_{\beta_{0,0}}^{rel} \times NF + \sum P_{\gamma}^{rel}(1 + \alpha_{total}) \times NF = 100$$

$$257 (14) \times NF + [164 (7) \times NF] = 100$$

$$NF = 100/421 (16) = 0.238 (9)$$

Both P_{β}^{abs} to the ground state and 293.56-keV nuclear level of ²¹⁵Po were simply calculated from their P_{β}^{rel} values and NF , and are coupled together on the basis of crude estimates of their uncertainties (i.e. arbitrary uncertainty of 20 % assigned to the value of $P_{\beta_{0,2}}^-$):

$$P_{\beta_{0,2}}^{abs} \text{ of } 30 (6) \%$$

$$\text{and } P_{\beta_{0,0}}^{abs} \text{ of } 61 (6) \%$$

These data should be treated with a high degree of caution. Their derivation also impacts significantly on the quantification of the other beta-particle emission probabilities.

Apart from the beta-particle emission directly to the ground state of ²¹⁵Po, the relative emission probabilities of all of the other beta-particle decays were calculated from population-depopulation balances of the relative gamma transition probabilities, as derived from the relative gamma-ray emission probabilities and internal conversion coefficients determined from the frozen orbital approximation of Kibédi *et al.* (2008Ki07) based on the theoretical model of Band *et al.* (2002Ba85, 2002Ra45). Direct beta population of the 271.228-keV nuclear level of ²¹⁵Po was calculated to be zero from the calculation of the known gamma transition probabilities populating and depopulating this particular excited state ((9/2⁻) to 7/2⁺ (1st forbidden non-unique)).

Beta-particle emission probabilities per 100 disintegrations of ²¹⁵Bi, transition type and log *ft*.

E_{β} (keV)	P_{β}	transition type [‡]	log <i>ft</i> [#]
	Recommended value		
790 (15)	2.8 (1) [*]	[1 st forbidden non-unique]	6.00
895 (15)	2.0 (2) [*]	[1 st forbidden non-unique]	6.34
1013 (15)	0.2 (1) [*]	[1 st forbidden non-unique]	7.5
1111 (15)	0.7 (1) [*]	[1 st forbidden non-unique]	7.1
1354 (15)	1.5 (1) [*]	[1 st forbidden non-unique]	7.10
1512 (15)	0.5 (1) [*]	[1 st forbidden non-unique]	7.8
1581 (15)	0.7 (1) [*]	(1 st forbidden non-unique)	7.7
1671 (15)	0.3 (2) [*]	(1 st forbidden non-unique)	8.1
1787 (15)	0.5 (1) [*]	(1 st forbidden unique)	9.0
1895 (15)	30 (6) ^{*†}	(1 st forbidden non-unique)	6.35
1918 (15)	–	(1 st forbidden non-unique)	–
2189 (15)	61 (6) [†]	(1 st forbidden non-unique)	6.28
	Σ 100 (8)		

^{*} Recommended absolute β^{-} emission probabilities derived from the relative gamma-ray emission probabilities, normalization factor of 0.238 (9), and theoretical internal conversion coefficients.

[†] Absolute emission probabilities calculated from fifth-power relationship of β^{-} end-point energies, with an arbitrary estimated uncertainty of 20 % assigned to the 1895-keV β^{-} emission probability.

[‡] Transition types within square brackets [] are not based on any spin-parity assignments – they have been assumed to be first forbidden non-unique as observed for the majority of the higher-energy β^{-} transitions.

[#] Log *ft* values calculated on the assumption of first forbidden non-unique transitions, apart from the 1787-keV beta emission (defined as most likely to be first forbidden unique).

The observed systematics of the two principle emissions in β^{-} decay for odd-even nuclides has been used in a quantitative manner to derive beta-particle emission probabilities in absolute terms. This approach is both approximate and of highly questionable merit – under these unsatisfactory circumstances, further experimental studies are required to determine direct β^{-} feeding to the ground state of daughter ²¹⁵Po with good accuracy.

Gamma rays

Energies

All gamma-ray transition energies were calculated from the structural details of the proposed decay scheme derived from 2001Br01 and 2003Ku26. The lower-energy nuclear level energies of 2001Br31 were adopted, along with higher-energy nuclear levels calculated from the gamma-ray studies of 2003Ku26. These data were subsequently used to re-determine the energies and associated uncertainties of the gamma-ray transitions between the various populated-depopulated levels.

Emission Probabilities

The only known experimental studies of relevance in defining the decay scheme of ²¹⁵Bi are the measurements by Ruchowska *et al.* (1990Ru02) in which the emission probabilities of seven gamma-ray transitions were quantified in terms of $P_{\gamma}(293.56 \text{ keV})$ of 1000 (redefined as 100 %), and the more extensive studies of Kurpeta *et al.* (2003Ku26) in which the emission probabilities of 19 gamma-ray transitions were quantified.

Table 3 and Fig. 6 of 2003Ku26 contain highly questionable absolute β -particle and γ -ray emission probabilities. While the resulting γ -ray transition probabilities populating the ²¹⁵Po ground state directly sum to only 57.6 %, no direct β^{-} decay is advocated to achieve a correct summation of 100 %. Private communications between Kurpeta (Institute of Experimental Physics, Warsaw University) and Kondev (ANL), April 2011, have clarified the caption of Table 3: γ intensities listed in this table are relative and not absolute (defined erroneously as %

per decay). Therefore, these γ -ray emission probabilities have been re-defined as relative to $P_{\gamma}(293.56 \text{ keV})$ of 100 %.

A number of unobserved low-intensity gamma rays have also been introduced by considering the equivalent gamma-ray studies of the α decay of ^{219}Rn – this process results in the introduction of the 130.58-, 224.04- and 405.43-keV gamma transitions, each with relative emission probabilities of less than 0.15 %.

Gamma-ray emission probabilities: as published, and relative to $P_{\gamma}(293.56 \text{ keV})$ of 100 %.

E_{γ} (keV)	P_{γ}^{rel}			Recommended value [*]
	1990Ru02	2003Ku26 [†] as published	adjusted	
130.58 (1)	–	–	–	0.039(4) [‡]
224.04 (7)	–	–	–	0.14 (2) [‡]
271.228 (10)	5.5 (5)	2.9 (1)	8.2 (3)	8.2 (3)
293.56 (4)	100 (7)	35.2 (11)	100 (3)	100 (3)
383.10 (8)	–	0.2 (1)	0.6 (3)	0.6 (3)
401.81 (1)	1.0 (4)	0.7 (1)	2.0 (3)	2.0 (3)
405.43 (7)	–	–	–	0.024 (4) [‡]
517.60 (6)	1.9 (3)	1.5 (1)	4.3 (3)	4.3 (3)
541.76 (22)	–	0.3 (1)	0.9 (3)	0.9 (3)
564.09 (22)	1.3 (3)	1.0 (1)	2.8 (3)	2.8 (3)
608.30 (7)	–	1.0 (1)	2.8 (3)	2.8 (3)
676.66 (7)	0.6 (2)	0.6 (1)	1.7 (3)	1.7 (3)
776.9 (1)	–	1.2 (2)	3.4 (6)	3.4 (6)
784 (2)	–	0.5 (1)	1.4 (3)	1.4 (3)
806.4 (20)	–	0.6 (1)	1.7 (3)	1.7 (3)
835.32 (22)	1.4 (3)	0.9 (1)	2.6 (3)	2.6 (3)
905 (2)	–	0.3 (1)	0.9 (3)	0.9 (3)
1023.3 (1)	–	0.9 (1)	2.6 (3)	2.6 (3)
1105.2 (4)	–	2.2 (1)	6.3 (3)	6.3 (3)
1127.6 (4)	–	0.7 (1)	2.0 (3)	2.0 (3)
1294.5 (1)	–	0.9 (1)	2.6 (3)	2.6 (3)
1398.8 (4)	–	1.2 (1)	3.4 (3)	3.4 (3)

[†] Published as absolute emission probabilities of doubtful overall pedigree (transition probabilities directly populating the ^{215}Po ground state only sum to 57.6 %, while direct β^- decay of zero is advocated); J. Kurpeta (Institute of Experimental Physics, Warsaw University), private communication to F.G. Kondev (ANL), 27 April 2011, concerning caption of Table 3 (2003Ku26): γ intensities are relative and not % per decay – therefore, emission probabilities have been adjusted to be relative to $P_{\gamma}(293.56 \text{ keV})$ of 100 %.

^{*} Recommended data biased completely towards the more extensive measurements of 2003Ku26.

[‡] Derived from equivalent γ -ray measurements of ^{219}Rn α decay.

Major disagreements are observed between the emission probability measurements of 1990Ru02 and 2003Ku26 that negate the merit of any form of weighted-mean analysis. Under these circumstances, the more comprehensive data of 2003Ku26 have been adopted relative to $P_{\gamma}(293.56 \text{ keV})$ of 100 %.

Multipolarities and Internal Conversion Coefficients

The decay scheme specified by 2001Br31 has been used to define the multipolarity of specific gamma transitions on the basis of the known spins and parities of the nuclear levels. Thus, the 224.04- and 401.81-keV gamma-ray emissions are adjudged to be E2 transitions. Multipolarity mixing ratios for the 130.58- and 271.228-keV gamma transitions of 0.60 (6) and 4.0 (4), respectively, were derived from the K/L and L sub-shell conversion-electron ratios determined by Davidson and Connor (1970Da09), while the 293.56- and 517.60-keV gamma-ray emissions were arbitrarily assigned mixing ratios of 1.0 (2) (i.e. 50 % M1 + 50 % E2). Recommended internal conversion coefficients have been determined from the frozen orbital approximation of Kibédi *et al.* (2008Ki07), based on the theoretical model of Band *et al.* (2002Ba85, 2002Ra45).

Gamma-ray emissions: multipolarities and theoretical internal conversion coefficients (frozen orbital approximation).

E_γ (keV)	Multipolarity	α_K	α_L	α_{M+}	α_{total}
130.58 (1)	73.5%M1 + 26.5%E2 $\delta = 0.60(6)$	3.19 (16)	0.94 (4)	0.31	4.44 (13)
224.04 (7)	(E2)	0.1296 (19)	0.1407 (20)	0.0487	0.319 (5)
271.228 (10)	6%M1 + 94%E2 $\delta = 4.0(4)$	0.111 (6)	0.0668 (11)	0.0232	0.201 (7)
293.56 (4)	(50%M1 + 50%E2) $\delta = 1.0(2)$	0.25 (4)	0.062 (4)	0.028	0.34 (5)
383.10 (8)	–	–	–	–	–
401.81 (1)	E2	0.0351 (5)	0.01528 (22)	0.00512	0.0555 (8)
405.43 (7)	–	–	–	–	–
517.60 (6)	50%M1 + 50%E2 $\delta = 1.0(2)$	0.058 (9)	0.0115 (11)	0.0035	0.073 (10)
541.76 (22)	–	–	–	–	–
564.09 (22)	–	–	–	–	–
608.30 (7)	(M1 + E2)	–	–	–	–
676.66 (7)	–	–	–	–	–
776.9 (1)	–	–	–	–	–
784 (2)	–	–	–	–	–
806.4 (20)	–	–	–	–	–
835.32 (22)	–	–	–	–	–
905 (2)	–	–	–	–	–
1023.3 (1)	–	–	–	–	–
1105.2 (4)	–	–	–	–	–
1127.6 (4)	–	–	–	–	–
1294.5 (1)	–	–	–	–	–
1398.8 (4)	–	–	–	–	–

While a decay scheme has been formulated from the gamma-ray emission probability measurements of Kurpeta *et al.* (2003Ku26), further studies are required to determine the absolute and relative gamma-ray emission probabilities and also quantify any direct β^- feeding to the ground state of daughter ²¹⁵Po with much greater confidence. Such work would assist greatly to remove the severe doubts associated with the proposed decay scheme.

Atomic Data

The x-ray data have been calculated using the evaluated gamma-ray data, and the atomic data from 1996Sc06, 1998ScZM and 1999ScZX. Both the x-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.01, 28 January 2003). This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

K and L X-ray emission probabilities per 100 disintegrations of ²¹⁵Bi.

			Energy (keV)	Photons per 100 disint.
XL	(Po)		9.658 – 16.213	2.7 (3)
	XL ₁	(Po)	9.658	0.065 (8)
	XL _α	(Po)	11.016 – 11.130	1.20 (13)
	XL _η	(Po)	12.085	0.022 (3)
	XL _β	(Po)	12.823 – 13.778	1.18 (11)
	XL _γ	(Po)	15.742 – 16.213	0.24 (2)
XK _α	XK _{α2}	(Po)	76.864 (4)	1.8 (3)
	XK _{α1}	(Po)	79.293 (5)	3.0 (5)
XK' _{β1}	XK _{β3}	(Po)	89.256)
	XK _{β1}	(Po)	89.807) 1.02 (16)
	XK _{β5}	(Po)	90.363)
XK' _{β2}	XK _{β2}	(Po)	92.263)
	XK _{β4}	(Po)	92.618) 0.32 (5)
	XKO _{2,3}	(Po)	92.983)

Electron energies were determined from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

Data Consistency

A Q_β-value of 2189 (15) keV has been adopted from the atomic mass evaluation of Audi *et al.* (2003Au03) while in the course of formulating the decay scheme of ²¹⁵Bi. This value has subsequently been compared with the Q-value calculated by summing the contributions of the individual emissions to the ²¹⁵Bi beta-decay process (i.e. β⁻, conversion electrons, γ, etc.):

$$\text{calculated Q-value} = \sum (E_i \times P_i) = 2190 (170) \text{ keV}$$

Percentage deviation from the Q-value of Audi *et al.* is (0 ± 8) %, which supports the derivation of a highly consistent decay scheme with a large variant.

References

- 1933Sa01 B.W. SARGENT, The Maximum Energy of the β-rays from Uranium X and Other Bodies, Proc. Royal Soc. (London) A139 (1933) 659-673. [β⁻ decay, 5th-power law]
- 1953Hy83 E.K. HYDE, A. GHIORSO, The Alpha-branching of AcK and the Presence of Astatine in Nature, Phys. Rev. 90 (1953) 267-270. [β⁻ decay, half-life]
- 1955Ev23 R.D. EVANS, The Atomic Nucleus, Tata McGraw-Hill Publishing Company Ltd., Bombay and New Delhi, India (1955) 559. [β⁻ decay, 5th-power law]

- 1963KaZZ I. KAPLAN, Nuclear Physics, 2nd Edition, Addison-Wesley Publishing Company Inc., Reading, Massachusetts, USA (1963) 364-365. [β^- decay, 5th-power law]
- 1965Nu03 M. NURMIA, D. GIESSING, W. SIEVERS, L. VARGA, Studies of the Natural Actinium Radioactive Series, Ann. Acad. Sci. Fenn. Ser. A VI, No.167 (1965).
[Half-life]
- 1970Da09 – W.F. DAVIDSON, R.D. CONNOR, The Decay of ^{223}Ra and its Daughter Products (II). The Decay of ^{219}Rn and ^{215}Po , Nucl. Phys. A149 (1970) 385-391.
[K/L and L sub-shell ratios, ICC]
- 1977La19 F.P. LARKINS, Semiempirical Auger-electron Energies for Elements $10 \leq Z \leq 100$, At. Data Nucl. Data Tables 20 (1977) 311-387. [Auger-electron energies]
- 1989Bu09 D.G. BURKE, H. FOLGER, H. GABELMANN, E. HAGEBØ, P. HILL, P. HOFF, O. JONSSON, N. KAFFRELL, W. KURCEWICZ, G. LØVHØIDEN, K. NYBØ, G. NYMAN, H. RAVN, K. RIISAGER, J. ROGOWSKI, K. STEFFENSEN, T.F. THORSTEINSEN, and the ISOLDE Collaboration, New Neutron-rich Isotopes of Astatine and Bismuth, Z. Phys. – Atomic Nuclei 333 (1989) 131-135. [Half-life]
- 1990Ru02 E. RUCHOWSKA, J. ZYLICZ, C.F. LIANG, P. PARIS, CH. BRIANÇON, The β^- decay of ^{215}Bi and a Firm Identification of ^{216}Bi , J. Phys. G: Nucl. Part. Phys. 16 (1990) 255-260. [E_γ , P_γ , half-life]
- 1991Ma16 M.J. MARTIN, Nuclear Data Sheets for A = 209, Nucl. Data Sheets 63 (1991) 723-844. [Nuclear structure, level energies]
- 1996Sc06 E. SCHÖNFELD, H. JANßEN, Evaluation of Atomic Shell Data, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527-533. [X_K , X_L , Auger electrons]
- 1998ScZM E. SCHÖNFELD, G. RODLOFF, Tables of the Energies of K-Auger Electrons for Elements with Atomic Numbers in the Range from Z = 11 to Z = 100, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX E. SCHÖNFELD, G. RODLOFF, Energies and Relative Emission Probabilities K X-rays for Elements with Atomic Numbers in the Range from Z = 5 to Z = 100, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2001Br31 E. BROWNE, Nuclear Data Sheets for A = 215, 219, 223, 227, 231, Nucl. Data Sheets 93 (2001) 763-1061. [Nuclear structure, level energies]
- 2002Ba85 I.M. BAND, M.B. TRZHASKOVSKAYA, C.W. NESTOR Jr., P.O. TIKKANEN, S. RAMAN, Dirac–Fock Internal Conversion Coefficients, At. Data Nucl. Data Tables 81 (2002) 1-334. [ICC]
- 2002Ra45 S. RAMAN, C.W. NESTOR Jr., A. ICHIHARA, M.B. TRZHASKOVSKAYA, How Good Are the Internal Conversion Coefficients Now? Phys. Rev. C66 (2002) 044312, 1-23. [ICC]
- 2003Ak06 Y.A. AKOVALI, Nuclear Data Sheets for A = 217, Nucl. Data Sheets 100 (2003) 141-178. [Nuclear structure, level energies]

- 2003Au03 G. AUDI, A.H. WAPSTRA, C. THIBAULT, The AME2003 Atomic Mass Evaluation (II). Tables, Graphs and References, Nucl. Phys. A729 (2003) 337-676. [Q-value]
- 2003Ku26 J. KURPETA, A. PŁOCHOCKI, A.N. ANDREYEV, J. ÄYSTÖ, A. DE SMET, H. DE WITTE, A.-H. EVENSEN, V. FEDOSEYEV, S. FRANCHOO, M. GÓRSKA, H. GRAWE, M. HUHTA, M. HUYSE, Z. JANAS, A. JOKINEN, M. KARNY, E. KUGLER, W. KURCEWICZ, U. KÖSTER, J. LETTRY, A. NIEMINEN, K. PARTES, M. RAMDHANE, H.L. RAVN, K. RYKACZEWSKI, J. SZERYPO, K. VAN DE VEL, P. VAN DUPPEN, L. WEISSMAN, G. WALTER, A. WÖHR, IS387 Collaboration and ISOLDE Collaboration, Isomeric and Ground-state Decay of ²¹⁵Bi, Eur. Phys. J. A18 (2003) 31-37. [E_γ, P_γ, P_β]
- 2004Br45 E. BROWNE, Nuclear Data Sheets for A = 211, Nucl. Data Sheets 103 (2004) 183-268. [Nuclear structure, energies]
- 2007Ba19 M.S. BASUNIA, Nuclear Data Sheets for A = 213, Nucl. Data Sheets 108 (2007) 633-680. [Nuclear structure, energies]
- 2008Ki07 T. KIBÉDI, T.W. BURROWS, M.B. TRZHASKOVSKAYA, P.M. DAVIDSON, C.W. NESTOR Jr., Evaluation of Theoretical Conversion Coefficients Using BrIcc, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202-229. [ICC]

**²¹⁵Po -Comments on evaluation of decay data
by V.P. Chechev**

This evaluation was done in November 2010 with a literature cut-off by the same date.

1. DECAY SCHEME

²¹⁵Po decays 100 % to levels of ²¹¹Pb by emission of α particles and $2.3 (2) \times 10^{-4}$ % to ²¹⁵At by emission of β^- particles. The structure of the adopted scheme of ²¹⁵Po decay is based on the experiment of 1998Li53 and the evaluation by E. Browne (2004Br45). The existence of the alpha-particle group with energy of 6950 keV, reported in 1962Wa18, 1971Gr17, was not confirmed in 1998Li53 and the relevant ²¹¹Pb level of 447 keV was omitted in this evaluation. Similarly, the questionable ²¹¹Pb level of 762 keV, determined by the alpha-particle group with energy of 6636 keV and intensity of $\sim 3 \times 10^{-4}$ %, has not been adopted.

The decay scheme of ²¹⁵Po is not completed as only approximate information is available for weak gamma transitions following α decay, their gamma-ray emission probabilities and multiplicities have not been determined, and, in fact, the ²¹¹Pb levels were deduced only from measurements of alpha-particle groups. In respect of ²¹⁵Po β^- decay, the β^- - spectrum has not been measured and a fine structure of β^- decay is unknown.

The current evaluated data are supported by the agreement between $Q(\text{calculated}) = 7526.2 (22)$ keV, deduced from the calculated average energies of all emissions, and $Q(\alpha) = 7526.3 (8)$ keV, adopted from 2003Au03. Percentage deviation of $Q(\text{calculated})$ from the $Q(\alpha)$ of Audi *et al.* (2003Au03) is (0.0 ± 0.3) %.

2. NUCLEAR DATA

$Q(\alpha)$ and $Q(\beta^-)$ values are from Audi *et al.* (2003Au03).

The ²¹⁵Po half-life is based on the experimental results given in Table 1.

Table 1. Experimental values of ²¹⁵Po half-life

Reference	Author(s)	Half-life (ms)	Method
1942Wa04	Ward	1.83 (4)	Observations with a single Geiger counter
1961Vo06	Volkov <i>et al.</i>	1.778 (5)	Measurements with ionization alpha-spectrometer equipped by time analyzer
1971Er02	Erlík <i>et al.</i>	1.785 (10)	Time interval analyzer method
1971Er02	Erlík <i>et al.</i>	1.784 (8)	Multichannel delay coincidence method

The set of the four experimental values is consistent. The weighted average for this data set is 1.781 with the internal uncertainty of 0.0039 and an external uncertainty of 0.0033 ($\chi^2/\nu = 0.72$).

The recommended value of the ^{215}Po half-life is **1.781 (4) ms**.

β^- branching of $2.3 (2) \times 10^{-4} \%$ was adopted from the measurement of 1950Av61. With this value the α branching is obtained to be 99.999 77 (2) %.

2.1. Alpha Transitions

The alpha transition energies have been obtained from the $Q(\alpha)$ value and ^{211}Pb level energies given in Table 2 from 2004Br45. The uncertainties in the energies of levels 2 - 7 have been adopted ± 3 keV taking into account the average discrepancy of experimental and calculated alpha-particle energies (Table 3) and as provided by uncertainties of gamma ray energies from 1998Li53 ≥ 1.0 keV for all γ rays, except for $\gamma_{438.9}$ keV.

Table 2. ^{211}Pb levels populated in ^{215}Po α -decay

Level	Energy (keV)	Spin and parity	Half-life	Probability of α - transition (%)
0	0.0	9/2+	36.1(4) min	99.934 (20)
1	438.9 (2)	(7/2)+		0.06 (2)
2	584 (3)			$4 (2) \times 10^{-4}$
3	598 (3)	(5/2+)		$1.6 (5) \times 10^{-3}$
4	643 (3)	11/2+		$8 (3) \times 10^{-4}$
5	733 (3)	(13/2+)		$8 (3) \times 10^{-4}$
6	815 (3)	(9/2+)		$2.0 (6) \times 10^{-3}$
7	894 (3)	(11/2+)		3×10^{-4}

The alpha transitions in ^{215}Po decay were observed in a number of works by study of an ^{223}Ra alpha emitting source (1962Wa18, 1965Va10, 1970Da09, 1998Li53). In 1962Wa18 the ^{215}Po alpha spectrum was measured with magnetic spectrometer. In 1965Va10 the coincidence of $\gamma_{1,0}$ (438.9 keV)-gamma ray with $\alpha_{0,1}$ (6.95 MeV) was observed. In 1970Da09 the alpha transition probability ($P(\alpha)$) was measured for $\alpha_{0,1}$ (6.95 MeV)-transition. Most accurate and detailed data were obtained by 1998Li53 with use of α - γ coincidences. These measurement results have been adopted for the recommended $P(\alpha)$ and compared in Table 3 with other available poor experimental data.

Table 3. Experimental ^{215}Po alpha transition probability values ($P(\alpha)$)

α -particle energy (keV)	1962Wa18	1970Da09	1998Li53
7386	100		99.93
6955	≈ 0.056	≈ 0.1	0.06 (2)
6813			$4 (2) \times 10^{-4}$
6799			$1.6 (5) \times 10^{-3}$
6755			$8 (3) \times 10^{-4}$
6667			$8 (3) \times 10^{-4}$
6586			$2.0 (6) \times 10^{-3}$
6509			$\sim 3 \times 10^{-4}$

The accurate $P(\alpha_{0,0})$ value has been deduced from $\Sigma P(\alpha_{0,i}) = 99.999\ 77\ (2)\ \%$, ($i = 0, 1, \dots, 7$) and, the individual adopted $P(\alpha_{0,i})$, ($i = 1 - 7$).

The α decay hindrance factors were calculated using the ALPHAD computer program from the ENSDF evaluation package with $r_0 (^{211}\text{Pb}) = 1.5393\ \text{fm}$ (2004Br45).

2.2. Gamma Transitions and Internal Conversion Coefficients

Information on the gamma-ray transition probabilities and the gamma-ray multiplicities is not available, except for $\gamma_{438.9\ \text{keV}}$ (1968Br17, 1970Da09, 1998Li53, see §6.2.2). The gamma-ray transition probability $P_{\gamma+ce}(\gamma_{1,0} - 438.9\ \text{keV})$ was then deduced from the probability balance: $P(\alpha_{0,1}) = P_{\gamma+ce}(\gamma_{1,0} - 438.9\ \text{keV})$. The multipolarity of this gamma-ray transition has been adopted as being E2. In 1998Li53 a multipolarity higher than a pure E2 was reported from the relative intensity $P(KX) / P_{\gamma}(438.9\ \text{keV}) = 0.034\ (10)$, then it was noted that a small amount of M1 cannot be ruled out.

ICCs have been interpolated using the BrIcc computer program, version v2.2a, data set BriccFO (2008Ki07).

3. ATOMIC DATA

The fluorescence yields, X-ray energies and relative probabilities, and Auger electrons energies and relative probabilities are from the SAISINUC software.

4. ALPHA EMISSIONS

The energy of the alpha-particle group $\alpha_{0,0}$ that populates the ^{211}Pb ground state is the absolute measurement result from 1971Gr17 adjusted in 1991Ry01 for change in calibration standards: $E(\alpha_{0,0}) = 7386.1\ (8)\ \text{keV}$. Latter coincides with the value deduced by the evaluator from the adopted $Q(\alpha)$ taking into account the recoil energy for ^{211}Pb .

The energy of alpha-particle group $\alpha_{0,1}$ of 6955.4 (8) keV has been deduced from the $Q(\alpha)$ value taking into account the level energy of 439.8 (2) keV and the recoil energy for ^{211}Pb . The above value of $E(\alpha_{0,1})$ can be compared to the measured $E(\alpha_{0,1})$ of 6956.7 keV (without uncertainty) by 1962Wa18, 1971Gr17 and of 6954 (3) keV by 1998Li53 with adjustment adopted in 2004Br45.

The energies of remaining alpha-particle groups have been deduced from $Q(\alpha)$ and the relevant ^{211}Pb level energies. In Table 4 the deduced (recommended) $E(\alpha)$ are compared with the experimental values from the measurements of 1998Li53 adjusted in 2004Br45 to the adopted $E(\alpha_{0,0}) = 7386.1\ (8)\ \text{keV}$.

Table 4. Experimental and deduced (recommended) ^{215}Po alpha-particle energies ($E(\alpha)$)

Level	Level energy (keV)	α -transition energy	Experimental $E(\alpha)$ (1998Li53) ^a	Deduced $E(\alpha)$ (recommended)
0	0.0	7526.3 (8)	7386.1 (8)	7386.1 (8)
1	438.9 (2)	7087.4 (10)	6954 (3)	6955.4 (8)
2	584 (3)	6942 (3)	6819 (15)	6813 (3)
3	598 (3)	6928 (3)	6803 (8)	6799 (3)
4	643 (3)	6883 (3)	6754 (10)	6755 (3)

Level	Level energy (keV)	α -transition energy	Experimental E(α) (1998Li53) ^a	Deduced E(α) (recommended)
5	733 (3)	6793 (3)	6671 (10)	6667 (3)
6	815 (3)	6711 (3)	6589 (8)	6586 (3)
7	894 (3)	6632 (3)	6519 (20)	6509 (3)

^a E(α) have been adjusted to the adopted E($\alpha_{0,0}$) = 7386.1 (8) keV.

5. ELECTRON EMISSIONS

The energies of the conversion electrons for the $\gamma_{438.9}$ keV transition have been obtained from the gamma-ray transition energy and the atomic electron binding energies.

The emission probabilities of the conversion electrons have been deduced using the P_γ and ICC values.

The absolute emission probabilities of K and L Auger electrons have been calculated using the EMISSION computer program.

6. PHOTON EMISSIONS

6.1 X - Ray emissions

The absolute emission probabilities of Pb KX- and LX-rays were calculated using the EMISSION computer program. The total emission probability of Pb KX-rays in decay of ^{215}Po was determined relatively to $P_\gamma(\gamma_{1,0} - 438.9 \text{ keV})$ (1998Li53). The experimental $P(\text{KX})/P_\gamma(\gamma_{1,0} - 438.9 \text{ keV}) = 0.034$ (10) agrees with the value of 0.029 (14) calculated with the EMISSION code.

The agreement between measured and calculated KX-ray emission probabilities supports the recommended γ -ray emission probability and assigned multipolarity for $\gamma_{1,0} - 438.9 \text{ keV}$.

6.2. Gamma emissions

6.2.1. Gamma ray energies

The gamma-ray energies (E_γ) have been taken from the measurements of 1998Li53. The uncertainties on the gamma-ray energies higher than 500 keV have been assumed being $\pm 3 \text{ keV}$ (see section 2.1). Other measurements of E ($\gamma_{1,0} - 438.9 \text{ keV}$) are reported in 1968Br17 (438.7 (3) keV) and in 1970Da09 (438.9 keV – without uncertainty).

6.2.2. Gamma ray emission probabilities

There is no available information on the gamma-ray emission probabilities, except for $P(\gamma_{438.9 \text{ keV}})$: 0.048 (5) % (1968Br17) and 0.064 (2) % (1970Da09). These discrepant values do not conflict with the recommended value of $P(\gamma_{438.9 \text{ keV}}) = 0.058$ (19) % deduced by the evaluator from the alpha transition probability $P(\alpha_{0,1}) = 0.06$ (2) % and total internal conversion coefficient $\alpha_T = 0.0405$ (6) under the assumption of E2 multipolarity.

7. REFERENCES

- 1942Wa04** A.G. Ward, Proc. Roy. Soc. (London) 181A, 183 (1942) (Half-life)
- 1950Av61** P. Avignon, J. Phys. Radium 11, 521 (1950) (β^- branching)
- 1961Vo06** Yu. M. Volkov, A.P. Komar, G.A. Korolev, G.E. Kocharov, Izvest. Akad. Nauk SSSR, Ser. Fiz. 25, 1188 (1961); Columbia Tech.Transl. 25, 1193 (1962) (Half-life)
- 1962Wa18** R.J. Walen, V. Nedovesov, G. Bastin-Scoffier, Nuclear Phys. 35, 232 (1962) (α -particle energies and emission probabilities)
- 1965Va10** K. Valli, J. Aaltonen, G. Graeffe, M. Nurmia, Ann. Acad. Sci. Fenn., Ser. A VI, No. 184 (1965) (α -particle energies and emission probabilities)
- 1968Br17** C. Briançon, C.F. Leang, R. Walen, Compt. Rend. 266B, 1533 (1968) (γ -ray energies and emission probabilities)
- 1970Da09** W.F. Davidson, R.D. Connor, Nucl. Phys. A149, 385 (1970) (γ -ray energies and emission probabilities)
- 1971Er02** A. Erlik, J. Felsteiner, H. Lindeman, M. Tatcher, Nucl. Instrum. Methods 92, 45 (1971) (Half-life)
- 1971Gr17** B. Grennberg, A. Rytz, Metrologia 7, 65 (1971) (α -particle energies)
- 1991Ry01** A. Rytz, At. Data Nucl. Data Tables 47, 205 (1991) (α -particle energies and emission probabilities)
- 1998Li53** C.F. Liang, P. Paris, R.K. Sheline, Phys. Rev. C58, 3223 (1998) (α -particle and γ -ray energies and emission probabilities)
- 2003Au03** G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A729, 337 (2003) (Q values)
- 2004Br45** E. Browne, Nucl. Data Sheets 103, 183 (2004) (^{215}Po α decay scheme, ^{211}Pb levels)
- 2008Ki07** T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr, Nucl. Instrum. Methods Phys. Res. A589, 202 (2008) (Band-Raman ICC for γ -ray transitions)

**²¹⁵At -Comments on evaluation of decay data
by V.P. Chechev**

This evaluation was done in December 2010 with a literature cut-off by the same date.

1. DECAY SCHEME

²¹⁵At decays 100 % to levels of ²¹¹Bi by emission of α particles. The adopted ²¹¹Bi levels populated in the ²¹⁵At decay are based on the experiment of 1966Gr07 and the evaluation by Browne (2004Br45).

The decay scheme of ²¹⁵At seems to be incomplete as the alpha decays to higher levels in daughter ²¹¹Bi, which are known from the β^- decay of ²¹¹Pb (see ²¹¹Bi Adopted Levels, Gammas of 2004Br45), are not observed yet.

The current evaluated data are supported by the agreement between $Q(\text{calculated}) = 8178 (5) \text{ keV}$, deduced from the calculated average energies of all emissions, and $Q(\alpha) = 8178 (4) \text{ keV}$, adopted from 2003Au03.

2. NUCLEAR DATA

$Q(\alpha)$ is from 2003Au03 where this value has been deduced from the measurement of α -particle energy $E(\alpha_{0,0}) = 8026 (4) \text{ keV}$ by 1982Bo04 recommended in 1991Ry01.

The ²¹⁵At half-life of 0.10 (2) ms is from the single measurement of 1951Me10.

2.1. Alpha Transitions

The alpha transition energies have been obtained from the $Q(\alpha)$ value and ²¹¹Bi level energies given in Table 1 from ²¹¹Bi Adopted Levels, Gammas of 2004Br45.

Table 1. ²¹¹Bi levels populated in ²¹⁵At α -decay

Level	Energy (keV)	Spin and parity	Half-life	Probability of α -transition (%)
0	0.0	9/2-	2.14 (2) min	99.95 (2)
1	404.854 (9)	7/2-	0.317 (11) ns	0.05 (2)

The alpha transition probability $P(\alpha_{0,1})$ is from the measurement of 1966Gr07 by means of α - γ coincidence technique with surface-barrier semi-conductor and NaI(Tl) detectors. The accurate $P(\alpha_{0,0})$ value has been deduced from the expression of $P(\alpha_{0,0}) + P(\alpha_{0,1}) = 100 \%$.

The α decay hindrance factors have been calculated using the ALPHAD computer program from the ENSDF evaluation package with $r_0(^{211}\text{Pb}) = 1.5443 \text{ fm}$ (2004Br45).

2.2. Gamma Transitions and Internal Conversion Coefficients

The 405-keV gamma-ray transition probability has been deduced from the intensity balance at the 405-keV level using the adopted alpha transition probability $P(\alpha_{0,1})$ and total internal conversion coefficient (ICC) α_T for $\gamma_{1,0}$ (405 keV). The multipolarity (M1+E2) and E2/M1 mixing ratio (δ) of -1.1 (1) have been taken from 2004Br45. These are based on the measurements of conversion electrons in ²¹¹Pb β^- decay and $\gamma(\theta)$ measurements with polarized ²¹¹Bi nuclei. ICCs $\alpha_T, \alpha_K, \alpha_L, \alpha_M$ have been interpolated using the BrIcc computer program, version v2.2a, data set BrIccFO (2008Ki07).

3. ATOMIC DATA

The fluorescence yields, X-ray energies and relative probabilities, and Auger electrons energies and relative probabilities are from the SAISINUC software.

4. ALPHA EMISSIONS

The energy of alpha-particle group $\alpha_{0,0}$ that populates the ²¹¹Bi ground state is the measured value from 1982Bo04 recommended in 1991Ry01. In 1966Gr07 the measured value of 8.00 (1) MeV was reported.

The energy of alpha-particle group $\alpha_{0,1}$ of 7628 (4) keV has been deduced from the $Q(\alpha)$ value taking into account the level energy of 404.854 (9) keV and the recoil energy for ²¹¹Bi. The above value of $E(\alpha_{0,1})$ can be compared to the value of 7626 (15) keV as measured by 1966Gr07 and adjusted by the evaluator to the adopted $E(\alpha_{0,0}) = 8026$ (4) keV (the original value of 1966Gr07 is 7.60 (1) MeV).

The earlier measured energy of α -emission in the decay of ²¹⁵At is 8.00 (2) MeV (1951Me10).

5. ELECTRON EMISSIONS

The energies of the conversion electrons for $\gamma_{1,0}$ (405 keV) have been obtained from the gamma-ray transition energy and the atomic electron binding energies.

The emission probabilities of the conversion electrons have been deduced using the P_γ and ICC values.

The absolute emission probabilities of K and L Auger electrons have been calculated using the EMISSION computer program.

6. PHOTON EMISSIONS

6.1 X - Ray emissions

The absolute emission probabilities of Pb KX- and LX-rays were calculated using the EMISSION computer program.

6.2. Gamma emissions

6.2.1. Gamma ray energies

The 405-keV gamma-ray energy has been adopted from the 405-keV level energy. In 1966Gr07 this energy was obtained from the ²¹⁵At α decay as ≈ 404 keV.

6.2.2. Gamma ray emission probabilities

The 405-keV gamma-ray emission probability has been deduced from the alpha transition probability $P(\alpha_{0,1}) = 0.05$ (2) % and total internal conversion coefficient $\alpha_T = 0.122$ (8).

7. REFERENCES

- 1951Me10 W.W. Meinke, A. Ghiorso, G.T. Seaborg, Phys. Rev. 81, 782 (1951)
(Half-life, energy of α -emission)
- 1966Gr07 G. Graeffe, P. Kauranen, J. Inorg. Nucl. Chem. 28, 933 (1966)
(α -particle energies and emission probabilities, ²¹¹Bi levels)
- 1982Bo04 J.D. Bowman, R.E. Eppley, E.K. Hyde, Phys. Rev. C25, 941 (1982)
(α -particle energies)
- 1991Ry01 A. Rytz, At. Data Nucl. Data Tables 47, 205 (1991)
(α -particle energies and emission probabilities)
- 2003Au03 G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A729, 337 (2003)
(Q value)
- 2004Br45 E. Browne, Nucl. Data Sheets 103, 183 (2004)
(²¹⁵At α decay scheme, ²¹¹Bi levels)
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr, Nucl. Instrum. Methods Phys. Res. A589, 202 (2008)
(Band-Raman ICC for γ -ray transitions)

**²¹⁶Po – Comments on evaluation of decay data
by A. L. Nichols**

Evaluated: July/August 2001

Re-evaluated: January 2004

Evaluation Procedures

Limitation of Relative Statistical Weight Method (LWM) was applied to average numbers throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

A simple decay scheme was derived from the gamma-ray studies of 1977Ku15, with an absolute emission probability of 0.0019(3)% for the single 804.9 keV gamma ray. This value and theoretical internal conversion coefficients were used to calculate the alpha -particle emission probabilities. Alpha -particle studies are required to confirm the validity of the proposed decay scheme.

Nuclear Data

The ²²⁸Th decay chain is important in quantifying the environmental impact of the decay of naturally-occurring ²³²Th.

Half-life

The recommended half-life is the weighted mean of three somewhat elderly measurements (1911Mo01, 1942Wa04 and 1963Di05). Further studies are merited to determine this value with greater confidence.

Reference	Half-life (s)
1911Mo01	0.145(15)
1942Wa04	0.158(8)
1963Di05	0.145(2)*
Recommended Value	0.150(5)

*Uncertainty adjusted to ± 0.007 to reduce weighting below 0.5.

Gamma Ray

Energy

The single gamma-ray energy was based on the nuclear level energy of 804.9(5) keV from 1992Ar05.

Emission Probability

The absolute emission probability of the 804.9(5) keV gamma ray was determined from the measurement of 1977Ku15, adjusted for the change from 3.95% (0.0395) to 4.12% (0.0412) of $P_{\gamma}(240.986 \text{ keV})$ for ²²⁴Ra.

Published Gamma-ray Emission Probabilities per 100 Disintegrations of ²¹⁶Po

E _g (keV)	P _g
	1977Ku15 [†]
804.9(5)	0.0018(3)

[†] Absolute value in measurements that include P_γ(240.986 keV) of 3.95% for ²²⁴Ra.

Absolute Gamma-ray Emission Probabilities per 100 Disintegrations of ²¹⁶Po

E _g (keV)	P _g ^{abs}	
	1977Ku15 [†]	Recommended Value
804.9(5)	0.0019(3)	0.0019(3)

[†] Adjusted with respect to evaluated P_γ(240.986 keV) of 4.12(3)% (0.0412) for ²²⁴Ra.

Multipolarity and Internal Conversion Coefficients

The decay scheme specified by 1992Ar05 has been used to define the multipolarity of the gamma transition on the basis of the known spins and parities of the two nuclear levels. Theoretical internal conversion coefficients were interpolated from the tabulations of 1978Ro22.

Alpha-particle EmissionsEnergies

Alpha-particle energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies of 1992Ar05 and the Q-value (1995Au04) were used to determine the energies and uncertainties of the alpha-particle transitions to the various levels, while allowing for the significant recoil components.

Emission Probabilities

Both alpha-particle emission probabilities were derived from the weighted mean emission probability of the single gamma transition and theoretical internal conversion coefficients.

Alpha-particle Emission Probabilities per 100 Disintegrations of ²¹⁶Po

E _a (keV)	P _a	
	1962Wa28	Recommended Values [*]
5988.6(10)	0.0021(4)	0.0019(3)
6778.6(5)	~ 100	99.9981(3)

^{*} Recommended emission probabilities derived from evaluated gamma-ray emission probability and theoretical internal conversion coefficients.

Atomic Data

The x-ray data have been calculated using the evaluated gamma-ray data, and the atomic data from 1996Sc06, 1998ScZM and 1999ScZX.

References

1911Mo01 - H. G. J. Moseley and K. Fajans, LIX. Radio -Active Products of Short Life, Phil. Mag. 22(1911)629. [Half-life]

1942Wa04 - A. G. Ward, A New Method of Determining Half -Value Periods from Observations with a Single Geiger Counter, Proc. Roy. Soc. (London) 181A(1942)183. [Half-life]

1962Wa28 - R. J. Walen, Spectrographie α du Radium 224 et de ses Dérivés, C. R. Acad. Sci. Paris 255(1962)1604. [P $_{\alpha}$]

1963Di05 - H. Diamond and J. E. Gindler, Alpha Half -Lives of ²¹⁶Po, ²¹⁷At and ²¹⁸Rn, J. Inorg. Nucl. Chem. 25(1963)143. [Half-life]

1977Ku15 - W. Kurcewicz, N. Kaffrell, N. Trautmann, A. Plochocki, J. Zylicz, M. Matul and K. Stryczniewicz, Collective States Fed by Weak α -transitions in the ²³²U Chain, Nucl. Phys. A289(1977)1. [P $_{\gamma}$]

1978Ro22 - F. Rösel, H. M. Fries, K. Alder and H. C. Pauli, Internal Conversion Coefficients for all Atomic Shells, ICC Values for Z = 68-104, At. Data Nucl. Data Tables 21(1978)291-514. [ICC]

1992Ar05 - A. Artna-Cohen, Nuclear Data Sheets for A = 212, Nucl. Data Sheets 66(1992)171. [Nuclear structure, energies]

1995Au04 - G. Audi and A. H. Wapstra, The 1995 Update to the Atomic Mass Evaluation, Nucl. Phys. A595(1995)409. [Q value]

1996Sc06 - E. Schönfeld and H. Janßen, Evaluation of Atomic Shell Data, Nucl. Instrum. Meth. Phys. Res. A369(1996)527. [X_K, X_L, Auger electrons]

1998ScZM - E. Schönfeld and G. Rodloff, Tables of the Energies of K -Auger Electrons for Elements with Atomic Numbers in the Range from Z = 11 to Z = 100, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]

1999ScZX - E. Schönfeld and G. Rodloff, Energies and Relative Emission Probabilities of K X-rays for Elements with Atomic Numbers in the Range from Z = 5 to Z = 100, PTB Report PTB-6.11-1999-1, February 1999. [X_K]

²¹⁷At - Comments on evaluation of decay data

Huang Xiaolong, Wang Baosong

This evaluation was completed in 2007. Literature available by December 2007 was included.

1 Decay Scheme

²¹⁷At disintegrates 99.9933 (24) % by α emission to levels in ²¹³Bi and 0.0067 (24) % by β^- emission to levels in ²¹⁷Rn. ²¹⁷At ground state has $J^\pi = 9/2^-$ (2007Ba19).

The α decay scheme of ²¹⁷At was built based on the measurement of 1997Ch19. The β^- decay scheme of ²¹⁷At has not been studied.

The recommended $Q(\alpha)$ value of 7201.3 (12) keV in Audi (2003Au03) agrees with the $Q(\alpha)$ value of 7197 (5) keV, calculated by the evaluator (using program RADLST) from average radiation energies. This agreement supports the completeness and correctness of the decay scheme.

2 Nuclear Data

The Q values are from the 2003Au03 evaluation.

Level energies have been obtained from a least-squares fit to γ -ray energies (GTOL computer code). Spin and parities are from 2007Ba19.

The measured and evaluated ²¹⁷At half-life values are listed in Table 1.

Table 1: Measured half-life values of ²¹⁷At and evaluated value.

$T_{1/2}$ (ms)	References	Measurement method
21	1947En03	
18 (2)	1950Ha52	Alpha pulse analyzer
32.3 (4)	1963Di05	
32.3 (4)	2007Ba19	NDS, from 1963Di05
32.3 (4)	Evaluated value	from 1963Di05

The adopted value is taken from the measurement of 1963Di05.

2.1 γ Transitions

The γ transition probabilities were calculated using the γ -ray emission intensities and the relevant internal conversion coefficients.

Multipolarities and mixing ratios of γ transitions are from 1997Ch19.

The internal conversion coefficients (ICC) and the associated uncertainties for the γ - transitions have been obtained using the BrIcc computer program (2008Ki07).

2.2 α Transitions

The measured and evaluated energies of alpha particles were listed in table 2. The evaluated values are from 1997Ch19, except as noted.

Table 2: Measured and evaluated value of α -particle energy for ²¹⁷At.

1967Dz02	1977Vy02	1982Bo04	1997Ch19	Evaluation
	6037 (3) ^b			6037 (3) ^c
			6322.0 (16)	6322.0 (16)
6422 (7) ^a				
6486 (7)			6484.7 (16)	6484.7 (16)
6541 (7) ^a				
6619 (7) ^a				
6772 (7) ^a				
6810 (7)			6813.8 (16)	6813.8 (16)
6849 (7) ^a				
7070 (8)	7062 (5)	7071 (2)	7066.9 (16)	7066.9 (16)

^a: the α transitions reported in 1967Dz02 only, were not confirmed in 1997Ch19 and 1997Ch53. So these alpha transitions have not been taken into account by the evaluators.

^b: 1977Vy02 assign this α transition to the ²²¹Rn decay;
1997Ch53 assign this α transition to the ²¹⁷At decay.

^c: from 1977Vy02.

The measured and evaluated alpha particle emission probabilities were listed in table 3. The evaluated alpha particle emission probabilities were deduced from the transition probability balance. These calculated and adopted values are in good agreement, for the main intensities, with the measured emission probabilities.

Table 3: Measured and adopted α -particle emission probabilities for ²¹⁷At.

E_{α} (keV)	P_{α} (%)				Adopted values
	1967Dz02	1969LeZW	1997Ch19	1997Ch53	
6037 (3)				< 0.002	< 0.002
6322.0 (16)			0.005 (1)	0.012 (6)	0.0049 (4)
6484.7 (16)	0.17 (3)	0.02 (1)	0.021 (2)	0.022 (2)	0.0167 (8)
6813.8 (16)	0.55 (9)	0.06 (2)	0.036 (3)	0.038 (4)	0.0384 (15)
7066.9 (16)	98.5 (10)	99.9 (1)	99.9	> 99.9	99.932 (3)

3. Atomic data

Atomic values ($\omega_K, \omega_L, \omega_M, \eta_{KL}$ and η_{LM}) are from Schönfeld (1996Sc06).

The X-ray and Auger electron emission probabilities have been deduced from γ -ray and conversion electron data by using the computer code RADLST.

4. Electron Emissions.

The conversion electron emission probabilities have been deduced from γ -ray transition data.

5. Photon Emissions

5.1 γ -ray energy values

The measured and evaluated γ -ray energies for ²¹⁷At α decay are listed in table 4. The evaluated values are from 1997Ch19. The 455 keV γ -ray is introduced by evaluators due to probabilities balance. This γ -ray was observed in 1964Va20, but not confirmed by 1997Ch19. 1997Ch53 assigned the 6037 keV α transition and introduced the 1050 keV level.

Table 4: Measured and evaluated value of γ -ray energy for ²¹⁷At.

1981Di14	1997Ch19	Evaluation
	165.8 ^a	165.8 ^a
258.5 (2)	257.88 (4)	257.88 (4)
	335.33 (10)	335.33 (10)
		455 ^b
	501.0 ^a	501.0 ^a
593.1 (2)	593.10 (10)	593.10 (10)
	758.9 (1)	758.9 (1)

^a: not placed in level scheme.

^b: from 1964Va20.

5.2 Absolute values of the γ -ray emission probabilities

The measured and evaluated γ -ray emission probabilities for ²¹⁷At α decay are listed in table 5. The evaluated values are from 1997Ch19, except as noted.

Table 5: Measured and evaluated γ -ray emission probabilities for ²¹⁷At.

E_{γ} (keV)	P_{γ}		
	1981Di14	1997Ch19	Evaluation
165.8 ^a		< 0.0002	< 0.0002
257.88 (4)	0.065 (5)	0.0287 (7)	0.0287 (7)
335.33 (10)		0.0062 (3)	0.0062 (3)
455			< 0.002 ^b
501.0 ^a		< 0.0002	< 0.0002
593.10 (10)	0.014 (1)	0.0115 (5)	0.0115 (5)
758.9 (1)		0.0049 (4)	0.0049 (4)

^a: not placed in level scheme.

^b: from intensity balance.

6. Branching Ratio

The measured and evaluated branching ratio for ²¹⁷At β^{-} decay are listed in table 6. The evaluated β^{-} decay branching ratio is from 1997Ch53, that's (% β^{-}) = 0.0067 (24) %. So (% α) = 99.9933 (24) %.

Table 6: Measured and evaluated branching ratio for ²¹⁷At β⁻ decay.

I_{β^-} (%)	References
0.0012 (6)	1969LeZW
0.005	1995Ch74
0.0067 (24)	1997Ch53
0.0067 (24)	Evaluated value, from 1997Ch53

7. References

- 1947En03 A.C. English, T.E. Cranshaw, P. Demers, J.A. Harvey, E.P. Hincks, J.V. Jelley, A.N. May, Phys. Rev. 72, 253 (1947) [T_{1/2}].
- 1950Ha52 F. Hagemann, L.I. Katzin, M.H. Studier, G.T. Seaborg, A. Ghiorso, Phys. Rev. 79, 435 (1950) [T_{1/2}].
- 1963Di05 H. Diamond, J.E. Gindler, J. Inorg. Nucl. Chem. 25, 143 (1963) [T_{1/2}].
- 1964Va20 K.Valli, Ann. Acad. Sci. Fennicae, Ser.A VI, No.165 (1964) [E_γ].
- 1967Dz02 B.S. Dzhelepov, R.B. Ivanov, M.A. Mikhailova, L.N. Moskvina, O.M. Nazarenko, V.F. Rodionov, Izv. Akad. Nauk SSSR, Ser. Fiz. 31, 568 (1967); Bull. Acad. Sci. USSR, Ser. Fiz. 31, 563 (1968) [E_α, I_α].
- 1969LeZW C.-F.Leang, Thesis, Univ. Paris (1969) [E_α, I_α, P_{β⁻}].
- 1977Vy02 T. Vylov, N.A. Golovkov, B.S. Dzhelepov, R.B. Ivanov, M.A. Mikhailova, Y.V. Narseev, V.G. Chumin, Izv. Akad. Nauk SSSR, Ser. Fiz. 41, 1635 (1977); Bull. Acad. Sci. USSR, Phys. Ser. 41, No.8, 85 (1977) [E_α].
- 1981Di14 J.K. Dickens, J.W. McConnell, Radiochem. Radioanal. Lett. 47, 331 (1981) [E_γ, P_γ].
- 1982Bo04 J.D. Bowman, R.E. Eppley, E.K. Hyde, Phys. Rev. C25, 941 (1982) [E_α].
- 1995Ch74 V.G. Chumin, S.S. Eliseev, K.Ya. Gromov, Yu.V. Narseev, V.I. Fominykh, V.V. Tsupko-Sitnikov, Bull. Rus. Acad. Sci. Phys. 59,1854(1995) [P_{β⁻}].
- 1996Sc06 E. Schönfeld, H.Janssen, Nucl. Instrum. Meth. Phys. Res. A369, 527 (1996) [Atomic data].
- 1997Ch19 V.G. Chumin, V.I. Fominykh, K.Ya. Gromov, M.Ya. Kuznetsova, V.V. Tsupko-Sitnikov, M.B. Yuldashev, Z. Phys. A358, 33 (1997) [E_α, I_α, E_γ, P_γ, P_{β⁻}, Multipolarity].
- 1997Ch53 V.G. Chumin, J.K. Jabber, K.V. Kalyapkin, S.A. Kudrya, V.V. Tsupko-Sitnikov, K.Ya. Gromov, V.I. Fominykh, T.A. Furyaev, Bull. Rus. Acad. Sci. Phys. 61, 1606 (1997) [P_α, P_{β⁻}].
- 2003Au03 G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A729, 129 (2003) [Q].
- 2007Ba19 M.S. Basunia, Nucl. Data Sheets 108, 633 (2007) [Spin, parity].
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr., Nucl. Instrum. Meth. Phys. Res. A589, 202 (2008) [Theoretical ICC].

**²¹⁷Rn - Comments on evaluation of the decay data
by Huang Xiaolong, Wang Baosong**

This evaluation was completed in 2007. Literature available by December 2007 was included.

1 Decay Scheme

²¹⁷Rn disintegrates 100 % by α emission to the levels in ²¹³Po. α decay of ²¹⁷Rn occurs directly to the ground state of ²¹³Po. ²¹⁷Rn ground state has $J^\pi = 9/2^+$ (2007Ba19).

2 Nuclear Data

The Q value is from the 2003Au03 evaluation.

The level energy, spin and parity are from 2007Ba19.

The measured and recommended ²¹⁷Rn half-life values are listed in Table 1.

Table 1 - Measured half-life values of ²¹⁷Rn and recommended value, in ms.

$T_{1/2}$ (ms)	References	notes
1.0 (1)	1951Me10	
0.54 (5)	1961Ru06	
0.54 (5)	2007Ba19	Nucl. Data Sheets, from 1961Ru06
0.54 (5)	Recommended value	from 1961Ru06

The recommended value is taken from the measurement of 1961Ru06.

2.1 α Transition

The measured alpha-particle energies are listed in table 2. The Q-value of 2003Au03 was used to determine the energy and uncertainty of the single alpha particle transition to the ground state of ²¹³Po.

An α transition of energy 7507 keV (no uncertainty) with ~ 0.1 % intensity was observed by 1961Ru06. The first excited state in ²¹³Po has been observed at 293 keV in ²¹³Bi decay. If the 7507 keV group belonged to ²¹⁷Rn, from the 7887 keV it would give 243 keV for the level energy of the first excited state. In addition 1961Ru06 did not observe any α - γ coincidence. The evaluator believes that the uncertain 7507 keV group reported by 1961Ru06 probably belongs to an impurity because no positive identification could be made by 1961Ru06.

Table 2 - Measured and recommended values of α -particle energy from ²¹⁷Rn decay

1961Ru06	1982Bo04	1991Ry01 ^a	Adopted value
7735 (4)	7739 (2)	7741 (2)	7742 (3)

a: Original energies of 1982Bo04 have been increased by 2 keV due to changes in calibration energies (1991Ry01).

So the evaluated alpha particle emission probability of the single alpha particle is 100 %.

The alpha hindrance factor $HF = 1.49$ was calculated using a parameter $R0 = 1.562$ (8) (2007Ba19), average of $R0(^{212}\text{Po}) = 1.5649$ (8) and $R0(^{214}\text{Po}) = 1.559$ (8) ; (1998Ak04).

3. References

- 1951Me10 W.W.Meinke, A.Ghiorso, G.T.Seaborg, Phys.Rev. 81, 782 (1951) [$T_{1/2}$].
1961Ru06 C.P.Ruiz, UCRL-9511 (1961) [$E_{\alpha}, T_{1/2}$].
1982Bo04 J.D.Bowman, R.E.Eppley, E.K.Hyde, Phys.Rev. C25, 941 (1982) [E_{α}].
1991Ry01 A.Rytz, At.Data Nucl.Data Tables 47, 205 (1991) [Evaluation].
2003Au03 G.Audi, A.H.Wapstra, C.Thibault, Nucl. Phys. A729(2003)129 [Q].
2007Ba19 M.S.Basunia, Nucl.Data Sheets 108, 633 (2007) [NDS]

**²¹⁸Po - Comments on evaluation of decay data
by V. Chisté and M. M. Bé**

This evaluation was completed in 2007. Literature available by January 2007 was included.

1 Decay Scheme

²¹⁸Po disintegrates by alpha emission mainly (99.978 (3) %) to the ground state level of ²¹⁴Pb. A weak beta minus emission (0.022 (3) %) to At -218 has been pointed out. Spin and parity are from the mass -chain evaluation of Y. A. Akevali (1987E112, 1995E108, 1998Ak04 for A = 218 and 1995E107 for A = 214) and A. K. Jain (2006Ja03 for A = 218).

A good agreement was found between the recommended Q value of Audi and the effective Q value of 6113.33 (22) keV for the α branch, calculated from the decay scheme data.

2 Nuclear Data

The Q values (α and β^-) are from the atomic mass evaluation of Audi *et al.* (2003Au03).

Experimental ²¹⁸Po half-life values (in minutes) are given in Table 1:

Table 1: Experimental values of ²¹⁸Po half-life.

Reference	Experimental value (min)	Comments
M. Curie (931Cu01)	3.05	Not used.
M. Blau (1924Bl02)	3.050 (9)	Not used.
J. R. Van Hise (1982Va09)	3.11 (2)	Uncertainty increased to take into account systematic uncertainty.
G. V. Potapov (1986Po17)	3.093 (6)	Original uncertainty corresponds to two standard deviations.
Recommended value	3.094 (6)	$\chi^2 = 0.66$

The recommended value was deduced from the two most recent values of ²¹⁸Po half-life (1982Va09 and 1986Po17), the value of M. Blau (1924Bl02) was omitted from analysis due to the difficulty in estimating a realistic uncertainty. The original uncertainty value given by Van Hise (1982Va09) is for 2σ , but it seems that they did not take into account the systematic uncertainties so the original uncertainty has been maintained.

A weighted average of 3.094 minutes has been calculated using Lweightcomputer program (version 3), with an internal uncertainty of 0.006 minutes. The reduced- χ^2 value is 0.66.

2.1 α Transitions and Emissions

The energies of the α -particle transitions given in Section 2.1 were calculated from Q_α (2003Au03) and level energies.

The energy of $\alpha_{0,0}$ emission given in section 4 was measured by 1971Gr17, and following the recommendations given by A. Rytz (1991Ry01) was decreased by 0.20 keV. The $\alpha_{0,1}$ emission energy is from R. J. Walen (1958Wa16).

The $\alpha_{0,1}$ emission probability is the measured value of R. J. Walen (1958Wa16) (0.0011 (11) %).

For the $\alpha_{0,0}$ emission probability and associated uncertainty, the following relation was applied:

$$P_{\alpha_{0,0}} + P_{\alpha_{0,1}} = 100 - P_{\beta}(264 \text{ keV}),$$

where $P_{\beta}(264 \text{ keV}) = 0.022 (3) \%$ (given by 1952Hi60, see **2.2**) and $P_{\alpha_{0,1}} = 0.0011 (11) \%$ (given by 1958Wa16). Taking into account these values, then $P_{\alpha_{0,0}} = 99.9769 (32) \%$.

2.2 β^- Transitions and Emissions

The maximum energy of the β^- transition in the decay of $^{218}\text{Po} \rightarrow ^{218}\text{At}$ has been taken from Audi (2003Au03) and, without any other information, is affected to a ground state to ground state transition.

The adopted 260-keV β^- transition probability was measured by F. Hiessberger (1952Hi60), 0.022(3) %, and is in agreement with the two values given by R. J. Walen : 0.0200 (5) % (1949Wa05) and 0.0185 % (1958Wa16), respectively.

2.3 g Transitions and Emissions

The $\gamma_{(1,0)}$ transition probability following the α -decay of $^{218}\text{Po} \rightarrow ^{214}\text{Pb}$ was deduced from the decay-scheme balance using the recommended experimental α -particle intensity value of 0.0011 (11) % given by R. J. Walen (1958Wa16). (see **2.1 a Transitions and Emissions**).

3 Atomic Data

Atomic values, ω_K , ω_L and n_{KL} and the X-ray relative probabilities are from Schönfeld and Janßen (1996Sc06).

4 References

- 1924B102 – M. Blau, Akad. Wiss. Wien, Berlin, 133(1924)17 [Half-life].
 1931Cu01 – M. Curie, A. Debierne, A. S. Eve, H. Geiger, O. Hahn, S. C. Lind, St. Meyer, E. Rutherford, E. Schweidler, Rev. Mod. Phys. 3(1931)427 [Half-life (^{214}Pb)].
 1949Wa05 – R. J. Walen, J. Phys. Radium 10(1949)95 [I_{β} , Half-life (^{218}At)].
 1952Hi60 – F. Hiessberger, B. Karlik, Stizber. Akad. Wiss. Wien, Math– Naturw. Kl. Abt. Ila 161(1952)51 [Branching ratio].
 1958Wa16 – R. J. Walen, G. Bastin, Comp. Rend. Int. Conf. Nucl. Phys., Paris, (1959)910 [E_{α} , I_{α} , I_{β}].
 1963Ba62 – G. Bastin-Scoffier, C. F. Leang, R. J. Walen, J. Phys. (Paris) 24(1963)854 [E_{α}].
 1971Gr17 – B. Grennberg, A. Rytz, Metrologia 7(1971)65 [E_{α}].
 1979Ry03 – A. Rytz, At. Data and Nucl. Data Tables 23(1979)205 [E_{α} , I_{α}].
 1982Va09 – J. R. Van Hise, D. E. Martin, R. A. Jackson, D. Y. Kunihira, E. Bolton, Phys. Rev. C25(1982)2802 [Half-life].
 1986Po17 – G. V. Potapov, P. S. Soloshenkov, Sov. At. Energ. 60(1986)345 [Half-life].
 1987El12 – Y. A. Akovali, Nucl. Data Sheets 52(1987)789 [Spin, parity and multipolarity].
 1991Ry01 – A. Rytz, At. Data and Nucl. Data Tables 47(1991)205 [E_{α} , I_{α}].
 1992Ba61 – P. Baltzer, K. G. Görden, A. Bäcklin, Nucl. Instrum. Meth. Phys. Res. A317(1992)357 [E_{α}].
 1995El07 – Y. A. Akovali, Nucl. Data Sheets 76(1995)127 [Energy level, spin, parity and multipolarity].
 1995El08 – Y. A. Akovali, Nucl. Data Sheets 76(1995)457 [Energy level and half-life].
 1996Sc06 – E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
 1998Ak04 – Y. A. Akovali, Nucl. Data Sheets 84(1998)1 [Branching ratio].
 2003Au03 – G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].
 2006Ja03 – A. K. Jain, B. Singh, Nucl. Data Sheets 107(2006)1027 [Spin, parity and multipolarity].

**²¹⁸At - Comments on evaluation of decay data
by V. Chisté and M. M. Bé**

This evaluation was completed in 2007. Literature available by January 2007 was included.

1 Decay Scheme

²¹⁸At disintegrates by alpha emission (99.9 (1) %) to ²¹⁴Bi mainly. The γ transitions between the ²¹⁴Bi levels have not been observed. However, a Q value of 6811 (12) keV is calculated in the disintegration of ²¹⁸At to ²¹⁴Bi from the decay scheme data compared to a value of 6867(3) keV from the Audi's tables. This deficiency in the calculated Q value suggests the possible existence of a weak gamma transition from the 62-keV to the ground state levels.

A weak beta minus emission (0.1 (1) %) to Rn-218 has been pointed out (1948Wa20). Spins and parities are from the mass-chain evaluation of Y. A. Akevali (1987E112, 1995E108 for A = 218 and 1995E107 for A = 214) and A. K. Jain (2006Ja03 for A = 218).

2 Nuclear Data

The Q values (α and β^-) are from the atomic mass evaluation of Audi *et al.* (2003Au03).

Experimental ²¹⁸At half-life values (in seconds) are given in Table 1:

Table 1: Experimental values of ²¹⁸At half-life.

Reference	Experimental value (s)	Comments
R. J. Walen (1949Wa05)	1.3 (2)	Uncertainty increased to take into account systematic uncertainty.
D. G. Burke (1989Bu09)	1.5 (3)	
Recommended value	1.4 (2)	$\chi^2 = 0.31$

The original uncertainty value given by R. J. Walen (1949Wa05) was multiplied by 2, in order to take into account the systematic uncertainties which were not considered by 1949Wa05. A weighted average has been calculated using Lweight computer program (version 3).

The recommended value of ²¹⁸At half-life is the weighted average of **1.4** second with an internal uncertainty of 0.2 second. The reduced- χ^2 value is 0.31.

2.1 α Transitions and Emissions

The energies of the α -particle transitions given in Section 2.1 were calculated from Q_α (2003Au03) and level energies.

The energy of $\alpha_{0,0}$, $\alpha_{0,1}$ and $\alpha_{0,2}$ emissions given in section 3 were measured by R.J. Walen (1963Wa29 (see 1964Hy02) and 1958Wa16), the adopted values are those recommended by A. Rytz (1991Ry01) where the original energy was decreased by 1 keV, due to a change in calibration energy (1995E107).

The $\alpha_{0,0}$, $\alpha_{0,1}$ and $\alpha_{0,2}$ emission probabilities are the measured values of R. J. Walen (1958Wa16), 3.6, 90.0 and 6.4 respectively, without uncertainties. From R. J. Walen, the total α decay is 99.9 (1) %. Since, there is no precision in the Walen's paper, the uncertainty of 0.1 % from propagation of the β^- transition probability (1948Wa20) has been assigned to each α line.

2.2 β^- Transitions and Emissions

The maximum energy of the β^- transition in the decay of $^{218}\text{At} \rightarrow ^{218}\text{Rn}$ is given by Audi (2003Au03) and, without any other information available, is affected to a ground state to ground state transition.

The adopted β^- transition probability was measured by R. J. Walen (1948Wa20) to be 0.1 (1) %

3 References

- 1948Wa20 R. J. Walen, *Comp. Rend. Acad. Sci. (Paris)* 227(1948)1090 [Branching ratio].
1949Wa05 R. J. Walen, *J. Phys. Radium* 10(1949)95 [I_β , Half-life].
1958Wa16 R. J. Walen, G. Bastin, *Comp. Rend. Int. Conf. Nucl. Phys., Paris, (1959)*910 [E_α , I_α , I_β].
1963Wa29 R. J. Walen, G. Bastin-Scoffier, *Priv. Comm. quoted by 1964Hy02 (1963)* [I_α].
1964Hy02 E. K. Hyde, I. Perlman, G. T. Seaborg, *The nuclear properties of heavy elements, Vol. II*, Prentice-Hall, Inc., Englewood Cliffs, N.J., (1964)460 [I_α].
1979Ry03 A. Rytz, *At. Data and Nucl. Data Tables* 23(1979)205 [E_α , I_α].
1987El12 Y. A. Ellis-Akovi, *Nucl. Data Sheets* 52(1987)789 [I_α , E_α , spin and parity].
1989Bu09 D. G. Burke, H. Folger, H. Gabelmann, E. Høegbø, P. Hill, P. Hoff, O. Jonsson, N. Kaffrell, W. Kurcewicz, G. Løvhøiden, K. Nybø, G. Nyman, H. Ravn, K. Riisager, J. Rogowski, K. Steffensen, T. F. Thorsteinsen and ISOLDE Collaboration, *Z. Phys. A*333(1989)131 [Half-life].
1991Ry01 A. Rytz, *At. Data and Nucl. Data Tables* 47(1991)205 [E_α , I_α].
1995El07 Y. A. Akovi, *Nucl. Data Sheets* 76(1995)127 [I_α , E_α , spin and parity].
1995El08 Y. A. Akovi, *Nucl. Data Sheets* 76(1995)457 [I_α , E_α , spin and parity].
2003Au03 G. Audi, A. H. Wapstra, C. Thibault, *Nucl. Phys. A*729(2003)129 [Q].
2006Ja03 A. K. Jain, B. Singh, *Nucl. Data Sheets* 107(2006)1027 [I_α , E_α , spin and parity].

**²¹⁸Rn - Comments on evaluation of decay data
by V. Chisté and M. M. Bé**

This evaluation was completed in 2007. Literature available by January 2007 was included.

1 Decay Scheme

²¹⁸Rn disintegrates by alpha emissions to the 609 -keV level (0.127 (7) %) and to the ground state (99.873 (7) %) of ²¹⁴Po. Spins and parities are from the mass-chain evaluation of Y. A. Akovali (1987E112, 1995E108, 1998Ak04 and 2006Ja03 for A = 218 and 1995E107 for A = 214).

A good agreement was found between the recommended Q value from Audi and the effective Q value (7262.5 (20) keV) calculated from decay scheme data.

2 Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2003Au03).

Experimental ²¹⁸Rn half-life values (in ms) are given in Table 1:

Table 1: Experimental values of ²¹⁸Rn half-life, in ms.

Reference	Experimental value (ms)	Comments
M.H. Studier (1948St42)	19	
P. A. Tove (1958To25)	39 (4)	
C. P. Ruiz (1961Ru06)	30 (3)	
H. Diamond (1963Di05)	35 (2)	Original uncertainty $\times 2$
A. Erlik (1971Er02)	39 (2)	
Recommended value	36.0 (19)	reduced $\chi^2 = 2.3$, critical $\chi^2 = 3.8$

The original uncertainty of Diamond includes statistical uncertainty only, it was multiply by 2 to try to take into account systematic components.

A weighted average has been calculated using Lweight computer program (version 3), then the recommended value of ²¹⁸Rn half-life is **36.0 ms** with an external uncertainty of **1.9 ms**.

2.1 a Transitions and Emissions

The energies of the α -particle transitions given in Section 2.1 were calculated from Q_α (2003Au03) and the level energies.

The energy of $\alpha_{0,0}$ emission given in section 4 is the weighted average of the two measured values of F. Asaro (1956As38) and J. D. Bowan (1982Bo04), with the recommendations given by A. Rytz (1991Ry0) where the original energy of 1956As38 was increased by 4 keV and the energy of 1982Bo04 was decreased by 4 keV, due to changes in calibration energies (1998Ak04). For the $\alpha_{0,1}$, the emission energy was calculated from Q_α (2003Au03), the level energy and taking the nucleus recoil into account.

The α emission probabilities were deduced from the level decay -scheme balance (see **2.2 Gamma Transitions**).

2.2 g Transitions

The 609-keV γ -ray transition probability was calculated using the γ -ray emission intensity and the relevant internal conversion coefficient (see **4.2 g Emissions**).

Multipolarity of this γ -ray transition (E2) is from 1995El04.

The internal conversion coefficient (ICC) for the 609keV γ -ray transition has been calculated using the BrIcc computer program (calculation for 'hole'), based on the theoretical values of I. M. Band (2002Ba85).

3 Atomic Data

Atomic values, ω_K , $\bar{\omega}_L$ and n_{KL} and the X-ray relative probabilities are from Schönfeld and Janßen (1996Sc06).

4 Photon Emissions

4.1 X-ray Emissions

The X-ray absolute intensities were calculated from γ -ray data and ICC using the EMISSION computer program.

4.2 g Emissions

The energy of the 609-keV γ -ray given in section 5.2 is from W. Kurcewicz (1976Ku08).

The emission intensity of the 609-keV γ -ray was calculated from the measured relative photon intensity of W. Kurcewicz (1976Ku08), who measured the U-230 decay chain, and from the absolute emission intensity of 2.77 (8) % for the 324.22-keV γ -ray of ²²²Ra decay, as measured by A. Peghaire (1969Pe17). This 609-keV emission intensity was then deduced being 0.124 (7) %.

This result can be compared with the less precise measured absolute intensities of 0.20 (5) (1956As38) and 0.16 (5) (1963Le17).

5 References

- 1948St42 – M.H. Studier, E.K. Hyde, Phys. Rev. 74 (1948) 591 [Half-life].
 1956As38 – F. Asaro, I. Perlman, Phys. Rev. 104(1956)91 [E_α].
 1958To25 – P. A. Tove, Ark. Fys. 13(1958)549 [Half-life].
 1961Ru06 – C. P. Ruiz, UCRL – 9511 (1961) [Half-life].
 1963Di05 – H. Diamond, J. E. Gindler, J. Inorg. Nucl. Chem. 25(1963)143 [Half-life].
 1963Le17 – C. M. Lederer, UCRL – 11028(1963) [I_γ].
 1969Pe17 – A. Peghaire, Nucl. Instrum. Meth. 75(1969)66 [I_γ].
 1971Er02 – A. Erlik, J. Felsteiner, H. Lindeman, M. Tatcher, Nucl. Instrum. Meth. 92(1971)45 [Half-life].
 1976Ku08 – W. Kurcewicz, N. Kaffrell, N. Trautmann, A. Plochocki, J. Kylicz, K. Stryczniewicz, I. Yutlandov, Nucl. Phys. A270(1976)175 [I_γ , E_γ , I_α].
 1979Ry03 – A. Rytz, At. Data and Nucl. Data Tables 23(1979)205 [E_α , I_α].
 1982Bo04 – J. D. Bowman, R. E. Eppley, E. K. Hyde, Phys. Rev. C25(1982)941 [E_α].
 1987El12 – Y. A. Ellis-Akovi, Nucl. Data Sheets 52(1987)789 [I_α , E_α , I_γ , E_γ , spin and parity].
 1991Ry01 – A. Rytz, At. Data and Nucl. Data Tables 47(1991)205 [E_α , I_α].
 1995El07 – Y. A. Akovi, Nucl. Data Sheets 76(1995)127 [I_α , E_α , I_γ , E_γ , spin and parity].
 1996Sc06 – E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
 1998Ak04 – Y. A. Akovi, Nucl. Data Sheets 84(1998)1 [I_α , E_α , I_γ , E_γ , spin and parity].
 2002Ba85 – I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr., P. O. Tikkanen, S. Raman, At. Data Nucl. Data Tables 81(2002)1 [Theoretical ICC].
 2003Au03 – G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].
 2006Ja03 – A. K. Jain, B. Singh, Nucl. Data Sheets 107(2006)1027 [Spin, parity and multipolarity].

²¹⁹At - Comments on evaluation of decay data by A. L. Nichols

Evaluated: September 2010

Evaluation Procedure

Limitation of Relative Statistical Weight Method (LWM) was applied to average numbers throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

Very little of substance can be gleaned from the literature concerning the decay characteristics of ²¹⁹At (2001Br31). Although no γ or β^- emissions have been observed, an alpha group at 6.27 MeV was assigned to ²¹⁹At by Hyde and Ghiorso (1953Hy83). A simple decay scheme has been constructed with little confidence from this early study. Alpha and β^- feeding directly to the ground states of daughter ²¹⁵Bi and ²¹⁹Rn were assumed, but these processes were neither observed satisfactorily nor quantified experimentally. Spin and parity of $7/2^-$ were tentatively assigned to the ground state of ²¹⁹At to align with $5/2^+$ identified with the ground state of daughter ²¹⁹Rn (2001Br31), in order to define the proposed single beta-particle emission as first forbidden non-unique. Further spectral studies are required to assemble and quantify the decay scheme of ²¹⁹At with much greater confidence.

Nuclear Data

Part of the $(4n + 3)$ naturally-occurring decay chain, and of relevance in quantifying the environmental impact of ²³⁵U and resulting decay-chain products. Specific radionuclides in this decay chain are noteworthy because of their distinctive decay characteristics (e.g. alpha decay of ²¹⁵Po, ²¹¹Bi and ²¹¹Po).

Half-life

The recommended half-life is the weighted mean of only two measurements (1953Hy83 and 1989Bu09).

Reference	Half-life (sec)
1953Hy83	54 (6)
1989Bu09	57 (4)
Recommended value	56 (4)*

*Uncertainty adjusted upwards from ± 3 to ± 4 in line with the most precise value of this limited data set.

Q values

Q^- of 1566 (3) keV and Q_α of 6324 (15) keV were adopted from the evaluated tabulations of Audi *et al.* (2003Au03, 2009AuZZ).

Alpha particleEnergy

The alpha-particle branch of ~ 97 % was assumed to populate the ground state of ²¹⁵Bi directly. Both the energy and uncertainty of this proposed alpha-particle emission were calculated to be 6208 (15) keV from the evaluated Q-value of 6324 (15) keV (2003Au03, 2009AuZZ).

Emission Probability

The alpha-particle emission probability was calculated from a quoted α/β^- ratio of approximately 30, as determined from measurements of the ²¹⁹At/²¹⁹Rn peak ratio (1953Hy83). An ill-defined alpha branch of ~ 97 % can be derived from this ratio without an assigned uncertainty.

Alpha-particle emission probability per 100 disintegrations of ²¹⁹At, and hindrance factor

$E_\alpha(\text{keV})$	P_α	HF
	Recommended value	
6208 (15)	~ 97	~ 1.07

An unweighted mean value of 1.547 (9) was adopted for the radius parameter $r_0(^{215}\text{Bi})$ as derived from the equivalent data for neighbouring nuclei (1998Ak04), and used in the calculation of the α -hindrance factor (HF):

$$r_0(^{215}\text{Bi}) = [r_0(^{214}\text{Pb}) + r_0(^{216}\text{Po})] / 2$$

$$= [1.5379 (7) + 1.5555 (2)] / 2 = 1.547 (9)$$

Beta particleEnergy

The beta-particle branch of ~ 3 % was assumed to populate the ground state of ²¹⁹Rn directly. Therefore, the recommended energy and uncertainty of this single beta-particle transition was adopted from the evaluated Q-value of 1566 (3) keV (2003Au03, 2009AuZZ).

Emission Probability

The beta-particle emission probability was calculated from a quoted α/β^- ratio of approximately 30, as determined from measurements of the ²¹⁹At/²¹⁹Rn peak ratio (1953Hy83). A single, ill-defined, first forbidden non-unique transition is proposed directly to the ground state of ²¹⁹Rn, with an emission probability of ~ 3 % and no assigned uncertainty.

Beta-particle emission probability per 100 disintegrations of ²¹⁹At, transition type and log ft

$E_\beta(\text{keV})$	P_β	transition type	log ft
	Recommended value		
1566 (3)	~ 3	(1 st forbidden non-unique)	~ 6.2

Data Consistency

An effective Q-value of 6181 (15) keV has been adopted from the atomic mass evaluation of Audi *et al.* (2003Au03, 2009AuZZ) while in the course of formulating the decay scheme of ²¹⁹At. This value has subsequently been compared with the Q-value calculated by summing the contributions of the individual emissions to the ²¹⁹At alpha- and beta-decay processes:

$$\text{calculated Q-value} = \sum (E_i \times P_i) = 6181 (15) \text{ keV}$$

Percentage deviation from the effective Q-value of Audi *et al.* is $(0.0 \pm 0.3) \%$, which supports the derivation of a highly consistent decay scheme.

References

- 1953Hy83 E.K. HYDE, A. GHIORSO, The Alpha-branching of AcK and the Presence of Astatine in Nature, *Phys. Rev.* 90 (1953) 267-270. [Half-life, E_α , α and β^- decay, α/β^- ratio]
- 1989Bu09 D.G. BURKE, H. FOLGER, H. GABELMANN, E. HAGEBØ, P. HILL, P. HOFF, O. JONSSON, N. KAFFRELL, W. KURCEWICZ, G. LØVHØIDEN, K. NYBØ, G. NYMAN, H. RAVN, K. RIISAGER, J. ROGOWSKI, K. STEFFENSEN, T.F. THORSTEINSEN, and the ISOLDE Collaboration, New Neutron-rich Isotopes of Astatine and Bismuth, *Z. Phys. – Atomic Nuclei* 333 (1989) 131-135. [Half-life]
- 1998Ak04 Y.A. AKOVALI, Review of Alpha-decay Data from Doubly-even Nuclei, *Nucl. Data Sheets* 84 (1998) 1-114. [alpha decay, r_0 parameters]
- 2001Br31 E. BROWNE, Nuclear Data Sheets for A = 215, 219, 223, 227, 231, *Nucl. Data Sheets* 93 (2001) 763-1061. [Nuclear structure, level energies]
- 2003Au03 G. AUDI, A.H. WAPSTRA, C. THIBAULT, The AME2003 Atomic Mass Evaluation (II). Tables, Graphs and References, *Nucl. Phys.* A729 (2003) 337-676. [Q-value]
- 2009AuZZ G. AUDI, W. MENG, D. LUNNEY, B. PFEIFFER, Atomic Mass Evaluation 2009, Atomic Mass Data Centre, Centre de Spectrométrie Nucléaire et de Spectrométrie de Mass (CSNSM), Orsay, France, private communication, 2009. [Q-value]

**²¹⁹Rn – Comments on evaluation of decay data
by A. L. Nichols**

Evaluated: October 2010

Evaluation Procedure

Limitation of Relative Statistical Weight Method (LWM) was applied to average numbers throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

A reasonably comprehensive decay scheme has been derived from the alpha-particle studies of 1962Wa18, 1991Ry01 and 1999Li05, and the gamma-ray measurements of 1967Da20, 1968Br17, 1970Da09, 1970Kr08 and 1999Li05.

Nuclear Data

Part of the (4n + 3) decay chain, and of relevance in quantifying the environmental impact of ²³⁵U and various decay-chain products. Specific radionuclides in this decay chain are noteworthy because of their distinctive and important decay characteristics (e.g. alpha decay of ²¹⁵Po, ²¹¹Bi and ²¹¹Po).

Half-life

The recommended half-life is the weighted mean of two measurements by 1961Ro14, and an additional independent study by 1966Hu20.

Reference	Half-life (s)
1961Ro14	4.01 (6)
	4.00 (5)
1966Hu20	3.96 (1) [#]
Recommended value	3.98 (3)

[#] Uncertainty adjusted to ± 0.04 to reduce weighting below 50 %.

Q value

Q_{α} of 6946.1 (3) keV was adopted from the evaluated tabulations of Audi *et al.* (2003Au03).

Alpha Particles

Energies

Alpha-particle energies were calculated from the structural details of the proposed decay scheme. While the energies of the alpha-particle emissions have been directly measured by 1962Wa18, 1971Gr07, 1991Ry01 and 1999Li05, the nuclear level energies of 2001Br31 and evaluated Q-value of 6946.1 (3) keV (2003Au03) were preferably used to determine the recommended energies and uncertainties of the alpha-particle transitions, taking into account the significant recoil.

Adopted nuclear levels of ²¹⁵Po: J^π and origins (2001Br31).

Nuclear level	Nuclear level energy (keV)	J ^π	Origins
0	0.0	9/2 +	²¹⁵ Bi β ⁻ decay, ²¹⁹ Rn α decay
1	271.228 ± 0.010	7/2 +	²¹⁵ Bi β ⁻ decay, ²¹⁹ Rn α decay
2	293.56 ± 0.04	(11/2) +	²¹⁵ Bi β ⁻ decay, ²¹⁹ Rn α decay
3	401.812 ± 0.010	5/2 +	²¹⁵ Bi β ⁻ decay, ²¹⁹ Rn α decay
4	517.60 ± 0.06	7/2 +, 9/2 +	²¹⁵ Bi β ⁻ decay, ²¹⁹ Rn α decay
5	608.30 ± 0.07	(11/2 +, 13/2 +)	²¹⁹ Rn α decay
6	676.66 ± 0.07		²¹⁵ Bi β ⁻ decay, ²¹⁹ Rn α decay
7	708.1 ± 0.5		²¹⁹ Rn α decay
8	732.7 ± 0.4		²¹⁹ Rn α decay
9	835.32 ± 0.22		²¹⁵ Bi β ⁻ decay, ²¹⁹ Rn α decay
10	877.2 ± 0.6		²¹⁹ Rn α decay
11	891.1 ± 0.3		²¹⁹ Rn α decay
12	930 ± 1		²¹⁹ Rn α decay
13	1073.7 ± 0.4	(5/2 +)	²¹⁹ Rn α decay
14	1094.2 ± 1.0		²¹⁹ Rn α decay

Measured and recommended energies of the alpha-particle emissions of ²¹⁹Ra.

	E _α (keV)				
	1962Wa18	1971Gr07	1991Ry01	1999Li05	Recommended value*
α _{0,14}	–	–	–	5744 (15)	5745.0 (10)
α _{0,13}	–	–	–	5764 (8)	5765.1 (5)
α _{0,12}	–	–	–	5900 (15)	5906.2 (10)
α _{0,11}	–	–	–	5944 (6)	5944.4 (4)
α _{0,10}	–	–	–	5958 (15)	5958.1 (7)
α _{0,9}	5999.3	–	–	6000 (6)	5999.2 (4)
α _{0,8}	6100.5	–	–	6100 (8)	6099.9 (5)
α _{0,7}	~ 6146.2	–	–	6124 (8)	6124.1 (6)
α _{0,6}	6157.1	–	–	6158 (4)	6154.9 (3)
α _{0,5}	6222.1	–	–	6223 (6)	6222.0 (3)
α _{0,4}	6310.3	–	–	6311 (3)	6311.1 (3)
α _{0,3}	6423.2	–	6425.0 (10)	6425 (1)	6424.8 (3)
α _{0,2}	6527.5	–	–	6530 (2)	6531.0 (3)
α _{0,1}	6551.3	–	6552.6 (10)	6553 (1)	6553.0 (3)
α _{0,0}	6817.5	6819.3 (3)	6819.1 (3)	6819.1 (3)	6819.2 (3)

* Determined from the nuclear level energies of 2001Br31 and evaluated Q-value of 6946.1 (3) keV (2003Au03).

Emission Probabilities

Alpha-particle emission probabilities were derived from the recommended relative emission probabilities of the gamma rays, a normalisation factor of 0.111 (5), and theoretical internal conversion coefficients (see below). The normalisation factor (F) was determined from the sum of the relative emission probabilities of the alpha particles calculated on the basis of α-γ population/depopulation balances of all the nuclear levels of ²¹⁵Po:

$$\sum[\text{calculated relative } P_{\alpha} \text{ to } ^{215}\text{Po excited states}]F + (\text{absolute } P_{\alpha} \text{ to } ^{215}\text{Po ground state}) = 100$$

An absolute P_α of 79.4 (10) % directly to the ²¹⁵Po ground state was adopted from 1991Ry01. Denoting F as the normalisation factor for the relative emission probabilities of both the gamma rays and alpha particles:

$$186.0626F + 79.4 (10) = 100$$

$$F = 20.6 (10) / 186.0626 = 0.1107 (54) \rightarrow 0.111 (5)$$

An unweighted mean value of 1.557 (2) was adopted for the radius parameter $r_0(^{215}\text{Po})$ as derived from the equivalent data for neighbouring nuclei (1998Ak04), and used in the calculation of α -hindrance factors (HF):

$$\begin{aligned} r_0(^{215}\text{Po}) &= [r_0(^{214}\text{Po}) + r_0(^{216}\text{Po})] / 2 \\ &= [1.559(8) + 1.5555(2)] / 2 = 1.557 (2) \end{aligned}$$

Alpha-particle emission probabilities per 100 disintegrations of ²¹⁹Rn, and hindrance factors.

$E_\alpha(\text{keV})$	P_α				HF
	1962Wa18	1991Ry01	1999Li05	Recommended value*	
5745.0 (10)	–	–	< 0.0001	0.000 09 (5)	245
5765.1 (5)	–	–	0.001	0.000 94 (19)	33
5906.2 (10)	–	–	–	0.000 09 (5)	1590
5944.4 (4)	–	–	0.002	0.002 1 (3)	103
5958.1 (7)	–	–	0.0001	0.000 3 (1)	830
5999.2 (4)	0.0044	–	0.003	0.003 2 (5)	120
6099.9 (5)	0.003	–	0.001	0.001 23 (12)	880
6124.1 (6)	~ 0.0026	–	0.001	0.000 64 (12)	2170
6154.9 (3)	0.0174	–	0.018	0.018 4 (22)	103
6222.0 (3)	0.0026	–	0.004	0.004 3 (10)	860
6311.1 (3)	0.054	–	0.054	0.048 (3)	184
6424.8 (3)	7.5	7.5 (6)	7.5	7.85 (24)	3.31
6531.0 (3)	0.12	–	0.12	0.098 (5)	710
6553.0 (3)	11.5	12.9 (6)	13	12.6 (3)	6.75
6819.2 (3)	81	79.4 (10)	79.3	79.4 (10)	11.2
Σ 100.02729 \rightarrow 100.0 (11)					

* Recommended alpha-particle emission probabilities have been determined by calculating the populating alpha-particle balances of the daughter nuclear levels of ²¹⁵Po through individual consideration of their gamma population-depopulation, along with the adoption of a normalisation factor of 0.111 (5) for the relative emission probabilities of the observed gamma rays and derived alpha particles.

Gamma Rays

Energies

While the energies of the main gamma-ray emissions have been directly measured by 1967Da20, 1968Br17, 1970Da09, 1970Kr08, 1976Bl13 and 1999Li05, all of the recommended gamma-ray energies were calculated from the nuclear level energies of daughter ²¹⁵Po as adopted from 2001Br31.

Measured and recommended energies of the main gamma-ray emissions of ²¹⁹Ra.

	E_γ (keV)						
	1967Da20	1968Br17	1970Da09	1970Kr08	1976Bl13	1999Li05	Recommended value*
$\gamma_{3,1}(Po)$	130.9 (6)	130.5 (3)	130.7 (1)	130.6 (2)	130.588 (29)	130.6 (1)	130.58 (1)
$\gamma_{1,0}(Po)$	271.2 (5)	271.0 (2)	271.20 (5)	271.4 (1)	271.233 (10)	271.23 (5)	271.228 (10)
$\gamma_{2,0}(Po)$	293.2 (6)	293.4 (4)	294.0 (3)	293.8 (2)	293.538 (44)	293.6 (1)	293.56 (4)
$\gamma_{3,0}(Po)$	401.7 (5)	401.7 (1)	401.8 (2)	402.0 (3)	401.811 (10)	401.81 (5)	401.81 (1)
$\gamma_{4,0}(Po)$	517.1 (8)	517.4 (3)	516.5 (5)	–	517.639 (55)	517.5 (1)	517.60 (6)
$\gamma_{6,0}(Po)$	–	676.6 (3)	677.0 (10)	–	676.645 (70)	676.7 (1)	676.66 (7)

* Determined from the recommended nuclear level energies of 2001Br31.

Emission Probabilities

The emission probabilities were determined from measurements of 1967Da20, 1968Br17, 1970Da09, 1970Kr08, 1976B113 and 1999Li05. Weighted mean values were calculated for the relative emission probabilities of the 130.58-, 293.56-, 401.81-, 517.60-, 608.30-, 676.66- and 891.1-keV gamma rays, while all others were adopted from the more comprehensive set of data measured by 1999Li05.

Some of the reported gamma-ray emissions were of highly questionable validity, and were not considered for placement in the recommended decay scheme because of their nature and doubtful origins:

- (a) 115.4-keV gamma ray was only observed by 1968Br17, and was furthermore labeled by these authors as ill-assigned – removed from consideration.
- (b) 221.5-keV gamma ray was judged to be a major gamma emission from the alpha-decay mode of ²²³Ra – removed from further consideration.
- (c) 324.9- and 337.7-keV gamma-ray emission probabilities were expressed only in terms of their upper limits by 1967Da20, and not observed by 1999Li05 – removed from consideration.
- (d) 370.9-keV gamma ray was observed by 1965Va10, an emission probability expressed only in terms of an upper limit by 1967Da20, and not observed by 1999Li05 – removed from consideration.
- (e) 380-keV gamma ray was only observed by 1965Va10 without a quantified emission probability – removed from consideration.
- (f) 438.2-keV gamma ray was judged to be a major gamma emission from the alpha-decay mode of ²¹⁵Po – removed from further consideration.
- (g) 538.2- and 1005-keV gamma rays were observed by 1965Va10, emission probabilities quantified by 1967Da20, and not observed by 1999Li05 – removed from consideration.
- (h) 665.5-keV gamma-ray emission probability was assigned an upper limit by 1967Da20 and identified as a possible doublet, and fully quantified as a singlet by 1999Li05 – retained as an unplaced gamma transition emitted in the decay of ²¹⁹Rn.

Although some of the other observed gamma-ray emissions possess similar origins to the above, these transitions could be more comfortably placed in the proposed decay scheme, lending support to their acceptance and inclusion in the recommended data set.

Published gamma-ray emission probabilities.

E _γ (keV)	P _γ						
	1965Va10*	1967Da20†	1968Br17‡	1970Da09¶	1970Kr08	1976Bi13	1999Li05
115.4 (5)	–	–	0.033 (15)	–	–	–	–
130.58 (1)	observed	1.40 (14)	1.30 (25)	1.05 (25)	1.21 (10)	1.16 (12)	1.7 (2)
221.5 (3)	–	–	–) 0.28 (7)	–	–	–
224.04 (7)	–	–	–)	–	–	0.013 (2)
271.228 (10)	observed	100	110 (15)	100.00	100.0	105.5 (40)	100 (2)
293.56 (4)	–	0.64 (6)	0.77 (15)	0.59 (15)	0.51 (27)	0.76 (5)	0.68 (4)
322 (1)	–	–	–	–	–	–	0.0008 (4)
324.9 (10)	–	< 0.06	–	–	–	–	–
330.9 (4)	–	–	–	–	–	–	0.0090 (10)
337.7 (10)	–	< 0.08	–	–	–	–	–
370.9 (15)	observed	< 0.1	–	–	–	–	–
373.5 (3)	–	–	–	–	–	–	0.0023 (3)
~ 380	observed	–	–	–	–	–	–
383.1 (1)	–	–	–	–	–	–	0.0040 (6)
401.81 (1)	observed	58 (6)	67 (4)	65.2 (65)	69.0 (30)	61.6 (28)	59.0 (20)
405.4 (1)	–	–	–	–	–	–	0.0023 (4)
436.9 (5)	–	–	–	–	–	–	0.0028 (5)
438.2 (6)	–	0.54 (10)	–	–	–	–	–
461.5 (4)	–	–	–	–	–	–	0.0015 (3)
489.3 (3)	–	–	–	–	–	–	0.0058 (8)
517.60 (6)	observed	0.44 (10)	0.48 (4)	0.22 (5)	–	0.43 (3)	0.40 (2)
538.2 (15)	observed	0.06 (3)	–	–	–	–	–
556.1 (4)	–	–	–	–	–	–	0.0005 (3)
564.1 (2)	observed	< 0.03	–	–	–	–	0.014 (3)
576.6 (10)	–	–	–	–	–	–	0.0008 (4)
608.30 (7)	observed	0.04 (2)	–	–	–	–	0.040 (10)
619.9 (3)	–	–	–	–	–	–	0.003 (1)
665.5 (10)	–	< 0.08 ^Δ	–	–	–	–	0.0008 (4)
671.9 (4)	observed	–	–	–	–	–	0.002 (1)
676.66 (7)	–	0.21 (3)	0.23 (2) ^Δ	0.06 (3)	–	0.16 (1)	0.16 (2)
708.1 (5)	–	–	–	–	–	–	0.003 (1)
732.7 (4)	–	–	–	–	–	–	0.0006 (3)
802.5 (4)	–	–	–	–	–	–	0.003 (1)
835.32 (22)	observed	–	–	–	–	–	0.015 (3)
877.2 (6)	–	–	–	–	–	–	0.003 (1)
891.1 (3)	observed	0.015 (7)	–	–	–	–	0.007 (2)
1055 (2)	observed	0.006 (3)	–	–	–	–	–
1073.7 (4)	–	–	–	–	–	–	0.003 (1)

* Quantified only in terms of percentage depopulation of a number of ill-defined nuclear levels of ²¹⁵Po – neither the gamma-ray energies nor this form of relative emission probability were adopted in the subsequent analyses.

† Quoted relative emission probabilities of 1967Da20 are based on P_γ(271.228 keV) of 1000, and have been adjusted to P_γ(271.228 keV) of 100.

‡ Quoted relative emission probabilities of 1968Br17 for ²¹⁹Rn decay are based on P_γ(271.228 keV) of 0.110 (15), and have been adjusted to P_γ(271.228 keV) of 110 (15).

¶ Uncertainties in the relative emission probabilities are not defined by 1970Da09, and have been derived from the quoted uncertainties of the absolute emission probabilities.

^Δ Evidence for the existence of a doublet.

Relative gamma-ray emission probabilities: Relative to P_γ(271.228 keV) of 100 %.

E _γ (keV)	P _γ							Recommended value
	1967Da20	1968Br17	1970Da09	1970Kr08	1976B113	1999Li05		
– 115.4 (5)*	–	0.030 (14)	–	–	–	–	–	–
γ _{3,1} 130.58 (1)	1.40 (14)	1.18 (23)	1.05 (25)	1.21 (10)	1.10 (11)	1.7 (2)	1.2 (1)	
– 221.5 (3)#	–	–) 0.28 (7)	–	–	–	–	
γ _{4,2} 224.04 (7)	–	–)	–	–	0.013 (2)	0.013 (2)	
γ _{1,0} 271.228 (10)	100	100	100.00	100.0	100	100.0 (20)	100 (2)	
γ _{2,0} 293.56 (4)	0.64 (6)	0.70 (14)	0.59 (15)	0.51 (27)	0.72 (5)	0.68 (4)	0.68 (3)	
γ _{12,5} 322 (1)	–	–	–	–	–	0.0008 (4)	0.0008 (4)	
– 324.9 (10)§	< 0.06	–	–	–	–	–	–	
γ _{8,3} 330.9 (4)	–	–	–	–	–	0.0090 (10)	0.0090 (10)	
– 337.7 (10)†	< 0.08	–	–	–	–	–	–	
– 370.9 (15)†	< 0.1	–	–	–	–	–	–	
γ _{11,4} 373.5 (3)	–	–	–	–	–	0.0023 (3)	0.0023 (3)	
– ~ 380§	–	–	–	–	–	–	–	
γ _{6,2} 383.1 (1)	–	–	–	–	–	0.0040 (6)	0.0040 (6)	
γ _{3,0} 401.81 (1)	58 (6)	61 (4)	65.2 (65)	69 (3)	58.4 (27)	59.0 (20)	61 (2)	
γ _{6,1} 405.4 (1)	–	–	–	–	–	0.0023 (4)	0.0023 (4)	
γ _{7,1} 436.9 (5)	–	–	–	–	–	0.0028 (5)	0.0028 (5)	
– 438.2 (6)‡	0.54 (10)	–	–	–	–	–	–	
γ _{8,1} 461.5 (4)	–	–	–	–	–	0.0015 (3)	0.0015 (3)	
γ _{11,3} 489.3 (3)	–	–	–	–	–	0.0058 (8)	0.0058 (8)	
γ _{4,0} 517.60 (6)	0.44 (10)	0.44 (4)	0.22 (5)	–	0.41 (3)	0.40 (2)	0.39 (3)	
– 538.2 (15)†	0.06 (3)	–	–	–	–	–	–	
γ _{13,4} 556.1 (4)	–	–	–	–	–	0.0005 (3)	0.0005 (3)	
γ _{9,1} 564.1 (2)	< 0.03	–	–	–	–	0.014 (3)	0.014 (3)	
γ _{14,4} 576.6 (10)	–	–	–	–	–	0.0008 (4)	0.0008 (4)	
γ _{5,0} 608.30 (7)	0.04 (2)	–	–	–	–	0.040 (10)	0.040 (9)	
γ _{11,1} 619.9 (3)	–	–	–	–	–	0.003 (1)	0.003 (1)	
γ _{-1,1} 665.5 (10)¶	< 0.08 ^Δ	–	–	–	–	0.0008 (4)	0.0008 (4)	
γ _{13,3} 671.9 (4)	–	–	–	–	–	0.002 (1)	0.002 (1)	
γ _{6,0} 676.66 (7)	0.21 (3)	0.21 (2)	0.06 (3)	–	0.15 (1)	0.16 (2)	0.16 (2)	
γ _{7,0} 708.1 (5)	–	–	–	–	–	0.003 (1)	0.003 (1)	
γ _{8,0} 732.7 (4)	–	–	–	–	–	0.0006 (3)	0.0006 (3)	
γ _{13,1} 802.5 (4)	–	–	–	–	–	0.003 (1)	0.003 (1)	
γ _{9,0} 835.32 (22)	–	–	–	–	–	0.015 (3)	0.015 (3)	
γ _{10,0} 877.2 (6)	–	–	–	–	–	0.003 (1)	0.003 (1)	
γ _{11,0} 891.1 (3)	0.015 (7)	–	–	–	–	0.007 (2)	0.008 (2)	
– 1055 (2)†	0.006 (3)	–	–	–	–	–	–	
γ _{13,0} 1073.7 (4)	–	–	–	–	–	0.003 (1)	0.003 (1)	

* Only observed by 1968Br17, and of doubtful origin – discarded.

Determined from the measurements of 1970Da09, but identified as a gamma-ray emission within the alpha-decay mode of ²²³Ra – discarded.

§ Only observed in a qualitative manner by 1965Va10 and 1967Da20, and of doubtful origin – discarded.

† Derived only from the measurements of 1967Da20, and of doubtful origin – discarded.

‡ Determined from the measurements of 1967Da20, but identified as a gamma-ray emission within the alpha-decay mode of ²¹⁵Po – discarded.

¶ Derived only from the measurements of 1967Da20 and 1999Li05, and of doubtful origin – unplaced within the proposed ²¹⁹Rn decay scheme.

^Δ Evidence for the existence of a doublet.

Multipolarity and Internal Conversion Coefficients

The decay scheme specified by 2001Br31 has been used to define the multipolarity of specific gamma transitions on the basis of the known spins and parities of the nuclear levels. Thus, the 224.04-, 401.81 and 1073.7-keV gamma-ray emissions are adjudged to be E2 transitions. Multipolarity mixing ratios for the 130.58- and 271.228-keV gamma transitions of 0.60 (6) and 4.0 (4), respectively, were derived from the K/L and L sub-shell conversion-electron intensities determined by Davidson and Connor (1970Da09), while the 293.56-, 517.60- and 556.1-keV gamma-ray emissions were arbitrarily assigned mixing ratios of 1.0 (2) (i.e. 50%M1 + 50%E2). Recommended internal conversion coefficients have been determined from the frozen orbital approximation of Kibédi *et al.* (2008Ki07), based on the theoretical model of Band *et al.* (2002Ba85, 2002Ra45).

Gamma-ray emissions: multipolarity and theoretical internal conversion coefficients (frozen orbital approximation).

E_γ (keV)	P_γ^{abs} × 100	Multipolarity	α_K	α_L	α_{M+}	α_{total}
130.58 (1)	0.133 (11)	73.5%M1 + 26.5%E2 δ = 0.60 (6)	3.19 (16)	0.94 (4)	0.31	4.44 (13)
224.04 (7)	0.001 4 (2)	(E2)	0.129 6 (19)	0.140 7 (20)	0.048 7	0.319 (5)
271.228 (10)	11.07 (22)	(6%M1 + 94%E2) δ = 4.0 (4)	0.111 (6)	0.066 8 (11)	0.023 2	0.201 (7)
293.56 (4)	0.075 (3)	(50%M1 + 50%E2) δ = 1.0 (2)	0.25 (4)	0.062 (4)	0.028	0.34 (5)
322 (1)	0.000 09 (5)	–	–	–	–	–
330.9 (4)	0.001 00 (11)	–	–	–	–	–
373.5 (3)	0.000 25 (3)	–	–	–	–	–
383.1 (1)	0.000 44 (7)	–	–	–	–	–
401.81 (1)	6.75 (22)	E2	0.0351 (5)	0.015 28 (22)	0.005 12	0.055 5 (8)
405.4 (1)	0.000 25 (4)	–	–	–	–	–
436.9 (5)	0.000 31 (6)	–	–	–	–	–
461.5 (4)	0.000 17 (3)	–	–	–	–	–
489.3 (3)	0.000 64 (9)	–	–	–	–	–
517.60 (6)	0.043 (3)	(50%M1 + 50%E2) δ = 1.0 (2)	0.058 (9)	0.011 5 (11)	0.003 5	0.073 (10)
556.1 (4)	0.000 06 (4)	(50%M1 + 50%E2) δ = 1.0 (2)	0.048 (7)	0.009 5 (9)	0.003 5	0.061 (8)
564.1 (2)	0.001 5 (3)	–	–	–	–	–
576.6 (10)	0.000 09 (5)	–	–	–	–	–
608.30 (7)	0.004 4 (10)	(M1 + E2)	–	–	–	–
619.9 (3)	0.000 33 (11)	–	–	–	–	–
665.5 (10)	0.000 09 (5)	–	–	–	–	–
671.9 (4)	0.000 22 (11)	M1 + E2	–	–	–	–
676.66 (7)	0.018 (2)	–	–	–	–	–
708.1 (5)	0.000 33 (11)	–	–	–	–	–
732.7 (4)	0.000 07 (4)	–	–	–	–	–
802.5 (4)	0.000 33 (11)	M1 + E2	–	–	–	–
835.32 (22)	0.001 7 (3)	–	–	–	–	–
877.2 (6)	0.000 33 (11)	–	–	–	–	–
891.1 (3)	0.000 9 (2)	–	–	–	–	–
1073.7 (4)	0.000 33 (11)	E2	0.005 10 (8)	0.001 002 (14)	0.000 308	0.006 41 (9)

Atomic Data

The x-ray data have been calculated using the evaluated gamma-ray data, and the atomic data from 1996Sc06, 1998ScZM and 1999ScZX. Both the x-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.01, 28 January 2003). This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

K and L X-ray emission probabilities per 100 disintegrations of ²¹⁹Rn.

			Energy (keV)	Photons per 100 disint.
XL	(Po)		9.658 – 16.213	1.01 (5)
	XL ₁	(Po)	9.658	0.0229 (9)
	XL _α	(Po)	11.016 – 11.130	0.420 (15)
	XL _η	(Po)	12.085	0.0095 (4)
	XL _β	(Po)	12.823 – 13.778	0.475 (13)
	XL _γ	(Po)	15.742 – 16.213	0.098 (3)
XK _α	XK _{α2}	(Po)	76.864	0.540 (24)
	XK _{α1}	(Po)	79.293	0.90 (4)
XK' _{β1}	XK _{β3}	(Po)	89.256)
	XK _{β1}	(Po)	89.807) 0.309 (15)
	XK _{β5}	(Po)	90.363)
XK' _{β2}	XK _{β2}	(Po)	92.263)
	XK _{β4}	(Po)	92.618) 0.096 (5)
	XKO _{2,3}	(Po)	92.983)

Electron energies were determined from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

Data Consistency

A Q_α-value of 6946.1 (3) keV has been adopted from the atomic mass evaluation of Audi *et al.* (2003Au03) while in the course of formulating the decay scheme of ²¹⁹Rn. This value has subsequently been compared with the Q-value calculated by summing the contributions of the individual emissions to the ²¹⁹Rn alpha-decay process (i.e. α, electron, γ, etc.):

$$\text{calculated Q-value} = \sum (E_i \times P_i) = 6945 (70) \text{ keV}$$

Percentage deviation from the Q-value of Audi *et al.* is (0.0 ± 1.0) %, which supports the derivation of a highly consistent decay scheme with a rather significant variant.

References

- 1961Ro14 H. RODENBUSCH, G. HERRMANN, Ein Verfahren zur Bestimmung von Halbwertszeiten kurzlebiger gasförmiger Radioisotope. Die Halbwertszeiten des Thorons (²²⁰Rn) und Actinons (²¹⁹Rn), Z. Naturforsch. 16a (1961) 577-582. [Half-life]
- 1962Wa18 R.J. WALLEN, V. NEDOVESOV, G. BASTIN-SCOFFIER, Spectrographie α de ²²³Ra (AcX) et de ses Dérivés, Nucl. Phys. 35 (1962) 232-252. [E_α, P_α]
- 1965Va10 K. VALLI, J. AALTONEN, G. GRAEFFE, M. NURMIA, An Alpha-Gamma and Alpha-Conversion Electron Coincidence Study of Rn-219, Po-215, and Bi-211, Ann. Acad. Sci. Fenn., Ser. A VI, No. 184 (1965). [E_γ, P_γ]

- 1966Hu20 J.B. HURSH, Thoron Half-life, *J. Inorg. Nucl. Chem.* 28 (1966) 2771-2776. [Half-life]
- 1967Da20 J. DALMASSO, H. MARIA, Sur le Rayonnement γ du Radon 219, du Polonium 215 et du Plomb 211, *C. R. Acad. Sci. Paris* 265B (1967) 822-825. [E_γ , P_γ]
- 1968Br17 C. BRIANÇON, CHIN FAN LEANG, R. WALEN, Étude du Spectre γ Émis par le Radium-223 et ses Dérivés, *C. R. Acad. Sci. Paris* 266B (1968) 1533-1536. [E_γ , P_γ]
- 1970Da09 W.F. DAVIDSON, R.D. CONNOR, The Decay of ²²³Ra and its Daughter Products (II). The Decay of ²¹⁹Rn and ²¹⁵Po, *Nucl. Phys.* A149 (1970) 385-391.
[E_γ , P_γ , P_{ce} , K/L and L sub-shell ratios, ICC]
- 1971Gr07 B. GRENNBERG, A. RYTZ, Absolute Measurements of α -ray Energies, *Metrologia* 7 (1971) 65-77 [E_α]
- 1970Kr08 K. KRIEN, M.J. CANTY, P. HERZOG, Determination of Spins in the Decay of ²¹⁹Rn \rightarrow ²¹⁵Po by Alpha-Gamma Linear Polarization Measurements, *Nucl. Phys.* A157 (1970) 456-470.
[E_γ , P_γ , P_{ce} , L sub-shell ratios, ICC]
- 1976Bl13 K. BLATON-ALBICKA, B. KOTLIŃSKA-FILIPEK, M. MATUL, K. STRYCZNIEWICZ, M. NOWICKI, E. RUCHOWSKA-LUKASIAK, Precision Gamma-ray Spectroscopy of the Decay of ²²³Ra and its Daughter Products, *Nukleonika* 21 (1976) 935-947. [E_γ , P_γ]
- 1977La19 F.P. LARKINS, Semiempirical Auger-electron Energies for Elements $10 \leq Z \leq 100$, *At. Data Nucl. Data Tables* 20 (1977) 311-387. [Auger-electron energies]
- 1991Ry01 A. RYTZ, Recommended Energy and Intensity Values of Alpha Particles from Radioactive Decay, *At. Data Nucl. Data Tables* 47 (1991) 205-239. [E_α , P_α]
- 1996Sc06 E. SCHÖNFELD, H. JANßEN, Evaluation of Atomic Shell Data, *Nucl. Instrum. Methods Phys. Res.* A369 (1996) 527-533. [X_K , X_L , Auger electrons]
- 1998Ak04 Y.A. AKOVALI, Review of Alpha-decay Data from Doubly-even Nuclei, *Nucl. Data Sheets* 84 (1998) 1-114. [alpha decay, r_0 parameters]
- 1998ScZM E. SCHÖNFELD, G. RODLOFF, Tables of the Energies of K-Auger Electrons for Elements with Atomic Numbers in the Range from $Z = 11$ to $Z = 100$, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999Li05 C.F. LIANG, P. PARIS, R.K. SHELINE, Configurations and Level Structure of ²¹⁵Po, *Phys. Rev.* C59 (1999) 648-654. [E_α , P_α , E_γ , P_γ]
- 1999ScZX E. SCHÖNFELD, G. RODLOFF, Energies and Relative Emission Probabilities of K X-rays for Elements with Atomic Numbers in the Range from $Z = 5$ to $Z = 100$, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2001Br31 E. BROWNE, Nuclear Data Sheets for $A = 215, 219, 223, 227, 231$, *Nucl. Data Sheets* 93 (2001) 763-1061. [Nuclear structure, level energies]
- 2002Ba85 I.M. BAND, M.B. TRZHASKOVSKAYA, C.W. NESTOR Jr., P.O. TIKKANEN, S. RAMAN, Dirac-Fock Internal Conversion Coefficients, *At. Data Nucl. Data Tables* 81 (2002) 1-334. [ICC]
- 2002Ra45 S. RAMAN, C.W. NESTOR Jr., A. ICHIHARA, M.B. TRZHASKOVSKAYA, How Good Are the Internal Conversion Coefficients Now? *Phys. Rev.* C66 (2002) 044312, 1-23. [ICC]
- 2003Au03 G. AUDI, A.H. WAPSTRA, C. THIBAULT, The AME2003 Atomic Mass Evaluation (II). Tables, Graphs and References, *Nucl. Phys.* A729 (2003) 337-676. [Q-value]
- 2008Ki07 T. KIBÉDI, T.W. BURROWS, M.B. TRZHASKOVSKAYA, P.M. DAVIDSON, C.W. NESTOR Jr., Evaluation of Theoretical Conversion Coefficients Using BrIcc, *Nucl. Instrum. Methods Phys. Res.* A589 (2008) 202-229. [ICC]

**²²⁰Rn – Comments on evaluation of decay data
by A. L. Nichols**

Evaluated: July/August 2001
Re-evaluated: January 2004

Evaluation Procedures

Limitation of Relative Statistical Weight Method (LWM) was applied to average numbers throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

A simple decay scheme has been derived from the gamma γ -ray studies of 1972DaZA, 1977Ku15, and 1984Ge07. The single 549.76 keV gamma ray had a weighted mean emission probability of 0.115(15)% (0.00115(15)), and this value and theoretical internal conversion coefficients were used to calculate the absolute emission probabilities of the 5748.46 and 6288.22 keV alpha particles to the 549.76 keV and ground states of ²¹⁶Po, respectively. Alpha-particle studies are required to confirm the validity of the proposed decay scheme.

Nuclear Data

²²⁸Th decay chain is important in quantifying the environmental impact of the decay of naturally occurring ²³²Th.

Half-life

The recommended half-life is the weighted mean of measurements by 1955Sc81, 1961Ro14, 1963Gi17 and 1966Hu20. Further studies are merited to confirm the most recent studies of 1963Gi17 and 1966Hu20.

Reference	Half-life (s)
1955Sc81	51.5(10)*
1961Ro14	56.6(8)
	56.3(2)
1963Gi17	55.3(3)
1966Hu20	55.61(4)#
Recommended Value	55.8(3)

* Defined as outlier.

Uncertainty adjusted to ± 0.16 to reduce weighting below 0.5.

Gamma Ray

Energy

The single gamma-ray energy was based on the nuclear level energy of 549.76(4) keV from 1997Ar04.

Emission Probability

The absolute emission probability of the 549.76(4) keV gamma ray was determined from measurements by 1972DaZA, 1977Ku15 and 1984Ge07. A weighted mean value of 0.115(15)% (0.00115(15)) was derived through LWEIGHT.

Published Gamma-ray Emission Probabilities

E _g (keV)	P _g			
	1956Ma28 [†]	1972DaZA [‡]	1977Ku15 [¶]	1984Ge07 [#]
549.76(4)	0.025	0.29(9)	0.0950(80)	0.43(4)

[†] Defined as accurate to within a factor of 2; rejected from evaluation.

[‡] Relative to P_γ(2614.511 keV) of ²⁰⁸Tl.

[¶] Absolute value in measurements that include P_γ(240.986 keV) of 3.95% for ²²⁴Ra.

[#] Relative to P_γ(583.19 keV) of ²⁰⁸Tl.

Absolute Gamma-ray Emission Probabilities per 100 Disintegrations of ²²⁰Rn

E _g (keV)	P _g ^{abs}			
	1972DaZA [†]	1977Ku15 [†]	1984Ge07 [†]	Recommended Value [*]
549.76(4)	0.104(32)	0.0991(83)	0.130(3)	0.115(15)

[†] Data adjusted on the basis of the footnotes given above.

^{*} Weighted mean value adopted.

Multipolarity and Internal Conversion Coefficients

The decay scheme specified by 1997Ar04 has been used to define the multipolarity of the gamma transition on the basis of the known spins and parities of the two nuclear levels. Theoretical internal conversion coefficients were interpolated from the tabulations of 1978Ro22.

Alpha-particle EmissionsEnergies

Alpha-particle energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies of 1997Ar04 and the Q-value (1995Au04) were used to determine the energies and uncertainties of the alpha-particle transitions to the various levels, while allowing for the significant recoil components.

Emission Probabilities

Both alpha-particle emission probabilities were derived from the weighted mean emission probability of the single gamma transition and theoretical internal conversion coefficients.

Alpha-particle Emission Probabilities per 100 Disintegrations of ²²⁰Rn

E _a (keV)	P _a		
	1962Wa28	1977Ku15 [#]	Recommended Values [*]
5748.46(14)	0.07(2)	0.097(8)	0.118(15)
6288.22(10)	~ 100	99.9	99.882(15)

[#] Data were deduced from gamma-ray studies.

^{*} Recommended emission probabilities derived from evaluated gamma-ray emission probability and theoretical internal conversion coefficients.

Atomic Data

The x-ray data have been calculated using the evaluated gamma-ray data, and the atomic data from 1996Sc06, 1998ScZM and 1999ScZX.

References

- 1955Sc81 - H. Schmied, R. W. Fink and B. L. Robinson, The Half-Life of Emanation-220, J. Inorg. Nucl. Chem. 1(1955)342. [Half-life]
- 1956Ma28 - L. Madansky and F. Rasetti, Decay of Rn²²⁰ and Rn²²², Phys. Rev. 102(1956)464. [P_γ]
- 1961Ro14 - H. Rodenbusch and G. Herrmann, Ein Verfahren zur Bestimmung von Halbwertszeiten kurzlebiger gasförmiger Radioisotope. Die Halbwertszeiten des Thorons (²²⁰Rn) und Actinons (²¹⁹Rn), Z. Naturforsch. 16a(1961)577. [Half-life]
- 1962Wa28 - R. J. Walen, Spectrographie α du Radium 224 et de ses Dérivés, C. R. Acad. Sci. Paris 255(1962)1604. [P_α]
- 1963Gi17 - J. E. Gindler and D. W. Engelkemeir, Half-Life Determination of ²²⁰Rn, Radiochim. Acta 2(1963)58. [Half-life]
- 1966Hu20 - J. B. Hursh, Thoron Half-Life, J. Inorg. Nucl. Chem. 28(1966)2771. [Half-life]
- 1972DaZA - J. Dalmasso, Recherches sur le Rayonnement Gamma de Quelques Radioelements Naturels Appartenant a la Famille du Thorium, PhD thesis, University of Nice (1972); J. Dalmasso, H. Maria and C. Ythier, Étude du Rayonnement γ du Thorium 228 et de ses Dérivés, et plus Particulièrement du Thallium 208 (Th C"), C. R. Acad. Sci. Paris 277B(1973)467. [P_γ]
- 1977Ku15 - W. Kurcewicz, N. Kaffrell, N. Trautmann, A. Plochocki, J. Zylicz, M. Matul and K. Stryczniewicz, Collective States Fed by Weak α-transitions in the ²³²U Chain, Nucl. Phys. A289(1977)1. [P_γ]
- 1978Ro22 - F. Rösel, H. M. Fries, K. Alder and H. C. Pauli, Internal Conversion Coefficients for all Atomic Shells, ICC Values for Z = 68-104, At. Data Nucl. Data Tables 21(1978)291-514. [ICC]
- 1984Ge07 - R. J. Gehrke, V. J. Novick and J. D. Baker, γ-ray Emission Probabilities for the ²³²U Decay Chain, Int. J. Appl. Radiat. Isot. 35(1984)581. [P_γ]
- 1995Au04 - G. Audi and A. H. Wapstra, The 1995 Update to the Atomic Mass Evaluation, Nucl. Phys. A595(1995)409. [Q value]
- 1996Sc06 - E. Schönfeld and H. Janßen, Evaluation of Atomic Shell Data, Nucl. Instrum. Meth. Phys. Res. A369(1996)527. [X_K, X_L, Auger electrons]
- 1997Ar04 - A. Artna-Cohen, Nuclear Data Sheets for A = 216, 220, Nucl. Data Sheets 80(1997)157. [Nuclear structure, energies]
- 1998ScZM - E. Schönfeld and G. Rodloff, Tables of the Energies of K-Auger Electrons for Elements with Atomic Numbers in the Range from Z = 11 to Z = 100, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX - E. Schönfeld and G. Rodloff, Energies and Relative Emission Probabilities of K X-rays for Elements with Atomic Numbers in the Range from Z = 5 to Z = 100, PTB Report PTB-6.11-1999-1, February 1999. [X_K]

²²¹Fr - Comments on evaluation of decay data by Huang Xiaolong and Wang Baosong

This evaluation was completed in 2007. Literature available by December 2007 was included.

1 Decay Scheme

²²¹Fr disintegrates 99.9952 (15) % by α emission to levels in ²¹⁷At and 0.0048 (15) % by β^- emission to levels in ²²¹Ra. ²²¹Fr ground state has $J^\pi=5/2^-$ (2003Ak06).

The α decay scheme of ²²¹Fr was built based on the measurement described in 1995Sh01, 1999Gr33 and 2002Gr36. A study of 1997Ch53 showed the existence of a possible weak β^- decay of $(4.8 \pm 1.5) 10^{-3}$ % to ²²¹Ra. The β^- decay scheme of ²²¹Fr has not been studied.

The recommended $Q(\alpha)$ value of 6457.8 (14) keV in Audi(2003Au03) agrees with the $Q(\alpha)$ value of 6461.5 (25) keV, calculated by the evaluator (using program RADLST) from average radiation energies. This agreement supports the completeness and correctness of the decay scheme.

2 Nuclear Data

The Q value is from the 2003Au03 evaluation.

Level energies, have been obtained from a least-squares fit to γ -ray energies (GTOL computer code). Spin and parities are from 2003Ak06.

The measured and our evaluated ²²¹Fr half-life values are listed in Table 1. Notice that uncertainties in tables referred to the least significant digits.

Table 1 - Measured half-life values of ²²¹Fr and recommended value, in minutes

T_{1/2} (min)	References	measurement method
5	1947En03	
4.8	1950Ha52	Alpha pulse analyzer
4.9 (2)	1967LoZZ	
4.79 (2)	2007Je07	From Si sample. Metallic samples(Au,W) give shorter value
4.9 (2)	2003Ak06	NDS, from 1967LoZZ
4.85 (6)		Unweighted mean of 1967LoZZ and 2007Je07
4.791 (20)		Weighted mean of 1967LoZZ and 2007Je07, $\chi^2=0.3$
4.79 (2)	2007	Recommended value, from 2007Je07

2007Je07 measured the half-life at room temperature in different materials and obtained an improved value. As the weighted mean of 4.9 (2) min (1967LoZZ) and 4.79 (2) min (2007Je07) is very close to this most precise measurement, the value of 2007Je07 is recommended here.

2.1 γ Transitions

The γ -ray transition probabilities were calculated using the γ -ray emission intensities and the relevant internal conversion coefficients.

Multipolarities and mixing ratios of γ -ray transitions are from 1968Le07 and 1995Sh01. Multipolarities in square brackets are from level scheme (they are not measured).

The internal conversion coefficients (ICC) and their associated uncertainties for γ -ray transitions have been obtained using the BrIcc computer program, which uses the "Frozen Orbital" approximation (2002Ba85).

Experimental and theoretical conversion coefficients are compared in Table 2.

Table 2 - Comparison of theoretical and measured conversion coefficients

E_{β}/keV	Multipolarity	a(theory)	a(exp.)	
			1995Sh01	1999Gr33
53.81	M1	$\alpha_T=14.17, \alpha_L=10.79$ (16)	$\alpha_L=8$ (4)	
96.3	M1+E2	$\alpha_T=5.6, \alpha_L=4.1$ (18)	$\alpha_L>2.5$	$\alpha_T=25$ (15)
100.25	M1	$\alpha_T=11.97, \alpha_L=1.758$ (25)	$\alpha_L=1.2$ (6)	
117.82	M1	$\alpha_T=7.58$		$\alpha_T=13.5$ (86)
150.21	M1	$\alpha_T=3.8, \alpha_K=3.08$ (5)	$\alpha_K=2.6$ (5)	$\alpha_T=3.5$ (9)
171.83	E2	$\alpha_T=0.863, \alpha_K=0.226$ (4)	$\alpha_K=0.3$ (1)	$\alpha_T=0.84$ (2)
218.12	E2	$\alpha_T=0.367, \alpha_K=0.1375$ (20)	$\alpha_K=0.14$	
324.10	M1	$\alpha_T=0.446, \alpha_K=0.362$ (5)	$\alpha_K=0.4$ (2)	
359.86	M1	$\alpha_T=0.335, \alpha_K=0.272$ (4)	$\alpha_K=0.4$ (1)	
382.34	M1	$\alpha_T=0.284, \alpha_K=0.231$ (4)	$\alpha_K=0.25$ (10)	
410.64	E2	$\alpha_T=0.0548, \alpha_K=0.0344$ (5)	$\alpha_K=0.03$ (1)	

2.2 a Transitions

Measured energies of alpha particles are listed in table 3. Our recommended values are from 1968Le07 and 2002Gr36.

Table 3 - Measured and recommended values of α -particle energies (in keV) from ²²¹Fr α decay

1967Dz02	1968Le07 ^a	2002Gr36	Recommended
		5500 (40)	5500 (40)
		5530 (25)	5530 (25)
	5689 (3)		5689 (3)
	5697 (4)		5697 (4)
	5776 (3)		5776 (3)
	5783 (4)		5783 (4)
	5813 (3)		5813 (3)
5930 (7)	5925 (3)		5925 (3)
5940 (6)	5938.9 (20)		5938.9 (20)
5966 (6)	5965.9 (25)		5965.9 (25)
5980 (6)	5979.9 (20)		5979.9 (20)
6075 (5)	6075.9 (20)		6075.9 (20)
6125 (5)	6126.3 (15)		6126.3 (15)
6241 (6)	6243.0 (20)		6243.0 (20)
6338 (5)	6341.0 (13)		6341.0 (13)

^a: Values were adjusted based on the calibration recommendation of 1991Ry01.

Experimental and recommended α -particle emission probabilities are listed in Table 4. Our recommended alpha particle emission probabilities are average values of measured α -particle intensities with those deduced from γ -transition probability balance at each decay scheme level.

Table 4 - Experimental, recommended α -particle emission probabilities from ²²¹Fr decay

$E_a(\text{keV})$	$P_a(\%)$				
	1967Dz02	1968Le07	2002Gr36	Deduced from Pg	Recommended [†]
5500 (40)			3.3 (9) E-4	0.000038 (10)	0.000038 (10)
5530 (25)			9.0 (20) E-4	0.00010 (2)	0.00010 (2)
5689 (3)		0.002 (1)		0.0026 (5)	0.0025 (5)
5697 (4)		~0.001		~0.004	~0.003
5776 (3)		0.06 (1)		0.065 (4)	0.064 (4)
5783 (4)		0.005(2)		0.0029 (6)	0.0031 (6)
5813 (3)		0.004 (2)		0.006 (1)	0.006 (1)
5925 (3)	0.05 (1)	0.03 (1)		0.0280 (16)	0.0285 (24)
5938.9 (20)	0.13 (1)	0.17 (3)		0.127 (3)	0.128 (3)
5965.9 (25)	0.12 (1)	0.08 (1)		0.053 (4)	0.064 (16)
5979.9 (20)	0.46 (5)	0.49 (3)		0.27 (3)	0.39 (7)
6075.9 (20)	0.13 (2)	0.15 (3)		0.30 (6)	0.15 (3)
6126.3 (15)	14.5 (7)	15.1 (2)		15.3 (3)	15.1 (2)
6243.0 (20)	1.35 (7)	1.34 (10)		0.9 (5)	1.34 (7)
6341.0 (13)	83.2 (20)	83.4 (8)		82.9 (5)	82.8 (2)*

[†] Weighted average of values from the first four columns, normalized to a total of 100 %.

* Value reduced by a covariance effect introduced by the normalization to 100 %.

3. Atomic data

Atomic fluorescence yields ($\omega_K, \omega_L, \omega_M, \eta_{KL}$ and η_{LM}) are from Schönfeld (1996Sc06).

The X-ray and Auger electron emission probabilities have been deduced from γ -ray and conversion electron data by using the computer code RADLST. The deduced K X-ray emission probabilities $P_{KX} = 2.77 (19) \%$ agree with the measured value of 2.23 (20) % in 1995Sh01, thus confirming the completeness of the decay scheme.

4. Electron Emissions

The conversion electron emission probabilities have been deduced from γ -ray transition data using theoretical internal conversion coefficients.

5. Photon Emissions

5.1 γ -ray energy values

The experimental and our recommended γ -ray energies from ²²¹Fr α decay are listed in table 5. Unless otherwise specified the later are averages (or weighted averages) from values given in 1968Le07, 1994Ar23, 1995Sh01, and 1999Gr33. γ -rays of 809.3 and 891.9 keV reported only in 2002Gr36 have also been included here. Several γ -rays observed in 1995Bu17 and 1994Ar23 were interpreted as sum peaks in 1999Gr33. Values from 1995Bu17 have not been included in this averaging because this reference seems to be an earlier publication of 1999Gr33 (notice that only these two references reported the 201.4 - and 208.3-keV γ -rays).

Table 5 - Measured and recommended values of γ -ray energies for ²²¹Fr α decay.

1968Le07	1994Ar23	1995Bu17	1995Sh01	1999Gr33	2002Gr36	LWM	Recommended
		53.54 (18)	53.8 (1)	53.81 (3)		53.80 (28)	53.81 (3)
		68.11 (15)					
		96.12 (18)	96.3 (3)	96.3 (3)		96.20 (14)	96.3 (3)
99.5	100.63 (2)	99.52 (6)	100.2 (1)	100.25 (2)		100.40 (24)	100.25 (2)
118.2 (2)	117.67 (5)	118.18 (9)	117.8 (2)	117.82 (3)		117.80 (9)	117.82 (3)
150.0 (2)	150.43 (5)	150.04 (4)	150.0 (1)	150.21 (3)		150.20 (8)	150.21 (3)
171.3	172.05 (5)	171.68 (4)	171.6 (1)	171.83 (3)		171.80 (8)	171.83 (3)
		201.44 (50)		201.4 (6)		201.4 (4)	201.4 (6) ^a
		208.3 (5)		208.3 (6)		208.3 (4)	208.3 (6)
217.99 (4)	218.30 (2)	218.14 (3)	218.0 (1)	218.12 (2)		218.20 (11)	218.12 (2)
	250.7 (2)						
	253.15 (15)						
		263.39 (14)					
	271.91 (5)						
282.8	282.25 (5)	282.36 (15)	281.9 (3)	282.12 (9)		282.20 (4)	282.12 (9)
		297.11 (40)					
		299.59 (14)					
	310.20 (5)	310.14 (16)					
		314.11 (17)					
324.1	323.99	323.99 (6)	324.0 (2)	324.10 (6)		324.00 (4)	324.10 (6)
359.1	359.90 (2)	359.92 (6)	359.0 (2)	359.86 (4)		359.90 (6)	359.86 (4)
	368.17 (2)	368.18 (10)					
381.8	382.36 (2)	381.81 (4)	381.1 (2)	382.34 (4)		382.20 (15)	382.34 (4)
409.1	410.73 (2)	409.93 (7)	410.4 (2)	410.64 (5)		410.60 (16)	410.64 (5)
	435.68 (10)		437.8 (5)	437.00 (5)		436.4 (6)	437.00 (5)
		445.07 (20)	446.3 (8)	446.30 (8)		445.7 (6)	446.30 (8)
		469.6 (2)	469.0 (5)	468.3 (7)		469.40 (18)	468.3 (7)
			496.2 (10)				
	538.02 (10)	537.0 (2)	537.5 (8)	537.8 (8)		537.5 (5)	537.8 (8)
			562.3 (12)				562.3 (12)
		568.5 (3)	568.4 (10)	568.5 (3)		568.50 (21)	568.5 (3)
	577.76 (6)	576.9 (4)	577.0 (8)	576.9 (4)		577.70 (6)	576.9 (4)
				652 (2)			652 (2)
				658 (2)			658 (2) ^a
				665 (2)			665 (2)
					809.3 (2)		809.3 (2)
					891.9 (3)		891.9 (3)

^a: not placed in level scheme.

5.2 Relative γ -ray emission probabilities

Measured relative γ -ray intensities from ²²¹Fr are listed together with our recommended values in Table 6. Several γ -rays observed in 1995Bu17 and 1994Ar23 were interpreted as sum peaks in 1999Gr33. Thus their relative intensities may not be accurate so they have not been recommended here.

Results in 1995Sh01 are in agreement with those in 1999Gr33 within their experimental uncertainties, but they are not as complete and accurate. However, their decay scheme differs only by some weak transitions. For example, 1995Sh01 did not observe the 652 -0, 578-368 γ -ray transitions, thus it did not propose the 652 keV level. 1999Gr33 is the most precise measurement among the available experimental data. Unless otherwise specified, the present recommended values are weighted averages (LWM) from values given in 1999Gr33, 1995Sh05, 1994Ar23, 1968Le07, and two γ -rays from 2002Gr36.

Table 6 - Measured and recommended relative γ -ray emission probabilities for ²²¹Fr

E_{γ} (keV)	1968Le07	1994Ar23	1995Sh01	1999Gr33	2002Gr36	Recommended ^{&}
53.81 (3)			0.15 (4)	0.116 (27)		0.127 (22)
96.3 (3)			<0.09	0.063 (27)		0.058 (23)
100.25 (2)	0.95 (34)	1.47 (9)	1.33 (18)	0.89 (27)		1.37 (11)
117.82 (3)	0.34 (17)	0.328 (17)	0.044 (18)	0.20 (12)		0.19 (14)
150.21 (3)	0.69 (26)	0.362 (17)	0.53 (9)	0.420 (18)		0.393 (21)
171.83 (3)	0.52 (26)	0.517 (17)	0.58 (11)	0.680 (18)		0.60 (8)
201.4 (6) ^a				0.0098 (9)		0.0098 (9) [†]
208.3 (6)				0.045 (9)		0.045 (9) [†]
218.12 (2)	100	100	100	100		100
282.12 (9)	0.086	0.056 (9)	0.071 (27)	0.063 (9)		0.060 (6)
324.10 (6)	0.17 (9)	0.138 (9)	0.106 (27)	0.170 (9)		0.152 (10)
359.86 (4)	0.34 (17)	0.319 (17)	0.32 (9)	0.358 (18)		0.337 (12)
382.34 (4)	0.34 (17)	0.302 (17)	0.27 (9)	0.295 (18)		0.298 (12)
410.64 (5)	1.21 (34)	1.034 (86)	0.97 (18)	1.055 (18)		1.054 (17)
437.00 (5)		0.034 (6)	~0.009	0.0083 (9)		0.0083 (9) [†]
446.30 (8)			~0.009	0.0152 (36)		0.0152 (36) [†]
468.3 (7)			0.018 (9)	0.0152 (27)		0.0154 (26)
537.8 (8)		0.034 (10)	0.018 (9)	0.0447 (45)		0.039 (7)
562.3 (12)			~0.044			~0.044
568.5 (3)			~0.009	0.0107 (36)		0.0107 (36) [†]
576.9 (4)		0.041 (6)	0.035 (9)	0.0259 (36)		0.030 (5)
652 (2)				~0.00358		~0.00358 [†]
658 (2) ^a				~0.00626		~0.00626 [†]
665 (2)				~0.00805		~0.00805 [†]
809.3 (2)					9.0E-4 (20)	9.0E-4 (20) [*]
891.9 (3)					3.3E-4 (9)	3.3E-4 (9) [*]

[&] Deduced using the LWM statistical method, unless otherwise specified.

^a not placed in level scheme.

[†] From 1999Gr33

^{*} Reported only in 2002Gr36

5.3 Absolute g-ray emission probabilities

Measurements of the absolute γ -ray emission probability of the 218.12 keV transition from ²²¹Fr α decay are listed in Table 7.

Values in 1968Le07, 1986He06 and 1995Sh01 are the only absolute independent measurements. Among these absolute measurements, the one given in 1986He06 is the most precise.

1986He06 measured the γ -ray emission probability in equilibrium with ²²⁹Th. ²²⁹Th α -decay emits a 218.15-keV γ -ray, therefore this contribution should be subtracted.

1987He28 and 2000Ga52 measured γ -ray emission probabilities from the α -decay of ²²⁹Th to ²²⁵Ra relatively to $I_\gamma = 4.3$ for the 193.5-keV transition. They obtained 0.18 (2) and 0.134 (20) for the 218.15-keV γ -ray, respectively.

The weighted average of these values is 0.146 (20) relative to $I_\gamma(193.5) = 4.3$. Using a conversion factor of 1.026 (14) as given by 1987He28, the absolute value is: $0.146 (20) \times 1.026 (14) = 0.150 (20) \%$.

Thus, the corrected absolute γ -ray emission probability of the 218.15 -keV γ -ray from ²²¹Fr α decay is $11.57 (15) - 0.150 (20) = 11.42 (15) \%$, which is our recommended value.

Taking into account the β - branching ratio (see §6), the normalization factor between relative and absolute emission probabilities is $N = 11.42 (15) / 0.999952 (15) = 0.1142 (15)$.

Table 7 - Measured and recommended absolute γ -ray emission probability of 218.12 keV for ²²¹Fr

P_γ (218.12 keV)	References	Experimental value and method
12.5 (4)	1968Le07	
13.44 (27)	1981Di14	Ge(Li)
11.57 (15)	1986He06	Ge(Li), Au-Si surface barrier, in equilibrium with ²²⁹ Th
11.3 (10)	1995Sh01	Ge(Li), α - γ -ce coincidence
11.18 (15)	1999Gr33	Ge(Li), α γ coincidence, using $I_\alpha(6126) = 15.1 (2) \%$
11.42 (15)	Recommended	Evaluated value, from 1986He06

^a: value corrected using the evaluation of 1990Ak05.

The recommended absolute γ -ray emission probabilities are the recommended relative values shown in table 6 multiplied by 0.1142 (15), as given in table 8.

Table 8 - Absolute γ -ray emission probabilities from ²²¹Fr α decay.

E_γ (keV)	P_γ (%)		E_γ (keV)	P_γ (%)
53.8	0.0145 (25)		446.3	0.0017 (4)
96.3	0.007 (3)		468.3	0.0018 (3)
100.2	0.156 (13)		537.8	0.0045 (8)
117.8	0.022 (16)		562.3	0.005 (5)
150.2	0.0449 (25)		568.5	0.0012 (4)
171.8	0.069 (9)		576.9	0.0030 (6)
201.4	0.0011 (1)		652	0.0004 (4)
208.3	0.0051 (10)		658	0.0007 (7)
218.1	11.42 (15)		665	0.0009 (9)
282.12	0.0069 (7)		809.3	0.00010 (2)
324.1	0.0174 (12)		891.9	0.000038 (10)
359.9	0.0385 (15)			
382.3	0.0340 (14)			
410.6	0.1204 (25)			
437	0.0010 (1)			

6. b- Branching Ratio

Measured and recommended branching ratios for ²²¹Fr β⁻ decay are listed in Table 9. Our recommended β⁻ decay branching ratio from 1997Ch53 is I_β = 0.0048 (15) %. Thus, I_α = 99.9952 (15) %.

Table 9 - Measured and recommended branching ratio for ²²¹Fr β⁻ decay.

I _β (%)	References
0.0011 (5)	1995Ch74
0.0048 (15)	1997Ch53
0.0048 (15)	Recommended value, from 1997Ch53

7. References

- 1947En03 A.C. English, T.E. Cranshaw, P. Demers, J.A. Harvey, E.P. Hincks, J.V. Jelley, A.N. May, Phys.Rev. 72, 253 (1947) [T_{1/2}].
- 1950Ha52 F. Hagemann, L.I. Katzin, M.H. Studier, G.T. Seaborg, A. Ghiorso, Phys.Rev. 79, 435 (1950) [T_{1/2}].
- 1967Dz02 B.S. Dzhelepov, R.B. Ivanov, M.A. Mikhailova, L.N. Moskvin, O.M. Nazarenko, V.F. Rodionov, Izv.Akad.Nauk SSSR, Ser.Fiz. 31, 568 (1967) [Eα, Iα].
- 1967LoZZ W. Lourens, Thesis, Technische Hogeschool Delft (1967) [T_{1/2}].
- 1968Le07 C.-F. Leang, G. Bastin-Scoffier, Compt.Rend. 266B, 629 (1968) [Eα, Iα, Eγ, Iγ].
- 1981Di14 J.K. Dickens, J.W. McConnell, Radiochem.Radioanal.Lett. 47, 331 (1981) [Eγ, Pγ].
- 1986He06 R.G. Helmer, C.W. Reich, M.A. Lee, I. Ahmad, Int.J.Appl.Radiat.Isotop. 37, 139 (1986) [Eγ, Pγ].
- 1987He28 R.G. Helmer, M.A. Lee, C.W. Reich, Nuclear Physics A474 (1987) 77
- 1990Ak05 Y.A. Akovali, Nucl.Data Sheets 61, 623(1990) [NDS].
- 1991Ry01 A. Rytz, At.Data Nucl.Data Tables 47, 205 (1991) [Evaluation].
- 1994Ar23 G. Ardisson, V. Barci, O. El Samad, Nucl.Instrum.Methods Phys.Res. A339, 168 (1994) [Eγ, Iγ].
- 1995Bu17 Yu.S. Butabaev, I. Adam, K.Ya. Gromov, S.S. Eliseev, R.A. Niyazov, Yu.V. Norseev, V.I. Fominykh, A.Kh. Kholmatov, V.V. Tsupko-Sitnikov, V.G. Chumin, M.B. Yuldashev, Bull.Rus.Acad.Sci.Phys. 59, 5(1995) [Eγ, Iγ].
- 1995Ch74 V.G. Chumin, S.S. Eliseev, K.Ya. Gromov, Yu.V. Norseev, V.I. Fominykh, V.V. Tsupko-Sitnikov, Bull.Rus.Acad.Sci.Phys. 59,1854(1995) [P_β].
- 1995Sh01 R.K. Sheline, C.F. Liang, P. Paris, Phys.Rev. C51, 1192 (1995) [Eγ, Iγ,Pγ].
- 1996Sc06 E. Schönfeld, H. Janssen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
- 1997Ch53 V.G. Chumin, J.K. Jabber, K.V. Kalyapkin, S.A. Kudrya, V.V. Tsupko-Sitnikov, K.Ya. Gromov, V.I. Fominykh, T.A. Furyaev, Bull.Rus.Acad.Sci.Phys. 61, 1606 (1997) [P_β].
- 1999Gr33 K.Ya. Gromov, J.K. Jabber, Sh.R. Malikov, V.I. Fominykh, Yu.V. Kholnov, V.V. Tsupko-Sitnikov, V.G. Chumin, Bull.Rus.Acad.Sci.Phys. 63, 685 (1999) [Eγ, Iγ,Pγ].
- 2000Ga52 J. Gasparro, G. Ardisson, V. Barci, R.K. Sheline, Phys.Rev. C62, 064305 (2000) [Pγ].
- 2002Gr36 K.Ya. Gromov, S.A. Kudrya, Sh.R. Malikov, V.A. Sergienko, V.I. Fominykh, V.V. Tsupko-Sitnikov, V.G. Chumin, Bull.Rus.Acad.Sci.Phys. 66, 1519 (2002) [Eα, Iα, Eγ, Iγ].
- 2003Ak06 Y.A. Akovali, Nucl.Data Sheets 100, 141 (2003) [NDS].
- 2003Au03 G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].
- 2007Ba19 M.S. Basunia, Nucl.Data Sheets 108, 633 (2007) [NDS].
- 2007Je07 H.B. Jeppesen, J. Byskov-Nielsen, P. Wright, J.G. Correia, L.M. Fraile, H.O.U. Fynbo, K. Johnston, K. Riisager, Eur.Phys.J. A 32, 31 (2007) [T_{1/2}].

**²²²Rn - Comments on evaluation of decay data
by V. Chisté and M. M. Bé**

1 Decay Scheme

²²²Rn disintegrates by alpha emission mainly to the ground state level of ²¹⁸Po. Spin and parity are from the mass-chain evaluation of Y. A. Akovali (1987E112, 1995E08 for A = 218 and 1996E101 for A = 222) and A. K. Jain (2006Ja03 for A = 218).

The calculated Q value of 5590.2 (6) keV deduced from the decay scheme data is in good agreement with the adopted value from Audi *et al.*

2 Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2003Au03).

The experimental ²²²Rn half-life values (in days) are given in Table 1:

Table 1: Experimental values of ²²²Rn half-life.

Reference	Experimental value (d)	Comments
W. Bothe (1923Bo**)	3.824 (4)	Ionization-chamber. Revised uncertainty by N.E. Holden (1990Ho28).
I. Curie (1924Cu**)	3.823 (2)	Ionization-chamber. Revised uncertainty by N.E. Holden (1990Ho28).
J. Tobailem (1951To25)	3.825 (5)	Ionization-chamber. Revised uncertainty by N.E. Holden (1990Ho28).
J. Robert (1956Ro31)	3.825 (4)	Calorimetry. Revised uncertainty by N.E. Holden (1990Ho28).
P. C. Marin (1956Ma64)	3.8229 (17)	Revised uncertainty by N.E. Holden (1990Ho28).
N. S. Shimanskaya (1958Sh69)	3.83 (3)	Calorimetry. Outlier
D. K. Butt (1972Bu33)	3.8235 (17)	Revised uncertainty by N.E. Holden 1990Ho28.
R. Collé (1995Co**)	3.8224 (18)	Liquid scintillator.
H. Schrader (2004Sc04)	3.8195 (30)	Ionization-chamber. Outlier
Recommended value	3.8232 (8)	$\chi^2 = 0.11$

For the half-life values in references from W. Bothe (1923Bo*) to D. K. Butt (1972Bu33), the retained values take into account the uncertainty recommendations given by N. E. Holden (1990Ho28). With this data set, a weighted average was calculated using LWEIGHT computer program (version 3). Based on the Chauvenet's criterion, the Shimanskaya (1958Sh69) and Schrader's (2004Sc04) values have been shown outlier and then omitted in the final calculation.

The recommended value of ²²²Rn half-life is the weighted average of **3.8232 days** with an internal uncertainty of **0.0008 day**. The reduced- χ^2 value is 0.11 and the critical χ^2 value is 2.8.

2.1 a Transitions and Emissions

The energies of the α -particle transitions given in Section 2.1 were calculated from Q_α (2003Au03) and level energies.

The energy of the $\alpha_{0,0}$ emission given in section 4 is from A. Rytz (1991Ry01). For the $\alpha_{0,1}$ and $\alpha_{0,2}$, the emission energies are given by R. J. Walen (1958Wa16).

The α -particle emission probabilities are recommended by A. Rytz (1991Ry01). For the $\alpha_{0,1}$ emission probability, the adopted value is the measured value of R. J. Walen (1958Wa16) (0.078). Existence of the $\alpha_{0,2}$ branch is questionable.

2.2 g Transitions

The $\gamma_{(1,0)}$ transition probability was deduced from the decayscheme balance using recommended experimental α -particle intensity value of 0.078 given by R.J. Walen (1958Wa16). (see **2.1 a Transitions and Emissions**). The multipolarity of the 510-keV γ -ray transition is from 2006Ja03.

510-keV γ -ray : [E2]

The internal conversion coefficients (ICC's) for this γ -ray transition have been calculated using the Br Icc computer program, which interpolates from theoretical values of I. M. Band (2002Ba85).

3 Atomic Data

Atomic values, ω_K , ω_L and n_{KL} and the X-ray relative probabilities are from Schönfeld and Janßen (1996Sc06).

4 a Emissions

See **2.1 a Transitions and Emissions**.

5 Photon emissions

5.1 g-ray Emissions

The energy of the 510 keV γ -ray given in Section 5.1 was measured by L. Madansky (1956Ma28).

The intensity of 0,076 deduced from alpha intensity measurements is in agreement with the measured value of 0,07 obtained by L. Madansky (1956Ma28).

6 References

- 1923Bo** W. Bothe, Z. Phys. 16(1923)266 [Half-life].
 1924Cu** I. Curie, C. Chamié, J. Phys. Radium 5(1924)238 [Half-life].
 1951To25 J. Tobailem, Compt. Rend. (Paris) 233(1951)1360 [Half-life].
 1956Ma28 L. Madansky, F. Rasetti, Phys. Rev. 102(1956)464 [E_γ].
 1956Ro31 J. Robert, J. Phys. Radium 17(1956)605 [Half-life].
 1956Ma64 P. C. Marin, Brit. J. Appl. Phys. 7(1956)188 [Half-life].
 1958Sh69 N. S. Shimanskaya, Instr. Experiments Techniques 2(1958)283 [Half-life].
 1958Wa16 R. J. Walen, G. Bastin, Comp. Rend. Int. Conf. Nucl. Phys., Paris, (1959)910 [E_α , I_α].
 1971Gr17 B. Grennberg, A. Rytz, Metrologia 7(1971)65 [E_α].
 1972Bu33 D. K. Butt, A. R. Wilson, J. Phys. (London) A5(1972)1248 [Half-life].
 1979Ry03 A. Rytz, At. Data and Nucl. Data Tables 23(1979)205 [E_α , I_α].
 1987El12 Y. A. Ellis-Akovali, Nucl. Data Sheets 52(1987)789 [Spin, parity and multipolarity].
 1990Ho28 N. E. Holden, Pure Appl. Chem. 62(1990)941 [Half-life].
 1991Ry01 A. Rytz, At. Data and Nucl. Data Tables 47(1991)205 [E_α , I_α].
 1995Co** R. Collé, Radioact. Radiochem. 6(1995)16 [Half-life].
 1995El08 Y. A. Akovali, Nucl. Data Sheets 76(1995)457 [Spin, parity and multipolarity].
 1996El01 Y. A. Akovali, Nucl. Data Sheets 77(1996)271 [Spin, parity and multipolarity].
 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
 2002Ba85 I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr., P. O. Tikkanen, S. Raman, At. Data Nucl. Data Tables 81(2002)1 [Theoretical ICC].
 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].
 2004Sc04 H. Schrader, Appl. Radiat. Isot. 60(2004)317 [Half-life].
 2006Ja03 A. K. Jain, B. Singh, Nucl. Data Sheets 107(2006)1027 [Spin, parity and multipolarity].

²²³Fr-Comments on evaluation of the decay data

Huang Xiaolong, Wang Baosong

This evaluation was completed in 2008. Literature available by December 2008 was included.

1 Decay Scheme

²²³Fr disintegrates 0.020 (4) % by α emission to levels in ²¹⁹At and 99.980 (4) % by β^- emission to levels in ²²³Ra. ²²³Fr ground state has $J^\pi=3/2^-$ (2001Br31).

The α decay scheme of ²²³Fr was built based on the measurement of 2001Li44. The β^- decay scheme of ²²³Fr was built based on the measurement of 1993Ab01.

The adopted $Q(\alpha)$ and $Q(\beta^-)$ values of Audi(2003Au03) are in good agreement with the $Q(\alpha)$ and $Q(\beta^-)$ values deduced from the decay scheme data.

2 Nuclear Data

The Q values are from the 2003Au03 atomic-mass adjustment.

Level energies have been deduced from a least-squares fit to γ -ray energies (GTOL computer code). Spin and parities are from 2001Br31 and 2001Li44.

The measured and our recommended ²²³Fr half-life values are listed in Table 1.

Table 1 Measured half-life values of ²²³Fr and recommended value, in minutes.

$T_{1/2}$ (min)	References	measurement method
22 (1)	1955Ad10	
21.8 (4)	1967Li17	G-M counter
22.00 (7)	1993Ab01	HPGe detector
22.00 (7)	2001Br31	NDS, weighted average of 1967Li17, 1993Ab01
21.93 (7)		Unweighted mean
21.99 (7)		LWM, $\chi^2=0.12$
22.00 (7)		Recommended value, from 1993Ab01

The recommended value is from the measurement of 1993Ab01.

2.1 γ Transitions

The γ -ray transition probabilities were deduced using the γ -ray emission intensities and relevant theoretical internal conversion coefficients.

Multipolarities and mixing ratios of γ -ray transitions for β^- decay are from 2001Br31, for α decay from 2001Li44. The mixing ratio of the 29.78keV γ -ray is from the experimental data of 1990Br23, the uncertainty was assumed to be 10 %.

The internal conversion coefficients (ICC) and their associated uncertainties have been obtained using the BrIcc computer program, which applies the ‘‘Frozen Orbital’’ approximation (2002Ba85).

2.2 α Transitions

The measured and evaluated energies of alpha particles were listed in table 2. The recommended values are from 2001Li44.

Table 2. Measured and recommended values of α -particle energies from ²²³Fr decay

1955Ad10	2001Li44	recommended
	5462 (3)	5462 (3)
	5403 (3)	5403 (3)
5340 (80)	5314 (4)	5314 (4)
	5291 (4)	5291 (4)
	5172 (5)	5172 (5)

The measured and evaluated alpha particle emission probabilities are listed in table 3. The recommended values are from 2001Li44.

Table 3. Measured and recommended α -particle emission probabilities from ²²³Fr decay

E_α/keV	$P_\alpha(10^{-4})$		
	2001Li44	Calc.	recommended
5462 (3)	33 (15)	0	33 (15)
5403 (3)	44 (20)	95 (40)	44 (20)
5314 (4)	53 (23)	70 (35)	53 (23)
5291 (4)	60 (26)	60 (30)	60 (26)
5172 (5)	9 (5)	8 (5)	9 (5)

2.3 β^- transition

The maximum energies of the β^- transitions in the decay of ²²³Fr have been deduced from the Q value (2003Au03) and the level energies.

The adopted ϵ and β^- transition probabilities and their associated uncertainties were deduced from the γ transition probability balance at each level of the decay scheme.

The electron capture subshell probabilities and $lg ft$ values were calculated using the LOGFT program.

3. Atomic data

Atomic fluorescence yields ($\omega_K, \omega_L, \omega_M, \eta_{KL}$ and η_{LM}) are from Schönfeld (1996Sc06).

The X-ray and Auger electron emission probabilities have been deduced from γ -ray and conversion electron data using the computer code RADLST. Measured and calculated X-ray emission probabilities are compared in Table 4.

Table 4 Comparison of calculated and measured Ra X-ray emission intensities

	1982AIZL	Adopted (deduced)
$K_{\alpha 1}$	2.4 (5)	2.3 (3)
$K_{\alpha 2}$	1.43 (28)	1.44 (19)

The radium KX-ray emission probabilities, deduced from γ -ray data, agree with the measured values of 1982AIZL, thus confirming the completeness of the decay scheme.

4. Electron Emissions.

The conversion electron emission probabilities have been deduced from γ -ray emission probabilities and theoretical conversion coefficients.

5. Photon Emissions

5.1 γ -ray energy values

There is one measurement of γ -ray energies from ²²³Fr α decay, that's 2001Li44. Our recommended γ -ray energies from ²²³Fr α decay are from 2001Li44. The 24.14keV γ -ray, which was observed in ²²³Fr decay was assigned by evaluators to ²²³Fr α decay.

The measured and recommended γ -ray energies from ²²³Fr β^- decay are listed in table 5. The recommended values are from 1993Ab01.

Table 5 Measured and recommended values of γ -ray energies from ²²³Fr β^- decay

1964Yt01	1967MA19	1982ALZL	1993Ab01	recommended
			20.27 (5)	20.27 (5)
			27.27 (3) ^b	27.27 (3) ^b
			29.78 (4)	29.78 (4)
			31.69 (5)	31.69 (5)
			43.5 (2)	43.5 (2)
			44.0 (1)	44.0 (1)
			49.80 (5)	49.80 (5)
50 (2)	50.8 (5)	50.087 (12)	50.10 (2)	50.10 (2)
	61.0 (15)		61.43 (5)	61.43 (5)
			62.31 (6)	62.31 (6)
			73.5 (1)	73.5 (1)
80 (2)	80.0 (4)	79.651 (13)	79.65 (2)	79.65 (2)
		88.483 (11)	89.08 (10)	89.08 (10)
100 (5)			93.88 (5)	93.88 (5)
			111.05 (3)	111.05 (3)
136 (5)	134.4 (4)	134.641 (22)	134.60 (2)	134.60 (2)
			150.6 (4) ^b	150.6 (4) ^b
			155.5 (5)	155.5 (5)
167 (8)	173.1 (5)	173.393 (38)	173.35 (5)	173.35 (5)
	184.5 (5)	184.693 (38)	184.65 (5)	184.65 (5)
191 (15)			200.7 (2)	200.7 (2)
205 (5)	204.8 (4)	204.948 (15)	204.85 (5)	204.85 (5)
			205.6 (2)	205.6 (2)
			210.60 (5)	210.60 (5)
			218.80 (5)	218.80 (5)
			222.9 (3) ^b	222.9 (3) ^b
234 (3)	234.6 (4)	234.796 (10)	234.70 (5)	234.70 (5)
			236.05 (5)	236.05 (5)
	246 (1)	245.56 (21)	245.60 (5)	245.60 (5)
	250.6 (10)	250.12 (12)	250.25 (5) ^a	250.25 (5) ^a
			254.6 (2)	254.6 (2)
	256 (1)	256.09 (18)	256.18 (5)	256.18 (5)

1964Yt01	1967MA19	1982ALZL	1993Ab01	recommended
			262.9 (2)	262.9 (2)
			269.6 (3) ^b	269.6 (3) ^b
			272.8 (2)	272.8 (2)
			280.7 (5) ^a	280.7 (5) ^a
	286.0 (15)	285.9 (6)	286.0 (2)	286.0 (2)
289 (10)	289.6 (15)	289.73 (10)	289.67 (5)	289.67 (5)
			293.2 (2) ^b	293.2 (2) ^b
			296.5 (2)	296.5 (2)
	300.0 (15)	299.92 (20)	299.95 (5)	299.95 (5)
	304.2 (15)		304.40 (5)	304.40 (5)
	307.3 (15)	307.63 (20)	307.93 (5) ^a	307.93 (5) ^a
	313.3 (15)	312.7 (7)	312.65 (5)	312.65 (5)
			314.6 (2)	314.6 (2)
318 (10)	319.0 (5)	319.266 (22)	319.25 (5)	319.25 (5)
	330.0 (15)		329.80 (5)	329.80 (5)
	333.1 (15)		334.30 (6)	334.30 (6)
	338.7 (10)		339.50 (5)	339.50 (5)
	343.0 (15)		342.50 (7)	342.50 (7)
355 (12)			350.5 (2)	350.5 (2)
	369.0 (5)	369.46 (6)	369.32 (5)	369.32 (5)
			382.3 (2) ^b	382.3 (2) ^b
			434.4 (1)	434.4 (1)
			439.6 (3)	439.6 (3)
			444.5 (3)	444.5 (3)
			452.9 (2) ^a	452.9 (2) ^a
			457.5 (2)	457.5 (2)
			469.3 (2) ^a	469.3 (2) ^a
			475.4 (1) ^a	475.4 (1) ^a
			480.9 (3)	480.9 (3)
			493.4 (2)	493.4 (2)
			506.9 (2)	506.9 (2)
			516.7 (2)	516.7 (2)
			524.8 (2)	524.8 (2)
			533.1 (3)	533.1 (3)
			537.2 (2) ^a	537.2 (2) ^a
			539.8 (2)	539.8 (2)
			545.4 (4)	545.4 (4)
			552.3 (2)	552.3 (2)
			556.3 (3)	556.3 (3)
		568.85 (15)	569.03 (8)	569.03 (8)
			576.1 (4)	576.1 (4)
			581.3 (4)	581.3 (4)
			592.3 (2)	592.3 (2)
			596.9 (4)	596.9 (4)
			600.7 (4)	600.7 (4)

1964Yt01	1967MA19	1982ALZL	1993Ab01	recommended
			607.6 (3)	607.6 (3)
			613.6 (4)	613.6 (4)
			632.7 (3)	632.7 (3)
			663.7 (3)	663.7 (3)
			671.9 (4)	671.9 (4)
			682.3 (3) ^b	682.3 (3) ^b
			694.6 (3) ^b	694.6 (3) ^b
			708.3 (3)	708.3 (3)
	723 (1)	723.7 (7)	722.65 (5)	722.65 (5)
			724.15 (5)	724.15 (5)
			737.4 (3)	737.4 (3)
			742.4 (3)	742.4 (3)
	746.5 (15)	746.3 (9)	746.30 (5)	746.30 (5)
			753.65 (5)	753.65 (5)
	756 (2)		757.20 (5)	757.20 (5)
			762.6 (2)	762.6 (2)
	766.5 (20)	764.7 (7)	766.64 (5)	766.64 (5)
	776.0 (6)	776.0 (7)	775.83 (5)	775.83 (5)
781 (10)			780.8 (1)	780.8 (1)
	784 (2)		784.93 (5)	784.93 (5)
			787.6 (2) ^a	787.6 (2) ^a
	793 (2)		792.2 (3)	792.2 (3)
	797.5 (20)		796.22 (5)	796.22 (5)
		803.7 (7)	803.77 (5)	803.77 (5)
	804 (1)		806.0 (2)	806.0 (2)
	813 (2)	812.0 (10)	812.40 (6)	812.40 (6)
			816.5 (2)	816.5 (2)
	821.5 (25)		823.20 (7)	823.20 (7)
	826 (1)	826.4 (11)	825.95 (7)	825.95 (7)
	835 (2)		833.9 (2)	833.9 (2)
			837.5 (1)	837.5 (1)
	840.5 (20)	842.0 (9)	842.2 (1)	842.2 (1)
	847 (1)	847.7 (10)	846.85 (10) ^a	846.85 (10) ^a
	860 (2)		863.6 (1)	863.6 (1)
	864 (2)		867.4 (1)	867.4 (1)
	876.5 (10)	876.2 (10)	876.5 (1)	876.5 (1)
			878.1 (2)	878.1 (2)
	892 (3)		893.1 (2)	893.1 (2)
	897.5 (20)	896.7 (10)	896.7 (2)	896.7 (2)
	908 (2)	907.7 (10)	907.6 (2)	907.6 (2)
			911.3 (2)	911.3 (2)
			913.6 (3)	913.6 (3)
			926.5 (3)	926.5 (3)
			941.2 (3)	941.2 (3)
			949.3 (4)	949.3 (4)

1964Yt01	1967MA19	1982ALZL	1993Ab01	recommended
			958.0 (7)	958.0 (7)
			969.2 (4)	969.2 (4)
			975.2 (5)	975.2 (5)
			978.7 (4)	978.7 (4)
			989.4 (5)	989.4 (5)
			994.3 (3)	994.3 (3)
			999.3 (5)	999.3 (5)
			1025.1 (5)	1025.1 (5)

a: multiply placed. b: not placed in level scheme.

5.2 Relative values of γ -ray intensities

The measured and recommended γ -ray emission probabilities from ²²³Fr α decay are listed in table 6. The recommended values are from 2001Li44.

Table 6. Measured and recommended values of γ -ray energies and emission probabilities from ²²³Fr α decay

E_γ (keV)		P_γ (10^{-4} %)	
2001Li44	recommended	2001Li44	recommended
	24.14 (3)		60 (26) ^a
58.9 (2)	58.9 (2)	8 (3)	8 (3)
150.9 (2)	150.9 (2)	56 (5)	56 (5)
145.3 (3)	145.3 (3)	2 (1)	2 (1)

^a: (γ +ce), from intensity balance.

Measured values of the relative γ -ray intensities, the 234.7 keV being the reference line, from ²²³Fr β^- decay are listed in Table 7. 1964Yt01 and 1967MA19 are replaced by 1993Ab01, these three references coming from the same group. There is no detailed experimental information in 1982ALZL and only the data table are given. It's noted that among 131 γ -rays, 87 γ -rays are new and observed in 1993Ab01. Compared to 1993Ab01, 1982ALZL did not provide the γ -rays with low energy; their γ -ray intensities are in agreement with those reported by 1993Ab01 for some γ -rays, and for most of the weak γ -rays quite different. Then, the present adopted values are from 1993Ab01.

Table 7 Measured and recommended relative γ -ray intensities from ²²³Fr β^- decay.

E_γ /keV	I_γ				
	1964Yt01	1967MA19	1982ALZL	1993Ab01	recommended
20.27 (5)				53 (5)	53 (5)
27.27 (3) ^b				2.3 (4)	2.3 (4)
29.78 (4)				2.6 (4)	2.6 (4)
31.69 (5)				0.05	0.05
43.5 (2)				0.08	0.08
44.0 (1)				0.05	0.05
49.80 (5)				93 (8)	93 (8)
50.10 (2)	100	1000	1200 (80)	1224 (50)	1224 (50)
61.43 (5)		< 8		0.13	0.13

E_{γ}/keV	I_{γ}				
	1964Yt01	1967MA19	1982ALZL	1993Ab01	recommended
62.31 (6)				0.6 (2)	0.6 (2)
73.5 (1)				0.05 (3)	0.05 (3)
79.65 (2)	32.8 (3)	240 (24)	290 (20)	335 (20)	335 (20)
89.08 (10)			88 (3)	2.0 (1)	2.0 (1)
93.88 (5)			31 (1)	2.2 (3)	2.2 (3)
111.05 (3)			10.6 (4)	0.18 (4)	0.18 (4)
134.60 (2)	1.0 (1)	16.0 (16)	17.3 (10)	18.5 (5)	18.5 (5)
150.6 (4) ^b				0.10 (3)	0.10 (3)
155.5 (5)				0.1	0.1
173.35 (5)	0.6 (1)	4.0 (4)	3.35 (25)	4.26 (5)	4.26 (5)
184.65 (5)		9.0 (9)	7.7 (6)	8.27 (5)	8.27 (5)
200.7 (2)				0.10 (3)	0.10 (3)
204.85 (5)	2.1 (2)	34.0 (34)	30.9 (17)	33.7 (3)	33.7 (3)
205.6 (2)				0.22	0.22
210.60 (5)				0.36 (2)	0.36 (2)
218.80 (5)				0.32 (2)	0.32 (2)
222.9 (3) ^b				0.08 (2)	0.08 (2)
234.70 (5)	8.9 (8)	100	100.0 (35)	100	100
236.05 (5)				1.0 (2)	1.0 (2)
245.60 (5)		1.3 (4)	1.1 (4)	0.71 (3)	0.71 (3)
250.25 (5) ^a		1.3 (4)	1.0 (4)	0.11	0.11
250.25 (5) ^a				0.58	0.58
254.6 (2)				0.21 (2)	0.21 (2)
256.18 (5)		1.3 (4)	1.2 (4)	0.75 (3)	0.75 (3)
262.9 (2)				0.13 (3)	0.13 (3)
269.6 (3) ^b				0.03 (1)	0.03 (1)
272.8 (2)				0.15 (2)	0.15 (2)
280.7 (5) ^a				0.02	0.02
280.7 (5) ^a				0.02	0.02
286.0 (2)		0.52 (15)	0.2	0.17 (2)	0.17 (2)
289.67 (5)	1.4 (2)	7.2 (7)	7.6 (4)	7.7	7.7
293.2 (2) ^b				0.14 (3)	0.14 (3)
296.5 (2)				0.05 (1)	0.05 (1)
299.95 (5)		1.3 (4)	0.57 (13)	0.75 (4)	0.75 (4)
304.40 (5)		0.67 (20)		0.32 (2)	0.32 (2)
307.93 (5) ^a		0.90 (27)	0.7 (3)	0.45 (5)	0.45 (5)
307.93 (5) ^a				0.05 (5)	0.05 (5)
312.65 (5)		0.75 (22)	0.36 (18)	0.60 (5)	0.60 (5)
314.6 (2)				0.08 (2)	0.08 (2)
319.25 (5)	1.9 (3)	16.2 (16)	15.4 (8)	17.0 (5)	17.0 (5)
329.80 (5)		1.0 (3)		0.90 (5)	0.90 (5)
334.30 (6)		0.45 (13)		0.31 (2)	0.31 (2)
339.50 (5)		2.0 (4)		2.3 (2)	2.3 (2)
342.50 (7)		0.90 (27)		0.43 (4)	0.43 (4)

E_γ/keV	I_γ				
	1964Yt01	1967MA19	1982ALZL	1993Ab01	recommended
350.5 (2)	0.3 (1)			0.10 (5)	0.10 (5)
369.32 (5)		3.2 (3)	3.40 (33)	3.3 (2)	3.3 (2)
382.3 (2) ^b				0.03 (1)	0.03 (1)
434.4 (1)				0.08 (2)	0.08 (2)
439.6 (3)				0.011 (3)	0.011 (3)
444.5 (3)				0.04 (1)	0.04 (1)
452.9 (2) ^a				0.03	0.03
452.9 (2) ^a				0.03	0.03
457.5 (2)				0.03	0.03
469.3 (2) ^a				0.04	0.04
469.3 (2) ^a				0.04	0.04
475.4 (1) ^a				0.11	0.11
475.4 (1) ^a				0.1	0.1
480.9 (3)				0.05 (1)	0.05 (1)
493.4 (2)				0.09 (2)	0.09 (2)
506.9 (2)				0.08 (2)	0.08 (2)
516.7 (2)				0.12 (2)	0.12 (2)
524.8 (2)				0.16 (3)	0.16 (3)
533.1 (3)				0.07 (2)	0.07 (2)
537.2 (2) ^a				0.07	0.07
537.2 (2) ^a				0.12	0.12
539.8 (2)				0.22 (5)	0.22 (5)
545.4 (4)				0.011 (3)	0.011 (3)
552.3 (2)				0.10 (2)	0.10 (2)
556.3 (3)				0.04 (1)	0.04 (1)
569.03 (8)			1.9 (3)	1.8 (2)	1.8 (2)
576.1 (4)				0.04 (1)	0.04 (1)
581.3 (4)				0.05 (1)	0.05 (1)
592.3 (2)				0.12 (3)	0.12 (3)
596.9 (4)				0.03 (1)	0.03 (1)
600.7 (4)				0.020 (5)	0.020 (5)
607.6 (3)				0.08 (2)	0.08 (2)
613.6 (4)				0.04 (1)	0.04 (1)
632.7 (3)				0.08 (2)	0.08 (2)
663.7 (3)				0.04 (1)	0.04 (1)
671.9 (4)				0.020 (5)	0.020 (5)
682.3 (3) ^b				0.03 (1)	0.03 (1)
694.6 (3) ^b				0.03 (1)	0.03 (1)
708.3 (3)				0.05 (1)	0.05 (1)
722.65 (5)		1.5 (3)		1.4 (2)	1.4 (2)
724.15 (5)			1.9 (8)	0.52 (8)	0.52 (8)
737.4 (3)				0.033 (8)	0.033 (8)
742.4 (3)				0.04 (1)	0.04 (1)
746.30 (5)		0.70 (15)	0.7 (2)	0.74 (8)	0.74 (8)

E_γ/keV	I_γ				
	1964Yt01	1967MA19	1982ALZL	1993Ab01	recommended
753.65 (5)				0.35 (5)	0.35 (5)
757.20 (5)		0.40 (8)		0.28 (5)	0.28 (5)
762.6 (2)			0.7 (2)	0.09 (2)	0.09 (2)
766.64 (5)		0.80 (16)		0.83 (8)	0.83 (8)
775.83 (5)		12.3 (12)	15.1 (10)	16.8 (5)	16.8 (5)
780.8 (1)	1.9 (2)			0.11 (3)	0.11 (3)
784.93 (5)		0.70 (15)		0.32 (5)	0.32 (5)
787.6 (2) ^a				0.09 (2)	0.09 (2)
787.6 (2) ^a				0.01 (1)	0.01 (1)
792.2 (3)		0.40 (6)		0.020 (5)	0.020 (5)
796.22 (5)		0.30 (6)		0.40 (5)	0.40 (5)
803.77 (5)		1.70 (25)	2.4 (6)	2.2 (3)	2.2 (3)
806.0 (2)				0.05 (1)	0.05 (1)
812.40 (6)		0.60 (9)	0.66 (22)	0.78 (8)	0.78 (8)
816.5 (2)				0.05 (1)	0.05 (1)
823.20 (7)		0.30 (9)		0.26 (3)	0.26 (3)
825.95 (7)		1.4 (2)	2.3 (4)	2.0 (3)	2.0 (3)
833.9 (2)				0.05 (1)	0.05 (1)
837.5 (1)		0.20 (6)		0.36 (4)	0.36 (4)
842.2 (1)		0.20 (6)	0.5	0.18 (2)	0.18 (2)
846.85 (10) ^a		1.4 (2)	2.4 (11)	1.8 (3)	1.8 (3)
846.85 (10) ^a				0.2 (1)	0.2 (1)
863.6 (1)		0.10 (3)		0.14 (2)	0.14 (2)
867.4 (1)				0.06 (1)	0.06 (1)
876.5 (1)		1.3 (2)	1.4 (4)	1.4 (2)	1.4 (2)
878.1 (2)				0.12 (2)	0.12 (2)
893.1 (2)		0.10 (3)		0.09 (2)	0.09 (2)
896.7 (2)		0.50 (8)	0.7 (3)	0.50 (5)	0.50 (5)
907.6 (2)		0.40 (7)	0.3	0.53 (5)	0.53 (5)
911.3 (2)				0.03 (1)	0.03 (1)
913.6 (3)				0.015 (5)	0.015 (5)
926.5 (3)				0.06 (1)	0.06 (1)
941.2 (3)				0.11 (2)	0.11 (2)
949.3 (4)				0.012 (3)	0.012 (3)
958.0 (7)				0.013 (3)	0.013 (3)
969.2 (4)				0.012 (3)	0.012 (3)
975.2 (5)				0.006 (2)	0.006 (2)
978.7 (4)				0.025 (5)	0.025 (5)
989.4 (5)				0.005 (1)	0.005 (1)
994.3 (3)				0.004 (1)	0.004 (1)
999.3 (5)				0.007 (2)	0.007 (2)
1025.1 (5)				0.005 (1)	0.005 (1)

^a: multiply placed. ^b: not placed in level scheme.

5.3 Absolute values of γ -ray emission intensities

The reference gamma-ray line, in the table above, is 234.70 keV. But the measured absolute gamma-ray intensity was given for the 204.8 keV gamma-ray. So the normalization factor N is deduced from the 204.8 keV gamma-ray.

The calculation is:

The measured absolute gamma-ray intensity for the 204.8 keV line (1981Va28) is: $P(204.8 \text{ keV}) = 0.92 (18) \%$, the recommended relative gamma-ray intensity is $I(204.8 \text{ keV}) = 33.7 (3)$.

So $N = P(204.8 \text{ keV}) / I(204.8 \text{ keV}) = 0.92 (18) / 33.7 (3) = 0.027 (5)$.

This value is very close to that calculated with the formula $N = 100 / \sum [I(\text{ce}+\gamma)(\text{g.s.})]$ assuming $I_{\beta^-}(\text{g.s.}) \leq 1 \%$.

So, N has been taken from 1981Va28, that's $N = 0.027 (5)$.

The recommended absolute γ -ray emission probabilities are equal to the relative values given in table 7 multiplied by 0.027 (5).

6. Branching Ratio

The measured and recommended total branching ratios from ²²³Fr decay are listed in table 8. The recommended value of $\% \alpha = 0.020 (4) \%$ is from 2001Li44. Thus, $\% \beta^- = 99.980 (4) \%$.

Table 8 Measured and recommended α -branching ratio from ²²³Fr decay.

$I_{\alpha} / \%$	References
0.006	1955Ad10
0.020 (4)	2001Li44
0.020 (4)	Recommended value, from 2001Li44

7. References

- 1954Hy26 E.K.Hyde, Phys.Rev. 94, 1221 (1954) [P _{γ}].
- 1955Ad10 J.P.Adloff, Compt.Rend. 240, 1421 (1955) [T_{1/2}, E _{α} , I _{α}].
- 1964Yt01 C.Ythier, G.Mazzone, P.W.F.Louwrier, Physica 30, 2143 (1964) [E _{γ} , I _{γ}].
- 1967Li17 K.H.Lieser, E.Kluge, Radiochim.Acta 7, 3 (1967) [T_{1/2}].
- 1967Ma19 H.Maria, C.Ythier, P.Polak, A.H.Wapstra, Physica 34, 571(1967), [E _{γ} , I _{γ}].
- 1981Va28 S.K.Vasilev, B.S.Dzhelepov, R.B.Ivanov, M.A.Mikhailova, A.V.Mozzhukhin, B.I.Shestakov, Izv.Akad.Nauk SSSR, Ser.Fiz. 45, 1895 (1981) [P _{γ}].
- 1982AIZL Yu.V.Aleksandrov, S.K.Vasilev, B.S.Dzhelepov, R.B.Ivanov, M.A.Mikhailova, A.V.Mozzhukhin, A.V.Saulsky, B.I.Shestakov, Program and Theses, Proc.32nd Ann. Conf. Nucl. Spectrosc. Struct. At. Nuclei, Kiev, p.135 (1982) [I_{KX}, E _{γ} , I _{γ}].
- 1990Br23 Ch.Briançon, S.Cwiok, S.A.Eid, V.Green, W.D.Hamilton, C.F.Liang, R.J.Walen, J.Phys. (London) G16, 1735 (1990) [ICC]
- 1993Ab01 A.Abdul-Hadi, V.Barci, B.Weiss, H.Maria, G.Ardisson, M.Hussonnois, O.Constantinescu, Phys.Rev. C47, 94 (1993) [T_{1/2}, E _{γ} , I _{γ}].
- 1996Sc06 E.Schönfeld, H.Janssen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
- 2001Br31 E.Browne, Nucl.Data Sheets 93, 763 (2001) [NDS]
- 2001Li44 C.F.Liang, P.Paris, R.K.Sheline, Phys.Rev. C64, 034310 (2001) [E _{α} , P _{α} , I _{α}].
- 2002Ba85 I.M.Band, M.B.Trzhaskovskaya, C.W.Nestor, Jr., P.O.Tikkanen, S.Raman, At. Data Nucl. Data Tables 81, 1 (2002) [calculated ICC]
- 2003Au03 G.Audi, A.H.Wapstra, C.Thibault, Nucl. Phys. A729(2003)129 [Q].

**²²³Ra -Comments on evaluation of decay data
by V.P. Chechev**

Evaluated in December 2010 with a literature cut-off by the same date.

1. DECAY SCHEME

²²³Ra decays 100 % to levels of ²¹⁹Rn by emission of α particles, with a very small branch of $6.4 (1) \times 10^{-8}$ % by emission of ¹⁴C (1991Ma16). The adopted ²¹⁹Rn levels populated in the ²²³Ra α decay are based on the measurement by Sheline *et al.* (1998Sh02) and the NDS evaluation by Browne (2001Br31). An intense population takes place only to levels in ²¹⁹Rn with energy less than 500 keV (11 excited levels and ground state) and, in this part, the decay scheme is well defined, though, at some levels, there is a certain discrepancy in the P(α) values measured and deduced from probability balance.

At the same time, for a number of levels with higher energy there is disagreement between measured probabilities of alpha-transitions and the values deduced from P(γ +ce) balance. Besides, the placement of some γ -rays in the level scheme is uncertain and some observed γ -ray transitions have not been placed. Therefore, in this part the decay scheme cannot be considered as fully completed. Further measurements are needed to determine the γ -ray transitions and ²²³Ra α decay scheme with greater precision.

The decay scheme overall consistency is verified by the comparison between Q(calc) = 6027 (133) keV, deduced from the evaluated average energies of all emissions, and Q(α) = 5978.99 (21) keV from the atomic mass evaluation of Audi *et al.* (2003Au03). Percentage deviation between Q(calc) and Q(α) is (0.8 ± 2.2) %. The deviation is not big but more than for other α decaying applied radionuclides. This indicates the need in new precise measurements of α -particle and γ -ray transitions in decay of ²²³Ra.

2. NUCLEAR DATA

Q(α) value is from Audi *et al.* (2003Au03).

The ²²³Ra half-life is based on the experimental results given in Table 1.

Table 1: Experimental values of ²²³Ra half-life (in days)

Reference	Author(s)	Original value	Re-estimated	Method
1954Ha60	Hagee <i>et al.</i>	11.685 (56)		Proportional α counting
1959Ro51	Robert	11.22 (5)		Microcalorimetry
1965Ki05	Kirby <i>et al.</i>	11.4347 (11) ^a	11.4347 (44) ^a	Microcalorimetry
1965Ki05	Kirby <i>et al.</i>	11.4267 (62) ^b	11.427 (17) ^b	Proportional α counting
1967JoZX	Jordan and Blanke	11.372 (45)		Calorimetry
1987Mi10	Miller <i>et al.</i>	11.444 (46)		From α -activity following ²²⁷ Th decay

- ^a The original value was deduced as a weighted average of the data from observations with two calorimeters: 11.4432 (57) and 11.4344 (11) days. The uncertainties are probable errors of a single observation. To take into account the contribution of possible unrecognized systematic errors associated with the method, the evaluator expanded the uncertainty to a half of the difference of the two experimental results (0.0044 day).
- ^b The original value was deduced as a weighted average of the data from observations with a 2π windowless proportional counter for 10 samples. The uncertainties of the 10 measurement results include only statistical errors. To take into account the contribution of possible unrecognized systematic errors associated with the method, for the re-estimated value the evaluator used the smallest uncertainty of 0.017 d stated in the measurements.

From the six values adopted in the data analysis, the LWEIGHT computer program increased the uncertainty in the value of 11.4347 (1965Ki05) to 0.0139 to adjust weights according to the LRSW method and used a weighted average of 11.429 and an external uncertainty of 0.028 ($\chi^2/\nu = 8.05$).

The recommended value for the ²²³Ra half-life is 11.43 (3) days.

2.1. Alpha Transitions

The energies of the alpha transitions have been obtained from the $Q(\alpha)$ value and ²¹⁹Rn level energies given in Table 2 from 2001Br31, where they were deduced from a least-squares fit to gamma-ray energies.

Table 2: ²¹⁹Rn levels populated in ²²³Ra α -decay

Level	Energy (keV)	Spin and parity	Half-life	Probability of α - transition (%)
0	0.0	$5/2^+$	3.96 (1) s	1.0 (2)
1	4.47 (1)	$(9/2)^+$	15.4 (13) ns	-
2	14.37 (1)	$(7/2)^+$	875 (30) ps	0.32 (4)
3	126.77 (2)	$(11/2)^+$	402 (20) ps	10.0 (3)
4	158.64 (1)	$(7/2)^+$	42.3 (50) ps	49.6 (12)
5	269.48 (1)	$3/2^+$	14.2 (23) ps	25.8 (11)
6	338.27 (1)	$(5/2)^+$	6.1 (28) ps	10.6 (10)
7	342.78 (2)	$(5/2, 7/2)^-$		-
8	376.26 (2)	$(9/2)^+$	6.9 (38) ps	0.74 (25)
9	377.33 (6)	$(7/2, 9/2)^-$		-
10	397.1 (4)			≈ 0.008
11	445.03 (1)	$(5/2)^+$	6.2 (31) ps	1.60 (24)
12	446.82 (3)	$(5/2)^-$		0.50 (8)
13	490.92 (2)	$(5/2, 7/2, 9/2)^-$		-
14	514.5 (1)	$(9/2^+)$		≈ 0.13
15	517.7 ?			-
16	541.99 (2)	$(7/2^+)$		≈ 0.13
17	594.1 (1)	$(7/2)^-$		0.16 (4)

18	598.72 (2)	(5/2, 7/2, 9/2) ⁺		0.093
19	623.68 (4)			0.042
20	646.1 (1)			0.041
21	672.6 (5)			0.0053
22	711.3 (1)			0.026
23	732.8 (1)			0.021
24	748			≈ 0.0017
25	773			≈ 6×10 ⁻⁴
26	800			≈ 3×10 ⁻⁴
27	830			≈ 2×10 ⁻⁴
28	851			≈ 4×10 ⁻⁴
29	861			≈ 6.3×10 ⁻⁴
30	873			≈ 4.4×10 ⁻⁴

The recommended values of α -transition probabilities ($P(\alpha)$) are based on the measurements of 1957Pi31, 1962Gi04, 1962Wa18, 1970Da08 and also on the $P(\alpha)$ values deduced by the evaluator from $P(\gamma+ce)$ balance at the corresponding ²¹⁹Rn levels (Table 3).

As the lower part of the decay scheme (²¹⁹Rn levels with energy less than 500 keV) is reasonably complete and well defined, the probabilities of the prominent α -transitions reaching them have been deduced from $P(\gamma+ce)$ balances. The uncertainties of the recommended values were expanded, where necessary, to cover the unweighted mean (UWM) of experimental $P(\alpha)$ values.

The probabilities of weak α -transitions ($P(\alpha) < 0.0015$) have been taken mainly from the measurements of 1962Wa18 with magnetic spectrometer and also from the measurements of 1970Da08 with semiconductor detector. The $P(\alpha)$ values reported in 1962Wa18 have been renormalized to a sum of 100 % by 1970Da08. The uncertainties reported in 1970Da08 are only statistical (from averaging data of three measurements) and comparable with the supposed uncertainties of 1962Wa18, 1962Gi04 and 1957Pi31.

Table 3: Experimental and recommended probabilities (per 100 decays) of alpha-transitions observed in ²²³Ra α decay

	α -particle energy	1957Pi31	1962Gi04	1962Wa18 ^a	1970Da08	UWM	Deduced from $P(\gamma+ce)$ balance	Recommended
$\alpha_{0,0}$	5871	0.96	1.5	0.85	0.85 (4)	1.04 (16)		1.0 (2)^c
$\alpha_{0,2}$	5858	0.3		0.31	0.32 (4)			0.32 (4)^d
$\alpha_{0,3}$	5747	10.5	10.2	8.85 (18) ^b	9.50 (58)	9.8 (4)	10.0 (3)	10.0 (3)^e
$\alpha_{0,4}$	5716	50.4	48.0	52.2 (11) ^b	52.5 (8)	50.8 (10)	49.6 (9)	49.6 (12)^e
$\alpha_{0,5}$	5607	23.6	25.7	25.3 (5) ^b	24.2 (4)	24.7 (5)	25.8 (6)	25.8 (11)^e
$\alpha_{0,6}$	5540	10.3	10.2	8.85 (18) ^b	9.16 (30)	9.6 (4)	10.60 (17)	10.6 (10)^e
$\alpha_{0,8}$	5502	0.86	1.3	0.78	1.00 (15)	0.99 (11)	0.74 (3)	0.74 (25)^e
$\alpha_{0,10}$	5481			≈ 0.008			0.0007 (4)	≈ 0.008
$\alpha_{0,11}+\alpha_{0,12}$	5434	2.4	2.5	2.24	2.27 (20)	2.35 (6)	2.10 (9)	2.10 (25)^e

	α -particle energy	1957Pi31	1962Gi04	1962Wa18 ^a	1970Da08	UWM	Deduced from P(γ +ce) balance	Recommended
$\alpha_{0,14}$	5366	0.20	} Σ 0.25	0.108	\approx 0.13	0.15 (3)	0.014 (7)	\approx 0.13 ^d
$\alpha_{0,16}$	5339	0.07		0.098	\approx 0.13	0.099 (17)	0.089 (6)	\approx 0.13 ^d
$\alpha_{0,17}$	5287	} Σ 0.3	} Σ 0.3	0.126	\approx 0.16		0.16 (4)	0.16 (4) ^e
$\alpha_{0,18}$	5283			0.093				
$\alpha_{0,19}$	5259			0.042			0.079 (8)	0.042
$\alpha_{0,20}$	5236			0.041			0.022 (4)	0.041
$\alpha_{0,21}$	5212			0.0053			0.0011 (6)	0.0053
$\alpha_{0,22}$	5173			0.026			0.013 (4)	0.026
$\alpha_{0,23}$	5152			0.021			0.0134 (27)	0.021
$\alpha_{0,24}$	5135			\approx 0.0017				\approx 0.0017
$\alpha_{0,25}$	5112			\approx 0.0006				\approx 0.0006
$\alpha_{0,26}$	5086			\approx 0.0003				\approx 0.0003
$\alpha_{0,27}$	5056			\approx 0.0002				\approx 0.0002
$\alpha_{0,28}$	5036			\approx 0.0004				\approx 0.0004
$\alpha_{0,29}$	5026			\approx 0.00063				\approx 0.00063
$\alpha_{0,30}$	5014			\approx 0.00044				\approx 0.00044

^a Authors did not report individual uncertainties for intensity of each α -particle group but stated the relative uncertainty of 2 % for intense α -lines and 10 % for weak α -lines.

^b Uncertainty given by Rytz (1991Ry01).

^c Value recommended by Rytz (1991Ry01)

^d Adopted from 1970Da08.

^e Deduced from P(γ +ce) balance. Uncertainties were expanded to cover UWM of experimental P_α values.

The α decay hindrance factors have been calculated using the ALPHAD computer program from the ENSDF evaluation package with $r_0(^{219}\text{Rn}) = 1.543$ fm (2001Br31).

2.2. Gamma Transitions and Internal Conversion Coefficients

The recommended energies of the gamma-ray transitions are the same as those of the gamma-ray energies corrected by the minor nuclear recoil of ²¹⁹Rn.

The gamma-ray transition probabilities (P(γ +ce)) have been deduced from their evaluated gamma-ray emission probabilities (P(γ)) and total internal conversion coefficients (ICCs).

ICCs have been interpolated using the BrIcc computer program, version v2.2a, with the “frozen orbital” approximation (2008Ki07). Multipolarities of the gamma-ray transitions and E2/M1 mixing ratios (δ) are those deduced by 2001Br31, on the basis of measurements of conversion electrons (ce) by 1970Da08, 1970Kr01, 1972HeYM, 1974Ri05, and 1998Sh02.

P(γ +ce) values for the gamma-ray transitions $\gamma_{1,0}$ (4.4 keV), $\gamma_{9,7}$ (34.5 keV) and $\gamma_{22,18}$ (112.6 keV) have been deduced from probability balances at the ²¹⁹Rn ground state (level ‘0’), level ‘7’ (342.8 keV) and level ‘18’ (598.7 keV), respectively.

3. ATOMIC DATA

The fluorescence yields, X-ray energies and relative probabilities, and Auger energies and relative probabilities are from the SAISINUC software.

4. ALPHA EMISSIONS

The recommended energies of alpha particles have been deduced from the Q(α) value, taking into account the recoil energies for ²¹⁹Rn.

The recommended α -particle energies are compared in Table 4 with the experimental results from spectrometric measurements by 1957Pi31, 1961Ry02, 1962Wa18, 1964Wa19, 1970Da08, and 1971Gr17.

Table 4: Experimental and recommended α -particle energies (keV) in the decay of ²²³Ra ^a

	1957Pi31	1961Ry02	1962Wa18 ^b	1964Wa19	1970Da08	1971Gr17	Recommended
$\alpha_{0,0}$	5870 (2)		5871.6 (10)	5869.5 (17)	5871 (3)		5871.63 (21)
$\alpha_{0,2}$	5856		5857.5 (10)		5857 (3)		5857.52 (21)
$\alpha_{0,3}$	5745 (2)	5745.5	5747.4		5747 (3)	5747.0 (4)	5747.14 (21)
$\alpha_{0,4}$	5715	5714.3	5716.1		5715 (3)	5716.23 (29)	5715.84 (21)
$\alpha_{0,5}$	5605	5605.3	5607.1		5606 (3)	5606.73 (30)	5606.99 (21)
$\alpha_{0,6}$	5537	5537.1	5539.6		5537 (3)	5539.8 (9)	5539.43 (21)
$\alpha_{0,8}$	5500		5501.6 (10)		5501 (3)		5502.12 (21)
$\alpha_{0,11}$	5432		5433.6 (5)		5435 (3)		5434.59 (21)
$\alpha_{0,14}$	5363		5365.6 (10)		5367 (3)		5366.37 (23)
$\alpha_{0,16}$	5337		5338.7 (10)		5339 (3)		5339.37 (21)
$\alpha_{0,17}$	5287		5287.3 (10)		5288 (3)		5288.19 (23)

^a Authors' experimental values have been adjusted for changes in calibration energies following 1977Ma31 and 1991Ry01.

^b Uncertainties of 1962Wa18 are the values estimated by 1977Ma31.

5. ELECTRON EMISSIONS

The energies of the conversion electrons have been obtained from the gamma-ray transition energies and the atomic electron binding energies from 1977La19. The emission probabilities of the conversion electrons have been deduced using the evaluated $P(\gamma)$ and ICC values. Measurements of the ²¹⁹Rn conversion electrons were carried out by 1969Be67, 1970Da08, 1970Kr01, 1972HeYM, 1974Ri05, and 1998Sh02.

The total absolute emission probabilities of K and L Auger electrons have been calculated using the EMISSION computer program (1996Sc06, 2000Sc47).

6. PHOTON EMISSIONS

6.1 X - Ray emissions

The absolute emission probabilities of Rn KX- and LX-rays were calculated using the EMISSION computer program (Table 5). In 1976B113 the emission probabilities of Rn KX-rays were measured relatively to $P\gamma$ ($\gamma_{5,0}$ 269.5 keV). The experimental absolute $P(KX)$ values are given in Table 5 using the evaluated $P\gamma$ ($\gamma_{5,0}$ 269.5 keV) = 14.23 (15) %.

Table 5: Experimental (1976B113) and calculated absolute Rn KX-ray emission probabilities (%)

	1976B113	Calculated
$K\alpha_2$ (Rn)	11.5 (13)	14.86 (23)
$K\alpha_1$ (Rn)	19.4 (24)	24.5 (4)
$K\beta'_1$ (Rn)	9.4 (7)	8.50 (18)
$K\beta'_2$ (Rn)	1.71 (14)	2.72 (7)

6.2. Gamma emissions

6.2.1. Gamma-ray energies

The gamma-ray energies (E_γ) for $\gamma_{1,0}$ (4.5 keV), $\gamma_{2,1}$ (9.9 keV), $\gamma_{2,0}$ (14.4 keV), $\gamma_{4,3}$ (31.9 keV), $\gamma_{12,7}$ (104.0 keV), $\gamma_{4,2}$ (144.3 keV), $\gamma_{8,3}$ (249.5 keV), $\gamma_{7,0}$ (342.8 keV), $\gamma_{8,2}$ (361.9 keV), $\gamma_{9,1}$ (372.9 keV), $\gamma_{8,0}$ (376.3 keV), $\gamma_{16,4}$ (383.4 keV), $\gamma_{12,2}$ (432.4 keV), $\gamma_{16,0}$ (542.0 keV), and $\gamma_{19,0}$ (623.7 keV) have been deduced directly from the adopted ²¹⁹Rn level energies.

The remaining gamma-ray energies have been taken mainly from 2001Br31. They are weighted averages of the experimental values from 1998Sh02, 1976B113, 1972HeYM, 1970Da08, 1970Kr01, and 1968Br17, except as specified otherwise in footnotes of Table 6. The most precise measurements of E_γ from 1976B113 with Ge(Li) detector dominate the weighted averages.

Less accurate measurements of E_γ were reported in 1957Pi31, 1957Pa07, 1966Po02, 1969Be67, they were not used in the evaluation.

It should be noted that in 2001Br31 many questionable unplaced gamma-ray transitions are given from some spectrometric measurement results published in the above references, but they have not been yet confirmed by other authors. Observation of such gamma-ray transitions may be assigned to daughters of ²²³Ra or other isotope impurities and most of these gamma-rays have not been included in the current evaluation. The criterion for their inclusion was an observation in α - γ high resolution coincidence with planar and high efficiency coaxial Ge detectors in the latest experiment by 1998Sh02.

6.2.2. Gamma-ray emission probabilities

The experimental and evaluated relative emission probabilities (I_γ) of gamma-rays in decay of ²²³Ra are presented in Table 6. The adopted values are the weighted means of the experimental values except when noted. The statistical processing was done using the LWEIGHT computer program. The uncertainties assigned in this evaluation to the recommended values are always greater than or equal to the smallest uncertainty in any of the experimental values used in the calculation.

The normalization factor (N) was obtained from the probability balance to the ²¹⁹Rn ground state (level '0') and excited levels '1' (4.5 keV) and '2' (14.4 keV):

$$\sum(1+\alpha_T)I_\gamma(\gamma_{i,0}, \gamma_{j,1}, \gamma_{k,2}) + P(\alpha_{0,0}) + P(\alpha_{0,2}) = 1$$

where $i = 4, 5, 6, 7, 8, 11, 16, 17, 18, 19, 20, 22, 23$;

$j = 3, 4, 6, 8, 9, 14, 16, 19, 20, 23$;

$k = 4, 5, 6, 7, 8, 9, 11, 12, 14, 16, 17, 18, 19, 20, 22, 23$.

$$N = P(\gamma)(269.5 \text{ keV}) = 0.1423 (15).$$

This adopted value can be compared with the measured P_γ (269.5 keV) of 0.140 (15) (1968Br17) and the value of 0.136 (10) (1970Da08) deduced from an α -feed of the 269 keV level in ²¹⁹Rn.

The absolute gamma-ray emission probabilities ($P(\gamma)$) have been deduced from the evaluated relative gamma-ray emission probabilities (Table 6) using the derived normalization factor of 0.1423 (15).

$P(\gamma)$ values for the gamma-ray transitions $\gamma_{2,1}$ (9.9 keV) and $\gamma_{2,0}$ (14.4 keV) have been obtained directly from probability balance at the ²¹⁹Rn level '2' (14.4 keV) and the ratio of $P(\gamma_{2,1} - 9.9 \text{ keV})/P(\gamma_{2,0} - 14.4 \text{ keV}) = 0.86 (9)$ deduced from measured ratio of intensities of conversion electrons (1974Ri05) and the ratio of theoretical ICCs.

P_γ value for the gamma-ray transition $\gamma_{1,0}$ (4.5 keV) has been estimated from $P_\gamma + ce$ using the total theoretical ICC α_T .

Table 6: Recommended energies (E_γ) and experimental and evaluated relative emission probabilities (I_γ) of gamma-rays in decay of ²²³Ra

	Recommended E_γ (keV)	1968Br17	1970Kr01	1970Da08	1972HeYM	1976Bl13	1998Sh02	Evaluated I_γ
$\gamma_{1.0}$	4.47 (1)^a							
$\gamma_{2.1}$	9.90 (2)^a							
$\gamma_{2.0}$	14.37 (1)^a							
$\gamma_{4.3}$	31.87 (2)^a		0.000 74 (15)				0.001	0.000 74 (15)
$\gamma_{9.7}$	34.5 (2)^b							
$\gamma_{12.9}$	69.5 (1)^b						0.05 (2)	0.05 (2)
$\gamma_{15.12}$	70.9 (2)^b						0.025 (8)	0.025 (8)
$\gamma_{11.7}$	102.2 (2)^b						0.006 (3)	0.006 (3)
$\gamma_{17.13}$	103.2 (2)^b	0.100 (14) ^e			0.12 (7) ^e		0.04 (2)	0.04 (2)
$\gamma_{12.7}$	104.04 (4)^a					0.134 (15)	0.14 (2)	0.136 (15)
$\gamma_{11.6}$	106.78 (3)	0.164 (29)	0.14 (3)	0.16 (4)	0.19 (6)	0.157 (15)	0.17 (1)	0.164 (10)
$\gamma_{12.6}$	108.5 (2)^b						0.04 (2)	0.04 (2)
$\gamma_{5.4}$	110.856 (10)	0.40 (6)	0.331 (29) ^f	0.41 (4)	0.21 (9) ^f	0.40 (4)	0.42 (3)	0.41 (3)
$\gamma_{22.18}$	112.6^c							
$\gamma_{13.8}$	114.7 (2)				0.07 (4)		0.07 (3)	0.07 (3)
$\gamma_{3.1}$	122.319 (10)	8.2 (11)	8.75 (15)	9.8 (10)	8.7 (4)	7.5 (8)	8.7 (1)	8.70 (10)
$\gamma_{20.14}$	131.6 (2)				0.037 (22)		0.04 (2)	0.04 (2)
$\gamma_{14.8}$	138.3 (3)^b						0.012 (5)	0.012 (5)
$\gamma_{4.2}$	144.27 (2)^a	22.1 (21)	23.8 (5)	23.0 (24)	27.4 (18) ^f	21.6 (22)	23.5 (5)	23.6 (5)
$\gamma_{17.12}$	147.2 (2)^b						0.04 (2)	0.04 (2)
$\gamma_{4.1}$	154.208 (10)	38.6 (29)	41.1 (8)	38 (4)	44.4 (26)	38 (4)	41 (1)	41.0 (8)
$\gamma_{4.0}$	158.635 (10)	5.0 (5)	5.02 (10)	5.6 (6)	5.3 (4)	4.6 (4)	5.0 (1)	5.01 (10)
$\gamma_{16.8}$	165.8 (2)				0.037 (22)		0.04 (2)	0.038 (20)

Comments on evaluation

²²³Ra

	Recommended E _γ (keV)	1968Br17	1970Kr01	1970Da08	1972HeYM	1976Bl13	1998Sh02	Evaluated I _γ
γ _{11.5}	175.65 (15)		0.10 (3)		0.15 (4)		0.14 (3)	0.12 (3)
γ _{12.5}	177.3 (1)	0.21 (7)	0.34 (3)		0.35 (6)		0.34 (3)	0.33 (3)
γ _{6.4}	179.54 (6)	1.07 (29)	1.07 (29)	1.10 (13)	1.16 (15)	1.01 (10)	1.1 (1)	1.08 (10)
γ _{20.12}	199.3 (3)				0.022 (15)		0.02 (1)	0.021 (10)
γ _{18.9}	221.32 (24)	0.25 (7)	0.22 (4)		0.25 (4)		0.26 (4)	0.25 (4)
γ _{19.8}	247.2 (5)				0.066 (22)		0.07 (2)	0.068 (20)
γ _{8.3}	249.49 (3)^a	0.26 (7)			0.29 (13)		0.28 (7)	0.27 (7)
γ _{17.7}	251.6 (3)	0.49 (11)	0.27 (7)	0.42 (15)	0.49 (15)	0.47 (7)	0.3 (1)	0.39 (7)
γ _{5.2}	255.2 (2)	0.43 (11)		0.37 (15)	0.24 (7)	0.33 (7)	0.38 (5)	0.34 (5)
γ _{17.6}	255.7 (3)				0.037 (22)		0.04 (2)	0.039 (20)
γ _{18.6}	260.4 (3)				0.044 (22)		0.05 (2)	0.047 (20)
γ _{5.0}	269.463 (10)	100 (11)	100 (2)	100 (7)	100 (4)	100 (4)	100 (2)	100 (2)
γ _{10.3}	270.3 (4)^b						0.005 (3)	0.005 (3)
γ _{23.12}	286.0 (4)^b						0.008 (4)	0.008 (4)
γ _{12.4}	288.18 (3)	1.14 (14)	1.16 (5)	1.08 (12)	0.93 (13) ^f	1.07 (5)	1.15 (3)	1.13 (3)
γ _{6.2}	323.871 (10)	26.5 (29)	29.4 (6)	26.5 (26)	26.8 (11)	26.8 (13)	28.7 (5)	28.5 (5)
γ _{7.2}	328.38 (3)^a	1.43 (14)	1.52 (7)	1.19 (24)	1.18 (18)	1.40 (8)	1.5 (5)	1.43 (7)
γ _{6.1}	334.01 (6)	0.61 (9)	0.76 (6)	0.91 (18)	0.69 (11)	0.54 (7)	0.73 (4)	0.70 (4)
γ _{6.0}	338.282 (10)	19.3 (18)	21 (5)	19.0 (20)	19.2 (7)	18.5 (9)	20.4 (4)	20.0 (4)
γ _{7.0}	342.78 (2)^a	1.43 (14)	1.70 (9)	1.5 (4)	0.71 (16) ^f	1.49 (12)	1.6 (1)	1.59 (9)
γ _{23.9}	355.5 (2)^b						0.03 (1)	0.03 (1)
γ _{14.4}	355.7 (2)^b						0.02 (1)	0.02 (1)
γ _{8.2}	361.89 (2)^a	0.29 (7)	0.34 (4)	0.37 (7)	0.24 (6)	0.298 (22)	0.20 (5)	0.20 (5)^g
γ _{9.2}	362.9 (2)^b						0.11 (5)	0.11 (5)
γ _{22.7}	368.56 (12)				0.06 (3)		0.06 (3)	0.06 (3)
γ _{8.1}	371.676 (15)	3.9 (4)	3.56 (7)	4.0 (6)	4.2 (4)	3.14 (16)	3.5 (1)	3.51 (7)

Comments on evaluation

²²³Ra

	Recommended E _γ (keV)	1968Br17	1970Kr01	1970Da08	1972HeYM	1976Bl13	1998Sh02	Evaluated I _γ
γ _{9,1}	372.86 (1) ^{a,b}	≈ 0.7				0.73 (8)	0.36	0.36 ⁱ
γ _{8,0}	376.26 (2) ^a			0.088 (29)			0.09 (3)	0.09 (3)
γ _{16,4}	383.35 (2) ^a	≈ 0.04		0.11 (4)	0.029 (22)		-	0.05 (3)
γ _{14,3}	387.7 (2)				0.10 (4)		0.11 (4)	0.11 (4)
γ _{23,7}	390.1 (2)	≈ 0.05			0.022 (15)		0.05 (2)	0.032 (15)
γ _{11,2}	430.6 (3)				0.14 (4)		0.14 (4)	0.14 (4)
γ _{12,2}	432.45 (3) ^a	0.24 (3)	0.26 (4)	≈0.22	0.24 (6)	0.186 (30) ^f	0.25 (2)	0.25 (2)
γ _{11,0}	445.033 (12)	8.7 (4)	11.0 (8) ^f	9.3 (10)	9.2 (7)	8.5 (4)	9.3 (3)	9.0 (3)
γ _{20,4}	487.5 (2)	0.071 (14)	0.10 (4)	≈0.11	0.08 (4)		0.08 (1)	0.08 (1)
γ _{-1,1}	490.8 (3) ^b						0.012 (5)	0.012 (5)
γ _{14,2}	500.0 (4) ^b						0.010 (4)	0.010 (4)
γ _{14,1}	510.0 (4) ^b						0.003 (2)	0.003 (2)
γ _{-1,2}	523.2 (4) ^b						0.010 (4)	0.010 (4)
γ _{16,2}	527.611 (13)	0.50 (5)	0.54 (5)	0.51 (10)	0.47 (11)	0.410 (22) ^f	0.51 (3)	0.51 (3)
γ _{-1,3}	532.9 (4) ^b						0.010 (4)	0.010 (4)
γ _{16,1}	537.6 (1) ^b						0.015 (5)	0.015 (5)
γ _{16,0}	541.99 (2) ^{a,b}						0.010 (4)	0.010 (4)
γ _{21,3}	545.8 (5) ^b						0.008 (4)	0.008 (4)
γ _{23,4}	574.1 (7) ^b						0.008 (4)	0.008 (4)
γ _{17,2}	579.6 (3) ^b						0.010 (4)	0.010 (4)
γ _{18,2}	584.3 (3) ^b						0.010 (4)	0.010 (4)
γ _{17,0}	594.0 (3) ^b						0.010 (4)	0.010 (4)
γ _{18,0}	598.721 (24)	0.57 (6)	0.76 (7)	0.68 (11)	0.66 (13)	0.626 (30)	0.68 (3)	0.65 (3)
γ _{19,2}	609.31 (4)	0.36 (4)	0.54 (7)	0.46 (7)	0.30 (11)	0.373 (22)	0.41 (2)	0.40 (2)
γ _{19,1}	619.1 (4) ^b						0.025 (8)	0.025 (8)
γ _{19,0}	623.68 (4) ^a	0.057 (29)					0.06 (3)	0.06 (3)

Comments on evaluation

²²³Ra

	Recommended E _γ (keV)	1968Br17	1970Kr01	1970Da08	1972HeYM	1976Bl13	1998Sh02	Evaluated I _γ
γ _{20,2}	631.7 (7)			0.22 (7)			0.003 (2)	0.003 (2)
γ _{20,1}	641.7 (4)^b						0.012 (5)	0.012 (5)
γ _{20,0}	646.1 (5)^b						0.003 (3)	0.003 (3)
γ _{22,2}	696.9 (7)^b						0.005 (2)	0.005 (2)
γ _{22,0}	711.3 (2)^b	0.025 (7)					0.026 (7)	0.026 (7)
γ _{23,2}	718.4 (4)^b						0.010 (4)	0.010 (4)
γ _{23,1}	728.4 (8)^b						0.002 (1)	0.002 (1)
γ _{23,0}	732.8 (6)^b						0.004 (2)	0.004 (2)
γ _{-1,4}	737.2 (8)^b						0.002 (1)	0.002 (1)

^a From the adopted ²¹⁹Rn level energies.

^b From 1998Sh02; new gamma-ray transition observed.

^c Reported only by 1998Sh02 without uncertainty in energy and without intensity value.

^d Not reported by 1998Sh02 but observed in 1968Br37, 1970Da08, 1972HeYM, 1976Bl13.

^e Reported γ-ray with energy of 103.7 keV that may be a sum of 103.2 keV and 104.0 keV γ-rays.

^f Omitted on Chauvenet's criterion.

^g Adopted from 1998Sh02 because of possible contribution of impurity Pb γ-rays in measurements of single γ-spectra.

^h Adopted from 1998Sh02 because of contribution of unplaced 373.3 keV γ-ray observed in measurements of single γ-spectra and not observed in α-γ coincidences.

7. REFERENCES

- 1954Ha60 G.R. Hagee, M.L. Curtis, G.R. Grove, Phys. Rev. 96, 817A (1954) (Half-life)
- 1957Pa07 H. Paul, H. Warhanek, Helv. Phys. Acta 30, 272 (1957) (γ -ray energies and emission probabilities)
- 1957Pi31 R.C. Pilger, Jr., Thesis, Univ. California (1957); UCRL-3877 (1957) (α -particle and γ -ray energies and emission probabilities)
- 1959Ro51 J. Robert, Ann. Phys. (Paris) 4, 89 (1959) (Half-life)
- 1961Ry02 A. Rytz, Helv. Phys. Acta 34, 240 (1961) (α -particle energies and emission probabilities)
- 1962Wa18 R.J. Walen, V. Nedovesov, G. Bastin-Scoffier, Nuclear Phys. 35, 232 (1962) (α -particle energies and emission probabilities)
- 1962Gi04 M. Giannini, D. Prospero, S. Sciuti, Nuovo Cimento 25, 1314 (1962) (α -particle energies and emission probabilities)
- 1964Wa19 A.H. Wapstra, Nucl. Phys. 57, 48 (1964) (α -particle energies and emission probabilities)
- 1965Ki05 H.W. Kirby, K.C. Jordan, J.Z. Braun, M.L. Curtis, M.L. Salutsky, J. Inorg. Nucl. Chem. 27, 1881 (1965) (Half-life)
- 1966Po02 P. Polak, A.H. Wapstra, C. Ythier, Priv. Comm. (1966). (γ -ray energies and emission probabilities)
- 1967JoZX K.C. Jordan, B.C. Blanke, Proc. Symp. Standardization of Radionuclides, Vienna, Austria (1966), Intern. At. Energy Agency, Vienna, p.567 (1967); CONF-661012-4 (1967) (Half-life)
- 1968Br17 C. Briançon, C.F. Leang, R. Walen, Compt. Rend. 266B, 1533 (1968) (γ -ray energies and emission probabilities)
- 1969Be67 D. Bertault, M. Vidal, G.Y. Petit, J. Phys.(Paris) 30, 909 (1969) (Conversion electron spectra, 269 keV γ -ray multipolarity)
- 1970Da08 W.F. Davidson, R.D. Connor, Nucl. Phys. A149, 363 (1970) (α -particle and γ -ray energies and emission probabilities, E2/M1 mixing ratios)
- 1970Kr01 K. Krien, C. Gunther, J.D. Bowman, B. Klemme, Nucl. Phys. A141, 75 (1970) (γ -ray energies and emission probabilities, E2/M1 mixing ratios)
- 1971Gr17 B. Grennberg, A. Rytz, Metrologia 7, 65 (1971) (α -particle energies and emission probabilities)
- 1972HeYM W.H.A. Hesselink, NP-19781 (1972) (γ -ray energies and emission probabilities)
- 1974Ri05 B. Richter, M.J. Canty, L. Ley, M.V. Banaschik, A. Neskakis, Nucl. Phys. A223, 234 (1974) (Conversion electron spectra, E2/M1 mixing ratios)
- 1976B113 K. Blaton-Albicka, B. Kotlinska-Filipek, M. Matul, K. Stryczniewicz, M. Nowicki, E. Ruchowska-Lukasiak, Nukleonika 21, 935 (1976) (γ -ray energies and emission probabilities)

- 1977Ma31 C. Maples, Nucl. Data Sheets 22, 243 (1977) (α -particle energies and emission probabilities)
- 1977La19 F.P. Larkins, At. Data Nucl. Data Tables 20, 311 (1977) (Atomic electron binding energies)
- 1987Mi10 G.J. Miller, J.C. McGeorge, I. Anthony, R.O. Owens, Phys. Rev. C36, 420 (1987) (Half-life)
- 1991Ma16 M.J. Martin, Nucl. Data Sheets 63, 723 (1991) (Branch of ²²³Ra decay by emission of ¹⁴C)
- 1991Ry01 A. Rytz, At. Data Nucl. Data Tables 47, 205 (1991) (α -particle energies and emission probabilities)
- 1996Sc06 E. Schönfeld, H. Janssen, Nucl. Instrum. Methods Phys. Res. A369, 527 (1996) (Atomic data)
- 1998Sh02 R.K. Sheline, C.F. Liang, P. Paris, Phys.Rev. C57, 104 (1998) (γ -ray energies and emission probabilities)
- 2000Sc47 E. Schönfeld, H. Janssen, Appl. Radiat. Isot. 52, 595 (2000) (Calculation of emission probabilities of X-rays and Auger electrons)
- 2001Br31 E. Browne, Nucl.Data Sheets 93, 763 (2001); Erratum Nucl. Data Sheets 96, 391 (2002) (²²³Ra α decay scheme and α decay data evaluation)
- 2003Au03 G. Audi, A.H. Wapstra, and C. Thibault, Nucl. Phys. A729, 337 (2003) (Q value)
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr., Nucl. Instrum. Methods Phys. Res. A589, 202 (2008) (Band-Raman ICC for γ -ray transitions)

**²²⁴Ra – Comments on evaluation of decay data
by A. L. Nichols**

Evaluated: July/August 2001

Re-evaluated: January 2004

Evaluation Procedures

Limitation of Relative Statistical Weight Method (LWM) was applied to average numbers throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

A relatively simple decay scheme was constructed from the alpha -particle studies of 1962Wa28, 1969Pe17, 1971So15 and 1984Bo15, and the gamma -ray measurements of 1969Pe17, 1972DaZA, 1977Ku15, 1982Sa36, 1983Sc13, 1983Va22, 1984Bo15, 1984Ge07 and 1992Li05. Only the gamma -ray studies of 1977Ku15 provide any detail beyond the 240.986 keV gamma ray; all other measurements are dedicated to the determination of the absolute emission probability of the 240.986 keV gamma ray. A weighted mean emission probability was determined for this transition, and the other emission probabilities as measured by 1977Ku15 were subsequently adjusted.

Cluster decay has been observed by 1985Pr01 and 1991Ho15, and reviewed by 1995Ar33 and 1997Tr17. ¹⁴C emissions were detected with a branching fraction of 5(1)E -11. However, this decay mode has not been included in the decay-data summary section.

Nuclear Data

²²⁸Th decay chain is important in quantifying the environmental impact of the decay of naturally occurring ²³²Th. Specific radionuclides in this decay chain are noteworthy because of their decay characteristics (²²⁴Ra alpha decay to ²²⁰Rn; ²¹²Bi and ²⁰⁸Tl gamma-ray emissions).

Half-life

The recommended half-life represents the unweighted mean of two somewhat elderly studies (1962Li02 and 1971Jo14) and a much more recent measurement (2004ScZZ). Further measurements are required to determine this half-life with greater confidence.

Reference	Half-life (d)
1962Li02	3.62(1)
1971Jo14	3.665(38)
2004ScZZ	3.6319(23)
Recommended Value	3.627(7)

Gamma Rays

Energies

All gamma-ray transition energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies of 1997Ar04 were adopted, and used to determine the energies and associated uncertainties of the gamma-ray transitions between the various populated-depopulated levels.

Emission Probabilities

Absolute emission probabilities were determined from measurements of the 240.986 keV gamma ray by 1969Pe17, 1972DaZA, 1982Sa36, 1983Sc13, 1983Va22, 1984Bo15, 1984Ge07 and 1992Li05. A weighted mean value of 4.12(3)% was derived through LWEIGHT, and the uncertainty was increased slightly to the lowest measured value of ± 0.04 to give 4.12(4)% (0.0412(4)).

Only 1977Ku15 has measured the emission probabilities of other low -intensity gamma transitions identified with ²²⁴Ra alpha decay; these data are reported relative to a value of 39500(1300) for the 240.986 keV gamma emission, as taken from 1969Pe17. Hence, the low -intensity emission probabilities have been subsequently adjusted on the basis of $P_\gamma(240.986 \text{ keV})$ of 4.12(4)% (0.0412(4)).

Absolute Gamma-ray Emission Probabilities per 100 Disintegrations of ²²⁴Ra

$E_g \text{ (keV)}$	P_g^{abs}				
	1969Pe17	1972DaZA [‡]	1977Ku15 [†]	1982Sa36	1983Sc13
240.986(6)	3.95(13)	3.9(7)	[3.95(13) →	3.9(2)	4.04(17)
292.70(11)	-	-	4.12(4)]	-	-
404.5(1)	-	-	0.0063(7)	-	-
422.04(11)	-	-	0.0022(5)	-	-
645.44(9)	-	~ 0.007	0.0030(5)	-	-
			0.0054(9)		

$E_g \text{ (keV)}$	P_g^{abs} (cont.)				
	1983Va22	1984Bo15	1984Ge07	1992Li05	Recommended Values [*]
240.986(6)	4.05(9)	4.05(9)	4.17(4)	4.11(12)	4.12(4)
292.70(11)	-	-	-	-	0.0063(7)
404.5(1)	-	-	-	-	0.0022(5)
422.04(11)	-	-	-	-	0.0030(5)
645.44(9)	-	-	-	-	0.0054(9)

[‡] Data expressed relative to $P_\gamma(2614.511 \text{ keV})$ of ²⁰⁸Tl have been adjusted.

[†] Data adjusted on the basis of $P_\gamma(240.986 \text{ keV})$ of 4.12(4)%.

^{*} Recommended gamma-ray emission probabilities above 241 keV taken from adjusted data of 1977Ku15.

Multipolarities and Internal Conversion Coefficients

The nuclear level scheme specified by 1997Ar04 has been used to define the multipolarities of the gamma transitions on the basis of known spins and parities. Recommended internal conversion coefficients have been interpolated from the theoretical tabulations of 1978Ro22.

Alpha-particle EmissionsEnergies

All alpha-particle energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies of 1997Ar04 and the Q-value of 1995Au04 were used to determine the energies and uncertainties of the alpha -particle transitions to the various levels, while allowing for the significant recoil components.

Emission Probabilities

Alpha-particle emission probabilities to the first excited states of ²²⁰Rn have been directly measured by 1969Pe17, 1971So15, 1984Bo15 and 1993Ba72, and these data can be used to calculate the alpha-particle emission probability directly to the ground state of ²²⁰Rn:

Alpha-particle emission probability data of 1969Pe17 are effectively normalised to 94.95(5)% and 5.05(5)%, similarly for the equivalent data of 1971So15, with normalised values of 95.1(4)% and 4.9(4)%, and 1984Bo15, with normalised values of 94.94(4)% and 5.06(4)%.

1993Ba72: two alpha -particle emissions are quantified that sum to 100.03%, and the two associated uncertainties are effectively inconsistent; data adjusted so that uncertainties correspond ($\pm 0.04\%$) to give:

Comments on evaluation

$P_{\alpha}(5685.50 \text{ keV})$ of 95.10%, and uncertainty of $\pm 0.04\%$;
and $P_{\alpha}(5448.81 \text{ keV})$ of 4.93%, and uncertainty of $\pm 0.04\%$.

A weighted mean value of 95.00(4)% (0.9500(4)) can be determined for $P_{\alpha}(5685.50 \text{ keV})$, and matched with a value of 5.01(4)% (0.0501(4)) for $P_{\alpha}(5448.81 \text{ keV})$. Thus, a discrepancy exists between measurements of the absolute emission probability of the 240.986 keV gamma ray and measurements of the direct alpha-particle emission probability to the ground state of Rn-220:

(i) assuming that the measured gamma -ray emission probabilities are absolute (as quoted in the various references) and $P_{\gamma}(240.986 \text{ keV})$ is 0.0412(4), $NF = 1.000$, $P_{\alpha}(5685.50 \text{ keV})$ of 0.9472(7) can be calculated taking into account the low -intensity gamma-ray transition probabilities populating the 240.986 keV nuclear level:

$$P_{\alpha}(5448.81 \text{ keV}) = P_{\gamma}(240.986 \text{ keV})(1 + \alpha_{\text{tot}}(240.986 \text{ keV})) - [\sum P_{\gamma_i} (1 + \alpha_i) \text{ populating nuclear level}] = [0.0412(4) \times 1.280(8)] - 0.000125(18) = 0.0526(7)$$

and $P_{\alpha}(5685.50 \text{ keV}) = 0.9472(7)$

(ii) if gamma -ray emission probabilities are judged to be not strictly absolute and $P_{\alpha}(5685.50 \text{ keV})$ of 0.9500(4) is adopted as the weighted mean of the alpha -particle measurements, $NF = 0.947(8)$ and $P_{\gamma}(240.986 \text{ keV})$ is 0.0390(3).

Although this problem cannot be resolved on the basis of the known measurements, the gamma -ray data were judged to be more reliable. Therefore, the recommended alpha -particle emission probabilities were determined from the gamma -ray data and theoretical internal conversion coefficients, rather than available alpha-particle measurements. These calculations resulted in an absolute emission probability of 0.0526(7) for the 5448.81 keV alpha particle (compared with a weighted mean value of 0.0501(4) from the alpha-particle measurements), and 0.9472(7) for the 5685.50 keV alpha particle. Further spectroscopic measurements are required to resolve the discrepancies between the alpha -particle and gamma-ray data (ie., decay-data studies involving the 240.986 keV and ground states of ²²⁰Rn).

Alpha-particle Emission Probabilities per 100 Disintegrations of ²²⁴Ra

$E_{\alpha}(\text{keV})$	P_{α}							
	1953As31	1962Wa28	1969Pe17	1971So15	1977Ku15 [#]	1984Bo15	1993Ba72	Recommended Values [*]
5034.31(25)	-	0.0031	-	-	0.0029(5)	-	-	0.0030(5)
5051.58(24)	-	0.0072	-	-	0.0073(10)	-	-	0.0076(14)
5161.34(25)	-	0.0073	-	-	0.0069(8)	-	-	0.0074(8)
5448.81(16)	4.9	5.5	5.05(5)	4.9(4)	[5.0(16)]	5.06(4)	[4.93(4)] [¶]	5.26(7)
5685.50(15)	95.1	94	[94.95(5)]	95.1(4)	94.98(16)	[94.94(4)]	[95.10(4)] [¶]	94.72(7)

[#] Data were deduced from gamma-ray studies.

[¶] Relative data are quoted as 4.93(3) and 95.1(6), and have been adjusted to give consistent uncertainties.

^{*} Recommended emission probabilities derived from evaluated gamma -ray emission probabilities and theoretical internal conversion coefficients.

Atomic Data

The x-ray data have been calculated using the evaluated gamma-ray data, and the atomic data from 1996Sc06, 1998ScZM and 1999ScZX.

References

1953As31 - F. Asaro, F. Stephens and I. Perlman, Complex Alpha Spectra of Radiothorium (Th ²²⁸) and Thorium-X (Ra²²⁴), Phys. Rev. 92(1953)1495. [P_{α}]
 1962Ll02 - R. D. Lloyd, C. W. Mays, D. R. Atherton and D. O. Clark, The Half -Period of Ra²²⁴ (Thorium X), COO 225(1962)88. [Half-life]
 1962Wa28 - R. J. Walen, Spectrographie α du Radium 224 et de ses Dérivés, C. R. Acad. Sci. Paris 255(1962)1604. [P_{α}]

- 1969Pe17 - A. Peghaire, Mesures Precises d'Intensities Absolues de Rayonnements γ pour des Emetteurs α , Nucl. Instrum. Meth. 75(1969)66. [P_α , P_γ]
- 1971Jo14 - K. C. Jordan, G. W. Otto and R. P. Ratay, Calorimetric Determination of the Half -Lives of ^{228}Th and ^{224}Ra , J. Inorg. Nucl. Chem. 33(1971)1215. [Half-life]
- 1971So15 - J. C. Soares, J. P. Ribeiro, A. Gonçalves, F. B. Gil and J. G. Ferreira, Sur les Intensités Relatives et Quelques Énergies des Spectres α de ^{238}Pu , ^{232}U et ^{224}Ra , C. R. Acad. Sci. Paris 273B(1971)985. [P_α]
- 1972DaZA - J. Dalmasso, Recherches sur le Rayonnement Gamma de Quelques Radioelements Naturels Appartenant la Famille du Thorium, PhD thesis, University of Nice (1972); J. Dalmasso, H. Maria and C. Ythier, Étude du Rayonnement γ du Thorium 228 et de ses Dérivés, et plus Particulièrement du Thallium 208 (Th C"), C. R. Acad. Sci. Paris 277B(1973)467. [P_γ]
- 1977Ku15 - W. Kurcewicz, N. Kaffrell, N. Trautmann, A. Plochocki, J. Zylicz, M. Matul and K. Stryczniewicz, Collective States Fed by Weak α -transitions in the ^{232}U Chain, Nucl. Phys. A289(1977)1. [P_γ]
- 1978Ro22 - F. Rösel, H. M. Fries, K. Alder and H. C. Pauli, Internal Conversion Coefficients for all Atomic Shells, ICC Values for $Z = 68-104$, At. Data Nucl. Data Tables 21(1978)291-514. [ICC]
- 1982Sa36 - S. Sadasivan and V. M. Raghunath, Intensities of Gamma Rays in the ^{232}Th Decay Chain, Nucl. Instrum. Meth. 196(1982)561. [P_γ]
- 1983Sc13 - U. Schötzg and K. Debertin, Photon Emission Probabilities per Decay of ^{226}Ra and ^{232}Th in Equilibrium with their Daughter Products, Int. J. Appl. Radiat. Isot. 34(1983)533. [P_γ]
- 1983Va22 - R. Vaninbroux and H. H. Hansen, Determination of γ -ray Emission Probabilities in the Decay of ^{228}Th and its Daughters, Int. J. Appl. Radiat. Isot. 34(1983)1395. [P_γ]
- 1984Bo15 - G. Bortels, D. Reher and R. Vaninbroux, Emission Probabilities for the 5.449 -MeV Alpha Particles and 241 keV Gamma Rays in ^{224}Ra - ^{220}Rn Decay, Int. J. Appl. Radiat. Isot. 35(1984)305. [P_α , P_γ]
- 1984Ge07 - R. J. Gehrke, V. J. Novick and J. D. Baker, γ -ray Emission Probabilities for the ^{232}U Decay Chain, Int. J. Appl. Radiat. Isot. 35(1984)581. [P_γ]
- 1985Pr01 - P. B. Price, J. D. Stevenson, S. W. Barwick and H. L. Ravn, Discovery of Radioactive Decay of ^{222}Ra and ^{224}Ra by ^{14}C Emission, Phys. Rev. Letts. 54(1985)297. [Cluster decay]
- 1991Ho15 - E. Hourani, L. Rosier, G. Berrier, Ronsin, A. Elayi, A. C. Mueller, G. Rappenecker, G. Rotbard, G. Renou, A. Lièbe, L. Stab and H. L. Ravn, Fine Structure in ^{14}C Emission from ^{223}Ra and ^{224}Ra , Phys. Rev. C44(1991)1424. [Cluster decay]
- 1992Li05 - W-J. Lin and G. Harbottle, Gamma -ray Emission Intensities of the ^{232}Th Chain in Secular Equilibrium of ^{235}U and the Progeny of ^{238}U , J. Radioanal. Nucl. Chem. 157(1992)367. [P_γ]
- 1993Ba72 - T. Babeliowsky and G. Bortels, ALFA: A Program for Accurate Analysis of Complex Alpha -particle Spectra on a PC, Appl. Radiat. Isot. 44(1993)1349. [P_α]
- 1995Ar33 - G. Ardisson and M. Hussonnois, Radiochemical Investigations of Cluster Radioactivities, Radiochim. Acta 70/71(1995)123. [Cluster decay]
- 1995Au04 - G. Audi and A. H. Wapstra, The 1995 Update to the Atomic Mass Evaluation, Nucl. Phys. A595(1995)409.[Q value]
- 1996Sc06 - E. Schönfeld and H. Janßen, Evaluation of Atomic Shell Data, Nucl. Instrum. Meth. Phys. Res. A369(1996)527. [X_K , X_L , Auger electrons]
- 1997Ar04 - A. Artina-Cohen, Nuclear Data Sheets for $A = 216, 220$, Nucl. Data Sheets 80(1997)157. [Nuclear structure, Energies]
- 1997Tr17 - S. P. Tretyakova and V. L. Mikheev, Experimental Investigation of the Cluster Radioactivity of Atomic Nuclei, Nuovo Cimento 110(1997)1043. [Cluster decay]
- 1998ScZM - E. Schönfeld and G. Rodloff, Tables of the Energies of K -Auger Electrons for Elements with Atomic Numbers in the Range from $Z = 11$ to $Z = 100$, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX - E. Schönfeld and G. Rodloff, Energies and Relative Emission Probabilities of K X-rays for Elements with Atomic Numbers in the Range from $Z = 5$ to $Z = 100$, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2004ScZZ - H. Schrader, Half -life Measurements with Ionization Chambers: A Study of Systematic Effects and Results, Appl. Radiat. Isot. 60(2004)317. [Half-life]

²²⁵Ra - Comments on evaluation of decay data by Huang Xiaolong and Wang Baosong

This evaluation was completed in 2007. Literature available by May 2007 was included.

1 Decay Scheme

²²⁵Ra disintegrates 100 % by β^- emission to levels in ²²⁵Ac. ²²⁵Ra ground state has $J^\pi = 1/2^+$ (1990Ak03).

The recommended $Q(\beta^-)$ value of 356 (5) keV in Audi (2003Au03) agrees with the $Q(\beta^-)$ value of 353 (8) keV, calculated by the evaluator (using program RADLST) from average radiation energies. This agreement supports the completeness and correctness of the decay scheme.

2 Nuclear Data

The $Q(\beta^-)$ value is from the mass adjustment in 2003Au03.

Level energies, spin and parities are from 1990Ak03.

The measured and recommended ²²⁵Ra half-life values are listed in Table 1.

Table 1: Measured half-life values of ²²⁵Ra and recommended value.

$T_{1/2}$ (d)	References	Measurement method
14	1947En03	
14.8 (2)	1950Ha52	Alpha pulse analyser, 10 $T_{1/2}$
15.02 (56)	1987Mi10	Solid-state detector, linear least squares fit
14.91 (11)		Unweighted mean
14.82 (19)		Weighted mean, $\chi^2=0.14$
14.82 (19)	Recommended value	From weighted mean

The half-life weighted average has been calculated using the LWM computer program. The recommended half-life is from LWM result. Further measurements are needed to determine this value with greater precision.

2.1 β^- Transitions

The maximum energies of the β^- transitions in the decay of ²²⁵Ra have been deduced from the $Q(\beta^-)$ value (2003Au03) and the level energies.

The adopted β^- transition probabilities and their associated uncertainties to the 40-keV level and to the ground state were deduced from $P(\gamma) = 30.0 (7) \%$ and $\alpha_T = 1.293 (19)$ for the 40-keV γ -ray. No β^- transitions to the 120.8- and 155.6- keV levels were observed. Based on Ac KX-ray intensities an upper limit of $< 0.01 \%$ for the respective β^- transitions to these levels was reported in 1984Ah01.

The $\log ft$ values and average β^- energies have been calculated with the program LOGFT.

2.2 γ Transitions

The transition probability of the 40-keV γ -ray was calculated using its γ -ray emission intensity and the relevant total internal conversion coefficient.

The multipolarity of this γ -ray transition is from 1990Ak03.

The internal conversion coefficient (ICC) (and its associated uncertainty) for the 40-keV γ -ray transition has been interpolated from theoretical values based on the “Frozen Orbital” approximation (2002Ba85) using the BrIcc computer program (2008Ki07).

3 Atomic Data

Atomic fluorescence yields ($\omega_K, \omega_L, \omega_M, \eta_{KL}$ and η_{LM}) are from Schönfeld (1996Sc06).

The X-ray and Auger electron emission probabilities have been deduced from γ -ray and conversion electron data by using the computer code RADLST.

4 Electron emissions

The conversion electron emission probabilities have been deduced from γ -ray transition data using theoretical internal conversion coefficients.

5 Photon emissions

5.1 γ -ray energy

Measurements of the 40-keV γ -ray energy from ²²⁵Ra are listed in Table 2 together with their weighted mean value. The recommended value is from the weighted mean value.

Table 2: Measured and recommended γ -ray energy from ²²⁵Ra β^- decay.

1955Ma61	1955St04	1981Di14	1987Ah05	LWM	Evaluation
41 (2)	40 (1)	40.12 (5)	40.09 (5)	40.11 (4)	40.11 (4)

5.2 Absolute values of the γ -ray emission probability

The measurements of the absolute γ -ray emission probabilities from ²²⁵Ra decay are listed in Table 3. The present recommended value is taken from a precise measurement in equilibrium with ²²⁹Th (1986He06).

Table 3: Measured and recommended absolute γ -ray emission probability of 40.09keV for ²²⁵Ra.

P_γ (40.09 keV) (%)	References	Measurement method
33	1955Ma61	Scintillation spectrometry
29	1955St04	
39.3 (12)	1981Di14	Ge(Li)
30.0 (7)	1986He06	Ge(Li) and Au-Si surface barrier, in equilibrium with ²²⁹ Th
30.0 (7)		Recommended value from 1986He06

6 References

- 1947En03 A. C. English, T. E. Cranshaw, P. Demers, J. A. Harvey, E. P. Hincks, J. V. Jelley, A. N. May, *Phys. Rev.* 72, 253(1947) [$T_{1/2}$].
- 1950Ha52 F. Hagemann, L. I. Katzin, M. H. Studier, G. T. Seaborg, A. Ghiorso, *Phys. Rev.* 79, 435 (1950) [$T_{1/2}$].
- 1955Ma61 L. B. Magnusson, F. Wagner, Jr., D. W. Engelkemeir, M. S. Freedman, ANL-5386 (1955) [E_γ , P_γ].
- 1955St04 F. S. Stephens, UCRL-2970 (1955) [E_γ , P_γ].
- 1981Di14 J. K. Dickens, J. W. McConnell, *Radiochem. Radioanal. Lett.* 47, 331 (1981) [E_γ , P_γ].
- 1984Ah01 I. Ahmad, R.R. Chasman, J.E. Gindler, A.M. Friedman, *Phys. Rev. Lett.* 52, 503 (1984) [Ac KX-ray].
- 1986He06 R. G. Helmer, C. W. Reich, M. A. Lee, I. Ahmad, *Int. J. Appl. Radiat. Isotop.* 37, 139(1986) [P_γ].
- 1987Ah05 I. Ahmad, J. E. Gindler, A. M. Friedman, R. R. Chasman, T. Ishii, *Nucl. Phys.* A472, 285 (1987) [E_γ].
- 1987Mi10 G. J. Miller, J. C. McGeorge, I. Anthony, R. O. Owens, *Phys. Rev.* C36, 420 (1987) [$T_{1/2}$].
- 1990Ak03 Y. A. Akovali, *Nucl. Data Sheets* 60, 617 (1990) [Level energies, spin and parity].
- 1996Sc06 E. Schönfeld, H. Janssen, *Nucl. Instrum. Meth. Phys. Res.* A369, 527 (1996) [Atomic data].
- 2002Ba85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, *At. Data Nucl. Data Tables* 81, 1 (2002) [Calculated ICC]
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, *Nucl. Phys.* A729, 129(2003)[Q].
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr. , *Nucl. Instrum. Meth. Phys. Res.* A589, 202(2008) [Theoretical ICC].

²²⁵Ac - Comments on evaluation of the decay data

Huang Xiaolong, Wang Baosong

This evaluation was completed in 2008. Literature available by December 2008 was included.

1 Decay Scheme

²²⁵Ac disintegrates 100 % by α emission to levels in ²²¹Fr. ²²⁵Ac ground state has $J^\pi=(3/2^-)$ (1990Ak03).

The ²²⁵Ac α decay scheme was built from the experimental conversion-electron data of 1971DzZP, 1972Dz14 and 2000Ar23, the α - γ coincidence data of 2003Ku44, the γ - γ coincidence data of 1990Ko14, and the experimental singles γ -rays data of 2000Ar23 and 2003Ku44.

The recommended $Q(\alpha)$ value of 5935.1 (14) keV in Audi (2003Au03) agrees with the $Q(\alpha)$ value of 5932.5 (16) keV, calculated by the evaluator (using program RADLST) from average radiation energies. This agreement supports the completeness and correctness of the decay scheme.

2 Nuclear Data

The Q value is from the mass adjustment in 2003Au03.

Level energies have been obtained from a least-squares fit to γ -ray energies (GTOL computer code). Spin and parities are from 1990Ak03, 2000Ar23 and 2003Ku44.

The measured and recommended ²²⁵Ac half-life values are listed in Table 1.

Table 1: Measured half-life values of ²²⁵Ac and recommended value, in days.

$T_{1/2}$ (d)	References	Measurement method
10	1947En03	
10.0 (1)	1950Ha52	Alpha pulse analyzer, 10 $T_{1/2}$
10.0 (1)	Recommended value	From 1950Ha52

The recommended value is taken from the measurement of 1950Ha52. Further measurements are merited to determine this value with greater confidence.

2.1 γ Transitions

The γ -ray transition probabilities were calculated using the γ -ray emission intensities and the relevant internal conversion coefficients.

Multipolarities and mixing ratios of γ -ray transitions are from 1971DzZP, 1972Dz14, 1977Vy02, 1990ArZZ and 2003Ku44. The multipolarity marked in square brackets for other γ transition are from the level scheme (they are not measured).

The internal conversion coefficients (ICC) and their associated uncertainties for γ -ray transitions have been obtained using the BrIcc computer program (2008Ki07), which uses the ‘‘Frozen Orbital’’ approximation (2002Ba85). Experimental and theoretical conversion coefficients are compared in Table 2.

Table 2: Comparison of theoretical and measured conversion coefficients.

E _γ (keV)	Multipolarity	α(theory)	α(exp.)
			(2003Ku44)
78.8	M1	α _T = 5.63, α _L = 4.27	α _T = 5.1 (11)
87.41	M1	α _T = 4.16, α _L = 3.16	α _T = 2.8 (6)
114	M1	α _T = 9.86, α _L = 7.93	α _T = 13.0 (17)
139.6	M1+E2	α _T = 3.9, α _K = 2.4	α _T = 3.2 (5)
145.16	(E1)	α _T = 0.191	α _T ≤ 0.1
153.92	E1	α _T = 0.166	α _T ≤ 0.35
197.5	E1	α _T = 0.0908	α _T ≤ 0.04

2.2 α Transitions

The level energies of ²²¹Fr are determined from the least-squares fit to the recommended γ-ray energies. The level energies of ²²¹Fr and Q-values (2003Au03) were used to determine the energies and uncertainties of the alpha particle transitions to the various levels.

The recommended energies of alpha particles were calculated from the proposed decay scheme and listed in table 3. The recommended values are in good agreement with the measurements of 1967Dz02. Other measurements are 1964Gr11, 1967Ba51, and 1972Go29.

Table 3: Measured and recommended value of α-particle energy for ²²⁵Ac (keV).

1964Gr11	1967Ba51 ^a	1967Dz02 ^b	1972Go29	1991Ry01	Recommended
5829 (5)	5829 (2)	5829 (2)		5829.6 (14)	5829.6 (14)
	5804 (2)				5804.2 (14)
5792 (5)	5793 (3)	5792 (3)	5792.5 (22)	5793.1 (21)	5793.1 (21)
	5791 (4)		5790.6 (22)		5791.7 (14)
5732 (5)	5731 (2)	5731 (3)		5731.9 (17)	5731.9 (17)
					5731.6 (14)
					5730.5 (14)
5724 (5)	5722.6 (25)	5723 (3)			5723.1 (14)
					5686.4 (14)
5683 (5)	5681 (2)	5681 (3)			5682.2 (14)
5638 (5)	5636.2 (20)	5637 (3)			5637.3 (14)
5610 (5)	5607.6 (30)	5608 (3)			5609.0 (14)
	5597.5 (40)				5599.3 (14)
5581 (5)	5579.1 (30)	5577 (3)			5580.5 (14)
	5561.6 (40)				5563.3 (14)
	5552.6 (40)				5555.3 (14)
	5544.1 (40)				5546.5 (14)
	5538.5 (40)	5540 (5)			5540.1 (14)
	5521.5 (70)	5526 (5)			5523.7 (14)
	5514.5 (7)	(5519)			5515.2 (14)
5494 (10)		5497 (4)			5497.4 (14)
		5489 (4)			5487.4 (14)
		(5468)			5468.4 (14)
5448 (10)	5441.1 (40)	5444 (3)			5443.3 (14)

1964Gr11	1967Ba51 ^a	1967Dz02 ^b	1972Go29	1991Ry01	Recommended
	5433.5 (40)	5437 (4)			5435.8 (14)
					5430.1 (14)
	5419 (7)	5427 (4)			5428.3 (14)
		5411 (4)			5414.5 (14)
5398		5391 (4)			5391.2 (14)
5367		(5377)			5379.0 (14)
		(5355)			5356.2 (14)
		(5342)			5341.9 (14)
5328 (10)	5318 (4)	5322 (3)			5321.2 (14)
5295 (10)	5285 (4)	5286 (3)			5287.6 (14)
	5266.5 (40)	5271 (4)			5269.1 (14)
	5229 (7)	5238 (4)			5239.3 (14)
5225	5209.3 (50)	5211 (3)			5210.2 (14)
	5205.5 (50)	5201 (5)			5203.3 (14)
		(5192)			5195.1 (14)
		5160 (5)			5162.1 (14)
		5130 (5)			5129.0 (14)
		5091 (4)			5094.1 (14)
					5076.8 (14)
		5066 (5)			5064.1 (14)
		(5030)			5035.5 (14)
					5025.5 (14)
		(5020)			5019.3 (14)
					4992.7 (14)
		4901 (5)			4903.6 (14)

^a: Original energies should be increased by 1 keV due to changes in calibration energies (recommended by 1979Ry03).

^b: Original energies should be decreased by 0.3 keV due to changes in calibration energies (recommended by 1979Ry03)

The evaluated alpha particle emission probabilities were deduced from the transition intensity balance and listed in table 4. These calculated results are in good agreement with the measured emission probabilities of the main alpha transitions. The measurements are from 1964Gr11, 1967Ba51, 1967Dz02, 1972Go29, and 2003Ku4.

Table 4: Measured and recommended α -particle emission probabilities for ²²⁵Ac.

E_α (keV)	P_α					
	1964Gr11	1967Ba51	1967Dz02	1972Go29	2003Ku44	Evaluation
5829.6 (14)	52 (3)	50.65 (15)	51.6 (15)			52.4 (24)
5804.2 (14)		0.3				0.3
5793.1 (21)	28 (3)	24.3 (1)	26.7 (10)	18.1 (20)	20.2 (11)	18.9 (20)
5791.7 (14)		2.50 (1)			8.4 (5)	6.2 (9)
5731.9 (17)	12 (2)	10.10 (3)	10.0 (1)	8.6 (9)	8.5 (4)	9.0 (5)
5731.6 (14)					1.6 (2)	1.24 (10)
5730.5 (14)					1.05 (8)	1.6 (3)

E_{α} (keV)	P_{α}					Evaluation
	1964Gr11	1967Ba51	1967Dz02	1972Go29	2003Ku44	
5723.1 (14)		3.40 (1)	2.9 (5)		3.77 (19)	2.03 (23)
5686.4 (14)					0.095 (4)	0.021 (14)
5682.2 (14)	1.3 (3)	1.250 (4)	1.4 (2)		1.08 (5)	1.31 (4)
5637.3 (14)	4.2 (3)	4.350 (13)	4.5 (3)		3.7 (1)	4.16 (23)
5609.0 (14)	1.0 (3)	1.20 (1)	1.1 (1)		0.86 (3)	1.09 (5)
5599.3 (14)		0.0410 (1)			0.099 (4)	0.114 (7)
5580.5 (14)	1.0 (3)	1.20 (4)	1.2 (1)		0.89 (3)	0.95 (4)
5563.3 (14)		0.0340 (1)			0.0034 (5)	0.017 (7)
5555.3 (14)		0.1000 (3)			0.089 (4)	0.084 (10)
5546.5 (14)		0.0310 (1)			0.075 (3)	0.055 (12)
5540.1 (14)		0.0150 (5)	0.04 (1)		0.0070 (7)	0.0072 (8)
5523.7 (14)		~ 0.005	0.010 (2)			0.013 (6)
5515.2 (14)		~ 0.005	≤ 0.02			0.0052 (19)
5497.4 (14)	~ 0.02		0.003 (1)			0.0022 (7)
5487.4 (14)			0.0020 (7)			0.0020 (3)
5468.4 (14)			≤ 0.001			0.00052 (18)
5443.3 (14)	0.15 (5)	0.150 (1)	0.13 (1)		0.086 (4)	0.098 (19)
5435.8 (14)		0.0710 (2)	0.07 (2)		0.029 (2)	0.0083 (6)
5430.1 (14)						0.0028 (8)
5428.3 (14)			0.008 (3)		0.0010 (1)	0.0023 (3)
5414.5 (14)		~ 0.003	0.0020 (5)			0.0030 (4)
5391.2 (14)	~ 0.01		0.0010 (5)			0.0006 (4)
5379.0 (14)	~ 0.01		≤ 0.001			0.0020 (5)
5356.2 (14)			≤ 0.001			9.7E-5 (2)
5341.9 (14)			≤ 0.001		0.0009 (3)	0.0027 (8)
5321.2 (14)	0.07 (3)	0.080 (2)	0.068 (8)		0.054 (2)	0.007 (7)
5287.6 (14)	0.2 (1)	0.300 (1)	0.23 (1)		0.17 (1)	0.214 (10)
5269.1 (14)		0.0180 (5)	0.009 (2)		0.0086 (8)	0.048 (19)
5239.3 (14)			0.0030 (8)		0.00019 (8)	0.0026 (5)
5210.2 (14)	~ 0.02	0.0250 (1)	0.003 (3)		0.022 (2)	0.022 (1)
5203.3 (14)		0.0130 (1)	0.0020 (5)		0.0044 (6)	0.0101 (10)
5195.1 (14)			≤ 0.002			0.00015 (5)
5162.1 (14)			0.0020 (8)			0.00066 (12)
5129.0 (14)			0.0020 (8)		0.0013 (3)	0.0058 (8)
5094.1 (14)			0.006 (1)		0.0054 (15)	0.015 (7)
5076.8 (14)						0.0038 (19)
5064.1 (14)			0.003 (1)		0.0014 (2)	0.00114 (18)
5035.5 (14)			≤ 0.001			0.0021 (3)
5025.5 (14)						0.00083 (21)
5019.3 (14)			≤ 0.001		~ 0.00004	0.00015 (5)
4992.7 (14)						0.0013 (3)
4903.6 (14)			0.0020 (5)			0.0011 (4)

3. Atomic data

Atomic fluorescence yields ($\omega_K, \omega_L, \omega_M, \eta_{KL}$ and η_{LM}) are from Schönfeld (1996Sc06).

The X-ray and Auger electron emission probabilities have been deduced from γ -ray and conversion electron data by using the computer code RADLST. Measured and calculated X-ray emission probabilities are compared in Table 5.

Table 5: Comparison of the calculated and measured X-ray emission probabilities.

	1972Dz14	Adopted (deduced)
$K_{\alpha 1}$	1.5 (2)	1.64 (12)
$K_{\alpha 2}$	1.0 (1)	1.00 (8)

The deduced KX-ray emission probabilities agree with the measured value of 1972Dz14, thus confirming the completeness of the decay scheme.

4. Electron Emissions.

The conversion electron emission probabilities have been deduced from γ -ray transition data.

5. Photon Emissions

5.1 γ -ray energy values

There are many measured γ -ray energies of ²²⁵Ac. The present evaluated values are taken from the LWM average value of 1972Dz14, 2000Ar23 and 2003Ku44. The measurements of 1990ArZZ were replaced by 2000Ar23. The experimental and our recommended γ -ray energies from ²²⁵Ac α decay are listed in table 6.

Table 6: Measured and recommended value of γ -ray energy for ²²⁵Ac (keV).

1972Dz14	1990ArZZ	2000Ar23	2003Ku44	LWM	Evaluation
		10.6	10.6		10.6
26.0 (1)	26.05 (10)	26.0 (1)	26.0	26.0 (1)	26.0 (1)
36.6 (1)	36.65 (3)	36.70 (3)	36.7 (1)	36.69 (3)	36.69 (3)
38.5 (1)	38.53 (3)	38.60 (4)	38.5 (1)	38.58 (4)	38.58 (4)
	46.24 (5)	46.24 (5)	46.2 (2)	46.24 (5)	46.24 (5)
49.0 (2)	49.09 (5)	49.13 (4)	49.1 (2)	49.12 (4)	49.12 (4)
		50.2			50.2
53.8 (1)		53.01 (5)		53.4 (4)	53.4 (4)
57.8 (1)	57.75 (5)	57.69 (4)	57.8 (2)	57.71 (4)	57.71 (4)
		62.6 (3)			62.6 (3)
62.90 (5)	62.95 (3)	62.96 (3)	62.9 (1)	62.94 (3)	62.94 (3)
		63.5 (3)			63.5 (3)
64.1 (1)	64.28 (5)	64.28 (3)	64.3 (1)	64.27 (3)	64.27 (3)
69.8 (1)	69.8 (2)	69.87 (5)		69.86 (5)	69.86 (5)
71.7 (1)	71.74 (3)	71.72 (4)	71.4 (3)	71.71 (4)	71.71 (4)
73.6 (1)	73.5 (1)	73.36 (20)	73.5	73.55 (9)	73.55 (9)
73.83 (5)	73.86 (2)	73.85 (4)	73.9 (1)	73.85 (3)	73.85 (3)
74.9 (2)	74.9 (2)	74.82 (5)	74.6 (4)	74.82 (5)	74.82 (5)

1972Dz14	1990ArZZ	2000Ar23	2003Ku44	LWM	Evaluation
			78.8		78.8
					82.6 ^{ab}
87.38 (5)	87.41 (3)	87.42 (3)	87.4 (1)	87.41 (3)	87.41 (3)
94.9 (2)	94.90 (5)	94.90 (3)	94.9 (1)	94.90 (2)	94.90 (2)
96.3 (2)	96.15 (5)	96.15 (5)	96.7 (5)	96.16 (5)	96.16 (5)
99.55 (10)	99.63 (5)	99.71 (6)	99.6	99.67 (5)	99.67 (5)
99.8 (1)	99.91 (5)	100.07 (10)	99.8 (1)	99.89 (6)	99.89 (6)
100.8 (1)	100.96 (5)	100.87 (4)	100.8 (2)	100.86 (4)	100.86 (4)
	103.46 (10)	103.44 (12)	103.6 (2)	103.48 (10)	103.48 (10)
108.4 (1)	108.41 (3)	108.38 (3)	108.4 (1)	108.38 (3)	108.38 (3)
111.5 (1)	111.54 (3)	111.52 (3)	111.5 (1)	111.52 (3)	111.52 (3)
	112.8 (2)	112.8 (2)	112.8	112.8 (2)	112.8 (2)
			114		114
		119.09 (6)			119.09 (6) ^b
119.9 (1)	119.87 (5)	119.84 (3)	119.9 (1)	119.85 (3)	119.85 (3)
		121.06 (7)			121.06 (7)
123.8 (1)	123.75 (5)	123.73 (4)	123.8 (1)	123.75 (4)	123.75 (4)
124.8 (1)	124.82 (5)	124.81 (3)	124.8 (1)	124.81 (3)	124.81 (3)
	126.15 (10)	126.09 (5)	126.2 (2)	126.10 (5)	126.10 (5)
	129.2 (2)	129.22 (7)	129.2 (2)	129.22 (7)	129.22 (7)
	133.64 (5)	133.60 (4)	133.6 (1)	133.60 (3)	133.60 (3)
134.8 (1)	134.86 (5)	134.85 (3)	134.9 (1)	134.85 (3)	134.85 (3)
		137.40 (10)	137.6		137.40 (10)
					138.2 ^{ab}
			139.6		139.6
		144.7 (2)	144.7		144.7 (2)
145.0 (2)	145.17 (5)	145.15 (3)	145.2 (1)	145.15 (3)	145.15 (3)
150.09 (5)	150.04 (2)	150.02 (4)	150.1 (1)	150.05 (3)	150.05 (3)
	152.63 (5)	152.64 (3)	152.6 (2)	152.64 (3)	152.64 (3)
154.0 (1)	153.92 (5)	153.91 (3)	153.9 (1)	153.92 (3)	153.92 (3)
157.25 (5)	157.26 (2)	157.24 (3)	157.3 (2)	157.25 (3)	157.25 (3)
		161.35 (7)			161.35 (7)
	169.1 (2)	169.18 (4)	169.1	169.18 (4)	169.18 (4)
			169.9		169.9
170.6 (1)	170.7 (2)	170.83 (6)	170.7 (2)	170.77 (5)	170.77 (5)
			173.4		173.4
	178.4 (1)	178.29 (3)	178.3 (2)	178.29 (3)	178.29 (3)
	179.8 (2)	179.78 (4)	179.8 (3)	179.78 (4)	179.78 (4)
			183		183
			186.1		186.1
186.1 (1)	186.2 (1)	186.31 (3)	186.3	186.29 (3)	186.29 (3)
			187.2		187.2
188.0 (1)	188.00 (5)	187.95 (3)	188.0 (1)	187.96 (3)	187.96 (3)
			193.2		193.2
195.69 (7)	195.78 (5)	195.74 (3)	195.8 (2)	195.74 (3)	195.74 (3)

1972Dz14	1990ArZZ	2000Ar23	2003Ku44	LWM	Evaluation
		197.50 (3)	197.4		197.50 (3)
	197.7 (1)		197.9		197.7 (1)
198.70 (7)	198.7 (1)	198.23 (8)	198.4 (3)	198.47 (23)	198.47 (23)
		205.12 (11)	204.7 (3)	205.07 (11)	205.07 (11)
	216.90 (5)	216.89 (3)	216.9 (2)	216.89 (3)	216.89 (3)
		220.43 (8)			220.43 (8)
224.56 (7)	224.64 (5)	224.58 (3)	224.7 (1)	224.59 (3)	224.59 (3)
			228.2 (4)		228.2 (4)
	231.3 (2)	231.14 (7)	231.3 (2)	231.16 (7)	231.16 (7)
			236.0 (6)		236.0 (6)
		238.64 (8)			238.64 (8)
	240.8 (1)	240.68 (3)	240.7 (2)	240.68 (3)	240.68 (3)
	243.2 (1)	243.11 (5)	243.2 (2)	243.12 (5)	243.12 (5)
	249.5 (2)	249.60 (3)	249.6 (2)	249.60 (3)	249.60 (3)
253.50 (7)	253.54 (5)	253.45 (3)	253.5 (1)	253.46 (3)	253.46 (3)
		256.0 (2)	256		256.0 (2)
	279.25 (10)	279.18 (3)	279.3 (3)	279.18 (3)	279.18 (3)
	282.1 (2)				282.1 (2)
	284.8 (1)	284.75 (3)	284.8 (3)	284.75 (3)	284.75 (3)
		298.32 (5)	298.6 (3)	298.33 (5)	298.33 (5)
		317.23 (18)	317.4		317.23 (18)
		321.77 (4)	321.8 (4)	321.77 (4)	321.77 (4)
	348.5 (1)	348.33 (4)	348.2 (4)	348.33 (4)	348.33 (4)
	354.8 (2)	354.54 (6)	354.9 (3)	354.56 (6)	354.56 (6)
			356.6		356.6
	362.5 (1)	362.38 (3)	362.2 (4)	362.38 (3)	362.38 (3)
		367.72 (12)	368.3 (6)	367.74 (12)	367.74 (12)
	375.2 (1)	374.98 (5)	375.0 (7)	374.98 (5)	374.98 (5)
		388.07 (7)			388.07 (7)
	403.1 (1)	403.1 (1)	403.4 (3)	403.13 (10)	403.13 (10)
	406.1 (1)	405.95 (3)	406.2 (3)	405.95 (3)	405.95 (3)
	418.1 (1)	417.90 (3)	417.9 (3)	417.90 (2)	417.90 (2)
		429.80 (18)			429.80 (18)
		434.81 (5)	435.0 (3)	434.82 (5)	434.82 (5)
		442.16 (8)			442.16 (8)
		443.43 (10)			443.43 (10)
		446.31 (10)			446.31 (10)
		451.04 (5)	450.1 (7)	451.04 (5)	451.04 (5)
452.4 (1)	452.4 (1)	452.21 (3)	452.4 (2)	452.23 (3)	452.23 (3)
	458.8 (2)	458.79 (8)	458.8 (4)	458.79 (8)	458.79 (8)
	462.4 (4)	462.43 (13)	462.4 (6)	462.43 (13)	462.43 (13)
	469.5 (3)	469.48 (5)	469.5 (3)	469.48 (5)	469.48 (5)
	481.05 (5)	480.84 (3)	481.1 (2)	480.85 (3)	480.85 (3)
		491.42 (10)	492.6 (6)	491.45 (10)	491.45 (10)
	496.9 (3)				496.9 (3)

1972Dz14	1990ArZZ	2000Ar23	2003Ku44	LWM	Evaluation
			498.6 (6)		498.6 (6)
			512.5 (7)		512.5 (7)
	515.40 (5)	515.12 (3)	515.3 (2)	515.13 (3)	515.13 (3)
	517.78 (5)	517.50 (3)	517.9 (2)	517.51 (3)	517.51 (3)
	522.3 (1)	522.14 (4)	522.1 (2)	522.14 (4)	522.14 (4)
	526.09 (5)	525.77 (3)	526.1 (1)	525.94 (17)	525.94 (17)
		527.29 (5)			527.29 (5) ^b
	529.9 (1)	529.59 (3)	529.7 (3)	529.59 (3)	529.59 (3)
	531.3 (1)	530.86 (4)	531.2 (3)	530.87 (4)	530.87 (4)
		532.11 (9)			532.11 (9)
	538.1 (1)				538.1 (1)
			545.8 (6)		545.8 (6)
	552.0 (1)	551.78 (3)	552.0 (2)	551.79 (3)	551.79 (3)
		564.31 (11)	565.6 (7)	564.34 (11)	564.34 (11)
		567.47 (5)	568.3 (6)	567.48 (5)	567.48 (5)
	571.0 (1)	570.68 (3)	571.0 (2)	570.69 (3)	570.69 (3)
		590.41 (5)	591.4 (7)	590.42 (5)	590.42 (5)
	594.2 (1)	593.86 (4)	594.6 (3)	593.87 (4)	593.87 (4)
	601.1 (1)	600.92 (3)	601.0 (3)	600.92 (3)	600.92 (3)
	603.3 (1)	603.09 (4)	603.5 (5)	603.09 (4)	603.09 (4)
		628.93 (10)	629.9 (7)	628.95 (10)	628.95 (10)
			637.1 (7)		637.1 (7)
		645.87 (13)	646.3 (3)	645.94 (12)	645.94 (12)
	649.2 (1)	649.01 (4)	649.5 (2)	649.03 (4)	649.03 (4)
			653.5 (4)		653.5 (4) ^b
		656.18 (11)			656.18 (11)
		657.88 (5)			657.88 (5)
		667.10 (8)	668.1 (4)	667.14 (8)	667.14 (8)
		675.51 (18)	674.3 (4)	674.9 (3)	674.9 (3)
	679.7 (1)	679.35 (6)	680.4 (6)	679.36 (6)	679.36 (6)
		697.54 (13)	698.4 (4)	697.62 (12)	697.62 (12) ^b
		702.00 (14)			702.00 (14)
	747.0 (1)	747.0 (1)	747	747.0 (1)	747.0 (1)
		752.46 (12)			752.46 (12)
	753.7 (3)	754.04 (13)	753.7	754.04 (13)	754.04 (13)
	758.7 (1)				758.7 (1) ^b
		767.6 (4)	768.4 (5)	767.9 (3)	767.9 (3)
			780.6 (6)		780.6 (6)
		808.48 (10)			808.48 (10)
			824.2 (7)		824.2 (7)

^a: from 1969Le09.

^b: not placed in level scheme.

5.2 Relative values of the γ -ray intensities

The results of measurements of the relative γ -ray intensities of ²²⁵Ac are listed in table 7. Compared to the old measurements of 1967Le23 and 1972Dz14, recently measurements of 2000Ar23 and 2003Ku44 have better energy resolutions and higher detector efficiency. On the other hand, some measurements of 1967Le23 and 1972Dz14 have no uncertainties. Thus the recommended values are taken from the LWM average of the measured values of 2000Ar23 and 2003Ku44. The measurements of 1990ArZZ were replaced by 2000Ar23; measurements of 1994Gr20 were replaced by 2003Ku44.

Table 7: Measured and recommended relative γ -ray intensities for ²²⁵Ac.

E_γ (keV)	I_γ							
	1967Le23	1972Dz14	1990ArZZ	1994Gr20	2000Ar23	2003Ku44	LWM	Evaluation
10.6								2.17 (28)*
26.0 (1)		~ 0.21	< 1.4		0.23 (3)	0.25 (8)	0.23 (3)	0.23 (3)
36.69 (3)	~ 4.1	~ 2.1	2.19 (27)	2.63 (36)	2.65 (33)	2.58 (27)	2.61 (21)	2.61 (21)
38.58 (4)		1.4	1.64 (14)	1.84 (50)	1.48 (23)	1.57 (16)	1.54 (13)	1.54 (13)
46.24 (5)			0.55 (27)		0.82 (17)	0.65 (11)	0.70 (9)	0.70 (9)
49.12 (4)		0.7	0.96 (27)	1.07 (36)	1.3 (2)	1.10 (13)	1.16 (11)	1.16 (11)
50.2					~ 0.09			~ 0.09
53.4 (4)		2.68 (56)			< 0.58			< 0.58
57.71 (4)		0.7	0.55 (27)	0.71 (36)	0.88 (19)	0.65 (14)	0.73 (11)	0.73 (11)
62.6 (3)					0.77 (17)			0.77 (17)
62.94 (3)	58 (7)	77.5 (70)	56.2 (27)	69.1 (52)	69.5 (87)	71.7 (49)	71.2 (42)	71 (4)
63.5 (3)					3.0 (4)			3.0 (4)
64.27 (3)		8.5 (28)	4.1 (4)	5.4 (5)	6.8 (7)	6.83 (75)	6.8 (5)	6.8 (5)
69.86 (5)		0.7	0.68 (27)	0.89 (36)	0.68 (17)		0.68 (17)	0.68 (17)
71.71 (4)		1.4	1.78 (14)	1.96 (48)	1.87 (20)	2.10 (43)	1.91 (18)	1.91 (18)
73.55 (9)		2.8	1.23 (27)		2.17 (72)	4.2 (12)	2.7 (6)	2.7 (6)
73.85 (3)	55 (10)	45.1 (42)	39.6 (18)	43.0 (34)	46.3 (58)	44.0 (36)	44.6 (31)	44.6 (31)
74.82 (5)		5.6	2.19 (27)		1.88 (43)	3.7 (12)	2.1 (4)	2.1 (4)
78.8				3.0 (13)		1.78 (27)		1.78 (27)
82.6 ^x	21 (5)							21 (5)
87.41 (3)	< 6.8	40.8 (42)	31.9 (15)	40.5 (29)	44.9 (58)	37.7 (29)	39.1 (26)	39.1 (26)
94.90 (2)		22.5 (85)	11.9 (11)	12.5 (14)	18.8 (27)	14.0 (15)	15.1 (13)	15.1 (13)
96.16 (5)	4 (1)	4.2 (14)	3.84 (41)		< 4.3	4.7 (9)		4.7 (9)
99.67 (5)	301 (55)	95.8 (99)	78.1 (41)	243 (2)	197 (27)	117 (12)	110 (7)	110 (7)
99.89 (6)		239 (28)	127.4 (68)		38 (14)	167 (20)	156 (11)	156 (11)
100.86 (4)		7.0	8.8 (14)	10.9 (27)	17.5 (19)	12.5 (12)	13.9 (10)	13.9 (10)
103.48 (10)	~ 1.4		0.55 (27)		0.94 (27)	0.38 (9)	0.44 (9)	0.44 (9)
108.38 (3)	38 (7)	39.4 (42)	31.5 (14)	37.9 (27)	39.1 (43)	36.0 (26)	36.8 (22)	36.8 (22)
111.52 (3)	44 (7)	45.1 (42)	39.9 (18)	48.0 (36)	49.2 (58)	44.0 (32)	45.2 (28)	45.2 (28)
112.8 (2)			0.27 (13)		< 0.43	0.30 (4)		0.30 (4)
114						0.125 (18)		0.125 (18)
119.09 (6) ^x					2.6 (4)			2.6 (4)
119.85 (3)	9.6 (27)	8.5 (14)	9.3 (8)	12.1 (13)	14.0 (14)	11.0 (7)	11.6 (6)	11.6 (6)
121.06 (7)					2.5 (7)			2.5 (7)
123.75 (4)		26.8 (28)	9.0 (8)	10.9 (14)	14.2 (14)	12.0 (9)	12.6 (8)	12.6 (8)

E_γ (keV)	I_γ							
	1967Le23	1972Dz14	1990ArZZ	1994Gr20	2000Ar23	2003Ku44	LWM	Evaluation
124.81 (3)	29 (7)	7.0 (14)	3.3 (3)	4.6 (9)	4.6 (4)	4.0 (3)	4.22 (24)	4.22 (24)
126.10 (5)			0.96 (27)		1.06 (20)	1.17 (12)	1.14 (10)	1.14 (10)
129.22 (7)			0.41 (14)		0.48 (16)	0.37 (7)	0.39 (7)	0.39 (7)
133.60 (3)			1.78 (27)	2.7 (4)	13.9 (27)	2.83 (22)		2.83 (22)
134.85 (3)	5.5 (27)	5.6 (14)	3.84 (41)	5.0 (5)	4.8 (7)	4.5 (4)	4.6 (4)	4.6 (4)
137.40 (10)					0.43 (19)	0.32 (4)	0.33 (4)	0.33 (4)
138.2 ^x	2.7 (14)							2.7 (14)
139.6						0.20 (3)		0.20 (3)
144.7 (2)	21 (4)				~ 0.07	0.067 (17)		0.067 (17)
145.15 (3)		18.3 (42)	18.4 (8)	21.8 (18)	21.4 (22)	21.0 (15)	21.1 (12)	21.1 (12)
150.05 (3)	100	100	100	100	100	100	100	100
152.64 (3)			2.2 (3)	2.7 (4)	2.39 (27)	3.17 (23)	2.84 (18)	2.84 (18)
153.92 (3)	23 (4)	26.8 (70)	23.6 (11)	27.7 (30)	28.2 (29)	30.3 (21)	29.6 (17)	29.6 (17)
157.25 (3)	51 (10)	43.7 (42)	45.2 (27)	55.4 (5)	50.7 (58)	53.3 (43)	52.4 (35)	52.4 (35)
161.35 (7)					0.52 (13)			0.52 (13)
169.18 (4)			2.33 (27)	2.86 (36)	2.29 (27)	1.17 (18)	1.7 (6)	1.7 (6)
169.9						2.0 (2)		2.0 (2)
170.77 (5)	5.5 (28)	1.4	0.96 (41)		1.06 (19)	2.83 (22)	1.9 (9)	1.9 (9)
173.4 ^x						1.67 (19)		1.67 (19)
178.29 (3)	2.7 (14)		1.78 (14)		2.32 (26)	2.33 (20)	2.33 (16)	2.33 (16)
179.78 (4)			0.96 (27)	1.25 (36)	1.53 (19)	1.57 (13)	1.56 (11)	1.56 (11)
183 ^x						1.22 (19)		1.22 (19)
186.1						1.83 (19)		1.83 (19)
186.29 (3)		2.8	2.47 (55)		2.74 (30)	0.60 (6)		0.60 (6) ^b
187.2						1.48 (9)		1.48 (9)
187.96 (3)	81 (8)	64.8 (70)	67.8 (34)	78.6 (5)	78.1 (87)	75 (5)	75.8 (44)	76 (4)
193.2 ^x						0.28 (5)		0.28 (5)
195.74 (3)	19 (4)	19.7 (28)	20.5 (14)	25.2 (13)	23.4 (23)	20.5 (14)	21.3 (12)	21.3 (12)
197.50 (3)				3.6 (7)		3.83 (39)		3.8 (4) ^b
197.7 (1)			7.53 (68)	4.1 (9)	7.8 (10)	5.5 (6)		5.5 (6) ^b
198.47 (23)	4.1 (12)	2.8	3.01 (68)	3.8 (9)	2.55 (26)	2.83 (22)	2.71 (17)	2.71 (17)
205.07 (11)					0.27 (10)	0.18 (7)	0.21 (6)	0.21 (6)
216.89 (3)	47 (14)		39.7 (82)	53 (10)	47.8 (43)	45.2 (33)	46.2 (27)	46.2 (27)
220.43 (8)					0.87 (26)			0.87 (26)
224.59 (3)	15 (4)	11.3 (14)	12.1 (12)	14.8 (14)	15.6 (17)	16.3 (12)	16.1 (10)	16.1 (10)
228.2 (4)						0.67 (17)		0.67 (17)
231.16 (7)			0.27 (13)		0.30 (7)	1.10 (12)	0.7 (4)	0.7 (4)
236.0 (6)						0.25 (4)		0.25 (4)
238.64 (8)					0.14 (4)			0.14 (4)
240.68 (3)	2.7 (13)		0.96 (27)		1.71 (19)	1.67 (19)	1.69 (14)	1.69 (14)
243.12 (5)			0.16 (7)		0.39 (7)	0.50 (6)	0.45 (5)	0.45 (5)
249.60 (3)	2.7 (13)		1.51 (68)		1.9 (2)	2.0 (2)	1.95 (14)	1.95 (14)
253.46 (3)	21 (5)	14.1 (14)	15.5 (7)	18.4 (9)	18.5 (19)	19.3 (14)	19.0 (11)	19.0 (11)
256.0 (2)					0.05 (1)	0.100 (34)	0.054 (10)	0.054 (10)

E_γ (keV)	I_γ							Evaluation
	1967Le23	1972Dz14	1990ArZZ	1994Gr20	2000Ar23	2003Ku44	LWM	
279.18 (3)	4.1 (12)		2.33 (27)		4.63 (43)	4.17 (39)	4.4 (3)	4.4 (3)
282.1 (2)			0.55 (27)					0.079 (6)*
284.75 (3)	~ 1.4		0.55 (27)	0.71 (36)	1.09 (13)	1.05 (10)	1.07 (8)	1.07 (8)
298.33 (5)					0.29 (4)	0.30 (9)	0.29 (4)	0.29 (4) ^c
317.23 (18)					0.06 (3)	> 0.018		0.06 (3) ^c
321.77 (4)					0.46 (7)	0.50 (7)	0.48 (5)	0.48 (5) ^c
348.33 (4)			0.41 (14)		0.46 (7)	0.42 (5)	0.43 (4)	0.43 (4)
354.56 (6)			0.21 (5)	0.25 (7)	0.19 (3)	0.38 (5)	0.29 (10)	0.29 (10)
356.6						0.037 (15)		0.037 (15)
362.38 (3)			0.82 (27)		0.9 (1)	0.70 (8)	0.78 (6)	0.78 (6)
367.74 (12)					0.05 (3)	0.10 (3)	0.075 (25)	0.075 (25)
374.98 (5)			0.41 (14)		0.027 (4)	0.28 (7)		0.28 (7)
388.07 (7)					0.18 (3)			0.18 (3)
403.13 (10)			0.18 (5)		< 0.29	0.027 (23)		0.027 (23)
405.95 (3)			0.96 (27)		1.14 (13)	1.12 (9)	1.13 (7)	1.13 (7)
417.90 (2)			0.68 (14)		0.82 (10)	0.80 (8)	0.81 (6)	0.81 (6)
429.80 (18)					0.055 (27)			0.055 (27)
434.82 (5)					0.46 (7)	0.40 (5)	0.42 (4)	0.42 (4)
442.16 (8)					0.65 (10)			0.65 (10)
443.43 (10)					~ 0.014			~ 0.014 ^d
443.43 (10)					0.20 (7)			0.20 (7) ^d
446.31 (10)					0.09 (5)			0.09 (5)
451.04 (5)					0.41 (7)	0.53 (14)	0.43 (6)	0.43 (6)
452.23 (3)	15 (5)	15.5 (14)	14.8 (12)		17.1 (19)	14.8 (11)	15.4 (10)	15.4 (10)
458.79 (8)			0.68 (27)		0.07 (2)	0.097 (37)	0.076 (18)	0.076 (18)
462.43 (13)			2.2 (11)		0.055 (16)	0.125 (45)	0.063 (15)	0.063 (15)
469.48 (5)			0.55 (14)		0.26 (10)	0.47 (6)	0.41 (5)	0.41 (5)
480.85 (3)	4.1 (12)		4.1 (4)		4.9 (6)	4.83 (41)	4.85 (34)	4.9 (3)
491.45 (10)					0.06 (2)	0.037 (23)	0.05 (2)	0.05 (2)
496.9 (3)			0.21 (10)					0.21 (10)
498.6 (6)						0.12 (3)		0.12 (3)
512.5 (7)						0.08 (3)		0.08 (3)
515.13 (3)	~ 1.4		2.47 (27)		2.95 (30)	3.17 (23)	3.09 (18)	3.09 (18)
517.51 (3)			1.78 (27)		2.1 (2)	2.5 (2)	2.30 (14)	2.30 (14)
522.14 (4)			0.21 (5)		0.30 (3)	0.30 (5)	0.30 (2)	0.30 (2)
525.94 (17)	~ 1.4		3.97 (41)		4.63 (43)	5.50 (43)	5.1 (3)	5.1 (3)
527.29 (5) ^x					0.27 (4)			0.27 (4)
529.59 (3)			0.82 (41)		1.01 (12)	1.18 (13)	1.09 (9)	1.09 (9)
530.87 (4)			0.55 (14)		0.68 (9)	0.67 (9)	0.68 (6)	0.68 (6)
532.11 (9)					0.11 (3)			0.11 (3)
538.1 (1)			0.55 (14)					0.55 (14)
545.8 (6)						0.077 (20)		0.077 (20)
551.79 (3)			0.55 (14)		0.56 (7)	0.93 (8)	0.75 (19)	0.75 (19)
564.34 (11)					~ 0.014	0.032 (13)		0.032 (13)

E_γ (keV)	I_γ							
	1967Le23	1972Dz14	1990ArZZ	1994Gr20	2000Ar23	2003Ku44	LWM	Evaluation
567.48 (5)					0.13 (2)	0.22 (5)	0.18 (5)	0.18 (5)
570.69 (3)			0.55 (14)		0.59 (7)	0.53 (9)	0.57 (6)	0.57 (6)
590.42 (5)					0.12 (2)	0.12 (3)	0.12 (2)	0.12 (2)
593.87 (4)			0.22 (11)		0.41 (4)	0.47 (10)	0.42 (4)	0.42 (4)
600.92 (3)			0.47 (14)		0.35 (6)	0.62 (15)		0.35 (6) ^{ad}
600.92 (3)					~ 0.87			~ 0.87 ^{ad}
603.09 (4)			0.27 (13)		0.25 (3)	0.27 (7)	0.25 (3)	0.25 (3)
628.95 (10)					0.049 (13)	0.043 (14)	0.046 (10)	0.046 (10)
637.1 (7)						~ 0.017		~ 0.017
645.94 (12)					0.032 (10)	0.017 (7)	0.022 (6)	0.022 (6)
649.03 (4)			0.18 (5)		0.27 (3)	0.20 (5)	0.25 (3)	0.25 (3)
653.5 (4) ^x						0.025 (7)		0.025 (7)
656.18 (11)					0.07 (3)			0.07 (3)
657.88 (5)					0.20 (4)			0.20 (4)
667.14 (8)					0.56 (13)	0.040 (12)	0.30 (26)	0.30 (26)
674.9 (3)					0.019 (9)	0.012 (7)	0.015 (6)	0.015 (6)
679.36 (6)			0.11 (3)		0.09 (2)	0.102 (26)	0.095 (16)	0.095 (16) ^c
697.62 (12) ^x					0.035 (13)	0.028 (8)	0.030 (7)	0.030 (7)
702.00 (14)					0.023 (10)			0.023 (10)
747.0 (1)			0.16 (5)		< 0.29	< 0.017		0.16 (5)
752.46 (12)					0.038 (10)			0.038 (10)
754.04 (13)			0.11 (3)		0.033 (10)	< 0.017		0.033 (10)
758.7 (1) ^x			0.68 (14)					0.68 (14)
767.9 (3)					0.049 (13)	0.040 (12)	0.044 (9)	0.044 (9)
780.6 (6)						0.008 (2)		0.008 (2)
808.48 (10)					0.30 (4)			0.30 (4)
824.2 (7)						~ 0.007		~ 0.007

^a: From 2000Ar23.

^b: From 2003Ku44.

^c: Multiply placed, intensity not divided.

^d: Multiply placed, intensity suitable divided.

^{*}: From intensity balance.

^x: Not placed in level scheme.

5.3 Absolute values of the γ -ray emission probabilities

Measured absolute γ -ray emission probabilities for the 150.04 keV line for ²²⁵Ac are compiled and listed in Table 8.

2000Ar23 gives the value 0.691 (16) %, which was obtained from correction of the intensity of 1986He06 using the measured value 0.053 (6) % (2000Ga52) for the 149.89 keV transition in ²²⁹Th α -decay and the measured value 0.051 (10) % (1995Sh01) for the 150.14 keV transition in ²²¹Fr α -decay.

Conversely, to correct the measured intensity of 1986He06, if using the measured value 0.053 (6) % (2000Ga52) for the 149.89 keV transition in ²²⁹Th α -decay and the evaluated value 0.0478 (23) % (1990Ak05) for the 150.14 keV transition in ²²¹Fr α decay, the value would be then 0.695 (13) %. These corrected values are in good agreement with the measured value in 1995Ch74.

The recommended absolute γ -ray emission probability of the 150.04 keV γ -ray is from the measurement of 1995Ch74 and adopted as the normalization factor N, with $N = 0.006\ 93\ (12)$. The recommended absolute γ -ray emission probabilities are the relative values evaluated in table 7 multiplied by 0.006 93 (12).

Table 8: Measured and recommended absolute γ -ray emission probability of 150.04 keV for ²²⁵Ac.

P_γ (150.04 keV) (%)	References	Measurement method
0.981 (3)	1981Di14	Ge(Li)
0.796 (11)	1986He06	Ge(Li), Au-Si surface barrier, in equilibrium with ²²⁹ Th.
0.693 (12)	1995Ch74	Ge(Li), $\alpha\gamma$ -coincidence.
0.691 (16)	2000Ar23	From 1986He06 corrected by 2000Ga52 and 1995Sh01.
0.693 (12)		Recommended value from 1995Ch74

6. References

- 1947En03 A.C. English, T.E. Cranshaw, P. Demers, J.A. Harvey, E.P. Hincks, J.V. Jelley, A.N. May, Phys. Rev. 72, 253 (1947) [$T_{1/2}$].
- 1950Ha52 F. Hagemann, L.I. Katzin, M.H. Studier, G.T. Seaborg, A. Ghiorso, Phys. Rev. 79, 435 (1950) [$T_{1/2}$].
- 1964Gr11 G. Graeffe, K. Valli, J. Aaltonen, Ann. Acad. Sci. Fenn., Ser. A VI, No.145 (1964) [E_α , I_α].
- 1967Ba51 G. Bastin-Scoffier, Compt. Rend. 265B, 863 (1967) [E_α , I_α].
- 1967Dz02 B.S. Dzhelepov, R.B. Ivanov, M.A. Mikhailova, L.N. Moskvina, O.M. Nazarenko, V.F. Rodionov, Izv. Akad. Nauk SSSR, Ser. Fiz. 31, 568 (1967) [E_α , I_α].
- 1967Le23 C.-F. Leang, Compt. Rend. 265B, 417 (1967) [E_γ , I_γ].
- 1969Le09 C.-F. Leang, F. Gautier, J. Phys. (Paris) 30, 296 (1969) [E_γ , ce]
- 1971DzZP B.S. Dzhelepov, A.V. Zolotavin, R.B. Ivanov, M.A. Mikhailova, V.O. Sergeev, M.I. Sovtsov, Proc.21st Ann. Conf. Nucl. Spectrosc. Struct. At. Nuclei, Moscow, Pt.1, p.140 (1971) [E_γ , I_γ , $I(\text{ce})$, Multipolarity].
- 1972Dz14 B.S. Dzhelepov, R.B. Ivanov, M.A. Mikhailova, V.O. Sergeev, Izv. Akad. Nauk SSSR, Ser. Fiz. 36, 2080 (1972) [E_γ , I_γ , $E(\text{ce})$, $I(\text{ce})$, Multipolarity].
- 1972Go29 N.A. Golovkov, B.S. Dzhelepov, R.B. Ivanov, M.A. Mikhailova, V.G. Chumin, Sov. J. Nucl. Phys. 15, 349 (1972) [I_α].
- 1977Vy02 T. Vylov, N.A. Golovkov, B.S. Dzhelepov, R.B. Ivanov, M.A. Mikhailova, Y.V. Norseev, V.G. Chumin, Bull. Acad. Sci. USSR, Phys. Ser. 41, No.8, 85 (1977) [E_γ , I_γ , Multipolarity].
- 1979Ry03 A. Rytz, At. Data Nucl. Data Tables 23, 507 (1979) [E_α , I_α].
- 1981Di14 J.K. Dickens, J.W. McConnell, Radiochem. Radioanal. Lett. 47, 331 (1981) [E_γ , I_γ].
- 1986He06 R.G. Helmer, C.W. Reich, M.A. Lee, I. Ahmad, Int. J. Appl. Radiat. Isotop. 37, 139(1986) [E_γ , I_γ , P_γ].
- 1990Ak03 Y.A. Akovali, Nucl. Data Sheets 60, 617(1990) [Spin, parity].
- 1990Ak05 Y.A. Akovali, Nucl. Data Sheets 61, 623(1990) [NDS].
- 1990ArZZ G. Ardisson, M.C. Kouassi, J. Dalmaso, Priv. Comm. (1990) [E_γ , I_γ , Multipolarity].
- 1990Ko14 M.C. Kouassi, J. Dalmaso, H. Maria, G. Ardisson, M. Hussonnois, J. Radioanal. Nucl. Chem. 144, 387 (1990) [E_γ , I_γ].
- 1991Ko12 M.C. Kouassi, J. Dalmaso, M. Hussonnois, V. Barci, G. Ardisson, J. Radioanal. Nucl. Chem. 153, 293 (1991) [E_γ , I_γ].

- 1994Gr20 K.Ya. Gromov, M.Ya. Kuznetsova, Yu.N. Norseev, N.I. Rukhadze, V.I. Fominykh, V.V. Tsupko-Sitnikov, V.G. Chumin, M.B. Yuldashev, Yu.S. Butabaev, R.A. Niyazov, Bull. Rus. Acad. Sci. Phys. 58, 29 (1994) [E_γ , I_γ].
- 1995Ch74 V.G. Chumin, S.S. Eliseev, K.Ya. Gromov, Yu.V. Norseev, V.I. Fominykh, V.V. Tsupko-Sitnikov, Bull. Rus. Acad. Sci. Phys. 59, 1854 (1995) [E_γ , I_γ , P_γ].
- 1995Sh01 R.K. Sheline, C.F. Liang, P. Paris, Phys. Rev. C51, 1192 (1995) [P_γ].
- 1996Sc06 E. Schönfeld, H. Janssen, Nucl. Instrum. Meth. Phys. Res. A369, 527 (1996) [Atomic data].
- 2000Ar23 G. Ardisson, J. Gasparro, V. Barci, R.K. Sheline, Phys. Rev. C62, 064306 (2000) [E_γ , I_γ].
- 2000Ga52 J. Gasparro, G. Ardisson, V. Barci, R.K. Sheline, Phys. Rev. C62, 064305 (2000) [E_γ , I_γ , P_γ].
- 2002Ba85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, At. Data Nucl. Data Tables 81, 1 (2002) [Calculated ICC]
- 2003Au03 G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A729, 129 (2003) [Q].
- 2003Ku44 S.A. Kudrya, V.M. Gorozhankin, K.Ya. Gromov, Sh.R. Malikov, L.A. Malov, V.A. Sergienko, V.I. Fominykh, V.V. Tsupko-Sitnikov, V.G. Chumin, E.A. Yakushev, Bull. Rus. Acad. Sci. Phys. 67, 7 (2003) [E_γ , I_γ , E_α , I_α , $E(\text{ce})$, $I(\text{ce})$, Multipolarity].
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr., Nucl. Instrum. Meth. Phys. Res. A589, 202 (2008) [Theoretical ICC].

²²⁶Ra - Comments on evaluation of decay data by V. Chisté and M. M. Bé

This evaluation was completed in 2006. This updated version was done in January 2007. The literature available by this date is included.

1 Decay Scheme

²²⁶Ra disintegrates by alpha emissions mainly to the 186 keV level and to the ground state level of ²²²Rn. Spin and parity are from the mass-chain evaluation of Y. A. Akovali (1996El01 and 1996Ak02).

A certain number of measurements of the 186 keV gamma intensity were carried out and the adopted data set is consistent, so the deduced intensity can be considered having a good level of confidence. Therefore, the decay scheme here was built from the gamma-ray intensity measurements.

A good agreement was found between the effective Q value (4870.5 (27) keV) calculated from the decay scheme data and the adopted and recommended value from Audi.

2 Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2003Au03).

Experimental ²²⁶Ra half-life values (in years) are given in Table 1:

Table 1: Experimental values of ²²⁶Ra half-life.

Reference	Experimental value (a)	Comments
S. W. Watson (1928Wa**)	1608	Not used: no uncertainty. Calorimetry.
H. J. J. Braddick (1928Br**)	1603	Not used: no uncertainty. α current.
I. Curie (1928Cu**)	1590	Not used: no uncertainty. Ion current.
F. A. B. Ward (1929Wa**)	1599	Not used: no uncertainty. Number α 's emitted.
L. Meitner (1930Me**)	1590	Not used: no uncertainty. Calorimetry.
E. Gleditsch (1935Gl02)	1691	Not used: no uncertainty. Growth rate.
P. Günther (1939Gü**)	1603	Not used: no uncertainty. He production.
T. P. Kohman (1949Ko01)	1622 (13)	Number α 's emitted.
W. Sebaoun (1956Se10)	1617 (12)	Number α 's emitted.
G. V. Gorshkov (1959Go80)	1577 (9)	Calorimetry.
G. Martin (1959Ma12)	1602 (8)	Calorimetry.
H. Ramthun (1966Ra13)	1599 (7)	Calorimetry.
Recommended value	1600 (7)	$\chi^2 = 2.87$

The weighted average was calculated with LWEIGHT computer program (version 3).

The evaluators have chosen to take into account the only five experimental values with uncertainty found in the literature: 1949Ko01, 1956Se10, 1959Go80, 1959Ma12 and 1966Ra13. With this data set, the largest contribution to the weighted average comes from the value of Ramthun amounting to 33 %. The weighted average of **1600 a** and the external uncertainty of **7 a** is the half-life adopted value. The reduced- χ^2 value is 2.87.

2.1 α Transitions

The transition energies of the α -particles given in Section 2.1 were calculated from Q_α (2003Au03) and level energies.

2.2 g Transitions

The transitions probabilities were calculated using the γ -ray emission intensities and the relevant internal conversion coefficients (see **6.2 Gamma Emissions**).

Multipolarities of these γ -ray transitions are from 1996E101.

186-keV γ -ray : E2
 262-keV γ -ray : [E2]
 414-keV γ -ray : [E1]
 449-keV γ -ray : [E1]
 600-keV γ -ray : [E1]

The internal conversion coefficients (ICC's) for these γ -ray transitions have been interpolated from theoretical values of I. M. Band (2002Ba85) using the BrIcc computer program (calculation for 'hole'). Theoretical values are compared with measured values below:

	De Pinho (1973De50)	Band (Icc99v3a computer program, no hole) ^a	BrIcc program (recommended values)
α_K	0.200 (9)	0.186 (6)	0.190 (3)
α_{L1}	0.031 (6)	0.0319 (10)	0.0321 (5)
α_{L2}	0.226 (16)	0.208 (6)	0.208 (3)
α_{L3}	0.124 (8)	0.1196 (36)	0.1196 (17)
α_L	0.380 (20)	0.360 (11)	0.360 (5)

^a The evaluators have used a fractional uncertainty of 3 % for all Band conversion coefficients.

The results of De Pinho (1973De50) and the theoretical values calculated with two different programs (Icc99v3a (calculation for 'no hole') and BrIcc) are consistent to each other. The recommended values are the BrIcc values for the all conversion coefficients.

3 Atomic Data

Atomic values, ω_K , ω_L and n_{KL} and the X-ray and Auger electron relative probabilities are from Schönfeld and Janßen (1996Sc06).

4 a Emissions

The α -particle energies for the $\alpha_{0,2}$, $\alpha_{0,3}$ and $\alpha_{0,4}$ are from G. Bastin-Scoffier (1963Ba62), with uncertainties given by A. Rytz (1991Ry01). For the $\alpha_{0,0}$ and $\alpha_{0,1}$ emissions, the energies are from A. Rytz (1991Ry01).

The emission intensities of the α -particles have been deduced from the $P(\gamma + ce)$ decay scheme balance at each level. In Table 2, the calculated and recommended values of the emission intensities are compared with the experimental results. For the two most important lines a slight agreement was found between the experimental results given by 2001La14 and the recommended values deduced from the decay scheme balance. For the three weak lines the calculated alpha emission intensities deduced from γ ray measurements are in good agreement with the measured values of Bastin-Scoffier.

Table 2: Experimental and recommended (deduced) values of the α -particles emission intensities.

Energy (keV)	G. Bastin-Scoffier (1963Ba62)	S. LaMont (2001La14)	Recommended values
4784.34 (25)	94.45 (5) ^a	93.84 (11)	94.038 (40)
4601 (1)	5.55 (5) ^a	6.16 (3)	5.950 (40)
4340 (1)	0.0065 (3)		0.0066 (22)
4191 (2)	0.0010 (1)		0.0008
4160 (2)	0.00027 (5)		0.0002

^a uncertainties as given by Rytz.

5 Electron Emissions

The conversion electrons emission intensities have been calculated from γ -ray data using the EMISSION computer program.

6 Photon emissions

6.1 X-rays

The X-ray absolute intensities have been calculated from γ -ray data and ICC using the EMISSION computer program. In Table 3, the recommended values of ²²²Rn X-ray emission intensities are compared with the experimental results.

Table 3: Experimental and recommended values of X-ray emission intensities.

	Delgado (2002De03)	Schötzig (1983Sc13)	De Pinho (1973De50) ^a	Recommended values
K α_1	0.215 (3)			0.317 (6)
K α_2	0.156 (39)			0.192 (4)
K α	0.371 (39)	0.418 (21)		0.509 (7)
K β_1	0.079 (5)			0.1098 (25)
K β_2	0.020 (4)			0.0351 (10)
K β	0.099 (6)	0.145 (9)		0.1449 (27)
XK	0.47 (4)	0.563 (23)	0.693 (26)	0.654 (8)
XL1			0.0181 (25)	0.0147 (4)
XL2			0.420 (28)	0.427 (10)
XL3			0.401 (14)	0.365 (9)
XL			0.839 (43)	0.807 (13)

^a Calculated with $I_\gamma(186) = 3.555 (19)$

The calculated recommended values and 1973De50 values, based on the assumption that $I_\gamma(186) = 3.555 (19)$, are significantly greater than those measured by Delgado (2002De03) or Schötzig (1983Sc13).

The recommended data are in agreement, within the uncertainty values, with the experimental ones of 1973De50, who used a ²²⁶Ra source from which the descendants were removed, since Schötzig and Delgado carried out measurements with sources in equilibrium with their daughters.

6.2 γ -ray Emissions

The energies of the γ -ray emissions given in Section 6.2 are from Y. A. Akovali (1996El01).

The experimental relative γ emission intensities in ²²²Rn are based on all available relative and absolute measurements of gamma-rays for the ²²⁶Ra decay chain. The normalization factor to convert the relative emission intensities to absolute intensities is the weighted average of the measured absolute gamma-ray emission intensities (Table 4) of the most intense line in ²²⁶Ra decay chain, presents in the ²¹⁴Pb disintegration namely the 609.3-keV line.

Table 4: Experimental 609.3 keV absolute gamma-ray emission intensities.

References	Experimental values (%)	Comments
E. W. A. Lingeman (1969Li10)	42.8 (40)	
D. G. Olson (1983Ol01)	45.0 (7)	
U. Schötzig (1983Sc13)	44.6 (5)	
W. –J. Lin (1991Li11)	46.1 (5)	
J. Morel (1998Mo14)	44.8 (6)	Superseded by 2004Mo07
J. Morel (2004Mo07)	45.57 (18)	
Recommended value	45.49 (19)	$\chi^2 = 1.45$

The recommended normalization factor is the weighted average of the five experimental values: 45.49 with an external uncertainty of 0.19.

The experimental relative γ emission intensities of 186- and 262-keV given in Table 5 are relative to the ²¹⁴Bi 609-keV γ -ray.

Table 5: Experimental data set of the 186- and 262- keV relative γ emission intensities.

References	186-keV γ -ray	262-keV γ -ray	Comments
K. Ya. Gromov (1969Gr33)	9.5 (10)		Not used by the evaluators.
G. Wallace (1969Wa27)	9.91 (31)		Not used by the evaluators.
R.S. Mowatt (1970Mo28)	8.20 (12)		outlier
V. S. Aleksandrov (1974AlZT)	8.87 (30)		outlier
V. Zobel (1977Zo01)	9.00 (10)		Not used by the evaluators.
M. A. Farouk (1982Fa10)	9.07 (14)		Not used by the evaluators.
D. G. Olson (1983Ol01)	7.69 (11)		
U. Schötzig (1983Sc13)	7.72 (14)		
G. Mouze (1990MoZP)	8.58 (5)	0.012 (4)	outlier
W. –J. Lin (1991Li11)	7.89 (14)		
D. Sardari (2000Sa32)	7.6 (8)	0.012 (4)	
J. U. Delgado (2002De03)	7.812 (31)		
G. L. Molnar (2002MoZP)	7.85 (5)		
J. Morel (2004Mo07)	7.812 (31)		Not used by the evaluators.
Recommended values	7.815 (25)	0.012 (4)	
χ^2	0.52		

Were omitted from analysis:

- four values: A. Hachem (1975Ha31), G. Mouze (1981Mo28), H. Akcay (1982Ak03) and O. Diallo (1993Di09), because these values come from the same laboratory of G. Mouze (1990MoZP).
- the sets of values from K. Ya. Gromov (1969Gr33), G. Wallace (1969Wa27) and M. A. Farouk (1982Fa10), because of lack in the articles concerning their experimental measurements.
- the set of values from V. Zobel (1977Zo01), because these values have changed the consistency of the data set when they were introduced in the preliminary calculation with Lweight program and produced inconsistent weighted average for gamma emission intensity.

For the 186-keV γ -ray, the evaluators have chosen to take into account the nine values with associated uncertainty for the calculation. The relative γ emission intensity value given by 2004Mo07 is the same one that those measured by J. U. Delgado (2002De03). In 2004Mo07 article, the author measured the 609.3 keV absolute emission probability (Table 4) and normalized the 2002De03 data set with this value of 45.57(18), so the value given in 2004Mo07 was omitted. The weighted average of the remaining values above was calculated using LWEIGHT computer program (version 3). Based on the Chauvenet's criterion, Mowatt (1970Mo28), Aleksandrov (1974AlZT) and Mouze (1990MoZP) were shown outlier values by the Lweight program, then

they have been omitted.

The adopted relative value is the weighted mean of the six remaining values: 7.815, with an internal uncertainty of 0.025 and a reduced χ^2 of 0.52, so this data set is consistent. The largest contribution comes from the value of Delgado (2002De03), amounting to 63 %.

For the 414-, 449- and 600-keV γ -rays, the evaluators used the measured ratios of Lourens (1971Lo19): $I_{414}/I_{186} = 0,00086$; $I_{449}/I_{186} = 5,5 \times 10^{-5}$; $I_{600}/I_{186} = 0,00014$ and the absolute value $I_{\gamma}(186) = 3.555$ (19) %, to determine their absolute emission intensities.

The evaluated relative and absolute γ -ray emission intensities are given in Table 6.

Table 6: Evaluated relative and absolute γ -ray emission intensities.

Energy (keV)	Relative emission intensity (%)	Absolute emission intensity (%)
186.211 (13)	7.815 (25)	3.555 (19)
262.27 (5)	0.012 (4)	0.0055 (18)
414.60 (5)		0.0003
449.37 (10)		0.0002
600.66 (5)		0.0005

6 References

- 1928Wa** S. W. Watson, M. C. Henderson, Proc. Roy. Soc. A118(1928)318 [Half-life].
 1928Br** H. J. J. Braddick, H. M. Cave, Proc. Roy. Soc. A121(1928)367 [Half-life].
 1928Cu** I. Curie, F. Joliot, Compt. Rend. (Paris) 187(1928)43 [Half-life].
 1929Wa** F. A. B. Ward, C. E. Wynn-Williams, H. M. Cave, Proc. Roy. Soc. A125(1929)713 [Half-life].
 1930Me** L. Meitner, W. Ortmann, Z. Phys. 60(1930)143 [Half-life].
 1935Gl02 E. Gleditsch, E. Foeyn, Am. J. Sci. 29(1935)253 [Half-life].
 1939Gü** P. Günther, Z. Phys. Chem. A185(1939)367 [Half-life].
 1949Ko01 T. P. Kohman, D. P. Ames, J. Sedet, NNES 14B(1949)1675 [Half-life].
 1956Se10 W. Sebaoun, Ann. Phys. (Paris) 1(1956)680 [Half-life].
 1959Ma12 G. R. Martin, D. C. Tuck, Int. J. Appl. Radiat. Isot. 5(1959)141 [Half-life].
 1959Go80 G. V. Gorshkov, Z. G. Gretchenko, A. T. Il'inskaya, B. S. Kuznetsov, N. S. Shimanskaya, At. Energ. (USSR) 7(1959)912 [Half-life].
 1960St20 F. S. Stephens, F. Asaro, I. Perlman, Phys. Rev. 119(1960)796 [I_{γ}].
 1963Ba62 G. Bastin-Scoffier, C. F. Leang, R. J. Walen, J. Phys. 24(1963)854 [I_{α}].
 1966Ra13 H. Ramthun, Nucleonik 8(1966)244 [Half-life].
 1969Gr33 K. Ya. Gromov, B. M. Sabirov, J. J. Urbanets, Bull. Acad. Sci. USSR, Phys. Ser. 33(1970)1510 [I_{γ}].
 1969Li10 E. W. A. Lingeman, J. Konijn, P. Polak, A. H. Wapstra, Nucl. Phys. A133(1969)630 [I_{γ}].
 1969Wa27 G. Wallace, G. E. Coote, Nucl. Instrum. Meth. 74(1969)353 [I_{γ}].
 1970Mo28 R. S. Mowatt, Can. J. Phys. 48(1970)2606 [I_{γ}].
 1971Lo19 W. Lourens, A. H. Wapstra, Z. Phys. 247(1971)147 [I_{γ}].
 1974AlZT V. S. Aleksandrov, JINR – PL 7308(1973) [I_{γ}].
 1973De50 A. G. de Pinho, M. Weskler, Z. Naturforsch. 28a(1973)1635 [X-ray emission intensities].
 1975Ha31 A. Hachem, Compt. Rend. (Paris) 281B(1975)45 [I_{γ}].
 1977Zo01 V. Zobel, J. Eberth, U. Eberth, E. Eube, Nucl. Instrum. Meth. 141(1977)329 [I_{γ}].
 1981Mo28 G. Mouze, Compt. Rend. (Paris) 292(1981)1243 [I_{γ}].
 1982Ak03 H. Akcay, G. Mouze, D. Maillard, Ch. Ythier, Radiochem. Radioanal. Lett. 51(1982)1 [I_{γ}].
 1982Fa10 M. A. Farouk, A. M. Al-Soraya, Nucl. Instrum. Meth. 200(1982)593 [I_{γ}].
 1983Ol01 D. G. Olson, Nucl. Instrum. Meth. 206(1983)313 [I_{γ}].
 1983Sc13 U. Schötzig, K. Debertain, Int. J. Appl. Radiat. Isot. 34(1983)533 [I_{γ}].
 1990Ho28 N. E. Holden, Pure Appl. Chem. 62(1990)941 [Half-life].

- 1990MoZP G. Mouze, C. Ythier, J. F. Comanducci, Rev. Roumaine Phys. 35(1990)337 [I_γ].
- 1991Li11 W. -J. Lin, G. Harbottle, J. Radioanal. Nucl. Chem. Lett. 153(1991)137 [I_γ].
- 1991Ry01 A. Rytz, At. Data and Nucl. Data Tables 47(1991)205 [E_α , I_α].
- 1993Di09 O. Diallo, G. Mouze, C. Ythier, J. F. Comanducci, Nuovo Cimento 106A(1993)1321 [I_γ].
- 1996Ak02 Y. A. Akovali, Nucl. Data Sheets 77(1996)433 [Spin, parity and multipolarity].
- 1996El01 Y. A. Akovali, Nucl. Data Sheets 77(1996)271 [Spin, parity and multipolarity].
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
- 1998Mo14 J. Morel, M. Etcheverry, J. L. Picolo, Appl. Radiat. Isot. 49(1998)1387 [I_γ].
- 2000Sa32 D. Sardari, T. D. MacMahon, J. Radioanal. Nucl. Chem. 244(2000)463 [I_γ].
- 2001La14 S. P. LaMont, R. J. Gehrke, S. E. Glover, R. H. Filby, J. Radioanal. Nucl. Chem. 248(2001)247 [I_α].
- 2002De03 J. U. Delgado, J. Morel, M. Etcheverry, Appl. Radiat. Isot. 56(2002)137 [I_γ].
- 2002MoZP G. L. Molnar, Z. S. Révay, T. Belgya, 11th Int. Symp. on capture gamma-ray spectroscopy, 2-6 Sep. 2002, Pruhonice (2003)522 [I_γ].
- 2002Ba85 I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, Jr., P. O. Tikkanen, S. Raman, AtData Nucl. Data Tables 81(2002)1 [Theoretical ICC].
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].
- 2004He** R. G. Helmer, IAEA – CRP Report to be published (2004) [I_γ].
- 2004Mo07 J. Morel, S. Speman, M. Rasko, E. Terechtchenko, J. U. Delgado, Appl. Radiat. Isot. 60(2004)341 [I_γ].

**²²⁷Ac – Comments on evaluation of decay data
by V. P. Chechev and N.K. Kuzmenko**

This evaluation was completed in June 2008 with a literature cut off by the same date. The SAISINUC software and associated supporting computer programs were used in assembling the data following the established protocol within DDEP (2002Be).

1. Decay Scheme

The ²²⁷Ac decay scheme is based on the evaluation of Browne (2001Br31). ²²⁷Ac disintegrates (1,380 (4) %) by alpha transitions to the ground state and excited states of ²²³Fr and (98,620 (4) %) by beta transitions to the ground state and excited states of ²²⁷Th. The decay scheme cannot be considered well established since only approximate values are available for beta and gamma transition probabilities in the β⁻-decay of ²²⁷Ac and the measurements of weak alpha transitions probabilities in the α-decay of ²²⁷Ac are not sufficiently accurate.

2. Nuclear Data

Q(α) value is from 2003Au03.

The evaluated half-life of ²²⁷Ac is based on the experimental results given in Table 1.

Table 1. Experimental values of the ²²⁷Ac half-life (in years)

Reference	Author(s)	Value
1950Ho79	Hollander and Leininger	22,0 (3)
1955To07	Tobailem	21,6 (4)
1956Sh43	Shimanskaya and Yashugina	21,2 (8)
1959Ro51	Robert	21,6 (3)
1963Ei10	Eichelberger et al.	21,7714 (+56 -33)
1967JoZX	Jordan and Blanke	21,7728 (+29 -32)

The weighted mean of the 6 values is 21,772. The internal uncertainty is 0,0022, if we use the smallest uncertainties from 1963Ei10 and 1967JoZX, and 0,0028, if we use the largest uncertainties from these measurements. $\chi^2/\nu = 0,34$ and $0,33$, respectively.

Our recommended value for the ²²⁷Ac half-life is 21,772 (3) years.

2.1 Alpha Transitions

The energies of the alpha transitions have been obtained from $Q(\alpha)$ value and the level energies given in Table 2 from 2001Br31 where they were deduced from a least squares fit to gamma-ray energies.

The comparison of the adopted energies of alpha particles for most intense transitions with the measured values is shown in Table 3 (columns 3 and 4). The measured energies of the alpha particles (Table 3) have been adjusted for changes in the calibration standards (1986Ry04, 1991Ry01): +3,5 keV correction for values from 1966Ba19, +5 keV correction for values from 1959No41.

Table 2. ^{223}Fr levels populated in the ^{227}Ac α -decay

Level	Level energy, keV	Spin and parity	Half-life	α -transition energy, keV	Probability of alpha transitions ($\times 100$)
0	0	$3/2^-$	22,00 (7) min	5042,19 (14)	0,658 (14)
1	12,89 (5)	$(5/2^-)$		5029,30 (15)	0,546 (17)
2	54,97 (7)	$1/2^-$		4987,22 (16)	0,0015
3	82,13 (6)	$(7/2^-)$		4960,06 (15)	0,087 (7)
4	99,63 (6)	$(3/2^-)$		4942,56 (15)	} 0,08 (1)
5	101,00 (6)	$(5/2^-)$		4941,19 (15)	
6	134,51 (6)	$(3/2^+)$		4907,68 (15)	0,001
7	149,3 (3)	$(1/2^+)$			
8	160,48 (7)	$(3/2^+)$		4881,71 (16)	0,014 (7)
9	172,08 (6)	$(5/2^+)$		4870,11 (15)	0,0011
10	187,18 (10)	$(5/2^-)$		4855,01 (17)	} 0,025 (7)
11	189,10 (7)	$(7/2^-)$		4853,09 (16)	
12	219,61 (9)	$(7/2^+)$		4822,58 (17)	} 0,0012
13	222,75 (10)	$(7/2^+)$		4819,44 (17)	
14	242,63 (7)	$(5/2)$		4799,56 (16)	} 0,006 (3)
15	243,85 (13)	$(5/2)$		4798,34 (19)	
16	244,66 (15)	$(7/2)$		4797,53 (21)	
17	298,7 (3)	$(9/2)$			
18	365,47 (10)			4676,72 (17)	$\approx 3 \cdot 10^{-4}$
19	379 (7)			4663 (7)	$\approx 4 \cdot 10^{-5}$
20	449 (5)			4593 (5)	$\approx 4 \cdot 10^{-5}$
21	503 (7)			4539 (7)	$\approx 7 \cdot 10^{-5}$
22	515,20 (22)	$3/2^-$		4526,99 (26)	$\approx 7 \cdot 10^{-4}$
23	540,74 (25)	$(5/2^+)$		4501,45 (29)	$\approx 8 \cdot 10^{-5}$
24	601 (7)	$(5/2^-)$		4441,19 (16)	$\approx 4 \cdot 10^{-5}$

The recommended probabilities of the $\alpha_{0,i}$ -transitions with $i = 0, 1, 3, 4, 8, 11, 14$ are from 1959No41. The remaining ones are from 1966Ba19. A comparison of the α -transition probabilities, taken directly from measurements of 1959No41, 1966Ba19 with those deduced from $P(\gamma+ce)$ intensity balance, is given in Table 3. The total probability of α -transitions is from 1970Ki12 (1,3800 (36) %), see also 1974Mo05 (1,359 (14) %). The α -decay hindrance factors have been calculated using the ALPHAD computer program from the ENSDF evaluation package with $r_0 = 1,538$ fm, average of $r_0(^{222}\text{Rn}) = 1,5397$ (4) fm, $r_0(^{222}\text{Ra}) = 1,5383$ (8) fm and $r_0(^{224}\text{Ra}) = 1,5332$ (8) fm, see 2001Br31.

Table 3. Energies and probabilities (×100) of most intense α-transitions in the ²²⁷Ac decay

Level	Level energy, keV	Energies of a-particles, obtained from Q(α), keV	Measured energies of a-particles, keV	Probabilities (×100), adopted from 1959No41, 1966Ba19	Probabilities (×100), deduced from intensity balance
0	0	4953,23 (14)	4953,26 (14)	0,658 (14)	0,48 (24)
1	12,89 (5)	4940,57 (15)	4940,7 (8)	0,546 (16)	0,63 (15)
3	82,13 (6)	4872,55 (15)	4872,7 (2)	0,087 (7)	0,09 (3)
4	99,63 (6)	4855,36 (15)	4855 (2)	} 0,08 (1)	} 0,10 (6)
5	101,00 (6)	4854,01 (15)			
6	134,51 (6)	4821,09 (15)	4822 (4)	0,014 (7)	0,0090 (26)
10	187,18 (10)			} 0,025 (7)	} 0,028 (10)
11	189,10 (7)	4767,47 (15)	4768 (3)		

2.2 Beta Transitions

The energies of β⁻ transitions have been obtained from Q⁻(²²⁷Ac) and ²²⁷Th level energies given in Table 4. The β⁻-emission probabilities per 100 β⁻ particles in ²²⁷Ac β⁻-decay have been taken from 1995Li04 . The value of ΣP_{β⁻}(i) has been obtained as (100 % - ΣPa (i)) = 98,620 (4) %. This is the total probability of beta transitions to the ground state and excited states of ²²⁷Th.

Table 4. ²²⁷Th levels populated in the ²²⁷Ac β⁻-decay

Level	Level Energy, keV	Spin and Parity	Half-life	β ⁻ -emission probability per 100 β ⁻ particles
0	0,0	1/2 +	18,68 (9) d	≈ 54
1	9,3	(5/2+)		≈ 35
2	24,5	3/2+		≈ 10
3	37,9	3/2-		0,3

2.3 Gamma Transitions and Internal Conversion Coefficients

The evaluated energies of the gamma -ray transitions are virtually the same as the gamma -ray energies because nuclear recoil is negligible.

The gamma-ray transition probabilities in ²²³Fr have been deduced from their gamma -ray emission probabilities and the calculated total ICCs. The gamma -ray transition probabilities in ²²⁷Th have been adopted from 199 5Li04. ICCs have been calculated by a program supplied with the SAISINUC software (2002Be). This code uses interpolated values of Band et al. (2002Ba85). The multipolarities and mixing ratios δ of the gamma -ray transitions in ²²³Fr and ²²⁷Th have been taken from 2001Br31. The uncertainties in the ICCs for pure multipolarities have been taken as 2 %.

3. Atomic Data

The atomic data (fluorescence yields, X-ray energies and relative probabilities, Auger electrons energies and relative probabilities) were obtained using the SAISINUC software (2002Be).

4.1 Alpha Emissions

Details are given in Section 2.1.

4.2 Beta Emissions

Details are given in Section 2.2.

5. Photon Emissions

5.1 X-Ray Emissions

The absolute emission probabilities of Fr KX and LX -rays and Th LX-rays have been calculated using the EMISSION code (2000Schönfeld). An experimental Fr KX -rays intensity value of 0,0136 (16) % (from 1995Sh03) agrees well with 0,0145 (24) %, deduced by the evaluators.

5.2 Gamma-Ray Emissions

Gamma-Ray Energies

The energies of gamma-rays in ^{223}Fr have been adopted from 1995Sh03. The energies of gamma-rays $\gamma_{1,0}$ and $\gamma_{2,1}$ in ^{227}Th have been adopted from 1959No41. The energies of gamma-rays $\gamma_{2,0}$ and $\gamma_{3,1}$ in ^{227}Th have been adopted from 1997Mu08.

Gamma-Ray Emission Probabilities

The absolute emission probabilities of gamma-rays in ^{223}Fr are from 1995Sh03. The absolute emission probabilities of gamma-rays in ^{227}Th have been deduced from the absolute β^- -emission probabilities in the ^{227}Ac β^- -decay and α_T using the ratio of $P(\gamma_{37,9\text{-keV}}) / P(\gamma_{28,6\text{-keV}}) = 9,0 (12) / 7,7 (10) = 1,17 (22)$ from ^{227}Pa EC decay (1995Li04), and the value of $P(\gamma_{24,3\text{-keV}}) / P(\gamma_{15,2\text{-keV}}) = 20 / 0,44 = 45,5$ from alpha decay of ^{231}U (2001Br31).

6. Electron Emissions

The energies of conversion electrons have been obtained from the gamma transition energies and atomic electron binding energies. The emission probabilities of conversion electrons have been deduced from the evaluated $P(\gamma)$ and ICC values.

The number of K- and L- Auger electrons per 100 disintegrations has been calculated using the EMISSION code (2000Schönfeld).

Average β^- energies have been calculated using the LOGFT computer program.

7. References

- 1950Ho79 J.M. Hollander and R.F. Leininger, Phys. Rev. 80, 915 (1950)
(Half-life)
- 1955To07 J. Tobailem, J. Phys. Radium 16, 48 (1955)
(Half-life)
- 1956Sh43 N.S. Shimanskaya and E.A. Yashugina, At.Energ. 1, 133 (1956); J.Nuclear Energy 5, 161 (1957)
(Half-life)
- 1959No41 G.I. Novikova, E.A. Volkova, L.I. Goldin, D.M. Ziv and L.I. Tretyakova, Zh. Eksp. Teor. Fiz. 37, 928(1959) (Sov. Phys. JETP 10, 663 (1960))
(Total α -transition probability, gamma-ray energies and α - energies, electron conversion)
- 1959Ro51 J. Robert, Ann. Phys.(Paris) 4, 89 (1959)
(Half-life)
- 1963Ei10 J.F. Eichelberger, G.R. Grove, L.V. Jones and E.A. Rembold, MLM-1155, p.12 (1963)
(Half-life)
- 1966Ba19 G. Bastin, C.-F. Leang, and R.J. Walen, Compt.Rend. 262B, 370 (1966)
(α -transition probabilities and α - energies)
- 1967JoZX K.C. Jordan and B.C. Blanke, Proc. Symp. Standardization of Radionuclides, Vienna, Austria (1966), IAEA, Vienna, p.567 (1967); CONF-661012-4 (1967)
(Half-life)
- 1970Ki12 H.W. Kirby, J. Inorg. Nucl. Chem. 32, 2823 (1970)
(Total α -transition probability)
- 1974Mo05 M. Monsecour, P. De Regge, A. Demildt, and L.H. Baetsle, J. Inorg. Nucl. Chem. 36, 719 (1974)
(Total α -transition probability)
- 1986Ry04 A. Rytz, R.A.P. Wiltshire, and M. King, Nucl. Instrum. Methods Phys. Res. A253, 47 (1986)
(α -transition energies)
- 1991Ry01 A. Rytz, At. Data Nucl. Data Tables 47, 205 (1991)
(α -transition energies)
- 1995Li04 C.F. Liang, P. Paris, R.K. Sheline, D. Nosek, and J. Kvasil, Phys. Rev. C51, 1199 (1995)
(Beta emission probabilities)
- 1995Sh03 R.K. Sheline, C.F. Liang, P. Paris, J. Kvasil, and D. Nosek, Phys. Rev. C51, 1708 (1995)
(Gamma-ray energies)
- 1997Mu08 U. Muller, P. Sevenich, K. Freitag, C. Gunther, P. Herzog, G.D. Jones, C. Kliem, J. Manns, T. Weber, B. Will, and the ISOLDE Collaboration, Phys. Rev. C55, 2267 (1997)
(Gamma-ray energies)
- 2000Schönfeld E. Schönfeld and H. Janßen, Nucl. Instr. Meth. Phys. Res. A369, 527 (2000)
(EMISSION computer code)
- 2001Br31 E. Browne, Nucl. Data Sheets 93, 763 (2001)
(Decay scheme of ^{227}Ac , gamma-ray multiplicities, admixture coefficients) p. 975 (Decay scheme of ^{231}U , gamma- ray emission probabilities of $P(\gamma_{2,0})$ and $P(\gamma_{2,1})$ in ^{227}Th) p. 978
- 2002Ba85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, P.O. Tikkanen and S. Raman, Atom. Data and Nucl. Data Tables 91, 1 (2002)
(Theoretical internal conversion coefficients)
- 2002Be M.M. Bé, R. Helmer and V. Chisté, J. Nucl. Sci. Tech., suppl.2, 481 (2002)
(Saisinuc software)
- 2003Au03 G. Audi, A.H. Wapstra and C. Thibault, Nucl. Phys. A729, 337 (2003)
(Q value)

²²⁷Th – Comments on evaluation of decay data by E. Browne

1) Evaluation Procedures

The *Limitation of Relative Statistical Weight* (LWM) [1985ZiZY] method, used for averaging numbers throughout this evaluation, provided a uniform approach for the analysis of discrepant data. The uncertainty assigned in this evaluation to the recommended value is always greater than or equal to the smallest uncertainty in any of the experimental values used in the calculation. This evaluation was completed in August 2001, with minor editing done in March 2002.

2) Decay Scheme

²²⁷Th decays 100% by emission of α particles, 24,2(9)% populates the ground state of ²²³Ra. Evaluator normalized the decay scheme using measured values of the absolute emission probability of the 50.13-keV γ -ray, as described here in Section 5. There are several low-energy γ -rays, many of them with very large and not well-known conversion coefficients that have limited the accuracy of their respective total transition probabilities. For this reason the individual feedings, deduced from transition-intensity balances at each level, are also inaccurate. Thus such feedings have not been shown here. The α -particle probabilities (in percent) to individual levels presented in the decay scheme are experimental values from α -spectroscopic measurements of 1964Ba33. α -hindrance factors given in the decay scheme are from 2001Br31, calculated by using a radius parameter r_0 (²²³Ra) = 1.536, average of r_0 (²²²Ra) = 1.5383(8) and r_0 (²²⁴Ra) = 1.5332(8) (1998Ak04). The level energies, spins, parities, as well as γ -ray multipolarities and mixing ratios shown in the decay scheme are from 2001Br31.

3) Nuclear Data

Table 1. ²²⁷Th measured half-life values

Half-life (days)	Reference
18.169 (84)	1954Ha60
18.729 (48)	1963Ei10
18.7176 (52)	1967JoZX
18.738 (54)	1987Mi10

The (unpublished) value given in 1954Ha60 significantly disagrees with the other measured values. The ²²⁷Th source used in 1954Ha60 contained several daughter radionuclides from the decay chain. Moreover, they used proportional counters to detect alpha particles, without any elemental discrimination. This situation may have introduced a systematic error in their half-life. Thus, the evaluator excluded this value from the statistical analysis. The recommended half-life of ²²⁷Th is the weighted average (LWM) ($\chi^2/\nu = 0.1$) of the other three measured values, 18.718(5) days.

$Q_\alpha = 6146.43(15)$ keV is from 1995Au05.

4) Alpha Particles

Alpha particle energies and absolute probabilities presented in Section 4 are evaluated values from 2001Br31. Most α -particle energies are from 1964Ba33, increased by 1.7 keV to correct them for a systematic deviation (2001Br31). The energies of $\alpha_{(0,12)}$, $\alpha_{(0,3)}$, and $\alpha_{(0,0)}$ are from 1971Gr17, as recommended by 1991Ry01. Absolute α -particle probabilities are from 1964Ba33.

5) Gamma Rays

Energies

The recommended γ -ray energies given in Sections 2.2 and 6.2 are weighted averages (LWM) of values from 1993Ab01, 1990Br23, 1972He18, and 1969Br27, unless otherwise specified in Table 2.

Emission Probabilities

The recommended relative γ -ray emission probabilities given in Table 2 are weighted averages (LWM) of values from 1993Ab01, 1972He18, and 1969Br27, unless otherwise specified in this table.

Excepting the 304.50-keV gamma ray, all the conversion coefficients given in Section 2.2 are theoretical values from 1978Ro22 interpolated by using program ICC [1] for the recommended γ -ray energies and multiplicities. The 304.50-keV gamma ray has an E0 component, thus the conversion coefficients given here for this transition are experimental values.

The γ -ray emission (and total transition) probabilities given in Sections 6.2 and 2.2, respectively, have been normalized to an absolute scale (per 100 α decays) using a normalization factor $N = 0.126(6)$. Evaluator deduced this value from $I_{\text{avg}}(50.13 \gamma) = 8.20(17)\%$, weighted average of the following measured absolute γ -ray emission probabilities: $I_{\gamma}(50.13) = 8.18(17)\%$ (1990Ko40) and $I_{\gamma}(50;13) = 8.4(6)\%$ (1969Pe17).

A normalization factor $N = 0.127(11)$ may be obtained by using the decay scheme and the sum of all the relative γ -ray transition probabilities (photons + electrons) to the ground state and to the first excited state at 29 keV, then equating this sum to 72.9(10)% (that is, to $100\% - I_{\alpha}(\text{gs} + 29\text{-keV level}) = 100\% - 27.1(10)\% = 72.9(10)\%$). This value, although less precise, is in good agreement with the one given before, and it confirms the correctness and consistency of the decay scheme.

6) Atomic Data

X-ray and Auger (relative and absolute) electron emission probabilities given in Sections 3, 6.1 and 5, respectively, have been calculated by means of the computer code EMISSION (version 3.01, Nov. 3, 1999) [2]), which makes use of the atomic data from 1996Sc06, from reference [3], and from the evaluated γ -ray data given in Sections 2.2 and 6.2. In addition, internal conversion electron energies and absolute emission probabilities for the strongest lines are presented in Section 5. Electron energies have been calculated using electron binding energies from 1977La19, and γ -ray energies from Section 2.2. Absolute electron emission probabilities have been calculated using absolute γ -ray emission probabilities given in Section 6.2 and conversion coefficients from Section 2.2.

7) References

[1] V. M. Gorozhankin, N. Coursol, E. A. Yakushev. ICC99: A computer program for interpolating internal conversion coefficients from Hager and Seltzer, Rösel et al., and from Band et al. (1999). (Internal conversion coefficients)

- [2] E. Schönfeld, H. Janssen. *Applied Radiation Isotopes* **52**, 595 (2000). (X-ray and Auger electron emission probabilities).
- [3] Eckart Schönfeld, Gisela Rodloff. Report PTB-6.11-98-1, Braunschweig, October 1998. (Auger electron energies).
- 1954Ha60 - G. R. Hagee, M. L. Curtis, G. R. Grove. *Phys. Rev.* **96**, 817A (1954). (Half-life)
- 1963Ei10 - J. F. Eichelberger, G. R. Grove, L. V. Jones, E. A. Rembold. *MLM-1155*, p.12 (1963). (Half-life)
- 1967JoZX - K. C. Jordan, B. C. Blanke. *Proc. Symp. Standardization of Radionuclides, Vienna, Austria* (1966), Intern. At. Energy Agency, Vienna, p.567 (1967); CONF-661012-4 (1967). (Half-life)
- 1969Pe17 - A. Peghaire. *Nucl. Instr. Methods* **75**, 66 (1969). (Absolute γ -ray emission probability)
- 1969Br27 - C. Briançon, R. Walen. *J. Phys. (Paris)* **30**, 753 (1969) (γ -ray energies and emission probabilities).
- 1971Gr17 - B. Grennberg, A. Rytz. *Metrologia* **7**, 65 (1971) (α -particle energies).
- 1972He18 - W. H. A. Hesselink, A. H. Wapstra, J. G. Kromme, E. J. Haighton, M. Van Kampen, W. Hutjes, K. E. M. Dijkman. *Nucl. Phys.* **A191**, 283 (1972). (γ -ray energies and emission probabilities).
- 1977La19 - F. P. Larkins. *At. Data Nucl. Data Tables* **20**, 311 (1977). (Atomic electron binding energies).
- 1978Ro22 - F. Rosel, H. M. Fries, K. Alder, H. C. Pauli. *At. Data Nucl. Data Tables* **21**, 92 (1978). (γ -ray theoretical internal conversion coefficients).
- 1985ZiZY - W. L. Zijp, Report ECN FYS/RASA-85/19 (1985). (Discrepant Data. Limited Relative Statistical Weight Method).
- 1987Mi10 - G. J. Miller, J. C. McGeorge, I. Anthony, R. O. Owens. *Phys. Rev.* **C36**, 420 (1987). (Half-life).
- 1990Br23 - Ch. Briancon, S. Cwiok, S. A. Eid, V. Gree, W. D. Hamilton, C. F. Liang, R. J. Walen. *J. Phys. (London)* **G16**, 1735 (1990). (γ -ray energies).
- 1990Ko40 - K. Komura, M. Yamamoto, K. Ueno. *Nucl. Instrum. Methods. Phys. Res.* **A295**, 461 (1990). (Atomic data, X-rays, Auger electrons).
- 1991Ry01- A. Rytz. *At. Data Nucl. Data Tables* **47**, 205 (1991). (α -particle energies).
- 1993Ab01 - A. Abdul-Hadi, V. Barci, B. Weiss, H. Maria, G. Ardisson, M. Hussonnois, O. Constantinescu. *Phys. Rev.* **C47**, 94 (1993). (γ -ray energies and emission probabilities).
- 1995Au04 - G. Audi, A. H. Wapstra. *Nucl. Phys.* **A595**, 409 (1995). (Q_α)
- 1996Sc06 - E. Schönfeld, H. Janssen. *Nucl. Instrum. Methods. Phys. Res.* **A369**, 527 (1996). (Atomic data, X-rays, Auger electrons).
- 1998Ak04 - Y.A. Akovali. *Nucl. Data Sheets* **84**, 1 (1998). (Alpha decay. Radius parameter of even-even nuclei).
- 2001Br31 - E. Browne. *Nucl. Data Sheets* **93**, 763 (2001). (Evaluated data (ENSDF) for nuclei with $A=227$. Corrected α -particle energies).

Table 2. ²²⁷ Th Alpha Decay - Gamma-Ray Energies and Relative Emission Probabilities											
1993Ab01(E _γ)	1993Ab01(I _γ)	1990Br23(E _γ)	1990Br23(I _γ)	1972He18(E _γ)	1972He18(I _γ)	1969Br27(E _γ)	1969Br27(I _γ)	Adopted E _γ ^a	Adopted I _γ ^b	χ ² /ν(E _γ)	χ ² /ν(I _γ)
6.5 (3)	0.7 (2)	6.3				6		6.5 (3)*	0.7 (2)*		
8.3 (2)	0.06 (2)	8				8.0 (2)		8.15 (20)	0.06 (2)*	1.1	
20.19 (5)	1.9 (2)	20.30 (5)	0.769			20.3 (2)	1.5 (5)	20.25 (5)	1.84 (20)	1.3	0.55
20.8 (2)	0.05 (2)	20.95 (5)						20.94 (5)	0.05 (2)*	0.53	
22.0 (2)	0.07 (7)							22.0 (2)*	0.07 (7)*		
24.13 (5)	0.68 (5)							24.13 (5)*	0.68 (5)*		
27.32 (5)	0.23 (4)	27.50 (10)	0.154					27.41 (9)	0.23 (4)*	1.6	
		29.60 (3)	0.046					29.60 (3)**	0.046**		
29.86 (5)	0.56 (8)	29.86 (1)	0.769			29.9 (2)	0.8 (2)	29.86 (1)	0.59 (8)	0.02	0.87
31.56 (5)	0.51 (8)	31.58 (1)	0.692			31.6 (2)	0.62 (17)	31.58 (1)	0.53 (8)	0.08	0.34
33.3 (2)	0.06 (2)	33.40 (8)	0.108					33.39 (8)	0.06 (2)*	0.22	
40.20 (3)	0.12 (3)	40.20 (10)	0.154	40.1				40.20 (3)	0.12 (3)*	0.01	
41.91 (5)	0.12 (3)	42.2 (3)	0.308	42.2 (5)	0.70 (26)	42.1 (3)	0.31 (6)	41.93 (5)	0.22 (10)	0.52	4.3
43.75 (5)	1.6 (1)	43.80 (5)	1.538	43.5 (5)	2.1 (6)	43.8 (2)	1.77 (21)	43.77 (5)	1.65 (10)	0.27	0.69
				43.8 (5)	0.43 (17)			43.8 (5)&	0.43 (17)&		
44.33 (5)	0.4 (1)	44.10 (5)	0.046	44.1	0.06 (3)	44.1		44.22 (12)	0.41 (10)	5.3	0.04
		44.40 (5)		44.3 (5)	0.11 (5)	44.4	0.15	44.40 (5)**			
		46.45 (5)						46.45 (5)**			
				48.3 (5)							
48.1 (2)	0.12 (3)	48.30 (3)	0.077	48.5 (5)	0.39 (10)	48.3	0.08 (1)	48.30 (3)	0.11 (4)	0.57	4.5
49.75 (5)	3.5 (5)	49.90 (7)	4.615	49.8 (3)	1.7 (13)	49.9	4.6 (14)	49.82 (5)	3.3 (7)	1.2	1.2
50.11 (2)	63 (2)	50.13 (1)	61.538	50.2 (2)	75.7 (52)	50.2 (2)	65 (3)	50.13 (1)	65 (3)	0.36	2.7
50.8 (2)	0.11 (5)	50.85 (5)	0.031	50.7 (5)	0.14 (6)			50.85 (5)	0.12 (5)	0.07	0.15
				51.2				51.2&			
54.1 (1)	0.05 (1)	54.20 (4)	0.008	54.2				54.19 (4)	0.05 (1)*	0.86	
		56.00 (6)	0.038	56.1	0.01 (1)	56.1	0.08 (2)	56.00 (6)**	0.038**		
56.3 (2)	0.12 (2)	56.55 (3)	0.077			56.6	0.13 (4)	56.42 (14)	0.07 (6)	0.78	10
				59.6 (5)	0.08 (3)			59.6 (5)&	0.08 (3)&		
61.42 (5)	0.70 (8)	61.44 (2)	0.846	61.5		61.5 (2)	0.69 (14)	61.44 (2)	0.70 (8)	0.12	
				62							

1993Ab01(E γ)	1993Ab01(I γ)	1990Br23(E γ)	1990Br23(I γ)	1972He18(E γ)	1972He18(I γ)	1969Br27(E γ)	1969Br27(I γ)	Adopted E γ ^a	Adopted I γ ^b	χ^2/ν (E γ)	χ^2/ν (I γ)
62.33 (5)	1.5 (2)	62.45 (5)	1.385	62.5 (3)	2.2 (5)	62.5 (2)	1.54 (23)	62.45 (5)	1.57 (20)	2.6	0.86
62.7 (2)	0.05 (2)	62.65 (4)	0.077	62.7 (3)	0.08 (3)			62.68 (3)	0.056 (20)	0.5	0.45
64.5 (2)	0.19 (3)	64.30 (10)	0.115	64.5 (5)	0.24 (9)			64.35 (10)	0.20 (3)	0.45	0.28
65.2 (1)	0.13 (3)	64.70 (10)	0.077					64.95 (25)	0.13 (3)*	12	
				66.2 (5)	0.05 (3)			66.2 (5)&	0.05 (3)&		
				66.4 (5)	0.06 (3)			66.4 (5)&	0.06 (3)&		
		68.70 (10)	0.046	68.7	0.01 (1)			68.70 (10)**	0.046**		
68.72 (5)	0.53 (4)	68.75 (3)	0.346	68.8 (5)	0.24 (10)	68.8 (2)	0.44 (7)	68.74 (3)	0.45 (8)	0.12	3.2
69.8 (3)	0.08 (3)			69.8 (5)	0.08 (3)			69.8 (3)	0.08 (3)		
72.85 (5)	0.32 (4)	72.80 (10)	0.231	72.9	0.03 (3)	72.9 (1)	0.22 (4)	72.85 (5)	0.19 (15)	0.25	12
73.8 (2)	0.07 (2)	73.60 (5)	0.077	73.7 (5)	0.15 (5)	73.7 (1)	0.15 (2)	73.63 (5)	0.11 (4)	0.53	4.3
75.00 (5)	0.29 (3)	75.1	0.154	75.3 (5)	0.08 (5)	75.1 (3)	0.18 (5)	75.01 (5)	0.21 (8)	0.23	7
						77.4 (4)	0.08	77.4 (4)#	0.08#		
79.66 (3)	15.1 (5)	79.72 (1)	15.385	79.7 (2)	15.7 (44)	79.8 (2)	15.4 (15)	79.69 (2)	15.1 (5)	0.78	0.04
		84									
89.17 (8)	0.03 (1)	90.0 (3)	0.031	89.9				89.6 (4)	0.03 (1)*	3.8	
93.86 (5)	11.9 (5)	93.90 (10)	10.000	94.0 (2)	11.7 (3)	94.0 (2)	10.8 (11)	93.88 (5)	11.7 (3)	0.31	0.28
94.9 (1)	0.30 (4)	94.99 (5)	0.092	95		95	0.09 (1)	94.97 (5)	0.19 (11)	0.65	14
96.02 (5)	0.6 (1)	96.1 (2)	0.462	96.1 (5)	0.39 (17)	96.1 (2)	0.54 (13)	96.03 (5)	0.54 (10)	0.1	0.57
99.5 (2)	0.20 (5)	99.60 (10)	0.100	99.5				99.58 (10)	0.20 (5)*	0.2	
		99.60 (20)	0.015	99.7				99.60 (20)**	0.1**		
100.2 (2)	0.7 (2)	100.27 (3)	0.731	100.4 (5)	0.7 (3)	100.3	0.62 (12)	100.27 (3)	0.65 (12)	0.1	0.08
		102.50 (10)	0.009	102.5				102.50 (10)**	0.009**		
106.1 (2)	0.03 (1)	105.20 (10)	0.077					105.20 (10)**	0.077**		
107.9 (2)	0.05 (2)	107.75 (7)	0.046	108	0.06 (3)	107.5 (5)	0.07 (2)	107.76 (7)	0.060 (20)	0.39	0.25
108.9 (3)	0.03 (1)	108.00 (10)	0.000	109.6 (5)	0.05 (2)			109.2 (4)	0.041 (12)	0.98	0.84
110.7 (2)	0.04 (1)	110.65 (5)	0.062	110.6	0.01 (1)			110.65 (5)	0.025 (16)	0.06	4.8
				112.6 (5)	0.07 (3)			112.6 (5)&	0.07 (3)&		
113.06 (2)	6.6 (3)	113.16 (2)	5.385	113.1 (2)	4.7 (6)	113.1 (2)	5.5 (6)	113.11 (5) ^c	5.9 (8)	4.2	3.6
117.20 (5)	1.7 (1)	117.20 (5)	1.308	117.0 (3)	1.4 (3)	117.2 (2)	1.38 (14)	117.20 (5)	1.54 (11)	0.15	1.6
		117.20 (5)	0.077	117.5 (5)	0.10 (3)			117.5 (5)&	0.10 (3)&		

1993Ab01(E γ)	1993Ab01(I γ)	1990Br23(E γ)	1990Br23(I γ)	1972He18(E γ)	1972He18(I γ)	1969Br27(E γ)	1969Br27(I γ)	Adopted E γ ^a	Adopted I γ ^b	χ^2/ν (E γ)	χ^2/ν (I γ)
123.6 (1)	0.14 (2)	123.5 (2)	0.154	123.6 (5)	0.07 (2)	123.6	0.08	123.58 (10)	0.11 (4)	0.1	6.1
124.4 (2)	0.04 (2)	125	0.023	124.4	0.01 (1)	124.4	0.02	124.44 (20)	0.032 (17)	0.31	0.28
				124.7 (5)	0.03 (2)						
128.02 (2)	0.025 (4)							128.02 (2)*	0.025 (4)*		
129.4 (2)	0.010 (5)							129.4 (2)*	0.010 (5)*		
134.6 (1)	0.30 (5)	134.5 (3)	0.154	134.2 (3)	0.26 (5)	134.6 (2)	0.23 (5)	134.6 (1)	0.26 (5)	0.56	0.49
138.4 (1)	0.11 (2)	138	0.018					138.4 (1)*	0.11 (2)*		
140.5 (3)	0.05 (2)	141.0 (5)	0.038	140.5 (3)	0.28 (5)			140.6 (3)	0.17 (2)	0.42	11
141.34 (5)	1.1 (1)	141.50 (5)	1.000	141.2 (3)	0.57 (13)	141.4 (2)	1.00 (15)	141.42 (5)	0.92 (18)	1.9	5.4
150.1 (2)	0.07 (3)	150.2 (2)	0.038	149.8 (5)	0.16 (3)	150.3 (5)	0.07 (2)	150.14 (20)	0.086 (24)	0.23	2.1
162.2 (1)	0.07 (2)	162.1 (3)	0.062	162.2 (5)	0.05 (2)	162.1 (5)	0.07	162.19 (10)	0.060 (20)	0.04	0.5
164.5 (1)	0.11 (2)	164.8	0.077			164.9 (5)	0.12 (3)	164.52 (10)	0.113 (20)	0.62	0.08
168.4 (1)	0.11 (2)	168.25 (15)	0.100	168.3 (3)	0.12 (3)	168.7 (5)	0.12 (3)	168.36 (10)	0.115 (20)	0.4	0.06
169.7 (2)	0.06 (2)	170.0 (1)	0.031	170.1 (5)	0.03 (2)			169.95 (10)	0.043 (17)	0.95	1.5
171.5 (2)	0.03 (1)			171.4				171.5 (2)*	0.03 (1)*		
173.45 (5)	0.16 (2)	173.40 (10)	0.123	173.4 (5)	0.10 (3)	173.5 (3)	0.12	173.45 (3)	0.135 (20)	0.09	1.5
				175.8 (3)	0.16 (3)			175.8 (3)&	0.16 (4)&		
181.1 (3)	0.02 (1)	181	0.015					181.1 (3)*	0.02 (1)*		
182.3 (2)	0.03 (1)							182.3 (2)*	0.03 (1)*		
184.65 (5)	0.29 (3)	184.65 (5)	0.262	184.7 (3)	0.23 (4)	184.7 (3)	0.31 (5)	184.65 (5)	0.28 (3)	0.02	0.73
197.5 (1)	0.07 (2)	197.60 (10)	0.077	197.6 (5)	0.09 (3)	197.8 (5)	0.12 (3)	197.56 (10)	0.10 (3)	0.25	0.4
200.5 (1)	0.17 (2)	200.5 (2)	0.154	200.5	0.02 (2)	201.0 (4)	0.25 (6)	200.50 (10)	0.10 (7)		23
201.7 (1)	0.16 (2)	201.60 (10)	0.138	201.8 (3)	0.19 (3)			201.64 (10)	0.184 (20)	1.1	0.88
				202.5 (5)	0.05 (2)			202.5 (5)&	0.05 (2)&		
204.2 (1)	1.7 (2)	204.14 (10)	1.538	204.2 (3)	2.0 (4)	204.3 (2)	1.6 (4)	204.14 (10)	1.76 (20)	0.19	0.44
204.9 (1)	1.2 (2)	205.02 (10)	0.769	205.2 (3)	1.5 (3)	205.0 (2)	1.2 (3)	204.98 (10)	1.27 (20)	0.45	0.38
206.05 (6)	1.9 (2)	206.10 (5)	1.538	206.1 (3)	2.3 (4)	206.2 (2)	1.7 (4)	206.08 (5)	1.97 (20)	0.25	0.89
		206.3	0.062	206.4	0.02 (2)			206.4&	0.02 (2)&		
210.58 (5)	9.4 (3)	210.65 (5)	8.462	210.6 (2)	11.0 (8)	210.7 (2)	8.5 (9)	210.62 (5)	9.7 (7)	0.39	2.4
212.76 (5)	0.63 (5)	212.65 (4)	0.615	212.6 (3)	0.74 (13)	212.2	0.38 (10)	212.70 (4)	0.61 (7)	1.3	1.3
				212.7 (3)	0.15 (4)	213	0.46 (12)	212.7 (3)&	0.15 (4)&		

216.0 (1)	0.002 (1)							216.0 (1)*	0.002 (1)*		
218.89 (5)	0.83 (8)	219.0 (2)	0.538	218.8 (3)	0.48 (9)	219.0 (2)	0.85 (10)	218.90 (5)	0.85 (8)	0.22	0.05
		219.0 (2)	0.231	219.0 (3)	0.39 (9)			219.0 (2)&	0.39 (9)&		
222.8 (2)	0.04 (1)	223.60 (15)	0.015					223.2 (4)*	0.04 (1)*	8	
225.9 (1)	0.07 (2)	225.5 (5)	0.015	224.7 (5)	0.13 (3)	225.5 (10)	0.03	225.5 (3)*	0.07 (2)*	1.3	
229.4 (2)	0.03 (1)	230.3 (3)	0.005	230.4				229.9 (5)*	0.03 (1)*	4.5	
234.7 (1)	3.4 (3)	234.80 (10)	3.615	234.9 (3)	5.0 (10)	234.9	3.1 (6)	234.76 (10)	3.5 (4)	0.37	1.4
235.94 (3)	100	235.97 (2)	100.000	236.0 (2)	100 (4)	236.0 (2)	100 (8)	235.96 (2)	100 (2)	0.26	
246.1 (1)	0.10 (3)	246.1 (3)	0.077	246.4 (5)	0.10 (3)	246.2 (3)	0.08 (3)	246.12 (10)	0.095 (17)	0.14	0.18
248.1 (1)	0.19 (4)							248.1 (1)*	0.19 (4)*		
				249.6 (5)	0.06 (2)			249.6 (5)&	0.06 (2)&		
250.1 (2)	0.08 (2)	250.15 (5)	3.231	250.2 (3)	2.4 (4)	250.2		250.15 (5)	0.069 (13)	0.04	0.52
250.19 (3)	4.0 (3)	250.35 (5)	1.077	250.4 (3)	0.61 (17)	250.4	3.1 (6)	250.27 (8)	3.5 (3)	2.7	1.6
252.50 (5)	0.9 (2)	252.6 (4)	0.769	252.5 (5)	1.0 (3)	252.6	0.77 (19)	252.50 (5)	0.86 (12)	0.03	0.21
254.62 (3)	5.6 (3)	254.67 (10)	5.385	254.7 (3)	7.9 (10)	254.7	3.9 (8)	254.63 (3)	5.5 (10)	0.15	4.9
256.22 (2)	54 (1)	256.25 (2)	56.154	256.2 (2)	55 (4)	256.3 (2)	57 (3)	256.23 (2)	54.3 (10)	0.42	0.46
260.6 (2)	0.04 (1)							260.6 (2)*	0.04 (1)*		
262.85 (5)	0.9 (1)	262.90 (10)	0.769	262.7 (5)	0.87 (17)	263.0 (2)	0.77 (9)	262.87 (5)	0.83 (6)	0.26	0.49
265.3 (2)	0.04 (1)							265.3 (2)*	0.04 (1)*		
267.0 (2)	0.08 (2)	267.1 (2)	0.019	267				267.05 (20)	0.08 (2)*	0.13	
267.7 (2)	0.06 (2)	268.0 (2)	0.077	267.9		268.0 (5)	0.05 (2)	267.86 (20)	0.055 (20)	0.6	0.13
				270.5							
270.6 (2)	0.16 (3)	270.5 (2)	0.062	270.7 (5)	0.28 (10)			270.56 (20)	0.22 (7)	0.1	0.72
272.90 (5)	3.9 (2)	272.90 (10)	3.846	273.0 (3)	4.3 (6)	273.0 (2)	3.9 (6)	272.91 (5)	3.94 (20)	0.11	0.2
279.7 (5)	0.35 (5)	279.7 (10)	0.462	279.7 (3)	0.78 (17)	279.8 (2)	0.38	279.80 (5)	0.42 (10)	0.03	2.7
280.4 (2)	0.02 (1)	281.0 (2)	0.054	281				280.7 (3)	0.02 (1)*	4.5	
281.42 (5)	1.4 (1)	281.40 (10)	1.231	281.4 (3)	1.3 (3)	281.4 (2)	1.3 (3)	281.42 (5)	1.38 (9)	0.01	0.09
284.2 (1)	0.4 (1)	284.4 (2)	0.385	284.3	0.22 (10)			284.24 (10)	0.31 (10)	0.8	1.6
285.6 (2)	0.25 (5)	285.50 (10)	0.385	285.6 (3)	0.48 (9)	285.4 (3)	0.38 (10)	285.52 (10)	0.34 (9)	0.14	2.2
286.06(2)	15 (1)	286.12 (2)	11.538	286.2 (2)	14.3 (7)	286.2 (2)	12.3 (6)	286.09 (2)	13.5 (12)	1.7	3.8
289.6 (1)	15 (3)	289.5 (3)	0.054	289.6	0.02 (2)			289.59 (10)	15 (3)*	0.1	

1993Ab01(E γ)	1993Ab01(I γ)	1990Br23(E γ)	1990Br23(I γ)	1972He18(E γ)	1972He18(I γ)	1969Br27(E γ)	1969Br27(I γ)	Adopted E γ ^a	Adopted I γ ^b	χ^2/ν (E γ)	χ^2/ν (I γ)
289.8 (1)	0.15 (3)	289.5 (3)	0.012					289.77 (10)	0.15 (3)*	0.9	
292.41 (5)	0.52 (6)	292.40 (10)	0.538	292.3 (5)	0.48 (10)	292.5 (3)	0.54 (13)	292.41 (5)	0.51 (6)	0.05	0.08
296.50 (5)	3.3 (3)	296.50 (5)	3.769	296.6 (3)	3.4 (6)	296.6 (2)	3.7 (5)	296.50 (5)	3.4 (3)	0.12	0.24
299.95 (3)	17.3 (5)	300.00 (3)	16.923	300.0 (2)	16.4 (12)	300.0 (2)	16.9 (17)	299.98 (3)	17.1 (5)	0.47	0.25
300.8 (2)	0.11 (2)	300.35 (3)	0.846	300.3 (3)	2.4 (4)			300.50 (16)	0.11 (2)*	1.8	
304.47 (3)	8.6 (5)	304.52 (2)	7.692	304.4 (3)	12 (1)	304.5 (2)	7.7 (8)	304.50 (2)	8.9 (10)	0.68	5.2
306.1 (3)	0.08 (3)							306.1 (3)*	0.08 (3)*		
308.40 (5)	0.14 (2)	308.40 (10)	0.108	308.5 (5)	0.13 (3)	308.4 (3)	0.11 (3)	308.40 (3)	0.131 (20)	0.01	0.35
312.69 (3)	4.0 (3)	312.70 (10)	3.846	312.6 (3)	4.5 (9)	312.7 (2)	3.9 (6)	312.69 (3)	4.0 (3)	0.03	0.16
		314.75 (10)	0.269	314.8 (5)	0.22 (9)			314.75 (10)	0.27**	0.01	
314.85 (4)	3.7 (3)	314.85 (10)	3.385	314.8 (3)	4.7 (9)	314.9 (2)	3.6 (5)	314.85 (4)	3.8 (3)	0.03	0.62
		318.4 (2)	0.046	318.8 (5)	0.05 (2)			318.46 (20)	0.052 (17)&	0.55	
319.24 (5)	0.30 (3)	319.2 (2)	0.231	319.2 (5)	0.16 (4)	319.2 (2)	0.26 (3)	319.24 (5)	0.25 (5)	0.03	4
324.8 (2)	0.08 (2)	324.9 (2)	0.046					324.88 (20)	0.08 (2)*	0.29	
325.7 (3)	0.07 (3)	326.10 (10)	0.231	325.2 (5)	0.04 (2)			325.99 (18)	0.049 (20)	0.89	0.69
326.7				326.2	0.01 (1)	326.4 (5)	0.23				
329.85 (2)	21.7 (5)	329.85 (3)	21.538	329.9 (2)	25.2 (14)	329.9 (2)	21.5 (19)	329.85 (2)	22.8 (12)	0.04	2.2
332.2 (2)	0.013							332.2 (2)*	0.013 (4)*		
334.36 (2)	8.2 (3)	334.38 (2)	8.462	334.4 (3)	10.0 (9)	334.5 (2)	8.5 (11)	334.37 (2)	8.8 (6)	0.31	1.3
339.6 (2)	0.03 (1)	339.80 (10)	0.012	339.8				339.76 (10)	0.03 (1)*	0.8	
342.56 (4)	3.4 (1)	342.50 (10)	3.231	342.5 (3)	1.7 (4)	342.5 (2)	3.2 (6)	342.55 (4)	2.7 (7)	0.13	9.3
346.48 (5)	0.10 (1)	346.45 (1)	0.077	346.3 (5)	0.07 (2)	346.5 (5)	0.08 (3)	346.45 (1)	0.093 (10)	0.15	1
				348.5 (5)	0.05 (2)			348.5 (5)&	0.052 (17)&		
350.66 (2)	0.9 (2)	350.40 (10)	0.923	350.5 (3)	0.70 (17)	350.5 (2)	0.92 (14)	350.54 (7)	0.85 (14)	1.3	0.54
		352.60 (10)	0.100	352.6 (5)	0.08 (2)	352.7 (3)	0.10 (3)	352.61 (10)	0.078 (17)&	0.01	
362.7 (1)	0.04 (1)	362.4 (2)	0.038	362.5 (5)	0.03 (2)	362.6 (2)	0.04 (1)	362.63 (10)	0.393 (10)	0.63	0.04
369.5 (5)	0.05 (1)	369.35 (5)	0.046	369.4 (5)	0.03 (2)	369.4	0.05 (1)	369.35 (5)	0.048 (10)	0.05	0.33
371.0 (1)	0.06 (2)	370.85 (5)	0.054	370.9	0.01 (1)	370.9	0.05 (1)	370.93 (8)	0.031 (21)	1.1	5.8
		374.8 (2)	0.012	375.1		374.5 (1)	0.01	374.8 (2)**	0.012**		
376.0 (3)	0.04 (1)	376.30 (10)	0.005					376.27 (10)	0.04 (1)*	0.9	
379.4 (1)	0.08 (2)							379.4 (1)*	0.08 (2)*		

1993Ab01(E γ)	1993Ab01(I γ)	1990Br23(E γ)	1990Br23(I γ)	1972He18(E γ)	1972He18(I γ)	1969Br27(E γ)	1969Br27(I γ)	Adopted E γ ^a	Adopted I γ ^b	χ^2/ν (E γ)	χ^2/ν (I γ)
381.9 (1)	0.05 (1)	382.4 (6)	0.046	382.4 (5)	0.05 (2)	382.5 (1)	0.05	382.2 (3)	0.050 (10)	6.1	
383.51 (4)	0.37 (5)	383.50 (10)	0.385	383.6	0.01 (1)	383.6 (2)	0.38 (8)	383.51 (4)	0.19 (18)	0.11	
				392.4 (5)	0.08 (2)			392.4 (5)&	0.078 (17)&		
398.2 (2)	0.011 (3)	399.0 (4)	0.015			399.0 (15)	0.02	398.6 (3)	0.011 (3)**	1.1	
401.9 (1)	0.10 (3)	402.50 (10)	0.092	402.5	0.02 (2)	402.6 (3)	0.09 (3)	402.2 (3)	0.06 (4)	9.8	3.4
415.1 (1)	0.016 (3)	415.1 (2)	0.014	415.2		415.2 (3)	0.01	415.11 (10)	0.011 (5)	0.05	3
432.4 (5)	0.030 (4)	432.30 (10)	0.038	432.5 (5)	0.03 (2)	432.4 (2)	0.04 (1)	432.33 (10)	0.032 (4)	0.12	0.45
						442.5 (10)	0.00046	442.5 (10)#	0.00046#		
						445	0.0039 (39)	445#	0.004 (4)#		
						448.0 (6)	0.00115	448.0 (6)#	0.0011#		
452.9 (3)	0.002 (5)					452.7 (6)	0.00077	452.9 (3)	0.002 (5)*	0.09	
						457.5 (1)	0.00054	457.5 (1)#	0.00054#		
						462 (1)	0.00038	462 (1)#	0.00038#		
466.8 (2)	0.004 (2)					466.5 (10)	0.00038	466.8 (2)	0.004 (2)*	0.09	
469.0 (2)	0.007 (2)							469.0 (2)*	0.007 (2)*		
						480 (1)	0.0023 (7)	480 (1)#	0.0023 (7)#		
						482 (1)	0.0011 (3)	482 (1)#	0.0011 (3)#		
						493.1 (2)	0.0042 (6)	493.1 (2)#	0.0042 (6)#		
507.5 (1)	0.007 (2)					507.4 (3)	0.0031 (6)	507.5 (1)	0.0051 (20)	0.1	1.9
516.7 (3)	0.003 (1)					516.4 (3)	0.0013 (3)	516.6 (3)	0.0022 (8)	0.5	1.3
521.8 (3)	0.003 (1)							521.8 (3)*	0.003 (1)*		
524.7 (4)	0.0018 (4)					524.3 (4)	0.00115 (23)	524.5 (4)	0.0015 (3)	0.5	1.3
534.5 (4)	0.001					535.0 (12)	0.00077 (23)	534.6 (4)	0.00077 (23)#	0.16	
536.9 (1)	0.013 (2)					537.0 (3)	0.085 (12)	536.9 (1)	0.0085 (13)#	0.1	
540.2 (3)	0.002 (1)							540.2 (3)*	0.002 (1)*		
						552.4 (5)	0.0018 (3)	552.4 (5)#	0.0018 (4)#		
556.0 (2)	0.004 (1)					556.5 (5)	0.0017 (3)	556.1 (2)	0.0029 (12)	0.86	2.7
565.4 (1)	0.011 (2)							565.4 (1)*	0.011 (2)*		
569.4 (5)	0.010 (2)					569.0 (3)	0.0046 (7)	569.1 (3)	0.0046 (7)#	0.47	
576.0 (2)	0.004 (1)					575.7 (7)	0.0010 (2)	576.0 (2)	0.0025 (15)	0.31	4.5

Comments on evaluation

²²⁷Th

1993Ab01(E _γ)	1993Ab01(I _γ)	1990Br23(E _γ)	1990Br23(I _γ)	1972He18(E _γ)	1972He18(I _γ)	1969Br27(E _γ)	1969Br27(I _γ)	Adopted E _γ ^a	Adopted I _γ ^b	χ ² /ν(E _γ)	χ ² /ν(I _γ)
579.0 (2)	0.006 (2)					578.5 (7)	0.0010 (2)	579.0 (2)	0.0035 (25)	0.47	3.1
585.8 (1)	0.007 (2)							585.8 (1)*	0.007 (2)*		
						589.0 (6)	0.00046 (12)	589.0 (6)#	0.00046 (12)#		
						596 (1)	0.00008	596 (1)#	0.00008#		
598.9 (2)	0.005 (1)							598.9 (2)*	0.005 (1)*		
607.8 (3)	0.002 (1)					607.5 (4)	0.0014 (3)	607.7 (3)	0.0014 (3)	0.36	0.33
						621.4 (5)	0.00046 (12)	621.4 (5)#	0.00046 (12)#		
623.8 (5)	0.002 (1)					623.8 (5)	0.0012 (3)	623.8 (5)	0.0013 (3)		0.59
						632.3 (7)	0.00108 (22)	632.3 (7)#	0.0011 (2)#		
						641.0 (5)	0.00015 (5)	641.0 (5)#	0.00015 (5)#		
644.3 (3)	0.0010 (3)					644.2 (5)	0.00038 (12)	644.3 (3)	0.0007 (3)	0.03	2.1
						648.5 (5)	0.00046 (14)	648.5 (5)#	0.00015 (5)#		
662.5 (3)	0.003 (1)					663.1 (5)	0.00046 (14)	662.8 (4)	0.00046 (14)#	0.72	
						692.0 (7)	0.00031 (9)	692.0 (7)#	0.00031 (9)#		
						704.3 (5)	0.00062 (12)	704.3 (5)#	0.00062 (12)#		
						707.2 (7)	0.00031 (9)	707.2 (7)#	0.00031 (9)#		
						718.5 (10)	0.00023 (9)	718.5 (10)#	0.00023 (9)#		
						722.1 (6)	0.0029 (9)	722.1 (6)#	0.0029 (9)#		
723.5 (1)	0.008 (2)					723.6 (6)	0.0029 (9)	723.5 (1)	0.0021 (8)#	0.03	
						734.4 (5)	0.0008 (3)	734.4 (5)#	0.0008 (3)#		
735.4 (2)	0.002 (1)					735.5 (5)	0.0012 (4)	735.4 (2)	0.0013 (4)	0.03	0.55
						738.4 (10)	0.00054 (13)	738.4 (10)#	0.00054 (13)#		
						746.4 (7)	0.0008 (3)	746.4 (7)#	0.0008 (3)#		
749.2 (1)	0.004 (1)					748.5 (5)	0.0023 (5)	748.8 (4)	0.0032 (9)	0.98	1.4
754.1 (2)	0.003 (1)					754.0 (6)	0.00077 (19)	754.1 (2)	0.0019 (11)	0.02	2.5
						756.9 (2)	0.0015 (4)	756.9 (2)#	0.0015 (4)#		
757.7 (1)	0.010 (2)					756.9 (2)	0.0062 (15)	757.3 (4)	0.0081 (19)	8	1.8
763.1 (2)	0.003 (1)					762.2 (5)	0.0020 (4)	762.6 (5)	0.0021 (4)	1.6	0.86
						766.3 (5)	0.0023 (5)	766.3 (5)#	0.0023 (5)#		
773.5 (4)	0.0013 (3)					773.0 (8)	0.0010 (4)	773.4 (4)	0.0012 (3)	0.31	0.36
776.3 (1)	0.013 (2)					775.3 (2)	0.0115 (12)	775.8 (5)	0.012 (1)	13	0.2

1993Ab01(E γ)	1993Ab01(I γ)	1990Br23(E γ)	1990Br23(I γ)	1972He18(E γ)	1972He18(I γ)	1969Br27(E γ)	1969Br27(I γ)	Adopted E γ ^a	Adopted I γ ^b	$\chi^2/\nu(\text{E}\gamma)$	$\chi^2/\nu(\text{I}\gamma)$
781.5 (2)	0.0025 (8)					780.5 (3)	0.0025 (5)	781.0 (5)	0.0025 (5)	5.6	
						784.2 (5)	0.00077 (19)	784.2 (5)#	0.00077 (19)#		
787.7 (5)	0.0011 (3)					787.4 (5)	0.00069 (17)	787.6 (5)	0.00089 (21)	0.18	0.9
						787.4 (5)	0.00031 (8)	787.4 (5)#	0.00031 (8)#		
						792.6 (6)	0.00031 (8)	792.6 (6)#	0.00031 (8)#		
						792.6 (6)	0.00023 (6)	792.6 (6)#	0.00023 (6)#		
797.7 (1)	0.008 (1)					796.8 (2)	0.0062 (6)	797.3 (5)	0.0071 (9)	10	1.6
804.2 (1)	0.009 (1)					803.5 (2)	0.0075 (8)	803.9 (4)	0.005 (4)	6.1	34
808.6 (4)	0.0006 (2)					807.5	0.00038	808.6 (4)#	0.0006 (2)#		
813.0 (1)	0.024 (5)					812.2 (2)	0.0208 (21)	812.6 (4)	0.013 (2)	8	9.6
818.1 (2)	0.0019 (5)					818.0 (10)	0.00077 (23)	818.1 (2)	0.0013 (6)	0.01	2.6
						818.0 (10)	0.00023 (9)	818.0 (10)#	0.00023 (9)#		
823.8 (1)	0.024 (5)					823.1 (2)	0.0192 (19)	823.4 (4)	0.020 (2)	6.1	0.86
826.9 (5)	0.0012 (4)					826.0 (10)	0.0015 (5)	826.7 (5)	0.0013 (4)	0.65	0.22
						828.5 (5)	0.0015 (4)	828.5 (5)#	0.0015 (4)#		
829.0 (2)	0.0060 (2)					828.5 (5)	0.00008 (3)	828.9 (2)	0.0060 (2)*	0.86	
838.2 (2)	0.005 (1)					837.3 (3)	0.0031 (3)	837.8 (5)	0.0041 (9)	4.5	1.8
842.8 (1)	0.007 (1)					842.2 (3)	0.0046 (5)	842.5 (3)	0.0069 (10)	2	0.15
						846.7 (5)	0.00115 (23)	846.7 (5)#	0.00115 (23)#		
847.8 (3)	0.003 (1)					848.7 (5)	0.00046 (14)	848.3 (6)	0.0021 (9)	0.4	1.6
						854.3 (5)	0.00054 (11)	854.3 (5)#	0.00054 (11)#		
						857.3 (7)	0.00046 (14)	857.3 (7)#	0.00046 (14)#		
858.9 (2)	0.003 (1)					858.8 (3)	0.0019 (3)	858.9 (2)	0.0020 (3)	0.08	1.1
						863 (1)	0.00015 (6)	863 (1)#	0.00015 (6)#		
867.1 (5)	0.004 (1)					867.5 (5)	0.00054 (11)	867.3 (5)	0.0023 (17)	0.32	6
876.5 (5)	0.0023 (6)					876.2 (4)	0.0012 (4)	876.3 (4)	0.0018 (6)	0.22	1.7
878.2 (4)	0.0015 (5)					878.2 (4)	0.0009 (3)	878.2 (4)	0.0011 (3)		1.1
						891 (1)	0.00015 (5)	891 (1)#	0.00015 (5)#		
						893 (1)	0.00010 (3)	893 (1)#	0.00010 (3)#		
						896.1 (5)	0.00085 (21)	896.1 (5)#	0.00085 (21)#		
908.9 (1)	0.021 (2)					908.2 (2)	0.0161 (24)	908.6 (4)	0.0185 (25)	6.1	3.1

1993Ab01(E γ)	1993Ab01(I γ)	1990Br23(E γ)	1990Br23(I γ)	1972He18(E γ)	1972He18(I γ)	1969Br27(E γ)	1969Br27(I γ)	Adopted E γ ^a	Adopted I γ ^b	χ^2/ν (E γ)	χ^2/ν (I γ)
						910 (1)	0.00012 (5)	910 (1)#	0.00012 (5)#		
						920.0 (5)	0.00009 (2)	920.0 (5)#	0.00009 (2)#		
						927 (1)	0.00005 (2)	927 (1)#	0.00005 (2)#		
						938.0 (8)	0.00008 (3)	938.0 (8)#	0.00008 (3)#		
						941.6 (3)	0.00055 (8)	941.6 (3)#	0.00055 (8)#		
						958.7 (3)	0.00048 (10)	958.7 (3)#	0.00048 (10)#		
970.3 (2)	0.0020 (5)					970.0 (4)	0.00023 (5)	970.2 (2)	0.0011 (9)		6.3
						971.7 (10)	0.00008 (4)	971.7 (10)#	0.00008 (4)#		
						988 (1)					
						990.0 (7)	0.00027 (7)	990.0 (7)#	0.00027 (7)#		
						995 (1)	0.00005	995 (1)#	0.00005 (3)#		
						999.8 (5)	0.00023 (6)	999.8 (5)#	0.00023 (6)#		
						1015.2 (7)	0.00012 (3)	1015.2 (7)#	0.00012 (3)#		
						1020 (1)	0.00015 (5)	1020 (1)#	0.00015 (5)#		
						1025 (1)	0.00012 (3)	1025 (1)#	0.00012 (3)#		
* From 93Ab01											
** From 90Br23											
& From 72He18											
# From 69Br27											
a Weighted average (LWM) of values from 93Ab01, 90Br23, 72He18, 69Br27, unless otherwise specified.											
b Weighted average (LWM) of values from 93Ab01, 72He18, 69Br27, unless otherwise specified.											
c Double											

²²⁸Ra – Comments on Evaluation of Decay Data by A. Luca

This evaluation was completed in June 2009. The literature available by December 31st, 2008 was included.

1. Evaluation Procedures

The Limitation of Relative Statistical Weight (LWM) method was applied for averaging numbers throughout this evaluation; this method was implemented by using the computer code LWEIGHT, ver. 4 (designed for Excel, MS Office). The uncertainty assigned to an average value in this evaluation is never lower than the lowest uncertainty of any of the experimental input values.

2. Decay Scheme

²²⁸Ra decays 100 % by beta minus particle emissions, populating the ²²⁸Ac excited states. The decay scheme was studied by a few authors (1961To10, 1972HeYY, 1995So11). The most recent evaluation of the ²²⁸Ra nuclear structure and decay data, published in Nuclear Data Sheets, was made by A. Artna-Cohen (1997). In the present evaluation, the spin, parity and energy of the ²²⁸Ac excited levels, and the multipolarities of the γ -ray transitions, have been adopted from the above mentioned A=228 ENSDF mass-chain evaluation (1997Ar08).

3. Nuclear Data

The adopted beta decay energy value $Q(\beta)=45.8$ (7) keV, is from 2003Au03. This value is in very good agreement with the effective $Q(\beta)$ value of 46 keV (with an uncertainty of 6 keV), calculated from the decay scheme data, by using the SAISINUC software, version 2008 April.

3.1. Half-life

In the literature, only two measured ²²⁸Ra half-life ($T_{1/2}$) values are reported; both measurements are very old (the most recent is from 1962), so new half-life measurements are needed to improve the quality of the evaluation. The half-life values and their uncertainties are presented in Table 1; the value recommended by Curie et al. (1931), with an estimated uncertainty added by the evaluator, was also included. A critical review of the half-life computation (weighted average of 7 values) from reference 1962Ma58, was done by using the computer code LWEIGHT, ver.4. The set of data is consistent and the recommended value, 5.75 years, with an uncertainty of 0.04 years, is the weighted average (LWM, $\chi^2_{\nu}=4.6$) of the three input values. The references are expressed as NSR (Nuclear Science References) type keynumbers.

Table 1: ²²⁸Ra Half-life values

$T_{1/2}$ (years)	Uncertainty of $T_{1/2}$ (years)	Reference
6.7	1	1931Cu01
5.7	0.2	1960Du11
5.75	0.04	1962Ma58

3.2. Beta transitions and emissions

In the literature, the most complete reference reporting measurements of energy and emission intensities for ²²⁸Ra beta minus transitions is 1995So11.

For this evaluation, the beta transitions energies were calculated from Q(β-) and the energies of the decay scheme levels. The intensities of the beta branches were deduced from γ-ray transition intensity balance at each level (using also the corresponding total ICC values, computed as described below, in section 3.3), with the exception of the lowest energy branch (12.7 keV maximum energy) which was adopted from the measurements reported by 1995So11; also, the intensity ratio of the two highest energy beta branches, 39.1 keV and 39.5 keV, was adopted from the same reference (the uncertainty of this ratio, not mentioned in 1995So11, was neglected in present computations).

The intensity balance equations – including the normalization for the ground state of ²²⁸Ac, together with the experimental data mentioned above and below in section 3.3, were assembled in a linear system of nine equations with nine unknown parameters to be determined. The system of equations is the following (the numbers between round parentheses associated to I_β, I_γ and α_T correspond to the energy values of the beta minus/gamma-ray emissions/transitions, expressed in keV):

$$\begin{aligned}
 I_{\beta}(12.7) &= [1 + \alpha_T(12.88)] \cdot I_{\gamma}(12.88) + [1 + \alpha_T(26.40)] \cdot I_{\gamma}(26.40) \\
 I_{\beta}(25.6) + [1 + \alpha_T(12.88)] \cdot I_{\gamma}(12.88) &= [1 + \alpha_T(13.52)] \cdot I_{\gamma}(13.52) \\
 I_{\beta}(39.1) + [1 + \alpha_T(13.52)] \cdot I_{\gamma}(13.52) + [1 + \alpha_T(26.40)] \cdot I_{\gamma}(26.40) &= [1 + \alpha_T(6.67)] \cdot I_{\gamma}(6.67) \\
 I_{\beta}(39.5) &= [1 + \alpha_T(6.28)] \cdot I_{\gamma}(6.28) \\
 [1 + \alpha_T(6.28)] \cdot I_{\gamma}(6.28) + [1 + \alpha_T(6.67)] \cdot I_{\gamma}(6.67) &= 100 \% \\
 I_{\beta}(12.7) &= 30 (10) \% \\
 I_{\beta}(39.1) / I_{\beta}(39.5) &= 4 \\
 I_{\gamma}(13.52) &= 1.6 (1) \% \\
 I_{\gamma}(12.88) &= 0.30 (6) \%
 \end{aligned}$$

Using the gamma-ray emission probabilities for the 13.52 keV and 12.88 keV photons measured by 1995So11, a new intensity value of the 25.6 keV beta branch was computed by the evaluator (see Table 2); this was done because the 20 % beta intensity gives a negative gamma-ray emission probability for the 12.88 keV photons, according to the intensity balance of the 20.19 keV ²²⁸Ac excited level. The normalization condition of the beta emissions (the sum of the all the beta transitions intensities must be 100 %) was checked. The adopted energy and intensity values of the beta transitions, as well as their Log ft values are shown in Table 2.

Table 2: ²²⁸Ra β⁻ Energies and Emission Probabilities

E _{β⁻} (keV)	Uncertainty E _{β⁻} (keV)	Emission probability (%)	Emission probability (%), from 1995So11	Log ft
12.7	0.7	30 (10)	30 (10)	5.11
25.6	0.7	8.7 (9)	20 (6)	6.2
39.1	0.7	49 (10)	40 (10)	6.45
39.5	0.7	12 (10)	10	7.07

3.3. γ- transitions: γ rays and internal conversion electrons

The single paper reporting measurements of the γ-ray energies and some emission intensities following the ²²⁸Ra decay (only for 13.52 keV and 12.88 keV) is 1995So11. Using the measured 13.52 keV gamma-ray emission probability of 1.6 % (with a 0.1 % estimated uncertainty, added by the evaluator), the 12.88 keV photons measured emission probability of 0.30 (6) % and the internal conversion coefficients, the corresponding absolute gamma-ray emission probabilities and their uncertainties were computed for all the γ rays, by solving the linear system of equations from section 3.2; the obtained data are given below in Table 3. The internal conversion coefficients were computed with the program BrIcc, version 2.2b/20-Jan-2009, using the “Frozen Orbitals” approximation.

Other possible gamma-ray transitions observed only by Sood et al. (1995), but not placed in the level scheme, are: 15.15 keV, 15.5 keV, 16.2 keV and 30.6 keV.

Table 3: ²²⁸Ra γ -ray Energies and Absolute Emission Probabilities

E_γ (keV)	Uncertainty E_γ (keV)	Absolute Emission Probability (%)	Uncertainty of absolute emission probability (%)	Total ICC (α_T) and uncertainty
6.28	0.03	$1.8 \cdot 10^{-6}$	$1.5 \cdot 10^{-6}$	$6.68 (19) \cdot 10^6$
6.67	0.02	$5.7 \cdot 10^{-5}$	$0.9 \cdot 10^{-5}$	$1.560 (40) \cdot 10^6$
12.88	0.11	0.30	0.06	6.67 (18)
13.52	0.04	1.6	0.1	5.86 (10)
26.40	0.11	0.14	0.05	201 (4)

4. Atomic data

The mean L-shell fluorescence yield (ω_L) and the relative probabilities of vacancies in the L-shell were given by the computer program EMISSION v.3.10, 28-Jan-2003.

4.1. Auger electrons and X-rays

Because the decay energy, Q, is very low, there are no electron emissions from the ²²⁸Ac K-shell (Auger electrons or internal conversion electrons).

The emission intensity of the L Auger electrons (energy from 0.1 keV to 19.69 keV), was computed using the EMISSION computer program: 12 (5) %.

The absolute emission intensity values of the different groups of L X-rays (L_L , L_α , L_η , L_β and L_γ) were determined using the EMISSION program; the total L X-rays emission intensity is 9.6 (19) %, for an energy range between 10.87 keV and 18.92 keV. Neither measurements of X-ray energies nor of emission intensities were found in the literature, in order to compare them with the results of this evaluation.

5. Main production mode

The main production mode of ²²⁸Ra is by alpha-particle decay of the ²³²Th nuclei (²²⁸Ra is the daughter of ²³²Th), present in important quantities in many natural ores.

6. References

- 1931Cu01 M. Curie, A. Debierne, A.S. Eve, H. Geiger, O. Hahn, S.C. Lind, St. Meyer, E. Rutherford and E. Schweidler, "The Radioactive constants as of 1930", Rev. Mod. Phys. 3, 427-445 (1931).
- 1960Du11 R.A. Dudley, "MsTh half-period", MIT Report, NYO-9504, 85-86 (1960).
- 1961To10 J. Tousset and A. Moussa, "La désintégration du mésothorium 1. Schémas proposés", J. Phys. Radium, tome 22, 683-685 (1961).
- 1962Ma58 C.W. Mays, D.R. Atherton, R.D. Lloyd, H.F. Lucas, B.J. Stover, F.W. Bruenger, "The half-period of Ra²²⁸ (Mesothorium)", Utah Univ. Report, COO-225, 92-105 (1962).
- 1972HeYY M. Herment and A. Gizon, Annual report ISN Grenoble, 115 (1972).
- 1995So11 P.C. Sood, A. Gizon, D.G. Burke, B. Singh, C.F. Liang, R.K. Sheline, M.J. Martin, R.W. Hoff, " β decay of ²²⁸Ra and possible level structures in ²²⁸Ac", Phys. Rev. C, vol. 52, no. 1, 88-92 (1995).
- 1997Ar08 A. Artna-Cohen, "Nuclear Data Sheets for A = 228", Nucl. Data Sheets 80, 723-785 (1997).
- 2003Au03 G. Audi, A.H. Wapstra and C. Thibault, "The AME2003 atomic mass Evaluation (II). Tables, graphs, and references", Nucl. Phys. A 729, 337-676 (2003).

²²⁸Ac - Comments on Evaluation Decay Data By Andy Pearce

This evaluation was completed in September 2009 drawing in part from the mass-chain evaluation of Artna-Cohen^[1]. The literature available up until January 2009 was included. There is some evidence the decay scheme is not complete based upon the calculated beta emissions.

1 Decay Scheme

²²⁸Ac decays almost entirely by beta- decay to excited states in ²²⁸Th. The decay scheme (energies, half lives, spins and parities of levels in ²²⁸Th) are based upon the adopted levels and gammas from Artna-Cohen^[1], which in turn are largely derived from the work of Dalmaso et al^[2]. The assignments of gammas to levels are also largely derived from Dalmaso et al.

Baltzer et al^[3] place gammas at 56.8 keV and 137.4 keV originating from a 1588 keV 4- level in ²²⁸Pa decay. Gamma emissions of similar energies have been observed in ²²⁸Ac decay but not placed, and it is assumed these are the same transitions. Inserting this level in the ²²⁸Ac decay allows for alternative placement of the 356 keV gamma, for which the predicted multipolarity (E2) did not match the measured conversion coefficient^[4] ($\alpha=0.3-4$). Placement from the 1945 keV level to the 1588 keV level indicates a multipolarity of E1+M2; the mixing ratio has been tentatively estimated at $\delta=0.5$ giving $\alpha_T=0.35$.

Decay via alpha emission to ²²⁴Fr has been reported^[5] with a probability of $(5.5 \pm 2.2) \times 10^{-6}$ per 100 decays, but this is unconfirmed.

2 Nuclear Data

For the purposes of this evaluation it is assumed ²²⁸Ac decays entirely to ²²⁸Th and any alpha branching present is negligibly small. The Q-value of beta decay is taken from Audi, Wapstra & Thibault^[6] and is 2123.8 (27) keV. The effective Q-value calculated from the individual decay rates and energies calculated with the RADLST^[7] program is 2010 (100) keV; this is low compared with the value from Audi, Wapstra & Thibault and serves to confirm that the decay scheme is incomplete.

There have been only two measurements of the half life reported in the open literature, Hahn and Erbacher^[8] at 6.13 hours (quoted in Curie^[9], however the corresponding publication could not be identified) and Skanemarg & Skalberg^[10] at 6.15 (3) hours. No uncertainty is given for the value of Hahn & Erbacher, but it serves to give confidence in the recommended value of 6.15 (3) hours taken from Skanemarg and Skalberg. From an evaluators' point of view more data are required to provide a definitive half life, however in practice this radionuclide will generally be encountered in secular equilibrium with either the ²²⁸Ra or ²³²Th decay parents, and the exact value of the half life will likely be of no great concern to the user.

3 Atomic Data

All values of atomic data (ω_K , ω_L , η_{KL} , relative probabilities of X-ray and Auger emissions) were derived from Schönfeld and Janßen^[11].

4 Gamma-ray Transitions and Internal Conversion Coefficients

Gamma-ray transition energies are calculated from the differences in level energies from Artna-Cohen^[1].

Internal conversion coefficients have been determined using the BrIcc code^[12], with the gamma-ray multiplicities and mixing ratios from the evaluation of Artna-Cohen^[1]. For many emissions the multipolarity is undetermined, and no conversion coefficients could be calculated. Where a multipolarity is available but no mixing ratio given, a default value of 1 is assumed and the uncertainty of the derived conversion coefficients is increased accordingly.

Internal conversion coefficients have been measured by Herment and Vieu^[4] and Mahajan and Bidarkundi^[13] and these are compared in table 3 with the recommended values calculated with BrIcc. The agreement of the total conversion coefficients for the K and L shells is generally good. The 204 keV transition has been reassigned to M2 based upon the measured K conversion coefficient in 1982Ma52; the authors assign 96 % M2 + 4% E1 however the theoretical coefficients calculated using BrIcc for pure M2 are very close to the measured values.

Devare and Devare^[24] have measured accurately the L_{II}/L_I conversion ratios for the 184 keV transition from the 1153 keV level and have assigned the multipolarity as predominantly E0 + M1 with a mixing ratio of $\delta \approx 0.3$. It is not possible with BrIcc to calculate conversion coefficients of mixed type transitions incorporating an E0 element, therefore the K-shell conversion coefficient has been estimated at 80 (30) from the measured data as the median of values measured by Herment and Vieu and Mahajan and Bidarkundi and the total coefficient of 100 (40) calculated from the calculated K/total ratio derived from BrIcc with the multipolarity and mixing ratio as above.

5 Electron Emissions

5.1 Beta-particle Emissions

There are no published measurements of the beta emissions in the open literature. Beta-particle transition energies have therefore been determined from the Q-value and level energies, and the emission probabilities from the balance of the decay scheme using the program 'GTOL'^[14].

A normalisation factor for the gamma emission probabilities of 0.0454 (11) has been used to determine absolute gamma emission intensities; however, applying this same value when deriving level feedings implies a ~7 % feeding direct to the ground state. Such a transition would be a 2nd forbidden unique decay and such a high branching seems unreasonable. By assuming negligible feeding to the ground state leads a normalisation factor of 0.0475 (11) is calculated, and this value is used to calculate beta particle emission intensities with GTOL. There is therefore an unexplained ~7 % discrepancy between the beta and gamma emissions in this decay; Artna-Cohen^[1] suggests there are missing gammas in the decay scheme. Given the available data it seems such gammas would need to decay direct to the ground state and be of such character as to be difficult to detect, for example low energy or E0 transitions. Further measurements of the gamma data, particularly at low energy, would be of benefit, as would coincidence studies to validate the placement of gammas in the level scheme.

The Q-value coupled with the presence of a level in the decay scheme at 2123.1 (3) keV implies a beta emission with an end-point energy of 0.7 keV; it seems unlikely such a low energy emission could have a significant emission probability. This may indicate a deficiency in either the placement of gammas in the decay scheme or in the Q-value.

5.2 Auger & Conversion Electron Emissions

Auger and conversion electron emissions per 100 decays were calculated from the gamma-ray data and conversion coefficients according to the method of Schönfeld and Janßen^[11] using version 3.10 of the code EMISSION.

6 Photon Emissions

6.1 X-ray Emissions

The X-ray intensities per 100 decays have been calculated from the gamma-ray data and conversion coefficients using version 3.10 of the code EMISSION. No measurements of the X-ray emissions have been published so it is not possible to make a comparison of calculated and measured data. A comparison with values in the NUDAT database is given in table 4.

6.2 Gamma-ray Emissions

The gamma-ray emission energies have been taken from Helmer^[15] where possible, in which precise measurements were made by measuring energy differences against accepted calibration standards. Only the directly measured values have been taken, as the decay scheme used to derive further values was incomplete. These values have been adjusted to reflect the updated calibration standards given in Helmer and van der Leun^[16]. Where gamma-ray lines are not present in Helmer^[15], weighted means of the values in Herment and Vieu^[4], Taylor^[17], Kurcewicz et al^[18], Borner et al^[19], Dalmaso et al^[2] and Baltzer et al^[3] were taken. The uncertainties of Borner et al were expanded based upon the detector resolution stated in the publication. The values in these publications were first rescaled by a least-squares fit to be compatible with Helmer^[15]. In most cases the energy shift incurred by doing so was very small.

Gamma emission probabilities were determined by a weighted mean of values in Arnoux and Gizon^[20], Herment and Vieu^[4], Mahajan and Bidarkundi^[13], Sadasivan and Raghunath^[21], Schötzig and Debertin^[22], Dalmaso et al^[2], Lin and Harbottle^[23] and Baltzer et al^[3]. Uncertainties were expanded to match the minimum input uncertainty where appropriate. Values were first renormalised to 100 for the 463 keV emission. Baltzer et al relates to ²²⁸Pa decay, however additional information could be obtained from this publication for relative gamma emission probabilities originating from the 1431 keV level. Measured and evaluated relative emission probabilities are compared in appendices II and III. While there are several publications covering some of the emissions the intensities of many transitions have been derived solely from Dalmaso et al; where alternative data exists the agreement is often poor.

Several of the gamma emission probabilities measured by Mahajan and Bidarkundi have been rejected on technical grounds. The radionuclides used for calibration stated by the authors were ¹⁵²Eu, ¹⁶⁰Tb and ¹⁹²Ir. The lowest energy of any gamma line which could reliably be used for efficiency calibration belongs to ¹⁵²Eu at 122 keV; therefore, gamma emission intensities reported below this energy were rejected. Furthermore, in the energy region 321 keV to 338 keV there seems to be a consistent high bias to the data (see appendix I); two out of three measurements were rejected by Chauvenet's criterion. The remaining measurement passed Chauvenet's criterion but appears high and was rejected due to the obvious trend.

It is not clear that the data in Arnoux and Gizon^[20] and Herment and Vieu^[4] are independent; in many cases, the data are numerically identical, and appear to correspond to the work of the same research group. In cases where the same emission has been reported by both authors, only the data from the latter publication have been used in the analysis.

Absolute gamma emissions were measured by Schötzig and Debertin^[22] and Lin and Harbottle^[23]. The intensities of the 463 keV emissions were used to derive normalisation factors and these values are 0.0450 (12) and 0.0441 (11) respectively. The weighted mean of these two values at 0.0445 (11) was used to convert relative intensities into absolute intensities. However, this value is not consistent with expected beta decay characteristics (see section 5.1), suggesting deficiencies in either the adopted decay scheme or the measured data.

There are twenty gammas which cannot be placed in the level scheme. The total intensity is less than 0.25 % (accounting for some internal conversion). These are listed in table 2. The intensity of these gamma emissions is insufficient to account for the discrepancies observed in the decay scheme.

The 18.4 keV gamma has been observed but the probability not directly measured in ²²⁸Ac decay; a nominal value of 0.14 (3) for the total transition probability has been derived based upon coincidence measurements on ²²⁸Pa^[16]. This implies a gamma emission probability of 0.019 (4).

Several doublets have been reported in ²²⁸Ac decay. The measured gamma emission intensities have been divided between the components where possible by comparing with ²²⁸Pa decay^[3]:

168.53 (12) keV

The 1344 keV level has not being reported in ²²⁸Pa decay, therefore the intensity of the 168 keV emission from the 1928 keV level was derived from the ratio of the 168 keV emission to the 1741 keV and 1870 keV emissions in ²²⁸Pa:

Intensity 168 keV (²²⁸Ac, 1928 keV) = intensity 168 keV ²²⁸Pa / intensity 1741 keV ²²⁸Pa × intensity 1741 keV ²²⁸Ac

and:

Intensity 168 keV (²²⁸Ac, 1928 keV) = intensity 168 keV ²²⁸Pa / intensity 1870 keV ²²⁸Pa × intensity 1870 keV ²²⁸Ac

The calculated values are 0.0715 (18) and 0.0407 (43) respectively; these are not consistent so a median of the two values is taken, with an uncertainty large enough to cover the difference.

The assigned relative intensities are 0.056 (15) from the 1928 keV level and 0.25 (6) from the 1344 keV level. The absolute intensities are therefore 0.0025 (7) and 0.0111 (27) respectively.

278.80 (15) keV

The total relative intensity is 5.28 (28) determined by LRSW weighted mean of the measured data^[2,13,20]. The intensity has been split between transitions from the 1153 keV and 1431 keV levels by comparing the emission intensities in ²²⁸Ac decay and ²²⁸Pa decay. The relative intensities are therefore 4.6 (5) and 0.69 (7) for transitions from the 1153 keV and 1431 keV levels respectively.

649.02 (12) keV

The measured relative intensity is 0.94 (10) from Dalmaso et al^[2]. The intensity has been split between transitions from the 1168 keV and 1617 keV levels by comparing the emission intensities in ²²⁸Ac decay and ²²⁸Pa decay. The relative intensities are therefore 0.75 (8) and 0.189 (20) for transitions from the 1168 keV and 1617 keV levels respectively.

666.451 (46) keV

The measured relative intensity is 2.4 (2) from Dalmaso et al^[2]. The intensity has been split between transitions from the 1645 keV and 1892 keV levels by comparing the emission intensities in ²²⁸Ac decay and ²²⁸Pa decay. The relative intensities are therefore 2.27 (23) and 0.128 (13) for transitions from the 1645 keV and 1892 keV levels respectively.

688.117 (42) keV

Dalmaso et al^[2] suggests this emission originates from the 874 keV level; Baltzer et al^[4] suggest dual placement from the 874 keV and 1016 keV levels. The measured relative intensity is 1.58 (14) from Dalmaso et al. The intensity has been split by comparing the emission intensities in ²²⁸Ac decay and ²²⁸Pa decay. The relative intensities are therefore 0.161 (16) and 1.42 (14) for transitions from the 1645 keV and 1892 keV levels respectively.

791.43 (8) keV

The measured relative intensity is 0.54 (17) from Dalmaso et al^[2]. The intensity has been split between transitions from the 1760 keV and 1944 keV levels by comparing the emission intensities in ²²⁸Ac decay

and ^{228}Pa decay. The relative intensities are therefore 0.23 (7) and 0.31 (10) for transitions from the 1760 keV and 1944 keV levels respectively.

853.96 (8) keV

Artna-Cohen^[1] indicates a doublet between an unplaced transition and a transition originating from the 1944 keV level. The energy measured by Dalmaso et al^[2] is 853.19 (10) keV however a transition of this energy does not readily fit in to the level scheme as it stands. In the absence of confirmatory measurements it is assumed this transition is the same as that observed at 853.96 keV by Baltzer et al^[4] in ^{228}Pa decay. The entire measured intensity of 0.0124 (20) is assigned to the 1944 keV level, and the energy is determined from ^{228}Pa decay.

921.87 (12) keV

Artna-Cohen indicates a doublet between the 979 keV and 1925 keV levels. However, the transition from the 1925 keV level was reported separately by Baltzer et al^[4] in ^{228}Pa decay at an energy of 922.5 keV. Based upon the reported energy, it is assumed the transition measured in ^{228}Ac decay is predominantly from the 979 keV level. This emission has therefore been assigned in its entirety to decay from the 979 keV level.

930.99 (6) keV

The total measured intensity is 0.0129 (20) from 1987Da28. The intensity has been split between transitions from the 1450 keV and 1899 keV levels by comparing the emission intensities in ^{228}Ac decay and ^{228}Pa decay. The absolute intensities are therefore 0.0040 (10) and 0.0025 (23) for transitions from the 1760 keV and 1944 keV levels respectively.

1016.44 (8) keV

Observed in both ^{228}Ac and ^{228}Pa decay. Dalmaso et al^[2] suggests a multiple placing, originating from both the 1016 keV and 1344 keV levels of ^{228}Th . Baltzer et al^[4] gives only a “less than” value for the 1016 keV transition. The 1344 keV level is not indicated as being fed in ^{228}Pa decay, and assuming the intensity in ^{228}Pa decay is half the “less than” value gives a branching ratio very similar to that observed in ^{228}Ac decay. The intensity has therefore been assigned in its entirety to the transition from the 1016 keV level.

1110.604 (9) keV

The total measured intensity is 0.311 (24) from a weighted mean of Arnoux and Gizon^[20] and Dalmaso et al^[2]. The intensity has been split between transitions from the 1168 keV and 1297 keV levels by comparing the emission intensities in ^{228}Ac decay and ^{228}Pa decay (1995Ba42). The relative intensities are therefore 6.4 (5) and 0.60 (5) for transitions from the 1168 keV and 1297 keV levels respectively.

Table 1. Comparison of beta transition probabilities calculated using the gamma normalisation factor derived from measurements using a normalisation factor of 0.0445 (11) [A] and from assuming zero feeding to the ground state with a normalisation factor of 0.0474 (11) [B]. Where the calculated beta transition probability is within one standard deviation of zero, a “less than” value is stated. In these cases the emission is not assumed to occur with significant probability. The values calculated assuming zero feeding to the ground state have been recommended.

Level energy /keV	Beta endpoint energy /keV	Feeding /per 100 decays		log <i>ft</i>	Transition type
		[A]	[B]		
ground state	2123.8 (27)	7 (3)	<6	-	2nd forbidden unique
57.759 (4)	2066.0 (27)	6 (4)	6 (4)	9.0 (4)	allowed
186.823 (4)	1937.0 (27)	0.6 (5)	0.6 (5)	10 (4)	allowed
328.003 (4)	1795.8 (27)	0.67 (22)	0.72 (23)	10.65 (20)	1st forbidden unique
378.179 (10)	1745.6 (27)	0.138 (19)	0.147 (21)	12.29 (16)	2nd forbidden unique
396.078 (5)	1727.7 (27)	11.6 (5)	12.4 (5)	8.40 (15)	1st forbidden unique
519.192 (6)	1604.6 (27)	<0.07	<0.07	-	1st forbidden unique
831.823 (10)	1292.0 (27)	<0.01	<0.05	-	1st forbidden unique
874.473 (18)	1249.3 (27)	0.16 (10)	0.17 (10)	9.7 (3)	allowed
938.58 (7)	1185.2 (27)	<0.01	<0.01	-	2nd forbidden unique
944.196 (13)	1179.6 (27)	0.081 (15)	0.087 (16)	9.95 (17)	allowed /1st forbidden
968.369 (20)	1155.4 (27)	0.17 (3)	0.18 (3)	9.60 (16)	1st forbidden
968.968 (5)	1154.8 (27)	29 (3)	31 (4)	7.37 (16)	allowed
979.499 (14)	1144.3 (27)	0.224 (19)	0.238 (20)	9.47 (15)	allowed
1016.406 (21)	1107.4 (27)	0.36 (6)	0.39 (6)	9.20 (16)	allowed/1st forbidden
1022.527 (6)	1101.3 (27)	2.8 (4)	3.0 (4)	8.31 (16)	allowed
1059.93 (3)	1063.9 (27)	0.093 (11)	0.099 (11)	9.74 (15)	1st forbidden
1091.017 (8)	1032.8 (27)	0.16 (7)	0.16 (7)	9.48 (24)	allowed
1122.951 (6)	1000.8 (27)	6.27 (17)	6.67 (18)	7.81 (15)	1st forbidden
1153.467 (10)	970.3 (27)	6 (3)	6 (3)	7.8 (3)	allowed
1168.375 (5)	955.4 (27)	3.18 (11)	3.39 (11)	8.04 (15)	1st forbidden
1174.508 (18)	949.3 (27)	<	<		allowed
1175.39 (5)	948.4 (27)	0.155 (18)	0.166 (19)	9.34 (15)	allowed
1226.565 (7)	897.2 (27)	0.63 (7)	0.67 (8)	8.65 (15)	1st forbidden
1297.423 (10)	826.4 (27)	1.37 (10)	1.46 (11)	8.18 (15)	1st forbidden unique
1344.078 (11)	779.7 (27)	0.208 (18)	0.208 (18)	8.94 (15)	1st forbidden
1416.11 (6)	707.7 (27)	0.060 (8)	0.060 (8)	9.34 (16)	allowed /1st forbidden
1431.979 (6)	691.8 (27)	1.6 (5)	1.6 (5)	7.88 (20)	allowed
1450.394 (10)	673.4 (27)	0.25 (8)	0.26 (9)	8.63 (21)	1st forbidden
1531.474 (6)	592.3 (27)	<3	<3	-	allowed
1539.21 (9)	584.6 (27)	<0.01	0.030 (6)	9.36 (17)	allowed
1588.335 (14)	535.5 (27)	8.2 (22)	8.8 (23)	6.77 (19)	1st forbidden
1617.78 (7)	506.0 (27)	0.067 (10)	0.071 (10)	8.78 (16)	allowed
1638.284 (9)	485.5 (27)	1.16 (6)	1.23 (6)	7.48 (15)	allowed
1643.125 (15)	480.7 (27)	0.82 (3)	0.82 (3)	7.64 (15)	1st forbidden
1645.954 (12)	477.8 (27)	4.12 (20)	4.12 (20)	6.94 (15)	allowed
1682.81 (3)	441.0 (27)	1.21 (4)	1.21 (4)	7.35 (15)	allowed
1683.82 (5)	440.0 (27)	0.20 (3)	0.20 (3)	8.13 (16)	1st forbidden
1688.394 (11)	435.4 (27)	2.50 (16)	2.50 (16)	7.02 (15)	allowed
1724.283 (6)	399.5 (27)	1.81 (8)	1.93 (8)	7.01 (15)	allowed
1735.45 (25)	388.4 (27)	0.140 (10)	0.149 (11)	8.08 (15)	allowed
1743.89 (3)	379.9 (27)	0.355 (15)	0.378 (16)	7.65 (15)	allowed
1758.24 (12)	365.6 (27)	0.056 (8)	0.06 (8)	8.39 (16)	allowed
1760.218 (24)	363.6 (27)	0.130 (11)	0.139 (12)	8.02 (15)	allowed
1795.90 (10)	327.9 (27)	0.033 (5)	0.035 (6)	8.48 (16)	allowed
1797.65 (8)	326.2 (27)	0.048 (8)	0.051 (8)	8.30 (16)	allowed
1892.996 (17)	230.8 (27)	0.102 (8)	0.109 (8)	7.50 (15)	allowed
1899.95 (4)	223.9 (27)	0.064 (7)	0.069 (8)	7.65 (15)	allowed

Level energy /keV	Beta endpoint energy /keV	Feeding /per 100 decays		log <i>ft</i>	Transition type
		[A]	[B]		
1906.64 (10)	217.2 (27)	0.023 (5)	0.025 (5)	8.05 (17)	allowed
1928.57 (6)	195.2 (27)	0.057 (7)	0.061 (7)	7.52 (16)	allowed
1937.16 (9)	186.6 (27)	0.050 (6)	0.053 (6)	7.52 (15)	allowed
1944.895 (11)	178.9 (27)	0.289 (20)	0.307 (22)	6.70 (15)	allowed
1958.72 (22)	165.1 (27)	0.0035 (8)	0.0038 (8)	8.50 (17)	allowed
1964.98 (7)	158.8 (27)	0.0124 (13)	0.0132 (14)	7.91 (15)	allowed
1987.46 (10)	136.3 (27)	0.07 (4)	0.07 (4)	7.0 (3)	allowed
2010.11 (5)	113.7 (27)	0.224 (14)	0.238 (15)	6.20 (15)	allowed
2013.6 (3)	110.2 (27)	0.0030 (9)	0.0032 (10)	8.03 (20)	allowed
2022.84 (10)	101.0 (27)	0.057 (6)	0.061 (6)	6.64 (16)	allowed/1st forbidden
2029.84 (16)	94.0 (27)	0.024 (4)	0.026 (4)	6.91 (16)	allowed
2036.99 (17)	86.8 (27)	0.0065 (11)	0.0069 (10)	7.38 (17)	allowed
2123.1 (3)	0.7 (27)	0.0044 (10)	0.0047 (11)	≤3.3	allowed

Table 2. Unplaced gamma emissions. The following gamma emissions have not been unambiguously placed in the level scheme. The energy lost from the decay scheme is insufficient to explain the deviation in Q_{eff} and the total probability is insufficient to explain the anomalous feeding to the ground state.

Energy /keV	Emission probability per 100 decays	Observed in	Comments
466.40 (10)	0.0299 (34)	1987Da28	-
481.5 (5)	0.024 (5)	1987Da28, 1995Ba42	Placed at 1450 keV level by 1995Ba42, however unreasonable multipolarity of M2 or E3 results (from Artna-Cohen).
634.18 (10)	0.0111 (22)	1987Da28	-
1337.33 (20)	0.0051 (16)	1987Da28	-
1378.23 (10)	0.0062 (19)	1987Da28	-
1385.39 (10)	0.0111 (22)	1987Da28	-
1434.22 (15)	0.0084 (25)	1987Da28	-
1438.01 (10)	0.0062 (17)	1987Da28	-
1480.38 (15)	0.0170 (34)	1987Da28, 1995Ba42	-
1529.01 (34)	0.059 (6)	1969Ar16, 1987Da28, 1995Ba42	Assigned by 1995Ba42 to 1925 keV level, this level not listed in ^{228}Ac decay. If present 1738 keV may be multiply placed.
1671.67 (15)	0.0043 (14)	1987Da28	-
1684.04 (20)	0.0154 (49)	1987Da28	-
1721.49 (30)	0.0059 (19)	1987Da28	-
1745.32 (20)	0.0067 (9)	1987Da28	-
1784.40 (30)	0.0062 (11)	1987Da28, 1995Ba42	-
1787.20 (20)	0.0013 (5)	1987Da28	May correspond to transition placed from 1974 keV level in ^{228}Pa decay
1916.34 (33)	0.00081 (27)	1987Da28, 1995Ba42	May correspond to transition placed from 1974 keV level in ^{228}Pa decay
1919.54 (30)	0.0022 (6)	1987Da28, 1995Ba42	-
1944.24 (20)	0.0022 (6)	1987Da28	-
2001.0 (5)	0.00108 (28)	1987Da28	-
Total Intensity	0.22 (7)		

Table 3. Comparison of experimental and calculated conversion coefficients for selected gamma transitions.

Transition energy /keV	Multipolarity	Subshell	Publication	Measured conversion coefficient	Calculated (BrIcc)
57.759 (4)	E2	L _I +L _{II}	1982Ma52	50 (4)	61.8 (9)
		L _{III}	1982Ma52	35 (3)	50.4 (7)
		L-total	1971He23	117 (6)	112.2 (16)
99.495 (8)	M1	L _I	1982Ma52	1.9 (1)	2.58 (4)
		L _{II}	1982Ma52	0.12 (1)	0.305 (5)
		L _{III}	1982Ma52	0.17 (3)	0.01646 (20)
		L-total	1971He23	2.8 (1)	2.90 (4)
129.064 (6)	E2	L _I	1982Ma52	0.17 (3)	0.1025 (15)
		L _{II}	1982Ma52	1.6 (1)	1.494 (21)
		L _{III}	1982Ma52	~0.97	0.943 (14)
		L-total	1971He23	2.45 (15)	2.54 (4)
137.941 (17)	M1	K	1971He23	4.1 (14)	6.00 (9)
		L-total	1971He23	0.8 (4)	1.146 (16)
204.038 (9)	M2	K	1982Ma52	7.5 (8)	7.26 (11)

Table 4 Comparison of X-rays calculated using EMISSION with those in the NUDAT database. Note X-ray emission probabilities for this evaluation are generally within uncertainties of, but consistently higher than the NUDAT values.

Transition	NUDAT		DDEP	
	Energy /keV	Probability /%	Energy /keV	Probability /%
K- α 1	93.35	3.1 (4)	93.351	4.0 (11)
K- α 2	89.957	1.9 (3)	89.954	2.5 (7)
L-total	~13.0	33.7 (21)	11.1-19.5	37 (4)

7 References

- [1]. A. Artna-Cohen *Nuclear Data Sheets* 80 (1997) p723.
- [2]. J. Dalmaso and H. Maria, G. Ardisson *Physical Review C* 36 (1987) p2510.
- [3]. H. Baltzer, K. Freitag, C. Gunther, P. Herzog, J. Manns, U. Muller, R. Paulsen, P. Sevenich, T. Weber and B. Will *Zeitschrift für Physik A* 352 (1995) p47.
- [4]. M. Herment, C. Vieu *Comptes Rendus - Académie de Science B* 273 (1971) 1058
- [5]. F. Lux and N. Kaubisch *Angewandte Chemie International Edition* 8 (1969) p911.
- [6]. G. Audi, A. H. Wapstra and C. Thibault *Nuclear Physics A* 729 (2003) p337 and A. H. Wapstra, G. Audi and C. Thibault *Nuclear Physics A* 729 (2003) p129.
- [7]. T. W. Burrows *The Program RADLST* Brookhaven National Laboratory Report BNL-NSC-52142 (1988)
- [8]. O. Hahn and O. Erbacher *Zeitschrift für Physik* 27 (1926) p531 (note: this reference is given in ref. [9] but the corresponding article could not be identified).
- [9]. M. Curie, A. Debierne, A. S. Eve, H. Geiger, O. Hahn, C. Lind, St. Meyer, E. Rutherford and E. Schweidler *Review of Modern Physics* 3 (1931) p427.
- [10]. G. Skarnemark and M. Skålberg *International Journal of Applied Radiation and Isotopes* 36 (1985) p439.
- [11]. E. Schönfeld and H. Janßen *Nuclear Instruments and Methods A* 369 (1996) p527.
- [12]. T. Kibédi, T. W. Burrows, M. B. Trzhaskovskoya and C. W. Nestor ANU-P/1684 (2004)
- [13]. A. S. Mahajan and M. S. Bidarkundi *Indian Journal of Pure and Applied Physics* 20 (1982) p701.
- [14]. GTOL, see for example www.nndc.bnl.gov/toolspublications/toolspublications.html
- [15]. R. G. Helmer *Nuclear Instruments and Methods* 164 (1979) p355.
- [16]. R. G. Helmer and C. van der Leun *Nuclear Instruments and Methods A* 450 (2000) p35.
- [17]. H. W. Taylor *International Journal of Applied Radiation and Isotopes* 24 (1973) p593.
- [18]. W. Kurcewicz, N. Kaffrell, N. Trautmann, A. Plochocki, J. Zylicz, M. Matul, K. Stryczniewicz *Nuclear Physics A* 289 (1977) p1.
- [19]. H. G. Borner, G. Barreau, W. F. Davidson, P. Jeuch, T. von Egidy, J. Almeida, D. H. White *Nuclear Instruments and Methods* 166 (1979) p251.
- [20]. M. Arnoux and A. Gizon *Comptes Rendus - Académie de Science B* 269 (1969) p317.
- [21]. S. Sadasivan and V. M. Raghunath *Nuclear Instruments and Methods* 196 (1982) p561.
- [22]. U. Schötzig and K. Debertin *International Journal of Applied Radiation and Isotopes* 34 (1983) p533.
- [23]. W. J. Lin and G. Harbottle *Journal of Radioanalytical and Nuclear Chemistry* 157 (1992) p367.
- [24]. S. H. Devare and H. G. Devare *Physical Review C* 9 (1974) p2297.

Appendix I. Tables of gamma emission energies

Energies have been taken from $^{1979}\text{He10}^{[15]}$ where possible, adjusted using the later reference energies of Helmer^[16]. All other energies have been taken from a weighted mean of data from $^{1971}\text{He23}^{[4]}$, $^{1973}\text{Ta25}^{[17]}$, $^{1977}\text{Ku01}^{[18]}$, $^{1979}\text{Bo30}^{[19]}$, $^{1987}\text{Da28}^{[2]}$ and $^{1995}\text{Ba42}^{[5]}$. All energies have been adjusted to be on the same energy scale as $^{1979}\text{He10}^{[15]}$ before taking means. Nominal energies given in table headings are taken from the evaluation of Artna-Cohen^[1].

Nominal Energy /keV	42.46	56.86	57.766	77.34	99.509
$^{1979}\text{He10}$	-	-	57.752 (13)	-	99.505 (12)
$^{1971}\text{He23}$	-	-	57.74 (8)	-	99.49 (11)
$^{1973}\text{Ta25}$	-	-	57.78 (6)	-	99.45 (8)
$^{1977}\text{Ku01}$	-	-	57.77 (7)	-	-
$^{1979}\text{Bo30}$	-	-	-	-	-
$^{1987}\text{Da28}$	42.457 (50)	56.96 (5)	57.761 (6)	77.338 (30)	99.497 (6)
$^{1995}\text{Ba42}$	42.46 (10)	56.852 (32)	57.752 (22)	77.35 (10)	99.461 (61)
LWEIGHT4	42.46 (5)	56.88 (5)	57.760 (13)	77.340 (30)	99.496 (6)
Adopted	42.46 (5)	56.88 (5)	57.752 (13)	77.34 (3)	99.505 (12)
Comments	wtd mean	wtd mean	$^{1979}\text{He10}$	wtd mean	$^{1979}\text{He10}$

Nominal Energy /keV	100.41	114.54	129.065	135.51	137.95
$^{1979}\text{He10}$	-	-	129.0652 (30)	-	-
$^{1971}\text{He23}$	100.39 (11)	-	129.09 (11)	135.49 (20)	-
$^{1973}\text{Ta25}$	-	-	-	-	-
$^{1977}\text{Ku01}$	-	-	129.07 (7)	-	-
$^{1979}\text{Bo30}$	-	-	129.07 (16)	-	-
$^{1987}\text{Da28}$	100.408 (30)	114.56 (7)	129.067 (7)	135.539 (50)	137.91 (5)
$^{1995}\text{Ba42}$	100.41 (10)	-	129.051 (22)	135.501 (22)	137.941 (22)
LWEIGHT4	100.410 (30)	-	129.071 (8)	135.507 (20)	137.936 (22)
Adopted	100.41 (3)	114.56 (7)	129.065 (3)	135.507 (22)	137.936 (22)
Comments	wtd mean	$^{1987}\text{Da28}$	$^{1979}\text{He10}$	wtd mean	wtd mean

Nominal Energy /keV	141.01	145.849	153.977	168.65	173.964
$^{1979}\text{He10}$	-	-	-	-	173.964 (26)
$^{1971}\text{He23}$	140.89 (20)	146.19 (20)	153.99 (20)	-	174.00 (20)
$^{1973}\text{Ta25}$	-	-	-	-	-
$^{1977}\text{Ku01}$	141.0 (5)	-	-	-	-
$^{1979}\text{Bo30}$	-	-	153.956 (19)	-	-
$^{1987}\text{Da28}$	141.019 (30)	145.848 (11)	153.977 (11)	168.650 (11)	173.980 (10)
$^{1995}\text{Ba42}$	140.991 (22)	145.811 (22)	153.941 (22)	168.41 (9)	174.011 (41)
LWEIGHT4	140.999 (20)	145.842 (20)	153.967 (8)	168.53 (12)	173.995 (28)
Adopted	140.999 (20)	145.842 (20)	153.967 (11)	168.53 (12)	173.96 (3)
Comments	wtd mean	wtd mean	wtd mean	wtd mean	wtd mean

Comments on evaluation

Nominal Energy /keV	184.54	191.353	199.407	204.026	209.253
1979He10	-	-	-	-	-
1971He23	184.50 (11)	192.10 (30)	-	-	209.20 (20)
1973Ta25	-	-	-	-	-
1977Ku01	-	190.99 (20)	-	-	209.50 (50)
1979Bo30	-	-	-	-	209.238 (21)
1987Da28	184.540 (20)	191.353 (11)	199.408 (15)	204.027 (11)	209.254 (7)
1995Ba42	184.60 (5)	191.341 (21)	199.391 (21)	204.041 (21)	209.251 (21)
LWEIGHT4	184.547 (19)	191.351 (17)	199.402 (12)	204.029 (9)	209.248 (5)
Adopted	184.547 (19)	191.351 (17)	199.402 (15)	204.029 (11)	209.248 (7)
Comments	wtd mean	wtd mean	wtd mean	wtd mean	wtd mean

Nominal Energy /keV	214.85	223.8	231.42	257.7	263.62
1979He10	-	-	-	-	-
1971He23	-	223.70	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	214.85 (10)	223.85 (10)	231.42 (10)	257.52 (10)	263.58 (10)
1995Ba42	214.92 (10)	223.791 (21)	231.49 (5)	257.481 (21)	-
LWEIGHT4	214.89 (7)	223.793 (21)	231.477 (45)	257.482 (21)	-
Adopted	214.89 (10)	223.793 (21)	231.42 (10)	257.482 (21)	263.58 (10)
Comments	wtd mean	wtd mean	1987Da28	wtd mean	1987Da28

The value of 231.49 (5) keV line observed by 1995Ba42 is believed to relate to a separate gamma transition and has not been used.

Nominal Energy /keV	270.245	278.95	282.0	321.646	326.04
1979He10	270.245 (7)	-	282.022 (40)	321.646 (8)	-
1971He23	270.20 (50)	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	270.19 (21)	-	-	-	-
1979Bo30	270.235 (14)	-	-	-	-
1987Da28	270.245 (7)	278.952 (50)	281.922 (50)	321.653 (50)	326.04 (20)
1995Ba42	270.241 (21)	278.651 (21)	282.001 (21)	321.701 (31)	-
LWEIGHT4	270.255 (7)	278.80 (15)	281.989 (28)	321.688 (26)	-
Adopted	270.245 (7)	278.80 (15)	282.02 (4)	321.646 (8)	326.04 (20)
Comments	1979He10	wtd mean	1979He10	1979He10	1987Da28

Nominal Energy /keV	327.44	328.00	332.37	338.32	340.98
1979He10	-	-	332.371 (6)	338.320 (5)	-
1971He23	-	-	-	338.31 (40)	-
1973Ta25	-	-	-	-	-
1977Ku01	-	327.89 (22)	332.29 (10)	338.09 (22)	-
1979Bo30	-	328.003 (11)	-	338.321 (10)	-
1987Da28	-	328.004 (7)	332.374 (50)	338.324 (6)	340.964 (50)
1995Ba42	327.45 (4)	328.02 (4)	332.360 (21)	338.310 (21)	340.970 (21)
LWEIGHT4	-	328.004 (7)	332.366 (35)	338.342 (18)	340.969 (19)
Adopted	-	328.004 (7)	332.371 (6)	338.320 (5)	340.969 (21)
Comments	not used	wtd mean	1979He10	1979He10	wtd mean

The 327.44 keV line is listed in Artna-Cohen based upon expected presence inferred from ²²⁸Pa EC decay. However, only lines directly observed in ²²⁸Ac decay are considered here.

Nominal Energy /keV	356.94	372.57	377.99	384.47	389.12
1979He10	-	-	-	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	356.94 (10)	372.57 (20)	377.99 (10)	384.63 (20)	389.12 (15)
1995Ba42	356.35 (10)	372.590 (31)	377.98 (10)	384.43 (10)	389.40 (10)
LWEIGHT4	356.65 (30)	372.590 (30)	377.99 (7)	384.47 (9)	389.32 (13)
Adopted	356.7 (3)	372.59 (3)	377.99 (10)	384.47 (9)	389.32 (13)
Comments	wtd mean	wtd mean	wtd mean	wtd mean	wtd mean

Nominal Energy /keV	397.94	399.62	409.462	416.3	419.4
1979He10	-	-	409.460 (13)	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	409.487 (33)	-	-
1987Da28	397.95 (10)	399.63 (10)	409.456 (10)	416.31 (20)	419.43 (10)
1995Ba42	-	399.93 (7)	409.440 (21)	415.90 (8)	419.34 (10)
LWEIGHT4	-	399.83 (14)	409.464 (25)	415.96 (14)	419.38 (7)
Adopted	397.95 (10)	399.83 (14)	409.460 (13)	415.96 (14)	419.38 (7)
Comments	1987Da28	wtd mean	1979He10	wtd mean	wtd mean

Nominal Energy /keV	440.44	449.21	452.51	457.35	463.004
1979He10	440.450 (24)	449.11 (6)	-	-	463.002 (6)
1971He23	-	-	-	-	463.33 (41)
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	463.002 (13)
1987Da28	440.446 (50)	449.26 (10)	452.47 (10)	457.18 (15)	463.023 (10)
1995Ba42	440.390 (40)	449.22 (30)	452.51 (6)	-	463.01 (50)
LWEIGHT4	440.418 (35)	449.24 (7)	452.50 (5)	-	463.048 (15)
Adopted	440.450 (24)	449.11 (6)	452.50 (6)	457.18 (15)	463.002 (6)
Comments	1979He10	1979He10	wtd mean	1987Da28	wtd mean

Nominal Energy /keV	466.4	470.2	471.76	474.79	478.4
1979He10	-	-	-	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	478 (1)
1979Bo30	-	-	-	-	-
1987Da28	466.40 (10)	470.26 (20)	471.77 (15)	474.76 (10)	478.337 (50)
1995Ba42	-	469.89 (50)	-	475.09 (30)	478.440 (40)
LWEIGHT4	-	470.21 (19)	-	474.79 (10)	478.399 (37)
Adopted	466.40 (10)	470.21 (20)	471.77 (15)	474.79 (10)	478.40 (5)
Comments	1987Da28	wtd mean	1987Da28	wtd mean	wtd mean

Comments on evaluation

Nominal Energy /keV	480.94	490.33	492.37	497.64	503.823
1979He10	-	-	-	-	503.819 (23)
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	503.60 (33)
1979Bo30	-	-	-	-	-
1987Da28	480.95 (20)	490.33 (15)	492.38 (10)	497.50 (15)	503.83 (50)
1995Ba42	482.03 (5)	-	492.21 (10)	497.70 (10)	503.69 (20)
LWEIGHT4	481.5 (5)	-	492.29 (8)	497.64 (9)	503.67 (18)
Adopted	481.5 (5)	490.33 (15)	492.29 (8)	497.64 (10)	503.819 (23)
Comments	wtd mean	1987Da28	wtd mean	wtd mean	1979He10

Nominal Energy /keV	508.959	515.06	520.151	523.131	540.76
1979He10	508.955 (13)	-	520.16 (3)	523.129 (22)	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	508.968 (50)	515.07 (10)	520.18 (5)	523.118 (50)	540.77 (10)
1995Ba42	509.12 (8)	515.19 (11)	520.16 (8)	523.15 (11)	540.650 (50)
LWEIGHT4	509.04 (8)	515.12 (7)	520.17 (6)	523.13 (8)	540.674 (48)
Adopted	508.955 (13)	515.12 (7)	520.16 (3)	523.129 (22)	540.67 (5)
Comments	1979He10	wtd mean	1979He10	1979He10	wtd mean

Nominal Energy /keV	546.45	548.73	555.12	562.5	570.91
1979He10	-	-	-	562.496 (7)	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	547 (1)	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	546.479 (50)	548.74 (15)	555.13 (10)	562.529 (30)	570.91 (10)
1995Ba42	546.440 (21)	548.73 (11)	554.59 (30)	562.490 (40)	570.870 (40)
LWEIGHT4	546.445 (19)	548.73 (9)	555.07 (16)	562.509 (29)	570.876 (37)
Adopted	546.445 (21)	548.73 (11)	555.07 (16)	562.496 (7)	570.88 (4)
Comments	wtd mean	wtd mean	wtd mean	1979He10	wtd mean

Nominal Energy /keV	572.14	583.41	590.4	610.64	616.2
1979He10	572.10 (5)	-	-	-	616.212 (30)
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	572.30 (10)	583.419 (50)	-	610.65 (10)	616.27 (10)
1995Ba42	572.290 (21)	583.390 (10)	590.64 (11)	-	616.139 (50)
LWEIGHT4	572.290 (20)	583.391 (10)	-	-	616.20 (7)
Adopted	572.10 (5)	583.391 (10)	-	610.65 (10)	616.21 (3)
Comments	1979He10	wtd mean	not used	wtd mean	1979He10

Nominal Energy /keV	620.38	623.27	627.23	629.4	634.18
1979He10	-	-	-	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	620.390 (50)	623.27 (20)	627.24 (20)	629.410 (50)	634.18 (10)
1995Ba42	620.259 (50)	623.7 (2)	626.69 (10)	629.39 (20)	-
LWEIGHT4	620.32 (7)	623.48 (22)	626.80 (22)	629.409 (49)	-
Adopted	620.32 (7)	623.48 (22)	626.80 (22)	629.41 (5)	634.18 (10)
Comments	wtd mean	wtd mean	wtd mean	wtd mean	1987Da28

Nominal Energy /keV	640.34	648.84	651.5	660.1	663.88
1979He10	640.317 (37)	-	651.526 (28)	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	640.371 (50)	648.85 (10)	651.461 (50)	660.11 (30)	663.83 (10)
1995Ba42	640.319 (50)	649.11 (7)	651.49 (20)	660.17 (30)	663.91 (8)
LWEIGHT4	640.345 (36)	649.02 (12)	651.48 (14)	660.14 (21)	663.88 (6)
Adopted	640.32 (4)	649.02 (12)	651.53 (3)	660.1 (3)	663.88 (8)
Comments	1979He10	wtd mean	1979He10	wtd mean	wtd mean

Nominal Energy /keV	666.47	672.0	674.76	677.07	684.0
1979He10	666.451 (46)	-	674.625 (40)	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	666.46 (10)	672.01 (15)	674.61 (10)	677.12 (10)	-
1995Ba42	666.459 (40)	671.93 (10)	674.76 (10)	676.89 (20)	683.99 (30)
LWEIGHT4	666.46 (7)	671.95 (8)	674.69 (7)	677.08 (9)	-
Adopted	666.45 (5)	671.95 (8)	674.63 (4)	677.08 (10)	-
Comments	1979He10	wtd mean	1979He10	wtd mean	not used

Nomina Energy /keV	688.11	692.47	699.08	701.747	707.41
1979He10	-	-	-	701.742 (15)	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	688.112 (50)	-	699.09 (15)	701.752 (50)	707.422 (50)
1995Ba42	688.13 (8)	692.46 (7)	698.94 (10)	701.709 (40)	707.39 (30)
LWEIGHT4	688.117 (42)	-	698.99 (8)	701.731 (35)	707.421 (49)
Adopted	688.12 (4)	-	698.99 (10)	701.742 (15)	707.42 (5)
Comments	wtd mean	not used	wtd mean	1979He10	wtd mean

Comments on evaluation

Nominal Energy /keV	718.48	726.863	737.72	755.315	770.2
1979He10	-	727.317 (15)	-	755.313 (9)	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	718.49 (15)	726.876 (15)	737.733 (50)	755.325 (15)	-
1995Ba42	718.299 (21)	726.89 (10)	737.79 (20)	755.309 (21)	770.2 (2)
LWEIGHT4	718.303 (26)	726.88 (7)	737.736 (49)	755.317 (15)	-
Adopted	718.30 (3)	726.88 (10)	737.74 (5)	755.313 (9)	-
Comments	wtd mean	wtd mean	wtd mean	1979He10	not used

The 684 keV, 692 keV and 770 keV emissions have not been observed directly in ²²⁸Ac decay and are not included in this evaluation.

There is a considerable discrepancy between data measured for the 727 keV line by 1979He10 and the mean of values measured by 1987Da28 and 1995Ba42. The value measured by 1979He10 is not consistent with the decay scheme; this is possibly due to interference from the ²¹²Bi decay daughter. The weighted mean of 1987Da28 and 1995Ba42 is used instead.

Nominal Energy /keV	772.291	774.1	776.52	778.1	782.142
1979He10	772.291 (7)	-	-	-	782.140 (6)
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	772.294 (10)	774.11 (20)	776.57 (10)	-	782.154 (50)
1995Ba42	772.269 (21)	774.06 (10)	776.509 (30)	778.09 (20)	782.09 (20)
LWEIGHT4	772.282 (25)	774.07 (9)	776.514 (29)	-	782.12 (14)
Adopted	772.291 (7)	774.07 (10)	776.51 (3)	-	782.140 (6)
Comments	1979He10	wtd mean	wtd mean	not used	1979He10

Nominal Energy /keV	791.44	792.8	794.947	813.77	816.62
1979He10	-	-	794.942 (14)	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	817 (1)
1979Bo30	-	-	794.94 (14)	-	-
1987Da28	791.50 (25)	792.8 (10)	794.940 (10)	813.78 (15)	816.71 (10)
1995Ba42	791.42 (9)	792.69 (10)	794.959 (20)	813.92 (10)	816.92 (10)
LWEIGHT4	791.43 (8)	792.69 (10)	794.951 (14)	813.88 (8)	816.82 (7)
Adopted	791.43 (9)	792.69 (10)	794.942 (14)	813.88 (10)	816.82 (10)
Comments	wtd mean	wtd mean	1979He10	wtd mean	wtd mean

Nominal Energy /keV	824.934	830.486	835.71	840.377	853.17
1979He10	824.931 (25)	830.481 (8)	835.704 (8)	840.372 (9)	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	824.87 (10)	830.476 (20)	835.708 (20)	840.370 (20)	853.19 (10)
1995Ba42	-	830.469 (30)	835.639 (21)	840.349 (40)	853.96 (8)
LWEIGHT4	-	830.472 (22)	835.674 (35)	840.366 (18)	-
Adopted	824.931 (25)	830.481 (8)	835.704 (8)	840.372 (9)	853.96 (8)
Comments	1979He10	1979He10	1979He10	1979He10	1995Ba42

The values of the 853 keV line from 1987Da28 and 1995Ba42 are clearly discrepant; the value of 1995Ba42 has been preferred as it better fits the level scheme.

Nominal Energy /keV	870.45	873.11	874.45	877.39	880.76
1979He10	870.47 (7)	-	874.45 (8)	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	870.456 (50)	873.17 (15)	874.41 (15)	877.48 (10)	880.76 (10)
1995Ba42	870.438 (21)	872.99 (20)	874.49 (20)	877.34 (7)	-
LWEIGHT4	870.441 (19)	873.10 (12)	874.44 (12)	877.38 (6)	-
Adopted	870.47 (7)	873.10 (15)	874.45 (8)	877.38 (7)	880.76 (10)
Comments	1979He10	wtd mean	1979He10	wtd mean	1987Da28

Nominal Energy /keV	887.33	901.26	904.19	911.204	919.01
1979He10	-	-	904.20 (5)	911.196 (6)	-
1971He23	-	-	-	911.27 (25)	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	911.166 (40)	-
1987Da28	887.34 (10)	901.25 (15)	904.197 (50)	911.233 (11)	918.99 (10)
1995Ba42	887.18 (10)	90.388 (31)	904.178 (31)	911.188 (21)	919.39 (30)
LWEIGHT4	887.26 (8)	901.383 (30)	904.183 (26)	911.221 (13)	919.03 (12)
Adopted	887.26 (10)	901.38 (3)	904.20 (5)	911.196 (6)	919.03 (12)
Comments	wtd mean	wtd mean	1979He10	1979He10	wtd mean

Nominal Energy /keV	921.98	924.03	930.93	939.87	944.196
1979He10	-	-	-	-	944.191 (30)
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	922.00 (10)	-	930.95 (10)	939.89 (15)	944.178 (50)
1995Ba42	921.75 (10)	924.29 (20)	931.01 (7)	-	944.30 (6)
LWEIGHT4	921.87 (12)	-	930.99 (6)	-	944.23 (6)
Adopted	921.87 (12)	-	930.99 (7)	939.89 (15)	944.19 (3)
Comments	wtd mean	not used	wtd mean	1987Da28	1979He10

Nominal Energy /keV	947.982	958.61	964.766	968.971	975.98
1979He10	947.976 (24)	958.591 (38)	964.786 (8)	968.960 (9)	-
1971He23	-	-	964.48 (43)	968.88 (34)	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	964.68 (9)	969.161 (34)	-
1987Da28	947.968 (50)	958.638 (50)	964.783 (11)	968.989 (11)	975.978 (50)
1995Ba42	-	958.68 (11)	964.788 (21)	968.668 (21)	975.987 (50)
LWEIGHT4	-	958.645 (46)	964.783 (36)	968.90 (10)	975.983 (36)
Adopted	947.976 (24)	958.59 (4)	964.786 (8)	968.960 (9)	975.98 (5)
Comments	1979He10	1979He10	1979He10	1979He10	wtd mean

Comments on evaluation

Nominal Energy /keV	979.48	987.88	988.63	1000.69	1013.58
1979He10	-	-	-	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	979.50 (10)	987.73 (10)	988.65 (20)	1000.71 (15)	1013.60 (20)
1995Ba42	979.39 (40)	987.91 (10)	-	1000.67 (10)	1013.53 (13)
LWEIGHT4	979.49 (10)	987.87 (9)	-	1000.68 (8)	1013.55 (11)
Adopted	979.49 (10)	987.87 (10)	988.65 (20)	1000.68 (10)	1013.55 (13)
Comments	wtd mean	wtd mean	1987Da28	wtd mean	wtd mean

Nominal Energy /keV	1016.44	1017.92	1019.86	1033.248	1039.84
1979He10	-	-	-	1033.244 (23)	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1016.46 (15)	1017.94 (20)	1019.88 (10)	1033.240 (20)	1039.67 (15)
1995Ba42	1016.4 (10)	-	-	1033.26 (7)	1039.86 (6)
LWEIGHT4	1016.44 (8)	-	-	1033.241 (19)	1039.83 (7)
Adopted	1016.44 (10)	1017.94 (20)	1019.88 (10)	1033.244 (23)	1039.83 (7)
Comments	wtd mean	1987Da28	1987Da28	1979He10	wtd mean

Nominal Energy /keV	1040.92	1053.09	1054.22	1062.55	1065.19
1979He10	-	-	-	-	1065.168 (15)
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1040.94 (15)	1053.11 (20)	1054.13 (20)	1062.57 (15)	1065.200 (50)
1995Ba42	-	-	-	-	1065.20 (7)
LWEIGHT4	-	-	-	-	1065.200 (41)
Adopted	1040.94 (15)	1053.11 (20)	1054.13 (20)	1062.57 (15)	1065.168 (15)
Comments	1987Da28	1987Da28	1987Da28	1987Da28	1979He10

Nominal Energy /keV	1074.71	1088.18	1095.679	1103.43	1110.61
1979He10	-	-	1095.671 (23)	-	1110.604 (9)
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1074.73 (15)	1088.20 (15)	1095.711 (50)	1103.43 (10)	-
1995Ba42	-	-	1095.73 (14)	-	1110.537 (51)
LWEIGHT4	-	-	1095.713 (47)	-	-
Adopted	1074.73 (15)	1088.20 (15)	1095.671 (23)	1103.43 (10)	1110.604 (9)
Comments	1987Da28	1987Da28	1979He10	1987Da28	1979He10

201Nominal Energy /keV	1117.63	1135.24	1142.85	1148.16	1153.52
1979He10	-	-	-	-	1153.266 (35)
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1117.65 (10)	1135.26 (15)	1142.87 (15)	1148.14 (15)	1153.502 (50)
1995Ba42	-	1135.4 (10)	1142.8 (10)	1148.19 (14)	1153.59 (30)
LWEIGHT4	-	1135.26 (15)	1142.87 (15)	1148.17 (10)	1153.505 (50)
Adopted	1117.65 (10)	1135.26 (15)	1142.87 (15)	1148.17 (14)	1153.27 (4)
Comments	1987Da28	wtd mean	wtd mean	wtd mean	1979He10

Nominal Energy /keV	1157.14	1164.55	1175.31	1190.83	1217.03
1979He10	-	-	-	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1157.16 (15)	1164.52 (7)	1175.33 (10)	1190.83 (20)	1217.03 (10)
1995Ba42	-	1164.57 (7)	-	1190.9 (10)	-
LWEIGHT4	-	1164.55 (7)	-	1190.83 (20)	-
Adopted	1157.16 (15)	1164.55 (7)	1175.33 (10)	1190.83 (20)	1217.03 (10)
Comments	1987Da28	wtd mean	1987Da28	wtd mean	1987Da28

Nominal Energy /keV	1229.4	1245.16	1247.08	1250.04	1276.69
1979He10	-	-	1247.10 (5)	1250.062 (44)	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1229.42 (15)	1245.07 (20)	1247.065 (50)	1249.77 (15)	1276.72 (10)
1995Ba42	-	1245.16 (6)	1247.056 (51)	1249.69 (20)	-
LWEIGHT4	-	1245.15 (6)	1247.061 (36)	1249.74 (12)	-
Adopted	1229.42 (15)	1245.15 (6)	1247.10 (5)	1250.06 (5)	1276.72 (10)
Comments	1987Da28	wtd mean	1979He10	1979He10	1987Da28

Nominal Energy /keV	1286.27	1287.78	1309.71	1315.31	1337.33
1979He10	-	-	-	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1286.30 (20)	1287.71 (20)	1309.74 (20)	1315.37 (10)	1337.33 (20)
1995Ba42	1286.29 (30)	1287.78 (8)	1310.2 (10)	1315.19 (20)	-
LWEIGHT4	1286.29 (17)	1287.77 (7)	1309.76 (20)	1315.33 (10)	-
Adopted	1286.29 (20)	1287.77 (8)	1309.76 (20)	1315.33 (10)	1337.33 (20)
Comments	wtd mean	wtd mean	wtd mean	wtd mean	1987Da28

Comments on evaluation

Nominal Energy /keV	1344.59	1347.5	1357.78	1365.71	1374.24
1979He10	-	-	-	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1344.62 (15)	1347.50 (15)	1357.81 (15)	1365.73 (15)	1374.24 (7)
1995Ba42	1344.6 (10)	-	-	1365.71 (12)	1374.25 (7)
LWEIGHT4	1344.62 (15)	-	-	1365.71 (12)	1374.24 (6)
Adopted	1344.62 (15)	1347.50 (15)	1357.81 (15)	1365.71 (12)	1374.24 (7)
Comments	wtd mean	1987Da28	1987Da28	wtd mean	1987Da28

Nominal Energy /keV	1378.23	1385.39	1401.49	1415.55	1430.95
1979He10	-	-	-	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1378.23 (10)	1385.39 (10)	1401.52 (10)	1415.69 (10)	1430.98 (10)
1995Ba42	-	-	-	1415.41 (10)	1432.0 (10)
LWEIGHT4	-	-	-	1415.55 (14)	1430.99 (10)
Adopted	1378.23 (10)	1385.39 (10)	1401.52 (10)	1415.55 (14)	1430.99 (10)
Comments	1987Da28	1987Da28	1987Da28	wtd mean	wtd mean

Nominal Energy /keV	1434.22	1438.01	1451.4	1459.138	1469.71
1979He10	-	-	-	1459.131 (22)	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1434.22 (15)	1438.01 (10)	1451.43 (15)	1459.15 (15)	1469.74 (15)
1995Ba42	-	-	1451.4 (10)	1459.19 (20)	-
LWEIGHT4	-	-	1451.43 (15)	1459.16 (12)	-
Adopted	1434.22 (15)	1438.01 (10)	1451.43 (15)	1459.131 (22)	1469.74 (15)
Comments	1987Da28	1987Da28	wtd mean	1979He10	1987Da28

Nominal Energy /keV	1480.37	1495.93	1501.57	1529.02	1537.87
1979He10	-	1495.904 (16)	-	1529.010 (34)	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1480.37 (15)	1495.80 (5)	1501.600 (51)	-	1537.92 (10)
1995Ba42	1480.4 (3)	1496.14 (6)	1501.49 (20)	1529.01 (6)	1537.79 (20)
LWEIGHT4	1480.38 (15)	1495.97 (17)	1501.59 (5)	-	1537.89 (10)
Adopted	1480.38 (15)	1495.904 (16)	1501.59 (5)	1529.01 (4)	1537.89 (10)
Comments	wtd mean	1979He10	wtd mean	1979He10	wtd mean

Nominal Energy /keV	1548.65	1557.1	1559.78	1571.52	1573.26
1979He10	1548.65 (6)	1557.13 (7)	-	-	1573.23 (8)
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1548.64 (10)	1557.102 (51)	1559.88 (20)	1571.55 (20)	1573.39 (10)
1995Ba42	-	1557.05 (6)	-	-	1573.29 (30)
LWEIGHT4	-	1557.079 (39)	-	-	1573.38 (10)
Adopted	1548.65 (6)	1557.13 (7)	1559.88 (20)	1571.55 (20)	1573.23 (8)
Comments	1979He10	1979He10	1987Da28	1987Da28	1979He10

Nominal Energy /keV	1580.53	1588.19	1609.41	1625.06	1630.627
1979He10	1580.531 (25)	1588.200 (25)	-	1625.092 (35)	1630.618 (20)
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1580.522 (51)	1588.202 (51)	1609.44 (15)	1625.023 (51)	1630.663 (51)
1995Ba42	1580.49 (30)	1588.136 (52)	-	1624.99 (20)	1630.62 (6)
LWEIGHT4	1580.52 (5)	1588.170 (36)	-	1625.021 (49)	1630.644 (39)
Adopted	1580.531 (25)	1588.200 (25)	1609.44 (15)	1625.09 (4)	1630.618 (20)
Comments	1979He10	1979He10	1987Da28	1979He10	1979He10

Nominal Energy /keV	1638.281	1666.523	1671.64	1677.67	1684.01
1979He10	1638.272 (23)	1666.514 (13)	-	1677.66 (6)	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1638.304 (51)	1666.55 (51)	1671.67 (15)	1677.704 (51)	1684.04 (20)
1995Ba42	1638.29 (7)	1666.52 (6)	-	-	-
LWEIGHT4	1638.297 (41)	1666.52 (6)	-	-	-
Adopted	1638.272 (23)	1666.514 (13)	1671.67 (15)	1677.66 (6)	1684.04 (20)
Comments	1979He10	1979He10	1987Da28	1979He10	1987Da28

Nominal Energy /keV	1686.12	1700.59	1702.44	1706.17	1713.49
1979He10	1686.22 (11)	-	1702.40 (8)	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1686.095 (51)	1700.62 (20)	1702.57 (10)	1706.23 (10)	1713.51 (20)
1995Ba42	1686.14 (7)	-	1702.59 (30)	1706.15 (7)	1713.1 (10)
LWEIGHT4	1686.11 (4)	-	1702.57 (10)	1706.17 (6)	1713.49 (20)
Adopted	1686.22 (11)	1700.62 (20)	1702.40 (8)	1706.17 (7)	1713.49 (20)
Comments	1979He10	1987Da28	1979He10	wtd mean	wtd mean

Comments on evaluation

Nominal Energy /keV	1721.4	1724.2	1738.22	1740.4	1742.09
1979He10	-	1724.188 (43)	-	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1721.49 (30)	1724.28 (10)	1738.26 (25)	1740.46 (30)	1742.09 (30)
1995Ba42	-	1723.99 (20)	1738.465 (52)	-	-
LWEIGHT4	-	1724.22 (12)	1738.46 (5)	-	-
Adopted	1721.5 (3)	1724.19 (5)	1738.46 (5)	1740.5 (3)	1742.1 (3)
Comments	1987Da28	1979He10	wtd mean	1987Da28	1987Da28

Nominal Energy /keV	1745.28	1750.54	1758.11	1772.2	1784.4
1979He10	-	-	-	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1745.32 (20)	1750.58 (20)	1758.15 (10)	1772.22 (30)	1784.40 (30)
1995Ba42	-	-	1758.095 (53)	-	-
LWEIGHT4	-	-	1758.106 (47)	-	-
Adopted	1745.32 (20)	1750.58 (20)	1758.11 (5)	1772.2 (3)	1784.4 (3)
Comments	1987Da28	1987Da28	wtd mean	1987Da28	1987Da28

Nominal Energy /keV	1787.2	1795.15	1797.5	1800.86	1823.21
1979He10	-	-	-	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1787.30 (50)	1795.14 (50)	1797.50 (50)	1800.90 (20)	1823.26 (10)
1995Ba42	1787.18 (20)	1795.13 (6)	-	-	1823.17 (10)
LWEIGHT4	1787.20 (20)	1795.13 (6)	-	-	1823.22 (7)
Adopted	1787.20 (20)	1795.13 (6)	1797.5 (5)	1800.90 (20)	1823.22 (10)
Comments	wtd mean	wtd mean	1987Da28	1987Da28	wtd mean

Nominal Energy /keV	1826.7	1835.29	1842.14	1850.13	1870.81
1979He10	-	-	-	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1826.78 (30)	1835.47 (10)	1842.17 (10)	1850.17 (20)	1870.87 (9)
1995Ba42	-	1835.244 (53)	1842.13 (8)	-	1870.78 (9)
LWEIGHT4	-	1835.29 (9)	1842.15 (6)	-	1870.82 (7)
Adopted	1826.8 (3)	1835.29 (10)	1842.15 (8)	1850.17 (20)	1870.82 (9)
Comments	1987Da28	wtd mean	wtd mean	1987Da28	wtd mean

Nominal Energy /keV	1879.6	1887.12	1900.14	1907.13	1916.6
1979He10	-	-	-	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1879.60 (30)	1887.139 (51)	1900.11 (20)	1907.22 (20)	1915.94 (40)
1995Ba42	-	1887.113 (53)	1900.28 (30)	1907.11 (11)	1916.58 (30)
LWEIGHT4	-	1887.127 (37)	1900.16 (17)	1907.14 (10)	1916.34 (33)
Adopted	1879.6 (3)	1887.13 (5)	1900.16 (20)	1907.14 (11)	1916.3 (4)
Comments	1987Da28	wtd mean	wtd mean	wtd mean	wtd mean

Nominal Energy /keV	1919.5	1929.78	1936.2	1944.2	1952.37
1979He10	-	-	-	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1919.54 (30)	1929.78 (20)	1936.34 (30)	1944.24 (20)	1952.37 (15)
1995Ba42	-	-	-	-	1952.37 (10)
LWEIGHT4	-	-	-	-	1952.37 (8)
Adopted	1919.5 (3)	1929.78 (20)	1936.3 (3)	1944.24 (20)	1952.37 (10)
Comments	1987Da28	1987Da28	1987Da28	1987Da28	wtd mean

Nominal Energy /keV	1955.9	1958.4	1965.22	1971.9	1979.3
1979He10	-	-	-	-	-
1971He23	-	-	-	-	-
1973Ta25	-	-	-	-	-
1977Ku01	-	-	-	-	-
1979Bo30	-	-	-	-	-
1987Da28	1955.94 (50)	1958.41 (30)	1965.28 (20)	1971.96 (30)	1979.32 (30)
1995Ba42	-	-	1965.20 (12)	-	-
LWEIGHT4	-	-	1965.22 (10)	-	-
Adopted	1955.9 (5)	1958.4 (3)	1965.22 (12)	1972.0 (3)	1979.3 (3)
Comments	1987Da28	1987Da28	wtd mean	1987Da28	1987Da28

Nominal Energy /keV	2000.9	2029.4
1979He10	-	-
1971He23	-	-
1973Ta25	-	-
1977Ku01	-	-
1979Bo30	-	-
1987Da28	2000.98 (50)	2029.39 (50)
1995Ba42	-	-
LWEIGHT4	-	-
Adopted	2001.0 (5)	2029.4 (5)
Comments	1987Da28	1987Da28

Appendix II. Relative emission probabilities

Normalised to 100 for the 463 keV emission. Values marked with an asterisk (*) have been rejected from the weighted mean based on statistical evidence or on technical grounds. Where multiplets occur, the intensity is listed here once (in italics) and is the total measured intensity. Normalised values are also given - note the normalisation factor of 0.0445 (11) has been applied before rounding.

Nominal Energy /keV	18.4	42.46	56.88	57.752	77.34	99.505
1969Ar16	-	-	-	10.5 (5)*	-	30.6 (5)*
1971He23	-	-	-	10.5 (3)	-	28.3 (3)
1982Ma52	-	-	-	6.2 (4)*	-	36.5 (8)*
1982Sa36	-	-	-	11.4 (9)	-	31.0 (22)
1983Sc13	-	-	-	-	-	-
1987Da28	-	0.212 (61)	0.454 (94)	10.2 (11)	0.60 (12)	26.7 (33)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	-	-	10.57 (28)	-	28.33 (30)
Adopted	calc.	0.21 (6)	0.45 (9)	10.6 (3)	0.60 (12)	28.3 (3)
Normalised	-	0.009 (3)	0.020 (5)	0.470 (17)	0.027 (6)	1.26 (4)
Comments	no data	1987Da28	1987Da28	wtd mean	1987Da28	wtd mean

Nominal Energy /keV	100.41	114.56	129.065	135.507	137.936	141.000
1969Ar16	-	-	56.4 (4)*	-	-	0.80 (20)
1971He23	2.60 (10)	-	56.4 (4)	0.70 (10)	0.70 (10)	0.80 (20)
1982Ma52	-	-	54.1 (11)	-	-	-
1982Sa36	-	-	64.3 (56)	-	-	-
1983Sc13	-	-	49.6 (34)	-	-	-
1987Da28	2.18 (32)	0.23 (5)	61.2 (67)	0.41 (9)	0.55 (11)	1.18 (19)
1992Li05	-	-	55.3 (43)	-	-	~0.94
1995Ba42	-	-	-	-	-	-
LWEIGHT4	2.56 (12)	-	56.1 (5)	0.54 (14)	0.63 (10)	1.00 (19)
Adopted	2.56 (12)	0.23 (5)	56.1 (5)	0.54 (14)	0.63 (10)	1.00 (19)
Normalised	0.114 (6)	0.0102 (22)	2.50 (7)	0.024 (6)	0.028 (4)	0.045 (9)
Comments	wtd mean	1987Da28	wtd mean	wtd mean	wtd mean	wtd mean

Nominal Energy /keV	145.82	153.967	168.53	173.964	184.547	191.351
1969Ar16	3.6 (3)*	18.5 (4)*	-	-	2.2 (10)	3.9 (7)
1971He23	3.80 (10)	17.1 (4)	-	0.80 (20)	1.9 (6)	3.1 (3)
1982Ma52	-	17.1 (4)	-	-	0.70 (20)	-
1982Sa36	3.81 (30)	18.8 (15)	-	-	-	2.86 (27)
1983Sc13	-	15.6 (8)	-	-	-	-
1987Da28	3.6 (4)	18.2 (20)	<i>0.30 (6)</i>	0.82 (13)	1.64 (20)	3.03 (34)
1992Li05	-	15.8 (13)	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	3.79 (9)	16.94 (40)	-	0.81 (11)	1.21 (43)	2.98 (27)
Adopted	3.79 (10)	16.9 (4)	0.30 (6)	0.81 (13)	1.2 (4)	3.0 (3)
Normalised	0.169 (6)	0.754 (23)	-	0.036 (5)	0.054 (19)	0.133 (8)
Comments	wtd mean	wtd mean	<i>1987Da28</i>	wtd mean	wtd mean	wtd mean

Nominal Energy /keV	199.402	204.029	209.248	214.890	223.793	231.477
1969Ar16	6.0 (5)	3.0 (5)	93 (4)	-	1.6 (5)	-
1971He23	-	-	93 (4)*	-	-	-
1982Ma52	-	2.4 (2)	-	-	-	-
1982Sa36	7.4 (6)	2.38 (26)	93 (6)	-	-	-
1983Sc13	-	-	84.7 (34)	-	-	-
1987Da28	7.4 (8)	3.09 (34)	98 (9)	0.69 (11)	1.27 (14)	0.58 (9)
1992Li05	-	-	89.1 (35)	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	6.7 (5)	2.56 (17)	89.3 (19)	-	1.30 (13)	-
Adopted	6.7 (5)	2.56 (20)	89 (4)	0.69 (11)	1.30 (14)	0.58 (9)
Normalised	0.299 (23)	0.114 (8)	3.97 (13)	0.031 (5)	0.058 (6)	0.026 (4)
Comments	wtd mean	wtd mean	wtd mean	1987Da28	wtd mean	1987Da28

Nominal Energy /keV	257.482	263.580	270.245	278.80	282.022	321.646
1969Ar16	-	-	76.0 (40)	5.6 (5)	-	5.0 (10)
1971He23	-	-	78.0 (40)	-	-	-
1982Ma52	-	-	82.3 (20)	6.0 (4)	-	7.9 (5)
1982Sa36	-	-	79 (6)	-	2.62 (27)	5.24 (54)
1983Sc13	-	-	76.4 (29)	-	-	5.44 (53)
1987Da28	0.70 (8)	0.94 (11)	78 (7)	4.49 (37)	1.46 (14)	5.09 (44)
1992Li05	-	-	80.0 (27)	-	-	6.08 (12)
1995Ba42	0.626 (44)	0.97 (7)	-	-	-	-
LWEIGHT4	0.642 (39)	0.96 (6)	79.9 (13)	5.28 (49)	2.0 (6)	5.22 (28)
Adopted	0.64 (5)	0.96 (7)	79.9 (20)	5.3 (5)	2.0 (6)	5.2 (5)
Normalised	0.0286 (19)	0.043 (3)	3.55 (10)	0.235 (22)	0.09 (3)	0.232 (14)
Comments	wtd mean	wtd mean	wtd mean	<i>wtd mean</i>	wtd mean	wtd mean

Nominal Energy /keV	326.040	328.004	332.370	338.320	340.969	356.65
1969Ar16	-	68.0 (20)	7.2 (5)	250 (5)	10 (4)	-
1971He23	-	-	-	255 (5)	-	-
1982Ma52	-	78.3 (20)	-	282 (6)	-	-
1982Sa36	-	69.0 (58)	9.3 (11)	255 (17)	9.0 (10)	-
1983Sc13	-	-	6.2 (7)	250 (9)	9.0 (6)	-
1987Da28	0.78 (13)	69.7 (50)	9.8 (8)	256 (18)	7.7 (7)	0.400 (47)
1992Li05	-	-	10.57 (56)	260 (9)	9.8 (18)	-
1995Ba42	-	-	-	-	9.2 (6)	-
LWEIGHT4	-	68.3 (18)	8.4 (12)	255.1 (38)	9.09 (39)	-
Adopted	0.78 (13)	68.3 (20)	8.4 (12)	255 (5)	9.1 (6)	0.40 (5)
Normalised	0.035 (6)	3.04 (11)	0.37 (6)	11.4 (4)	0.405 (20)	0.0178 (21)
Comments	1987Da28	wtd mean	wtd mean	wtd mean	wtd mean	1987Da28

Comments on evaluation

Nominal Energy /keV	372.590	377.99	384.47	389.32	397.95	399.83
1969Ar16	-	-	-	-	-	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	0.158 (37)	0.576 (67)	0.158 (37)	0.242 (38)	0.642 (68)	0.691 (75)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	-	-	-	-	-
Adopted	0.16 (4)	0.58 (7)	0.16 (4)	0.24 (4)	0.64 (7)	0.69 (8)
Normalised	0.0070 (17)	0.026 (3)	0.0070 (17)	0.0108 (17)	0.029 (3)	0.031 (4)
Comments	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28

Nominal Energy /keV	409.46	415.96	419.38	440.45	449.11	452.50
1969Ar16	44 (2)	-	-	3.0 (5)	-	-
1971He23	-	-	-	-	-	-
1982Ma52	46.5 (10)	-	-	-	-	-
1982Sa36	42.9 (31)	-	-	-	-	-
1983Sc13	43.3 (19)	-	-	-	-	-
1987Da28	44.8 (33)	0.309 (51)	0.48 (8)	2.85 (23)	1.12 (13)	0.36 (12)
1992Li05	45.1 (23)	-	-	-	-	-
1995Ba42	44.7 (32)	-	-	-	-	0.456 (42)
LWEIGHT4	45.3 (7)	-	-	2.87 (21)	-	0.466 (40)
Adopted	45.3 (10)	0.31 (5)	0.48 (8)	2.87 (23)	1.12 (13)	0.47 (4)
Normalised	2.02 (6)	0.0138 (23)	0.022 (3)	0.128 (10)	0.050 (6)	0.0199 (19)
Comments	wtd mean	1987Da28	1987Da28	wtd mean	wtd mean	wtd mean

Nominal Energy /keV	457.18	466.40	470.21	471.77	474.79
1969Ar16	-	-	-	-	-
1971He23	-	-	-	-	-
1982Ma52	-	-	-	-	-
1982Sa36	-	-	-	-	-
1983Sc13	-	-	-	-	-
1987Da28	0.352 (57)	0.67 (8)	0.30 (6)	0.76 (8)	0.52 (8)
1992Li05	-	-	-	-	-
1995Ba42	-	-	-	-	-
LWEIGHT4	-	-	-	-	-
Adopted	0.35 (6)	0.67 (8)	0.30 (6)	0.76 (8)	0.52 (8)
Normalised	0.016 (3)	0.30 (4)	0.014 (3)	0.034 (4)	0.023 (4)
Comments	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28

Nominal Energy /keV	478.399	481.50	490.34	492.29	497.64	503.819	508.955
1969Ar16	5.7 (10)	-	-	-	-	3.0 (5)	12.0 (10)
1971He23	-	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	4.2 (9)	-
1987Da28	4.91 (44)	0.55 (11)	0.261 (56)	0.552 (61)	0.139 (43)	4.24 (37)	10.7 (12)
1992Li05	-	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-	-
LWEIGHT4	5.04 (40)	-	-	-	-	3.85 (41)	11.5 (8)
Adopted	5.0 (5)	0.55 (11)	0.26 (6)	0.55 (6)	0.14 (4)	3.9 (4)	11.5 (10)
Normalised	0.224 (19)	0.024 (5)	0.0116 (25)	0.025 (3)	0.0062 (19)	0.171 (19)	0.51 (4)
Comments	wtd mean	wtd mean	1987Da28	1987Da28	1987Da28	wtd mean	wtd mean

Nominal Energy /keV	515.12	520.159	523.129	540.674	546.445	548.73	555.07
1969Ar16	-	-	3.0 (1)	-	4.0 (5)	-	-
1971He23	-	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-	-
1987Da28	1.14 (13)	1.58 (14)	2.42 (22)	0.61 (7)	4.73 (38)	0.54 (8)	1.08 (12)
1992Li05	-	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-	-
LWEIGHT4	-	-	2.90 (22)	-	4.46 (35)	-	-
Adopted	1.13 (13)	1.58 (14)	2.90 (22)	0.61 (7)	4.5 (4)	0.54 (8)	1.08 (12)
Normalised	0.051 (6)	0.070 (7)	0.129 (10)	0.027 (3)	0.199 (16)	0.024 (4)	0.048 (6)
Comments	1987Da28	1987Da28	wtd mean	wtd mean	wtd mean	wtd mean	1987Da28

Nominal Energy /keV	562.496	570.876	572.10	583.391	610.65	616.212
1969Ar16	21 (2)	6.5 (10)	-	3.0 (5)	-	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	21.4 (26)	-	-	-	-	-
1983Sc13	19.8 (12)	-	-	-	-	-
1987Da28	18.8 (15)	3.82 (41)	3.52 (40)	2.61 (27)	0.54 (11)	1.88 (15)
1992Li05	20.3 (10)	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	20.0 (6)	4.2 (9)	-	2.70 (24)	-	-
Adopted	20.0 (10)	4.2 (9)	3.5 (4)	2.7 (3)	0.54 (11)	1.88 (15)
Normalised	0.89 (4)	0.19 (5)	0.156 (18)	0.120 (11)	0.024 (5)	0.084 (7)
Comments	wtd mean	wtd mean	1987Da28	wtd mean	1987Da28	1987Da28

Comments on evaluation

Nominal Energy /keV	620.32	623.48	626.80	629.409	634.18	640.317
1969Ar16	-	-	-	-	-	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	1.88 (15)	0.26 (6)	0.33 (7)	1.06 (12)	0.248 (50)	1.27 (14)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	-	-	-	-	-
Adopted	1.88 (15)	0.26 (6)	0.33 (7)	1.06 (12)	0.25 (5)	1.27 (14)
Normalised	0.084 (7)	0.012 (3)	0.015 (3)	0.047 (5)	0.0111 (22)	0.057 (6)
Comments	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28

Nominal Energy /keV	649.02	651.526	660.14	663.88	666.451	671.95
1969Ar16	-	-	-	-	-	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	0.94 (10)	2.12 (21)	0.121 (6)	0.66 (14)	1.45 (14)	0.61 (18)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	-	-	-	-	-
Adopted	0.94 (10)	2.12 (21)	0.121 (6)	0.66 (14)	1.45 (14)	0.61 (18)
Normalised	0.042 (5)	0.094 (10)	0.0054 (3)	0.029 (6)	0.065 (6)	0.027 (8)
Comments	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28

Nominal Energy /keV	674.625	677.08	688.117	698.99	701.742	707.421
1969Ar16	~3	-	-	-	~2.7	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	2.36 (22)	1.45 (14)	1.58 (14)	0.86 (13)	4.06 (31)	3.64 (40)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	-	-	-	-	-
Adopted	2.36 (22)	1.45 (14)	1.58 (14)	0.86 (13)	4.1 (3)	3.6 (4)
Normalised	0.105 (10)	0.065 (6)	0.070 (7)	0.038 (6)	0.181 (15)	0.162 (18)
Comments	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28

Nominal Energy /keV	718.303	727.317	737.736	755.313	772.291	774.07
1969Ar16	-	18 (4)	-	22 (3)	40 (4)	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	23.8 (26)	38.1 (30)	-
1983Sc13	-	-	-	23.6 (15)	32.2 (16)	-
1987Da28	0.44 (9)	14.5 (20)	0.87 (9)	21.8 (16)	33.9 (25)	1.39 (7)
1992Li05	-	-	-	23.4 (11)	34.0 (12)	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	15.2 (18)	-	23.1 (7)	34.1 (10)	-
Adopted	0.44 (9)	15.2 (20)	0.87 (9)	23.1 (11)	34.1 (12)	1.39 (7)
Normalised	0.019 (4)	0.68 (8)	0.039 (5)	1.03 (4)	1.52 (6)	0.062 (4)
Comments	1987Da28	wtd mean	1987Da28	wtd mean	wtd mean	1987Da28

Nominal Energy /keV	776.514	782.14	791.43	792.69	794.942	813.88
1969Ar16	-	17 (3)	-	-	105 (10)	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	100 (7)	-
1983Sc13	-	10.7 (14)	-	-	96.4 (35)	-
1987Da28	0.44 (14)	11.3 (8)	0.54 (17)	1.82 (9)	98 (7)	0.164 (37)
1992Li05	-	10.8 (13)	-	-	95.2 (34)	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	11.3 (7)	-	-	96.9 (22)	-
Adopted	0.44 (14)	11.3 (8)	0.54 (17)	1.82 (9)	97 (4)	0.16 (4)
Normalised	0.020 (6)	0.50 (4)	0.024 (7)	0.081 (5)	4.31 (14)	0.0073 (17)
Comments	1987Da28	1987Da28	1987Da28	1987Da28	wtd mean	wtd mean

Nominal Energy /keV	816.82	824.931	830.481	835.481	840.372	853.57
1969Ar16	-	-	16.5 (10)	39.5 (10)	23.0 (10)	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	15.0 (20)	42.9 (31)	22.1 (20)	-
1983Sc13	-	-	11.1 (12)	34.0 (22)	21.6 (13)	-
1987Da28	0.70 (8)	1.18 (13)	12.7 (9)	37.6 (26)	20.0 (16)	0.279 (45)
1992Li05	-	-	-	35.4 (20)	18.6 (24)	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	-	13.7 (12)	38.3 (12)	21.7 (7)	-
Adopted	0.70 (8)	1.18 (13)	13.7 (12)	38.3 (12)	21.7 (10)	0.28 (5)
Normalised	0.031 (4)	0.053 (6)	0.61 (6)	1.70 (7)	0.97 (4)	0.0124 (20)
Comments	1987Da28	1987Da28	wtd mean	wtd mean	wtd mean	wtd mean

Comments on evaluation

Nominal Energy /keV	870.47	872.99	874.45	877.38	880.76	887.26
1969Ar16	-	-	-	-	-	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	1.03 (11)	0.73 (15)	1.12 (25)	0.32 (6)	0.145 (43)	0.64 (7)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	-	-	-	-	-
Adopted	1.03 (11)	0.73 (15)	1.12 (25)	0.32 (6)	0.15 (5)	0.64 (7)
Normalised	0.046 (5)	0.032 (7)	0.050 (11)	0.014 (3)	0.0065 (19)	0.029 (3)
Comments	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28

Nominal Energy /keV	901.383	904.2	911.196	919.03	921.87	930.99
1969Ar16	-	-	590 (30)	-	-	-
1971He23	-	-	580 (20)	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	19.0 (25)	605 (42)	-	-	-
1983Sc13	-	17.6 (10)	591 (22)	-	-	-
1987Da28	0.38 (8)	17.0 (15)	606 (29)	0.64 (7)	<i>0.345 (51)</i>	<i>0.291 (45)</i>
1992Li05	-	-	573 (20)	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	17.5 (8)	588 (12)	-	-	-
Adopted	0.38 (8)	17.5 (10)	588 (20)	0.64 (7)	0.35 (5)	0.29 (5)
Normalised	0.017 (4)	0.78 (4)	26.2 (8)	0.028 (3)	<i>0.0154 (23)</i>	<i>0.0129 (20)</i>
Comments	1987Da28	wtd mean	wtd mean	wtd mean	<i>1987Da28</i>	<i>1987Da28</i>

Nominal Energy /keV	939.89	944.191	947.976	958.591	964.786	968.96
1969Ar16	-	-	-	-	100 (10)	360 (20)
1971He23	-	-	-	-	100 (10)	350 (10)
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	3.8 (14)	124 (9)	360 (26)
1983Sc13	-	-	-	-	112.2 (43)	361 (13)
1987Da28	0.21 (6)	2.24 (21)	2.49 (22)	6.8 (6)	118 (8)	372 (26)
1992Li05	-	-	-	-	110.4 (42)	349 (13)
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	-	-	6.4 (11)	112.1 (26)	357 (7)
Adopted	0.21 (6)	2.24 (21)	2.49 (22)	6.4 (11)	112 (5)	357 (10)
Normalised	0.009 (3)	0.100 (10)	0.111 (10)	0.29 (5)	4.99 (17)	15.9 (5)
Comments	1987Da28	1987Da28	1987Da28	wtd mean	wtd mean	wtd mean

Nominal Energy /keV	975.983	979.49	987.87	988.65	1000.68	1013.55
1969Ar16	-	-	4.3 (3)	-	-	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	1.16 (13)	0.62 (7)	1.82 (32)	1.82 (32)	0.121 (6)	0.109 (31)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	-	3.1 (12)	-	-	-
Adopted	1.16 (13)	0.62 (7)	3.1 (12)	1.8 (4)	0.121 (6)	0.11 (3)
Normalised	0.052 (6)	0.028 (3)	0.14 (6)	0.081 (14)	0.0054 (3)	0.0049 (14)
Comments	1987Da28	1987Da28	wtd mean	1987Da28	1987Da28	1987Da28

Nominal Energy /keV	1016.44	1017.94	1019.88	1033.244	1039.83	1040.94
1969Ar16	-	1.3 (3)	-	4.5 (3)	2.0 (4)	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	0.44 (6)	0.133 (31)	0.49 (10)	4.73 (38)	1.05 (22)	1.05 (22)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	-	-	-	1.27 (40)	-
Adopted	0.44 (6)	0.7 (6)	0.49 (10)	4.59 (24)	1.3 (4)	1.05 (22)
Normalised	0.019 (3)	0.03 (3)	0.022 (5)	0.204 (12)	0.056 (18)	0.047 (10)
Comments	1987Da28	wtd mean	1987Da28	wtd mean	wtd mean	1987Da28

Nominal Energy /keV	1053.11	1054.13	1062.57	1065.168	1074.73	1088.20
1969Ar16	~1	-	-	3.0 (2)	-	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	0.32 (9)	0.42 (13)	0.24 (7)	3.09 (29)	0.24 (7)	0.139 (31)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	-	-	3.03 (16)	-	-
Adopted	0.32 (9)	0.42 (13)	0.24 (7)	3.03 (20)	0.24 (7)	0.14 (3)
Normalised	0.014 (4)	0.019 (6)	0.011 (4)	0.135 (8)	0.011 (4)	0.0062 (14)
Comments	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28

Comments on evaluation

Nominal Energy /keV	1095.671	1103.43	1110.604	1117.65	1135.26	1142.87
1969Ar16	2.6 (3)	-	6.5 (10)	1.6 (3)	-	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	3.03 (28)	0.35 (6)	7.2 (6)	1.27 (19)	0.230 (38)	0.242 (50)
1992Li05	-	-	-	-	-	-
1995Ba42	-	0.224 (11)	-	-	-	-
LWEIGHT4	2.83 (22)	0.228 (24)	6.98 (51)	1.37 (16)	-	-
Adopted	2.8 (3)	0.228 (24)	7.0 (6)	1.37 (19)	0.23 (4)	0.24 (5)
Normalised	0.126 (10)	0.0102 (11)	0.311 (24)	0.061 (7)	0.0102 (17)	0.0108 (22)
Comments	wtd mean	wtd mean	wtd mean	wtd mean	1987Da28	1987Da28

Nominal Energy /keV	1148.17	1153.266	1157.16	1164.55	1175.33	1190.83
1969Ar16	-	4.0 (10)	-	~1.5	-	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	0.139 (31)	3.27 (29)	0.164 (31)	1.52 (14)	0.56 (8)	0.146 (37)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	3.33 (28)	-	-	-	-
Adopted	0.14 (3)	3.3 (3)	0.16 (3)	1.52 (14)	0.56 (8)	0.15 (4)
Normalised	0.0062 (14)	0.148 (13)	0.0073 (14)	0.067 (7)	0.025 (4)	0.0065 (17)
Comments	1987Da28	wtd mean	1987Da28	1987Da28	1987Da28	1987Da28

Nominal Energy /keV	1217.03	1229.42	1245.15	1247.10	1250.062	1276.72
1969Ar16	-	-	-	-	-	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	0.50 (8)	0.176 (55)	2.24 (44)	11.21 (55)	1.46 (14)	0.33 (7)
1992Li05	-	-	-	11.5 (14)	-	-
1995Ba42	-	-	2.50 (18)	-	-	-
LWEIGHT4	-	-	2.46 (17)	11.78 (46)	-	-
Adopted	0.50 (8)	0.18 (6)	2.46 (18)	11.8 (5)	1.46 (14)	0.33 (7)
Normalised	0.022 (4)	0.0078 (25)	0.110 (8)	0.524 (24)	0.065 (6)	0.015 (3)
Comments	1987Da28	1987Da28	wtd mean	wtd mean	1987Da28	1987Da28

Nominal Energy /keV	1286.29	1287.77	1309.76	1315.33	1337.33	1344.62
1969Ar16	-	3.0 (2)	-	-	-	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	1.17 (24)	1.88 (38)	0.44 (15)	0.35 (7)	0.115 (37)	0.212 (44)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	2.44 (56)	-	-	-	-
Adopted	1.17 (24)	2.4 (6)	0.44 (15)	0.35 (7)	0.12 (4)	0.21 (5)
Normalised	0.052 (11)	0.109 (25)	0.020 (7)	0.015 (3)	0.0051 (16)	0.0094 (20)
Comments	1987Da28	wtd mean	1987Da28	1987Da28	1987Da28	1987Da28

Nominal Energy /keV	1347.50	1357.81	1365.71	1374.24	1378.23	1385.39
1969Ar16	-	-	-	-	-	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	0.36 (8)	0.48 (10)	0.32 (7)	0.32 (9)	0.139 (43)	0.248 (50)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	0.447 (22)	-	-
LWEIGHT4	-	-	-	0.440 (29)	-	-
Adopted	0.36 (8)	0.48 (10)	0.32 (7)	0.44 (3)	0.14 (5)	0.25 (5)
Normalised	0.016 (4)	0.021 (5)	0.014 (3)	0.0196 (14)	0.0062 (19)	0.0111 (22)
Comments	1987Da28	1987Da28	1987Da28	wtd mean	1987Da28	1987Da28

Nominal Energy /keV	1401.52	1415.55	1430.99	1434.22	1438.01	1451.43
1969Ar16	-	-	-	-	-	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	0.29 (6)	0.50 (10)	0.82 (17)	0.188 (55)	0.139 (37)	0.248 (50)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	-	-	-	-	-
Adopted	0.29 (6)	0.50 (10)	0.82 (17)	0.19 (6)	0.14 (4)	0.25 (5)
Normalised	0.013 (3)	0.022 (5)	0.037 (8)	0.0084 (25)	0.0062 (17)	0.0111 (22)
Comments	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28

Comments on evaluation

Nominal Energy /keV	1459.131	1469.74	1480.38	1495.904	1501.59	1529.01
1969Ar16	20.0 (5)	-	-	21.0 (4)	11.6 (2)	~1
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	17.3 (10)	-	-	18.7 (16)	9.6 (11)	-
1987Da28	18.8 (15)	0.47 (9)	0.38 (8)	21.2 (16)	11.2 (10)	1.33 (14)
1992Li05	24.3 (17)	-	-	18.9 (13)	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	19.5 (11)	-	-	20.73 (43)	11.53 (25)	-
Adopted	19.5 (11)	0.47 (9)	0.38 (8)	20.7 (5)	11.53 (25)	1.33 (14)
Normalised	0.87 (5)	0.021 (5)	0.017 (4)	0.92 (3)	0.513 (17)	0.059 (6)
Comments	wtd mean	1987Da28	1987Da28	wtd mean	wtd mean	1987Da28

Nominal Energy /keV	1537.89	1548.65	1557.13	1559.88	1571.55	1573.23
1969Ar16	~0.8	~0.7	3.8 (2)	-	-	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	1.10 (12)	0.89 (10)	4.18 (37)	0.48 (10)	0.133 (37)	0.77 (9)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	-	3.89 (18)	-	-	-
Adopted	1.10 (12)	0.89 (10)	3.89 (20)	0.48 (10)	0.13 (4)	0.77 (9)
Normalised	0.049 (6)	0.040 (5)	0.173 (9)	0.021 (5)	0.0059 (17)	0.034 (4)
Comments	1987Da28	1987Da28	wtd mean	1987Da28	1987Da28	1987Da28

Nominal Energy /keV	1580.531	1588.20	1609.44	1625.092	1630.618	1638.272
1969Ar16	17 (3)	71 (3)	-	7.0 (20)	33.0 (20)	10.0 (10)
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	11.6 (20)	72.4 (29)	-	-	34.0 (22)	10.2 (9)
1987Da28	14.5 (14)	76.4 (52)	0.182 (37)	6.00 (51)	38.8 (31)	11.0 (10)
1992Li05	13.7 (14)	66.0 (17)	-	-	33.8 (12)	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	13.9 (9)	68.8 (20)	-	6.06 (50)	34.1 (9)	10.43 (55)
Adopted	13.9 (14)	68.8 (20)	0.182 (37)	6.1 (5)	34.1 (12)	10.4 (9)
Normalised	0.62 (4)	3.06 (12)	0.0081 (17)	0.270 (23)	1.52 (6)	0.46 (3)
Comments	wtd mean	1987Da28	1987Da28	wtd mean	wtd mean	wtd mean

Nominal Energy /keV	1666.514	1671.67	1677.66	1684.04	1686.22	1700.62
1969Ar16	3.8 (2)	-	~1.2	-	2.0 (2)	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	4.18 (37)	0.091 (31)	1.27 (14)	0.35 (11)	2.24 (21)	0.236 (56)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	3.89 (18)	-	-	-	2.11 (15)	-
Adopted	3.89 (20)	0.9 (3)	1.27 (14)	0.35 (11)	2.11 (20)	0.24 (6)
Normalised	0.173 (9)	0.0043 (14)	0.057 (6)	0.015 (5)	0.094 (7)	0.0105 (25)
Comments	wtd mean	1987Da28	1987Da28	1987Da28	wtd mean	1987Da28

Nominal Energy /keV	1702.40	1706.17	1713.49	1721.49	1724.188	1738.46
1969Ar16	1.5 (2)	-	-	-	~0.5	~0.6
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	1.13 (12)	0.200 (26)	0.127 (25)	0.133 (43)	0.68 (8)	0.41 (9)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	1.23 (17)	-	-	-	-	-
Adopted	1.23 (17)	0.20 (3)	0.127 (25)	0.13 (5)	0.68 (8)	0.41 (9)
Normalised	0.055 (7)	0.0089 (12)	0.0057 (11)	0.0059 (19)	0.030 (4)	0.018 (4)
Comments	wtd mean	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28

Nominal Energy /keV	1740.46	1742.09	1745.32	1750.58	1758.11	1772.22
1969Ar16	-	~0.5	-	-	~0.8	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	0.26 (8)	0.188 (55)	0.152 (20)	0.188 (20)	0.81 (9)	0.042 (12)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	-	-	-	-	-
Adopted	0.26 (8)	0.19 (6)	0.152 (20)	0.188 (20)	0.81 (9)	0.042 (12)
Normalised	0.011 (4)	0.0084 (25)	0.0067 (9)	0.0084 (9)	0.036 (4)	0.0019 (5)
Comments	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28

Comments on evaluation

Nominal Energy /keV	1784.40	1787.20	1795.13	1797.50	1800.90	1823.22
1969Ar16	-	-	-	-	-	1.00 (10)
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	0.139 (25)	0.030 (12)	0.048 (18)	0.048 (18)	0.103 (19)	1.03 (12)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	-	-	-	-	1.01 (8)
Adopted	0.139 (25)	0.030 (12)	0.048 (18)	0.048 (18)	0.103 (19)	1.01 (10)
Normalised	0.0062 (11)	0.0013 (5)	0.0022 (8)	0.0022 (8)	0.0046 (8)	0.046 (5)
Comments	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28	wtd mean

Nominal Energy /keV	1826.78	1835.29	1842.15	1850.17	1870.82	1879.60
1969Ar16	-	0.80 (10)	0.70 (10)	-	0.60 (10)	-
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	0.048 (18)	0.89 (10)	0.98 (11)	0.103 (19)	0.57 (6)	0.030 (12)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	0.84 (7)	0.83 (14)	-	0.578 (52)	-
Adopted	0.048 (18)	0.84 (10)	0.83 (14)	0.103 (19)	0.58 (5)	0.030 (12)
Normalised	0.0022 (8)	0.038 (4)	0.037 (6)	0.0046 (8)	0.0257 (24)	0.0013 (5)
Comments	1987Da28	wtd mean	wtd mean	wtd mean	wtd mean	1987Da28

Nominal Energy /keV	1887.13	1900.16	1907.14	1916.34	1919.54	1929.78
1969Ar16	2.1 (2)	-	~0.3	-	-	~0.4
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	2.12 (21)	0.067 (13)	0.279 (28)	0.018 (6)	0.048 (12)	0.467 (54)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	2.11 (14)	-	-	-	-	-
Adopted	2.11 (20)	0.067 (13)	0.28 (3)	0.018 (6)	0.048(12)	0.47 (6)
Normalised	0.094 (7)	0.0030 (6)	0.0124 (13)	0.008 (3)	0.0022 (6)	0.0208 (24)
Comments	wtd mean	1987Da28	1987Da28	1987Da28	1987Da28	1987Da28

Nominal Energy /keV	1936.34	1944.24	1952.37	1955.90	1958.41	1965.22
1969Ar16	-	-	1.4 (2)	-	-	0.60 (10)
1971He23	-	-	-	-	-	-
1982Ma52	-	-	-	-	-	-
1982Sa36	-	-	-	-	-	-
1983Sc13	-	-	-	-	-	-
1987Da28	0.048 (12)	0.048 (12)	1.39 (14)	0.018 (6)	0.036 (12)	0.478 (48)
1992Li05	-	-	-	-	-	-
1995Ba42	-	-	-	-	-	-
LWEIGHT4	-	-	1.40 (11)	-	-	0.502 (48)
Adopted	0.048 (12)	0.048 (12)	1.40 (14)	0.018 (6)	0.036 (12)	0.50 (5)
Normalised	0.0022 (6)	0.0022 (6)	0.062 (5)	0.008 (3)	0.0016 (5)	0.0223 (22)
Comments	1987Da28	1987Da28	wtd mean	1987Da28	1987Da28	wtd mean

Nominal Energy /keV	1971.96	1979.32	2001.0	2029.4
1969Ar16	-	-	-	-
1971He23	-	-	-	-
1982Ma52	-	-	-	-
1982Sa36	-	-	-	-
1983Sc13	-	-	-	-
1987Da28	0.085 (19)	0.042 (12)	0.024 (6)	0.042 (12)
1992Li05	-	-	-	-
1995Ba42	-	-	-	-
LWEIGHT4	-	-	-	-
Adopted	0.085 (19)	0.042 (12)	0.024 (6)	0.042 (12)
Normalised	0.0038 (8)	0.0019 (5)	0.0011 (3)	0.0019 (5)
Comments	1987Da28	1987Da28	1987Da28	1987Da28

**²²⁸Th – Comments on evaluation of decay data
by A. L. Nichols**

Evaluated: July/August 2001

Re-evaluated: January 2004 and April 2010

Evaluation Procedures

Limitation of Relative Statistical Weight Method (LWM) was applied to average numbers throughout the evaluation. The uncertainty assigned to the average value was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

²²⁸Th ($T_{1/2} = 698.6$ days) decays 100 % by alpha-particle emission ($Q(\alpha) = 5520.08$ (22) keV) to various excited levels and the ground state of ²²⁴Ra ($T_{1/2} = 3.631$ days). A reasonably well-defined decay scheme was derived from the alpha-particle studies of 1969Pe17, 1970Ba20, 1976BaZZ and 1993Ba72, and the gamma-ray measurements of 1977Ku15 and 1984Ge07. An additional gamma transition was added to the proposed decay scheme from equivalent studies of ²²⁴Fr decay by 1981Ku02: 908.28-keV gamma ray depopulating the 992.65-keV nuclear level of ²²⁴Ra. Weighted mean relative emission probabilities were calculated for the 131.612-, 166.410-, 205.99- and 215.985-keV gamma rays, while equivalent data for the other gamma transitions were adopted from the measurements of 1977Ku15; all of these relative emission probabilities were defined in terms of the 84.373-keV gamma ray (100 %).

²⁰O cluster decay has been observed by 1993Bo20 to be 1.13 (22) E-13. Subsequent reviews by 1995Ar33 and 1997Tr17 list a cluster-decay branching fraction of 1.13 (22) E-13 and 1.1 (2) E-13, respectively, based primarily on the earlier measurement.

Nuclear Data

The ²²⁸Th decay chain is important in quantifying the environmental impact of the decay of naturally-occurring ²³²Th. Certain radionuclides in this decay chain are noteworthy because of their decay characteristics: ²²⁴Ra alpha decay to ²²⁰Rn; ²¹²Bi and ²⁰⁸Tl gamma-ray emissions. ²⁰⁸Tl in particular emits high-energy gamma rays that represent a well-defined spectroscopic signature for this decay chain.

Half-life

The measurements of 1956Ki16, 1962Ma57, 1971Jo14 and 2002Un02 were adopted to give a least-squares weighted mean half-life of 698.55 (32) days. ²²⁸Th half-life quoted in 2002Un02 is also listed within 1992Un01. Woods has recommended a half-life of 698.60 (23) days (2007BeZP), but without due consideration of the calculated uncertainty with respect to the measured values, see relevant footnotes.

Reference	Half-life (d) [*]
1918Me01	695.8 [1.905 a] [†]
1956Ki16	697.6 (7) [1.910 (2) a]
1962Ma57	696.9 (15) [1.908 (4) a]
1962Ma57	703 (7) [1.924 (20) a] [‡]
1971Jo14	698.77 (32)
2002Un02	698.60 (36)
Recommended value	698.55 (32) [§] or 1.9126 (9) a

^{*} Conversion factor: 1 tropical year \equiv 365.2422 days.

[†] Uncertainty not specified – not included in weighted mean analysis of the data set.

[‡] Defined as an outlier.

[§] Recommended uncertainty adjusted from ± 0.22 to ± 0.32 , in alignment with the smallest uncertainty of the values used to calculate the average value.

Alpha Particles

Energies

All alpha-particle energies were derived from the structural details of the proposed decay scheme. While the energies of the main alpha-particle emissions have been directly measured by 1953As31, 1970Ba20, 1971Gr07, 1976BaZZ and 1991Ry01, the nuclear level energies of 1997Ar05 and evaluated Q-value of 5520.08 (22) keV (2003Au03) were used to determine the recommended energies and uncertainties of the alpha-particle emissions, while allowing for the significant recoil components.

Adopted nuclear levels of ²²⁴Ra: J^π and origins (1997Ar05).

Nuclear level	Nuclear level energy (keV)	J ^π	Origins
0	0.0	0 +	²²⁴ Fr β ⁻ decay, ²²⁴ Ac EC decay, ²²⁸ Th α decay, ²²⁶ Ra(p,t), ²²⁶ Ra(α,α'2nγ), ²²⁶ Ra(⁵⁸ Ni, ⁶⁰ Ni)
1	84.373 ± 0.003	2 +	²²⁴ Fr β ⁻ decay, ²²⁴ Ac EC decay, ²²⁸ Th α decay, ²²⁶ Ra(p,t), ²²⁶ Ra(α,α'2nγ), ²²⁶ Ra(⁵⁸ Ni, ⁶⁰ Ni)
2	215.985 ± 0.004	1 -	²²⁴ Fr β ⁻ decay, ²²⁴ Ac EC decay, ²²⁸ Th α decay, ²²⁶ Ra(p,t), ²²⁶ Ra(α,α'2nγ), ²²⁶ Ra(⁵⁸ Ni, ⁶⁰ Ni)
3	250.783 ± 0.005	4 +	²²⁴ Fr β ⁻ decay, ²²⁴ Ac EC decay, ²²⁸ Th α decay, ²²⁶ Ra(α,α'2nγ), ²²⁶ Ra(⁵⁸ Ni, ⁶⁰ Ni)
4	290.36 ± 0.04	(3) -	²²⁴ Fr β ⁻ decay, ²²⁴ Ac EC decay, ²²⁸ Th α decay, ²²⁶ Ra(α,α'2nγ), ²²⁶ Ra(⁵⁸ Ni, ⁶⁰ Ni)
5	433.07 ± 0.10	(5) -	²²⁴ Fr β ⁻ decay, ²²⁸ Th α decay, ²²⁶ Ra(α,α'2nγ), ²²⁶ Ra(⁵⁸ Ni, ⁶⁰ Ni)
6	479.20 ± 0.18	(6 +)	²²⁸ Th α decay, ²²⁶ Ra(p,t), ²²⁶ Ra(α,α'2nγ), ²²⁶ Ra(⁵⁸ Ni, ⁶⁰ Ni)
7	916.34 ± 0.07	0 +	²²⁴ Fr β ⁻ decay, ²²⁸ Th α decay, ²²⁶ Ra(p,t)
8	992.65 ± 0.06	(2 +)	²²⁴ Fr β ⁻ decay, ²²⁸ Th α decay

Measured and recommended energies of the main alpha-particle emissions of ²²⁸Th.

E _α (keV)	1953As31					Recommended value*
	1953As31	1970Ba20	1971Gr07	1976BaZZ	1991Ry01	
α _{0,8}	-	-	-	-	-	4448.00 (23)
α _{0,7}	-	-	-	-	-	4522.97 (23)
α _{0,6}	-	-	-	-	-	4952.5 (3)
α _{0,5}	-	-	-	-	-	4997.76 (24)
α _{0,4}	-	5136.1	-	-	-	5137.97 (22)
α _{0,3}	5173	5171.5	-	-	-	5176.86 (22)
α _{0,2}	5208	5208.9	-	-	-	5211.05 (22)
α _{0,1}	5388.5 (10)	5338.6	5340.54 (15)	5339.2 (10)	5340.36 (15)	5340.35 (22)
α _{0,0}	5421 (1)	5420.0	5423.33 (22)	5420.6 (10)	5423.15 (22)	5423.24 (22)

* Determined from the nuclear level energies of 1997Ar05 and evaluated Q-value of 5520.08 (22) keV (2003Au03).

Emission Probabilities

An alpha-particle emission probability of 73.4 (5) % was derived for alpha decay directly to the ground state of ²²⁴Ra, based on the various alpha-particle studies. This value and the gamma-ray data were used in conjunction with the theoretical internal conversion coefficients to determine a normalisation factor of 0.0119 (3) per 100 disintegrations for the relative emission probabilities of the gamma rays (see below).

Published alpha-particle emission probabilities per 100 disintegrations of ²²⁸Th.

E _α (keV)	P _α				
	1953As31	1969Pe17	1970Ba20	1976BaZZ	1993Ba72
4448.00 (23)	-	-	-	-	-
4522.97 (23)	-	-	-	-	-
4952.5 (3)	-	-	-	-	-
4997.76 (24)	-	-	-	-	-
5137.97 (22)	-	-	~ 0.05	-	-
5176.86 (22)	0.2	-	0.18	-	-
5211.05 (22)	0.4	-	0.36	-	-
5340.35 (22)	28	26.7 (2)	26.7	26.6 (5)	26.0 (8)
5423.24 (22)	71	[73.3 (2)]	72.7	72.4 (10)	74.0 (6)

Alpha-particle emission probability data of $^{1969}\text{Pe}17$ are effectively normalised to 73.3 (2) % and 26.7 (2) %.

$^{1976}\text{BaZZ}$ measurements require re-normalisation to $(100 - 0.36 - 0.18 - 0.05) = 99.41$ %
 $(72.4 + 26.6) N = 99.41$
 $N = 1.00414$ to give $P_{\alpha}(5423.24 \text{ keV})$ of 72.7 %, and uncertainty of ± 1.0 ;
 and $P_{\alpha}(5340.35 \text{ keV})$ of 26.7 %, and uncertainty of ± 0.5 .

$^{1993}\text{Ba}72$ studies also require re-normalisation to give $P_{\alpha}(5423.24 \text{ keV})$ of 73.6 % and uncertainty of ± 0.6 ; and $P_{\alpha}(5340.35 \text{ keV})$ of 25.8 %, and uncertainty of ± 0.8 .

A weighted mean value of 73.4 (5) % (0.734 (5)) was determined for $P_{\alpha}(5423.24 \text{ keV})$ from the data of $^{1976}\text{BaZZ}$ and $^{1993}\text{Ba}72$, which has been matched against a value of 26.0 (5) % (0.260 (5)) for $P_{\alpha}(5340.35 \text{ keV})$.

The absolute emission probabilities of the majority of the other alpha particles were calculated from population-depopulation of the nuclear level of ^{224}Ra and the gamma-ray normalisation factor. Although a consistent decay scheme was derived, further detailed alpha-particle measurements are required to develop and support the overall correctness of the proposed decay scheme. A hindrance factor (HF) of 1.000 for the 5423.24-keV alpha-particle emission yields $r_0(^{224}\text{Ra})$ of 1.5339 (3) fm, whereas the recommended value is 1.5332 (8) fm (1998Ak04).

Adjusted alpha-particle emission probabilities per 100 disintegrations of ^{228}Th , and hindrance factors.

$E_{\alpha}(\text{keV})$	P_{α}						HF
	$^{1953}\text{As}31$	$^{1969}\text{Pe}17$	$^{1970}\text{Ba}20$	$^{1976}\text{BaZZ}$	$^{1993}\text{Ba}72$	Recommended value*	
4448.00 (23)	-	-	-	-	-	$4.5 (7) \times 10^{-6}$	7.2
4522.97 (23)	-	-	-	-	-	$1.7 (3) \times 10^{-5}$	7.0
4952.5 (3)	-	-	-	-	-	$2.4 (5) \times 10^{-5}$	4600
4997.76 (24)	-	-	-	-	-	$1.0 (2) \times 10^{-5}$	21400
5137.97 (22)	-	-	~ 0.05	-	-	0.036 (6)	44
5176.86 (22)	0.2	-	0.18	-	-	0.218 (4)	12.5
5211.05 (22)	0.4	-	0.36	-	-	0.408 (7)	10.7
5340.35 (22)	28	26.7 (2)	26.7	26.7 (5)	25.8 (8)	26.0 (5)	0.958
5423.24 (22)	71	[73.3 (2)]	72.7	72.7 (10)	73.6 (6)	73.4 (5) [‡]	1.000
						$\Sigma 100.1 (7)$	

* Recommended emission probabilities of the low-intensity α transitions were derived from the evaluated gamma-ray emission probabilities and theoretical internal conversion coefficients.

[‡] $P_{\alpha}(5423.24 \text{ keV})$ of 73.4 (5) % is effectively the weighted mean of the re-normalised studies ($^{1976}\text{BaZZ}$, $^{1993}\text{Ba}72$), which has been subsequently matched with $P_{\alpha}(5340.35 \text{ keV})$ of 26.0 (5) %.

Gamma Rays

Energies

Although energies of the gamma-ray emissions have been measured by 1968Da21 and 1997Ku15 in particular, the well-defined nuclear level energies of 1997Ar05 were used to determine the recommended energies and associated uncertainties of the gamma-ray emissions between the various populated-depopulated levels because of their more extensive origins.

Measured and recommended gamma-ray energies.

E_{γ} (keV)			
1968Da21	1977Ku15	1977Ku25	Recommended value*
-	74.4 (1) [‡]	-	74.38 (4)
-	84.371 (3) [†]	84.371 (3)	84.373 (3)
131.6 (8)	131.610 (4) [†]	131.610 (4)	131.612 (5)
-	142.0 (5) [‡]	-	142.71 (11)
166.5 (8)	166.407 (4) [†]	166.407 (4)	166.410 (6)
-	182.2 (2) [‡]	-	182.29 (10)
-	205.93 (5)	-	205.99 (4)
216.1 (6)	215.979 (5) [†]	215.979 (5)	215.985 (4)
-	228.5 (2)	-	228.42 (18)
-	700.5 (5) [‡]	-	700.36 (7)
-	742.2 (5)	-	741.87 (6)
-	832.0 (2)	-	831.97 (7)
-	-	-	908.28 (6)
-	992.9 (10)	-	992.65 (6)

[†] Identical value and uncertainty also reported by 1977Ku25.

[‡] Data derived from coincidence measurements.

* Determined from the nuclear level energies of 1997Ar05.

Emission Probabilities

Gamma-ray emission probabilities have been partially or fully determined in the measurements of 1977Ku15, 1982Sa36 and 1984Ge07. However, the data derived by 1982Sa36 are significantly lower by 20 % to 30 % compared with the equivalent values measured by 1977Ku15 and 1984Ge07, and therefore they were set aside from in the weighted mean analysis. Weighted mean relative emission probabilities were calculated for the 131.612-, 166.410-, 205.99- and 215.985-keV gamma rays, while equivalent data for the other gamma emissions were directly adopted from the measurements of 1977Ku15. An additional gamma transition was added to the proposed decay scheme from the equivalent studies of ²²⁴Fr decay by 1981Ku02 as a 908.28-keV gamma ray depopulating the 992.65-keV nuclear level of ²²⁴Ra - this gamma transition may have been observed in the α decay of ²²⁸Th by 1977Ku15, but was adjudged by them to be background radiation (within the 911.2-keV peak). All of these relative emission probabilities were defined in terms of the emission probability of the 84.373-keV gamma ray (100.0 %).

Published gamma-ray emission probabilities.

E_{γ} (keV)	P_{γ}			
	1969Pe17*	1977Ku15 [†]	1982Sa36 [‡]	1984Ge07 [§]
74.38 (4)	-	4.0 (14)	-	-
84.373 (3)	1.21(6)	12100 (600)	1.9 (1)	100.0 (16)
131.612 (5)	-	1240 (60)	0.17 (2)	10.70 (15)
142.71 (11)	-	0.013 (4)	-	-
166.410 (6)	-	960 (50)	0.13 (1)	8.49 (12)
182.29 (10)	-	0.052 (18)	-	-
205.99 (4)	-	184 (9)	-	-
215.985 (4)	-	2390 (130)	0.30 (2)	1.61 (5)
228.42 (18)	-	0.18(3)	-	20.78 (25)
700.36 (7)	-	~ 0.03	-	-
741.87 (6)	-	0.014 (4)	-	-
831.97 (7)	-	0.14 (2)	-	-
908.28 (6)	-	-	-	-
992.65 (6)	-	~ 0.015	-	-

* Emission probability expressed in terms of photons per 100 disintegrations.

[†] Emission probabilities expressed in terms of photons per 10⁶ disintegrations.

[‡] Emission probabilities published relative to $P_{\gamma}(238.63 \text{ keV})$ for ²¹²Pb of 43.0 %.

[§] Emission probabilities published relative to $P_{\gamma}(84.373 \text{ keV})$ of 100.0 %.

Measured and recommended gamma-ray emission probabilities relative to $P_\gamma(84.373 \text{ keV})$ of 100 %.

E_γ (keV)	P_γ^{rel}			
	1977Ku15	1982Sa36	1984Ge07	Recommended value*
74.38 (4)	0.033 (12)	-	-	0.033 (12)
84.373 (3)	100 (5)	100 (5)	100.0 (16)	100.0 (16)
131.612 (5)	10.25 (50)	8.9 (10) [†]	10.70 (15)	10.7 (2)
142.71 (11)	0.000 11 (3)	-	-	0.000 11 (3)
166.410 (6)	7.9 (4)	6.8 (5) [†]	8.49 (12)	8.44 (12)
182.29 (10)	0.000 43 (15)	-	-	0.000 43 (15)
205.99 (4)	1.52 (7)	-	1.61 (5)	1.58 (4)
215.985 (4)	19.8 (11)	15.8 (11) [†]	20.78 (25)	20.7 (3)
228.42 (18)	0.001 5 (3)	-	-	0.001 5 (3)
700.36 (7)	~ 0.000 25	-	-	0.000 25 (8)
741.87 (6)	0.000 12 (3)	-	-	0.000 12 (3)
831.97 (7)	0.001 2 (2)	-	-	0.001 2 (2)
908.28 (6)	-	-	-	0.000 14 (4)
992.65 (6)	~ 0.000 12	-	-	0.000 12 (3)

* Weighted mean values adopted when judged appropriate.

[†] Significantly lower than equivalent data of 1977Ku15 and 1984Ge07 by 20 % to 30 %; judged to be an outlier, and therefore not considered in any weighted mean analysis.

Multipolarities and Internal Conversion Coefficients

The nuclear level scheme specified by Artna-Cohen has been used to define the multipolarities of the gamma transitions on the basis of known spins and polarities (1997Ar05). Limited studies of the internal conversion coefficients support the proposed transition types: E2 for both the 84.373- and 166.410-keV gamma rays (1953As31, 1966Co40, 1968Du06, 1969Pe17 and 1970SpZW).

Internal conversion coefficients as determined by measurement.

Reference	E_γ (keV)	α					Measurement technique
		α_L	α_{L2}	α_{L3}	α_{M+}	α_{total}	
1953As31	84.373 (3)	-	-	-	-	16	deduced from measured P_γ and P_α populating 84.37-keV nuclear level of ²²⁴ Ra
	166.410 (4)	-	-	-	-	1.2	deduced from measured P_γ and P_α populating 251-keV nuclear level of ²²⁴ Ra
1966Co40	84.373 (3)	14 (3)	7.6	6.3	3.8 (9)	18 (4)	P_{ce} measured by means of photographic emulsion technique
1968Du06	84.373 (3)	-	-	-	-	19.6 (14)	deduced from α -gated γ -ray spectra
1969Pe17	84.373 (3)	-	-	-	-	21.4 (9)	deduced from α -gated γ -ray spectra

Conversion electron spectra: Measurements of L- and M-subshell internal conversion ratios (1970SpZW).

E_γ (keV)	L_1/L_2	L_1/L_3	L_2/L_3	M_1/M_2	M_1/M_3	M_2/M_3
84.373 (3)	0.0388 (19)	0.0519 (21)	1.343 (10)	0.0471 (46)	0.0571 (57)	1.2187 (85)

The 908.28-keV gamma ray was identified as the only mixed multipolarity (M1 + E2), and was arbitrarily assigned a mixing ratio of 1.0 with an uncertainty of ± 0.2 . Recommended internal conversion coefficients have been determined from the frozen orbital approximation of Kibédi *et al.* (2008Ki07), based on the theoretical model of Band *et al.* (2002Ba85, 2002Ra45). Uncertainties of ± 1.5 % were adopted for all of the E1 and E2 gamma transitions (with minor upward adjustments associated with the significant figures for α_L and α_{M+}).

Gamma-ray emissions: multipolarities and theoretical internal conversion coefficients (frozen orbital approximation).

E_γ (keV)	Multipolarity	α_K	α_L	α_{M+}	α_{total}
74.38 (4)	[E2]	–	28.3 (4)	10.3	38.6 (6)
84.373 (3)	E2	–	15.57 (22)	5.63	21.2 (3)
131.612 (5)	E1	0.194 (3)	0.0406 (6)	0.0124	0.247 (4)
142.71 (11)	[E2]	0.279 (4)	1.368 (20)	0.493	2.14 (3)
166.410 (6)	E2	0.225 (4)	0.691 (10)	0.248	1.164 (17)
182.29 (10)	[E1]	0.089 4 (13)	0.017 57 (25)	0.005 63	0.112 6 (16)
205.99 (4)	(E1)	0.067 1 (10)	0.012 92 (18)	0.004 08	0.084 1 (12)
215.985 (4)	E1	0.060 0 (9)	0.011 48 (16)	0.003 72	0.075 2 (11)
228.42 (18)	[E2]	0.124 4 (18)	0.178 (3)	0.063 6	0.366 (6)
700.36 (7)	E1	0.005 02 (7)	0.000 834 (12)	0.000 256	0.006 11 (9)
741.87 (6)	[E2]	0.011 96 (17)	0.003 22 (5)	0.001 07	0.016 25 (23)
831.97 (7)	E2	0.009 70 (14)	0.002 40 (4)	0.000 79	0.012 89 (18)
908.28 (6)	[50%M1 + 50%E2] $\delta = 1.0$ (2)	0.019 0 (24)	0.003 6 (4)	0.001 4	0.024 (3)
992.65 (6)	[E2]	0.007 05 (10)	0.001 569 (22)	0.000 511	0.009 13 (13)

The normalisation factor was calculated for the gamma-ray emission probabilities by averaging the values determined by three different routes:

(i) direct population of the ^{224}Ra ground state

$$[\sum P_{\gamma_i} (1 + \alpha_i) \text{ to ground state}] \text{NF} + 0.734 (5) = 1.00$$

$$\text{NF} = 0.000 119 (3)$$

(ii) population/depopulation of the 84.373-keV nuclear level of ^{224}Ra

$$[P_\gamma(84.373 \text{ keV})(1 + \alpha(84.373 \text{ keV})) - \sum P_{\gamma_i} (1 + \alpha_i) \text{ to 84.373-keV level}] \text{NF} = 0.260 (5)$$

$$\text{NF} = 0.000 119 (3)$$

(iii) all α emissions

$$\sum P_\alpha \text{NF} = 1.00, \text{ and adopting } \alpha\text{-particle emission probability to } ^{224}\text{Ra ground state of } 0.734 (5)$$

(see section on alpha-particle emissions)

$$\text{NF} = 0.000 119 (3)$$

Thus, a normalization factor of 0.000 119 (3) has been adopted in the determination of the absolute gamma-ray emission probabilities.

Atomic Data

The x-ray and Auger-electron data have been calculated using the evaluated gamma-ray data, and atomic data from 1996Sc06, 1998ScZM and 1999ScZX. Both the x-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.01, 28 January 2003). This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

K and L X-ray emission probabilities per 100 disintegrations of ²²⁸Th.

			Energy (keV)	Photons per 100 disint.
XL	(Ra)		10.622 – 18.412	8.6 (4)
	XL ₁	(Ra)	10.622	0.166 (6)
	XL _α	(Ra)	12.196 – 12.339	2.86 (9)
	XL _η	(Ra)	13.662	0.109 (4)
	XL _β	(Ra)	14.236 – 15.447	4.67 (15)
	XL _γ	(Ra)	17.848 – 18.412	1.09 (4)
XK _α	XK _{α2}	(Ra)	85.43	0.018 0 (3)
	XK _{α1}	(Ra)	88.47	0.029 5 (5)
XK' _{β1}	XK _{β3}	(Ra)	99.432)
	XK _{β1} '	(Ra)	100.13) 0.010 34 (21)
	XK _{β5}	(Ra)	100.738)
XK' _{β2}	XK _{β2}	(Ra)	102.89)
	XK _{β4}	(Ra)	103.295) 0.003 39 (9)
	XKO _{2,3}	(Ra)	103.74)

Electron energies were determined from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

Data Consistency

A Q_{α} -value of 5520.08 (22) keV has been adopted from the atomic mass evaluation of Audi *et al.* (2003Au03) while in the course of formulating the decay scheme of ²²⁸Th. This value has subsequently been compared with the Q-value calculated by summing the contributions of the individual emissions to the ²²⁸Th alpha-decay process (i.e. α , electron, γ , etc.):

$$\text{calculated Q-value} = \sum (E_i \times P_i) = 5523 (40) \text{ keV}$$

Percentage deviation from the Q-value of Audi *et al.* is $-(0.1 \pm 0.7) \%$, which supports the derivation of a highly consistent decay scheme with a significant variant.

References

- 1918Me01 L. Meitner, Die Lebensdauer von Radiothor, Mesothor und Thorium, Physik. Zeitschr. 19 (1918) 257-263. [Half-life]
- 1953As31 F. Asaro, F. Stephens, Jr., I. Perlman, Complex Alpha Spectra of Radiothorium (Th²²⁸) and Thorium-X (Ra²²⁴), Phys. Rev. 92 (1953) 1495-1500. [E_α, P_α, P_γ, ICC]
- 1956Ki16 H.W. Kirby, G.R. Grove, D.L. Timma, Neutron-capture Cross Section of Actinium-227, Phys. Rev. 102 (1956) 1140-1141. [Half-life]
- 1962Ma57 C.W. Mays, D.R. Atherton, R.D. Lloyd, D.O. Clark, The Half-period of Th²²⁸ (RdTh), AEC Chicago Operations Office Report COO-225 (1962) 90-91 [Half-life]
- 1966Co40 M.O. Costa, M.R.S. Grade, Spectre d'Électrons de Conversion Interne Associés à la Transmutation du Thorium-228 en Radium-224, Port. Phys. 4 (1966) 267-279. [P_{ce}, ICC]

- 1968Da21 J. Dalmaso, C. Marsol, Sur le Rayonnement γ de Basse Énergie du Thorium 228 et de ses Dérivés et le Caractère Multipolaire des Transitions de ^{212}Po , C.R. Acad. Sci. (Paris) 267B (1968) 1366-1369. [E $_{\gamma}$]
- 1968Du06 C.L. Duke, W.L. Talbert, Jr., Total Internal-conversion Coefficients for Low-energy E2 Transitions in Ra 224 , Th 228 , U 234 , U 236 and Pu 240 , Phys. Rev. 173 (1968) 1125-1132. [ICC]
- 1969Pe17 A. Peghaire, Mesures Précises d'Intensités Absolues de Rayonnements γ pour des Emetteurs α , Nucl. Instrum. Methods 75 (1969) 66-70. [P $_{\alpha}$, P $_{\gamma}$, ICC]
- 1970Ba20 S.A. Baranov, V.M. Shatinskii, V.M. Kulakov, Yu.F. Rodionov, Investigation of the α Decay of the Two Isotopes Th 228 and Th 229 , Sov. J. Nucl. Phys. 11 (1970) 515-518. [E $_{\alpha}$, P $_{\alpha}$]
- 1970SpZW D.L. Spenny, A.A. Bartlett, Beta Ray Spectrometer Studies of E2 Conversion Electron Subshell Ratios, AEC Chicago Operations Office Report COO-535-620 (1970) 102-105. [L- and M-subshell ratios]
- 1971Gr07 B. Grennberg, A. Rytz, Absolute Measurements of α -ray Energies, Metrologia 7 (1971) 65-77. [E $_{\alpha}$]
- 1971Jo14 K.C. Jordan, G.W. Otto, R.P. Ratay, Calorimetric Determination of the Half-lives of ^{228}Th and ^{224}Ra , J. Inorg. Nucl. Chem. 33 (1971) 1215-1219. [Half-life]
- 1976BaZZ S.A. Baranov, A.G. Zelenkov, V.M. Kulakov, The Experimental Investigation of the Alpha Decay of Transactinium Isotopes, IAEA-186, Vol III (1976) 249-263, IAEA Vienna. [E $_{\alpha}$, P $_{\alpha}$]
- 1977Ku15 W. Kurcewicz, N. Kaffrell, N. Trautmann, A. Plochocki, J. Zylicz, M. Matul, K. Stryczniewicz, Collective States Fed by Weak α -transitions in the ^{232}U Chain, Nucl. Phys. A289 (1977) 1-14. [E $_{\gamma}$, P $_{\gamma}$]
- 1977Ku25 W. Kurcewicz, E. Ruchowska, N. Kaffrell, N. Trautmann, Precise Energies of Gamma Rays from the ^{230}Th and ^{228}Th Decay, Nucl. Instrum. Methods, 146 (1977) 613-614. [E $_{\gamma}$]
- 1977La19 F.P. Larkins, Semiempirical Auger-electron Energies for Elements $10 \leq Z \leq 100$, At. Data Nucl. Data Tables 20 (1977) 311-387. [Auger-electron energies]
- 1981Ku02 W. Kurcewicz, E. Ruchowska, N. Kaffrell, T. Björnstad, G. Nyman, Collective Excitations in the Transitional Nuclei $^{224,226}\text{Ra}$, Nucl. Phys. A356 (1981) 15-25. [908.28 keV P $_{\gamma}$]
- 1982Sa36 S. Sadasivan, V. M. Raghunath, Intensities of Gamma Rays in the ^{232}Th Decay Chain, Nucl. Instrum. Methods 196 (1982) 561-563. [P $_{\gamma}$]
- 1984Ge07 R.J. Gehrke, V.J. Novick, J.D. Baker, γ -ray Emission Probabilities for the ^{232}U Decay Chain, Int. J. Appl. Radiat. Isot. 35 (1984) 581-589. [P $_{\gamma}$]
- 1991Ry01 A. Rytz, Recommended Energy and Intensity Values of Alpha Particles from Radioactive Decay, At. Data Nucl. Data Tables 47 (1991) 205-239. [E $_{\alpha}$, P $_{\alpha}$]
- 1992Un01 M.P. Unterweger, D.D. Hoppes, F.J. Schima, New and Revised Half-life Measurements Results, Nucl. Instrum. Methods Phys. Res. A312 (1992) 349-352. [Half-life]
- 1993Ba72 T. Babeliowsky, G. Bortels, ALFA: A Program for Accurate Analysis of Complex Alpha-particle Spectra on a PC, Appl. Radiat. Isot. 44 (1993) 1349-1358. [P $_{\alpha}$]

- 1993Bo20 R. Bonetti, C. Chiesa, A. Guglielmetti, C. Migliorino, A. Cesana, M. Terrani, Discovery of Oxygen Radioactivity of Atomic Nuclei, Nucl. Phys. A556 (1993) 115-122. [Cluster decay]
- 1995Ar33 G. Ardisson, M. Hussonnois, Radiochemical Investigations of Cluster Radioactivities, Radiochim. Acta 70/71 (1995) 123-133. [Cluster decay]
- 1996Sc06 E. Schönfeld, H. Janßen, Evaluation of Atomic Shell Data, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527-533. [X_K , X_L , Auger electrons]
- 1997Ar05 A. Artna-Cohen, Nuclear Data Sheets for $A = 224$, Nucl. Data Sheets 80 (1997) 227-262. [Nuclear structure, level energies]
- 1997Tr17 S.P. Tretyakova, V.L. Mikheev, Experimental Investigation of the Cluster Radioactivity of Atomic Nuclei, Nuovo Cimento 110A (1997) 1043-1048. [Cluster decay]
- 1998Ak04 Y.A. Akovali, Review of Alpha-decay Data from Doubly-even Nuclei, Nucl. Data Sheets 84 (1998) 1-114. [alpha decay, r_0 parameters]
- 1998ScZM E. Schönfeld, G. Rodloff, Tables of the Energies of K-Auger Electrons for Elements with Atomic Numbers in the Range from $Z = 11$ to $Z = 100$, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX E. Schönfeld, G. Rodloff, Energies and Relative Emission Probabilities of K X-rays for Elements with Atomic Numbers in the Range from $Z = 5$ to $Z = 100$, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2002Ba85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, Dirac-Fock Internal Conversion Coefficients, At. Data Nucl. Data Tables 81 (2002) 1-334. [ICC]
- 2002Ra45 S. Raman, C.W. Nestor, Jr., A. Ichihara, M.B. Trzhaskovskaya, How Good are the Internal Conversion Coefficients Now? Phys. Rev. C66 (2002) 044312, 1-23. [ICC]
- 2002Un02 M.P. Unterweger, Half-life Measurements at the National Institute of Standards and Technology, Appl. Radiat. Isot. 56 (2002) 125-130. [Half-life]
- 2003Au03 G. Audi, A.H. Wapstra, C. Thibault, The AME2003 Atomic Mass Evaluation (II). Tables, Graphs and References, Nucl. Phys. A729 (2003) 337-676. [Q-value]
- 2007BeZP M.M. Bé, V.P. Chechev, R. Dersch, O.A.M. Helene, R.G. Helmer, M. Herman, S. Hlaváč, A. Marcinkowski, G.L. Molnár, A.L. Nichols, E. Schönfeld, V.R. Vanin, M.J. Woods, Half-life evaluations by M. J. Woods for "Update of X-ray and Gamma-ray Decay Data Standards for Detector Calibration and Other Applications", IAEA Scientific and Technical Report STI/PUB/1287, Volumes 1 and 2, May 2007, International Atomic Energy Agency, Vienna, Austria, ISBN 92-0-113606-4. [Half-life evaluation]
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr., Evaluation of Theoretical Conversion Coefficients using BrIcc, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202-229. [ICC]

**²³¹Th -Comments on evaluation of decay data
by Huang Xiaolong , Wang Baosong**

This evaluation was completed in 2007. Literature available by May 2007 was included.

1 Decay Scheme

²³¹Th disintegrates 100 % by β^- emission to levels in ²³¹Pa.

²³¹Th ground state has $J^\pi = 5/2^+$ (2001Br31).

The adopted $Q(\beta^-)$ value of 391.6 (15) keV from Audi(2003Au03) is good in agreement with the $Q(\beta^-)$ value of 372 (59) keV, calculated by the evaluator (using program RADLST) from average radiation energies and decay scheme data.

2 Nuclear Data

The Q value is from the 2003Au03 evaluation.

Level energies, spin and parities are from 2001Br31.

Measured and evaluated ²³¹Th half-life values are listed in Table 1.

Table 1: Measured half-life values of ²³¹Th and recommended value.

$T_{1/2}$ (h)	References	Measurement method
25.51 (23)	1949Kn09	Geiger counters, weighted average of 5 samples, 10 $T_{1/2}$
25.64 (10)	1951Ja17	G-M tube, unweighted average of 2 samples, 6 $T_{1/2}$
25.52 (1)	1958Ca19	$4\pi\beta$ counter, unweighted average of 18 sources, 4 $T_{1/2}$
25.7 (2)	1971Ko48	Ge(Li), γ -rays
25.76 (21)	1983Ch06	Ge(Li), 84keV γ -ray, 6 $T_{1/2}$
25.63 (5)		Unweighted mean
25.522 (10)		Weighted mean with all experimental values, $\chi^2=0.88$
25.522 (10)	Recommended value	Weighted mean

The weighted half-life average has been calculated using the LWM program.

2.1 β^- transitions

The maximum energies of the β^- transitions in the decay of ²³¹Th have been deduced from the Q value (2003Au03) and the level energies.

The adopted β^- transition probabilities and their associated uncertainties were deduced from the γ transition probability balance at each level of the decay scheme, using a normalization factor $N = 0.0670$ (7) (See Section 5.3). The $I_{\beta^-}(\text{g.s.} + 9.2 \text{ keV}) = 0.022$ (7) and $I_{\beta^-}(58.6 \text{ keV}) < 0.33$ are the experimental values from a β^- Kurie plot (1975Ho14). Measured and recommended β^- transition probabilities are given in Table 2.

Table 2: Measured and recommended β^- transition probabilities (%).

Level energy/keV	1975Ho14	Adopted value
0	0.022 (7)	0.022 (7)
58.6	< 0.33	< 0.33
77.7	< 0.33	0.43 (2)
84.2		29 (18)
101.4		41 (16)
102.3		13 (8)
174.2		1.36 (24)
183.5		12.2 (15)
218.2		0.31 (23)
247.3		2.7 (4)
318		0.00078 (5)
320.2		0.066 (2)
351.8		0.0032 (2)

The values of $lg ft$ and average β^- energies have been calculated with the program LOGFT.

2.2 γ -Ray Transitions

The γ -ray transition probabilities were calculated using the γ -ray emission intensities and the relevant internal conversion coefficients.

Multipolarities and mixing ratios of γ -ray transitions are from 1975Ho14 and 2001Br31.

The internal conversion coefficients (ICC) and their associated uncertainties for γ -ray transitions have been obtained using the BrIcc computer program (2008Ki07), which uses the ‘‘Frozen Orbital’’ approximation (2002Ba85). The mixing ratios of the 18- and 63-keV gamma transitions have asymmetric uncertainties, $\delta = 0.14 (+12, -4)$ and $0.52 (+20, -32)$, respectively. The ICC of the 84.214 keV γ -ray has been taken from a measurement of 1975Ho14 because it has an anomalous conversion coefficient. Experimental and theoretical conversion coefficients are compared in Table 3.

Table 3: Comparison of theoretical and measured conversion coefficients.

E_γ (keV)	Multipolarity	α (theory)	α (exp.)	
			1960As02	1975Ho14
18.07	M1+E2	$\alpha_T = 757$		$\alpha_{M3} > 9$
25.64	E1	$\alpha_T = 4.37, \alpha_L = 3.26, \alpha_M = 0.84$	$\alpha_T = 4.8 (10)$	$\alpha_{L3} = 1.6 (3), \alpha_M = 0.96 (9)$
58.57	E2	$\alpha_T = 155.5, \alpha_L = 113.6, \alpha_M = 31.3$		$\alpha_L = 115.9, \alpha_M = 29.9 (30)$
63.86	M1+E2	$\alpha_T = 34, \alpha_L = 25, \alpha_M = 7$		$\alpha_{L1} = 9.1 (16)$
68.5	E2	$\alpha_T = 73.3, \alpha_L = 53.5, \alpha_M = 14.8$		$\alpha_L = 57 (11)$
81.228	M1(+E2)	$\alpha_T = 8.1, \alpha_L = 6.1, \alpha_M = 1.5$		$\alpha_{L1} = 4.7 (8), \alpha_M = 1.3 (3)$
82.087	M1(+E2)	$\alpha_T = 7.9, \alpha_L = 5.9, \alpha_M = 1.4$		$\alpha_{L1+L3} = 5.7 (11), \alpha_M = 1.6 (4)$
84.214	E1	$\alpha_T = 0.19, \alpha_L = 0.14$	$\alpha_T = 2.8 (4)$	$\alpha_T = 2.50 (25), \alpha_M = 0.57 (10)$
99.278	M1+E2	$\alpha_T = 6, \alpha_L = 4.4, \alpha_M = 1.1$		$\alpha_M = 1.13 (14), \alpha_N = 0.35 (10)$
135.664	M1(+E2)	$\alpha_T = 8, \alpha_K = 6.1, \alpha_L = 1.4$		$\alpha_K = 6.5 (11), \alpha_L = 1.1 (3)$
145.94	M1+E2	$\alpha_T = 5.1, \alpha_K = 3.4, \alpha_L = 1.3$		$\alpha_K = 3.6 (8), \alpha_L = 0.8 (3)$
163.101	M1(+E2)	$\alpha_T = 4.9, \alpha_K = 3.9, \alpha_L = 0.78$		$\alpha_K = 4.1 (5), \alpha_L = 0.6 (1)$
217.94	E1	$\alpha_T = 0.079, \alpha_K = 0.062, \alpha_L = 0.01$		$\alpha_K < 0.12, \alpha_L < 0.09$
311	M1+E2	$\alpha_T = 0.6, \alpha_K = 0.5, \alpha_L = 0.1$		$\alpha_L = 0.11 (3), \alpha_M = 0.04 (1)$

3. Atomic data

Atomic fluorescence yields ($\omega_K, \omega_L, \omega_M, \eta_{KL}$ and η_{LM}) are from Schönfeld (1996Sc06).

The X-ray and Auger electron emission probabilities have been deduced from γ -ray and conversion electron data by using the computer code RADLST. Measured and calculated X-ray emission probabilities are compared in Table 4.

Table 4: Comparison of the calculated and measured X-ray emission probabilities.

	1973Br12	1999Ch12	Recommended (deduced)
$K_{\alpha 1}$	0.69 (8)	0.64 (4)	0.59 (7)
$K_{\alpha 2}$	0.40 (5)	0.376 (24)	0.37 (4)
K_{β}	0.332 (25)	0.310 (14)	0.28 (3)

The deduced KX-ray emission probabilities agree with the measured values of 1999Ch12 and 1973Br12, thus confirming the completeness of the decay scheme.

4. Electron Emissions.

The conversion electron emission probabilities have been deduced from γ -ray transition data using theoretical internal conversion coefficients.

5. Photon Emissions

5.1 γ -ray energy values

Measurements of γ -ray energy values from ²³¹Th β^- decay are listed in Table 5. The recommended values are taken from the measurements of 1975Ho14 and 1979Bo30, except as noted in Table 5.

It should be noticed that some uncertain weak γ -rays: 26.55, 29.30, 32.73, 33.32, 38.90, 41.55, 42.22, 45.34, 85.80, 97.55, 106.85, 173.0, 224.1 and 237.8 keV, were observed only by 1977Ba72. These γ -rays have not been considered in the present evaluation.

Table 5: Measured and recommended γ -ray energies for ²³¹Th.

1973Br12	1973Te06	1975Ho14	1977Ba72	1979Bo30	Recommended
					9.2 ^a
					10.25 ^a
		17.2	17.21		17.2
		18.07	18.05		18.07
(25.65)		25.64 (2)	25.64 (5)		25.64 (2)
42.80 (6)		42.86 (7)	42.22 (5)		42.86 (7)
44.1 (3)		44.08 (17)	45.34 (5)		44.08 (17)
58.47 (5)		58.57 (2)	58.54 (5)	58.5700 (24)	58.5700 (24) ^b
63.7 (2)		63.86 (3)	63.65 (5)		63.86 (3)
		68.5 (1)	68.55		68.5 (1)
72.66 (6)	72.74 (5)	72.78 (2)	72.70 (5)	72.7510 (25)	72.7510 (25) ^b
			76		77.69 ^c
81.18 (5)	81.20 (6)	81.24 (2)	81.16 (5)	81.2280 (14)	81.2280 (14) ^b
82.02 (6)	82.06 (7)	82.11 (2)	82.02 (5)	82.0870 (14)	82.0870 (14) ^b
(84.17)	84.20	84.21 (2)	84.16 (5)	84.2140 (13)	84.2140 (13) ^b
89.94 (5)	89.95 (4)	89.95 (2)	89.94 (5)		89.95 (2)

1973Br12	1973Te06	1975Ho14	1977Ba72	1979Bo30	Recommended
93.0 (1)	92.91 (10)	93.02 (4)			93.02 (4)
99.30 (5)	99.33 (5)	99.28 (2)	99.33 (5)	99.278 (3)	99.278 (3) ^b
102.30 (5)	102.32 (4)	102.27 (2)	102.23 (5)	102.2700 (13)	102.2700 (13) ^b
105.73 (10)	105.74 (10)	105.81 (3)			105.81 (3)
106.58 (10)	106.66 (8)	106.61 (3)	106.65 (10)		106.61 (3)
115.5 (2)		115.63 (3)	115.83 (10)		115.63 (3)
116.91 (5)		116.82 (2)	116.80 (10)		116.82 (2)
125.10 (5)		124.93 (2)	125.00 (10)	124.914 (17)	124.914 (17) ^b
134.14 (8)		134.03 (2)	134.00 (5)		134.03 (2)
135.77 (6)		135.68 (2)	135.66 (5)	135.664 (11)	135.664 (11) ^b
136.78 (20)		136.75 (7)			136.75 (7)
		140.54 (4)			140.54 (4)
145.15 (30)		145.06 (4)			145.06 (4)
146.00 (7)		145.94 (2)	145.90 (5)		145.94 (2)
163.16 (6)		163.12 (2)	163.15 (5)	163.101 (4)	163.101 (4) ^b
164.94 (10)		165.00 (5)	164.70 (10)		165.00 (5)
169.58 (10)		169.66 (3)			169.66 (3)
174.19 (8)		174.15 (2)	174.1 (10)		174.15 (2)
					177.66
183.47 (7)		183.50 (2)	183.4 (10)	183.480 (25)	183.480 (25) ^b
188.77 (20)		188.76 (2)	188.7 (10)		188.76 (2)
218.00 (7)		217.94 (3)	218.0 (5)		217.94 (3)
236.17 (7)		236.01 (3)	236.1 (10)		236.01 (3)
240.4 (2)		240.27 (5)			240.27 (5)
242.6 (1)		242.50 (4)			242.50 (4)
249.8 (3)		249.60 (7)	249.8		249.60 (7)
250.5 (3)		250.45 (7)			250.45 (7)
267.80 (7)		267.62 (8)	267.8		267.62 (8)
		274.10 (10)			274.10 (10)
308.9 (3)		308.78 (7)			308.78 (7)
311.0 (1)		311.00 (5)	312.3 (25)		311.00 (5)
318.0 (4)		317.87 (8)			317.87 (8)
320.2 (3)		320.15 (8)			320.15 (8)
		351.80 (10)			351.80 (10)

a: Expected but as yet unobserved.

b: From 1979Bo30 curved crystal.

c: From 1999Ch12.

5.2 Relative γ -ray intensities

Experimental γ -ray intensities from ²³¹Th β^- decay are listed in Table 6. The recommended values are from a LWM average of values reported in 1999Ch12, 1983BaZZ, 1975Ho14, 1973Te06 and 1973Br12.

1977Ba72 observed some uncertain weak γ -rays with measured relative γ -ray intensities different from those given in other measurements. These relative intensities may not be accurate and thus have not been considered here.

Table 6: Measured and evaluated relative γ -ray intensities for ²³¹Th.

E_γ (keV)	I_γ									Evaluation
	1953 Fr37	1971 Ko48	1973 Br12	1973 Te06	1975 Ho14	1977 Ba72	1983 BaZZ	1999 Ch12	LWM	
(9.2)										7.44 ^a
(10.25)										11.0 ^a
17.2										680 (230) ^b
18.07					≤ 5.1					310 (110) ^b
25.64	170	119 (25)	202 (20)		228 (15)	331.92 (56)	230 (16)	210 (10)	217 (7)	207 (10) ^c
42.86			0.87 (10)		0.89 (6)	0.469 (19)		0.89 (2)	0.89 (2)	0.89 (2)
44.08			0.06 (4)		0.011 (3)	0.527 (20)			0.011 (3)	0.011 (3)
58.5700		8.4 (6)	7.2 (7)		7.4 (3)	8.748 (82)	6.8 (6)	6.8 (2)	6.98 (16)	7.17 (22) ^c
63.86	< 40		0.68 (14)		0.35 (3)			0.29 (5)	0.35 (3)	0.35 (3)
68.5					0.088 (22)			0.088 (4)	0.088 (2)	0.088 (2)
72.7510		4.4 (4)	4.0 (4)	3.8 (2)	3.86 (23)	4.046 (59)	7.8 (8)	3.8 (1)	3.88 (24)	3.88 (24)
77.69								0.063 (10)	0.063 (10)	0.063 (10)
81.2280		1.03 (3)	14.2 (14)	13.5 (9)	13.7 (8)	11.69 (10)	13.2 (5)	13.5 (5)	13.5 (3)	13.5 (3)
82.0870		21.5 (13)	7.2 (7)	6.8 (4)	6.2 (5)	4.675 (67)	6.0 (3)	6.0 (3)	6.24 (17)	6.24 (17)
84.2140	100	100	100	100	100	100	100	100	100	100
89.95		13.9 (13)	15.3 (15)	15.3 (8)	14.5 (9)	13.25 (12)		15.0 (5)	15.0 (4)	15.0 (4)
93.02			0.50 (5)	0.9 (2)	0.69 (8)			0.71 (8)	0.60 (4)	0.60 (4)
99.278		1.03 (10)	2.1 (2)	2.2 (2)	1.85 (11)	1.555 (43)		2.0 (1)	2.05 (8)	2.05 (8)
102.2700		4.6 (4)	6.7 (7)	6.8 (4)	6.3 (5)	5.424 (82)	6.5 (3)	6.6 (2)	6.58 (14)	6.58 (14)
105.81	6 (5)		0.14 (2)	0.13 (8)	0.11 (1)			0.12 (1)	0.118 (7)	0.118 (7)
106.61		3.04 (25)	0.34 (4)	0.33 (10)	0.262 (15)	0.482 (25)		0.264 (11)	0.267 (9)	0.267 (9)
115.63			0.04 (1)		0.015 (3)	0.267 (20)		0.015 (4)	0.0164 (23)	0.0164 (23)
116.82			0.39 (4)		0.318 (20)	0.367 (21)		0.34 (2)	0.336 (13)	0.336 (13)
124.914	2		0.95 (9)		0.86 (5)	1.014 (43)	0.89 (12)	0.88 (2)	0.88 (2)	0.88 (2)
134.03			0.42 (5)		0.37 (2)	0.562 (24)	0.29 (14)	0.38 (1)	0.38 (1)	0.38 (1)
135.664			1.3 (1)		1.20 (8)	1.704 (28)	1.30 (23)	1.17 (4)	1.19 (3)	1.19 (3)
136.75			0.09 (3)		0.065 (3)			0.067 (3)	0.066 (2)	0.066 (2)
140.54					0.011 (1)			0.011 (1)	0.011 (1)	0.011 (1)
145.06			0.12 (3)		0.089 (6)			0.084 (6)	0.087 (4)	0.087 (4)
145.94			0.58 (6)		0.49 (3)	0.571 (25)		0.47 (2)	0.484 (16)	0.484 (16)
163.101	1.8		2.6 (3)		2.38 (14)	2.754 (64)		2.30 (8)	2.33 (7)	2.33 (7)
165.00			0.06 (3)		0.060 (6)	0.200 (11)		0.051 (2)	0.052 (2)	0.052 (2)
169.66			0.03 (1)		0.0185 (15)			0.021 (1)	0.021 (1)	0.021 (1)
174.15			0.31 (3)		0.278 (17)	0.704 (21)		0.26 (1)	0.268 (8)	0.268 (8)
177.66 ^x								0.00095 (20)	0.00095 (20)	0.00095 (20)
183.480			0.57 (6)		0.506 (20)	1.005 (26)		0.49 (2)	0.50 (1)	0.50 (1)
188.76			0.08 (1)		0.049 (3)	0.084 (8)		0.049 (1)	0.050 (4)	0.050 (4)
217.94	0.3		0.67 (7)		0.62 (5)	0.960 (29)	0.57 (2)	0.60 (1)	0.60 (1)	0.60 (1)
236.01	0.1		0.18 (2)		0.14 (1)	1.465 (28)		0.138 (5)	0.140 (4)	0.140 (4)
240.27			0.0050 (5)		0.0043 (5)			0.0040 (5)	0.0043 (6)	0.0043 (6)

E_γ (keV)	I_γ									
	1953 Fr37	1971 Ko48	1973 Br12	1973 Te06	1975 Ho14	1977 Ba72	1983 BaZZ	1999 Ch12	LWM	Evaluation
242.50			0.0130 (6)		0.013 (1)			0.011 (1)	0.0123 (6)	0.0123 (6)
249.60			0.010 (2)		0.012 (1)			0.012 (1)	0.012 (1)	0.012 (1)
250.45			0.011 (2)		0.010 (1)			0.010 (1)	0.010 (1)	0.010 (1)
267.62			0.0230 (6)		0.018 (2)			0.019 (1)	0.021 (2)	0.021 (2)
274.10					0.00046 (15)			0.0006 (2)	0.0005 (2)	0.0005 (2)
308.78			0.008 (1)		0.0060 (6)			0.0053 (2)	0.0054 (2)	0.0054 (2)
311.00			0.054 (5)		0.045 (3)			0.046 (2)	0.047 (2)	0.047 (2)
317.87			0.0020 (2)		0.00123 (15)			0.0013 (2)	0.0015 (2)	0.0015 (2)
320.15			0.0035 (3)		0.0017 (2)			0.0020 (2)	0.0022 (4)	0.0022 (4)
351.80					0.0011 (2)			0.0010 (2)	0.0010 (2)	0.0010 (2)

a: $I(\gamma+ce)$, from γ -ray transition intensity balance.

b: $I(\gamma+ce)$, from ce measurements(1975Ho14).

c: Adjusted value from intensity balance.

x: Not placed in level scheme.

5.3 Absolute values γ -ray emission probabilities

Measurements of the absolute emission probability of the 84.21keV γ -ray from ²³¹Th β^- decay and the LWM results are listed in Table 7. The measurement of 1973Br12 is an average of two α - γ coincidence measurements (6.7 (5) and 7.3 (4)). This value and the measurement of 1960As02 are higher than other measurements and not adopted in the calculation.

The recommended absolute γ -ray emission probability of the 84.21keV γ -ray is from the LWM calculation, and has been used here to produce a recommended normalization factor $N = 0.0670$ (7).

Table 7: Measured and recommended absolute γ -ray emission probability of 84.21keV for ²³¹Th.

P_γ (84.21 keV) (%)	References	measurement method
7.2 (1)	1960As02	Not used
7.9 (5)	1971Ko48	Ge(Li). Replaced by 1999Ch12
7.0 (3)	1973Br12	Ge(Li). Not used
6.5 (4)	1975Ho14	Ge(Li)
6.6 (3)	1982Va04	Si(Li). Weighted average of 3 sources
6.52 (13)	1983BaZZ	
7.25 (41)	1983Ch06	Ge(Li). Replaced by 1999Ch12
6.84 (10)	1984He12	Ge detector. Weighted average of 5 measurements
6.60 (25)	1999Ch12	LEPS. Secular equilibrium with ²³⁵ U
6.71 (10)	1986LoZT	CRP evaluation in 1986
6.89 (31)		LWM of all measurements
6.70 (7)		LWM, $\chi^2=1.1$
6.70 (7)		Recommended value

The recommended absolute γ -ray emission probabilities are the relative values evaluated in Table 6 multiplied by 0.0670 (7).

6. References

- 1949Kn09 G.B. Knight, R.L. Macklin, Phys. Rev. 75, 34 (1949) [$T_{1/2}$].
- 1951Ja17 A. Jaffey, J. Lerner, S. Warshaw, Phys. Rev. 82, 498 (1951) [$T_{1/2}$].
- 1953Fr37 M.S. Freedman, A.H. Jaffey, F. Wagner, Jr., J. May, Phys. Rev.89, 302 (1953) [I_γ].
- 1958Ca19 M.J. Cabell, Can. J. Phys. 36, 989 (1958) [$T_{1/2}$].
- 1960As02 F. Asaro, F.S. Stephens, J.M. Hollander, I. Perlman, Phys. Rev. 117, 492 (1960) [P_γ].
- 1971Ko48 K. Kobayashi, T. Hashimoto, I. Kimura, J. Nucl. Sci. Technol. 8, 492 (1971) [$T_{1/2}, E_\gamma, I_\gamma, P_\gamma$].
- 1973Br12 E. Browne, F. Asaro, Phys. Rev. C7, 2545 (1973) [$E_\gamma, I_\gamma, P_\gamma$].
- 1973Te06 W. Teoh, Nucl. Instrum. Methods 109, 509 (1973) [E_γ, I_γ].
- 1975Ho14 P. Hornshoj, P. Tidemand-Petersson, R. Kaczarowski, B. Kotlinska, J. Zylicz, Nucl. Phys. A248, 406 (1975) [$E_\gamma, I_\gamma, I(\text{ce}), I_\beta, \text{Multipolarity}$].
- 1977Ba72 S.A. Baranov, V.M. Shatinskii, A.G. Zelenkov, V.A. Pchelina, Sov. J. Nucl. Phys. 26, 486 (1977) [E_γ, I_γ].
- 1979Bo30 H.G. Borner, G. Barreau, W.F. Davidson, P. Jeuch, T. von Egidy, J. Almeida, D.H. White, Nucl. Instrum. Methods 166, 251 (1979) [E_γ].
- 1982Va04 R. Vaninbrouckx, B. Denecke, Nucl. Instrum. Methods 193, 191 (1982) [P_γ].
- 1983BaZZ C. Baktash, E. der Mateosian, O.C. Kistner, A.W. Sunyar, D. Horn, C.J. Lister, Bull. Am. Phys. Soc. 28, No.1, 41, HE7 (1983) [I_γ, P_γ].
- 1983Ch06 H. Chatani, Nucl. Instrum. Methods 205, 501 (1983) [$T_{1/2}, P_\gamma$].
- 1984He12 R.G. Helmer, C.W. Reich, Int. J. Appl. Radiat. Isotop. 35, 783 (1984) [P_γ].
- 1996Sc06 E. Schönfeld, H. Janssen, Nucl. Instrum. Meth. Phys. Res. A369, 527(1996) [Atomic data].
- 1999Ch12 H. Chatani, Nucl. Instrum. Meth. Phys. Res. A425, 277 (1999) [$E_\gamma, I_\gamma, P_\gamma$].
- 2001Br31 E. Browne, Nucl. Data Sheets 93, 763 (2001) [Level energies, spin and parity].
- 2002Ba85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, At. Data Nucl. Data Tables 81(2002)1 [Calculated ICC]
- 2003Au03 G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A729,129(2003) [Q].
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr. , Nucl. Instrum. Meth. Phys. Res. A589, 202(2008) [Theoretical ICC].

²³¹Pa – Comments on evaluation of decay data

A. Arinc

Evaluation completed: February 2010
Literature cut-off date: June 2009

Evaluation procedure

Weighted mean analyses were applied to determine recommended values throughout the evaluation when the data were in statistical agreement. When the data were not in statistical agreement, the Limitation of Relative Statistical Weights (LRSW) was used. Uncertainties were expanded to match the minimum input uncertainty where appropriate.

1. Decay scheme

²³¹Pa disintegrates by alpha emission to various excited levels and the ground state of ²²⁷Ac. The spin, parity, half-life of first excited state, multipolarities, mixing ratios and level energies of ²²⁷Ac are based on the mass-chain evaluation of Browne (2001Br31).

A lack of experimental data for low-energy gamma transitions and imprecise alpha spectrometry measurements has adversely affected the construction of the decay scheme. The strongest transition of the decay scheme $\gamma_{1,0}$ at 27.370 (10) keV has a transition probability with an uncertainty of 12 %. Further measurements are required in order to build a more reliable decay scheme.

2. Nuclear data

The $Q(\alpha)$ value of 5149.9 (8) keV is taken from the evaluation of Audi et al (2003Au03). The Q -value calculated with Saisinuc is 5100 (120) keV.

$$\begin{aligned} \% \text{ Deviation} &= [Q(\text{Audi } et \text{ al.}) - Q(\text{calculated}) / Q(\text{Audi } et \text{ al.})] \times 100 \\ &= [5149.9 (8) - 5100 (120) / 5149.9 (8)] \times 100 \\ &= [(49.9 \pm 120.0) / (5149.9 \pm 0.8)] \times 100 = (1.0 \pm 2.3) \% \end{aligned}$$

The experimental half-life values used for calculating the mean are given in Table 1. The half-life value of 32 000 (3 200) years from Van Grosse (1932Grosse) was omitted from the analysis due to its inaccuracy. The published values from 1969Ro33 and 1961Ki05 were adjusted by 2001Br31 to take into account the change in the adopted decay scheme.

The AveTool computer code was used to calculate the average using three statistical methods: Limitation of Relative Statistical Weights (LRSW), Normalised Residual Methods (NRM) and the Rajeval Technique (RT).

Table 1: Experimental half-life values of ²³¹Pa.

Reference	Half-life (years)	Comments
1949Va02	34 300 (300)	
1961Ki05	32 643 (260)	
1968Br04	32 340 (115)	
1969Ro33	32 765 (110)	
LRSW	32 670 (260)	
NRM	32 705 (93)	reduced- $\chi^2 = 2.40$
RT	32 718 (97)	reduced- $\chi^2 = 1.09$
Recommended value	32 670 (260)	

The data set is discrepant with a reduced- $\chi^2 = 12.84$ on the LRSW which is larger than the critical reduced- $\chi^2 = 3.78$ (99 % confidence level). Although the value from 1949Va02 is not in good agreement with the other three values it was not excluded by Chauvenet's criterion. The published uncertainty of 300 years of 1949Va02 was adjusted to 800 years by NRM and to 1300 years by RT, while the published uncertainty of 115 years of 1968Br04 was adjusted to 400 years by RT. The recommended value is the LRSW mean of 32 670 (260) years. This value was chosen as it includes the two most precise values.

Overall the half-life data set is unsatisfactory and there is a strong need for new half-life measurements.

3. Atomic data

The values of ω_K , ω_L , n_{KL} and relative probabilities of the X-ray and Auger emissions were derived from Schönfeld and Janßen (1996Sc06).

The energies and relative emission probabilities of the X-ray and Auger electrons have been calculated using the computer code EMISSION. A summary of the results is given in Tables 2 and 3. The calculated L X-ray and K X-ray subshell ratios were in good agreement with the published data from De Pinho (1974De11).

Table 2: Calculated L X-ray emission energies and probabilities.

L X-ray	Energy (keV)	Calculated value
Ll	10.87	1.10 (4)
L α	12.50 – 12.65	18.7 (7)
L η	14.08	0.303 (19)
L β	14.60 – 16.63	19.7 (7)
L γ	17.81 – 18.92	4.45 (16)
LX total		44.3 (13)

Table 3: Calculated K X-ray emission energies and probabilities.

K X-ray	Energy (keV)	Calculated value
K α_2	87.768	0.715 (23)
K α_1	90.885	1.16 (4)
K β_1'	102.10 – 103.46	0.410 (15)
K β_2'	105.68 – 106.56	0.136 (6)
KX total		2.42 (8)

4. Alpha particles

4.1 Alpha particle energies

The alpha transition energies have been calculated from the $Q(\alpha)$ value (2003Au03), and the level energies were adopted from Browne (2001Br31) and are given in Table 4.

Adopted alpha emission energies have been calculated from the transition energies taking into account the recoil energy of the daughter nucleus. The theoretically calculated values were compared to the published data where available (Table 5). The data from 1961Ba42, 1968Ba25 and 1976BaZZ are from the same main author (Baranov). Experimental alpha emission energies were taken from the compilation of 1991Ry01 when available; otherwise primarily from 1976BaZZ and then from 1961Ba42.

Alpha hindrance factors were calculated using the ALPHAD computer program. The radius parameter of $r_0(^{227}\text{Ac}) = 1.5323$ (14) was calculated as the average of $r_0(^{226}\text{Ra}) = 1.5331$ (13), $r_0(^{226}\text{Th}) = 1.531$ (5), $r_0(^{228}\text{Th}) = 1.5289$ (5) and $r_0(^{228}\text{Ra}) = 1.5361$ (22) from 1998Ak04. A summary of the adopted levels and theoretical and experimental alpha emission values is presented in Table 6.

Table 4: Adopted nuclear levels of ²²⁷Ac.

Nuclear level number	Nuclear level energy (keV)	Spin and parity	Half-life
0	0.0	3/2-	21.772 (3) a
1	27.37 (1)	3/2+	38.3 (3) ns
2	29.98 (1)	5/2-	
3	46.35 (1)	5/2+	
4	74.14 (1)	(7/2)-	
5	84.55 (1)	(7/2)+	
6	109.94 (2)	(9/2)+	
7	126.86 (2)	(9/2)-	
8	160 (2)		
9	187.32 (3)	(11/2+)	
10	198.71 (4)	(11/2-)	
11	210.78 (5)	(13/2+)	
12	271.29 (6)	(13/2-)	
13	273.14 (3)	(5/2)-	
14	304.73 (5)	(5/2+)	
15	330.04 (1)	3/2-	<70 ps
16	354.50 (4)	1/2-	
17	387.23 (2)	7/2-	

Nuclear level number	Nuclear level energy (keV)	Spin and parity	Half-life
18	425.59 (3)	5/2+	
19	435.19 (2)	(1/2)+	
20	437.96 (4)	(5/2-)	
21	469.24 (6)	(9/2+)	
22	501.28 (7)	(3/2-,5/2-)	
23	537.0 (1)	(3/2+)	
24	562.8 (1)	(3/2+,5/2+)	
25	656.4 (3)	(7/2+)	

Table 5: Experimental alpha emission energies (keV).

Transition	1961Ba42 ¹	1966Ba04	1968Ba25 ²	1976BaZZ	1991Ry01
$\alpha_{0,0}$	5058.9 (21)	5058.5 (15)	5057.5 (10)	5058.1 (10)	5058.6 (15)
$\alpha_{0,1}$	5032.0 (21)	-	5030.8	-	-
$\alpha_{0,2}$	5029.9 (21)	-	5028.3	-	5028.4 (10)
$\alpha_{0,3}$	5012.7 (20)	5013.5 (15)	5012.7	5013.3 (10)	5013.8 (14)
$\alpha_{0,4}$	4985.5 (20)	-	4985.8 (10)	4986.4 (10)	-
$\alpha_{0,5}$	4974.8 (20)	-	-	-	-
$\alpha_{0,6}$	4951.2 (20)	4951.0 (15)	4950.3 (10)	4950.9 (10)	4951.3 (14)
$\alpha_{0,7}$	4933.8 (21)	-	-	-	-
$\alpha_{0,8}$	4900.0 (21)	-	-	-	-
$\alpha_{0,9}$	-	-	-	-	-
$\alpha_{0,10}$	-	-	-	-	-
$\alpha_{0,11}$	4852.2 (21)	-	-	-	-
$\alpha_{0,12}$	4794.3 (22)	-	-	-	-
$\alpha_{0,13}$	-	-	-	-	-
$\alpha_{0,14}$	-	-	-	-	-
$\alpha_{0,15}$	4736.4 (23)	4733.5 (15)	-	4736.1 (10)	4736.0 (8)
$\alpha_{0,16}$	4712.0 (24)	-	-	-	-
$\alpha_{0,17}$	4679.7 (24)	-	-	-	-
$\alpha_{0,18}$	4642.2 (25)	-	-	-	-
$\alpha_{0,19}$	-	-	-	-	-
$\alpha_{0,20}$	4631 (3)	-	-	-	-
$\alpha_{0,21}$	4598 (3)	-	-	-	-
$\alpha_{0,22}$	4565 (3)	-	-	-	-
$\alpha_{0,23}$	-	-	-	-	-
$\alpha_{0,24}$	4506 (3)	-	-	-	-
$\alpha_{0,25}$	-	-	-	-	-

¹ Published value was adjusted to recommended values by 1991Ry01 and 4986.4 (10) keV by 1976BaZZ due to changes in calibration energy.

² Additional values, which were not placed in the decay scheme, were reported at 5026.6 keV (population of 32-keV energy level of ²²⁷Ac) and at 5009.0 keV (population of 49-keV energy level of ²²⁷Ac).

Table 6: Adopted levels, theoretical and experimental alpha particle emission energies and hindrance factors.

Transition	Level energy (keV)	Theoretical alpha emission energy ¹ (keV)	Experimental alpha emission energy (keV)	HF
$\alpha_{0,0}$	0.0	5060.7 (8)	5058.6 (15)	250
$\alpha_{0,1}$	27.37 (1)	5033.8 (8)	5032.0 (21)	707
$\alpha_{0,2}$	29.98 (1)	5031.2 (8)	5028.4 (10)	95
$\alpha_{0,3}$	46.35 (1)	5015.1 (8)	5013.8 (14)	59.5
$\alpha_{0,4}$	74.14 (1)	4987.8 (8)	4986.4 (10)	629
$\alpha_{0,5}$	84.55 (1)	4977.6 (8)	4974.8 (20)	2 160
$\alpha_{0,6}$	109.94 (2)	4952.6 (8)	4951.3 (14)	26.5
$\alpha_{0,7}$	126.86 (2)	4936.0 (8)	4933.8 (21)	160
$\alpha_{0,8}$	160 (2)	4903.4 (22)	4900.0 (21)	141 000
$\alpha_{0,9}$	187.32 (3)	4876.6 (8)	-	-
$\alpha_{0,10}$	198.71 (4)	4865.4 (8)	-	-
$\alpha_{0,11}$	210.78 (5)	4853.5 (8)	4852.2 (21)	94
$\alpha_{0,12}$	271.29 (6)	4794.1 (8)	4794.3 (22)	1 300
$\alpha_{0,13}$	273.14 (3)	4792.3 (8)	-	-
$\alpha_{0,14}$	304.73 (5)	4761.2 (8)	-	9 600
$\alpha_{0,15}$	330.04 (1)	4736.3 (8)	4736.0 (8)	2.46
$\alpha_{0,16}$	354.50 (4)	4712.3 (8)	4712.0 (24)	11.7
$\alpha_{0,17}$	387.23 (2)	4680.1 (8)	4679.7 (24)	4.6
$\alpha_{0,18}$	425.59 (3)	4642.5 (8)	4642.2 (25)	56
$\alpha_{0,19}$	435.19 (2)	4633.0 (8)	-	75.8
$\alpha_{0,20}$	437.96 (4)	4630.3 (8)	4631 (3)	47
$\alpha_{0,21}$	469.24 (6)	4599.6 (8)	4598 (3)	146
$\alpha_{0,22}$	501.28 (7)	4568.1 (8)	4565 (3)	160
$\alpha_{0,23}$	537.0 (1)	4533.0 (8)	-	930
$\alpha_{0,24}$	562.8 (1)	4507.6 (8)	4506 (3)	126
$\alpha_{0,25}$	656.4 (3)	4415.6 (9)	-	43

¹ Calculated from alpha transition energy, taking into account the recoil energy of the daughter nucleus.

4.2 Alpha particle emission probabilities

The alpha emission probabilities have been determined from published data measurements when available; otherwise they are calculated from the balance of the decay scheme. All available experimental measurements were derived from magnetic spectrometers (¹⁹⁵⁶Hu96, ¹⁹⁶¹Ba42 and ¹⁹⁷⁶BaZZ). Data from Baranov (¹⁹⁶¹Ba42 and ¹⁹⁷⁶BaZZ) and Hummel (¹⁹⁵⁶Hu96) are in good agreement, with the exception of the $\alpha_{0,15}$ emission. For the recommended alpha emission probabilities, the evaluator has used values with uncertainties when available, adjusting the uncertainty as necessary. Otherwise the average of the values from Baranov (¹⁹⁶¹Ba42) and Hummel (¹⁹⁵⁶Hu96) was used, with the uncertainty being estimated on the basis of the decay scheme and individual values.

The theoretical emission probabilities were calculated from the P(γ +ce) balances using the GTOL software. There are large uncertainties associated with the theoretical calculations at the lower energy levels (see Table 7). These large uncertainties arise as a consequence of the

incomplete decay scheme, which in turn is due to the difficulties experienced in measuring low-energy gamma transitions.

There is a discrepancy between the alpha particle and gamma ray feeding and the gamma ray depopulating the first excited state. This is very probably due to the dominant $\gamma_{1,0}$ transition for which the emission probability and ICC value are not very well known.

Weak alpha particle emissions to levels 14, 19, 23 and 25 were expected, but not observed experimentally. These emissions were added to the decay scheme.

Table 7: Alpha particle emission energies, published and recommended probabilities.

Transition	Adopted emission energy (keV)	1961Ba42	1956Hu96	1976BaZZ ¹	Calculated ² emission probability	Adopted emission probability
$\alpha_{0,0}$	5060.7 (8)	11.0	10	11.7 (1)	11 (8)	11.7 (5)
$\alpha_{0,1}$	5033.8 (8)	~ 2.5	} 23	-	10 (8)	2.8 (3)
$\alpha_{0,2}$	5031.2 (8)	≤ 20.0		-	16 (4)	20 (2)
$\alpha_{0,3}$	5015.1 (8)	25.4	24	25.3 (2)	26 (5)	25.3 (5)
$\alpha_{0,4}$	4987.8 (8)	1.4	} 2.3	1.60 (5)	0.97 (20)	1.60 (20)
$\alpha_{0,5}$	4977.6 (8)	0.4		-	-1 (4)	0.4 (1)
$\alpha_{0,6}$	4952.6 (8)	22.8	22	22.5 (2)	21.4 (14)	22.5 (5)
$\alpha_{0,7}$	4936.0 (8)	3.0	2.8	-	2.51 (12)	2.9 (3)
$\alpha_{0,8}$	4903.4 (22)	0.002	-	-	0	0.002 (1)
$\alpha_{0,9}$	4876.6 (8)	-	-	-	-0.46 (16)	-
$\alpha_{0,10}$	4865.4 (8)	-	-	-	0.012 (10)	-
$\alpha_{0,11}$	4853.5 (8)	1.4	1.4	-	1.41 (15)	1.40 (15)
$\alpha_{0,12}$	4794.1 (8)	0.04	-	-	0.066 (8)	0.040 (15)
$\alpha_{0,13}$	4792.3 (8)	-	-	-	0.00 (5)	-
$\alpha_{0,14}$	4761.2 (8)	-	-	-	0.003 2 (9)	0.003 2 (9)
$\alpha_{0,15}$	4736.3 (8)	8.4	11	8.35 (8)	9.1 (5)	8.4 (4)
$\alpha_{0,16}$	4712.3 (8)	~ 1	1.4	-	1.20 (22)	1.20 (22)
$\alpha_{0,17}$	4680.1 (8)	1.5	2.1	-	1.8 (4)	1.8 (3)
$\alpha_{0,18}$	4642.5 (8)	~ 0.1	-	-	0.080 (6)	0.080 (6)
$\alpha_{0,19}$	4633.0 (8)	-	-	-	0.050 4 (11)	0.050 4 (11)
$\alpha_{0,20}$	4630.3 (8)	~ 0.1	-	-	0.078 (21)	0.078 (21)
$\alpha_{0,21}$	4599.6 (8)	0.015	-	-	0.003 65 (22)	0.015 (7)
$\alpha_{0,22}$	4568.1 (8)	0.008	-	-	0.001 5 (5)	0.008 (4)
$\alpha_{0,23}$	4533.0 (8)	-	-	-	0.000 76 (20)	0.000 76 (20)
$\alpha_{0,24}$	4507.6 (8)	0.003	-	-	0.003 6 (3)	0.003 6 (3)
$\alpha_{0,25}$	4415.6 (9)	-	-	-	0.002 1 (5)	0.002 1 (5)

¹Authors have reported only type A uncertainties.

²Emission probabilities calculated from balance of the decay scheme.

5. Gamma rays

5.1 Gamma-ray transitions and internal conversion coefficients

All gamma-ray transition energies were calculated from the differences in level energies as adopted from Browne (2001Br31).

Theoretical internal conversion coefficients (ICCs) were calculated using the BrIcc code (Kibédi et al., 2008Ki07) with the “frozen orbital” approximation, which uses interpolated values of Band et al. (2002Ba85).

The agreement between theoretical and measured ICC values was poor for $\gamma_{1,0}$ – under these circumstances, the experimental ICC data was adopted.

ICCs for some low-energy gamma transitions

$\gamma_{3,2}$: 16.370 (14) keV

The transition energy for this gamma ray is within 1 keV of the L3 shell binding energy of 15.971 keV. Since the model may be inaccurate close to the binding energy, the BrIcc code cannot be used to calculate theoretical ICCs. Therefore, the theoretical ICCs were calculated by Kibédi using the RAINE code, resulting in a value of 5.06 (7) for the L3 shell conversion and a total conversion coefficient of 8.58 (12) for this transition.

$\gamma_{1,0}$: 27.370 (10) keV

Disagreement between theoretically derived and experimentally measured data has been observed for this low-energy E1 transition (Table 8).

Table 8: Experimental and calculated values of α_L for the $\gamma_{1,0}$ transition of 27.370 (10) keV and E1 multipolarity.

Reference	α_L	Comments
1960As02	2.8 (3)	Not used - same author as 1974De11
1961Ba42	3.6 (4)	
1970De19	3.0 (3)	
1974De11	3.7 (3)	
Experimental mean	3.3 (4)	Weighted mean of 3 values
BrIcc code	2.66 (4)	

Asaro et al. suggest that the disagreement observed for this E1 transition can be explained by a small M2 contribution (1960As02). Assuming a multipolarity of E1+M2, the mixing ratio that agrees with the recommended value of $\alpha_L = 3.3$ (4) is $\delta = 0.007$.

$\gamma_{2,0}$: 29.980 (10) keV

The mixing ratio of 0.22 (2) from the evaluation of Browne (2001Br31) was derived from the measurements of De Pinho (1974De11). De Pinho derives the mixing ratio from an experimental

value of $\alpha_L = 220$ (20); the author suggests that a value of δ^2 from approximately 0.042 to 0.053 can explain the experimental α_L coefficients observed. Changing the mixing ratio value within the limits indicated above varies the $\gamma_{2,0}$ transition probability significantly from 24.4 to 27.6. More precise measurements are necessary to clarify the decay scheme at this level.

Summary of ICCs

A summary of the ICCs for the low-energy gamma-ray transitions is given in Table 9.

Table 9: Energies, multiplicities and internal conversion coefficients for low-energy gamma-ray transitions. Data within square parentheses [] are unconfirmed.

Transition	Transition energy (keV)	Multipolarity	Mixing ratio	α_T	α_L	α_M
$\gamma_{3,2}$	16.370 (14)	[E1]	-	8.58 (12)	5.06 (7)	2.68 (4)
$\gamma_{3,1}$	18.980 (14)	[M1]	-	113.2 (16)	2.35 (4)	82.7 (12)
$\gamma_{11,9}$	23.46 (6)	[M1]	-	241 (4)	182 (3)	44.1 (7)
$\gamma_{16,15}$	24.46 (4)	[M1]	-	214 (4)	161.3 (24)	39.0 (6)
$\gamma_{6,5}$	25.390 (22)	[M1]	-	191 (3)	144.6 (21)	34.9 (5)
$\gamma_{1,0}$	27.370 (10)	E1 [+M2]	[0.007]	4.5 (6)	3.3 (4)	0.87 (13)
$\gamma_{2,0}$	29.980 (10)	M1+E2	0.22 (2)	270 (30)	202 (21)	52 (6)
$\gamma_{6,4}$	35.800 (22)	[E1]	-	1.746 (25)	1.313 (19)	0.327 (5)
$\gamma_{5,3}$	38.200 (14)	M1+E2	0.18 (5)	89 (19)	66 (14)	17 (4)
$\gamma_{4,2}$	44.160 (14)	[M1]	-	37.4 (6)	28.3 (4)	6.79 (10)
$\gamma_{3,0}$	46.350 (10)	[E1]	-	0.879 (13)	0.663 (10)	0.1634 (23)
$\gamma_{20,17}$	50.73 (5)	[M1]	-	24.9 (4)	18.8 (3)	4.52 (7)
$\gamma_{7,4}$	52.720 (22)	[M1]	-	22.2 (4)	16.81 (24)	4.03 (6)
$\gamma_{5,2}$	54.570 (14)	[E1]	-	0.569 (8)	0.430 (6)	0.1053 (15)
$\gamma_{15,13}$	56.90 (3)	[M1+E2]	[0.41 (7)]	37 (6)	28 (5)	7.1 (12)
$\gamma_{5,1}$	57.180 (14)	E2	-	148.1 (21)	108.6 (16)	29.6 (5)
$\gamma_{17,15}$	57.190 (22)	E2	-	148.0 (21)	108.5 (16)	29.6 (5)
$\gamma_{9,7}$	60.46 (4)	[E1]	-	0.433 (7)	0.327 (5)	0.0800 (12)
$\gamma_{6,3}$	63.590 (22)	E2	-	88.8 (13)	65.1 (10)	17.8 (3)
$\gamma_{10,7}$	71.85 (5)	[M1]	-	8.98 (13)	6.79 (10)	1.630 (23)
$\gamma_{12,10}$	72.58 (7)	[M1]	-	8.71 (13)	6.59 (10)	1.582 (23)
$\gamma_{4,0}$	74.140 (10)	[E2]	-	42.6 (6)	31.2 (5)	8.53 (12)
$\gamma_{9,6}$	77.38 (4)	[M1]	-	7.23 (11)	5.47 (8)	1.313 (19)
$\gamma_{7,2}$	96.880 (22)	E2	-	12.02 (17)	8.81 (13)	2.41 (4)
$\gamma_{11,6}$	100.84 (5)	[E2]	-	9.97 (15)	7.30 (11)	2.00 (3)
$\gamma_{9,5}$	102.77 (3)	[E2]	-	9.12 (13)	6.69 (10)	1.83 (3)

5.2 Gamma-ray emission energies

There are a total of 9 sets of measurements for the gamma-ray emission energies. The recommended values were calculated from the differences in level energies as adopted from Browne (2001Br31) and were compared to the experimental values calculated from the weighted means (calculated with LWEIGHT4 code) of Lange (1969La04), De Pinho (1970De19), Leang (1970Le11), Börner (1979Bo30) and Teoh (1979Te02). The measurements from Falk-Vairant (1953Fa08), Foucher (1960Fo05), Baranov (1961Ba42) and Abou-Leila (1963Ab04) were not taken into account as they either do not have uncertainties, or are imprecise (uncertainties of a few keV). Experimental results and recommended values can be seen in Table 1 of Appendix 1.

Unplaced gamma rays

Below 45 keV

In the region 30-45 keV, various authors have reported 7 unplaced gamma rays. See Table 10 below for the reported energies.

Table 10: Experimental gamma-ray emission energies for unplaced gamma rays below 45 keV.

Reference	1961Ba42	1969La04	1970De19	1979Te02
Energy (keV)	34.0	30.7 (5)	31.00 (5)	30.87 (4)
			31.54 (4)	31.55 (5)
		39.6 (5)	39.57 (4)	39.73 (3)
			39.97 (2)	40.00 (3)
			42.48 (5)	42.41 (4)
			43.05 (5)	43.08 (4)

Baranov (1961Ba42), De Pinho (1970De19), Teoh (1970Te02) and Banham (1983Banham) have reported gamma-ray emission probabilities for these energies.

De Pinho et al. mention in their later paper (1974De19) that the six transitions reported in their earlier paper (1970De19) were not confirmed by later measurements and were the result of X-ray summing effects.

The evaluator has decided not to include these 7 transitions in the final table of evaluated gamma rays because their genuine existence is questionable.

Above 45 keV

With the exception of the 59.4 keV and 512.2 keV gamma-ray emissions reported by Lange (1969La04), the 318.1 keV gamma ray reported by Leang (1970Le11) and the 536.6 keV gamma ray reported by Teoh (1979Te02), the other unplaced gamma rays have been listed in the table.

The unplaced gamma-ray transition at 56.78 (4) keV detected by De Pinho (1970De19) and Teoh (1979Te02) was placed in the decay scheme based on the energy difference that constitutes the $\gamma_{15,13}$ transition. The energy for $\gamma_{15,13}$ calculated from the difference in level energies is 56.90 (3) keV which is in good agreement with the experimental value. If this gamma transition was absent, the balance of the decay scheme at level 13 would result in an alpha emission ($\alpha_{0,13}$) with an intensity of 0.19 (4) %; since weaker alpha emissions were detected in this region, it seems unlikely such an emission could be missed, which lends support to the placement of $\gamma_{15,13}$. With a transition from level 3/2- to level 5/2- and assuming a probability for $\alpha_{0,13} = 0$, a multipolarity M1+E2 with a mixing ratio of $\delta = 0.41$ (7) has been tentatively deduced.

5.3 Gamma-ray emission probabilities

There are a total of 9 sets of measurements for the gamma-ray emission probabilities. Four of the authors (1970De19, 1970Le11, 1979Te02 and 1983Banham) have measured over a wide energy range. Two of the publications are from the same authors (1970De19 and 1974De11); the later publication was favoured for emissions reported in both papers. Values were first normalised

such that the intensity of the 283.7 keV peak was set to be 100. The scaling factor used for the two data sets from De Pinho et al. was derived from their most recent publication.

The recommended values are the weighted mean of De Pinho (1970De19, 1974De11), Leang (1970Le11), Teoh (1979Te02), Aničin (1982An02) and Banham (1983Banham, 1984BAYS). The measurements from Foucher (1960Fo05), Baranov (1961Ba42) and Lange (1969La04) were not taken into account as they have no reported uncertainties. The experimental results and recommended values can be seen in Table 2 of Appendix 1.

Normalisation factor

Two experimental values were reported:
0.016 49 (27) from Banham (1984BAYS)
0.016 (2) from Leang (1970Le11)

The theoretical value obtained from the balance of the decay scheme to the ground state is 0.016 3 (14), which is in good agreement with the experimental values. As the theoretical normalisation factor is strongly influenced by the dominant $\gamma_{1,0}$ transition for which the theoretical and experimental ICC values do not agree, the evaluator has decided to use the experimental normalisation factor of 0.016 5 (3) derived from Banham (1984BAYS).

Low-energy gamma-ray emission probabilities

The emission probabilities for many of the gamma-ray transitions of 25 keV and below were either missing or imprecise, and had to be calculated using the balance of the decay scheme.

$\gamma_{3,2}$: 16.370 (14) keV

One measured value of $I_{\gamma_{3,2}} = 13.4$ (5) from Banham (1984BAYS) is available and was adopted as the recommended value. De Pinho (1974De11) has measured a transition ratio between $I_{\gamma+ce(19\text{ keV})} / I_{\gamma+ce(16.4\text{ keV})} \approx 18$ (5). Calculating the same ratio with the evaluated data gives 20.1 (15) which is in good agreement with the value of De Pinho et al.

$\gamma_{3,1}$: 18.980 (14) keV

The lack of coherent experimental data reported for this transition, due to the intense L X-rays observed in this part of the spectrum, made it impossible to calculate a weighted mean. This transition was calculated from the balance of the decay scheme at level 3. The resulting relative emission probability is $I_{\gamma_{3,1}} = 22.2$ (16).

$\gamma_{11,9}$: 23.46 (6) keV

No values have been reported for this transition, and the emission probability has been calculated from the balance of the decay scheme to level 11. The resulting gamma-ray emission probability is $I_{\gamma_{11,9}} = 0.288$ (35).

$\gamma_{16,15}$: 24.46 (4) keV

Two measurements are available for this gamma ray: 0.7 (3) from Teoh (1979Te02), and ~ 0.59 from De Pinho (1970De19). Neither of these values is very precise, and therefore the evaluator decided to evaluate this gamma-ray emission probability by means of the balance of the decay scheme to level 16. The resulting relative emission probability is $I_{\gamma_{16,15}} = 0.30$ (6).

$\gamma_{6,5}$: 25.390 (22) keV

There are four reported measurements for this transition. The measurements are not in very good agreement and fall into two ranges. Measurements of 7.6 (12) from De Pinho (1974De11) and 6.9 (10) from Teoh (1979Te02) contrast with the equivalent data from Leang (1970Le11) and Banham (1984BAYS) of 18.75 and 16.5 (9), respectively. The evaluator has decided to calculate the gamma-ray emission probability using the mean value of the balance of the decay scheme to levels 5 and 6 to give $I\gamma_{6,5} = 5.8$ (4). This value is in agreement with the lower set of values from De Pinho and Teoh.

Multiple placement - doublets

The gamma-ray transitions $\gamma_{15,1}$ (302.7 keV) and $\gamma_{17,5}$ (302.7 keV), $\gamma_{5,1}$ (52.7 keV) and $\gamma_{17,15}$ (52.7 keV) have been placed twice in the decay scheme; their individual emission probabilities have been suitably divided as follows.

$\gamma_{15,1}$ and $\gamma_{17,5}$: 302.7 keV

The combined evaluated relative probability for this gamma ray is 149.4 (12). Two authors have reported values for the separated doublet and the agreement between authors is poor:

Transition	1979Te02	1982An02
$\gamma_{15,1}$	100 (10)	138 (20)
$\gamma_{17,5}$	40 (5)	10.6 (26)

Only $\gamma_{15,1}$ is observed in the decay of ²²⁷Ra, so using the ratios between this transition and $\gamma_{15,0}$, $\gamma_{15,2}$ and $\gamma_{15,3}$ the expected transition probability for the doublet in the ²³¹Pa decay was calculated:

²⁷ Ra transition	Calculated $\gamma_{15,1}$ in ²³¹ Pa decay
$\gamma_{15,0}$	132 (16)
$\gamma_{15,2}$	137 (16)
$\gamma_{15,3}$	141 (16)
Unweighted mean	137 (16)

The recommended emission probability for $\gamma_{15,1}$ is 137 (16), and therefore the calculated $\gamma_{17,5}$ probability is 13 (6).

These values are in good agreement with the value of $\gamma_{15,1}=138$ (20) and $\gamma_{17,5}=10.6$ (26) from Aničin (1982An02).

$\gamma_{5,1}$ and $\gamma_{17,15}$: 52.7 keV

The combined evaluated relative probability for this gamma ray is 2.16 (14). One author (1979Te02) has reported values for the separated doublet. The calculated values using the balance of the decay scheme at level 17 are as follows:

Transitions	1979Te02	Calculated
$\gamma_{5,1}$	1.58 (16)	1.88 (19)
$\gamma_{17,15}$	0.96 (10)	0.28 (13)

The agreement between the calculated and measured values is poor for $I\gamma_{17,15}$. The evaluator has decided to adopt the calculated values.

6. References

- 1932Grosse A.V. Grosse, *Naturwiss.*, 20 (1932) 505. [half-life]
- 1949Va02 Q. Van Winkle, R.G. Larson, L.I. Katzin, *J. Am. Chem. Soc.* 71 (1949) 2585. [half-life]
- 1953Fa08 P. Falk-Vairant, M. Riou, *J. Phys. Radium* 14 (1953) 65. [gamma-ray emission probabilities and energies]
- 1956Hu96 J.P. Hummel, Thesis, Univ. California (1956); UCRL-3456 (1956). [alpha-particle emission probabilities]
- 1960As02 F. Asaro, F.S. Stephens, J.M. Hollander, I. Perlman, *Phys. Rev.* 117 (1960) 492. [L- and M-shell conversion coefficients]
- 1960Fo05 R. Foucher, *C. R. Acad. Sci., Paris.* 250 (1960) 1249. [gamma-ray emission probabilities]
- 1961Ba42 S.A. Baranov, V.M. Kulakov, P.S. Samoilo, A.G. Zelenkov, Yu.F. Rodionov, S.V. Pirozhkov, *Zhur. Eksptl. i Teoret. Fiz.* 41 (1961) 1475; *Soviet Phys. JETP* 14 (1962) 1053. [alpha-particle emission energies and probabilities, experimental conversions]
- 1961Br32 F. Bragança Gil, G.Y. Petit, *J. Phys. Radium* 22 (1961) 680. [half-life first excited level]
- 1961Ki05 H.W. Kirby, *J. Inorg. Nucl. Chem.* 18 (1961) 8. [half-life]
- 1963Ab04 H. Abou-Leila, R. Foucher, A.G. De Pinho, N. Perrin, M. Valadares, *J. Phys.* 24 (1963) 857. [spin and parity, multipolarity]
- 1963Su10 V.B. Subrahmanyam, Thesis, Univ. California (1963); UCRL-11082 (1963). [alpha-particle emission probabilities]
- 1966Ba14 G. Bastin, C.F. Leang, R.J. Walen, *C. R. Acad. Sci., Paris.* 262B (1966) 89. [alpha-particle emission energies]
- 1968Ba25 S.A. Baranov, V.M. Kulakov, V.M. Shatinskii, *Yadern. Fiz.* 7 (1968) 727; *Sov. J. Nucl. Phys.* 7 (1968) 442. [alpha-particle emission energies]
- 1968Br04 D. Brown, S.N. Nixon, K.M. Glover, F.J.G. Rogers, *J. Inorg. Nucl. Chem.* 30 (1968) 19. [half-life]
- 1968Ha22 G.R. Hagee, R.C. Lange, A.G. Barnett, A.R. Campbell, C.R. Cothorn, D.F. Griffing, H.J. Hennecke, *Nucl. Phys.* A115 (1968) 157. [spin and parity, conversion electron emission probabilities]
- 1969Ba20 A.G. Barnett, A.R. Campbell, G.R. Hagee, *J. Inorg. Nucl. Chem.* 31 (1969) 1553. [multipolarity, mixing ratio, conversion electron emission probabilities]
- 1969La04 R.C. Lange, G.R. Hagee, *Nucl. Phys.* A124 (1969) 412. [gamma-ray emission energies and probabilities]
- 1969Ro33 J. Robert, C.F. Miranda, R. Muxart, *Radiochim. Acta* 11 (1969) 104 [half-life]
- 1970De19 A.G. De Pinho, E.F. da Silveira, N.L. da Costa, *Phys. Rev.* C2 (1970) 572. [gamma-ray emission energies and probabilities, ICC]
- 1970Le11 C.F. Leang, *J. Phys. Paris*, 31 (1970) 269. [gamma-ray emission energies and probabilities]
- 1971Le10 C.F. Leang, *J. Phys. Paris*, 32 (1971), 95. [spin and parity]

- 1972Ga39 R.K. Garg, S.D. Chauhan, S. Sanyal, S.C. Pancholi, S.L. Gupta, N.K. Saha, Z. Phys. 257 (1972) 124. [half-life first excited level]
- 1974De11 A.G. De Pinho, L.T. Auler, A.G. da Silva, Phys. Rev. C9 (1974) 2056. [gamma-ray and X-ray emission probabilities, ICC]
- 1976Baranov S.A. Baranov, A.G. Zelenkov, V.M. Kulakov, At. Energy (Sov. J. At. Energy) 41 (1976) 342. [alpha-particle emission energies and probabilities]
- 1976BaZZ S.A. Baranov, A.G. Zelenkov, V.M. Kulakov, Proc. Advisory Group Meeting on Transactinium Nucl. Data, Karlsruhe, IAEA-186, Vol. III (1976) p.249. [alpha-particle emission energies and probabilities]
- 1979Bo30 H.G. Borner, G. Barreau, W.F. Davidson, P. Jeuch, T. von Egidy, J. Almeida, D.H. White, Nucl. Instrum. Methods 166 (1979) 251. [gamma-ray emission energies]
- 1979Te02 W. Teoh, R.D. Connor, R.H. Betts, Nucl. Phys. A319 (1979) 122. [gamma-ray emission energies and probabilities, ICC, multipolarity]
- 1982An02 I. Anicin, I. Bikit, C. Girit, H. Guven, W.D. Hamilton, A.A. Yousif, J. Phys.(London) G8 (1982) 369. [gamma-ray emission probabilities]
- 1983Banham M.F. Banham, R. Jones, Int. J. Appl. Radiat. Isot.34 (1983) 1225. [gamma-ray emission probabilities]
- 1984BAYS M.F. Banham, private communication quoted by 1986LoZT (1984) [gamma-ray emission probabilities]
- 1985Is03 T. Ishii, I. Ahmad, J.E. Gindler, A.M. Friedman, R.R. Chasman, S.B. Kaufman, Nucl. Phys. A444 (1985) 237. [half-life first excited level]
- 1986LoZT IAEA Technical Reports Series No.261 (1986). [evaluated gamma-ray emission energies and probabilities and alpha-particle emission energies and probabilities]
- 1990Ho28 N.E. Holden, Pure Appl. Chem. 62 (1990) 941. [half-life evaluation]
- 1991Ry01 A. Rytz, At. Data Nucl. Data Tables 47 (1991) 205. [alpha-particle probability and energy evaluation]
- 1996Sc06 E. Schönfeld, H. Janssen, Nucl. Instrum. Methods Phys. Res. A369 (1996) 527. [atomic data]
- 1998Ak04 Y.A. Akovali, Nucl. Data Sheets 84 (1998) 1. [r_0 radius parameter]
- 2002Ba85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor Jr., P.O. Tikkanen, S. Raman, At. Data Nucl. Data Tables 81 (2002) 1. [ICC]
- 2001Br31 E. Browne, Nucl. Data Sheets 93 (2001) 763. [spin, parity, energy level, multipolarity]
- 2003Au03 G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A729 (2003) 337. [Q value]
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor Jr., Nucl. Instrum. Methods Phys. Res. A589 (2008) 202. [theoretical ICC]

Appendix 1: Experimental and recommended gamma-ray emission energies and probabilities.

Table 1. Experimental and recommended gamma-ray emission energies (keV).

	1969La04	1970De17	1970Le11	1979Bo30 ^a	1979Te02	Calculated from experimental data	Recommended values
$\gamma_{3,2}$	-	16.5 (1)	-	-	-	16.5 (1)	16.370 (14)
$\gamma_{3,1}$	-	18.88 ^b	-	-	-	18.88	18.980 (14)
$\gamma_{11,9}$	-	-	-	-	-	-	23.46 (6)
$\gamma_{16,15}$	-	24.5 (1)	-	-	24.6 (5)	24.50 (10)	24.46 (4)
$\gamma_{6,5}$	25.3 (5)	25.54 (6)	25.2 (2)	-	25.36 (8)	25.46 (6)	25.390 (22)
$\gamma_{1,0}$	27.3 (5)	27.35 (2)	27.3 (2)	-	27.38 (2)	27.365 (20)	27.370 (10)
$\gamma_{2,0}$	29.8 (5)	29.95 (2)	29.9 (2)	-	30.01 (3)	29.968 (20)	29.980 (10)
$\gamma_{6,4}$	35.6 (5)	35.82 (3)	35.8 (3)	-	35.86 (4)	35.834 (30)	35.800 (22)
$\gamma_{5,3}$	38.0 (5)	38.20 (2)	38.1 (2)	-	38.19 (2)	38.194 (20)	38.200 (14)
$\gamma_{4,2}$	43.9 (5)	44.16 (2)	44.1 (2)	-	44.13 (2)	44.145 (20)	44.160 (14)
$\gamma_{3,0}$	46.1 (5)	46.37 (2)	46.2 (2)	-	46.32 (2)	46.344 (20)	46.350 (10)
$\gamma_{20,17}$	-	50.98 (5)	-	-	50.68 (6)	50.83 (15)	50.73 (5)
$\gamma_{7,4}$	52.4 (5)	52.74 (2)	52.6 (2)	-	52.66 (3)	52.658 (30)	52.720 (22)
$\gamma_{5,2}$	54.8 (5)	54.61 (2)	54.5 (2)	-	54.56 (3)	54.594 (20)	54.570 (14)
$\gamma_{15,13}$	-	56.76 (4)	-	-	56.79 (4)	56.78 (4)	56.90 (3)
$\gamma_{5,1}$	-	57.19 (3)	57.0 (2)	-	57.19 (3)	57.188 (30)	57.180 (14)
$\gamma_{17,15}$	-	57.19 (3)	57.0 (2)	-	57.19 (3)	57.188 (30)	57.190 (22)
$\gamma_{9,7}$	-	60.50 (3)	60.2 (3)	-	60.47 (8)	60.494 (30)	60.46 (4)
$\gamma_{6,3}$	63.3 (5)	63.67 (3)	63.5 (2)	-	63.60 (4)	63.642 (30)	63.590 (22)
$\gamma_{-1,1}$	-	70.50 (5)	-	-	70.45 (8)	70.49 (5)	70.49 (5) ^c

	1969La04	1970De17	1970Le11	1979Bo30 ^a	1979Te02	Calculated from experimental data	Recommended values
$\gamma_{10,7}$	-	71.9 (1)	-	-	71.9 (1)	71.9 (1)	71.85 (5)
$\gamma_{12,10}$	-	72.5 (1)	-	-	72.78 (8)	72.67 (14)	72.58 (7)
$\gamma_{4,0}$	-	74.18 (4)	74.1 (3)	-	74.08 (6)	74.15 (4)	74.140 (10)
$\gamma_{9,6}$	77.1 (5)	77.36 (3)	77.2 (2)	-	77.30 (4)	77.336 (30)	77.38 (4)
$\gamma_{7,2}$	-	96.88 (3)	96.7 (2)	-	96.80 (3)	96.838 (30)	96.880 (22)
$\gamma_{11,6}$	-	100.92 (4)	100.5 (5)	-	100.77 (4)	100.84 (5)	100.84 (5)
$\gamma_{9,5}$	102.5 (5)	-	102.5 (4)	-	102.6 (5)	102.5 (4)	102.77 (3)
$\gamma_{10,4}$	-	124.6 (1)	124.4 (5)	-	124.56 (8)	124.57 (8)	124.57 (4)
$\gamma_{12,7}$	-	144.5 (1)	144.4 (5)	-	144.33 (8)	144.40 (8)	144.43 (6)
$\gamma_{13,4}$	-	199 (1)	198.7 (6)	-	198.89 (10)	198.89 (10)	199.00 (3)
$\gamma_{14,4}$	-	-	-	-	230.0 (10)	230.0 (10)	230.59 (5)
$\gamma_{-1,2}$	-	242.2 (1)	-	-	242.16 (8)	242.18 (8)	242.18 (8) ^c
$\gamma_{13,2}$	243.0 (5)	243.0 (1)	242.9 (4)	-	243.15 (9)	243.08 (9)	243.16 (3)
$\gamma_{15,5}$	-	245.4 (1)	245.3 (5)	-	245.77 (9)	245.60 (13)	245.490 (14)
$\gamma_{13,1}$	-	246.0 (2)	-	-	246.05 (9)	246.04 (9)	245.77 (3)
$\gamma_{15,4}$	256.1 (5)	255.78 (7)	255.9 (3)	-	255.76 (8)	255.78 (7)	255.900 (14)
$\gamma_{14,3}$	-	258.4 (1)	-	-	258.54 (15)	258.44 (10)	258.38 (5)
$\gamma_{17,7}$	260.2 (5)	260.14 (8)	260.2 (3)	-	260.23 (8)	260.19 (8)	260.37 (3)
$\gamma_{13,0}$	273.5 (5)	273.08 (9)	273.2 (3)	273.237 (117)	273.15 (9)	273.14 (9)	273.14 (3)
$\gamma_{17,6}$	277.7 (5)	276.99 (9)	277.2 (3)	277.322 (15)	277.10 (9)	277.19 (7)	277.29 (3)
$\gamma_{15,3}$	283.9 (5)	283.56 (6)	283.7 (3)	283.690 (16)	283.65 (5)	283.679 (16)	283.690 (14)

	1969La04	1970De17	1970Le11	1979Bo30 ^a	1979Te02	Calculated from experimental data	Recommended values
$\gamma_{-1,3}$	-	286.55 (10)	-	-	286.60 (10)	286.58 (10)	286.58 (10) ^c
$\gamma_{15,2}$	300.5 (5)	299.94 (6)	300.1 (2)	300.069 (12)	300.02 (5)	300.062 (15)	300.060 (14)
$\gamma_{15,1}$	303.2 (5)	302.52 (6)	302.7 (2)	302.669 (11)	302.65 (5)	302.664 (15)	302.670 (14)
$\gamma_{17,5}$	303.2 (5)	302.52 (6)	302.7 (2)	302.669 (11)	302.65 (5)	302.664 (15)	302.680 (22)
$\gamma_{-1,4}$	-	310.0 (1)	-	-	310.0 (5)	310.0 (1)	310.0 (1) ^c
$\gamma_{17,4}$	313.0 (5)	312.88 (8)	312.9 (3)	-	312.94 (5)	312.92 (5)	313.090 (22)
$\gamma_{16,1}$	-	327.02 (10)	327.2 (4)	327.130 (188)	327.26 (10)	327.14 (10)	327.13 (4)
$\gamma_{15,0}$	330.2 (5)	329.89 (6)	330.0 (2)	330.057 (18)	330.06 (5)	330.045 (22)	330.040 (10)
$\gamma_{17,3}$	341.0 (5)	340.61 (7)	340.8 (2)	-	340.77 (6)	340.71 (6)	340.880 (22)
$\gamma_{18,4}$	-	351.4 (1)	-	-	351.6 (1)	351.50 (10)	351.45 (3)
$\gamma_{16,0}$	-	354.38 (8)	354.6 (2)	354.474 (76)	354.57 (8)	354.48 (8)	354.50 (4)
$\gamma_{17,2}$	356.6 (5)	356.96 (7)	357.2 (2)	-	357.21 (6)	357.10 (7)	357.250 (22)
$\gamma_{17,1}$	-	359.25 (10)	358.6 (4)	-	359.57 (10)	359.39 (15)	359.860 (22)
$\gamma_{20,4}$	364.2 (5)	363.74 (10)	363.9 (4)	-	363.93 (10)	363.84 (10)	363.82 (4)
$\gamma_{-1,5}$	-	374.9 (1)	374.9 (4)	-	375.01 (10)	374.95 (10)	374.95 (10) ^c
$\gamma_{18,3}$	379.5 (5)	379.09 (8)	379.2 (3)	-	379.41 (6)	379.29 (9)	379.24 (3)
$\gamma_{21,5}$	-	384.7 (1)	384.8 (3)	-	384.7 (1)	384.71 (10)	384.69 (6)
$\gamma_{17,0}$	-	387.0 (1)	-	-	-	387.0 (1)	387.230 (20)
$\gamma_{20,3}$	392.5 (5)	391.5 (1)	391.7 (3)	-	391.67 (9)	391.61 (9)	391.61 (4)
$\gamma_{18,2}$	-	395.5 (1)	395.7 (4)	-	395.49 (10)	395.50 (10)	395.61 (3)
$\gamma_{18,1}$	398.4 (5)	398.10 (8)	398.1 (3)	-	398.19 (9)	398.14 (8)	398.22 (3)

	1969La04	1970De17	1970Le11	1979Bo30 ^a	1979Te02	Calculated from experimental data	Recommended values
$\gamma_{19,1}$	408.1 (5)	407.71 (6)	407.7 (3)	407.829 (31)	407.80 (5)	407.802 (31)	407.820 (22)
$\gamma_{20,1}$	410.5 (5)	410.5 (1)	410.3 (10)	-	410.1 (1)	410.30 (12)	410.59 (4)
$\gamma_{22,4}$	-	-	-	-	427.0 (10)	427.0 (10)	427.14 (7)
$\gamma_{19,0}$	-	435.1 (1)	434.9 (8)	-	435.0 (1)	435.05 (10)	435.190 (20)
$\gamma_{20,0}$	437.9 (5)	437.9 (1)	437.9 (8)	-	438.10 (9)	438.01 (9)	437.96 (4)
$\gamma_{-1,6}$	-	438.7 (1)	-	-	438.8 (2)	438.72 (10)	438.72 (10) ^c
$\gamma_{24,4}$	487.2 (5)	486.7 (3)	486.6 (10)	486.827 (27)	486.8 (10)	486.826 (27)	488.66 (10)
$\gamma_{23,3}$	-	491.0 (6)	491 (2)	-	491.0 (10)	491.0 (6)	490.65 (10)
$\gamma_{22,0}$	-	501.6 (5)	501 (1)	-	501.0 (10)	501.4 (5)	501.28 (7)
$\gamma_{23,1}$	-	509 (1)	510 (1)	-	510.0 (10)	509.7 (10)	509.63 (10)
$\gamma_{24,3}$	516.2 (5)	516.2 (6)	516 (1)	-	516.1 (10)	516.2 (5)	516.45 (10)
$\gamma_{24,1}$	-	535.3 (7)	535 (1)	-	-	535.2 (7)	535.43 (10)
$\gamma_{25,6}$	-	546.6 (7)	546 (1)	-	546.6 (10)	546.5 (7)	546.5 (3)
$\gamma_{25,5}$	-	572.1 (8)	571 (2)	-	571.0 (10)	571.6 (8)	571.9 (3)
$\gamma_{25,4}$	-	-	583 (2)	-	-	583 (2)	582.3 (3)
$\gamma_{25,3}$	-	-	609 (2)	-	-	609 (2)	610.1 (3)

^{a)} Uncertainty on energy calibration of detectors was added to published data

^{b)} Obtained from private communication.

^{c)} Unplaced gamma.

Table 2. Experimental and recommended relative gamma-ray emission probabilities.

	$E_\gamma(\text{keV})$	P_γ^{rel}					Recommended values
		1970De19 ^a	1970Le11	1974De11 ^a	1979Te02	1984BAYS ^b	
$\gamma_{3,2}$	16.370 (14)	-	-	-	-	13.4 (5)	13.4 (5)
$\gamma_{3,1}$	18.980 (14)	-	-	-	-	76.7 (15)	22.2 (16) ^c
$\gamma_{11,9}$	23.46 (6)	-	-	-	-	-	0.29 (4) ^c
$\gamma_{16,15}$	24.46 (4)	~ 0.59	-	-	0.7 (3)	-	0.30 (6) ^c
$\gamma_{6,5}$	25.390 (22)	~ 5.9	~ 18.75	7.6 (12)	6.9 (10)	16.5 (9)	5.8 (4) ^c
$\gamma_{1,0}$	27.370 (10)	588 (28)	440 (130)	588 (28)	640 (50)	673 (13)	655 (22)
$\gamma_{2,0}$	29.980 (10)	5.8 (5)	6.3 (19)	5.88 (24)	6.5 (5)	5.63 (30)	5.87 (24)
$\gamma_{6,4}$	35.800 (22)	1.00 (12)	0.94 (31)	1.15 (9)	0.94 (5)	-	0.99 (6)
$\gamma_{5,3}$	38.200 (14)	9.4 (9)	6.3 (19)	8.6 (6)	9.4 (5)	8.59 (33)	8.8 (4)
$\gamma_{4,2}$	44.160 (14)	3.8 (4)	2.8 (9)	3.41 (24)	3.77 (40)	2.7 (5)	3.36 (24)
$\gamma_{3,0}$	46.350 (10)	13.18 (12)	8.1 (25)	11.1 (5)	12.97 (64)	10.6 (7)	11.5 (6)
$\gamma_{20,17}$	50.73 (5)	0.09 (4)	-	0.12 (3)	0.3 (1)	-	0.14 (5)
$\gamma_{7,4}$	52.720 (22)	5.4 (5)	3.8 (13)	4.41 (22)	4.85 (34)	5.4 (6)	4.60 (22)
$\gamma_{5,2}$	54.570 (14)	5.1 (5)	3.8 (13)	4.12 (19)	4.33 (35)	4.44 (32)	4.22 (19)
$\gamma_{15,13}$	56.90 (3)	0.35 (6)	-	0.31 (5)	0.27 (4)	-	0.29 (4)
$\gamma_{5,1}$	57.180 (14)	} 2.47 (24)	} 1.9 (6)	} 1.94 (11)	1.58 (16)	} 2.34 (15)	} 2.16 (14)
$\gamma_{17,15}$	57.190 (22)				0.96 (10)		
$\gamma_{9,7}$	60.46 (4)	0.41 (6)	0.19 (13)	0.36 (4)	0.3 (1)	0.29 (5)	0.32 (4)
$\gamma_{6,3}$	63.590 (22)	3.2 (3)	1.9 (6)	2.82 (21)	2.7 (3)	2.70 (9)	2.70 (9)
$\gamma_{-1,1}$	70.49 (5)	0.41 (6)	-	0.29 (6)	0.6 (2)	0.30 (5)	0.31 (5)
$\gamma_{10,7}$	71.85 (5)	0.12 (6)	-	0.12 (4)	0.1 (1)	-	0.12 (4)

	E _γ (keV)	P _V ^{rel}						Recommended values
		1970De19 ^a	1970Le11	1974De11 ^a	1979Te02	1982An02	1984BAYS ^b	
γ _{12,10}	72.58 (7)	0.24 (12)	-	0.18 (4)	0.2 (1)	-	-	0.18 (4)
γ _{4,0}	74.140 (10)	1.59 (18)	1.25 (44)	1.41 (12)	1.24 (20)	-	1.35 (5)	1.35 (5)
γ _{9,6}	77.38 (4)	4.3 (5)	2.5 (6)	3.53 (24)	4.31 (20)	-	3.45 (7)	3.67 (25)
γ _{7,2}	96.880 (22)	5.6 (6)	4.1 (9)	5.5 (4)	5.62 (28)	-	5.00 (10)	5.08 (14)
γ _{11,6}	100.84 (5)	2.0 (3)	0.75 (31)	1.35 (12)	1.66 (25)	-	1.38 (4)	1.37 (5)
γ _{9,5}	102.77 (3)	~ 1.2	2.8 (9)	1.35 (24)	<0.8	-	0.9 (2)	1.13 (25)
γ _{10,4}	124.57 (4)	0.29 (12)	0.13 (6)	-	0.29 (9)	0.23 (13)	0.259 (24)	0.261 (24)
γ _{12,7}	144.43 (6)	0.76 (24)	0.25 (13)	-	0.64 (30)	0.70 (6)	0.69 (5)	0.70 (5)
γ _{13,4}	199.00 (3)	0.35 (12)	0.06 (3)	-	0.23 (10)	0.28 (5)	0.246 (29)	0.18 (7)
γ _{14,4}	230.59 (5)	-	-	-	0.10 (5)	-	-	0.10 (5)
γ _{-1,2}	242.18 (8)	0.53 (6)	-	-	0.5 (2)	0.44 (8)	0.70 (4)	0.60 (6)
γ _{13,2}	243.16 (3)	2.18 (18)	2.5 (6)	-	2.97 (24)	2.51 (43)	1.87 (4)	2.2 (3)
γ _{15,5}	245.490 (14)	0.47 (6)	0.44 (13)	-	0.48 (12)	0.44 (8)	0.382 (31)	0.41 (3)
γ _{13,1}	245.77 (3)	-	-	-	0.7 (2)	0.70 (20)	-	0.70 (20)
γ _{15,4}	255.900 (14)	6.4 (4)	8.1 (13)	-	6.34 (41)	7.00 (48)	6.41 (6)	6.42 (6)
γ _{14,3}	258.38 (5)	0.15 (4)	-	-	0.15 (5)	0.13 (4)	0.06 (2)	0.093 (24)
γ _{17,7}	260.37 (3)	10.9 (6)	11.3 (19)	-	11.39 (57)	11.03 (14)	10.97 (10)	11.00 (10)
γ _{13,0}	273.14 (3)	3.65 (18)	4.4 (9)	-	3.48 (24)	3.48 (18)	3.50 (4)	3.51 (4)
γ _{17,6}	277.29 (3)	4.24 (24)	5.0 (9)	-	3.88 (25)	4.59 (58)	4.12 (5)	4.12 (5)

	E _γ (keV)	P _V ^{rel}						Recommended values
		1970De19 ^a	1970Le11	1974De11 ^a	1979Te02	1982An02	1984BAYS ^b	
γ _{15,3}	283.690 (14)	100.0	100.0	100.0	100.0	100.0	100.0 (8)	100.0
γ _{-1,3}	286.58 (10)	0.59 (6)	-	-	0.8 (3)	0.68 (10)	0.632 (30)	0.63 (3)
γ _{15,2}	300.060 (14)	144 (8)	144 (13)		149.6 (75)	143.2 (55)	146.3 (13)	146.2 (13)
γ _{15,1}	302.670 (14)	} 148 (8)	} 144 (13)	} 294 (8)	100 (10)	138 (20)	} 149.6 (12)	} 149.4 (12)
γ _{17,5}	302.680 (22)				40 (5)	10.6 (26)		
γ _{-1,4}	310.0 (1)	0.088 (29)	-	-	0.07 (3)	0.03 (2)	0.058 (12)	0.056 (12)
γ _{17,4}	313.090 (22)	6.0 (4)	6.9 (13)	-	7.05 (56)	5.93 (17)	5.97 (5)	5.98 (5)
γ _{16,1}	327.13 (4)	1.88 (12)	2.5 (13)	-	2.27 (28)	2.19 (44)	2.22 (4)	2.19 (5)
γ _{15,0}	330.040 (10)	82.4 (41)	81 (13)	82.4 (29)	81.9 (65)	82.1 (12)	82.4 (7)	82.3 (7)
γ _{17,3}	340.880 (22)	10.5 (5)	10.0 (25)	-	10.9 (13)	10.62 (16)	10.80 (9)	10.75 (9)
γ _{18,4}	351.45 (3)	0.224 (24)	-	-	0.15 (6)	0.44 (7)	0.102 (4)	0.17 (7)
γ _{16,0}	354.50 (4)	6.00 (35)	6.3 (13)	-	5.07 (56)	5.92 (16)	5.81 (6)	5.83 (6)
γ _{17,2}	357.250 (22)	10.9 (6)	9.4 (19)	-	9.67 (82)	10.35 (46)	10.14 (9)	10.16 (9)
γ _{17,1}	359.860 (22)	0.57 (5)	0.38 (19)	-	0.41 (18)	0.42 (8)	0.512 (14)	0.512 (14)
γ _{20,4}	363.82 (4)	0.47 (4)	0.38 (19)	-	0.42 (15)	0.45 (5)	0.488 (14)	0.483 (14)
γ _{-1,5}	374.95 (10)	0.294 (24)	0.19 (6)	-	0.24 (10)	0.21 (3)	0.282 (15)	0.270 (16)
γ _{18,3}	379.24 (3)	3.12 (24)	2.5 (9)	-	2.89 (23)	2.96 (10)	3.03 (4)	3.02 (4)
γ _{21,5}	384.69 (6)	0.259 (24)	0.13 (6)	-	0.18 (4)	0.18 (5)	0.221 (12)	0.221 (13)
γ _{17,0}	387.230 (20)	0.029 (12)	-	-	-	0.01 (1)	0.018 (6)	0.018 (6)

Comments on evaluation

	$E_{\gamma}(\text{keV})$	P_{γ}^{rel}					Recommended values
		1970De19 ^a	1970Le11	1979Te02	1982An02	1984BAYS ^b	
$\gamma_{20,3}$	391.61 (4)	0.43 (4)	0.31 (13)	0.52 (8)	0.35 (5)	0.408 (11)	0.408 (11)
$\gamma_{18,2}$	395.61 (3)	0.165 (18)	0.06 (3)	0.11 (2)	0.12 (2)	0.148 (10)	0.137 (13)
$\gamma_{18,1}$	398.22 (3)	0.59 (5)	0.44 (19)	0.49 (9)	0.49 (12)	0.574 (14)	0.572 (14)
$\gamma_{19,1}$	407.820 (22)	2.29 (18)	1.3 (6)	2.13 (18)	2.07 (16)	2.156 (24)	2.160 (24)
$\gamma_{20,1}$	410.59 (4)	0.118 (12)	0.06 (3)	0.19 (4)	0.21 (6)	0.099 (11)	0.109 (13)
$\gamma_{22,4}$	427.14 (7)	-	-	0.04 (2)	-	-	0.04 (2)
$\gamma_{19,0}$	435.190 (20)	0.212 (24)	0.125 (60)	0.12 (3)	0.18 (1)	0.177 (10)	0.178 (10)
$\gamma_{20,0}$	437.96 (4)	0.259 (24)	0.25 (13)	0.20 (6)	0.28 (3)	0.283 (16)	0.273 (16)
$\gamma_{-1,6}$	438.72 (10)	0.094 (24)	-	0.07 (2)	-	-	0.080 (20)
$\gamma_{24,4}$	488.66 (10)	0.112 (24)	0.063 (30)	0.15 (5)	0.10 (3)	0.091 (9)	0.100 (10)
$\gamma_{23,3}$	490.65 (10)	0.0294	0.006	< 0.04	0.04 (2)	0.023 (6)	0.024 (6)
$\gamma_{22,0}$	501.28 (7)	0.035 (12)	0.0125	0.05 (2)	0.07 (7)	0.053 (11)	0.046 (11)
$\gamma_{23,1}$	509.63 (10)	0.018 (6)	0.031	0.05 (2)	0.10 (4)	-	0.022 (10)
$\gamma_{24,3}$	516.45 (10)	0.082 (18)	0.050	0.06 (2)	0.06 (2)	0.093 (9)	0.083 (9)
$\gamma_{24,1}$	535.43 (10)	0.029 (12)	0.031	0.05 (2)	0.04 (2)	0.038 (7)	0.037 (6)
$\gamma_{25,6}$	546.5 (3)	0.035 (12)	0.025	0.04 (2)	0.06 (2)	0.056 (8)	0.050 (8)
$\gamma_{25,5}$	571.9 (3)	0.029 (12)	0.019	0.04 (2)	0.02 (2)	-	0.029 (12)
$\gamma_{25,4}$	582.3 (3)	-	0.019	-	0.26 (1)	-	0.019 (10) ^d
$\gamma_{25,3}$	610.1 (3)	-	0.031	-	0.43 (2)	-	0.031 (20) ^d

^{a)} Same author for both publications - data from (1974De11) used when available.

^{b)} Data originally published as 1983Banham, then as private communication (1984BAYS) within 1986LoZT.

^{c)} Calculated from balance of decay scheme.

^{d)} Values taken from 1970Le11; uncertainties were evaluated.

**²³²Th – Comments on evaluation of decay data
by A. Arinc**

This evaluation was completed in September 2008 and has a literature cut off date of April 2008. The weighted mean was applied to determine recommended values throughout the evaluation where the data were in statistical agreement. Where the data were not in statistical agreement, the Limitation of Relative Statistical Weights (LRSW) was used.

1. Decay Scheme

The nuclide ²³²Th disintegrates by alpha emission to two excited levels and to the ground state of ²²⁸Ra. The spin, parity, half-life of first excited state, multipolarities and level energies of ²²⁸Ra are based on the mass-chain evaluation of A. Artna-Cohen (1997Ar08).

Spontaneous fission and cluster decay of ²⁴⁻²⁶Ne have been observed by R. Bonetti (1995Bo18) with a partial half-life of $1.22 \cdot 10^{21}$ years for the spontaneous fission and a partial half-life greater than $5.04 \cdot 10^{21}$ years for the cluster decay. However, these decay modes were not taken into account in this evaluation.

2. Nuclear data

The Q(a) value of 4081.6 (14) keV is taken from the evaluation of Audi *et al.* (2003Au03). The effective Q-value calculated from decay scheme data is 4070 (70) keV.

The experimental half-life values are given in table 1.

Table 1. Experimental half-life values of ²³²Th

Reference	Half-life (10 ¹⁰ years)	Comments
1963Le21	1.401 (7)	Rejected by Chauvenet's criterion
1960Fa07	1.410 (14)	
1956Ma43	1.45 (5)	
1956Pi42	1.39 (3)	
1956Se17	1.42 (7)	
1938Ko01	1.39 (3)	
Recommended value	1.402 (6)	

The value of R. Macklin (1956Ma43) was excluded from the data analysis by Chauvenet's criterion. The data set is consistent and the recommended value, which is the weighted average of 5 remaining values, is 1.402 (6) 10¹⁰ years. The reduced chi-square value is 0.18 which is smaller than the critical value 3.32.

2.1 Alpha Transitions and emissions

The alpha transition and emission energies have been determined from the Q-value and level energies. Published alpha emission energies are given in table 2.

Table 2. Published alpha emission energies (keV)

Transition	$a_{0,0}$	$a_{0,1}$	$a_{0,2}$
1954Philbert ¹	4014 (20)	3939 (20)	
1957Ha08 ²	4012.3 (50)		
1961Ko11 ²	4013.6 (50) ⁴	3950 (8)	3825 (10)
1962Ko12 ²	4013.4 (50)		
1989Sa01	4012.3 (14)	3947.2 (20)	
Mean experimental emission values	4012.4 (14)	3947.3 (20)	3825 (10)
Calculated values ³	4011.2 (14)	3948.5 (14)	3810.0 (14)
Recommended Values	4011.2 (14)	3948.5 (14)	3810.0 (14)

¹ The values were adjusted by the evaluator for changes in the calibration energy.

² The values were adjusted as suggested by A. Rytz (1991Ry01)

³ Calculated from alpha transition energies taking into account the recoil of the alpha particle

⁴ For the $a_{0,0}$ transition, the value from 1961Ko11 was not taken into account as the same author published an updated value in 1962Ko12

Alpha hindrance factors were calculated using the ALPHAD computer program. A summary of the adopted level, alpha transition and emission values is presented in table 3.

Table 3. Adopted level, alpha particle transition and emission energies

Transition	Level Energy (keV)	Alpha Transition Energy (keV)	Alpha Emission Energy (keV)	HF
$a_{0,0}$	0.0	4081.6 (14)	4011.2 (14)	1.000
$a_{0,1}$	63.823 (20)	4017.8 (14)	3948.5 (14)	1.02 (7)
$a_{0,2}$	204.68 (3)	3876.9 (14)	3810.0 (14)	16 (5)

2.2 Gamma Transitions and Internal Conversion Coefficients

The recommended $\gamma_{1,0}$ transition energy of 63.811 (10) keV was calculated by taking the weighted mean of 63.81 (7) keV (1973Ta25), 63.81 (1) keV (1983Mi30) and 63.84 (6) keV (1989Sa01). The recommended $\gamma_{2,1}$ transition energy of 140.880 (10) keV was calculated by taking the weighted mean of 140.88 (1) keV (1983Mi30) and 140.83 (15) keV (1989Sa01).

Internal conversion coefficients were calculated using the BrIcc code (T.Kibédi, 2005KiZW), which uses interpolated values of Band *et al.* (2002Ba85).

The γ -ray transition energies, multipolarities and electron internal conversion coefficients are presented in table 4.

Table 4. Energies, multipolarities and electron internal conversion coefficients for gamma transitions

Transition	Transition Energy (keV)	Multipolarity	a_T	a_K	a_L	a_M
$g_{1,0}$	63.811 (10)	E2	80.4 (12)	-	59.1 (9)	16.05 (23)
$g_{2,1}$	140.880 (10)	E2	2.26 (4)	0.283 (4)	1.450 (21)	0.394 (6)

3. Alpha particle emissions

The alpha particle emission intensities were deduced from the decay scheme and can be viewed in table 5.

Table 5. Alpha particle emission energies and probabilities

Transition	Emission Energy (keV)	Emission intensity (%)
$\alpha_{0,0}$	4012.4 (14)	78.9 (13)
$\alpha_{0,1}$	3947.3 (20)	21.0 (13)
$\alpha_{0,2}$	3810.0 (14)	0.068 (20)

The values calculated using the balancing of the decay scheme are in good agreement with the experimental values (table 6) but the former values have been used as they are more precise.

Table 6: Reported values on alpha particle emission intensities

Reference	$\alpha_{0,0}$	$\alpha_{0,1}$	$\alpha_{0,2}$	Comments
1952Du12		24 (3)		See note 1)
1956Al30		22 (2)		See note 1)
1959Ko58		23 (3)	0.20 (8)	See note 2)
1961Ko11	77	23	0.2	No uncertainties. See note 2)
1983Mi30	77 (3)	23 (2)	0.066 (7)	See 3)
1989Sa01	100	33 (5)		

Notes:

- 1) The values found in the publications of D. Dunlavy (1952Du12) and G. Albouy (1956Al30) represent the percentage of conversion electron accompanying alpha decays ($\alpha_{0,1}$ and $\alpha_{0,2}$).
- 2) The values published by G. Kocharov in 1959Ko58 and 1961Ko11 appear to be from the same experiment.
- 3) The values from T. Mitsugashira (1983Mi30) are deduced by the author from the gamma emission probabilities measured by the author.

4. Gamma-ray emissions

The published data for the gamma-ray emissions can be viewed in table 7.

Table 7: Experimental data on gamma-ray emission probabilities

Reference	Absolute values (%)		Ratio of 140 keV/63 keV
	63 keV	140 keV	
1982Sa36	0.29 ¹ (2)		
1983Mi30	0.24 (3)	0.018 (2)	0.075 (13)
1983Ro23	0.247 ² (15)		0.102 (9)
1989Sa01			0.055 (10)

¹Value recalculated using the new DDEP recommended value for $\gamma_{1,0}$ (84 keV) of ²²⁸Th decay.

²Value recalculated using the new DDEP recommended value for $\gamma_{2,0}$ (238 keV) of ²¹²Pb decay.

The recommended 63 keV emission intensity of 0.259 (15) % was calculated by taking the weighted mean of 0.29 (2) % (1982Sa36), 0.24 (3) % (1983Mi30) and 0.247 (15) % (1983Ro23).

The recommended ratio 140 keV/63 keV of 0.080 (22) was calculated by taking the weighted mean of 0.075 (13) (1983Mi30), 0.102 (9) (1983Ro23) and 0.055 (10) (1989Sa01). The spread in the results is

quite significant and the reduced chi -square is larger than the critical chi -square. This may be due to the low probability of the gamma combined with the low specific activity of ²³²Th. The recommended emission probability for the 140 keV line, calculated from the above ratio and the 63 keV emission probability, is 0.021 (6) %.

Transition	Recommended Values	Gamma-ray emission intensity (%)	a _T
g _{1,0}	63.811 (10)	0.259 (15)	80.4 (12)
g _{2,1}	140.880 (10)	0.021 (6)	2.26 (4)

5. Atomic data

The values of ω_K , ω_L and n_{KL} relative probabilities of the X-ray and Auger emissions are from Schönfeld and Janßen (1996Sc06).

The energies and relative emission probabilities of the X-ray and Auger electrons have been calculated by using the computer code EMISSION.

References

- 1938Ko01 A.F.Kovarik, N.I.Adams, Jr., Phys.Rev. 54 (1938), 413
[half-life]
- 1952Du12 D.C.Dunlavey, G.T.Seaborg, Phys.Rev. 87 (1952), 165
[alpha probability]
- 1954Philbert G.Philbert, J.Genin, L.Vignerou, J.phys. et radium, 15 (1954), 16
[alpha energy]
- 1956Al30 G.Albouy, Ann.Phys. 1 (1956), 99
[alpha probability]
- 1956Ma43 R.L.Macklin, H.S.Pomerance, J.Nuclear Energy 2 (1956), 243
[half-life]
- 1956Pi42 E.Picciocto, S.Wilgain, Nuovo Cimento 4 (1956), 1525
[half-life]
- 1956Se17 F.E.Sentfle, T.A.Farley, N.Lazar, Phys.Rev. 104 (1956), 1629
[half-life]
- 1957Ha08 B.G.Harvey, H.G.Jackson, T.A.Eastwood, G.C.Hanna, Can.J.Phys.35 (1957), 258
[alpha energy]
- 1959Ko58 G.E.Kocharov, A.P.Komar, G.A.Korolev, Zhur.Eksptl.i Teoret.Fiz. 36 (1959), 68; Soviet Phys.JETP 9 (1959), 48
[alpha probability]
- 1960Be25 R.E.Bell, S.Bjornholm, J.C.Severiens, Kgl.Danske Videnskab.Selskab, Mat.-fys.Medd. 32 (1960), No.12
[half-life]
- 1960Fa07 T.A.Farley, Can.J.Phys. 38 (1960), 1059
[half-life]
- 1961Ko11 G.E.Kocharov, G.A.Korolev, Izv.Akad.Nauk SSSR, Ser.Fiz. 25 (1961), 237; Columbia Tech.Transl. 25 (1962), 227
[alpha energy, alpha probability]
- 1962Ko12 G.A.Korolev, G.E.Kocharov, Izv.Akad.Nauk SSSR, Ser.Fiz. 26 (1962), 235; Columbia Tech.Transl. 26 (1963), 233
[alpha energy]
- 1963Le21 L.J.LeRoux, L.E.Glendenin, Natl.Conf.Nucl.Energy, Application of Isotopes and Radiation, Pretoria, South Africa, F.L.Warren, Ed., Atomic Energy Board, Pelindaba, South Africa (1963), p.83
[half-life]

- 1973Ta25 H.W.Taylor, *Int.J.Appl.Radiat.Isotop.* 24 (1973), 593
[gamma probability]
- 1982Sa36 S.Sadasivan, V.M.Raghunath, *Nucl.Instrum.Methods* 196 (1982), 561
[gamma probability]
- 1983Mi30 T.Mitsugashira, M.Maki, S.Suzuki, Y.Shiokawa, *Radiochem.Radioanal.Lett.* 58 (1983), 199
[gamma probability and energy, alpha probability]
- 1983Ro23 J.-C.Roy, L.Breton, J.-E.Cote, J.Turcotte, *Nucl.Instrum.Methods* 215 (1983), 409
[gamma probability]
- 1989Sa01 S.K.Saha, S.M.Sahakundu, *J.Phys.(London)* G15 (1989), 73
[gamma probability and energy, alpha probability and energy]
- 1990Ho28 N.E.Holden, *Pure Appl.Chem.* 62 (1990), 941
[half-life evaluation]
- 1991Ry01 A.Rytz, *At.Data Nucl.Data Tables* 47 (1991), 205
[alpha probability and energy evaluation]
- 1995Bo18 R.Bonetti, C.Chiesa, A.Guglielmetti, R.Matheoud, G.Poli, V.L.Mikheev, S.P.Tretyakova, *Phys.Rev.* C51 (1995), 2530
[Spontaneous fission probability, cluster decay]
- 1996Sc06 E.Schonfeld, H.Janssen, *Nucl.Instrum.Methods Phys.Res.* A369 (1996), 527
[atomic data]
- 1997Ar08 A.Artna-Cohen, *Nucl.Data Sheets* 80 (1997), 723
[spin, parity, energy level, multipolarity]
- 2002Ba85 I.M.Band, M.B.Trzhaskovskaya, C.W.Nestor, Jr., P.O.Tikkanen, S.Raman, *At.Data Nucl.Data Tables* 81 (2002), 1
[ICC]
- 2003Au03 G.Audi, A.H. Wapstra, C. Thibault, *Nucl. Phys.* A729 (2003) 337.
[Q value]
- 2005KiZW T.Kibedi, T.W.Burrows, M.B.Trzhaskovskaya, C.W.Nestor, Jr.,
Proc.Intern.Conf.Nuclear Data for Science and Technology, Santa Fe, New Mexico, 26 September-1 October, 2004, R.C.Haight, M.B.Chadwick, T.Kawano, P.Talou, Eds., *AIP Conf.Proc.* (2005) 769
[ICC]

²³²U - Comments on Evaluation of Decay Data by Andy Pearce

This evaluation was completed in August 2008 drawing in part on the mass β -chain evaluation of Artna-Cohen^[1]. Some references not in the NSR database were identified by cross-referencing with the evaluation of Nichols^[2]. The literature available up until January 2008 was included.

1 Decay Scheme

The decay scheme (nuclear level energies, half lives and spins of ²²⁸Th) are based upon the adopted levels and gammas from Artna-Cohen^[1].

2 Nuclear Data

Uranium-232 decays primarily by alpha decay to excited states in ²²⁸Th. A small branching of exotic decay via ²⁴Ne emission and a smaller branching of spontaneous fission have been reported^[3-5]. The Q-value for alpha decay is taken from Audi, Wapstra and Thibault^[6-7]. The alpha decay branching is reported as essentially 100 %.

Seven published values of the half life were found in literature from which three independent values with uncertainties were used for analysis. The value of 1964Ch05^[8] determined by calorimetry has been adjusted taking into account the Q-values of Audi, Wapstra and Thibault^[6-7]. The authors of 1979Ag04^[9] measured the half life by two methods and both are stated here, although an arithmetic mean of the two has been used in analysis. Similarly the authors of 1964Ch05^[8] performed measurements by two methods and an arithmetic mean is taken for subsequent analysis. Sufficient experimental details were published in 1964Ch05 to allow the values to be recalculated using, for example, current values of Q, however doing so has no significant effect on the data. The adopted value has been determined by a LRSW weighted mean of the values from 1954Se26^[10], 1964Ch05^[8] and 1979Ag04^[9]. Overall the data are not consistent, however no valid reason could be found to exclude or prefer any of the three values. The discrepancies probably reflect the difficulties in measuring half lives of the order of several decades, and the uncertainty of the adopted value is large. The available data are presented in table 1.

There have been several publications on the spontaneous fission/cluster decay of ²³²U^[3-5] suggesting that ²⁴Ne cluster decay has been misidentified in earlier work as spontaneous fission. This leads to significantly lower values for branching to spontaneous fission than in previous evaluations. The value quoted here for spontaneous fission is that from 1990Bo16^[4] and that for cluster decay is a weighted mean of the values from 1985Ba18^[3] and 1990Bo16^[4]. The earlier data from Jaffey and Hirsch^[10] has not been published in open literature. Analysis of their data in the light of recent work would appear to confirm the cluster decay branching ratio at approximately 2×10^{-10} per 100 decays. The available data are presented in table 2.

Table 1. Measured half lives of alpha decay of ^{232}U . The reports 1949Go01 and 1949Ja01 were not used in analysis as they were presented without uncertainties.

Reference	Value (days)	Uncertainty (days)	Method
1949Go01 ^[11]	10 957	-	Ingrowth from ^{232}Pa
1949Ja01 ^[12]	25 567	-	Ingrowth from ^{236}Pu
1954Se26	26 880	370	Isotope dilution mass spec and proportional counting
1964Ch05 [A]	26 080	110	Calorimetry
1964Ch05 [B]	26 330	110	Alpha counting
1964Ch05 [mean]	26 130	110	-
1979Ag04 [A]	25 200	150	Isotope dilution mass spec and LS/proportional counting
1979Ag04 [B]	25 170	140	Relative activity vs. ^{233}U
1979Ag04 [mean]	25 090	140	-
1986Ag01	25 170	140	Relative activity vs. ^{233}U ; same data as half of 1979Ag01, republished
LRSW/expanded	25 800	390	-
Median (all values)	25 600	400	
Adopted	25 800	400	LWM/expanded

Table 2. Branching ratios for cluster decay and spontaneous fission, calculated where necessary from the partial decay constants using the recommended half life. The cluster decay value of Jaffey and Hirsh has been calculated by doubling the spontaneous fission value (in cluster decay one fragment will be detected compared with two in spontaneous fission).

Reference	Spontaneous Fission (%)	Cluster Decay (^{24}Ne) (%)
2000Bo46	$2.8 (6) \times 10^{-12}$	-
1990Bo16	$<10^{-12}$	$8.7 (8) \times 10^{-10}$
1985Ba18	-	$2.0 (5) \times 10^{-10}$
Jaffey & Hirsh (unpublished)	$9 (3) \times 10^{-11}$	$1.8 (12) \times 10^{-10}$
Adopted	$2.8 (6) \times 10^{-12}$	$5 (3) \times 10^{-10}$

2.1 Alpha-particle Transitions

The energies of the alpha -particle transitions have been determined from the Q-value and the adopted levels from Artna-Cohen^[1]. Alpha-particle hindrance factors were calculated using ALPHAD^[13]. The values so obtained are presented in table 3.

Table 3. Adopted level and alpha-particle transition energies

Transition	Level Energy (keV)	Transition Energy (keV)	Alpha-particle Emission Energy (keV)	HF
α_0	0	5413.63 (9)	5320.24 (9)	1
α_{58}	57.759 (4)	5355.87 (9)	5263.48 (9)	1.04 (3)
α_{187}	186.823 (4)	5226.81 (9)	5136.64 (9)	16.4 (4)
α_{328}	328.003 (4)	5085.63 (9)	4997.90 (9)	112.0 (24)
α_{378}	378.179 (10)	5035.45 (9)	4948.59 (9)	6490 (80)
α_{396}	396.078 (5)	5017.55 (9)	4931.00 (9)	5270 (50)
α_{519}	519.192 (6)	4894.44 (9)	4810.01 (9)	710 (50)
α_{831}	831.823 (10)	4581.81 (9)	4502.77 (9)	10.6 (8)
α_{874}	874.473 (18)	4539.16 (9)	4460.86 (9)	33 (9)

2.2 Gamma-ray Transitions and Internal Conversion Coefficients

Gamma-ray transition energies (Table 5) are calculated from the differences in level energies from Artna-Cohen^[1]. Transition energies calculated from the level scheme are compared with those derived from measured values in table 6. No precise measurements have been reported for the energy of the 831 keV E0 transition.

Table 5. Recommended gamma -ray emission energies, rescaled to be compatible with the values of 1971He23. Values from 1971He23 have been recalculated based on improved calibration standards from 2000He14. The uncertainties of the recalculated values have been increased to be not less than those in the original publication.

Nominal energy (keV)	57	129	141	191	209	270	328
1971He23	57.78 (6)	129.1 (1)	-	-	-	270.2 (5)	-
1973Ta25	57.78 (6)	-	-	-	-	-	-
1977Ku15	57.77 (6)	129.07 (6)	-	-	-	270.2 (2)	-
1979Bo30	-	129.070 (16)	-	-	209.238 (21)	270.235 (21)	328.004 (11)
1979He10	57.752 (13)	129.065 (3)	-	-	-	270.245 (7)	-
1987Da28	57.758 (7)	129.067 (7)	141.02 (3)	191.353 (11)	209.254 (7)	270.245 (8)	328.004 (7)
1995Ba42	57.75 (2)	129.05 (2)	140.99 (2)	191.34 (2)	209.25 (2)	270.24 (2)	328.02 (4)
LWEIGHT4	57.757 (6)	129.0655 (27)	140.999 (17)	191.351 (9)	209.252 (6)	270.2441 (13)	328.004 (6)
Adopted	57.752 (13)	129.065 (3)	140.999 (20)	191.351 (11)	209.252 (6)	270.245 (7)	328.004 (7)
Comments	From 1979He10	From 1979He10	LWEIGHT, uncert inc.	LWEIGHT, uncert inc.	LWEIGHT	From 1979He10	LWEIGHT

Table 5 (Cont.)

Nominal energy (keV)	332	338	478	503	546	773	817
1971He23	-	338.3 (4)	-	-	-	-	-
1973Ta25	-	-	-	-	-	-	-
1977Ku15	332.3 (3)	338.1 (2)	-	503.6 (3)	-	773.4 (5)	817 (1)
1979Bo30	-	338.321(10)	-	-	-	-	-
1979He10	332.371 (6)	338.320 (5)	-	503.819(23)	-	-	-
1987Da28	332.37 (5)	338.324 (7)	478.34 (5)	503.83 (5)	546.48 (5)	774.1 (2)	816.7 (1)
1995Ba42	332.36 (2)	338.31 (2)	478.45 (4)	503.69 (20)	546.45 (2)	774.06 (10)	816.49 (12)
LWEIGHT4	332.370 (6)	338.3209 (37)	478.41 (5)	503.818 (21)	546.454 (19)	774.05 (9)	816.62 (7)
Adopted	332.371 (6)	338.320 (5)	478.41 (5)	503.819 (23)	546.454 (21)	774.05 (9)	816.62 (7)
Comments	From 1979He10	From 1979He10	LWEIGHT	From 1979He10	LWEIGHT, uncert. inc.	LWEIGHT, uncert inc.	LWEIGHT, uncert inc.

Comments on evaluation

Table 6. Recommended gamma -ray transition energies and internal conversion coefficients. Measured transition energies are those obtained from gamma -ray emission energies via the recoil correction, whereas derived transition energies are those determined from the level scheme.

Measured Energy (keV)	Transition Energy (keV)		Multi-polarity from ENSDF	Conversion Coefficients			
	Measured	Derived		a_K	a_L	a_{M+}	a_T
57.752 (13)	57.752 (13)	57.759 (4)	E2	-	112.2 (16)	41.1 (5)	153.2 (22)
129.065 (3)	129.065 (3)	129.064 (6)	E2	0.264 (4)	2.54 (4)	0.933 (41)	3.74 (6)
140.999 (20)	140.999 (20)	141.013 (12)	E1	0.1689 (24)	0.0362 (5)	0.01169 (14)	0.217 (3)
191.351 (11)	191.350 (11)	191.356 (11)	E2	0.1710 (24)	0.443 (7)	0.162 (7)	0.776 (11)
209.252 (6)	209.252 (6)	209.255 (7)	E1	0.0672 (10)	0.01333 (19)	0.00429 (5)	0.0848 (12)
270.245 (7)	270.245 (7)	270.244 (6)	E1	0.0376 (6)	0.00716 (10)	0.002297 (25)	0.0470 (7)
328.004 (7)	328.005 (7)	328.003 (4)	E1	0.0245 (4)	0.00455 (7)	0.001458 (16)	0.0305 (5)
332.371 (6)	332.372 (6)	332.369 (7)	E1	0.0238 (4)	0.00441 (7)	0.001414 (16)	0.0297 (5)
338.320 (5)	338.321 (5)	338.319 (7)	E1	0.0229 (4)	0.00424 (6)	0.001358 (16)	0.0285 (4)
478.41 (5)	478.41 (5)	478.395 (18)	E1	0.01118 (16)	0.0198 (3)	0.000631 (7)	0.01379 (20)
503.819 (23)	503.820 (23)	503.820 (11)	E1	0.01009 (15)	0.001775 (25)	0.000565 (6)	0.01243 (18)
546.454 (21)	546.455 (21)	546.470 (18)	E1	0.00861 (12)	0.001500 (21)	0.000478 (5)	0.01058 (15)
774.05 (9)	774.05 (9)	774.064 (11)	E2	0.01204 (17)	0.00333 (5)	0.001199 (13)	0.01649 (23)
816.62 (7)	816.62 (7)	816.714 (18)	M1+E2 (d=1)	0.028 (18)	0.006 (3)	0.0019 (7)	0.036 (21)
-	-	831.823 (10)	E0	-	-	-	-

Internal conversion coefficients have been determined using the BrIcc code ^[14], using the gamma -ray multiplicities and mixing ratios from the evaluation of Artna -Cohen^[1]. No mixing ratio could be found in literature for the 817 keV transition and a mixing ratio of 1 has been assumed. Measured and adopted conversion coefficients are compared in table 9.

Table 9. Comparison of available measured conversion coefficients with the values calculated with the BrIcc code. Adopted values are from the BrIcc code in all cases.

Energy (keV)	BrIcc		1971He23 ^[25]		1982Ma52 ^[35]	
	a_K	a_L	a_K	a_L	a_K	a_L
57.752 (13)	-	112.2 (16)	-	117 (3)	-	85 (5)
129.065 (3)	0.264 (4)	2.54 (4)	0.23 (1)	2.45 (8)	-	2.74 (12)
140.999 (20)	0.1689 (24)	0.0362 (5)	0.11 (5)	-	-	-
191.350 (11)	0.1710 (24)	0.443 (7)	0.20 (2)	-	-	-
209.252 (6)	0.0672 (10)	0.01333 (19)	0.058 (1)	-	-	-
270.245 (7)	0.0376 (6)	0.00716 (10)	0.025 (3)	-	0.042 (3)	-
338.320 (5)	0.0229 (4)	0.00424 (6)	0.008 (1)	-	0.030 (2)	-

3 Atomic Data

All values of atomic data (ω_K , ω_L , n_{KL} , relative probabilities of the X-ray and Auger emissions) were derived from Schönfeld and Janßen^[15].

4 Alpha-particle Emissions

The alpha-particle emission probabilities were calculated from the balance of the gamma -ray decay scheme using GTOL^[16]. The adopted emission probabilities of the three strongest transitions a_0 , a_{58} & a_{187} are in good agreement with a weighted mean of the available measured data^[17-21], and those of a_{328} & a_{831} are in agreement with the measured values of 1964Le17^[19]. However, there are significant unexplained differences between the recommended values and the values measured by Baranov^[21] for the emission probabilities of a_{328} , a_{381} and a_{396} . Further measurements of the weak alpha -particle and gamma -ray transitions would be necessary to fully resolve these issues.

Table 4. Alpha -particle emission probabilities. Note the value quoted in the table may not match the published value exactly, as the values have been adjusted to a common scale (by dividing by the probability of the most intense emission) to take into account undetected alpha-particle emissions.

Trans.	Alpha-particle emissions per 100 decays							Adopted values (%)
	1955As28	1955Go32	1963Le17	1965Be15	1966Ba49	LWEIGHT	GTOL	
a_0	68 (1)	68.0	-	67.8 (7)	68.6 (6)	68.0 (4)	69.1 (6)	69.1 (6)
a_{58}	32 (1)	34.1	-	32.2 (3)	31.2 (4)	31.7 (7)	30.6 (6)	30.6 (6)
a_{187}	0.32 (3)	-	-	0.30 (9)	0.28 (2)	0.294 (23)	0.325 (6)	0.325 (6)
a_{328}	-	-	$6 (2) \times 10^{-3}$	-	$2.9 (2) \times 10^{-4}$	$6 (2) \times 10^{-3}$	$6.22 (9) \times 10^{-3}$	$6.22 (9) \times 10^{-3}$
a_{378}	-	-	-	-	$1.7 (3) \times 10^{-4}$	$1.7 (4) \times 10^{-4}$	$5.1 (6) \times 10^{-5}$	$5.1 (6) \times 10^{-5}$
a_{396}	-	-	-	-	$2.1 (3) \times 10^{-4}$	$2.1 (4) \times 10^{-4}$	$4.8 (4) \times 10^{-5}$	$4.8 (4) \times 10^{-5}$
a_{519}	-	-	-	-	-	-	$5.4 (4) \times 10^{-5}$	$5.4 (4) \times 10^{-5}$
a_{831}	-	-	$2.4 (7) \times 10^{-5}$	-	-	$2.4 (7) \times 10^{-5}$	$2.14 (16) \times 10^{-5}$	$2.14 (16) \times 10^{-5}$
a_{874}	-	-	-	-	-	-	$3.3 (9) \times 10^{-6}$	$3.3 (9) \times 10^{-6}$

5 Electron Emissions

Auger and conversion electron emissions per 100 decays were calculated from the gamma -ray data and conversion coefficients according to the method of Schönfeld and Janßen^[22] using version 3.10 of the code EMISSION.

6 Photon Emissions

6.1 X-ray Emissions

The X-ray intensities per 100 decays have been calculated from the gamma -ray data and conversion coefficients using version 3.10 of the code EMISSION.

6.2 Gamma-ray Emissions

The gamma-ray emission energies have been taken from 1979He10 [23] where possible, in which precise measurements were made by measuring energy differences against accepted calibration standards. Only the directly measured values have been taken as the decay scheme used to derive further values was incomplete. These values have been adjusted to reflect the updated calibration standards given in 2000He14[24]. Where gamma-ray lines are not present in 1979He10, weighted means of the values in 1971He23[25], 1973Ta25[26], 1977Ku15[27], 1979Bo30[28], 1987Da28[29] and 1995Ba42[30] were taken. The values in these publications were first rescaled by a least-squares fit to be compatible with 1979He10. In most cases the energy shift incurred by doing so was very small.

Relative gamma-ray emission probabilities were determined by a weighted mean of values in 1966Ah02[31], 1977Ku15[27], 1984Ge07[32] and Banham & McChrohon [33]. Data for many of the less intense gamma-ray emissions have only been reported in 1977Ku15. In determining means, values were normalised to the 129 keV gamma-ray transition rather than the most intense 60 keV transition due to the experimental difficulties in measuring gamma-ray emissions below 100 keV. There were three absolute emission probability measurements, two by 1984Ge07 [32] and one by Banham & McChrohon [34]. The reference value of the normalisation factor was determined from the weighted mean of the absolute values of the 129 keV line and is $6.86(7) \times 10^{-4}$ per 100 decays. The normalisation factor was also calculated with the code GABS[34] and by balance of the feeding to the 1st excited state and the figures thus obtained were $7.0(3) \times 10^{-4}$ per 100 decays and $7.08(16) \times 10^{-4}$ per 100 decays respectively. These values are statistically compatible with the reference value.

The intensity of the 831 keV E0 transition is given by 1963Le17 as $2(1) \times 10^{-6}$ per 100 decays. The 831 keV transition is E0, thus, it emits only electrons.

Table 7. Relative gamma-ray emission probabilities, normalised to 100 emissions for the 129 keV line. Note one additional significant figure is quoted in columns 2-6 over that which would normally be quoted; this is intentional to allow statistics to be calculated. The 817 keV line is quoted by 1977Ku15 as ~ 0.0011 ; the uncertainty assumed is a relative uncertainty of $\pm 100\%$ at 3 s, giving a relative emission probability of 0.0011 ± 0.0004 .

Energy (keV)	Gamma-ray emissions per 100 emissions at 129 keV				
	1966Ah02	1977Ku15	1984Ge07	Banham 1986	Adopted
57.752 (13)	256 (26)	298.9 (118)	291.5 (65)	292.5 (42)	292 (4)
129.065 (3)	100	100	100	100	100
140.999 (20)	-	0.00453 (189)	-	-	0.0045 (19)
191.351 (11)	-	0.0453 (40)	-	-	0.0453 (40)
209.252 (6)	-	0.0155 (38)	-	-	0.0155 (38)
270.245 (7)	4.62 (90)	4.264 (198)	-	4.660 (68)	4.62 (9)
328.004 (7)	4.10 (88)	3.774 (161)	-	4.168 (62)	4.12 (9)
332.371 (6)	-	0.0717 (44)	-	-	0.0717 (44)
338.320 (5)	-	0.05396 (249)	-	-	0.0540 (25)
478.41 (5)	-	0.00208 (80)	-	-	0.0021 (8)
503.819 (23)	-	0.02113 (130)	-	-	0.0211 (13)
546.454 (21)	-	0.00147 (91)	-	-	0.0015 (9)
774.05 (9)	-	0.00679 (115)	-	-	0.0068 (12)
816.62 (7)	-	~ 0.0011	-	-	0.0011 (4)
831.823 (10)	-	E0	-	-	E0

Table 8. Recommended gamma-ray emission probabilities.

Energy (keV)	Multipolarity	Gamma-ray Emission Probability per 100 decays
57.752 (13)	E2	0.200 (4)
129.065 (3)	E2	0.0686 (7)
140.999 (20)	E1	$3.1 (13) \times 10^{-6}$
191.351 (11)	E2	$3.1 (3) \times 10^{-5}$
209.252 (6)	E1	$1.1 (3) \times 10^{-5}$
270.245 (7)	E1	0.00317 (7)
328.004 (7)	E1	0.00283 (7)
332.371 (6)	E1	$4.9 (3) \times 10^{-5}$
338.320 (5)	E1	$3.70 (18) \times 10^{-5}$
478.41 (5)	E1	$1.4 (6) \times 10^{-6}$
503.819 (23)	E1	$1.45 (9) \times 10^{-5}$
546.454 (21)	E1	$1.0 (6) \times 10^{-6}$
774.05 (9)	E2	$4.7 (8) \times 10^{-6}$
816.62 (7)	M1+E2 (d=1)	$8 (3) \times 10^{-7}$
831.823 (10)	E0	0 [TI 2 (1) $\times 10^{-6}$]

7 References

- [1]. Artna-Cohen, A. *Nuclear Data Sheets* 80 (1997) p. 723 [1997NDS]
- [2]. Nichols A. L. and James M. F. *Radioactive Heavy Element Decay Data for Reactor Calculations* AEE Winthirth Report 1407 (1981)
- [3]. Barwick S. W. Price P. B. and Stevenson J. D. *Phys Rev C* 31 (1985) p. 1984 [1985Ba18]
- [4]. Bonetti R. Fioretto E. Migliorino C. Pasinetti A. Barranco F. Vigezzi E. and Broglia R. A. *Phys Lett. B* 241 (1990) p. 179 [1990Bo16]
- [5]. Bonetti R. and Guglielmetti A. *Phys Rev. C* 62 (2000) 047304 [2000Bo46]
- [6]. Audi G. Wapstra A. H. and Thibault C. *Nucl. Phys. A* 729 (2003) p. 337 [2003Au03]
- [7]. Wapstra A. H. Audi G. and Thibault C. *Nucl. Phys. A* 729 (2003) p. 129 [2003Wa32]
- [8]. Chilton J. M. Gilbert R. A. Leuze R. E. and Lyon W. S. *J Inorg. Nucl. Chem.* 26 (1964) p. 395 [1964Ch05]
- [9]. Aggarwal S. K. Manohar S. B. Acharya S. N. Prakash S. and Jain H. C. *Phys. Rev. C* 20 (1979) p. 1533 [1979Ag04]
- [10]. Sellers P. A. Stevens C. M. and Studier M. H. *Phys. Rev.* 94 (1954) p. 952. [1954Se26]
- [11]. Gofman J. W. Seaborg G. T. *The Transuranium Elements: Research Papers* National Nuclear Energy Series 14B p. 1427 [1949Go01]
- [12]. James R. A. Florin A. E. Hopkins H. H. Jr. and Ghiorso A. *The Transuranium Elements: Research Papers* National Nuclear Energy Series 14B p. 1604 [1949Ja01]
- [13]. ALPHAD, see for example www.nndc.bnl.gov/toolspublications/toolspublications.html

Comments on evaluation

- [14]. Kibédi T. Burrows T. W. Trzhaskovskoya M. B. Nestor C. W. ANU-P/1684 (2004)
- [15]. Schönfeld E. and Janßen H. *Nucl. Instrum. Meth. A* 369 (1996) p. 527
- [16]. GTOL, see for example www.nndc.bnl.gov/toolspublications/toolspublications.html
- [17]. Asaro F. and Perlman I. *Phys. Rev.* 99 (1955) p. 37 [**1955As28**]
- [18]. Scharff-Goldhaber G. Mateosian E. Harbottle G and McKeown M. *Phys. Rev.* 99 (1955) p. 180. [**1955Go32**]
- [19]. Lederer C. M. Thesis, University of California (1963) UCRL-11028 [**1963Le17**]
- [20]. Bertolini G. Cappellani F. Restelli G. Scherff H. L. *Nucl. Phys.* 68 (1965) p. 170 [**1965Be15**]
- [21]. Baranov S. A. Aliev L. G. Kulakov V. M. and Shatinskii V. M. *Soviet J. Nucl. Phys.* 4 (1967) p. 673 [**1966Ba49**]
- [22]. Schönfeld E. and Janßen H. *Appl. Radiat. Isot.* 52 (2000) p. 596
- [23]. Helmer, R. G. *Nucl. Instrum. Meth.* 164 (1979) p. 355 [**1979He10**]
- [24]. Helmer R. G. and van der Leun C. *Nucl. Instrum. Meth. A* 450 (2000) p. 35 [**2000He14**]
- [25]. Herment M. and Vieu C. *C. R. Acad. Sci. B* 273 (1971) p. 1058 [**1971He23**]
- [26]. Taylor H. W. *Appl. Radiat. Isot.* 24 (1973) p. 594 [**1973Ta25**]
- [27]. Kurcewicz W. Kaffrell N. Trautmann N. Plochocki A. Zylicz J. Matul M. Stryczniewicz K. *Nucl. Phys. A* 289 (1977) p. 1 [**1977Ku15**]
- [28]. Borner H. G. Barreau G. Davidson W. F. Jeuch P. von Egidy T. Almeida J. White J. H. *Nucl. Instrum. Meth.* 166 (1979) p. 251 [**1979Bo30**]
- [29]. Dalmaso J. Maria H. and Ardisson G. *Phys. Rev. C* 36 (1987) p. 2510 [**1987Da28**]
- [30]. Baltzer H. Freitag K. Gunther C. Herzog P. Manns J. Muller U. Paulsen R. Sevenich P. Weber T and Will B. Z. *Phys. A* 352 (1995) p. 47 [**1995Ba42**]
- [31]. Ahmad I. Thesis, University of California (1966) [**1966Ah02**]
- [32]. Gehrke R. J. Novick V. J. and Baker J. D. *Appl. Radiat. Isot.* 35 (1984) p. 581 [**1984Ge07**]
- [33]. Banham M. F. and McChrohon R. *The Measurement of Gamma-ray Emission Probabilities for the Nuclides ²³¹Pa, ²³³Pa, ²³²U, ²³⁵U, ²³⁷U and ²³⁷Np* AERE Report 11353
- [34]. GABS, see for example www.nndc.bnl.gov/toolspublications/toolspublications.html
- [35]. Mahajan A. S. and Bidarkundi M. S. *Ind. J. Pure Appl. Phys.* 20 (1982) p. 701 [**1982Ma52**]

²³³Th– Comments on evaluation of decay data
by V.P.Chechev and N.K.Kuzmenko

This evaluation was done originally in 2004 and then updated and revised in January 2009 with a literature cut-off by the same date.

1. DECAY SCHEME

The decay scheme is based on 2005Si15. Some gamma-ray transitions were not observed directly in ²³³Th decay but have been adopted from ²³⁷Np α -decay. There are no precise measurements of beta transitions from the decay of ²³³Th available. Data on gamma-ray emission probabilities have been taken mainly from measurements in 2008De31.

Several unplaced gamma rays were observed. These gamma rays carry ≤ 3 % of the total intensity of all the gamma rays placed in the decay scheme.

2. NUCLEAR DATA

Q^- value is from 2003Au03.

The recommended half-life of ²³³Th is based on the experimental results given in Table 1.

Table 1. Experimental values of ²³³Th half-life (in minutes)

Reference	Author(s)	Original value	Re-estimated	Measurement method
1952Ru10	Rutledge et al.	23,6 (6)		β -counting
1955Je26	Jenkins	22,12 (5) ^a	22,12 (7)	β -counting, good purification of the thorium sample
1957Dr46	Dropesky and Langer	22,4 (1)		β -counting
1969HoZY	Hoekstra	22,3 (1)		Gamma-ray counting
1989Ab05	Abzouzi et al.	22,30 (2) ^b	22,30 (10)	Gamma-ray counting
1998Us01	Usman et al.	21,83 (4) ^c	21,83 (10)	Gamma-ray counting
2008De31	DeVries and Griffin	21,99 (5) ^d	21,99 (9)	Liquid scintillation counting, multiple purifications of the thorium sample

^a Original value was deduced as the mean value from two experiments with the same result of 22,12 (7) min. As these experiments were correlated the evaluators used the value of 22,12 (7) min.

^b Uncertainty may include only statistical errors. The evaluators have taken into account the contribution of possible systematic errors associated with the gamma-ray counting method (see below).

^c Possible systematic errors associated to the gamma-ray counting method may have been caused by the use of a different shape of pulser and gamma-ray peaks, and by contamination of the gamma-ray spectrum with ²³³Pa and other radionuclides. Based on data scattering in three experiments for the strongest 459 keV and 669 keV gamma ray peaks (with a half-life ranging from 21,748 to 21,945 min) the evaluators have estimated the overall uncertainty of 0.10 min, which includes possible systematic errors.

^d Authors reported only statistical errors $\leq (0,2 - 0,3)$ %. Assuming possible systematic errors of the same order of magnitude ($\sim 0,3$ %), the evaluators have estimated an overall uncertainty of 0,09 min.

The value from 1952Ru10 has been omitted because it is an outlier. The unweighted mean of the 6 remaining values from Table 1 is 22,16 (9), the weighted mean is 22,15, the internal uncertainty is 0,037, the external uncertainty is 0,082. The LWEIGHT computer program recommended the weighted mean and its external uncertainty. Therefore, the recommended value of ²³³Th half-life is 22,15 (8) minutes.

2.1. Beta-transitions

The energies of β^- transitions have been obtained from the Q^- value and the ²³³Pa level energies given in Table 2, taken mainly from 2005Si15. The adopted level energies include also available data from ²³⁷Np alpha-decay. The energies of the levels "5", "10" and "12" have been obtained directly from the energies of the $\gamma_{5,0}$ (94,65 keV), $\gamma_{10,0}$ (237,86 keV) and $\gamma_{12,0}$ (447,762 keV) gamma rays, respectively.

The comparison of measured and recommended energies of β^- transitions is given in Table 3.

The emission probabilities of β^- transitions have been deduced from the P($\gamma+e$) balance at each level of ²³³Pa. The accurate combined β^- intensity of the $\beta_{0,0}$ and $\beta_{0,1}$ transitions is 84,0 (5) %, using 100 % for the total intensity of the beta decay from ²³³Th.

Table 2. ²³³Pa levels populated in ²³³Th decay

Level	Energy (keV)	Spin and Parity	Half-life	Probabilities of β^- -transitions (%)
0	0	3/2 ⁻	26,98 (2) d	34 (6)
1	6,65 (5)	1/2 ⁻		50 (6)
2	57,10 (2)	7/2 ⁻		-
3	70,49 (10)	5/2 ⁻		-
4	86,477 (10)	5/2 ⁺		-
5	94,65 (5)	3/2 ⁺		10,4 (4)
6	103,8 (1)	7/2 ⁺		-
7	169,159 (10)	1/2 ⁺		0,692 (12)
8	201,62 (5)	3/2 ⁺		0,074 (8)
9	212,34 (5)	5/2 ⁺		-
10	237,86 (6)	5/2 ⁺		-
11	257,30 (15)	5/2 ⁻		0,60 (3)
12	447,762 (20)	3/2 ⁻		0,821 (14)
13	454,40 (7)	3/2 ⁺		0,217 (13)
14	553,88 (6)	1/2 ⁺ , 3/2 ⁺		1,23 (3)
15	585,50 (5)	3/2 ⁺		0,15 (3)
16	669,9 (5)	(3/2 ⁻)		0,0174 (22)
17	764,55 (6)	1/2 ⁺ , 3/2 ⁺		1,19 (3)
18	811,6 (2)	(3/2 ⁺)		0,385 (4)
19	984,8 (5)	(3/2 ⁺)		0,205 (2)
20	1018,7 (5)	(3/2)		0,0434 (9)

Table 3. Measured and recommended energies of β^- -transitions

	1957Dr46	1957Fr55	Recommended
$\beta_{0,0}$	1230 (10)	1245 (3)	1243,1 (14)
$\beta_{0,5}$		1158	1148,4 (14)
$\beta_{0,7}$		1073	1073,9 (14)
$\beta_{0,11}$		880	985,8 (14)
$\beta_{0,12}$		790	795,3 (14)
$\beta_{0,13}$			788,7 (14)
$\beta_{0,14}$			689,2 (14)
$\beta_{0,15}$			657,6 (14)
$\beta_{0,16}$		580	573,2 (14)
$\beta_{0,17}$			478,5 (14)
$\beta_{0,18}$			431,5 (14)

2.2. Gamma Transitions and Internal Conversion Coefficients

The recommended energies of gamma-ray transitions are virtually the same as the gamma-ray energies because nuclear recoil is negligible for ²³³Pa.

The gamma-ray transition probabilities [P(γ +ce)] have been obtained using the gamma-ray emission probabilities and total conversion coefficients (ICC). The ICC have been interpolated using the BrIcc package with the so called “Frozen Orbital” approximation (2008Ki07). The uncertainties in the ICC for pure multiplicities have been taken as 2 %.

P(γ +ce)(8,22 keV) has been obtained from the intensity imbalance at the level “4” (86,477 keV) assuming negligible beta transition probability to this level [P($\beta_{0,4}$)=0]. The obtained value of P(γ +ce)(8,22 keV)=12,3 (4) % differs from \approx 19 % estimated in 2005Si15 using the intensity of N conversion electrons measured in 1976JeZU. It should be noted that the E2/M1 mixing ratio for the 8,22 keV gamma ray transition has not been measured. However, the experimental P_{ce} (N2) \approx P_{ce} (N3) indicates a large contribution of E2 multipolarity for this transition.

The ICC for the anomalous E1 gamma-ray transition $\gamma_{4,0}$ (86,477 keV) has been taken from 1988Wo01. The value of the total internal conversion coefficient of 1,43 (8) measured in 1988Wo01 agrees well with the theoretical assessment of 1,49 (18) (which includes the effect of nuclear penetration) obtained in 2008Go10 for this anomalous E1 gamma-ray transition.

The conversion electron data of 1988Wo01 indicate that the gamma-transition $\gamma_{4,2}$ (29,37 keV) also may be an anomalous E1. However, the evaluators have been adopted (following 2005Si15) the theoretical ICC for this transition since the detector efficiency was not completely reliable for energies lower than 50 keV, as pointed out in 1988Wo01.

Multipolarities and E2/M1 mixing ratios have been adopted from conversion electron measurements of 1972SeZI, 1976JeZU, and from data on ²³⁷Np alpha-decay (see 2005Si15).

3. ATOMIC DATA

The atomic data (fluorescence yields, X-ray energies and relative probabilities, and Auger electrons energies and relative probabilities) have been deduced by using the SAISINUC software (2002Be).

4. ELECTRON EMISSIONS

The energies of the conversion electrons have been obtained from the gamma-ray transition energies and the electron binding energies.

The absolute emission probabilities of conversion electrons have been obtained using the recommended P_γ and ICC.

The absolute emission probabilities of K- and L- Auger electrons have been deduced from the emission probabilities of KX- and LX- rays measured in 2008De31. Their values, given in Table 4, are compared to the results of calculations using the evaluated P_γ , ICC with the EMISSION computer program.

Table 4. Absolute emission probabilities of K- and L- Auger electrons from the decay of ²³³Th

	Calculated using recommended P_γ , ICC (EMISSION code)	Deduced from absolute intensities of LX-, KX- rays measured in 2008De31	Recommended
e_{AL} (Pa)	6,71 (26)	8,6 (10)	8,6 (10)
e_{AK} (Pa)	0,037 (6)	0,041 (5)	0,041 (5)
KLL	0,022 (4)	0,024 (3)	0,024 (3)
KLX	0,013 (2)	0,014 (2)	0,014 (2)
KXY	0,0020 (3)	0,0021 (3)	0,0021 (3)

β^- average energies have been obtained using the LOGFT computer program.

5. PHOTON EMISSIONS

5.1. X-Ray Emissions

The recommended absolute emission probabilities of Pa KX- and LX- rays are from the measurements of 2008De31, which include a contribution from possible systematic errors in the uncertainties of photon intensities (see section 5.2).

In Tables 5 and 6, a comparison of measured and calculated emission probabilities for specific groups of Pa KX- and LX- rays is given. The calculated values have been obtained with the EMISSION computer program using the adopted atomic data for Pa and the recommended total absolute emission probabilities of K- and L- conversion electrons from ²³³Th → ²³³Pa decay.

Table 5. Experimental and calculated values of absolute Pa KX- ray emission probabilities in the decay of ²³³Th

	Energy (keV)	1969HoZY (measured)	2008De31 (measured)	Calculated	Recommended
K α_2	92,288	0,54 (7)	0,39 (1)	0,357 (20)	0,39 (1)
K α_1	95,869	1,01 (7)	0,615 (10)	0,57 (4)	0,615 (13)
K β'_1	107,60-109,07	0,28	0,235 (5)	0,206 (12)	0,235 (6)
K β'_2	111,40-112,38	0,09	0,079 (3)	0,070 (5)	0,079 (3)

Table 6. Experimental and calculated values of absolute Pa LX- ray emission probabilities in the decay of ²³³Th

	Energy (keV)	2008De31 (measured)	Calculated	Recommended
Ll	11,366	0,14 (2)	0,151 (5)	0,14 (2)
L α	13,122 – 13,291	2,84 (32)	2,48 (7)	2,84 (32)
L η	14,946		0,0626 (20)	
L β	15,3 – 16,7	4,3 (5)	3,07 (8)	4,3 (5)
L γ	19,9 – 21,6	0,95 (11)	0,706 (17)	0,95 (11)

5. 2. Gamma-Ray Emissions

The energies of the gamma-rays $\gamma_{7,5}$ (74 keV), $\gamma_{9,6}$ (109 keV), $\gamma_{8,4}$ (115 keV), $\gamma_{11,6}$ (153 keV), $\gamma_{17,15}$ (179 keV), $\gamma_{10,2}$ (181 keV), $\gamma_{11,3}$ (187 keV), $\gamma_{8,1}$ (195 keV), $\gamma_{17,14}$ (211 keV), $\gamma_{13,10}$ (216 keV), $\gamma_{12,8}$ (246 keV), $\gamma_{11,1}$ (251 keV), $\gamma_{13,8}$ (253 keV), $\gamma_{13,7}$ (285 keV), $\gamma_{15,10}$ (348 keV), $\gamma_{12,4}$ (361 keV), $\gamma_{13,4}$ (368 keV), $\gamma_{12,3}$ (377 keV), $\gamma_{14,4}$ (467 keV), $\gamma_{17,10}$ (527 keV), $\gamma_{17,9}$ (552 keV), $\gamma_{7,8}$ (563 keV), $\gamma_{17,7}$ (595 keV), $\gamma_{18,9}$ (599 keV), $\gamma_{18,7}$ (642 keV), $\gamma_{16,0}$ (670 keV), $\gamma_{17,4}$ (678 keV), $\gamma_{18,6}$ (708 keV), $\gamma_{18,5}$ (717 keV), $\gamma_{18,4}$ (725 keV), $\gamma_{18,3}$ (741 keV), $\gamma_{17,1}$ (758 keV), $\gamma_{19,8}$ (783 keV), $\gamma_{18,1}$ (805 keV), $\gamma_{20,7}$ (849 keV), $\gamma_{19,4}$ (898 keV), $\gamma_{20,3}$ (948 keV), $\gamma_{19,1}$ (978 keV) have been deduced from the adopted ²³³Pa level energies (Table 2).

The energies of the gamma-rays $\gamma_{7,1}$ (162 keV), $\gamma_{7,0}$ (169 keV), $\gamma_{12,11}$ (190 keV), $\gamma_{13,5}$ (360 keV), $\gamma_{12,1}$ (441 keV), $\gamma_{12,0}$ (448 keV), $\gamma_{14,5}$ (459 keV), $\gamma_{15,5}$ (491 keV), $\gamma_{17,5}$ (670 keV) are from precise measurements performed with a crystal spectrometer (1979Bo30).

The following gamma-rays, $\gamma_{6,2}$ (46 keV), $\gamma_{3,0}$ (70 keV), $\gamma_{8,4}$ (115 keV), $\gamma_{10,6}$ (134 keV), $\gamma_{9,3}$ (141 keV), $\gamma_{9,2}$ (155 keV), $\gamma_{10,2}$ (181 keV), $\gamma_{9,0}$ (212 keV), $\gamma_{10,0}$ (238 keV) have not been observed in ²³³Th decay. These gamma-rays have been adopted from the decay scheme on the basis of the available data on ²³⁷Np α -decay.

In Table 7 various experimental energies for a number of prominent gamma-rays in the decay of ²³³Th are compared with evaluated results.

The recommended energies of the remaining gamma-rays are from 1969HoZY, 1972SeZI, 1972Vo08 following the evaluation by 2005Si15. See also 1968Br25, 1968Da24, 1969Va06, 1970Se06 and 1972De67.

Table 7. Experimental and evaluated gamma-ray energies in the decay of ²³³Th

	1976Sk01	1979Bo30	1979Go12	1988Wo01 (Ge detector)	1988Wo01 (LEPS detector)	Evaluated (recommended)
γ _{4,2}	29,373 (10)		29,374 (20)	29,5 (17)	29,18 (21)	29,373 (10)
γ _{6,2}	46,534 (40)		46,53 (6)	46,7 (11)	46,28 (18)	46,53 (4)
γ _{2,0}	57,15 (4)	57,11 (5)	57,104 (20)	57,15 (80)	56,88 (17)	57,10 (2)
γ _{3,0}						70,49 (10) ^{a, b}
γ _{4,0}	86,503 (20)	86,48 (6)	86,477 (10)	86,50 (48)	86,26 (14)	86,477 (10)
γ _{5,1}	88,04 (16)		87,988 (30)			87,99 (3)
γ _{5,0}	94,66 (5)		94,638 (50)			94,65 (5)
γ _{8,4}	115,40 (35)		115,40 (35)			115,14 (5) ^a
γ _{9,5}	117,681 (30)		117,702 (20)	117,72 (50)	117,41 (15)	117,692 (20)
γ _{8,3}	131,043 (30)		131,101 (25)	131,09 (52)	130,62 (15)	131,101 (25)
γ _{10,6}	134,23 (4)		134,285 (20)	134,27 (53)		134,285 (20)
γ _{9,3}			141,74 (10)			141,74 (10)
γ _{10,5}	143,208 (25)		143,249 (20)	143,27 (56)	142,96 (16)	143,230 (20)
γ _{10,4}	151,375 (35)		151,414 (20)	151,42 (60)	151,06 (17)	151,409 (20)
γ _{9,2}	155,22 (4)		155,239 (20)	155,28 (63)		155,239 (20)
γ _{7,1}	162,50 (6)	162,504 (12)	162,41 (8)	162,45 (68)		162,504 (12)
γ _{7,0}	169,17 (5)	169,162 (10)	169,156 (20)	169,18 (73)		169,159 (10)
γ _{11,4}	170,63 (8)		170,59 (6)			170,60 (6)
γ _{10,2}	180,80 (8)		180,81 (10)	180,87 (85)		180,76 (3) ^a
γ _{11,3}	186,8 (5)		186,86 (35)			186,80 (18) ^a
γ _{12,11}		190,552 (14)				190,552 (14)
γ _{8,0}	201,72 (5)		201,62 (5)	201,8 (11)		201,62 (5)
γ _{9,0}	212,415 (25)	212,4 (12)	212,290 (50)			212,34 (5) ^a
γ _{10,0}	238,04 (4)		237,862 (60)	238,0 (14)		237,86 (6)
γ _{13,5}		359,745 (40)				359,74 (4)
γ _{12,1}		440,943 (40)				440,94 (4)

^a deduced from level energies

^b observed by 1969HoXY (71,0 keV) and 1974HeYW (70,75 (10) keV)

The gamma-ray transitions with energies (keV) of 80, 105,147, 211, 242, 310, 383, 409, 418, 454, 465, 474, 497, 505, 513, 517, 532, 554, 555, 579, 583, 681, 690, 698, 704, 728, 745, 752, 767, 774, 784, 832, 847, 871, 874, 919, 935, 942, 943, 955, 961, 963, 968, 994, 1001, 1007, 1011, 1026, 1092, 1132, 1139, 1144 and 1201 have not been placed in the ²³³Th decay scheme.

The gamma-ray transitions $\gamma_{7,1}$ (162 keV) and $\gamma_{11,5}$ (162 keV), $\gamma_{16,0}$ (670 keV) and $\gamma_{17,5}$ (670 keV) are doublets, and have been placed twice in the decay scheme; their intensities have been suitably divided (2005Si15).

The absolute gamma-ray emission probabilities have been adopted from 2008De31. In 2008De31 absolute photon intensities were measured using multiple purifications of stock solutions of ²³³Th produced by the ²³²Th(n, γ) reaction. The measurement consisted of liquid scintillation counting (LSC) and γ -ray spectroscopy with HPGe detectors. As the authors of 2008De31 quoted only statistical uncertainties for their intensity values, the evaluators have considered an additional contribution from possible systematic errors when estimating the overall uncertainties in the absolute photon intensities. This contribution was estimated on the basis of data scattering for LSC measurements and detection efficiency uncertainties for γ -ray spectroscopy discussed in 2008De31 and 2008De10. The estimations of detection uncertainties ($\sim 11\%$ for ≤ 20 keV, $\sim 1\%$ for 29 keV, and $\sim 0,7\%$ for energies ≥ 50 keV) have been adopted from 2008De31, 2008De10 and combined with the statistical uncertainties. In particular, the systematic uncertainty due to the absolute LSC measurements of the effective number of disintegrations in 2008De31 has been estimated as $\sim 1\%$ on the basis of measured data scattering, and it has been used here.

$P(\gamma)(6,65$ keV) has been deduced from the absolute intensity of N1-conversion electrons of 9 (1) per 100 decays measured in 1976JeZU using the theoretical conversion coefficient $\alpha(N1) = 545$ (11) for an M1 multipolarity.

The recommended absolute gamma-ray emission probability for $\gamma_{2,0}(57,1$ keV) (0,0498 (15) %) agrees well with 0,057 (11) % but is much more precise. The latter was deduced from the absolute intensity of L-conversion electrons measured in 1976JeZU and the theoretical ICC (see 2005Si15).

In Table 8 the relative gamma-ray emission probabilities measured in 2008De31 (scaling to 100 for the 57,1-keV γ -ray) are compared to the early experimental results reported without uncertainties. Such a comparison shows that the intensities in 2008De31 for the major transitions are $\sim (10-30)\%$ lower than results in 1969HoZY, 1972SeZI. This may be due to the use of not sufficiently purified ²³³Th samples in the early measurements.

Table 8. Measured relative gamma-ray emission probabilities in the decay of ²³³Th

	Energy (keV)	2008De31	1969HoZY, 1972SeZI (see 2005Si15)
$\gamma_{1,0}$	6,65 (5)	--	29,6
$\gamma_{4,2}$	29,373 (10)	$4,34 (5) \cdot 10^3$	$4,6 \cdot 10^3$
$\gamma_{2,0}$	57,10 (2)	100	100
$\gamma_{3,1}$	63,92 (6)	--	1,5
$\gamma_{3,0}$	70,49 (10)	--	1,5
$\gamma_{7,5}$	74,51 (5)	80,4 (9)	96
$\gamma_{4,0}$	86,477 (10)	$3,68 (4) \cdot 10^3$	$5,0 \cdot 10^3$
$\gamma_{5,1}$	87,99 (3)	340 (4)	333
$\gamma_{5,0}$	94,66 (5)	$1,55 (2) \cdot 10^3$	$1,5 \cdot 10^3$
$\gamma_{9,6}$	108,5 (1)	--	1,1
$\gamma_{8,4}$	115,14 (5)	0,6 (13)	4,1
$\gamma_{9,5}$	117,692 (20)	5,8 (6)	2,8
$\gamma_{8,3}$	131,101 (25)	101 (2)	122
$\gamma_{10,6}$	134,285 (20)	3,6 (9)	4,1
$\gamma_{10,5}$	143,230 (20)	22,8 (15)	26
$\gamma_{10,4}$	151,409 (20)	13,4 (8)	17
$\gamma_{11,6}$	153,49 (18)	81,4 (9)	122
$\gamma_{9,2}$	155,239 (20)	4,6 (1)	1,7
$\gamma_{7,1}$	162,504 (12)	335 (4)	278
$\gamma_{11,5}$	162,504	--	315
$\gamma_{7,0}$	169,162 (10)	502 (6)	630
$\gamma_{11,4}$	170,60 (6)	101 (2)	241
$\gamma_{17,15}$	179,05 (8)	55,6 (7)	70

	Energy (keV)	2008De31	1969HoZY, 1972SeZI (see 2005Si15)
$\gamma_{10,2}$	180,76 (3)	2,2 (6)	1,3
$\gamma_{11,3}$	186,80 (18)	41,8 (13)	63
$\gamma_{12,11}$	190,552 (14)	172 (2)	241
$\gamma_{8,1}$	194,97 (7)	214 (3)	296
$\gamma_{8,0}$	201,62 (5)	44,2 (18)	57
$\gamma_{17,14}$	210,67 (8)	35,6 (18)	65
$\gamma_{-1,4}$	211,3 (2)	40 (2)	35
$\gamma_{9,0}$	212,34 (5)	13 (1)	2,8
$\gamma_{13,10}$	216,54 (8)	26 (3)	28
$\gamma_{18,15}$	226,1 (2)	34 (2)	43
$\gamma_{10,0}$	237,86 (2)	3,8 (8)	3,9
$\gamma_{12,8}$	246,14 (6)	8,2 (14)	--
$\gamma_{11,1}$	250,65 (16)	9,4 (7)	8,7
$\gamma_{13,8}$	252,78 (9)	13,2 (7)	22
$\gamma_{11,0}$	257,30 (15)	105 (2)	126
$\gamma_{12,7}$	278,7 (4)	9,4 (11)	14
$\gamma_{13,7}$	285,24 (7)	31 (2)	39
$\gamma_{15,10}$	347,64 (6)	29 (2)	22
$\gamma_{13,5}$	359,74 (4)	174 (2)	222
$\gamma_{12,4}$	361,285 (22)	43,6 (9)	70
$\gamma_{13,4}$	367,92 (7)	7,4 (15)	8,7
$\gamma_{12,3}$	377,27 (11)	55 (2)	70
$\gamma_{19,15}$	398,8 (5)	22,2 (15)	26
$\gamma_{-1,8}$	408,8 (5)	1 (1)	7,0
$\gamma_{16,11}$	412,5 (5)	16,6 (3)	24
$\gamma_{-1,9}$	418,4 (5)	18,2 (18)	22
$\gamma_{19,14}$	430,9 (4)	35,6 (7)	42
$\gamma_{20,15}$	433,2 (4)	23,4 (8)	28
$\gamma_{12,1}$	440,94 (4)	382 (4)	426
$\gamma_{12,0}$	447,762 (20)	208 (3)	278
$\gamma_{14,5}$	459,222 (7)	1,98 (2)·10 ³	2,6·10 ³
$\gamma_{14,4}$	467,40 (6)	28,8 (9)	33
$\gamma_{-1,12}$	473,9 (5)	6,6 (12)	6,5
$\gamma_{15,5}$	490,80 (6)	215 (4)	315
$\gamma_{-1,13}$	497,1 (4)	25,6 (8)	39
$\gamma_{15,4}$	499,02 (4)	315 (3)	389
$\gamma_{-1,14}$	505,5 (6)	11,0 (6)	9,1
$\gamma_{-1,15}$	513,4 (4)	26,6 (10)	37
$\gamma_{-1,16}$	517,0 (4)	9,2 (1)	13
$\gamma_{17,10}$	526,69 (6)	92,6 (19)	12
$\gamma_{-1,17}$	531,8 (4)	14 (2)	7,8
$\gamma_{17,9}$	552,21 (8)	33 (1)	44
$\gamma_{-1,18}$	554,9 (5)	6,2 (6)	6,5
$\gamma_{17,8}$	562,93 (8)	109 (2)	130
$\gamma_{18,10}$	573,7 (4)	66,4 (14)	78
$\gamma_{17,7}$	595,39 (6)	236 (20)	296
$\gamma_{18,9}$	599,3 (2)	58,8 (12)	87
$\gamma_{18,8}$	610,0 (3)	113 (2)	157
$\gamma_{18,7}$	642,4 (2)	40,4 (8)	52

	Energy (keV)	2008De31	1969HoZY, 1972SeZI (see 2005Si15)
$\gamma_{16,1}$	663,3 (5)	7,4 (10)	4,4
$\gamma_{17,5}$	669,901 (16)	1,00 (3)·10 ³	1,3·10 ³
$\gamma_{17,4}$	678,04 (10)	129 (2)	161
$\gamma_{-1,22}$	681,2 (6)	28,6 (8)	30
$\gamma_{-1,23}$	698,5 (6)	21,2 (1)	22
$\gamma_{-1,24}$	703,7 (6)	18,2 (1)	20
$\gamma_{18,6}$	707,8 (3)	18,2 (1)	22
$\gamma_{18,5}$	717,0 (2)	84,2 (17)	104
$\gamma_{18,4}$	725,1 (2)	126 (2)	161
$\gamma_{18,3}$	741,1 (2)	47,2 (9)	57
$\gamma_{-1,27}$	744,9 (5)	10,6 (4)	13
$\gamma_{-1,28}$	751,6 (6)	4,6 (6)	4,4
$\gamma_{17,1}$	757,90 (7)	64,8 (13)	78
$\gamma_{17,0}$	764,55 (6)	178 (2)	222
$\gamma_{-1,30}$	774,0 (4)	21,6 (12)	26
$\gamma_{19,8}$	783,2 (5)	11,2 (7)	11
$\gamma_{-1,31}$	784,2 (5)	4,4 (6)	9,1
$\gamma_{18,1}$	805,0 (2)	42,8 (13)	57
$\gamma_{20,9}$	806,4 (5)	24,6 (14)	24
$\gamma_{18,0}$	811,6 (2)	12,0 (5)	14
$\gamma_{19,7}$	815,9 (4)	39 (2)	52
$\gamma_{20,8}$	817,0 (6)	19 (1)	30
$\gamma_{20,7}$	849,5 (5)	7,8 (5)	8,7
$\gamma_{-1,34}$	870,7 (7)	6,2 (5)	3,9
$\gamma_{-1,35}$	874,0 (5)	24,0 (8)	11
$\gamma_{19,6}$	880,9 (5)	19,4 (8)	14
$\gamma_{19,5}$	890,1 (5)	210 (8)	259
$\gamma_{19,4}$	898,3 (5)	4,4	6,1
$\gamma_{-1,37}$	935,2 (7)	73,8 (15)	91
$\gamma_{-1,38}$	941,9 (8)	9,6 (1)	14
$\gamma_{20,3}$	948,3 (5)	12,0 (7)	14
$\gamma_{-1,40}$	955 (1)	0,4 (5)	10
$\gamma_{-1,41}$	960,8 (8)	8,2 (3)	13
$\gamma_{-1,42}$	962,8 (9)	3,0 (1)	2,6
$\gamma_{-1,43}$	968,2 (9)	16,6 (7)	20
$\gamma_{19,1}$	978,2 (5)	11,6 (7)	14
$\gamma_{19,0}$	984,8 (5)	20,4 (6)	2,6
$\gamma_{-1,44}$	994 (1)	1,2 (2)	1,7
$\gamma_{-1,45}$	1001 (1)	1,6 (4)	2,2
$\gamma_{-1,46}$	1007 (1)	2,8 (4)	5,2
$\gamma_{-1,47}$	1011 (1)	3,8 (4)	7,4

6. CONSISTENCY OF RECOMMENDED DATA

The most accurate Q value, Q(M), is taken from the atomic mass adjustment table of Audi et al. (2003Au03). Comparison of Q(eff)(deduced as the sum of average energies per disintegration ($\sum E_i \times P_i$) for all emissions accompanying 233Th β - decay) with the tabulated decay energy Q(M) allows to check a consistency of the recommended decay-scheme parameters obtained in this evaluation.

Here E_i and P_i are the evaluated energies and emission probabilities of the i-th alpha particle, beta particle, gamma ray, X-ray, etc..

Consistency (percentage deviation) is determined by $\{[Q(M) - Q(\text{eff})] / Q(M)\} \times 100$. “Percentage deviations above 5 % would be regarded as high and imply a poorly defined decay scheme; a value of less than 5 % indicates the construction of a reasonably consistent decay scheme” (quoted from the article by A.L. Nichols in Appl. Rad. Isotopes 55 (2001) 23-70).

For the above ²³³Th decay data evaluation we have $Q(M) = 1243,1 (14) \text{ keV}$ and $Q(\text{eff}) = 1247 (2) \text{ keV}$, i.e. consistency is better than 1 %.

7. REFERENCES

- 1952Ru10 W.C. Rutledge, J.M. Cork, and S.B. Burson, Phys. Rev. 86, 775 (1952)
(Half-life)
- 1955Je26 E.N. Jenkins, Analyst 80, 301 (1955)
(Half-life)
- 1957Dr46 B.J. Dropesky and L.M. Langer, Phys. Rev. 108, 90 (1957)
(Half-life, energy of $\beta_{0,0}$ -transition)
- 1957Fr55 M.S. Freedman, D.W. Engelkemeir, F.T. Porter, F. Wagner, Jr., and P. Day, Priv. Comm., unpublished (1957), quoted in 1967Le24: C.M. Lederer, J.M. Hollander, and I. Perlman, Table of Isotopes, Sixth Edition, John Wiley and Sons, Inc., New York (1967)
(Gamma ray emission probabilities, β -transition energies)
- 1968Br25 E. Browne and F. Asaro, UCRL-17989, p 1. (1968)
(Gamma-ray energies)
- 1968Da24 R. Dams and F. Adams, Radiochim. Acta 10, 1(1968)
(Gamma-ray energies)
- 1969HoZY W. Hoekstra, Thesis, Technische Hogeschool, Delft. (1969)
(Half-life, KX- - ray emission probabilities , gamma - ray relative probabilities)
- 1969Va06 J.M. Vara and R. Gaeta, Nucl. Phys. A130, 586 (1969)
(Gamma-ray energies)
- 1970Se06 C. Sebillle, G. Bastin, C.F. Leang, R. Piepenbring, and M.F. Perrin, Compt. Rend. 270A, 354 (1970)
(Gamma-ray energies)
- 1972De67 M. de Bruin and P.J.M. Korthoven, , J. Radioanal. Chem. 10, 125. (1972):
(Gamma-ray energies)
- 1972SeZI C. Sebillle-Schuck, Thesis, Paris Univ. (1972); FRNC-TH-255 (1972)
(Gamma - ray relative probabilities, gamma-ray multipolarities, conversion electron characteristics)
- 1972Vo08 T von Egidy, O.W.B. Schult, D. Rabenstein, J.R. Erskine, O.A. Wasson, R.E. Chrien, D. Breitig, R.P. Sharma, H.A. Baader, and H.R. Koch, Phys. Rev. C6, 266(1972)
(Gamma-ray energies)
- 1976JeZU P. Jeuch, Thesis, Tech. Univ. Munchen. (1976)
(Gamma-ray multipolarities, conversion electron characteristics)
- 1976Sk01 M. Skalsey and R.D. Connor, Can. J. Phys. 54, 1409 (1976)
(Gamma-ray energies)
- 1979Bo30 H.G. Borner, G. Barreau, W.F. Davidson, P. Jeuch, T. von Egidy, J. Almeida, and D.H. White, Nucl. Instrum. Methods 166, 251 (1979)
(Gamma-ray energies)
- 1979Go12 L. Gonzalez, R. Gaeta, E. Vano, and J.M. Los Arcos, Nucl. Phys. A324, 126 (1979)
(Gamma-ray energies)
- 1988Wo01 S.A. Woods, P. Christmas, P. Cross, S.M. Judge, and W. Gelletly, Nucl. Instrum. Methods Phys. Res. A264, 333 (1988); Addendum Nucl. Instrum. Methods Phys. Res. A272, 924 (1988)
(Gamma ray energies, ICC for $\gamma_{4,0}$)
- 1989Ab05 A. Abzouzi, M.S. Antony, and V.B. Ndocko Ndongue, J. Radioanal. Nucl. Chem. 135, 1 (1989)
(Half-life)
- 1998Us01 K. Usman, T.D. Macmahon, and S.I. Kafala, Appl. Radiat. Isot. 49, 1329 (1998)
(Half-life)

- 2002Be M.M. Bé, R. Helmer, V. Chisté, J. Nucl. Sci. Tech., suppl.2, 481 (2002)
(SAISINUC software)
- 2003Au03 G. Audi, A.H. Wapstra, and C. Thibault, Nucl. Phys. A729, 337 (2003)
(Q value)
- 2005Si15 B. Singh and J. K. Tuli, Nuclear Data Sheets105, 109 (2005)
(Decay data evaluation, decay scheme, ²³³Pa level energies, multipolarities)
- 2008De10 D.J. DeVries and H.C. Griffin, Appl. Rad. Isotop., 66, 668 (2008)
(Uncertainties of absolute photon emission probabilities)
- 2008De31 D.J. DeVries and H.C. Griffin, Appl. Radiat. Isot. 66, 1999 (2008)
(Absolute and relative gamma ray and X-ray emission probabilities)
- 2008Go10 V.M. Gorozhankin and M.M. Bé, Appl. Radiat. Isot. 66, 722 (2008)
(ICC for anomalous E1 gamma-ray transitions)
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, and C.W.Nestor, Jr., Nucl.
Instrum. Methods Phys. Res. A589, 202 (2008)
(Theoretical ICC)

²³³Pa - Comments on evaluation of decay data by V. P. Chechev and K. N. Kuzmenko

This evaluation was done originally in September 2004, corrected in December 2004 and in March 2006, and then updated in April 2009 with a literature cut-off by the same date.

1 Decay Scheme

The decay scheme is based on the experimental results of Kouassi et al. (1990Ko41) and NDS evaluations of 1990Ak02 and 2005Si15. In addition to the nuclear transitions well studied (1990Ak02), Singh and Tuli (2005Si15) list a large number of weak transitions and γ rays from unpublished work of de Bettencourt (1985DeZR), defining them as tentative. Latter ones have not been considered in this evaluation. The list of the tentative gamma rays is given in section 5.2.1. These gamma rays carry $\leq 1,2$ % of the total intensity of all the gamma rays placed in the decay scheme.

2 Nuclear Data

Q^- value is from 2003Au03.

The recommended half-life of ²³³Pa is based on the experimental results given in Table 1.

Table 1. Experimental values of ²³³Pa half-life (in days)

Reference	Author(s)	Value	Measurement method
1941Gr03	Grosse et al.	27,4 (4)	β -counting
1956Mc60	Mc Isaac and Freiling	27,0 (1)	$4\pi\gamma$ ionization chamber (4 $T_{1/2}$) and β proportional counter (2 $T_{1/2}$)
1957Wr37	Wright et al.	26,95 (6)	Gamma ionization chamber and β proportional counter (2 $T_{1/2}$)
1986Jo07	Jones et al.	26,967 (2)	Gamma ionization chamber (11 $T_{1/2}$)
1999Popov	Popov and Timofeev	26,9 (1)	Ge(Li) γ -ray spectrometer
2000Us01	Usman and MacMahon	27,02 (3)	HPGe gamma-ray spectrometer (8 gamma lines, 5 $T_{1/2}$)

The weighted mean of the six values from Table 1 of 26,967(2) is dominated by the very accurate value from 1986Jo07. The LWEIGHT computer program uses a limitation of relative statistical weights (LRSW method), and increased the uncertainty of 1986Jo07 from 0,002 to 0,025 to give a weighted mean of 26,984(18). The evaluation technique from 2000Ch01 also uses the LRSW method and some additional criteria to give the same value of 26,984 (18). The Rajeval data evaluation technique (1992Ra08) uses different criteria to adjust the uncertainties, and has increased the uncertainties of 1986Jo07 and 2000Us01 to give the same value of 26,984 (18).

Huang et al. (2005Hu06) used the analogous procedures for their statistical analysis, and adopted the mean of the normalized residuals and Rajeval technique to give the value for the ²³³Pa half-life of 26,971 (13) d. However, they did not take into account the measurement of Popov and Timofeev (1999).

Thus, taking into account the accuracy of most of the available measurements, the best estimate of the ²³³Pa half-life is believed to be a recommended value of 26,98 (2) days.

2.1 b- Transitions

The energies of β^- transitions have been obtained from the Q^- value and the ²³³U level energies given in Table 2 from 2005Si15.

Table 2. ²³³U levels populated in ²³³Pa β^- -decay

Level	Energy (keV)	Spin and Parity	Half-life	Probability of β^- transitions (%)
0	0,0	5/2 ⁺	1,592 (2) × 10 ⁵ a	6,3 (23)
1	40,350 (4)	7/2 ⁺	0,11 (8) ns	0,3 (19)
2	92,16 (4)	9/2 ⁺		
3	298,810 (4)	5/2 ⁻		0,12 (5)
4	301,94 (9)	5/2 ⁻		0,010 (2)
5	311,904 (4)	3/2 ⁺	0,120 (15) ns	26,6 (32)
6	320,83 (4)	7/2 ⁻		0,020 (3)
7	340,477 (4)	5/2 ⁺	52 (10) ps	25,9 (32)
8	380,43 (8)	7/2 ⁺		0,020 (3)
9	398,496 (4)	1/2 ⁺	55 (20) ps	15,4 (8)
10	415,758 (4)	3/2 ⁺	≤ 30 ps	25,4 (16)
11	456,114 (6)	5/2 ⁺		0,0011 (2)

The recommended probabilities of β^- -transitions have been deduced from the P($\gamma+ce$) balance at each level of ²³³U.

The accurate sum of intensities of β^- -transitions to the ground and first excited states [$P(\beta_{0,0})+P(\beta_{0,1})$]×100 has been deduced as (100 % – $\Sigma P_{i,j}(\gamma+ce)(j=0,1,2)$), where the latter value includes only the intensities of the gamma-ray transitions feeding the ground state and the 40,3- and 92,2-keV levels. The 92,2-keV level (9/2⁺) cannot be fed directly in the β^- decay of ²³³Pa ground state (3/2⁻). This forbiddenness allows the accurate combine β^- intensity of the $\beta_{0,0}$ and $\beta_{0,1}$ transitions to be evaluated as 100% – 93,4 (22) % = 6,6 (22) % to be compared with a value of 8,8 (14) % as measured by Browne et al. (1989Br24) and deduced from the decay scheme in 1990Ko41 (6,9 (15) %) and in 2005Hu06 (7,4 (6) %), respectively.

Measured and recommended β^- -transition energies and probabilities are given in Tables 3 and 4, respectively.

Table 3. Measured and recommended energies of β^- transitions (keV)

	1954Br37	1955On05	1960Un01	1963Bj03	Recommended
$\beta_{0,10}$	140 (14)	145 (10)	155 (7)	154 (5)	154,3 (20)
$\beta_{0,9}$			175 (8)		171,5 (20)
$\beta_{0,5}$	256 (4)	257 (5)	250 (5)	254 (5)	258,2 (20)
$\beta_{0,0}$	568 (5)	568 (5)		578 (10)	570,1 (20)

Table 4. Measured and evaluated probabilities (%) of β^- transitions

	1954Br37	1955On05	1963Bj03	Evaluated
$\beta_{0,10}$	50	37	32	25,4 (16)
$\beta_{0,5}$	45	58	56	26,6 (32)
$\beta_{0,0}$	5	5	12	6,3 (23)

2.2 Gamma-Ray Transitions and Internal Conversion Coefficients

The recommended energies of gamma-ray transitions are virtually the same as the gamma-ray energies because nuclear recoil is negligible.

The gamma-ray transition probabilities have been obtained from the gamma-ray emission probabilities and the total internal conversion coefficients (ICCs). Multipolarities of gamma-ray transitions have been taken from 2005Si15. The ICCs have been interpolated using the BrIcc package with the so called “Frozen Orbital” approximation (2008Ki07). The relative uncertainties of α_K , α_L , α_M , α_T for pure multipolarities have been taken as 2 %.

E2 admixtures for M1+E2 gamma -ray transitions in ²³³Pa decay were determined in a whole number of measurements (see Table 5).

For the ICC evaluation the E2 admixture of Q0166 from 1962Sc03 has been used for the gamma ray of 17,2 keV ($\gamma_{10,9}$). The best set of E2/M1 mixing ratios obtained by Krane (1986Kr10) from the angular correlation measurements has been used to determine the ICCs for the gamma rays of 28,6 -keV ($\gamma_{7,5}$), 75,3-keV ($\gamma_{10,7}$), 86,6-keV ($\gamma_{9,5}$), and 103,9-keV($\gamma_{10,5}$), respectively. This set agrees mainly with the conversion electron data from 1961Al19, 1962Sc03, 1963Bi03, 1966Ze02, 1973Va33, 1985DeZR, 1988Wo01, and 1990Pe16 (Table 5). Use of the BrIcc package for correction of the above conversion electron data does not change this conclusion.

The evaluators have adopted the value of 0,54 (4) from 1962Sc03 for the 40,3 -keV gamma ray ($\gamma_{1,0}$) E2 admixture that coincides with the values obtained by Zender (1966Ze02) and Krane (1986Kr10). This ratio produces a better P(γ +ce) balance for the 40,3-keV level (“1”). If the smaller value of 0,3 reported by Albridge et al. (1961 Al19) and Bisgard et al. (1963 Bi03) was used, the intensity of the β^- transition to the level “1” would have been negative (2006Ch39).

The ICC values measured by Browne et al. (1989Br24) have been adopted for the most intense, predominantly M1, 300,1- ($\gamma_{7,1}$), 311,9- ($\gamma_{5,0}$), and 340,5-keV ($\gamma_{7,0}$) transitions affected by nuclear penetration effects.

The E2/M1 mixing ratio $\delta \approx 0,62$ has been taken from 2005Si15 for the gamma ray of 51,8-keV ($\gamma_{2,1}$).

Table 5. Experimental and recommended E2 γ -ray admixtures

E γ (keV)	1961 Al19	1962 Sc03	1963 Bi03	1966 Ze02	1986 Kr10	1973 Va33	1985 DeZR	1988 Wo01	1990 Pe16	Recommended admixture & δ
28,6	0,030 (5)	0,0102 (8)	0,02 (1)	0,024 (2)	0,0244 (15)	0,02	0,019 (2)	0,03 (1)	0,022 (3)	0,0244 (15) δ 0,158 (10)
40,3	0,30 (10)	0,54 (4)	0,31 (2)	0,54 (5)	0,54 (8)	0,43	0,46 (5)			0,54 (4) δ 1,08 (12)
75,3	0,01 (1)	< 0,0005	0	< 0,005	0,022 (16)	0	0,008 (4)	0		0,022 (16) δ 0,15 (8)
86,6	0,020 (5)	< 0,002	0,01 (1)	< 0,006	0,0031 (3)	0,01	0,0049 (7)	0,046 (27)		0,0031 (3) δ 0,056 (5)
103,9	0,04 (1)	< 0,03	0,01 (1)	0,020 (15)	0,010 (14)	0,01	0,022 (2)	0,073 (9)		0,010 (14) δ 0,1 (1)
300,1	0,12 (10)	0,03	0	0	0,006 (2)	0	0,025 (3)	0		0,006 (2) δ 0,08 (3)
311,9	< 0,02	< 0,03	0	< 0,016	0,010 (1)	0	0,063 (6)	0		0,010 (1) δ 0,10 (1)
415,8	0,82 (7)	0,96 (4)	0,78 (11)	0,76 (8)		0,84				0,83 (7) δ 2,2 (9)

^a Weighted average of 1961Al19, 1962Sc03, 1963Bi03, and 1966Ze02.

3 Atomic Data

The atomic data are from Schönfeld (1996Sc06).

4 Electron Emissions

The energies of the conversion electrons have been obtained from the gamma transition energies and the electron binding energies.

The emission probabilities of the conversion electrons have been deduced using evaluated P_{γ} and ICC values.

The absolute emission probabilities of K Auger electrons have been deduced using the $P(c_K)$ values and the adopted ω_K given in section 3.

The absolute emission probabilities of L Auger electrons have been deduced using the $P(c_L)$ and $P(c_{eL})$ values and the adopted ω_L , n_{KL} given in section 3.

β^- average energies have been calculated using the LOGFT computer program.

5 Photon emissions

5.1 X-ray emissions

The absolute emission probabilities of U KX-rays have been deduced using the adopted value of ω_K (U) and the total evaluated absolute emission probability of K conversion electrons in the $^{233}\text{Pa} \rightarrow ^{233}\text{U}$ decay. In Table 6 the deduced values are compared to measured results. The total absolute U KX-ray emission probability of 30,7 (9) % agrees with the experimental values of 29,3 (28) % from 2008De10 and 30,0 (4) % from 2004Sh07.

The absolute emission probabilities of U LX-rays have been calculated with the EMISSION computer program using the adopted values of ω_L (U), ω_K (U), n_{KL} (U) and the evaluated absolute emission probabilities of L_1 , L_2 , L_3 , and K- conversion electrons in ^{233}Pa β^- -decay.

As authors of 2008De10 quote only statistical uncertainties for their intensity values, the evaluators have considered additionally a contribution of possible systematic errors to obtain the overall uncertainties in the absolute photon intensities. This contribution was estimated on the basis of detection efficiency uncertainties for γ -ray spectroscopy discussed in 2008De10: 1 % for U KX-ray emission probabilities and 20 % for U LX-ray emission probabilities.

Table 6. Experimental and recommended (calculated) absolute U KX- ray emission probabilities in decay of ²³³Pa

	Energy (keV)	1979 Ge08	1984Va28	1990Ko41	2000 Smith	2000Sc04	2002Lu01	2004Sh07	2008De10	Recommended (calculated)
Kα ₂	94,666	8,8 (5)	8,8 (4)	8,3 (4)		8,78 (10)	8,77 (9)	8,78 (10)	8,50 (14)	9,10 (26)
Kα ₁	98,440	14,4 (8)	14,3 (5)	13,4 (7)	14,3 (3)	14,4 (4)	14,17 (14)	14,22 (17)	14,02 (24)	14,6 (4)
Kβ ₃	110,421	} 5,2 (3)	1,78 (9)	1,60 (8)	1,89 (4)	1,90 (5)	1,708 (25)	1,708 (38)	1,694 (31)	
Kβ ₁	111,298		3,27 (15)	3,11 (15)			3,35 (5)	3,34 (8)	3,24 (6)	} 5,25 (18)
Kβ ₅	111,964			0,15 (1)			0,1230 (17)	0,1239 (28)	0,139 (18)	
Kβ ₂	114,407 (11)			0,52 (7)		} 1,59 (9)	1,293 (23)	1,34 (5)	} 1,317 (21)	} 1,49 (10)
Kβ ₄	115,012	} 1,74 (8)	} 1,71 (8)	0,78 (7)	} 1,73 (8)		0,0380 (6)	0,0388 (13)		
KO _{2,3}	115,377						} 0,39 (2)	0,332 (10)	} 0,391 (9)	} 0,399 (18)
KP _{2,3}	115,580									

Table 7. Experimental and recommended (calculated) absolute U XL- ray emission probabilities in decay of ²³³Pa

	Energy (keV)	2000Sc04	2004Sh07	2008De10	Recommended (calculated)
L1	11,62	1,18 (7)	0,78 (11)	1,19 (25)	1,05 (4)
Lα	13,93	15,7 (7)	12,7 (13)	21,5 (43)	16,9 (6)
Lβ	15,73-17,45			16,9 (34)	18,1 (6)
Lγ ₁	20,17-20,84			3,2 (6)	2,25 (13)

5.2 Gamma-ray emissions

The energies of gamma rays have been taken from 1990Ko41 (see also 2005Si15) except for $\gamma_{10,9}$ (17,26 keV) and $\gamma_{2,0}$ (92,16 keV) which were deduced from the adopted ²³³U levels. A comparison of the recommended γ -ray energies with early experimental results is given in Table 8.

In Table 9 the experimental and recommended absolute gamma ray emission probabilities (P) are presented. All the values given in Table 8 given in Table 9 are absolute measurement results (per 100 disintegrations).

The original values from 1973Va33 and 1990Ko41 have been renormalized by the evaluators to $P_{\gamma_{2,0}}$ (311,9 keV) = 38,3 (5). Values given in the last column are weighted averages (LRSW) of individual results taking into account the LRSW procedure and sometimes increasing the uncertainty to cover the most precise input value (2006Ch39).

$P_{\gamma_{10,9}}$ (17,26 keV) = 0,0041 has been deduced from the value of $P_{ce}(M1) = 0,0054$ (1962Sc03) and the ICC value of $\alpha_{M1} = 132,3$ (1993Ba60) calculated for this conversion line using an E2/M1 admixture of 0,016 ($\delta=0,13$) from 1990Ak02.

$P_{\gamma_{2,1}}$ (51,8 keV) and $P_{\gamma_{2,0}}$ (92,2 keV) have been obtained from the $P_{\gamma+ce}$ balance at the 92,2-keV level and the ratio $P_{\gamma_{2,1}}/P_{\gamma_{2,0}} = 0,21$ (4) taken from 1990Ak02.

The contribution of 1 % estimated on the basis of detection efficiency uncertainty for γ -ray spectroscopy discussed in 2008De10 has been added to the overall uncertainties for the recommended γ -ray emission probabilities.

Table 8. Experimental and recommended gamma-ray energies in decay of ²³³Pa, in keV

	1952Br84	1961Al19	1967Br20	1968Ma13	1971Vo02	1972De67	1973Va33	1988Wo01	1990Ko41	Recommended
$\gamma_{10,9}$						17,2 (1)				17,262 (6)
$\gamma_{7,5}$	28,67 (2)					28,6 (1)		28,375 (5)	28,559 (10)	28,559 (10)
$\gamma_{1,0}$	40,47 (10)	40,35 (1)				40,5 (1)			40,349 (5)	40,349 (5)
$\gamma_{7,3}$		41,65 (2)							41,663 (10)	41,663 (10)
$\gamma_{10,7}$	75,4 (2)	75,28 (1)				75,27 (3)		75,354 (4)	75,269 (10)	75,269 (10)
$\gamma_{9,5}$	87,0 (3)	86,59 (1)				86,58 (3)		86,814 (3)	86,595 (10)	86,595 (5)
$\gamma_{2,0}$							92,0 (5)		92,1 (5)	92,16 (4)
$\gamma_{10,5}$		103,86 (2)						103,971 (9)	103,860 (10)	103,860 (10)
$\gamma_{6,2}$									228,57 (5)	228,57 (5)
$\gamma_{7,2}$			248,3 (3)	248,69 (24)			248,0 (2)		248,38 (4)	248,38 (4)
$\gamma_{3,1}$							258,292)		258,45 (2)	258,45 (2)
$\gamma_{5,1}$		271,62 (23)			271,48 (8)				271,555 (10)	271,555 (10)
$\gamma_{6,1}$									280,61 (5)	280,61 (5)
$\gamma_{8,2}$									288,42 (10)	288,42 (10)
$\gamma_{3,0}$									298,81 (2)	298,81 (2)
$\gamma_{7,1}$		300,20 (24)						300,34 (2)	300,129 (5)	300,129 (5)
$\gamma_{4,0}$									301,99 (10)	301,99 (10)
$\gamma_{5,0}$		311,91 (13)						312,17 (12)	311,904 (5)	311,904 (5)

	1952Br84	1961Al19	1967Br20	1968Ma13	1971Vo02	1972De67	1973Va33	1988Wo01	1990Ko41	Recommended
$\gamma_{6,0}$									320,73 (10)	320,73 (10)
$\gamma_{7,0}$		340,51 (18)						340,81 (3)	340,476 (5)	340,476 (5)
$\gamma_{10,1}$		375,35 (32)			375,45 (4)				375,404 (5)	375,404 (5)
$\gamma_{8,0}$									380,28 (10)	380,28 (10)
$\gamma_{9,0}$		398,57 (40)			398,62 (8)				398,492 (5)	398,492 (5)
$\gamma_{10,0}$		415,87 (42)			415,76 (4)				415,764 (5)	415,764 (5)
$\gamma_{11,0}$									455,96 (10)	455,96 (10)

Table 9. Experimental and recommended absolute gamma-ray emission probabilities (%) in decay of ²³³Pa, in %.

E γ (keV)	1973Va33	1978Poenitz	1979Ge08	1984Va27	1985DeZR	1988Wo01	1990Ko41	2000Wo01	2000Sc04	2002Lu01 2000Lu01	2004Sh07	2006Ha53	2008De10	Recommended
17,2														
28,56	0,069 (8)			0,15 (1)	0,096 (35)	0,068 (9)	0,074 (8)			0,034 (10)	0,019 (2)			0,071 (8) ^a
40,35	0,039 (8)						0,024 (4)	0,0215 (16)		0,028 (4)	0,032 (4)			0,024 (2)
41,66	0,013 (4)						0,014 (3)							0,014 (3)
75,27	1,25 (8)		1,39 (8)	1,30 (4)		1,25 (9)	1,25 (9)	1,401 (25)	1,38 (4)	1,270 (8)				1,30 (3)
86,60	1,87 (23)		1,97 (12)			1,87 (25)	1,93 (11)			2,61 (23)				1,99 (10)
92,1	< 0,004						< 0,002							0,002 (1)
103,86	0,73 (8)		0,87 (3)	0,87 (3)		0,73 (9)	0,847 (60)	0,853 (8)	0,844 (17)	0,855 (6)	0,825 (25)			0,853 (6)
228,57					0,0058 (8)		0,0042 (7)							0,0042 (7)

E _γ (keV)	1973Va33	1978Poenitz	1979Ge08	1984Va27	1985DeZR	1988Wo01	1990Ko41	2000Wo01	2000Sc04	2002Lu01 2000Lu01	2004Sh07	2006Ha53	2008De10	Recommended
248,38	0,0039 (12)		0,059 (2)	0,06 (1)			0,058 (4)	0,0607 (12)	0,0618 (11)	0,057 (6)				0,0609 (11)
258,45	0,0039 (16)				0,031 (4)		0,027 (2)		0,0274 (6)					0,0274 (6)
271,56	0,30 (3)		0,33 (1)	0,32 (1)			0,334 (17)	0,3227 (29)	0,323 (4)	0,323 (5)	0,290 (56)			0,323 (3)
280,61					0,0116 (13)		0,011 (2)							0,011 (2)
288,42					0,0164 (5)		0,016 (3)							0,016 (3)
298,81	0,035						0,085 (7)			0,147 (29)				0,12 (5)
300,13	6,57 (31)		6,62 (10)	6,64 (6)		6,57 (46)	6,76 (7)	6,66 (6)	6,55 (7)	6,39 (6)			6,47 (8)	6,60 (21)
301,99					0,027 (4)		0,010 (2)							0,010 (2)
311,90		38,6 (15)	38,6 (5)	38,65 (39)				38,7 (4)	38,5 (4)	37,80 (23)	37,5 (24)	41,6 (9)	38,08 (51)	38,3 (5)
320,73					0,039 (12)		0,0051 (4)							0,0051 (4)
340,48	4,47 (46)		4,47 (6)	4,52 (5)		4,48 (51)		4,52 (4)	4,50 (5)	4,41 (3)	4,36 (44)		4,436 (56)	4,47 (3)
375,40			0,68 (1)	0,69 (1)				0,690 (6)	0,686 (7)	0,687 (6)	0,58 (8)			0,684 (7)
380,28					0,0039 (8)		0,0037 (9)							0,0037 (9)
398,49			1,39 (2)	1,43 (2)				1,407 (11)	1,406 (15)	1,39 (1)	1,33 (10)			1,408 (14)
415,76			1,74 (2)	1,74 (2)				1,771 (14)	1,765 (18)	1,740 (7)	1,59 (10)		1,724 (23)	1,747 (7)
455,96							0,0011 (2)							0,0011 (2)

^a Weighted average of the values from 1988Wo01 and 1990Ko41 (see discussion in 2006Ch39).

5.2.1 Tentative gamma-ray

This section is given only for information on measurements done in the thesis of 1985DeZR. These results require confirmation and do not consider for evaluation (as well as by Singh and Tuli in Nucl. Data Sheets (2005Si05)).

Energy (keV)	P γ (%)	Level energy (keV)
18,7 (2)	0,023 (8)	320,83
18,7 (2)	0,023 (8)	330,67
19,7 (2)	0,046 (15)	340,478
22,0 (3)		320,83
23,9 (2)	0,0031 (12)	344,56 ?
24,7 (2)	0,0031 (12)	571,36 ?
28,7 (1)		330,67
31,9 (2)	0,0023 (8)	330,67
35,3 (2)	0,0015 (4)	432,81 ?
35,8 (2)	0,0019 (8)	380,48
38,5 (2)	0,0032 (12)	340,478
38,5 (2)	0,0031 (12)	392,25 ?
39,9 (3)		380,48
40,4 (1)		432,81 ?
40,4 (1)		494,75 ?
40,7 (3)		456,113
40,7 (3)		496,65 ?
41,7 (1)	< 0,019	432,81 ?
42,7 (2)	0,0019 (8)	344,56 ?
45,8 (2)	\approx 0,0004	344,56 ?
46,7 (2)	\approx 0,0008	391,09 ?
47,7 (2)	\approx 0,0008	392,25 ?
48,8 (2)	\approx 0,0008	393,33 ?
48,8 (2)	\approx 0,0008	475,69 ?
49,7 (2)	\approx 0,0008	380,48
51,8 (2)	0,0012 (4)	353,71 ?
51,8 (2)	0,0012 (4)	392,25 ?
52,5 (2)	\approx 0,0008	393,33 ?
52,5 (2)	\approx 0,0008	432,81 ?
53,2 (2)	0,0012 (4)	397,71 ?
55,0 (2)	\approx 0,0008	353,71 ?
58,0 (2)	0,0012 (4)	398,495
59,2 (2)	\approx 0,0008	563,00 ?
59,6 (2)	0,0008 (8)	380,48
60,6 (2)	\approx 0,008	391,09 ?
60,6 (2)	\approx 0,008	414,37 ?
60,6 (2)	\approx 0,008	441,20 ?
61,6	\approx 0,0008	392,25 ?
61,6	\approx 0,0008	494,75 ?
63,2 (2)	\approx 0,0004	155,35 ?
63,2 (2)	\approx 0,0004	454,29 ?
63,6 (2)	\approx 0,0008	496,65 ?
65,5 (2)	0,0027 (8)	410,13 ?
66,4 (2)	0,0019 (8)	570,27 ?
67,5 (2)	0,0019 (8)	571,36 ?
68,5 (2)	0,0027 (8)	380,48

Energy (keV)	P γ (%)	Level energy (keV)
69,6 (2)	0,0046 (12)	410,13 ?
70,3 (2)	0,0027 (8)	391,09 ?
71,3 (2)	0,0039 (12)	392,25 ?
71,3 (2)	0,0039 (12)	565,90 ?
74,0 (2)	0,0035 (8)	414,37 ?
74,4 (2)		229,79 ?
75,3 (1)		456,113
75,3 (1)		473,04 ?
77,0 (2)	0,0019 (8)	388,68 ?
77,0 (2)	0,0019 (8)	397,71 ?
77,9 (2)	\approx 0,0012	475,69 ?
78,4 (2)	0,0077 (19)	380,48
79,1 (3)	\approx 0,0008	432,81 ?
80,8 (2)	0,0015 (4)	473,04 ?
81,8 (2)	0,0015 (4)	380,48
81,8 (2)	0,0015 (4)	393,33 ?
81,8 (2)	0,0015 (4)	473,04 ?
82,5 (2)	0,0015 (4)	427,08 ?
84,8 (2)	< 0,0131	475,69 ?
85,2	< 0,0131	315,06 ?
85,2	< 0,0131	415,758
86,6 (1)		388,68 ?
86,6 (1)		427,08 ?
87,5 (3)		441,20 ?
89,0 (3)	< 0,0147	391,09 ?
89,3 (3)	< 0,0147	410,13 ?
90,0 (2)	0,0012 (4)	388,68 ?
91,0 (2)	0,0012 (4)	546,83
91,5 (2)	0,0012 (4)	393,33 ?
92,2 (2)	0,0035 (12)	391,09 ?
92,5 (2)		432,81 ?
92,7 (2)	0,0012 (8)	473,04 ?
93,0 (2)	< 0,0015	565,90 ?
93,5 (2)	< 0,0015	414,37 ?
94,5 (3)		393,33 ?
94,5 (3)		570,27 ?
95,3 (3)		475,69 ?
95,3 (3)		571,36 ?
96,7 (2)	0,0040 (12)	441,20 ?
97,0 (2)	0,0050 (12)	494,75 ?
98,0 (2)		410,13 ?
100,6 (2)	0,0031 (12)	454,29 ?
102,1 (2)	0,0019 (4)	432,81 ?
102,5 (2)	0,0023 (8)	414,37 ?
102,5 (2)	0,0023 (8)	494,75 ?
103,8 (1)		494,75 ?
104,5 (3)		496,65 ?

Energy (keV)	P γ (%)	Level energy (keV)
105,7 (3)	$\approx 0,0008$	496,65 ?
106,3 (2)	0,0008	427,08 ?
106,3 (2)	0,0008	503,90 ?
108,1 (2)	0,0012 (4)	410,13 ?
110,0 (3)		565,90 ?
111,5 (3)		410,12 ?
112,1 (3)		432,81 ?
112,4 (3)		414/37 ?
113,0 (3)	0,0035 (12)	503,90 ?
114,9 (3)		155,35 ?
115,3 (3)		427,08 ?
115,3 (3)		456,113
116,5 (3)	0,0058 (8)	496,65 ?
116,9 (1)	0,0058 (8)	415,758
119,6 (2)	$\approx 0,0008$	473,04 ?
122,0 (2)	0,0015 (4)	475,69 ?
125,1 (3)	0,0015 (4)	427,08 ?
125,1 (3)	0,0015 (4)	456,113
128,3 (2)	0,0012 (4)	427,08 ?
130,0 (2)	0,0012 (4)	571,36 ?
131,0 (2)	0,0015 (4)	475,69 ?
131,0 (2)	0,0015 (4)	546,83
132,9 (2)	0,0012 (4)	565,90 ?
135,2 (3)	0,0023 (8)	456,113
135,2 (3)	0,0023 (8)	475,69 ?
135,8 (2)	0,0015 (4)	563,00 ?
136,5 (2)	0,0019 (8)	546,83
139,3 (2)	0,0023 (8)	441,20 ?
139,3 (2)	0,0023 (8)	454,29 ?
142,7 (2)	0,0023 (8)	496,65 ?
143,1 (2)	0,0015 (4)	570,27 ?
144,4 (2)	0,0035 (8)	456,113
144,4 (2)	0,0035 (8)	571,36 ?
148,5 (2)	0,0027 (8)	546,83
150,5 (2)	0,0023 (8)	503,90 ?
153,7 (2)	0,0039 (8)	546,83
154,7 (2)	0,0023 (4)	475,69 ?
154,7 (2)	0,0023 (4)	546,83
156,1 (2)	0,0023 (8)	496,65 ?
157,0 (2)	0,0027 (8)	571,36 ?
157,9 (2)	0,0023 (8)	473,04 ?
159,1 (2)	0,0039 (8)	503,90 ?
160,0 (2)	0,0031 (8)	570,27 ?
161,2 (2)	0,0027 (8)	571,36 ?
162,4 (2)	0,0023 (8)	392,25 ?
163,3 (2)	0,0023 (8)	503,90 ?
166,6 (3)	0,0012 (4)	546,83
168,0 (2)	0,0031 (8)	397,71 ?
170,6 (2)	0,0050 (8)	563,00 ?
172,8 (2)	0,0027 (8)	503,90 ?
173,7 (2)	0,0042 (12)	475,69 ?
173,7 (2)	0,0042 (12)	565,90 ?

Energy (keV)	P γ (%)	Level energy (keV)
174,7 (2)	0,0042 (12)	565,90 ?
175,2 (2)	0,0012 (4)	330,67
178,0 (2)	0,0027 (8)	570,27 ?
178,0 (2)	0,0027 (8)	571,36 ?
180,1 (2)	0,0027 (8)	571,36 ?
182,7 (2)	0,0027 (8)	571,36 ?
183,3 (3)	0,0012 (4)	503,90 ?
184,8 (2)	0,0031 (8)	496,65 ?
185,7 (3)	0,0035 (8)	565,90 ?
198,5 (2)	0,0031 (8)	353,71 ?
202,1 (2)	0,0031 (8)	503,90 ?
202,1 (2)	0,0031 (8)	546,83
205,3 (2)	0,0031 (8)	503,90 ?
206,4 (2)	0,0027 (8)	546,83
209,2 (2)	0,0023 (8)	563,00 ?
215,8 (2)	0,0027 (8)	546,83
217,8 (2)	0,0031 (8)	571,36 ?
224,4 (2)	0,0023 (8)	454,29 ?
225,2 (2)	0,0046 (12)	380,48
225,2 (2)	0,0046 (12)	565,90 ?
226,1 (2)	0,0027 (8)	546,83
226,1 (2)	0,0027 (8)	570,27 ?
226,8 (2)	0,0031 (8)	571,36 ?
232,1 (2)	0,0027 (8)	563,00 ?
235,0 (2)	0,0012 (4)	546,83
236,0 (2)	0,0023 (8)	391,09 ?
238,5 (2)	0,0054 (12)	330,67
239,8 (2)	0,0031 (8)	570,27 ?
242,3 (2)	0,0027 (8)	397,71 ?
242,3 (2)	0,0027 (8)	563,00 ?
243,4 (2)	0,0023 (8)	473,04 ?
244,6 (2)	0,0027 (8)	546,83
248,1 (1)		546,83
249,6 (2)	0,0031 (8)	570,27 ?
250,4 (2)	0,0031 (8)	571,36 ?
252,3 (2)	0,0039 (8)	344,56 ?
261,4 (2)	0,0039 (12)	302,00
261,4 (2)	0,0039 (12)	353,71 ?
264,4 (2)	0,0035 (8)	563,00 ?
268,1 (2)	0,0031 (8)	570,27 ?
269,3 (2)	0,0031 (8)	571,36 ?
271,4 (1)		570,27 ?
272,8 (3)	0,0039 (8)	571,36 ?
290,1 (1)	0,0035 (8)	330,67
298,7 (2)		391,09 ?
298,7 (2)		454,29 ?
300,0 (1)		392,25 ?
304,0 (2)	0,0046 (12)	344,56 ?
305,4 (2)	0,0050 (12)	397,71 ?
313,5 (2)	0,0139 (23)	353,71 ?
317,6 (3)	0,0023 (8)	473,04 ?
330,5 (3)	0,0023 (4)	330,67

Energy (keV)	P _γ (%)	Level energy (keV)
335,9 (3)	0,0027 (8)	565,90 ?
339,5 (5)		380,48
339,5 (5)		494,75 ?
340,5 (1)		432,81 ?
341,4 (5)		496,65 ?
344,5 (3)	0,0015 (4)	344,56 ?
351,8 (3)	0,0046 (8)	392,25 ?
363,9 (3)	0,0035 (8)	456,113
374,0 (3)	0,0073 (19)	414,37 ?
386,8 (3)	0,0031 (8)	427,08 ?
393,3 (3)	0,0050 (12)	393,33 ?
400,5 (3)	0,0031 (8)	441,20 ?
402,9 (2)	0,0023 (8)	494,75 ?
404,5 (3)	0,0035 (8)	496,65 ?
410,0 (3)	0,0069 (12)	410,13 ?
414,3 (3)	0,0054 (19)	414,37 ?

Energy (keV)	P _γ (%)	Level energy (keV)
415,764 (5)		456,113
427,0 (3)	0,0019 (8)	427,08 ?
432,8 (3)	≈ 0,0008	432,81 ?
435,1 (3)	0,0012 (4)	475,69 ?
441,1 (3)	0,0019 (8)	441,20 ?
454,2 (3)	0,0012 (8)	494,75 ?
454,2 (3)	0,0012 (8)	546,83
463,6 (3)	≈ 0,0008	503,90 ?
471,1 (3)	0,0012 (4)	563,00 ?
473,8 (3)	0,0019 (8)	565,90 ?
475,6 (3)	0,0019 (8)	475,69 ?
478,0 (3)	0,0012 (4)	570,27 ?
496,9 (3)	0,0012 (8)	496,65 ?
503,7 (3)	0,0012 (4)	503,90 ?
506,3 (3)	0,0012 (8)	546,83

6. Consistency

The most accurate Q value, Q(M), is taken from the atomic mass adjustment table of Audi et al. (2003Au03). Comparison of Q(eff) (deduced as the sum of average energies per disintegration ($\sum E_i \times P_i$) for all emissions accompanying ²³³Pa β⁻ decay) with the tabulated decay energy Q(M) allows to check a consistency of the recommended decay-scheme parameters obtained in this evaluation.

Here E_i and P_i are the evaluated energies and emission probabilities of the i-th alpha particle, beta particle, gamma-ray, X-ray, etc. Consistency (percentage deviation) is determined by $\{[Q(M)-Q(\text{eff})]/Q(M)\} \times 100$. "Percentage deviations above 5 % would be regarded as high and imply a poorly defined decay scheme; a value of less than 5 % indicates the construction of a reasonably consistent decay scheme" (quoted from the article by A.L. Nichols in Appl. Rad. Isotopes 55 (2001) 23-70).

For the above ²³³Pa decay data evaluation we have Q(M) = 570,1 (20) keV and Q(eff) = 572 (20) keV, i.e. consistency is 0,35 %.

7. References

- 1941Gr03 A. V. Grosse, E. T. Booth, J. R. Dunning, Phys. Rev. 59(1941)322 (Half-life)
- 1952Br84 C. I. Browne, Jr., Thesis, Univ. California (1952); UCRL-1764(1952) (Gamma-ray energies)
- 1954Br37 W. D. Brodie, Proc. Phys. Soc.(London) 67A(1954)397 (Measured energies and probabilities of β-transitions)
- 1955On05 Ong Ping Hok, P. Kramer, Physica 21(1955)676 (Measured energies and probabilities of β-transitions)
- 1956Mc60 L. D. Mc Isaac and E. C. Freiling, Nucleonics 14(10)(1956)65 (Half-life)
- 1957Wr37 H. W. Wright, E. T. Wyatt, S. A. Reynolds, W. S. Lyon, T. H. Handley, Nucl. Sci. Eng. 2(1957)427 (Half-life)
- 1960Un01 J. P. Unik, Thesis, Univ. California (1960); UCRL -9105(1960) (Measured energies and probabilities of β-transitions)

- 1961Al19 R. G. Albridge, J. M. Hollander, C. J. Gallagher, J. H. Hamilton, Nucl. Phys. 27(1961)529 (Gamma-ray energies and multipolarities, E2 admixtures)
- 1962Sc03 G. Schultze, J. Ahlf, Nucl. Phys. 30(1962)163 (Multipolarities, E2 admixtures)
- 1963Bi03 K. M. Bisgard, P. Dahl, P. Hornshoj, A. B. Knutsen, Nucl. Phys. 41(1963)21 (Multipolarities, E2 admixtures)
- 1963Bj03 S. Bjørnholm, M. Lederer, F. Asaro, and I. Perlman, Phys. Rev. 130(1963)2000 (Energies and probabilities of β -transitions)
- 1966Ze02 M. J. Zender, Thesis, Vanderbilt Univ. (1966) (Multipolarities, E2 admixtures)
- 1967Br20 C. Briancon, C. -F. Leang, P. Paris, Compt. Rend. 264B(1967)1522 (Gamma-ray energies)
- 1968Ma13 S. G. Malmskog and M. Hojeberg, Ark. Fys. 35(1968)197 (Gamma-ray energies)
- 1971Vo02 Z. T. von Egidy, O. W. B. Schult, W. Kallinger, D. Breitig, R. P. Sharma, H. R. Koch, H. A. Baader, Naturforsch. 26a(1971)1092 (Gamma-ray energies)
- 1972De67 M. de Bruin, P. J. M. Korthoven, J. Radioanal. Chem. 10(1972)125 (Gamma-ray energies)
- 1973Va33 T. Valkeapaa, A. Siivola, G. Graeffe, Phys. Fenn. 9(1973)43 (Gamma-ray energies and emission probabilities)
- 1978Poenitz W. P. Poenitz, D. I. Smith, United States Dept. of Energy, Washington DC, Rep. ANL/NDM 42 (March 1978) (Gamma-ray emission probabilities)
- 1979Ge08 R. J. Gehrke, R. G. Helmer, C. W. Reich, Nucl. Sci. Eng. 70(1979)298 (X- and gamma-ray emission probabilities)
- 1984Va27 R. Vaninbrouckx, G. Bortels, B. Denecke, Int. J. Appl. Radiat. Isotop. 35(1984)905 (X- and gamma-ray emission probabilities)
- 1985DeZR M. J. de Bettencourt, Thesis, Univ. Paris-Sud(Orsay) (1985) (Tentative gamma-rays)
- 1986Jo07 R. T. Jones, J. S. Merritt, A. Okazaki, Nucl. Sci. Eng. 93(1986)171 (Half-life)
- 1986Kr10 K. S. Krane, Nucl. Phys. A459(1986)1 (Multipolarities, E2 admixtures)
- 1988Wo01 S. A. Woods, P. Christmas, P. Cross, S. M. Judge, W. Gelletly, Nucl. Instrum. Meth. Phys. Res. A264(1988) 333; Addendum Nucl. Instrum. Meth. Phys. Res. A272(1988)924 (Gamma-ray energies)
- 1989Br24 E. Browne, B. Sur, E. B. Norman et al, Nucl. Phys. A501(1989)477 (Experimental ICC, gamma multipolarities, beta transition probabilities)
- 1990Ak02 Y. A. Akevali, Nucl. Data Sheets 59(1990)263 (A=233 NDS evaluation, gamma-ray multipolarities, E2 admixtures)
- 1990Ko41 M. C. Kouassi, C. Ardisson-Marsol, G. Ardisson, J. Phys. (London) G16(1990)1881. (Level scheme, multipolarities, absolute KX-ray emission probability and gamma-ray energies)
- 1990Pe16 J. Pearcey, S. A. Woods, P. Christmas, Nucl. Instrum. Meth. Phys. Res. A294(1990)516 (E2 γ -ray admixtures)
- 1992Ra08 M. U. Rajput, T. D. Mac Mahon, Nucl. Instrum. Meth. Phys. Res. A312(1992)289 (Evaluation technique)
- 1996SC06 E. Schönfeld, H. Janßen, Nucl. Instr. Meth. Phys. Res. A369(1996)527 (atomic data)
- 1999Popov Yu. S. Popov and G. A. Timofeev, Radiokhimiya 41(1999)27 (in Russian) (Half-life)
- 2000Sc04 U. Schötzig, E. Schönfeld, H. Jarßen, Appl. Rad. Isotop. 52(2000)883 (Gamma-ray and X-ray emission probabilities)
- 2000Smith D. Smith, M. I. Woods, D. H. Woods, Preliminary Report, NPL, Teddington, 2000 (Gamma-ray and X-ray emission probabilities)
- 2000Us01 K. Usman, T. D. MacMahon, Appl. Radiat. Isot. 52(2000)585 (Half-life)
- 2000Ch01 V. P. Chechev, A. G. Egorov, Appl. Rad. Isot. 52(2000)601 (Evaluation technique)
- 2000Lu01 A. Luca, M. Etcheverry, J. Morel, Appl. Rad. Isot. 52(2000)481 (Gamma-ray emission probabilities)
- 2000Wo01 S. A. Woods, D. H. Woods, P. de Lavison, S. M. Jerome, J. L. Makepeace, M. J. Woods, L. J. Husband, S. Lineham, Appl. Radiat. Isot. 52(2000)475 (Gamma-ray emission probabilities)

- 2002Be M. M. Bé, R. Helmer, V. Chisté, J. Nucl. Sci. Tech., suppl.2(2002)481 (SAISINUC software)
- 2002Lu01 A. Luca, S. Sepman, K. Iakovlev, G. Shchukin, M. Etcheverry, J. Morel, Appl. Rad. Isot. 56(2002)173 (Gamma-ray and X-ray emission probabilities)
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)337 (Q value)
- 2004Sh07 G. Shchukin, K. Iakovlev, J. Morel, Appl. Rad. Isot. 60(2004)239 (Gamma -ray emission probabilities)
- 2005Hu06 X. Huang, P. Liu, B. Wang, Appl. Radiat. Isot. 62(2005)797 (Evaluation of ^{233}Pa Decay Data)
- 2005Si15 B. Singh, J. K. Tuli, Nucl. Data Sheets 105(2005)109(A=233 NDS evaluation, ^{233}U level energies, gamma-ray energies and multipolarities)
- 2006Ch39 V. P. Chechev, N. K. Kuzmenko, Appl. Radiat. Isot. 64 (2006)140 3 (^{233}Pa decay data evaluation)
- 2006Ha53 H. Harada, S. Nakamura, M. Ohta, T. Fujii, H. Yamana, J. Nucl. Sci. Technol.(Tokyo) 43 (2006)1289 (Gamma-ray emission probabilities)
- 2008De10 D. J. DeVries and H. C. Griffin, Appl. Rad. Isotop., 66(2008)1999 (Uncertainties of LX-ray absolute emission probabilities)
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, and C. W. Nestor, Jr., Nucl. Instrum. Meth. Phys. Res. A589(2008)202 (Theoretical ICC)

²³⁴Th – Comments on Evaluation of Decay Data by A. Luca

This evaluation was completed in May 2009. The literature available by December 31st, 2008 was included.

1. Evaluation Procedures

The Limitation of Relative Statistical Weight (LWM) method was applied for averaging numbers throughout this evaluation; this method was implemented by using the computer code LWEIGHT, ver. 4 (designed for Excel, MS Office). The uncertainty assigned to an average value in this evaluation is never lower than the lowest uncertainty of any of the experimental input values.

2. Decay Scheme

²³⁴Th decays 100 % by beta minus particle emissions, mainly to ²³⁴Pa^m - the 1.159 min. half-life metastable state of ²³⁴Pa (the first experimentally established case of nuclear isomerism, by O. Hahn, in 1921). The decay scheme was studied by many authors, since early '60s (1961Ge13, 1962Br05, 1963Bj02, 1964Ab04, 1965Fo12 and 1973Go40). The first recommended values for the main ²³⁴Th nuclear decay data were published in the evaluation of Coursol et al., in 1990 (1990Co08); other important evaluation can be found in 1998Ad08. In the present evaluation, the spin, parity, energy and half-life values of the ²³⁴Pa excited levels, and the multiplicities of the γ -ray transitions, have been adopted from the most recent A=234 ENSDF mass-chain evaluation, published by E. Browne and J.K. Tuli (2007Br04). The very important low energy and intensity isomeric transition (maximum energy of less than 10 keV) from ²³⁴Pa^m to the first excited level of ²³⁴Pa (explaining the 73.92 keV gamma-ray transition to the ²³⁴Pa ground state), was not observed yet, probably because the conversion lines are obscured by intense Auger M and Coster-Kronig electrons (according to Godart and Gizon, 1973); as a consequence, the energies of all the ²³⁴Pa excited levels decaying to ²³⁴Pa^m are known to be upheld 10 keV at most with a systematic uncertainty (usually considered as "x" keV, in 2007Br04 and other evaluations; in the present evaluation, this quantity is not written in the decay scheme, but it should be added to the energy of the excited levels, respectively subtracted from the reported beta transitions energies). A more detailed decay scheme of ²³⁴Th can be found in 2007Br04. The decay of ²³⁴Pa^m (by alpha-particle emission and isomeric transition) is not studied in this evaluation.

3. Nuclear Data

The adopted beta decay energy value $Q(\beta^-)=272(10)$ keV, is based on the energy measurements of Godart and Gizon (1973Go40): 198.5 (15) keV for the maximum energy of the beta minus particle emissions and 73.92 (2) keV for the isomeric transition; an uncertainty of 10 keV was assigned to the result, according to the above-mentioned considerations. The adopted value of $Q(\beta^-)$ is in agreement with the value from 2003Audi03: 273.1 (32) keV (based on some older energy measurements of the beta minus particle emissions). The value adopted by this evaluation is also in good agreement with the effective $Q(\beta^-)$ value of 273 keV (with an uncertainty of 11 keV), calculated from the decay scheme data, by using the SAISINUC software.

3.1. Half-life

In the literature, only a few measured ²³⁴Th half-life ($T_{1/2}$) values are reported; these measurements are very old (the most recent is from 1948), so new half-life measurements are needed to improve the quality of the evaluation. The half-life values and their uncertainties are presented in Table 1; the value recommended by Curie et al. (1931), with an estimated uncertainty added by the evaluator, was also included. The set of data is consistent and the recommended value, 24.10 days, with an uncertainty of 0.03 day, is the weighted average (LWM, $\chi^2_{\nu}=3.78$) of the four input values. The references are expressed as NSR (Nuclear Science References) type keynumbers:

Table 1 : ²³⁴Th Half-life values

$T_{1/2}$ (days)	Uncertainty of $T_{1/2}$ (days)	Reference
23.8	0.7	1920Ki01
24.5	0.5	1931Cu01
24.1	0.2	1939Sa11
24.101	0.025	1948Kn23

3.2. Beta transitions and emissions

In the literature, the most complete reference reporting measurements of energy and emission intensities for ²³⁴Th beta minus transitions is 1973Go40.

For this evaluation, the beta transitions energies were calculated from $Q(\beta^-)$ and the energies of the decay scheme levels; the high energy uncertainty (10 keV) is explained by the possible low energy and intensity isomeric transition (as described above, in section 2, Decay Scheme). The intensities of the beta branches were deduced from γ -ray transition intensity balance at each level, with the exception of the main branch; its intensity was deduced from the normalization condition of the beta emissions (the sum of the all the beta transitions intensities must be 100 %). The existence of the weakest beta decay branch (95.8 keV) is questionable (2007Br04). The energy and intensity values of the beta transitions, as well as their Log ft values are shown in Table 2.

Table 2: ²³⁴Th β^- Energies and Emission Probabilities

E_{β^-} (keV)	Uncertainty E_{β^-} (keV)	Transition intensity (%)	Transition intensity (%), from 1973Go40	Log ft
85	10	1.6 (6)	1.3 (7)	7.0
95	10	0.016 (5)	-	9.1
105	10	6.5 (7)	5.4 (10)	6.7
106	10	14.1 (12)	20.7 (10)	6.3
198	10	77.8 (15)	72.5 (20)	6.4

3.3. γ - transitions: γ rays and internal conversion electrons

Many measurements of the γ -ray energies and emission intensities following the ²³⁴Th decay were published by different authors: 1973Go40, 1973Sa33, 1973Ta25, 1978Ch06, 1982Mo30, 1990Sc09, 1993Su37, 2004Ab03 and 2006Al28. The interest for high quality data of photon emission probabilities is justified especially in the field of environmental radioactivity monitoring. Table 3 presents measured values of the 63.30 (2) keV γ -ray emission probability following the decay of ²³⁴Th. The set of data is consistent and the recommended value, 3.75 (8) %, is the weighted average (LWM, $\chi^2_{\nu}=3.32$) of the five input values. The references are expressed as NSR type keynumbers.

Table 3 : Absolute Emission Intensity Results (in %) for the 63.30-keV γ ray.

Gamma-ray emission probability	Uncertainty of the gamma-ray emission probability	Reference
3.3	0.3	1973Go40
4.05	0.20	1982Mo30
3.6	0.2	1990Sc09
3.99	0.20	1993Su37
3.73	0.07	2004Ab03
3.75	0.08	Adopted

Using this evaluated value and the relative photon intensity values from the measurements of Chu and Scharff-Goldhaber (1978), the corresponding absolute gamma-ray emission probabilities and their uncertainties were computed for all the γ rays and are given below in Table 4. The relative photon intensities measured by Chu and Scharff-Goldhaber were preferred to those of Godart and Gizon (1973), mainly because in this case the U KX-rays contributions were resolved from the gamma-ray peaks situated in the (90-115) keV energy range of the spectra; no other references reporting relative photon intensities measurements were found in the literature.

The intensity balance for level 3 (103.42 keV) was used to compute the emission probability for the 73.85 keV photons, but the obtained value was negative (-0.011 %); as the placement of this transition in the level scheme is uncertain (2007Br04), this low probability photon emission was not considered in this evaluation. Other possible gamma-ray transitions neither confirmed nor placed in the level scheme (proposed / observed only by some authors) are: 57.75 keV, 87.02 keV, 92.00 keV, 103.71 keV, 108.00 keV, 132.9 keV and 184.8 keV.

The internal conversion coefficients were computed with the program BrIcc, version 2.2/2008, using the "Frozen Orbitals" approximation. A difficult case is the computation of the ICC for the 112.81 keV gamma-ray transition, because this energy is too close to the K-shell binding energy for protactinium (112.6 keV) and the software can not be used directly for this purpose. Following Browne and Tuli (2007), a limit on $\alpha(K)$ (≤ 0.29) has been obtained from extrapolation of $\alpha(K)$'s for energies higher than 113.6 keV; however, this procedure introduced a large uncertainty of the total ICC value (see Table 4).

Table 4: ²³⁴Th γ -ray Energies and Absolute Emission Probabilities

E_γ (keV)	Uncertainty E_γ (keV)	Absolute Emission Probability (%)	Uncertainty of absolute emission probability (%)	Total ICC (α_T)
20.01	0.02	0.005 1	0.002 1	240 (70)
29.50	0.02	0.001 23	0.000 14	4390 (70)
62.88	0.02	0.016 4	0.002 8	25 (5)
63.30	0.02	3.75	0.08	0.405 (6)
73.92	0.02	0.013 3	0.001 4	10.6 (4)
83.31	0.05	0.061	0.005	0.196 (3)
92.38	0.01	2.18	0.19	5.27 (8)
92.80	0.02	2.15	0.19	0.1472 (21)
103.35	0.10	0.003 2	0.001 0	3.81 (6)
112.81	0.05	0.215	0.022	0.23 (14)

4. Atomic data

The K-shell fluorescence yield (ω_K), the mean L-shell fluorescence yield (ω_L) and the mean number of vacancies in the L-shell produced by one vacancy in the K-shell (η_{KL}) were determined using the computer program EMISSION v3.10, 28-Jan-2003: 0.970 (4), 0.488 (18) and 0.795 (5) respectively.

4.1. Auger electrons and X-rays

The relative probability values of the K Auger electron emissions (KLL, KLX, KXY) normalized to the KLL value, were computed using the same EMISSION computer program. The total K Auger electron emission probability (absolute) and the emission probability of the L Auger electrons were also calculated. The energy ranges for K and L Auger electrons were filled-in by the SAISINUC program, version 2008 April.

The relative probability (normalized to $K_{\alpha 1}$ X-rays emission) and the absolute emission probability values of the different groups of K and L X-rays were determined using the same EMISSION program. The energy range values of the K and L X-rays are from the tables linked to SAISINUC. Neither measurements of X-ray energies nor of emission probabilities were found in the literature, in order to compare them with the results of this evaluation.

5. Main production mode

The main production mode of ²³⁴Th is by alpha-particle decay of the ²³⁸U nuclei (²³⁴Th is the daughter of ²³⁸U), present in important quantities in many natural ores.

6. References

- 1920Ki01 G. Kirsch, Mitt. Ra. Inst. 127, Wien. Ber. 11a, 129, 309 (1920), M-Sch. P. 377.
- 1931Cu01 M. Curie, A. Debierne, A.S. Eve, H. Geiger, O. Hahn, S.C. Lind, St. Meyer, E. Rutherford and E. Schweidler, "The Radioactive constants as of 1930", Rev. Mod. Phys. 3, 427-445 (1931).
- 1939Sa11 B. W. Sargent, Can. J. Research A17, 103 (1939).
- 1948Kn23 G.B. Knight and R.L. Macklin, "Half-Life of UX₁ (Th₂₃₄)", Phys. Rev. 74, 1540-1541 (1948).
- 1961Ge13 J.S. Geiger, R.L. Graham and T.A. Eastwood, "The Decay of Th²³⁴", AECL-1472, PR-P-52, 26-27 (1961).
- 1962Br05 J.-P. Briand, "Sur la nature de la transition de 29 keV du Protactinium 234 (UX₂)", Comp. Rend. Acad. Sci. (Paris) 254, 84-86 (1962).
- 1963Bj02 S. Bjornholm and O.B. Nielsen, "The decay of the 1.14 min Isomer of Pa²³⁴ (UX₂)", Nucl. Phys. 42, 642-659 (1963).
- 1964Ab04 H. Abou-Leila, "Mesures des périodes des niveaux du protactinium 234 désexcités par les transitions γ de 63 et 93 keV", Comp. Rend. Acad. Sci. (Paris) 258, 5632-5635 (1964).
- 1965Fo12 R. Foucher, "New Data on the Level Diagram of the Odd-Odd Nucleus Pa²³⁴", Izv. Akad. Nauk SSSR, Ser.Fiz. 29, 100 (1965); Bull. Acad. Sci. USSR, Phys. Ser. 29, 99-100 (1966).
- 1973Sa33 T.E. Sampson, "Precision Measurement of Gamma Ray Energies from ²³⁸U Daughters, II", Nucl. Instrum. Methods 111, 209-211 (1973).
- 1973Ta25 Harry W. Taylor, "Gamma Rays Emitted by Uranium and Thorium in the Energy Range 10-120 keV", Int. J. Appl. Radiat. Isotop. 24, 593-597 (1973).
- 1973Go40 J. Godart and A. Gizon, "Niveaux de ²³⁴Pa Atteints par la Désintégration de ²³⁴Th", Nucl. Phys. A217, 159-176 (1973).
- 1978Ch06 Y.Y. Chu and G. Scharff-Goldhaber, "Decay of ²³⁴Th to the ²³⁴Pa isomers", Phys. Rev. C 17, 1507-1509 (1978).
- 1982Mo30 Michael H. Momeni, "Analyses of Uranium and Actinium Gamma Spectra: An Application to Measurements of Environmental Contamination", Nucl. Instrum. Methods 193, 185-190 (1982).
- 1990Sc09 H.L. Scott and K.W. Marlow, "Gamma-Ray Emission Probabilities of the Daughters of ²³⁸U", Nucl. Instrum. Methods A 286, 549-555 (1990).
- 1990Co08 N. Coursol, F. Lagoutine and B. Duchemin, "Evaluation of Non-Neutron Nuclear Data for the Uranium-238 Decay Chain", Nucl. Instrum. Methods A 286, 589-594 (1990).

Comments on evaluations

- 1993Su37 G.A. Sutton, S.T. Napier, M. John and A. Taylor, "Uranium-238 Decay Chain Data", *The Science of the Total Environment* 130/131, 393-401 (1993).
- 1998Ad08 I. Adsley, J.S. Backhouse, A.L. Nichols and J. Toole. "U-238 Decay Chain: Resolution of Observed Anomalies in the Measured Secular Equilibrium Between Th-234 and Daughter Pa-234m", *Appl. Rad. Isotopes* 49, 1337-1344 (1998).
- 2003Au03 G. Audi, A.H. Wapstra and C. Thibault, "The AME2003 atomic mass Evaluation (II). Tables, graphs, and references", *Nucl. Phys. A* 729, 337-676 (2003).
- 2004Ab03 S. Abousahl, P. van Belle, B. Lynch and H. Ottmar, "New measurement of the emission probability of the 63.290 keV ²³⁴Th γ -ray from ²³⁸U α decay", *Nucl. Instrum. Methods A* 517, 211-218 (2004).
- 2006Al28 F.S. Al-Saleh, Al-J.H. Al-Mukren and M.A. Farouk, "Precise determination of the absolute intensities of the gamma-ray lines of ²³⁵U and some ²³⁸U daughters", *Nucl. Instrum. Methods A* 568, 734-738 (2006).
- 2007Br04 E. Browne and J.K. Tuli, "Nuclear Data Sheets for A = 234", *Nucl. Data Sheets* 108, 681-772 (2007).

²³⁴Pa-Comments on evaluation of decay data

Huang Xiaolong, Wang Baosong

This evaluation was completed in 2009. Literature available by January 2009 was included.

1 Decay Scheme

²³⁴Pa disintegrates 100 % by β^- emissions to levels in ²³⁴U. ²³⁴Pa ground state has $J^\pi = 4^+$ (2007Br04).

The β^- decay scheme of ²³⁴Pa is based on the measurement results of 1986Ar05. 28 observed γ -rays were not placed in the current decay scheme. These gamma rays carry about 3.2 % of the total intensity of all the gamma rays placed in the decay scheme.

The $Q(\beta^-)$ value of 2195 (4) keV adopted from 2003Au03 is not in good agreement with the effective $Q(\beta^-)$ value of 2336 (70) keV, calculated by the evaluators from average radiation energies using the RADLST computer program. The total intensity $\Sigma I(\beta^-)$ deduced by the evaluators from intensity balance at each level is about 110 %.

These results suggest that the γ -ray intensity balance for some levels may be incomplete and the decay scheme has some inconsistency. Further measurements are strongly needed to determine the γ transitions and the decay scheme with greater precision.

2 Nuclear Data

The $Q(\beta^-)$ value is from the mass adjustment in 2003Au03.

Level energies, have been obtained from a least-squares fit to γ -ray energies (GTOL computer code). Spin and parities are from 2007Br04.

The measured and recommended ²³⁴Pa half-life values are listed in Table 1.

Table 1: Measured half-life values of ²³⁴Pa and recommended value

$T_{1/2}$ (h)	References	Comments
6.7	1931Cu01	Not used
6.658 (12)	1954Zi02	
6.75 (3)	1962Bj01	
6.704 (46)		Unweighted mean
6.671 (11)		Weighted mean
6.704 (46)		LWEIGHT weighted mean, $\chi^2 = 4.7$
6.70 (5)		Recommended value

The weighted average for this data set of the 2 discrepant experimental values is dominated by the accurate value of 1954Zi02. The LWEIGHT computer program, which uses a Limitation of Relative Statistical Weights (LRSW method), has increased the 1954Zi02 uncertainty from 0.012 to 0.030 and used a weighted mean and an external uncertainty for recommended average.

Thus, the adopted value of the ²³⁴Pa half-life is 6.70 (5) hours.

2.1 β^- transitions

The maximum energies of the β^- transitions in the decay of ²³⁴Pa have been deduced from the $Q(\beta^-)$ value (2003Au03) and the ²³⁴U level energies (Table 2), obtained from a least-squares fit to recommended γ -ray energies (GTOL computer code).

Table 2: ²³⁴U levels populated in ²³⁴Pa β⁻ decay

Level energy (keV)	Spin & parity	Half-life	β ⁻ transition probabilities (%)
0.0	0+	2.455 (6) × 10 ⁵ a	
43.481 (15)	2+	0.252 (7) ns	
143.375 (21)	4+		< 5
296.075 (24)	6+		
497.05 (4)	8+		
786.295 (15)	1-		
809.92 (8)	0+	< 0.1 ns	
849.265 (23)	3-		< 0.8
851.73 (5)	2+	> 1.74 ps	
926.744 (21)	2+	1.38 (17) ps	
947.59 (5)	4+		< 0.8
962.55 (3)	5-		< 0.4
968.45 (3)	3+		< 2.5
989.444 (20)	2-	0.76 (4) ns	< 3.1
1023.795 (24)	3-		< 5
1023.92 (3)	4+		1.5 (13)
1069.297 (22)	4-		< 8
1085.07 (10)	2+		
1090.89 (4)	5+		0.69 (20)
1096.12 (9)	6+		
1125.29 (5)	7-		
1126.65 (3)	2+		
1127.535 (25)	5-		1.9 (10)
1165.41 (4)	3+		
1172.03 (3)	6+		
1194.761 (23)	6-		< 1.5
1214.70 (5)	4+		0.30 (12)
1237.24 (3)	1-		
1261.77 (3)	7+		
1274.32 (9)	(5+)		
1277.45 (3)	7-		
1312.20 (9)	3-		0.109 (18)
1341.33 (8)	(6+)		
1421.252 (24)	6-	33.5 (20) μs	
1447.89 (10)	5-		0.11 (3)
1456.54 (7)	(2-)		
1486.17 (12)	(3-)		0.117 (25)
1496.14 (3)	3+		< 2.7
1502.38 (8)	3,4+		0.25 (4)
1533.37 (5)	(4-)		0.21 (4)
1537.25 (3)	4+		< 0.9
1543.71 (5)	4+		0.10 (9)
1548.10 (8)	(5)		0.078 (20)
1552.554 (24)	5+	2.20 (25) ns	19.6 (18)
1581.67 (10)	(5-)		0.05 (3)
1588.84 (3)	5+		< 0.7
1619.46 (9)	(6+)		0.035 (20)
1649.99 (12)	(6-)		0.18 (4)

Level energy (keV)	Spin & parity	Half-life	β^- transition probabilities (%)
1653.35 (7)	(3+)		0.95 (13)
1693.42 (3)	5-		6.9 (8)
1722.89 (4)	3-		8.4 (9)
1723.424 (24)	4+		36 (5)
1737.42 (7)	3+		1.16 (14)
1738.18 (6)	(3+)		0.78 (19)
1761.86 (6)	(4-)		2.8 (4)
1770.79 (9)	(3+)		0.129 (17)
1782.58 (3)	5+		8 (3)
1784.19 (13)	4+		0.061 (11)
1793.05 (6)	4+		0.41 (8)
1811.62 (6)	4+		1.43 (15)
1843.88 (17)	3,4,5-		0.17 (3)
1863.08 (15)	(5+)		0.029 (7)
1881.75 (7)	4+		0.25 (3)
1916.28 (9)	3,4+		0.21 (3)
1927.51 (7)	4+		0.22 (4)
1940.52 (9)	4+		0.35 (5)
1958.75 (4)	3-		0.44 (19)
1968.84 (10)	4+,5		0.044 (12)
1981.22 (7)	4+		0.59 (8)
2000.45 (13)	(4+)		0.122 (16)
2019.82 (13)	4+		0.112 (16)
2033.54 (5)	3+,4+		0.90 (15)
2037.06 (17)	4+,5		0.055 (8)
2066.24 (10)			0.140 (24)
2068.82 (11)	3,4,5+		0.40 (7)
2101.42 (9)	5+		0.064 (11)
2115.71 (11)	4+		0.21 (3)
2144.04 (9)	3+,4+		0.42 (5)

Table 3: Measured and evaluated β^- energies (keV) and probabilities (%) in the ²³⁴Pa decay

1956On07		1959De30		1968Bj06		Evaluated	
E $_{\beta^-}$	P $_{\beta^-}$	E $_{\beta^-}$	P $_{\beta^-}$	E $_{\beta^-}$	P $_{\beta^-}$	E $_{\beta^-}$	P $_{\beta^-}$
155	28	141 (10)	35.5			158 (4)	0.055 (8)
		274 (10)	21.4	280 (70)	12	279 (4)	0.21 (3)
320 (20)	32					313 (4)	0.25 (3)
		363 (10)	10.3			383 (4)	1.43 (15)
		477 (10)	16.0			472 (4)	36 (5)
530 (20)	27			550 (100)	63	545 (4)	0.18 (4)
		576 (10)	13.2			576 (4)	0.035 (20)
				790 (100)	19	747 (4)	0.11 (3)
		1042 (20)	3.6			1067 (4)	1.9 (10)
1130 (50)	13					1126 (4)	< 8
				1190 (100)	5	1171 (4)	< 5
				1510 (200)	≤ 1	1346 (4)	< 0.8

The adopted β^- transition probabilities and the associated uncertainties were deduced from the γ transition probability balance at each ²³⁴U level.

The values of *logft* and average β^- energies have been calculated with the LOGFT computer program.

2.2 γ Transitions

The γ -ray transition probabilities were deduced using the γ -ray emission intensities and the relevant internal conversion coefficients.

Multipolarities and mixing ratios of γ -ray transitions are from 1967Wa26, 1968Bj06 and 2007Br04.

The internal conversion coefficient (ICC) (and its associated uncertainty) for γ -ray transitions have been interpolated from theoretical values based on the “Frozen Orbital” approximation (2002Ba85) using the BrIcc computer program (2008Ki07).

3. Atomic data

Atomic fluorescence yields ($\omega_K, \omega_L, \omega_M, \eta_{KL}$ and η_{LM}) are from Schönfeld (1996Sc06).

The X-ray and Auger electron emission probabilities have been deduced from γ -ray and conversion electron data by using the RADLST computer code.

The deduced total KX-ray emission probability is 35.7 (12) %. Measured KX-ray emission probability is 50.9 % (from $I(KX\text{-ray})/I_\gamma(131\text{keV } \gamma\text{-ray}) = 2.8$ in 1967Wa26). The 30 % deviation suggests a problem with the decay scheme.

4. Electron Emissions.

The conversion electron emission probabilities have been deduced from γ -ray transition data using theoretical internal conversion coefficients.

5. Photon Emissions

5.1 γ -ray energies

Measured results for the energies of γ -rays from ²³⁴Pa are listed in Table 4. The recommended values are taken from the precise measurements of 2000Ni13, 1986Ar05 and 1972Sa06, except as noted in the table.

Table 4: Measured and recommended γ -ray energy values from ²³⁴Pa β^- decay

1968Bj06	1968Go20	1972Sa06	1975Ar24	1986Ar05	2000Ni13	Recommended
34.30 (4)						34.30 (4)
						41.82 (11) ^a
43.40 (5)				43.49 (2)		43.49 (2)
45.19 (5)				45.45 (5)		45.45 (5)
				54.96 (10)		54.96 (10)
				55.45 (5)		55.45 (5)
58.20 (6)				58.2 (1)		58.20 (6)
				59.19 (5)		59.19 (5)
63.40 (7)			63.0 (5)	62.70 (1)		62.70 (1)
67.10 (7)				67.25 (10)		67.25 (10)
69.90 (7)				69.46 (5)		69.46 (5)
				75.0 (3) ^a		75.0 (3) ^a
79.69 (8)			80.5 (5)	79.84 (2)		79.84 (2)
				97.17 (10)		97.17 (10)
99.67 (10)				99.86 (2)		99.86 (2)
				100.89 (2)		100.89 (2)
103.41 (11)				103.77 (2)		103.77 (2)

Comments on evaluation

1968Bj06	1968Go20	1972Sa06	1975Ar24	1986Ar05	2000Ni13	Recommended
				106.68 (5)		106.68 (5)
125.20 (13)			125.3 (2)	125.46 (1)		125.46 (1)
131.00 (13)			131.28 (10)	131.30 (1)		131.30 (1)
134.37 (14)				134.61 (2)		134.61 (2)
			137.7 (5)	137.23 (5)		137.23 (5)
139.97 (14)			140.3 (2)	140.15 (2)		140.15 (2)
				140.91 (3)		140.91 (3)
144.35 (15)			143.6 (5)	143.78 (2)		143.78 (2)
~ 150.2				149.88 (3)		149.88 (3)
152.46 (16)			153.0 (2)	152.71 (2)		152.71 (2)
159.10 (16)			159.2 (3)	159.48 (2)		159.48 (2)
				164.94 (5)		164.94 (5)
			166.3 (10)	165.61 (5)		165.61 (5)
170.77 (18)			170.6 (3)	170.85 (2)		170.85 (2)
			174.6 (8)	174.55 (3)		174.55 (3)
				179.80 (8)		179.80 (8)
185.95 (19)			186.2 (5)	186.15 (2)		186.15 (2)
193.4 (2)			193.5 (5)	193.73 (3)		193.73 (3)
196.4 (2)			196.5 (10)	196.80 (5)		196.80 (5)
199.7 (2)				199.95 (5)		199.95 (5)
200.6 (2)			200.9 (3)	200.97 (3)		200.97 (3)
202.9 (2)			202.9 (3)	203.12 (3)		203.12 (3)
219.60 (22)			220.8 (5)	220.00 (8)		220.00 (8)
				221.15 (10)		221.15 (10)
				221.83 (10)		221.83 (10)
226.15 (23)			226.87 (10)	226.50 (3)		226.50 (3)
227.00 (23)				227.25 (3)		227.25 (3)
				232.21 (3)		232.21 (3)
						233.6 (2) ^b
				235.11 (3)		235.11 (3)
						235.9 (3) ^b
				240.2 (1)		240.2 (1)
245.00 (25)			245.2 (3)	245.37 (2)		245.37 (2)
						247.79 (7) ^b
248.80 (25)			249.1 (3)	249.22 (1)		249.22 (1)
				257.2 (1)		257.2 (1)
			267.1 (8)	267.12 (5)		267.12 (5)
271.85 (27)			272.1 (3)	272.28 (5)		272.28 (5)
				275.04 (10)		275.04 (10)
			277.9 (8)	278.3 (1)		278.3 (1)
293.5 (3)			293.7 (2)	293.79 (5)		293.79 (5)
				295.91 (8)		295.91 (8)
				298.7 (2)		298.7 (2)
			309.6 (8)	308.6 (2)		308.6 (2)
				310.2 (1)		310.2 (1)
						310.52 (10) ^b
312.5 (3)				313.5 (1)		313.5 (1)
316.8 (3)			316.3 (8)	316.7 (1)		316.7 (1)
			320.7 (8)	320.4 (1)		320.4 (1)
328.3 (3)			330.3 (5)	330.40 (5)		330.40 (5)

1968Bj06	1968Go20	1972Sa06	1975Ar24	1986Ar05	2000Ni13	Recommended
330.6 (3)				331.4 (1)		331.4 (1)
				340.2 (1)		340.2 (1)
				343.8 (2)		343.8 (2)
351.6 (4)			351.8 (3)	351.9 (1)		351.9 (1)
				357.9 (1)		357.9 (1)
				360.6 (3)		360.6 (3)
				365.0 (3)		365.0 (3)
369.3 (4)			369.6 (3)	369.50 (5)		369.50 (5)
371.8 (4)			372.2 (3)	372.0 (1)		372.0 (1)
				379.1 (1)		379.1 (1)
				385.4 (1)		385.4 (1)
						387.94 (6) ^b
				394.1 (1)		394.1 (1)
				397.7 (3)		397.7 (3)
				401.8 (2)		401.8 (2)
			409.8 (5)	409.8 (1)		409.8 (1)
				416.1 (1)		416.1 (1)
				425.3 (2)		425.3 (2)
427.0 (4)			426.8 (5)	426.95 (5)		426.95 (5)
						427.4 (4) ^b
			432.6 (5)	433.1 (1)		433.1 (1)
			446.9 (5)	446.6 (1)		446.6 (1)
						450.93 (4) ^b
				452.4 (3)		452.4 (3)
458.6 (5)			458.6 (3)	458.68 (5)		458.68 (5)
			461.8 (10)	461.5 (1)		461.5 (1)
				464.2 (1)		464.2 (1)
468.0 (5)			467.5 (10)	468.0 (1)		468.0 (1)
			472.1 (10)	472.3 (1)		472.3 (1)
474.0 (5)			473.5 (10)	474.2 (2)		474.2 (2)
			478.7 (10)	478.6 (1)		478.6 (1)
			480.5 (8)	481.0 (1)		481.0 (1)
			498.9 (10)	498.0 (1)		498.0 (1)
				502.0 (1)		502.0 (1)
506.0 (5)			506.8 (5)	506.75 (5)		506.75 (5)
513.6 (5)			513.7 (5)	513.4 (1)		513.4 (1)
			520.2 (5)	519.6 (1)		519.6 (1)
521.0 (5)			521.0 (5)	521.4 (1)		521.4 (1)
527.6 (5)			528.0 (5)	527.9 (1)		527.9 (1)
				529.1 (3)		529.1 (3)
			533.2 (10)	534.1 (1)		534.1 (1)
			537.1 (10)	537.2 (1)		537.2 (1)
				543.8 (1)		543.8 (1)
				553.7 (1)		553.7 (1)
			557 (1)	558.0 (2)		558.0 (2)
				559.2 (2)		559.2 (2)
				562.8 (3)		562.8 (3)
565.1 (6)			566.3 (10)	565.2 (1)		565.2 (1)
568.7 (6)				568.9 (2)		568.9 (2)
569.5 (6)			569.26 (10)	569.5 (1)		569.5 (1)

Comments on evaluation

1968Bj06	1968Go20	1972Sa06	1975Ar24	1986Ar05	2000Ni13	Recommended
574.0 (6)				575.5 (1)		575.5 (1)
				584.1 (1)		584.1 (1)
			586.1 (8)	586.3 (1)		586.3 (1)
				590.3 (10)		590.3 (10)
				595.4 (2)		595.4 (2)
			596.5 (5)	596.9 (1)		596.9 (1)
			602.7 (5)	602.6 (1)		602.6 (1)
				604.6 (3)		604.6 (3)
612.0 (6)			611.4 (10)	612.0 (1)		612.0 (1)
			616.2 (5)	617.0 (2)		617.0 (2)
				619.0 (2)		619.0 (2)
623.8 (6)			623.6 (5)	624.2 (1)		624.2 (1)
629.1 (6)			627.5 (5)	628.1 (1)		628.1 (1)
			630.6 (10)	629.4 (1)		629.4 (1)
				632.6 (2)		632.6 (2)
			634.5 (10)	634.3 (2)		634.3 (2)
			643.2 (10)	643.2 (2)		643.2 (2) ^x
646.0 (7)			646.2 (10)	646.5 (1)		646.5 (1)
653.7 (7)			653.2 (8)	653.7 (1)		653.7 (1)
			655.0 (8)	655.2 (2)		655.2 (2)
657.0 (7)			658.0 (5)	657.4 (1)		657.4 (1)
			660.6 (10)	659.8 (1)		659.8 (1) ^x
			664.6 (10)	663.9 (1)		663.9 (1)
667.0 (7)			666.7 (6)	666.5 (1)		666.5 (1)
			669.8 (5)	669.7 (1)		669.7 (1)
				675.1 (1)		675.1 (1)
			683.3 (8)	683.9 (2)		683.9 (2)
687.0 (7)			685.5 (10)	685.1 (2)		685.1 (2)
692.8 (7)			692.5 (5)	692.6 (1)		692.6 (1)
699.0 (7)			699.1 (3)	699.03 (5)		699.03 (5)
706.8 (7)			706.0 (2)	705.9 (1)		705.9 (1)
						708.3 (2) ^b
			711.2 (8)	711.5 (1)		711.5 (1) ^x
				713.7 (1)		713.7 (1)
				716.5 (2)		716.5 (2)
				727.8 (2)		727.8 (2)
				730.9 (2)		730.9 (2)
732.9 (7)			733.0 (2)	733.39 (5)		733.39 (5)
737.5 (7)			738.4 (5)	738.0 (1)		738.0 (1)
~ 743.4	742.8 (6)	742.814 (22)	742.67 (20)	742.81 (3)	742.813 (5)	742.813 (5)
			746.5 (15)	745.9 (1)		745.9 (1)
				748.1 (3)		748.1 (3)
756.6 (8)			754.8 (6)	755.0 (1)		755.0 (1)
				758.9 (1)		758.9 (1)
			760 (1)	761.0 (2)		761.0 (2)
				764.8 (2)		764.8 (2)
767.0 (8)	765.0 (7)	766.358 (20)	865.7 (8)	766.4 (2)		766.4 (2)
			768.7 (10)	769.1 (1)		769.1 (1)
				772.4 (2)		772.4 (2)
			777.9 (10)	778.6 (2)		778.6 (2) ^x

1968Bj06	1968Go20	1972Sa06	1975Ar24	1986Ar05	2000Ni13	Recommended
~ 780.9			780.5 (6)	780.4 (2)		780.4 (2)
			783.1 (10)	783.4 (1)		783.4 (1)
~ 787.0	786.3 (5)	786.272 (22)	786.2 (6)	786.27 (3)		786.27 (3)
			793.6 (10)	792.8 (3)		792.8 (3)
				794.9 (2)		794.9 (2)
797.2 (8)			796.2 (5)	796.1 (1)		796.1 (1)
						799.7 (2) ^b
				802.3 (2)		802.3 (2)
804.2 (8)			804.5 (10)	804.1 (1)		804.1 (1)
806.8 (8)			805.5 (5)	805.8 (5)		805.8 (5)
808.0 (8)				808.4 (3)		808.4 (3)
810.0 (8)				810.0 (7)		810.0 (7)
			812.5 (15)	811.5 (1)		811.5 (1)
				814.2 (1)		814.2 (1)
~ 820.2			819.4 (5)	819.2 (1)		819.2 (1)
824.0 (8)			824.7 (5)	824.2 (2)		824.2 (2) ^x
826.3 (8)				825.1 (2)		825.1 (2)
				829.3 (2)		829.3 (2)
832.4 (24)			831.1 (5)	831.5 (1)		831.5 (1)
			841.9 (10)	839.5 (1)		839.5 (1)
~ 845.4			844.1 (10)	844.1 (1)		844.1 (1)
				846.1 (2)		846.1 (2) ^x
				848.9 (2)		848.9 (2)
				851.8 (1)		851.8 (1)
				857.7 (2)		857.7 (2)
				863.2 (2)		863.2 (2)
				869.7 (1)		869.7 (1)
872.0 (26)			872.9 (10)	874.0 (3)		874.0 (3)
876.4 (26)			876.7 (7)	876.0 (1)		876.0 (1)
880.2 (27)	880.0 (7)	880.514 (36)	880.6 (5)	880.5 (1)		880.5 (1)
883.0 (27)	883.0 (6)	883.237 (33)	883.5 (5)	883.24 (4)		883.24 (4)
				890.1 (4)		890.1 (4)
899.3 (27)			898.6 (5)	898.67 (5)		898.67 (5)
905.2 (28)			904.2 (10)	904.2 (1)		904.2 (1)
				916.5 (2)		916.5 (2)
				918.4 (1)		918.4 (1)
				920.5 (2)		920.5 (2) ^x
926 (3)			924.6 (10)	925.0 (1)		925.0 (1)
						926.0 (2) ^a
927.1 (28)			926.7 (5)	926.7 (1)		926.7 (1)
				935.8 (2)		935.8 (2)
				942.0 (3)		942.0 (3)
946.3 (28)	945.8 (3)	946.002 (28)	945.78 (10)	946.00 (3)		946.00 (3)
949.6 (28)				947.7 (2)		947.7 (2)
				952.7 (1)		952.7 (1)
			959 (1)	960.0 (1)		960.0 (1)
966.4 (29)			965.9 (10)	965.8 (1)		965.8 (1)
				975.1 (1)		975.1 (1)
			978.8 (10)	978.2 (3)		978.2 (3)
980.8 (29)			980.5 (5)	980.3 (1)		980.3 (1)

1968Bj06	1968Go20	1972Sa06	1975Ar24	1986Ar05	2000Ni13	Recommended
				981.6 (3)		981.6 (3)
984.5 (29)			983.4 (10)	984.2 (1)		984.2 (1)
				989.5 (1)		989.5 (1)
				992.0 (2)		992.0 (2) ^x
				994.6 (3)		994.6 (3)
				997.7 (3)		997.7 (3)
				1009.9 (3)		1009.9 (3)
				1019.5 (4)		1019.5 (4)
				1021.8 (2)		1021.8 (2)
1023.1 (30)			1022.7 (8)	1023.6 (2)		1023.6 (2) ^x
				1025.3 (2)		1025.3 (2) ^x
1028.6 (30)			1028.1 (8)	1028.7 (1)		1028.7 (1)
				1032.8 (2)		1032.8 (2)
				1035.9 (2)		1035.9 (2) ^x
				1037.9 (2)		1037.9 (2)
				1041.1 (2)		1041.1 (2)
1044.9 (31)				1044.4 (2)		1044.4 (2)
				1051.4 (2)		1051.4 (2)
				1057.8 (3)		1057.8 (3)
				1065.1 (1)		1065.1 (1)
1073 (3)			1074.4 (10)	1073.6 (2)		1073.6 (2)
1084 (3)			1082.5 (6)	1083.2 (1)		1083.2 (1)
				1085.3 (3)		1085.3 (3)
			1108.5 (6)	1106.9 (2)		1106.9 (2)
				1110.6 (1)		1110.6 (1)
1121.9 (33)			1122.3 (6)	1121.7 (1)		1121.7 (1)
				1125.2 (1)		1125.2 (1)
1126.8 (33)			1126.0 (6)	1126.8 (1)		1126.8 (1)
				1151.4 (3)		1151.4 (3)
			1153.1 (6)	1153.5 (3)		1153.5 (3)
			1171.3 (8)	1171.3 (1)		1171.3 (1)
				1173.1 (1)		1173.1 (1)
				1182.1 (2)		1182.1 (2)
		1193.767 (30)		1194.0 (2)		1194.0 (2)
1217 (4)			1217.5 (8)	1217.3 (1)		1217.3 (1)
				1220.4 (2)		1220.4 (2) ^x
1239 (4)				1237.3 (3)		1237.3 (3)
			1240.9 (8)	1241.2 (1)		1241.2 (1)
				1247.8 (2)		1247.8 (2)
				1252.6 (2)		1252.6 (2)
				1256.5 (1)		1256.5 (1)
			1277.1 (8)	1277.7 (2)		1277.7 (2)
1292 (4)			1292.8 (8)	1292.8 (1)		1292.8 (1)
				1296.4 (2)		1296.4 (2) ^x
				1301.2 (2)		1301.2 (2) ^x
				1327.0 (2)		1327.0 (2) ^x
				1342.9 (2)		1342.9 (2)
			1353.0 (6)	1352.9 (1)		1352.9 (1)
1354 (4)				1354.6 (2)		1354.6 (2)
			1358.4 (10)	1359.0 (1)		1359.0 (1)

1968Bj06	1968Go20	1972Sa06	1975Ar24	1986Ar05	2000Ni13	Recommended
				1389.6 (2)		1389.6 (2)
1394 (4)			1394.1 (5)	1393.9 (1)		1393.9 (1)
			1399.7 (10)	1397.5 (2)		1397.5 (2)
				1400.3 (1)		1400.3 (1)
				1409.1 (2)		1409.1 (2)
				1414.4 (2)		1414.4 (2)
			1427 (1)	1426.9 (1)		1426.9 (1)
				1442.8 (2)		1442.8 (2)
1446 (4)			1446.1 (8)	1445.4 (1)		1445.4 (1)
1453 (4)			1452.6 (15)	1452.7 (1)		1452.7 (1)
				1458.9 (1)		1458.9 (1)
				1475.8 (2)		1475.8 (2)
				1485.4 (2)		1485.4 (2)
				1488.0 (2)		1488.0 (2)
1493 (4)			1493.7 (10)	1493.6 (1)		1493.6 (1)
				1496.0 (2)		1496.0 (2)
				1500.0 (2)		1500.0 (2)
				1507.3 (2)		1507.3 (2) ^x
				1510.1 (2)		1510.1 (2)
1516 (5)				1515.6 (2)		1515.6 (2)
				1520.7 (2)		1520.7 (2) ^x
				1538.8 (2)		1538.8 (2) ^x
~1552			1549.2 (10)	1550.1 (1)		1550.1 (1)
				1567.0 (2)		1567.0 (2)
			1580.1 (10)	1579.9 (1)		1579.9 (1)
			1585.4 (10)	1585.9 (1)		1585.9 (1)
1595 (5)			1593.8 (8)	1594.0 (1)		1594.0 (1)
				1618.3 (2)		1618.3 (2)
			1628 (1)	1627.3 (1)		1627.3 (1)
			1638.2 (10)	1638.1 (1)		1638.1 (1)
1640 (5)				1640.5 (3)		1640.5 (3)
				1644.9 (2)		1644.9 (2)
1653 (5)				1650.2 (2)		1650.2 (2)
				1655.7 (1)		1655.7 (1) ^x
				1664.8 (3)		1664.8 (3) ^x
			1668.5 (8)	1668.4 (1)		1668.4 (1)
1671 (5)				1672.8 (1)		1672.8 (1)
				1679.5 (1)		1679.5 (1)
1688 (5)			1686.3 (10)	1685.7 (1)		1685.7 (1)
			1694.0 (8)	1693.8 (2)		1693.8 (2)
1695 (5)				1695.0 (3)		1695.0 (3)
				1700.5 (2)		1700.5 (2)
				1719.7 (2)		1719.7 (2)
				1723.2 (2)		1723.2 (2)
				1727.8 (2)		1727.8 (2)
1736 (5)			1737.9 (10)	1737.7 (2)		1737.7 (2)
				1741.1 (2)		1741.1 (2)
				1743.2 (2)		1743.2 (2) ^x
				1750.0 (1)		1750.0 (1)
1756 (5)				1757.5 (1)		1757.5 (1) ^x

1968Bj06	1968Go20	1972Sa06	1975Ar24	1986Ar05	2000Ni13	Recommended
			1768.4 (15)	1768.0 (3)		1768.0 (3)
				1770.8 (2)		1770.8 (2)
1775 (5)			1772.2 (15)	1773.0 (2)		1773.0 (2)
				1783.7 (2)		1783.7 (2)
			1796.9 (10)	1797.1 (1)		1797.1 (1)
1802 (5)				1805.8 (3)		1805.8 (3)
				1815.3 (3)		1815.3 (3)
				1819.8 (3)		1819.8 (3)
				1825.1 (3)		1825.1 (3)
1828 (5)				1830.8 (3)		1830.8 (3) ^x
			1838.20 (8)	1838.0 (2)		1838.0 (2)
1849 (6)			1850 (1)	1849.8 (2)		1849.8 (2) ^x
			1872.8 (10)	1872.8 (2)		1872.8 (2)
				1884.1 (3)		1884.1 (3)
			1891.1 (10)	1890.1 (2)		1890.1 (2)
				1893.4 (3)		1893.4 (3)
			1897.5 (10)	1896.7 (2)		1896.7 (2)
1905 (6)				1915.5 (3)		1915.5 (3)
			1926.5 (6)	1925.4 (2)		1925.4 (2)
				1927.9 (4)		1927.9 (4) ^x
				1935.2 (4)		1935.2 (4) ^x
1940 (6)			1937.8 (10)	1937.7 (3)		1937.7 (3)
				1958.0 (4)		1958.0 (4)
				1971.2 (4)		1971.2 (4)
				1977.4 (4)		1977.4 (4)
				1989.6 (4)		1989.6 (4)
				2072.2 (4)		2072.2 (4)

a: Expected but as yet unobserved, energy from level scheme.

b: Expected but as yet unobserved, energy from adopted gammas.

x: Not placed in level scheme.

5.2 Relative values of the γ -ray intensities

Measured results for the relative γ -ray intensities from ²³⁴Pa are listed in table 5. The recommended values are from the measurements of 1986Ar05, except as noted in the footnotes of the table.

The values from 1975Ar24 were superseded by the same group in 1986Ar05. The uncertainties of 1968Bj06 are large (~ 20-30 %), and not listed in table. Some γ -ray intensities from 1990Sc09 are also not listed in table because these intensities contain the contributions from ^{234m}Pa decay.

Table 5: Measured and recommended relative γ -ray intensities in decay of ²³⁴Pa

E_γ /keV	I_γ							Recommended
	1967Wa09	1968Bj06 ¹	1975Ar24	1986Ar05	1990Sc09 ¹	2006Al28 ¹	LWEIGHT	
34.30								0.0036 ^f
41.82 ^a								0.27 (7) ^d
43.49		0.123		0.12 (3)				0.12 (3)
45.45		0.009		0.026 (8)				0.026 (8)
54.96 ^b				~ 0.009				~ 0.009
54.96 ^b								~ 0.009
55.45				0.026 (8)				0.026 (8)
58.20		0.0026		< 0.009				0.0026 (8)

E_γ/keV	I_γ							Recommended
	1967Wa09	1968Bj06 ¹	1975Ar24	1986Ar05	1990Sc09 ¹	2006Al28 ¹	LWEIGHT	
59.19				0.031 (10)				0.031 (10)
62.70	3.2	2.45	3.6	1.5 (4)				1.5 (4)
67.25				0.035 (10)				0.035 (10)
69.46				0.017 (7)				0.017 (7)
75.0 ^a								0.030 (6) ^d
79.84		0.11		0.06 (2)				0.06 (2)
97.17				0.23 (8)				0.23 (8)
99.86		4.64		3.1 (5)				3.1 (5)
100.89				0.12 (2)				0.12 (2)
103.77		0.114		0.23 (3)				0.23 (3)
106.68				0.035 (10)				0.035 (10)
125.46	1.2	0.79	0.61	0.76 (9)				0.76 (9)
131.30	18	17.5	17.5	17.5	17.5	17.5		17.5
134.61		0.13		0.11 (2)				0.11 (2)
137.23				0.026 (8)				0.026 (8)
140.15	0.9			0.49 (5)				0.49 (5)
140.91				0.30 (3)				0.30 (3)
143.78	0.2		0.32	0.31 (3)				0.31 (3)
149.88				0.07 (2)				0.07 (2)
152.71	6	5.25	5.78	5.8 (4)	5.08 (19)		5.2 (2)	5.8 (4)
159.48	0.6	0.44	0.61	0.63 (7)				0.63 (7)
164.94				0.05 (2)				0.05 (2)
165.61				0.07 (2)				0.07 (2)
170.85	0.4	0.44	0.55	0.49 (5)				0.49 (5)
174.55			0.21	0.16 (2)				0.16 (2)
179.80				0.043 (15)				0.043 (15)
186.15	1.8	1.5	2.02	1.71 (10)				1.71 (10)
193.73	0.5	0.6	0.51	0.48 (6)				0.48 (6)
196.80		< 0.44	0.06	0.07 (2)				0.07 (2)
199.95				0.07 (2)				0.07 (2)
200.97		0.9	0.96	0.87 (9)				0.87 (9)
203.12	2.1		1.14	1.19 (10)				1.19 (10)
220.00	0.4		0.1	0.14 (2)				0.14 (2)
221.15				0.05 (2)				0.05 (2)
221.83				0.07 (2)				0.07 (2)
226.50	10	5.6	10.1	4.1 (3)	10.25 (15)			4.7 (3) [#]
227.25		5.25		5.6 (3)				5.6 (3)
232.21				0.17 (2)				0.17 (2)
233.6 ^a								~ 0.018 ^d
235.11				0.11 (2)				0.11 (2)
235.9 ^a								0.0044 (25) ^e
240.2				0.05 (2)				0.05 (2)
245.37	0.8	0.9	0.66	0.73 (8)				0.73 (8)
247.79 ^a								3.6 (3)E-4 ^e
249.22	2.5	2.19	2.45	2.4 (3)	2.14 (10)		2.2 (1)	2.4 (3)
257.2				0.05 (2)				0.05 (2)
267.12			0.15	0.17 (2)				0.17 (2)
272.28	1	0.9	0.88	1.05 (10)	1.1 (1)		1.08 (7)	1.05 (10)

E_γ/keV	I_γ							Recommended
	1967Wa09	1968Bj06 ¹	1975Ar24	1986Ar05	1990Sc09 ¹	2006Al28 ¹	LWEIGHT	
275.04				0.09 (2)				0.09 (2)
278.3			0.06	0.04 (1)				0.04 (1)
293.79	3.7		3.4	2.9 (2)	3.0 (1)		2.98 (9)	2.9 (2)
295.91				0.14 (2)				0.14 (2)
298.7				0.013 (5)				0.013 (5)
308.6				0.020 (5)				0.020 (5)
310.2				0.07 (1)				0.07 (1)
310.52 ^a								1.30 (14)E-4 ^e
313.5				0.10 (1)				0.10 (1)
316.7			0.11	0.10 (1)				0.10 (1)
320.4				0.050 (6)				0.050 (6)
330.4 ^b	1.1		0.75	0.75 (5)				0.75 (5)
330.4 ^b								
331.4				0.07 (1)				0.07 (1)
340.2				0.039 (8)				0.039 (8)
343.8				0.033 (7)				0.033 (7)
351.9	0.5	0.6	0.53	0.40 (3)				0.40 (3)
357.9				0.035 (10)				0.035 (10)
360.6				0.017 (6)				0.017 (6)
365.0 ^b				0.017 (6)				0.017 (6)
365.0 ^b								
369.50	3.5	2.63	2.49	2.40 (15)	2.69 (10)		2.60 (8)	2.40 (15)
372.0	1	0.96	1.23	1.18 (8)	1.41 (10)		1.27 (6)	1.18 (8)
379.1				0.04 (1)				0.04 (1)
385.4				0.04 (1)				0.04 (1)
387.94 ^a								6.9 (4)E-4 ^e
394.1				0.09 (1)				0.09 (1)
397.7				0.026 (6)				0.026 (6)
401.8 ^x				0.035 (10)				0.035 (10)
409.8	0.4		0.48	0.33 (3)				0.33 (3)
416.1	0.1			0.035 (10)				0.035 (10)
425.3 ^x				0.035 (10)				0.035 (10)
426.95	0.8		0.47	0.44 (3)				0.44 (3)
427.4 ^a								3.0 (8)E-5 ^e
433.1			0.05	0.09 (1)				0.09 (1)
446.6 ^b			0.11	0.11 (1)				0.11 (1)
446.6 ^b								
450.93 ^a								3.8 (18)E-3 ^e
452.4				0.026 (8)				0.026 (8)
458.68	1.3		1.26	1.10 (6)	1.22 (10)		1.13 (5)	1.10 (6)
461.5 ^b				0.033 (10)				0.033 (10)
461.5 ^b								
464.2				0.03 (1)				0.03 (1)
468.0				0.21 (2)				0.21 (2)
472.3			0.21	0.35 (2)				0.35 (2)
474.2				0.035 (10)				0.035 (10)
478.6 ^b			0.26	0.12 (1)				0.12 (1)
478.6 ^b								
481.0	0.4		0.42	0.30 (2)				0.30 (2)

E_{γ}/keV	I_{γ}							
	1967Wa09	1968Bj06 ¹	1975Ar24	1986Ar05	1990Sc09 ¹	2006Al28 ¹	LWEIGHT	Recommended
498.0 ^b			0.09	0.06 (1)				0.06 (1)
498.0 ^b								
502.0				0.026 (8)				0.026 (8)
506.75	1.5		1.4	1.25 (8)	2.14 (12)		1.7 (5)	1.25 (8)
513.4 ^c	1.3		1.3	1.10 (7)				~ 0.73
513.4 ^c								~ 0.37
519.6				0.38 (3)				0.38 (3)
521.4	1.1	0.9	0.81	0.72 (5)				0.72 (5)
527.9		0.6	0.61	0.38 (3)				0.38 (3)
529.1 ^b	0.3			0.09 (3)				0.09 (3)
529.1 ^b								
534.1				0.08 (1)				0.08 (1)
537.2			0.14	0.08 (1)				0.08 (1)
543.8				0.13 (2)				0.13 (2)
553.7				0.043 (15)				0.043 (15)
558.0 ^b				0.09 (2)				0.09 (2)
558.0 ^b								
559.2				0.07 (2)				0.07 (2)
562.8				0.035 (10)				0.035 (10)
565.2 ^b		0.9		1.00 (6)				1.00 (6)
565.2 ^b								
568.9		2.63		3.5 (4)				3.5 (4)
569.5	14.5	8.75	12.1	8.0 (8)	12.42 (16)	12.9 (38)		8.9 (8) ⁻
575.5				0.026 (8)				0.026 (8)
584.1				0.17 (2)				0.17 (2)
586.3			0.09	0.07 (1)				0.07 (1)
590.3				0.035 (10)				0.035 (10)
595.4				0.09 (2)				0.09 (2)
596.9 ^b			0.31	0.19 (2)				0.19 (2)
596.9 ^b								
602.6	0.4		0.76	0.52 (3)				0.52 (3)
604.6				0.05 (2)				0.05 (2)
612.0		0.6	0.61	0.37 (3)				0.37 (3)
617.0 ^b				0.05 (2)				0.05 (2)
617.0 ^b								
619.0				0.035 (10)				0.035 (10)
624.2		0.35	0.54	0.34 (3)				0.34 (3)
628.1				0.23 (4)				0.23 (4)
629.4		0.35		0.34 (5)				0.34 (5)
632.6				0.035 (10)				0.035 (10)
634.3 ^b				0.13 (2)				0.13 (2)
634.3 ^b								
643.2 ^x				0.026 (8)				0.026 (8)
646.5		0.9	0.19	0.11 (1)				0.11 (1)
653.7 ^b		0.44	0.58	0.45 (6)				0.45 (6)
653.7 ^b								
655.2	0.7			0.13 (2)				0.13 (2)
657.4	0.7	0.9		0.38 (3)				0.38 (3)
659.8 ^x				0.26 (2)				0.26 (2)

E_γ/keV	I_γ						LWEIGHT	Recommended
	1967Wa09	1968Bj06 ¹	1975Ar24	1986Ar05	1990Sc09 ¹	2006Al28 ¹		
663.9			0.9	0.52 (7)				0.52 (7)
666.5	2.2		1.49	1.13 (7)	0.92 (9)		1.05 (6)	1.13 (7)
669.7 ^c	2.0		1.14	0.96 (5)	1.04 (10)		0.98 (5)	0.96 (5)
669.7 ^c								< 0.0005
675.1				0.097 (10)				0.097 (10)
683.9				0.15 (3)				0.15 (3)
685.1 ^b			0.24	0.14 (3)				0.14 (3)
685.1 ^b								
692.6	1.3	1.5	1.4	1.20 (7)				1.20 (7)
699.03 ^b	4.1	3.5	4.16	3.5 (2)	3.61 (10)		3.59 (9)	3.5 (2)
699.03 ^b								
705.9	2.9	3.1	2.14	2.2 (1)				2.2 (1)
708.3 ^a								0.022 (8) ^e
711.5 ^x			0.18	0.15 (2)				0.15 (2)
713.7 ^b				0.14 (2)				0.14 (2)
713.7 ^b								
716.5				0.030 (8)				0.030 (8)
727.8				0.11 (1)				0.11 (1)
730.9				0.61 (8)				0.61 (8)
733.39	9.2	7	7.5	6.7 (4)	7.04 (11)		7.02 (11)	6.7 (4)
738.0		1.75	1.14	1.12 (7)	1.29 (11)		1.17 (6)	1.12 (7)
742.813	2.5		2.0	2.0 (1)				2.0 (1)
745.9			0.11	0.31 (3)				0.31 (3)
748.1				0.10 (2)				0.10 (2)
755.0 ^b	0.6		1.4	1.18 (6)	1.29 (11)		1.21 (5)	1.18 (6)
755.0 ^b								
758.9				0.24 (2)				0.24 (2)
761.0				0.07 (2)				0.07 (2)
764.8				0.19 (4)				0.19 (4)
766.4	0.4	0.26		0.25 (4)				0.25 (4)
769.1				0.18 (1)				0.18 (1)
772.4				0.07 (2)				0.07 (2)
778.6 ^x				0.044 (8)				0.044 (8)
780.4	0.7		0.88	0.87 (4)				0.87 (4)
783.4			0.44	0.29 (3)				0.29 (3)
786.272	1		1.4	1.16 (6)				1.16 (6)
792.8				0.043 (10)				0.043 (10)
794.9				0.65 (8)				0.65 (8)
796.1	3.8		2.9	2.5 (2)	3.31 (15)		2.9 (4)	2.5 (2)
799.7 ^a								
802.3				0.030 (8)				0.030 (8)
804.1			0.35	0.6 (2)				0.6 (2)
805.8	3	2.9	2.71	2.45 (15)				2.45 (15)
808.4				0.035 (10)				0.035 (10)
810.0								0.19 (6) ^f
811.5				0.12 (1)				0.12 (1)
814.2				0.30 (2)				0.30 (2)
819.2	2.8		2.01	1.83 (10)	2.26 (9)		2.05 (22)	1.83 (10)
824.2 ^x			3.23	1.2 (1)				1.2 (1)

E_{γ}/keV	I_{γ}							
	1967Wa09	1968Bj06 ¹	1975Ar24	1986Ar05	1990Sc09 ¹	2006Al28 ¹	LWEIGHT	Recommended
825.1	4.3			1.83 (10)	4.16 (11)			1.83 (10)
829.3				0.35 (10)				0.35 (10)
831.5	5.3		4.46	4.0 (2)	4.77 (9)	3.8 (23)	4.38 (28)	4.0 (2)
839.5				0.030 (7)				0.030 (7)
844.1	0.4		0.44	0.41 (3)				0.41 (3)
846.1 ^x				0.05 (1)				0.05 (1)
848.9				0.026 (7)				0.026 (7)
851.8				0.07 (2)				0.07 (2)
857.7				0.035 (7)				0.035 (7)
863.2				0.07 (2)				0.07 (2)
869.7				0.19 (2)				0.19 (2)
874.0				0.035 (7)				0.035 (7)
876.0	7		3.19	2.45 (2)	2.57 (8)		2.46 (2)	2.45 (2)
880.52 ^c	18		11.64	10.1 (6)	12.97 (12)		11.6 (15)	4.1 (4)
880.52 ^c								6.0 (5)
883.24	4		10.85	9.3 (6)				9.3 (6)
890.1				0.026 (7)				0.026 (7)
898.67	4.3	3.6	3.15	3.15 (20)	3.61 (8)		3.55 (8)	3.15 (20)
904.2	0.5		0.41	0.33 (2)				0.33 (2)
916.5				0.023 (6)				0.023 (6)
918.4				0.096 (10)				0.096 (10)
920.5 ^x				0.028 (7)				0.028 (7)
925.0		8.8		7.6 (5)	8.69 (11)		8.64 (11)	7.6 (5)
926.0 ^a								1.7 (12) ^g
926.7	22	8.75	14.7	8.7 (5)				7.0 (9) ^g
935.8				0.064 (7)				0.064 (7)
942.0				0.044 (7)				0.044 (7)
946.00	19	13.1	16.1	13.0 (8)				13.0 (8)
947.7				1.57 (15)	1.90 (9)		1.81 (8)	1.57 (15)
952.7				0.08 (1)				0.08 (1)
960.0	0.2		0.09	0.07 (1)				0.07 (1)
965.8	0.4	0.7	0.09	0.46 (3)				0.46 (3)
975.1				0.026 (7)				0.026 (7)
978.2				0.087 (20)				0.087 (20)
980.3 ^c	3.8		~ 2.6	1.92 (10)	2.75 (9)			~ 2.6 ^h
980.3 ^c			~ 1.7					~ 1.7 ^h
981.6				0.7 (2)				0.7 (2)
984.2	1.5		1.49	1.57 (15)	1.84 (8)		1.78 (7)	1.57 (15)
989.5				0.10 (1)				0.10 (1)
992.0 ^x				0.08 (2)				0.08 (2)
994.6				0.06 (2)				0.06 (2)
997.7				0.044 (10)				0.044 (10)
1009.9 ^b				0.064 (10)				0.064 (10)
1009.9 ^b								
1019.5				0.026 (7)				0.026 (7)
1021.8	0.4			0.14 (3)				0.14 (3)
1023.6 ^x				0.06 (2)				0.06 (2)
1025.3 ^x				0.05 (2)				0.05 (2)
1028.7	0.8	0.8	0.44	0.55 (3)				0.55 (3)

E_γ/keV	I_γ							
	1967Wa09	1968Bj06 ¹	1975Ar24	1986Ar05	1990Sc09 ¹	2006Al28 ¹	LWEIGHT	Recommended
1032.8				0.017 (4)				0.017 (4)
1035.9 ^x				0.025 (9)				0.025 (9)
1037.9				0.017 (6)				0.017 (6)
1041.1				0.031 (10)				0.031 (10)
1044.4		0.44		~ 0.030				~ 0.030
1051.4				0.06 (1)				0.06 (1)
1057.8				~ 0.017				~ 0.017
1065.1				0.026 (7)				0.026 (7)
1073.6		0.21	0.17	0.10 (1)				0.10 (1)
1083.2	0.6	0.7		0.49 (3)				0.49 (3)
1085.3				0.026 (7)				0.026 (7)
1106.9				0.08 (1)				0.08 (1)
1110.6				0.06 (1)				0.06 (1)
1121.7	0.4		0.44	0.24 (3)				0.24 (3)
1125.2	0.8			0.35 (7)				0.35 (7)
1126.8				0.29 (3)				0.29 (3)
1151.4 ^b				0.031 (9)				0.031 (9)
1151.4 ^b								
1153.5				0.044 (7)				0.044 (7)
1171.3				0.087 (10)				0.087 (10)
1173.1				0.044 (7)				0.044 (7)
1182.1				~ 0.009				~ 0.009
1193.77				0.020 (5)				0.020 (5)
1217.3		0.9	0.32	0.21 (2)				0.21 (2)
1220.4 ^x				0.06 (1)				0.06 (1)
1237.3				< 0.009				< 0.009
1241.2				0.22 (2)				0.22 (2)
1247.8				0.021 (5)				0.021 (5)
1252.6				0.017 (7)				0.017 (7)
1256.5				0.057 (6)				0.057 (6)
1277.7			0.24	0.043 (7)				0.043 (7)
1292.8	0.6	0.7	0.45	0.45 (3)	0.55 (6)		0.47 (3)	0.45 (3)
1296.4 ^x				0.028 (6)				0.028 (6)
1301.2 ^x				0.017 (4)				0.017 (4)
1327.0 ^x				0.017 (4)				0.017 (4)
1342.9				0.012 (4)				0.012 (4)
1352.9	1.7	1.84	1.10	1.12 (5)	1.17 (5)		1.15 (4)	1.12 (5)
1354.6				0.13 (3)				0.13 (3)
1359.0			0.11	0.15 (2)				0.15 (2)
1389.6				0.07 (2)				0.07 (2)
1393.9	2.8	2.2	2.1	2.0 (1)	2.39 (6)		2.2 (2)	2.0 (1)
1397.5				0.08 (2)				0.08 (2)
1400.3				0.17 (2)				0.17 (2)
1409.1				0.043 (8)				0.043 (8)
1414.4				< 0.0026				< 0.0026
1426.9			0.17	0.16 (2)				0.16 (2)
1442.8				0.030 (6)				0.030 (6)
1445.4	0.3			0.31 (3)				0.31 (3)
1452.7	0.9	1	0.7	0.78 (5)	0.74 (6)		0.76 (4)	0.78 (5)

E_{γ}/keV	I_{γ}							LWEIGHT	Recommended
	1967Wa09	1968Bj06 ¹	1975Ar24	1986Ar05	1990Sc09 ¹	2006Al28 ¹			
1458.9				0.09 (2)				0.09 (2)	
1475.8				0.008 (3)				0.008 (3)	
1485.4				0.029 (6)				0.029 (6)	
1488.0				0.013 (5)				0.013 (5)	
1493.6		0.26	0.17	0.10 (1)				0.10 (1)	
1496.0				0.035 (8)				0.035 (8)	
1500.0				0.011 (3)				0.011 (3)	
1507.3 ^x				0.019 (4)				0.019 (4)	
1510.1				< 0.009				< 0.009	
1515.6		0.35		0.07 (1)				0.07 (1)	
1520.7 ^x				~ 0.009				~ 0.009	
1538.8 ^x				0.013 (3)				0.013 (3)	
1550.1			0.09	0.07 (1)				0.07 (1)	
1567.0				0.011 (2)				0.011 (2)	
1579.9			0.15	0.07 (2)				0.07 (2)	
1585.9	0.3		0.26	0.14 (1)				0.14 (1)	
1594.0	0.8	0.6	0.46	0.30 (2)				0.30 (2)	
1618.3				0.009 (3)				0.009 (3)	
1627.3			0.09	0.073 (8)				0.073 (8)	
1638.1	0.3		0.19	0.20 (1)				0.20 (1)	
1640.5		0.6		0.010 (3)				0.010 (3)	
1644.9				0.010 (3)				0.010 (3)	
1650.2				< 0.005				< 0.005	
1655.7 ^x				0.025 (3)				0.025 (3)	
1664.8 ^x				0.017 (6)				0.017 (6)	
1668.41	1		0.33	0.74 (5)	0.74 (5)			0.74 (5)	
1672.8				0.033 (10)				0.033 (10)	
1679.5				0.074 (16)				0.074 (16)	
1685.7				0.30 (2)				0.30 (2)	
1693.8			0.80	0.67 (7)				0.67 (7)	
1695.0	1.1	1.4		0.26 (6)				0.26 (6)	
1700.5				0.10 (1)				0.10 (1)	
1719.7				0.017 (5)				0.017 (5)	
1723.2				0.015 (3)				0.015 (3)	
1727.8				0.019 (4)				0.019 (4)	
1737.7		0.19	0.07	0.072 (8)				0.072 (8)	
1741.1				0.047 (6)				0.047 (6)	
1743.2 ^x				0.032 (7)				0.032 (7)	
1750.0				0.062 (7)				0.062 (7)	
1757.5 ^x				0.023 (5)				0.023 (5)	
1768.0	0.2		0.05	0.019 (4)				0.019 (4)	
1770.8				0.065 (15)				0.065 (15)	
1773.0				0.065 (15)				0.065 (15)	
1783.7				0.024 (6)				0.024 (6)	
1797.1	0.3		0.19	0.23 (2)				0.23 (2)	
1805.8				0.005 (2)				0.005 (2)	
1815.3				0.009 (3)				0.009 (3)	
1819.8				0.004 (1)				0.004 (1)	
1825.1				0.009 (3)				0.009 (3)	

E_{γ}/keV	I_{γ}							
	1967Wa09	1968Bj06 [!]	1975Ar24	1986Ar05	1990Sc09 [!]	2006Al28 [!]	LWEIGHT	Recommended
1830.8 [×]				0.004 (1)				0.004 (1)
1838.0 ^b			0.08	0.040 (9)				0.040 (9)
1838.0 ^b								
1849.8 [×]		0.044		0.027 (6)				0.027 (6)
1872.8				0.034 (8)				0.034 (8)
1884.1				0.015 (4)				0.015 (4)
1890.1	0.4			0.14 (1)				0.14 (1)
1893.4				~ 0.006				~ 0.006
1896.7				0.10 (2)				0.10 (2)
1915.5				0.019 (4)				0.019 (4)
1925.4	0.6		0.28	0.29 (4)	0.31 (3)		0.30 (2)	0.29 (4)
1927.9 [×]				0.052 (10)				0.052 (10)
1935.2 [×]				~ 0.009				~ 0.009
1937.7			0.04	0.04 (1)				0.04 (1)
1958.0				0.0096 (25)				0.0096 (25)
1971.2				~ 0.0026				~ 0.0026
1977.4				0.016 (4)				0.016 (4)
1989.6				0.007 (3)				0.007 (3)
2072.2				0.004 (2)				0.004 (2)

!: Normalized to $I(\gamma_{131.3}) = 17.5$.

#: From $I(\gamma_{227.25}) = 5.6 (3)$ in 1986Ar05 and $I(\gamma_{226.5+\gamma_{227.25}}) = 10.25 (15)$ in 1990Sc09.

~: From $I(\gamma_{568.9}) = 3.5 (4)$ in 1986Ar05 and $I(\gamma_{569.5+568.9}) = 12.42 (16)$ in 1990Sc09.

a: Expected but as unobserved yet.

b: Multiply placed, intensity not divided.

c: Multiply placed, intensity suitably divided.

d: $I(\gamma+ce)$, from γ -ray transition intensity balance.

e: From adopted γ branching.

f: From $I(\gamma+ce)$, from ce measurements(1968Bj06).

g: From $I_{\gamma}(926+926.7) = 8.7 (5)$ and $I_{\gamma}(926.7)/I_{\gamma}(883.2) = 0.75 (8)$ in ²³⁸Pu α decay.

h: From $\gamma\gamma$ coincidence measurements(1968Bj06).

×: Not placed in level scheme.

5.3 Absolute values of the γ -ray emission probabilities

There is no measured absolute γ -ray emission probability in the ²³⁴Pa β^- decay. The normalization factor N for translation of the relative intensities to the absolute emission probabilities has been obtained from the relation of $\Sigma I(\gamma+ce)(g.s.) + \Sigma I(\gamma+ce)(43.5\text{keV level}) = 100 \%$, excluding the 43.5-keV transition and supposing no β^- feeding to the above-mentioned two states. $N = 1.04 (9)$.

The recommended absolute γ -ray emission probabilities (photons per 100 disintegrations) are the relative values recommended in table 5 multiplied by 1.04 (9).

6. References

- 1931Cu01** M.Curie, A.Debierne, A.S.Eve, H.Geiger, O.Hahn, S.C.Lind, S.Meyer, E.Rutherford, E.Schweidler, *Revs.Modern Phys.* 3, 427 (1931) [$T_{1/2}$]
- 1954Zi02** W.L.Zijp, S.Tom, G.J.Sizoo, *Physica* 20, 727 (1954) [$T_{1/2}$]
- 1962Bj01** S.Bjornholm, O.B.Nielsen, *Nuclear Phys.* 30, 488 (1962) [E_γ , I_γ]
- 1967Wa09** A.H.Wapstra, *Nucl.Phys.* A97, 641 (1967) [E_γ , I_γ]
- 1967Wa26** A.H.Wapstra, *Physica* 37, 261 (1967) [ce, X-ray]
- 1968Bj06** S.Bjornholm, J.Borggreen, D.Davies, N.J.S.Hansen, J.Pedersen, H.L.Nielsen, *Nucl.Phys.* A118, 261(1968) [E_γ , E_β , I_γ , I(ce), I_β]
- 1968Go20** J.Godart, A.Gizon, J.Boutet, R.Henck, *Compt.Rend.* 267B, 300 (1968) [E_γ , I_γ]
- 1972Sa06** T.E.Sampson, *Nucl.Instrum.Methods* 98, 37 (1972) [E_γ]
- 1975Ar24** G.Ardisson, C.Ardisson, *Radiochem.Radioanal.Lett.* 21, 357 (1975) [E_γ , I_γ]
- 1986Ar05** C.Ardisson, J.Dalmaso, G.Ardisson, *Phys.Rev.* C33, 2132 (1986) [E_γ , I_γ]
- 1990Sc09** H.L.Scott, K.W.Marlow, *Nucl.Instrum.Methods Phys.Res.* A286, 549 (1990) [P_γ]
- 1996Sc06** E.Schönfeld, H.Janssen, *Nucl. Instrum. Meth. Phys. Res.* A369, 527(1996) [Atomic data].
- 2002Ba85** I.M.Band, M.B.Trzhaskovskaya, C.W.Nestor, Jr., P.O.Tikkanen, S.Raman, *At. Data Nucl. Data Tables* 81, 1 (2002) [ICC]
- 2000Ni13** Y.Nir-El, *Radiochim.Acta* 88, 83 (2000) [E_γ , I_γ]
- 2003Au03** G.Audi, A.H.Wapstra, C.Thibault, *Nucl. Phys.* A729(2003)129 [Q].
- 2006Al28** F.S.Al-Saleh, Al-J.H.Al-Mukren, M.A.Farouk, *Nucl.Instrum.Methods Phys.Res.* A568, 734 (2006) [E_γ , P_γ]
- 2007Br04** E.Browne, J.K.Tuli, *Nucl.Data Sheets* 108, 681 (2007) [NDS]
- 2008Ki07** T.Kibédi, T.W.Burrows, M.B.Trzhaskovskaya, P.M.Davidson, C.W.Nestor, Jr., *Nucl.Instrum.Methods Phys.Res.* A589, 202 (2008) [Theoretical ICC]

^{234m}Pa - Comments on evaluation of the decay data

Huang Xiaolong, Wang Baosong

This evaluation was completed in 2009. Literature available by January 2009 was included.

1 Decay Scheme

^{234m}Pa disintegrates 99.85 (1) % by β^- emissions to levels in ²³⁴U and also 0.15 (1) % through IT decay to ²³⁴Pa. ^{234m}Pa isomer state has $J^\pi = (0)^-$ (2007Br04).

Measured and recommended branching ratios for ^{234m}Pa IT decay are listed in Table 1.

Table 1: Measured and recommended branching ratio for ^{234m}Pa IT decay.

IT (%)	References	Comments
0.150 (25)	1938Fe02	
0.12	1945Br05	Not used
0.63	1954Zi02	Not used
0.18 (2)	1960Fo15	
0.13 (3)	1963Bj02	Deduced by comparing $I_{\text{ce's}}$, $I_{\gamma\text{'s}}$, and β^- disintegration rates from ^{234g} Pa following ^{234m} Pa decay
0.15 (5)	1973Go40	
0.19 (6)		
0.19 (5)	1978Ch06	Deduced from measured $I_\gamma(73.9 \text{ keV})$
0.157 (14)	1990Sc09	Deduced from measured $P_\gamma(131 \text{ keV})$
0.126 (16)	2006Al28	Deduced from measured $P_\gamma(131 \text{ keV})$
0.151 (8)		LWEIGHT
0.15 (1)		Adopted

Statistical processing was performed with the LWEIGHT computer program.

Our recommended IT decay branching ratio is $I_{\text{IT}} = 0.15 (1) \%$ which taken from LWEIGHT result. Thus, $I_{\beta^-} = 99.85 (1) \%$.

The ^{234m}Pa β^- decay scheme was built based mainly on measurement results from 1963Bj02, 1967Wa09 and 1975Ar23. 16 γ -rays were not placed in the current decay scheme. The total photon intensity of these γ transitions is about 0.018 %.

The adopted $Q(\beta^-)$ value of $2269(4) + x \text{ keV}$ has been obtained from $Q(\beta^-) = 2195 (4) \text{ keV}$ for ²³⁴Pa β^- decay (2003Au03), the energy of γ -ray transition 73.92 keV and the estimate of isomeric transition energy $x < 10 \text{ keV}$ deduced from the limit on experimental detection (1973Go40) in ²³⁴Th β^- decay. The adopted $Q(\beta^-)$ is in certain agreement with the effective $Q(\beta^-)$ value of 2259.7 (24) keV, calculated by the evaluators from average radiation energies using the RADLST computer program. This agreement supports the completeness and correctness of the decay scheme.

2 Nuclear Data

The $Q(\beta^-)$ value is from the mass adjustment in 2003Au03 and the energies of γ -ray transitions in ^{234m}Pa IT decay (see above).

Level energies, have been obtained from a least-squares fit to γ -ray energies (GTOL computer code). Spin and parities are from 2007Br04.

The measured and recommended ^{234m}Pa half-life values are listed in Table 2.

Table 2: Measured half-life values of ^{234m}Pa and recommended value

T _{1/2} (min)	References	Comments
1.175 (3)	1951Ba83	
1.25 (10)	1956On07	
1.14 (1)	1963Bj02	
1.183 (37)	1969SaZR	
1.175	1969DeZX	Not used
1.159 (16)	2004Wo02	Evaluated value
1.187 (23)		Unweighted mean
1.159 (11)		LWEIGHT weighted mean, $\chi^2=2.54$
1.159 (11)		Recommended value

The weighted average of 1.15946 for this data set of the 4 values is dominated by the accurate value of 1951Ba83. The LWEIGHT computer program, which uses a Limitation of Relative Statistical Weights (LRSW method), has increased the 1951Ba83 uncertainty from 0.003 to 0.0096 and used a weighted mean and an external uncertainty for recommended average.

Thus, the adopted value of the ^{234m}Pa half-life is 1.159 (11) minute.

2.1 β^- transitions

The maximum energies of the β^- transitions in the decay of ^{234m}Pa have been deduced from the Q(β^-) value (2003Au03), and the level energies which given in Tables 3 and 4.

Table 3: ²³⁴Pa levels populated in ^{234m}Pa IT decay

Level energy (keV)	Spin & parity	Half-life
0.0	4+	6.70 (5) h
73.92 (2)	(3+)	
73.92+x	(0-)	1.159 (11) min

Table 4: ²³⁴U levels populated in ^{234m}Pa β^- decay

Level energy (keV)	Spin & parity	Half-life	β^- transition probabilities (%)
0.0	0+	2.455 (6) $\times 10^5$ a	97.599 (24)
43.428 (14)	2+	0.252 (7) ns	
143.279 (24)	4+		
786.243 (14)	1-		0.049 (3)
809.786 (23)	0+	< 0.1 ns	0.945 (12)
849.18 (7)	3-		
851.56 (4)	2+	> 1.74 ps	
926.659 (20)	2+	1.38 (17) ps	
989.359 (19)	2-	0.76 (4) ns	
1044.469 (15)	0+		1.006 (13)
1085.04 (4)	2+		

Level energy (keV)	Spin & parity	Half-life	β^- transition probabilities (%)
1126.32 (4)	2+		
1174.2 (4)	(1,2+)		0.004 6 (3)
1237.23 (3)	1-		0.012 1 (11)
1435.05 (5)	1-		0.009 2 (11)
1457.40 (8)	(2-)		
1500.8 (3)	(1)		0.013 1 (6)
1553.62 (6)	(1)		0.032 0 (6)
1570.53 (4)	1+		0.002 31 (19)
1591.64 (7)	(1)		0.024 9 (5)
1601.68 (4)	1+		0.001 27 (23)
1666.77 (5)	(1-)		0.006 1 (3)
1693.7? (6)	(1-)		0.002 4 (3)
1781.19 (8)	(0+,1)		0.035 7 (18)
1796.4 (6)	(1)		0.002 1 (3)
1808.97 (7)	(1-)		0.014 6 (7)
1863.11 (7)	(1)		0.003 11 (19)
1874.86 (8)	(1)		0.025 8 (3)
1911.04 (5)	(1-)		0.045 2 (8)
1936.68 (7)	(1)		0.010 8 (3)
1970.0 (5)	(1-)		0.003 89 (22)

The adopted β^- transition probabilities and the associated uncertainties were deduced from the γ transition probability balance at each level of the decay scheme.

The values of $\log ft$ and average β^- energies have been calculated with the program LOGFT.

2.2 γ Transitions

The γ -ray transition probabilities were deduced using the γ -ray emission intensities and the relevant internal conversion coefficients.

Multipolarities and mixing ratios of γ -ray transitions are from 1963Bj02 and 2007Br04.

The internal conversion coefficient (ICC) (and its associated uncertainty) for γ -ray transitions have been interpolated from theoretical values based on the ‘‘Frozen Orbital’’ approximation (2002Ba85) using the BrIcc computer program (2008Ki07).

3. Atomic data

Atomic fluorescence yields ($\omega_K, \omega_L, \omega_M, \eta_{KL}$ and η_{LM}) are from Schönfeld (1996Sc06).

The X-ray and Auger electron emission probabilities have been deduced from γ -ray and conversion electron data by using the computer code RADLST.

The deduced total KX-ray emission probability of 0.67 ± 0.01 %, is in agreement with the measured value of 0.72 (1963Bj02), thus confirming the completeness of the decay scheme.

4. Electron Emissions.

The conversion electron emission probabilities have been deduced from γ -ray transition data using theoretical internal conversion coefficients.

5. Photon Emissions

5.1 γ -ray energies

Measured results for the energies of γ -rays from ^{234m}Pa decay are listed in Table 5. The recommended values were obtained mainly from measurements of 2004Br43, 2000Ni13, 1975Ar23, 1972Sa06 and 1967Wa09 using the LWEIGHT computer program, except as noted in the table.

Table 5: Measured and recommended γ -ray energy values from ^{234m}Pa decay

1963Bj02	1967Wa09	1972Sa06	1975Ar23	2000Ni13	2004Br43	LWEIGHT	Recommended
							< 10 [#]
							41.82 ^a
43.5							43.49 (2) ^b
							62.70 (1) ^a
							73.92 (2) [#]
							99.86 (2) ^{ab}
							135.32 (8) ^a
							137.23 (5) ^a
			140.1 (10)				140.1 (10)
							166.5 (1) ^a
	185.2 (5)		184.7 (5)			185.0 (4)	185.0 (4)
			193.4 (8)				193.4 (8)
							197.91 (15) ^a
			199.9 (10)				199.9 (10)
			203.3 (8)				203.3 (8)
			209.9 (4)				209.9 (4)
							233.6 (2) ^a
							235.9 (3) ^{ab}
236 (1)							236 (1)
			243.5 (8)				243.5 (8) ^x
			247.7 (8)				247.7 (8)
255 (5)	258.0 (5)		258.26 (3)	258.227 (3)		258.227 (3)	258.227 (3)
			275.5 (8)				275.5 (8)
			299.0 (10)				299.0 (10)
			311.0 (10)				311.0 (10)
							316.7 (1) ^a
			338.1 (8)				338.1 (8)
							340.2 (1) ^a
			357.5 (10)				357.5 (10)
			362.8 (10)				362.8 (10)
			387.6 (8)				387.6 (8)
							427.4 (2) ^a
							445.91 (10) ^a
	451.4 (6)		450.97 (10)			450.98 (10)	450.98 (10)
			453.58 (10)				453.58 (10)
			456.7 (10)				456.7 (10)
			468.43 (10)				468.43 (10)
			475.74 (10)				475.74 (10)

Comments on evaluation

1963Bj02	1967Wa09	1972Sa06	1975Ar23	2000Ni13	2004Br43	LWEIGHT	Recommended
							485.44 (7) ^a
			507.5 (10)				507.5 (10)
			509.2 (8)				509.2 (8)
							516.60 (6) ^a
							526.02 (10) ^a
			543.98 (10)				543.98 (10)
							557.24 (6) ^a
			557.3 (10)				557.3 (10) ^x
			572.0 (10)				572.0 (10)
							581.19 (10) ^a
			624.6 (10)				624.6 (10)
			647.7 (8)				647.7 (8) ^x
			649.0 (10)				649.0 (10)
			655.3 (10)				655.3 (10)
			670.8 (10)				670.8 (10)
			673.9 (10)				673.9 (10)
			683.4 (10)				683.4 (10)
			691.0 (3)				691.0 (3)
			695.5 (10)				695.5 (10)
			699.02 (10)				699.02 (10)
			702.0 (1)				702.0 (1)
			705.94 (12)				705.94 (12)
			708.2 (10)				708.2 (10)
							719.01 (7) ^a
			732.5 (10)				732.5 (10)
			740.10 (8)				740.10 (8)
746 (5)	742.7 (6)	742.814 (22)	742.77 (8)	742.813 (5)		742.813 (5)	742.813 (5)
							750.12 (6) ^a
			760.3 (10)				760.3 (10) ^x
							760.53 (15) ^a
765	766.5 (6)	766.358 (20)	766.42 (10)			766.361 (20)	766.361 (20)
			781.75 (10)				781.75 (10)
							783.4 (1) ^a
	786.3 (8)	786.272 (22)	786.28 (10)			786.272 (22)	786.272 (22)
790 (5)							791.94 (5) ^b
806			805.75 (10)				805.75 (10)
			808.2 (1)				808.2 (1)
811							810.0 (7) ^b
			818.2 (5)				818.2 (5)
	825.5 (2)		825.6 (5)			825.5 (2)	825.5 (2)
			844.1 (8)				844.1 (8)
	852.1 (12)		851.58 (10)			851.6 (1)	851.6 (1)
			866.8 (10)				866.8 (10)
		880.514 (36)	880.9 (5)			880.52 (4)	880.52 (4)
		883.237 (33)	883.22 (10)			883.24 (3)	883.24 (3)

1963Bj02	1967Wa09	1972Sa06	1975Ar23	2000Ni13	2004Br43	LWEIGHT	Recommended
			887.29 (10)				887.29 (10) ^x
			921.72 (10)				921.72 (10)
			926.61 (10)				926.61 (10)
			936.3 (10)				936.3 (10)
			941.96 (10)				941.96 (10)
		946.002 (28)	945.94 (2)			945.961 (16)	945.961 (16)
			960.0 (10)				960.0 (10)
			996.1 (20)				996.1 (20)
1001	1001.3 (5)	1001.025 (22)	1000.99 (10)		1001.03 (3)	1001.026 (18)	1001.026 (18)
1045			1041.70 (10)				1041.70 (10)
			1059.4 (8)				1059.4 (8)
			1061.86 (10)				1061.86 (10)
			1081.9 (10)				1081.9 (10)
			1084.25 (10)				1084.25 (10)
			1120.6 (8)				1120.6 (8)
	1125.2 (8)		1124.93 (10)			1124.93 (10)	1124.93 (10)
1160			1174.2 (10)				1174.2 (10)
	1194.2 (6)	1193.767 (30)	1193.73 (12)			1193.77 (3)	1193.77 (3)
			1220.37 (10)				1220.37 (10) ^x
	1238.0 (7)		1237.26 (10)			1237.28 (10)	1237.28 (10)
			1353.0 (15)				1353.0 (15) ^x
	1392 (2)		1392.7 (10)			1392.6 (9)	1392.6 (9)
	1414.7 (10)		1413.88 (10)			1413.89 (10)	1413.89 (10)
1440	1435.5 (8)		1434.14 (10)			1434.16 (10)	1434.16 (10)
			1458.5 (15)				1458.5 (15)
			1501 (2)				1501 (2)
	1510.9 (7)		1510.21 (10)			1510.22 (10)	1510.22 (10)
	1528.2 (12)		1527.27 (10)			1527.28 (10)	1527.28 (10)
			1550.0 (10)				1550.0 (10)
	1554.7 (8)		1553.75 (10)			1553.77 (10)	1553.77 (10)
			1558.4 (10)				1558.4 (10)
	1570.6 (12)		1570.67 (10)			1570.67 (10)	1570.67 (10)
	1593.4 (7)		1593.8 (10)			1593.5 (6)	1593.5 (6)
			1601.8 (15)				1601.8 (15)
			1667.6 (10)				1667.6 (10)
			1694.1 (10)				1694.1 (10)
			1720.5 (15)				1720.5 (15) ^x
			1732.2 (15)				1732.2 (15) ^x
	1738.5 (7)		1737.75 (10)			1737.77 (10)	1737.77 (10)
1750	1759 (2)		1759.81 (10)			1759.81 (10)	1759.81 (10)
	1765.5 (6)		1765.44 (10)			1765.44 (10)	1765.44 (10)
	1796.5 (20)		1796.2 (10)			1796.3 (9)	1796.3 (9)
	1809.4 (7)		1809.04 (10)			1809.05 (10)	1809.05 (10)
			1819.69 (10)				1819.69 (10)
	1831.9 (10)		1831.36 (10)			1831.37 (10)	1831.37 (10)

1963Bj02	1967Wa09	1972Sa06	1975Ar23	2000Ni13	2004Br43	LWEIGHT	Recommended
			1863.09 (10)				1863.09 (10)
	1868.6 (8)		1867.69 (10)			1867.7 (1)	1867.7 (1)
	1876.3 (8)		1874.88 (10)			1874.9 (1)	1874.9 (1)
	1893.5 (8)		1893.51 (11)			1893.51 (11)	1893.51 (11)
	1911.5 (7)		1911.19 (11)			1911.20 (11)	1911.20 (11)
			1926.5 (10)				1926.5 (10)
	1937.5 (7)		1937.04 (13)			1937.01 (13)	1937.01 (13)
	1970.4 (10)		1970.0 (15)			1970.3 (8)	1970.3 (8)
					2022.24 (12)		2022.24 (12) ^x
					2041.23 (13)		2041.23 (13) ^x
					2065.80 (13)		2065.80 (13) ^x
					2093.19 (38)		2093.19 (38) ^x
					2102.14 (15)		2102.14 (15) ^x
					2136.69 (14)		2136.69 (14) ^x

#: IT decay, energy from 1973Go40.

a: Expected but as yet unobserved, energy from adopted gammas.

b: Energy from ²³⁸Pu α decay.

x: Not placed in level scheme.

5.2 Relative values of the γ-ray intensities

Measurements of the relative γ-ray intensities from ^{234m}Pa are listed in table 6. The recommended values have been obtained with the LWEIGHT computer program using measurement results from 2006Al28, 2004Br43, 2000Ni13, 1992Si17, 1990Sc09, 1986Mo09, 1975Ar23 1971GuZQ and 1967Wa09.

As the measured results of 1990Sc09 and 1971GuZQ contained the contributions from ^{234g}Pa β⁻ decay, these contributions had to be estimated and removed from the values cited in 2007Br04. Also the measurement results of 1963Bj02 have been rejected and not listed in table as the associated uncertainties are not given.

Table 6: Measured and recommended relative γ-ray intensities from ^{234m}Pa decay

<i>E_γ</i> /keV	<i>I_γ</i>										Recommended
	1967Wa09	1971GuZQ	1975Ar23	1986Mo09	1990Sc09	1992Si17	2000Ni13	2004Br43!	2006Al28!	LWEIGHT	
< 10											17.7 (12) [#]
41.82 ^a											1.61 (8) ^b
43.49											166.8 (4) ^b
62.70 ^a											0.15 (4) ^d
73.92											1.53 (12) [#]
99.86 ^a											0.96 (7) ^b
135.32 ^a											0.000 50 (6) ^d
137.23 ^a											0.0057 (21) ^d
140.1			< 0.15								< 0.15
166.5 ^a											0.000 028 (6) ^d
185.0	0.2 (1)		0.203 (17)							0.203 (17)	0.203 (17)
193.4			0.085 (17)								0.085 (17)
197.91 ^a											0.003 2 (7) ^d

E_{γ}/keV	I_{γ}										Recommended
	1967Wa09	1971GuZQ	1975Ar23	1986Mo09	1990Sc09	1992Si17	2000Ni13	2004Br43!	2006Al28!	LWEIGHT	
199.9			0.068 (14)								0.068 (14)
203.3			0.122 (24)		0.145 (12) ^c				0.14 (1)		0.14 (1)
209.9			0.156 (17)								0.156 (17)
233.6 ^a			0.059 (12)								≈ 0.1 ^e
235.9 ^a											0.010 (4) ^d
236											8.7 (9) ^f
243.5 ^x			0.059 (10)								0.059 (10)
247.7			0.114 (26)								0.114 (26)
258.227	6.7 (17)	8.82 (24)	9.66 (39)		8.70 (4)	8.6 (6)	9.08 (24)		8.46 (33)	8.72 (4)	8.72 (4)
275.5			0.037 (7)								0.037 (7)
299.0			0.076 (15)								0.076 (15)
311.0			0.061 (12)								0.061 (12)
316.7 ^a											0.022 (5) ^d
338.1			0.134 (27)								0.134 (27)
340.2 ^a											0.008 5 (25) ^d
357.5			0.095 (20)								0.095 (20)
362.8			0.081 (17)								0.081 (17)
387.6 ^g			0.170 (17)								0.115 (17)
387.6 ^g											0.056 (4) ^d
427.4 ^a											0.002 4 (6) ^d
445.91 ^a											0.003 6 (8) ^d
450.98	0.42 (10)	0.42 (5)	0.356 (34)		0.358 (19)	0.39 (8)			0.366 (15)		0.366 (15)
453.58		0.254 (24)	0.288 (34)		0.23 (2)	0.31 (6)			0.251 (14)		0.251 (14)
456.7			0.085 (17)								0.085 (17)
468.43		0.204 (41) ^c	0.280 (27)		0.237 (16) ^c			0.19 (10) ^c	0.243 (13)		0.243 (13)
475.74		0.209 (42)	0.339 (34)		0.274 (18)	0.34 (9)			0.280 (15)		0.280 (15)
485.44 ^a											0.002 2 (2) ^d
507.5			0.187 (17)								0.187 (17)
509.2			0.254 (34)								0.254 (34)
516.60 ^a											0.001 44 (19) ^d
526.02 ^a											0.001 06 (14) ^d
543.98		0.40 (8) ^c	0.441 (51)		0.404 (19) ^c	0.46 (6)		0.32 (21) ^c	0.412 (17)		0.412 (17)
557.24 ^a											0.000 98 (13) ^d
557.3 ^x			0.085 (19)								0.085 (19)
572.0			0.103 (20)								0.103 (20)
581.19 ^a											0.009 4 (11) ^d
624.6			0.170 (17)								0.013 7 (14) ^d
647.7 ^x			0.187 (17)								0.187 (17)
649.0 ^g			0.127 (25)								0.007 (1)
649.0 ^g											0.12 (3)
655.3			0.164 (17)								0.164 (17)
670.8			0.044 (10)								0.044 (10)
673.9			0.076 (15)								0.076 (15)
683.4			0.068 (14)								0.068 (14)

Comments on evaluation

E_γ/keV	I_γ										Recommended
	1967Wa09	1971GuZQ	1975Ar23	1986Mo09	1990Sc09	1992Si17	2000Ni13	2004Br43!	2006Al28!	LWEIGHT	
691.0		1.09 (6)	0.932 (85)		1.073 (23)	0.92 (10)				1.06 (2)	1.06 (2)
695.5			0.187 (17)			0.28 (6)				0.194 (16)	0.194 (16)
699.02		0.70 (7)	0.095 (19)		0.68 (3)					0.68 (3)	0.68 (3)
702.0		0.85 (8)	0.915 (85)		0.846 (20)	0.93 (10)			0.67 (34)	0.852 (17)	0.852 (17)
705.94		0.72 (7) ^c	0.481 (51)		0.656 (16) ^c	0.47 (12)				0.61 (6)	0.61 (6)
708.2			< 0.085								< 0.085
719.01 ^a											0.003 02 (24) ^d
732.5			0.154 (17)								0.154 (17)
740.10		1.33 (12)	1.20 (12)		1.41 (3)	1.26 (12)				1.39 (3)	1.39 (3)
742.813	13.3 (17)	11.12 (24) ^c	9.59 (39)	11.3 (7)	10.93 (8) ^c	10.4 (5)	12.27 (23) ^c			11.13 (28)	11.13 (28)
750.12 ^a											0.002 02 (27) ^d
760.3 ^x			0.187 (17)								0.187 (17)
760.53 ^a											0.000 5 (1) ^d
766.361	36.7 (67)	37.8 (4) ^c	35.1 (14)	39.91 (84)	38.36 (25) ^c	37.6 (11)			35.7 (16)	38.2 (2)	38.2 (2)
781.75		0.845 (85)	0.898 (85)		0.93 (2)	0.86 (12)				0.923 (19)	0.923 (19)
783.4 ^a											0.004 6 (8) ^d
786.272	5 (1)	6.41 (12) ^c	5.80 (22)	6.36 (46)	6.37 (6) ^c	5.97 (33)				6.33 (5)	6.33 (5)
791.94											0.001 17 (15) ^d
805.75		0.718 (38) ^c	0.509 (51)		0.820 (15) ^c	0.49 (15)				0.73 (9)	0.73 (9)
808.2		0.34 (4) ^c	0.356 (34)		0.303 (30) ^c	0.39 (10)				0.332 (19)	0.332 (19)
810.0											85 ^e
818.2			0.119 (34)								0.119 (34)
825.5	0.42 (25)	0.489 (25) ^c	0.168 (34)		0.547 (14) ^c					0.46 (9)	0.17 (4)
844.1			0.129 (27)								0.129 (27)
851.6	0.67 (25)	0.879 (48) ^c	0.746 (68)		0.820 (17) ^c	0.83 (9)				0.822 (16)	0.822 (16)
866.8		0.145 (24)	0.127 (26)							0.137 (18)	0.137 (18)
880.52		0.438 (9) ^c	0.458 (51)		0.468 (4) ^c	0.52 (16)				0.463 (4)	0.463 (4)
883.24		0.428 (13) ^c	0.424 (34)		0.453 (4) ^c	0.50 (16)			0.38 (10)	0.450 (4)	0.450 (4)
887.29 ^x		0.761 (36)	0.882 (85)		0.846 (15)	0.90 (12)				0.836 (14)	0.836 (14)
921.72		1.51 (7)	1.41 (14)		1.51 (2)	1.40 (15)			1.34 (45)	1.506 (19)	1.506 (19)
926.61		0.215 (11) ^c	0.148 (15)		0.213 (3) ^c					0.202 (18)	0.148 (15)
936.3		0.091 (23)	0.22 (5)							0.12 (2)	0.12 (2)
941.96		0.282 (24) ^c	0.356 (34)		0.289 (12) ^c	0.33 (6)				0.295 (10)	0.295 (10)
945.961		1.33 (3) ^c	1.19 (12)	1.27 (15)	1.242 (11) ^c	1.25 (37)			1.18 (31)	1.252 (10)	1.252 (10)
960.0			0.102 (34)								0.102 (34)
996.1		0.90 (5)	0.492 (85)			0.51 (10)				0.7 (2)	0.7 (2)
1001.026	100	100	100	100	100	100	100	100	100		100
1041.70		0.111 (22)	0.170 (17)		0.137 (11)					0.141 (9)	0.141 (9)
1059.4			0.131 (26)								0.131 (26)
1061.86		0.290 (24)	0.237 (17)		0.274 (15)	0.25 (10)				0.264 (10)	0.264 (10)
1081.9			0.107 (22)								0.107 (22)
1084.25			0.058 (10)		0.136 (11) ^c					0.10 (4)	0.10 (4)
1120.6			0.204 (17)								0.204 (17)
1124.93 ^g	0.50 (17)	0.495 (21) ^c	0.475 (51)		0.436 (14) ^c	0.48 (8)				0.456 (11)	0.046 (1)

E_{γ}/keV	I_{γ}										Recommended
	1967Wa09	1971GuZQ	1975Ar23	1986Mo09	1990Sc09	1992Si17	2000Ni13	2004Br43!	2006Al28!	LWEIGHT	
1124.93 ^{&}											0.41 (1)
1174.2			0.227 (22)								0.227 (22)
1193.77	1.33 (33)	1.615 (36) ^c	1.525 (85)		1.606 (16) ^c	1.58 (14)			1.67 (78)	1.605 (15)	1.605 (15)
1220.37 ^x		0.106 (23)	0.119 (34)		0.107 (11)					0.108 (10)	0.108 (10)
1237.28	0.50 (17)	0.592 (24)	0.610 (68)		0.632 (12)	0.58 (13)				0.623 (11)	0.623 (11)
1353.0 ^x		0.271 (27)	0.075 (15)		0.226 (10)					0.18 (6)	0.18 (6)
1392.6	0.50 (25)	0.447 (48)	0.187 (17)		0.465 (5)					0.34 (12)	0.34 (12)
1413.89	0.2 (1)	0.279 (17)	0.254 (17)		0.274 (12)					0.270 (9)	0.270 (9)
1434.16	1.17 (33)	1.09 (6)	0.99 (10)		1.156 (15)	1.12 (17)				1.149 (15)	1.149 (15)
1458.5			0.220 (51)								0.22 (5)
1501			0.153								0.153
1510.22	1.83 (33)	1.57 (4)	1.54 (10)		1.538 (19)	1.59 (15)				1.545 (17)	1.545 (17)
1527.28	0.33 (10)	0.263 (16)	0.254 (34)		0.286 (11)					0.277 (9)	0.277 (9)
1550.0		0.153 (11) ^c	0.220 (17)		0.151 (9) ^c					0.162 (17)	0.162 (17)
1553.77	1.0 (2)	0.990 (24)	1.068 (85)		0.966 (16)	1.07 (18)				0.976 (13)	0.976 (13)
1558.4		0.085 (12)	0.090 (19)							0.086 (10)	0.086 (10)
1570.67	0.10 (4)	0.127 (19)	0.146 (34)		0.131 (11)					0.130 (9)	0.130 (9)
1593.5	1.33 (33)	0.284 (3) ^c	0.458 (51)		0.253 (9) ^c	0.45 (19)				0.278 (13)	0.278 (13)
1601.8			0.056 (25)								0.056 (25)
1667.6		0.145 (12)	0.098 (21)		0.143 (9)					0.139 (7)	0.139 (7)
1694.1		0.044 (4) ^c	0.054 (10)		0.044 (3) ^c					0.0445 (23)	0.0445 (23)
1720.5 ^x			0.039 (17)								0.039 (17)
1732.2 ^x			0.220 (34)								0.220 (34)
1737.77	3.0 (4)	2.545 (24) ^c	2.41 (10)		2.51 (3) ^c	2.45 (25)				2.528 (18)	2.528 (18)
1759.81 ^x	0.33 (17)	0.174 (7)	0.271 (34)		0.167 (7)					0.173 (5)	0.173 (5)
1765.44	1.17 (33)	0.918 (24)	1.04 (10)		1.037 (15)	1.01 (25)				0.99 (6)	0.99 (6)
1796.3	0.10 (7)	0.036 (6) ^c	0.037 (7)							0.037 (5)	0.037 (5)
1809.05	0.4 (1)	0.447 (12)	0.508 (51)		0.441 (9)	0.46 (9)				0.444 (7)	0.444 (7)
1819.69		0.103 (7) ^c	0.141 (31)		0.106 (8)					0.105 (5)	0.105 (5)
1831.37	2.33 (33)	2.114 (24)	1.90 (7)		2.05 (3)	2.09 (21)				2.077 (18)	2.077 (18)
1863.09		0.139 (11)	0.144 (29)		0.143 (6)					0.142 (5)	0.142 (5)
1867.7	1.33 (33)	1.105 (11)	0.90 (9)		1.097 (16)	1.15 (17)				1.101 (9)	1.101 (9)
1874.9	1.17 (33)	0.942 (24)	0.932 (85)		0.977 (15)	0.97 (14)				0.967 (13)	0.967 (13)
1893.51	0.33 (10)	0.256 (11) ^c	0.254 (17)		0.260 (8) ^c	0.26 (7)				0.258 (6)	0.258 (6)
1911.20	0.83 (17)	0.737 (12)	0.627 (68)		0.751 (12)	0.74 (11)				0.742 (8)	0.742 (8)
1926.5		0.057 (5) ^c	0.053 (10)		0.049 (5) ^c					0.053 (4)	0.053 (4)
1937.0	0.4 (1)	0.336 (7) ^c	0.356 (34)		0.335 (8) ^c	0.38 (9)				0.336 (5)	0.336 (5)
1970.3	0.033 (33)	0.0483 (36)	0.066 (14)							0.049 (4)	0.049 (4)
2022.24 ^x									0.022 (2)		0.022 (2)
2041.23 ^x									0.013 (1)		0.013 (1)
2065.80 ^x									0.008 4 (12)		0.008 4 (12)
2093.19 ^x									0.002 4 (7)		0.002 4 (7)
2102.14 ^x									0.007 2 (10)		0.007 2 (10)
2136.69 ^x									0.008 4 (5)		0.008 4 (5)

- #: I($\gamma+ce$), from IT decay.
 a: Expected but as yet unobserved.
 b: From γ -ray transition intensity balance.
 c: Removed the contributions from ^{234g}Pa β^- decay.
 d: Deduced from adopted γ branching in 2007Br04.
 e: I($\gamma+ce$), from $I(\gamma+ce)(\gamma234)/I(\gamma1042) \approx 0.7$ in ^{234}Np ϵ decay.
 f: I($\gamma+ce$), from measured $I_{ce}(K) = 70$.
 g: I($\gamma+ce$), from $I_{ce}(810)/I(\gamma1001) = 0.51 / 0.6$ in 1963Bj02.
 &: Multiply placed, intensity suitably divided.
 x: Not placed in level scheme.

5.3 Absolute values of the γ -ray emission probabilities

Measurements of the absolute γ -ray emission probability of 1001.026 keV per 100 disintegrations of ^{234m}Pa β^- -decay and three weighted average results are listed in Table 7.

It should be noted that the uncertainties quoted in 1990Sc09, 1986Mo09, and 1971GuZQ are questionable (perhaps, only statistical errors were included) when compared with the data of 1992Si17 who used a purified ^{234m}Pa source. Thus 2 % systematic uncertainty was added by the evaluators to those measurement results.

Table 7: Measured and recommended absolute emission probability of the 1001.026 keV γ -ray per 100 disintegrations of ^{234m}Pa β^- decay

$P_\gamma(1001.026 \text{ keV})$ (%)	References	Comments
0.59 (10)	1963Bj02	scintillation spectrometers
0.828 (18)	1971GuZQ	
0.92	1982Mo30	Not used
0.834 (21)	1986Mo09	Ge(Li)
0.839 (20)	1990Sc09	HPGe
0.818 (30)	1992Ja17	
0.788 (43)	1992Li05	
0.845 (21)	1992Si17	HPGe, 0.844 104 with another method
0.910 (25)	1993Su37	
0.924 (17)	1999An40	HPGe
0.861 (15)	2003Yu06	n-type Ge detector
0.923 (30)	2006Al28	HPGe, from extended sample
0.835 (11)	1998Ad08	Evaluation
0.835 (4)	1999Nz01	Evaluation
0.862 (13)		Average of all measurements with LWEIGHT program, $\chi^2 = 3.7$
0.856 (12)		Average of all measurements with Normalised residuals method
0.848 (8)		Average of all measurements with Rajput and MacMahon method
0.848 (8)		Recommended value

The recommended value of the absolute γ -ray emission probability of the 1001.026 keV γ -ray is obtained with the method of averaging discrepant data of Rajput and MacMahon (1992Ra08) and adopted as the normalization factor N, with $N = 0.008\ 48\ (8) \times 0.998\ 5\ (1)$.

Thus, the recommended absolute γ -ray emission probabilities are the relative values recommended in Table 6 multiplied by 0.008 47 (8).

6. References

- 1938Fe02 N.Feather, E.Bretscher, Proc.Roy.Soc.(London) 165A, 530 (1938) [IT Branching Ratio]
- 1945Br05 H.Bradt, P.Scherrer, Helv.Phys.Acta 18, 405 (1945) [IT Branching Ratio]
- 1951Ba83 F.Barendregt, S.Tom, Physica 17, 817 (1951) [$T_{1/2}$]
- 1954Zi02 W.L.Zijp, S.Tom, G.J.Sizoo, Physica 20, 727 (1954) [$T_{1/2}$]
- 1956On07 Ong Ping Hok, J.T.Verschoor, P.Born, Physica 22, 465 (1956) [$T_{1/2}$]
- 1960Fo15 J.H.Forrest, S.J.Lyle, G.R.Martin, J.J.Maulden, J.Inorg.Nucl.Chem. 15, 210 (1960) [IT Branching Ratio]
- 1963Bj02 S.Bjornholm, O.B.Nielsen, Nucl.Phys. 42, 642 (1963) [E_γ , I_γ , I_{KX} , P_γ , $T_{1/2}$, IT Branching Ratio]
- 1967Wa09 A.H.Wapstra, Nucl.Phys. A97, 641 (1967) [E_γ , I_γ]
- 1969DeZX R.Denig, N.Trautmann, N.Kaffrell, G.Herrmann, Proc.Int.Conf. Protactinium, 3rd, Schloss Elmau, Germany (1969) [$T_{1/2}$]
- 1969SaZR M.Saeki, K.Kimura, T.Ishimori, JAERI-1178, p.25 (1969) [$T_{1/2}$]
- 1971GuZQ R.Gunnink, J.F.Tinney, UCRL-51086 (1971) [E_γ , I_γ , P_γ]
- 1972Sa06 T.E.Sampson, Nucl.Instrum.Methods 98, 37 (1972) [E_γ]
- 1973Go40 J.Godart, A.Gizon, Nucl.Phys. A217, 159 (1973) [IT Branching Ratio]
- 1975Ar23 G.Ardisson, C.Marsol, Nuovo Cim. 28A, 155 (1975) [E_γ , I_γ]
- 1978Ch06 Y.Y.Chu, G.Scharff-Goldhaber, Phys.Rev. C17, 1507 (1978) [IT Branching Ratio]
- 1982Mo30 M.H.Momeni, Nucl.Instrum.Methods 193, 185 (1982) [P_γ]
- 1986Mo09 C.E.Moss, Radiat.Eff. 94, 81 (1986) [P_γ]
- 1990Sc09 H.L.Scott, K.W.Marlow, Nucl.Instrum.Methods Phys.Res. A286, 549 (1990) [P_γ]
- 1992Ja17 P.Jagam, J.J.Simpson, J.Radioanal.Nucl.Chem. 166, 393 (1992) [P_γ]
- 1992Li05 W.-J.Lin, G.Harbottle, J.Radioanal.Nucl.Chem. 157, 367 (1992) [P_γ]
- 1992Ra08 M.U.Rajput, T.D.Mac Mahon, Nucl. Instrum. Methods Phys.Res. A312, 289 (1992) [Techniques for Evaluating Discrepant Data]
- 1992Si17 K. Siemon, R.A. Esterlund, J. Van Aarle, M. Knaack, W. Westmeier, P. Patzelt, Appl. Radiat. Isot. 43, 873 (1992) [P_γ]
- 1993Su37 G.A.Sutton, S.T.Napier, M.John, A.Taylor, Sci.Total Environ. 130/131, 393 (1993) [P_γ]
- 1996Sc06 E.Schönfeld, H.Janben, Nucl. Instrum. Meth. Phys. Res. A369, 527 (1996) [Atomic data].
- 1998Ad08 I.Adsley, J.S.Backhouse, A.L.Nichols, J.Toole, Appl.Radiat.Isot. 49, 1337 (1998) [Evaluated P_γ]
- 1999An40 S.Anilkumar, N.Krishnan, M.C.Abani, Appl.Radiat.Isot. 51, 725 (1999) [P_γ]
- 1999Nz01 A.C.Nzuruba, Nucl.Instrum.Methods Phys.Res. A424, 425 (1999) [Evaluated data]
- 2000Ni13 Y.Nir-El, Radiochim.Acta 88, 83 (2000) [E_γ , I_γ]
- 2002Ba85 I.M.Band, M.B.Trzhaskovskaya, C.W.Nestor, Jr., P.O.Tikkanen, S.Raman, At. Data Nucl. Data Tables 81, 1 (2002) [ICC]
- 2003Au03 G.Audi, A.H.Wapstra, C.Thibault, Nucl. Phys. A729(2003)129 [Q].
- 2003Yu06 H.Yucel, H.Karadeniz, M.A.Cetiner, H.Demirel, S.Turhan, J.Radioanal.Nucl.Chem. 258, 445 (2003) [P_γ]
- 2004Br43 V.B.Brudanin, K.Ya.Gromov, S.I.Vasiliev, A.A.Klimenko, A.A.Smolnikov, V.I.Fominykh, V.G.Chumin, Part. and Nucl., Lett. 122, 84 (2004) [E_γ , I_γ]
- 2004Wo02 M.J.Woods, S.M.Collins, Appl.Radiat.Isot. 60, 257 (2004) [Evaluated $T_{1/2}$]
- 2006Al28 F.S.Al-Saleh, Al-J.H.Al-Mukren, M.A.Farouk, Nucl. Instrum. Methods Phys. Res. A568, 734 (2006) [P_γ]
- 2007Br04 E.Browne, J.K.Tuli, Nucl.Data Sheets 108, 681 (2007) [NDS]
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr., Nucl.Instrum.Methods Phys.Res. A589, 202 (2008) [Theoretical ICC]

²³⁴U - Comments on evaluation of decay data by V. Chisté and M.M. Bé

This evaluation was completed in 2005. Literature available by September 2005 was included.

1 Decay Scheme

²³⁴U disintegrates by alpha emission to excited and ground state levels of ²³⁰Th. Spin and half-lives of excited states are from the mass-chain evaluation of Y.A. Akovali (1993Ak02 to A = 230, and 1994Ak05 to A = 234).

2 Nuclear Data

The Q value is from atomic mass evaluation of Audi et al. (2003Au03).

The experimental ²³⁴U half-life values (in years) are given in Table 1:

Table 1: Experimental values of ²³⁴U half-life.

Reference	Original value (10 ⁵ a)	Revised Value by Holden (1981HoZI and 1989Ho24)	Comments
Nier (1939Ni03)	2.70 (27)		Not used.
Chamberlain (1946Ch02)	2.29 (14)		Not used. Measurements of relative abundance of ²³⁴ U and ²³⁸ U.
Chamberlain (1946Ch02)	2.35 (14)		Not used. Measurements of α -activity of ²³⁴ U.
Baldinger (1949Ba41)	2.33 (10)		Not used.
Goldin (1949Go18)	2.67 (4)		Not used.
Kienberger (1949Ki26)	2.552 (8)		Not used. Superseded 1952Ki19
Fleming (1952Fl20)	2.475 (16)	2.475 (24)	Not used. Uncertainty increased for missing details.
Kienberger (1952Ki19)	2.520 (8)		Not used.
White (1965Wh05)	2.47 (3)		Not used.
Meadows (1970MeZN)	2.439 (14)	2.439 (18)	Not used. Uncertainty increased for missing details.
de Bievre (1972DeYN)	2.446 (7)	2.450 (9) *	Revised by author (see 1989Ho24)
Lounsbury (1972LoZL)	2.444 (6)	2.458 (13) *	Revised by author (see 1989Ho24)
Geidel'man (1980Ge13)	2.4604 (45)	2.459 (9) *	4 $\pi\alpha$ - x coincidence. Revised uncertainty for missing details.
	2.4570 (45)		Liquid scintillator. Revised uncertainty for missing details.
Poenitz (1983 and 1985 Poenitz)	2.457 (5)		Not used.
Davideenam (1984Davideenam)	2.457 (5)		Not used. Evaluated value.
Recommended value		2.455 (6)	reduced $\chi^2 = 0.28$

The first six and less precise values (1940's) were omitted from analysis. For remaining values, the evaluators have chosen to take into account the recommendations given by N.E. Holden (1989Ho24), thus the only three experimental values (*) with associated uncertainties used to the weighted average are 1972DeYN, 1972LoZL and 1980Ge13. For the data in 1980Ge13, the evaluators have chosen to use the average value of 2.459 (9) $\times 10^5$ a, calculated from two experimental values given in the paper to produce a single DDEP value from each laboratory. A weighted average has been calculated using LWEIGHT computer program (version 3). However, the treatment of uncertainties in 1989Ho24 ("... when detailed information on the uncertainties was available in each of these experiments, the standard deviation for the experiment was combined with one third of the systematic error to provide the uncertainty quoted in the table: $\sigma_{\text{tot}} = \sigma_{\text{statistical}} + 1/3 \sigma_{\text{systematic}}$ ") seemed more realistic, so the evaluators recommend a half-life of 2.455×10^5 a with a final uncertainty of 0.006×10^5 a. The reduced - χ^2 value is 0.28.

The experimental ^{230}Th half-life values (in years) are given in Table 2:

Table 2: Experimental values of ^{230}Th half-life.

Reference	Value (a)	Uncertainty (a)
M. Curie (1930Cu02)	82300	2469
E.K. Hyde (1949Hy03)	80000	3000
R.W. Attree (1961At01)	75200	1600
J.W. Meadows (1980Me10)	75381	295
Recommend value	75500	500

The recommended value is the weighted average (calculated with LWEIGHT computer program) of $75.5 \cdot 10^3 a$ with an external uncertainty of $0.5 \cdot 10^3 a$. The reduced χ^2 value is 3.3.

The evaluated spontaneous fission partial half-life of ^{234}U is based on the experimental results given in Table 3.

Table 3: Experimental values of ^{234}U spontaneous fission half-life (in 10^{16} years).

Reference	Value	Uncertainty	Comments
A. Ghiorso (1952Gh27)	2	1	Not used.
H.R. von Gunten (1981Vo02)	1.42	0.08	
S. Wang (1987Sh27)	1.90	0.15	
Recommend value	1.5	0.2	reduced $\chi^2 = 5.12$

The evaluators have not use the value given in 1952Gh27, as recommended in 1989Ho24.

Evaluators' recommended value is the weighted average of the two remaining values: $1.5 \cdot 10^{16} a$ with an external uncertainty of $0.2 \cdot 10^{16} a$. The reduced χ^2 value is 5.12.

This value produces a spontaneous fission branching of $1.6 (2) \cdot 10^{-9} \%$.

2.1 a Transitions

The energies of the α -particle transitions given in Section 2.1 have been calculated from the Q_α (2003Au03) and level energies deduced by the evaluators from a least-squares fit to γ -ray energies.

2.2 g Transitions

The transition probabilities have been calculated using the γ -ray emission intensities and the relevant internal conversion coefficients (see **4.2 Gamma Emissions**).

For the 634-keV γ -ray (E0 transition), $P_{(\gamma+ce)} = 1.4 (7) \cdot 10^{-5} \%$ has been deduced from decay scheme balance.

Multipolarities of γ -ray transitions in decay of ^{230}Th are from 1993Ak02:

53-keV γ -ray : E2	581-keV γ -ray: E2
120-keV γ -ray : E2	624-keV γ -ray: E0 + E2 + M1
454-keV γ -ray : E1	634-keV γ -ray: E0
503-keV γ -ray: [E2]	677-keV γ -ray: [E2]
508-keV γ -ray: E1	

The internal conversion coefficients (ICC's) have been calculated using the Icc99v3a computer program (GETICC dialog), which uses interpolated values from new tables of Band et al (2002Ba85). The evaluators have used a fractional uncertainty of 3 % for all conversion coefficients.

3 Atomic Data

Atomic values, ω_K , ω_L and n_{KL} , X-ray and Auger electrons relative probabilities are from Schönfeld and Jaßén (1996Sc06).

4 a Emissions

α -particle energies are from Q_α (2003Au03) and level energies (see section 2.1). For the $\alpha_{0,0}$ and $\alpha_{0,1}$ emissions, the energies are from A. Rytz (1991Ri01).

The measured α -emission intensities are given in Table 4.

Table 4: Measured α -emission intensities, in %.

Energy (keV)	1955Go57	1960Ba44	1961Ko11	1963Bj03	1984Va41	1987Bo25	Recommended Value
4774.6 ($\alpha_{0,0}$)	72	72.5 (30)	73		71.38 (5)	71.37 (2)	71.37 (2)
4722.4 ($\alpha_{0,1}$)		27.15 (15)	27		28.42 (5)	28.42 (2)	28.42 (2)
4603.5 ($\alpha_{0,2}$)		= 0.37 (11)	0.3		0.206 (4)	0.199 (2)	0.210 (2)
4275.2 ($\alpha_{0,3}$)				4 (1) 10^{-5}			4 (1) 10^{-5}
4150.6 ($\alpha_{0,4}$)				1.2 (5) 10^{-5}			2.6 10^{-5}
4108.6 ($\alpha_{0,5}$)				0.3 10^{-5}			7.0 10^{-6}

The U-234 spectrum was recorded by 1984Va41, a second analysis of the same data was done by 1987Bo25, these latest values are the adopted results for the 4774 - and 4722-keV α -emissions intensity. The 4603-keV intensity is deduced from the decay scheme, the tree others being negligible.

The 4275-, 4150-, 4108- keV emission intensities are deduced from 1963Bj03 and decay scheme transition probability balance (§6.2).

6 Photon Emissions

6.1 X-rays

The X-ray and Auger electrons absolute intensities have been calculated from γ -ray data and ICC by using the EMISSION computer program.

In the Table 5 the recommended values of ^{230}Th X-ray emission probabilities are compared with the experimental results. Good agreement was found between the experimental results given by 1977Bemis, 1984Va41 and 1995Jo23 and the recommended values calculated from the decay scheme data set. This agreement confirms the completeness and consistency of the decay scheme.

Table 5: Experimental and recommended (calculated) values of ^{230}Th X-ray emission intensities.

Reference	1977Bemis	1984Va41	1995Jo23	Recommended value
11.118 – 19.504 (L X-ray)	9.81 (13)	10.35 (14)	10.02 (7)	10.2 (4)
L λ - 11.118			0.206 (3)	0.209 (12)
L α - 12.808 – 12.967			3.42 (2)	3.48 (17)
L η - 14.509				0.118 (7)
L β - 14.972 – 16.425			5.17 (4)	5.16 (26)
L γ - 18.363 – 19.504			1.22 (1)	1.21 (6)

Reference	1977Bemis	1984Va41	1995Jo23	Recommended value
89.95 (X K _{α2})		2.53 (7) 10 ⁻³		2.69 (25) 10 ⁻³
93.35 (X K _{α1})		4.15 (10) 10 ⁻³		4.4 (4) 10 ⁻³

6.2 Gamma emissions

The energies of the γ -ray emissions given in Section 6 are from Y.A. Akovali (1993Ak02). The experimental intensity of the 120-keV γ emission given in Table 6 is relative to the 53-keV γ -ray.

Table 6: Experimental relative γ emission intensity (P_{rel}) in %.

γ Energy (keV)	1966Ah02	1974HeYW	1984Va41	Recommended value
53.20	100	100 (5)	100	100.0 (25)
120.90	34 (4)	34.2 (18)	27.5 (5)	30.8 (24)

The recommended values are the weighted averages of the three values given with uncertainties. The normalization factor to convert the relative emission intensities to absolute emission intensities is calculated with the formula:

$$\text{Normalization factor} = \frac{(100\% - 71.371(19)\%)}{\sum [(1 + a_T)P_{rel}]} = 0.001253 (40),$$

where the sum is over all they transitions to the ground state and α_T is the relevant conversion coefficient. In this case, the contribution of 508 - (see next), 634 - and 677 -keV γ transitions are considered negligible. The uncertainty was calculated through the propagation on the formula given above.

For the 454- and 508-keV absolute emission probabilities, the evaluators have following relations:
 $P_\gamma(454) + P_\gamma(508) = 4 (1) 10^{-5}$ (from 1963Bj03) and
 $P_\gamma(508) = 0.60 (4) \times P_\gamma(454)$ (from average value of measured ratios in ²³⁰Pa and ²³⁰Ac decays. See 1993Ak02). Then the evaluator obtains $P_\gamma(454) = 0.000025 (6) \%$ and $P_\gamma(508) = 0.0000150 (39) \%$. For the others γ rays, the evaluators present the experimental absolute emission values given in 1993Ak02. The evaluated relative and absolute γ -rays emission intensities are given in Table 7.

Table 7: Evaluated relative and absolute γ -ray emission intensities.

Energy (keV)	Relative emission intensity (%)	Absolute emission intensity (%)
53.20 (2)	100.0 (25)	0.1253 (40)
120.90 (4)	30.8 (24)	0.0386 (32)
454.96 (5)		0.000025 (6)
503.5 (1)		0.00000095
508.16 (5)		0.0000150 (39)
581.7 (1)		0.000012 (5)
624.4 (1)		0.00000082
677.6 (1)		0.000001

7 References

- 1939Ni03 – O. Nier, Phys. Rev. 55(1939)150 [$T_{1/2}$ (U-234)].
- 1930Cu02 – M. Curie, S. Cotelle, Comp. Rend. Acad. Sci. (Paris) 190(1930)1289 [$T_{1/2}$ (Th-230)].
- 1946Ch02 – O. Chamberlain, D. Willians, P. Yuster, Phys. Rev. 70(1946)580 [$T_{1/2}$ (U-234)].
- 1949Ba41 – E. Baldinger, P. Huber, Helv. Phys. Acta 22(1949)365 [$T_{1/2}$ (U-234)].
- 1949Go18 – A.S. Goldin, G.B. Knight, P.A. Macklin, R.L. Macklin, Phys. Rev. 76(1949)336 [$T_{1/2}$ (U-234)].
- 1949Hy03 – E.K. Hyde, NNS 14B(1949)1435 [$T_{1/2}$ (Th-230)].
- 1949Ki26 – A.C. Kienberger, Phys. Rev. 76(1949)1561 [$T_{1/2}$ (U-234)].
- 1952Fl20 – E.H. Fleming Jr., A. Ghiorso, B.B. Cunningham, Phys. Rev. 88(1952)642 [$T_{1/2}$ (U-234)].
- 1952Ki19 – A.C. Kienberger, Phys. Rev. 87(1952)520 [$T_{1/2}$ (U-234)].
- 1952Gh27 – A. Ghiorso, G.H. Higgins, A.E. Larsh, G.T. Seaborg, S.G. Thompson, Phys. Rev. 87(1952)1963 [S.F. Half-life].
- 1960Ba44 – S.A. Baranov, A.G. Zelenkov, V.M. Kulakov, Bull. Acad. Sci. USSR, Phys. Ser. 24(1960)1045 [α emission].
- 1961At01 – R.W. Attree, M.J. Cabell, R.L. Cushing, J.J. Pieroni, Can. J. Phys. 40(1961)194 [$T_{1/2}$ (Th-230)].
- 1961Ko11 – G.E. Kocharov, G.A. Korolev, Bull. Acad. Sci. USSR, Phys. Ser. 25(1961)227 [α emission].
- 1963Bj03 – S. Bjørnholm, M. Lederer, F. Asaro, I. Perlman, Phys. Rev. 130(1963)2000 [α and γ emission intensities].
- 1965Ne03 – W.R. Neal, H.W. Kraner, Phys. Rev. 137(1965)B1164 [53- and 174-keV half-lives].
- 1965Wh05 – P.H. White, G.J. Wall, F.R. Pontet, J. Nucl. En. A/B 19(1965)33 [$T_{1/2}$ (U-234)].
- 1966Ah02 – I. Ahmad, UCRL – 16888(1966) [γ energy and intensity].
- 1969Hanna – G.C. Hanna, C.H. Westcott, H.D. Lemmel, B.R. Leonard Jr., J.S. Story, P.M. Attree, At. Energy Rev. 7, vol. 4(1969)3 [$T_{1/2}$ (U-234)].
- 1970DeYN – P. de Bievre, K.F. Lauer, Y. de Duigou, H. Moret, G. Muschenborn, J. Spaepen, A. Spornol, R. Vaninbrouckx, V. Verdingh, Chem. Nucl. Data (Canterbury) (1971)221 [$T_{1/2}$ (U-234)].
- 1970LoZL – M. Lounsbury, R.W. Durham, Chem. Nucl. Data (Canterbury) (1971)215 [$T_{1/2}$ (U-234)].
- 1970MeZN – J.W. Meadows, ANL – 7610(1970)44 [$T_{1/2}$ (U-234)].
- 1972Sc01 – M. Schmorak, C.E. Bemis Jr., M.J. Zender, N.B. Grove, P.F. Dittner, Nucl. Phys. A178(1970)410 [γ energy].
- 1973Ta25 – H.W. Taylor, Int. J. Appl. Radiat. Isotop. 24(1973)593 [γ energy].
- 1974HeYW – R.L. Heath, ANCR – 1000-2(1974)14 [γ energy and intensity].
- 1977Bemis – C.E. Bemis, Jr., L. Tubbs, ORNL – 5297(1977)93 [X-ray intensity].
- 1980Me10 – J.W. Meadows, R.J. Armani, E.L. Calis, A.M. Essling, Phys. Rev. C22(1980)750 [$T_{1/2}$ (Th-230)].
- 1980Ge13 – A.M. Geidel'man, Yu. S. Egorov, A.V. Lovtysyus, V.I. Orlov, L.D. Preobrazhenskaya, M.V. Ryzhinskii, A.V. Stepanov, A.A. Lipovskii, Yu.V. Khol'nov, B.N. Belyaev, M.K. Adbullakhatov, G.A. Akopov, V.S. Belykh, E.A. Gromova, V.Ya. Mishin, L.F. Solntseva, Bull. Acad. Sci. USSR, Phys. Serv. 44,5(1980)23 [$T_{1/2}$ (U-234)].
- 1981HoZI – N.E. Holden, BNL – NCS 51320(1981)111 [$T_{1/2}$ (U-234)].
- 1981Vo02 – H.R. von Gunten, A. Grütter, H.W. Reist, M. Baggenstos, Phys. Rev. C23(1981)1110 [S.F. Half-life].
- 1983Ak12 – Y.A. Akovali, Nucl. Data Sheets 40(1983)523 [Spin, parity, energy level, multipolarity].
- 1983Poenitz – W.P. Poenitz, J.W. Meadows, ANL – NDM 84(1983)33 [$T_{1/2}$ (U-234)].
- 1984Di08 – M. Divadeenam, J.R. Stehn, Ann. Nucl. Energy 11(1984)375 [$T_{1/2}$ (U-234)].
- 1984Va41 – R. Vaninbrouckx, G. Bortels, B. Denecke, Int. J. Appl. Radiat. Isotop. 35(1984)1081 [X-ray, α and γ emission intensities].
- 1985Poenitz – W.P. Poenitz, J.W. Meadows, IAEA – TECDOC 335(1985)485 [$T_{1/2}$ (U-234)].
- 1985Axton – E.J. Axton, IAEA – TECDOC 335(1985)214 [$T_{1/2}$ (U-234)].
- 1986LoZT – A. Lorentz, A.L. Nichols, IAEA – Tech. Rep. 261(1986)63 [$T_{1/2}$ (U-234), [α and γ emission intensities].
- 1987Bo25 – G. Bortels, P. Collaers, Appl. Rad. Isotopes 38(1987)831 [α emission].
- 1987Sh27 – S. Wang, P.B. Brice, S.W. Barmick, K.J. Moody, E.K. Hulet, Phys. Rev. C36(1987)2717 [S.F. Half-life].

- 1989Ho24 – N.E. Holden, Pure and Appl. Chem. 61(1989)1483 [$T_{1/2}$ (U-234)].
1991Ry01 – A. Rytz, Atomic Data and Nuclear Data Tables 47(1991)205 [E_{α}].
1993Ak02 – Y.A. Akovali, Nucl. Data Sheets 69(1993)155 [Spin, parity, energy level, multipolarity].
1994Ak05 – Y.A. Akovali, Nucl. Data Sheets 71(1993)181 [Spin, parity, energy level, multipolarity].
1995Jo23 – P.N. Johnston, P.A. Burns, Nucl. Instrum. Meth. Phys. Res A361(1995)229 [X-ray intensity] .
1996Sc06 – E. Schönfeld, H. Janssen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
2002Ba85 – I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, Atomic Data and Nuclear Data Tables 81(2002)1 [α].
2003Au03 – G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].

²³⁵U - Comments on evaluation of the decay data

Huang Xiaolong, Wang Baosong

This evaluation was completed in 2008, and data available in the literature by June 2008 was included.

1 Decay Scheme

²³⁵U disintegrates 100 % by α emission to levels in ²³¹Th. ²³⁵U ground state has $J^\pi = 7/2^-$ (2003Br12). The spontaneous fission branching ratio is $7.0 (20) \times 10^{-9}$ % (from $T_{1/2}(\text{SF}) = 1.0 (3) \times 10^{19}$ a (2000Ho27) and $T_{1/2} = 7.04 (1) \times 10^8$ a.)

The α decay scheme of ²³⁵U was built based on the measurements described in 1974Te03, 1975Va11 and 1977Ba72. A study of 2004Da24 showed the existence of weak α decay branches to some levels in ²³¹Th.

2 Nuclear Data

A Q value of 4678.3 (7) keV is given in 2003Au03 atomic mass adjustment.

Level energies have been obtained from a least-squares fit to γ -ray energies (GTOL computer code). Spin and parities are from 2003Br12.

The measured and evaluated ²³⁵U half-life values are listed in Table 1. Notice that the uncertainties in all the tables are in the two least significant digits.

Table 1: Measured half-life values of ²³⁵U and recommended value, in 10^8 a.

References	Original value (10 ⁸ a)	Materials	Revised value by 2004Sc03	Comments
1939Ni03	7.13 (16)	Natural U	6.97 (24)	Pb/U activity ratio, Mass spectrometry
1950Kn17	7.53 (22)	Enriched U	7.11 (14)	Specific activity, Ionisation chamber
1951Sa30	7.07	Natural U	6.77 (21)	²³⁵ U/ ²³⁸ U activity ratio, Ionisation chamber
1952Fl20	7.13 (16)	Enriched U	7.12 (16)	Specific activity, proportional counter
1957Cl16	7.67	Natural U	7.64 (43)	activity ratio, Ionisation chamber
1957Wu39	6.84 (15)	Natural U	6.95 (16)	²³⁵ U/ ²³⁴ U activity ratios, Ionization chamber
1965De06	6.92 (9)	Natural U		²³⁵ U/ ²³⁸ U activity ratio, Solid-state detector, Updated by 1974De19
1965Wh05	7.13 (9)	Enriched U	7.12 (9)	Specific activity, Solid-state detector
1971Ja07	7.0381 (48)	Highly enriched U	7.04 (1)	Specific activity, proportional counter
1974De19	6.85 (9)	Highly enriched U	6.79 (13)	²³⁵ U central peak branching ratio, Solid-state detector
1993Bu10	7.04 (1)	Enriched U		Specific activity, (gas + NaI scintillator) Systematic error excluded
2003Br12	7.04 (1)			NDS weighted average with 1993Bu10, 1974De19, 1971Ja07, 1965Wh05, 1965De06 and 1957Wu39
			7.06 (9)	Unweighted mean
			7.04 (1)	Weighted mean, $\chi^2=1.12$. Recommended value

The evaluators have chosen to follow the recommendations given by R. Schön (2004Sc03), who studied in detail various problems with the measurements of the half-life of ^{235}U and decided to recommend the half-life given by 1971Ja07, but multiplied by 2 its original uncertainty in order to include the systematic uncertainties that had not been considered in 1971Ja07. The weighted mean is the same as this precise measurement given in 1971Ja07.

The measured and evaluated ^{235}U spontaneous fission half-life values are listed in Table 2. The value in 1981Vo02 is recommended here.

Table 2: Measured spontaneous fission half-life values of ^{235}U and recommended value, in 10^{19}a .

$T_{1/2}$ (10^{19}a)	References	measurement method
0.018	1952Se67	Ionisation chamber; not used
0.035 (9)	1966Al23	Fission track detectors; not used
> 0.18	1974GrZA	Rotating bubble chamber; no corrections; not used
0.98 (28)	1981Vo02	99.76 % enriched; rotating bubble chamber; corrected for the (α , n, f) reaction
1.0 (3)	2003Br12	NDS, from evaluation of 2000Ho27
0.98 (28)	Recommended value	From 1981Vo02

2.1 γ -Ray Transitions

The γ -ray transition probabilities were deduced from the γ -ray emission probabilities and the relevant internal conversion coefficients.

Multipolarities and mixing ratios of γ -ray transitions are from 2003Br12.

Theoretical internal conversion coefficients (ICC) and their associated uncertainties for γ -ray transitions have been obtained using the BRICC computer program (2008Ki07), which uses the ‘‘Frozen Orbital’’ approximation (2002Ba85).

2.2 α -Particle Transitions

Measured energies of alpha particles are listed in Table 3. Our recommended values are from 1975Va11, 1991Ry01, 2004Da24, Q_α (2003Au03) and level energies.

Table 3: Measured and recommended values of α -particle energies (in keV) from ^{235}U α decay.

1960Ba44	1962Pi06	1966Ga03	1975Va11	1991Ry01	2004Da24	Calc. from level energy and $Q(\alpha)$	Recommended
		3977 (10)			3976 (5)	3897.2 (7)	3897.2 (7)
						3975.3 (7)	3976 (5)
						3990.5 (9)	3990.5 (9)
						4013.2 (8)	4013.2 (8)
						4053.9 (7)	4053.9 (7)
		4069 (10)			4077	4077.5 (7)	4077.5 (7)
	4153	4140 (3)	4145 (6)		4152 (5)	4154.2 (7)	4152 (5)
4214	4210	4210 (3)	4209 (4)	4214.7 (19)	4215.8 (5)	4217.4 (7)	4214.7 (19) ^b

1960Ba44	1962Pi06	1966Ga03	1975Va11	1991Ry01	2004Da24	Calc. from level energy and Q(α)	Recommended
			4219 (6)			4219.6 (7)	4219.6 (7)
						4227.6 (7)	4227.6 (7)
		4240 (10)			4248 (5)	4252.6 (7)	4248 (5)
	4261				4266 (5)	4270 (4)	4266 (5)
		4267 (10)				4279.3 (7)	4279.3 (7)
			4280		4282 (5) ^a	4286.9 (7)	4286.9 (7)
		4289 (10)	4295			4302.1 (7)	4302.1 (7)
4320	4318	4319 (3)	4322 (4)		4322.9 (6) ^a	4325.4 (7)	4322 (4)
4326						4327.9 (7)	4327.9 (7)
					4364.3 (4) ^a	4361.9 (7)	4361.9 (7)
4368	4361	4362 (3)	4358 (4)	4366.1 (20)		4365.8 (7)	4366.1 (20) ^b
		4368 (5)				4381.1 (7)	4381.1 (7)
4394	4391	4394 (3)	4392 (3)	4397.8 (13)	4395.3 (4)	4396.8 (7)	4397.8 (13) ^b
4412	4414	4411 (5)	4411 (5)		4414.9 (5)	4416.1 (7)	4414.9 (5)
4438	4440	4424 (5)	4435 (5)		4437.9 (40)	4439.3 (7)	4437.9 (40)
4496	4497	4496 (3)	4501 (4)		4502.4 (7)	4504.2 (7)	4502.4 (7)
4550	4551	4550 (3)	4555 (3)		4556.0 (4)	4557.4 (7)	4556.0 (4)
4592	4592	4592 (3)	4597 (3)	4596.4 (13)	4597.3 (4)	4598.7 (7)	4596.4 (13) ^b

^a: May be a multiplet.

^b: From 1991Ry01.

Experimental and recommended α -particle emission probabilities are listed in Table 4. Our recommended alpha particle emission probabilities are LWM average values of measured α -particle intensities given in 1975Va11, 2004Da24 and 2005Ga36. Other recommended values are from results deduced from γ -ray transition intensity balance at each nuclear level.

Table 4: Measured and recommended values of α -particle emission probabilities from ²³⁵U decay.

E_α (keV)	P_α (%)							Deduced from I_γ	LWM	Recommended ^f
	1960Ba44	1962Pi06	1966Ga03	1975Va11	2004Da24	2005Ga36				
3976					~0.007			≈0.0011		≈0.0011
4013.2								0.040 (1)		0.0396 (10)
4077.5						0.016 (12)		0.0177 (3)		0.016 (12)
4152		~ 0.3	1.0	0.9 (2) ^a	0.31 (2)	0.286 (18)		0.506 (14)	0.297 (13)	0.294 (13)
4214.7	5.5	5.5	6.2	5.7 (6)	6.28 (11)	5.91 (7)		6.0 (4)	6.01 (12)	5.95 (12)
4219.6								0.0175 (2)		0.01732 (12)
4227.6				~ 0.9				0.123 (6)		0.122 (6)
4248?			< 0.5		0.07 (1)			0.07 (1)		0.069 (10)?
4266					0.26 (2)	0.200 (16)		0.22 (8)	0.22 (3)	0.22 (3)
4279.3			< 0.3					0.0332 (4)		0.0329 (5)

E_{α} (keV)	P_{α} (%)							Deduced from I_{γ}	LWM	Recommended [†]
	1960Ba44	1962Pi06	1966Ga03	1975Va11	2004Da24	2005Ga36				
4286.9		0.6			0.14 (1) ^a	0.066 (13)	0.096 (12)		0.065 (13)	
4302.1			< 0.5				0.00969 (12)		0.00959 (13)	
4322	3	2.9	3.5	4.7 (5) ^a	3.78 (8) ^a	3.37 (6)	3.3 (7)		3.33 (6)	
4327.9	11						0.409 (13)		0.405 (13)	
4361.9							0.208 (21)		0.206 (21)	
4366.1	6	19	12.3	17 (2) ^a	18.8 (2) ^a	19.00 (13)	19 (5)		18.80 (13)	
4381.1			6.1				0.107 (16)		0.106 (16)	
4397.8	62	58	53.0 (13)	54 (3)	57.11 (41)	57.98 (22)	58 (5)	57.8 (3)	57.19 (20)	
4414.9	2	~ 4	2.3	2.1 (2)	3.07 (7)	3.11 (6)	3.5 (22)	3.04 (16)	3.01 (16)	
4437.9	3	~ 0.6	1.8	~ 0.7	0.27 (2)	0.219 (16)	0.206 (16)	0.239 (25)	0.236 (25)	
4502.4	1	1.2	1.4	1.7 (2)	1.32 (5)	1.25 (4)	1.23 (24)	1.29 (5)	1.28 (5)	
4556.0	3	3.7	1.7	4.5 (5)	3.74 (8)	3.87 (6)	3 (3)	3.83 (6)	3.79 (6)	
4596.4	< 1	4.7	1.2	5.4 (5)	4.84 (9)	4.74 (7)	4 (4)	4.79 (6)	4.74 (6)	

[†]: Normalized to a total of 100 %.

^a: May be a multiplet.

3. Atomic data

Atomic fluorescence yields ($\omega_K, \omega_L, \omega_M, \eta_{KL}$ and η_{LM}) are from Schönfeld (1996Sc06).

The X-ray and Auger electron emission probabilities have been deduced from γ -ray and conversion electron data by using the computer code RADLST. The deduced K X-ray emission probabilities $P_{K\alpha 1} = 5.75$ (14) agree with the measured value of 5.55 (14) in 1996Ru11, thus confirming the completeness of the decay scheme.

4. Electron Emissions.

The conversion electron emission probabilities have been deduced from γ -ray transition data using theoretical internal conversion coefficients.

5. Photon Emissions

5.1 γ -ray energy values

The experimental and our recommended γ -ray energies from ^{235}U α decay are listed in table 6. Our recommended values are mainly from the LWM averages based on measurements of 1971Cl03, 1974Te03, 1975Va11, 1977Ba72 and 1984He12 unless otherwise specified. Values in 1986LoZT are from the CRP evaluations done in 1986.

5.2 Absolute γ -ray emission probabilities

Measured relative and absolute γ -ray intensities from ^{235}U are listed together with evaluated values in Table 7. Among these measurements, 1966Ga03, 1971Cl03, 1971KrZH, 1974Te03, 1975Va11, 1977Ba72 and 1996Ru11 are measured relative γ -ray intensities. Other values reported in 1982Va04, 1983BaZZ, 1983OI01, 1984He12, 1992Li05 and 2006Al28 are measured absolute γ -ray intensities. Thus we evaluated and recommended the γ -ray emission probability of the 185.7-keV reference line firstly.

There are 7 independent measurements of the absolute γ -ray emission probability of the 185.7-keV reference line. Among these absolute measurements, 1982Va04, 1983BaZZ and 1984He12 belong to CRP measurements. The measurement reported in 2006Al28 has not been recommended because of interference with gamma rays from a ^{226}Ra impurity.

The CRP evaluations done in 1986 are reported in 1986LoZT where a recommended $P_{\gamma}(185.7) = 57.2$ (2) is given. We re-calculated $P_{\gamma}(185.7)$ and found that the LWM average value based on CRP measurements reported in 1982Va04, 1983BaZZ and 1984He12 is 57.3 (4), and LWM of 1982Va04, 1983BaZZ, 1983OI01, 1984He12 is 57.1 (3). Our recommended value is taken from the LWM average of values given (Table 5) in 1982Va04, 1983BaZZ, 1983OI01, 1984He12, 1992Li05 and 1999Ch12, that is, $P_{\gamma}(185.7) = (57.1 \pm 0.3) \%$.

Table 5: Experimental 185.7-keV absolute gamma-ray emission probabilities.

References	Experimental values (%)	Comments
R. Vaninbroukx (1982Va04)	57.5 (9)	
C. Baktash (1983BaZZ)	57.3 (6)	
D. G. Olson (1983OI01)	56.1 (8)	
R. G. Helmer (1984He12)	57.2 (5)	
W. -J. Lin (1992Li05)	56.8 (13)	
H. Chatani (1999Ch12)	58 (2)	
Recommended value	57.1 (3)	$\chi^2 = 0.43$

Results for most γ rays given in 1966Ga03 and 1977Ba72 were not used because they did not have uncertainties, unless these were the only measurements for such γ -rays. Relative γ -ray intensities reported in 1971Cl03, 1971KrZH, 1974Te03, 1975Va11 and 1996Ru11 have been normalized using the present recommended $P_{\gamma} = (57.1 \pm 0.3)$ for the 185.7 keV reference line.

Our “best” recommended absolute γ -ray emission probabilities are mainly from LWM averages of measurements reported in 1971Cl03, 1971KrZH, 1974Te03, 1975Va11, 1982Va04, 1983BaZZ, 1983OI01, 1984He12, 1992Li05 and 1996Ru11 unless otherwise specified.

Table 6: Measured and recommended values of γ -ray energies for ^{235}U α decay.

1966Ga03	1971Cl03	1974Te03	1975Va11	1977Ba72	1984He12	1986LoZT	LWM	Recommended
			19.59	19.55 (5)				19.55 (5)
		31.50 (20)	31.59 (14)	31.60 (5)			31.60 (5)	31.60 (5)
				34.7 (1)				34.7 (1) ^x
		41.70 (15)	41.1			41.4 (3)		41.4 (3) ^a
		41.96 (15)	42.1 (1)	41.95 (10)		41.96 (15)	42.01 (6)	42.01 (6)
		51.20 (10)	51.7 (4)	51.20 (5)			51.21 (4)	51.21 (4)
				54.1 (1)				54.1 (1)
			54.1	54.25 (5)				54.25 (5)
				64.45 (5)				64.45 (5)
			72.7 (2)			72.7 (2)		72.7 (2)
				73.72 (5)				73.72 (5)
	74.923 (23)	74.76 (20)	75.02 (5)			75.02 (5)	74.94 (3)	74.94 (3)
			95.7					95.7
		96.09 (2)	96.1	96.2				96.09 (2)
97 (4)								97 (4)
109 (4)	109.120 (8)	109.145 (10)	109.25 (5)	109.25 (5)		109.16 (2)	109.19 (7)	109.19 (7)
115 (4)	115.2 (3)		115.5 (2)	115.45 (5)			115.45 (5)	115.45 (5)
			120.0	120.35 (5)				120.35 (5)
			136.6	136.55 (5)				136.55 (5)
	140.75 (10)	140.758 (20)	140.80 (8)	140.75 (5)		140.76 (4)	140.76 (2)	140.76 (2)
				142.40 (5)				142.40 (5)
144 (2)	143.776 (10)	143.753 (8)	143.77 (2)	143.75 (5)	143.768 (3)	143.76 (2)	143.767 (3)	143.767 (3)
			147.0					147
151 (4)	150.960 (33)	150.939 (20)	150.94 (3)	150.85 (5)		150.93 (2)	150.936 (15)	150.936 (15)
	163.363 (10)	163.349 (9)	163.36 (2)	163.25 (5)	163.357 (3)	163.33 (2)	163.356 (3)	163.356 (3)
			173.0 (10)					173 (1)

1966Ga03	1971Cl03	1974Te03	1975Va11	1977Ba72	1984He12	1986LoZT	LWM	Recommended
			182.1					182.1
	182.72 (20)	182.65 (15)	182.7 (2)	182.60 (5)		182.61 (5)	182.62 (5)	182.62 (5)
184 (2)	185.718 (11)	185.712 (10)	185.72 (2)	185.65 (5)	185.722 (4)	185.715 (5)	185.720 (4)	185.720 (4)
196 (4)	194.941 (9)	194.938 (10)	194.94 (2)	194.95 (5)		194.94 (1)	194.940 (6)	194.940 (6)
	198.91 (15)	198.898 (15)	198.88 (6)	198.75 (10)		198.90 (2)	198.894 (14)	198.894 (14)
				199.6 (1)				199.6 (1) ^x
	202.133 (14)	202.105 (12)	202.12 (2)	202.05 (5)		202.11 (2)	202.12 (1)	202.12 (1)
	205.311 (12)	205.312 (10)	205.31 (2)	205.25 (5)	205.318 (4)	205.311 (10)	205.316 (4)	205.316 (4)
		215.26 (20)	215.28 (5)	215.3 (1)			215.28 (4)	215.28 (4)
	221.375 (40)	221.397 (25)	221.38 (2)	221.40 (5)		221.38 (2)	221.386 (14)	221.386 (14)
			228.78 (5)	228.7 (1)			228.76 (5)	228.76 (5)
	233.53 (4)	233.49 (3)	233.50 (3)	233.55 (10)		233.50 (3)	233.50 (2)	233.50 (2)
	240.93 (4)	240.95 (4)	240.87 (3)	240.75 (5)		240.87 (3)	240.88 (4)	240.88 (4)
	246.83 (4)	246.59 (10)	246.84 (2)	246.85 (5)		246.84 (4)	246.83 (2)	246.83 (2)
				251.5 (1)				251.5 (1) ^x
	266.44 (8)	266.40 (10)	266.50 (5)				266.47 (4)	266.47 (4)
		275.35 (15)						275.35 (15)
			275.24 (20)	275.50 (5)			275.49 (6)	275.49 (6)
				279.50 (5)				279.50 (5) ^x
			281.42 (5)					281.42 (5)
285 (5)			282.92 (5)	283.0 (1)			282.94 (5)	282.94 (5)
			289.56 (4)					289.56 (4)
			291.2					291.2
		291.58 (15)	291.65 (3)	291.65 (5)			291.65 (3)	291.65 (3)
				294.3 (1)				294.3 (1) ^x
			301.7 (1)					301.7 (1)
			310.69 (6)					310.69 (6)
			317.10 (8)					317.10 (8)

1966Ga03	1971Cl03	1974Te03	1975Va11	1977Ba72	1984He12	1986LoZT	LWM	Recommended
				325.8 (1)				325.8 (1)
			343.5 (2)					343.5 (2)
				345.4 (1)				345.4 (1) ^x
		345.84 (15)	345.93 (3)	345.90 (5)			345.92 (3)	345.92 (3)
350 (5)								350 (5)
			356.03 (5)					356.03 (5)
				368.5 (1)?				368.5 (1)?
				371.8 (1)				371.8 (1) ^x
		387.79 (15)	387.84 (3)	387.85 (10)			387.84 (3)	387.84 (3)
		390.27 (20)						390.27 (20)
			410.29 (4)					410.29 (4)
430 (5)				~433.0 (5)				433.0 (5)
			448.40 (6)					448.40 (6)
			455.1 (1)					455.1 (1) ^x
			517.9 (2)					517.9 (2) ^x
			742.5 (2)					742.5 (2) ^x
			794.7 (1)					794.7 (1) ^x

^x: Not placed in level scheme.

^a: From 1986LoZT.

Table 7: Measured and recommended absolute γ -ray emission probabilities for ²³⁵U.

E_γ (keV)	1966Ga03 ^a	1971Cl03 ^a	1971KrZH ^a	1974Te03 ^a	1975Va11 ^a	1977Ba72 ^a	1982Va04	1983BaZZ	1983OI01	1984He12	1986LoZT	1992Li05	1996Ru11 ^a	LWM	Adopted [*]
19.55															60 (1) [#]
31.60				0.017 (6)		0.046									0.017 (6)
34.7 [*]						0.037									0.037
41.4				0.029 (11)											0.029 (11)
42.01			0.053	0.04 (2)	0.0169	0.063		0.06 (1)			0.06 (1)			0.056 (9)	0.056 (9)
51.21				0.004 (2) ^b	0.034 (7)	0.017									0.034 (7)
54.1						0.03?									\approx 0.00115 [#]
54.25						0.03?									\approx 0.0285 [#]
64.45						0.018									0.018
72.7					0.116										0.116
73.72						0.01									0.01
74.94		0.0012 (1) ^b	0.137	0.051 (6)	0.074			0.51 (5) ^b			0.06 (1)				0.051 (6)
95.7															
96.09				0.091 (11)											0.091 (11)
97	<1														0.016 (4) [#]
109.19	5.1	1.60 (12)	1.59 (21)	1.77 (17)	1.48 (21)	1.03		1.53 (5)			1.54 (5)	2.17 (17)	1.80 (6)	1.66 (13)	1.66 (13)
115.45	<1	0.14 (1) ^b	0.12 (3) ^b		0.033 (12)	0.017									0.03 (1)
120.35						0.026									0.026
136.55						0.012									0.012
140.76		0.183 (13)	0.18 (2)	0.26 (3)	0.22 (3)	0.171		0.214 (15)			0.22 (2)			0.200 (12)	0.20 (1)
142.40						0.0051									0.0051
143.767	11.7	10.3 (8)	10.3 (6)	11.2 (11)	11.1 (12)	9.92	10.9 (2)	10.7 (2)	10.93 (15)	11.01 (8)	10.96 (8)	10.99 (61)	10.9 (2)	10.94 (6)	10.94 (6)
147															
150.936	<1	0.114 (9)	0.116 (32)	0.080 (11)	0.080 (11)	0.074		0.066 (10)			0.08 (1)			0.088 (26)	0.09 (3)
163.356		4.9 (4)	4.9 (3)	4.99 (51)	5.1 (5)	4.16	5.0 (1)	4.97 (10)	5.07 (8)	5.12 (4)	5.08 (4)	4.98 (12)	5.08 (5)	5.076 (26)	5.08 (3)
173			0.016		0.006 (5)										0.006 (5)

E_γ (keV)	1966Ga03 ^a	1971Cl03 ^a	1971KrZH ^a	1974Te03 ^a	1975Va11 ^a	1977Ba72 ^a	1982Va04	1983BaZZ	1983Ol01	1984He12	1986LoZT	1992Li05	1996Ru11 ^a	LWM	Adopted*
182.1															
182.62		0.43 (3)	0.42 (4)	0.42 (14)	0.44 (10)	0.312		0.339 (17)			0.34 (2)	0.803 (103)	0.43 (5)	0.39 (5)	0.39 (5)
185.720							57.5 (9)	57.3 (6)	56.1 (8)	57.2 (5)	57.2 (2)	56.8 (13)		57.1 (3)	57.1 (3)
194.940	4.7	0.69 (5)	0.69 (6)	0.61 (9)	0.62 (6)	0.67		0.626 (13)			0.63 (1)	0.618 (48)	0.61 (2)	0.626 (10)	0.63 (1)
198.894		0.032 (3)	0.032	0.046 (6)	0.033 (5)	0.097?		0.047 (6)			0.42 (6)			0.036 (2)	0.036 (2)
199.6 ^x						0.097?									~0.06 ^{&}
202.12		1.06 (8)	1.1 (5)	1.07 (11)	1.07 (11)	1.25		1.08 (2)			1.08 (2)	1.16 (7)	1.06 (4)	1.080 (17)	1.08 (2)
205.316		5.3 (4)	5.18 (32)	4.9 (4)	5.0 (5)	5.51	5.0 (2)	5.05 (5)	5.03 (9)	4.96 (5)	5.01 (5)	4.98 (14)	5.03 (5)	5.015 (26)	5.02 (3)
215.28			0.42	0.029 (6)	0.029 (3)	0.025								0.029 (3)	0.029 (3)
221.386		0.126 (9)	0.08	0.12 (3)	0.116 (11)	0.125		0.114 (6)			0.12 (1)			0.118 (5)	0.118 (5)
228.76			0.0085		0.0074	0.0011									0.0074
233.50		0.042 (3)	0.021	0.034 (11)	0.032			0.029 (5)			0.029 (5)			0.038 (4)	0.038 (4)
240.88		0.074 (6)	0.0032	0.063 (17)	0.085	0.089		0.076 (6)			0.075 (6)			0.074 (4)	0.074 (4)
246.83		0.063 (5)	0.021	0.046 (17)	0.085	0.067?		0.053 (3)			0.053 (3)			0.055 (3)	0.055 (3)
251.5 ^x						0.067?									~0.012 [^]
266.47		0.0080 (6)	0.0053	0.0063 (17)	0.0095									0.0078 (6)	0.0078 (6)
275.35				0.051 (6)											0.051 (6)
275.49			0.042		0.032	0.114									0.032
279.5 ^x						0.264									0.264
281.42					0.0063										0.0063
282.94	0.001		0.0032		0.0063	0.004									0.0063
289.56					0.0074										0.0074
291.2															
291.65			0.021	0.040 (6)	0.032	0.095									0.040 (6)
294.3 ^x						0.033									0.033
301.7					0.0053										0.0053
310.69			0.0017		0.0053										0.0053
317.10					0.0011										0.0011

E_{γ} (keV)	1966Ga03 ^a	1971Cl03 ^a	1971KrZH ^a	1974Te03 ^a	1975Va11 ^a	1977Ba72 ^a	1982Va04	1983BaZZ	1983O101	1984He12	1986LoZT	1992Li05	1996Ru11 ^a	LWM	Adopted*
325.8						0.004									0.004
343.5					0.0032										0.0032
345.4 ^x						0.072?									~0.03 ⁺
345.92			0.0017	0.040 (6)	0.074	0.072?									0.040 (6)
350	0.006														0.006
356.03					0.0053										0.0053
371.8 ^x						0.069?									
387.84				0.040 (6)	0.0085	0.159									0.040 (6)
390.27				0.040 (1)											0.040 (1)
410.29					0.0032										0.0032
433.0	0.001					0.004									0.004
448.40					0.0011										0.0011
455.1 ^x					0.0085										0.0085
517.9 ^x					0.00042										0.00042
742.5 ^x					0.00042										0.00042
794.7 ^x					0.00063										0.00063

x: Not placed in level scheme.

#: From intensity balance.

&: From $P_{\gamma}(198.9 + 199.6) = 0.097\%$.

^: From $P_{\gamma}(246.8 + 251.5) = 0.067\%$.

+: From $P_{\gamma}(345.4 + 345.9) = 0.072\%$.

*: Deduced using the LWM statistical method, unless otherwise specified.

a: The P_{γ} values have been deduced from the measured relative intensities and normalized to $P_{\gamma} = (57.1 \pm 0.3)\%$ for the 185.7 keV reference line.

b: This value, which deviates by a factor of about 10 from the results of the other measurements, was not used in the calculation of the recommended value.

6. References

- 1939Ni03 A.O. Nier, Phys. Rev. 55(1939)150 [Half-life]
- 1950Kn17 G.B. Knight, Report ORNL K-663 (1950) [Half-life]
- 1951Sa30 G.J. Sayag, Comp. Rend. Acad. Sci. (Paris) 232 (1951) 2091 [Half-life]
- 1952Fl20 E.H. Fleming, Jr., A. Ghiorso, B.B. Cunningham, Phys. Rev. 88(1952)642 [Half-life]
- 1952Se67 E. Segre, Phys. Rev. 86(1952)21 [Half-life]
- 1957C116 F.L. Clark, H.J. Spencer-Palmer, R.N. Woodward, J.S. African Chem. Inst. 10(1957)62 [Half-life]
- 1957Wu39 E. Wurger, K.P. Meyer, P. Huber, Helv. Phys. Acta 30(1957)157 [Half-life]
- 1960Ba44 S.A. Baranov, A.G. Zelenkov, V.M. Kulakov, Izvest. Akad. Nauk SSSR, Ser. Fiz. 24(1960)1035 [Alpha energies and intensities]
- 1962Pi06 R.C. Pilger, F.S. Stephens, F. Asaro, I. Perlman, Priv. Comm., quoted by 1964Hy02 unpublished (1962) [Alpha energies and intensities]
- 1965De06 A.J. Deruytter, I.G. Schroder, J.A. Moore, Nucl. Sci. Eng. 21(1965)325 [Half-life]
- 1965Wh05 P. H. White, G.J. Wall, F.R. Pontet, J. Nucl. Energy A/B19(1965)33 [Half-life]
- 1966Al23 B.M. Aleksandrov, A.S. Krivokhatskii, L.Z. Malkin, K.A. Petrzhak, At. Energ. 20(1966)315 [Half-life]
- 1966Ga03 R. Gaeta, M.A. Vigon, Nucl. Phys. 76(1966)353 [Alpha energies and intensities, gamma-ray energies and intensities]
- 1971Cl03 J.E. Cline, IN-1448 Rev. (1971) [Gamma-ray energies and intensities]
- 1971Ja07 A.H. Jaffey, K.F. Flynn, L.E. Glendenin, W.C. Bentley, A.M. Essling, Phys. Rev. C4(1971)1889 [Half-life]
- 1971KrZH L.A. Kroger, C.W. Reich, J.E. Cline, ANCR-1016(1971)75 [Gamma-ray energies and intensities]
- 1974De19 A.J. Deruytter, G. Wegener-Penning, Phys. Rev. C10, 383 (1974)383 [Half-life]
- 1974GrZA A. Grutter, H.R. von Gunten, V. Herrnberger, B. Hahn, U. Moser, H.W. Reist, G. Sletten, I.A.E.A. Vienna, Vol.1, p.305 (1974) [Half-life]
- 1974Te03 W. Teoh, R.D. Connor, R.H. Betts, Nucl. Phys. A228(1974)432 [Gamma-ray energies and intensities]
- 1975Va11 E. Vano, R. Gaeta, L. Gonzalez, C.F. Liang, Nucl. Phys. A251(1975)225 [Gamma-ray energies and intensities]
- 1977Ba72 S.A. Baranov, V.M. Shatinskii, A.G. Zelenkov, V.A. Pchelina, Sov. J. Nucl. Phys. 26(1977)486 [Gamma-ray energies and intensities]
- 1981Vo02 H.R. von Gunten, A. Grutter, H.W. Reist, M. Baggenstos, Phys. Rev. C23(1981)1110 [Half-life]
- 1982Va04 R. Vaninbroux, B. Denecke, Nucl. Instrum. Methods 193(1982)191 [Gamma-ray energies and emission probabilities]
- 1983BaZZ C. Baktash, E. der Mateosian, O.C. Kistner, A.W. Sunyar, D. Horn, C.J. Lister, Bull. Am. Phys. Soc. 28, No.1, HE7 (1983)41 [Gamma-ray emission probabilities]
- 1983Ol01 D.G. Olson, Nucl. Instrum. Methods 206(1983)313 [Gamma-ray emission probabilities]
- 1984He12 R.G. Helmer, C.W. Reich, Int. J. Appl. Radiat. Isotop. 35(1984)783 [Gamma-ray energies and emission probabilities]
- 1986LoZT A. Lorenz, IAEA Tech.Rept.Ser., No.261 (1986) [evaluated gamma-ray energies and emission probabilities]
- 1991Ry01 A. Rytz, At. Data Nucl. Data Tables 47(1991)205 [Evaluated alpha intensities]

- 1992Li05 W.-J. Lin, G. Harbottle, J. Radioanal. Nucl. Chem. 157(1992)367 [Gamma-ray emission probabilities]
- 1993Bu10 C.C. Bueno, M.D.S. Santos, Appl. Radiat. Isotop. 44(1993)567 [Half-life]
- 1996Ru11 H. Ruellan, M.C. Lépy, M. Etcheverry, J. Plagnard, J. Morel, Nucl. Instrum. Meth. Phys. Res. A369(1996)651 [Gamma-ray and X-ray intensities]
- 1996Sc06 E. Schönfeld, H.Janssen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data]
- 1999Ch12 H. Chatani, Nucl. Instrum. Meth. Phys. Res. A425(1999)277 [Gamma-ray emission probabilities]
- 2000Ho27 N.E. Holden, D.C. Hoffman, Pure Appl. Chem. 72(2000)1525 [Evaluated half-life]
- 2002Ba85 I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, At. Data Nucl. Data Tables 81(2002)1 [Calculated ICC]
- 2003Au03 G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q value]
- 2003Br12 E. Browne, Nucl. Data Sheets 98(2003)665 [Spin, parity, multipolarity, mixing ratios]
- 2004Da24 F. Dayras, N. Chauvin, Nucl. Instrum. Meth. Phys. Res. A530(2004)391 [Alpha energies and intensities]
- 2004Sc03 R. Schön, G. Winkler, W. Kutschera, Appl. Radiat. Isotop. 60(2004)263 [Evaluated half-life]
- 2005Ga36 E. Garcia-Toraño, M.T. Crespo, M. Roteta, G. Sibbens, S. Pommé, A.M. Sanchez, M.P.R. Montero, S. Woods, A. Pearce, Nucl. Instrum. Meth. Phys. Res. A550(2005)581 [Alpha energies and intensities]
- 2006AL28 F.S. Al-Saleh, Al-J.H. Al-Mukren, M.A. Farouk, Nucl. Instrum. Meth. Phys. Res. A568(2006)734 [gamma-ray emission probabilities]
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr. , Nucl. Instrum. Meth. Phys. Res. A589(2008)202 [Theoretical ICC].

²³⁶U – Comments on Evaluation of Decay Data by A. Luca

This evaluation was completed in April 2008. The literature available by February 2008 was included.

1. Evaluation Procedures

The Limitation of Relative Statistical Weight (LWM) (1988WoZO) method was applied for averaging numbers throughout this evaluation; this method was implemented by using the computer code LWEIGHT, ver. 4 (designed for Excel, MS Office), [1] [2]. The uncertainty assigned to an average value in this evaluation is never lower than the lowest uncertainty of any of the experimental input values.

2. Decay Scheme

²³⁶U decays 100 % by alpha-particle emissions, mainly to the ground state and to the 49 keV excited level of ²³²Th. ²³⁶U decays also by spontaneous nuclear fission, with a weak branch (about 9 · 10⁻⁸ %). According to Tretyakova et al. (1994Tr12), a very weak cluster decay of ²³⁶U (~10⁻¹³ probability relative to the alpha emission), consisting of Ne and Mg emission, was observed. The spin, parity, energy and half-life of the ²³²Th excited levels, and the multiplicities of the γ -ray transitions have been adopted from the A=232 ENSDF mass-chain evaluation of E. Browne (2006Br19).

3. Nuclear Data

The adopted alpha-decay energy value $Q(\alpha) = 4573.1(9)$ keV, is from 2003Au03. This value is in agreement with the effective $Q(\alpha)$ value of 4570 keV (with an uncertainty of 260 keV), calculated from the decay scheme data, by using the SAISINUC software. This agreement proves the consistency and correctness of the decay scheme.

3.1. Half-life

The measured half-life ($T_{1/2}$) values, with the reviewed uncertainties (1989Ho24), are shown below in Table 1. After a new critical review (based on the most precise modern activity measurements by using the defined solid angle α -particle counting method, according to the Bureau International des Poids et Mesures (BIPM), Key Comparison Database, section “Calibration and Measurement Capabilities” (CMCs) - Ionizing Radiation, <http://kcdb.bipm.org/AppendixC/>), the uncertainty of the most recent half-life value (1972F103) was increased from about 0.06 % to 0.25 %; accordingly, the half-life was rounded from 2.3415 · 10⁷ to 2.342 · 10⁷ years. The set of data is consistent and the recommended value, 2.343 · 10⁷ years, with the uncertainty of 0.006 · 10⁷ years, is the weighted average (LWM, $\chi^2_{\nu}=0.72$) of the three input values. The references are expressed as NSR (Nuclear Science References) type keynumbers:

Table 1

$T_{1/2}$ (10 ⁷ years)	Uncertainty of $T_{1/2}$ (10 ⁷ years)	Reference
2.46	0.14	1951Ja09
2.391	0.057	1952F120
2.3415	0.0039	1972F103

The measured half-life ($T_{1/2}$) values for the ²³⁶U spontaneous fission are presented below in Table2:

Table 2

$T_{1/2\text{ sf}} (10^{16} \text{ years})$	Uncertainty of $T_{1/2\text{ sf}} (10^{16} \text{ years})$	Reference
2.0	1.6	Jaffey and Hirsch, 1949 [3]
2.7	0.3	1971Co35
2.43	0.13	1981Vo02
2.7	0.4	1983Be66

The value mentioned in ref. [3] was unpublished, but it is cited in E.K. Hyde, 1964 [4]. This data set is consistent, and the recommended value, $2.49 \cdot 10^{16}$ years, with the uncertainty of $0.13 \cdot 10^{16}$ years, is the weighted average (LWM, $\chi^2_{\nu} = 0.36$) of the four input values from the first column.

3.2. Alpha transitions and emissions

In the literature, only one reference about measurements of energy and emission probability for ^{236}U alpha transitions was found: 1960Ko04. In another reference (1992It01), the measured energy of the main alpha-particle emission (4.49 MeV) was reported.

For this evaluation, the energies and the intensities of α_0 and α_{49} are from 1960Ko04. The energy of α_{162} is also from 1960Ko04, but its intensity is from γ -ray transition intensity balance. The energy of α_{333} is from $Q(\alpha) = 4573.1 (9) \text{ keV}$ and $E(\text{level}) = 333.40 \text{ keV}$; its intensity is from γ -ray transition intensity balance (2006Br19). These values, as well as their α hindrance factors (HF) are shown in Table 3.

Table 3

$E_{\alpha} (\text{keV})$	Uncertainty $E_{\alpha} (\text{keV})$	Emission intensity (%)	α Hindrance Factor (HF)
4494	3	73.8 (40)	1.0
4445	5	26.1 (40)	1.2
4332	8	0.149 (22)	27.3
4168	-	0.000 14 (5)	1160

3.3. g- transitions: g rays and internal conversion electrons

Measurements of the two main γ -ray transition energies are presented in a paper by Schmorak *et al.*, 1972Sc01. Their uncertainties may have been somewhat underestimated for the detection system that they used. Measurements of the energies and relative intensities for the γ rays following the decay of ^{236}U were published only by Gehrke *et al.* (2002Ge02), as shown in Table 4.

The decay-scheme normalization condition applied for the ^{232}Th ground state, allowed the determination of the absolute emission probability for the 49.46 keV γ ray ($I_{\gamma 49}$, expressed in %):

$(\alpha_{49}^T + 1) \cdot I_{\gamma}(49) + I_{\alpha}(4494) = 100 \%$, where $\alpha_{49}^T = 324.4$ is the theoretical internal conversion coefficient (program BrIcc v2.0a, [5]) for the 49 -keV γ ray and $I_{\alpha}(4494) = 73.8 (40) \%$. The resulting value for the absolute emission probability of the main γ ray following the ^{236}U alpha decay, is $I_{\gamma}(49) = 0.081 (12) \%$. Using this value and the relative intensity values of the 112 keV and 171 keV γ -ray emissions measured by Gehrke *et al.*, the corresponding absolute emission probabilities and their uncertainties were computed and are given below in Table 4.

Table 4:

$E_{\gamma} (\text{keV})$	Uncertainty $E_{\gamma} (\text{keV})$	Relative Emission probability (%)	Absolute Emission probability (%)	Total ICC (α_T)
49.46	0.10	100	0.081 (12)	324
112.79	0.10	24.1 (1)	0.019 5 (31)	6.67
171.15	0.20	0.080 (24)	0.000 065 (22)	1.186

4. Atomic data

The K-shell fluorescence yield (ω_K), the mean L-shell fluorescence yield (ν_L) and the mean number of vacancies in the L-shell produced by one vacancy in the K-shell (η_{KL}) were determined using the computer program EMISSION v3.10, 28-Jan-2003 [6]: 0.969 (4), 0.476 (18) and 0.797 (5) respectively.

4.1. Auger electrons and X-rays

The relative probability values of the K Auger electron emissions (KLL, KLX, KXY) normalized to the KLL value, were computed using the same EMISSION computer program. The total K Auger electron emission probability (absolute) and the emission probability of the L Auger electrons were also calculated. The energy ranges for K and L Auger electrons were filled-in by the SAISINUC program [7]. The relative probability (normalized to $K_{\alpha 1}$ X-rays emission) and the absolute emission probability values of the different groups of K and L X-rays were determined using the same EMISSION program. The energy range values of the K and L X-rays are from the tables linked to SAISINUC. The results for absolute emission probabilities of LX rays ($I(LX) = 9.4$ (10) %) agrees with $I(LX) = 9.4$ (13) % given in the Table of Radioactive Isotopes [8]. The KX ray emission probabilities are so weak that are not given in reference [8].

Neither measurements of X-ray energies nor of emission probabilities were found in the literature.

5. Main production mode

The main production mode of ^{236}U is by irradiating ^{235}U nuclei with thermal neutrons in nuclear reactors; the ^{236}U is produced by thermal neutron captures: $^{235}\text{U}(n,\gamma)^{236}\text{U}$. The neutron-capture cross section is 98.3 (8) barn [9].

6. References

- [1] MacMahon and E. Browne, LWEIGHT, A Computer Program to Calculate Averages, version 1.3, March 2000 (electronic file LWEIGHT.doc distributed with the program).
- [2] M.M. Bé, C. Dulieu, LWEIGHT4 (instructions for use the MS Excel version of the program, electronic file), CEA/Laboratoire National Henri Becquerel, Saclay, France, 2005.
- [3] A.H. Jaffey, A. Hirsch, unpublished data, 1949 cited in ref. [4].
- [4] E.K. Hyde, "The Nuclear Properties of the Heavy Elements", Prentice-Hall, Englewood Cliffs, vol. III, page 75 (1964).
- [5] T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, C.W. Nestor Jr., "BrIcc Program Package v 2.0" (BrIcc Manual, pdf file), ANU-P/1684, December 2005.
- [6] E. Schönfeld, H. Janssen, "Calculation of emission probabilities of X-rays and Auger electrons emitted in nuclear disintegration processes", Appl. Radiat. Isot. 52, 595-600 (2000).
- [7] M.M. Bé, "SAISI NUC 2000 Manual", English version, LNE - LNHB/CEA -2006, Saclay, France.
- [8] E. Browne and R.B. Firestone, "Table of Radioactive Isotopes," John Wiley & Sons, Inc. (1986).
- [9] S.F. Mughabghab, M. Divadeenam and N.E. Holden, "Neutron cross sections from neutron resonance parameters and thermal cross sections", Academic Press, 1981.
- 1951Ja09 A. H. Jaffey, H. Diamond, A. Hirsch and J. Mech, "Half-Life and Alpha-Particle Energy of U^{236} ", Phys. Rev. 84, 785-786 (1951).
- 1952FI20 E.H. Fleming Jr., A. Ghiorso, B.B. Cunningham, "The Specific Alpha-Activities and Half-Lives of U^{234} , U^{235} , and U^{236} ", Phys. Rev. 88, 642-652 (1952).
- 1960Ko04 A.P. Komar, G.A. Korolev, G.E. Kocharov, "Investigation of the alpha decay of ^{236}U ", Zh. Eksp. Teor. Fiz. 38, 1436-1438 (1960).
- 1971Co35 H. Conde and M. Holmberg, "Prompt nuBar in Spontaneous and Neutron Induced Fission of ^{236}U and its Half-Life for Spontaneous Fission", J. Nucl. Energy 25, 331-338 (1971).
- 1972FI03 K.F. Flynn, A.H. Jaffey, W.C. Bentley, A.M. Essling, "Precision measurement of half-life and specific activity of ^{236}U ", J. Inorg. Nucl. Chem. 34, 1121-1129 (1972).

- 1972Sc01 M. Schmorak, C.E. Bemis Jr., M.J. Zender, N.B. Gove and P.F. Dittner, "Ground State Rotational Bands in Doubly Even Actinide Nuclei", Nuclear Physics A178, 410-416 (1972).
- 1981Vo02 H.R. von Gunten, A. Gruetter, H.W. Reist, M. Baggenstos, "Ground state spontaneous-fission half-lives of uranium isotopes", Phys. Rev. C23, 1110-1112 (1981).
- 1983Be66 S.N. Belenky, M.D. Skorokhvatov and A.V. Etenko, "Measurement of the Characteristics of Spontaneous Fission of ^{238}U and ^{236}U ", Sov. At. Energy 55, 528-531 (1983).
- 1988WoZO M.J. Woods and A.S. Munster, Evaluation of Half-Life Data, National Physical Laboratory, Teddington, UK, Rep. RS(EXT) 95 (1988).
- 1989Ho24 N.E. Holden, "Total and spontaneous fission half-lives for Uranium, Plutonium, Americium and Curium nuclides", Pure Appl. Chem. 61, 1483-1504 (1989).
- 1992It01 J.L. Iturbe, "Identification of ^{236}U in commercially available uranium compounds by alpha particle spectrometry", Appl. Radiat. Isot. 43, 817-818 (1992).
- 1994Tr12 S.P. Tretyakova, V.L. Mikheev, V.A. Ponomarenko, A.N. Golovchenko, A.A. Ogloblin, V.A. Shigin, "Cluster decay of ^{236}U ", Pisma Zh. Eksp. Teor. Fiz. 59, 368 (1994).
- 2002Ge02 R.J. Gehrke, J.D. Baker, C.L. Riddle, "Feeding of the ^{232}Th levels from the decay of ^{236}U ", Appl. Radiat. Isot. 56, 567-568 (2002).
- 2003Au03 G. Audi, A.H. Wapstra and C. Thibault, The AME2003 atomic mass Evaluation (II). Tables, graphs, and references, Nucl. Phys. A729, 337-676 (2003).
- 2006Br19 E. Browne, "Nuclear Data Sheets for A=232", Nucl. Data Sheets 107, 2579-2648 (2006).

²³⁶Np – Comments on evaluation of decay data by V.P. Chechev and N.K. Kuzmenko

This evaluation was done originally in June 2006 and then updated in April 2009 with a literature cut-off by the same date.

1. DECAY SCHEME

From the systematics of the isomer levels it has been assumed in 1981Li30 (see also the analysis carried out in 1991Sc08) that the short-lived state of ²³⁶Np (22,5 h) lies higher in energy than the long-lived state of ²³⁶Np ($1,55 \times 10^5$ a). In line with this assumption we consider the long-lived state of ²³⁶Np as the ground state. Using Q values for electron capture decay of the isomer and ground state and a close energy cycle we can estimate the energy level spacing between these states as 60 (50) keV.

The decay scheme of the long-lived ²³⁶Np includes three decay modes: β^- decay to ²³⁶Pu, electron capture decay (EC) to ²³⁶U and α decay to ²³²Pa (see 2006Br20). A favored α -particle branch to the (6^-) level at ≈ 400 keV is expected in ²³²Pa from α systematics (1972El21, 1980Sc26, 2006Br20). However, this decay was not observed experimentally.

The β^- -decay branching, $\Sigma P(\beta^-)$, and alpha-decay branching, $\Sigma P(\alpha)$, have been deduced by the evaluators from the partial half-lives $T_{1/2}(\beta^-)$ and $T_{1/2}(\alpha)$, respectively, measured in 1981Li30. The EC-decay branching, $\Sigma P(\text{EC})$, has been obtained as the difference of $1 - \Sigma P(\beta^-) - \Sigma P(\alpha)$.

2. NUCLEAR DATA

Q^- , Q_{EC} , $Q(\alpha)$ values are from 2003Au03.

The total half-life of ²³⁶Np is based on the evaluated partial half-lives $T_{1/2}(\alpha)$, $T_{1/2}(\beta^-)$, $T_{1/2}(\text{EC})$ measured in 1981Li30.

The evaluated $T_{1/2}(\alpha) = 9,5 (35) \times 10^7$ years has been obtained as an average of the two measurements of 1981Li30 (specific activity, ²³²U gamma-ray of 894 keV was measured): $9,4 (35) \times 10^7$ and $9,6 (35) \times 10^7$ years. A standard deviation of the individual measurement has been adopted for the uncertainty of the evaluated alpha-decay half-life using a rule that the uncertainty assigned to the recommended value should be greater than or equal to the smallest uncertainty in any experimental value.

$T_{1/2}(\beta^-) = 1,29 (3) \times 10^6$ years has been adopted here from the ²³⁶Pu growth measurement of 1981Li30. The result of this measurement is independent of the decay scheme, and it is equal to the weighted average of 1,34 (15), 1,29 (3), 1,32 (9), 1,69 (30), 1,29 (3), 1,31 (8) (in 10^6 years) given in 1981Li30. The uncertainties of these measurements do not include any estimation of uncertainties from the decay scheme parameters. It agrees well with an earlier measurement in 1972En06 ($1,29 (+ 0,07, - 0,05) \times 10^5$ a).

The evaluated $T_{1/2}(\text{EC}) = 1,77 (10) \times 10^5$ years has been obtained as an average of the two ²³⁶U/²³⁵U mass ratio measurements in 1981Li30: $1,75 (10) \times 10^5$ and $1,79 (10) \times 10^5$ years. These ²³⁶U growth measurement results are independent of the decay scheme. A standard deviation of the individual measurement has been adopted for the uncertainty of the evaluated partial EC-decay half-life. The specific gamma-ray activity method (²³⁶U 160,3-keV gamma-ray was measured) was used in other measurements presented in 1981Li30 (in 10^5 years): 1,60 (4), 1,73 (2), 1,77 (11), 1,75 (10), 1,79 (10),

1,74 (1), 1,78 (10). The uncertainties of these measurements do not include an estimation of uncertainties from the decay scheme parameters.

Thus, the recommended value of the total ²³⁶Np half-life obtained from the relation $T_{1/2} = [(T_{1/2}(\alpha))^{-1} + (T_{1/2}(\beta^-))^{-1} + (T_{1/2}(\text{EC}))^{-1}]^{-1}$ is $1,55 (8) \times 10^5$ years.

2.1.1. Electron Capture Transitions

The energies of the electron capture transitions have been deduced from the Q_{EC} value and the level energies given in Table 1 from 2006Br20 where they were deduced from a least squares fit to gamma-ray energies.

Table 1. ²³⁶U levels populated in ²³⁶Np electron capture decay

Level number	Energy (keV)	Spin and parity	Half-life	Probability of ε - transition (x 100)
0	0,0	0 ⁺	2,342·10 ⁷ a	-
1	45,2440 (20)	2 ⁺	234 (6) ps	-
2	149,477 (6)	4 ⁺	124 (7) ps	0,0 (44)
3	309,785 (7)	6 ⁺	58 (3) ps	87,8 (43)
4	687,59 (4)	1 ⁻	3,78 (9) ns	-
5	744,18 (7)	3 ⁻	< 0,1 ns	-
6	848,1 (8)	5 ⁻		~ 0,09

The probabilities of the electron capture transitions $P(\text{EC}_{0,2})$ and $P(\text{EC}_{0,3})$ have been deduced from the correlations of:

$$P(\text{EC}_{0,2}) + P(\text{EC}_{0,3}) = 100 \% - \sum P(\beta^-) - \sum P(\alpha) = 87,8 (6) \% \text{ and } P(\text{EC}_{0,3}) = P(\gamma_{3,2} + \text{ce})(160\text{-keV}).$$

The upper limit of $P(\text{EC}_{0,2}) < 4,4 \%$ has been obtained from the level intensity balance: $P(\text{EC}_{0,2}) = 0,0 (44) \%$. The estimate of $P(\text{EC}_{0,6}) \sim 0,1 \%$ is given in 1996FiZX.

2.1.2. Beta Transitions

The energies of the β^- transitions have been deduced from the Q^- value and the level energies given in Table 2 from 2006Br20 where they were deduced from a least squares fit to gamma-ray energies.

Table 2. ²³⁶Pu levels populated in ²³⁶Np β^- -decay

Level number	Energy (keV)	Spin and parity	Half-life	Probability of β^- - transition (x100)
0	0,0	0 ⁺	2,858 a	-
1	44,63 (10)	2 ⁺		-
2	147,45 (10)	4 ⁺		0,2 (14)
3	305,80 (11)	6 ⁺		11,8 (12)

The β^- transition probability $P(\beta_{0,3}) = P(\gamma_{3,2} + \text{ce})(158\text{-keV})$ and $P(\beta_{0,2}) = 12,0 (6) \% - P(\beta_{0,3}) = 0,2 (14) \%$. An upper limit of $P(\beta_{0,2}) < 1,6 \%$ follows this result.

2.2. Gamma Transitions and Internal Conversion Coefficients

The evaluated energies of gamma-ray transitions are virtually the same as the photon energies because nuclear recoil is negligible.

The gamma-ray transition probabilities have been obtained from the gamma-ray emission probabilities and the total internal conversion coefficients (ICCs).

Multipolarities of gamma-ray transitions have been taken from 2006Br20. The ICCs have been interpolated using the BrIcc package with the so called “*Frozen Orbital*” approximation (2008Ki07) except for $\gamma_{4,1}$ (642,3-keV) and $\gamma_{4,0}$ (687,6-keV). The relative uncertainties of the ICCs for pure multipolarities have been taken as 2 %.

For $\gamma_{4,1}$ (642,3-keV) and $\gamma_{4,0}$ (687,6-keV) the ICC values of α_K and α_L are experimental results from ²⁴⁰Pu α -decay study (1969Le05, 1977Po05). The ICC values of α_M and α_T for these transitions have been deduced using α_M/α_L and α_{NO}/α_M from 1971Dr11. More accurate ICC measurements for these E1 anomalously converted gamma-ray-transitions are required.

3. ATOMIC DATA

The atomic data are from Schönfeld and Janßen (1996Sc06).

4. ELECTRON EMISSIONS

The energies of the conversion electrons have been obtained from the gamma transition energies and the atomic electron binding energies.

The emission probabilities of the conversion electrons have been obtained using evaluated $P(\gamma)$ and ICC values.

The absolute emission probabilities of K and L Auger electrons have been obtained using the EMISSION computer program.

β^- average energies have been obtained using the LOGFT computer program.

5. PHOTON EMISSIONS

5.1. X-Ray Emissions

The absolute emission probabilities of U and Pu KX- and LX-rays have been deduced using the EMISSION computer program.

For U LX-ray intensity calculations the theoretical fractional ratios $P_{EC}(L2)/P_{EC}(L1) = 0,115$ and $P_{EC}(L3)/P_{EC}(L1) = 0$ for the EC-transition to the level “3” (309 keV) of ²³⁶U have been used (1972Dzhelepov). The calculated relative intensities of U KX- rays accompanying the electron capture of ²³⁶Np are in a good agreement with the experimental results (Table 3).

Table 3. Intensities of U KX- rays (relatively to $P(\gamma_{3,2}-160,3 \text{ keV})$) accompanying ²³⁶Np electron capture

	1983Ah03 (experimental)	Adopted (deduced)
X_K		
$K\alpha_2$	0,61 (2)	0,64 (3)
$K\alpha_1$	0,99 (3)	1,02 (5)
$K\beta_1'$	0,38 (2)	0,368 (19)
$K\beta_2'$	0,131 (7)	0,126 (7)

5.2. Gamma Ray Emissions

5.2.1. Gamma Ray Energies (²³⁶U)

The energies of gamma-rays in ²³⁶Np electron capture have been taken from 2006Br20.

5.2.2. Gamma Ray Energies (²³⁶Pu)

The energies of gamma-rays $\gamma_{1,0}$ (44,6 keV), $\gamma_{2,1}$ (102,8 keV), $\gamma_{3,2}$ (158,3 keV) accompanying β^- decay of ²³⁶Np have been adopted from measurements given in 1983Ah02 (see also 2006Br20).

5.2.3. Gamma-Ray Emission Probabilities (²³⁶U)

The evaluated gamma-ray emission probabilities $P(\gamma)$ have been deduced using the relative gamma-ray intensities from 1983Ah02 (Table 4), the relation of $\sum P(EC_{0,i}) = 87,8$ (6) % = $P(\gamma_{2,1} + ce)(104,23$ keV) and the intensity balance at ²³⁶U each level. We have assumed that the populations to the two lower levels ("0" and "1") in the ²³⁶Np electron capture decay are negligible and have taken into account the intensity balance correlation for the gamma-ray transitions to these levels, that is $P(\gamma_{1,0} + ce)(45,2$ keV) = $P(\gamma_{2,1} + ce)(104,2$ keV).

The recommended gamma-ray emission probabilities for γ -rays de-exciting level "4" ($\gamma_{4,2}(538,1$ keV), $\gamma_{4,1}(642,3$ keV), and $\gamma_{4,0}(687,6$ keV)) have been deduced from the correlation:

$P(\gamma_{5,4} + ce)(56,6$ keV) = $P(\gamma_{4,2} + ce)(538,1$ keV) + $P(\gamma_{4,1} + ce)(642,3$ keV) + $P(\gamma_{4,0} + ce)(687,6$ keV) using the relative intensities for these γ -rays evaluated from the ²⁴⁰Pu α -decay (Table 5) and assuming $P(EC_{0,4}) = 0$.

Table 4. Gamma rays in decay of the long-lived ²³⁶Np measured in 1983Ah02

	Energy (keV)	Relative intensity
$\gamma_{1,0}$ (²³⁶ U)	45,23 (3)	0,4 (1)
$\gamma_{2,1}$ (²³⁶ Pu)	102,82 (2)	2,9 (2)
$\gamma_{2,1}$ (²³⁶ U)	104,23 (2)	23 (1)
$\gamma_{3,2}$ (²³⁶ Pu)	158,35 (2)	13,5 (7)
$\gamma_{3,2}$ (²³⁶ U)	160,33 (2)	100

Table 5. Experimental and evaluated absolute emission probabilities of gamma-rays de-exciting the ²³⁶U level with energy of 687,6 keV in decay of ²⁴⁰Pu (per 10⁸ α -decays) and the deduced relative intensities of these gamma-rays

	Energy (keV)	1969Le05	1971GuZY	1975OtZX	1975Dr05	1976GuZN	Evaluated	Evaluated relative intensities
$\gamma_{4,2}$	538,1	$\approx 0,23$ ^a		0,147 (12)			0,147 (12)	1,17 (10)
$\gamma_{4,1}$	642,3	14,5 ^a	14,5 (5) ^b	12,6 (4)	13 (1)	12,45 (30)	12,6 (3) ^c	100 (3)
$\gamma_{4,0}$	687,6	3,77 (11)	3,70 (15) ^b	3,30 (13)		3,55 (9)	3,56 (15) ^d	28,3 (13)

^a Omitted from averaging as uncertainty is not quoted

^b Omitted from averaging as the data of 1971GuZY have been revised in 1976GuZN

^c Weighted average of 3 experimental values; the uncertainty is the smallest quoted uncertainty

^d Weighted average of 3 experimental values; the uncertainty is external

5.2.4. Gamma-Ray Emission Probabilities (²³⁶Pu)

The recommended gamma-ray emission probabilities $P(\gamma)$ have been deduced using the relative gamma-ray intensities from 1983Ah02 (Table 4), the quantity $\sum P(\beta^-) = 12,05 (60) \% = P(\gamma_{2,1+ce})(102,8 \text{ keV})$ and the intensity balance at each ²³⁶Pu level. We have assumed that the populations to the two lower levels (“0” and “1”) in the ²³⁶Np beta minus decay are negligible and have taken into account the intensity balance of the gamma-ray transitions to these levels, that is $P(\gamma_{1,0+ce})(44,6 \text{ keV}) = P(\gamma_{2,1+ce})(102,8 \text{ keV})$.

6. CONSISTENCY OF RECOMMENDED DATA

The most accurate Q value, $Q(M)$, is taken from the atomic mass adjustment table of Audi et al. (2003Au03). Comparison of $Q(\text{eff})$ (deduced as the sum of average energies per disintegration ($\sum E_i \times P_i$) for all emissions accompanying ²³⁶Np β^- -decay or ²³⁶Np electron capture) with the tabulated decay energy $Q^-(M) \times P(\beta^-)$ for β^- -decay or $Q_{EC}(M) \times P(EC)$ for electron capture allows to check a consistency of the recommended ²³⁶Np decay-scheme parameters obtained in this evaluation.

Here E_i and P_i are the evaluated energies and emission probabilities of the i -th beta particle, gamma ray, X-ray, etc. The values of $P(\beta^-)$, $P(EC)$ are β^- -decay and EC branching, respectively. Consistency (percentage deviation) is determined by $\{[Q(M) - Q(\text{eff})]/Q(M)\} \times 100$. “Percentage deviations above 5 % would be regarded as high and imply a poorly defined decay scheme; a value of less than 5 % indicates the construction of a reasonably consistent decay scheme” (quoted from the article by A. L. Nichols in Appl. Rad. Isotopes 55(2001)23-70).

For the above ²³⁶Np decay data evaluation we have for β^- -decay $Q^-(M) \times P(\beta^-) = 58 (7) \text{ keV}$ and $Q(\text{eff}, \beta^-) = 57 (7) \text{ keV}$, i.e. consistency is less than 2 % if we do not take into account the uncertainties, and the exact percentage deviation is $(1.7 \pm 17) \%$ if we consider the uncertainties. Similarly, for ²³⁶Np electron capture we have $Q_{EC}(M) \times P(EC) = 817 (44) \text{ keV}$ and $Q(\text{eff}, EC) = 817 (50) \text{ keV}$ and the percentage deviation is $0 \pm 8 \%$. These values indicate the right evaluation and inaccurate measurements of ²³⁶Np decay-scheme parameters.

7. REFERENCES

- 1969Le05 C. M. Lederer, J. M. Jaklevic, S. G. Prussin, Nucl. Phys. A135(1969)36 (Relative intensities of gamma rays).
- 1971GuZY R. Gunnink, R. J. Morrow, UCRL 51087(1971) (Emission probabilities of gamma-rays in the decay of ²⁴⁰Pu).
- 1971Dr11 O. Dragoun, Z. Plajner, F. Schmutzler, NDT A9(1971)119 (α_M / α_L and α_{NO} / α_M).
- 1972Dzhelepov B. S. Dzhelepov, L. N. Zyryanova, Yu. P. Suslov, Beta-processes, 1972, Nauka, Leningrad (Fractional probabilities in L-electron capture).
- 1972El21 Y. A. Ellis, M. R. Schmorak, Nucl. Data Sheets B8(1972)345 (Systematics of nuclear level properties).
- 1972En06 D. W. Engelkemeir, J. E. Gindler, J. Inorg. Nucl. Chem. 34(1972)1799 (Half-life).
- 1975OtZX H. Ottmar, P. Matussek, I. Piper, Proc Int. Symp. Neutron Capture Gamma Ray Spectroscopy and Related Topics, 2nd, Petten, The Netherlands (1974), K. Abrahams, F. Stecher Rasmussen, P. Van Assche, Eds, Reactor Centrum Nederland, p 658(1975) (Emission probabilities of gamma-rays in decay of ²⁴⁰Pu).
- 1975Dr05 T. Dragnev, K. Scharf, Intern. J. Appl. Radiat. Isotop. 26(1975)125 (Gamma ray emission probabilities in decay of ²⁴⁰Pu).
- 1976GuZN R. Gunnink, J. E. Evans and A. L. Prindle, UCRL-52139(1976) (Emission probabilities of gamma-rays in decay of ²⁴⁰Pu).
- 1977Po05 W. L. Posthonus, K. E. G. Löbner, I. Piper et al., Z. Phys. A281(1977)717 (ICC measurements).

- 1980Sc26 M. R. Schmorak, Nucl. Data Sheets 31(1980)283 (Systematics of nuclear level properties).
- 1981Li30 M. Lindner, R. J. Dupzyk, R. W. Hoff, R. J. Nagle, J. Inorg. Nucl. Chem. 43(1981)3071 (Half-life, partial half-lives).
- 1983Ah02 I. Ahmad, J. Hines, J. E. Gindler, Phys. Rev. C27(1983)2239 (Gamma-ray relative intensities and energies, KX-ray energies)
- 1991Sc08 M. R. Schmorak, Nucl. Data Sheets 63(1991)139 (Analysis of isomer levels in Np-236).
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 (Atomic data).
- 1996FiZX R. B. Firestone, Table of Isotopes, Eighth Edition, Volume II: A=151-272, V. S. Shirley (Editor), C. M. Baglin, S. Y. F. Chu and J. Zipkin (Assistant Editors), 1996, 1998, 1999 (β^- -transition probabilities).
- 2003Au03 G. Audi, A. H. Wapstra and C. Thibault, Nucl. Phys. A729(2003)337 (Q values).
- 2006Br20 E. Browne and J. K. Tuli, Nucl. Data Sheets 107(2006)2579 and 2649 (Decay scheme, level energies, gamma-ray multipolarities).
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson and C. W. Nestor, Jr., Nucl. Instrum. Methods Phys. Res. A589(2008)202 (Theoretical ICC).

²³⁶Np^m – COMMENTS ON EVALUATION OF DECAY DATA

by V.P.Chechev and N.K.Kuzmenko

This evaluation was completed in June 2006. The literature available by May 2006 was included.

1. DECAY SCHEME

From the systematics of isomer levels it was assumed in 1981L i30 (see also the analysis carried out in 1991Sc08) that the short-lived state of ²³⁶Np (22,5 h) lies higher in energy than the long-lived state of ²³⁶Np (1,55 · 10⁵ y). In line with this assumption we have considered the long-lived state of ²³⁶Np as the ground state. Using Q values for electron capture decays of the isomer and ground states together with closed energy cycles we can estimate the energy level spacing between these states as 60(50) keV.

The decay scheme of the isomer ²³⁶Np^m includes two decay modes: β⁻ decay to ²³⁶Pu and electron capture decay (EC) to ²³⁶U (see evaluations of 1991Sc08, 1996FiZX). The β⁻-decay branching, ΣP(β⁻), has been adopted from 1969Le05. The EC -decay branching, ΣP(EC), has been obtained as the difference of 1-ΣP(β⁻).

2. NUCLEAR DATA

Q⁻(²³⁶Np^m) is from 1969Le05 (the end-point energy of the β⁻ spectrum was measured). Q_{EC}(²³⁶Np^m) has been calculated from the closed energy cycle of decays ending in ²³²Th. The values of Q⁻(²³⁶Np^m), Q_α(²³⁶Pu), Q⁻(²³²Pa), Q_{EC}(²³²Pa) and Q_α(²³⁶U) from 2003Au03 were used in this calculation.

The half-life of ²³⁶Np^m is from 1969Le05. This result agrees with other (less accurate) measurements (1949Ja01 – 22 h, 1984Gr33 – 22,5 h).

2.1. Electron Capture Transitions

The energies of the electron capture transitions have been deduced from the Q_{EC} value and the level energies (Table 1) obtained from the evaluated gamma-ray energies.

Table 1. ²³⁶U levels populated in the ²³⁶Np^m electron capture decay

Level number	Energy, keV	Spin and parity	Half-life	Probability of EC - transition (×100)
0	0,0	0 ⁺	2,342·10 ⁷ yr	43,1(32)
1	45,242(3)	2 ⁺	234 ps	8,3(30)
2	149,476(15)	4 ⁺	124 ps	-
4	687,60(5)	1 ⁻	3,8 ns	1,64(9)

The individual EC- transition probabilities P(EC_{1,i}) have been deduced from the intensity balance for each level and the total EC -decay probability ΣP(e_{1,i}).

2.2. Beta Transitions

The β⁻- transition energies have been deduced from the Q⁻ value and the level energies (Table 2) obtained from the evaluated gamma-ray energies.

Table 2. ²³⁶Pu levels populated in the ²³⁶Np^m β⁻-decay

Level number	Energy, keV	Spin and parity	Half-life	Probability of β ⁻ - transition (× 100)
0	0,0	0 ⁺	2,858 yr	36(4)
1	44,63(10)	2 ⁺		11(4)

The β^- transition probabilities $P(\beta_{1,0})$, $P(\beta_{1,1})$ have been obtained from the ratio $P(\beta_{1,0})/P(\beta_{1,1}) = 38(7)/12(5)$ measured in 1959Gi58 and the total β^- -decay probability $\Sigma P(\beta_{1,i})$.

2.3. Gamma Transitions and Internal Conversion Coefficients (²³⁶U)

The evaluated transition energies are virtually the same as the photon energies because nuclear recoil is negligible.

The gamma-ray transition probabilities have been obtained from the gamma-ray emission probabilities and the total internal conversion coefficients (ICC's). Multipolarities of gamma-ray transitions have been taken from 1991Sc08 and 1996FiZX. The gamma-ray transition probability $P(\gamma_{1,0} + \text{ce})(44,6\text{-keV})$ has been deduced from the relation of $P(\gamma_{1,0} + \text{ce})(44,6\text{-keV}) = P(\beta_{0,1})$.

ICC's have been interpolated using the BRICC computer program, except for $\gamma_{4,1}$ (642,3-keV) and $\gamma_{4,0}$ (687,6-keV) because of nuclear penetration effects. The relative uncertainties of α_K , α_L , α_M , α_T for pure multipolarities have been taken as 2%.

α_K and α_L for $\gamma_{4,1}$ (642,3-keV) and $\gamma_{4,0}$ (687,6-keV) are experimental values from data in ²⁴⁰Pu α -decay (1969Le05 and 1977Po05, see also the evaluation of 2004Be). α_M and α_T for these transitions have been evaluated using α_M/α_L and α_{NO}/α_M from 1971Dr11. More accurate ICC measurements for these transitions are required.

3. ATOMIC DATA

3.1. Fluorescence yields

The fluorescence yield data are from 1996Sc06 (Schönfeld and Janßen).

3.2. X Radiations

The LX-ray energies are from 1996FiZX. The KX-ray energies and the relative KX-ray emission probabilities are from 1999Schönfeld.

The X-ray energies are based on the wavelengths given in the compilation of 1967Be65 (Bearden).

The relative KX-ray emission probabilities have been taken from 1999Schönfeld.

3.3. Auger Electrons

The ratios $P(\text{KLX})/P(\text{KLL})$, $P(\text{KXY})/P(\text{KLL})$ are taken from 1996Sc06.

4. ELECTRON EMISSIONS

The energies of the conversion electrons have been deduced from the gamma transition energies and the electron binding energies.

The emission probabilities of the conversion electrons have been deduced using evaluated P_γ and ICC values.

The absolute emission probabilities of K and L Auger electrons have been obtained with the EMISSION computer program.

β^- average energies have been obtained using the LOGFT computer program.

5. PHOTON EMISSIONS

5.1. X-Ray Emissions

The absolute emission probabilities of KX- and LX-rays have been obtained with the EMISSION computer program.

For U LX-ray calculations the ratios $P_{\text{EC}}(\text{L2})/P_{\text{EC}}(\text{L1}) = 0,115$ and $P_{\text{EC}}(\text{L3})/P_{\text{EC}}(\text{L1}) = 0$ from the theoretical calculations of 1972Dzheleпов were used for all levels populated in the ²³⁶Np^m electron capture decay.

5.2. Gamma Ray Emissions

5.2.1. Gamma Ray Energies (²³⁶U)

The energies of gamma rays accompanying the ²³⁶Np^m electron capture decay have been adopted from the evaluated DDEP data in ²⁴⁰Pu α -decay (2004Be).

5.2.2. Gamma Ray Energies (²³⁶Pu)

The energy of $\gamma_{1,0}$ (44,6 keV) accompanying the β^- - decay of ²³⁶Np^m has been adopted from measurements in 1983Ah02.

5.2.3. Gamma-Ray Emission Probabilities (²³⁶U)

The gamma-ray emission probability $P(\gamma)$ for $\gamma_{1,0}$ (45,2 keV) has been obtained from the ratio $\Sigma P(e_i)(45,2 \text{ keV}) / P(\gamma_{4,1})(642,3 \text{ keV}) = 9(3)$ measured in 1969Le05.

The evaluated gamma ray emission probability $P(\gamma_{4,1})(642,3 \text{ keV}) = 0,96(20)\%$ has been deduced using the following values:

- 1) $\Sigma P(e_{1,i})=53(1)\%$;
- 2) measured ratio $P(XK\alpha) / P(\gamma_{3,1})(642,3 \text{ keV})=27,6(10)$ from 1969Le05;
- 3) theoretical value of the ratio $P(XK\alpha)/P(XK\beta)=0,298(5)$;
- 4) relative (partial) intensities of gamma rays de -exciting level "4" [$\gamma_{4,2}$ (538,1 keV), $\gamma_{4,1}$ (642,3 keV), $\gamma_{4,0}$ (687,5 keV)], which have been deduced from the absolute gamma -ray emission probabilities evaluated in the ²⁴⁰Pu α -decay (Table 5), and a_K for these gamma-rays;
- 5) the measured ratio $\Sigma P_K(i) P(EC_{1,i}) / \Sigma P(\beta_{1,i})=0,75(15)$ from 1956Gr11, which can be represented as $P_K^{(average)} = \Sigma P_K(i) P(EC_{1,i}) / \Sigma P(\beta_{1,i})=0,67(13)$.

The most accurate evaluation of $P_K^{(average)}$ (and also the new evaluation of $P(\gamma_{4,1})$ (642,3 keV) and other values) may be obtained by using the theoretical $P_K(i)$, the values of $P(EC_{1,i})$ deduced from $P(\gamma_{4,1})(642,3 \text{ keV}) = 0,96(20)\%$, and the fact that a contribution of the third term (with $P(EC_{1,4})$) to $P_K^{(average)}$ comprises $\sim 2,5\%$. This value has been taken as a fractional uncertainty for the $P_K^{(average)} = 0,75(2)$. Using the latter and the relations 1) - 4) we have deduced a more accurate evaluation of $P(\gamma_{4,1})(642,3 \text{ keV}) = 1,08(6)\%$, and correspondingly a more accurate evaluation for other decay data.

The gamma-ray emission probability $P(\gamma_{2,1})$ (104,2 keV) has been calculated from $P(\gamma_{2,1} + ce)$ (104,2 keV) = $P(\gamma_{4,2} + ce)(538,1 \text{ keV})$ assuming that the electron capture feeding of level "2" is negligible.

Table 5. Experimental and evaluated absolute emission probabilities of gamma rays de -exciting the ²³⁶U level with energy of 687,6 keV in the decay of ²⁴⁰Pu (per 10⁸ a-decays) and the deduced relative intensities of these gamma rays

	Energy, keV	1969Le05	1971GuZY	1975OtZX	1975Dr05	1976GuZN	Evaluated	Evaluated relative intensities
$\gamma_{4,2}$	538,1	$\approx 0,23^a$		0,147(12)			0,147(12)	1,17(10)
$\gamma_{4,1}$	642,3	14,5 ^a	14,5(5) ^b	12,6(4)	13(1)	12,45(30)	12,6(3) ^c	100 (3)
$\gamma_{4,0}$	687,6	3,77(11)	3,70(15) ^b	3,30(13)		3,55(9)	3,56(15) ^d	28,3(13)

^a Omitted from averaging as uncertainty is not quoted

^b Omitted from averaging as the data of 1971GuZY have been revised in 1976GuZN

^c Weighted mean of 3 experimental values; the uncertainty is the smallest quoted uncertainty

^d Weighted mean of 3 experimental values; the uncertainty is external

5.2.4. Gamma-Ray Emission Probability (²³⁶Pu)

The gamma-ray emission probability $P(\gamma)$ for $\gamma_{1,0}$ (44,6 keV) has been obtained from $P(\beta_{1,1})$ and the adopted α_T for this gamma-ray transition.

6. REFERENCES

1949Ja01 R.A.James, A.E.Florin, H.H.Hopkins, Jr., and A.Ghiorso, NNS 14B (1949) 1604 (Half-life)
 1956Gr11 P.R. Gray, Phys.Rev. 101(1956) 1306. (Relative probability of K-electron capture in the decay of ^{236m}Np)
 1959Gi58 J.E. Gindler, R.K. Sjoblom, J. Inorg. Nucl. Chem. 12 (1959) 8. (Probabilities of beta transitions)
 1967Be65 J.A. Bearden, Rev. Mod. Phys. 39 (1967) 78. (X-ray energies)

- 1969Le05 C.M. Lederer, J.M. Jaklevic, S.G. Prussin, Nucl. Phys. A135 (1969) 36.
(Relative intensities of gamma rays)
- 1971GuZY R. Gunnink, R.J. Morrow, In: UCRL 51087 (1971).
(Emission probabilities of gamma-rays in the decay of ²⁴⁰Pu)
- 1971Dr11 O. Dragoun, Z. Plajner, F. Schmutzler, NDT A9 (1971) 119.
(α_M / α_L and α_{NO} / α_M)
- 1972Dzhelepov B.S. Dzhelepov, L.N. Zyryanova, Yu.P. Suslov, Beta-processes, 1972, Nauka, Leningrad
(Fractional probabilities in L-electron capture)
- 1975OtZX H. Ottmar, P. Matussek, I. Piper In: Proc Int Symp Neutron Capture Gamma Ray Spectroscopy and Related Topics, 2nd, Petten, The Netherlands (1974), K. Abrahams, F Stecher-Rasmussen, P Van Assche, Eds, Reactor Centrum Nederland, p 658 (1975).
(Emission probabilities of gamma-rays in the decay of ²⁴⁰Pu)
- 1975Dr05 T. Dragnev, K. Scharf, Intern. J. Appl. Radiat. Isotop. 26 (1975) 125.
(Gamma ray emission probabilities in the decay of ²⁴⁰Pu)
- 1976GuZN R. Gunnink, J.E. Evans and A.L. Prindle, UCRL-52139 (1976).
(Emission probabilities of gamma-rays in the decay of ²⁴⁰Pu)
- 1977Po05 W.L. Posthonus, K.E.G. Löbner, I. Piper e.a., Z. Phys. A281 (1977) 717
(ICC measurements)
- 1984Gr33 E.A.Gromova, S.S.Kovalenko, Yu.A.Nemilov, Yu.A.Selitsky, A.V.Stepanov, A.M.Fridkin, V.B.Funshtein, V.A.Yakovlev, G.V.Valsky and G.A.Petrov, At.Energ. 56 (1984) 212;
Sov.At.Energy 56 (1984) 230
(Half-life)
- 1991Sc08 M.R.Schmorak, Nucl.Data Sheets 63 (1991) 139.
(Decay scheme, gamma ray multipolarities)
- 1996Sc06 E. Schönfeld and H. Janßen, Nucl. Instrum. Methods Phys. Res. A369, 527 (1996)
(Atomic data)
- 1996FiZX R.B. Firestone, Table of Isotopes, Eighth Edition, Volume II: A=151-272, V.S. Shirley (Editor), C.M. Baglin, S.Y.F. Chu, and J. Zipkin (Assistant Editors), 1996, 1998, 1999
(Decay scheme, LX ray energies, multipolarities)
- 1999Schonfeld E. Schonfeld and G. Rodloff - PTB-6.11-1999-1999-1, Braunschweig, February 1999
(KX-ray energies and relative emission probabilities)
- 2003Au03 G. Audi, A.H. Wapstra, and C. Thibault, Nucl. Phys. A729, 337 (2003)
(Q values)
- 2004Be M. M. Bé, V. Chisté, C. Dulieu, E. Browne, V. Chechev, N. Kuzmenko, R. Helmer, A. Nichols, E. Schönfeld, and R. Dersch. Table of Radionuclides, Vol.2. A = 151 to 242. ²⁴⁰Pu.
– Bureau International des Poids et Mesures, 2004. See also: Recommended Data by the Decay Data Evaluation Project working group. ²⁴⁰Pu.
http://www.nucleide.org/DDEP_WG/DDEPData.htm

²³⁷U - Comments on evaluation of decay data by V.P. Chechev and N.K. Kuzmenko

This evaluation was done originally in September 2005 and then revised in April 2009 with a literature cut-off by the same date.

1 Decay Scheme

The decay scheme is based on 2006Ba41.

2 Nuclear Data

Q⁻ value is from 2003Au03.

The recommended half-life of ²³⁷U is based on the experimental results given in Table 1.

Table 1. Experimental values of the ²³⁷U half-life (in days)

Reference	Author(s)	Value
1949Me43	Melander and Slatis	6,63 (5)
1953Wa05	Huizenga and Flynn	6,75 (1)
1958Ca16	Cabell et al.	6,752 (2)

The weighted mean of the 3 values from the Table 1 of 6,752 (2) is dominated by the very accurate value of 1958Ca16. The EV1NEW computer program, which uses the limitation of relative statistical weights by 0,5 (LRSW method), increased the 1958Ca16 uncertainty from 0,002 to 0,0098 and gave 6,749 (16).

Therefore, the recommended value of ²³⁷U half-life is 6,749 (16) days.

2.1 Beta Transitions

The energies of β⁻ transitions have been obtained from the Q⁻ value and the level energies given in Table 2 from 2006Ba41.

Table 2. ²³⁷Np levels populated in ²³⁷U β⁻ decay

Level	Energy, keV	Spin and Parity	Half-life	Probability of β ⁻ transitions (×100)
0	0,0	5/2 ⁺	2,144 (7)×10 ⁶ a	-
1	33,196 29 (22)	7/2 ⁺	54 (24) ps	-
2	59,540 92 (10)	5/2 ⁻	67 (2) ns	6,7 (42)
3	75,899 (5)	9/2 ⁺	≈ 28 ps	-
4	102,959 (3)	7/2 ⁻	80 (40) ps	-
5	267,556 (12)	3/2 ⁻	5,2 (2) ns	40,9 (31)
6	281,356 (18)	1/2 ⁻	-	48,2 (25)
7	332,376 (16)	1/2 ⁺	≤ 1,0 ns	2,9 (9)
8	368,602 (20)	5/2 ⁺	-	-
9	370,928 (23)	3/2 ⁺	-	1,3 (9)

The probabilities of β^- transitions have been deduced from the $P(\gamma + ce)$ balance at each level of ^{237}Np .

The 459,1 keV $\beta^-_{0,2}$ transition probability of 7 (4) % has been obtained using the relation of $100 - \sum P_i(\beta^-)$. The value deduced from the $P(\gamma + ce)$ balance is 7 (6) %.

Some experimental estimations of the β^- transition energies and probabilities are given in 1949Me43, 1953Wa05 and 1957Ra04. More precise measurements would prove beneficial.

2.2 Gamma-ray Transitions and Internal Conversion Coefficients

The recommended energies of the gamma-ray transitions are mainly the same as the gamma-ray energies because nuclear recoil is negligible for ^{237}Np .

The gamma-ray transition probabilities have been obtained from the gamma-ray emission probabilities and the total internal conversion coefficients (ICCs). Multipolarities of gamma-ray transitions have been taken from 2006Ba41. The ICCs have been interpolated using the BrIcc package with the so called “*Frozen Orbital*” approximation (2008Ki07). The relative uncertainties of the ICC for pure multipolarities have been taken as 2 %.

The ICC for the intense E1 anomalously converted gamma-ray-transitions $\gamma_{2,1}$ (26,3-keV) and $\gamma_{2,0}$ (59,5-keV) have been obtained from a joint analysis of the gamma-ray and L-, M- conversion electron probabilities measured in ^{241}Am α decay and ^{237}U β^- decay (1996Jo28, 2006Ba41). The experimental conversion electron data are given in 1959Sa10, 1964Wo03, 1966Ko06, 1966Le13, 1966Ya05, and 1998Ko61. For discussion of E1 anomalously converted gamma transitions see 1960As02, 1966Ya05, 1967Pa23, 1970Gr36, and 1996Jo28.

The E2/M1 mixing ratio of 16,6 (25) % for $\gamma_{4,2}$ (43,4-keV) has been obtained by averaging the four measurement results from 1964Wo03 (17,6 (19) %), 1966Ko06 (13 (2) %), 1966Ya05 (11 (4) %), and 1998Ko61 (21,2 (22) %).

The E2/M1 mixing ratio of 15 (8) % for $\gamma_{9,7}$ (38,5-keV) has been deduced using the ratio $P_{ce}(L_2; \gamma_{9,7}) / P_{ce}(M_3; \gamma_{9,7}) = 10$ (5) from 1966Ya05 and the theoretical values from the BrIcc package. $P_{\gamma+ce}(\gamma_{9,8}, 2,3\text{-keV})$ has been deduced assuming that there is no β^- feeding to the 368,59-keV level.

$P_{\gamma+ce}(\gamma_{3,1}, 42,7\text{-keV})$ and $P_{\gamma+ce}(\gamma_{3,0}, 75,8\text{-keV})$ have been deduced from $P_{\gamma_{3,0}}/P_{\gamma_{3,1}} = 3/28$ (see 2006Ba41) assuming that there is no β^- feeding to the 75,92-keV level.

The gamma-ray transitions with energies 114,09 keV and 340,45 keV have not been placed in the level scheme.

3 Atomic Data

The atomic data are from Schönfeld and Janßen (1996Sc06).

4 Electron Emissions

The energies of the conversion electrons have been obtained from the gamma-ray transition energies and the electron binding energies.

The absolute emission probabilities of the conversion electrons have been calculated using recommended P_γ and ICC values.

The total absolute emission probabilities of K and L Auger electrons have been calculated using the EMISSION computer program.

β^- average energies have been calculated using the LOGFT computer program.

5. Photon Emissions

5.1 X-ray Emissions

The absolute emission probabilities of U KX and LX-rays have been calculated using the EMISSION computer program.

In Table 3 the calculated values are compared to the experimental data. The uncertainty in the detector efficiency (2 %) was added to the uncertainties listed in 1976GuZN.

Table 3. Experimental and recommended Np KX - ray emission probabilities in decay of ²³⁷U

	Energy (keV)	1966Ya05	1976GuZN	Recommended (calculated)
K α_2	97,069	16,2 (17)	15,8 (7)	14,8 (4)
K α_1	101,059	22,6 (24)	25,2 (9)	23,5 (6)
K β_1	113,944	9,8 (10)	9,22 (32)	8,57 (27)
K β_2	117,463	3,1 (4)	2,3 (5)	2,95 (10)

5.2 Gamma-rays emissions

The energies of gamma rays $\gamma_{2,1}$ (26,3-keV) and $\gamma_{2,0}$ (59,5-keV) are from 2000He14. $E_{\gamma_{1,0}}$ (33,2 keV) has been calculated as the difference $E_{\gamma_{2,0}} - E_{\gamma_{2,1}}$. The energies of gamma rays $\gamma_{4,3}$, $\gamma_{3,1}$, $\gamma_{4,2}$ have been taken from 1998Ko61. The rest gamma-ray energies have been adopted from 2006Ba41 based on experimental data of 1996Ya05, and 1976GuZN. Other measurements: 1957Ra04, 1963Ak04, 1968Da24, 1971Cl03. The uncertainty in the detector efficiency (2 %) was added to the uncertainties listed in 1976GuZN.

In Table 4 the experimental and evaluated absolute gamma ray emission probabilities (P_γ) are presented.

Table 4. Experimental and evaluated absolute gamma-ray emission probabilities (%) in decay of ²³⁷U.

E_γ , keV	1966Ya05	1971Cl03	1976GuZN	1982BuZF	1984BaYS	1985He02	1985Wi04	Evaluated
51,01	0,21 (10)		0,340 (14)		0,44 (6)			0,340 (14)
59,54	32,9 (40)	32,8 (25)	34,5 (8)		33,8 (9)			34,1 (9)
64,83	1,15 (16)	1,19 (9)	1,30 (3)		1,31 (5)		1,282 (17)	1,286 (17)
164,61	1,80 (9)	1,82 (14)	1,84 (5)		1,85 (5)	1,865 (23)	1,853 (23)	1,855 (23)
208,00			21,7 (5)	21,5 (14)		21,2 (3)	21,2 (3)	21,28 (30)
221,80	0,0199 (18)	0,0182 (14)	0,0212 (8)		0,0199 (25)			0,0204 (8)
234,40	0,0190 (18)	0,0273 (20)	0,0205 (8)		0,0224 (40)			0,0205 (8)
267,54	0,698 (30)	0,755 (20)	0,740 (18)		0,723 (25)	0,714 (22)	0,711 (10)	0,721 (10)
332,36	1,18 (8)	1,19 (9)	1,21 (3)		1,18 (4)		1,200 (16)	1,199 (16)
335,38	0,094 (9)	0,109 (9)	0,097 (3)		0,092 (5)		0,0951 (22)	0,0958 (22)
368,59	0,045 (4)	0,044 (3)	0,043 (2)		0,042 (3)		0,0392 (17)	0,0416 (17)
370,94	0,109 (9)	0,125 (10)	0,110 (4)		0,109 (6)		0,1073 (17)	0,109 (2)

The measurement results for gamma ray emission probabilities given in 1976GuZN, 1982BuZF, 1985He02, 1985Wi04 are absolute. The measurements results given in 1966Ya05, 1971Cl03, 1984BaYS are relative. The latter ones have been renormalized by evaluators at $P_\gamma(208 \text{ keV}) = 21,3 (3) \%$.

$P_{\gamma_{6,5}}$ has been deduced from $P_{ce}(M1) = 29,9 (3) \%$, as measured by 1966Ya05, and $ICC \alpha_{M1} = 281 (9)$.

$P_{\gamma_{4,1}}$ has been deduced from $P_{\gamma_{4,1}} / P_{\gamma_{4,2}} = 2,9 (4) / 73 (8)$, as measured in ^{241}Am α -decay (see 2006Ba41).

$P_{\gamma_{4,0}}$ has been deduced from $P_{\gamma_{4,0}} / P_{\gamma_{4,2}} = 19,5 (1) / 73 (8)$, as measured in ^{241}Am α -decay (see 2006Ba41).

$P_{\gamma_{8,2}}$ has been deduced from $P_{\gamma_{8,2}} / P_{\gamma_{8,1}} = 10,14 / 49,6$ as measured in ^{241}Am α -decay (see 2006Ba41).

$P_{\gamma_{8,3}}$ has been deduced from $P_{\gamma_{8,3}} / P_{\gamma_{5,2}} = 0,000 12 (3)$, as measured by 1966Ya05.

$P_{\gamma_{9,7}}$ has been deduced by evaluators from the ratio $P_{ce}(L_2; \gamma_{9,7}) / P_{ce}(K; \gamma_{5,2}) = 0,0056 (20)$ from 1966Ya05 and total ICC's.

$P_{\gamma}(340,4\text{-keV})$ has been adopted from 1976GuZN.

6. Consistency of Recommended Data

The most accurate Q value, $Q(M)$, is taken from the atomic mass adjustment table of Audi et al. (2003Au03). Comparison of $Q(\text{eff})$ (deduced as the sum of average energies per disintegration ($\sum E_i \times P_i$) for all emissions accompanying ^{237}U β^- decay) with the tabulated decay energy $Q(M)$ allows to check a consistency of the recommended decay-scheme parameters obtained in this evaluation.

Here E_i and P_i are the evaluated energies and emission probabilities of the i -th alpha particle, beta particle, gamma ray, X-ray, etc. Consistency (percentage deviation) is determined by $\{[Q(M) - Q(\text{eff})] / Q(M)\} \times 100$. "Percentage deviations above 5 % would be regarded as high and imply a poorly defined decay scheme; a value of less than 5 % indicates the construction of a reasonably consistent decay scheme" (quoted from the article by A.L. Nichols in Appl. Rad. Isotopes 55 (2001) 23-70).

For the above ^{237}U decay data evaluation we have $Q(M) = 518,6 (6) \text{ keV}$ and $Q(\text{eff}) = 519 (23) \text{ keV}$, i.e. consistency is not worse than 4,4 %.

7. References

- 1949Me43 L. Melander, H. Slatis, Arkiv Mat. Astron. Fysik 36A(1948)15 (Half-life, energies and probabilities of β -transitions).
- 1953Wa05 F. Wagner, Jr., M. S. Freedman, D. W. Engelkemeir, J. R. Huizenga, Phys. Rev. 89(1953)502 (Half-life, energies and probabilities of β -transitions).
- 1957Ra04 J. O. Rasmussen, F. L. Canavan, J. M. Hollander, Phys. Rev. 107(1957)141 (Energies and probabilities of β -transitions).
- 1958Ca16 M. J. Cabell, T. A. Eastwood, P. J. Champion, J. Nucl. Energy 7(1958)81 (Half-life).
- 1959Sa10 P. S. Samoïlov, Izvest, Akad. Nauk SSSR, Ser. Fiz. 23(1959)1416 (Gamma-ray multipolarities).
- 1960As02 F. Asaro, F. S. Stephens, J. M. Hollander, I. Perlman, Phys. Rev. 117(1960)492 (ICC for the anomalously converted gamma-transitions).
- 1963Ak04 E. Akatsu, T. Kuroyanagi, T. Ishimori, Radiochim. Acta 2(1963)1 (Gamma-ray energies).
- 1964Wo03 J. L. Wolfson, J. J. H. Park, Can. J. Phys. 42(1964)1387; Erratum Can. J. Phys. 48(1970)2782 (E2/M1 mixing ratios).
- 1966Ko06 L. N. Kondratev, E. F. Tretyakov, Bull. Acad. Sci. USSR, Phys. Ser. 30(1967)393 (E2/M1 mixing ratios).
- 1966Le13 C. M. Lederer, J. K. Poggenburg, F. Asaro, J. O. Rasmussen, I. Perlman, Nucl. Phys. 84(1966)481 (Conversion electron data).

- 1966Ya05 T. Yamazaki, J. M. Hollander, Nucl. Phys. 84(1966)505 (Gamma-ray and X-ray energies and multipolarities, E2 admixtures, relative probability of conversion electrons).
- 1967Pa23 H. -C. Pauli, K. Alder, Z. Physik 202(1967)255 (Anomalously converted E1 gamma-ray transitions).
- 1968Da24 R. Dams, F. Adams, Radiochim. Acta 10(1968)1 (Gamma-ray energies).
- 1970Gr36 V. N. Grigorev, A. P. Feresin, Yad. Fiz. 12(1970)665; Sov. J. Nucl. Phys. 12(1971)361 (Anomalously converted E1 gamma-ray transitions).
- 1971Cl03 J. E. Cline. IN-1448 Rev. (1971) (Gamma-ray energies and emission probabilities).
- 1976GuZN R. Gunnink, J. E. Evans and A. L. Prindle, UCRL-52139(1976) (Gamma-ray energies and emission probabilities).
- 1982BuZF A. V. Bushuev, O. V. Matveev, V. N. Ozerkov, V. V. Chachin, INDC(CCP)-193/G(1982)30 (Gamma-ray emission probabilities).
- 1984BaYS M. F. Banham, Priv. Comm. (1984), quoted by 1986LoZT (Gamma-ray emission probabilities).
- 1985He02 R. G. Helmer, C. W. Reich, Int. J. Appl. Radiat. Isotop. 36(1985)117 (Gamma-ray emission probabilities).
- 1985Wi04 H. Willmes, T. Ando, R. J. Gehrke, Int. J. Appl. Radiat. Isotop. 36(1985)123 (X-ray and gamma-ray emission probabilities, Gamma-ray emission probabilities).
- 1986LoZT A. Lorenz, IAEA Tech. Rept. Ser. 261(1986) (Gamma-ray emission probabilities).
- 1996Jo28 P. N. Johnston, Nucl. Instrum. Meth. Phys. Res. A369(1996)107 (ICC for the anomalously converted gamma-transitions).
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instr. Meth. Phys. Res. A369(1996)527 (Atomic data).
- 1996Ya05 R. Yanez, W. Loveland, D. J. Morrissey, K. Aleklett, J. O. Liljenzin, E. Hagebo, D. Jerrestam, L. Westerberg, Phys. Lett. 376B(1996)29 (Gamma-ray energies).
- 1998Ko61 A. Kovalik, E. A. Yakushev, V. M. Gorozhankin, A. F. Novgorodov, M. Rysavy, J. Phys.(London) G24(1998)2247 (Gamma-ray transition energies and multipolarities).
- 2000He14 R. G. Helmer, C. van der Leun, Nucl. Instrum. Meth. Phys. Res. A450(2000)35 (Gamma-ray energies).
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)337 (Q value).
- 2006Ba41 M. S. Basunia, Nucl. Data Sheets 107(2006)3323 (Decay data evaluation, gamma-ray energies and multipolarities, decay scheme).
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, and C. W. Nestor, Jr., Nucl. Instrum. Meth. Phys. Res. A589(2008)202 (Theoretical ICC).

²³⁷Np – Comments on evaluation of decay data
by V. P. Chechev and N.K. Kuzmenko

This evaluation was done originally in October 2007 and then updated in April 2009 with a literature cut-off by the same date. The Saisinuc software (2002Be) and associated supporting programs were used in assembling the data following the established protocol within DDEP.

1. DECAY SCHEME

Decay scheme is based on 2005Si15. It cannot be considered complete since the α -feedings measured directly in ²³⁷Np α -decay and those deduced from the level gamma-ray intensity balances are not always in good agreement as shown in Table 1 (see also 2005Si15).

Table 1. Comparison of the prominent α -feedings ($P_\alpha \times 100$) measured directly in ²³⁷Np α -decay with those deduced from the level gamma-ray intensity balances

Level	Level energy (keV)	$P_\alpha \times 100$ Adopted from measurements	$P_\alpha \times 100$ Deduced from γ -ray intensity balance
0	0		
1	6.654 (25)	}2.92 (4)	1 (3)
2	57.101 (14)	2.430 (17)	8 (4)
3	70.510 (25)	2.02 (2)	1.4 (3)
4	86.469 (9)		
6	103.636 (20)	}80.1 (5)	}79.1 (24)
7	109.04 (5)		
13	212.342 (18)	3.46 (3)	2.8 (9)
14	237.895 (13)	6.43 (3)	5.1 (7)

2. NUCLEAR DATA

$Q(\alpha)$ value is from 2003Au03.

The recommended half-life of ²³⁷Np is based on the experimental results given in Table 2.

Table 2. Experimental values of ²³⁷Np half-life (in 10⁶ years)

Reference	Author(s)	Value	Comments and method
1949Ma01	Magnusson and LaChapelle	2.20 (11)	First isolation of the element 93 and a determination of ²³⁷ Np half-life
1960Br12	Brauer et al.	2.14 (1)	Specific activity
1992Lo03	Lowles et al.	2.144 (7)	Specific activity, many sources, known geometry gas flow proportional counters for α -particle counting

Comments on evaluation

The weighted mean of the 3 values is 2.143 with the internal uncertainty of 0.0057 and external uncertainty of 0.0025 and $\chi^2/\nu = 0.19$. The unweighted mean is 2.161 (19). *

The recommended value of ^{237}Np half-life of $2.144 (7) \times 10^6$ years has been adopted from the most accurate measurement of 1992Lo03.

The recommended ^{237}Np spontaneous fission half-life $T_{1/2}(\text{SF}) \geq 1 \times 10^{18}$ years is from 1961Dr04. The theoretical values of $T_{1/2}(\text{SF})$ are about 10^{18} yr (1988Io05) and 10^{14} yr (1992Gr16).

2.1 Alpha Transitions

The energies of the alpha transitions have been deduced from the Q value and the level energies given in Table 3 from 2005Si15 where they were deduced from a least squares fit to gamma-ray energies. The energies of the gamma rays adopted from 2005Si15 are given below, in Table 7.

Table 3. ^{233}Pa levels populated in ^{237}Np α -decay

Level	Level energy (keV)	Spin and parity	Half-life	Energy of α -particles (keV)	Probability of alpha transition (%)
0	0	$3/2^-$	26.98 (2) d	4872.7 (14)	2.41 (3)
1	6.654 (25)	$1/2^-$		4866.4 (14)	0.51 (3)
2	57.101 (14)	$7/2^-$		4816.8 (10)	2.430 (17)
3	70.510 (25)	$5/2^-$		4803.5 (10)	2.02 (2)
4	86.469 (9)	$5/2^+$	35.8 (4) ns	4788.0 (9)	47.64 (6)
5	94.645 (16)	$3/2^+$			
6	103.636 (20)	$7/2^+$		4771.4 (8)	23.0 (3)
7	109.04 (5)	$9/2^+$		4766.5 (8)	9.5 (3)
8	133.2 (10)	$(11/2^+)$		4741.3 (20)	0.019
9	163.34 (10)	$(11/2^-)$		4712.3 (20)	
10	169.152 (20)	$1/2^+$		4708.3 (20)	1.174 (13)
11	179.1 (4)	$(9/2^-)$		4698.2 (8)	0.535 (10)
12	201.594 (19)	$3/2^+$		4676.4	0.38 (2)
13	212.342 (18)	$5/2^+$		4665.0 (9)	3.46 (3)
14	237.895 (13)	$5/2^+$		4640.0 (10)	6.43 (3)
15	257.1 (4)	$5/2^-$		4619.7 (21)	0.032 (8)
16	279.71 (3)	$(7/2^+)$		4599.1 (18)	0.37 (1)
17	300.48 (3)	$7/2^+$		4578.6 (14)	0.39 (2)
18	303.59 (7)	$(9/2^+)$		4573 (3)	0.048 (23)
19	306.05 (10)	$(7/2^+)$			
20	365.93 (8)	$9/2^+$		4515.1 (19)	0.038 (4)

The recommended α -transition probabilities have been obtained by averaging the experimental results (see Table 4). The probabilities of the $\alpha_{0,8}$ - and $\alpha_{0,12}$ - transitions have been deduced from the decay scheme. The α -decay hindrance factors have been calculated using the ALPHAD computer program from the ENSDF evaluation package with $r_0 = 1.517 (4)$ fm (see 2005Si15).

New α -transition with energy of 4550.5 (22) keV and intensity of 0.011 (3) % unplaced in ^{237}Np decay scheme was seen by 2002Wo03 (also 2000Si02).

Table 4. Experimental and recommended probabilities of α -transitions ($\times 100$) from ²³⁷Np α -decay

Level	Level energy (keV)	Energy of α -particles (keV)	1961Ba44	1969Br12	1990Bo44	2002Wo03	Recommended $P_\alpha \times 100$
0	0	4872.7 (14)	0.925	2.6 (2)	2.43 (3)	2.39 (4)	2.41 (3)
1	6.654 (25)	4866.4 (14)	0.24		0.49 (3)	0.53 (4)	0.51 (3)
2	57.101 (14)	4816.8 (10)		2.5 (4)	2.47 (2)	2.430 (17)	2.430 (17)
3	70.510 (25)	4803.5 (10)	2.014 (17)		2.06 (5)		2.014 (17)
4	86.469 (9)	4788.0 (9)		47 (9)	47.75 (20)	47.64 (6)	47.64 (6)
5	94.645 (16)						
6	103.636 (20)	4771.4 (8)		25 (6)	22.7 (4)	23.2 (3)	23.0 (3)
7	109.04 (5)	4766.5 (8)		8 (3)	9.7 (3)	9.3 (3)	9.5 (3)
8	133.2 (10)	4741.3 (20)					0.019
9	163.34 (10)	4712.3 (20)				< 1.17	< 1.17
10	169.152 (20)	4708.3 (20)				< 1.17	< 1.17
11	179.1 (4)	4698.2 (8)		0.48 (20)	0.54 (4)	0.535 (10)	0.535 (10)
12	201.594 (19)	4676.4					0.38 (2)
13	212.342 (18)	4665.0 (9)		3.32 (10)	3.43 (4)	3.478 (24)	3.46 (3)
14	237.895 (13)	4640.0 (10)		6.18 (12)	6.45 (4)	6.43 (3)	6.43 (3)
15	257.1 (4)	4619.7 (21)				0.032 (8)	0.032 (8)
16	279.71 (3)	4599.1 (18)		0.34 (4)	0.39 (2)	0.371 (9)	0.373 (9)
17	300.48 (3)	4578.6 (14)		0.40 (4)	0.41 (2)	0.369 (23)	0.393 (23)
18	303.59 (7)						
19	306.05 (10)	4573 (3)	0.048 (23)				0.048 (23)
		4550.5 (22)				0.011 (3)	0.011 (3)
20	365.93 (8)	4515.1 (19)		0.04 (2)	0.041 (4)	0.035 (4)	0.038 (4)

2.2. Gamma Transitions and Internal Conversion Coefficients

The energies of the gamma-ray transitions are virtually the same as the gamma-ray energies because nuclear recoil is negligible.

The gamma-ray transition probabilities have been deduced from their gamma-ray emission probabilities and total internal conversion coefficients (ICCs) deduced with a computer program supplied with the Saisinuc software (2002Be). The ICCs have been interpolated using the BrIcc package with the so called “Frozen Orbital” approximation (2008Ki07, see also 2002Ba85). The multipolarities and admixture coefficients δ have been taken from 2005Si15. The uncertainties in the ICCs for pure multipolarities have been taken as 2 %.

ICCs for the anomalously converted E1 gamma-ray transition $\gamma_{4,0}$ (86.477 keV) have been adopted from 1988Wo01 (see also 1960As02 and 1969Br12).

The conversion electron data of 1988Wo01 indicate that the gamma-transition $\gamma_{4,2}$ (29.374 keV) may be an anomalous E1. However the evaluators have been adopted the theoretical ICCs since the detector efficiency was not completely reliable for such energy as pointed out in 1988Wo01.

3. Atomic Data

The atomic data (fluorescence yields, X-ray energies and relative probabilities, and Auger electrons energies and relative probabilities) are from Schönfeld and Janßen (1996Sc06).

4. Alpha Emissions

The alpha particle energies have been taken from 2002Wo03 (see also 2000Si02). They are somewhat different (in limits of uncertainties) from those obtained from alpha transition energies taking into account nuclear recoil for ²³³Pa.

Details of alpha transition probability evaluation are given in Section 2.1.

5. Photon Emissions

5.1. X-Ray Emissions

The absolute X-ray emission probabilities (per 100 disintegrations) have been evaluated using the experimental data, see Tables 5, 6.

Table 5. Experimental and recommended absolute Pa KX- ray emission probabilities (×100)

	1984Va27	2000Sc04	2002Lu01	2004Sh07	2008De10	Recommended
Kα ₂	1.90 (10)	1.82 (5)	1.80 (20)	1.80 (3)	1.813 (20)	1.813 (20)
Kα ₁	3.00 (15)	2.98 (7)	2.89 (2)	2.89 (4)	2.932 (30)	2.906 (20)
Kβ ₁	1.03 (5)	0.86 (2)	1.06 (2)	1.02 (4)	1.154 (14)	1.06 (10)
Kβ ₂	0.35 (2)		0.373 (10)	0.38 (2)	0.380 (9)	0.380 (9)

Table 6. Experimental and recommended absolute Pa LX- ray emission probabilities (×100)

	2000Sc04	2004Sh07	2008De10	Recommended
Ll	1.55 (8)	1.31 (20)	1.33 (27)	1.32 (8)
Lα	26 (3)	23.3 (24)	23.1 (47)	24.0 (24)
Lβ	29.5 (20) ^a	24.3 (31) ^b	28 (6)	28.0 (20)
Lη	0.64 (6)	0.50 (4)		0.54 (4)
Lγ	5.8 (4) ^c	5.4 (8) ^d	7.8 (16)	5.8 (4)

^a Obtained by the evaluators from the sum absolute intensity (Pa Lβ + U Lβ) of 47.5 (19) % using the intensities of Pa Lβ-components measured in 2000Sc04 and the U Lβ-intensity of 18.0 (6) % from ²³³Pa decay data evaluation (2006Ch39) revised in April 2009.

^b Obtained by the evaluators from the sum absolute intensity (Pa Lβ + U Lβ) of 42.3 (30) % using the intensities of Pa Lβ-components measured in 2000Sc04 and the U Lβ-intensity of 18.0 (6) % from ²³³Pa decay data evaluation (2006Ch39) revised in April 2009.

^c Obtained by the evaluators from the sum absolute intensity (Pa Lγ + U Lγ) of 10.0 (4) % using the intensities of Pa Lγ-components measured in 2000Sc04 and the U Lβ-intensity of 4.18 (13) % from ²³³Pa decay data evaluation (2006Ch39) revised in April 2009.

^d Obtained by the evaluators from the sum absolute intensity (Pa Lγ + U Lγ) of 9.6 (8) % using the intensities of Pa Lγ-components measured in 2000Sc04 and the U Lβ-intensity of 4.18 (13) % from ²³³Pa decay data evaluation (2006Ch39) revised in April 2009.

5.2. Gamma-Ray Emissions

Energies

The gamma-ray energies have been adopted from 2005Si15. The gamma ray energy for γ_{7,6} (5.18 keV) has been adopted from 1990Lo04. The energies for γ_{1,0} (6.65 keV), γ_{5,4} (8.22 keV) and γ_{7,4} (17.4 keV) are from ²³³Th decay. For γ_{13,12} (10.7 keV) and γ_{8,7} (21.4 keV) the energies are from 1979Go12. The gamma-ray energies of γ_{6,5} (9.0 keV) and γ_{7,4} (22.6 keV) have been deduced from ²³³Pa level scheme (details of information of these and other gamma-ray transitions see in 2005Si15). Table 7 contains the experimental and adopted energies of the remaining gamma rays.

Table 7. Experimental and adopted energies (in keV) of gamma rays from ²³⁷Np decay

1969Br12	1969HoXY	1971Cl03	1974HeYW	1976Sk01	1979Go12	1988Wo01 (Ge- detector)	1988Wo01 (LEPS- detector)	Adopted
29.29 (10)	29.30 (5)	29.38 (2)	29.375 (20)	29.373 (10)	29.374 (20)	29.5 (17)	29.18 (21)	29.374 (20)
46.46 (10)	46.6 (1)	-	46.60 (10)	46.53 (4)	46.53 (6)	46.7 (11)	46.28 (18)	46.53 (6)
57.15 (10)	57.1 (1)	57.11 (2)	57.112 (20)	57.15 (4)	57.104 (20)	57.15 (80)	56.88 (17)	57.104 (20)
	62.9	-	62.5 (5)					62.59 (10)
			63.92 (8)					63.90 (10)
	71.0		70.75 (10)					70.49 (10)
86.49 (10)	86.40 (5)	86.49 (2)	86.486 (10)	86.503 (20)	86.477 (10)	86.50 (48)	86.26 (14)	86.477 (10)
		-		88.04 (16)				87.99 (3)
			94.66 (10)	94.66 (5)				94.64 (5)
106.22 (10)	106.30 (8)	106.30 (20)	106.15 (25)	106.12 (5)		106.17 (48)		106.15 (25)
				108.6				108.7
				115.45 (20)	115.40 (35)			115.40 (35)
117.65 (7)	117.5 (1)	117.72 (2)	117.718 (20)	117.681 (30)	117.702 (20)	117.72 (50)	117.41 (15)	117.702 (20)
131.11 (7)	131.2 (1)	131.11 (2)	131.11 (2)	131.11 (7)	131.101 (25)	131.09 (52)	130.62 (15)	131.101 (25)
134.23 (7)	134.4 (1)	134.28 (2)	134.28 (3)	134.23 (4)	134.285 (20)	134.27 (53)		134.285 (20)
				140.60 (10)	-			139.9 (1)
143.26 (7)	143.35 (5)	143.25 (1)	143.254 (10)	143.208 (25)	143.249 (20)	143.27 (56)	142.96 (16)	143.249 (20)
151.31 (7)	151.5 (1)	151.41 (1)	151.410 (15)	151.37 (4)	151.414 (20)	151.42 (60)	151.06 (17)	151.414 (20)
				153.52				153.37 (10)
155.20 (7)	155.4 (1)	155.25 (2)	155.25 (2)	155.22 (4)	155.239 (20)	155.28 (63)		155.239 (20)
162.38 (7)	162.7 (1)	162.52 (3)	162.52 (3)	162.50 (6)	162.41 (8)	162.45 (68)		162.41 (8)
169.09 (7)	169.4 (1)	169.16 (3)	169.16 (3)	169.17 (5)	169.156 (20)	169.18 (73)		169.156 (20)
170.56 (10)	171.2 (3)	170.64 (5)	170.64 (5)	170.63 (8)	170.59 (6)			170.59 (6)
175.93 (10)	176.1 (1)	176.06 (5)	176.06 (5)	176.09 (7)	176.12 (6)	176.17 (80)		176.12 (6)
180.66 (10)	180.8 (1)	180.78 (5)	180.78 (5)	180.80 (8)	180.81 (10)	180.87 (85)		180.81 (10)
186.86 (30)				186.8 (5)	186.86 (35)			186.86 (35)
191.34 (10)		191.42 (3)	191.42 (3)	191.45 (6)	191.46 (5)	191.46 (97)		191.46 (5)
193.05 (10)		193.22 (3)	193.22 (3)	193.26 (4)	193.26 (5)	193.24 (98)		193.26 (5)
				194.67 (20)				194.67 (20)
194.91 (7)	195.00 (5)	194.97 (2)	194.97 (2)	195.096 (20)	194.95 (3)	195.1 (10)		194.95 (3)
196.81 (10)	-	196.80 (10)	196.80 (10)	196.84 (6)	196.86 (5)	196.9 (10)		196.86 (5)
			199.9 (1)	200.17 (10)	199.95 (6)			199.95 (6)
201.68 (8)	201.75 (10)	201.67 (20)	201.670 (25)	201.72 (5)	201.62 (5)	201.8 (11)		201.62 (5)
			202.9 (2)	202.69 (25)				202.9 (2)
209.07 (8)	209.1 (2)	209.18 (3)	209.18 (3)	209.25 (5)	209.19 (5)	209.2 (12)		209.19 (5)
212.28 (7)	212.4 (1)	212.33 (2)	212.33 (2)	212.42 (5)	212.29 (5)	212.4 (12)		212.29 (5)
213.92 (10)	-	213.96 (4)	213.96 (4)	214.09 (5)	214.01 (5)	214.1 (12)		214.01 (5)
				222.52 (25)				222.6 (2)
229.84 (10)	229.9 (1)	229.90 (10)	229.90 (10)	230.01 (10)	229.94 (5)			229.94 (5)
237.91 (7)	238.2 (1)	237.91 (2)	237.908 (10)	238.04 (4)	237.862 (60)	238.0 (14)		237.86 (2)
248.6 (4)	248.8 (1)	248.8 (5)	248.8 (5)	248.9 (1)	248.95 (10)			248.95 (10)
257.14 (40)	257.3 (2)	257.15 (50)	257.15 (50)	257.20 (20)	257.09 (20)			257.09 (20)
262.48 (40)	262.6 (2)	262.42 (50)	262.42 (50)	262.44 (15)	262.44 (20)			262.44 (20)

Emission Probabilities

The value P_{γ_{14,12}} (36.32 keV) of 0.000 05 (1) has been adopted from 1990Lo04. The values P_{γ_{-1,1}} (21.5 keV) of 0.003 56 (13) and P_{γ_{-1,2}} (27.7 keV) of 0.008 4 (7) have been adopted from 2004Sh07. The values P_{γ_{17,14}} (62.59 keV) of 0.000 06 (2), P_{γ_{3,1}} (63.9 keV) of 0.000 108 (4) and P_{γ_{10,5}} (74.54 keV) of 0.000 12 (3) have been adopted from 1981Ba68. The value P_{γ_{9,2}} (106.15 keV) of 0.000 49 (1) has been adopted from 2002Lu01. For absolute gamma-ray emission probabilities see 1981Ba68, 1984Va27, 2000Sc04, 2000Wo01, 2002Wo03, 2004Sh07. The remaining relative emission probabilities are listed in Table 9. These have been renormalized by the evaluators to P_γ (86.48 keV) = 12.26 (12) % obtained as a

Comments on evaluation

weighted average of 1984Banham, 1984Va27, 2000Sc04, 2000Wo01, 2002Wo03, 2002Lu01, 2004Sh07, 2008De10.

There are significant unexplained (as stated in 2002Wo03) discrepancies in the intensities of several gamma rays with the following energies: 29.4, 46.5, 88.0, 117.7, 169.2, 193.3, 195.0, 257.1 and 279.6 keV.

The value of $P_{\gamma_{4,0}}$ (86.48 keV) used for normalization of the decay scheme is itself discrepant since this gamma ray and the gamma ray with the energy 86.6 keV from the decay of its daughter ^{233}Pa become apparent as a complex peak, and the separated intensities in various studies are not always in good agreement. Table 8 contains the experimental and evaluated values of the absolute emission probability of gamma ray $\gamma_{4,0}$ (86.48 keV). The results of 2000Sc04, 2002Lu01 and 2004Sh07 given in Table 8 have been corrected taking into account the intensity of gamma ray with the energy 86.6 keV from the decay of ^{233}Pa : $P_{\gamma} (^{233}\text{Pa}, 86.6 \text{ keV}) = 1.99 (11) \%$, see 2006Ch39.

Table 8. Experimental and recommended values of the 86.48 keV γ ray emission probability

1984Banham	1984Va27	2000Wo01 2002Wo03	2000Sc04	2002Lu01	2004Sh07	2008De10	Recommended
12.20 (12)	12.44 (33)	12.86 (21)	12.1 (3)	12.02 (12) [#]	11.6 (5)	12.38 (13)	12.26 (12)

[#] Although the $P_{\gamma_{4,0}}$ (86.48 keV) = 11.40 (24) % is given in 2002Lu01, the evaluators used more accurate value of 14.01 (6) % measured in 2002Lu01 for P_{γ} (86.48+86.6 from ^{233}Pa decay) to deduce $P_{\gamma_{4,0}}$ (86.48 keV) = 12.02 (12) %.

The recommended values of the gamma ray emission probabilities given in Table 9 have been obtained by averaging experimental data using the LWEIGHT computer program. The uncertainty assigned in this evaluation to the recommended value is always greater than or equal to the smallest uncertainty in any of the experimental values used in the statistical processing.

The systematic uncertainties (1 %) of U KX-ray emission probability from 2008De10 have been added to statistic uncertainties measured in 2008De10.

The systematic uncertainties (20 %) of U LX-ray emission probability from 2008De10 have been added to statistic uncertainties measured in 2008De10.

Table 9 (part 1). Experimental and recommended emission probabilities of gamma rays in ^{237}Np decay

E_{γ}	1969Br12	1976Sk01	1979Go12	1981Ba68 1984Banham	1984Va27	1988Wo01 (Ge- detector)	1988Wo01 (LEPS- detector)
29.37	13.7 (20)	16.2 (9)	10.1 (10)	15.4 (2)	15.03 (40)	-	19.2 (9)
46.53	0.137 (20)	0.12 (2)	0.10 (1)	0.104 (6)	0.10 (1)	0.12 (1)	0.14 (2)
57.10	0.412 (38)	0.433 (25)	0.37 (4)	0.373 (11)	0.39 (1)	0.34 (1)	0.43 (3)
62.6		0.012		0.006 (2)			
63.9				0.0108 (4)			
86.48	12.6	12.3	12.3	12.20 (12)	12.44 (33)	12.3	12.3
87.99	0.157 (20)	0.14 (4)	0.12 (1)	0.138 (3)	0.14 (1)	-	-
94.64		0.62 (4)	0.54 (5)				
106.15		0.044 (9)	0.05 (5)				
108.7							
115.40		0.26 (8)					
117.70	0.167 (20)	0.180 (12)	0.148 (15)	0.175 (2)	0.168 (5)	0.16 (7)	0.15 (2)
131.1	0.087 (9)	0.10 (1)	0.079 (8)	0.086 (1)	-	0.091 (5)	0.09 (2)
134.28	0.069 (8)	0.081 (16)	0.062 (6)	0.071 (1)	-	0.080 (5)	
143.25	0.412 (40)	0.462 (28)	0.40 (4)	0.430 (4)	0.434 (10)	0.43 (1)	0.42 (3)
151.41	0.244 (30)	0.249 (16)	0.223 (23)	0.236 (2)	0.232 (6)	0.248 (7)	0.20 (3)
153.4		0.007 (2)					

Comments on evaluation

155.24	0.095 (9)	0.097 (7)	0.085 (9)	0.0917 (10)	-	0.086 (6)	-
162.4		0.041 (7)	0.027 (4)			0.032 (4)	-
169.16	0.074 (8)	0.082 (9)	0.072 (7)	0.0711 (7)	-	0.057 (4)	-
170.59		0.016 (2)	0.024 (5)				
176.12		0.017 (3)				0.02 (4)	-
180.8		0.022 (5)	0.021 (4)			0.015 (2)	-
186.86		0.003 (3)					
191.46		0.017 (3)	0.026 (5)			0.014 (5)	-
193.26		0.043 (4)	0.05 (5)			0.049 (3)	-
194.67		0.05 (2)					
194.95	0.206 (20)	0.169 (21)	0.16 (2)	0.184 (2)	0.188 (5)	0.191 (6)	-
196.86		0.023 (3)	0.019 (4)			0.021 (2)	-
201.6		0.044 (5)	0.044 (5)			0.041 (4)	-
209.2		0.019 (2)	0.016 (3)			0.010 (2)	-
212.3	0.157 (20)	0.166 (11)	0.157 (16)	0.150 (2)	0.155 (5)	0.156 (4)	-
214.0		0.047 (4)	0.06 (4)			0.034 (1)	
222.6		0.002 (2)					
229.94		0.011 (3)	0.018 (4)				
237.86	0.067 (6)	0.075 (9)	0.062 (7)	0.0586 (12)	-	0.059 (3)	-
248.95		0.005 (2)	0.05 (1)				
257.09		0.007 (3)	0.019 (6)				
262.44		0.008 (2)	0.007 (1)				
279.65		0.002 (2)	0.011 (4)				
288.3							

Table 9 (part 2). Experimental and recommended emission probabilities of gamma rays in ²³⁷Np decay

E _γ	1990Lo04	2000Sc04	2000Wo01	2002Lu01	2004Sh07	2008De10	Recommended
29.37	13.7 (1)	14.1 (15)	13.2 (4)	13.51 (16)	13.15 (36)	15.08 (16)	14.3 (6)
46.53	0.112 (1)	0.104 (4)	0.1067 (19)	0.163 (5)	0.100 (13)	0.114 (3)	0.109 (4)
57.10	0.360 (2)	0.354 (8)	0.360 (5)	0.366 (3)	0.356 (16)	0.458 (6)	0.381 (21)
62.6							0.006 (2)
63.9	0.0090 (9)						0.0107 (4)
86.48	12.3	14.1 (3)&	12.86 (21)	14.01 (6) &	13.6 (5)&	12.38 (13)&	12.26 (12)
87.99	0.143 (1)			0.167 (4)	0.134 (13)	0.144 (5)	0.143 (3)
94.64				0.615 (23)	0.575 (19)	0.730 (10)	0.66 (7)
106.15	0.048 (1)				0.0509 (26)	0.0573 (28)	0.0509 (29)
108.7		0.0864 (19)		0.070 (3)	0.0723 (36)		0.071 (3)
115.40	0.47 (11)*	0.332 (10)*					0.0026 (8)#
117.70	0.168 (1)	0.169 (4)	0.188 (3)	0.184 (12)	0.169 (17)	0.1660 (29)	0.171 (4)
131.1	0.079 (1)	0.0857 (22)		0.088 (3)	0.075 (5)		0.084 (5)
134.28	0.064 (1)	0.0670 (28)		0.075 (3)	0.073 (6)		0.069 (5)
143.25	0.387 (2)	0.443 (8)	0.439 (5)	0.428 (3)	0.394 (24)	0.423 (6)	0.42 (4)
151.41		0.232 (24)	0.228 (3)	0.244 (3)	0.223 (14)	0.234 (4)	0.234 (2)
153.4							0.007 (2)
155.24	0.080 (1)	0.0889 (18)		0.091 (6)	0.087 (6)		0.088 (8)
162.4		0.0327 (12)					0.033 (1)
169.16		0.0633 (19)		0.092 (11)			0.0672 (3)
170.59							0.020 (4)
176.12		0.012 (4)					0.015 (3)
180.8		0.0158 (10)					0.016 (1)
186.86							0.003 (3)
191.46		0.0192 (12)		0.015 (4)	0.023 (5)		0.019 (1)
193.26		0.0437 (10)		0.030 (5)	0.041 (8)		0.044 (1)
194.67	0.033 (1)			0.033 (8)	0.03 (1)		0.033 (1)

194.95	0.156 (2)	0.177 (5)	0.161 (4)	0.164 (7)	0.161 (34)		0.174 (20)
196.86		0.0208 (12)		0.024 (5)	0.020 (4)		0.0208 (1)
201.6		0.0393 (9)					
209.2		0.0142 (9)		0.019 (2)	< 0.02		0.0150 (15)
212.3		0.151 (3)	0.148 (3)	0.150 (4)			0.17 (1)
214.0	0.132 (2)	0.0362 (8)		0.039 (2)			0.037 (2)
222.6							0.002 (2)
229.94							0.014 (3)
237.86		0.0569 (6)		0.056 (3)	0.067 (4)		0.0573 (6)
248.95		0.0050 (14)		0.006 (3)			0.005 (1)
257.09							0.02 (1)
262.44		0.00471 (18)					0.0048 (2)
279.65		0.0109 (4)					0.0108 (4)
288.3		0.0164 (5)					0.0162 (5)

* Sum intensity of $\gamma_{12,14}$ and KX(Pa)

Adopted from 2005Si15

& Measured $P_{\gamma 86.48+86.6}$ keV from ²³³Pa decay)

6. Electron Emissions

The energies of the conversion electrons have been obtained from the gamma transition energies and the electron binding energies. The emission probabilities of conversion electrons have been deduced from the evaluated P(γ) and ICC values.

The number of K- and L- Auger electrons per 100 disintegrations has been deduced using the evaluated XK- and XL- emission probabilities.

7. Consistency of recommended data

The most accurate Q value, Q(M), is taken from the atomic mass adjustment table of Audi et al. (2003Au03). Comparison of Q(eff) (deduced as the sum of average energies per disintegration ($\sum E_i \times P_i$) for all emissions accompanying ²³⁷Np α - decay) with the tabulated decay energy Q(M) allows to check a consistency of the recommended decay-scheme parameters obtained in this evaluation.

Here E_i and P_i are the evaluated energies and emission probabilities of the i-th alpha particle, beta particle, gamma ray, X-ray, etc. Consistency (percentage deviation) is determined by $\{[Q(M) - Q(\text{eff})] / Q(M)\} \times 100$. "Percentage deviations above 5 % would be regarded as high and imply a poorly defined decay scheme; a value of less than 5 % indicates the construction of a reasonably consistent decay scheme" (quoted from the article by A.L. Nichols in Appl. Rad. Isotopes 55 (2001) 23-70).

For the above ²³⁷Np decay data evaluation we have Q(M) = 4958.3 (12) keV and Q(eff) = 4966 (21) keV, i.e. consistency is not superior, but better than 1 %.

8. References

- 1949Ma01 L. Magnusson and T. LaChapelle, NNES 14B(1949)39 (Half-life).
- 1960Br12 F. P. Brauer, R. W. Stromatt, J. D. Ludwick, F. P. Roberts, W. L. Lyon, J. Inorg. Nuclear Chem. 12(1960)234 (Half-life).
- 1961Dr04 V. A. Druin, V. P. Perelygin and G. I. Khlebnikov, Zhur. Eksptl. i Teoret. Fiz. 40(1961) 1296; Soviet Phys. JETP 13(1961)913 (Spontaneous fission half-life).
- 1960As02 F. Asaro, F. S. Stephens, J. M. Hollander and I. Perlman, Phys.Rev. 117(1960)492 (Gamma-ray energies and emission probabilities, ICC for the 86.5 keV gamma-ray).
- 1961Ba44 S. A. Baranov, V. M. Kulakov, P. S. Samoilov, A. G. Zelenkov and Y. F. Rodionov, Zhur. Eksptl. i Teoret. Fiz., 41(1961)1733; Soviet Phys. JETP 14(1962)1232 (α -transition probabilities).

Comments on evaluation

- 1968Br12 E. Browne and F. Asaro, Priv. Comm. (October 1969). Quoted in 1968Br25 (α -transition energies and probabilities, gamma-ray emission probabilities).
- 1968Br25 E. Browne and F. Asaro, Report UCRL-17989, p. 1(1968) (α -transition energies and probabilities, gamma-ray emission probabilities).
- 1969HoXY W. Hoekstra, Thesis, Technische Hogeschool, Delft (1969) (Gamma-ray energies).
- 1971Cl03 J. E. Cline, IN-1448 Rev. (1971) (Gamma-ray energies).
- 1974HeYW R. L. Heath, ANCR-1000-2 (1974) (Gamma-ray energies).
- 1976Sk01 M. Skalsey and R. D. Connor, Can. J. Phys. 54(1976)1409 (Gamma-ray energies and emission probabilities).
- 1979Go12 L. Gonzalez, R. Gaeta, E. Vano and J. M. Los Arcos, Nucl. Phys. A324(1979)126 (Gamma-ray energies and probabilities).
- 1981Ba68 M. F. Banham and A. J. Fudge, J. Radioanal. Chem. 64(1981)167 (Gamma-ray probabilities).
- 1984Va27 R. Vaninbroukx, G. Bortels and B. Denecke, Int. J. Appl. Radiat. Isotop. 35(1984)905 (X- and gamma- ray emission probabilities).
- 1984BaYS M. F. Banham, Priv. Comm. (1984). Quoted in 1986LoZT (Gamma-ray probabilities).
- 1986LoZT A. Lorentz, Techn. Report Ser. No. 261, IAEA, Vienna (1986) (Gamma-ray probabilities).
- 1988Io05 D. B. Ion, R. Ion-Mihai and M. Ivascu, Rev. Roum. Phys. 33(1988)1075 (Spontaneous fission half-life).
- 1988Wo01 S. A. Woods, P. Christmas, P. Cross, S. M. Judge and W. Gelletly, Nucl. Instrum. Methods Phys. Res. A264(1988)333; Addendum Nucl. Instrum. Methods Phys. Res. A272(1988)924 (Gamma ray energies and emission probabilities, ICC for the 86,5 keV gamma-ray).
- 1990Bo44 G. Bortels, D. Mouchel, R. Eykens, E. Garcia-Torano, M. L. Acena, R. A. P. Wiltshire, M. King, A. J. Fudge and P. Burger, Nucl. Instrum. Methods Phys. Res. A295(1990)199 (α -transition probabilities).
- 1990Lo04 I. M. Lowles, T. D. Mac Mahon, M. F. Banham, A. J. Fudge and R. A. P. Wiltshire, Nucl. Instrum. Meth. Phys. Res. A286(1990)556 (Gamma-ray energies and probabilities).
- 1992Gr16 A. F. Grashin and A. D. Efimenko, Bull. Rus. Acad. Sci. Phys. 56(1992)66 (Spontaneous fission half-life).
- 1992Lo03 I. M. Lowles, T. D. Mac Mahon, R. A. P. Wiltshire, D. Crossley and A. J. Fudge, Nucl. Instrum. Meth. Phys. Res. A312(1992)339 (Half-life).
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 (Atomic data).
- 2000Sc04 U. Schötzgig, E. Schönfeld and H. Janßen, Appl. Radiat. Isot. 52(2000)883 (X- and gamma-ray emission probabilities).
- 2000Schönfeld E. Schönfeld, H. Janßen, Nucl. Instr. Meth. Phys. Res. A369(2000)527 (EMISSION computer code).
- 2000Si02 G. Sibbens and B. Denecke, Appl. Radiat. Isot. 52(2000)467 (α -transition probabilities, gamma-ray energies).
- 2000Wo01 S. A. Woods, D. H. Woods, P. de Lavison, S. M. Jerome, J. L. Makepeace, M. J. Woods, L. J. Husband and S. Lineham, Appl. Radiat. Isot. 52(2000)475 (Gamma-ray emission probabilities).
- 2002Ba85 I. M. Band, M. B. Trzhaskovskaya, C. W. Nestor, P. O. Tikkanen and S. Raman, Atom. Data and Nucl. Data Tables 91(2002)1 (Theoretical internal conversion coefficients).
- 2002Be M.M. Bé, R. Helmer, V. Chisté, J. Nucl. Sci. Tech., suppl.2(2002)481 (Saisinuc software).
- 2002Lu01 A. Luca, S. Sepman, K. Iakovlev, G. Shchukin, M. Etcheverry and J. Morel, Appl. Radiat. Isot. 56(2002)173 (KX - ray and gamma-ray emission probabilities).
- 2002Wo03 M. J. Woods, D. H. Woods, S. A. Woods, L. J. Husband, S. M. Jerome, C. Michotte, G. Ratel, M. Crespo, E. Garcia-Torano, L. Rodriguez, A. Luca, B. Denecke, G. Sibbens, J. Morel, M. Etcheverry, D. Santry, H. Janßen, E. Schönfeld and U. Schötzgig, Appl. Radiat. Isot. 56(2002)415 (α -particle energies and α -transition probabilities and X-, gamma- ray emission probabilities).
- 2003Au03 G. Audi, A. H. Wapstra and C. Thibault, Nucl. Phys. A729(2003)337 (Q value).
- 2004Sh07 G. Shchukin, K. Iakovlev and J. Morel, Appl. Radiat. Isot. 60(2004)239 (X- and gamma- ray emission probabilities).
- 2005Si15 B. Singh and K. Tuli, Nucl. Data Sheets 105(2005)109 (Decay scheme, gamma-ray multipolarities, admixture coefficients)

Comments on evaluation

- 2006Ch39 V. P. Chechev and N. K. Kuzmenko, Appl. Radiat. Isot. 64(2006)1403 (Gamma-ray emission probabilities in ^{233}Pa decay).
- 2008De10 D. J. deVries and H. C. Griffin, Appl. Rad. Isotop., 66(2008)1999 (Gamma-ray, KX-ray and LX-ray emission probabilities, and uncertainties of gamma-ray, KX-ray and LX-ray absolute emission probabilities).
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, and C. W. Nestor, Jr., Nucl. Instrum. Methods Phys. Res. A589(2008)202 (Theoretical ICC).

²³⁸U - Comments on evaluation of decay data by V. Chisté and M.M. Bé

This evaluation was completed in January 2006, and the literature available at this date has been included here.

1 Decay Scheme

²³⁸U disintegrates by alpha emission to two excited levels and to the ground state of ²³⁴Th. Spin and half-lives of excited states are from the mass-chain evaluation of Y.A. Akovali (1983E111 and 1994Ak05 for A = 234) and F.E. Chukreev (2002Ch52 for A = 238).

2 Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2003Au03).

Experimental ²³⁸U half-life values (in years x 10⁹) are given in Table 1:

Table 1: Experimental values of ²³⁸U half-life.

Reference	Original value (10 ⁹ a)	Revised Value by Schön (2004Sc03)	Comments
Kovarik (1932Ko01)	4.52		Not used. Natural U.
Schiedt (1935Schiedt)	4.42 (3)	4.46 (3) (a) 4.41 (5) (b)	Not used. Natural U. Corrected for ²³⁵ U. (a) ²³⁴ U and ²³⁸ U assumed to be in equilibrium. (b) ²³⁴ U and ²³⁸ U assumed to be not in equilibrium.
Curtis (1941Curtis)	4.514 (9)		Not used. Natural U. Lacking details.
Kienberger (1949Ki26)	4.490 (10)	4.495 (18)	Not used. Enriched U.
Kovarik (1955Ko13)	4.507 (9)	4.51 (2) (a) 4.46 (5) (b)	Not used. Natural U. (a) ²³⁴ U and ²³⁸ U assumed to be in equilibrium. (b) ²³⁴ U and ²³⁸ U assumed to be not in equilibrium.
Lechman (1957Le21)	4.56 (3)		Not used. Enriched U.
Steyn (1959St45)	4.460 (10)	4.457 (4) (a) 4.41 (4) (b)	Not used. Natural U. (a) ²³⁴ U and ²³⁸ U assumed to be in equilibrium. (b) ²³⁴ U and ²³⁸ U assumed to be not in equilibrium.
Jaffey (1971Ja07)	4.4683 (24)	4.468 (5)	Highly enriched U.
Recommended value		4.468 (5)	

The evaluators have chosen to follow the recommendations given by R. Schön (2004Sc03), who studied in detail various problems with the measurements of the half-life of ²³⁸U. So, the recommended value is the half-life obtained by Jaffey (1971Ja07), but its original uncertainty was multiplied by 2 (as suggested by Schön (2004Sc03)) in order to take into account the systematic uncertainties which were not considered by 1971Ja07.

Experimental ²³⁴Th half-life values (in days) are given in Table 2:

Table 2: Experimental values of ^{234}Th half-life.

Reference	Value (d)	Uncertainty (d)
M. Curie (1931Cu01)	24.5	
B.W. Sargent (1939Sa11)	24.1	0.2
G.B. Knight (1948Kn23)	24.101	0.025
Recommended value is (from 1994Ak05)	24.10	0.03

The recommended value is $24.10 d$ with an uncertainty of $0.03 d$, from Y. A. Akovali (1994Ak05).

The evaluated spontaneous fission partial half-life of ^{238}U is based on the experimental results given in Table 3.

Table 3: Experimental values of spontaneous fission decay rate of ^{238}U (λ^{238} , in 10^{-17} years $^{-1}$).

Reference	Value	Uncertainty	Comments by Holden (2000Ho27)
W.J. Withehouse (1950Whitehouse)	8.38	0.52	Ionization chamber.
E. Sègres (1952Se67)	8.60	0.29	Ionization chamber.
R.L. Fleischer (1964Fl07)	6.85	0.20	Not used. Mica-uranium sandwich.
A. Spadavecchia (1967Sp12)	8.42	0.10	Rotating bubble chamber.
J.H. Roberts (1968Ro15)	7.03	0.11	Not used. Mica-uranium sandwich.
H.R. von Gunten (1969Vo24)	8.66	0.22	Fission products of ^{238}U .
D. Galliker (1970Ga27)	8.46	0.06	Rotating bubble chamber.
D. Storzer (1970Storzer)	8.49	0.76	Fission tracks in dated uranium glass.
J.D. Kleeman (1971Kl14)	6.8	0.6	Not used. Lexan-uranium sandwich.
W.M. Thury (1971Th17)	8.66	0.43	Third order coincidence.
M.P.T. Leme (1971Le11)	7.30	0.16	Not used. Mica-uranium sandwich.
H.A. Khan (1973Kh10)	6.82	0.55	Not used. Mica-uranium sandwich.
K.N. Ivanov (1974Iv01)	7.12	0.32	Not used. Mica-uranium sandwich.
V. Emma (1975Em03)	7.2	0.2	Not used. Mica-uranium sandwich.
G.A. Wagner (1975Wa37)	8.7	0.6	Fission tracks in dated uranium glass.
K. Thiel (1976Th12)	8.57	0.42	Fission tracks in dated uranium glass.
M. Kase (1978Ka40)	8.22	0.20	Ionization chamber.
A.G. Popeko (1980Po09)	7.9	0.4	Multiple neutron coincidence.
E.R.V. Spaggiari (1980Sp10)	9.26	0.17	Not used. Mica-uranium sandwich.
Z.N.R. Baptista (1981Ba70)	6.6	0.2	Not used. Mica-uranium sandwich.
J.C. Hadler (1981Hadler)	8.6	0.4	Not used. Mica-uranium sandwich.
H.G. de Carvalho (1982De22)	11.8	0.7	Not used. Fission tracks in ordinary glass.
S.N. Belenky (1983Be66)	8.35	0.40	Multiple neutron coincidence.
B. Vartanian (1984Va34)	8.23	0.43	Not used. Fissions tracks (plastic, uranium foils).
M.P. Ivanov (1985Iv01)	8.29	0.27	Double ionization chamber.
S.S. Liu (1991Liu)	7.03	0.21	Not used. Solid-state track detectors.
Recommended value of λ^{238} (in 10^{-17} years $^{-1}$)	8.451	0.060	reduced $\chi^2 = 0.30$
Recommended half-life value (in 10^{15} years)	8.202	0.060	

The evaluators, following the recommendations of N.E. Holden (2000Ho27), have not used in their calculations the measurements with fission tracks in mica-uranium, lexan-uranium sandwiches or ordinary glass, because they significantly disagree with the rest (for more details see 2000Ho27). Thus the experimental values with associated uncertainties used in the weighted average calculation are those from 1950Whitehouse, 1952Se67, 1967Sp12,

1969Vo24, 1970Ga27, 1970Storzer, 1971Th17, 1975Wa37, 1976Th12, 1978Ka40, 1980Po09, 1983Be66 and 1985Iv01. A weighted average has been calculated using LWEIGHT computer program (version 3). Based on the Chauvenet's criterion, Popeko's value (1980Po09) has been shown to be an outlier.

The recommended value of λ^{238} is the weighted average (calculated with LWEIGHT computer program) of $8.451 \cdot 10^{-17} a^{-1}$ with an internal uncertainty of $0.046 \cdot 10^{-17} a^{-1}$. However, evaluators have adopted an uncertainty of $0.060 \cdot 10^{-17} a^{-1}$, minimum input value.

Using this value of λ^{238} and the formula:

$$t_{1/2} = \frac{\ln(2)}{\lambda^{238}},$$

the evaluators have deduced a partial spontaneous fission halflife of $8.202(60) \cdot 10^{15} a$ for ^{238}U and a spontaneous fission branching of $5.45(4) \cdot 10^{-05} \%$.

2.1 a Transitions and Emissions.

The energies of the α -particle transitions given in Section 2.1 have been calculated from Q_{α} (2003Au03) and level energies.

The energies of $\alpha_{0,0}$, $\alpha_{0,1}$ and $\alpha_{0,2}$ emissions given in Section 4 are from A. Rytz (1991Ri01).

Measured α -emission intensities are given in Table 4.

Table 4: Measured α -emission intensities, in %.

Energy (keV)	1959Ko58	2000Ga05	Recommended Value
4198 ($\alpha_{0,0}$)	77 (4)	77.54 (50)	77.54 (50)
4151 ($\alpha_{0,1}$)	23 (4)	22.33 (50)	22.33 (50)
4038 ($\alpha_{0,2}$)	0.23 (7)	0.13 (3)	0.13 (3)

The results of these two intensity measurements (1959Ko58 and 2000Ga05) are consistent with each other. Evaluators have adopted the most recent and precise results of Garcia-Toraño (2000Ga05).

2.2 g Transitions

The γ -ray probabilities of the 49- and 113-keV transitions have been deduced from decay-scheme balance by using the recommended experimental alpha emission intensity values (2000Ga05). (see **2.1 a Transitions and Emissions**).

Multipolarities of γ -ray transitions in the decay of ^{234}Th are from 1994Ak05:

49-keV γ -ray : E2

113-keV γ -ray: [E2]

The internal conversion coefficients (ICC's) have been calculated using the Icc99v3a computer program (GETICC dialog), which uses the new tables of Band et al (2002Ba85) (results of calculation for "hole" and "no hole" are the same). The evaluators have used a fractional uncertainty of 3 % for all conversion coefficients.

3 Atomic Data

Values of atomic values quantities ω_K , ω_L and n_{KL} , are from Schönfeld and Janßen (1996Sc06).

3.1 X rays and Auger electrons

The relative probabilities of X-ray and Auger electrons have been calculated from γ -ray data using the EMISSION computer program.

4 a Emissions

See 2.1 a Transitions and Emissions.

5 Electron emissions

The Auger electrons emission probabilities have been calculated from γ -ray data using the EMISSION computer program.

6 Photon Emissions

6.1 K x-rays

X-ray emission probabilities have been calculated from γ -ray data using the EMISSION computer program.

6.2 g-ray emissions

The energies of the γ -ray emissions given in Section 6 are from Y.A. Akovali (1994Ak05).

The absolute γ -ray emission intensities have been deduced from the absolute γ -ray transition probabilities and the internal conversion coefficients (ICC's). (see 2.2 g Transitions.).

Table 5 shows the recommended absolute γ -ray (photon) emission intensities of the 49- and 113-keV emissions as well as the experimental results obtained from direct measurements of emission intensities.

The agreement is not good, maybe due to experimental difficulties (many peaks of different contaminant isotopes in this energy region) when measuring these weak γ -ray intensities.

Table 5: Experimental absolute γ emission intensity in %.

γ Energy (keV)	1984Ro21	1990Ko40	1996Ru11	Recommended value
49.55	0.064 (8)	0.059 (2)		0.0698 (26)
113.5	0.0102 (15)		0.07 (1)	0.0174 (47)

A fair agreement has been found between the results given by J-C. Roy (1984Ro21) and the evaluators' recommended value for the 49-keV γ -ray.

For the 113-keV γ -ray, there is no good agreement either between results of direct experimental measurements or between those latter and the recommended value. In this energy region the experimental difficulties are associated with presence of many small peaks from different isotopes in the γ -ray spectrum.

7 References

- 1932Ko01 – A.F. Kovarik, N.I. Adams Jr., Phys. Rev. 40(1932)718 [$T_{1/2}$ (U-238)].
- 1935Schiedt – R. Schiedt, Österr. Akad. d. Wiss., Sitzungsberichte d. mathem. – naturw. Kl., Abt. IIa, 144Bd., Heft 5 und 6(1935)192 [$T_{1/2}$ (U-238)].
- 1941Curtis – L.F. Curtis, L.L. Stockman, B.W. Brown, US National Bureau of Standards Report No. A80 US GPO (1941), Washington DC [$T_{1/2}$ (U-238)].
- 1949Ki26 – A.C. Kienberger, Phys. Rev. 76(1949)1561 [$T_{1/2}$ (U-238)].
- 1950Whitehouse – W.J. Whitehouse, W. Galbraith, Phil. Mag. 41(1950)429 [S.F. half-life].
- 1952Se67 – E. Segrè, Phys. Rev. 86(1952)21 [S.F. half-life].
- 1955Ko13 – A.F. Kovarik, N.I. Adams Jr., Phys. Rev. 98(1955)46 [Half-life].
- 1956Kuroda – P. Kuroda, R.R. Edwards, F.T. Ashizawa, J. Chem. Phys. 25(1956)603 [S.F. half-life].
- 1957Kuroda – P. Kuroda, R.R. Edwards, J. Inorg. Nucl. Chem. 3(1957)345 [S.F. half-life].
- 1957Le21 – R.B. Leachman, H.W. Schmitt, J. Nucl. Energy 1(1957)38 [Half-life].
- 1957Cl16 – F.L. Clark, H.J. Spencer-Palmer, R.N. Woodward, J.S. Africain Chem. Inst. 10(1957)62 [Half-life].
- 1958Parker – P.L. Parker, P.K. Kuroda, J. Inorg. Nucl. Chem. 5(1958)153 [S.F. half-life].
- 1959St45 – J. Steyn, F.W.E. Strelow, Proc. Symp. Metrology Radionucl., Vienna, Austria, (1959)155 [Half-life].
- 1959Ku81 – B.D. Kuz'minov, L.S. Kutsaeva, V.G. Nesterov, L.I. Prokhorova, G.P. Smirenkin, Sov. Phys. JETP 37(1959)290 [S.F. half-life].
- 1959Ko58 – G.E. Kocharov, A.P. Komar, G.A. Korolev, Sov. Phys. JETP 36(1959)48 [Alpha probability].
- 1961Ko11 – G.E. Kocharov, G.A. Korolev, Bull. Acad. Sci. USSR, Phys. Ser. 25(1961)227 [Alpha probability].
- 1964Fl07 – R.L. Fleischer, P.B. Price, Phys. Rev. 133(1964)B63 [S.F. half-life].
- 1967Sp12 – A. Spadavecchia, B. Hahn, Helv. Phys. Acta 40(1967)1063 [S.F. half-life].
- 1968Ro15 – J.H. Roberts, R. Gold, R.J. Armani, Phys. Rev. 174(1968)1482 [S.F. half-life].
- 1969Vo24 – H.R. von Gunten, Actinides Rev. 1(1969)275 [S.F. half-life].
- 1970Ga27 – D. Galliker, E. Hugentobler, B. Hahn, Helv. Phys. Acta 43(1970)593 [S.F. half-life].
- 1970Storzer – D. Storzer, Thesis Universität Heidelberg (1970) [S.F. half-life].
- 1971Ja07 – A.H. Jaffey, K.F. Flynn, L.E. Glendenin, W.C. Bentley, A.M. Essling, Phys. Rev. C4(1971)1889 [Half-life].
- 1971Le11 – M.P.T. Leme, C. Renner, M. Cattani, Nucl. Instrum. Meth. 91(1971)577 [S.F. half-life].
- 1971Kl14 – J.D. Kleeman, J.F. Lovering, Geochimica et Cosmochimica Acta 35(1971)637 [S.F. half-life].
- 1971Th17 – W.M. Thury, Acta Physica Austriaca 33(1971)375 [S.F. half-life].
- 1973Kh10 – H.A. Khan, S.A. Durrani, Radiation Effects 17(1973)133 [S.F. half-life].
- 1974Iv01 – K.N. Ivanov, K.A. Petrzhak, Sov. At. Energ. 36(1974)514 [S.F. half-life].
- 1975Em03 – V. Emma, S. Lo Nigro, Nucl. Instrum. Meth. 128(1975)355 [S.F. half-life].
- 1975Wa37 – G.A. Wagner, G.M. Reimer, B.S. Carpenter, H. Faul, R. Van der Linden, R. Gijbels, Geochimica et Cosmochimica Acta 39(1975)1279 [S.F. half-life].
- 1976Th12 – K. Thiel, W. Herr, Earth and Planetary Sci. Lett. 30(1976)50 [S.F. half-life].
- 1978Ka40 – M. Kase, J. Kikuchi, T. Doke, Nucl. Instrum. Meth. 154(1978)335 [S.F. half-life].
- 1978Ri07 – D.M.C. Rizzo, An. Acad. Bras. Ciênc. 50(1978)303 [S.F. half-life].
- 1980Sp10 – E.R.V. Spaggiari, An. Acad. Bras. Ciênc. 52(1980)213 [S.F. half-life].
- 1980Po09 – A.G. Popeko, G.M. Ter-Akopian, Nucl. Instrum. Meth. 178(1980)163 [S.F. half-life].
- 1981Ba70 – Z.N.R. Baptista, M.S.M. Mantovani, F.B. Ribeiro, An. Acad. Bras. Ciênc. 53(1981)437 [S.F. half-life].
- 1981Hadler – J.C. Hadler, C.M.G. Lattes, A. Marques, M.D.D. Marques, D.A.B. Serra, G. Bigazzi, Nucl. Tracks 5(1981)46 [S.F. half-life].
- 1982De22 – H.G. de Carvalho, J.B. Martins, E.L. Medeiros, O.A.P. Tavares, Nucl. Instrum. Meth. 197(1982)417 [S.F. half-life].
- 1983El11 – Y.A. Akovali, Nucl. Data Sheets 40(1983)523 [Spin, parity, energy level, multipolarity].
- 1983Be66 – S.N. Belen'kii, M.D. Skorokhvatov, A.V. Etenko, Sov. At. Energ. 55(1983)528 [S.F. half-life].
- 1984Va35 – B. Vartanian, Helv. Phys. Acta 57(1984)416 [S.F. half-life].
- 1984Ro21 – J.-C. Roy, L. Breton, J.-E. Côté, J. Turcotte, Int. J. Appl. Radiat. Isotop. 35(1984)899 [Gamma probability].
- 1984Va34 – B. Vartanian, Helv. Phys. Acta 57(1984)292 [S.F. half-life].

- 1985Iv01 – M.P. Ivanov, G.M. Ter -Akopian, B.V. Fefilov, A.S. Voronin, Nucl. Instrum. Meth. Phys. Res. A234(1985)152 [S.F. half-life].
- 1987Al28 – B. Al-Bataina, J. Jänecke, Radiochimica Acta 42(1987)159 [Half-life].
- 1989Ho24 – N.E. Holden, Pure and Appl. Chem. 61(1989)1483 [$T_{1/2}$ (U-234)].
- 1990Ko40 – K. Komura, M. Yamamoto, K. Ueno, Nucl. Instrum. Meth. Phys. Res. A295(1990)461 [Gamma probability].
- 1991Liu – S.S. Liu, F. Zhang, Science in China 34, 9(1991)1120 [S.F. half-life].
- 1991Ry01 – A. Rytz, Atomic Data and Nuclear Data Tables 47(1991)205 [E_{α}].
- 1994Du15 – B. Duchemin, N. Coursol, M.-M. Bé, Nucl. Instrum. Meth. Phys. Res. A339(1994)146[Alpha and gamma probabilities].
- 1994Ak05 – Y.A. Akovali, Nucl. Data Sheets 71(1993)181 [Spin, parity, energy level, multipolarity].
- 1996Ru11 – H. Ruellan, M.-C. Lépy, M. Etcheverry, J. Plagnard, J. Morel, Nucl. Instrum. Meth. Phys. Res. A369(1996)651 [Gamma probability].
- 1996Sc06 – E. Schönfeld, H. Janssen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
- 1998Ad08 – I. Adsley, J.S. Backhouse, A.L. Nichols, J. Toole, Appl. Rad. Isotopes 49(1998)1337 [Gamma probability].
- 2000Ga05 – E. Garcia-Toraño, Appl. Rad. Isotopes 52(2000)591 [Alpha probability].
- 2000Ho27 – N.E. Holden, D.C. Hoffman, Pure Appl. Chem. 72(2000)1525 [S.F. half-life].
- 2002Ba85 – I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, Atomic Data and Nuclear Data Tables 81(2002)1 [α].
- 2002Ch52 – F.E. Chukreev, V.E. Makarenko, M.J. Martin, Nucl. Data Sheets 97(2002)123 [Spin, parity, energy level, multipolarity].
- 2003Ha06 – J.C. Hadler, G. Bigazzi, S. Guedes, P.J. Iunes, M. Oddone, C.A. Tello, S.R. Paulo, J. Radioanal. Nucl. Chem. 256(2003)155 [S.F. half-life].
- 2003Au03 – G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)129 [Q].
- 2004Sc03 – R. Schön, G. Winkler, W. Kustchera, Appl. Rad. Isotopes 60(2004)263 [Half-life].

**²³⁸Np - Comments on evaluation of decay data
by V. P. Chechev and N.K. Kuzmenko**

This evaluation was completed in November 2006 with a literature cut off by the same date.

1. Decay Scheme

The decay scheme is based on the evaluation of Chukreev *et al.* (2002Ch52) and can be basically considered completed.

2. Nuclear Data

Q^- value is from 2003Au03.

The evaluated half-life of ²³⁸Np is based on the experimental results given in Table 1.

Table 1. Experimental values of the ²³⁸Np half-life (in days)

Reference	Author(s)	Value
1950Fr53	Freedman <i>et al.</i>	2,10 (1)
1958A192	Albridge <i>et al.</i>	2,16 (15)
1966Qa01	Qaim	2,117 (2)
1990Ch35	Chang <i>et al.</i>	2,0980 (3)*
2006Re09	Rengan <i>et al.</i>	2,1024 (5)*

* Only statistical uncertainty

The evaluators increased the relative uncertainties of 1990Ch35 and 2006Re09 to 0,05% to take into account possible systematic uncertainties. The LWEIGHT computer program has omitted the outlier of 1958A192 and used a weighted average of 2,1024 with the expanded uncertainty of 0,0044 to give a recommended value.

The adopted value of the ²³⁸Np half-life is 2,102 (5) days.

2.1. Beta Transitions

The energies of β^- transitions have been calculated from the Q^- value and the level energies given in Table 2 from 2006Re09. The probabilities of β^- -transitions have been deduced from the $P(\gamma+ce)$ balance for each level of ²³⁸Pu.

The β transition probability to the 44 -keV level has been deduced from the 44 -keV level intensity balance using $P(\gamma_{1,0} + ce)(44,07\text{-keV})$ obtained from the intensity balance for the ground state (see 2.2)

Table 2. ^{238}Pu levels populated in the ^{238}Np β^- -decay

Level number	Level Energy, keV	Spin and parity	Half-life	Probability of β^- -transition (%)
0	0,0	0^+	87,74 (3) a	-
1	44,08 (2)	2^+	177 (5) ps	41,0 (25)
2	145,95 (2)	4^+		-
3	303,38 (6)	6^+		-
4	605,14 (4)	1^-		0,103 (3)
5	661,40 (6)	3^-		0,036 (3)
6	763,24 (11)	5^-		-
7	941,46 (8)	0^+		-
8	962,78 (2)	1^-		1,25 (1)
9	968,2 (4)	(2^-)		0,082 (6)
10	983,09 (7)	2^+		0,27 (3)
11	985,45 (5)	2^-		0,49 (1)
12	1028,54 (2)	2^+		44,75 (19)
13	1069,94 (2)	3^+		11,50 (7)
14	1082,56 (6)	$(4)^-$		-
15	1202,46 (8)	$(3)^-$		0,51 (6)

Table 3. Measured and evaluated β^- energies (keV) and probabilities (%) in the ^{238}Np decay

1955Ra28		1956Ba95	1962Bo03		Evaluated	
Eb $^-$	Pb $^-$	Pb $^-$	Eb $^-$	Pb $^-$	Eb $^-$	Pb $^-$
			200	8	221,6 (4)	11,50 (7)
			250 (10)	31		
258	53	55			263,0 (4)	44,75 (19)
			280 (10)	20		
			1133	2,8		
1272	47	45	1236 (5)	38	1247,4 (4)	41,0 (25)

2.2. Gamma Transitions and Internal Conversion Coefficients

The evaluated energies of gamma-ray transitions are essentially the same as the gamma-ray energies because nuclear recoil is negligible.

The $P(\gamma+ce)$ values have been calculated from the gamma -ray emission probabilities and the total internal conversion coefficients (ICC's).

For E0- gamma transition 941,5-keV ($\gamma_{7,0}$) the value $P(ce) = 0,0106$ (9) is based on measurements $P(ceK)$ of 1981Le15 and ICC ratios from the BrIcc package.

The experimental values of ICC's (from 1981Le15) have been adopted for the following gamma-ray transitions: 120,11-keV ($\gamma_{15,14}$), 220,9-keV ($\gamma_{-1,6}$), 923,9-keV ($\gamma_{13,2}$), (E0+E2) gamma-ray transition 939-keV ($\gamma_{10,1}$) (see also 1960Al29), 983,0-keV ($\gamma_{10,0}$) and 984,5-keV ($\gamma_{12,1}$). ICC's have been interpolated from the BrIcc package. The relative uncertainties of α_K , α_L , α_M , α_T for pure multiplicities have been taken as 2 %. The multiplicities and E2/M1, M2/E1 mixing ratios have been taken from 2002Ch52. These are based on

conversion electron measurements of 1952Du12, 1956Ba95, 1956Sm18, 1960As10, and 1965Ak02.

$P(\gamma_{1,0+ce})(44,08\text{-keV})$ has been deduced from the intensity balance for the ground state assuming that there is no beta-feeding to the $''0''$ -level. The second forbidden beta-transition is expected to the ground state with $\lg ft > 15$ which implies $< 0,01\%$ (2006Re09).

3. Atomic Data

3.1. Fluorescence yields

Fluorescence yield data are from 1996Sc06 (Schönfeld and Janßen).

3.1.1. X rays

The Pu KX-ray relative emission probabilities have been taken from 1999ScZX

3.1.2. Auger Electrons

The energies of Auger electrons have been calculated from atomic electron binding energies. The $P(KLX)/P(KLL)$, $P(KXY)/P(KLL)$ ratios have been taken from 1996Sc06.

5. Electron Emissions

The energies of the conversion electrons have been calculated from the gamma transition energies and the electron binding energies.

The emission probabilities of the conversion electrons have been calculated using evaluated P_γ and ICC values.

The absolute emission probabilities of K and L Auger electrons have been calculated using the EMISSION computer program.

β^- average energies have been calculated using the LOGFT computer program.

6. Photon emissions

6.1. X-Ray Emissions

The absolute emission probabilities of Pu KX- and LX-rays have been calculated using the EMISSION computer program.

Table 4. Measured and evaluated probabilities of Pu KX in the decay of ²³⁸Np.

	1972Wi22	1981Le15	Evaluated
K α_2	0,18(1)		0,210 (8)
K α_1	0,272(12)		0,332 (12)
K β'_1		0,11	0,122 (5)
K β'_2		0,050	0,042 (2)

6.2. Gamma Emissions

The gamma ray energies have been evaluated from experimental data (Table 3)

Table 5. The measured and recommended gamma ray energies in the ^{238}Np β^- -decay (keV).

1970Lederer	1972Wi22	1981Le15	2006Re09	Recommended
44	44,08 (3)		44,06 (2)	44,07 (2)
101,93 (4)	101,88 (2)		101,88 (3)	101,88 (2)
			103,74 (2)	103,74 (2)
			116,27 (8)	116,27 (8)
			117,27 (8)	117,27 (8)
119,9 (1)	120,14 (5)		120,09 (5)	120,11 (5)
			120,5	120,5
			120,70 (8)	120,70 (8)
			121,70 (8)	121,70 (8)
132,49 (11)	132,6 (6)		132,8 (5)	132,5 (1)
157,4 (3)		157,42 (5)	157,42	157,42 (5)
173,78 (11)	174,06 (8)		174,08 (5)	174,08 (5)
220,87 (11)			220,87	220,87 (11)
301,19 (12)	301,81 (19)		301,37 (7)	301,37 (7)
319,29 (11)			319,96 (20)	319,29 (11)
321,75 (20)			321,75	321,75 (20)
323,98 (9)	324,08 (17)		324,07 (15)	324,02 (9)
357,60 (9)	357,64 (7)		357,68 (9)	357,64 (7)
378,05 (13)			378,0 (10)	378,05 (13)
380,28 (13)	380,33 (22)		380,33 (10)	380,31 (10)
421,15 (11)	421,12 (16)		421,05 (10)	421,10 (10)
459,8 (2)		459,80 (22)	459,8 (2)	459,8 (2)
515,58 (12)	515,47 (17)	515,25 (19)	515,53 (7)	515,51 (7)
561,09 (10)	561,15 (7)	561,02 (10)	561,17 (5)	561,14 (5)
605,24 (13)	605,14 (9)	605,04 (10)	605,18 (5)	605,16 (5)
617,45 (12)	617,39 (11)	617,22 (12)	617,41 (5)	617,39 (5)
837,18 (15)	837,0 (4)	837,01 (15)	836,88 (7)	836,96 (7)
882,65 (7)	882,63 (3)		882,63 (3)	882,63 (3)
897,28 (20)		897,33 (10)	897,55 (30)	897,34 (10)
918,70 (7)	918,69 (4)	918,7 (2)	918,70 (4)	918,70 (4)
923,99 (6)	923,98 (2)		923,99 (2)	923,99 (2)
936,57 (9)	936,61 (6)		936,60 (5)	936,60 (5)
939,00 (10)	938,6 (5)	938,91 (10)	938,85 (30)	938,94 (10)
941,39 (6)	941,38 (5)		941,41 (4)	941,40 (4)
941,5 (3)				941,5 (3)
962,80 (7)	962,77 (3)	962,8 (2)	962,76 (2)	962,76 (2)
984,46 (7)	984,45 (2)	984,5 (1)	984,45	984,45 (2)
1025,87 (6)	1025,87 (2)		1025,87 (2)	1025,87 (2)
1028,54 (6)	1028,54 (2)	1028,5 (2)	1028,53 (2)	1028,54 (2)

The absolute emission probabilities for gamma -rays have been deduced from the evaluated relative intensities (see Table 6) using the weighted mean $P(\gamma_{12,1})(984,5\text{-keV}) = 0,2518$ (13) of the two absolute measurement results: 0,2517 (13) from 2006Re09 and 0,2519 (21) from 1990Ch15.

It should be noted that in 1981Le15 the differing absolute value of $P(\gamma_{12,1})(984,5\text{-keV}) = 0,278$ (8) was deduced from an intensity balance for the ground state of ^{238}Pu .

Using the value of 0,397(6) from 2006Re09 for the relative gamma ray intensity of $\gamma_{1,0}$ (44,07-keV) and the evaluated relative intensities for the remaining gamma -rays from Table 4, we obtain from the ground state intensity balance the value of $P(\gamma_{12,1})(984,5\text{-keV}) = 0,257$ (6) which supports our above more exact value and disagree with 1981Le15.

The absolute gamma ray intensity for $\gamma_{1,0}$ (44,07-keV) has been deduced from the evaluated $P(\gamma_{1,0} + \text{c.e.})(44,07\text{ keV})$ and the adopted total ICC.

The absolute gamma ray intensities for $\gamma_{5,1}$ (617,36-keV) and $\gamma_{6,2}$ (617,36-keV) have been deduced using the

ratio $P(\gamma_{5,1})(617,36\text{-keV})/P(\gamma_{6,2})(617,36\text{-keV}) = 65/9$ adopted from 1981Le15.

The relative gamma ray intensity ($P'(\gamma)$) and energy for $\gamma_{9,4}$ (924-keV) have been adopted from 1970Be57.

The recommended $P'(\gamma)$ for $\gamma_{1,0}$ (44,07-keV) has been obtained as a ratio of the evaluated $P(\gamma_{1,0})(44,07\text{-keV})$ to $P(\gamma_{12,1})(984,5\text{-keV})$ and it has also been compared to measured values.

Table 6. Measured and evaluated relative gamma-ray intensities.

Energy (keV)	1972Wi22	1981Le15*	1990Ch35	2006Re09	Recommended
44,07	≈0,2	0,32 (4) ^a	0,35 (4)	0,397 (6)	0,406 (9)
99,53				0,771 (8)	0,771 (8)
101,9	0,88 (2)	0,97 (4)	1,01 (3)	1,01 (1)	1,00 (3)
103,7				1,24 (1)	1,24 (1)
116,3				0,158	0,158
117,3				0,295	0,295
120,1	0,41 (2)	0,37 (3)		0,453 (9)	0,40 (2)
120,5				0,079	0,079
120,7					
121,7				0,040 (4)	0,040 (4)
132,5	0,013 (7)	0,0101 (7)		0,0056 (3)	0,0056 (3)
157,4		≈0,004			≈0,004
174,0	0,11 (1)	0,094 (4)	0,091 (3)	0,088 (6)	0,091 (3)
220,9		0,0122 (14)		0,007 (6)	0,012 (2)
301,4	0,05 (1)	0,043 (4)	0,040 (4)	0,054 (11)	0,042 (4)
319,3		0,032 (4)		0,038 (12)	0,033 (4)
321,8		0,0047 (22)		0,008 (8)	0,005 (2)
324,0	0,070 (11)	0,058 (4)	0,057 (3)	0,061 (10)	0,058 (3)
336,4					0,0009 (5)
357,6	0,22 (2)	0,191 (11)	0,200 (5)	0,20 (1)	0,200 (5)
378,0		0,012 (2)		0,008 (8)	0,012 (2)
380,3	0,05 (1)	0,043 (2)		0,064 (12)	0,044 (2)
421,1	0,096 (15)	0,083 (4)	0,087 (4)	0,079 (12)	0,085 (4)
459,8		≈0,011		0,009 (6)	0,009 (6)
515,5	0,14 (2)	0,155 (7)	0,148 (5)	0,14 (1)	0,150 (5)
561,1	0,43 (2)	0,41 (2)	0,416 (7)	0,461 (16)	0,423 (7)
605,2	0,31 (3)	0,284 (14)	0,318 (9)	0,29 (2)	0,306 (9)
617,39 (5) } 617,4	0,29 (3)	0,266 (14)	0,270 (9)	0,262 (12)	0,268 (9)
837,0	0,076 (22)	0,101 (7)		0,079 (3)	0,082 (3)
882,6	3,19 (16)	3,13 (11)	3,23 (3)	3,17 (2)	3,19 (2)
885,0				0,16 (2)	0,16 (2)
897,3		0,029 (4)	0,029 (4)	0,032 (8)	0,029 (4)
918,7	2,16 (11)	2,12 (7)	2,11 (2)	2,09 (2)	2,10 (2)
923,99	10,4 (5)	10,3 (3)	10,4 (1)	10,32 (6)	10,34 (6)
924					0,26
936,6	1,39 (7)	1,44 (4)	1,46 (2)	1,41 (11)	1,45 (2)
938,9	0,13 (6)	0,10 (3)	0,13 (1)	0,13 (1)	0,13 (1)
941,4	1,91 (10)	1,98 (7)	2,04 (2)	1,97 (2)	2,00 (2)
941,5					
962,8	2,56 (13)	2,52 (7)	2,56 (3)	2,56 (3)	2,56 (3)
968,5	0,06 (2)	-	-	0,004	0,06 (2)
983,0					0,27 (8)
984,4	100	100	100	100	100
1025,9	34,5 (17)	34,9 (22)	34,59 (50)	34,82 (18)	34,79 (18)
1028,5	72,5 (36)	73,0 (29)	72,61 (70)	72,42 (37)	72,47 (37)

* Absolute gamma-ray emission probabilities cited in 1981Le15 (normalized to 27,8 for the 984,5-keV gamma- ray) have been converted to the relative gamma-ray intensities.

^a Measured value. In 1981Le15 it is noted that the value deduced from an intensity balance is 0,36 (2).

7. References

- 1950Fr53 M.S. Freedman, A.H. Jaffey and F. Wagner, Jr, Phys.Rev. 79, 410 (1950) (Half-life)
- 1952Du12 D.C. Dunlavy and G.T. Seaborg, Phys. Rev. 87, 165 (1952)
(Conversion electron measurements, gamma ray multipolarities)
- 1955Ra28 J.O.Rasmussen, F.S.Stephens, D.Strominger and B.Astrom, Phys.Rev. 99, 47 (1955)
(Measured β^- energies and probabilities)
- 1956Ba95 S.A. Baranov and K.N. Shlyagin, At. Energ. USSR 1, 52 (1956); J. Nuclear Energy 3, 132(1956) (Conversion electron measurements, gamma-ray multipolarities, measured β^- probabilities)
- 1956Sm18 W.G. Smith and J.M. Hollander, Phys. Rev. 101, 746 (1956)
(Conversion electron measurements, gamma-ray multipolarities)
- 1958Al92 R.G. Albridge, J.C. Hubbs and R. Marrus, Phys.Rev. 111, 1137 (1958) (Half-life)
- 1960Al29 R.G. Albridge and J.M. Hollander, Nucl. Phys. 21, 438 (1960)
(Conversion electron measurements, gamma-ray multipolarities)
- 1960As10 F. Asaro and I. Perlman, UCRL-9566, p.50 (1960)
(Conversion electron measurements, gamma-ray multipolarities)
- 1962Bo03 J.Borggreen, O.B.Nielsen and H.Nordby, Nuclear Phys. 29, 515 (1962)
(Measured β^- energies and probabilities)
- 1965Ak02 G.G. Akalaev, N.A. Vartanov and P.S. Samoilov, NP-14688 (1965)
(Conversion electron measurements, gamma-ray multipolarities)
- 1966Qa01 S.M.Qaim, Nucl.Phys. 84, 411 (1966) (Half-life)
- 1970Be57 B.Bengtson, J.Jensen, M.Moszynski, H.L.Nielsen Nucl.Phys. A159, 249 (1970)
(924-keV gamma ray energy and relative emission probability)
- 1970Lederer C.M. Lederer, Priv. Comm. 1970, see C.M. Lederer, V.Shirley, Table of Isotopes, N.Y., John Wiley and Sons,1978. (Gamma ray energies)
- 1972Wi22 W.J.B. Winter, A.H. Wapstra, P.F.A. Goudsmit and J. Konijn, Nucl. Phys. A197, 417 (1972) (Relative gamma ray intensities)
- 1981Le15 C.M. Lederer, Phys. Rev. C24, 1175 (1981) (Relative gamma ray intensities)
- 1990Ch35 Y. Chang, B. Zhu, C. Yan, G. Shi and J. Chin, Nucl.Phys. 12, No 1, 65 (1990)
(Relative gamma ray intensities, absolute 984-keV gamma ray emission probability)
- 1996Sc06 E. Schönfeld and H. Janßen, Nucl. Instrum. Methods Phys. Res. A369, 527 (1996)
(Atomic data)
- 1999ScZX E. Schönfeld and G. Rodloff, PTB-6.11-1999-1999-1, Braunschweig, February1999
(KX rays relative emission probabilities)
- 2002Ch52 F.E. Chukreev, V.E. Makarenko and M.J. Martin, Nucl. Data Sheets 97, 129 (2002)
(Nuclear data evaluation for A=238)
- 2003Au03 G. Audi, A.H. Wapstra and C. Thibault, Nucl. Phys. A729, 337 (2003) (Q value)
- 2006Re09 K. Rengan, D. Devries, H. Griffin, Nucl. Instrum. Methods Phys. Res. A565, 612 (2006).
(Gamma-ray energies, relative gamma ray intensities, absolute 984-keV gamma ray emission probability)

²³⁸Pu – Comments on evaluation of decay data by V. P. Chechev

This evaluation was done originally in March 2003, corrected in June 2004, and then updated in June 2009 with a literature cut-off by the same date.

1. DECAY SCHEME

The decay scheme is based on 2007Br04. Some expected weak gamma-ray transitions were not observed directly in ²³⁸Pu α -decay but have been adopted from decay of ²³⁴Pa and ²³⁴Np.

2. NUCLEAR DATA

Q(α) value is from 2003Au03.

The recommended half-life of ²³⁸Pu is based on the experimental results given in Table 1.

Table 1. Experimental values of ²³⁸Pu half-life (in years)

Reference	Author(s)	Original value ^a	Re-estimated value ^a	Measurement method	Used for final averaging
1950Jaffey	Jaffey and Lerner	89.59 (37)	89.3 (9) ^b	Direct decay (4 samples)	No
1951Jaffey-1	Jaffey and Magnusson	77	-	Growth of ²³⁸ Pu from ²³⁸ Np	No
1951Jaffey-2	Jaffey	89 (9)	-	Direct decay	No
1951Seaborg	Seaborg et al.	92 (2)	-	Growth of ²³⁸ Pu from ²⁴² Cm	No
1954Jo10	Jones et al.	89	-		No
1957Ho71	Hoffman et al.	86.41 (30)	86.4 (5) ^b	Growth of ²³⁸ Pu from ²⁴² Cm	No
1965Eichelber	Eichelberger et al.	87.60 (6)	-	Calorimetry	No
1967Jordan	Jordan	87.22 (52)	-	Calorimetry	No
1969Benson	Benson	87.75 (5)	-	Calorimetry	No
1974StYG	Strohm and Jordan	87.77(3)	-	Calorimetry	Yes
1976Po08	Polyukhov et al.	86.98 (20)	87.0 (7) ^c	Specific activity	Yes
1977Di04	Diamond et al.	87.71 (3)	-	Growth of ²³⁸ Pu from ²⁴² Cm	Yes
1981Ag06	Aggarwal et al.	87.98 (51)	-	Relative activity ²³⁸ Pu/ ²³⁹ Pu	Yes
1981 Sevastyanov	Sevastyanov and Yarina	86.51 (30)	86.5 (9) ^d	Direct decay (1 sample)	No

^a Uncertainty at the level of 1 σ .

^b Re-estimated in 1977Di06.

^c Re-estimated by the evaluator using analysis of 1977Di06.

^d Re-estimated by the evaluator.

By omitting two values reported without uncertainties, the weighted average of the remaining 12 values is 87.73 with an internal uncertainty of 0.019 and $\chi^2/\nu = 2.0$. The average value of 87.73 (3) could be adopted for half-life of ²³⁸Pu. However several calorimetric results obtained in the same laboratory (MLM) may be correlated. In fact, the value 87.77 (3) (1974StYG) comes from the latest calorimetric measurement at this laboratory. Also, the early inaccurate experimental results published in 1950 – 1957 may be omitted, as they were obtained with samples of low isotopic purity. Besides, there are grounds for omitting the result of 1981Sevastyanov (V. D. Sevastyanov and V. P. Jarina, *Voprosi Atomnoi Nauki i Tekhniki, seriya Jadernie Konstanti*. 5(44)(1981)21), as it was obtained only from one sample using an inaccurate method of direct decay.

Therefore, the four best experimental results obtained by different methods were used for the final statistical analysis. These are 87.77 (3) – 1974StYG; 87.0 (7) – 1976Po08; 87.71 (3) – 1977Di04 and 87.98 (51) – 1981Ag06. The weighted average of these data sets is 87.74 with an internal uncertainty of

0.021 and $\chi^2/\nu = 1.1$. The recommended value of ²³⁸Pu half-life is 87.74 (3) years where the uncertainty is the smallest experimental uncertainty.

The evaluated spontaneous fission half-life of ²³⁸Pu has been based on the experimental results given in Table 2. The weighted average of 5 selected values (with reported uncertainties) is 4.74 with an internal uncertainty 0.081 and $\chi^2/\nu = 0.72$.

The recommended value of ²³⁸Pu spontaneous fission is 4.74 (12)·10¹⁰ years where the uncertainty is the smallest experimental uncertainty.

Table 2. Experimental values of ²³⁸Pu spontaneous fission half-life (in 10¹⁰ years)

Reference	Author(s)	Original value ^a	Re-estimated value ^a	Measurement method	Used for final averaging
1949Jaffey	Jaffey and Hirsch	4.9 (4)	4.7 (6) ^b	Ioniz. chamber	Yes
1952Se67	Segre	2.6	3.9 ^b	Ioniz. chamber	No
1961Dr04	Druin et al.	5.0 (6)	5.1 (6) ^b	Photoemulsion	Yes
1972Ha11	Hastings and Strohm	4.77 (14)	-	Si(Au)	Yes
1975GaZX	Gay and Sher	4.63 (12)	-	Fission fragm. coincid. in mica	Yes
1988SeZY	Selitsky et al.	5.01 (21)	-	2 π ioniz. chamber	Yes

^a Uncertainty at the level of 1 σ .

^b Adjusted in 1972Ha11 to ²³⁸Pu half-life of 87.77 yr. See also 2000Ho27.

2.1. Alpha Transitions

The energies of the alpha transitions have been obtained from the Q value and the level energies given in Table 3 from 2007Br04.

Table 3. ²³⁴U levels populated in ²³⁸Pu α decay

Level number	Energy, keV	Spin and parity	Half-life	Probability of α -transition (x100)
0	0,0	0+	2.455 (6) 10 ⁵ yr	71.04 (6)
1	43.4981 (10)	2+	0.252 (7) ns	28.85 (6)
2	143.352 (4)	4+		0.104 (3)
3	296.072 (4)	6+		0.00292 (4)
4	497.04 (3)	8+		6.80 (23)·10 ⁻⁶
5	786.288 (16)	1-		8.21 (16)·10 ⁻⁶
6	809.907 (18)	0+	< 0.1 ns	1.0·10 ⁻⁴
7	849.266 (18)	3-		7.5 (22)·10 ⁻⁸
8	851.74 (3)	2+	> 1.74 ps	8.1·10 ⁻⁶
9	926.720 (15)	2+	1.38 (17) ps	1.30 (5)·10 ⁻⁶
10	947.64 (6)	4+		2.3·10 ⁻⁷
11	989.430 (13)	2-	0.76 (4) ns	1.50 (15)·10 ⁻⁷
12	1023.77 (3)	4+		~ 2.0·10 ⁻⁷
13	1044.536 (23)	0+		1.17(7)·10 ⁻⁶
14	1085.26 (4)	2+		~ 1.2·10 ⁻⁶

The probabilities of the most intense transitions $\alpha_{0,0}$ and $\alpha_{0,1}$ have been obtained by averaging experimental data (Table 4). The probabilities of all the remaining α -transitions have been deduced from the P(γ +ce) balances at relevant levels in ²³⁴U.

Table 4. Experimental and recommended values of α -transition probabilities ($\times 100$) in the decay of ²³⁸Pu

	Energy keV	1954 As07	1957 Ko33	1970 Ba72	1971 So15	1984 Ah06	1984 Bo41	1984 Burns	1987 Bo25	1998 Ya17	Recommended
$\alpha_{0,0}$	5499	72 ^a	71.1 (12)	72.2 ^a	70.7 (2)	70.9 (1)	70.91 (10)	71.11 (4)	71.3 (6)	71.14 (10)	71.04 (6) ^b
$\alpha_{0,1}$	5456	28 ^a	28.7 (12)	27.8 ^a	29.3 (2)	29.0 (1)	28.98 (10)	28.78 (4)	28.6 (4)	28.74 (10)	28.85 (6) ^b
$\alpha_{0,2}$	5358		0.13 (1)	0.068 ^a	0.1 ^a	0.106 (3)	0.105 (5)	0.1002 (17)		0.114 (10)	0.104 (3) ^c
$\alpha_{0,3}$	5208		0.005 (1)	0.0018 ^a		0.036 (5)	0.0030 (1)				0.00292 (4) ^{d,e}
$\alpha_{0,4}$	5010			$\sim 4 \cdot 10^{-6}$							$6.80 (23) \cdot 10^{-6}$ ^e
$\alpha_{0,5}$	4726			$2.2 \cdot 10^{-5}$							$8.21 (16) \cdot 10^{-6}$ ^e
$\alpha_{0,6}$	4703			$5 \cdot 10^{-5}$							$1.0 \cdot 10^{-4}$ ^{e,f}
$\alpha_{0,7}$	4664										$7.5 (22) \cdot 10^{-8}$ ^e
$\alpha_{0,8}$	4662			$< 2 \cdot 10^{-5}$							$8.1 \cdot 10^{-6}$ ^e
$\alpha_{0,9}$	4588			$(1.2 \cdot 10^{-5})$							$1.30 (5) \cdot 10^{-6}$ ^e

^a Omitted from averaging because no uncertainty was reported.

^b Weighted average of 7 experimental values; uncertainty is external.

^c Weighted average of 5 experimental values (with quoted uncertainties) is 0.104 (3); the value deduced from P(γ +ce) balance is 0.1030 (24); the recommended value is 0.104 (3).

^d Agrees well with the experimental value from 1984Bo41

^e Evaluated from P(γ +ce) balance.

^f Value of $1.2 (4) \cdot 10^{-4}$ was obtained by α - γ and α -ce coincidences in 1963Bj03.

2.2. Gamma Transitions and Internal Conversion Coefficients

The recommended energies of gamma-ray transitions are virtually the same as the gamma-ray energies because nuclear recoil is negligible for ²³⁴U.

Gamma-ray transition probabilities [P(γ +ce)] have been deduced from the gamma-ray emission probabilities and total internal conversion coefficients (ICCs). The ICCs have been interpolated using the BrIcc package with the so called “Frozen Orbital” approximation (2008Ki07). The uncertainties in the ICCs for pure multipolarities have been taken as 2 %.

The emission probabilities of E0- and (E0+E2)- transitions have been obtained by using experimental conversion electron intensities from ²³⁴Pa and ²³⁴Np decays (see 2007Br04) and data from ²³⁸Pu α -decay of 1963Bj03, 1964Le17, 1964Le22.

3. ATOMIC DATA

3.1. Fluorescence yields

Fluorescence yield data are from 1996Sc06 (Schönfeld and Janßen).

3.2. X Radiations

The U KX-ray energies have been taken from 1999Schönfeld where the calculated values based on X-ray wavelengths from 1967Be65 (Bearden). In Table 5 the recommended values of U KX-ray energies are compared with experimental values.

The relative K X-ray emission probabilities have been taken from 1999Schönfeld.

Table 5. Experimental and recommended (calculated) values of U KX-ray energies (keV)

	1976GuZN	1982Ba56	1983Ah02	Recommended
K α_2	94.655 (5)	94.656 (2)	94.67 (2)	94.666
K α_1	98.442 (5)	98.435 (2)	98.45 (2)	98.440
K β_3	110.42 ^a	110.416 (3)	110.42 (3)	110.421
K β_1	111.30 ^a	111.300 (2)	111.31 (2)	111.298
K β_5	-	111.868 (5)- K β_5 , 112.043 (5)- K β_5	112.01 (5)	111.964
K $\beta_{2,4}$	114.54 ^a	-	114.50 (3)	114.46
KO $_{2,3}$	115.40 ^a	-	115.40 (5)	115.377

The energies of U LX-rays taken from the SAISINUC software supporting programs agree with the measurements of 1994Le37 where the fine structure of LX-radiation was measured in decays of ²³⁹Pu and ²⁴⁰Pu.

3.3. Auger Electrons

The energies of Auger electrons are from the SAISINUC software supporting programs.

The ratios P(KLX)/P(KLL), P(KXY)/P(KLL) are from 1996Sc06.

4. ALPHA EMISSIONS

The energy of alpha particles corresponding to the alpha transition to the ground state of ²³⁴U, E($\alpha_{0,0}$), has been adopted from the absolute measurement of 1971Gr17 with a correction of - 0.18 keV recommended by A. Rytz in 1991Ry01 because of changes in calibrations energies.

The energies of all other alpha particles have been calculated from Q(α), E($\alpha_{0,0}$) and the level energies taking into account the recoil energies.

In Table 6 the deduced (recommended) values of α -particle energies are compared with the experimental results obtained by using magnetic and semiconductor spectrometry.

Table 6. Experimental and recommended values of α -particle energies (keV) in decay of ²³⁸Pu.

	Measured ^a						Recommended
	1954As07	1957Ko33	1962Le11	1968Ba25	1970Ba72	1971Gr17	
$\alpha_{0,0}$	5499	5497.7 (10)	5499.2 (8)	5499.2 (10)	5499.2 (8) ^c	5499.03 (20) ^b	5499.03 (20) ^b
$\alpha_{0,1}$	5456	5454.7 (10)	5456.3 (8)	5456.1 (10)	5456.1	5456.3 (4)	5456.3 (2)
$\alpha_{0,2}$	5358	5358.6 (10)	5362 (1)		5357.7		5358.1 (2)
$\alpha_{0,3}$		5215 (5)			5205.6		5208.0 (2)
$\alpha_{0,4}$					≈5015		5010.4 (2)
$\alpha_{0,5}$					4724		4726.0 (2)
$\alpha_{0,6}$					4704		4702.8 (2)
$\alpha_{0,7}$					-		4664.1 (2)
$\alpha_{0,8}$					4661		4661.7 (2)
$\alpha_{0,9}$					≈4590		4587.9 (2)

^a Original values have been adjusted for changes in calibration energies as suggested in 1991Ry01.

^b Absolute measurement; this value is recommended in 1991Ry01 and used in 2003Au03 for obtaining Q(α).

^c Value is from 1962Le11; adopted in 1970Ba72 as calibration energy.

5. ELECTRON EMISSIONS

The energies of conversion electrons have been obtained from the gamma-ray transition energies and atomic-electron binding energies.

The emission probabilities of conversion electrons have been deduced from the evaluated $P(\gamma)$ and ICC values. Below the experimental L1:L2:L3 conversion electron sub-shell intensities from 1969Am02 are compared with theoretical values for the most intense E2 transition of $\gamma_{1,0}$ (43.498 keV).

Theoretical	Measured
3.85 (11) : 113 (3) : 100	3.99 (22) : 114.7 (20) : 100

The total absolute emission probabilities of K Auger electrons have been deduced using the evaluated total $P(XK)$ and the adopted fluorescence yield ω_K .

The total absolute emission probability of L Auger electrons have been deduced using the total evaluated $P(XL)$ and the adopted fluorescence yield ω_L .

6. PHOTON EMISSIONS

6.1. X-Ray Emissions

6.1.1. M X-Rays

The total absolute emission probability of MX-rays is based on the measurement (1990Po14) of the relative emission probability $P(MX)/P(LX) = 0.194$ (24).

6.1.2. L X-Rays

The calculation of the total absolute emission probability of LX-rays [$P(XL)$], using the EMISSION computer program (2000Schönfeld), gives $P(XL) = 10.55$ (25) %. The available experimental results for $P(XL)$ are discrepant: 13 % - 1954As07; 10.6 (3) % - 1964Ha14; 12.83 (14) % - 1968By01; 9.2 (1) % - 1968Salgueiro; 11.2 (4) % - 1968Swinth; 11.4 (3) % - 1971Swinth; 14.18 (11) % - 1976Va23; 11.38 (10) % - 1977Bemis; 11.55 (18) % - 1984Bo41; 10.62 (32) % - 1984DrZX and 1984BaYT; 10.63 (8) % - 1995Jo23.

The result of the most accurate and latest measurement (1995Jo23) agrees well with the calculated values and with the value from 1984DrZX where the fine structure of LX-radiation was measured. The value from 1995Jo23 has been adopted as the recommended absolute emission probability of U LX-rays from decay of ²³⁸Pu: $P(XL) = 10.63$ (8) %.

For the evaluation of emission probabilities of the LX-ray components L_I , L_α , $L_{\beta\eta}$, L_γ the measured values given in Table 7 were renormalized by the evaluator to the adopted value $P(XL) = 10.63$ (8) % and then averaged. In Table 8 the evaluated emission probabilities are compared with values calculated in 1995Jo23 from alpha-branching ratios, theoretical ICC and theoretical atomic branching ratios.

Table 7. Experimental absolute emission probabilities of U LX-rays from α decay of ²³⁸Pu

	1976Va23	1977Bemis	1984Bo41	1995Jo23
L_I	-	0.26 (1)	0.260 (7)	0.231 (3)
L_α	5.05 (6)	4.15 (7)	4.06 (6)	3.81 (3)
$L_{\beta\eta}$	7.41 (9)	5.61 (7)	5.85 (9)	5.31 (4)
L_γ	1.48 (2)	1.36 (2)	1.38 (2)	1.29 (1)

Table 8. Renormalized experimental, evaluated, and calculated absolute emission probabilities of U LX-rays from α decay of ²³⁸Pu

	1976Va23 (measured)	1977Bemis (measured)	1984Bo41 (measured)	1995Jo23 (measured)	Adopted (averaged)	Calculated (1995Jo23)	Calculated (EMISSION code)
Ll	-	0.24 (1)	0.239 (7)	0.231 (3)	0.235 (4) ^b	0.234	0.232 (8)
L α	3.77 (5)	3.88 (7)	3.74 (6)	3.81 (3)	3.80 (3) ^c	3.78	3.73 (12)
L $\beta\eta$	5.53 (7) ^a	5.24 (7)	5.38 (8)	5.31 (4)	5.31 (4) ^c	5.42	5.23 (16)
L γ	1.10 (2) ^a	1.27 (2)	1.27 (2)	1.29 (1)	1.28 (1) ^c	1.26	1.23 (4)

^a Omitted from averaging based on statistical considerations.

^b Weighted average; uncertainty is internal.

^c Weighted average; uncertainty is the smallest experimental one.

6.1.3. KX-Rays

The absolute X-ray emission probability of U K α_2 with energy 98.44 keV (P(K α_2)) has been adopted from 1976GuZN. The absolute emission probabilities of all other X-rays have been deduced from their relative emission probabilities using the adopted P(K α_2) = 1.69 (4) $\cdot 10^{-4}$ %. (The uncertainty of this value includes an additional 2 % detector efficiency uncertainty).

The total absolute KX-ray emission probability P(XK) = 3.56 (11) $\cdot 10^{-4}$ %, obtained using P(K α_2) and the ratio of P(XK) / P(K α_2), exceeds the value calculated from ω_K and the total emission probability of K-conversion electrons P^(ce)(XK) = 2.6 $\cdot 10^{-4}$ %. This disagreement may be due to an inaccurate estimation of K-conversion electron intensities from E0 and (E0 + E2) transitions in decay of ²³⁸Pu.

6.2. Gamma-Ray Emissions

6.2.1. Gamma-Ray Energies

The energies of prominent gamma-rays $\gamma_{1,0}$ (43.5 keV), $\gamma_{2,1}$ (99.9 keV) and $\gamma_{3,2}$ (152.7 keV) have been taken from 1984He19, with a correction of 5.8 ppm in the gamma-ray energy scale as provided by 2000He14. The energies of gamma-rays $\gamma_{13,5}$ (258.2 keV) and $\gamma_{5,1}$ (742.8 keV) are from 2000Ni13. The remaining gamma-ray energies have been taken from 2007Br04 based on the measurements of 1969LeZX and also 1954As07, 1955Ch02, 1956Ne17, 1971Cl03, 1971GuZY, 1971Ma68, 1976GuZN, 1984Ov01. Several of gamma-rays were not observed in ²³⁸Pu α -decay and their energies have been taken from the decay of ²³⁴Pa and ²³⁴Np (2007Br04). The experimental and recommended gamma-ray energies are given in Table 9.

Table 9. Experimental and recommended gamma-ray energies (keV) from ²³⁸Pu α decay ^a

	1969LeZX	1971GuZY	1972Sc01	1976GuZN	1984He19	Recommended
$\gamma_{1,0}$		43.492 (10)	43.491 (9)	43.477 (5)	43.498 (1)	43.498 (1)
$\gamma_{2,1}$	99.84 (4)	99.871 (10)	99.85 (1)	99.864 (5)	99.853 (3)	99.852 (3)
$\gamma_{3,2}$	152.71 (5)	152.77 (3)	152.719 (19)	152.68 (2)	152.720 (2)	152.719 (2)
$\gamma_{4,3}$	200.9 (2)	200.98	201.017 (30)	200.98		200.97 (3)
$\gamma_{14,7}$	235.9 (3)					235.9 (3)
$\gamma_{13,5}$	258.3 (2)	258.23				258.227 (3)
$\gamma_{14,5}$	299.2 (2)					299.1 (2)
$\gamma_{7,2}$	706.1 (3)	705.6		705.6		705.9 (1)
$\gamma_{8,2}$	708.4 (2)	708.4		708.4		708.3 (2)
$\gamma_{5,1}$	742.77 (10)	742.82		742.82		742.813 (5)
$\gamma_{6,1}$	766.39 (10)	766.41 (2)		766.41		766.38 (2)
$\gamma_{5,0}$	786.30 (10)	786.30		786.30		786.27 (3)
$\gamma_{7,1}$	805.8 (3)	805.42		805.4		805.80 (5)
$\gamma_{8,1}$	808.25 (15)	808.23		808.2		808.20 (10)
$\gamma_{8,0}$	851.70 (10)	851.73		851.7		851.70 (10)
$\gamma_{12,2}$	880.5 (3)					880.5 (1)

	1969LeZX	1971GuZY	1972Sc01	1976GuZN	1984He19	Recommended
$\gamma_{9,1}$	883.23 (10)	883.21				883.24 (4)
$\gamma_{10,1}$	904.37 (15)	904.34				904.37 (15)
$\gamma_{9,0}$	926.72 (15)	926.73				926.72 (10)
$\gamma_{14,2}$	941.9 (2)	942.02				941.94 (10)
$\gamma_{11,1}$	946.0 (3)	946.12				946.00 (3)
$\gamma_{13,1}$	1001.03 (15)	1001.10				1001.03 (3)
$\gamma_{14,1}$	1041.8 (3)	1041.90				1041.7 (2)
$\gamma_{14,0}$	1085.4 (3)	1085.40				1085.4 (2)

^a Other much more inaccurate measurement results can be found in 1954As07, 1955Ch02, 1956Ne17, 1971Cl03 and 1971Ma68. They agree with those given in Table 9.

6.2.2. Gamma-Ray Emission Probabilities

The experimental and recommended absolute gamma-ray emission probabilities $P(\gamma)$ for prominent γ -rays (with energies < 200 keV) are given in Table 10. The recommended $P(\gamma)$ values have been obtained by averaging several experimental results. They agree well with the values deduced from intensity balances at relevant ²³⁴U levels using $P(\alpha)$ and total ICCs.

Table 10. Experimental and recommended absolute emission probabilities (per 10^4 α -decays) for prominent gamma-rays from the decay of ²³⁸Pu

	E_γ (keV)	1976GuZN	1976Um01	1979 Vaninbr oukx	1984Bo41	1984He19	1984Ov01	1994Ba91	Recommended (averaged) ^a	Deduced ^b
$\gamma_{1,0}$	43.5	3.93 (8)	4.11 (8)	3.93 (12)	3.96 (10)	3.82 (8)			3.97 (8)	4.06 (8)
$\gamma_{2,1}$	99.8	0.724 (14)			0.730 (11)	0.743 (8)	0.631 (38) ^c		0.735 (8)	0.741 (25)
$\gamma_{3,2}$	152.7	0.0956 (20)			0.0928 (14)	0.0936 (10)	0.086 (4) ^c	0.0923(7)	0.0930 (7)	0.095 (4)

^a Weighted averages; uncertainties are the smallest experimental values.

^b Deduced from $P(\alpha)$ values and total ICCs.

^c Omitted based on statistical considerations.

The relative emission probabilities of $\gamma_{14,7}$ (235.9 keV), $\gamma_{13,8}$ (258.2 keV) and $\gamma_{14,5}$ (299.1 keV) have been adopted from 1969LeZX. The absolute emission probability of $\gamma_{10,2}$ (804.4 keV) has been deduced using the ratio of $P(\gamma_{804.4 \text{ keV}}) / P(\gamma_{904.4 \text{ keV}}) = 1.8$ (7) measured in ²³⁴Pa β^- -decay (2007Br04). $P(\gamma)$ values for other gamma-rays, which were also not observed in the ²³⁸Pu α -decay, have been deduced from decay of ²³⁴Pa and ²³⁴Np (2007Br04) using experimental relative gamma-ray emission probabilities.

The absolute emission probabilities of all other weak gamma-rays (with energies more than 200 keV) have been deduced from their evaluated relative emission probabilities given in Table 11.

The value $P(\gamma_{766}) = 2.19$ (5)· 10^{-7} measured in 1976GuZN (the uncertainty includes an additional 2 % detector efficiency uncertainty) was used as a normalization factor. This value agrees well with the value of 2.19 (9)· 10^{-7} , deduced from the measured in 1979Ce04 $P(\gamma_{786}) = 3.16$ (9)· 10^{-8} and the relative intensity $P(\gamma_{786}) / P(\gamma_{766}) = 0.144$ (4), as well as with the value of 2.21 (15)· 10^{-7} measured in 1984Ov01. The latter value has been obtained by the evaluator from authors' P_γ renormalized to $P(\gamma_{152.7\text{-keV}}) = 9.30$ (7)· 10^{-6} .

Table 11. Experimental and recommended relative emission probabilities of gamma-rays with energy more than 200 keV from decay of ²³⁸Pu

		1969LeZX	1971GuZY	1971Ma68	1976GuZN	1979Ce04	1984Ov01	Recommended
$\gamma_{4,3}$	201.0	15 (3)	17.8 (3)		18.6 (4)	17.0 (5)		17.9 (4)
$\gamma_{14,7}$	235.9	0.04 (2)						0.04 (2)
$\gamma_{13,5}$	258.2	0.35 (5)	0.28 (6)					0.32 (5)
$\gamma_{14,5}$	299.1	0.20 (5)						0.20 (5)
$\gamma_{7,2}$	705.9	0.42 (6) ^a	0.225 (23)		0.23 (10)		0.25 (10)	0.23 (5)
$\gamma_{8,2}$	708.3	1.15 (9) ^a	2.24 (23)	2.5 (6)	2.29 (23)	2.5 (6)	1.7 (3)	2.22 (14)
$\gamma_{5,1}$	742.8	23.2 (4)	23.1 (2)	25.7 (15)	23.6 (5)	23.8 (4)	22.6 (12)	23.3 (2)
$\gamma_{6,1}$	766.4	100	100	100	100	100	100	100
$\gamma_{5,0}$	786.3	14.5 (3)	14.7 (2)	14.9 (10)	15.0 (3)	14.4 (4)	13.7 (5)	14.6 (2)
$\gamma_{7,1}$	805.8	0.56 (6)	0.56 (6)		0.59 (3)		0.7 (2)	0.58 (3)
$\gamma_{8,1}$	808.2	3.40 (8)	3.57 (10)	3.2 (5)	3.65 (13)	3.52 (18)	4.0 (4)	3.50 (8)
$\gamma_{8,0}$	851.7	5.79 (20)	5.79 (11)	6.6 (6)	5.89 (17)		4.9 (5)	5.81 (11)
$\gamma_{12,2}$	880.5	0.7 (2)					0.65 (16)	0.68 (16)
$\gamma_{9,1}$	883.2	3.43 (15)	2.72 (27)	3.3 (5)		3.54 (25)	3.2 (6)	3.30 (17)
$\gamma_{10,1}$	904.4	0.30 (4)	0.26 (8)				0.25 (10)	0.28 (5)
$\gamma_{9,0}$	926.7	2.53 (10)	2.56 (10)	2.7 (6)		2.58 (13)	2.4 (3)	2.55 (10)
$\gamma_{14,2}$	941.9	2.06 (9)	2.19 (9)	2.2 (6)		2.23 (27)	1.9 (4)	2.13 (9)
$\gamma_{11,1}$	946.0	0.40 (6)	0.43 (9)					0.42 (6)
$\gamma_{13,1}$	1001.0	4.39 (14)	5.42 (33) ^a	4.0 (7)		4.61 (18)	4.1 (5)	4.46 (14)
$\gamma_{14,1}$	1041.7	0.84 (7)	0.95 (10)	0.7 (3)			1.3 (3)	0.90 (7)
$\gamma_{14,0}$	1085.4	0.34 (4)	0.95 (10) ^a	1.1 (4) ^a			0.5 (2)	0.35 (4)

^a Omitted on the basis of statistical considerations.

7. Consistency of recommended data

The most accurate Q value, Q(M), is taken from the atomic mass adjustment table of Audi et al. (2003Au03). Comparison of Q(eff)(deduced as the sum of average energies per disintegration ($\sum E_i \times P_i$) for all emissions accompanying ²³⁸Pu α - decay) with the tabulated decay energy Q(M) allows to check a consistency of the recommended decay-scheme parameters obtained in this evaluation.

Here E_i and P_i are the evaluated energies and emission probabilities of the i-th alpha particle, beta particle, gamma-ray, X-ray, etc. Consistency (percentage deviation) is determined by $\{[Q(M) - Q(\text{eff})]/Q(M)\} \times 100$. "Percentage deviations above 5 % would be regarded as high and imply a poorly defined decay scheme; a value of less than 5 % indicates the construction of a reasonably consistent decay scheme" (quoted from the article by A.L. Nichols in Appl. Rad. Isotopes 55 (2001) 23-70).

For the above ²³⁸Pu decay data evaluation we have Q(M) = 5593.20 (19) keV and Q(eff) = 5593(5) keV. Thereafter, the percentage deviation is $(0.00 \pm 0.09) \%$, i.e. consistency is superior.

8. References

- 1949Jaffey A. H. Jaffey and A. Hirsch, Report ANL-4286 (1949) [Spontaneous fission half-life].
 1950Jaffey A. H. Jaffey and J. Lerner, Report ANL-4411 (1950) [Half-life].
 1951Jaffey 1 A. H. Jaffey and L. B. Magnusson, Paper No. 14.2. National Nuclear Energy Plutonium Project Record Div. IV. Vol. 14B. Part II. p. 978. McGraw-Hill, New York (1951) [Half-life].
 1951Jaffey-2 A. H. Jaffey, Paper No. 2.2. National Nuclear Energy Plutonium Project Record Div. IV. Vol. 14B. Part I. p. 89. McGraw-Hill, New York (1951) [Half-life].
 1951Seaborg G. T. Seaborg, R. A. James and A. Giorso, The Transuranium Elements (Edited by G. T. Seaborg, J. J. Katz and W. M. Manning). Paper No. 14.2. National Nuclear Energy Series, Plutonium Project Record, Div. IV. Vol. 14B. Part II, p. 978. McGraw-Hill, New York (1951) [Half-life].
 1952Se67 E. Segre, Phys. Rev. 86(1952)21 [Spontaneous fission half-life].

- 1954As07 F. Asaro and I. Perlman, Phys. Rev. 94(1954)381[Alpha-particle energies and emission probabilities].
- 1954Jo10 K. W. Jones, R. A. Douglas, M. T. McEllistrem and H. T. Richards, Phys. Rev. 94(1954)947 [Half-life].
- 1955Ch02 E. L. Church and A. W. Sunyar, Phys. Rev. 98(1955)1186A [Gamma-ray energies].
- 1956Ne17 J. O. Newton, B. Rose and J. Milsted, Phil. Mag. 1(1956)981 [Gamma-ray energies].
- 1957Ho71 D. C. Hoffman, G. P. Ford and F. O. Lawrence, J. Inorg. Nucl. Chem. 5(1957)6 [Half-life].
- 1957Ko33 L. N. Kondratev, G. I. Novikova, V. B. Dedov and L. L. Goldin, Izv. Akad. Nauk SSSR, Ser Fiz 21(1957)907 [Alpha-particle energies and emission probabilities].
- 1961Dr04 V. A. Druin, V. P. Perelygin and G. I. Khlebnikov, Soviet Phys. JETP 13(1961)913; Zhurn. Eksptl. i Teoret. Fiz. 40(1961)1296 [Spontaneous fission half-life].
- 1962Le11 C. F. Leang, Compt. Rend. 255(1962)3155 [Alpha-particle energies and emission probabilities].
- 1963Bj03 S. Bjornholm, C. M. Lederer, F. Asaro and I. Perlman, Phys. Rev. 130(1963)2000 [Alpha transition probabilities].
- 1964Ha14 J. W. Halley, D. Engelkemeir, Phys. Rev. 134(1964)A24 [L X-ray emission probabilities].
- 1964Le17 F. Les, Acta. Phys. Polon. 26(1964)951 [E0+E2 transition probabilities].
- 1964Le22 C. M. Lederer, Priv. Comm. Quoted by 67Le24, unpublished(1964) [E0+E2 transition probabilities].
- 1965Eichelberger J. F. Eichelberger, G. R. Grove and L. V. Jones, MLM-1238(1965) [Half-life].
- 1967Jordan K. C. Jordan, Part of Mound Laboratory Progress Report for Chemistry, Report No. MLM-1443, July–September (1967),p. 11 [Half-life].
- 1968Ba25 S. A. Baranov V. M. Kulakov and V. M. Shatinskii, Nucl. Phys. 7(1968)442.; Yadern. Fiz. 7(1968)727 [Alpha-particle energies and emission probabilities].
- 1968By01 J. Byrne, W. Gelletly, M. A. S. Ross and F. Shaikh, Phys. Rev. 170(1968)80 [L X-ray emission probabilities].
- 1968Salgueiro L. Salgueiro et al., C. R. Acad. Sci. 267B(1968)1293 [L X-ray emission probabilities].
- 1968Swinth K. L. Swinth, Nucleonics in Aerospace, Ed.P. Polishuk, N.Y. Plenum Press, 1968, p. 279 [L X-ray emission probabilities]
- 1969Am02 S. R. Amtey, J. H. Hamilton, A. V. Ramayya et al., Nucl. Phys. A126(1969)201 [Conversion electron relative intensities]
- 1969Benson D. Benson, Priv. Comm. (1969). Cited in 1981Ag06 [Half-life].
- 1969LeZX C. M. Lederer, F. Asaro and I. Perlman, UCRL-18667(1969)3 [Gamma-ray energies and emission probabilities].
- 1970Ba72 S. A. Baranov V. M. Kulakov, V. M. Shatinskii and Z. S. Gladkikh, Soviet J. Nucl. Phys. 12(1971)604; Yad. Fiz. 12 (1970) 1105 [Alpha-particle energies and emission probabilities].
- 1971Cl03 J. E. Cline, IN-1448 Rev. (1971) [Gamma-ray energies and emission probabilities].
- 1971Gr17 B. Grennberg and A. Rytz, Metrologia 7(1971)65 [Alpha-particle energies].
- 1971GuZY R. Gunnink, R. J. Morrow, UCRL-51087(1971) [Gamma-ray energies and emission probabilities].
- 1971Ma68 A. I. Makarenko, L. A. Ostretsov and N. V. Forafontov, Izv. Akad. Nauk SSSR, Ser. Fiz. 35(1971)2335; Bull. Acad. Sci. USSR, Phys. Ser. 35(1972)2118 [Gamma-ray energies and emission probabilities].
- 1971So15 J. C. Soares, J. P. Ribeiro, A. Gonçalves, F. B. Gil and J. C. Ferreira, Compt. Rend. 273B(1971)985 [Alpha-particle energies and emission probabilities].
- 1971Swinth K. L. Swinth, IEEE Transactions Nuclear Science 18(1971)125 [L X-ray emission probabilities]
- 1972Ha11 J. D. Hastings and W. W. Strohm, J. Inorg. Nucl. Chem. 34(1972)25 [Spontaneous fission half-life].
- 1972Sc01 M. Schmorak, C. E. Bemis, Jr., M. J. Zender, N. B. Gove and P. F. Dittner, Nucl. Phys. A178 (1972)410 [Gamma-ray energies].
- 1974StYG W. W. Strohm and K. C. Jordan, Nucl. Soc. 18(1974)185 [Half-life].
- 1975GaZX R. R. Gay and R. Sher, Bull. Am. Phys. Soc. 20(2)(1975)160,GB13 [Spontaneous fission half-life].

- 1976GuZN R. Gunnink, J. E. Evans and A. L. Prindle, UCRL-52139(1976) [Gamma-ray energies and emission probabilities].
- 1976Po08 V. G. Polyukhov, G. A. Timofeev, P. A. Privalova, V. Y. Gabeskiriya and A. P. Chetverikov, Soviet. J. At. Energy 40(1976)66 ; At. Energ. 40(1976)61 [Half-life]
- 1976Um01 H. Umezawa, T. Suzuki and S. Ichikawa, J. Nucl. Sci. Technol. 13(1976)327 [Gamma-ray and emission probabilities].
- 1976Va23 D. G. Vasilik and R. W. Martin, Nucl. Instrum. Methods 135(1976)405 [L X-ray emission probabilities].
- 1977Bemis C. E. Bemis Jr. and L. Tubbs, Report ORNL-5297(1977)93 [L X-ray emission probabilities]
- 1977Di04 H. Diamond, W. C. Bentley, A. H. Jaffey and K. F. Flynn, Phys. Rev. C15(1977)1034 [Half-life].
- 1977La19 F. P. Larkins, At. Data Nucl. Data Tables 20(1977)313 [Auger electron energies]
- 1978Ro22 F. Rosel, H. M. Friess, K. Alder and H. C. Pauli At. Data Nucl. Data Tables 21(1978)92 [Theoretical ICC].
- 1979Vaninbroukx R. Vaninbroukx, G. Grosse and W. Zehner, Report CBNM/RN/45/79(1979) [Gamma-ray emission probabilities].
- 1979Ce04 A. Cesana, G. Sandrelli, V. Sangiust and M. Terrani, Energia Nucl. (Milan) 26(1979)526 [Gamma-ray energies and emission probabilities].
- 1981Ag06 S. K. Aggarwal, A. V. Jadhav, S. A. Chitambar, K. Raghuraman, S. N. Acharya, A. R. Parab, C. K. Sivaramakrishnan and H. C. Jain, Radiochem. Radioanal. Lett. 46(1981)69 [Half-life].
- 1981Sevastyanov V. D. Sevastyanov and V. P. Jarina, Voprosi Atomnoi Nauki i Tekhniki, Seriya Jadernie Konstanti 5(44)(1981)21 [Half-life].
- 1982Ba56 G. Barreau, H. G. Borner, T. von Egidy, R. W. Hoff., Z. Phys. A308(1982)209 [K X-ray energies].
- 1983Ah02 I. Ahmad, J. Hines, J. E. Gindler, Phys. Rev. C27(1983)2239 [K X-ray energies].
- 1984Ah06 I. Ahmad, Nucl. Instrum. Meth. 223(1984)319 [Alpha-particle energies and emission probabilities].
- 1984BaYT L. M. Bak, P. Dryak, V. G. Nedovesov, S. A. Sidorenko, G. E. Shukin, K. P. Yakovlev, Program and Theses, Proc. 34th Ann. Conf. Nucl. Spectrosc. At. Nuclei, Alma-Ata, (1984), p. 541 [L X-ray emission probabilities].
- 1984Bo41 G. Bortels, B. Denecke, R. Valninbroukx, Nucl. Instrum. Meth. 223(1984)329 [Alpha-particle, gamma-ray and L X-ray energies and emission probabilities].
- 1984Burns P.A. Burns, P.N. Johnston and J.R. Moroney, Priv. Comm. (1984).Cited in 1986LoZT: A. Lorenz, IAEA Tech. Report, Ser. No 261(1986)147 [Alpha-particle energies and emission probabilities].
- 1984DrZX P. Dryak, Yu. S. Egorov, V. G. Nedovesov, I. Plkh, G. E. Shukin, Program and Theses, Proc. 34th Ann. Conf. Nucl. Spectrosc. At. Nuclei, Alma-Ata, (1984), p. 540 [L X-ray emission probabilities].
- 1984He19 R. G. Helmer and C. W. Reich, Int. J. Appl. Radiat. Isotop. 35(1984)1067 [Gamma-ray energies and emission probabilities].
- 1984Ov01 V. V. Ovechkin, V. M. Chesalin and I. A. Shkabura, Izv. Akad. Nauk. SSSR, Ser. Fiz. 48(1984)1029 [Gamma-ray energies and emission probabilities].
- 1987Bo25 G. Bortels and P. Collaers, Appl. Radiat. Isot. 38(1987)831 [Alpha-particle energies and emission probabilities].
- 1988SeZY Yu. A. Selitsky, V. B. Funshtein, V. A. Yakovlev, Program and Theses, Proc.38th Ann. Conf. Nucl. Spectrosc. Struct. At. Nuclei, Baku, p. 131(1988) [Spontaneous fission half-life].
- 1990Po14 Yu. S. Popov, I. B. Makarov, D. Kh. Srurov, E. A. Erin, Sov. Radiochem. 32(1990)425; Radiokhimiya 32(1990)2 [M X-ray emission probability].
- 1991Jo02 P. N. Johnston, J. R. Moroney and P. A. Burns, Appl. Radiat. Isot. 42(1991)245 [Alpha-particle energies].
- 1991Ry01 A. Rytz, At. Data Nucl. Data Tables 47(1991)205 [Alpha-particle energies].
- 1994Ba91 D. T. Baran, Appl. Radiat. Isot. 45(1994)1177 [Gamma-ray emission probabilities].
- 1994Le37 M. C. Lépy, B. Duchemin, J. Morel, Nucl. Instrum. Meth. Phys. Res. A353(1994)10 [L X-ray energies and emission probabilities].

Comments on evaluation

- 1995Jo23 P. N. Johnston and P. A. Burns, Nucl. Instrum. Meth. Phys. Res. A361(1995)229 [L X-ray energies and emission probabilities].
- 1996Sc06 E. Schönfeld and H. Janßen, Nucl. Instrum. Meth. Phys. Res. A369(1996)527 [Atomic data].
- 1998Ya17 J. Yang and J. Ni, Nucl. Instrum. Meth. Phys. Res. A413(1998)239 [Alpha-particle energies and emission probabilities].
- 1999Schönfeld E. Schönfeld and G. Rodloff PTB-6.11-1999-1999-1, Braunschweig, February 1999 [K X-ray energies and relative emission probabilities].
- 2000He14 R. G. Helmer and C. van der Leun, Nucl. Instrum. Meth. Phys. Res. A450(2000)35 [Gamma-ray energies].
- 2000Ho27 N. E. Holden, D. C. Hoffman, Pure Appl. Chem. 72(2000)1525 [Spontaneous fission half-life].
- 2000Ni13 Y. Nir-El, Radiochim. Acta 88(2000)83 [Gamma-ray energies].
- 2000Schönfeld E. Schönfeld and H. Janßen. Appl. Rad. Isot. 52(2000)595 [L X-ray and Auger electron emission probabilities].
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)337 [Q value].
- 2007Br04 E. Browne, J. K. Tuli, Nuclear Data Sheets 108 (2007)681 [Level energies and data from ²³⁴Pa and ²³⁴Np decays].
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor, Jr., Nucl. Instrum. Meth. Phys. Res. A589(2008)202 [Theoretical ICC].

²³⁹U – COMMENTS ON EVALUATION OF DECAY DATA

by V.P.Chechev and N.K.Kuzmenko

This evaluation was completed in October 2008 and updated in March 2009 with a literature cut-off by the same date.

1. DECAY SCHEME

Decay scheme is based on 2003Br12. Most (99 %) of ²³⁹U beta decay feeds the well-studied ²³⁹Np levels below 118 keV. However more than 30 excited states of ²³⁹Np have been associated with weak ²³⁹U beta transitions and in this part the decay scheme cannot be considered as completed.

Several unplaced gamma rays were observed in 2006Wo03. These gamma rays carry ≤ 2 % of the total intensity of all the gamma rays placed in the decay scheme.

Wong and Griffin (2006Wo03), based on the energies of many of these gamma rays, suggested different versions of their placement including alternative with respect to 2003Br12. These suggestions require an additional careful analysis. Therefore the evaluators have been accepted only small change in the decay scheme from 2003Br12 associated with moving the 1197 -keV level off and adding the new 849,45-keV level.

The 1197-keV level stated in 2003Br12 has been deleted from the level scheme since the 535 -, 1122- and 1197 - keV gamma transitions previously reported were not observed in 2006Wo03 and attributed to possible impurities. The new (declared in 2006Wo03) 849,45-keV level de-exciting via 502-, 608-, 728-, 775- and 849- keV gamma rays has been placed to the decay scheme.

It should be noted that a number of ²³⁹Np levels reported only from nuclear reactions may be populated (according to the data of 2006Wo03) in ²³⁹U β^- -decay.

Several gamma rays previously reported were not observed with high reliability in 2006Wo03. They were ascribed to fission product impurities in their study.

2. NUCLEAR DATA

Q^- value is from 2003Au03.

The recommended half-life of ²³⁹U is based on the experimental results given in Table 1.

Table 1. Experimental values of ²³⁹U half-life (in minutes)

Reference	Author(s)	Original value	Re-estimated	Measurement method
1943Mi10	Mitchell et al.	23,54 (5)		β -counting
1947Fe05	Feather and Krishnan	23,5 (7)		β - and gamma-ray counting
1969Hu21	Hunt et al.	23,40 (5)		β -counting
1989Ab05	Abzouzi et al.	23,44 (2) ^a	23,44 (11)	Gamma-ray counting
2008Griffin	Griffin	23,37 ^b	23,37 (10)	Liquid scintillation counting

^a Uncertainty may include only statistical errors. The evaluators have taken into account the contribution of possible systematic errors (uncertainty of the Type B) associated with the gamma-ray counting method (see Comments on evaluation of ²³³Th half-life).

^b Author did not report the uncertainty. Possible statistical and systematic errors associated with the used LSC method were discussed in 2008De10 under the measurements of ²³³Th half-life (22 min). The evaluators have estimated an overall relative uncertainty of $\sim 0,4$ % (see Comments on evaluation of ²³³Th half-life).

The unweighted mean of the 6 values from Table 1 is 23,45 (3), the weighted mean is 23,46, the internal uncertainty is 0,032, the external uncertainty is 0,035. The LWEIGHT computer program recommended the weighted mean and its internal uncertainty. The smallest experimental uncertainty is 0,05. Therefore, the recommended value of ²³⁹U half-life is 23,46 (5) minutes.

2.1. Beta Transitions

The energies of β^- transitions have been obtained from the Q^- value and the ²³⁹Np level energies given in Table 2 from 2003Br12.

Table 2. ²³⁹Np levels populated in ²³⁹U β^- -decay

Level	Energy (keV)	Spin and Parity	Half-life	
0	0,0	5/2+	2,356 (3) d	14,4 (22)
1	31,1310 (12)	7/2+		9,4 (15)
2	71,210 (2)	9/2+		
3	74,664 (1)	5/2-	1,39 (3) ns	72,8 (19)
4	117,727 (20)	7/2-	= 40 ps	2,2 (4)
5	122,5 (10)	(11/2+)		
6	173,10 (4)	9/2-		
7	241,36 (5)	(11/2-)		
8	260,799 (17)	(3/2-)		
9	438,83 (5)	(11/2+)		
10	448,178 (16)	(3/2-)		
11	452,736 (2)	(5/2+,7/2-)		
12	474,36 (6)			0,0033 (4)
13	517,998 (20)	(7/2-)		0,063 (2)
14	530,29 (6)			0,0029 (4)
15	563,89 (4)			0,0247 (7)
16	579,40 (4)	(9/2-)		
17	662,282 (17)	(5/2-)		0,261 (6)
18	695,229 (23)	(7/2-)		0,0118 (11)
19	781,93 (4)			
20	784,94 (5)			
21	819,26 (3)	(7/2)		0,228 (3)
22	844,10 (3)	(5/2,7/2)		0,215 (3)
23	849,44 (9)			0,0264 (4)
24	863,46 (6)	(3/2,5/2,7/2)		0,0005 (2)
25	959,18 (3)			0,0284 (7)
26	964,234 (20)	(7/2-)		0,211 (3)
27	966,55 (5)	(7/2,9/2-)		0,0008 (2)
28	992,158 (22)	(7/2-)		0,0262 (9)
29	1013,64 (8)			0,0074 (4)
30	1040,37 (4)	(5/2-,7/2)		0,0077 (4)
31	1049,24 (4)	(9/2-)		0,0059 (4)
32	1096,99 (3)			0,0060 (5)

The emission probabilities of β^- -transitions have been deduced from the $P(\gamma+ce)$ balance at each level of ²³⁹Np. β^- -transitions with $P(\beta) < 0,5\%$ are tentative because of unplaced γ -ray transitions (see 2006Wo03).

2.2. Gamma-ray Transitions and Internal Conversion Coefficients

The recommended energies of the gamma-ray transitions are virtually the same as the gamma-ray energies because nuclear recoil is negligible for ^{239}Np .

The gamma-ray transition probabilities have been obtained from the gamma-ray emission probabilities and the total internal conversion coefficients (ICC). Multipolarities of gamma-ray transitions have been taken from 2003Br12. The ICC have been interpolated using the BrIcc package with the so called “Frozen Orbital” approximation (2008Ki07). The relative uncertainties of the ICC for pure multipolarities have been taken as 2 %.

$P(\gamma_{2,0} + \text{ce})(71,2\text{-keV})$ and $P(\gamma_{7,2} + \text{ce})(170,2\text{-keV})$ have been obtained from the level P($\gamma + \text{ce}$) balance assuming that there is no beta-feeding to the 2- and 7- levels, respectively.

The M1/E2 mixing ratios for $\gamma_{1,0} - 31,1\text{ keV}$ (0,028), $\gamma_{4,3} - 43,1\text{ keV}$ (0,126) and $\gamma_{6,4} - 55,2\text{ keV}$ (0,26) have been taken from ^{243}Am α decay (2003Br12).

The remaining gamma transition multipolarities and M1/E2 mixing ratios have been adopted from ^{239}U β^- -decay (see 2003Br12) based on measurements of 1957Ho07, 1964Bl11, 1969En02.

3. ATOMIC DATA

The atomic data (fluorescence yields, X-ray energies and relative probabilities, and Auger electrons energies and relative probabilities) have been deduced by using the SAISINUC software (2002Be).

4. ELECTRON EMISSIONS

The energies of the conversion electrons have been calculated from the gamma-ray transition energies and the electron binding energies.

The absolute emission probabilities of the conversion electrons have been obtained using the recommended P_γ and ICC values.

The absolute emission probability of K Auger electrons has been deduced from the recommended $\Sigma\text{KX} = 0,305$ (10) %. The absolute emission probability of L Auger electrons has been obtained using the recommended P_γ and ICC values with the EMISSION computer program.

β^- average energies have been calculated using the LOGFT computer program.

5. PHOTON EMISSIONS

5.1. X-Ray Emissions

The recommended absolute emission probabilities of KX-rays have been obtained using the total number of K vacancies of 0,314 (10) % deduced in 2008Griffin from their KX-ray measurements.

The recommended absolute emission probabilities of LX-rays have been obtained using the recommended P_γ , ICC values and the total number of K vacancies with the EMISSION computer program. The calculated total absolute intensity of LX-rays of 16,1 (5) % can be compared with the value of 17 (4) % measured in 2008Griffin (here the author's value of 18,1 % has been corrected to the evaluated $P_{\gamma_{2,0}}(74,7\text{ keV}) = 51,6$ (13) % used instead of 53,9 (5) % measured in 2008Griffin). The uncertainty of the measured LX-ray intensity was not given in 2008Griffin. It has been accepted by the evaluators using the relative uncertainty of the detection efficiency for energies at and below 20 keV ~ 20 % estimated in 2008De10.

5.2. Gamma-Ray Emissions

The gamma ray energies < 120 keV have been obtained from the adopted level energies.

The gamma ray energies > 120 keV have been adopted from 2006Wo03. They agree mainly with the values from 2003Br12 based on experimental data of 1964Bl11, 1969Cl12, 1971Ar47, 1975Pa04, 1979Bo30, 1982Ah04 and data from nuclear reactions. The exceptions comprise the gamma-ray transitions feeding the ^{239}Np ground state and the gamma rays with energies from 2006Wo03 different

from 2003Br12. In such cases the recommended gamma ray energies have been obtained from the adopted level energies.

Several unplaced gamma rays observed in decay of ²³⁹U, α -decay of ²⁴³Am, and the particle transfer reactions were discussed in 2006Wo03 in detail. The transfer reactions and α -spectroscopy give direct information on level energies, but with uncertainties ~ 3 keV. Gamma ray spectroscopy, unsupported by coincidence correlations, gives relatively precise energies, but often placements are ambiguous. Therefore, all such gamma rays have been qualified by the evaluators as unplaced in the decay scheme.

The absolute emission probabilities for most intense gamma rays have been evaluated from experimental data (Table 4). The results of 1984Holloway are superseded by the same group in 1996Sa23 and have not been included in the procedure of averaging.

Table 4. Experimental and evaluated absolute emission probabilities (%) for most intense gamma-rays in decay of ²³⁹U.

E_γ (keV)	1964 Bl11	1965 Yurova	1968 Ma06	1969 Cl12	1984 Holloway	1996 Sa23	2008 Griffin	Evaluated
31,1					0,065 (7)	0,064 (7)	0,075 (4)	0,072 (4)
43,5	4,1 (2)			4,45 (60)	4,18 (13)	4,07 (11)	4,93 (15)	4,35 (28)
74,7		47 (4)	62 (9)	50 (5)	48,2 (10)	49,2 (12)	53,9 (5)	51,6 (13)
86,7				0,060 (6)	0,052 (6)	0,053 (6)	0,054 (5)	0,055 (5)
117,7				0,145 (15)	0,13 (4)	0,14 (3)	0,099 (9)	0,113 (9)

$P_{\gamma_{2,0}}$ (71,2 keV) and $P_{\gamma_{7,2}}$ (170,2 keV) have been obtained from the $P(\gamma+ce)$ and α_T . The value of $P_{\gamma_{4,3}}$ (43,1 keV) has been deduced using the ratio $P_{\gamma_{4,3}} / P_{\gamma_{4,0}} = 0,115$ (12) from 1969En02.

The remaining absolute gamma ray emission probabilities for gamma rays with energy more than 120 keV have been deduced from relative gamma ray emission probabilities P_γ^{rel} (2006Wo03). There the evaluators have used the coefficient $k = P_\gamma^{rel}(74,7 \text{ keV}) / 0,539$ (5) given in 2008Griffin. It was corrected to the evaluated $P_\gamma(74,7 \text{ keV}) = 0,516$ (13) taking also into account the detection efficiency uncertainty (0,7 %): $k = 218,1$ (48). The obtained P_γ agree with the values from 2003Br12 based on experimental data of 1964Bl11, 1965Yurova, 1968Ma06, 1969Cl12, 1971Ar47 and 1984Holloway.

6. CONSISTENCY OF RECOMMENDED DATA

The most accurate Q value, $Q(M)$, is taken from the atomic mass adjustment table of Audi et al. (2003Au03). Comparison of $Q(eff)$ (deduced as the sum of average energies per disintegration ($\sum E_i \times P_i$) for all emissions accompanying ²³³Th β - decay) with the tabulated decay energy $Q(M)$ allows to check a consistency of the recommended decay-scheme parameters obtained in this evaluation.

Here E_i and P_i are the evaluated energies and emission probabilities of the i -th alpha particle, beta particle, gamma ray, X-ray, etc. Consistency (percentage deviation) is determined by $\{|Q(M) - Q(eff)| / Q(M)\} \times 100$. "Percentage deviations above 5 % would be regarded as high and imply a poorly defined decay scheme; a value of less than 5 % indicates the construction of a reasonably consistent decay scheme" (quoted from the article by A.L. Nichols in Appl. Rad. Isotopes 55 (2001) 23-70).

For the above ²³⁹U decay data evaluation we have $Q(M) = 1261,5$ (16) keV and $Q(eff) = 1263$ (36) keV, i.e. consistency is better than 2 %.

7. REFERENCES

- 1943Mi10 A.C.G. Mitchell, L. Slotin, J. Marshall, V.A. Nedzel, L.J. Brown, J.R. Pruett, CP-597 (1943)
(Half-life)
- 1947Fe05 N. Feather, R.S. Krishnan, Proc. Cambridge Phil.Soc. 43, 267 (1947)
(Half-life)
- 1957Ho07 J.M. Hollander, Priv. Comm., quoted by 1960As02 (1960)
(Gamma transition multipolarities)
- 1964B111 K.J. Blinowska, P.G. Hansen, H.L. Nielsen, O. Schult, K. Wien, Nucl. Phys. 55, 331 (1964)
(Gamma transition multipolarities, energies and absolute emission probabilities)
- 1965Yurova L.N. Yurova, A.V. Bushuev, V.G. Bortsov, Soviet. J. At. Energy 18, 75 (1965)
(Gamma ray absolute emission probabilities)
- 1968Ma06 D.R. MacKenzie, R.D. Connor, Nucl. Phys. A108, 81 (1968)
(Gamma ray absolute emission probabilities)
- 1969C112 J.E. Cline, D.A. Tripp, Priv. Comm. (November 1969)
(Gamma ray energies and absolute emission probabilities)
- 1969En02 D. Engelkemeir, Phys.Rev. 181, 1675 (1969)
(Gamma transition multipolarities)
- 1969Hu21 J.B. Hunt, J.C. Robertson, T.B. Ryves, J. Nucl. Energy 23, 705 (1969)
(Half-life)
- 1971Ar47 A. Artna-Cohen, Nucl. Data Sheets B6, 577 (1971)
(Gamma ray energies)
- 1975Pa04 J.C.Pate, K.R.Baker, R.W.Fink, D.A.McClure, N.S.Kendrick, Jr., Z.Phys. A272, 169 (1975)
(Gamma ray energies)
- 1979Bo30 H.G.Borner, G.Barreau, W.F.Davidson, P.Jeuch, T.von Egidy, J.Almeida, D.H.White, Nucl.Instrum.Methods 166, 251 (1979)
(Gamma ray energies)
- 1982Ah04 I.Ahmad, Nucl. Instrum. Methods 193, 9 (1982)
(Gamma ray energies)
- 1984Holloway S.P.Holloway, J.B. Olomo, T.D. MacMahon, B.W. Hooton. Nuclear Data for Science and Technology(K.H. Bockhoff, Ed.) Reidel, Dordrecht (1983) 287, S.P.Holloway, PhD thesis, University of London (1983), T.D. MacMahon, Imperial College at Silwood Park, re-assessment and private communication (1984).Cited in: Decay Data of the Transactinium Nuclides, IAEA, Vienna, Tec. Rep. Ser. 261, 1986
(Gamma ray absolute emission probabilities)
- 1989Ab05 A. Abzouzi, M.S. Antony, V.B. Ndocko Ndongue, J. Radioanal. Nucl. Chem. 135, 1 (1989)
(Half-life)
- 1996Sa23 D. Sardari, T.D. Mac Mahon, S.P. Holloway, Nucl. Instrum. Methods Phys. Res. A369, 486 (1996)
(Gamma ray absolute emission probabilities)

- 2002Be M.M. Bé, R. Helmer, V. Chisté, J. Nucl. Sci. Tech., suppl.2, 481 (2002)
(SAISINUC software)
- 2003Au03 G. Audi, A.H. Wapstra, and C. Thibault, Nucl. Phys. A729, 337 (2003)
(Q value)
- 2003Br12 E. Browne, Nucl. Data Sheets 98, 665 (2003)
(Decay data evaluations, multipolarities, decay scheme)
- 2006Wo03 E.L. Wong and H.C. Griffin, Nucl. Instrum. Methods Phys. Res. A558, 441 (2006)
(Gamma ray emission probabilities and energies)
- 2008Griffin J.H. Hamilton, E.L. Wong, and H.C. Griffin, In: "Fission and properties of neutron
rich nuclei", World Sci. 2008, p.264 (2008)
(X-ray and low energy gamma ray absolute emission probabilities)
- 2008De10 D.J. DeVries and H.C. Griffin, Appl. Rad. Isotop., 66, 1999 (2008)
(Uncertainties of LX-ray absolute emission probabilities)
- 2008Ki07 T. Kibedi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, and C.W. Nestor, Jr.,
Nucl. Instrum. Methods Phys. Res. A589, 202 (2008)
(Theoretical ICC)

**²³⁹Np – Comments on evaluation of decay data
by V.P. Chechev and N.K. Kuzmenko**

This evaluation was completed in June 2006. The literature available by May 2006 was included.

1. Decay Scheme

Decay scheme has been taken from 2003Br12.

2. Nuclear Data

Q⁻ value is from 2003Au03.

The evaluated half-life of ²³⁹Np is based on the experimental results given in Table 1.

Table 1. Experimental values of the ²³⁹Np half-life (in days)

Reference	Author(s)	Value
1956Wi25	Wish	2,346 (4)
1959Co63	Connor and Fairweather	2,34 (2)
1959Co93	Cohen <i>et al.</i>	2,366 (3)
1966Qa01	Qaim	2,354 (8)
1969Bi12	Bigham <i>et al.</i>	2,346 (4)
1990Ab06	Abzouzi <i>et al.</i>	2,3565 (4)

The weighted average of 2,3564 for this discrepant data set of the 6 values is dominated by the very accurate value of 1990Ab06. The LWEIGHT computer program, which uses a limitation of relative statistical weights (LSW method), has increased the 1990Ab06 uncertainty from 0,0004 to 0,0020 and used a weighted average and an external uncertainty having led to 2,356 (3) as a recommended value.

Thus, the adopted value of the ²³⁹Np half-life is **2,356 (3) days**.

2.1. Beta Transitions

The energies of β⁻ transitions have been calculated from the Q⁻ value and the level energies given in Table 2 from 2003Br12 where they have been deduced from a least squares fit to gamma-ray energies (see also 1996FiZX).

Table 2. ²³⁹Np levels populated in the ²³⁹Np β⁻-decay

Level	Energy (keV)	Spin and parity	Half-life	Probability of β ⁻ -transition (%)
0	0	1/2+	24100 (11) a	-
1	7,861 (2)	3/2+	36 (3) ps	6,5 (10)
2	57,276 (2)	5/2+	101 (5) ps	0,4 (72)
3	75,706 (3)	7/2+	83 (8) ps	-
4	163,76 (2)	9/2+	73 (4) ps	-
5	285,460 (2)	5/2+	1,12 (5) ns	43,0 (22)
6	330,125 (4)	7/2+		9,4 (14)
7	387,41 (2)	9/2+		-
8	391,586 (3)	7/2-	193 (4) ns	38,8 (9)
9	469,8 (4)	(1/2-)		0,0027

Level	Energy (keV)	Spin and parity	Half-life	Probability of β^- -transition (%)
10	492,2 (3)	3/2-		0,02
11	505,2	(5/2-)		0,0074
12	511,81 (6)	7/2+		1,56 (16)
13	556,2	(7/2-)		0,0026

The probabilities of β^- -transitions have been deduced from the P(γ +ce) balance for each level of ²³⁹Np. Measured and evaluated β^- -transition probabilities are given in Table 3.

Table 3. Measured and evaluated probabilities (%) of β^- -transitions

	1952Fr25	1956Ba95	1959SCo63	Adopted
$\beta_{0,8}$	52	45	28	38,8 (9)
$\beta_{0,6}$	10	27	13,5	9,4 (14)
$\beta_{0,5}$	31	21	48	43,0 (22)
$\beta_{0,2}$	1,7	}	4	0,4 (72)
$\beta_{0,1}$	4,8	}7	6,5	6,5 (10)

2.2. Gamma-ray Transitions and Internal Conversion Coefficients

The evaluated energies of gamma -ray transitions are virtually the same as the photon energies because nuclear recoil is negligible.

The gamma-ray transition probabilities, P(γ +ce), have been calculated from the gamma -ray emission probabilities and the total internal conversion coefficients (ICC's). Multipolarities of gamma -ray transitions have been taken from 2003Br12 (see also 1996FiZX). ICC's have been interpolated from the BrIcc package. The relative uncertainties of α_K , α_L , α_M , α_T for pure multipolarities have been taken as 2 %. The transition $\gamma_{8,5}$ is anomalously converted, ICC's for this transition have been taken from the measurements of 1959Ew90.

P($\gamma_{1,0}$ +ce)(7,86-keV) has been deduced from the intensity balance for the ground state assuming that there is no beta -feeding to the ''0'' -level. P($\gamma_{3,2}$ +ce) (18,43-keV) has been deduced from the intensity balance for the level ''3'' (75,70-keV) assuming that there is no beta-feeding to the ''3''-level.

The mixing ratios (d) for gamma -ray transitions have been taken from 2003Br12 based on measurements of 1959Ew90, 1972Kr07, 1990Si12 and 1991Sh06.

3. Atomic Data

3.1. Fluorescence yields

The fluorescence yield data are from 1996Sc06 (Schönfeld and Janßen).

3.2. X Radiations

The LX-ray energies are from 1996FiZX. The KX-ray energies and the relative KX-ray emission probabilities are from 1999Schönfeld .

The ratios P(KLX)/P(KLL), P(KXY)/P(KLL) are from 1996Sc06.

4. Electron Emissions

The energies of the conversion electrons have been calculated from the gamma transition energies and the electron binding energies.

The emission probabilities of the conversion electrons have been calculated using evaluated P_γ and ICC values.

The absolute emission probabilities of K and L Auger electrons have been calculated using the EMISSION computer program.

β^- average energies have been calculated using the LOGFT computer program.

5. Photon Emissions

5.1. X-Ray Emissions

The absolute emission probabilities of Pu KX- and LX-rays have been calculated using the EMISSION code.

Measured and calculated absolute emission probabilities of Pu KX-rays are given in Tables 4.

Table 4. Measured and calculated absolute emission probabilities (%) of Pu KX-rays.

	1972Ah02	1982Ah04	Calculated
K α_2 (Pu)	14,4 (6)	12,8 (4)	13,5 (4)
K α_1 (Pu)	22,2 (6)	20,4 (6)	21,4 (6)
K β'_1 (Pu)	-	7,3 (3)	7,84 (25)
K β'_2 (Pu)	2,8 (1)	2,6 (1)	2,72 (10)

5.2. Gamma-Ray Emissions

The gamma ray energies, E_γ , for $\gamma_{1,0}$ (7,86-keV), $\gamma_{2,1}$ (49,4-keV) and $\gamma_{4,2}$ (106,5-keV) were calculated from the level energies. The gamma ray energies with $E_\gamma > 334,3$ keV have been taken from 1974HeYW. The other gamma energies were adopted from 2003Br12 based on experimental data of 1959Ew90, 1965Ma17, 1972Po04, 1979Bo30 and 1982Ah04.

$P(\gamma_{1,0})(7,86\text{-keV})$ has been deduced from $P(\gamma_{1,0} + \text{ce})(7,86\text{-keV})$ and the adopted α_T .

$P(\gamma_{3,2})(18,43\text{-keV})$ has been deduced from $P(\gamma_{3,1})(67,84\text{-keV})$ and the ratio of $P(\gamma_{3,2} + \text{ce})(18,43\text{-keV})/P(\gamma_{3,1})(67,88\text{-keV}) < 0,2$ from 1996FiZX.

$P(\gamma_{2,0})(57,273\text{-keV}) = 0,12$ (3) % has been deduced from $P(\gamma_{2,1})(49,41\text{-keV})$ and $P(\gamma_{2,1})(49,41\text{-keV})/P(\gamma_{2,0})(57,27\text{-keV}) = 0,85$ (12) from 1996FiZX.

$P(\gamma_{7,6})(57,29\text{-keV}) \sim 0,012$ % has been deduced from $P(\gamma_{7,6})(57,3\text{-keV}) + P(\gamma_{2,0})(57,273\text{-keV}) = 0,135$ (7) % and $P(\gamma_{2,0})(57,273\text{-keV})$.

$P(\gamma_{8,6})(61,88\text{-keV})$ and $P(\gamma_{3,1})(67,84\text{-keV})$ have been taken from 1974HeYW.

$P(\gamma_{7,5})(101,96\text{-keV})$ has been taken from ^{239}Am e decay (see 2003Br02).

$P(\gamma_{8,4})(227,83\text{-keV})$ has been taken from the decay scheme (see 2003Br02).

$P(\gamma_{6,1})(322,3\text{-keV})$ has been deduced from the P_γ branching in ^{239}Am e decay and ^{243}Cm a decay (see 2003Br02).

$P(\gamma_{4,3})(88,06\text{-keV})$, $P(\gamma_{4,2})(106,50\text{-keV})$ and $P(\gamma_{6,4})(166,39\text{-keV})$ have been calculated from the conversion data of 1959Ew90 and the adopted α_T .

$P(\gamma_{7,3})(311,70\text{-keV}) = 0,002$ (2) % has been deduced from $P(\gamma_{7,3})(311,70\text{-keV})/P(\gamma_{7,6})(57,29\text{-keV}) = 0,34$ (14) from 1996FiZX.

The absolute emission probabilities of the other gamma-rays have been evaluated from experimental data (Table 5).

6. References

- 1952Fr25 M.S.Freedman, F.Wagner, Jr., D.W.Engelkemeir. Phys.Rev. 88, 1155 (1952)
(Probability of β —transitions)
- 1956Ba95 S.A.Baranov, K.N.Shlyagin, At.Energ.USSR 1, 52 (1956); J.Nuclear Energy 3, 132 (1956)
(Probability of β —transitions)
- 1956Wi25 L.Wish, Nucleonics 14, 105 (1956)
(Half-life)
- 1959Co63 R.D.Connor, I.L.Fairweather, Proc. Phys. Soc.(London) 74, 161 (1959)
(Probability of β —transitions, half-life)
- 1959Ew90 G.T.Ewan, J.S.Geiger, R.L.Graham, D.R.MacKenzie. Phys.Rev. 116, 950 (1959)
(Gamma ray energies)
- 1965Ma17 B.P.K. Maier. Z. Phys. 184, 143 (1965)
(Gamma ray energies)
- 1966Qa01 M. Qaim. Nucl. Phys. 84, 411 (1966)
(Half-life)
- 1969Bi12 C.B.Bigham, Can. J. Phys. 47, 1317 (1969)
(Half-life)
- 1972Po04 F.T. Porter, Phys.Rev. C5, 1738 (1972)
(Gamma ray energies)
- 1972Kr07 L.S.Krane, Phys.Rev. C5, 1671 (1972)
(Gamma transition multipolarities)
- 1972Ah02 I.Ahmad, M.Wahlgren, Nucl. Instrum. Methods 99, 333 (1972)
(Gamma ray absolute emission probabilities)
- 1974HeYW R.L.Heath, ANCR-1000-2 (1974)
(Gamma ray energies and absolute emission probabilities)
- 1974Yu04 L.N.Yurova, A.V.Bushuev, V.I. Petrov, At.Energ. 436, 51 (1974); Sov.At.Energy 36, 52 (1974)
(Gamma ray absolute emission probabilities)
- 1977St35 D.I.Starozhukov, Y.S.Popov, P.A.Privalova , At.Energ. 42, 319 (1977);
Sov.At.Energy 42, 355 (1977)
(Gamma ray absolute emission probabilities)
- 1979Bo30 H.G.Borner, G.Barreau, W.F.Davidson, P.Jeuch, T.von Egidy, J.Almeida, D.H.White,
Nucl.Instrum.Methods 166, 251 (1979)
(Gamma ray energies)
- 1979Mo25 V.K.Mozhaev, V.A.Dulin, Y.A.Kazanskii , At.Energ. 47, 55 (1979);
Sov.At.Energy 47, 566 (1979)
(Gamma ray absolute emission probabilities)
- 1982Ah04 I.Ahmad, Nucl.Instrum.Methods 193, 9 (1982)
(Gamma ray energies and absolute emission probabilities)
- 1984Va41 R.Vaninbroucx, G.Bortels, B.Denecke , Int.J.Appl.Radiat.Isotop. 35, 1081 (1984)
(Gamma ray absolute emission probabilities)
- 1986Ch17 Y.Chang, Z.Cheng, C.Yan, G.Shi, D.Qiao Radiat.Eff. 94, 97 (1986)
(Gamma ray absolute emission probabilities)
- 1990Ab06 A.Abzouzi, M.S.Antony, V.B.Ndocko Ndongue, D.Oster, J.Radioanal.Nucl.Chem. 145, 361 (1990)
(Half-life)
- 1990Si12 E.Simeckova, P.Cizek, M.Finger, J.John, P.Malinsky, V.N.Pavlov, Hyperfine Interactions 59,
185 (1990)
(Gamma transition multipolarities)
- 1991Sh06 Y.Shiokawa, M.Yagi , J.Radioanal.Nucl.Chem. 149, 51 (1991)
(Gamma transition multipolarities, ICC)
- 1991Po17 Yu.S.Popov, D.Kh.Surov, I.B.Makarov, E.A.Erin, G.A.Timofeev. Radiokhimiya 33, 3 (1991);
Sov.J.Radiochemistry 33, 1 (1991)
(Gamma ray absolute emission probabilities)
- 1992Ha02 M.A.Hammed, I.M.Lowles, T.D.Mac Mahon. Nucl.Instrum.Methods Phys.Res. A312, 308 (1992)
(Gamma ray absolute emission probabilities)

Comments on evaluation

- 1996FiZX R.B. Firestone, Table of Isotopes, Eighth Edition, Volume II: A=151-272, V.S. Shirley (Editor), C.M. Baglin, S.Y.F. Chu, and J. Zipkin (Assistant Editors), 1996, 1998, 1999 (LX-energies, gamma ray relative intensities, multipolarities)
- 1996Wo05 S.A.Woods, D.H.Woods, M.J.Woods, S.M.Jerome, M.Burke, N.E.Bowles, S.E.M.Lucas, C.Paton Walsh , Nucl. Instrum. Methods Phys. Res. A369, 472 (1996)
(Gamma ray absolute emission probabilities)
- 1996Sc06 E. Schönfeld and H. Janßen, Nucl. Instrum. Methods Phys. Res. A369, 527 (1996)
(Atomic data)
- 1999Schönfeld E. Schönfeld and G. Rodloff, PTB-6, 11-1999-1999-1, Braunschweig, February (1999)
(KX ray energies and relative emission probabilities)
- 2003Au03 G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys, A729, 337 (2003)
(Q value)
- 2003Br12 E.Browne, Nucl,Data Sheets 98, 665 (2003)
(Gamma ray and level energies, gamma ray multipolarities, decay scheme)

Table 5. Experimental and evaluated absolute emission probabilities (%) for gamma-rays in the decay of ^{239}Np .

E_γ (keV)	1972Ah02	1974Yu04	1974HeYW	1977St35 1991Po17	1979Mo25	1982Ah04	1984Va41	1986Ch17	1986Wo05	1992Ha02	Adopted
44,66						0,13 (1)					0,13 (1)
49,41			0,18 (3)			0,11 (1)					0,145 (35)
57,273						0,135 (7)					0,12 (3)
57,3											~0,012
61,46						1,29 (6)	1,29 (2)		1,40 (7)	1,27 (3)	1,29 (2)
106,12	27,8 (9)			26,6 (10)		26,4 (8)	27,50 (40)	26,08 (38)	25,23 (28)	25,6 (2)	25,9 (3)
181,69	0,075 (8)					0,083 (4)	0,07 (1)		0,085 (5)	0,088 (2)	0,086 (2)
209,75	3,42 (10)			3,36 (14)		3,30 (10)	3,46 (5)	3,28 (5)	3,43 (7)	3,47 (3)	3,42 (3)
226,38				0,24 (3)		0,290 (16)	0,28 (2)		0,230 (14)	0,25 (1)	0,255 (14)
228,18	11,4 (3)			11,78 (44)		11,2 (3)	11,21 (18)	11,05 (14)	10,91 (16)	11,54 (5)	11,32 (22)
254,41	0,11 (1)					0,110 (6)	0,12 (1)		0,1078 (27)	0,113 (4)	0,110 (3)
272,84	0,08 (1)					0,077 (4)	0,08 (1)		0,0762 (24)		0,077 (3)
277,60	14,5 (5)	14,1 (4)		15,0 (4)	14,30 (24)	14,5 (4)	14,38 (21)	14,21 (13)	14,53 (17)	14,46 (10)	14,4 (1)
285,46	0,76 (2)			0,93 (6)		0,790 (25)	0,77 (2)	0,765 (9)	0,797 (10)	0,80 (1)	0,78 (1)
315,88	1,52 (5)			1,63 (7)		1,60 (5)	1,60 (3)	1,55 (2)	1,604 (20)	1,60 (1)	1,59 (1)
334,31	1,95 (7)			2,1 (1)		2,06 (6)	2,08 (3)	1,99 (2)	2,050 (25)	2,05 (2)	2,04 (2)
392,4			0,0016								0,0016
429,5			0,0039								0,0039
434,7			0,013								0,013
447,6			0,00026								0,00026
454,2			0,00082								0,00082
461,9			0,0016								0,0016
469,8			0,0011								0,0011
484,3			0,001								0,001
492,3			0,006								0,006
497,8			0,0032								0,0032
498,7			0,001								0,001
504,2			0,00078								0,00078

**²³⁹Pu – Comments on evaluation of decay data
by V. P. Chechev**

This evaluation was originally done in October 2005 and then revised in January 2007. The literature available by January 2007 has been included.

1. Decay Scheme

The decay scheme is based on the evaluation of Browne (2003Br12). It can be considered as basically completed though there are weak gamma rays observed in experiment and unplaced in the decay scheme. Besides several weak gamma transitions expected from the decay scheme have not been observed in ²³⁹Pu alpha decay yet. They have been taken from data on nuclear reactions, in particular, from ²³⁴U(n,γ)-reaction (1979Al03), and also from ²³⁵Pa β⁻ decay (1986Mi10).

Many alpha transitions to ²³⁵U excited levels with energy more than 600 keV were not observed either. They are expected from data on level spins and gamma rays de-excited these levels (see 2003Br12).

2. Nuclear Data

Q(α) value is from 2003Au03.

The evaluated half-life of ²³⁹Pu is based on the experimental results given in Table 1. Re-estimated values and uncertainties were used for averaging where necessary.

Table 1. Experimental values of the ²³⁹Pu half-life (in years)

Reference	Author(s)	Value	Measurement method
1970OeZZ	Oetting	24 048 (25) ^{a, b}	Calorimetry
1975Al15	Alexandrov <i>et al.</i>	24 060 (19) ^b	Specific activity
1975GlZQ	Glover <i>et al.</i>	24 115 (80)	Specific activity
1977Ja08	Jaffe <i>et al.</i>	24 124 (14)	Specific activity
1977Ja08	Jaffe <i>et al.</i>	24 139 (13)	Mass spectrometry
1978Se12	Seabaugh <i>et al.</i>	24 101 (10) ^b	Calorimetry
1978Gunn	Gunn	24 102 (10) ^b	Calorimetry
1978Lu10	Lucas <i>et al.</i>	24 112 (33) ^c	Specific activity
1978Ma45	Marsch <i>et al.</i>	24 164 (17) ^b	Mass spectrometry
1978Pr07	Prindle <i>et al.</i>	24 019 (15) ^d	Specific activity
1978Pr07	Prindle <i>et al.</i>	24 089 (19) ^d	Mass spectrometry
1981Brown	Brown	24 088 (25) ^b	Specific activity

^a Value corrected in 1977Ja08 is given.

^b Uncertainty quoted by authors for the 95 % confidence level has been reduced by a factor 2.

^c Uncertainty combined from a standard deviation of 16 yr and a systematic error of 50 yr by Holden (1989Ho24) is given.

^d Uncertainty corrected by Holden (1989Ho24) is given.

The weighted mean of the 12 values is 24 100 with the internal uncertainty of 4,5 and external uncertainty of 11 and $\chi^2/\nu = 5,9$. The unweighted mean is 24 097 (12). The LWEIGHT computer program has chosen the weighted mean and the external uncertainty of 11.

Thus, the recommended value of the ²³⁹Pu half-life is 24 100 (11) years. It agrees well with the value of 24 101 (12) years deduced from constant matching in a least-squares fit of thermal data for fissile nuclei (1984Di08) and can be compared to the recommended values from the Russian handbook

(1988ChZL) of 24 100 (20) years and from the critical review by Glover and Nichols (1990GIZZ) of 24 113 (11) years.

The adopted ²³⁹Pu spontaneous fission half-life of $8 (2) \times 10^{15}$ years is the value recommended in 2000Ho27. It is based on the experimental results given in Table 2.

Table 2. Experimental values of the spontaneous fission ²³⁹Pu half-life (in 10^{15} years)

Reference	Author(s)	Value	Measurement method
1952Se67	Segre	5,5 (16)	Ionization chamber
1985Dr09	Druzhinin <i>et al.</i>	7,8 (16)	$\lambda_{SF} / \lambda_{\alpha} = 3,1 (6) \cdot 10^{-12}$

2.1 Alpha Transitions

The energies of the alpha transitions have been deduced from the Q value and the level energies given in Table 3 from 2003Br12. The latter ones were deduced from a least squares fit to γ ray energies from ²³⁹Pu α decay. The energies of the gamma rays adopted from 2003Br12 are given below, in Table 9.

Table 3. ²³⁵U levels populated in the ²³⁹Pu α -decay

Level number	Energy, keV	Spin and parity	Half-life	Probability of α -transition (%)
0	0	7/2-	$7,04(1) \cdot 10^8$ y	$\sim 0,03^b$
1	0,0765 (4)	1/2+	≈ 26 min	70,79 (10)
2	13,0400 (21)	3/2+	0,50(3) ns	17,14 (4)
3	46,207 (10)	9/2-		< 0,02
4	51,7007 (11)	5/2+	191(5) ps	11,87 (3)
5	81,741 (4)	7/2+		0,052 (8)
6	103,035 (10)	11/2-		0,0375 (12)
7	129,2961 (10)	5/2+		0,013 (4)
8	150,467 (15)	9/2+		0,0182 (27)
9	170,708 (14)	13/2-		
10	171,388 (5)	7/2+		0,0034 (10)
11	197,119 (14)	11/2+		0,007 (1)
12	225,423 (8)	9/2+		0,0050 (7)
13	249,130 (12)	15/2-		0,0030 (16)
14	291,144 (19)	11/2+		0,0007 (3)
15	294,669 (15)	13/2+		0,0018 (5)
16	332,845 (4)	5/2+		0,00354 (7)
17	338,52 (6)	17/2-		$\approx 2 \cdot 10^{-5}$
18	357,30 (6) ?	(15/2+)		$1,7 (4) \cdot 10^{-5}$
19	367,069 (8)	7/2+		0,000944 (17)
20	393,225 (6)	3/2+		0,00125 (3)
21	414,779 (11)	9/2+		0,00075 (11)
22	426,755 (3)	5/2+		0,00570 (5)
23	445,716 (20)	7/2+		$4,00 (11) \cdot 10^{-5}$
24	474,297 (13)	7/2+		0,00056 (5)
25	509,92 (17)	(9/2+)		$3,3 (7) \cdot 10^{-6}$
26	533,228 (10)	9/2+		0,00086 (3)
27	608,08 (5)	11/2+		$1,2 (4) \cdot 10^{-5}$
28	633,17 (6)	(5/2)-		$2,84 (7) \cdot 10^{-6}$
29	637,81 (5)	3/2-		$3,22 (21) \cdot 10^{-6}$
30	658,97 (4)	1/2-		$2,64 (6) \cdot 10^{-5}$
31	664,541 (23)	(5/2)-		$6,31 (11) \cdot 10^{-6}$
32	670,99 (4)	(7/2)-		$< 3,4 \cdot 10^{-8}$
33	701,02 (3)	(7/2)-		$7,07 (13) \cdot 10^{-6}$
34	703,757 (19)	3/2-		$1,14 (3) \cdot 10^{-5}$
35	720,25 (3)	(9/2)-		$2,13 (9) \cdot 10^{-6}$
36	750,07 (16)	(9/2)-		$3,4 (4) \cdot 10^{-7}$
37	761,04 (5)	(1/2)-		$1,03 (17) \cdot 10^{-7}$

Level number	Energy, keV	Spin and parity	Half-life	Probability of α -transition (%)
38	769,27 (6)	1/2+		$2,7 (3) \cdot 10^{-5}$
39	769,5 (3)	3/2-		$1,03 (12) \cdot 10^{-5}$
40	777,59 (19)	(11/2)-		$2,47 (19) \cdot 10^{-7}$
41	779,51 (3)	3/2+		$1,01 (11) \cdot 10^{-6}$
42	805,72 (6)	3/2-		$8,4 (14) \cdot 10^{-8}$
43	821,25 (4)	5/2+		$3,0 (3) \cdot 10^{-7}$
44	843,859 (10)	(1/2)+		$2,28 (12) \cdot 10^{-7}$
45	845,3 (10) ?	(7/2+)		$\sim 4,2 \cdot 10^{-8}$
46	865,20 (2) ^a	3/2+		$9,8 (13) \cdot 10^{-8}$
47	891,89 (15)	5/2+		$1,99 (12) \cdot 10^{-7}$
48	968,451 (20)	3/2+		$6,1 (15) \cdot 10^{-8}$
49	970,52 (22) ?	(5/2,7/2)		$4,1 (4) \cdot 10^{-8}$
50	986,65 (17)	(13/2-)		$7,7 (7) \cdot 10^{-8}$
51	992,72 (22)	(5/2+)		$2,0 (3) \cdot 10^{-7}$
52	1057,58 (13)	(7/2)		$9,3 (9) \cdot 10^{-8}$
53	1116,20 (20) ?	(5/2-)		$2,1 (5) \cdot 10^{-8}$

^a Obtained as a sum of E(level '10') and E($\gamma_{46,10}$)

^b Value based on systematics (see 2003Br12 and comments therein)

The probabilities of the most intense transitions $\alpha_{0,1}$, $\alpha_{0,2}$ and $\alpha_{0,4}$ have been obtained by averaging experimental results from measurements with semi-conductor detectors of 1987Bo25, 1992B113, 1993Ga28, 1994Ra27, 1996Sa24, 1996Vi07 and 2002Da21 (see Table 4). They agree with each other and disagree with early measurements with magnetic spectrometers of 1961Dz05, 1963Ba09, 1976BaZZ (Table 4) and 1952As28, 1957As83, 1957No15. The values evaluated from the above experimental results have been recommended as more precise than those that are deduced from γ -ray transition intensity balances.

The probabilities of the transitions $\alpha_{0,k}$ ($k=5\div 8, 10, 13, 15, 16, 19\div 22, 24, 26$) evaluated from all the available experimental data reported with uncertainties are compared in Table 4 with the values deduced from intensity balances. The latter ones were recommended as more precise. The experimental P(α)-values have been recommended in those cases ($\alpha_{0,11}, \alpha_{0,12}, \alpha_{0,14}$) where the intensity balances were used for obtaining P($\gamma+ce$)-values (see several γ -ray transitions with deduced ICC and (E2/M1)-admixture ratios in section 2.2).

The probabilities of the remaining α -transitions including unobserved but expected from the decay scheme have been evaluated from the P($\gamma+ce$) balances for corresponding levels of ^{235}U .

The values of hindrance factors were calculated using ALPHAD code and $r_0(^{235}\text{U}) = 1,5122$, average of $r_0(^{234}\text{U}) = 1,5075$ and $r_0(^{236}\text{U}) = 1,5168$ from 1998Ak04.

Table 4. Experimental and recommended probabilities (%) of most intense α -transitions observed in ²³⁹Pu decay *

	α -part. energy	1961 Dz05	1963Ba09 1976BaZZ	1965 Ho04	1966 Ah02	1987 Bo25	1992 Bl13	1993 Ga28**	1994 Ra27	1996 Sa24	1996 Vi07	2002 Da21**	Evaluated from data of the measurements	Deduced from P(γ +ce) balance	Recommended
$\alpha_{0,1}$	5156	72	73,3 (8)			71,2 (7)	70,73 (46)	70,77 (14)	71,6 (2)	70,91 (11)	71 (5)	70,71 (10)	70,79 (10) ^a	70,8 (4)	70,79 (10)
$\alpha_{0,2}$	5144	17	15,1 (8)			16,7 (5)	17,56 (28)	17,11 (14)	16,6 (2)	17,12 (9)	18 (4)	17,16 (4)	17,14 (4) ^b	17,1 (3)	17,14 (4)
$\alpha_{0,4}$	5106	11	11,5 (8)	11,5		12,1 (2)	11,80 (19)	11,94 (7)	11,8 (1)	11,84 (5)	11,1 (15)	11,88 (3)	11,87 (3) ^c	11,9 (3)	11,87 (3)
$\alpha_{0,5}$	5076	0,038	0,036 (3)	0,043			0,03 (1)	0,078 (8)		0,054 (6)		0,057 (2)	0,050 (7) ^d	0,052 (8)	0,052 (8)
$\alpha_{0,6}$	5055	0,030	0,025 (5)	$\geq 0,0033$				0,047 (13)		0,036 (4)		0,044 (2)	0,038 (4) ^e	0,0375 (12)	0,0375 (12)
$\alpha_{0,7}$	5029		0,005 (1)	0,0038	0,005			0,009 (3)		0,016 (2)		0,023 (1)	0,014 (9) ^f	0,013 (4)	0,013 (4)
$\alpha_{0,8}$	5009	0,018	0,013 (5)	0,011				0,017 (2)		0,021 (6)		0,034 (2)	0,017 (2) ^g	0,0182 (27)	0,0182 (27)
$\alpha_{0,10}$	4988	0,008	0,007 (2)	0,0041	0,006			0,013 (2)				0,018 (1)	0,010 (2) ^h	0,0034 (10)	0,0034 (10)
$\alpha_{0,11}$	4963	0,008	0,006 (3)	0,0044				0,007 (1)				0,0157 (12)	0,007 (1) ^h		0,007 (1)
$\alpha_{0,12}$	4935	0,008	0,0040 (10)	0,0029	0,003			0,0060 (10)				0,0135 (11)	0,0050 (7) ^h		0,0050 (7)
$\alpha_{0,13}$	4912	$\sim 0,003$	0,0005 (3)					0,0024 (9)				0,0097 (9)	0,0007 (3) ^h	0,0030 (16)	0,0030 (16)
$\alpha_{0,14}$	4870		0,0007 (3)									0,0089 (9)	0,0007 (3) ⁱ		0,0007 (3)
$\alpha_{0,15}$	4867	0,004	0,002 (2)	0,0007	0,0008			0,0019 (7)				0,011 (1)	0,0019 (7) ^h	0,0018 (5)	0,0018 (5)
$\alpha_{0,16}$	4829		0,0015	0,0021	0,0021			0,0024 (7)					0,0024 (7)	0,00354 (7)	0,00354 (7)
$\alpha_{0,19}$	4796		0,0007 (2)	0,0008	0,0007			0,0012 (6)					0,0075 (19) ^j	0,000944 (17)	0,000944 (17)
$\alpha_{0,20}$	4770		0,0008 (3)	$\geq 0,001$	0,0006			0,0015 (6)					0,00094 (27) ^j	0,00125 (3)	0,00125 (3)
$\alpha_{0,21}$	4749		$\approx 0,0006$		0,0004							0,0059 (8)	$\approx 0,0005$ ^k	0,00075 (11)	0,00075 (11)
$\alpha_{0,22}$	4737	0,007	0,0045 (10)	0,003	0,005			0,0051 (8)				0,0109 (10)	0,0045 (10) ^h	0,00570 (5)	0,00570 (5)
$\alpha_{0,24}$	4690				0,0005 (2)								0,0005 (2)	0,00056 (5)	0,00056 (5)
$\alpha_{0,26}$	4632				0,0007 (2)								0,0007 (2)	0,00086 (3)	0,00086 (3)

* Other measurements: 1957No15, 1963Bj03, 1981AhZV, 1984Ah06, 1990An33. The 1957No15 results are from measurements with magnetic spectrometer. In 1963Bj03 the $\alpha_{0,30}$ and $\alpha_{0,38}$ probabilities (%) were measured: 0,00008(3) and 0,000025(8), respectively. These values have been adopted as recommended $\alpha_{0,30}$ and $\alpha_{0,38}$ probabilities. The value of α_{30} probability (%) calculated from γ -ray transition intensity balance of 0,000 026 4 (6) disagrees with 1963Bj03 and the calculated value of α_{38} probability (%) of 0,000 027 (4) agrees well with 1963Bj03. In 1984Ah06 the ($\alpha_{0,1} + \alpha_{0,2}$)- probability (%) was measured as 88,0 (6) in agreement with all the available measurements. In 1990An33 the $\alpha_{0,1}$, $\alpha_{0,2}$, $\alpha_{0,4}$ -probabilities (%) were measured: 73 (1), 15 (1), 12 (1), respectively.

** 2002Da21 analyzed α spectrum of 1993Ga28. The values of 1993Ga28 are combined results from measurements at CIEMAT (Spain) and IRMM (Belgium).

^a The LWEIGHT computer program has identified one after another 1996Vi07, 1994Ra27 and 1987Bo25 values as outliers and recommended a weighted average (70,79) of the 4 remaining values and an internal uncertainty of 0,064. The smallest experimental uncertainty of 0,10 is adopted for the evaluated value.

^b The LWEIGHT computer program has identified 1996Vi07 as outlier and (after omitting this value) recommended a weighted average (17,14) of the 6 remaining values and an internal uncertainty of 0,034. The smallest experimental uncertainty of 0,04 is adopted for the evaluated value.

^c The LWEIGHT computer program has identified one after another 1996Vi07 and 1987Bo25 values as outliers and (after omitting these values) recommended a weighted average (11,87) of the 5 remaining values and an internal uncertainty of 0,023. The smallest experimental uncertainty of 0,03 is adopted for the evaluated value.

^d The LWEIGHT computer program has increased the uncertainty of 2002Da21 to 0,00247 and recommended a weighted average (0,050) of the 5 discrepant experimental values (1976BaZZ, 1992B113, 1993Ga28, 1996Sa24, 2002Da21) with the expanded uncertainty of 0,007.

^e The LWEIGHT computer program has increased the uncertainty of 2002Da21 to 0,00304 and recommended a weighted average (0,038) of the 4 experimental values (1976BaZZ, 1993Ga28, 1996Sa24, 2002Da21) with an external uncertainty (0,004).

^f The LWEIGHT computer program has recommended a weighted average (0,014) of the 4 highly discrepant experimental values (1976BaZZ, 1993Ga28, 1996Sa24 and 2002Da21) and expanded the uncertainty to 0,009.

^g A weighted average of the 3 experimental values (1976BaZZ, 1993Ga28, 1996Sa24). The value of 0,034 (2) from 2002Da21 has been omitted as outlier. This big value leads to the appreciable intensity disbalance for the level "8" (150,5 keV).

^h A weighted average of the 2 experimental values (1976BaZZ, 1993Ga28). The value from 2002Da21 has been omitted as this big value leads to the considerable intensity imbalance. Reported experimental data are discrepant.

ⁱ Value from 1976BaZZ. The value from 2002Da21 has been omitted as this big value leads to the considerable intensity imbalance.

^j A weighted average of the values from 1976BaZZ and 1993Ga28.

^k An unweighted average of the values from 1976BaZZ and 1966Ah02. The value from 2002Da21 has been omitted as this big value leads to the considerable intensity imbalance

2.2. Gamma Transitions and Internal Conversion Coefficients

The gamma-ray transition probabilities and total internal conversion coefficients (ICC's) for (M1+E2)-transitions $\gamma_{2,1}$ (12,98 keV), $\gamma_{3,0}$ (46,21 keV), $\gamma_{4,2}$ (38,66 keV), $\gamma_{12,10}$ (54,04 keV), $\gamma_{11,8}$ (46,68 keV) and $\gamma_{14,12}$ (65,71 keV) were deduced from intensity balances for the corresponding levels ("2", "3", "4", "10", "11" and "14", respectively). The total internal conversion coefficients (ICC's) and (E2/M1)-admixture ratios for these transitions were obtained using the α -transition probabilities and γ -ray emission probabilities evaluated from experimental data. For the gamma-ray transition $\gamma_{3,0}$ (46,21 keV) the values of $P(\gamma+ce)$, total ICC and (E2/M1)-admixture ratio have been deduced supposing a negligible intensity of the questionable α -transition to the level "3" (1/2+ \rightarrow 9/2-).

For gamma-ray transition $\gamma_{5,4}$ (30,04 keV) the value $P(\gamma+ce) = 0,033$ (11) % is obtained from the intensity balance for the level "5" by use of the value $P(\alpha_{0,5}) = 0,050$ (7) % evaluated directly from α -spectrometric experimental data. This corresponds to the adopted M1 multipolarity for $\gamma_{5,4}$ -transition: $P(\gamma_{5,4}+ce) = 0,0346$ (14) % has been deduced using the theoretical $\alpha_T(M1) = 58,6$ (12).

The multipolarity of the gamma-ray transition $\gamma_{10,7}$ (41,93 keV) has also been adopted as M1 because even small E2 admixture leads to larger total ICC disturbing $P(\gamma+ce)$ - balance for the level "7" (129,3 keV).

The transition probabilities for the remaining gamma-rays have been deduced from their gamma-ray emission probabilities and total ICC's interpolated from theoretical values of 2002Ba85 using the BrIcc package (Table 11). The multiplicities and admixture coefficients $\delta(E2/M1)$ have been taken from 2003Br12 (see comments therein and in footnotes to Table 11). The uncertainties of α_K , α_L , α_M , α_T for pure multiplicities have been taken as 2 %.

The total ICC for E0+M1 transitions are experimental values from (n, γ) reaction data of 1979Al03 (see 2003Br12 and comments therein).

3. Atomic Data

3.1. Fluorescence yields

The fluorescence yield data are from 1996Sc06 (Schönfeld and Janßen).

3.2. X Radiations

The energies of U LX-rays were deduced from 1994Le28 and 1994Le37 where the fine structure of LX radiation was measured in the decay of ^{239}Pu . Other measurements of U LX-rays can be found in 1983Ah02, 1984Bo41, 1992Ba08 and 1995Jo23.

The U KX-ray energies were taken from 1999ScZX where the calculated values based on X-ray wavelengths from 1967Be65 (Bearde n). In Table 5 the adopted values of U KX-ray energies are compared with experimental values.

Table 5. Experimental and adopted (calculated) values of U KX-ray energies (keV)

	1976GuZN	1982Ba56	1983Ah02	Adopted
$K\alpha_2$	94,655 (5)	94,656 (2)	94,67 (2)	94,666
$K\alpha_1$	98,442 (5)	98,435 (2)	98,45 (2)	98,440
$K\beta_3$	110,42	110,416 (3)	110,42 (3)	110,421
$K\beta_1$	111,30	111,300 (2)	111,31 (2)	111,298
$K\beta_5$	-	111,868 (5)- $K\beta_5$ '' 111,868 (5)- $K\beta_5$	112,01 (5)	111,964
$K\beta_{2,4}$	114,54	-	114,50 (3)	114,46
$KO_{2,3}$	115,40	-	115,40 (5)	115,377

3.3. Auger Electrons

The ratios $P(KLX)/P(KLL)$, $P(KXY)/P(KLL)$ are taken from 1996Sc06.

4. Alpha emissions

The energy of the alpha particles corresponding to the alpha transition to the first excited state of ²³⁵U, $E(\alpha_{0,1})$, has been adopted from the absolute measurement of 1980RyZX taking into account the correction of $-0,11$ keV recommended by A. Rytz in 1991Ry01.

The energies of all other α -emission energies have been deduced from the alpha transition energies taking into account the recoil energies.

In Table 6 the deduced (evaluated) values of α -emission energies are compared with the experimental results obtained with alpha spectrometers.

Table 6. Experimental and evaluated α -emission energies in ²³⁹Pu decay (keV)

	Measured ^a					Recommended in 1991Ry01	Evaluated
	1962Le11	1963Ba09	1966Ho09	1968Ba25	1981AhZV		
$\alpha_{0,1}$	5156,7 (6)	5156,6 (8)	5157	5156,6 (8)		5156,59 (14) ^b	5156,59 (14)
$\alpha_{0,2}$	5144,0 (7)	5144	5144	5144,3 (8)		5144,3 (8)	5143,82 (21)
$\alpha_{0,4}$	5106,0 (7)	5106	5105	5105,8 (8)		5105,8 (8)	5105,81 (21)
$\alpha_{0,5}$		5077	5075		5076 (5)		5076,28 (21)
$\alpha_{0,6}$		5055	5055		5054 (5)		5055,34 (21)
$\alpha_{0,7}$		5030	5029		5028 (3)		5029,51 (21)
$\alpha_{0,8}$		5009	5007		5006 (5)		5008,70 (21)
$\alpha_{0,10}$		4987	4988		4987 (3)		4988,13 (21)
$\alpha_{0,11}$		4962	4960		4960 (5)		4962,83 (21)
$\alpha_{0,12}$		4936	4932		4934 (3)		4935,00 (21)
$\alpha_{0,13}$		4913			4912 (5)		4911,69 (21)
$\alpha_{0,14}$		4872			4871 (5)		4870,38 (21)
$\alpha_{0,15}$		4867	4864		4866 (5)		4866,91 (21)
$\alpha_{0,16}$		4829	4829		4828 (3)		4829,38 (21)
$\alpha_{0,19}$		4800	4794		4795 (4)		4795,73 (21)
$\alpha_{0,20}$			4769		4769 (5)		4770,01 (21)
$\alpha_{0,21}$					4749 (5)		4748,81 (21)
$\alpha_{0,22}$		4738	4739		4736 (3)		4737,05 (21)
$\alpha_{0,24}$		4694	4694		4691 (3)		4690,29 (21)
$\alpha_{0,26}$ ^c		4635	4639		4632 (3)		4632,35 (21)

^a Original values have been adjusted taking into account changes in calibration energies as suggested in 1991Ry01.

^b Absolute measurement; the value has been adopted as recommended in 1991Ry01 (see text above).

^c Other measurements: 1963Bj03, 1975Ba65, 1992Fr04, 1999Sa19. In 1963Bj03 the $\alpha_{0,38}$ and $\alpha_{0,30}$ energies were measured: ≈ 4380 keV and 4510 (20) keV, respectively. In 1975Ba65 the measurement value of the $\alpha_{0,1}$ energy (5156,77 (41) keV) is reported. In 1992Fr04 the $\alpha_{0,1}$ energy was measured by time-of-flight method: 5155,36 (19) keV. In 1999Sa19 alpha peak fitting parameters for analysis of the complex alpha spectrum ²³⁹Pu + ²⁴⁰Pu (keV) were deduced and the following alpha energies were used: $\alpha_{0,1}$ -5156,59; $\alpha_{0,2}$ -5143,90; $\alpha_{0,4}$ -5105,80; $\alpha_{0,5}$ -5076,00.

5. Electron Emissions

The energies of the conversion electrons have been calculated from the gamma transition energies and the electron binding energies. The emission probabilities of conversion electrons have been deduced from the evaluated $P(\gamma)$ and ICC values. The experimental spectrum of the conversion electrons in the decay of ²³⁹Pu is given in 1965Tr03. The conversion electrons were measured also in 1979Al03.

The total absolute emission probability of K Auger electrons has been calculated using the evaluated total emission probability of U KX-rays and the adopted $\omega_K = 0,970$ (4).

The absolute total emission probability of L Auger electrons were computed using the evaluated total absolute emission probability of U LX-rays and the adopted $\omega_L = 0,500$ (19).

6. Photon Emissions

6.1. X-Ray Emissions

6.1.1. LX-Rays

The evaluated absolute emission probabilities of U LX γ -rays have been obtained as weighted means of measurement values from 1992B107 (and 1994Mo36 by the same group), 1994Le28 and 1994Le37 (Table 7). The uncertainties of the evaluated values are not less than the smallest quoted experimental uncertainties.

Table 7. Experimental and evaluated values of absolute LX-ray emission probabilities in the decay of ²³⁹Pu (per 100 disintegrations)

LX-ray	Energy, keV	1992B107, 1994Mo36	1994Le28	1994Le37	Evaluated
L α_1	11,62	0,0996 (11)	0,1027 (21)	0,1016 (17)	0,1008 (11)
L α_2	11,90	-	0,00214 (18)	-	0,00214 (18)
L α_3	13,44	- ^a	0,143 (5)	0,150 (18)	0,146 (13)
L α_4	13,62	- ^a	1,507 (19)	1,498 (31)	1,503 (22)
L η	15,40	0,0566 (10)	0,0498 (10)	0,0544 (9)	0,0537 (19)
L β	17,06	2,301 (23) ^b	2,27 (4) ^b	2,28 (5) ^b	2,288 (23)
L γ	20,30	0,568 (6) ^b	0,564 (10) ^b	0,579 (14) ^b	0,569 (6)
LX total		4,67 (5)	4,63 (5)	4,66 (6)	4,66 (5)

^aIn 1992B107 the total L α -ray intensity of 1,649 (20) was measured in agreement with the value of 1,649 (18) from 1994Le28 and the value of 1,648 (36) from 1994Le37.

^bIn all the three quoted works the intensities of individual L β and L γ components were also measured.

The evaluated P(XL) = 4,66 (5) % exceeds slightly the value of 4,5 (1) % calculated using the evaluated total absolute emission probability of L conversion electrons and the adopted value $\omega_L = 0,500$ (19).

Other measurement results of P(XL) are: 5,3 (5) % (1966Ah02), 4,76 (12) % (1968Swinth), 4,60 (10) % (1971Swinth), 4,50 (14) % (1984Geidelman).

6.1.2. KX-Rays

The evaluated absolute emission probabilities of U KX γ -rays have been obtained as weighted means of measurement values from 1976GuZN and 1994Mo36 (Table 8). Uncertainty in detector efficiency (2 %) was added to the uncertainties listed in 1976GuZN and their values were renormalized to the adopted absolute emission probability of the γ -ray $\gamma_{7,0}$ (129,3 keV) of $6,31 (4) \times 10^{-3}$.

Table 8. Experimental and evaluated values of absolute U KX-ray emission probabilities in the decay of ²³⁹Pu (per 100 disintegrations)

KX-ray	Energy, keV	1976GuZN	1994Mo36	Evaluated
K α_2	94,666	0,004 25 (9)	0,004 17 (4)	0,004 18 (4)
K α_1	98,440	0,006 81 (14)	0,006 52 (9)	0,006 61 (9)
K β_3	110,421	0,000 801 (16)	0,000 797 (6)	0,000 798 (6)
K β_1	111,298	0,001 56 (3)	0,001 536 (12)	0,001 536 (20)
K β_5	111,964	0,000 031 (3)	0,000 054 (11)	0,000 033 (3)
K $\beta_{2,4}$	114,46	0,000 633 (18)	0,000 629 (7)	0,000 629 (7)
K α_{OP}	115,37-115,58	0,000 654 (16)	0,000 708 (9)	0,000 68 (3)
KX total		0,014 74 (29)	0,014 41 (14)	0,014 47 (14)

6.2. Gamma-Ray Emissions

The recommended γ -ray energies have been adopted from 2003Br12 based on experimental data of 1979Al03 ((n, γ)-results) and 1968Cl02, 1971GuZY, 1976GuZN, 1982He02, 1992B107, 1994Mo36 (²³⁹Pu α -decay). Other measurements: 19 65Tr03, 1966Ah02, 1966Ho09 (Table 9). For several weak

Comments on evaluation

transitions γ -ray the energies have been deduced directly from the level energies or adopted from 1979Al03 (see footnotes to Table 9).

The absolute γ -ray emission probabilities have been deduced using the evaluated γ -ray relative probabilities and the absolute emission probability of the γ -ray $\gamma_{7,0}$ (129,3 keV) of $6,31 (4) \times 10^{-5}$ obtained as a weighted average of the 5 absolute measurement results (per 10^5 disintegrations): 6,26 (13) from 1976GuZN, 6,23 (4) from 1980Despres, 6,41 (5) from 1982He02, 6,48 (10) from 1984Iw02 and 6,31 (4) from 1994Mo36. The uncertainty (0,04) of the evaluated value is the smallest experimental uncertainty.

The relative experimental and evaluated γ -ray emission probabilities are given in Table 10. The evaluated values have been obtained by averaging experimental values listed in Table 10 or have been adopted from one of the experimental works, in most cases from 1976GuZN. The averaging -out has been done using the LWEIGHT computer program. The uncertainties are not less than the smallest experimental uncertainties.

In Table 11 the multipolarities, E2/M1 mixing ratios and ICC are shown for soft gamma rays with energy less than 120 keV and comments of deducing multipolarities (with uncertainties for E2/M1 mixing ratios where possible) are given. The δ -mixing ratios for other gamma rays (with energy more than 120 keV) are given in the footnote at the bottom of Table 11.

Table 9. Experimental and adopted energies of gamma rays in ²³⁹Pu decay (keV)

	1965 Tr03	1966 Ah02	1966 Ho09	1968 Cl02	1971 GuZY	1976 GuZN	1979 Al03	1982 He02	1994 Mo36	Adopted
$\gamma_{1,0}$										0,0765 (4)
$\gamma_{2,1}$					13,0				12,975 (10)	12,975 (10)
$\gamma_{-1,1}$									14,22 (3)	14,22 (3)
$\gamma_{5,4}$					30,09	30,04 (10)		30,251 (10)	30,03 (10)	30,04 (2)
$\gamma_{4,2}$		38,7 (1)	37		38,69			38,660 (2)		38,661 (2)
$\gamma_{-1,2}$					40,57	40,41 (5)				40,41 (5)
$\gamma_{10,7}$				41,99 (10)		42,06 (3)			41,93 (5)	41,93 (5)
$\gamma_{3,0}$		46,2 (1)			46,23			46,218 (10)		46,21 (5)
$\gamma_{11,8}$						46,69 (10)			46,68 (3)	46,68 (3)
$\gamma_{7,5}$					47,56				47,60 (3)	47,60 (3)
$\gamma_{4,1}$		51,6 (1)	52		51,628	51,629 (10)	51,628 (4)	51,624 (1)		51,624 (1)
$\gamma_{12,10}$					54,05	54,040	54,026 (5)	54,039 (8)		54,039 (8)
$\gamma_{6,3}$		56,8 (2)			56,828	56,838		56,825 (3)		56,828 (3)
$\gamma_{14,12}$					65,69	65,74 (10)		65,675 (20)		65,708 (30)
$\gamma_{9,6}$					67,69	67,67		67,674 (12)		67,674 (12)
$\gamma_{5,2}$		68,3 (2)	69		68,73	68,72	68,697 (3)	68,696 (6)		68,696 (6)
$\gamma_{8,5}$										68,73 (2) ^b
$\gamma_{-1,3}$										74,96 (10)
$\gamma_{7,4}$		77,6 (2)		77,60 (5)		77,607	77,599 (2)	77,592 (14)		77,592 (14)
$\gamma_{13,9}$				78,48 (5)	78,38	78,42		78,44 (3)		78,43 (2)
$\gamma_{17,13}$										89,39 (6) ^b
$\gamma_{10,5}$				89,59		89,59		89,73 (4)	89,64 (3)	89,64 (3)
$\gamma_{12,7}$						96,13 (5)			96,14 (3)	96,14 (3)
$\gamma_{15,11}$			97,4 (6)		97,6 (3)					97,6 (3)
$\gamma_{8,4}$			98,7 (5)		98,81	98,78 (2)				98,78 (2)
$\gamma_{6,0}$		103,0	102,8 (8)		103,03	103,02 (2)		103,086 (14)		103,06 (3)
$\gamma_{11,5}$			117,6 (11)		115,35	115,38 (5)				115,38 (5)
$\gamma_{7,2}$		116,0			116,24	116,26 (2)	116,262 (3)			116,26 (2)
$\gamma_{10,4}$					119,72	119,708		119,73 (3)	119,70 (3)	119,70 (3)

	1965 Tr03	1966 Ah02	1966 Ho09	1968 Cl02	1971 GuZY	1976 GuZN	1979 Al03	1982 He02	1994 Mo36	Adopted
Y _{14,10}										119,76 (2) ^b
Y _{12,6}				122,35 (12)						122,35 (12)
Y _{37,29}							123,228 (5)			123,228 (5)
Y _{21,14}					123,67	123,62 (5)				123,62 (5)
Y _{9,3}			124,3 (15)		124,52	124,51 (3)				124,51 (3)
Y _{10,3}		125,0			125,17	125,21 (10)				125,21 (10)
Y _{7,0}		129,3 (2)	129,3 (3)		129,28	129,294 (10)	129,302 (2)	129,296 (1)		129,296 (1)
Y _{19,12}		141,7 (3)			141,64	141,657 (20)		141,62 (4)		141,657 (20)
Y _{12,5}					143,4		143,655 (6)			143,35 (20)
Y _{15,8}		144,2	144,1 (8)		144,19	144,211		144,201 (3)		144,201 (3)
Y _{13,6}		146,0			146,05	146,077		146,094 (6)		146,094 (6)
Y _{10,2}					158,3	158,1 (3)				158,1 (3)
Y _{18,11}				159,6 (2)		160,19 (5)				160,19 (5)
Y _{16,10}			160,3 (11)	160,07 (13)	161,45		161,449 (3)	161,482 (12)		161,450 (15)
Y _{17,9}					168,1	167,81 (5)				167,81 (5)
Y _{10,0}		171,4	171,3 (5)		171,34	171,344	171,370 (11)	171,393 (6)		171,393 (6)
Y _{42,28}							172,560 (11)			172,560 (8)
Y _{12,4}					173,6	173,70 (5)				173,70 (5)
Y _{12,3}		179,2 (2)	178,6 (8)		179,17	179,19		179,220 (12)		179,220 (12)
Y _{-1,4}					184,3	184,55 (5)				184,55 (5)
Y _{14,6}					188,27	188,23 (10)				188,23 (10)
Y _{21,12}		189,1	189,2 (16)		189,34	189,32		189,360 (10)		189,360 (10)
Y _{-1,5}				193,13 (12)		193,13 (12)	195,220 (12)			193,13 (12)
Y _{19,10}		195,6	195,7 (8)		195,65	195,66	195,70 (2)	195,679 (8)		195,679 (8)
Y _{-1,6}					197,98	196,87 (5)	196,872 (7)			196,87 (5)
Y _{16,7}		203,5	203,5 (8)	203,34 (8)	203,52	203,537	203,553 (7)	203,550 (5)		203,550 (5)
Y _{21,11}										218,0 (5)
Y _{12,0}			224,9 (15)		225,43	225,37		225,384 (15)		225,42 (4)
Y _{19,7}				238,2 (2)	237,77	237,38	237,774 (6)	237,77 (10)		237,77 (10)
Y _{26,14}			241,2 (20)		242,09	242,08 (3)				242,08 (3)
Y _{21,10}					243,33	243,38		243,38 (3)		243,38 (3)
Y _{14,3}					244,80	244,95 (5)	244,583 (8)			244,92 (5)
Y _{24,12}					248,95	248,95		248,95 (5)		248,95 (5)
Y _{22,10}		255,5	255,1 (5)	258,20 (10)	255,33	255,38		255,384 (15)		255,384 (15)
Y _{20,7}		264,0			263,93	263,93	263,916 (4)	263,97 (3)		263,95 (3)
Y _{30,20}					265,54	265,7 (3)				265,7 (3)
Y _{16,4}					281,2	281,2 (2)				281,2 (2)
Y _{19,5}					285,3	285,3 (2)				285,3 (2)
Y _{22,7}		297,6	297,8 (8)		297,43	297,49	297,42 (3)	297,46 (3)		297,46 (3)
Y _{24,10}					302,87	302,87		302,87 (5)		302,87 (5)
Y _{26,12}					307,81	307,85		307,85 (5)		307,85 (5)
Y _{21,6}		311,8	312,8 (15)		311,69	311,74		311,78 (4)		311,78 (4)
Y _{23,7}					316,35	316,41	316,444 (6)	316,41 (4)		316,41 (3)
Y _{16,2}					319,7	319,68 (10)				319,68 (10)

	1965 TrO3	1966 Ah02	1966 Ho09	1968 Cl02	1971 GuZY	1976 GuZN	1979 Al03	1982 He02	1994 Mo36	Adopted
$\gamma_{19,3}$		321,1			320,8	320,88		320,862 (20)		320,862 (20)
$\gamma_{24,8}$	324	323,9	322,8 (8)		323,76	323,81	323,853 (4)	323,841 (29)		323,84 (3)
$\gamma_{16,0}$	331,1 (5)	333,0	333,2 (5)		332,80	332,838	332,841 (2)	332,845 (5)		332,845 (5)
$\gamma_{26,11}$	336,1 (7)	336,3			336,06	336,107		336,120 (12)		336,113 (12)
$\gamma_{20,4}$	342,6 (7)	341,7	340,0 (20)		341,48	341,510 (2)	341,510 (2)	341,502 (19)		341,506 (10)
$\gamma_{24,7}$										345,001 (13) ^b
$\gamma_{22,5}$	345,6 (7)	345,1 (3)	345,2 (5)		344,96	345,014	345,003 (4)	345,013 (4)		345,013 (4)
$\gamma_{-1,7}$						350,8 (3)				350,8 (3)
$\gamma_{19,2}$					354,1	354,0 (5)				354,0 (5)
$\gamma_{26,10}$	363,5 (10)		363,4 (20)		361,9	361,89		361,90 (6)		361,89 (5)
$\gamma_{19,0}$		367,4			367,02	367,050		367,096 (26)		367,073 (25)
$\gamma_{21,3}$		368,7	369,3 (15)		368,53	368,550		368,557 (27)		368,554 (20)
$\gamma_{22,4}$	375,2 (3)	375,2 (2)	376,3 (5)		375,02	375,042	375,043 (7)	375,054 (3)		375,054 (3)
$\gamma_{20,2}$	380,7 (7)	380,4	381,3 (15)		380,16	380,166	380,173 (3)	380,191 (6)		380,191 (6)
$\gamma_{26,8}$	383,2 (7)	382,9	382,7 (15)		382,72	382,751		382,698 (16)		382,75 (5)
$\gamma_{24,5}$	392,5 (7)				392,45	392,53	392,552 (6)	392,53 (3)		392,53 (3)
$\gamma_{20,1}$	393,4 (7)	393,4 (3)	393,5 (8)		393,06	393,14	393,138 (6)	393,14 (3)		393,14 (3)
$\gamma_{23,3}$					399,44	399,51	399,530 (12)	399,54 (9)		399,53 (6)
$\gamma_{25,6}$	406,2 (5)		408,0 (15)		406,2 (5)	406,9		406,77 (25)		406,8 (2)
$\gamma_{27,11}$					410,77	411,15 (30)				411,2 (3)
$\gamma_{42,20}$										412,49 (6) ^b
$\gamma_{22,2}$	414,0 (3)	413,7	414,2 (5)		413,69	413,712	413,710 (13)	413,713 (5)		413,713 (5)
$\gamma_{24,4}$	422,8 (7)	422,6	423,4 (8)		422,57	422,586	422,596 (8)	422,598 (19)		422,598 (19)
$\gamma_{22,1}$		426,7			426,67	426,68 (8)				426,68 (3)
$\gamma_{24,3}$						428,4 (3)				428,4 (3)
$\gamma_{26,6}$					430,0	430,08 (10)				430,08 (10)
$\gamma_{23,0}$			445,8 (8)		445,78	445,72 (3)	445,740 (17)	445,81 (10)		445,72 (3)
$\gamma_{-1,8}$						446,82 (20)				446,82 (20)
$\gamma_{26,5}$	452,0 (7)	451,6	451,9 (5)		451,45	451,474		451,481 (10)		451,481 (10)
$\gamma_{27,8}$					457,57	457,61 (5)				457,61 (5)
$\gamma_{24,2}$					461,29	461,25 (5)				461,25 (5)
$\gamma_{25,3}$					463,8	463,9				463,9 (3)
$\gamma_{24,0}$					474,4	473,9				473,9 (5)
$\gamma_{26,4}$			480,7 (20)		481,55	481,54		481,78 (12)		481,66 (12)
$\gamma_{26,3}$					487,0	487,06				487,06 (10)
$\gamma_{31,10}$					493,1	493,08 (5)				493,08 (5)
$\gamma_{-1,9}$						497,0				497,0 (5)
$\gamma_{27,5}$						526,4				526,4 (4)
$\gamma_{-1,10}$					538,9	538,8 (2)				538,8 (2)
$\gamma_{33,8}$					550,6	550,5 (2)				550,5 (2)
$\gamma_{-1,11}$					557,7	557,3 (5)				557,3 (5)
$\gamma_{36,10}$						579,4 (3)				579,4 (3)
$\gamma_{31,5}$						582,89	582,75 (8)			582,89 (10)
$\gamma_{29,4}$					586,4	586,3	586,940 (14)			586,3 (3)

	1965 Tr03	1966 Ah02	1966 Ho09	1968 Cl02	1971 GuZY	1976 GuZN	1979 Al03	1982 He02	1994 Mo36	Adopted
Υ _{43,12}						596,0				596,0 (5)
Υ _{33,6}					598,4	597,99 (5)				597,99 (5)
Υ _{36,8}						599,6 (2)				599,6 (2)
Υ _{40,10}					607,3	606,9 (2)				606,9 (2)
Υ _{-1,12}						608,9 (2)				608,9 (2)
Υ _{31,4}					612,9	612,83 (3)	612,838 (6)			612,83 (3)
Υ _{35,6}					617,4	617,10 (10)	617,212 (7)			617,10 (10)
Υ _{31,3}					618,9	618,28 (6)	618,335 (6)			618,28 (6)
Υ _{33,5}						619,21 (6)				619,21 (6)
Υ _{29,2}							624,75 (10)			624,78 (5)
Υ _{32,3}					624,8	624,78 (5)				624,78 (3)
Υ _{28,0}					633,19	633,15 (6)	633,088 (6)			633,15 (6)
Υ _{29,1}										637,73 (5) ^b
Υ _{29,0}			636,0 (30)		637,97	637,84 (6)	637,77 (1)			637,80 (5)
Υ _{38,7}					640,15	640,075		639,99 (10)		639,99 (10)
Υ _{30,2}			645,5 (30)		646,02	645,969	645,894 (5)	645,98 (3)		645,94 (4)
Υ _{33,4}					649,5	649,32 (6)				649,32 (6)
Υ _{-1,13}						650,529 (60)				650,529 (60)
Υ _{34,4}					652,19	652,074	652,052 (5)	651,79 (10)		652,05 (2)
Υ _{33,3}					654,86	654,88 (8)	654,80 (2)			654,88 (8)
Υ _{30,1}					658,99	658,929	658,862 (5)	658,63 (15)		658,86 (6)
Υ _{31,0}					664,67	664,58 (5)	664,520 (12)			664,58 (5)
Υ _{36,5}						668,2 (5)				668,2 (5)
Υ _{43,4}						670,8				670,8 (5)
Υ _{32,0}										670,99 (4)
Υ _{40,6}					674,2	674,05 (3)				674,05 (3)
Υ _{40,5}										674,4 (5)
Υ _{-1,14}					686,16	685,97 (11)	685,861 (6)			685,97 (11)
Υ _{-1,15}						688,1 (3)				688,1 (3)
Υ _{34,2}					690,85	690,81 (8)	690,730 (22)			690,81 (8)
Υ _{-1,16}						693,2 (5)				693,2 (5)
Υ _{46,10}							693,81 (1)			693,81 (1) ^c
Υ _{41,5}						697,8				697,8 (5)
Υ _{-1,17}						699,6 (5)				699,6 (5)
Υ _{33,0}					701,00	701,1 (2)				701,1 (2)
Υ _{34,1}					703,79	703,68 (5)	703,680 (22)			703,68 (5)
Υ _{-1,18}						712,96 (5)				712,96 (5)
Υ _{44,7}						714,71	714,57 (1)			714,71 (14)
Υ _{39,4}					717,76	717,72	718,23 (1)	718,0 (5)		718,0 (5)
Υ _{35,0}						720,3 (5)				720,3 (5)
Υ _{47,10}							720,550 (25)			720,56 (3)
Υ _{41,4}					727,81	727,9	727,860 (25)			727,9 (2)
Υ _{46,7}						736,5	735,910 (15)			736,5 (5)
Υ _{-1,19}						742,7 (5)				742,7 (5)

	1965 Tr03	1966 Ah02	1966 Ho09	1968 Cl02	1971 GuZY	1976 GuZN	1979 Al03	1982 He02	1994 Mo36	Adopted
$\gamma_{37,2}$						747,4	747,97 (1)			747,4 (5)
$\gamma_{38,2}$					}	}756,4 (2)	756,190 (35)			756,23 (6) ^b
$\gamma_{39,2}$			756,0 (30)		}756,40	}	756,87 (6)			756,4 (4)
$\gamma_{47,7}$							762,6 (2)			762,6 (2)
$\gamma_{45,5}$						763,7	763,60 (15)			763,60 (15) ^c
$\gamma_{41,2}$			766,8 (30)			766,6	766,53 (4)			766,47 (3)
$\gamma_{51,12}$							767,29 (4)			767,29 (4)
$\gamma_{38,1}$							769,15 (8)		769,19 (4) ^a	769,15 (8)
$\gamma_{39,1}$					769,38	769,4 (5)	769,59			769,4 (5)
$\gamma_{43,4}$							769,87 (2)			769,54 (4)
$\gamma_{-1,20}$						777,1				777,1 (3)
$\gamma_{41,1}$					779,5	779,61	779,42 (2)			779,43 (3) ^b
$\gamma_{-1,21}$					787,3	786,9 (2)	786,90 (2)			786,9 (2)
$\gamma_{-1,22}$					793,0	788,5 (3)				788,5 (3)
$\gamma_{42,2}$						792,9	792,58 (5)			792,68 (6) ^b
$\gamma_{-1,23}$					796,5	796,9 (3)				796,9 (3)
$\gamma_{-1,24}$					803,3	803,2 (2)				803,2 (2)
$\gamma_{42,1}$						805,9	805,65 (1)			805,65 (6) ^b
$\gamma_{43,2}$					808,2	808,4	808,19 (4)			808,21 (4) ^b
$\gamma_{46,4}$					813,9	813,7	813,510 (17)			813,7 (2)
$\gamma_{50,9}$						816,0 (2)				816,0 (2)
$\gamma_{43,0}$					821,1					821,25 (4) ^b
$\gamma_{51,10}$						821,3 (2)				821,3 (2)
$\gamma_{-1,25}$						826,8 (3)				826,8 (3)
$\gamma_{-1,26}$					828,8	828,9 (2)	828,82 (4)			828,9 (2)
$\gamma_{52,12}$					832,1	832,5				832,2 (2)
$\gamma_{-1,27}$						837,3 (2)				837,3 (2)
$\gamma_{47,4}$					839,0	840,4	840,26 (10)			840,4 (2)
$\gamma_{44,1}$					843,8	844,0	843,78 (1)			843,780 (10)
$\gamma_{47,2}$					879,0	879,2				879,2 (3)
$\gamma_{47,1}$						891,0				891,0 (3)
$\gamma_{-1,28}$						895,4 (3)				895,4 (3)
$\gamma_{-1,29}$						898,1 (3)				898,1 (3)
$\gamma_{-1,30}$						905,5 (3)				905,5 (3)
$\gamma_{-1,31}$						911,7 (3)				911,7 (3)
$\gamma_{49,4}$						918,7 (3)				918,7 (3)
$\gamma_{-1,32}$						931,9 (3)				931,9 (3)
$\gamma_{50,3}$					940,1	940,3 (3)				940,3 (3)
$\gamma_{48,2}$					956,4	955,6	955,390 (21)			955,41 (2) ^b
$\gamma_{49,2}$						957,6 (3)				957,6 (3)
$\gamma_{48,1}$							968,390 (34)			968,37 (2)
$\gamma_{51,2}$					979,5	979,7				979,7 (3)
$\gamma_{-1,33}$						982,7 (3)				982,7 (3)
$\gamma_{53,7}$					986,7	986,9	986,920 (35)			986,92 (4) ^c

	1965 Tr03	1966 Ah02	1966 Ho09	1968 Cl02	1971 GuZY	1976 GuZN	1979 Al03	1982 He02	1994 Mo36	Adopted
$\gamma_{51,1}$					992,5	992,7	992,639 (33)			992,64 (3) ^c
$\gamma_{52,4}$					1005,5	1005,7				1005,7 (3)
$\gamma_{-1,34}$						1009,4 (3)				1009,4 (3)
$\gamma_{52,0}$					1057,3					1057,3 (2)

^a Measured in 1980Despres

^b Obtained as a level energy difference

^c Adopted from 1979Al03

Table 10. Experimental and evaluated relative emission probabilities of gamma rays in decay of ²³⁹Pu &

	Energy, keV	1966 Ah02	1976 GuZN	1980 Despres	1982 He02	1984 Iw02	1992 Bl07	1994 Mo36	Evaluated
$\gamma_{1,0}$	0,077								~0,00016 ^a
$\gamma_{2,1}$	12,98						540 (14)	540 (14)	540 (14)
$\gamma_{-1,1}$	14,22							87 (6)	87 (6) [*]
$\gamma_{5,4}$	30,04		3,47 (13)		15,4 (4)			4,4 (13)	3,47 (13)
$\gamma_{4,2}$	38,66	152 (15)	168 (4)		157,0 (4)		165,8 (24)	165,5 (21)	166 (3)
$\gamma_{-1,2}$	40,41		2,58 (26)						2,58 (26) [*]
$\gamma_{10,7}$	41,93		2,64 (10)		4,07 (10)			2,31 (24)	2,59 (12)
$\gamma_{3,0}$	46,21	16 (2)	11,8 (12)		14,6 (7)			11,43 (17)	11,5 (2)
$\gamma_{11,8}$	46,68		0,93 (6)		1,2 (1)			0,74 (4)	0,80 (9)
$\gamma_{7,5}$	47,60							0,99 (4)	0,99 (4)
$\gamma_{4,1}$	51,62	410 (40)	431 (9)		422 (3)		434 (6)	431 (4)	427 (3)
$\gamma_{12,10}$	54,04		3,19 (8)		3,01 (7)			3,08 (4)	3,08 (4)
$\gamma_{6,3}$	56,83	16 (2)	18,0 (4)		17,4 (4)			18,26 (21)	18,0 (2)
$\gamma_{14,12}$	65,71		0,72 (4)		0,72 (6)			0,82 (5)	0,75 (4)
$\gamma_{9,6}$	67,67		2,57 (7)		2,70 (11)			2,40 (4)	2,50 (8)
$\gamma_{5,2}$	68,70	}14 (2)	8,15 (18)		7,9 (2)			7,69 (10)	5,7 (16) ^b
$\gamma_{8,5}$	68,73	}							2,1 (10) ^b
$\gamma_{-1,3}$	74,96								0,60 (10) ^{c *}
$\gamma_{7,4}$	77,59	11,2	6,23 (13)		6,8 (2)			6,02 (8)	6,08 (9)
$\gamma_{13,9}$	78,43		2,43 (6)		2,1 (2)			2,44 (4)	2,43 (4)
$\gamma_{17,13}$	89,39								~0,03 ^d
$\gamma_{10,5}$	89,64				0,47 (8)			0,43 (3)	0,43 (3)
$\gamma_{12,7}$	96,14		0,36 (7)					0,60 (3)	0,60 (3)
$\gamma_{15,11}$	97,6								1,4 (10) ^{e, a}
$\gamma_{8,4}$	98,78		19,5 (7)					23,2 (11)	21,4 (18)
$\gamma_{6,0}$	103,06		3,47 (9)					3,42 (9)	3,44 (9)
$\gamma_{11,5}$	115,38		7,27 (18)						7,3 (8) ^f
$\gamma_{7,2}$	116,26		9,54 (24)					8,99 (17)	9,2 (3)
$\gamma_{10,4}$	119,70		}0,479 (14)		}0,53 (2)			0,479 (29)	0,33 (4) ^g
$\gamma_{14,10}$	119,76		}		{				0,15 (2) ^{g, i}
$\gamma_{12,6}$	122,35		0,05 (3)					0,015 (2)	0,015 (2) ⁱ
$\gamma_{37,29}$	123,23								0,000025 (6) ^h
$\gamma_{21,14}$	123,62		0,315 (20)					0,376 (14)	0,376 (14)
$\gamma_{9,3}$	124,51		0,98 (4)					1,08 (3)	1,08 (3)
$\gamma_{10,3}$	125,21		1,13 (3)					0,892 (24)	0,892 (24)
$\gamma_{7,0}$	129,30	100	100	100	100	100		100	100

	Energy, keV	1966 Ah02	1976 GuZN	1980 Despres	1982 He02	1984 Iw02	1992 Bl07	1994 Mo36	Evaluated
$\gamma_{19,12}$	141,66	0,6 (1)	0,511 (15)	0,45 (7)	0,46 (8)	0,63 (18)			0,509 (15)
$\gamma_{12,5}$	143,35		0,276 (12)	0,45	}4,80 (9)	}4,75 (13)			0,276 (12)
$\gamma_{15,8}$	144,20	5 (1)	4,52 (10)	4,75 (24)	}	}			4,52 (10)
$\gamma_{13,6}$	146,09	2,1 (2)	1,90 (4)	1,80 (18)	2,00 (10)	1,91 (10)			1,91 (4)
$\gamma_{10,2}$	158,1		0,0160 (16)						0,0160 (16)
$\gamma_{18,11}$	160,19		0,099 (20)						0,099 (20) ⁱ
$\gamma_{16,10}$	161,45		1,92 (4)	2,00 (12)	1,96 (4)	1,91 (10)			1,94 (10)
$\gamma_{17,9}$	167,81		0,047 (12)						0,047 (12)
$\gamma_{10,0}$	171,39	1,8 (2)	1,76 (5)	1,69 (10)	1,74 (4)	1,70 (9)			1,74 (4)
$\gamma_{42,28}$	172,56								~0,00005 ^h
$\gamma_{12,4}$	173,70		0,049 (12)						0,049 (12)
$\gamma_{12,3}$	179,22	1,2 (2)	1,05 (3)	1,04 (8)	1,04 (3)	1,00 (5)			1,04 (3)
$\gamma_{-1,4}$	184,55		0,034 (10)						0,034 (10) *
$\gamma_{14,6}$	188,23		0,174 (18)						0,174 (18)
$\gamma_{21,12}$	189,36	1,5 (2)	1,33 (4)	1,33 (12)	1,30 (2)	1,28 (3)			1,30 (2)
$\gamma_{-1,5}$	193,13		0,142 (15)						0,142 (15) *
$\gamma_{19,10}$	195,68	1,9 (2)	1,70 (4)	1,64 (11)	1,68 (3)	1,68 (4)			1,68 (3)
$\gamma_{-1,6}$	196,87		0,059 (7)						0,059 (7) *
$\gamma_{16,7}$	203,55	9 (1)	8,95 (18)	8,94 (42)	8,90 (13)	8,95 (14)			8,93 (13)
$\gamma_{21,11}$	218,0								0,019 (16) ⁱ
$\gamma_{12,0}$	225,42		0,249 (11)	0,22 (2)	0,23 (2)	0,23 (2)			0,238 (11)
$\gamma_{19,7}$	237,77		0,230 (10)	0,23 (2)		0,32 (2)			0,230 (10)
$\gamma_{26,14}$	242,08		0,117 (8)	}	}	}			0,117 (8)
$\gamma_{21,10}$	243,38		0,404 (11)	}0,41	}0,38 (3)	}0,61 (4)			0,404 (11)
$\gamma_{14,3}$	244,92		0,081 (8)	}	}	}			0,081 (8)
$\gamma_{24,12}$	248,95		0,115 (12)	0,112 (11)	0,11 (1)	0,106 (20)			0,111 (10)
$\gamma_{22,10}$	255,38	1,6 (2)	1,29 (4)	1,27 (10)	1,27 (3)	1,23 (3)			1,26 (3)
$\gamma_{20,7}$	263,95	0,6 (1)	0,417 (15)	0,40 (4)	0,42 (4)	0,39 (3)			0,411 (15)
$\gamma_{30,20}$	265,7		0,025 (6)						0,025 (6)
$\gamma_{16,4}$	281,2		0,035 (5)	0,033 (10)		0,025 (13)			0,034 (5)
$\gamma_{19,5}$	285,3		0,030 (6)	0,03					0,030 (6)
$\gamma_{22,7}$	297,46	0,9 (1)	0,802 (23)	0,77 (8)	0,78 (2)	0,77 (2)			0,78 (2)
$\gamma_{24,10}$	302,87		0,081 (7)	0,070 (12)	0,075 (10)	0,074 (12)			0,077 (7)
$\gamma_{26,12}$	307,85		0,088 (6)	0,076 (12)	0,08 (2)	0,073 (12)			0,083 (6)
$\gamma_{21,6}$	311,78	0,5 (1)	0,412 (12)	0,39 (4)	0,40 (3)	0,36 (8)			0,408 (12)
$\gamma_{23,7}$	316,41		0,217 (8)	0,21 (4)	0,20 (4)	0,196 (14)			0,211 (8)
$\gamma_{16,2}$	319,7		0,077 (8)			}0,85 (2)			0,077 (8)
$\gamma_{19,3}$	320,86	0,8 (1)	0,856 (19)	0,86 (8)	0,86 (3)	}			0,856 (19)
$\gamma_{24,8}$	323,84	0,9 (1)	0,866 (19)	0,82 (8)	0,84 (2)	0,81 (2)			0,84 (2)
$\gamma_{16,0}$	332,85	8 (1)	8,08 (16)	7,64 (32)	7,70 (11)	7,64 (11)			7,74 (11)
$\gamma_{26,11}$	336,11	1,8 (2)	1,81 (4)	1,72 (13)	1,73 (4)	1,75 (4)			1,76 (4)
$\gamma_{20,4}$	341,51	1,2 (1)	1,058 (22)	1,05 (10)	1,00 (4)	1,02 (2)			1,03 (2)
$\gamma_{24,7}$	345,00	}	}						<0,8 ⁱ
$\gamma_{22,5}$	345,013	}8,7 (9)	}8,93 (18)	8,75 (30)	8,67 (13)	8,61 (11)			8,69 (11)
$\gamma_{-1,7}$	350,8		0,028 (6)						0,028 (6) *
$\gamma_{19,2}$	354,0		0,012 (5)						0,012 (5)
$\gamma_{26,10}$	361,89		0,195 (11)	0,18 (2)	0,22 (2)	0,17 (1)			0,185 (11)

	Energy, keV	1966 Ah02	1976 GuZn	1980 Despres	1982 He02	1984 Iw02	1992 BI07	1994 Mo36	Evaluated
$\gamma_{19,0}$	367,07	1,6 (2)	1,38 (3)	1,38 (6)	1,44 (3)	1,35 (2)			1,38 (3)
$\gamma_{21,3}$	368,55	1,4 (2)	1,44 (3)	1,39 (6)	1,37 (3)	1,38 (2)			1,39 (2)
$\gamma_{22,4}$	375,05	25 (3)	25,1 (5)	24,9 (8)	24,2 (3)	24,2 (3)			24,4 (3)
$\gamma_{20,2}$	380,19	5 (1)	4,87 (10)	4,78 (26)	4,75 (7)	4,77 (6)			4,78 (6)
$\gamma_{26,8}$	382,75	4 (1)	4,13 (8)	4,08 (32)	4,02 (6)	4,04 (5)			4,05 (5)
$\gamma_{24,5}$	392,53		}8,83 (18)	}8,72 (35)	}8,55 (13)	1,91 (25)			1,91 (25)
$\gamma_{20,1}$	393,14	10 (1)	}	}	}	6,64 (26)			6,64 (26)
$\gamma_{23,3}$	399,53		0,097 (4)		0,09 (1)	0,103 (17)			0,097 (4)
$\gamma_{25,6}$	406,8		0,010 (4)		0,046 (11)				0,010 (4)
$\gamma_{27,11}$	411,2		0,11 (5)						0,11 (5)
$\gamma_{42,20}$	412,49					}23,0 (3)			-0,00029 ^j
$\gamma_{22,2}$	413,71	25 (3)	23,8 (5)	23,8 (8)	23,0 (3)	}			23,2 (3)
$\gamma_{24,4}$	422,60	2,0 (3)	1,90 (4)	1,91 (14)	1,88 (4)	1,90 (3)			1,90 (3)
$\gamma_{22,1}$	426,68	0,3 (1)	0,372 (9)	0,36 (4)		0,42 (2)			0,379 (9)
$\gamma_{24,3}$	428,4		0,0160 (16)						0,0160 (16)
$\gamma_{26,6}$	430,1		0,069 (3)	0,068 (7)		0,065 (6)			0,068 (3)
$\gamma_{23,0}$	445,72		0,139 (4)	0,146 (15)		0,13 (11)			0,139 (4)
$\gamma_{-1,8}$	446,8		0,0135 (20)						0,0135 (20) *
$\gamma_{26,5}$	451,48	3,4 (5)	3,02 (7)	3,08 (19)	2,96 (4)	2,93 (4)			2,96 (4)
$\gamma_{27,8}$	457,61		0,0238 (5)	0,026 (3)		0,023 (6)			0,0239 (5)
$\gamma_{24,2}$	461,25		0,0363 (8)						0,0363 (8)
$\gamma_{25,3}$	463,9		0,0044 (5)						0,0044 (5)
$\gamma_{24,0}$	473,9		0,0009 (5)						0,0009 (5)
$\gamma_{26,4}$	481,7		0,0735 (15)	0,077 (8)		0,069 (4)			0,0731 (15)
$\gamma_{26,3}$	487,1		0,042 (3)						0,042 (3)
$\gamma_{31,10}^?$	493,08		0,0139 (5)	0,014 (2)		0,013 (3)			0,0139 (5)
$\gamma_{-1,9}$	497,0		0,0007 (4)						0,0007 (4) *
$\gamma_{27,5}$	526,4		0,0009 (3)						0,0009 (3)
$\gamma_{-1,10}$	538,8		0,0049 (3)						0,0049 (3) *
$\gamma_{33,8}$	550,5		0,0067 (4)	0,0074 (8)		0,0079 (31)			0,0069 (4)
$\gamma_{-1,11}$	557,3		0,0006 (3)						0,0006 (3) *
$\gamma_{36,10}$	579,4		0,0014 (3)						0,0014 (3)
$\gamma_{31,5}$	582,9		0,0098 (4)						0,0098 (4)
$\gamma_{29,4}$	586,3		0,00244 (25)						0,00244 (25)
$\gamma_{43,12}$	596,0		0,00062 (19)						0,00062 (19)
$\gamma_{33,6}$	597,99		0,0267 (10)	0,032 (3)		0,030 (3)			0,0275 (10)
$\gamma_{36,8}$	599,6		0,0032 (4)						0,0032 (4)
$\gamma_{40,10}$	606,9		0,00192 (20)						0,00192 (20)
$\gamma_{-1,12}$	608,9		0,00185 (19)						0,00185 (19) *
$\gamma_{31,4}$	612,83		0,0151 (8)	0,025		0,016 (4)			0,0151 (8)
$\gamma_{35,6}$	617,10		0,0214 (12)	}0,08 (1)	}0,09 (1)	}0,069 (5)			0,0214 (12)
$\gamma_{31,3}$	618,28		0,0326 (12)	}	}	}			0,0326 (12)
$\gamma_{33,5}$	619,21		0,0193 (12)						0,0193 (12)
$\gamma_{29,2}$	624,78		0,0073 (3) }						0,0073 (3) ^k
$\gamma_{32,3}$	624,78		}						<0,0003 ^k
$\gamma_{28,0}$	633,15		0,0404 (9)	0,043 (4)		0,036 (3)			0,0404 (9)
$\gamma_{29,1}$	637,73		}0,0409 (10)	}0,047 (5)		}0,047 (4)			0,0101 (10) ^k

	Energy, keV	1966 Ah02	1976 GuZN	1980 Despres	1982 He02	1984 Iw02	1992 Bl07	1994 Mo36	Evaluated
$\gamma_{29,0}$	637,80		}	}		}			0,0304 (30) ^k
$\gamma_{38,7}$	639,99		0,131 (3)	0,139 (14)	0,16 (2)	0,142 (5)			0,134 (3)
$\gamma_{30,2}$	645,94		0,238 (5)	0,25 (3)	0,21 (2)	0,236 (6)			0,236 (5)
$\gamma_{33,4}$	649,32		0,0114 (8)						0,0114 (8)
$\gamma_{-1,13}$	650,53		0,0043 (7)						0,0043 (7) [*]
$\gamma_{34,4}$	652,05		0,105 (3)	0,105 (11)	0,125 (15)	0,102 (5)			0,105 (3)
$\gamma_{33,3}$	654,88		0,0359 (8)	0,029 (7)		0,023 (5)			0,0359 (8)
$\gamma_{30,1}$	658,86		0,155 (4)	0,159 (16)	0,125 (14)	0,150 (5)			0,152 (4)
$\gamma_{31,0}$	664,58		0,0265 (6)	0,027 (3)		0,026 (3)			0,0265 (6)
$\gamma_{36,5}$	668,2		0,00063 (19)						0,00063 (19)
$\gamma_{43,4}^?$	670,8		}0,00014 (4)						<0,00014 (4) ^{l,i}
$\gamma_{32,0}^?$	670,99		}						<0,00014 (4) ^{l,i}
$\gamma_{40,6}$	674,05		0,0082 (3)	}0,0096 (10)		0,0080 (3)			0,0080 (3) ^k
$\gamma_{40,5}$	674,4			}					0,0016 (2) ^k
$\gamma_{-1,14}$	685,97		0,0199 (5)	0,0158 (16)		0,023 (4)			0,0199 (5) [*]
$\gamma_{-1,15}$	688,1		0,00177 (18)						0,00177 (18) [*]
$\gamma_{34,2}$	690,81		0,0089 (5)	0,0104 (10)		0,014 (3)			0,0093 (7)
$\gamma_{-1,16}$	693,2		}0,00080 (24)						0,0005 (2) ^{g,*}
$\gamma_{46,10}$	693,81		}						0,0003 (1) ^g
$\gamma_{41,5}$	697,8		0,00117 (24)						0,00117 (24)
$\gamma_{-1,17}$	699,6		0,00126 (25)						0,00126 (25) [*]
$\gamma_{33,0}$	701,1		0,0082 (3)	0,0095 (10)		0,0106 (34)			0,0083 (3)
$\gamma_{34,1}$	703,68		0,063 (2)	0,067 (7)		0,070 (4)			0,065 (2)
$\gamma_{-1,18}$	712,96		0,00082 (10)						0,00082 (10) [*]
$\gamma_{44,7}$	714,7		0,00125 (13)						0,00125 (13)
$\gamma_{39,4}$	718,0		0,0438 (9)	0,048 (5)		0,042 (3)			0,0438 (9)
$\gamma_{35,0}$	720,3		}0,00078 (8)						0,00046 (5) ^g
$\gamma_{47,10}$	720,56		}						0,00032 (3) ^g
$\gamma_{41,4}$	727,9		0,00198 (11)						0,00198 (11)
$\gamma_{46,7}$	736,5		0,00048 (14)						0,00048 (14)
$\gamma_{-1,19}$	742,7		0,00060 (18)						0,00060 (18) [*]
$\gamma_{37,2}$	747,4		0,00129 (26)						0,00129 (26)
$\gamma_{38,2}$	756,23		}0,0554 (11)	}0,061 (6)		}0,054 (4)			0,044 (8) ^g
$\gamma_{39,2}$	756,4		}	}		}			0,011 (3) ^g
$\gamma_{47,7}$	762,6								~0,00016 ^g
$\gamma_{45,5}$	763,60		0,00052 (26)						0,00035 ^g
$\gamma_{41,2}$	766,47		}0,00439 (24)						0,0021 (3) ^g
$\gamma_{51,12}$	767,29		}						0,0022 (5) ^{g,1}
$\gamma_{38,1}$	769,15		}0,179 (4)	}0,20 (2)		}0,187 (5)			0,081 (16) ^g
$\gamma_{39,1}$	769,4		}	}		}			0,108 (19) ^g
$\gamma_{43,4}$	769,54		}	}		}			- ^m
$\gamma_{-1,20}$	777,1		0,00044 (11)						0,00044 (11) [*]
$\gamma_{41,1}$	779,43		0,00217 (14)						0,00217 (14)
$\gamma_{-1,21}$	786,9		0,00138 (14)						0,00138 (14) [*]
$\gamma_{-1,22}$	788,5		0,00056 (11)						0,00056 (11)
$\gamma_{42,2}$	792,68		0,00032 (6)						0,00032 (6)
$\gamma_{-1,23}$	796,9		0,00024 (5)						0,00024 (5) [*]

	Energy, keV	1966 Ah02	1976 GuZN	1980 Despres	1982 He02	1984 Iw02	1992 Bl07	1994 Mo36	Evaluated
$\gamma_{1,24}$	803,2		0,00102 (7)						0,00102 (7) *
$\gamma_{42,1}$	805,65		0,00044 (7)						0,00044 (7)
$\gamma_{43,2}$	808,21		0,00193 (10)						0,00193 (10)
$\gamma_{46,4}$	813,7		0,00072 (7)						0,00072 (7)
$\gamma_{50,9}$	816,0		0,00039 (6)						0,00039 (6)
$\gamma_{43,0}$	821,25		}0,00088 (9)						0,00079 (17) ⁿ
$\gamma_{51,10}$	821,3		}						-0,00009 ⁿ
$\gamma_{1,25}$	826,8		0,00029 (10)						0,00029 (10) *
$\gamma_{1,26}$	828,9		0,00212 (13)						0,00212 (13) *
$\gamma_{52,12}$	832,2		0,00047 (6)						0,00047 (6)
$\gamma_{1,27}$	837,3		0,00031 (6)						0,00031 (6) *
$\gamma_{47,4}$	840,4		0,00077 (8)						0,00077 (8)
$\gamma_{44,1}$	843,78		0,00214 (12)						0,00214 (12)
$\gamma_{47,2}$	879,2		0,00058 (6)						0,00058 (6)
$\gamma_{47,1}$	891,0		0,00119 (13)						0,00119 (13)
$\gamma_{1,28}$	895,4		0,00012 (4)						0,00012 (4) *
$\gamma_{1,29}$	898,1		0,00028 (6)						0,00028 (6) *
$\gamma_{1,30}$	905,5		0,00012 (4)						0,00012 (4) *
$\gamma_{1,31}$	911,7		0,00022 (5)						0,00022 (5) *
$\gamma_{49,4}$	918,7		0,00014 (5)						0,00014 (5)
$\gamma_{1,32}$	931,9		0,00020 (7)						0,00020 (7) *
$\gamma_{50,3}$	940,3		0,00079 (8)						0,00079 (8)
$\gamma_{48,2}$	955,41		0,00049 (5)						0,00049 (5)
$\gamma_{49,2}$	957,6		0,00051 (5)						0,00051 (5)
$\gamma_{48,1}$	968,37								-0,00044 ^h
$\gamma_{51,2}$	979,7		0,00044 (7)						0,00044 (7)
$\gamma_{1,33}$	982,7		0,00017 (4)						0,00017 (4) *
$\gamma_{53,7}$	986,92		0,00033 (7)						0,00033 (7)
$\gamma_{51,1}$	992,64		0,00042 (6)						0,00042 (6)
$\gamma_{52,4}$	1005,7		0,00028 (4)						0,00028 (4)
$\gamma_{1,34}$	1009,4		0,00022 (4)						0,00022 (4) *
$\gamma_{52,0}$	1057,3								0,00071 (11) ^j

[&] Other measurements for some γ rays: 1965Tr03, 1966Ho09, 1968Cl02, 1971GuZY, 1981UmZZ, 1992Ba08, 1992Co10, 1997Bu23, 1997Ko52.

* Unplaced in level scheme.

^a Deduced from P(γ +ce) and total ICC.

^b Intensity suitably divided for doublet in 2003Br12 (see comments therein).

^c From 1971GuZY. Reported also in Coulomb excitation, see comments in 2003Br12.

^d Intensity suitably divided for doublet in 2003Br12 using systematics.

^e Seen in conversion electron spectrum only (1965Tr03).

^f From 1976GuZN and corrected for X-ray component in 2003Br12.

^g Intensity suitably divided for doublet in 2003Br12 based on (n, γ) data (1979Al03).

^h From (n, γ) data (1979Al03). See 2003Br12.

ⁱ Placement of this transition in the level scheme is uncertain (2003Br12).

^j From 2003Br12.

^k Intensity suitably divided for doublet in 1996Firestone.

^l Multiply placed, undivided intensity given.

^m E0-transition.

ⁿ Possible doublet (see 2003Br12); multiply placed.

Table 11. Energies, multiplicities, E2/M1 mixing ratios and ICC for soft gamma rays (< 120 keV) in decay of ^{239}Pu *

Energy (keV)	Multiplicity	δ -mixing ratio	K	L1	L2	L3	L	M	TOT
0,0765 (4)	E3								
12,975 (10)	M1+0,19 (2) %E2 ^a	0,0436 (23) ^a						451 (13)	607 (17) ^a
14,22 (3)									
30,04 (2)	(M1) ^a			104,9 (21)	12,42 (25)	0,687 (14)	118,0 (24)	28,7 (6)	157 (3)
38,661 (2)	M1+22,2 (16) %E2 ^a	0,534 (24) ^a		42,3 (8)	110 (7)	96 (7)	249 (14)	67 (4)	339 (19)
40,41 (5)									
41,93 (5)	(M1) ^a			39,3 (8)	4,66 (9)	0,249 (5)	44,2 (9)	10,71 (21)	58,6 (12)
46,21 (5)	M1+1,8 (5) %E2 ^a	0,134 (19) ^a		29,0 (6)	7,0 (11)	3,3 (9)	39,4 (19)	9,8 (5)	52,6 (27) ^a
46,68 (3)	M1+9 (5) %E2 ^a	0,32 (9) ^a		26,6 (12)	21 (10)	16 (9)	63 (17)	17 (5)	86 (24) ^a
47,60 (3)	(M1)			27,0 (5)	3,22 (6)	0,170 (3)	30,4 (6)	7,37 (15)	40,4 (8)
51,624 (1)	E2			4,20 (8)	120,4 (24)	101,8 (20)	226 (5)	62,6 (13)	310 (6)
54,039 (8)	M1 ^a			18,6 (4)	2,22 (4)	0,1154 (23)	21,0 (4)	5,08 (10)	27,8 (6)
56,828 (3)	M1+5,0 (8) %E2 ^b	0,23 (2) ^b		15,4 (3)	5,7 (7)	3,3 (5)	24,3 (11)	6,14 (30)	32,6 (15)
65,708 (30)	M1+4 (6) %E2 ^a	0,21 (16) ^a		10,1 (7)	2,8 (29)	1,35 (24)	14 (5)	3,6 (13)	19 (6) ^a
67,674 (12)	M1+3,63 (11) %E2 ^b	0,194 (3) ^b		9,33 (19)	2,34 (5)	1,01 (3)	12,7 (4)	3,15 (9)	16,9 (5)
68,696 (6)	E2			1,19 (24)	31,6 (6)	24,5 (5)	57,3 (11)	15,9 (3)	78,6 (16)
68,73 (2)	(M1+20 %E2) ^c	0,5 ^c		7,6	7,2	4,9	20	5,2	27
74,96 (10)									

Energy (keV)	Multipolarity	δ -mixing ratio	K	L1	L2	L3	L	M	TOT
77,592 (14)	M1(+20 (32) %E2) ^d	0,5 (5) ^d		5,3 (2)	4 (5)	2,7 (40)	12 (7)	3,2 (21)	17 (10)
78,43 (2)	M1(+20 (32) %E2) ^d	0,5 (5) ^d		5,2 (17)	4 (5)	2,6 (40)	12 (7)	3,1 (20)	16 (10)
89,39 (6)	[M1]			4,28 (9)	0,519 (10)	0,0253 (5)	4,82 (10)	1,167 (23)	6,40 (13)
89,64 (3)	(M1+E2)						11 (6)	2,8 (17)	14 (8)
96,14 (3)	[E2]			0,318 (6)	6,72 (14)	4,63 (9)	11,67 (23)	3,24 (7)	16,0 (3)
97,6 (3)	M1+20 (19) %E2 ^d	0,5 (3) ^d		2,71 (6)	1,6 (11)	0,9 (8)	5,2 (14)	1,3 (4)	7,0 (19)
98,78 (2)	E2			0,289 (6)	5,94 (12)	4,05 (8)	10,28 (21)	2,85 (6)	14,1 (3)
103,06 (3)	E2			0,250 (5)	4,90 (10)	3,29 (7)	8,44 (17)	2,34 (5)	11,58 (23)
115,38 (5)	E2			0,172 (3)	2,95 (6)	1,88 (4)	5,00 (10)	1,39 (3)	6,87 (14)
116,26 (2)	M1(+24 (36) %E2) ^d	0,56 (56) ^d	8,4 (18)	1,5 (6)	0,9 (9)	0,5 (6)	2,9 (6)	0,74 (16)	12,2 (26)
119,70 (3)	(M1+E2)		5 (5)				3,1 (11)	0,8 (3)	9 (4)
119,76 (2)	[E2]		0,200 (4)	0,154 (3)	2,49 (5)	1,57 (3)	4,22 (8)	1,169 (23)	5,99 (12)

* For gamma rays with energies more than 120 keV the multiplicities are taken from 2003Br12 based on conversion electron data of 1965Tr03, experimental (n, γ) results of 1979Al03 or assigned from the decay scheme (in square brackets). The δ -mixing ratios are: 1,0 (10) for $\gamma_{26,5}$ (451,5 keV), < 1 for $\gamma_{40,10}$ (606,9 keV), < 0,5 for $\gamma_{28,0}$ (633,2 keV), 1,2 (2) for $\gamma_{46,7}$ (736,5 keV), 0,6 (2) for $\gamma_{46,7}$ (955,4 keV) and 0,6 (3) $\gamma_{46,7}$ (968,4 keV).

^a Deduced from intensity balance.

^b From muonic ²³⁵U atom.

^c From systematics.

^d From conversion electron data of 1965Tr03.

7. References

- 1952As28 F. Asaro, I. Perlman, Phys.Rev. 88, 828 (1952)
(α -transition energies and probabilities)
- 1952Se67 E. Segre, Phys.Rev. 86, 21 (1952)
(SF half-life)
- 1957As83 F.Asaro, S.G.Thompson, F.S.Stephens, Jr., I.Pperlman, Priv.Comm., quoted in 1964Hy02 (1964)
(α -transition energies and probabilities)
- 1957No15 G.I.Novikova, L.N.Kondratev, Y.P.Sobolev, L.L.Goldin, Zhur.Eksptl.i Teoret.Fiz. 32, 1018 (1957);
Soviet Phys.JETP 5, 832 (1957)
(α -transition energies and probabilities)
- 1961Dz05 B.S.Dzhelepov, R.B.Ivanov, V.G.Nedovesov, Zhur.Eksptl.i Teoret.Fiz. 41, 1725 (1961); Soviet
Phys.JETP 14, 1227 (1962)
(α -transition energies and probabilities)
- 1962Le11 C.F.Leang, Compt.Rend. 255, 3155 (1962)
(α -transition energies)
- 1963Ba09 S.A.Baranov, V.M.Kulakov, S.N.Belenky, Nucl.Phys. 41, 95 (1963)
(α -transition energies and probabilities)
- 1963Bj03 S. Bjornholm, C.M. Lederer, F. Asaro, I. Perlman, Phys. Rev. 130, 2000 (1963)
(α -transition energies and probabilities)
- 1965Ho04 F.Horsch, Z.Physik 183, 352 (1965)
(α -transition energies and probabilities)
- 1965Tr03 E.F. Tretyakov, L.N. Kondratev, Bull. Acad. Sci. USSR, Phys. Ser. 29, 243 (1966); Izv Akad Nauk
SSSR, Ser Fiz 29, 242 (1965)
(Gamma-ray and conversion electron energies and emission probabilities)
- 1966Ah02 I. Ahmad, In: Thesis, Univ California (1966); UCRL-16888
(α -transition and gamma-ray energies and emission probabilities)
- 1966Ho09 F.Horsch, Z.Physik 194, 405 (1966)
(Gamma-ray energies and emission probabilities)
- 1967Be65 J.A. Bearden, Rev. Mod. Phys. 39 (1967) 78.
(X-ray energies)
- 1968Ba25 S.A.Baranov, V.M.Kulakov, V.M.Shatinskii, Yadern.Fiz. 7, 727 (1968); Soviet J.Nucl.Phys. 7, 442
(1968)
(α -transition energies)
- 1968Cl02 J.E. Cline, Nucl. Phys. A106, 481 (1968)
(Gamma-ray energies and emission probabilities)
- 1968Swinth K.L. Swinth, Nucleonic in Acrospace, Ed. P.Polyshuk.N.Y.:Plenum Press, p. 279, (1968). Quoted in
1988ChZL
(LX-ray emission probabilities)
- 1970OeZZ F.L.Oetting, Proc.Int.Conf.on Plutonium and Other Actinides, 4th, Santa Fe, New Mexico,
M.A.Musil, Ed., The Metallurgical Soc., New York, Pt.1, p.154 (1970)
(Half-life)
- 1971GuZY R. Gunnink, R.J. Morrow, In: UCRL 51087 (1971).
(Gamma ray energies and emission probabilities)
- 1971Swinth K.L. Swinth, IEEE Trans. Nucl. Sci. 18(1), 125 (1971). Quoted in 1988ChZL
(LX-ray emission probabilities)
- 1975Al15 B.M.Aleksandrov, V.T.Antsiferov, L.S.Bulyanitsa, A.M.Geidelman, Y.S.Egorov, L.M.Krizhanskii,
A.A.Lipovskii, V.G.Nedovesov, L.D.Preobrazhenskaya, L.A.Razumovskii, V.M.Smirnov,
Y.V.Kholnov, Y.L.Chereshevich, V.I.Sharalapov, G.E.Shchukin, K.I.Yakovlev, G.M.Yanchilenko,
Izv.Akad.Nauk SSSR, Ser.Fiz. 39, 482 (1975); Bull.Acad.Sci.USSR, Phys.Ser. 39, No.3, 20 (1975)
(Half-life)
- 1975Ba65 S.A.Baranov, V.M.Shatinsky, Yad.Fiz. 22, 670 (1975); Sov.J.Nucl.Phys. 22, 346 (1976)
(α -transition energies)
- 1975GIZQ K.M.Glover, R.A.P.Wiltshire, F.J.G.Rogers, M.King, UKNDC(75)-P-71, p.55 (1975)
(Half-life)

- 1976BaZZ S.A. Baranov, A.G. Zelenkov, V.M. Kulakov, In: Proc Advisory Group Meeting on Transactinium Nucl. Data, Karlsruhe; Vol III, (1976)249; IAEA-186.
(α -transition probabilities)
- 1976GuZN R. Gunnink, J.E. Evans and A.L. Prindle, UCRL-52139 (1976).
(Gamma-ray energies and emission probabilities)
- 1977Ja08 A.H.Jaffey, H.Diamond, W.C.Bentley, K.F.Flynn, D.J.Rokop, A.M.Essling, J.Williams, Phys.Rev. C16, 354 (1977)
(Half-life)
- 1978Gunn S.R. Gunn, Int.J.Appl.Radiat.Isotop. 29, 497 (1978)
(Half-life)
- 1978Lu10 L.L.Lucas, J.R.Noyce, B.M.Coursey, Int.J.Appl.Radiat.Isotop. 29, 501 (1978)
(Half-life)
- 1978Ma45 S.F.Marsh, R.M.Abernathey, R.J.Beckman, R.K.Zeigler, J.E.Rein, Int.J.Appl.Radiat.Isotop. 29, 509 (1978)
(Half-life)
- 1978Pr07 A.Prindle, J.Evans, R.Dupzyk, R.Nagle, R.Newbury, Int.J.Appl.Radiat.Isotop. 29, 517 (1978)
(Half-life)
- 1978Se12 P.W.Seabaugh, K.C.Jordan, Int.J.Appl.Radiat.Isotop. 29, 489 (1978)
(Half-life)
- 1979Al03 J.Almeida, T.von Egidy, P.H.M.van Assche, H.G.Borner, W.F.Davidson, K.Schreckenbach, A.I.Namenson, Nucl.Phys. A315, 71 (1979)
(Gamma-ray and conversion electron energies)
- 1980Despres M. Despres, Rep. CEA-R-5065 (1980), quoted in:Decay Data of the Transactinium Nuclides, IAEA, Vienna, Technical Reports, Ser. No.261, 1986
(Gamma-ray energies and emission probabilities)
- 1980RyZX A. Rytz, In: Proc Intern Conf Atomic Masses and Fundamental Constants, 6th, East Lansing (1979), J A Nolen,Jr, W Benensen Eds, Plenum Press, New York, p 249, (1980).
(Absolute α -particle energy measurement)
- 1981AhZV I.Ahmad, INDC(NDS)-126/NE, p.28 (1981)
(α -transition energies and probabilities)
- 1981Brown F. Brown, Priv. Comm., 1981. Quoted in N.E. Holden, BNL-NCS-35514, p.1, (1984). See Nucl. Stand. Ref. Data, IAEA- TECDoc-335, Vienna, (1985)
(Half-life)
- 1981UmZZ H. Umezawa, In: INDC(NDS)-126/NE, 38 (1981)
(Gamma-ray emission probabilities)
- 1982Ba56 G.Barreau, H.G.Borner, T.von Egidy, R.W.Hoff, Z.Phys. A308, 209 (1982)
(KX-ray energies)
- 1982He02 R.G.Helmer, C.W.Reich, R.J.Gehrke, J.D.Baker, Int.J.Appl.Radiat.Isotop. 33, 23 (1982)
(Gamma-ray energies and emission probabilities)
- 1983Ah02 I. Ahmad, J. Hines, J.E. Gindler, Phys. Rev. C27 (1983) 2239.
(KX-ray energies)
- 1984Ah06 I. Ahmad, Nuclear Instrum. Methods 223 (1984) 319.
(α -transition probabilities)
- 1984Bo41 G.Bortels, B.Denecke, R.Vaninbroukx, Nucl.Instrum.Methods 223, 329 (1984)
(U LX-ray energies)
- 1984Di08 M.Divadeenam, J.R.Stehn, Ann.Nucl.Energy 11, 375 (1984)
(Half-life)
- 1984Geidelman A.M. Geidelman, P. Dryak, Yu. S. Egorov et al. In: Proc. II Int. Symp. "Methods of Production and Measurement of Standard Sources and Solutions", Chopak, Hungary, vol. II, p. 381, (1984).
Quoted in 1988ChZL.
(U LX-ray energies)
- 1984Iw02 Y. Iwata, Y. Yoshizawa, T. Suzuki, S. Ichikawa, S. Okazaki, Intern. J. Appl. Radiat. Isotop. 35(1984)1.
(Gamma-ray emission probabilities)
- 1985Dr09 A.A.Druzhinin, V.N.Polynov, A.M.Korochkin, E.A.Nikitin, L.I.Lagutina, At.Energ. 59, 68 (1985);
Sov.At.Energy 59, 628 (1985)
(SF Half-life)

- 1986Mi10 S.Mirzadeh, Y.Y.Chu, S.Katcoff, L.K.Peker, Phys.Rev. C33, 2159 (1986)
(²³⁵U level energies, ²³⁵Pa β⁻ decay)
- 1987Bo25 G.Bortels, P.Collaers, Appl.Radiat.Isot. 38, 831 (1987)
(α-transition probabilities)
- 1988ChZL V.P.Chechev, N.K.Kuzmenko, V.O.Sergeev, K.P.Artamonova. Evaluated Decay Data of
Transuranium Radionuclides, Handbook, Publishing House Energoatomizdat, Moscow (1988)
(Evaluation of ²³⁹Pu decay data)
- 1989Ho24 N.E.Holden, Pure Appl.Chem. 61, 1483 (1989)
(Half-life)
- 1990An33 S.V.Anichenkov, Yu.S.Popov, Radiokhimiya 32, 109 (1990); Sov.J.Radiochemistry 32, 401 (1991)
(α-transition probabilities)
- 1990GIZZ K.M.Glover, A.L.Nichols, AERE-R-13822 (1990)
(Half-life)
- 1991Ry01 A. Rytz, At. Data Nucl. Data Tables. 47 (1991) 205.
(α-transition energies and probabilities)
- 1992Ba08 G. Barci-Funel, J. Dalmaso, G. Ardisson, Appl. Radiat. Isot. 43, 37 (1992)
(LX and gamma-ray emission probabilities)
- 1992Bl07 C.J. Bland, J. Morel, E. Etcheverry, M.C. Lepy, Nucl. Instrum. Methods Phys. Res. A312, 323 (1992)
(LX and gamma-ray emission probabilities)
- 1992Bl13 C.J. Bland, J. Truffy, Appl. Radiat. Isot. 43, 1241 (1992)
(α-transition probabilities)
- 1992Co10 N.Coursol, N.Coron, D.Masse, H.Stroke, J.W.Zhou, P.de Marcillac, J.Lebianc, G.Artzner,
G.Dambier, J.Bouchard, G.Jegoudez, J.P.Lepeltier, G.Nollez, C.Golbach, J.-L.Piccolo,
Nucl.Instrum.Methods Phys.Res. A312, 24 (1992)
(Gamma-ray emission probabilities)
- 1992Fr04 E.A. Frolov, Appl. Radiat. Isot. 43, 211 (1992)
(α-transition energies)
- 1993Ga28 E.Garcia-Toraño, M.L.Acena, G.Bortels, D.Mouchel, Nucl.Instrum.Methods Phys.Res. A334, 477
(1993)
(α-transition probabilities)
- 1993Sc22 M.R. Schmorak, Nucl. Data Sheets 69, 375 (1993)
(Decay Scheme)
- 1994Ba91 D.T.Baran, Appl.Radiat.Isot. 45, 1177 (1994)
(α-transition probabilities)
- 1994Le28 M.C.Lépy and K. Debertain, Nucl.Instrum.Methods Phys.Res. A339, 218 (1994)
(LX-ray emission probabilities)
- 1994Le37 M.C.Lépy, B.Duchemin, J.Morel, Nucl.Instrum.Methods Phys.Res. A353, 10 (1994)
(LX-ray emission probabilities)
- 1994Mo36 J. Morel, E. Etcheverry, M. Vallee, Nucl. Instrum. Methods Phys. Res. A339, 232 (1994)
(X- and gamma-ray energies and emission probabilities)
- 1994Ra27 W. Raab, J.L. Parus Nucl. Instrum. Methods Phys. Res. A339, 116 (1994)
(α-transition probabilities)
- 1995Jo23 P.N.Johnston, P.A.Burns, Nucl.Instrum.Methods Phys.Res. A361 (1995) 229.
(U LX ray energies and emission probabilities)
- 1996Firestone R.B. Firestone, V.S. Shirley, C.M. Baglin, S.Y.F. Chu, J. Zipkin, 1996. Table of Isotopes. Eighth
Edition, Volume II: A=151-272.
(Decay scheme, gamma ray energies and multipolarities)
- 1996Sa24 A.M. Sanchez, P.R. Montero, and F.V. Tome, Nucl.Instrum.Methods Phys.Res. A369, 593 (1996)
(α-transition probabilities)
- 1996Sc06 E. Schönfeld and H. Janßen, Nucl. Instrum. Methods Phys. Res. A369, 527 (1996)
(Atomic data)
- 1996Vi07 L.L.Vintro, P.I.Mitchell, O.M.Condren, M.Moran, J.Vives i Batlle, J.A.Sanchez-Cabeza,
Nucl.Instrum.Methods Phys.Res. A369(1996)597
(α-transition probabilities)
- 1997Bu23 A.V.Bushuev, V.N.Zubarev, E.V.Petrova et al. At.Energ. 82, 117 (1997);
(Gamma-ray emission probabilities)

Comments on evaluation

- 1997Ko52 R.O.Korob, S.L.Figueroa, *Radiochim.Acta* 77, 161 (1997)
(Gamma-ray emission probabilities)
- 1999Sa19 A.M.Sanchez, P.R.Montero, *Nucl.Instrum.Methods Phys.Res.* A420, 481 (1999)
(α -transition probabilities)
- 1999ScZX E. Schönfeld and G. Rodloff - PTB-6.11-1999-1999-1, Braunschweig, February 1999
(KX-ray energies and relative emission probabilities)
- 2002BA85 I.M.Band, M.B.Trzhaskovskaya, C.W.Nestor, Jr., P.O.Tikkanen, S.Raman, *At.Data Nucl.Data Tables*
81, 1 (2002)
(Theoretical ICC)
- 2002Da21 F. Dayras, *Nucl.Instrum.Methods Phys.Res.* A490, 492 (2002)
(α -transition probabilities)
- 2003Au03 G.Audi, A.H.Wapstra, and C.Thibault, *Nucl.Phys.* A729, 337 (2003)
(Q value)
- 2003Br12 E.Browne, *Nucl.Data Sheets* 98, 665 (2003)
(Evaluation of ²³⁹Pu decay data, ²³⁵U level energies, gamma-ray emission probabilities, α -transition probabilities)

²⁴⁰Pu – Comments on evaluation of decay data

by V. P. Chechev

This evaluation was done originally in 2004 (2004BeZQ, 2005ChZU) and then updated in June 2009 with a literature cut-off by the same date.

1. DECAY SCHEME

The decay scheme is based on 2006Br20. Some expected weak gamma-ray transitions have not been observed directly in ²⁴⁰Pu alpha decay but were adopted from decay of ²³⁶Pa and ²³⁶Np and from data on nuclear reactions.

The alpha transitions to ²³⁶U highly excited levels with energy of 958.960 and 967 keV were not observed. They are expected from data on level spins and gamma-rays de-excited these levels.

2. NUCLEAR DATA

Q(α) value is from 2003Au03.

The recommended half-life of ²⁴⁰Pu is based on the experimental results given in Table 1. Re-estimated values were used for averaging where necessary.

Table 1. Experimental values of ²⁴⁰Pu half-life (in years)

Reference	Author(s)	Original value	Re-estimated value	Measurement method	
1951In03	Inghram et al.	6580 (40)	6500 (45) ^{b, c}	Mass-Spectrometry	
1951We21	Westrum	6300 (600)			
1954Farwell	Farwell et al.	6760	6610 (55) ^b	α-Particle Counting	
1956Bu92	Butler et al.	6600 (100)		α-Particle Counting	
1959Dokuchaev	Dokuchaev	6620 (50)		α-Particle Counting	
1968Oe02	Oetting	6524 (10)		6537 (15) ^c	Calorimetry
1978Ja11	Jaffey et al.	6569 (6)		6569 (7) ^c	α-Particle Counting
1984Be19	Beckmann et al.	6574 (6) ^a		6574 (7) ^c	Mass-Spectrometry
1984St06	Steinkruger et al.	6571 (9) ^a		6552.2 (66) ^c	α-Particle Counting
1984Lu04	Lucas and Noyce	6552.2 (20)			α-Particle Counting
1984Ru04	Rudy et al.	6552.4 (17)			Calorimetry
2007Ah05	Ahmad et al.	6545 (19)			Ingrowth of ²⁴⁰ Pu in ²⁴⁴ Cm source, ²⁴⁰ Pu/ ²⁴⁴ Cm activity ratio measurement

^a Quoted uncertainties, corresponding to 95 % confidence level, have been reduced by a factor 2.

^b Re-estimated in 1978Ja11.

^c Re-estimated in 1986LoZT.

With omitting the value of 1954Farwell reported without uncertainty the weighted average of the remaining 11 values is 6561 yr with the internal uncertainty 3.1 yr and external uncertainty 3.8 yr.

According to the criterion adopted by the members of the CRP (1986LoZT) a minimum uncertainty of the recommended ²⁴⁰Pu half-life should be attributed as 7 years.

Therefore, the adopted value of the ²⁴⁰Pu half-life is 6561 (7) years.

The recommended of ²⁴⁰Pu spontaneous fission half-life is based on the experimental results given in Table 2.

Table 2. Experimental values of ²⁴⁰Pu spontaneous fission half-life (in 10¹¹ years)

Reference	Author(s)	Measurement value	Measurement method	Used for final averaging
1953Ki72	Kinderman	1.314 (26)	Low geometry α -counting	No
1954Ba14	Barclay et al.	1.225 (30)	Low geometry α -counting	No
1954Ch74	Chamberlain et al.	1.20	Low geometry α -counting	No
1959Mi90	Mikheev et al.	1.20	Low geometry α -counting	No
1962Wa13	Watt et al.	1.340 (15)	Low geometry α -counting	No
1963Ma50	Malkin et al.	1.45 (2)	Low geometry α -counting	No
1967White	White	1.27 (5)	No details available	No
1967Fi13	Fieldhouse et al.	1.176 (25) ^a	SF neutron emission rates	Yes
1979BuZC	Budtz-Jorgensen et al.	1.15 (3)	Fragment spectra, ionization chamber	Yes
1984An25	Androsenko et al.	1.15 (3)	SF neutron emission rates	Yes
1988SeZY	Selickij et al.	1.17 (3)	Fragment detection in 2π geometry	Yes
1989Dy01	Dytlewski et al.	1.12 (2)	Neutron coincidences and low geometry α -counting	Yes
1991Iv01	Ivanov et al.	1.15 (2)	$\lambda_{SF}/\lambda\alpha$ in ²⁴⁰ Pu standards	Yes

^a Re-estimated in 2000Ho27. Original value is 1.170 (25).

Early measurement values have been omitted from averaging according to analysis of Holden and Hoffman (2000Ho27). The weighted average of 6 selected values is 1.15 with the internal uncertainty 0.010 and external uncertainty 0.0087.

The recommended value of the ²⁴⁰Pu spontaneous fission is 1.15 (2) 10¹¹ years where the uncertainty is the smallest quoted uncertainty.

2.1 Alpha Transitions

The energies of the alpha transitions have been obtained from the Q value and the level energies given in Table 3 from 2006Br20.

Table 3. ²³⁶U levels populated in ²⁴⁰Pu α -decay

Level number	Energy, keV	Spin and parity	Half-life	Probability of α -transition ($\times 100$)
0	0	0 ⁺	2.343 (6)·10 ⁷ yr	72.74 (18)
1	45.2440 (20)	2 ⁺	234 (6) ps	27.16 (19)
2	149.477 (6)	4 ⁺	124 (7) ps	0.0863 (18)
3	309.785 (7)	6 ⁺	58 (3) ps	0.001082 (18)
4	522.25 (5)	8 ⁺	24 (2) ps	4.7 (5)·10 ⁻⁵
5	687.59 (4)	1 ⁻	3.78 (9) ns	1.93 (4)·10 ⁻⁵
6	744.18 (7)	3 ⁻	< 0.1 ns	
7	919.14 (17)	0 ⁺		$\approx 6.5 \cdot 10^{-7}$
8	957.90 (17)	(2 ⁺)		< 1.7·10 ⁻⁷
9	960.3 (3)	(2 ⁺)		< 1.3·10 ⁻⁷
10	966.62 (9)	1 ⁻		< 1·10 ⁻⁷

The probabilities of the most intense transitions $\alpha_{0,0}$ and $\alpha_{0,1}$ were obtained by averaging experimental data (Table 4). The probabilities of all the remaining α -transitions have been deduced from the $P(\gamma+ce)$ balances at relevant levels in ^{236}U . The $\alpha_{0,6}$ -transition probability of $1.3(7) \cdot 10^{-8} \%$ has been taken from 2006Br20.

Table 4. Experimental and recommended values of α -transition probabilities ($\times 100$) in ^{240}Pu decay

	α -particle energy keV	1956 Ko67	1956 Go43	1952 As28 1957 As83	1969 Le05	1977 Ba69	1984 Ah06	1990 An33	1992 Bl13	1994 Ra27	1994 Sa63	1996 Vi07	2004 Si03	Recommended
$\alpha_{0,0}$	5168	75.5	75.5	76		73.51 (36)	72.8 (1)	73.0 (5)	72.55 (20)	73.1 (1)	72.5 (11)	74 (2)	72.56 (6)	72.74 (18) ^a
$\alpha_{0,1}$	5124	24.4	24.5	24		26.39 (21)	27.1 (1)	27.0 (5)	27.35 (10)	26.8 (1)	27.5 (11)	26 (2)	27.35 (7)	27.16 (19) ^b
$\alpha_{0,2}$	5021	0.091 (6)	0.085 (15)	0.1		0.096 (5)	0.090 (5)		0.10 (2)					0.0863 (18) ^c
$\alpha_{0,3}$	4864	0.0032 (1)				0.001								0.001082 (18) ^c
$\alpha_{0,4}$	4655													4.7 (5) $\cdot 10^{-5c}$
$\alpha_{0,5}$	4492				2.1(4) 10^{-5}									1.93 (4) $\cdot 10^{-5c}$

^a LWEIGHT computer program has increased the uncertainty of 2004Si03 to 0.0649 and recommended a weighted average (72.74) with the expanded uncertainty of 0.18 so range includes the most precise value of 72.56.

^b LWEIGHT computer program has recommended a weighted average (27.16) with the expanded uncertainty of 0.19 so range includes the most precise value of 27.35.

^c Deduced from $(\gamma+ce)$ -intensity balance at relevant levels.

2.2. Gamma Transitions and Internal Conversion Coefficients

The recommended energies of gamma-ray transitions are virtually the same as the gamma-ray energies because nuclear recoil is negligible for ^{234}U .

The gamma-ray transition probabilities were deduced from the gamma-ray emission probabilities and total internal conversion coefficients (ICCs). The ICCs have been interpolated using the BrIcc package with the so called “Frozen Orbital” approximation (2008Ki07). The uncertainties in the ICCs for pure multiplicities have been taken as 2 %. The multiplicities have been taken from 2006Br20.

The experimental values of ICC have been adopted for the E1 anomalously converted gamma-ray transitions $\gamma_{5,1}$ (642.4 keV) and $\gamma_{5,0}$ (687.6 keV).

3. ATOMIC DATA

3.1. Fluorescence yields

The fluorescence yield data are from 1996Sc06 (Schönfeld and Janßen).

3.2. X-Rays

The energies of U LX-rays taken from the SAISINUC software supporting programs agree with the measurements of 1994Le28 and 1994Le37 where the fine structure of LX-radiation was measured in decays of ^{239}Pu and ^{240}Pu . Other measurements of U LX-rays can be found in 1983Ah02, 1984Bo41, 1992Ba08 and 1995Jo23.

The U KX-ray energies have been taken from 1999Schönfeld where the calculated values based on X-ray wavelengths from 1967Be65 (Bearden). In Table 5 the adopted values of U KX-ray energies are compared with experimental values.

The relative KX-ray emission probabilities were taken from 1999Schönfeld.

Table 5. Experimental and recommended (calculated) values of U KX-ray energies (keV)

	1976GuZN	1982Ba56	1983Ah02	Adopted
K α_2	94.655 (5)	94.656 (2)	94.67 (2)	94.666
K α_1	98.442 (5)	98.435 (2)	98.45 (2)	98.440
K β_3	110.42	110.416 (3)	110.42 (3)	110.421
K β_1	111.30	111.300 (2)	111.31 (2)	111.298
K β_5	-	111.868 (5) - K β_5 '' 112/043 (5) - K β_5 '	112.01 (5)	111.964
K $\beta_{2,4}$	114.54	-	114.50 (3)	114.46
KO $_{2,3}$	115.40	-	115.40 (5)	115.377

3.3. Auger Electrons

The energies of Auger electrons are from the SAISINUC software supporting programs.

The ratios P(KLX)/P(KLL), P(KXY)/P(KLL) are taken from 1996Sc06.

4. ALPHA EMISSIONS

The energy of alpha particles corresponding to the alpha transition to a ground state of ²³⁶U, E($\alpha_{0,0}$), has been adopted from the absolute measurement of 1972Go33 taking into account the correction of - 0.17 keV recommended by A.Rytz in 1991Ry01.

The energies of all other alpha particles have been deduced from Q(α), E($\alpha_{0,0}$) and the level energies taking into account the ²³⁶U recoil energies.

In Table 6 the deduced (recommended) values of α -particle energies are compared with the experimental results.

Table 6. Experimental and recommended α -particle energies in decay of ²⁴⁰Pu, keV

	Measured ^a						Recommended
	1956 Ko67	1956 Go43	1952As28 1957As83	1962 Le11	1972 Go33	1977 Ba69	
$\alpha_{0,0}$	5166	5165	5168 (4)	5167.7 (7)	5168.13 (15) ^b	5168.13 (15) ^b	5168.13 (15) ^b
$\alpha_{0,1}$	5122	5121	5123 (5)	5123.3 (7)	5123.26 (23)	5123.45 (25)	5123.6 (2)
$\alpha_{0,2}$	5021 (2)	5020	5019			5021.3 (5)	5021.1 (2)
$\alpha_{0,3}$	4858 (5)	4856				4863.4 (5)	4863.5 (2)

^a Original values have been adjusted taking into account changes in calibration energies as suggested in 1991Ry01.

^b Absolute measurement; the value was adopted as recommended in 1991Ry01 and used in 2003Au03 for obtaining Q(α).

It should be noted that Sibbens and Pommé (2004Si03) measured (using a 50 mm² high-resolution planar silicon detector) the energies of ²⁴⁰Pu alpha particles relatively to reference peaks of ²³⁸Pu and ²³⁹Pu for a ^{238,239,240}Pu mixture. They obtained E($\alpha_{0,0}$) = 5168.54 (14) keV and E($\alpha_{0,1}$) = 5124.10 (15) keV discrepant with other published data.

5. ELECTRON EMISSIONS

The energies of the conversion electrons have been obtained from the gamma transition energies and the atomic-electron binding energies.

The emission probabilities of conversion electrons have been deduced from the evaluated $P(\gamma)$ and ICC values. The experimental spectrum of the conversion electrons in decay of ²⁴⁰Pu is given in 1958Sa21.

The absolute emission probabilities of K Auger electrons have been calculated using the EMISSION computer program (2000Schönfeld).

The total absolute emission probability of L Auger electrons has been deduced using the adopted total absolute emission probability of U LX-rays and fluorescence yield $\omega_L = 0.500$ (19).

6. PHOTON EMISSIONS

6.1. X-Ray Emissions

The absolute emission probabilities of U LX-rays have been obtained as weighted averages of measurement results from 1994Le28 and 1994Le37. The uncertainties are the smallest quoted uncertainties.

The total absolute emission probability of U LX-rays $P(XL) = 10.34$ (15) %, adopted from measurements of 1994Le28, 1994Le37, agrees well with the value of $P(XL) = 10.14$ (23) %, calculated with using the EMISSION computer program (2000Schönfeld). The measurement result of 1970Swinth (11.5 (3) %) disagrees with the adopted and calculated values.

The absolute KX-ray emission probabilities have been calculated using the EMISSION computer program (2000Schönfeld).

6.2. Gamma-Ray Emissions

The energies of gamma-rays have been adopted from 2006Br20 based on the available experimental data from ²⁴⁰Pu α -decay (Table 7) and data from decay of ²³⁶Pa and ²³⁶Np.

Table 7. Measured in ²⁴⁰Pu α -decay ^a and recommended values of gamma-ray energies (keV)

	1969Le05	1971GuZY	1972Sc01	1974HeYW	1975OtZX	1976GuZN	1981He16	Recommended
$\gamma_{1,0}$		45.235 (20)	45.242 (6)			45.232 (5)	45.244 (3)	45.2440 (20)
$\gamma_{2,1}$		104.233 (10)	104.233 (5)	104.15 (2)		104.244 (5)	104.234 (6)	104.233 (5)
$\gamma_{3,2}$		160.35 (50)	160.310 (8)	160.27 (2)	160.312 (10)	160.280 (15)	160.308 (3)	160.308 (3)
$\gamma_{4,3}$			212.4 (1)		212.48 (5)			212.46 (5)
$\gamma_{5,2}$	538.05 (30)				538.09 (15)			538.10 (10)
$\gamma_{5,1}$	642.43 (10)			642.48 (15)	642.33 (10)	642.48		642.34 (5)
$\gamma_{5,0}$	687.77 (15)			688.01 (15)	687.57 (10)	687.7		687.56 (10)
$\gamma_{7,1}$	873.91 (20)				873.92 (15)			874.0 (2)

^a. For other much more inaccurate measurements results, see in 1958Sa21, 1959Tr37 and 1972CiZS.

The experimental and recommended gamma-ray emission probabilities for γ -rays with energy less than 200 keV are given in Table 8. The recommended $P(\gamma)$ values have been obtained by averaging several experimental results (except for $P(\gamma_{1,0})$ that calculated from intensity balance).

Table 8. Experimental and recommended emission probabilities of gamma-rays in ²⁴⁰Pu decay with energy less than 200 keV (per 10⁴ α-decays)

	Energy (keV)	1971 GuZY	1972 Sc01	1975 OtZX	1976 GuZN	1976 Um01	1981 He16	1981 Morel	1994 Ba91	Recommended
γ _{1,0}	45.24	4.50 (10) ^a	4.50 ^b		4.53 (9) ^d	4.61 (14) ^e	4.35 (9)			4.62 (9) ^f
γ _{2,1}	104.23	0.700 (14) ^a	0.91 (5) ^c	0.70 ^b	0.698 (14) ^d		0.718 (7)			0.714 (7) ^g
γ _{3,2}	160.31	0.0420 (8) ^a	0.049 (12) ^c	0.0408 (10)	0.0402 (8) ^d		0.0402 (4)	0.0402 (7)	0.04065 (17)	0.04045 (22) ^h

^a Omitted from averaging as the results of 1971GuZY were superseded in 1976GuZN.

^b Omitted from averaging as an uncertainty is not quoted.

^c Omitted on statistical considerations (using Chauvenet's criterion).

^d The uncertainty quoted in 1976GuZN was re-estimated in 1986LoZT to include a 2 % detector efficiency uncertainty.

^e The uncertainty quoted in 1976Um01 was re-estimated in 1986LoZT to include a 2 % detector efficiency uncertainty and 1 % from the sample isotopic composition.

^f Deduced from intensity balance at level 45,24 keV using P(α_{0,1}) = 27,16 (19) % and total ICC α_T(γ_{1,0}) = 589 (12). The recommended value agrees with the measurement of 1976Um01 and differs from the measurement result of 1981He16.

^g Weighted average of 1976GuZN and 1981He16; the uncertainty is the smallest quoted uncertainty.

^h LWEIGHT computer program identified an outlier (1972Sc01). With the five remained experimental values for processing the program increased the uncertainty of 1994Ba91 to 0.00030 and recommended a weighted average; the uncertainty is internal.

The emission probabilities of γ_{4,3}(212 keV) and γ_{5,2}(538 keV) have been adopted from absolute measurements of 1975OtZX. The emission probabilities of γ_{5,1}(642 keV) and γ_{5,0}(687 keV) have been obtained by averaging experimental data (Table 9).

Table 9. Experimental and recommended emission probabilities of gamma-rays de-exciting the ²³⁶U level with energy of 687.6 keV in ²⁴⁰Pu decay (per 10⁸ α-decays)

	Energy, keV	1969Le05	1971GuZY	1975OtZX	1975Dr05	1976GuZN	Recommended
γ _{5,2}	538.1	≈ 0.23 ^a		0.147 (12)			0.147 (12)
γ _{5,1}	642.4	14.5 ^a	14.5 (5) ^b	12.6 (4)	13 (1)	12.45 (30)	12.6 (3) ^c
γ _{5,0}	687.6	3.77 (11)	3.70 (15) ^b	3.30 (13)		3.55 (9)	3.56 (9) ^c

^a Omitted from averaging as an uncertainty is not quoted.

^b Omitted from averaging as the results of 1971GuZY were superseded in 1976GuZN.

^c Weighted average of 3 experimental values; the uncertainty is the smallest quoted uncertainty.

The emission probability of γ_{7,1} (874 keV) was obtained as a weighted average of measurement results from 1969Le05 and 1975OtZX.

The weak gamma-rays with energy more than 900 keV were reported in 1969Le05 and 1976GuZN. They are expected from the decay scheme but their emission probabilities (<10⁻⁷ per 100 decays) were determined with a great inaccuracy.

7. CONSISTENCY OF RECOMMENDED DATA

The most accurate Q value, Q(M), is taken from the atomic mass adjustment table of Audi et al. (2003Au03). Comparison of Q(eff)(deduced as the sum of average energies per disintegration (ΣE_i × P_i) for all emissions accompanying ²⁴⁰Pu α- decay) with the tabulated decay energy Q(M) allows to check a consistency of the recommended decay-scheme parameters obtained in this evaluation.

Here E_i and P_i are the evaluated energies and emission probabilities of the i-th alpha particle, beta particle, gamma-ray, X-ray, etc. Consistency (percentage deviation) is determined by {[Q(M) - Q(eff)] / Q(M)} × 100. "Percentage deviations above 5 % would be regarded as high and imply a poorly defined decay scheme; a value of less than 5 % indicates the construction of a reasonably

consistent decay scheme” (quoted from the article by A. L. Nichols in Appl. Rad. Isotopes 55(2001) 23-70).

For the above ²⁴⁰Pu decay data evaluation we have $Q(M) = 5255.75 (14) \text{ keV}$ and $Q(\text{eff}) = 5255 (9) \text{ keV}$. Thereafter, the percentage deviation is $(0.00 \pm 0.17) \%$, i.e. consistency is superior.

8. REFERENCES

- 1951In03 M. G. Inghram, D. C. Hess, P. R. Fields, G. L. Pyle., Phys. Rev. 83(1951)1250. (Half-life)
- 1951We21 E. F. Westrum, Phys. Rev. 83(1951)1249. (Half-life)
- 1952As28 F. Asaro, I. Perlman, Phys. Rev. 88(1952)828. (α -transition energies and probabilities)
- 1953Ki72 E. M. Kinderman, Hanford Lab. Report HW-27660(1953). (SF half-life)
- 1954Ba14 F. R. Barclay, W. Galbraith, K. M. Glover, G. R. Hall, W. J. Whitehouse, Proc. Phys. Soc. (London) 67A(1954)646. (SF half-life)
- 1954Ch74 O. Chamberlain, G. W. Farwell, E. Segre, Phys. Rev. 94(1954)156. (SF half-life)
- 1954Farwell G. Farwell., J. E. Roberts, A. C. Wahl, Phys. Rev. 94(1954)363. (Half-life)
- 1956Bu92 J. P. Butler, T. A. Eastwood, T. L. Collins, M. E. Jones, F. M. Rourke, R. P. Schuman, Phys. Rev. 103(1956) 634. (Half-life)
- 1956Go43 L. L. Goldin, G. I. Novikova, E. F. Tretyakov, Phys. Rev. 103(1956)1004. (α -transition energies and probabilities)
- 1956Ko67 L. M. Kondratev, G. I. Novikova, Y. P. Sobolev, L. L. Goldin, Zh. Eksp. Teor. Fiz. 31 (1956) 771. (Soviet Phys. JETP 4(1956)645). (α -transition energies and probabilities)
- 1957As83 F. Asaro, S. G. Thompson, F. S. Stephens, Jr., I. Perlman, Priv. Comm (1957), quoted by 1964Hy02. (α -transition energies and probabilities)
- 1958Sa21 P. S. Samoilov, Atomnaya Energ. 4(1958)81 Soviet. J. At. Energy 4(1958)102. (Gamma-ray energies)
- 1959Dokuchaev Ya. P. Dokuchaev, Atomnaya Energ. 6(1959)74. (Half-life)
- 1959Mi90 V. L. Mikheev, N. K. Skobelev, V. A. Druin, G. N. Flerov, Zhur. Eksptl. i Teoret. Fiz. 37(1959)859 Soviet Phys. JETP 10 (1960) 612. (Half-life)
- 1959Tr37 E. F. Tretyakov, L. N. Kondratev, G. I. Khlebnikov, L. L. Goldin, Zh. Eksp. Teor. Fiz. 36(1959)362 Soviet Phys. JETP 9(1959)250. (Gamma-ray energies)
- 1962Le11 C. F. Leang, Compt. Rend. 255(1962)3155. (α -transition energies)
- 1962Wa13 D. E. Watt, F. J. Bannister, J. B. Laidler, F. Brown, Phys. Rev. 126(1962)264. (SF half-life)
- 1963Ma50 L. Z. Malkin, I. D. Alkhazov, A. S. Krivokhatsky, K. A. Petrzhak, At. Energ. USSR 15(1963)158 Soviet. J. At. Energy 15(1964)851. (SF half-life)
- 1964Hy02 E. K. Hyde, I. Perlman, G. T. Seaborg (1964): The Nuclear Properties of the Heavy Elements, Vol II. Prentice-Hall, Inc., Englewood Cliffs, N J. (α -transition energies and probabilities)
- 1967Be65 J. A. Bearden, Rev. Mod. Phys. 39(1967)78. (X-ray energies)
- 1967Fi13 P. Fieldhouse, D. S. Mather, E. R. Culliford, J. Nucl. Energy 21(1967)749. (SF half-life)
- 1967White P. H. White, priv. comm., cited in J.Nucl.Energy 21(1967)749 and in 2000Ho27. (SF half-life)
- 1968Oe02 F. L. Oetting, Proc. Symp. Thermodyn. Nucl. Mater. With Emphasis on Solution Syst., Vienna, Austria (1967), IAEA, Vienna, p. 55 (1968). (Half-life)
- 1969Le05 C. M. Lederer, J. M. Jaklevic, S. G. Prussin, Nucl. Phys. A135(1969)36. (α -transition energies and probabilities)
- 1970Swinth K. L. Swinth, IEEE Nuclear Science Symp.,1970, Vol. 4, p.125. (LX-ray emission probabilities)
- 1971GuZY R. Gunnink, R. J. Morrow, UCRL 51087(1971). (Gamma-ray energies and emission probabilities)
- 1972CIZS J. E. Cline, R. J. Gehrke, L. D. McIsaac, ANCR1069(1972). (Gamma-ray energies)
- 1972Go33 D. J. Gorman, A. Rytz, H. V. Michel, Compt. Rend. B275(1972)291. (α -transition energies)

- 1972Sc01 M. Schmorak, C. E. Bemis Jr, M. J. Zender, N. B. Gove, P. F. Dittner, Nucl. Phys. A178(1972)410. (Gamma-ray energies and emission probabilities)
- 1974HeYW R. L. Heath, Gamma-Ray Spectrum Catalogue, ANCR 1000(1974)2. (Gamma-ray energies)
- 1975Dr05 T. Dragnev, K. Scharf, Intern. J. Appl. Radiat. Isotop. 26(1975)125. (Gamma-ray emission probabilities)
- 1975OtZX H. Ottmar, P. Matussek, I. Piper, Proc. Int. Symp. Neutron Capture, Gamma-Ray Spectroscopy and Related Topics, 2nd, Petten, The Netherlands (1974), K. Abrahams, F. Stecher-Rasmussen, P. Van Assche, Eds, Reactor Centrum Nederland, p. 658 (1975). (Gamma-ray energies and emission probabilities)
- 1976GuZN R. Gunnink, J. E. Evans, A. L. Prindle, UCRL-52139(1976). (Gamma-ray energies and emission probabilities)
- 1976Um01 H. Umezawa, T. Suzuki, S. Ichikawa, J. Nucl. Sci. Technol. 13(1976)327. (Gamma-ray emission probabilities)
- 1977Ba69 S. A. Baranov, V. M. Shatinskii, Yad. Fiz. 26(1977)461 Soviet J. Nucl. Phys. 26(1977)244. (α -transition energies and probabilities)
- 1978Ja11 A. H. Jaffey, H. Diamond, W. C. Bentley, D. G. Graczyk, K. P. Flynn, Phys. Rev. C18(1978)969. (Half-life)
- 1979BuZC C. Budtz-Jorgensen, H. -H. Knitter, NEANDC(E) 202U, Vol III, p. 9 (1979). (SF half-life)
- 1981He16 R. G. Helmer, C. W. Reich, Intern. J. Appl. Radiat. Isotop. 32(1981)829. (Gamma-ray energies and emission probabilities)
- 1981Morel J. Morel et al., LMRI, Saclay, private communication, 1981 Cited in IAEA, Vienna, Tec. Rep. 261, 1986. (Gamma-ray emission probabilities)
- 1982Ba56 G. Barreau, H. G. Borner, T. von Egidy, R. W. Hoff. Z. Phys. A308(1982)209. (KX-ray energies)
- 1983Ah02 I. Ahmad, J. Hines, J. E. Gindler, Phys. Rev. C27(1983)2239. (KX-ray energies)
- 1984Ah06 I. Ahmad, Nuclear Instrum. Methods 223(1984)319. (α -transition probabilities)
- 1984An25 A. A. Androsenko, P. A. Androsenko, Yu. V. Ivanov, A. E. Konyaev, V. F. Kositsyn, E. M. Tsenter, V. T. Shchebolev, At. Energ. 57(1984)357 Sov. At. Energy 57(1984)788. (SF half-life)
- 1984Be19 R. J. Beckman, S. F. Marsh, R. M. Abernathy, J. E. Rein, Intern. J. Appl. Radiat. Isotop. 35(1984)163. (Half-life)
- 1984Bo41 G. Bortels, B. Denecke, R. Vaninbroux, Nuclear Instrum. Methods 223(1984)329. (LX-ray energies)
- 1984Lu04 L. L. Lucas, J. R. Noyce, Intern. J. Appl. Radiat. Isotop. 35(1984)173. (Half-life)
- 1984Ru04 C. R. Rudy, K. C. Jordan, R. Tsugawa, Intern. J. Appl. Radiat. Isotop. 35(1984)177. (Half-life)
- 1984St06 F. J. Steinkruger, G. M. Matlack, R. J. Beckman, Intern. J. Appl. Radiat. Isotop. 35(1984)171. (Half-life)
- 1986LoZT A. Lorentz, Decay Data of the Transactinium Nuclides, IAEA, Vienna, Tec. Rep. Ser. 261(1986). (Gamma-ray emission probabilities)
- 1988SeZY Yu. A. Selitsky, V. B. Funshtein, V. A. Yakovlev, Program and Theses, Proc. 38th Ann. Conf. Nucl. Spectrosc. Struct. At. Nuclei, Baku, Acad. Sci. USSR, (1988)131. (SF half-life)
- 1989Dy01 N. Dytlewski, M. G. Hines, J. W. Boldeman, Nucl. Sci. Eng. 102(1989)423. (SF half-life)
- 1990An33 S. V. Anichenkov, Yu. S. Popov, Radiokhimiya 32(1990)109 Sov. Radiochem. 32(1991)401. (α -transition probabilities)
- 1991Iv01 Yu. V. Ivanov, A. E. Konyaev, V. F. Kositsyn, E. A. Kholnova, V. T. Shchebolev, M. F. Yudin, At. Energ. 70(1991)396 Sov. At. Energy 70(1991)491. (SF half-life)
- 1991Ry01 A. Rytz, At. Data Nucl. Data Tables 47(1991)205. (α -transition energies)
- 1992Ba08 G. Barci-Funel, J. Dalmasso, G. Ardisson, Appl. Radiat. Isot. 43(1992)37. (X-ray energies)
- 1992Bl13 C. J. Bland, J. Truffy, Appl. Radiat. Isot. 43(1992)1241. (α -transition probabilities)
- 1994Ba91 D. T. Baran, Appl. Radiat. Isot. 45(1994)1177. (α -transition probabilities)
- 1994Le28 M. C. Lépy, K. Debertain. Nucl. Instrum. Meth. Phys. Res. A339(1994)218. (LX-ray energies and emission probabilities)

- 1994Le37 M. C. Lépy, B. Duchemin, J. Morel, Nucl. Instrum. Meth. Phys. Res. A353(1994)10. (LX-ray energies and emission probabilities)
- 1994Ra27 W. Raab, J. L. Parus, Nucl. Instrum. Meth. Phys. Res. A339(1994)116. (α -transition probabilities)
- 1994Sa63 A. M. Sanchez, F. V. Tome, J. D. Bejarano, Nucl. Instrum. Meth. Phys. Res. A340(1994)509. (α -transition probabilities)
- 1995Jo23 P. N. Johnston, P. A. Burns, Nucl. Instrum. Meth. Phys. Res. A361(1995)229. (U LX-ray energies and emission probabilities)
- 1996Vi07 L. L. Vintro, P. I. Mitchell, O. M. Condren, M. Moran, J. Vives i Batlle, J. A. Sanchez-Cabeza, Nucl. Instrum. Meth. Phys. Res. A369(1996)597. (α -transition probabilities)
- 1999Schönfeld E. Schönfeld and G. Rodloff PTB-6.11-1999-1999-1, Braunschweig, Februar 1999 (KX-ray energies and relative emission probabilities)
- 2000Ho27 N. E. Holden, D. C. Hoffman, Pure Appl. Chem. 72(2000)1525. (SF half-life)
- 2000Schönfeld E. Schönfeld, H. Janßen, Appl. Rad. Isot. 52(2000)595. (X-ray and Auger electron emission probabilities, EMISSION code)
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)337. (Q value)
- 2004BeZQ M. M. Bé, V. Chisté, C. Dulieu, E. Browne, V. Chechev, N. Kuzmenko, R. Helmer, A. Nichols, E. Schönfeld, and R. Dersch, Table of Radionuclides (Vol.2 A = 151 to 242), Monographie BIPM-5, Vol. 2, p. 247 – 255. Bureau International des Poids et Mesures(2004)
- 2004Si03 G. Sibbens, S. Pommé, Appl. Radiat. Isot. 60(2004)155. (α -transition energies and probabilities)
- 2005ChZU V. P. Chechev, Proc. Intern. Conf. Nuclear Data for Science and Technology, Santa Fé, New Mexico, 26 September-1 October, 2004, R. C. Haight, M. B. Chadwick, T. Kawano, P. Talou, Eds., Vol. 1, p. 91 (2005); AIP Conf. Proc. 769(2005). (²⁴⁰Pu Decay Data Evaluation)
- 2006Br20 E. Browne, J. K. Tuli, Nuclear Data Sheets 107(2006)2649. (Decay scheme, ²³⁶U level energies, gamma-ray multipolarities, data from ²³⁶Pa and ²³⁶Np decays)
- 2007Ah05 I. Ahmad, F. G. Kondev, J. P. Greene, M. A. Kellett, A. L. Nichols, Nucl. Instrum. Meth. Phys. Res. A579(2007)458. (Half-life)
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr., Nucl. Instrum. Meth. Phys. Res. A589(2008)202. (Theoretical ICC)

**²⁴¹Pu – Comments on evaluation of decay data
by V.P.Chechev and N.K. Kuzmenko**

This evaluation was completed in November 2005 and corrected in September 2006. The literature available by September 2006 was included.

1. Decay Scheme

The decay scheme is based on the evaluation of 2006Ba41 (see also the evaluations of 1995Ak01 and 1978El02). It can be considered as basically completed though some very weak gamma transitions were not observed in ²⁴¹Pu alpha decay.

It should be noted there is an ambiguity in the placement of 121,2 keV γ -transition in ²³⁷U level scheme due to doublet (7/2+, 11/2+) near 204 keV. Following 2006Ba41 we show the above γ -transition in Pu-241 α -decay as going from the level 7/2+ while Fotiades *et al.* (2004Fo01) observed this transition in (n,2n)-reaction as going from the level 11/2+.

The upper limit of SF decay is from 1985Dr09.

2. Nuclear Data

Q(α) value is from 2003Au03.

The evaluated ²⁴¹Pu half-life is based on the experimental data given in Table 1. A detailed review of half-life measurements up to 1985 can be found in 1987Ag03. References to earlier measurements are listed in 1978El02. Discrepancies in the measurements were examined by 1986Ha06 and 1987Ba84 in terms of chemical dependency of low-energy β^- decay. In 1986Ha06 a conclusion is drawn that chemical variations (~ 0,3 %) cannot be accountable completely for half-life discrepancies (≥ 1 %).

Table 1. Experimental values of the ²⁴¹Pu half-life (in years)

Reference ^a	Author(s)	Measurement method	Stated value	Revised value	Comments
1953Ma19	MacKenzie <i>et al.</i>	Ingrowth of ²⁴¹ Am by α counting	13,0 (2)	14,1 (2)	Re-estimated for the ²⁴¹ Am half-life of 432,6 (6) a
1956Ro26	Rose and Milstead	Ingrowth of ²⁴¹ Am by 60-keV γ counting	12,77 (28)	13,87 (30)	Re-estimated for the ²⁴¹ Am half-life of 432,6 (6) a. OMITTED: outlier
1960Br15	Brown <i>et al.</i>	Ingrowth of ²⁴¹ Am by α counting	13,24 (24)	14,12 (26)	Re-estimated for the ²⁴¹ Am half-life of 432,6 (6) a
1961Sm03	Smith	Ingrowth of ²⁴¹ Am α -emission	13,0 (3)	14,1 (3) 13,3 (3)	Re-estimated for the ²⁴¹ Am half-life of 432,6 (6) a
1966French	French <i>et al.</i>	Change in ²⁴¹ Pu/Pu ratio by MS	13,59 (46)		Quoted in 1987Ag03 OMITTED: outlier
1966Stepan	Stepan and Nisle	Change in ²⁴¹ Pu reactivity with time	13,63 (36)		OMITTED: updated in 1970Ni02
1967Shields	Shields	Change in ²⁴¹ Pu/Pu ratio in a Pu isotopic standard in 2 years by MS	14,4 (2)		Quoted in 1967Oe01. Stated uncertainty at 0,95 C.L. OMITTED: updated in 1970Sh18

1968Ca19	Cabell	Change in ²⁴¹ Pu/ ^{240,242} Pu ratios in 4,5 years by MS	14,98 (33)		OMITTED: updated in 1971Ca15, outlier
1970Ni02	Nisle and Stepan	Change in ²⁴¹ Pu reactivity with time (in 2,5 yr)	14,63 (27)		
1970Sh18	Shields	Change in ²⁴¹ Pu/Pu ratio in a Pu isotopic standard in 4 years by MS	14,6 (4)	14,6 (2)	Stated uncertainty at 0,95 C.L. For statistical analysis it has been multiplied by 0,5
1971Ca15	Cabel and Wilkins	Change in ²⁴¹ Pu/ ^{240,242} Pu ratios in 6,65 years by MS	15,16 (19)		OMITTED: outlier
1972 Whitehead	Whitehead <i>et al.</i>	Ingrowth of ²⁴¹ Am by 60-keV γ counting	14,91 (15)	14,96 (15)	Re-estimated for the ²⁴¹ Am half-life of 432,6 (6) a OMITTED: updated in 1977Whitehead, outlier
1973JoYT	Jordan	Calorimetric determination of power decay	14,355 (7)		Quoted in 1974StYG
1973Ze02	Zeigler and Ferris	Change in ²⁴¹ Pu/ ²⁴⁰ Pu ratio by MS	14,89 (11)		OMITTED: outlier
1975WiYM	Wilkins	Change in ²⁴¹ Pu/ ^{240,242} Pu ratio by MS	15,02 (10)		OMITTED: outlier
1976McZB	McKean and Crouch	Change in ²⁴¹ Pu/ ^{240,242} Pu ratio by MS	14,35 (6)		
1977Crouch	Crouch and McKean	Change in ²⁴¹ Pu/ ^{240,242} Pu ratio by MS	14,41 (12)		Average of measurement results from 1976-1977 series of experiments
1977 Whitehead	Whitehead	Ingrowth of ²⁴¹ Am 60-keV γ ray	14,56 (15)		
1978 Vaninbroukx	Vaninbroukx	Ingrowth of ²⁴¹ Am by α and 60-keV γ ray counting	14,60 (10)		
1978 Vaninbroukx	Vaninbroukx	Change in ²⁴¹ Pu/ ²⁴⁰ Pu ratio by MS	14,30 (14)		
1979Garner	Garner and Machlan	Change in ²⁴¹ Pu/ ^{240,242} Pu ratio by MS	14,38 (7)		
1980Ag02	Aggarwal and Jane	Ingrowth of ²⁴¹ Am by α spectrometry	14,42 (9)		80 α -spectrometric measurements in 457 days
1980Ma45	Marsch <i>et al.</i>	Change in ²⁴¹ Pu/ ²⁴² Pu ratio in 3,6 yr by MS	14,38 (6)	14,38 (3)	Stated uncertainty at 0,95 C.L. For statistical analysis it has been multiplied by 0,5
1981Ag01	Aggarwal <i>et al.</i>	Ingrowth of ²⁴¹ Am by IDAS	14,52 (8)		
1981Ag07	Aggarwal <i>et al.</i>	Ingrowth of ²⁴¹ Am by α spectrometry and APS	14,44 (6)		Average of the measurement results from two independent series of experiments
1982Ag01	Aggarwal <i>et al.</i>	Ingrowth of ²⁴¹ Am by IDMS	14,32 (11)	14,32 (6)	Revised uncertainty, see 1989Ho24

1982Hiyama	Hiyama <i>et al.</i>	Change in ²⁴¹ Pu/ ²⁴⁰ Pu ratio by MS	14,29 (15)		Quoted in 1989Ho24
1983DeZX	De Bievre <i>et al.</i>	Change in ²⁴¹ Pu/ ²⁴⁰ Pu ratio in 6 years by MS	14,33 (2)		OMITTED: superseded in 1997DeZY
1985Ag02	Aggarwal <i>et al.</i>	Changes in ²⁴¹ Pu/ ²⁴⁰ Pu, ²⁴¹ Pu/ ²³⁹ Pu, ²⁴¹ Pu/ ²⁴² Pu ratios in 5 years by MS	14,38 (2)		In 1985Ag02 it is noted that values from 1980Ag02, 1981Ag01, 1981Ag07, 1982Ag01 were obtained in independent sets of experiments
1986Ti04	Timofeev <i>et al.</i>	Ingrowth of ²⁴¹ Am by IDMS	14,57 (10)	14,57 (5)	Stated uncertainty at 0,95 C.L. For statistical analysis it has been multiplied by 0,5
1989Pa21	Parker <i>et al.</i>	Change in ²⁴¹ Pu/ ²³⁹ Pu ratio by high resolution γ -spectrometry	14,355 (40)		156 sets of normalized spectral full energy peak-area ratios from 13 plutonium samples during 10 years
1997DeZY	De Bievre and Verbruggen	Change in ²⁴¹ Pu/ ²⁴⁰ Pu ratio by precision MS	14,290 (6)	14,290 (3)	Stated uncertainty at 0,95 C.L. For statistical analysis it has been multiplied by 0,5

MS=Mass Spectrometry, IDMS=Isotope Dilution Mass Spectrometry, IDAS=Isotope Dilution Alpha Spectrometry

^a In 1978El02 two more experimental values of are quoted from the private communications of 1977RGZZ and 1978RGZZ. These values are intermediate results of experiments and not discussed later on including the review of 1987Ag03.

After omitting the five superseded values from 1966Stepan, 1967Shields, 1968Ca19, 1972Whitehead and 1983DeZX the data set for statistical processing includes the 24 values. The LWEIGHT computer program using the LRSW analysis has identified the four outliers of 1971Ca15, 1975WiYM, 1973Ze02 and 1956Ro26 and increased the uncertainty of 1997DeZY by 2,04 times. The weighted average of the remaining twenty three values is 14,327, with an internal uncertainty of 0,037, a reduced χ^2 of 5,34, and an external uncertainty of 0,010. The unweighted average is 14,371 (34). The LWEIGHT program has chosen the weighted average and expanded the final uncertainty to 0,037 so range includes the most precise value of 14,290.

The adopted value of the ²⁴¹Pu half-life is 14,33 (4) years, or 5234 (15) days.

Possible chemical effects do not exceed or about the stated relative uncertainty of the half-life.

2.1. Beta Transition

²⁴¹Pu decays by β^- emission to the ground state of ²⁴¹Pu (Table 2).

Table 2. ²⁴¹Am level populated in the ²⁴¹Pu β^- -decay

Level	Energy, (keV)	Spin and parity	Half-life	Probability (%)
0	20,8 (2)	5/2 ⁻	432,6 (6) a	99,997 56 (2)

The experimental and evaluated values of the β^- transition energy are given in Table 3.

The value $Q^- = 20,78 (20)$ keV from 1999YaZX was superseded by the same group in 1999Dr13 and 2000Dr02. Audi *et al.* (2003Au03) give $Q^- = 20,78 (13)$ keV taking into account the value from 1999YaZX (see also 2005Ma88).

Table 3. Experimental values of the ²⁴¹Pu β⁻ transition energy (keV)

Level	1952Fr25	1956Sh31	1999Dr13 2000Dr02	Evaluated
0	20,5 (12)	20,8 (2)	20,7 (3)	20,8 (2)

The probability of the β⁻-transition was deduced from the evaluated α branching (Table 4).

Table 4. Experimental and evaluated values of α branching (α/β⁻), per decay, in the ²⁴¹Pu decay

1961Sm03	1968Ah01	1976GuZN	1977VaYR	Evaluated
2,44 (10)·10 ⁻⁵	2,45 (8)·10 ⁻⁵	2,46 (1)·10 ⁻⁵	2,42 (2)·10 ⁻⁵	2,44 (2)·10 ⁻⁵

2.2. Alpha Transitions

The energies of the alpha transitions have been deduced from Q_α value and the level energies given in Table 5. The level energies were calculated from the gamma -ray energies except for the levels “8”, “9” and “10” the energies of which were taken from 1996FiZX.

Table 5. ²³⁷U levels populated in the ²⁴¹Pu α decay

Level number	Energy, (keV)	Spin and parity	Half-life	Experimental probability of α transition (%) 1965Ba26	Experimental probability of α transition (%) 1968Ah01	Adopted probability of α transition (%)
0	0,0	1/2 ⁺	6,752 (2) d	8,6·10 ⁻⁶		8,6 (10)·10 ⁻⁶
1	11,39 (2)	3/2 ⁺		2,5·10 ⁻⁵		2,5 (2)·10 ⁻⁵
2	56,30 (12)	5/2 ⁺		0,88·10 ⁻⁵	1,00 (12)·10 ⁻⁵	1,00 (12)·10 ⁻⁵
3	82,97 (13)	7/2 ⁺		2,73·10 ⁻⁵	3,2 (3)·10 ⁻⁵	3,2 (3)·10 ⁻⁵
4	159,96 (2)	5/2 ⁺	3,1 (1) ns	2,04·10 ⁻³	2,03 (4)·10 ⁻³	2,03 (4)·10 ⁻³
5	204,19 (14)	7/2 ⁺		3,00·10 ⁻⁴	2,95 (8)·10 ⁻⁴	2,95 (8)·10 ⁻⁴
6	260,95 (17)	9/2 ⁺	-	2,88·10 ⁻⁵		2,9 (3)·10 ⁻⁵
7	274,0 (10)	(7/2) ⁻	155 (6) ns		0,5 (2)·10 ⁻⁵	0,5 (2)·10 ⁻⁵
8	316 (5)	(9/2) ⁻	-		≈1,7·10 ⁻⁶	≈1,7·10 ⁻⁶
9	327 (3)	11/2 ⁺	-	≈7·10 ⁻⁷		≈7·10 ⁻⁷
10	367 (3)	(11/2) ⁻			≈7·10 ⁻⁷	≈7·10 ⁻⁷

The absolute alpha transition probabilities, P(α_i), were calculated using the value of 2,44 (2)·10⁻⁵ for the ²⁴¹Pu alpha decay branching. The uncertainties of P(α_{0,0}) and P(α_{0,1}) have been estimated using the relative uncertainty of the sum of P(α_{0,0}) and P(α_{0,1}) (equal to 1/15) from 1968Ah01.

The probabilities of α-transitions (per 100 α decays) are from the measurements of 1965Ba26 and 1968Ah01. Other measurements: 1976BaZZ. The values of hindrance factors have been calculated using ALPHAD code and r₀ = 1,5156 (9) from 1998Ak04.

2.3. Gamma-ray Transitions and Internal Conversion Coefficients

The evaluated energies of gamma -ray transitions are virtually the same as the photon energies because nuclear recoil is negligible.

The gamma-ray transition probabilities, $P_{\gamma+ce}$, were deduced from the gamma -ray emission probabilities and the total internal conversion coefficients (ICC's) interpolated from the BrIcc package. The relative uncertainties of α_K , α_L , α_M , α_T for pure gamma ray multiplicities have been taken as 2 %.

$P_{\gamma+ce}(\gamma_{1,0} 11,39\text{-keV})$, $P_{\gamma+ce}(\gamma_{3,2} 26,6\text{-keV})$, $P_{\gamma+ce}(\gamma_{5,4} 44,18\text{-keV})$, $P_{\gamma+ce}(\gamma_{2,1} 44,86\text{-keV})$ and $P_{\gamma+ce}(\gamma_{6,5} 56,76\text{-keV})$ were derived from the intensity balances using the adopted probabilities of α -transitions to the corresponding levels. The E2/M1 mixing ratios for $\gamma_{5,4}$ (44,18-keV), $\gamma_{2,1}$ (44,86-keV) and $\gamma_{6,5}$ (56,76-keV) have been deduced from the calculated total conversion coefficients. The gamma transition multiplicities and the E2/M1 mixing ratios for the remaining gamma transitions have been adopted from the analysis of the ²³⁷U level scheme in 1995Ak01.

The transition $\gamma_{6,4}$ (100,94 keV) was not observed experimentally; it is obscured by U KX -rays. This transition is given in 1995Ak01.

3. Atomic Data

3.1. Fluorescence yields

The fluorescence yield data are from 1996Sc06 (Schönfeld and Janßen).

3.2. X Radiations

The relative KX-ray emission probabilities are from 1999ScZX.

3.3. Auger Electrons

The ratios $P(KLX)/P(KLL)$, $P(KXY)/P(KLL)$ are from 1996Sc06.

4. Electron Emissions

The energies of the conversion electrons have been calculated from the gamma transition energies and the electron binding energies.

The emission probabilities of the conversion electrons have been calculated using the evaluated P_γ and ICC values.

The total absolute emission probabilities of K and L Auger electrons have been calculated using the EMISSION computer program.

β^- average energy was adopted from the measurement of 1968Oe01. The calculated value is discrepant: 5,23(5) keV.

5. Alpha Emissions

In Table 6 the experimental and adopted energies of α particles (in keV) are given. The original values of 1965Ba26, 1968Ba25 were increased by 0,4 keV and the values of 1968Ah01 by 0,6 keV because of changes in calibration energies, as recommended by Rytz in 1991Ry01. Other measurements: 1953As40, 1964Dz03, 1976BaZZ, 1984Gl03.

The adopted energies of α particles have been obtained from Q_α value and the level energies given in Table 5 taking into account the relevant recoil energies.

Table 6. α - particle energies in the ²⁴¹Pu decay (keV)

	1965Ba26 1968Ba25	1968Ah01	Adopted (calculated from Q_α)
$\alpha_{0,10}$		4693 (6)	4694 (3)
$\alpha_{0,9}$	4732		4733 (3)
$\alpha_{0,8}$		4743 (5)	4744 (5)
$\alpha_{0,7}$		4784 (5)	4785,1 (11)
$\alpha_{0,6}$	4798	4798 (3)	4798,0 (5)
$\alpha_{0,5}$	4853,3 (12)	4853 (3)	4853,8 (5)
$\alpha_{0,4}$	4896,3 (12)	4896 (3)	4897,3 (5)
$\alpha_{0,3}$	4971	4973 (3)	4973,1 (5)
$\alpha_{0,2}$	4998	5000 (4)	4999,2 (5)
$\alpha_{0,1}$	5041	5043 (3)	5043,4 (5)
$\alpha_{0,0}$	5051	5056 (5)	5054,6 (5)

6. Photon Emissions

6.1. X-Ray Emissions

The absolute emission probabilities of U KX and LX γ -rays have been calculated using the EMISSION code.

		Energy, (keV)	Number of photons per 100 disintegrations
X _K	K α_2 (U)	94,666	3,00 (7)·10 ⁻⁴
	K α_1 (U)	98,440	4,79 (10)·10 ⁻⁴
	K β_3 (U)	110,421	}
	K β_1 (U)	111,298	} 1,79 (5)·10 ⁻⁴
	K β_5 (U)	111,964	}
	K $\beta_{2,4}$ (U)	114,46	} 0,59 (2)·10 ⁻⁴
	KO _{2,3} (U)	115,377	}
X _L	L α_1 (U)	11,619	0,336 (12)·10 ⁻⁴
	L α_2 (U)	13,438	0,556 (19)·10 ⁻⁴
	L α_3 (U)	13,615	4,87 (17)·10 ⁻⁴
	L η (U)	15,399	0,0444 (13)·10 ⁻⁴
	L β (U)	15,727 – 18,206	4,77 (8)·10 ⁻⁴
	L γ (U)	19,507 – 20,714	1,09 (2)·10 ⁻⁴

6.2. Gamma-Ray Emissions

In Table 7 the experimental and adopted energies of gamma γ -rays are given (see also the evaluation of 1988ChZL). Other measurements: 1952Fr25, 1965Ba35, 1976Um01, 1979Ce04, 1993Dr05.

The energies of $\gamma_{1,0}$ (11,39 keV), $\gamma_{3,2}$ (26,67 keV) and $\gamma_{6,4}$ (100,94 keV) have been calculated from the level scheme: $E\gamma_{1,0}$ (11,39 keV) = $E\gamma_{4,0}$ - $E\gamma_{4,1}$; $E\gamma_{3,2}$ (26,67 keV) = $E\gamma_{4,2}$ - $E\gamma_{4,3}$; $E\gamma_{6,4}$ (100,94 keV) = $E\gamma_{5,4}$ + $E\gamma_{6,5}$.

Table 7. Experimental and evaluated gamma-ray energies in the ²⁴¹Pu decay (keV)

	1968Ah01	1971GuZN 1976GuZN	1972Cline	Adopted
$\gamma_{1,0}$		11,39		11,39 (2)
$\gamma_{3,2}$				26,67 (4)
$\gamma_{5,4}$		44,19 (3)	44,175 (30)	44,18 (3)
$\gamma_{2,1}$	44,7 (3)	44,86 (10)		44,86 (10)
$\gamma_{2,0}$	56,6 (2)	56,30 (12)	56,412 (30)	56,30 (12)
$\gamma_{6,5}$		56,76 (10)		56,76 (10)
$\gamma_{3,1}$		71,60 (7)	71,672 (40)	71,64 (9)
$\gamma_{4,3}$	76,9 (2)	76,96 (10)	77,014 (40)	77,01 (4)
$\gamma_{6,4}$				100,94 (11)
$\gamma_{4,2}$	103,5 (2)	103,680 (5)	103,540 (40)	103,680 (5)
$\gamma_{7,4}$	114,0 (10)		115,342 (40)	114,0 (10)
$\gamma_{5,3}$	120,7 (5)	121,2 (10)	121,220 (30)	121,22 (5)
$\gamma_{4,1}$	148,5 (2)	148,567 (10)	148,560 (20)	148,567 (10)
$\gamma_{4,0}$	160,0 (2)	160,00 (4)	159,960 (20)	159,96 (2)

In Table 8 the experimental and evaluated absolute gamma -ray emission probabilities are given. The evaluated values have been obtained using the LWEIGHT computer program. The uncertainty assigned in this evaluation to the recommended value is always greater than or equal to the smallest uncertainty in any of the experimental values used in the statistical processing.

Table 8. Experimental and evaluated absolute emission probabilities of gamma rays in the ²⁴¹Pu decay per 10⁶ disintegrations

E γ (keV)	1968Ah01	1976GuZN	1976U m01	1978DiZU	1985He02	1985Wi04	1994Ba91	Evaluated
44,18		0,042 (2)						0,042 (2)
44,86		0,0084 (10)						0,0084 (10)
56,30		0,025 (2)						0,025 (2)
56,76		0,010 (1)						0,010 (1)
71,64		0,029 (2)						0,029 (2)
77,0	0,18 (2)	0,220 (8)			0,211 (5)	0,203 (4)		0,207 (4)
100,94		0,00072						0,00072
103,68	1,10 (12)	1,03 (3)		1,04 (5)	1,02 (3)	1,032 (12)		1,03 (2)
114,0		0,062 (12)						0,062 (12)
121,22		0,0070 (7)						0,0070 (7)
148,6	2,20 (22)	1,86 (3)	1,91 (4)	1,85 (7)	1,863 (17)	1,855 (16)	1,863 (8)	1,863 (8)
159,9	0,078 (8)	0,0671 (15)			0,0654 (19)	0,0651 (14)	0,06321 (40)	0,0645 (9)

The absolute emission probability of $\gamma_{6,4}$ (100,94 keV) has been deduced from the ratio of $P_{\gamma}(\gamma_{6,4}; 100,94 \text{ keV}) / P_{\gamma}(\gamma_{6,5}; 56,76 \text{ keV}) = 5,87$ which has been calculated in 1995Ak01 by using the Alaga rule.

The absolute emission probabilities of the remaining gamma rays have been adopted from 1976GuZN.

7. References

- 1952Fr25 M.S. Freedman, F. Wagner, Jr., D.W. Engelkemeir, Phys. Rev. 88, 1155 (1952) (β -transition energy, gamma-ray energies)
- 1953As40 F. Asaro, Thesis, Univ. California (1953); UCRL-2180 (1953) (α -transition energies)
- 1953Ma19 D.R. MacKenzie, M. Lounsbury, A.W. Boyd, Phys. Rev. 90, 327 (1953) (Half-life)
- 1956Ro26 B. Rose and J. Milsted, J. Nuclear Energy 2, 264 (1956) (Half-life)
- 1956Sh31 K.N. Shliagin, Izvest. Akad. Nauk SSSR, Ser. Fiz. 20, 891 (1956); Columbia Tech. Transl. 20, 810 (1957) (β -transition energy)
- 1957Ha10 G.R. Hall, T.L. Markin, J. Inorg. Nuclear Chem. 4, 137 (1957). Quoted in 1969Hanna (Half-life)
- 1960Br15 F. Brown, G.G. George, D.E. Green, D.E. Watt, J. Inorg. Nuclear Chem. 13, 192 (1960) (Half-life)
- 1961Sm03 H.L. Smith, J. Inorg. Nuclear Chem. 17, 178 (1961) (Half-life, β -transition probability)
- 1964Dz03 B.S. Dzhelepov, R.B. Ivanov, V.G. Nedovesov, Zh. Eksperim. i Teor. Fiz. 46, 1517 (1964); Soviet Phys. JETP 19, 1027 (1964) (α -transition energies)
- 1965Ba26 S.A. Baranov, M.K. Gadzhiev, V.M. Kulakov, V.M. Matinskii, Yadern. Fiz. 1, 557 (1965); Soviet J. Nucl. Phys. 1, 397 (1965) (α -transition energies)
- 1965Ba35 I.A. Baranov, V.V. Berdikov, A.S. Krivokhatskii, A.N. Silantev, Izv. Akad. Nauk SSSR, Ser. Fiz. 29, 163 (1965); Bull. Acad. Sci. USSR, Phys. Ser. 29, 161 (1966) (α -transition energies and probabilities, gamma-ray energies)
- 1966French R.J. French, F.L. Langford, Jr., W.D. Leggett, and R.J. Nodvik, Yankee Core Evaluation Program Quarterly Progress Report for the Period ending 30 September 1966. Quoted in 1987Ag03 (Half-life)
- 1966Stepan I.E. Stepan and R.G. Niske, Trans. Am. Nucl. Soc. 9, 451 (1966) (Half-life)
- 1967Shields W.R. Priv. Comm., 1967. Quoted in 1967Oe01 (Half-life)
- 1968Ah01 I. Ahmad, A.M. Friedman, J.P. Unik, Nucl. Phys. A119, 27 (1968) (α - and gamma transition energies, α/β -branching)
- 1968Ba25 S.A. Baranov, V.M. Kulakov, V.M. Shatinskii, Yadern. Fiz. 7, 727 (1968); Soviet J. Nucl. Phys. 7, 442 (1968) (α -transition energies)
- 1968Ca19 M.J. Cabell, J. Inorg. Nucl. Chem. 30, 2583 (1968) (Half-life)
- 1968Oe01 F.L. Oetting, Phys. Rev. 168, 1398 (1968) (Average β -transition energy)
- 1970Ni02 R.G. Nisle, I.E. Stephan, Nucl. Sci. Eng. 39, 257 (1970) (Half-life)
- 1970Sh18 W.R. Shields, NBS Tech. Note 546, p.25 (1970) (Half-life)
- 1971Ca15 M.J. Cabell and M. Wilkins, J. Inorg. Nucl. Chem. 33, 903 (1971) (Half-life)
- 1971GuZN R. Gunnink, R.J. Morrow, UCRL-51087 (1971) (Gamma-ray energies)
- 1972Cline J.E. Cline, R.J. Gehrke, L.D. McIsaak, ANCR-1069 (1972) (Gamma-ray energies)
- 1972Whitehead C. Whitehead, A.C. Sherwood, and B. Rose, AERE-PR/NP 18 (1972) (Half-life)
- 1973JoYT K.C. Jordan, Priv. Comm. (1973). Quoted in 1974StYG (Half-life)
- 1973Ze02 R.K. Zeigler, Y. Ferris, J. Inorg. Nucl. Chem. 35, 3417 (1973) (Half-life)
- 1974STYG W.W. Strohm and K.C. Jordan, Trans. Amer. Nucl. Soc. 18, 185 (1974) (Half-life)
- 1975WiYM M. Wilkins, UKNDC (75)-P71, p.56 (1975) (Half-life)
- 1976BaZZ S.A. Baranov, A.G. Zelenkov, V.M. Kulakov, Proc. Advisory Group Meeting on Transactinium Nucl. Data, Karlsruhe, Vol. III, p.249 (1976); IAEA -186 (1976) (α -transition energies and probabilities)

- 1976GuZN R. Gunnink, J.E. Evans, A.L. Prindle, UCRL -52139 (1976) (Gamma-ray energies and emission probabilities, α/β -branching)
- 1976McZB I.E. McKean and E.A.C. Crouch, UKNDC (76)-P80, p.41 (1976) (Half-life)
- 1976Um01 H. Umezawa, T. Suzuki, S. Ichikawa, J. Nucl. Sci. Technol. 13, 327 (1976) (Gamma - ray energies and emission probabilities)
- 1977Crouch E.A. Crouch and I.C. McKean, UKNDC (78) -P88, AERE Harwell, Oxfordshire, p.96 (1978) (Half-life)
- 1977RGZZ Priv.Comm. (June 1977) Research Groups, Atlantic Richfield Hanford Lab., Savannah River Lab., Livermore Lab., Mound Lab., Rocky Flats Lab., LASL (1977) (Half-life)
- 1977VaYR R. Vaninbroux, J. Broothaerts, P. De Bievre, B. Denecke, M. Gallet, NEANDC(E) - 192U, Vol.3, p.55 (1977) (α/β -branching)
- 1977Whitehead C. Whitehead, UK Prog. Rep.NEANDC(E)-182-8, 41 (1977) (Half-life)
- 1978DiZU J.K. Dickens, J.S. Emery, R.M. Freestone, T.A. Love, J.W. McConnell, K.J. Northcutt, R.W. Peelle, ORNL/NUREG/TM-223 (1978) (Gamma-ray emission probabilities)
- 1978El02 Y.A. Ellis, Nucl. Data Sheets 23, 123 (1978) (Half-life, decay data evaluation)
- 1978RGZZ REPT COO-535-766, p100, Research Group (1978) (Half-life)
- 1978VaZC R. Vaninbroux, Proc. Intern. Conf. Neutron Physics and Nuclear Data, Harwell, p.235 (1978) (Half-life)
- 1979Ce04 A. Cesana, G. Sandrelli, V. Sangiust, M. Terrani, Energ.Nucl.(Milan) 26, 526 (1979) (Gamma-ray energies)
- 1979Garner E.I. Garner and L.A. Machian, Trans. Amer. Nucl. Soc. 33, suppl.1, 3 (1979) (Half - life)
- 1980Ag02 S.K. Aggarwal and H.C. Jain, Phys. Rev. C21, 2033 (1980) (Half-life)
- 1980Ma45 S.F. Marsh, R.M. Abernathy, R.J. Beckman, J.E. Rein, Int .J. Appl. Radiat. Isotop. 31, 629 (1980) (Half-life)
- 1981Ag01 S.K. Aggarwal, S.N. Acharya, A.R. Parab, H.C. Jain, Phys. Rev. C23, 1748 (1981) (Half-life)
- 1981Ag07 S.K. Aggarwal, S.N. Acharya, A.R. Parab, H.C. Jain, Radiochim. Acta 29, 65 (1981) (Half-life)
- 1982Ag01 S.K. Aggarwal, S.A. Chitambar, A.R. Parab, H.C. Jain, Radi ochem. Radioanal. Lett. 54, 83 (1982) (Half-life)
- 1982Hiyama T. Hiyama, Y. Wada, K. Onishi, Ann. Prog. Rep. of Power Reactor and Nucl. Fuel Develop. Corp., Tokai Works, April 1981 - March 1982. PNCT -N-831-82-01. p.96 (1982) (Half-life)
- 1983DeZX P. De Bievre, M. Gallet, R. Werz, Int. J. Mass Spectrometry and Ion Physics 51, 111 (1983) (Half-life)
- 1984Gl03 K.M. Glover, Int. J. Appl. Radiat. Isotop. 35, 239 (1984) (α -transition energies)
- 1985Ag02 S.K. Aggarwal, A.R. Parab, S.A. Chitambar, H.C. Jain, Phys. Rev. C31, 1885 (1985) (Half-life)
- 1985Dr09 A.A. Druzhinin, V.N. Polynov, A.M. Korochkin, E.A. Nikitin, L.I. Lagutina, At. Energ. 59, 68 (1985); Sov. At. Energy 59, 628 (1985) (Half -life of the spontaneous fission)
- 1985He02 R.G. Helmer, C.W. Reich, Int. J. Appl. Radiat. Isotop. 36, 117 (1985) (Gamma -ray energies and probabilities)
- 1985Wi04 H. Willmes, T. Ando, R.J. Gehrke, Int. J. Appl. Radiat. Isotop. 36, 123 (1985) (Gamma-ray energies and emission probabilities)
- 1986Ha06 M.R. Harston and N.C. Pyper, Phys. Rev. Lett. 56, 1790 (1986) (Half -life, chemical dependence)
- 1986Ti04 G.A. Timofeev, V.V. Kalygin, P.A. Privalova, At. Energ. 60, 287 (1986); Sov. At. Energy 60, 343 (1986) (Half-life)
- 1987Ag03 S.K. Aggarwa and H.C.Jain, J. Radioanal. Nucl.Chem. 109, 183 (1987) (Half -life, review)

- 1987Ba84 I.M. Band, M.A. Listengarten and M.B. Trzhaskovskaya, *Izv. Akad. Nauk SSSR, Ser. Fiz.* 51, 1998 (1987); *Bull. Acad. Sci. USSR, Phys. Ser.* 51, No.11, 112 (1987) (Half-life, chemical dependence)
- 1988ChZL V.P. Chechev, N.K. Kuzmenko, V.O. Sergeev and K.P. Artamonova, *Evaluated Decay Data of Transuranium Radionuclides, Handbook*, Publishing House Energoatomizdat, Moscow (1988) (Evaluation of ²⁴¹Pu decay data)
- 1989Ho24 N.E. Holden, *Pure Appl. Chem.* 61, 1483 (1989) (Half-life, compilation)
- 1989Pa21 J.L. Parker, R.N. Likes, A. Goldman, *Appl. Radiat. Isot.* 40, 793 (1989) (Half-life)
- 1991Ry01 A. Rytz, *At. Data Nucl. Data Tables* 47, 205 (1991) (α -transition energies)
- 1993Dr05 T. Draznev, *Appl. Radiat. Isot.* 44, 613 (1993) (Gamma-ray energies)
- 1994Ba91 D.T. Baran, *Appl. Radiat. Isot.* 45, 1177 (1994) (Gamma-ray emission probabilities)
- 1995Ak01 Y.A. Akevali, *Nucl. Data Sheets* 74, 461 (1995) (Decay scheme, multipolarities)
- 1996FiZX R.B. Firestone, *Table of Isotopes, Eighth Edition, Volume II: A=151 -272*, V.S. Shirley (Editor), C.M. Baglin, S.Y.F. Chu, and J. Zipkin (Assistant Editors), 1996, 1998, 1999 (Decay scheme, gamma ray energies, multipolarities and level energies)
- 1996Sc06 E. Schönfeld and H. Janßen, *Nucl. Instrum. Methods Phys. Res.* A369, 527 (1996) (Atomic data)
- 1997DeZY P. De Bievre, A. Verbruggen, *Proc. Intern. on Nuclear Data for Science and Technology, Trieste, Italy, 19 -24 May, 1997*, G. Reffo, A. Ventura, C. Grandi, Eds., Editrice Compositori, Italy, Pt.1, p.839 (1997) (Half-life)
- 1998Ak04 Y.A. Akevali, *Nucl. Data Sheets* 84, 1 (1998) (r_0 of ²³⁷U)
- 1999Dr13 O. Dragoun, A. Spalek, M. Rysavy, A. Kovalik, E.A. Yakushev, V. Brabec, A.F. Novgorodov, N. Dragounova, J. Rizek, *J. Phys.(London) G25*, 1839 (1999) (β -transition energy)
- 1999ScZX E. Schönfeld and G. Rodloff - PTB-6.11-1999-1999-1, Braunschweig, February 1999 (KX ray energies and relative emission probabilities)
- 1999YaZX E.A. Yakushev, V.M. Gorozhankin, O. Dragoun, A. Kovalik, A.F. Novgorodov, M. Rysavy, A. Shpalek, *Program and Thesis, Proc. 49th Ann. Conf. Nucl. Spectrosc. Struct. At. Nuclei, Dubna*, p.118 (1999) (β -transition energy)
- 2000Dr02 O. Dragoun, A. Spalek, M. Rysavy, A. Kovalik, E.A. Yakushev, V. Brabec, J. Frana, D. Venos, *Appl. Radiat. Isot.* 52, 387 (2000) (β -transition energy)
- 2003Au03 G. Audi, A.H. Wapstra, and C. Thibault, *Nucl. Phys.* A729, 337 (2003) (Q value)
- 2004Fo01 N. Fotiades, G. D. Johns, R. O. Nelson, M. B. Chadwick, M. Devlin, M. S. Wilburn, P. G. Young, J. A. Becker, D. E. Archer, L. A. Bernstein, P. E. Garrett, C. A. McGrath, D. P. McNabb, and W. Younes, *Phys. Rev. C* 69, 024601 (2004) (Placement of 121.2 keV γ -ray transition)
- 2005Ma88 M.J. Martin, *Nucl. Data Sheets* 106, 89 (2005) (Evaluation of β -transition energy, α/β -branching)
- 2006Ba41 M.S. Basania, *Nucl. Data Sheets* 107, 2323 (2006) (Decay scheme, multipolarities)

²⁴¹Am - Comments on evaluation of decay data by V. P. Chechev and N. K. Kuzmenko

This evaluation was done originally in October 2002, revised in January 2004 and then updated in September 2009 with a literature cut-off by the same date.

1 Decay Scheme

The scheme of ²⁴¹Am decay is rather complex. It contains more than forty excited levels in ²³⁷Np populated by alpha- and gamma-ray transitions (2006Ba41, 1995Ak01). The intense population takes place only for lower levels with the energy less than 230 keV (8 excited levels and ground state in ²³⁷Np) and in this part the decay scheme is mainly defined. Nevertheless here there are some gamma-ray transitions scarcely studied and expected but not certainly observed such as 27-keV, 54-keV, 97-keV that leads to not so good intensity balance for some levels. Additional difficulties are due to anomalous internal conversion of the 26-keV and 59-keV gamma ray transitions because of “penetration effects” (1996Jo28, 2008Go10).

For high levels the decay scheme has not been completed yet since many observed gamma-ray transitions were not placed and some expected gamma transitions were not observed. The population of these levels does not exceed 0,1 %.

The unplaced gamma rays carry $\leq 0,6$ % of the total intensity of all the gamma rays placed in the decay scheme.

2 Nuclear Data

Q value is from Audi et al. (2003Au03).

The recommended ²⁴¹Am half-life is based on the experimental results given in Table 1.

Table 1. Experimental values of ²⁴¹Am half-life (in years).

Reference	Author(s)	Original value	Measurement method
1967Oe01	Oetting and Gunn	432,7 (7)	Calorimetry
1968Br22	Brown and Propst	433 (7)	Specific Activity Determination
1968St02	Stone and Hulet	436,6 (30)	Specific Activity Determination
1972Jo07	Jove and Robert	426,3 (21)	Calorimetry
1974StYG	Strohm and Jordan	432,5 (7)	Calorimetry
1974StYZ		435,0 (7)	Specific Activity Determination
1974Po16	Polyukhov et al.	432,8 (16)	Specific Activity Determination
1975Ra35	Ramthun and Muller	432,0 (2)	Calorimetry

The values before 1967 have been omitted due to their large systematic uncertainties (those values lead to the ²⁴¹Am half-life of 458 years).

The eight values were used for statistical processing. The uncertainty of 1975Ra35 was increased to 0,38 a to adjust weights according to the LRSW method.

Statistical processing of the final data set with the reduced χ^2 of 3,58 gives the unweighted mean of 432,6 (11) years and the weighted mean of 432,6 with an internal uncertainty of 0,27 and an external uncertainty of 0,51.

The LWEIGHT computer program has used the weighted mean and expanded the uncertainty to 0,6 so range includes the most precise value of 432,0 (1975Ra35). Therefore, the recommended value of ²⁴¹Am half-life is 432,6 (6) years.

The value of $1,2 (3) 10^{14}$ years has been adopted for ²⁴¹Am spontaneous fission half-life as recommended in 2000Ho27.

2.1 α Transitions

The energies of the alpha transitions have been deduced from the Q value and ²³⁷Np level energies given in Table 2 from 2006Ba41 where they were deduced from a least-squares fit to gamma ray energies.

Table 2. ²³⁷Np levels populated in ²⁴¹Am α -decay.

Level number	Energy (keV)	Spin and parity	Half-life	Probability of α -transition ($\times 100$)
0	0,0	5/2 ⁺	2,144 (7) 10 ⁶ yr	0,38 (1)
1	33,19629 (22)	7/2 ⁺	54 (24) ps	0,23 (1)
2	59,54092 (10)	5/2 ⁻	67 (2) ns	84,45 (10)
3	75,899 (5)	9/2 ⁺	~ 56 ps	< 0,04
4	102,959 (3)	7/2 ⁻	80 (40) ps	13,23 (10)
5	129,99 (3)	11/2 ⁺		~ 0,01
6	158,497 (11)	9/2 ⁻		1,66 (3)
7	191,53 (6)	13/2 ⁺		
8	225,957 (16)	11/2 ⁻		0,014 (3)
9	267,556 (17)	3/2 ⁻	5,2 (2) ns	5 10 ⁻⁴
10	281,356 (20)	1/2 ⁻		
11	305,05 (3)	13/2 ⁻		0,0022 (3)
12	316,8 (2) ?			
13	324,420 (23)	(7/2 ⁻)		0,0013
14	332,376 (16)	1/2 ⁺	≤ 1 ns	
15	359,7 (1)	(5/2 ⁻)		6 10 ⁻⁴
16	368,602 (20)	5/2 ⁺		9 10 ⁻⁴
17	370,928 (23)	3/2 ⁺		3 10 ⁻⁴
18	395,53 (4)	15/2 ⁻		7 10 ⁻⁴
19	418,2 (1) ?			
20	434,12 (5)	(11/2 ⁻)		4 10 ⁻⁴
21	444,78 (10) ?			
22	452,545 (22)	9/2 ⁺		~ 4 10 ⁻⁴
23	459,693 (24)	7/2 ⁺		~ 4 10 ⁻⁴
24	486,21 (9)	(9/2 ⁻)		1,1 10 ⁻⁴
25	497,01 (5)	17/2 ⁻		
26	514,19 (4)	(3/2 ⁻)		
27	546,12 (6)	(5/2 ⁻)		1 10 ⁻⁴
28	590,09 (4)	(7/2 ⁻)		
29	592,33 (7)	13/2 ⁺		
30	597,99 (9)	11/2 ⁺		
31	646,03 (17)	(9/2 ⁻)		
32	666,19 (10)	(5/2 ⁺ , 7/2 ⁻)		
33	721,961 (13)	5/2 ⁻		7 10 ⁻⁴

Level number	Energy (keV)	Spin and parity	Half-life	Probability of α -transition ($\times 100$)
34	755,685 (19)	7/2 ⁻		8,6 10 ⁻⁵
35	770,57 (5)			
36	799,82 (4)	9/2 ⁻		4 (3) 10 ⁻⁵
37	805,77 (12)	(7/2 ⁺ , 9/2 ⁺)		
38	853,36 (15)	11/2 ⁻		
39	861,65 (19)	(5/2 ⁺ , 7/2)		
40	920,88 (20)			
41	946 (2)			
42	962 (3) ?			
43	1014 (3) ?			

The probabilities of the alpha transitions $\alpha_{0,0}$, $\alpha_{0,1}$, $\alpha_{0,2}$, $\alpha_{0,4}$ and $\alpha_{0,6}$ have been obtained by averaging experimental values from the spectrometric measurements carried out for the last twenty five years (Table 3). Earlier measurements for these alpha transitions see in 2006Ba41.

Table 3. Experimental and recommended probabilities (%) of the most intense alpha transitions.

	α -particle energy (keV)	1984Ah06 1993Ahmad	1987Bo25	1994B112	1996 Bueno	1996 Sanchez	1998Ya17	Recommended
$\alpha_{0,0}$	5544	0,36 (1)	0,34 (5)	0,36 (5)	0,5 (2)	0,36 (3)	0,394 (9)	0,38 (1)
$\alpha_{0,1}$	5511	0,23 (1)	0,22 (3)	0,22 (6)	-	0,28 (3)	0,224 (7)	0,23 (1)
$\alpha_{0,2}$	5486	84,6 (2) ^a	84,7 (9)	84,69 (28)	84,5 (8)	84,5 (3)	84,30 (7)	84,45 (10)
$\alpha_{0,4}$	5443	13,1 (1) ^a	13,0 (3)	13,08 (24)	12,5 (3)	13,2 (3)	13,40 (8)	13,23 (10)
$\alpha_{0,6}$	5388	1,65 (8)	1,6 (1)	1,66 (6)	1,6 (2)	1,65 (7)	1,67 (2)	1,66 (3)

^aThe $\alpha_{0,2}$ and $\alpha_{0,4}$ probabilities from 1984Ah06 were superseded by the same author in 1993Ahmad. The latter values are given in Table 3.

The probabilities of the alpha transitions $\alpha_{0,3}$, $\alpha_{0,5}$, $\alpha_{0,9}$, $\alpha_{0,13}$, $\alpha_{0,15}$, $\alpha_{0,33}$ have been adopted from the magnetic spectrometer measurements of 1964Ba26. The probabilities of the $\alpha_{0,8}$ and $\alpha_{0,11}$ transitions have been obtained from measurements of 1955Go57, 1964Ba26 and 1965Mi06. The probabilities of the $\alpha_{0,34}$ and $\alpha_{0,36}$ transitions have been deduced from the intensity balance of gamma transitions.

2.2 γ Transitions

The recommended energies of the gamma-ray transitions are virtually the same as the gamma-ray energies because nuclear recoil is negligible for ²³⁷Np.

The gamma-ray transition probabilities have been deduced from their gamma-ray emission probabilities and the evaluated total ICC's.

ICC's for the intense E1 anomalously converted gamma-ray transitions $\gamma_{2,1}$ (26,3 keV) and $\gamma_{2,0}$ (59,5 keV) have been obtained from a joint analysis of the gamma ray and L-, M- conversion electron probabilities measured in ²⁴¹Am α decay and ²³⁷U β^- decay (1996Jo28, 2006Ba41). Experimental conversion electron data are given in 1959Sa10, 1964Wo03, 1966Ko06, 1966Le13, 1966Ya05, and 1998Ko61. For discussion of anomalous electric dipole gamma-ray transitions see 1960As02, 1966Ya05, 1967Pa23, 1970Gr36, 1996Jo28, and 2008Go10. In 2008Go10 an assessment of ICCs for a number of such transitions was made. In particular, the total ICCs for gamma-ray transitions $\gamma_{2,1}$ (26,3 keV) and $\gamma_{2,0}$ (59,5 keV) in ²³⁷Np have been assessed as 7,9 (8) and 0,99 (9), respectively.

ICC's for other gamma transitions have been interpolated using the BrIcc computer program, version v2.2a, data set BriccFO (2008Ki07). Multipolarities of the gamma-ray transitions and E2/M1 mixing ratios

have been adopted from 2006Ba41 based on the measurements of 1959Sa10, 1964Wo03, 1966Ko06, 1966Ya05, 1998Ko61.

The E2 admixture of 16,6 (25) % for M1+E2 gamma-ray transition $\gamma_{4,2}(43,4\text{-keV})$ has been obtained by averaging the four measurement results from 1964Wo03 (17,6 (19) %), 1966Ko06 (13 (2) %), 1966Ya05 (11 (4) %), and 1998Ko61 (21,2 (22) %).

3 Atomic Data

The atomic data (fluorescence yields, X-ray energies and relative probabilities, and Auger electrons energies and relative probabilities) were deduced by using the Saisinuc software (2002Be). The fluorescence yield ω_M is from 1989Hubbell.

The XL -ray energies are taken from 2001Sc08.

The XK -ray energies are taken from 1999Schönfeld. Below these calculated (adopted) values are compared with the experimental results of 1982Ba56 and 1983Ah02:

	Calculated (1999Schönfeld)	Measured in 1982Ba56	Measured in 1983Ah02
$K\alpha_2$	97,069	97,069 (3)	97,08 (2)
$K\alpha_1$	101,059	101,057 (3)	101,07 (2)
$K\beta_3$	113,303	113,308 (4)	113,30 (2)
$K\beta_1$	114,234	114,244 (3)	114,24 (2)
$K\beta_5$	114,912	-	114,95 (2)
$K\beta_2$	117,463		} 117,51 (3)
$K\beta_4$	117,876		
$KO_{2,3}$	118,429	-	118,45 (5)

4 α Emissions

The recommended energies of alpha particles have been deduced from the energies of alpha transitions taking into account the recoil energies for ^{237}Np .

The experimental values of the alpha particle energies from spectrometric measurements are given in 1971Gr17, 1968Ba25, 1968Ka09, 1965Mi06, 1964Ba26, 1962Le11, 1957Ro20, 1955Go57 (see also 2006Ba41). Most of them have lesser accuracy in comparison with the recommended values.

5 Electron emissions

The energies of the conversion electrons have been obtained from the gamma-ray transition energies and the atomic electron binding energies.

The emission probabilities of the conversion electrons have been deduced using the evaluated P_γ and ICC values. The total absolute emission probabilities of K and L Auger electrons have been calculated using the EMISSION computer program.

6 Photon emissions

6.1 X-ray emissions

The total absolute emission probability of Np MX - rays is the experimental result of 1971Ka48.

The recommended absolute emission probabilities of Np LX - rays have been obtained by averaging of experimental results (per 100 disintegrations) shown in Table 4.

Table 4. Experimental and recommended absolute Np LX-ray emission probabilities (%) ^a.

	1971 Ge11	1971 Wa28	1974 Ca16	1976 GuZN	1980 Cohen	1988 Co07	1992 Bl07	1994 Le37	2008 Le07	Recom- mended	2001Sc08 (calculated)
L1	0,81 (7)	0,87 (6)	0,86 (2)	0,806 (40)	0,87 (3)	0,83 (3)	0,837 (10)	0,864 (12)	0,837 (9)	0,844 (9) ^b	0,842 (27)
L α	12,6 (9)	13,5 (12)	13,20 (25)	13,2 (7)	13,2 (3)	12,7 (4)	13,01 (10)	13,03 (13)	13,00 (12)	13,02 (10) ^b	13,3 (4)
L η	} 19,1 (14)	19,1 (14)	19,25 (40)	19,2 (10)	19,78 (36)	0,368 (5)	0,377 (15)	0,369 (12)	0,404 (5)	0,384 (20) ^c	0,383 (16)
L β											
L γ	4,75 (35)	4,75 (35)	4,85 (15)	4,94 (25)	4,96 (20)	4,8 (2)	4,815 (38)	4,74 (8)	4,84 (3)	4,83 (3) ^b	5,17 (14)

^a In addition to given references the value of 19,46 (16) for L η +L β was obtained in 1974Ga40.

^b The smallest uncertainty of the experimental results.

^c The LWEIGHT computer program has used the weighted mean of 0,3843 and expanded the uncertainty so range includes the most precise value of 2008Le07.

The experimental results of 1993Lépy (per 100 disintegrations) are quoted in 2001Sc08: L1 - 0,875 (18), L α - 13,10 (21), L η - 0,354 (8), L β - 18,5 (4), L γ - 4,84 (8). These results were superseded in 1994Le37 and were not used by the evaluators for statistical processing.

The evaluated total absolute emission probability of LX - rays P(XL) = 37,66 (17) % can be compared with the value of 36,8 (21) % calculated using the EMISSION computer program.

The absolute emission probabilities of Np XK -rays have been calculated using the EMISSION computer program. The recommended value of the total absolute emission probability P(XK) = 0,003 82 (10) % can be compared with measurements of 1976GuZN which give P(XK) = 0,004 01 (10) %.

Below the experimental data of 1976GuZN are compared with the calculated values of absolute emission probability for KX-ray components:

	1976GuZN (measured) ^a	Recommended (calculated)
K α_2	0,001 18 (4)	0,001 134 (30)
K α_1	0,001 89 (6)	0,001 81 (5)
K β_1	7,1 (3) 10 ⁻⁴	6,58 (21) 10 ⁻⁴
K β_2	2,29 (15) 10 ⁻⁴	2,26 (8) 10 ⁻⁴

^a The uncertainties quoted in 1976GuZN have been increased by 2 % to allow for the uncertainty of the detector calibration.

6.2 Gamma-ray emissions

6.2.1 Gamma-ray energies

The gamma ray energies have been taken mainly from 2006Ba41 (see also the evaluation of 1988ChZL). Some gamma ray energies have been deduced directly from the adopted ²³⁷Np level energies.

The recommended gamma ray energy values are based on measurements of 1955Da02, 1959Sa10, 1964Wo03, 1966Ko06, 1966Ya05, 1968Je01, 1968Ka09, 1970Ne11, 1976GuZN, 1978Ge06, 1978Ge17, 1978Ov01, 1979Ar11, 1984Ov02, and 1998Ab43.

The energies of gamma rays $\gamma_{2,1}$ (26,3 keV) and $\gamma_{2,0}$ (59,5 keV) have been adopted from 2000He14. The energy of gamma ray $\gamma_{1,0}$ (33,2 keV) has been deduced as the difference of E $\gamma_{2,0}$ - E $\gamma_{2,1}$. The energies of gamma rays $\gamma_{3,1}$ (42,7 keV), $\gamma_{4,2}$ (43,4 keV), and $\gamma_{8,4}$ (123,0 keV) have been taken from 1998Ko61. The gamma ray with energy of 32,183 keV has been adopted from 1976GuZN and was not reported by others.

The energies of gamma rays $\gamma_{27,26}$ (31,9 keV), $\gamma_{17,14}$ (38,5 keV), $\gamma_{14,10}$ (51,0 keV), $\gamma_{5,3}$ (54,1 keV), $\gamma_{13,9}$ (56,9 keV), $\gamma_{7,5}$ (61,6 keV), $\gamma_{14,9}$ (64,8 keV), $\gamma_{36,33}$ (77,9 keV), $\gamma_{11,8}$ (79,0 keV), $\gamma_{15,9}$ (92,4 keV) and $\gamma_{5,1}$ (96,8 keV) have been deduced from the adopted ²³⁷Np level energies. These gamma ray transitions were not observed in the ²⁴¹Am α -decay; they are expected from the decay scheme (see 2006Ba41).

The gamma rays $\gamma_{20,11}$ (129,1 keV), $\gamma_{23,13}$ (135,3 keV), $\gamma_{30,23}$ (138,3 keV) and unplaced in decay scheme gamma rays with energies of 128,05 keV and 136,7 keV have been adopted from 1979Ar11 and were not observed by others.

Many unplaced gamma rays are reported only in 1998Ab43.

6.2.2 Gamma-ray emission probabilities

The recommended absolute emission probabilities ($P\gamma$) of the most intense gamma rays $\gamma_{1,0}$ (26,3 keV), $\gamma_{2,1}$ (33,2 keV), $\gamma_{4,2}$ (43,4 keV) and $\gamma_{2,0}$ (59,5 keV) have been deduced from the available experimental data (Table 5).

Table 5. Experimental and recommended values of the most intense gamma rays in ²⁴¹Am α -decay.

Reference	$P\gamma_{1,0}$ (26,3 keV) $\times 100$	$P\gamma_{2,1}$ (33,2 keV) $\times 100$	$P\gamma_{4,2}$ (43,4 keV) $\times 100$	$P\gamma_{2,0}$ (59,5 keV) $\times 100$
1952Be24	2,8 (3)			40,0 (15)
1957Ma17	2,5 (2)		0,073 (7)	35,9 (6)
1964Mc12				34,6 (7)
1965Mi06				38,0 (6)
1969Pe17				35,3 (6)
1971Ge11	2,23 (18)	0,104 (11)	0,057 (18)	
1974Ca16	2,4 (1)			
1975Le09				36,3(4)
1976GuZN	2,45 (5)			
1976Pl05				35,5 (3)
1978Ge06	2,54 (26)	0,106 (11)	0,073 (7)	
1983Ah02		0,125 (8)		
1983De11	2,41 (5)			
1983Hu04				35,82 (17) ^d
1984Ov02		0,12 (1)	0,066 (5)	
1987De22				36,36 (17)
1992Bl07	2,395 (19)	0,1233 (28)	0,0654 (29)	36,03 (25)
1992Ma16				35,6 (2)
2005Iw01	2,06 (3)			35,87 (17)
Recommended	2,31 (8)^a	0,1215 (28)^b	0,0669 (29)^c	35,92 (17)^e

^a The LWEIGHT computer program has used the weighted mean of 2,31 and expanded the uncertainty so range includes the most precise value of 1992Bl07.

^b The LWEIGHT computer program has used the weighted mean of 0,12148 and external uncertainty of 0,0028. The smallest value of experimental uncertainties is also 0,0028.

^c The LWEIGHT computer program has used the weighted mean of 0,0669 and internal uncertainty of 0,0022. The smallest value of experimental uncertainties is 0,0029.

^d Uncertainty quoted by authors (0,12) has been increased to 0,17 by the evaluators to include possible systematic errors in correction factors to 59,5-keV-peak counting rate.

^e The LWEIGHT computer program has identified one by one the three outliers of 1952Be24, 1965Mi06 and 1964Mc12 and used the weighted mean of 35,92 (8). The smallest value of experimental uncertainties of 0,17 has been adopted as the uncertainty.

The absolute emission probabilities of gamma rays $\gamma_{3,1}$ (42,7 keV), $\gamma_{6,4}$ (55,6 keV), $\gamma_{57,8}$ (57,8 keV), $\gamma_{8,6}$ (67,5 keV), and $\gamma_{4,1}$ (69,8 keV) have been adopted from the measurements of 1978Ge06.

The absolute emission probabilities of gamma rays $\gamma_{6,2}$ (99,0 keV), $\gamma_{4,0}$ (103,0 keV), $\gamma_{8,4}$ (123,0 keV), and $\gamma_{6,1}$ (125,3 keV) have been adopted from the measurements of 1976GuZN.

The remaining weak gamma ray emission probabilities ($P_\gamma < 10^{-5}$) have been adopted from the evaluations of 2006Ba41 and 1988ChZL, based mainly on the measurements of 1976GuZN and 1978Ge17 with Ge(Li) detectors, and (for gamma rays with energy more than 200 keV) from the measurements of 1998Ab43 with 40 % HPGe detector and intense purified sources. The uncertainties quoted in 1998Ab43 have been increased by 1 % to allow for the uncertainty of the detector calibration.

Other measurements of P_γ are given in 19840v02, 1983Hu04, 1983De11, 1983Ah02, 1979Ce04, 1978Ge06, 1976Pl05, 1975Le09, 1974Ca16, 1974HeYW, 1971Ge11, 1971Cl03, 1967Gu08, 1967Br26, 1966Ko06, 1965Mc12, 1965Be38, 1957Ro20, 1957Ma17, 1956Ho38, 1955Tu13, 1955Ja01, 1955Da02, 1955Ba31, and 1952Be24.

The gamma ray emission probabilities quoted in 1976GuZN and also in 19840v02, 1978Ge06, 1974Ca16, 1971Ge11, 1967Gu08 have been normalized to P_γ (59,54 keV) = 0,3592. The gamma ray emission probabilities from 1971Cl03, 1978Ge17 have been normalized to P_γ (208,00 keV) = $7,86 \cdot 10^{-6}$.

7 Consistency of recommended data

The most accurate Q value, Q(M), is taken from the atomic mass adjustment table of Audi et al. (2003Au03). Comparison of Q(eff) (deduced as the sum of average energies per disintegration ($\sum E_i \times P_i$) for all emissions accompanying ^{241}Am α - decay) with the tabulated decay energy Q(M) allows to check a consistency of the recommended decay-scheme parameters obtained in this evaluation.

Here E_i and P_i are the evaluated energies and emission probabilities of the i-th alpha particle, gamma ray, X-ray, etc. Consistency (percentage deviation) is determined by $\{|Q(M) - Q(\text{eff})\} / Q(M)\} \times 100$. "Percentage deviations above 5 % would be regarded as high and imply a poorly defined decay scheme; a value of less than 5 % indicates the construction of a reasonably consistent decay scheme" (quoted from the article by A.L. Nichols in Appl. Rad. Isotopes 55 (2001) 23-70).

For the above ^{241}Am decay data evaluation we have Q(M) = 5637,82 (12) keV and Q(eff) = 5638 (8) keV, i.e. consistency is better than 0,15 %.

8 References

- 1952Be24 J. K. Beling, J. O. Newton, B. Rose, Phys. Rev. 86(1952)797; Erratum Phys. Rev. 87(1952) 1144 (gamma-ray emission probabilities).
- 1955Da02 R. B. Day, Phys. Rev. 97(1955)689 (gamma-ray emission probabilities).
- 1955Go57 L. L. Goldin, G. I. Novikova, E. F. Tretyakov, Conf. Acad. Sci. USSR Peaceful Uses of Atomic Energy, Session Div. Phys. Math. Sci., Moscow, p 226 (1955); Consultants Bureau Transl., p. 167 (energies of alpha-particles, alpha-particle emission probabilities).
- 1955Ja01 H. Jaffe, T. O. Passell, C. I. Browne, I. Perlman, Phys. Rev. 97(1955)142 (gamma-ray emission probabilities).
- 1955TU13 J. F. Turner, Phil. Mag. 46(1955)687 (gamma-ray emission probabilities).
- 1956Ho38 J. M. Hollander, W. G. Smith, J. O. Rasmussen, Phys. Rev. 102(1956)1372 (gamma-ray emission probabilities).
- 1957Ma17 L. B. Magnusson, Phys. Rev. 107(1957)161 (gamma-ray energies and emission probabilities).
- 1957Ro20 S. Rosenblum, M. Valadares, J. Milsted, J. Phys. Radium 18(1957)609 (energies of alpha-particles).
- 1959Sa10 P. S. Samoilov, Columbia Tech. Transl. 23(1959)1401 (gamma-ray energy, gamma transition probabilities and multipolarities).

- 1960As02 F. Asaro, F. S. Stephens, J. M. Hollander, I. Perlman, Phys. Rev. 117(1960)492 (anomalous electric dipole gamma-ray transitions).
- 1962Le11 C. F. Leang, Compt. Rend. 255(1962)3155 (energies of alpha-particles).
- 1964Ba26 S. A. Baranov, V. M. Kulakov, V. M. Shatinsky, Nucl. Phys. 56(1964)252 (alpha-particle energies and emission probabilities).
- 1964Wo03 J. L. Wolfson, J. J. H. Park, Can. J. Phys. 42(1964)1387. Erratum. Can. J. Phys. 48(1970)2782 (gamma-ray energies and multipolarities).
- 1965Be38 G. Bertolini, F. Cappellani, G. Restelli, Nucl. Instr. Methods 32(1965)86 (gamma-ray emission probabilities).
- 1965Mc12 L. D. McIsaac, IDO Report 17052, p. 31 (gamma-ray emission probabilities).
- 1965Mi06 W. Michaelis, Z. Phys. 186(1965)42 (alpha particle energies and emission probabilities).
- 1966Ko06 L. N. Kondratev, E. F. Tretyakov, Izv. Akad. Nauk. SSSR, Ser. Fiz. 30(1966)386; Bull. Acad. Sci. USSR, Phys. Ser. 30(1967)393 (internal conversion probabilities).
- 1966Le13 C. M. Lederer, J. K. Poggenburg, F. Asaro, J. O. Rasmussen, I. Perlman, Nucl. Phys. 84(1966)481 (internal conversion coefficients).
- 1966Ya05 T. Yamazaki, J. M. Hollander, Nucl. Phys. 84(1966)505 (internal conversion probabilities).
- 1967Br26 C. Briancon, M. Valadares, R. J. Walen, Compt. Rend. 265B(1967)1496 (gamma-ray emission probabilities).
- 1967Gu08 C. Gunther, D. R. Parsignault, Nucl. Phys. A104(1967)588 (XK-ray emission probabilities).
- 1967Oe01 F. L. Oetting, S. R. Gunn, J. Inorg. Nucl. Chem. 29(1967)26595 (half-life).
- 1967Pa23 H. -C. Pauli, K. Alder, Z. Physik 202(1967)255 (anomalous electric dipole gamma-ray transitions).
- 1968Ba25 S. A. Baranov, V. M. Kulakov, V. M. Shatinskii, Soviet J. Nucl. Phys. 7(1968)442 (energies of alpha-particles).
- 1968Br22 L. C. Brown, R. C. Propst, Inorg. Nucl. Chem. 30(1968)2591 (half-life).
- 1968Je01 R. W. Jewell, W. John, R. Massey, B. G. Saunders, Nuclear Instrum. Methods 62(1968)68 (gamma-ray energies).
- 1968Ka09 R. Kamoun, R. Ballini, S. Bergstrom-Rohlin, J.-M. Kuchly, P. Siffert, Compt. Rend. 266B(1968)1241 (energies of alpha-particles).
- 1968St02 R. E. Stone, E. K. Hulet, J. Inorg. Nucl. Chem. 30(1968)2003 (half-life).
- 1969Pe17 A. Peghaire, Nucl. Instr. Methods 75(1969)66 (gamma-ray emission probabilities).
- 1970Gr36 V. N. Grigorev, A. P. Feresin, Yad. Fiz. 12(1970)665; Sov. J. Nucl. Phys. 12(1971)361 (anomalous electric dipole gamma-ray transitions).
- 1970Ne11 G. C. Nelson, B. G. Saunders, Nuclear Instrum. Methods 84(1970)90 (gamma-ray energies).
- 1971Cl03 J. E. Cline, IN-1448(1971) (gamma -ray emission probabilities).
- 1971Ge11 R. J. Gehrke, R. A. Lokken, Nuclear Instrum. Methods 97(1971)219 (XL- and gamma -ray emission probabilities).
- 1971Gr17 B. Grennberg, A. Rytz, Metrologia 7(1971)65 (energies of alpha-particles).
- 1971Ka48 E. Karttunen, H. U. Freund, R. W. Fink, Phys. Rev. A4(1971)1695 (MX-ray emission probability)
- 1971Wa28 R. L. Watson, T. K. Li, Nucl. Phys. A178(1971)201 (LX-ray emission probabilities).
- 1972Jo07 J. Jove, R. Robert, Radiochem. Radioanal. Letters 10(1972)139 (half-life).
- 1974Ca16 J. L. Campbell, L. A. McNelles, Nuclear Instrum. Methods 117(1974)519 (LX- and gamma -ray emission probabilities).
- 1974Ga40 W. J. Gallagher, S. J. Cipolla, Nuclear Instrum. Methods 122(1974)405 (LX- ray emission probabilities).
- 1974HeYW R. L. Heath, ANCR 1000(1974)2 (gamma - ray energies and emission probabilities).
- 1974Po16 V. G. Polyukhov, G. A. Timofeev, P. A. Privalova, P. F. Baklanova, At. Energ. 36(1974)319; J. At. Energy 36(1974)402 (half-life).
- 1974StYG W. W. Strohm, K. C. Jordan, Trans. Am. Nucl. Soc. 18(1974)185 (half-life).
- 1975Le09 J. Legrand, J. P. Perolat, C. Bac, J. Gorry, Intern. J. Appl. Radiat. Isotop. 26(1975)179 (gamma - ray emission probabilities).
- 1975Ra35 H. Ramthun, W. Muller, Intern. J. Appl. Radiat. Isotop. 26(1975)589 (half-life).
- 1976GuZN R. Gunnink, J. E. Evans, A. L. Prindle, Report UCRL-52139 (1976) (LX-, KX- and gamma-ray emission probabilities).
- 1976PI05 J. Plch, J. Zderadicka, L. Kokta, Czech. J. Phys. 26B(1976)1344 (gamma-ray emission probability).

- 1978Ge06 A. Genoux-Lubain, G. Ardisson, Radiochem. Radioanal. Letters 33(1978)59 (gamma-ray energies and emission probabilities).
- 1978Ge17 A. Genoux-Lubain, G. Ardisson, Compt. Rend., B 287(1978)13 (gamma-ray energies and emission probabilities).
- 1978Ov01 V. V. Ovechkin, Izv. Akad. Nauk. SSSR, Ser. Fiz. 42(1978)101; Bull. Acad. Sci. USSR, Phys. Ser. 42(1)(1978)82 (gamma-ray energies and emission probabilities).
- 1979Ar11 C. Ardisson, A. Genoux-Lubain, V. Barci, G. Ardisson, Radiochem. Radioanal. Letters 40(1979)207 (gamma-ray energies).
- 1980Cohen D. D. Cohen, Nuclear Instrum. Meth. 178(1980)481 (LX-ray emission probabilities).
- 1982Ba56 G. Barreau, H. G. Borner, T von Egidy, R. W. Hoff, Z. Phys. A308, 209 (1982) (KX-ray energies)
- 1983Ah02 I. Ahmad, J. Hines, J.E. Gindler, Phys. Rev. C27, 2239 (1983) (LX-, KX-ray energies and KX-, gamma-ray emission probabilities)
- 1983De11 K. Debertin, W. Pessara, Intern. J. Appl. Radiat. Isotop. 34(1983)515 (gamma-ray emission probabilities).
- 1983Hu04 J. M. R. Hutchinson, P. A. Mullen, Intern. J. Appl. Radiat. Isotop. 34(1983)543 (gamma-ray emission probabilities).
- 1984Ah06 I. Ahmad, Nuclear Instrum. Methods 223(1984)319 (alpha-particle emission probabilities).
- 1984Ov02 V. V. Ovechkin, A. E. Khokhlov, Izv. Akad. Nauk SSSR, Ser. Fiz. 48(1984)1032 (gamma-ray energies and emission probabilities).
- 1987Bo25 G. Bortels, P. Collaers, Appl. Radiat. Isot. 38(1987)831 (alpha-particle emission probabilities).
- 1987De22 B. Denecke, Appl. Radiat. Isot. 38(1987)823 (gamma-ray emission probabilities).
- 1988ChZL V. P. Chechev, N. K. Kuzmenko, V. O. Sergeev, K. P. Artamonova, (1988) Transuranium Radionuclides, Handbook. Publishing House Energoatomizdat, Moscow (gamma-ray energies).
- 1988Co07 D. D. Cohen, Nucl. Instrum. Meth. Phys. Res. A267(1988)492 (LX-ray emission probabilities).
- 1989Hubbell J. H. Hubbell, NIST Internal Report 89-4144(1989).
- 1992B107 C. J. Bland, J. Morel, E. Etcheverry, M. C. Lépy, Nucl. Instrum. Meth. Phys. Res. A312(1992)323 (gamma- and LX-ray emission probabilities).
- 1992Ma16 L. J. Martin, P. A. Burns, Nucl. Instrum. Meth. Phys. Res. A312(1992)146 (gamma-ray emission probabilities).
- 1993Ahmad I. Ahmad, Private Communication, cited in 1994B112 (alpha-particle emission probabilities).
- 1993Lépy M. C. Lépy, K. Debertin, H. Janssen, and U. Schötzig, PTB report PTB-Ra-31 (LX-ray emission probabilities).
- 1994B112 C. J. Bland, Nucl. Instrum. Meth. Phys. Res. A339(1994)180 (alpha-particle emission probabilities).
- 1994Le37 M.C. Lépy, B. Duchemin, J. Morel, Nucl. Instr. Meth. Phys. Res., A353(1994)10 (LX-ray emission probabilities).
- 1995Ak01 Y. A. Akovali, Nucl. Data Sheets 74(1995)461 (Decay scheme).
- 1996Bueno C. C. Bueno, J. A. C. Gonçalves, M. D. S. Santos, Nucl. Instr. Meth. Phys. Res., A371(1996)460 (alpha-particle emission probabilities).
- 1996Jo28 P. N. Johnston Nucl. Instr. Meth. Phys. Res., A369(1996)107 (evaluated gamma-ray emission probabilities and internal conversion coefficients).
- 1996Sanchez A. M. Sanchez, P. R. Montero, F. V. Tome, Nucl. Instr. Meth. Phys. Res. A369(1996)593 (alpha-particle emission probabilities).
- 1998Ab43 A. Abdul-Hadi, J. Radional. Nucl. Chem. 231(1998)147 (gamma-ray energies and emission probabilities).
- 1998Ko61 A. Kovalik, E. A. Yakushev, V. M. Gorozhankin, M. Novgorodov, M. Rysavy, J. Phys. (London) G24(1998)2247 (energies and emission probabilities of conversion electrons).
- 1998Ya17 J. Yang, J. Ni, Nucl. Instrum. Meth. Phys. Res. A413(1998)239 (alpha-particle emission probabilities).
- 1999Schönfeld E. Schönfeld, G. Rodloff, PTB-6. 11-1999-1, Braunschweig, Februar 1999 (XK-ray energies and emission probabilities).
- 2000He14 R. G. Helmer, C. van der Leun, Nucl. Instr. Meth. Phys. Res., A450(2000)35 (gamma-ray energies).

- 2000Ho27 N. E. Holden, D. C. Hoffman, Pure Appl. Chem. 72(2000)1525 (²⁴¹Am spontaneous fission half-life).
- 2001Sc08 E. Schönfeld, U. Schötzig, Appl. Radiat. Isot. 54(2001)785 (calculated absolute emission probabilities of LX-rays).
- 2002Bé M.M. Bé, R. Helmer, V. Chisté, J. Nucl. Sci. Tech., 2(2002)481 (Saisinuc software).
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)337 (Q value)
- 2005Iw01 A. Iwahara, M. A. L. da Silva, A. E. Carvalho Filho, E. M. de Oliveira Bernardes, J. U. Delgado, Appl. Radiat. Isot. 63(2005)107 (absolute emission probabilities of gamma rays).
- 2006Ba41 M. S. Basunia, Nucl. Data Sheets 107(2006)2323 (²⁴¹Am decay scheme, ²³⁷Np level energies and γ -ray transition multipolarities).
- 2008Go10 V. M. Gorozhankin, M. -M. Bé, Appl. Radiat. Isot., 66(2008)722 (ICC for anomalous E1 γ -ray transitions).
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr, Nucl. Instrum. Meth. Phys. Res. A589(2008)202 (Band-Raman ICC for γ -ray transitions).
- 2008Le07 M. C. Lépy, J. Plagnard, L. Ferreux, Appl. Radiat. Isot. 66(2008)715 (absolute emission probabilities of LX-rays).

²⁴²Pu - Comments on evaluation of decay data by V. P. Chechev

This evaluation was done originally in 2004 (2004BeZQ, 2005ChZU) and then updated in June 2009 with a literature cut-off by the same date.

1 Decay Scheme

The decay scheme can be basically considered completed though weak alpha transitions to some highly excited ²³⁸U levels (with energy more than 307 keV, see 2002Ch52) are possible but have not been observed yet. They are expected from data on level spins and $Q(\alpha)$ value and cannot appreciably influence intensity balances at the four lower levels well established.

2 Nuclear Data

$Q(\alpha)$ value is from 2003Au03.

The recommended half-life of ²⁴²Pu is based on the experimental results given in Table 1. Re-estimated values were used for averaging where necessary.

Table 1. Experimental values of ²⁴²Pu half-life (in 10⁵ years).

Reference	Author(s)	Original value	Re-estimated value	Measurement method
1956Bu64	Butler et al.	3.73 (5)	3.65 (5) ^a	²⁴² Pu/ ²³⁸ Pu, mass- and α -spectrometry
1956Bu92	Butler et al.	3.79 (5)		Specific activity, ionization chamber
1956Me37	Metch et al.	3.88 (10)	3.85 (10) ^a	²⁴² Pu/ ²⁴⁰ Pu, mass- and α -spectrometry
1969Be06	Bemis et al.	3.869 (16)	3.82 (3) ^b	²⁴² Pu/ ²³⁹ Pu, mass- and α -spectrometry
1970Du02	Durham and Molson	3.66 (7)	3.67 (7) ^a	²⁴² Pu/ ²³⁸ Pu, mass- and α -spectrometry
1976Bu23	Bulaynitsa et al.	3.702 (7) ^c		Specific activity, 4 π -X coincidences
1976Os05	Osborne and Flotov	3.763 (9)		Calorimetry
1978MeZL	Meadows	3.736 (25)	3.708 (29) ^a	²⁴² Pu/ ²³⁹ Pu, mass- and α -spectrometry
1979Ag03	Aggarwal et al.	3.742 (24)		²⁴² Pu/ ²³⁹ Pu, mass- and α -spectrometry
1979Ag03	Aggarwal et al.	3.766 (25)		²⁴² Pu/ ²³⁸ Pu, mass- and α -spectrometry

^a Re-estimated in 1979Ag03 using the values of 87.74 yr for ²³⁸Pu half-life and 24110 yr for ²³⁹Pu half-life.

^b Re-estimated in 1976Bu23 as a result of analysis of systematic uncertainties in 1969Be06 and using better values of auxiliary half-lives (see also 1979Ag03).

^c Quoted uncertainty, corresponding to 95 % confidence level, has been reduced by a factor 2.

The weighted average of the ten values is 3.7304 with the internal uncertainty 0.0051 and external uncertainty 0.0116 and $\chi^2/\nu = 3.16$. The uncertainty of 1976Bulaynitsa was increased to 0.007 24 to adjust weights according to the Limitation of Relative Statistical Weight method.

The LWRIGHT computer program has used the weighted average and expanded the uncertainty to 0.0284 so range includes the most precise value of 3.702 (1976Bu23).

The recommended value of ²⁴²Pu half-life is 3.73 (3) 10⁵ years.

The recommended spontaneous fission half-life of ²⁴²Pu is based on the experimental results given in Table 2.

Table 2. Experimental values of the spontaneous fission ²⁴²Pu half-life (in 10¹⁰ years).

Reference	Author(s)	Original value	Re-estimated value ^a	Measurement method
1956Studier	Studier and Hirsch	6.7 (7)		Quoted by Mech et al.(1956); no details available
1956Me37	Mech et al.	7.06 (19)	6.79 (19)	α /SF; low geometry α -counting and Ar-CH ₃ counter for SF
1956Bu92	Butler et al.	6.64 (10)	6.65 (10)	α /SF; ionization chamber
1961Dr04	Druin et al.	6.6 (7)		Gas scintillator; relative to α half-life of ²³⁸ Pu
1963Ma50	Malkin et al.	7.45 (17)		Gas scintillator; specific activity
1978MeZL	Meadows	6.80 (5)	6.74 (5)	α /SF; relative to half-life of ²³⁹ Pu
1980Kh05	Khan et al.	7.43		Mica fission track detector
1988SeZY	Selickij et al.	6.86 (26)		Fission fragment detection in 2 π geometry

^a Re-estimated in 2000Ho27.

Omitting the value of 1980Kh05 reported without uncertainty, the weighted average of the seven remaining values is 6.79 with the internal uncertainty 0.032 and external uncertainty 0.090 and $\chi^2/\nu = 2.94$.

The adopted value of the ²⁴²Pu spontaneous fission is 6.79 (10) 10¹⁰ years where the uncertainty is the smallest quoted experimental uncertainty.

2.1 α Transitions

The energies of the alpha transitions were obtained from the Q value and the level energies given in Table 3 from 2002Ch52.

Table 3. ²³⁸U levels populated in the ²⁴²Pu α -decay.

Level number	Energy, keV	Spin and parity	Half-life	Probability of α -transition ($\times 100$)
0	0,0	0 ⁺	4.468 (5)·10 ⁹ yr	76.53 (17)
1	44.915 (13)	2 ⁺	206 (3) ps	23.44 (17)
2	148.39 (3)	4 ⁺		0.030 4 (13)
3	307.19 (8)	6 ⁺		0.000 84 (6)

The probabilities of the transitions of $\alpha_{0,0}$, $\alpha_{0,1}$ and $\alpha_{0,2}$ have been obtained by averaging the direct alpha-emission measurement results (the most accurate of them are from 1986Va33) and the values deduced from the gamma-ray transition probability (P(γ +ce)) balances at the corresponding ²³⁸U levels. The deduced values are based on the measurements of absolute gamma-ray emission probabilities (P(γ)) from 1986Va33 (see Table 6) and adopted total internal conversion coefficients (ICCs).

Such averaging is possible as in 1986Va33 the independent measurements were carried out for alpha-emission intensities (with Si(Au) detector) and gamma-ray intensities (with two Ge detectors). The correlation between these measurements can be only due to the same sources used but it is negligible taking into account a large difference between the methods and detectors. Determination of the ²⁴²Pu disintegration rates for six sources required for the absolute gamma intensity measurements was made in 1986Va33 using absolute alpha particle counting under well-defined low solid angles, i.e. out of connection with the alpha - emission intensity measurements with Si(Au) detector.

The probability of the $\alpha_{0,3}$ -transition has been deduced from the P(γ +ce) balance at the ²³⁸U level of 307.19 keV (Table 4).

Table 4. Experimental, deduced and recommended values of α -transition probabilities ($\times 100$) in ^{242}Pu decay.

	α -particle energy (keV)	1953Asaro	1956Hu96	1976Barano v	1986Va33	Deduced from P(γ) measured in 1986Va33	Recommended
$\alpha_{0,0}$	4902	80 (6) ^a	74 (4) ^a	79.7 (20) ^b	76.45 (17)	77.3 (6)	76.53 (17) ^c
$\alpha_{0,1}$	4858	20 (6) ^a	26 (4) ^a	20.2 (20) ^b	23.52 (17)	22.7 (6)	23.44 (17) ^c
$\alpha_{0,2}$	4756	-	-	-	0.029 0 (14)	0.031 7 (13)	0.030 4 (13)
$\alpha_{0,3}$	4600	-	-	-	-	0.000 84 (6)	0.000 84 (6)

^a No uncertainties were quoted by the authors. The uncertainties adopted here were estimated by R. Vaninbroukx from the spectra shown in the papers (1986LoZT).

^b The uncertainties of 2.7 for 79.7 and 1.1 for 20.2 quoted by the authors were re-estimated by R. Vaninbroukx (1986LoZT).

^c Weighted average of the five values including direct measurement results and deduced value, uncertainty is the smallest quoted one.

^d Weighted average of the two values including direct $\alpha_{0,2}$ -transition measurement result and deduced value, uncertainty is the smallest quoted one.

2.2 γ Transitions

The recommended energies of gamma-ray transitions are virtually the same as the gamma-ray energies because nuclear recoil is negligible for ^{234}U .

The gamma-ray transition probabilities have been deduced from the gamma-ray emission probabilities and total internal conversion coefficients (ICCs). The ICCs have been interpolated using the BrIcc package with the so called “*Frozen Orbital*” approximation (2008Ki07). The uncertainties in the ICCs for pure multipolarities have been taken as 2 %. The multipolarities have been taken from 2002Ch52.

3 Atomic Data

3.1. Fluorescence yields

The fluorescence yield data are from 1996Sc06 (Schönfeld and Janßen).

3.2 X-rays and Auger electrons

The energies of U LX-rays taken from the SAISINUC software supporting programs agree with the measurements of 1994Le37 where the fine structure of LX radiation was measured in decay of ^{240}Pu .

The U KX-ray energies have been taken from 1999Schönfeld where the calculated values based on X-ray wavelengths from 1967Be65.

The relative KX-ray emission probabilities have been taken from 1999Schönfeld.

The energies of Auger electrons are from the SAISINUC software supporting programs. The ratios P(KLX)/P(KLL), P(KXY)/P(KLL) are taken from 1996Sc06.

4 Alpha Emissions

The α -emission energies have been obtained from Q value and ^{238}U level energies taking into account the ^{238}U recoil energies. In Table 5 the recommended values of α -emission energies are compared with the experimental results from alpha-spectrometric measurements and also with the evaluated data by A. Rytz (1991Ry01).

Table 5. Experimental and recommended α -emission energies in decay of ²⁴²Pu (keV).

	Measured ^a				1991Ry01	Recommended
	1953Asaro	1956Hu96	1956Ko67	1968Ba25		
$\alpha_{0,0}$	4904.6 (20)	4903.7 (30)	4907.2 (30)	4900.4 (12)	4902.3 (14)	4902.3 (10)
$\alpha_{0,1}$	4860.6 (20)	4859.7 (30)	4863.2 (30)	4856.1 (12)	4858.1 (15)	4858.2 (10)
$\alpha_{0,2}$	-	-	-	-	-	4756.2 (10)
$\alpha_{0,3}$	-	-	-	-	-	4600.1 (10)

^a Original values have been adjusted taking into account changes in calibration energies as suggested in 1991Ry01.

5 Electron Emissions

The energies of conversion electrons have been obtained from the gamma-ray transition energies and the atomic-electron binding energies. The emission probabilities of the conversion electrons have been deduced from the evaluated $P(\gamma)$ and ICC values.

The absolute emission probabilities of K and L Auger electrons have been calculated using the EMISSION computer program (2000Schönfeld).

6 Photon emissions

6.1 X-ray Emissions

The absolute emission probability of U MX-rays ($\alpha\beta$) in decay of ²⁴²Pu has been deduced from the relative intensity $P(XM\alpha\beta)/P(XL\eta\beta) = 0.41$ (4) measured in 1990Po14.

The absolute emission probabilities of U KX- and U LX-rays in decay of ²⁴²Pu have been calculated using the EMISSION computer program (2000Schönfeld).

6.2 Gamma-ray Emissions

The energies of gamma-rays have been adopted from 1972Sc01.

The absolute emission probabilities of the gamma-rays $\gamma_{1,0}$ (44.915 keV) and $\gamma_{2,1}$ (103.50 keV) have been deduced from the recommended $P(\alpha)$ values (Table 4) and the adopted total ICCs on the basis of intensity balances at the corresponding ²³⁸U levels. The absolute emission probability of the gamma-ray $\gamma_{3,2}$ (158.80 keV) has been adopted from the direct measurement of 1986Va33 (Table 6).

Table 6. Experimental and recommended absolute emission probabilities of gamma-rays ($\times 100$) in ²⁴²Pu decay.

	Energy (keV)	1972Sc01	1986Va33	Recommended
$\gamma_{1,0}$	44.915	-	0.037 2 (7)	0.038 4 (8)
$\gamma_{2,1}$	103.50	0.008 1 (9) ^a	0.002 63 (9)	0.002 53 (12)
$\gamma_{3,2}$	158.80	0.005 (2) ^a	0.000 298 (20)	0.000 298 (20)

^a Not used in the evaluation as considered in 1986LoZT.

7 Consistency of recommended data

The most accurate Q value, Q(M), is taken from the atomic mass adjustment table of Audi et al. (2003Au03). Comparison of Q(eff) (deduced as the sum of average energies per disintegration ($\sum E_i \times P_i$) for all emissions accompanying ²⁴²Pu α -decay) with the tabulated decay energy Q(M) allows to check a consistency of the recommended decay-scheme parameters obtained in this evaluation.

Here E_i and P_i are the evaluated energies and emission probabilities of the i -th alpha-particle, beta particle, gamma-ray, X-ray, etc. Consistency (percentage deviation) is determined by $\{[Q(M) - Q(\text{eff})] / Q(M)\} \times 100$. "Percentage deviations above 5 % would be regarded as high and imply a poorly defined decay scheme; a value of less than 5 % indicates the construction of a reasonably consistent decay scheme" (quoted from the article by A.L. Nichols in *Appl. Rad. Isotopes* 55 (2001) 23-70).

For the above ²⁴²Pu decay data evaluation we have $Q(M) = 4984.5$ (10) keV and $Q(\text{eff}) = 4984$ (13) keV. Thereafter, the percentage deviation is (0.01 ± 0.26) %, i.e. consistency is superior.

8 References

- 1953Asaro F. Asaro, Thesis, Univ. of California, Livermore, CA, Rep. UCRL-2180(1953) (Alpha-particle energies and emission probabilities).
- 1956Bu64 J. P. Butler, M. Lounsbury, J. Merritt, *Can. J. Chem.* 34(1956)253 (Half-life).
- 1956Bu92 J. P. Butler, T. A. Eastwood, T. L. Collins, M. E. Jones, F. M. Rourke, R. P. Schuman, *Phys. Rev.* 103(1956)634 (Half-life, SF half-life).
- 1956Hu96 J. P. Hummel, Thesis, Univ. California, Livermore, CA, Rep. UCRL-3456(1956) (Alpha-particle energies and emission probabilities).
- 1956Ko67 L. M. Kondratev, G. I. Novikova, Y. P. Sobolev, L. L. Goldin, *Zh. Eksp. Teor. Fiz.* 31(1956)771 - *Soviet Phys. JETP* 4(1956)645 (Alpha-particle energies and emission probabilities).
- 1956Me37 J. F. Mech, H. Diamond, M. H. Studier, P. R. Fields, A. Hirsch, C. M. Stephens, R. F. Barnes, D. J. Henderson, J. R. Huizenga, *Phys. Rev.* 103(1956)340 (Half-life, SF half-life).
- 1956Studier M. H. Studier, A. Hirsch, Private Communication. Quoted in 1956Me37 (SF half-life).
- 1961Dr04 V. A. Druin, V. P. Perelygin and G. I. Khlebnikov, *Soviet Phys. JETP* 13(1961)913 - *Zhurn. Eksptl. i Teoret. Fiz.* 40(1961)1296 (SF half-life).
- 1963Ma50 L. Z. Malkin, I. D. Alkhozov, A. S. Krivokhatsky, K. A. Petrzhak, *At. Energ. USSR* 15(1963)158 - *Soviet. J. At. Energy* 15(1964)851 (SF half-life).
- 1967Be65 J. A. Bearden, *Rev. Mod. Phys.* 39(1967)78 (X-ray energies).
- 1968Ba25 S. A. Baranov, V. M. Kulakov, V. M. Shatinskii, *Nucl. Phys.* 7(1968)442 - *Yadern. Fiz.* 7(1968)727 (Alpha-particle energies).
- 1969Be06 C. E. Bemis Jr., J. Halperin, R. Eby, J. Inorg. Nucl. Chem. 31(1969)599 (Half-life).
- 1970Du02 R. W. Durham, F. Molson, *Can. J. Phys.* 48(1970)716 (Half-life).
- 1972Sc01 M. Schmorak, C. E. Bemis Jr, M. J. Zender, N. B. Gove, P. F. Dittner, *Nucl. Phys.* A178(1972)410 (Gamma-ray energies and emission probabilities).
- 1976Baranov S. A. Baranov, A. G. Zelenkov; V. M. Kulakov, *Sov. At. Energy* 41(1976)987 (Alpha-emission probabilities).
- 1976Bu23 L. S. Bulyanitsa, A. M. Geidelman, Y. S. Egorov, L. M. Krizhanskii, A. A. Lipovskii, L. D. Preobrazhenskaya, A. V. Lovtsyus, Y. V. Kholnov, *Bull. Akad. Sci. USSR, Phys. Ser.* 40(10)(1976)42 - *Izv Akad Nauk SSSR, Ser. Fiz.* 40(1976)2075 (Half-life).
- 1976Os05 D. W. Osborne, H. E. Flotow, *Phys. Rev. C* 14(1976)1174 (Half-life).
- 1978MeZL J. W. Meadows, BNL-NCS-24273(1978)10 (A830926) (Half-life, SF half-life).
- 1979Ag03 S. K. Aggarwal, S. N. Acharya, A. R. Parab, H. C. Jain, *Phys. Rev. C* 20(1979)1135 (Half-Life).
- 1980Kh05 N. A. Khan, H. A. Khan, K. Gul, M. Anwar, G. Hussain, R. A. Akbar, A. Waheed, M. S. Shaikh, *Nucl. Instrum. Methods* 173(1980)163 (SF half-life).
- 1986LoZT A. Lorenz, IAEA Tech. Rep. Ser., No 261(1986) (A871001 M881119 Part 2) (Evaluated decay data).
- 1986Va33 R. Vaninbrouckx, G. Bortels, B. Denecke, *Int. J. Appl. Radiat. Isotop.* 37(1986)1167 (Alpha-, gamma-ray emission probabilities).
- 1988SeZY Yu. A. Selitsky, V. B. Funshtein, V. A. Yakovlev, *Proc. 38th Ann. Conf. Nucl. Spectrosc. Struct. At. Nuclei, Baku, Acad. Sci. USSR* (1988)131 (SF half-life).
- 1990Po14 Yu. S. Popov, I. B. Makarov, D. Kh. Srurov, E. A. Erin, *Radiokhimiya.* 32(1990)2 - *Sov. J. Radiochemistry* 32(1990)425 (MX-, LX-ray relative emission probabilities).
- 1991Ry01 A. Rytz, *At. Data Nucl. Data Tables.* 47(1991)205 (Alpha-emission energies).
- 1994Le37 M.C. Lépy, B. Duchemin, J. Morel, *Nucl. Instrum. Meth. Phys. Res.* A353(1994)10 (LX-ray energies and emission probabilities).
- 1996Sc06 E. Schönfeld, H. Janßen, *Nucl. Instrum. Meth. Phys. Res.* A369(1996)527 (Atomic data).

- 1999Schönfeld E. Schönfeld, G. Rodloff, PTB-6.11-1999-1999-1, Braunschweig, Februar 1999 (KX-ray energies and relative emission probabilities).
- 2000Ho27 N. E. Holden, D. C. Hoffman, Pure Appl. Chem. 72(2000)1525 (SF half-life).
- 2000Schönfeld E. Schönfeld, H. Janßen, Appl. Rad. Isotop. 52(2000)595 (X-ray and Auger electron emission probabilities, EMISSION code).
- 2002Ch52 F. E. Chukreev, V. E. Makarenko, M. J. Martin, Nucl. Data Sheets 97(2002)129 (Decay Scheme, ²³⁸U level energies, gamma-ray multiplicities).
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729(2003)337 (Q value).
- 2004BeZQ M.M. Bé, V. Chisté, C. Dulieu, E. Browne, V. Chechev, N. Kuzmenko, R. Helmer, A. Nichols, E. Schönfeld, and R. Dersch, Table of Radionuclides (Vol.2 - A = 151 to 242), Monographie BIPM-5, Vol. 2, p. 277 – 281. Bureau International des Poids et Mesures (2004) (²⁴²Pu Decay Data Evaluation).
- 2005ChZU V. P. Chechev, Proc. Intern. Conf. Nuclear Data for Science and Technology, Santa Fé, New Mexico, 26 September-1 October, 2004, R. C. Haight, M. B. Chadwick, T. Kawano, P. Talou, Eds., Vol. 1, p. 91 (2005); AIP Conf. Proc. 769 (2005) (²⁴²Pu Decay Data Evaluation).
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor Jr., Nucl. Instrum. Meth. Phys. Res. A589(2008)202 (Theoretical ICC).

²⁴²Am - Comments on evaluation of decay data

by A. L. Nichols

Evaluated: March 2007/September 2008**Evaluation Procedure**

Limitation of Relative Statistical Weight Method (LWM) was applied to average the decay data when appropriate.

Decay Scheme

A relatively simple decay scheme was constructed from the β^- /EC ratio and branching fraction measurements of Hoff *et al.* (1955Ho67, 1959Ho02), Baranov and Shlyagin (1955Ba31), Asaro *et al.* (1960As05), Gasteiger *et al.* (1969Ga17), Aleksandrov *et al.* (1969Al20) and Gabeskiriya (1972Ga35). There are no known well-defined gamma-ray spectroscopic studies.

Some confusion arose during the course of the 1950s as to the correct identity of the ground and metastable states of ²⁴²Am. This problem was resolved in 1960 by Asaro *et al.* (1960As05) when the 16-hour half-life activity was shown to be the ground state. The possible existence of an alpha branch has been extensively considered by Barnes *et al.* (1959Ba22) and Aleksandrov *et al.* (1969Al20). While Barnes *et al.* found such a branching fraction ($BF_\alpha = 0.004\ 76\ (14)$), subsequent studies have shown no evidence for this particular decay mode, and Aleksandrov *et al.* were only able to set a limit of less than 10^{-7} of the total ²⁴²Am decay.

Nuclear Data

²⁴²Am needs to be better characterized for improved quantification of the production and decay heat contribution of ²⁴²Cm.

Half-life

The recommended half-life of 16.01 (2) hours has been adopted from three known sets of measurements (1953Ke38, 1969Al20, 1982Wi05). Five independent half-life measurements were individually reported by Aleksandrov *et al.* (1969Al20) from which a value of 16.07 (14) h was calculated (LWM). A limited data set of effectively three studies is rather unsatisfactory, and further measurements are required to determine the half-life with much greater confidence.

Half-life measurements

Reference	Half-life (hours)
1953Ke38	16.01 ± 0.02
1969Al20	16.07 ± 0.14
1982Wi05	16.1 ± 0.1
Recommended value	16.01 ± 0.02

Gamma Rays

Energies

All gamma-ray transition energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies of Akovali were adopted (2002Ak06), and used to determine the energies and associated uncertainties of the gamma-ray transitions that depopulate the first excited states of ²⁴²Pu and ²⁴²Cm.

Emission Probabilities

There are no known dedicated measurements of the gamma-ray emission probabilities. Under these unsatisfactory circumstances, the proposed gamma-ray decay data were derived from the tabulated P_{ce}/P_{β^-} data of Baranov and Shlyagin (1955Ba31) and the BF_{β} measurements (1959Ba22, 1959Ho02, 1969Al20, 1969Ga17, 1972Ga35). A BF_{β} of 0.831 (3) was derived in terms of LWM, with the uncertainty extended to the minimum value measured (± 0.003); this parameter was adopted in preference to the equivalent LWM calculation for the β^-/EC ratio (i.e. 4.88 (8) compared with a value of 4.92 (9) calculated from the weighted mean BF_{β}).

β^-/EC ratio and BF_{β} (Branching fraction).

Reference	BF_{β}	β^-/EC
1955Ba31	0.82	4.6
1955Ho67	0.81	4.2
1959Ba22	$0.836 \pm 0.008^*$	5.1 ± 0.2
1959Ho02	0.836 ± 0.003	$5.1 \pm 0.1^*$
1960As05	0.836^*	5.1
1969Al20	$0.82 \pm 0.01^*$	4.6 ± 0.3
1969Ga17	0.828 ± 0.004	$4.8 \pm 0.1^*$
1972Ga35	$0.827 \pm 0.003^*$	4.78 ± 0.08
Recommended value	0.831 ± 0.003	[4.88 \pm 0.08]

* Emphasis of the publication, and assumed to be the primary measurement.

Baranov and Shlyagin determined the conversion-electron emission intensities separately for both the electron-capture and beta decay processes, along with the β^- decay in equivalent units (1955Ba31) to furnish the following ratios:

$$P_{ce}(EC \text{ component})/P_{\beta^-} = 153.5/1200, \text{ and}$$

$$P_{ce}(\beta^- \text{ component})/P_{\beta^-} = 661/1200.$$

One problem involves the assignment of uncertainties to the P_{ce}/P_{β^-} values as determined by Baranov and Shlyagin. Both parameters are the ratios of two equivalent measurements, and the resulting uncertainty for each of these ratios was assumed to be approximately 5 %:

$$P_{ce}(EC \text{ component})/P_{\beta^-} = 153.5/1200 = 0.128 (6)$$

$$P_{ce}(\beta^- \text{ component})/P_{\beta^-} = 661/1200 = 0.551 (28).$$

Using these data and BF_{β} of 0.831 (3):

$$P_{ce}(\beta^-) = 0.551 (28) \times 0.831 (3) = 0.458 (23) \text{ for the 42.13-keV gamma ray,}$$

$$\text{and } P_{ce}(EC) = 0.128 (6) \times 0.831 (3) = 0.106 (5) \text{ for the 44.54-keV gamma ray.}$$

These values were then used in conjunction with the theoretical internal conversion coefficients to calculate the absolute gamma-ray emission probabilities.

Quite remarkably, the resulting gamma-ray emission probabilities are in good agreement with the tabulated spectroscopic data of Vylov *et al.* (1980VyZZ) which are listed as 42.129 (7) keV

and 0.039 (5) %, and 44.542 (25) keV and 0.015 (3) %. Accurate, high-resolution gamma-ray measurements are required to confirm the validity of the proposed decay scheme.

Gamma-ray emissions: recommended energies, emission probabilities, multiplicities and theoretical internal conversion coefficients (frozen orbital approximation).

	E_γ (keV)	P_γ^{abs}	Multi	α_K	α_L	α_{M+}	α_{tot}
$\gamma_{1,0}$ (Cm)	42.13 (5)	0.040 ± 0.002	E2	-	836 (12)	319 (5)	1155 (17)
$\gamma_{1,0}$ (Pu)	44.54 (2)	0.014 ± 0.001	E2	-	544 (8)	204 (3)	748 (11)

Multipolarities and Internal Conversion Coefficients

The nuclear level scheme specified by Akovali has been used to define the multiplicities of the gamma transitions on the basis of known spins and parities (2002Ak06). Recommended internal conversion coefficients have been determined from the theoretical tabulations of Band *et al.* (2002Ba25, 2002Ra45) by means of the methodology of Kibedi *et al.* (2008Ki07).

Beta-particle Emissions

Energies and emission probabilities

Beta-particle energies were calculated from the nuclear level energies of Akovali (2002Ak06) and a Q_{β^-} value of 664.5 ± 0.4 keV taken from Audi *et al.* (2003Au03).

Assuming virtually full internal conversion of the 42.13-keV gamma transition, the beta-particle emission probabilities were calculated from BF_β of 0.831 (3) and $P_{ce}(\beta^-)$ of 0.458 (23):

Beta-particle Emission Probabilities per 100 Disintegrations of ²⁴²Am.

	E_β (keV)	av. E_β (keV)	P_β	Transition type	log <i>ft</i>
$\beta_{0,1}^-$	622.4 ± 0.4	185.92 ± 0.14	45.8 ± 2.3	1 st forbidden non-unique	6.84
$\beta_{0,0}^-$	664.5 ± 0.4	200.17 ± 0.14	37.3 ± 2.3	1 st forbidden non-unique	7.03

EC Transitions

Energies and transition probabilities

EC transition energies were calculated from the nuclear level energies of Akovali (2002Ak06) and a Q_{EC} value of 751.3 ± 0.7 keV from Audi *et al.* (2003Au03).

Assuming virtually full internal conversion of the 44.54-keV gamma transition, the EC transition probabilities were calculated from BF_{EC} of 0.169 (3) and $P_{ce}(EC)$ of 0.106 (5):

EC Transition Probabilities per 100 Disintegrations of ²⁴²Am.

	E_{EC} (keV)	P_{EC}	Transition type	log <i>ft</i>	P_K	P_L	P_M
$EC_{0,1}$	706.8 ± 0.7	10.6 ± 0.5	1 st forbidden non-unique	7.26	0.7261 (23)	0.2016 (15)	0.0532 (10)
$EC_{0,0}$	751.3 ± 0.7	6.3 ± 0.6	1 st forbidden non-unique	7.55	0.7303 (22)	0.1987 (15)	0.0522 (10)

Atomic Data

The x-ray and Auger-electron data have been calculated using the evaluated gamma-ray data, and atomic data from 1996Sc06, 1998ScZM and 1999ScZX. Both the x-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.01, 28 January 2003, with the emission.101 database extended to $Z = 96$ to calculate component L x-ray data of daughter Cm). This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

K and L X-ray Emission Probabilities per 100 Disintegrations of ²⁴²Am.

			Energy keV	Photons per 100 disint.
XL		(Pu)	12.124 – 22.153	10.8 (5)
	XL ₁	(Pu)	12.124	0.293 (11)
	XL _α	(Pu)	14.087 – 14.282	4.56 (16)
	XL _η	(Pu)	16.333	0.084 (4)
	XL _β	(Pu)	16.498 – 18.541	4.64 (15)
	XL _γ	(Pu)	21.420 – 22.153	1.03 (4)
XK _α	XK _{α2}	(Pu)	99.525	3.55 (17)
	XK _{α1}	(Pu)	103.734	5.6 (3)
XK _{β1}	XK _{β3}	(Pu)	116.244)
	XK _{β1'}	(Pu)	117.228) 2.06 (11)
	XK _{β5}	(Pu)	117.918)
XK _{β2}	XK _{β2}	(Pu)	120.540)
	XK _{β4}	(Pu)	120.969) 0.72 (4)
	XKO _{2,3}	(Pu)	121.543)
XL		(Cm)	12.633 – 23.527	18.0 (11)
	XL ₁	(Cm)	12.633	0.451 (22)
	XL _α	(Cm)	14.746 – 14.961	6.8 (3)
	XL _η	(Cm)	17.314	0.194 (11)
	XL _β	(Cm)	17.286 – 19.688	8.7 (4)
	XL _γ	(Cm)	22.735 – 23.527	2.09 (10)

Electron energies were determined from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

References

- 1953Ke38 T.K. KEENAN, R.A. PENNEMAN, B.B. McINTEER, A new determination of the half-life of Am^{242m}: the problem of counting short-lived activities, J. Chem. Phys. 21 (1953) 1802-1803. [half-life]
- 1955Ba31 S.A. BARANOV, K.N. SHLYAGIN, Energy levels of the U²³⁷ nucleus and the decay of Am^{242m}, Peaceful Use of Atomic Energy, Conf. Acad. Sci. USSR, Moscow, 1955; Consultants Bureau translation (1956) 183-194. [E_γ, P_{cc}/P_β, BF_{EC}, BF_β, β⁻/EC ratio]
- 1955Ho67 R.W. HOFF, H. JAFFE, T.O. PASSELL, F.S. STEPHENS, E.K. HULET, S.G. THOMPSON, Radioactive decay of the isomers of americium-242, Phys. Rev. 100 (1955) 1403-1406. [β⁻/EC ratio]

- 1959Ba22 R.F. BARNES, D.J. HENDERSON, A.L. HARKNESS, H. DIAMOND, The alpha and electron capture partial half-lives of ²⁴²Am, *J. Inorg. Nucl. Chem.* 9 (1959) 105-107. [BF_{EC}]
- 1959Ho02 R.W. HOFF, E.K. HULET, M.C. MICHEL, Branching ratio of ^{242m}Am decay, *J. Nucl. Energy* 8 (1959) 224-228. [β⁻/EC ratio]
- 1960As05 F. ASARO, I. PERLMAN, J.O. RASMUSSEN, S.G. THOMPSON, Isomers of Am²⁴⁴, *Phys. Rev.* 120 (1960) 934-943. [β⁻/EC ratio]
- 1961Ma27 R. MARRUS, J. WINOCUR, Hyperfine structure and nuclear moments of americium-242, *Phys. Rev.* 124 (1961) 1904-1906. [spin state]
- 1969Al20 B.M. ALEKSANDROV, M.A. BAK, V.V. BERDIKOV, R.B. IVANOV, A.S. KRIVOKHATSKII, V.G. NEDOVESOV, K.A. PETRZHAK, Yu.G. PETROV, Yu.F. ROMANOV, É.A. SHLYAMIN, The decay of Am²⁴², *Sov. At. Energy* 27 (1969) 724-728. [half-life, β⁻/EC ratio, α decay]
- 1969Ga17 R. GASTEIGER, G. HÖHLEIN, W. WEINLÄNDER, Bestimmung des Zerfallsverhältnisses β⁻/EC des ²⁴²Am, *Radiochim. Acta* 11 (1969) 158-161. [β⁻/EC ratio]
- 1972Ga35 V.Ya. GABESKIRIYA, Relative probability of Am²⁴² beta decay, *Sov. At. Energy* 32 (1972) 201-202. [β⁻/EC ratio]
- 1977La19 F.P. LARKINS, Semiempirical Auger-electron energies for elements 10 ≤ Z ≤ 100, *At. Data Nucl. Data Tables* 20 (1977) 311-387. [Auger-electron energies]
- 1980VyZZ Ts. VYLOV, G.-J. BEYER, V.M. GOROZHANKIN, Zh. ZHELEV, A.I. IVANOV, R.B. IVANOV, V.G. KALINNIKOV, M.Ya. KUZNETSOVA, N.A. LEBEDEV, M.A. MIKHAILOVA, A.I. MUMINOV, A.F. NOVGORODOV, Yu.V. NORSEEV, Sh. OMANOV, B.P. OSIPENKO, E.K. STEPANOV, K. THIEME, V.G. CHUMIN, A.F. SHCHUS, Yu.V. YUSHKEVICH, *Spektren der Strahlung Radioaktiver Nuklide, Gemessen mit Halbleiterdetektoren, Zentralinstitut für Kernforschung, Rossendorf bei Dresden, und Vereinigtes Institut für Kernforschung Dubna, ZfK-399* (1980); Ts. VYLOV, V.M. GOROZHANKIN, Zh. ZHELEV, A.I. IVANOV, R.B. IVANOV, V.G. KALINNIKOV, M.Ya. KUZNETSOVA, N.A. LEBEDEV, M.A. MIKHAILOVA, A.I. MUMINOV, A.F. NOVGORODOV, Yu.V. NORSEEV, Sh. OMANOV, B.P. OSIPENKO, E.K. STEPANOV, V.G. CHUMIN, A.F. SHCHUS, Yu.V. YUSHKEVICH, *Spectra of Radiations of Radioactive Nuclides*, Editor: K.Ya. GROMOV, FAN Publishing, Tashkent, USSR (1980). [E_X, P_X, E_γ, P_γ]
- 1982Wi05 K. WISSHAK, J. WICKENHAUSER, F. KÄPPELER, G. REFFO, F. FABBI, The isomeric ratio in thermal and fast neutron capture of americium-241, *Nucl. Sci. Eng.* 81 (1982) 396-417. [half-life]
- 1996Sc06 E. SCHÖNFELD, H. JANßEN, Evaluation of atomic shell data, *Nucl. Instrum. Meth. Phys. Res.* A369 (1996) 527-533. [X_K, X_L, Auger electrons]
- 1998ScZM E. SCHÖNFELD, G. RODLOFF, Tables of the energies of K-Auger electrons for elements with atomic numbers in the range from Z = 11 to Z = 100, *PTB Report PTB-6.11-98-1*, October 1998. [Auger electrons]
- 1999ScZX E. SCHÖNFELD, G. RODLOFF, Energies and relative emission probabilities of K X-rays for elements with atomic numbers in the range from Z = 5 to Z = 100, *PTB Report PTB-6.11-1999-1*, February 1999. [X_K]

- 2002Ak06 Y.A. AKOVALI, Nuclear data sheets for A = 242, Nucl. Data Sheets 96 (2002) 177-239. [nuclear levels]
- 2002Ba85 I.M. BAND, M.B. TRZHASKOVSKAYA, C.W. NESTOR, Jr., P.O. TIKKANEN, S. RAMAN, Dirac–Fock internal conversion coefficients, At. Data Nucl. Data Tables 81 (2002) 1-334. [ICC]
- 2002Ra45 S. RAMAN, C.W. NESTOR, Jr., A. ICHIHARA, M.B. TRZHASKOVSKAYA, How good are the internal conversion coefficients now? Phys. Rev. C66 (2002) 044312, 1-23. [ICC]
- 2003Au03 G. AUDI, A.H. WAPSTRA, C. THIBAUT, The AME2003 atomic mass evaluation (II). Tables, graphs and references, Nucl. Phys. A729 (2003) 337-676. [Q-value]
- 2008Ki07 T. KIBÉDI, T.W. BURROWS, M.B. TRZHASKOVSKAYA, P.M. DAVIDSON, C.W. NESTOR, Jr., Evaluation of theoretical conversion coefficients using BrIcc, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202-229. [ICC]

^{242m}Am - Comments on evaluation of decay data

by A. L. Nichols

Evaluated: April 2007/April 2010

Evaluation Procedure

Limitation of Relative Statistical Weight Method (LWM) was applied to average the decay data when appropriate.

Decay Scheme

A simple IT-decay mode dominates the decay scheme of ^{242m}Am. The small α branch is complex, and many features of this decay mode remain unresolved despite the extensive study of Hoff *et al.* (1990Ho02).

Some confusion arose during the course of the 1950s as to the correct identity of the ground and metastable states of ²⁴²Am. This problem was resolved in 1960 by Asaro *et al.* (1960As05) when the 16-hour half-life activity was shown to be the ground state and the longer-lived 140-year isomer was defined as the metastable state. The α branch has been determined by Barnes *et al.* (1959Ba22) and Zelenkov *et al.* (1979Ze05) to be 0.46 (1) %. Hoff *et al.* have studied the emissions from the α -decay mode in considerable detail (1990Ho02), and the more modest measurements of Baranov *et al.* (1979Ba67) and Vylov *et al.* (1980VyZZ) show reasonable agreement with this extensive data set. A small spontaneous fission branch of $1.5 (6) \times 10^{-8}$ % has been quantified by Caldwell *et al.* (1967Ca04), while an upper limit of 4.8×10^{-9} % has been specified by Zelenko *et al.* (1986Ze06).

Nuclear Data

The decay characteristics of ^{242m}Am need to be better defined for improved quantification of the production and decay heat contributions of ²⁴²Cm and ²⁴⁴Cm.

Half-life

A recommended half-life of 143 (2) years has been adopted from the two known measurements (1959Ba22, 1979Ze05). This limited data set is unsatisfactory, and further studies are required to determine the half-life with much greater confidence.

Half-life measurements

Reference	Half-life (years)
1959Ba22	152 \pm 7
1979Ze05	141.9 \pm 1.7
Recommended value	143 \pm 2

Branching Fractions

Barnes *et al.* and Zelenkov *et al.* have determined the α branching fraction for ^{242m}Am (1959Ba22, 1979Ze05), and these data were used to derive an α branch of 0.46 (1) % and IT branch of 99.54 (1) %.

Reference	BF _{α}
1959Ba22	0.004 76 \pm 0.000 14
1979Ze05	0.004 5 \pm 0.000 1
Recommended value	0.004 6 \pm 0.000 1
α branch	(0.46 \pm 0.01) %

A spontaneous fission branch of $1.5 (6) \times 10^{-8} \%$ can be determined from the recommended total half-life of 143 (2) years and measured spontaneous fission half-life of $9.5 (35) \times 10^{11}$ years (1967Ca04). Similarly, an upper limit of $4.8 \times 10^{-9} \%$ for the spontaneous fission branch can be derived from equivalent studies of the spontaneous fission half-life of $> 3 \times 10^{12}$ years (1986Ze06). Under these uncertain circumstances, a recommended value of $< 4.8 \times 10^{-9} \%$ has been adopted for the spontaneous fission branch of ^{242m}Am.

Q values

Q_{IT} of 48.60 (5) keV and Q_α of 5637.10 (25) keV were adopted from the evaluated tabulations of Audi *et al.* (2003Au03).

Alpha Particles

Alpha-particle measurements reveal a relatively complex α-decay mode (1979Ba67, 1980VyZZ, 1990Ho02). The Q_α of 5637.10 (25) keV (2003Au03) and nuclear level energies as defined by Chukreev *et al.* (2002Ch52) were used to calculate the alpha-particle energies, while the alpha-particle emission probabilities were primarily adopted from the measurements of Hoff *et al.* (1990Ho02) and fortified by the introduction of a number of minor transitions observed by Baranov *et al.* (1979Ba67), all expressed in terms of decay per 100 alphas. Small adjustments were made to some of the low-intensity alpha-particle emission probabilities after consideration of the observed differences between the two sets of measurements (i.e., 5091.9-, 5248.15/5248.21- and 5272.96-keV alpha-particle emission probabilities). Some of the proposed daughter nuclear levels of comparable energy were also judged to be populated by alpha-particle transitions that were not experimentally resolved (i.e., alpha-particle transitions to nuclear levels with energies of 297.03/299.23, 300.68/300.743 and 374.7/376.7 keV). Under these circumstances, the observed alpha-particle emission was arbitrarily shared between the two nuclear levels of relevance. An unweighted mean value of 1.508 (5) was adopted for the radius parameter r₀(²³⁸Np) as derived from the equivalent data for neighboring doubly-even nuclei (1998Ak04), and used in the calculation of α-hindrance factors (HF):

$$r_0(^{238}\text{Np}) = [r_0(^{240}\text{Pu}) + r_0(^{242}\text{Pu}) + r_0(^{242}\text{Cm}) + r_0(^{244}\text{Cm})] / 4$$

$$= [1.5168 (3) + 1.5143 (9) + 1.5013 (10) + 1.4979 (7)] / 4 = 1.508 (5)$$

All of the available alpha-particle decay data were assessed in conjunction with the gamma-ray measurements of Hoff *et al.* These extremely significant gamma-ray studies do not furnish gamma-ray energy and emission probability data that can be adopted to depopulate the major alpha-populating nuclear levels of ²³⁸Np in a consistent and satisfactory manner (i.e., nuclear levels at 407.59 keV and 342.439 keV populated by alpha particles with emission probabilities of 5.6 (2) % and 89.0 (7) % per 100 alphas, respectively). These two seriously incomplete features within the decay scheme are also observed to impact in various ways throughout the gamma-ray decay to the ground state of ²³⁸Np. While the recommended alpha-particle emissions are believed to be reasonably sound, the related gamma-ray data remain significantly incomplete.

Alpha-particle emissions: energies, emission probabilities and hindrance factors.

1979Ba67		1980VyZZ		1990Ho02		Recommended		
E _α (keV)	P _α (x100α)	E _α (keV)	P _α (x100α)	E _α (keV)	P _α (x100α)	E _α (keV)	P _α (x100α)	HF
4974.9	~ 0.002	-	-	-	-	4975 (3)	0.002 (1)	2400
5027.1	0.02	-	-	5031 (5)	0.02 (1)	5027.3 (15)	0.02 (1)	540
5064.2	0.22	5065 (5)	0.23	5072 (3)	0.25 (7)	5068 (3)	0.25 (7)	81
5082	0.03	5082 (5)) 0.34	-	-	5082.6 (12)	0.03 (1)	840
5088.4	0.19))	5093 (4)	0.21 (7)	5091.9 (7)	0.20 (7)	146
5141.6 (5)	5.82	5142.35 (104)	6.601 (163)	5144.4 (9)	5.6 (2)	5143.07 (26)	5.6 (2)	11.2

1979Ba67		1980VyZZ		1990Ho02		Recommended		
E_α (keV)	P_α (x100 α)	E_α (keV)	P_α (x100 α)	E_α (keV)	P_α (x100 α)	E_α (keV)	P_α (x100 α)	HF
5153.3	0.02	-	-	-	-	5153.2 (15)	0.02 (1)	3600
~ 5173) 0.04	-	-	-	-	5173.45 (26)	0.02 (1)	4900
5173.7)	-	-	-	-	5175.4 (10)	0.02 (1)	5000
5206.8 (5)	89.84	5205.92 (72)	100.00 (167)	5208.4 (8)	89.0 (7)	5207.15 (25)	89.0 (7)	1.80
5214.7 ?	0.03	-	-	-	-	5215.4 (7)	0.03 (1)	6000
5248.2	~ 0.11	5248 (5)	0.67	5248.4 (22)	1.0 (1)) 5248.15 (25)	0.4 (1)	730
) 5248.21 (26)	0.4 (1)	730
5250.0	0.04	-	-	-	-) 5249.64 (26)	0.02 (1)	14800
) 5251.80 (25)	0.02 (1)	15300
~ 5273	0.86	5284	~ 0.34	5271 (3)	1.1 (1)	5272.96 (25)	1.0 (1)	414
5313.5	0.69	5312 (5)	0.90	5316 (3)	0.6 (1)	5314.95 (25)	0.6 (1)	1250
-	-	-	-	5331 (5)	0.15 (10)	5331.97 (25)	0.15 (10)	6400
5367.2	1.17	5364 (5)	1.67	5369.1 (18)	1.1 (2)	5367.73 (25)	1.1 (2)	1430
5409.3	1.04	5408 (5)	1.35	5412.4 (21)	1.0 (2)	5410.13 (25)	1.0 (2)	2820
5458.2	0.14	-	-	-	-	5458.68 (25)	0.14 (4)	39000
5517.3	0.006	-	-	-	-	5517.93 (25)	0.003 (3)	4000000

Σ 100.025

Gamma Rays

Energies

All gamma-ray transition energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies of Akovali and Chukreev *et al.* were adopted (2002Ak06, 2002Ch52), and used to determine the energies and associated uncertainties of the gamma-ray transitions that populate and depopulate the excited nuclear levels of ^{238}Np and $^{242\text{m}}\text{Am}$.

Emission Probabilities

Dedicated measurements of the gamma-ray emission probabilities of $^{242\text{m}}\text{Am}$ are limited to the significant studies of Hoff *et al.* (1990Ho02). Under these rather unsatisfactory circumstances, these gamma-ray decay data were adopted wholesale, although some comparison could be made with the limited data set of Vylov *et al.* (1980VyZZ). Hoff *et al.* have directly identified over 60 gamma-rays with the α -decay mode which has a branch of only 0.46 (1) %, while the major IT decay mode involves only one highly-converted gamma transition (48.60 keV).

Hoff *et al.* report the emission probabilities of a number of unresolved gamma rays in terms of what is believed to be the upper limit for each: 4.0 (3) per 100 α for both 109.61 (1) and 109.618 (3) keV; 0.024 (7) per 100 α for both 139.05 (3) and 139.11 (2) keV; 0.150 (8) per 100 α for both 152.70 (2) and 152.73 (1) keV; 3.50 (10) per 100 α for both 163.1 (5) and 163.29 (1) keV; and 0.122 (10) per 100 α for both 250.33 (3) and 250.37 (2) keV. These data have been adopted and both entries carried forward as values less or equal (\leq) to the specified emission probability, as well as being incorporated into the database of recommended emission probabilities and transition probabilities.

The 26.427-keV gamma transition is particularly problematic, with a measured emission probability of 1.36 (1) per 100 α (1990Ho02). Combining this value with the internal conversion coefficients for an E2 transition generates an unrealistic absolute transition probability for this single gamma from the

26.427-keV nuclear level to the ground state of 2.12 %. There is a good possibility that this particular γ line arises partially from the alpha decay of ²⁴¹Am (isotopic content of 0.79 %, and γ -ray emission energy of 26.345 keV), and/or consists of a number of unresolved transitions that could not be identified nor located elsewhere within the incomplete decay scheme of ^{242m}Am. Therefore, the emission probability per 100 α of the 26.427-keV gamma transition has been significantly reduced to < 0.154, and is primarily based on α and γ transition probabilities per 100 α that are known to populate the 26.427-keV nuclear level.

Np K X-rays complicate the interpretation of the gamma-ray emission probability data over the energy ranges from 97 to 101 keV (K α) and 113.3 to 118.5 keV (K β). Hoff *et al.* observed gamma rays with energies in the vicinity of 97 keV and between 113.7 and 118 keV. While many of these particular emissions can be incorporated into the proposed decay scheme, their emission probabilities and existence in this form are doubtful.

A limited number of the gamma rays observed by Hoff *et al.* could not be placed in the proposed and incomplete alpha-decay scheme: 89.60 (5) keV with P_γ per 100 α of 0.29 (7), 160.61 (2) keV with P_γ per 100 α of 0.09 (4), 165.97 (15) keV with P_γ per 100 α of 0.010 (5), and 233.69 (10) keV with P_γ per 100 α of 0.028 (7).

Gamma-ray emissions: measured and recommended energies and emission probabilities.

1980VyZZ			1990Ho02				Adopted	
E_γ (keV)	P_γ^{rel}	P_γ per 100 α	E_γ (keV) α decay	P_γ per 100 α	E_γ (keV) (n, γ)	P_γ per 100n	E_γ (keV)	P_γ per 100 α
-	-	-	-	-	24.37 (2)	0.0064 (10)	24.34 (1)	-
-	-	-	26.32 (3)	1.36 (10)	26.43 (2)	0.104 (15)	26.427 (2)	< 0.154*
-	-	-	-	-	32.67 (3)	0.026 (4)	32.64 (1)	-
-	-	-	-	-	34.97 (3)	0.082 (11)	34.97 (1)	-
-	-	-	-	-	35.90 (2)	0.0109 (15)	35.90 (1)	-
-	-	-	-	-	43.11 (3)	0.044 (7)	43.11 (1)	-
-	-	-	-	-	43.32 (3)	0.0046 (7)	43.33 (1)	-
-	-	-	-	-	43.84 (3)	0.075 (11)	43.83 (2)	-
-	-	-	-	-	43.98 (4)	0.058 (8)	43.89 (2)	-
-	-	-	-	-	46.84 (3)	0.111 (17)	46.833 (3)	-
-	-	-	-	-	-	-	48.60 (5)	IT decay
49.367 (4)	100	[29.1]	49.35 (2)	29.1 (9)	49.372 (2)	8.57 (14)	49.371 (3)	29.1 (9)
-	-	-	-	-	52.98 (7)	0.029 (4)	53.2 (6)	-
-	-	-	53.69 (3)	0.45 (6)	53.70 (7)	0.038 (6)	53.67 (1)	0.45 (6)
-	-	-	-	-	53.88 (4)	0.017 (3)	53.85 (2)	-
-	-	-	57.54 (6)	0.21 (5)	-	-	57.51 (1)	0.21 (5)
-	-	-	-	-	59.31 (4)	0.0082 (13)	59.32 (1)	-
-	-	-	60.13 (6)	1.19 (11)	60.243 (4)	0.68 (5)	60.247 (3)	1.19 (11)
-	-	-	-	-	62.33 (3)	0.0100 (15)	62.330 (4)	-
66.808 (20)	11.2	3.26	66.89 (2)	3.25 (10)	66.919 (5)	0.23 (3)	66.92 (1)	3.25 (10)
67.9	3.9	1.1	67.93 (3)	0.87 (7)	-	-	67.92 (2)	0.87 (7)
73.3	3.2	0.93	73.66 (2)	1.71 (12)	73.715 (4)	0.49 (15)	73.72 (1)	1.71 (12)
-	-	-	-	-	75.97 (7)	0.0051 (8)	75.98 (1)	-
-	-	-	-	-	79.483 (17)	0.17 (3)	79.48 (1)	-
-	-	-	-	-	79.74 (3)	0.019 (3)	79.73 (2)	-
-	-	-	84.9 (2)	0.21 (7)	-	-	85.16 (7)	0.21 (7)
86.680 (36)	19.5	5.67	86.65 (2)	4.97 (15)	86.676 (2)	2.3 (4)	86.674 (2)	4.97 (15)
-	-	-	89.60 (5)	0.29 (7)	-	-	89.60 (5)	0.29 (7)#
92.5	2.2	0.64	92.52 (3)	0.61 (7)	92.486 (7)	0.19 (4)	92.48 (1)	0.61 (7)
-	-	-	93.82 (3)	0.79 (9)	93.67 (5)	0.19 (4)	93.88 (1)	0.79 (9)
-	-	-	-	-	95.22 (2)	0.046 (6)	95.22 (1)	-
-	-	-	95.7 (6)	-	-	-	96.204 (3)	-
-	-	-	-	-	96.82 (5)	0.025 (4)	96.78 (1)	-
97.077	K α_2 (Np)	-	-	-	-	-	-	-
-	-	-	98.0 (6)	-	97.22 (5)	0.046 (6)	97.18 (2)	-
101.068	K α_1 (Np)	-	-	-	-	-	-	-

Comments on evaluation

1980VyZZ			1990Ho02				Adopted	
E_γ (keV)	P_γ^{rel}	P_γ per 100 α	E_γ (keV) α decay	P_γ per 100 α	E_γ (keV) (n,γ)	P_γ per 100 n	E_γ (keV)	P_γ per 100 α
109.6	12.9	3.75	109.61 (2)	4.0 (3)	109.614 (4)	1.09 (24)	109.61 (1)	≤ 4.0 (3)
			109.62 (2)	4.0 (3)	109.614 (4)	1.09 (24)	109.618 (3)	≤ 4.0 (3)
111.1	1.5	0.44	111.16 (5)	0.55 (9)	111.197 (15)	0.19 (4)	111.18 (1)	0.55 (9)
113.3-114.9	K_β (Np)	-	-	-	-	-	-	-
-	-	-	113.7 (6)	-	-	-	113.9 (5)	-
			114.3 (6)	-	-	-	-	-
117.5-118.4	K_β (Np)	-	-	-	-	-	-	-
-	-	-	117.2 (6)	-	-	-	117.2 (6)	-
-	-	-	117.8 (6)	-	-	-	117.80 (7)	-
-	-	-	117.85 (60)	-	-	-	117.85 (7)	-
-	-	-	121.3 (6)	-	-	-	121.59 (2)	-
-	-	-	-	-	121.69 (4)	0.076 (11)	121.645 (9)	-
-	-	-	122.5 (6)	-	122.76 (7)	0.019 (8)	122.81 (1)	-
-	-	-	126.83 (5)	0.028 (14)	-	-	126.92 (1)	0.028 (14)
-	-	-	131.49 (8)	0.059 (14)	-	-	131.50 (5)	0.059 (14)
-	-	-	132.6 (6)	-	-	-	132.07 (6)	-
135.17 (6)	5.6	1.63	135.19 (2)	1.47 (8)	-	-	135.21 (2)	1.47 (8)
137.02 (6)	5.1	1.48	136.03 (2)	2.05 (6)	136.045 (10)	0.68 (11)	136.045 (2)	2.05 (6)
-	-	-	139.05 (2)	0.024 (7)	-	-	139.05 (3)	≤ 0.024 (7)
-	-	-	139.05 (2)	0.024 (7)	-	-	139.11 (2)	≤ 0.024 (7)
-	-	-	151.07 (5)	0.018 (4)	-	-	151.01 (3)	0.018 (4)
152.75 (6)	0.7	0.2	152.70 (2)	0.150 (8)	152.69 (3)	0.29 (4)	152.70 (2)	≤ 0.150 (8)
-	-	-	152.73 (2)	0.150 (8)	152.69 (3)	0.29 (4)	152.73 (1)	≤ 0.150 (8)
			-	-	153.192 (12)	0.43 (5)	153.19 (1)	-
153.84 (6)	2.4	0.70	153.85 (2)	0.721 (22)	153.870 (9)	0.45 (10)	153.87 (1)	0.721 (22)
-	-	-	156.46 (2)	0.059 (10)	156.452 (2)	4.23 (22)	156.451 (3)	0.059 (10)
-	-	-	160.61 (2)	0.09 (4)	-	-	160.61 (2)	0.09 (4) [#]
163.24 (4)	12.7	3.70	163.25 (2)	3.50 (10)	163.29 (5)	0.28 (4)	163.1 (5)	≤ 3.50 (10)
-	-	-	163.25 (2)	3.50 (10)	-	-	163.29 (1)	≤ 3.50 (10)
-	-	-	164.67 (7)	-	-	-	164.64 (7)	-
-	-	-	165.97 (15)	0.010 (5)	-	-	165.97 (15)	0.010 (5) [#]
-	-	-	170.50 (1)	0.136 (10)	-	-	170.7 (8)	0.136 (10)
-	-	-	174.76 (6)	0.038 (10)	-	-	174.76 (6)	0.038 (10)
-	-	-	176.68 (15)	0.006 (3)	176.62 (5)	0.17 (3)	176.66 (2)	0.006 (3)
-	-	-	182.86 (2)	0.199 (7)	182.876 (2)	13.9 (16)	182.878 (2)	0.199 (7)
-	-	-	189.01 (3)	0.059 (10)	189.099 (6)	0.44 (4)	189.10 (1)	0.059 (10)
-	-	-	190.88 (5)	0.023 (5)	-	-	190.88 (5)	0.023 (5)
194.63 (5)	-	-	194.61 (2)	0.308 (10)	-	-	194.59 (2)	0.308 (10)
-	-	-	196.46 (10)	0.021 (10)	-	-	196.52 (1)	0.021 (10)
206.34 (5)	2.0	0.58	206.37 (2)	0.34 (4)	-	-	206.39 (1)	0.34 (4)
-	-	-	213.20 (14)	0.012 (4)	-	-	213.19 (1)	0.012 (4)
-	-	-	215.52 (2)	0.129 (21)	215.517 (5)	0.81 (4)	215.522 (4)	0.129 (21)
-	-	-	232.40 (3)	0.122 (7)	232.433 (8)	0.29 (5)	232.43 (1)	0.122 (7)
-	-	-	233.69 (10)	0.028 (7)	233.650 (6)	0.243 (22)	233.69 (10)	0.028 (7) [#]
-	-	-	237.02 (10)	0.010 (5)	-	-	236.90 (6)	0.010 (5)
-	-	-	238.53 (5)	0.0035 (18)	-	-	238.35 (7)	0.0035 (18)
-	-	-	250.33 (3)	0.122 (10)	-	-	250.33 (3)	≤ 0.122 (10)
-	-	-	250.33 (3)	0.122 (10)	250.40 (4)	0.35 (6)	250.37 (2)	≤ 0.122 (10)
-	-	-	270.55 (6)	0.0063 (18)	-	-	270.55 (7)	0.0063 (18)
-	-	-	272.75 (7)	0.0081 (18)	-	-	272.80 (6)	0.0081 (18)
-	-	-	280.04 (5)	0.0130 (14)	-	-	280.11 (1)	0.0130 (14)
-	-	-	299.20 (14)	0.006 (3)	-	-	299.23 (6)	0.006 (3)

* emission probability per 100 α has been significantly reduced to < 0.154 , based on the α branching fraction and γ transition probabilities per 100 α of the γ transitions populating the 26.427-keV nuclear level.

[#] not placed in the proposed partial decay scheme.

Placements of gamma-ray transitions.

Adopted E_γ (keV)	Proposed location in decay scheme (^{238}Np nuclear levels)	Adopted E_γ (keV)	Proposed location in decay scheme (^{238}Np nuclear levels)
24.34 (1)	86.674 (2) – 62.330 (4)	121.645 (9)	121.645 (9) – 0
26.427 (2)	26.427 (2) – 0	122.81 (1)	258.853 (8) – 136.045 (2)
32.64 (1)	215.522 (4) – 182.878 (2)	126.92 (1)	342.439 (8) – 215.522 (4)
34.97 (1)	121.645 (9) – 86.674 (2)	131.50 (5)	297.03 (5) – 165.532 (15)
35.90 (1)	62.330 (4) – 26.427 (2)	132.07 (6)	407.59 (6) – 275.519 (9)
43.11 (1)	179.154 (7) – 136.045 (2)	135.21 (2)	300.743 (16) – 165.532 (15)
43.33 (1)	258.853 (8) – 215.522 (4)	136.045 (2)	136.045 (2) – 0
43.83 (2)	106.155 (15) – 62.330 (4)	139.05 (3)	300.743 (16) – 161.69 (2)?
43.89 (2)	165.532 (15) – 121.645 (9)	139.11 (2)	165.532 (15) – 26.427 (2)
46.833 (3)	182.878 (2) – 136.045 (2)	151.01 (3)	312.70 (2) – 161.69 (2)?
49.371 (3)	136.045 (2) – 86.674 (2)	152.70 (2)	258.853 (8) – 106.155 (15)
53.2 (6)	218.7 (6) – 165.532 (15)	152.73 (1)	179.154 (7) – 26.427 (2)
53.67 (1)	232.828 (8) – 179.154 (7)	153.19 (1)	215.522 (4) – 62.330 (4)
53.85 (2)	312.70 (2) – 258.853 (8)	153.87 (1)	275.519 (9) – 121.645 (9)
57.51 (1)	179.154 (7) – 121.645 (9)	156.451 (3)	182.878 (2) – 26.427 (2)
59.32 (1)	121.645 (9) – 62.330 (4)	160.61 (2)	not placed in decay scheme
60.247 (3)	86.674 (2) – 26.427 (2)	163.1 (5)	328.6 (5) – 165.532 (15)
62.330 (4)	62.330 (4) – 0	163.29 (1)	342.439 (8) – 179.154 (7)
66.92 (1)	342.439 (8) – 275.519 (9)	164.64 (7)	300.68 (7) – 136.045 (2)
67.92 (2)	300.743 (16) – 232.828 (8)	165.97 (15)	not placed in decay scheme
73.72 (1)	136.045 (2) – 62.330 (4)	170.7 (8)	389.4 (5) – 218.7 (6)
75.98 (1)	258.853 (8) – 182.878 (2)	174.76 (6)	407.59 (6) – 232.828 (8)
79.48 (1)	215.522 (4) – 136.045 (2)	176.66 (2)	312.70 (2) – 136.045 (2)
79.73 (2)	106.155 (15) – 26.427 (2)	182.878 (2)	182.878 (2) – 0
85.16 (7)	300.68 (7) – 215.522 (4)	189.10 (1)	215.522 (4) – 26.427 (2)
86.674 (2)	86.674 (2) – 0	190.88 (5)	297.03 (5) – 106.155 (15)
89.60 (5)	not placed in decay scheme	194.59 (2)	300.743 (16) – 106.155 (15)
92.48 (1)	179.154 (7) – 86.674 (2)	196.52 (1)	258.853 (8) – 62.330 (4)
93.88 (1)	215.522 (4) – 121.645 (9)	206.39 (1)	342.439 (8) – 136.045 (2)
95.22 (1)	121.645 (9) – 26.427 (2)	213.19 (1)	275.519 (9) – 62.330 (4)
96.204 (3)	182.878 (2) – 86.674 (2)	215.522 (4)	215.522 (4) – 0
96.78 (1)	232.828 (8) – 136.045 (2)	232.43 (1)	258.853 (8) – 26.427 (2)
97.18 (2)	312.70 (2) – 215.522 (4)? X-ray?	233.69 (10)	not placed in decay scheme
109.61 (1)	342.439 (8) – 232.828 (8)	236.90 (6)	299.23 (6) – 62.330 (4)
109.618 (3)	136.045 (2) – 26.427 (2)	238.35 (7)	300.68 (7) – 62.330 (4)
111.18 (1)	232.828 (8) – 121.645 (9)	250.33 (3)	250.33 (3) – 0
113.9 (5)	389.4 (5) – 275.519 (9); X-ray?	250.37 (2)	312.70 (2) – 62.330 (4)
114.3 (6)	not placed in decay scheme; X-ray?	270.55 (7)	376.70 (7) – 106.155 (15)
117.2 (6)	459.6 (6) – 342.439 (8); X-ray?	272.80 (6)	299.23 (6) – 26.427 (2)
117.80 (7)	300.68 (7) – 182.878 (2)? X-ray?	280.11 (1)	342.439 (8) – 62.330 (4)
117.85 (7)	376.70 (7) – 258.853 (8)? X-ray?	299.23 (6)	299.23 (6) – 0
121.59 (2)	300.743 (16) – 179.154 (7)		

Measurements have also been carried out by Hoff *et al.* on the gamma-ray emissions following thermal-neutron capture on ^{237}Np – when judged appropriate, some of these (n, γ) data have been used to develop the proposed decay scheme of $^{242\text{m}}\text{Am}$. The (n, γ) data were inspected in detail, and the opportunity taken to utilize these gamma-ray emission probabilities in a relative sense if their equivalent gamma-ray data

had also been detected and quantified in the $^{242\text{m}}\text{Am}$ studies. For example, consider the depopulation of the 258.853-keV nuclear level:

Proposed depopulating γ -ray transition (keV)	P_γ per 100n (1990Ho02)	P_γ per 100 α (1990Ho02)	P_γ per 100 α calculated	P_γ per 100 α adopted
43.33	0.0046 (7)	–	0.0019 (3)	0.0019 (3)
75.98	0.0051 (8)	–	0.0021 (3)	0.0021 (3)
122.81	0.019 (8)	–	0.008 (4)	0.008 (4)
152.70	≤ 0.29	≤ 0.150	–	≤ 0.150
196.52	–	0.021 (10)	–	0.021 (10)
232.43	0.29 (5)	0.122 (7)	–	0.122 (7)

Unobserved P_γ per 100 α data can be calculated on the reasonable assumption that the relative emission probabilities of these six depopulating gamma rays would be the same irrespective of the mode of feeding to that nuclear level. The emission probability of the 232.43-keV gamma ray has been measured for both the $^{237}\text{Np}(n,\gamma)$ reaction and the α decay of $^{242\text{m}}\text{Am}$, and this ratio can be used to determine the equivalent relative emission probabilities of the 43.33-, 75.98- and 122.81-keV gamma-ray transitions in the α -decay mode. Thus, P_γ per 100 α for the 75.98-keV gamma ray can be calculated:

$$(0.122/0.29) \times 0.0051 (8) = 0.0021 (3)$$

This approach was adopted for a number of specific gamma transitions, as noted in the relevant footnote of the table below.

Despite the introduction of gamma-ray data as outlined above, additional gamma transitions are required to create a more comprehensive and consistent decay scheme for the $^{242\text{m}}\text{Am}$ alpha-decay mode. While some of these possibilities can be gleaned from the $^{237}\text{Np}(n,\gamma)$ reaction data, they cannot be quantified in terms of P_γ per 100 α because of commensurate limitations in the (n, γ) measurements – these particular gamma rays are denoted by a dash (–) within the column entitled “Adopted P_γ per 100 α ” in the table below.

Gamma-ray emissions: multipolarities and theoretical internal conversion coefficients (frozen orbital approximation).

Adopted E_γ (keV)	Adopted P_γ per 100 α *	Multipolarity	α_K	α_L	α_{M+}	α_{total}	
24.34 (1)	0.014 (2) [#]	M1+0.01%E2 $\delta = 0.01$	–	242 (4)	80	322 (5)	α
26.427 (2)	$< 0.154^\ddagger$	M1+1%E2 $\delta = 0.10$	–	252 (4)	86	338 (5)	α
32.64 (1)	0.0041 (6) [#]	M1+0.02%E2 $\delta = 0.014$	–	102.6 (15)	33.8	136.4 (20)	α
34.97 (1)	–	M1+1%E2 $\delta = 0.10$	–	98.7 (14)	33.2	131.9 (19)	α
35.90 (1)	–	M1+1.8%E2 $\delta = 0.135$	–	101.5 (15)	34.5	136.0 (19)	α
43.11 (1)	0.014 (3) [#]	M1+0.2%E2 $\delta = 0.045$	–	46.1 (7)	15.2	61.3 (9)	α
43.33 (1)	0.0019 (3) [#]	M1+9.1%E2 $\delta = 0.32$	–	93.5 (14)	33.2	126.7 (18)	α
43.83 (2)	–	M1+0.9%E2 $\delta = 0.095$	–	47.3 (7)	15.8	63.1 (9)	α
43.89 (2)	–	M1+1.3%E2 $\delta = 0.115$	–	49.2 (7)	16.5	65.7 (10)	α
46.833 (3)	0.0016 (3) [#]	M1+0.4%E2 $\delta = 0.063$	–	36.7 (6)	12.1	48.8 (7)	α
48.60 (5)	IT decay	E4	–	$3.33 (5) \times 10^5$	$3.71 (6) \times 10^5$	$7.04 (8) \times 10^5$	IT
49.371 (3)	29.1 (9)	E1	–	0.615 (9)	0.206	0.821 (12)	α
53.2 (6)	–	(M1+E2)	–	–	–	–	α

Adopted E_γ (keV)	Adopted P_γ per 100 α^*	Multipolarity	α_K	α_L	α_{M+}	α_{total}	
53.67 (1)	0.45 (6)	M1+5.9%E2 $\delta = 0.25$	–	34.2 (5)	11.8	46.0 (7)	α
53.85 (2)	0.0006 (3) [#]	M1+2.4%E2 $\delta = 0.16$	–	27.8 (4)	9.4	37.2 (6)	α
57.51 (1)	0.21 (5)	E1	–	0.412 (6)	0.137	0.549 (8)	α
59.32 (1)	–	M1+E2	–	–	–	–	α
60.247 (3)	1.19 (11)	M1+0.5%E2 $\delta = 0.07$	–	17.34 (25)	5.76	23.1 (4)	α
62.330 (4)	–	E2	–	98.9 (14)	37.1	136.0 (19)	α
66.92 (1)	3.25 (10)	E1	–	0.277 (4)	0.091	0.368 (6)	α
67.92 (2)	0.87 (7)	M1+11%E2 $\delta = 0.35$ (6)	–	18 (2)	6	24 (3)	α
73.72 (1)	1.71 (12)	E1	–	0.214 (3)	0.071	0.285 (4)	α
75.98 (1)	0.0021 (3) [#]	E2	–	38.4 (6)	14.4	52.8 (8)	α
79.48 (1)	0.027 (5) [#]	M1+50%E2 $\delta = 1.0$ (2)	–	19 (3)	7	26 (4)	α
79.73 (2)	–	E2	–	30.6 (5)	11.5	42.1 (6)	α
85.16 (7)	0.21 (7)	M1+50%E2 $\delta = 1.0$ (2)	–	14 (2)	5	19 (3)	α
86.674 (2)	4.97 (15)	M1+1%E2 $\delta = 0.10$	–	5.98 (9)	1.97	7.95 (12)	α
89.60 (5)	0.29 (7)	–	–	–	–	–	α
92.48 (1)	0.61 (7)	E1	–	0.1184 (17)	0.0390	0.1574 (22)	α
93.88 (1)	0.79 (9)	E1	–	0.1138 (16)	0.0375	0.1513 (22)	α
95.22 (1)	–	M1+E2	–	–	–	–	α
96.204 (3)	–	E1	–	0.1068 (15)	0.0352	0.1420 (20)	α
96.78 (1)	0.072 (12) [#]	E2	–	12.28 (18)	4.62	16.90 (24)	α
97.18 (2)	0.0016 (8) [#]	E2	–	12.05 (17)	4.53	16.58 (24)	α
109.61 (1)	≤ 4.0 (3)	M1+50%E2 $\delta = 1.0$ (2)	–	4.9 (5)	1.8	6.7 (7)	α
109.618 (3)	≤ 4.0 (3)	E1	–	0.0760 (11)	0.0250	0.1010 (15)	α
111.18 (1)	0.55 (9)	E1	–	0.0733 (11)	0.0241	0.0974 (14)	α
113.9 (5)	–	E2	–	5.77 (15)	2.17	7.94 (20)	α
114.3 (6)	–	–	–	–	–	–	α
117.2 (6)	–	E1	–	0.0639 (13)	0.0211	0.0850 (17)	α
117.80 (7)	–	(M1+E2)	–	–	–	–	α
117.85 (7)	–	E2	–	4.93 (7)	1.85	6.78 (10)	α
121.59 (2)	–	E2	0.178 (3)	4.27 (6)	1.612	6.06 (9)	α
121.645 (9)	–	E2	0.179 (3)	4.27 (6)	1.601	6.05 (9)	α
122.81 (1)	0.008 (4) [#]	M1+50%E2 $\delta = 1.0$ (2)	5.4 (12)	3.11 (22)	1.09	9.6 (9)	α
126.92 (1)	0.028 (14)	E2	0.196 (3)	3.51 (5)	1.324	5.03 (7)	α
131.50 (5)	0.059 (14)	E1	0.205 (3)	0.0475 (7)	0.0155	0.268 (4)	α
132.07 (6)	–	E1	0.203 (3)	0.0470 (7)	0.0150	0.265 (4)	α
135.21 (2)	1.47 (8)	E1	0.192 (3)	0.0443 (7)	0.0147	0.251 (4)	α
136.045 (2)	2.05 (6)	E1	0.190 (3)	0.0436 (6)	0.0137	0.247 (4)	α
139.05 (3)	≤ 0.024 (7)	E1	0.180 (3)	0.0412 (6)	0.0135	0.235 (4)	α
139.11 (2)	≤ 0.024 (7)	E2	0.211 (3)	2.32 (4)	0.869	3.40 (5)	α
151.01 (3)	0.018 (4)	E1	0.1495 (21)	0.0334 (5)	0.0111	0.194 (3)	α
152.70 (2)	≤ 0.150 (8)	E1	0.1458 (21)	0.0325 (5)	0.0107	0.189 (3)	α
152.73 (1)	≤ 0.150 (8)	E1	0.1457 (21)	0.0324 (5)	0.0107	0.189 (3)	α
153.19 (1)	0.068 (8) [#]	E1	0.1447 (21)	0.0322 (5)	0.0101	0.187 (3)	α
153.87 (1)	0.721 (22)	M1+1.9%E2 $\delta = 0.14$	5.53 (8)	1.123 (16)	0.367	7.02 (10)	α
156.451 (3)	0.059 (10)	E1	0.1379 (20)	0.0305 (5)	0.0100	0.1784 (25)	α
160.61 (2)	0.09 (4)	–	–	–	–	–	α
163.1 (5)	≤ 3.50 (10)	M1+50%E2 $\delta = 1.0$ (2)	2.5 (5)	1.04 (3)	0.36	3.9 (5)	α

Adopted E_γ (keV)	Adopted P_γ per 100 α *	Multipolarity	α_K	α_L	α_{M+}	α_{total}	
163.29 (1)	≤ 3.50 (10)	M1+50%E2 $\delta = 1.0$ (2)	2.5 (5)	1.04 (3)	0.36	3.9 (5)	α
164.64 (7)	–	(M1 + E2)	–	–	–	–	α
165.97 (15)	0.010 (5)	–	–	–	–	–	α
170.7 (8)	0.136 (10)	(M1+50%E2) $\delta = 1.0$ (2)	2.2 (5)	0.882 (23)	0.318	3.4 (5)	α
174.76 (6)	0.038 (10)	M1+50%E2 $\delta = 1.0$ (2)	2.1 (5)	0.809 (17)	0.191	3.1 (4)	α
176.66 (2)	0.006 (3)	E2	0.181 (3)	0.804 (12)	0.300	1.285 (18)	α
182.878 (2)	0.199 (7)	E1	0.0965 (14)	0.0206 (3)	0.0067	0.1238 (18)	α
189.10 (1)	0.059 (10)	E1	0.0894 (13)	0.0190 (3)	0.0062	0.1146 (16)	α
190.88 (5)	0.023 (5)	E1	0.0875 (13)	0.0185 (3)	0.0061	0.1121 (16)	α
194.59 (2)	0.308 (10)	E1	0.0837 (12)	0.01768 (25)	0.00582	0.1072 (15)	α
196.52 (1)	0.021 (10)	E1	0.0819 (12)	0.01725 (25)	0.00565	0.1048 (15)	α
206.39 (1)	0.34 (4)	E2	0.1454 (21)	0.412 (6)	0.1536	0.711 (10)	α
213.19 (1)	0.012 (4)	M1+50%E2 $\delta = 1.0$ (2)	1.19 (24)	0.401 (11)	0.139	1.73 (25)	α
215.522 (4)	0.129 (21)	E1	0.0664 (10)	0.01376 (20)	0.00454	0.0847 (12)	α
232.43 (1)	0.122 (7)	E1	0.0560 (8)	0.01145 (16)	0.00375	0.0712 (10)	α
233.69 (10)	0.028 (7)	–	–	–	–	–	α
236.90 (6)	0.010 (5)	M1+50%E2 $\delta = 1.0$ (2)	0.89 (18)	0.280 (12)	0.100	1.27 (19)	α
238.35 (7)	0.0035 (18)	E1	0.0530 (8)	0.01078 (16)	0.00352	0.0673 (10)	α
250.33 (3)	≤ 0.122 (10)	(M1+50% E2) $\delta = 1.0$ (2)	0.77 (15)	0.233 (12)	0.077	1.08 (16)	α
250.37 (2)	≤ 0.122 (10)	E1	0.0475 (7)	0.00958 (14)	0.00312	0.0602 (9)	α
270.55 (7)	0.0063 (18)	E1	0.0400 (6)	0.00798 (12)	0.00262	0.0506 (7)	α
272.80 (6)	0.0081 (18)	M1+50%E2 $\delta = 1.0$ (2)	0.61 (12)	0.176 (11)	0.064	0.85 (13)	α
280.11 (1)	0.0130 (14)	E1	0.0371 (6)	0.00735 (11)	0.00235	0.0468 (7)	α
299.23 (6)	0.006 (3)	M1+50%E2 $\delta = 1.0$ (2)	0.48 (9)	0.131 (9)	0.039	0.65 (10)	α

* gamma rays with emission probabilities denoted by a dash (–) are believed to be relevant to the α -decay scheme, but could not be quantified in terms of P_γ per 100 α , and are omitted from the final recommendations.

not observed in γ -ray measurements of the α -decay mode of ^{242m}Am – derived from equivalent (n, γ) studies (1990Ho02).

‡ reported emission probability of 1.36 (10) per 100 α decays exceeds expectations considerably from the point of view of the proposed decay scheme and the internal conversion coefficients of this (M1 + 1 % E2) transition – value was reduced to < 0.154 to maintain a satisfactory balance for depopulation of the 26.427(2)-keV nuclear level and population to the ground state of ^{238}Np .

Multipolarities and Internal Conversion Coefficients

The nuclear level schemes specified by Akovali and Chukreev *et al.* have been used to define the multipolarities of the gamma transitions on the basis of known spins and parities (2002Ak06, 2002Ch52). Mixing ratios for many of the (M1 + E2) transitions up to gamma-ray energies of 153.87-keV are the assignments proposed by Hoff *et al.* (1990Ho02), while others were arbitrarily assigned mixing ratios of 1.0 with an uncertainty of 20 %. Others were derived from consideration of population-depopulation of the relevant nuclear levels. Recommended internal conversion coefficients have been determined from the frozen orbital approximation of Kibédi *et al.* (2008Ki07), based on the theoretical model of Band *et al.* (2002Ba85, 2002Ra45).

Significant conflict and inconsistencies remain in the proposed decay scheme for the rather small α branch, despite the impressive work of Hoff *et al.* (1990Ho02). Emphasis has been placed on the validity and comprehensive nature of the α -spectroscopy studies of Baranov *et al.* and Hoff *et al.* (1979Ba67, 1990Ho02) that may be of questionable merit. There are also very strong indications that the known gamma-ray data are unable to support the significant γ depopulation of the 407.59- and 342.439-keV nuclear levels of ^{238}Np . Arguably, further accurate high-resolution gamma-ray spectroscopy studies are required to develop and complete the rather complex α -decay mode with much greater confidence.

Atomic Data

The x-ray and Auger-electron data have been calculated using the evaluated gamma-ray data, and atomic data from 1996Sc06, 1998ScZM and 1999ScZX. Both the x-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.01, 28 January 2003, with the emission.101 database extended to $Z = 96$ to calculate component L x-ray data for IT decay to ^{242}Am). This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data. A number of the gamma-ray emission probabilities can only be quantified in terms of recommended upper limits (\leq and $<$) – EMISSION calculations have been undertaken with these values reduced by a factor of approximately two and assigned uncertainties of the same magnitude.

K and L X-ray emission probabilities per 100 disintegrations of ^{242}mAm .

			Energy (keV)	Photons per 100 disint.
XL		(Np)	11.871 – 21.491	0.37 (4)
	XL ₁	(Np)	11.871	0.0090 (9)
	XL _α	(Np)	13.761 – 13.946	0.143 (13)
	XL _η	(Np)	15.861	0.0022 (4)
	XL _β	(Np)	16.109 – 17.992	0.164 (13)
	XL _γ	(Np)	20.784 – 21.491	0.040 (3)
	XK _α	XK _{α2}	(Np)	97.069
XK _{α1}		(Np)	101.059	0.030 (14)
XK' _{β1}	XK _{β3}	(Np)	113.303) 0.011 (5)
	XK _{β1}	(Np)	114.234	
	XK _{β5}	(Np)	114.912	
XK' _{β2}	XK _{β2}	(Np)	117.463) 0.0037 (17)
	XK _{β4}	(Np)	117.876	
	XKO _{2,3}	(Np)	118.429	
XL		(Am)	12.377 – 22.836	25.0 (11)
	XL ₁	(Am)	12.377	0.608 (18)
	XL _α	(Am)	14.414 – 14.620	9.33 (24)
	XL _η	(Am)	16.819	0.274 (9)
	XL _β	(Am)	16.890 – 19.110	12.2 (3)
	XL _γ	(Am)	22.072 – 22.836	2.90 (8)

Electron energies were determined from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

Data Consistency

An effective Q-value of 74.31 (5) keV has been adopted from the atomic mass evaluation of Audi *et al.* (2003Au03) while in the course of formulating the decay scheme of ^{242}mAm . This value has subsequently been compared with the Q-value calculated by summing the contributions of the individual emissions to the ^{242}mAm alpha- and IT-decay processes (i.e. α , conversion electrons, γ , etc.):

$$\text{calculated Q-value} = \sum (E_i \times P_i) = 72.9 (12) \text{ keV}$$

Percentage deviation from the effective Q-value of Audi *et al.* is $(1.9 \pm 1.6) \%$, which supports the derivation of a reasonably consistent decay scheme with a large variant. Much of this deviation can be attributed to the incompleteness of the recommended alpha- and gamma-transition data within the small alpha-decay mode.

References

- 1950St61 K. STREET, Jr., A. GHIORSO, G.T. SEABORG, The isotopes of americium, *Phys. Rev.* 79 (1950) 530-531. [approximate half-life]
- 1959Ba22 R.F. BARNES, D.J. HENDERSON, A.L. HARKNESS, H. DIAMOND, The alpha and electron capture partial half-lives of ^{242}Am , *J. Inorg. Nucl. Chem.* 9 (1959) 105-107. [BF $_{\alpha}$]
- 1960As05 F. ASARO, I. PERLMAN, J.O. RASMUSSEN, S.G. THOMPSON, Isomers of Am^{242} , *Phys. Rev.* 120 (1960) 934-943. [resolution of isomers]
- 1967Ca04 J.T. CALDWELL, S.C. FULTZ, C.D. BOWMAN, R.W. HOFF, Spontaneous fission half-life of $\text{Am}^{242\text{m}}$, *Phys. Rev.* 155 (1967) 1309-1313. [sf half-life]
- 1977La19 F.P. LARKINS, Semiempirical Auger-electron energies for elements $10 \leq Z \leq 100$, *At. Data Nucl. Data Tables* 20 (1977) 311-387. [Auger-electron energies]
- 1979Ba67 S.A. BARANOV, V.M. SHATINSKII, L.V. CHISTYAKOV, New data on the alpha decay of $^{242\text{m}}\text{Am}$, *Sov. At. Energy* 47 (1980) 1022-1024. [E $_{\alpha}$, P $_{\alpha}$]
- 1979Ze05 A.G. ZELENKOV, V.A. PCHELIN, Yu.F. RODIONOV, L.V. CHISTYAKOV, V.M. SHUBKO, New measurements of the partial half-lives of an isomeric state of $^{242\text{m}}\text{Am}$, *Sov. At. Energy* 47 (1980) 1024-1025. [half-life, BF $_{\alpha}$]
- 1980VyZZ Ts. VYLOV, G.-J. BEYER, V.M. GOROZHANKIN, Zh. ZHELEV, A.I. IVANOV, R.B. IVANOV, V.G. KALINNIKOV, M.Ya. KUZNETSOVA, N.A. LEBEDEV, M.A. MIKHAILOVA, A.I. MUMINOV, A.F. NOVGORODOV, Yu.V. NORSEEV, Sh. OMANOV, B.P. OSIPENKO, E.K. STEPANOV, K. THIEME, V.G. CHUMIN, A.F. SHCHUS, Yu.V. YUSHKEVICH, *Spektren der Strahlung Radioaktiver Nuklide, Gemessen mit Halbleiterdetektoren, Zentralinstitut für Kernforschung, Rossendorf bei Dresden, und Vereinigtes Institut für Kernforschung Dubna, ZfK-399* (1980); Ts. VYLOV, V.M. GOROZHANKIN, Zh. ZHELEV, A.I. IVANOV, R.B. IVANOV, V.G. KALINNIKOV, M.Ya. KUZNETSOVA, N.A. LEBEDEV, M.A. MIKHAILOVA, A.I. MUMINOV, A.F. NOVGORODOV, Yu.V. NORSEEV, Sh. OMANOV, B.P. OSIPENKO, E.K. STEPANOV, V.G. CHUMIN, A.F. SHCHUS, Yu.V. YUSHKEVICH, *Spectra of Radiations of Radioactive Nuclides*, Editor: K.Ya. GROMOV, FAN Publishing, Tashkent, USSR (1980). [E $_X$, P $_X$, E $_{\gamma}$, P $_{\gamma}$]
- 1986Ze06 A.G. ZELENKOV, V.A. PCHELIN, Yu.F. RODIONOV, L.V. CHISTYAKOV, V.S. SHIRYAEV, V.M. SHUBKO, Measurements of the spontaneous-fission half-lives of ^{242}Cm and $^{242\text{m}}\text{Am}$, *Sov. At. Energy* 60 (1986) 492-493. [sf half-life]
- 1990Ho02 R.W. HOFF, S. DRISSI, J. KERN, W. STRASSMANN, H.G. BÖRNER, K. SCHRECKENBACH, G. BARREAU, W.D. RUHTER, L.G. MANN, D.H. WHITE, J.H. LANDRUM, R.J. DUPZYK, R.F. CASTEN, W.R. KANE, D.D. WARNER, Nuclear structure of ^{238}Np from neutron-capture and α -decay measurements, *Phys. Rev.* C41 (1990) 484-512. [E $_{\alpha}$, P $_{\alpha}$, E $_{\gamma}$, P $_{\gamma}$]
- 1996Sc06 E. SCHÖNFELD, H. JANßEN, Evaluation of atomic shell data, *Nucl. Instrum. Methods Phys. Res.* A369 (1996) 527-533. [X $_K$, X $_L$, Auger electrons]
- 1998Ak04 Y.A. AKOVALI, Review of alpha-decay data from doubly-even nuclei, *Nucl. Data Sheets* 84 (1998) 1-114. [alpha decay, r_0 parameters]

- 1998ScZM E. SCHÖNFELD, G. RODLOFF, Tables of the energies of K-Auger electrons for elements with atomic numbers in the range from $Z = 11$ to $Z = 100$, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX E. SCHÖNFELD, G. RODLOFF, Energies and relative emission probabilities of K X-rays for elements with atomic numbers in the range from $Z = 5$ to $Z = 100$, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2002Ak06 Y.A. AKOVALI, Nuclear data sheets for $A = 242$, Nucl. Data Sheets 96 (2002) 177-239. [nuclear levels]
- 2002Ba85 I.M. BAND, M.B. TRZHASKOVSKAYA, C.W. NESTOR, Jr., P.O. TIKKANEN, S. RAMAN, Dirac–Fock internal conversion coefficients, At. Data Nucl. Data Tables 81 (2002) 1-334. [ICC]
- 2002Ch52 F.E. CHUKREEV, V.E. MAKARENKO, M.J. MARTIN, Nuclear data sheets for $A = 238$, Nucl. Data Sheets 97 (2002) 129-240. [nuclear levels]
- 2002Ra45 S. RAMAN, C.W. NESTOR, Jr., A. ICHIHARA, M.B. TRZHASKOVSKAYA, How good are the internal conversion coefficients now? Phys. Rev. C66 (2002) 044312, 1-23. [ICC]
- 2003Au03 G. AUDI, A.H. WAPSTRA, C. THIBAUT, The AME2003 atomic mass evaluation (II). Tables, graphs and references, Nucl. Phys. A729 (2003) 337-676. [Q-value]
- 2008Ki07 T. KIBÉDI, T.W. BURROWS, M.B. TRZHASKOVSKAYA, P.M. DAVIDSON, C.W. NESTOR, Jr., Evaluation of theoretical conversion coefficients using BrIcc, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202-229. [ICC]

²⁴²Cm - Comments on evaluation of decay data by V.P. Chechev

This evaluation was completed in February 2005 (see 2006Ch34) and then corrected in October 2009 with a literature cut-off by the same date.

1 Decay Scheme

The decay scheme is based on the evaluation of Chukreev *et al.* (2002Ch52) and can be considered essentially complete although some weak gamma-ray transitions have not been observed in ²⁴²Cm alpha decay. Such gamma rays were taken from ²³⁸Am→²³⁸Pu, ²³⁸Np→²³⁸Pu decays and have been included in the decay scheme.

2 Nuclear Data

Q(α) is from 2003Au03.

The evaluated half-life of ²⁴²Cm is based on the experimental results given in Table 1. Re-estimated values were used for averaging when needed.

Table 1. Experimental values of the ²⁴²Cm half-life (in days)

Reference	Author(s)	Original value	Re-estimated value	Measurement method
1950Ha14	Hanna et al.	162.5 (20)	-	α-counting with low geometry counter
1954Gl37	Glover and Milsted	162.46 (14) ^a	162.46 (32) ^c	α-counting with low geometry counter
1954Hu32	Hutchinson and White	163.0 (18)	-	Calorimetry
1957Treiman	Treiman et al.	162.7 (1)	-	Calorimetry
1965Fl02	Flynn et al.	164.4 (4)	163.1 (4) ^d	2π α counting
1975Ke02	Kerrigan and Banick	163.2 (2) ^b	-	Calorimetry
1977Di04	Diamond et al.	162.76 (4)	162.76 (8) ^c	Intermediate geometry α-counting
1979Ch41	Chang et al.	163.02 (11)	163.02 (18) ^c	α-counting with low geometry counter
1980Jadhav	Jadhav et al.	162.13 (215)	162.13 (225)	α-spectrometry with solid state detector
1981Us03	Usuda and Umezawa	161.35 (20)	161.35 (30) ^c	α-counting with 2π proportional counter
1982Ag02	Aggarwal et al.	163.17 (6)	163.17 (11) ^c	α-counting with proportional counter
1982Ag02	Aggarwal et al.	162.82 (21)	162.82 (26) ^c	α-spectrometry with solid state detector
1984Wi14	Wiltshire et al.	163.0 (2)	-	α-counting with low geometry counter

^a The uncertainty of 0.27 quoted by authors, which corresponds to 95 % confidence level, has been reduced by a factor 2.

^b The uncertainty of 0.4 quoted by authors, which corresponds to 95 % confidence level, has been reduced by a factor 2.

^c Quoted uncertainties have been re-estimated in 1986LoZT.

^d The value has been recalculated in 1977Di04.

The LWEIGHT and EV1NEW computer programs identified two outliers in the above data set. These are the values from 1981Us03 and 1980Jadhav. Omitting these values in the calculation and using the remaining 11 results produced a weighted mean of 162.86 with an internal uncertainty of 0.05 and an external uncertainty of 0.06 ($\chi^2/\nu = 1.6$). The EV1NEW program has chosen the smallest experimental uncertainty of 0.08 as the uncertainty of the weighted average.

Thus the recommended value of the ²⁴²Cm half-life is 162.86 (8) days.

The evaluated spontaneous fission partial half-life of ²⁴²Cm is based on the experimental results given in Table 2. Re-estimated values were used for averaging when needed.

Table 2. Experimental values of the ²⁴²Cm spontaneous fission half-life (in 10⁶ years)

Reference	Author(s)	Original value	Re-estimated value ^a	Measurement method
1951Ha87	Hanna et al.	7.2 (2)	-	Fission fragment counting, ionization chamber
1967Ar09	Armani and Gold	6.09 (18)	6.82 (18)	Fission neutron counting, LiI detector
1979Ch41	Chang et al.	7.46 (6)	-	Mica fission track detector
1982Ra33	Raghuraman et al.	7.15 (15)	-	Solid state detector
1982UmZZ	Umezawa et al.	6.89 (17)	-	Mica fission track detector
1986Ze06	Zelenkov et al.	6.9 (3)	6.98 (33)	α /SF, Si(Au) detectors
1989Us04	Usuda et al.	6.96 (18)	-	Absolute fission track counting

^a Recalculated in 2000Ho27

Omitting the value of 1979Ch41 (outlier) the weighted mean of the six remaining values becomes 7.005 with an internal uncertainty of 0.076 and an external uncertainty of 0.063 ($\chi^2/\nu = 0.69$).

The recommended value of the ²⁴²Cm spontaneous fission half-life is 7.01 (15) 10⁶ years, where the uncertainty is the smallest quoted uncertainty of 6 experimental results.

2.1 α Transitions

The energies of the alpha-particle transitions given in Section 2.1 have been deduced from the Q value and the level energies given in Table 3 from 2002Ch52.

Table 3. ²³⁸Pu levels populated in the ²⁴²Cm α -decay

Level number	Energy (keV)	Spin and parity	Half-life	Probability of α -transition (%)
0	0.0	0 ⁺	87.74 (3) a	74.06 (7)
1	44.08 (3)	2 ⁺	177 (5) ps	25.94 (7)
2	146.00 (5)	4 ⁺		0.034 (2)
3	303.42 (7)	6 ⁺		0.0046 (5)
4	513.62 (16)	8 ⁺		2×10^{-5}
5	605.08 (7)	1 ⁻		$2.5 (5) \times 10^{-4}$
6	661.28 (11)	3 ⁻		$1.3 (3) \times 10^{-5}$
7	763.22 (12)	5 ⁻		$\leq 2.2 \times 10^{-7}$
8	941.44 (9)	0 ⁺		$3.5 (7) \times 10^{-5}$
9	962.72 (8)	1 ⁻		$1.13 (21) \times 10^{-6}$
10	983.00 (9)	2 ⁺		$1.7 (5) \times 10^{-6}$
11	1018.6 (3)	1 ⁻		$\leq 2 \times 10^{-7}$
12	1028.62 (5)	2 ⁺		$3.7 (10) \times 10^{-6}$
13	1125.79 (17)	(4 ⁺)		$3.1 (10) \times 10^{-7}$
14	1228.69 (22)	0 ⁺		$5.5 (15) \times 10^{-7}$
15	1264.29 (22)	2 ⁺		$5.2 (14) \times 10^{-7}$

The emission probabilities of the most intensive transitions $\alpha_{0,i}$ ($i = 0$ to 4) have been obtained by averaging experimental data (Table 4). The emission probabilities of the remaining α -particle transitions have been deduced either from the $P(\gamma+ce)$ decay-scheme balances or by averaging experimental and deduced values (for example, $\alpha_{0,5}$).

Table 4. Experimental, calculated and recommended α -transition probabilities (%) in the ²⁴²Cm decay

	α -particle energy (keV)	1953As14	1958Ko87	1963Dz07	1966Ba07	1998Ya17	Deduced from decay-scheme balance c	Recommended
$\alpha_{0,0}$	6113	73.7 (5)	73.5 (5)	74 (2)	74.2 (5) ^a	74.08 (7)		74.06 (7) ^d
$\alpha_{0,1}$	6069	26.3 (5)	26.5 (5)	26.0 (9)	25.8 (5) ^a	25.92 (6)		25.94 (7) ^d
$\alpha_{0,2}$	5969	0.035 (2) ^a	0.030 (2) ^b	0.035 (2)	0.036 (2) ^a			0.034 (2) ^e
$\alpha_{0,3}$	5816		0.0046 (5)		0.0046			0.0046 (5) ^f
$\alpha_{0,4}$	5608			1963Bj01	$2 \cdot 10^{-5}$			$2 \cdot 10^{-5}$ ^g
$\alpha_{0,5}$	5518			$2.8 (5) \cdot 10^{-4}$	$2.5 (6) \cdot 10^{-4}$		$2.6 (7) \cdot 10^{-4}$	$2.5 (5) \cdot 10^{-4}$ ^e
$\alpha_{0,6}$	5462						$1.3 (3) \cdot 10^{-5}$	$1.3 (3) \cdot 10^{-5}$ ^c
$\alpha_{0,7}$	5366						$2.2 \cdot 10^{-7}$	$2.2 \cdot 10^{-7}$ ^c
$\alpha_{0,8}$	5187			$3.4 (8) \cdot 10^{-5}$	$2.5 (8) \cdot 10^{-5}$		$3.5 (7) \cdot 10^{-5}$	$3.5 (7) \cdot 10^{-5}$ ^c
$\alpha_{0,9}$	5166						$1.13 (21) \cdot 10^{-6}$	$1.13 (21) \cdot 10^{-6}$ ^c
$\alpha_{0,10}$ ^h	5146				$\leq 5 \cdot 10^{-6}$		$1.7 (5) \cdot 10^{-6}$	$1.7 (5) \cdot 10^{-6}$ ^c

^aNo uncertainties are quoted by the authors. The uncertainties have been adopted by the evaluator based on the similarity of the spectra measured with magnetic spectrometers in 1953As14, 1958Ko87 and 1966Ba07.

^bThe uncertainty of 0.001 quoted by authors has been increased by a factor of 2 by the evaluator (see ^a).

^cDeduced from $P(\gamma+ce)$ decay-scheme balances for corresponding ²³⁸Pu levels.

^dWeighted average of experimental values. The experimental data of 1998Ya17 have been obtained by the most accurate method (using a semiconductor detector).

^eWeighted average of experimental and deduced values.

^fAdopted experimental value from 1958Ko87.

^gAdopted experimental value from 1966Ba07.

^hThe probabilities of remaining alpha-transitions ($\alpha_{0,11}$ and $\alpha_{0,15}$) have been deduced from $P(\gamma+ce)$ decay-scheme balances.

2.2 γ Transitions and Internal Conversion Coefficients

The recommended energies of gamma-ray transitions are essentially the same as the gamma-ray energies because nuclear recoil is negligible.

The probabilities, $P(\gamma+ce)$, for gamma-ray transitions of 44-($\gamma_{1,0}$), 102-($\gamma_{2,1}$), 157-($\gamma_{3,2}$), and 210-keV ($\gamma_{4,3}$) have been deduced from transition- probability balances, using the emission intensities of α -transitions directly measured.

For E0- gamma transitions 941-($\gamma_{8,0}$) and 1229-keV ($\gamma_{14,0}$) the $P(\gamma+ce)$ values have been taken from data on the electron capture decay ²³⁸Am \rightarrow ²³⁸Pu (see 2002Ch52 and references therein).

The remaining $P(\gamma+ce)$ values have been obtained from the gamma-ray emission probabilities and the total internal conversion coefficients (ICC's). The experimental values of ICC's have been adopted for (E0+E2) gamma-ray transitions 939-($\gamma_{10,1}$) and 1220-keV ($\gamma_{15,1}$). The remaining ICC's have been interpolated using the BrIcc package with the so called "Frozen Orbital" approximation (2008Ki07). The relative uncertainties of α_K , α_L , α_M , α_T for pure multiplicities have been taken as 2 %.

The multiplicities and E2/M1, M2/E1 mixing ratios have been taken from 2002Ch52. These are based on conversion electron measurements of 1952Du12, 1956Ba95, 1956Sm18, 1960As10, and 1965Ak02 made in the ²⁴²Cm α -decay.

3 Atomic Data

3.1 Fluorescence yields

Fluorescence yield data are from 1996Sc06 (Schönfeld and Janßen).

3.1.1 X rays

The Pu KX-ray energies and relative emission probabilities have been taken from 1999Schönfeld, where the calculated energy values are based on X-ray wavelengths from 1967Be65 (Bearden). In Table 5 the recommended values of Pu KX-ray energies are compared with experimental results.

Table 5. Experimental and recommended values of Pu KX-ray energies (keV)

	1980Di13	1982Ba56	Recommended
K α_2	99.55 (3)	99.530 (2)	99.525
K α_1	103.76 (3)	103.741 (2)	103.734
K β_3	116.27	116.242 (2)	116.244
K β_1	117.26	117.233 (2)	117.228
K $\beta_{2,4}$	120.60 (15)	-	120.553
KO $_{2,3}$	121.55 (6)	-	121.543

The Pu KX-ray energies in 1980Di13 were measured in the alpha decay of ²⁴⁵Cm. The relative emission probabilities of KX-rays were given as:

$$K\alpha_2 : K\alpha_1 : K\beta_3 : K\beta_1 : K\beta_{2,4} = 64.7 (23) : 100.0 (33) : 12.9 (7) : 23.1 (10) : 8.9 (5).$$

3.1.2 Auger Electrons

The energies of Auger electrons have been calculated from atomic electron binding energies.

The P(KLX)/P(KLL), P(KXY)/P(KLL) ratios have been taken from 1996Sc06.

4 α Emissions

The energy of the alpha-particle group to the ground state of ²³⁸Pu, E($\alpha_{0,0}$) is from the absolute measurement of 1971Gr17, with a correction of -0.20 keV recommended by A. Rytz in 1991Ry01.

The energies of all other α particles have been deduced from Q $_{\alpha}$ and the ²³⁸Pu level energies including the recoil energy corrections (see 2002Ch52).

In Table 6 the recommended values of α -particle energies are compared with the experimental results obtained with magnetic alpha spectrometers.

Table 6. Experimental ^a and recommended α -emission energies in the decay of ²⁴²Cm (keV)

	1953As14	1958Ko87	1963Dz07	1966Ba07 1971Bb10	1971Gr17	Recommended
$\alpha_{0,0}$	6113	6114	6113 (1)	6112.9 (3)	6112.72 (8)	6112.72 (8)
$\alpha_{0,1}$	6069	6070	6069 (1)	6069.5 (5)	6069.43 (12)	6069.37 (9)
$\alpha_{0,2}$	5968	5968 (2)	5969 (3)	5970		5969.24 (9)
$\alpha_{0,3}$	-	5816 (2)	-	5817		5816.39 (11)
$\alpha_{0,4}$	-	-	-	5609		5607.76 (16)
$\alpha_{0,5}$	-	-	-	5514		5517.75 (11)
$\alpha_{0,8}$	-	-	-	5189		5186.95 (12)
$\alpha_{0,10}$	-	-	-	5146		5146.07 (12)

^a Authors' values have been adjusted for changes in calibration energies (see 1991Ry01)

5 Electron emissions

The energies of conversion electrons have been obtained using gamma-ray transition energies and electron binding energies. The emission probabilities of conversion electrons have been deduced from the evaluated $P(\gamma)$ and ICC values.

The absolute emission probabilities of K and L Auger electrons have been calculated using the EMISSION computer program (2000Schönfeld).

6 Photon emissions

6.1. X-Ray emissions

The absolute emission probabilities of U KX- and U LX-rays in decay of ²⁴²Pu have been calculated using the EMISSION computer program (2000Schönfeld).

The calculated total absolute emission probability of LX-rays $P(XL) = 9.92$ (23) % agrees well with the experimental value of 9.70 (14) % from 1970By01 and disagrees with the value of 11.7 (3) % measured in 1971Swinth.

The relative Pu LX-ray emission probabilities in ²⁴²Cm α -decay measured in 1990Po14 [4.9 (8)-L β ; 66 (7)-L α ; 100-L $\eta\beta$; 23 (3)-L γ] agree well with the values calculated using the EMISSION computer program with the exception of $L\alpha/L\eta\beta^{\text{calc.}} = 79$ (4) / 100. The latter agrees well with the experimental result from 1995Jo23, $L\alpha/L\eta\beta$ (Pu) = 80.9 (9) / 100, obtained for LX-rays in the decay of other even-even curium isotope – ²⁴⁴Cm.

6.2 Gamma emission

6.2.1. Gamma-ray energies

The energy of the 44-keV gamma ray ($\gamma_{1,0}$) is from ²³⁸Np \rightarrow ²³⁸Pu β^- decay (1972Wi22); it agrees with the less accurate measurements in ²⁴²Cm α -decay (44.11 (5) keV - 1956Sm18) and in ²³⁸Am ϵ -decay (44.1 (1) keV - 1972Ah04).

The energies of the 102-($\gamma_{2,1}$), 157-($\gamma_{3,2}$), 336-($\gamma_{8,5}$), 358-($\gamma_{9,5}$), 605-($\gamma_{5,0}$), 940-($\gamma_{10,1}$), and 941-($\gamma_{8,0}$) keV gamma rays have been obtained from the available experimental data of 1981Le15 (²⁴²Cm α -decay and ²³⁸Np β^- -decay), 1972Wi22, 1972Ah04, 1956Sm18, and 1971Po09 (²³⁸Am ϵ -decay) using the adopted ²³⁸Pu level energies.

The energies of the 210-($\gamma_{4,3}$), 617-($\gamma_{7,2}$), and 883-($\gamma_{12,2}$) keV gamma rays, which were not observed in the ²⁴²Cm α -decay, have been deduced from the adopted level energies. The energies of the remaining gamma rays have been taken from the measurements of 1981Le15 (²⁴²Cm α -decay).

6.2.2. Gamma-ray emission probabilities

The absolute emission probabilities for gamma-rays of 44-($\gamma_{1,0}$), 102- ($\gamma_{2,1}$), 157-($\gamma_{3,2}$), and 210-($\gamma_{4,3}$) keV have been deduced from decay-scheme probability balances using the intensities of α -transitions evaluated directly from experimental data.

The absolute emission probabilities of > 300 keV gamma-rays (except for 883- and 1229-keV γ -rays) have been obtained from relative gamma-ray emission probabilities $P(\gamma)/P(\gamma\ 561\text{keV})$ measured in 1981Le15. The normalization factor $P(\gamma\ 561\text{keV}) = 1.5 \cdot 10^{-4}$ per 100 disintegrations, which was used here, was estimated in 1981Le15 using a previous $\alpha\gamma$ coincidence measurement of the sum of the absolute emission probabilities of the 515-, 561-, 605-, and 617-keV gamma-rays (1963Le17).

$P(\gamma\ 883\text{keV})$ and $P(\gamma\ 1229\text{keV})$ are from 2002Ch52, using the experimental data on ²³⁸Np β^- -decay and ²³⁸Am ε -decay, respectively.

7. Consistency of recommended data

The most accurate Q value, Q(M), is taken from the atomic mass adjustment table of Audi et al. (2003Au03). Comparison of Q(eff)(deduced as the sum of average energies per disintegration ($\sum E_i \times P_i$) for all emissions accompanying ²⁴²Cm α - decay) with the tabulated decay energy Q(M) allows to check a consistency of the recommended decay-scheme parameters obtained in this evaluation.

Here E_i and P_i are the evaluated energies and emission probabilities of the i-th alpha particle, beta particle, gamma ray, X-ray, etc. Consistency (percentage deviation) is determined by $\{|Q(M) - Q(\text{eff})\}/Q(M)\} \times 100$. "Percentage deviations above 5 % would be regarded as high and imply a poorly defined decay scheme; a value of less than 5 % indicates the construction of a reasonably consistent decay scheme" (quoted from the article by A.L. Nichols in Appl. Rad. Isotopes 55 (2001) 23-70).

For the above ²⁴²Cm decay data evaluation we have $Q(M) = 6215.56(8)$ keV and $Q(\text{eff}) = 6217(6)$ keV, i.e. consistency is better than 0.12 %.

8. References

- 1950Ha14 G. C. Hanna, B. G. Harvey, N. Moss, Phys. Rev. 78, 617 (1950) [Half-life]
- 1951Ha87 G. C. Hanna, B. G. Harvey, N. Moss, P. R. Tunncliffe, Phys. Rev. 81, 466 (1951) [SF half-life]
- 1952Du12 D. C. Dunlavey, G. T. Seaborg, Phys. Rev. 87, 165 (1952) [Conversion electron measurements, gamma ray multipolarities]
- 1953As14 F. Asaro, S. G. Thompson, I. Perlman, Phys. Rev. 92, 694 (1953) [α -transition energies and probabilities]
- 1954Gl37 K. M. Glover, J. Milsted, Nature 173, 1238 (1954) [Half-life]
- 1954Hu32 W. P. Hutchinson, A. G. White, Nature 173, 1238 (1954) [Half-life]
- 1956Ba95 S. A. Baranov, K. N. Shlyagin, At. Energ. USSR 1, 52 (1956); J. Nuclear Energy 3, 132 (1956) [Conversion electron measurements, gamma-ray multipolarities]
- 1956Sm18 W. G. Smith, J. M. Hollander, Phys. Rev. 101, 746 (1956) [Gamma ray energies and multipolarities]
- 1957Treiman L. N. Treiman, R. A. Penneman, B. Bevan, unpublished, cited in Ref.: J. Inorg. Nucl. Chem., 5, 6(1957) [Half-life]

- 1958Ko87 L. N. Kondratev, V. B. Dedov, L. L. Goldin, *Izvest. Akad. Nauk SSSR, Ser. Fiz.* 22, 99 (1958); *Columbia Tech. Transl.* 22, 97 (1959) [α -emission energies and probabilities]
- 1960As10 F. Asaro, I. Perlman, UCRL-9566, p.50 (1960)
[Conversion electron measurements, gamma-ray multipolarities]
- 1963Dz07 B. S. Dzhelepov, R. B. Ivanov, V. G. Nedovesov, V. P. Chechev, *Zh. Eksperim. i Teor. Fiz.* 45, 1360 (1963); *Soviet Phys. JETP* 18, 937 (1964) [α -emission energies and probabilities]
- 1963Le17 C. M. Lederer (1963), Thesis, Univ California (1963); UCRL-11028
[Absolute gamma-ray emission probabilities]
- 1965Ak02 G. G. Akalaev, N. A. Vartanov, P. S. Samoilov, NP-14688 (1965)
[Conversion electron measurements, gamma-ray multipolarities]
- 1965Fl02 K. F. Flynn, L. E. Glendenin, E. P. Steinberg, *Nucl. Sci. Eng.* 22, 416 (1965) [Half-life]
- 1966Ba07 S. A. Baranov, Y. F. Rodionov, V. M. Kulakov, V. M. Shatinskii, *Yadern. Fiz.* 4, 1108 (1966); *Soviet J. Nucl. Phys.* 4, 798 (1967) [α -energies and transition probabilities]
- 1967Ar09 R. J. Armani, R. Gold, (1967). *Proc Symp Standardization of Radionuclides, Vienna, Austria* (1966), Intern. At. Energy Agency, Vienna, p. 621 [SF half-life]
- 1967Be65 J. A. Bearden, *Rev. Mod. Phys.* 39, 78(1967) [X-ray energies]
- 1970By01 J. Byrne, R. J. D. Beattie, S. Benda, I. Collingwood, *J. Phys.* B3, 1166 (1970)
[Experimental LX-ray absolute emission probability]
- 1971Bb10 S. A. Baranov, V. M. Shatinskii, V. M. Kulakov, *Yad. Fiz.* 14, 1101 (1971); *Sov. J. Nucl. Phys.* 14, 614 (1972) [α -particle energies]
- 1971Gr17 B. Grennberg, A. Rytz, *Metrologia* 7, 65 (1971) [α -particle energies]
- 1971Po09 J. C. Post, A. H. W. Aten, Jr, *Radiochim Acta* 15, 205 (1971) [Gamma-ray energies]
- 1971Swinth K. L. Swinth, *IEEE Trans. Nucl. Sci.* 18, No.1, Part 1, 125 (1971)
[Experimental LX-ray absolute emission probability]
- 1972Ah04 I. Ahmad, R. K. Sjoblom, R. F. Barnes, F. Wagner, Jr, P. R. Fields, *Nucl. Phys.* A186, 620 (1972) [Gamma-ray energies]
- 1972Wi22 W. J. B. Winter, A. H. Wapstra, P. F. A. Goudsmit, J. Konijn, *Nucl. Phys.* A197, 417 (1972) [Gamma ray energies]
- 1975Ke02 W. J. Kerrigan, C. J. Banick, *J. Inorg. Nucl. Chem.* 37, 641(1975) [Half-life]
- 1977Di04 H. Diamond, W. C. Bentley, A. H. Jaffey, K. F. Flynn, *Phys. Rev.* C15, 1034 (1977) [Half-life]
- 1979Ch41 Huan-Qiao Chang, Jin-Cheng Xu, Tong-Qing Wen, *Chin. J. Nucl. Phys.* 1, 21 (1979) [SF half-life]
- 1980Di13 J. K. Dickens, J. W. McConnell, *Phys. Rev.* C22, 1344 (1980) [Experimental X-ray energies]
- 1980Jadhav A. V. Jadhav, K. A. Mathew, K. Raghuraman, C. K. Sivaramakrishnan, *Proc. of the Nucl. Chem. and Radiochem. Symp., Waltair, 1980*, 184 [Half-life]

- 1981Le15 C. M. Lederer, Phys. Rev. C24, 1175 (1981) [Gamma-ray energies and probabilities]
- 1981Us03 S. Usuda, H. Umezawa, J. Inorg. Nucl. Chem. 43, 3081 (1981) [Half-life]
- 1982Ag02 S. K. Aggarwal, A. V. Jadhav, S. A. Chitambar et al., Radiochem. Radioanal. Letters 54, 99 (1982) [Half-life]
- 1982Ba56 G. Barreau, H. G. Borner, T. von Egidy, R. W. Hoff, Z. Phys. A308, 209 (1982) [Experimental X-ray energies]
- 1982Ra33 K. Raghuraman, N. K. Chaudhuri, A. V. Jadhav, C. K. Sivaramakrishnan, R. H. Iyer, Radiochem. Radioanal. Letters 55, 1 (1982) [SF half-life]
- 1982UmZZ H. Umezawa, Progress Report to the International Atomic Energy Agency on the Measurement of Nuclear Decay Data of Curium-242. In: INDC(NDS)-138/GE, p. 32 (1982) [SF half-life]
- 1984Wi14 R. A. P. Wiltshire, Nuclear Instrum. Methods 223, 535 (1984) [Half-life]
- 1986LoZT A. Lorenz, IAEA Tech. Rept. Ser., No.261 (1986) [Half-life evaluation]
- 1986Ze06 A. G. Zelenkov, V. A. Pchelin, Yu. F. Rodionov, L. V. Chistyakov, V. S. Shiryaev, V. M. Shubko, Sov. At. Energy 60, 492 (1986) [SF half-life]
- 1989Us04 S. Usuda, H. Umezawa, Int. J. Radiat. Appl. Instr. D (Nucl. Tracks Radiat. Meas.) 16, 247 (1989) [SF half-life]
- 1990Po14 Yu. S. Popov, I. B. Makarov, D. Kh. Srurov, E. A. Erin, Radiokhimiya 32, 2 (1990); Sov. J. Radiochemistry 32, 425 (1990) [Experimental relative LX-ray emission probabilities]
- 1991Ry01 A. Rytz, At. Data Nucl. Data Tables. 47, 205 (1991) [Alpha-emission energies]
- 1995Jo23 P. N. Johnston, P. A. Burns, Nucl. Instrum. Methods Phys. Res. A361, 229 (1995) [Experimental relative LX-ray emission probabilities]
- 1996Sc06 E. Schönfeld, H. Janßen, Nucl. Instrum. Methods Phys. Res. A369, 527 (1996) [Atomic data]
- 1998Ya17 J. Yang, J. Ni, Nucl. Instrum. Methods Phys. Res. A413, 239 (1998) [α-transition probabilities]
- 1999Schönfeld E. Schönfeld, G. Rodloff, PTB-6.11-1999-1999-1, Braunschweig, February 1999 [KX ray energies and relative emission probabilities]
- 2000Ho27 N. E. Holden, D. C. Hoffman, Pure Appl. Chem. 72, 1525 (2000) [SF half-life]
- 2000Schönfeld E. Schönfeld, H. Janßen, Appl. Rad. Isot. 52(2000)595. [X-ray and Auger electron emission probabilities, EMISSION code]
- 2002Ch52 F. E. Chukreev, V. E. Makarenko, M. J. Martin, Nucl. Data Sheets 97, 129 (2002) [Nuclear data evaluation for A=238]
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, Nucl. Phys. A729, 337 (2003) [Q value]
- 2006Ch34 V. P. Chechev, Phys. Atomic Nuclei 69, 1188 (2006) [²⁴²Cm decay data evaluation-2005]
- 2008Ki07 T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor, Jr., Nucl. Instrum. Methods Phys. Res. A589, 202 (2008) [Theoretical ICC]

**²⁴³Am - Comments on evaluation of decay data
by E. Browne, M.-M. Bé, R.G. Helmer**

This evaluation was completed in September 2004 and reviewed in 2009. The literature available by April 2009 was included. Half-life and conversion coefficients have been updated.

Several measurements of the α emission intensities were carried out and their results are in good agreement. However, the available experimental γ -ray emission intensities are mostly imprecise and in poor agreement with each other.

The decay scheme overall consistency is supported by the agreement between $Q(\text{eff}) = 5439.6 (40)$ keV, deduced from average radiation energies and intensities, and $Q(\alpha) = 5438.8 (10)$ keV, from the atomic mass adjustment (2003Au03).

Evaluation Procedures

The *Limitation of Relative Statistical Weight* (LWM) [1985ZiZY] method, used for averaging numbers throughout this evaluation, provided a uniform approach for the analysis of discrepant data.

1 Decay Scheme

²⁴³Am decays 100 % by emission of α particles, with a minute branch of $3.8 (7) \times 10^{-9}$ % (2002Sa53) by spontaneous fission. Other value: $3.7 (9) \times 10^{-9}$ % (1966Gv01). The α -particle intensities (in percent) to individual levels presented in the decay scheme are experimental values from α -spectroscopic measurements. α -hindrance factors given in the decay scheme have been calculated by using a radius parameter r_0 (²³⁹Np) = 1.505, average of r_0 (²³⁸U) = 1.5143 (9), r_0 (²⁴⁰U) = 1.5062 (10), r_0 (²³⁸Pu) = 1.5013 (10), and r_0 (²⁴⁰Pu) = 1.4979 (7) (1998Ak04). The level energies, spins, parities, as well as γ -ray multiplicities shown in the decay scheme are recommended values from the evaluation 2003Br12.

Levels at 71- and 122 keV are based on α - γ coincidence experiments with γ rays (169-, 50.6-, and 195 keV) that feed such levels. The de-excitations of these two levels, however, have not been observed. The expected γ rays may have been masked by more intense ones, which de-excite other levels.

2 Nuclear Data

The recommended half-life of ²⁴³Am is 7367 *a*, a weighted average of the values given in Table 1, the most accurate value (from 2007Ag02) contributes 54 % to the statistical weight. The calculated internal uncertainty is 17 *a*. However, the recommended uncertainty is the smallest uncertainty in the input values, i.e., 23 *a*. This half-life compares well with other recommended values such as 7370 (40) *a* (1992Ak06), 7366 (20) *a* (1991BaZS), and 7370 (15) *a* (1986LoZT).

$Q_\alpha = 5438.8 (10)$ keV is from the atomic mass adjustment 2003Au03.

Table 1. ²⁴³Am measured half-life values

Reference	Method	$T_{1/2}({}^{243}\text{Am})/T_{1/2}({}^{241}\text{Am})$	$T_{1/2}({}^{243}\text{Am})$ (a)	u_c	Remarks	
1959Ba22	Relative activity	16.85 (35)	7289.3 *	151.7	An uncertainty of 1.4 % (100 a) from 1960Be10 is mostly systematic. Thus, dividing this value by the square root of the number of measurements (5) is questionable and was not done in the evaluation of 1986LOZT. Omitted from analysis.	
1960Be10	Relative activity	16.70 (10)	7224.4 *	100.0		
1968Br22	Relative activity	16.96 (13)	7336.9 *	57.2	This value is the weighted mean result from specific $T_{1/2}({}^{243}\text{Am})$ determination and from measurements relative to $T_{1/2}({}^{241}\text{Am})$	
	Specific activity		7390	50		
1974Po17			7380	34		
1980Ag05	Relative activity	17.010 (95)	7359 *	42		Superseded by 2007Ag02
2007Ag02	Relative activity	17.022 (27)	7363.7 *	23		$\chi^2/n-1 = 0.2$, χ^2 crit = 3.3, int. uc. = 17 Weighted average.
	LWM		7367	17		
Recommended value			7367	23	Some results depend on the $T_{1/2}({}^{241}\text{Am})$ and then are not independent so the uncertainty is the minimum value from input.	

* Relative to $T_{1/2}({}^{241}\text{Am}) = 432.6$ (6) a (Chechev in 2004BeZQ).

3 Atomic Data

X-ray and Auger (relative and absolute) electron emission probabilities given in Sections 3 and 5, respectively, have been calculated by means of the computer code EMISSION (version 3,01, Nov. 3, 1999) [1], which makes use of the atomic data from 1996Sc06, from reference [2], and from the evaluated γ -ray data given in Sections 2.1 and 4.2. In addition, internal conversion electron energies and absolute emission probabilities for the strongest lines are presented in Section 5. Electron energies have been calculated using electron binding energies from 1977La19, and γ -ray energies from Section 2.1. Absolute electron emission probabilities have been calculated using absolute γ -ray emission probabilities given in Section 4.2 and conversion coefficients from Section 2.1.

4 Alpha Particles

α -Particle Energies

Most of the recommended α -particle energies in this evaluation are weighted averages (*Limited Relative Statistical Weight* method, LWM) of values from 1964Ba26 and 1968Ba25 (magnetic spectrograph), and from 1996Sa24 and 2002Da21 (semiconductor detectors). Values reported by 2002Da21 are from the analysis of an α -particle spectrum measured by 1992Ga01.

A. Rytz (1991Ry01) has critically evaluated the α -particle groups at 5233, 5275, and 5379 keV. His energies, also recommended in this evaluation, are virtually the same as the weighted average energies given in Table 2. This table shows the results of various measurements as well as the values recommended in this evaluation.

Table 2. ²⁴³Am Alpha-Particle Energies

1964Ba26	1968Ba25	1996Sa24	2002Da21 ^{&}	W. Average	Rec. Values
4695 (3)			[4697]#		4695 (3)
4919 (3)					4919 (3)
4930 (3)			[4936]#		4930 (3)
4946 (3)			[4951]#		4946 (3)
4997 (3)			[5001]#		4997 (3)
5008 (3)		5002(5)	5012 (5)	5008 (3)	5008 (3)
5029 (3)		5030 (5)		5029 (3)	5029 (3)
5035 (3)			5037 (5)	5035 (3)	5035 (3)
5088 (3)		5083 (5)	5091 (5)	5088 (5)	5088 (5)
5113 (1)		5109 (5)	5113 (5)	5113 (1)	5113 (1)
5181 (1)		5177 (5)	5178 (5)	5181 (1)	5181 (1)
5234 (1)	5232.9 (10)	5232 (5)	5233 (5)	5233.4 (10)	5233.3 (10)*
5276 (1)	5274.8 (10)	5275 (5)	5275 (5)	5275.3 (10)	5275.3 (10)*
5321 (1)		5319 (5)	5318 (5)	5321 (1)	5321 (1)
5350 (1)		5350 (5)	5349 (5)	5350 (1)	5349.4 (23)*

2002Da21 did not measure the alpha spectrum of ²⁴³Am. The alpha spectrum used was from 1992Ga01, who had not identified these very weak peaks. 2002Da21 reported for these peaks, intensities ranging from 2 to 13 times those given by 1964Ba26. Evaluators have interpreted this discrepancy as possibly caused by *spurious peaks* produced in the spectral peak-shape analysis of 2002Da21. Thus, they did not use these α -particle energies in the averaging process.

* From 1991Ry01.

& Rounded values. Uncertainties assigned by evaluators are typical values for spectra measured with semiconductor detectors.

α -Particle Emission Intensities

Table 3 shows the emission intensities measured by various authors. The uncertainties given by all of them (except one, 1996Sa24) are statistical values deduced from spectral peak-shape analysis. Such uncertainties do not include a constraint imposed by normalizing the sum of the emission probabilities to 100,

that is, to absolute emission intensities ($p_i(\%)$) per 100 α -particle disintegrations of the parent nuclide. The following formula (1988Br07) may be used to convert uncertainties (dI_i) in relative α -particle emission intensities (I_i) to values in the absolute emission intensities ($dp_i(\%)$):

$$dp_i(\%)/p_i(\%) = [(dI_i/I_i)^2 (1 - 2 I_i/\Sigma I_k) + \Sigma dI_k^2/(\Sigma I_k)^2]^{1/2} \quad (1)$$

The uncertainties given by 1996Sa24 (see Table 3) are those in the absolute α -emission intensities ($dp_i(\%)$), whereas the other authors give uncertainties only in the relative α -emission intensities (dI_i). This situation significantly affects only the two most intense α -particle groups for which 1996Sa24 give the same uncertainty of 0.03.

The energies and absolute emission intensities recommended in this evaluation are given in Section 2.2. The following description shows the procedure used here for determining these recommended absolute emission intensities:

1. Changing the uncertainty in the 5275-keV α -particle group before averaging from its absolute value of $dp(\%) = 0.03$ (1996Sa24) to a relative value (estimated by evaluators) of $dI = 0.06$.
2. Averaging (i.e., weighted averages, LWM) the relative emission intensities given by various authors (1955St98, 1956Hu96, 1964Ba26, 1966Le13, 1992Ga01, 1996Sa24, 2002Da21) and depicted in Table 3. Relative emission probabilities from 1998Ya17 (also shown in Table 3) are in disagreement with those from these authors, thus significantly increasing χ^2/ν for most averages. Their uncertainties include a “non-statistical component.” Unfortunately, 1998Ya17 give neither their values for these components nor the criteria used for estimating them. Therefore, data from 1998Ya17 have not been used for averaging.
3. Converting uncertainties in the recommended emission intensities (Table 3, column 9) to uncertainties in the absolute α -particle emission intensities by using formula (1). It should be noticed that only the uncertainties in the two most intense α -particle groups have been affected by this procedure.

Table 3. ²⁴³Am Alpha particle emission intensities

E α (keV)	1955St98	1956Hu96	1964Ba26	1966Le13	1992Ga01	1998Ya17	1996Sa24 ^{##}	2002Da21 ^{\$}	I α (avg) ^{&&}	χ^2/ν	Rec. I α ^{&&&}
4695			0.000 6	0.001 7 (5) ^{***}				0.003 8 (4) ^{^^}			0.001 7 (5)
4919			0.000 085								0.000 085
4930			0.000 18					0.002 6 (3) ^{^^}			0.000 18
4946			0.000 34					0.002 8 (3) ^{^^}			0.000 34
4997			0.001 6 [#]		0.001 6 (5) [#]		0.002 0 (4) [#]	0.003 1 (4) ^{^^}	0.001 8 (3)	0.39	0.001 8 (4) [#]
5008								0.005 2 (4) ^{^^}			
5029			0.002 2 [^]		0.003 3 (5) [^]		0.004 4 (5) [^]	0.008 2 (5) ^{^^}	0.003 9 (4)	2.4	0.003 9 (6) [^]
5035											
5088			0.004		0.005 6 (7)		0.005 5 (6)	0.011 2 (6) ^{^^}	0.005 5 (5)	0.01	0.005 5 (6)
5113			0.005 4		0.010 (1)		0.010 1 (10)	0.019 (1) ^{^^}	0.010 0 (7)	0	0.010 (1)
5181	1.1 (3) ^{&}	1.3 (2)	1.1		1.36 (1)	0.98 (2)	1.388 (8)	1.391 (7)	1.383 (5)	2.0	1.383 (7)
5233	11.5 (3) [*]	11.5 (3)	10.6 (2) ^{**}		11.46 (3)	11.04 (7)	11.37 (3)	11.52 (2)	11.46 (6)	7.1	11.46 (5) ^{\$\$}
5275	87.1 (4) [*]	86.9 (4)	87.9 (3) ^{**}		86.74 (6)	87.42 (8)	86.79 (3)	86.60 (7)	86.74 (4)	4.1	86.74 (5) ^{\$\$}
5321	0.16	0.16	0.12		0.190 (7)	0.270 (6)	0.194 (3)	0.190 (3)	0.192 (2)	0.48	0.192 (3)
5349	0.17	0.17	0.16		0.230 (7)	0.298 (8)	0.243 (3)	0.240 (3)	0.240 (2)	1.5	0.240 (3)

^{\$} 2002Da21 analyzed an α spectrum of 1992Ga01.

[&] Uncertainty assumed by evaluator.

^{*} From 1955St98, quoted in 1991Ry01; uncertainties are from 1991Ry01.

[#] 4997 α + 5008 α

[^] 5029 α + 5035 α

^{**} From 1964Ba26, quoted in 1991Ry01; uncertainties are from 1991Ry01.

^{##} Uncertainties include the effect of covariances when normalizing $\Sigma I\alpha = 100$.

^{^^} α -particle intensities are at least about twice those found by other authors, which suggest a possible systematic bias in the analysis of the spectrum. These values were not used for averaging.

^{***} Agrees well with I α =0.001 48 (3) % from γ -ray transition intensity balance.

^{&&} Weighted average using the Limitation of Relative Statistical Weights method. Data from 1998Ya17 have not been included. See text.

^{\$\$} Normalization of I α to $\Sigma I\alpha=100$ requires same values for these uncertainties. See text.

^{&&&} Uncertainty is always greater than or equal to the smallest uncertainty in any of the experimental values used in the calculation

5 Gamma Rays

Energies

The recommended γ -ray energies given in Sections 2.1 and 4.2 are weighted averages (LWM) of values given in 1982Ah04 and 1975Pa04, complemented with values from 1996Sa23, 1969En02, and 1968Va09 (See table 4).

E_γ (keV)	E_γ (keV)	E_γ (keV)	E_γ (keV)	E_γ (keV)	E_γ (keV)		E_γ (keV)
1996Sa23	1982Ah04	1975Pa04	1969En02	1968Va09	W. Avg.*	χ^2/ν	Rec. E_γ
31.13	31.14 (3)	31.10 (15)		31.2	31.14 (3)	0.068	31.14 (3)
		43.1	43.1				43.1#
43.53	43.53 (2)	43.53 (15)		43.6	43.53 (2)		43.53 (2)
50.6				50.6			50.6&\$
55.18			55.4	55.4			55.18&
74.66	74.66 (2)	74.67 (15)	74.7	74.8	74.66 (2)	0.004	74.66 (2)
86.71	86.71 (2)	86.79 (15)	86.7	86.7	86.71 (2)	0.27	86.71 (2)
98.5			98.5				98.5^
117.84		117.60 (15)	117.8	117.8			117.60 (15)#
141.89	141.89 (3)	142.18 (15)	142	142	141.90 (3)	3.6	141.90 (6)
169				169			169\$
195				195			195\$

* Weighted average of values in 1982Ah04 and 1975Pa04.

From 1975Pa04

& From 1996Sa23

\$ From 1968Va09

^ From 1969En02

The recommended absolute γ -ray emission (photons) and transition (photons + electrons) intensities given in Sections 4.2 and 2.2, respectively, are weighted averages (LWM) of values in 1996Sa23, 1996Wo05, 1984Va41, 1982Ah04, 1979Po20, 1977St35, 1975Pa04, 1972Ah02, 1969Al14 and 1960As02 (see Table 5).

The conversion coefficients used for deducing absolute transition probabilities (see section 2.2) are theoretical values interpolated from the Band's tables (2002Ba85) by using the computer code BrIcc (2008Ki07) with the so called "Frozen orbital" approximation.

The M1/E2 mixing ratio for $\gamma_{3,0}$ (31.1 keV) $\delta = 0.17$ was deduced from probability balance in ²⁴³Am α -decay and in ²³⁹U β^- -decay

The M1/E2 mixing ratios for $\gamma_{4,3}$ (43.1 keV) $\delta = 0.38$ (4) and $\gamma_{6,4}$ (55.2 keV) $\delta = 0.75$ (10) have been taken from Engelkemeir (1969En02).

The remaining M1/E2 mixing ratios are from 2003Br12 based on measurements of 1957Ho07, 1964Bl11, 1969En02.

Table 5. ²⁴³Am γ -ray Absolute Emission Probabilities

E_γ (keV)	I_γ	$I_\gamma^{\textcircled{a}}$	I_γ	I_γ	I_γ	I_γ	I_γ	I_γ	I_γ	I_γ	I_γ	W. Avg.	χ^2/ν	I_γ^{a}
Rec. Value	1960As02	1968Va09	1969Al14	1972Ah02	1975Pa04	1977St35	1979Po20	1982Ah04	1984Va41	1996Wo05	1996Sa23			Rec. Value
31.14 (3)								0.069 (7)			0.0477 (13)	0.0484 (13)	9	0.048 (4)
43.1		0.03												0.065 [^]
43.53 (2)	4 (1)	5.3	5 (1)	5.5 (3)			5.3 (12)	6.20 (30)	6.04 (13)	5.93 (10)	5.72 (17)	5.89 (7)	1.4	5.89 (10)
50.6		0.0027									0.0062 (10)			0.0062 (10)#
55.18		0.0094									0.0168 (11)			0.0168 (11)#
74.66 (2)	69 (3)	61		66 (3)		59.1 (40)	60 (4)	68.0 (20)	68.5 (15)	66.7 (12)	68.4 (13)	67.2 (7)	1.4	67.2 (12)
86.71 (2)		0.37						0.340 (15)	0.35 (1)	0.342 (15)	0.344 (9)	0.346 (6)	0.2	0.346 (9)
98.5											0.0151 (21)			0.0151 (21)#
117.60 (15)		0.75			0.56 (8)						0.57 (5)			0.57 (5)#
141.90 (6)		0.13						0.128 (6)	0.13 (1)	0.117 (5)	0.1068 (26)	0.115 (2)	3.8	0.115 (8)
169		0.0012												0.0012 [^]
195		0.00085												0.00085 [^]

a Recommended absolute emission probabilities are weighted averages (LWM) of experimental values, unless otherwise noted. Uncertainty is always greater than or equal to the smallest uncertainty in any of the experimental values used in the calculation.

* From Table 4

From 1996Sa23

& From 1968Va09

[^] Estimated by 2003Br12 from $\alpha_M(43.1\gamma, \text{exp.}) = 31$, $I_M(\text{ce}, 43.1\gamma) / I_\gamma(117) = 3.56$ (1969En02), and $I_\gamma(117) = 0.57$.

[@] Uncertainties are at least 10 %.

6 References

- [1] E. Schönfeld, H. Janßen. Applied Radiation Isotopes **52**, 595 (2000). (X-ray and Auger electron emission probabilities).
- [2] E. Schönfeld, G. Rodloff. Report PTB-6.11-98-1, Braunschweig, October 1998. (Auger electron energies).
- 1955St98 F. Stephens, J. Hummel, F. Asaro. Phys. Rev. **98**, 261 (1955) (²⁴³Am α -particle emission probabilities).
- 1956Hu96 J.P. Hummel. Thesis. Univ. of California (1956): UCRL-3456 (1956) (²⁴³Am α -particle emission probabilities).
- 1959Ba22 R.F. Barnes, D.J. Henderson, A.L. Harkness, H. Diamond. J. Inorg. Nuclear Chem. **9**, 105 (1959) (²⁴³Am half-life).
- 1960As02 F. Asaro, F.S. Stephens, J.M. Hollander, I. Perlman. Phys. Rev. **117**, 492 (1960) (²⁴³Am γ -ray emission probabilities).
- 1960Be10 A.B. Beadle, D.F. Dance, K.M. Glover, J. Milsted. J. Inorg. Nuclear Chem. **12**, 359 (1960) (²⁴³Am half-life).
- 1964Ba26 S.A. Baranov, V.M. Kulakov, V.M. Shatinsky. Nucl. Phys. **56**, 252 (1964) (²⁴³Am α -particle energies and emission probabilities).
- 1966Gv01 B.A. Gvozdev, B.B. Zakhvataev, V.I. Kuznetsov, V.P. Perelygin, S.V. Pirozkov, E.G. Chudinov, I.K. Shvetsov. Radiokhimiya **8**, 493 (1966); Sov. Radiochem. **8**, 459 (1966).
- 1966Le13 C.M. Lederer, J.K. Poggenburg, F. Asaro, J.O. Rasmussen, I. Perlman. Nucl. Phys. **84**, 481 (1966) (²⁴³Am α -particle emission probabilities).
- 1968Ba25 S.A. Baranov, V.M. Kulakov, V.M. Shatinskii. Yadern. **7**, 727 (1968); Sov. J. Nucl. Phys. **7**, 442 (1968) (²⁴³Am α -particle energies).
- 1968Be22 G. Berzins, M.E. Bunker, J.W. Starnier. Nucl. Phys. **A114**, 512 (1968) (²⁴³Am half-life).
- 1968Va09 J.R. Van Hise, D. Engelkemeir. Phys. Rev. **171**, 1325 (1968) (²⁴³Am γ -ray energies and emission probabilities).
- 1969Al14 B.M. Aleksandrov, O.I. Grigorev, N.S. Shimanskaya. Yadern. Fiz. **10**, 14 (1969); Soviet J. Nucl. Phys. **10**, 8 (1970) (²⁴³Am γ -ray emission probabilities).
- 1969En02 D. Engelkemeir. Phys. Rev. **181**, 1675 (1969) (²⁴³Am γ -ray energies, δ).
- 1972Ah02 I. Ahmad, M. Wahlgren. Nucl. Instrum. Methods **99**, 333 (1972) (²⁴³Am γ -ray emission probabilities).
- 1974Po17 V.G. Polyukov, G.A. Timofeev, P.A. Privalova, V.Y. Gabeskiriya, A.P. Chetverikov. At. Energ. **37**, 357 (1974); Sov. At. Energ. **37**, 1103 (1975) (²⁴³Am half-life).
- 1975Pa04 J.C. Pate, K.R. Baker, R.W. Fink, D.A. McClure, N.S. Kendrick, Jr. Z. Phys. **A272**, 169 (1975) (²⁴³Am γ -ray energies and emission probabilities).
- 1977La19 F.P. Larkins. At. Data Nucl. Data Tables **20**, 311 (1977) (Atomic electron binding energies).
- 1977St35 D.I. Starozhukov, Y.S. Popov, P.A. Privalova. At. Energ. **42**, 319 (1977); Sov. At. Energy **42**, 355 (1977) (²⁴³Am γ -ray emission probabilities).
- 1978Ro22 F. Rosel, H.M. Fries, K. Alder, H.C. Pauli. At. Data Nucl. Data Tables **21**, 92 (1978) (γ -ray theoretical internal conversion coefficients).
- 1979Po20 Y.S. Popov, D.I. Starozhukov, V.B. Mishenev, P.A. Privalova, A.I. Mishchenko. At. Energ. **46**, 111 (1979); Sov. At. Energy **46**, 123 (1979) (²⁴³Am γ -ray emission probabilities).
- 1980Ag05 S.K. Aggarwal, A.R. Parab, H.C. Jain. Phys. Rev. **C22**, 767 (1980) (²⁴³Am half-life).
- 1982Ah04 I. Ahmad. Nucl. Instrum. Methods **193**, 9 (1982) (²⁴³Am γ -ray energies and emission probabilities).
- 1984Va41 R. Vaninbrouckx, G. Bortels, B. Denecke. Int. J. Appl. Radiat. Isotop. **35**, 1081 (1984) (²⁴³Am γ -ray emission probabilities).
- 1985ZiZY W.L. Zijp, Report ECN FYS/RASA-85/19 (1985) (Discrepant Data. Limited Relative Statistical Weight Method).
- 1986LoZT A. Lorenz. IAEA Tech. Rept. Ser., No. **261** (1986) (²⁴³Am half-life: ²⁴³Am recommended half-life).
- 1988Br07 E. Browne. Nucl. Instrum. Methods Phys. Res. **A265**, 541 (1988) (Uncertainties in α -particle

- emission probabilities).
- 1991BaZS W. Bambynek, T. Barta, R. Jedlovsky, P. Christmas, N. Coursol, K. Debertin, R.G. Helmer, A.L. Nichols, F.J. Schima, Y. Yoshizawa. IAEA-TECDOC-619 (1991) (^{241}Am half-life as standard: ^{243}Am recommended half-life).
- 1991Ry01 A. Rytz. At. Data Nucl. Data Tables **47**, 205 (1991) (^{243}Am α -particle energies).
- 1992Ak06 Y.A. Akovali. Nucl. Data Sheets **66**, 897 (1992) (^{243}Am recommended half-life).
- 1992Ga01 E. Garcia-Torano, M.L. Acena, G. Bortels, D. Mouchel. Nucl. Instrum. Methods Phys. Res. **A312**, 317 (1992) (^{243}Am α -particle energies and emission probabilities).
- 1996Sa23 D. Sardari, T.D. Mac Mahon, S.P. Holloway. Nucl. Instrum. Methods Phys. Res. **A369**, 486 (1996) (^{243}Am γ -ray energies and emission probabilities).
- 1996Sa24 A.M. Sanchez, P.R. Montero, F.V. Tome. Nucl. Instrum. Methods Phys. Res. **A369**, 593 (1996) (^{243}Am α -particle energies and emission probabilities).
- 1996Sc06 E. Schönfeld, H. Janßen. Nucl. Instrum. Methods Phys. Res. **A369**, 527 (1996) (Atomic data, X-rays, Auger electrons).
- 1996Wo05 S.A. Woods, D.H. Woods, M.J. Woods, S.M. Jerome, M. Burke, N.E. Bowles, S.E.M. Lucas, C. Paton Walsh. Nucl. Instrum. Methods Phys. Res. **A369**, 472 (1996) (^{243}Am γ -ray emission probabilities).
- 1998Ya17 Jichun Yang, Jianzhong Ni. Nucl. Instrum. Methods Phys. Res. **A413**, 239 (1998). (^{243}Am α -particle emission probabilities).
- 1998Ak04 Y.A. Akovali. Nucl. Data Sheets **84**, 1 (1998) (Alpha decay. Radius parameter of even-even nuclei).
- 1998Ya17 Jichun Yang, Jianzhong Ni. Nucl. Instrum. Methods Phys. Res. **A413**, 239 (1998). (^{243}Am α -particle emission probabilities).
- 2002Ba85 I.M. Band, M.B. Trzhaskovskaya. At. Data. Nucl. Data Tables **88**, 1 (2002). Theoretical ICC
- 2002Da21 F. Dayras. Nucl. Instrum. Methods Phys. Res. **A490**, 492 (2002) (^{243}Am α -particle energies and emission probabilities).
- 2002Sa53 R. Sampathkumar, P.C. Kalsi, A. Ramaswami. J. Radioanal. Nucl. Chem. **253**, 523 (2002) (^{243}Am spontaneous fission branching).
- 2003Au03 G. Audi, A.H. Wapstra, C. Thibault. Nucl. Phys. **A729**, 337 (2003) (2003 Atomic Mass Adjustment).
- 2003Br12 E. Browne. Nucl. Data Sheets **98**, 665 (2003) (Evaluated data (ENSDF) for nuclei with A=239).
- 2004BeZQ M.M. Bé, V. Chisté, C. Dulieu, E. Browne, V. Chechev, N. Kuzmenko, R. Helmer, A. Nichols, E. Schönfeld, R. Dersch. *Table of Radionuclides (Vol.2 - A = 151 to 242)* Monographie BIPM-5, ISBN 92-822-2207-1. (^{241}Am half-life)
- 2007Ag02 S.K. Aggarwal *et al.* Nucl. Instrum. Methods Phys. Res. **A571**, 663 (2007) (^{243}Am half-life)
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, and C.W. Nestor, Jr., Nucl. Instrum. Methods Phys. Res. **A589**, 202 (2008) (Theoretical ICC)

**²⁴³Cm -Comments on evaluation of decay data
by V.P. Chechev**

This evaluation was done in October 2010 with a literature cut-off by the same date.

1. DECAY SCHEME

The structure of the adopted scheme of ²⁴³Cm decay is based on the evaluations by E. Browne (2003Br12) and Y.A. Akovali (2004Ak21). The decay scheme includes two decay modes: α decay to ²³⁹Pu (99.71 (3) %) and electron capture decay (EC) to ²⁴³Am (0.29 (3) %).

EC branching was obtained from the EC decay half-life of $1.0 (1) \times 10^4$ a, as determined in 1958Ch38 from a ratio of ²⁴³Am and ²⁴³Cm α activities (correction for the half-lives of ²⁴³Am and ²⁴³Cm adopted by DDEP does not change this value), and from the total ²⁴³Cm half-life of 29.1 (1) a (correction for the recommended below value of the ²⁴³Cm half-life does not change the EC branching). The EC decay occurs 100 % to the ground state 5/2- of ²⁴³Am (1958Ch38, 2004Ak21).

In the ²⁴³Cm α decay to ²³⁹Pu the intense population takes place only to levels in ²³⁹Pu with energy less than 400 keV (8 excited levels and ground state) and in this part the decay scheme is well defined. Nevertheless, a number of gamma-ray transitions with energy less than 200 keV were not observed in the ²⁴³Cm α decay, such as 7.86-keV, 49.4-keV, 61.5-keV, 67.8-keV, 88.1-keV, 102.0-keV, 106.1-keV, 106.5-keV, and 166.4-keV. These transitions, included in the ²⁴³Cm decay scheme, have been derived from measurements of ²³⁹Np and ²³⁹Am decays.

For levels with higher energy, the decay scheme has not been completed since many gamma-ray transitions have not been observed yet. In addition, many levels were placed in the ²⁴³Cm α decay scheme based only on questionable weak α transitions. Therefore, further measurements are needed to determine the γ transitions and ²⁴³Cm α decay scheme with greater precision.

2. NUCLEAR DATA

$Q(\alpha)$ and $Q(\text{EC})$ values are from Audi *et al.* (2003Au03).

The recommended ²⁴³Cm half-life is based on the experimental results given in Table 1.

Table 1. Experimental values of ²⁴³Cm half-life (in years)

Reference	Author(s)	Original value	Re-estimated value	Comments
1950Th52	Thompson et al.	Roughly 100		Not used
1953As40	Asaro	35		Not used
1958Ch38	Choppin and Thompson	29.0 (8)	28.5 (2) ^a	Relative activity to ²⁴⁴ Cm
1986Ti03	Timofeev et al.	29.20 (12)	29.20 (13) ^a	Relative activity to ²⁴⁴ Cm

^a Re-estimated by the evaluator to the ²⁴⁴Cm half-life of 18.11 (3) a

The weighted average of 29.0 for this two re-estimated discrepant experimental data set is dominated by the accurate value of 1986Ti03. The LWEIGHT computer program, which uses a *Limitation of Relative Statistical Weights* (LRSW method), has expanded the 1986Ti03 uncertainty from 0.13 to 0.20 and used a weighted mean (28.85) and an external uncertainty (0.35) for the average of the adjusted data set ($\chi^2/\nu = 6.13$).

The recommended value of the ²⁴³Cm half-life is **28.9 (4) years**.

The value of $5.5 (9) \times 10^{11}$ years was adopted for ²⁴³Cm spontaneous fission (SF) half-life from the measurement of 1987Po19. SF branching of $5.3 (9) \times 10^{-9}$ % has been obtained using the adopted values of SF half-life and total half-life of 28.9 (4) a.

2.1. Alpha Transitions

The energies of the alpha transitions have been obtained from the $Q(\alpha)$ value and ²³⁹Pu level energies given in Table 2 from 2003Br12 where they were deduced from a least-squares fit to gamma ray energies.

Table 2. ²³⁹Pu levels populated in ²⁴³Cm α -decay

Level	Energy (keV)	Spin and parity	Half-life	Probability of α -transition (%)
0	0	1/2+	24100 (11) a	1.3 (2)
1	7.861 (2)	3/2+	36 (3) ps	4.4 (2)
2	57.275 (2)	5/2+	101 (5) ps	1.05 (12)
3	75.705 (3)	7/2+	83 (8) ps	5.7 (2)
4	163.76 (3)	9/2+	73 (4) ps	0.1
5	192.8 (10)	11/2+		0.7
6	285.460 (2)	5/2+	1.12 (5) ns	73.4 (4)
7	330.124 (4)	7/2+		11.3 (2)
8	387.42 (2)	9/2+		1.6 (1)
9	391.584 (3)	7/2-	193 (4) ns	0.2
10	427 (3) ?			0.03
11	434 (3)	(9/2-)		0.14
12	451 (5) ?			0.06
13	462 (3)	(11/2+)		0.03
14	469.8 (4)	(1/2-)		≤ 0.01
15	481 (3) ?			0.01
16	487 (3)	(11/2-)		0.02
17	492.1 (3)	3/2-		0.009
18	499 (3)			0.007
19	505.6 (2)	(5/2-)		0.007
20	538 (3)			0.002
21	543 (3) ?			0.006
22	556.2 (5)	(7/2-)		0.002
23	746 (3)			0.003
24	756 (3)			0.003
25	763 (3)			0.001
26	813 (3)			0.0015
27	850 (15)			0.00039

The probabilities of the most intense α -transitions ($I_\alpha > 1$ %) have been obtained by averaging experimental values from the spectrometric measurements (Table 3). Probabilities of the rest of alpha

transitions have been adopted from the magnetic spectrometer measurements of 1966Ba07. The probability of the $\alpha_{0,9}$ transition (0.2 %) from 1966Ba07 disagrees with the value deduced from the gamma-ray transition intensity balance (> 0.4 %).

The α -decay hindrance factors have been calculated using the ALPHAD computer program from the ENSDF evaluation package with $r_0(^{239}\text{Pu}) = 1.4996$ fm (see 2003Br12).

Table 3. Experimental (per 100 α decays) and recommended probabilities (per 100 decays) of the most intense α -transitions ($I_\alpha > 1$ %) observed in ²⁴³Cm α decay

	α -part. energy	1957 As83	1963 Dz07	1966Ba07	1973 Ah04	2009 KoZV	Evaluated from α measurement results (per 100 α decays)	Deduced from $P(\gamma+ce)$ balance	Recommended (per 100 (α +EC) decays)
$\alpha_{0,0}$	6066	1.0 (2) ^a		1.5 (2) ^a			1.3 (2) ^a		1.3 (2)
$\alpha_{0,1}$	6058	5		4.7 (3) ^a	4.3 (2)	4.5 (3)	4.4 (2) ^a		4.4 (2)
$\alpha_{0,2}$	6010		0.95		1.05 (5)	1.1 (2)	1.05 (12) ^b		1.05 (12)
$\alpha_{0,3}$	5992	6.0 (2) ^a	5.4 (2)	5.63 (20) ^a	5.6 (2)	5.8 (2)	5.7 (2) ^a		5.7 (2)
$\alpha_{0,6}$	5785	73 (4) ^a	73 (4)	73.54 (40) ^a	74.2 (8)	72.9 (12)	73.6 (4) ^a	73.8 (26)	73.4 (4)
$\alpha_{0,7}$	5742	11.5 (6) ^a	12.3 (6)	10.65 (60) ^a	11.1 (2)	11.6 (4)	11.3 (2) ^a	11	11.3 (2)
$\alpha_{0,8}$	5686		1.7	1.6	1.52 (5)	1.8 (1)	1.6 (1) ^a		1.6 (1)

^a Weighted average, uncertainty is the smallest experimental one.

^b Weighted average, uncertainty is external.

2.2. Gamma Transitions and Internal Conversion Coefficients

The recommended energies of the gamma-ray transitions are virtually the same as those of the gamma-ray energies because the nuclear recoil is negligible for ²³⁹Pu.

The intensities of gamma-ray transitions $\gamma_{1,0}$ (7.86 keV), $\gamma_{3,2}$ (18.43 keV), $\gamma_{2,0}$ (57.27 keV), $\gamma_{3,1}$ (67.84 keV), $\gamma_{4,2}$ (106.47 keV), and $\gamma_{5,3}$ (117.1 keV) have been deduced from intensity balances at the ²³⁹Pu levels “0” (0 keV), “3” (75.7 keV), “2” (57.3 keV), “1” (7.86 keV), “4” (163.8 keV), and “5” (192.8 keV), respectively.

The rest of gamma-ray transition intensities (P_γ) have been deduced from their evaluated gamma-ray emission probabilities and total internal conversion coefficients (ICCs).

ICCs have been interpolated using the BrIcc computer program, version v2.2a, data set BrIccFO (2008Ki07). Multipolarities of the gamma-ray transitions and E2/M1 mixing ratios (δ) are based on the measurements of conversion electrons (ce) in ²³⁹Am, ²³⁹Np and ²⁴³Cm decays and $\gamma(\theta)$ measurements by 1972Kr07, 1990Si12 from polarized ²³⁹Np. The multipolarities have been taken from 2003Br12. The δ values have been adopted mainly from 2003Br12, except as noted below.

The δ values for $\gamma_{6,3}$ (209.75 keV), $\gamma_{6,2}$ (228.18 keV), and $\gamma_{6,1}$ (277.60 keV) have been taken from $\gamma(\theta)$ measurements of 1972Kr07, 1990Si12. Asymmetric uncertainties of 1972Kr07, 1990Si12 were symmetrized by transformation to equivalent symmetric normal distribution using a method described in 2003Au03 (p. 21): for $\gamma_{6,3}$ $\delta = -0.004 (+1 -24) \rightarrow \delta = -0.019 (15)$ and for $\gamma_{6,2}$ $\delta = +0.004 (+9 -1) \rightarrow \delta = +0.009 (6)$. The value of $\delta = 0.24 (4)$ has been adopted for $\gamma_{7,6}$ (44.66 keV) to provide more accurate probability balances at the levels “6” (285.46 keV) and “7” (330.12 keV).

ICCs for the E1(+M2) anomalously converted 106.1-keV gamma-ray transition are from conversion electron measurements of 1959Ew90.

3. ATOMIC DATA

The fluorescence yields, X-ray energies and relative probabilities, and Auger electrons energies and relative probabilities are from the SAISINUC software.

4. ALPHA EMISSIONS

The recommended energies of alpha particles have been deduced from the energies of alpha transitions taking into account the recoil energies for ²³⁹Pu.

In Table 4 the deduced (recommended) values of α -particle energies for the four intense α -transitions are compared with the experimental results from spectrometric measurements of 1957As83, 1963Dz07 and 1966Ba07. Other measurement results can be found in 1953As14, 1957As70, 1962Iv01, 1963Le17, 1970By01, 1971Bb10, 1976BaZZ, and 1977VaZW. Most of them have lesser accuracy in comparison with the recommended values.

Table 4. Experimental and recommended values of α -particle energies (keV) in the decay of ²⁴³Cm

	Measured ^a			Evaluated in 1991Ry01	Recommended
	1957As83	1963Dz07	1966Ba07		
$\alpha_{0,0}$	6061 (3) ^b		6067 (2) ^b	6066.2 (17)	6067.2 (10)
$\alpha_{0,3}$	5987 (3) ^b	5992 (3)	5993 (2) ^b	5991.8 (15)	5992.7 (10)
$\alpha_{0,6}$	5780 (3) ^b	5785 (3)	5784.5 (10)	5785.2 (9)	5786.4 (10)
$\alpha_{0,7}$	5736 (3) ^b	5740 (3)	5741.6 (10)	5742.1 (9)	5742.5 (10)

^a Original values have been adjusted for changes in calibration energies as suggested in 1991Ry01.

^b Uncertainty deduced or guessed by A. Rytz.

5. ELECTRON EMISSIONS

The energies of the conversion electrons have been obtained from the gamma-ray transition energies and the atomic electron binding energies.

The emission probabilities of the conversion electrons were deduced using the evaluated P_γ and ICC values. The total absolute emission probabilities of K and L Auger electrons were calculated using the EMISSION computer program.

6. PHOTON EMISSIONS

6.1 X - Ray emissions

The absolute emission probabilities of Pu KX- and LX-rays were calculated using the EMISSION computer program (Table 5).

Table 5. Measured (1972Ah02) and calculated absolute Pu KX-ray emission probabilities (%).

	1972Ah02	Calculated
K α_2 (Pu)	13.5 (5)	13.34 (28)
K α_1 (Pu)	20.8 (8)	21.1 (5)
K β'_1 (Pu)	7.6 (3)	7.75 (21)
K β'_2 (Pu)	2.6 (1)	2.69 (8)

The good agreement between measured and calculated KX-ray emission probabilities supports the recommended γ -ray emission probabilities and assigned multiplicities.

6.2. Gamma emissions

6.2.1. Gamma ray energies

The gamma ray energies have been taken mainly from 2003Br12. They are based on measurements of γ -ray transitions observed in ^{239}Np and ^{239}Am decays by 1959Ew90, 1964Ba31, 1965Ma17, 1972Po04, 1979Bo30, and 1982Ah04. The energies of gamma rays $\gamma_{9,8}$ (4.16 keV), $\gamma_{1,0}$ (7.86 keV), $\gamma_{3,2}$ (18.43 keV), $\gamma_{2,1}$ (49.41 keV), $\gamma_{8,7}$ (57.30 keV), $\gamma_{5,3}$ (117.1 keV), and $\gamma_{22,2}$ (498.9 keV) have been deduced directly from the adopted ^{239}Pu level energies.

Earlier measurement results can be found in 1953As14, 1955Sc08, and 1956Ne17.

The gamma rays reported in 1963Le17 only, have not been placed in the decay scheme.

6.2.2. Gamma ray emission probabilities

The absolute gamma ray emission probabilities (P_γ) were adopted from the experimental values given by 1972Ah02 multiplied by 0.9971 (3), except when noted below.

P_γ of gamma rays $\gamma_{1,0}$ (7.86 keV), $\gamma_{3,2}$ (18.43 keV), $\gamma_{2,0}$ (57.27 keV), $\gamma_{3,1}$ (67.84 keV), $\gamma_{4,2}$ (106.47 keV), and $\gamma_{5,3}$ (117.1 keV) have been obtained from $P(\gamma+ce)$ values deduced from intensity balances (see section 2.2).

P_γ of gamma ray $\gamma_{7,6}$ (44.66 keV) has been deduced using the ratio $I_\gamma(44.66 \text{ keV}) / I_\gamma(254.4 \text{ keV}) = 0.13 (1) / 0.1091 (22)$ in ^{239}Np β^- decay (2005Tr08, 2008BeZV) and $P_{\gamma_{7,3}}$ (254.4 keV) = 0.11 (1) % measured in 1972Ah02.

P_γ of gamma ray $\gamma_{8,7}$ (57.30 keV) has been deduced from the total intensity of doublet $P_{\gamma_{2,0}}$ (57.27 keV) + $P_{\gamma_{8,7}}$ (57.30 keV) = 0.14 (1) % measured in 1972Ah02 and $P_{\gamma_{2,0}}$ (57.27 keV) = 0.06 % from transition probability balance at the 57 keV level.

P_γ of gamma ray $\gamma_{9,7}$ (61.46 keV) has been deduced using the ratio $I_\gamma(61.46 \text{ keV}) / I_\gamma(334.31 \text{ keV}) = 1.29 (2) / 2.05 (2)$ in ^{239}Np β^- decay (2005Tr08, 2008BeZV) and $P_{\gamma_{9,2}}$ (334.31 keV) = 0.024 (2) % measured in 1972Ah02.

P_γ of gamma ray $\gamma_{4,3}$ (88.06 keV) has been deduced using from the ratio $I_\gamma(88.06 \text{ keV}) / I_\gamma(106.47 \text{ keV}) = 0.006 (2) / 0.049 (2)$ in ²³⁹Np β^- decay (2005Tr08, 2008BeZV) and $P_{\gamma_{4,2}}(106.47 \text{ keV}) = 0.015 \%$.

P_γ of gamma ray $\gamma_{8,6}$ (101.96 keV) has been deduced using data from ²³⁹Am ϵ decay (1972Po04).

P_γ of gamma ray $\gamma_{9,6}$ (106.12 keV) has been deduced using data from ²³⁹Np β^- decay of the relative intensity $I_\gamma(106.12 \text{ keV}) / I_\gamma(334.31 \text{ keV}) = 25.32 (17) / 2.055 (13)$ as measured in 2005Tr08 and $P_{\gamma_{9,2}}(334.31 \text{ keV}) = 0.024 (2) \%$ measured in 1972Ah02.

P_γ of gamma ray $\gamma_{7,4}$ (166.39 keV) has been deduced using the ratio $I_\gamma(166.39 \text{ keV}) / I_\gamma(254.4 \text{ keV}) = 0.016 (7) / 0.1091 (22)$ in ²³⁹Np β^- decay (2005Tr08, 2008BeZV) and $P_{\gamma_{7,3}}(254.4 \text{ keV}) = 0.11 (1) \%$ measured in 1972Ah02.

7. CONSISTENCY OF RECOMMENDED DATA

The most accurate $Q(\alpha)$ value, $Q_\alpha(M)$, is taken from the atomic mass adjustment table of Audi et al. (2003Au03). Comparison of $Q_\alpha(\text{eff})$ (deduced as the sum of average energies per disintegration ($\sum E_i \times P_i$) for all emissions accompanying ²⁴³Cm α - decay) with the tabulated decay energy $Q_\alpha(M) \times 0.9971$ allows to check a consistency of the recommended decay-scheme parameters obtained in this evaluation.

Here E_i and P_i are the evaluated energies and emission probabilities of the i -th alpha particle, γ - ray, X-ray, etc. Consistency (percentage deviation) is determined by $\{[Q_\alpha(M) \times 0.9971 - Q(\text{eff})] / Q_\alpha(M) \times 0.9971\} \times 100$. "Percentage deviations above 5 % would be regarded as high and imply a poorly defined decay scheme; a value of less than 5 % indicates the construction of a reasonably consistent decay scheme" (quoted from the article by A.L. Nichols in Appl. Rad. Isotopes 55 (2001) 23-70).

For the current ²⁴³Cm decay data evaluation we have $Q_\alpha(M) \times 0.9971 = 6150.9 (10) \text{ keV}$ and $Q(\text{eff}) = 6171 (35) \text{ keV}$, i.e. consistency is better than 1.0 %.

8. REFERENCES

- 1950Th52 S.G. Thompson, A. Ghiorso and G.T. Seaborg, Phys. Rev. 80, 781 (1950)
(Half-life)
- 1953As14 F. Asaro, S.G. Thompson, I. Perlman, Phys. Rev. 92, 694 (1953)
(γ -ray transitions energies)
- 1953As40 F. Asaro, Thesis, Univ. California (1953); UCRL-2180 (1953)
(Half-life)
- 1955Sc08 J.F. Schooley, J. Rasmussen, UCRL-2932, p.63 (1955)
(α -particle and γ -ray energies)
- 1956Ne17 J.O. Newton, B. Rose, J. Milsted, Phil. Mag. 1, 981 (1956)
(γ -ray energies)
- 1957As70 F. Asaro, S.G. Thompson, F.S. Stephens, Jr., I. Perlman, Bull. Am. Phys. Soc. 2, No.8, 393 R1 (1957)
(α -particle energies)

- 1957As83 F. Asaro, S.G. Thompson, F.S. Stephens, Jr., I. Perlman, Priv. Comm. (1957). Quoted by 1964Hy02 (1964)
(α -particle energies and emission probabilities)
- 1958Ch38 G.R. Choppin and S.G. Thompson, J. Inorg. Nucl. Chem. 7, 197 (1958)
(Total and EC decay half-life, alpha-particle energies and emission probabilities)
- 1959Ew90 G.T. Ewan, J.S. Geiger, R.L. Graham, D.R. MacKenzie, Phys. Rev. 116, 950 (1959)
(Conversion electron intensities and ICC)
- 1962Iv01 R.B. Ivanov, A.S. Krivokhatskii, V.G. Nedovesov, Izvest. Akad. Nauk SSSR, Ser. Fiz. 26, 976 (1962); Columbia Tech. Transl. 26, 984 (1963)
(α -particle energies)
- 1963Dz07 B.S. Dzhelepov, R.B. Ivanov, V.G. Nedovesov, V.P. Chechev, Zh. Eksperim. i Teor. Fiz. 45, 1360 (1963); Soviet Phys. JETP 18, 937 (1964)
(Alpha-particle energies and emission probabilities)
- 1963Le17 C.M. Lederer, Thesis, Univ. California (1963); UCRL-11028 (1963)
(α -particle energies)
- 1964Ba31 I.A. Baranov, A.S. Krivokhatskii, A.N. Silantev, Izv. Akad. Nauk SSSR, Ser. Fiz. 28, 1255 (1964); Bull. Acad. Sci. USSR, Phys. Ser. 28, 1154 (1965).
(γ -ray energies)
- 1964Hy02 E.K. Hyde, I. Perlman, G.T. Seaborg, The Nuclear Properties of the Heavy Elements, Vol. II, Prentice-Hall, Inc., Englewood Cliffs, N.J. (1964)
(α -particle energies)
- 1965Ma17 B.P.K. Maier, Z. Physik 184, 143 (1965).
(γ -ray energies)
- 1966Ba07 S.A. Baranov, Y.F. Rodionov, V.M. Kulakov, V.M. Shatinskii, Yadern. Fiz. 4, 1108 (1966); Soviet J. Nucl. Phys. 4, 798 (1967)
(Alpha-particle energies and emission probabilities)
- 1970By01 J. Byrne, R.J.D. Beattie, S. Benda, I. Collingwood, J. Phys. B 3, 1166 (1970)
(α -particle energies)
- 1971Bb10 S.A. Baranov, V.M. Shatinskii, V.M. Kulakov, Yad. Fiz. 14, 1101 (1971); Sov. J. Nucl. Phys. 14, 614 (1972)
(α -particle energies)
- 1972Ah02 I. Ahmad, M. Wahlgren, Nucl. Instrum. Methods 99, 333 (1972)
(KX-ray and gamma ray emission probabilities)
- 1972Kr07 K.S. Krane, C.E. Olsen, W.A. Steyert, Phys. Rev. C5, 1671 (1972)
(E2/M1 mixing ratios)
- 1972Po04 F.T. Porter, I. Ahmad, M.S. Freedman, R.F. Barnes, R.K. Sjoblom, F. Wagner, Jr., P.R. Fields, Phys. Rev. C5, 1738 (1972)
(γ -ray energies)
- 1973Ah04 I. Ahmad, H. Diamond, J. Milsted, J. Lerner, R.K. Sjoblom, Nucl. Phys. A208, 287 (1973). Quoted in 2009KoZV
(Alpha-transition probabilities)
- 1976BaZZ S.A. Baranov, A.G. Zelenkov, V.M. Kulakov, Proc. Advisory Group Meeting on Transactinium Nucl. Data, Karlsruhe, Vol. III, p.249 (1976); IAEA-186 (1976).
(α -particle energies)
- 1977VaZW V.I. Vakarov, H. Sodan, R. Kalpakchieva, Y.T. Oganessian, Y.E. Penionzhkevich, V.N. Polyanskii, L.P. Chelnokov, JINR-P7-10123 (1977)
(α -particle energies)

- 1979Bo30 H.G. Borner, G. Barreau, W.F. Davidson, P. Jeuch, T. von Egidy, J. Almeida, D.H. White, Nucl. Instrum. Methods 166, 251 (1979).
(γ -ray energies)
- 1982Ah04 I. Ahmad, Nucl. Instrum. Methods 193, 9 (1982).
(γ -ray energies)
- 1986Ti03 G.A. Timofeev, V.V. Kalygin, and P.A. Privalova, Atom. Energiya 60, 286 (1986); Sov. At. Energy 60, 341 (1986)
(Half-life)
- 1987Po19 V.N. Polynov, A.A. Druzhinin, A.M. Korochkin, E.A. Nikitin, V.A. Bochkarev, V.N. Vyachin, V.G. Lapin, and M.Yu. Maksimov, Atom. Energiya 62, 277 (1987); Sov. At. Energy 62, 336 (1987)
(SF half-life)
- 1990Si12 E. Simeckova, P. Cizek, M. Finger, J. John, P. Malinsky, V.N. Pavlov, Hyperfine Interactions 59, 185 (1990).
(E2/M1 mixing ratios)
- 1991Ry01 A. Rytz, At. Data Nucl. Data Tables 47, 205 (1991)
(α -particle energies)
- 2003Au03 G. Audi, A.H. Wapstra, and C. Thibault, Nucl. Phys. A729, 337 (2003)
(Q value)
- 2003Br12 E. Browne, Nucl. Data Sheets 98, 665 (2003)
(^{243}Cm α decay scheme and α decay data evaluation)
- 2004Ak21 Y.A. Akovali, Nucl. Data Sheets 103, 515 (2004)
(^{243}Cm ε decay scheme and ε decay data evaluation)
- 2005Tr08 A.Trkov, G.L. Molnar, Zs. Revay, S.F. Mughabghab, R.B. Firestone, V.G. Pronyaev, A.L. Nichols, M.C. Moxon, Nucl. Sci. Eng. 150, 336 (2005)
(^{239}Pu gamma-ray emission probabilities in decay of ^{239}Np)
- 2008BeZV V.P. Chechev, N.K. Kuzmenko. ^{239}Np . In: Monographie BIPM-5, Vol.4. *Table of Radionuclides (Vol.4– A = 133 to 252)* by M.-M. Be, V. Chisté, C. Duié, E. Browne, V. Chechev, N. Kuzmenko, F. Kondev, A. Luca, M. Galan, A. Pearce, and X. Huang. / Sevres: Bureau International des Poids et Mesures, 2008.
(^{239}Pu gamma-ray emission probabilities in decay of ^{239}Np)
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr, Nucl. Instrum. Methods Phys. Res. A589, 202 (2008)
(Band-Raman ICC for γ -ray transitions)
- 2009KoZV F.G. Kondev, I. Ahmad, M.P. Carpenter, C.J. Chiara, J.P. Greene, R.V.F. Janssens, M.A. Kellett, T.L. Khoo, T. Lauritsen, C.J. Lister, E.F. Moore, A.L. Nichols, D. Seweryniak, S. Zhu, Proc. 13th Intern. Symposium on Capture Gamma-Ray Spectroscopy and Related Topics, Cologne, Germany, 25-29 Aug.2008, J. Jolie, A. Zilges, N. Warr, A. Blazhev, Eds., p.199 (2009); AIP Conf. Proc. 1090 (2009)
(Alpha-transition probabilities)

²⁴⁴Am - Comments on evaluation of decay data by A. L. Nichols

Evaluated: January 2007/February 2009

Evaluation Procedure

Limitation of Relative Statistical Weight Method (LWM) was applied to average the decay data when appropriate (but see below).

Decay Scheme

A relatively simple decay scheme was constructed from the gamma-ray studies of 1962Va08, 1963Ha29, 1967Sc34 and 1984Ho02. Only the gamma-ray measurements of Hoff *et al.* provide any estimates of the uncertainties in the gamma-ray probabilities expressed in terms of their relative intensity per 100 neutron captures in a high-flux reactor (1984Ho02). All other studies contained no information with respect to their overall uncertainties. Thus, no weighted mean data could be derived, and the data of 1984Ho02 were adopted wholesale and re-adjusted when seemed necessary (expressed in terms of the 743.977-keV gamma-ray emission probability (100 %)). Further measurements are merited to quantify the gamma-ray emission probabilities and decay scheme with greater certainty.

Nuclear Data

²⁴⁴Am is an important actinide for high burn-up fuel within the reactor core, and needs to be better characterized for improved assessments of accelerator-driven systems (ADS) and ²⁴⁴Cm decay heat contribution.

Half-life

The recommended half-life has been adopted from the single known measurement of Vandenbosch and Day (1962Va08). Further measurements are required to determine this half-life with much greater confidence.

Half-life measurement.

Reference	Half-life (hours)
1962Va08	10.1 ± 0.1

Gamma Rays

Energies

All gamma-ray transition energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies of Akovali were adopted (2003Ak04), and used to determine the energies and associated uncertainties of the gamma-ray transitions between the various populated-depopulated levels. However, Akovali recommended the gamma-ray energies determined by Hoff *et al.* (1984Ho02) by means of two curved-crystal spectrometers – minor differences do occur between the calculated energies of the higher energy transitions (538.402 (16), 743.977 (5) and 897.840 (7) keV) and those observed by Hoff *et al.*

Emission Probabilities

Relative emission probabilities and their uncertainties were determined from measurements of Hoff *et al.* (1984Ho02). These data were estimated to be in reasonably good agreement with the earlier measurements of Vandenbosch and Day, and Schuman (1962Va08, 1967Sc34), although these latter two sets of data possessed no uncertainties. Under these unsatisfactory circumstances, the data of Hoff *et al.* had to be adopted wholesale as the only suitable starting point in the attempted construction of a consistent decay scheme. Adjustments were made to the relative emission probabilities of the 99.383-, 153.863- and 205.575-keV gamma rays (adjusted from 7.0 (12) to 7.5 (13), 25 (5) to 28.6 (60), and 0.52 (12) to 0.53 (12), respectively) to conform with respect to the expected population-depopulation balance for the 501.79-, 296.21- and 142.35-keV nuclear levels of ²⁴⁴Cm. Furthermore, a relative emission probability had to be calculated for the 42.96-keV gamma ray for which there were no data at all (from a population-depopulation balance of the 42.96-keV nuclear level of ²⁴⁴Cm (populated by the 99.38-keV gamma ray and depopulated by the 42.96-keV gamma ray). Downward adjustments were made to the uncertainties of specific gamma-ray transitions and emissions through consideration of these and other data that are judged to be heavily correlated (99.383- and 153.863-keV gamma rays compared with 743.977-keV gamma ray and each other).

Measured relative gamma-ray emission probabilities.

	E_γ (keV)	1962Va08	1967Sc34	1984Ho02
		$P_\gamma^{Abs} \rightarrow P_\gamma^{rel}$	$P_\gamma^{Abs} \rightarrow P_\gamma^{rel}$	$P_\gamma^{Abs} \rightarrow P_\gamma^{rel}$
$\gamma_{1,0}$ (Cm)	42.965 (10)	-	-	-
$\gamma_{2,1}$ (Cm)	99.383 (4)	-	-	0.23 (4) \rightarrow 7.0 (12)
$\gamma_{3,2}$ (Cm)	153.863 (2)	72 \rightarrow 100	-	0.82 (16) \rightarrow 25 (5)
$\gamma_{4,3}$ (Cm)	205.575 (4)	0.4 \rightarrow 0.6	-	0.017 (4) \rightarrow 0.52 (12)
$\gamma_{9,4}$ (Cm)	538.402 (16)	0.4 \rightarrow 0.6	-	0.033 (7) \rightarrow 1.0 (2)
$\gamma_{9,3}$ (Cm)	743.977 (5)	72 \rightarrow 100	66.2 \rightarrow 100	3.3 (9) \rightarrow 100 (27)
$\gamma_{9,2}$ (Cm)	897.840 (7)	28 \rightarrow 39	27.6 \rightarrow 42	1.4 (4) \rightarrow 42 (12)

Gamma-ray emissions: recommended energies, relative emission probabilities, multipolarities and theoretical internal conversion coefficients (frozen orbital approximation).

γ (keV)	P_γ^{rel}	Multipolarity	α_K	α_L	α_{M+}	α_{tot}
42.965 (10)	0.145 (12)*	E2	-	760 (11)	290 (4)	1050 (15)
99.383 (4)	7.5 (13) [§]	E2	-	13.9 (2)	5.4 (1)	19.3 (3)
153.863 (2)	28.6 (60) [§]	E2	0.174 (3)	1.90 (3)	0.74 (1)	2.81 (4)
205.575 (4)	0.53 (12) [§]	E2	0.141 (2)	0.541 (8)	0.205 (3)	0.887 (13)
538.402 (16)	1.0 (2)	E2	0.0292 (4)	0.0149 (2)	0.0054 (1)	0.0495 (7)
743.977 (5)	100 (27)	M1 + E2 $\delta = -0.92$ (8)	0.059 (4)	0.0130 (7)	0.0050 (3)	0.077 (5)
897.840 (7)	42 (12)	E2	0.0122 (2)	0.00358 (5)	0.00124 (2)	0.0170 (3)

* Determined from the calculated theoretical internal conversion coefficients and the transition probability of the 99.383-keV gamma ray feeding the 42.965-keV nuclear level of ²⁴⁴Cm.

[§] Adjusted to conform with respect to the expected population-depopulation balances for the 501.79-, 296.21- and 142.35-keV nuclear levels of ²⁴⁴Cm.

A normalisation factor of 0.66 (14) was calculated from the relative emission probabilities of the three gamma rays that depopulate the 1040.188-keV nuclear level:

$$\sum_3 P_\gamma (1 + \alpha_{tot}) \times F = 100\%$$

$$\begin{aligned}
 & [P^{rel}(897.84\text{keV})(1 + \alpha_{tot}) + P^{rel}(743.97\text{keV})(1 + \alpha_{tot}) + P^{rel}(538.40\text{keV})(1 + \alpha_{tot})] \times F \\
 & = 100 \\
 F & = 100/151 (32) = 0.66 \pm 0.08
 \end{aligned}$$

Multipolarities and Internal Conversion Coefficients

The nuclear level scheme specified by Akovali has been used to define the multipolarities of the gamma transitions on the basis of known spins and parities (2003Ak04). Hansen *et al.* undertook angular correlation measurements to confirm the assignment of the 1040.2-keV nuclear level as the only ²⁴⁴Cm nuclear level populated directly by β^- decay (1963Ha29), in which the depopulating 743.977-keV gamma ray was defined as (46 \pm 4) % quadrupole [E2] and (54 \pm 4) % dipole [M1] to give a mixing ratio (δ) of -0.92 (8) for this transition. Recommended internal conversion coefficients have been determined from the theoretical tabulations of Band *et al.* (2002Ba25, 2002Ra45) by means of the methodology of Kibédi *et al.* (2008Ki07).

Beta-particle Emission

Energy and emission probability

The single beta-particle energy was calculated from the structural detail of the proposed decay scheme.

A nuclear level energy of 1040.188 (12) keV from Akovali (2003Ak04) and a Q_{β^-} value of 1427.3 \pm 1.0 keV from Audi *et al.* (2003Au03) were used to determine the energy and uncertainty of the beta-particle transition. By definition, this single beta transition was assigned an emission probability of 100 %.

Beta-particle Emission Probability per 100 Disintegrations of ²⁴⁴Am.

	E_{β} (keV)	P_{β}	Transition type	$\log ft$
$\beta_{0,9}^-$	387.1 \pm 1.0	100	(1 st forbidden non-unique)	5.63

Atomic Data

The x-ray and Auger-electron data have been calculated using the evaluated gamma-ray data, and atomic data from 1996Sc06, 1998ScZM and 1999ScZX. Both the x-ray and Auger-electron emission probabilities were determined by means of the EMISSION computer program (version 4.01, 28 January 2003, with the emission.101 database extended to Z = 96 to calculate component L x-ray data of daughter Cm). This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

K and L X-ray Emission Probabilities per 100 Disintegrations of ²⁴⁴Am.

			Energy (keV)	Photons per 100 disint.
XL		(Cm)	12.633 – 23.527	100 (10)
	XL _{L1}	(Cm)	12.633	2.36 (24)
	XL _{α}	(Cm)	14.746 – 14.961	36 (4)
	XL _{η}	(Cm)	17.314	1.15 (15)
	XL _{β}	(Cm)	17.286 – 19.688	51 (5)
	XL _{γ}	(Cm)	22.735 – 23.527	12.5 (13)
XK _{α}	XK _{α2}	(Cm)	104.590	2.2 (3)
	XK _{α1}	(Cm)	109.271	3.4 (4)
XK _{β1}	XK _{β3}	(Cm)	122.304)
	XK _{β1}	(Cm)	123.403) 1.29 (16)
	XK _{β5}	(Cm)	124.124)
XK _{β2}	XK _{β2}	(Cm)	126.889)
	XK _{β4}	(Cm)	127.352) 0.45 (6)
	XKO _{2,3}	(Cm)	127.970)

Electron energies were determined from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

References

- 1962Va08 S.E. VANDENBOSCH, P. DAY, The decay scheme of 10.1-h Am²⁴⁴, Nucl. Phys. 30 (1962) 177-190. [half-life, P_β, P_{ce}, relative P_γ]
- 1963Ha29 P.G. HANSEN, K. WILSKY, C.V.K. BABA, S.E. VANDENBOSCH, Decay of an isomeric state in Cm²⁴⁴, Nucl. Phys. 45 (1963) 410-416. [nuclear levels, δ]
- 1967Sc34 R.P. SCHUMAN, Nuclear chemistry: resonance activation integral measurements, IN-1126 (1967) 19. [relative P_γ]
- 1977La19 F.P. LARKINS, Semiempirical Auger-electron energies for elements 10 ≤ Z ≤ 100, At. Data Nucl. Data Tables 20 (1977) 311-387. [Auger-electron energies]
- 1984Ho02 R.W. HOFF, T. VON EGIDY, R.W. LOUGHEED, D.H. WHITE, H.G. BÖRNER, K. SCHRECKENBACH, G. BARREAU, D.D. WARNER, Levels of ²⁴⁴Cm populated by the beta decay of 10-h ²⁴⁴Am^g and 26-min ²⁴⁴Am^m, Phys. Rev. C29 (1984) 618-622. [relative P_γ, multipolarity]
- 1996Sc06 E. SCHÖNFELD, H. JANßEN, Evaluation of atomic shell data, Nucl. Instrum. Meth. Phys. Res. A369 (1996) 527-533. [X_K, X_L, Auger electrons]
- 1998ScZM E. SCHÖNFELD, G. RODLOFF, Tables of the energies of K-Auger electrons for elements with atomic numbers in the range from Z = 11 to Z = 100, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]
- 1999ScZX E. SCHÖNFELD, G. RODLOFF, Energies and relative emission probabilities of K X-rays for elements with atomic numbers in the range from Z = 5 to Z = 100, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2002Ba25 I.M. BAND, M.B. TRZHASKOVSKAYA, C.W. NESTOR, Jr., P.O. TIKKANEN, S. RAMAN, Dirac–Fock internal conversion coefficients, At. Data Nucl. Data Tables 81 (2002) 1-334. [ICC]
- 2002Ra45 S. RAMAN, C.W. NESTOR, Jr., A. ICHIHARA, M.B. TRZHASKOVSKAYA, How good are the internal conversion coefficients now? Phys. Rev. C66 (2002) 044312, 1-23. [ICC]
- 2003Ak04 Y.A. AKOVALI, Nuclear data sheets for A = 244, Nucl. Data Sheets 99 (2003) 197-273. [nuclear levels]
- 2003Au03 G. AUDI, A.H. WAPSTRA, C. THIBAULT, The AME2003 atomic mass evaluation (II). Tables, graphs and references, Nucl. Phys. A729 (2003) 337-676. [Q-value]
- 2008Ki07 T. KIBÉDI, T.W. BURROWS, M.B. TRZHASKOVSKAYA, P.M. DAVIDSON, C.W. NESTOR, Jr., Evaluation of theoretical conversion coefficients using BrIcc, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202-229. [ICC]

²⁴⁴Am^m - Comments on evaluation of decay data

by A. L. Nichols

Evaluated: January 2007/February 2009

Evaluation Procedure

Limitation of Relative Statistical Weight Method (LWM) was applied to average the decay data when appropriate (but see below).

Decay Scheme

A relatively simple decay scheme was constructed from the branching fraction measurements of Fields *et al.* and Gabeskiya *et al.* (1955Fi36, 1976Ga31) and the gamma-ray studies of Hoff *et al.* (1984Ho02). Only the gamma-ray studies of Hoff *et al.* provide estimates of the gamma-ray emission probabilities and their uncertainties per 100 neutron captures. Thus, no weighted mean data could be derived, and the data of 1984Ho02 were adopted as published.

Nuclear Data

²⁴⁴Am^m is an important actinide for high burn-up fuel within the reactor core, and needs to be better characterized for assessments of accelerator-driven systems (ADS) and ²⁴⁴Cm production and decay heat contribution.

Half-life

The recommended half-life has been adopted from two known measurements that did not quantify the uncertainties (1950St61, 1954Gh24). Thus, the assigned uncertainty is a crude estimate of ~ 10 %. This situation is extremely unsatisfactory, and further measurements are required to determine the half-life and uncertainty with much greater confidence.

Half-life measurements.

Reference	Half-life (min)
1950St61	~ 25
1954Gh24	26
Recommended value	26 ± 3

Branching Fractions (BF)

Fields *et al.* and Gabeskiya *et al.* have determined the EC/ β^- ratio (1955Fi36, 1976Ga31).

Reference	EC/ β^-
1955Fi36	0.000 38 ± 0.000 03*
1976Ga31	0.000 361 ± 0.000 013
Recommended value	0.000 36 ± 0.000 01

* Adjusted from 0.000 39 (3) on consideration of ²⁴⁴Cm half-life (18.11 (3) years).

Recommended EC/ β^- ratio was used to derive BF $_{\beta^-}$ of 0.999 64 (1) and BF_{EC} of 0.000 36 (1).

Gamma Rays

Energies

All gamma-ray transition energies were calculated from the structural details of the proposed decay scheme. The nuclear level energies of 2003Ak04 were adopted, and used to determine the energies and associated uncertainties of the gamma-ray transitions between the various populated-depopulated levels.

Emission Probabilities

Relative emission probabilities and their uncertainties were determined from the studies of Hoff *et al.* (1984Ho02). There are no other known measurements of these important decay characteristics. Under such unsatisfactory circumstances, the data of Hoff *et al.* had to be adopted wholesale, and further measurements are required to confirm the validity of the proposed decay scheme.

Measured gamma-ray emission probabilities per 100 neutron captures.

	E _γ (keV)	P _γ	Multipolarity
		1984Ho02	
γ _{1,0} (Cm)	42.965 (10)	(0.029)*	E2
γ _{6,1} (Cm)	941.95 (3)	0.33 (11)	E2
γ _{7,1} (Cm)	977.80 (4)	not detected	E0 (+ M1 + E2)
γ _{6,0} (Cm)	984.91 (2)	not detected	E0
γ _{10,1} (Cm)	1041.22 (3)	0.18 (6)	(M1 + E2)
γ _{11,1} (Cm)	1062.95 (3)	0.26 (8)	E1
γ _{10,0} (Cm)	1084.181 (14)	0.34 (11)	(E2)
γ _{11,0} (Cm)	1105.91 (2)	0.04 (2)	(E1)

* Calculated from experimental electron emission probabilities and theoretical internal conversion coefficients

Vandenbosch *et al.* have measured the ²⁴³Am(*n*,γ) cross-section ratio for ²⁴⁴Am^m and ²⁴⁴Am production (1964Va04), and this value has been used to convert the P_γ per 100 neutron captures to P_γ per 100 disintegrations of ²⁴⁴Am^m:

$$\frac{\sigma(^{243}\text{Am}(n,\gamma)^{244}\text{Am}^m)}{\sigma(^{243}\text{Am}(n,\gamma)^{244}\text{Am})} = 18.6(19)$$

$$^{244}\text{Am}^m = 18.6(19) \times ^{244}\text{Am} \quad (1)$$

Consider (*n*,γ) reaction to produce ²⁴⁴Am, and expressing the generation of ²⁴⁴Am and ²⁴⁴Am^m in the following manner:

$$\sum(^{244}\text{Am} + ^{244}\text{Am}^m) = 100\% \quad (2)$$

Substituting Eqn. (1) in (2):

$$^{244}\text{Am} = 100/19.6(19) = (5.1 \pm 0.5)\%$$

and $^{244}\text{Am}^m = (94.9 \pm 0.5)\%$

Absolute P_γ per 100 disintegrations of $^{244}\text{Am}^m$ were obtained by multiplying the P_γ per 100 neutron capture data of Hoff *et al.* by a factor of 1/0.949 (5).

There is considerable ambivalence in the quantification of the transition probabilities of the E0 977.80- and 984.91-keV gammas that cannot be satisfactorily resolved on the basis of the available measurements. While Hoff *et al.* found no evidence for any gamma-ray emissions with these particular energies (1984Ho02), von Egidy *et al.* observed a 977.92-keV gamma ray in their neutron capture studies with the following emission probability ratio (1984Vo07):

$$\frac{P_\gamma(977.92 \text{ keV})}{P_\gamma(1084.18 \text{ keV})} = \frac{0.12(4)}{0.52(16)},$$

and substituting $P_\gamma(1084.18 \text{ keV}) = 0.36(12)$ in this equation from the β^- decay of $^{244}\text{Am}^m$,

$$P_\gamma(977.92 \text{ keV}) = \frac{0.12(4)}{0.52(16)} \times 0.36(12) = 0.083(28),$$

with the recommended uncertainty reflecting only the uncertainty in $P_\gamma(1084.18 \text{ keV})$. This value is in good agreement with equivalent calculations involving the 1041.22- and 1062.95-keV gamma rays (0.084 (27) and 0.081 (24), respectively) that were also observed by von Egidy *et al.* (1984Vo07).

Gamma-ray emissions: recommended energies, absolute emission probabilities, multipolarities and theoretical internal conversion coefficients (frozen orbital approximation).

E_γ (keV)	P_γ^{abs}	Multipolarity	α_K	α_L	α_{M+}	α_{tot}
42.965 (10)	0.030 (9)*	E2	-	760 (11)	290 (4)	1050 (15)
941.95 (3)	0.35 (12)	E2	0.011 20 (16)	0.003 18 (5)	0.001 09	0.015 47 (22)
977.80 (4)	-	E0 (+ M1 + E2)	-	-	-	-
984.91 (2)	-	E0	-	-	-	-
1041.22 (3)	0.19 (6)	(M1 + E2)	-	-	-	-
1062.95 (3)	0.27 (8)	anomalous E1 ⁺	0.09 (3)	0.015 (4)	0.005	0.11 (3)
1084.181 (14)	0.36 (12)	anomalous (E2) ⁺	0.030 (8)	0.008 (2)	0.003	0.041 (11)
1105.91 (2)	0.04 (2)	anomalous (E1) ⁺	0.14 (3)	0.024 (6)	0.006	0.17 (4)

* Uncertainty of 30 % assigned on the basis of total transition probability of 30 (9), as defined by Hoff *et al.* (1984Ho02).

+ Anomalous internal conversion coefficients derived from the measurements of Hoff *et al.* (1984Ho02), with the components adjusted to match theoretical data on a relative basis.

Hoff *et al.* used a beta spectrometer to study the conversion electrons and determine the internal conversion coefficients of the various gamma transitions (1984Ho02). Total transition probability (TP_{total}) per 100 neutron captures were also derived by Hoff *et al.* for the two gamma transitions: 977.80-keV TP_γ per 100 disintegrations of $^{244}\text{Am}^m$ approximated to 0.08 (2), and 984.91-keV TP_γ per 100 disintegrations of $^{244}\text{Am}^m$ approximated to 1.0 (1). Anomalous internal conversion coefficients were observed for the 1062.95-, 1084.181- and 1105.91-keV gamma rays. A combination of the TP_γ and TP_{total} measurements of Hoff *et al.* and von Egidy *et al.* were

adopted (1984Ho02, 1984Vo07), while complete sets of anomalous internal conversion coefficients were determined on the basis of the theoretical data derived from Kibédi *et al.* (2008Ki07) and adjusted in terms of the studies of Hoff *et al.* (1984Ho02). The emission probability of the 42.965-keV gamma ray was estimated from Hoff *et al.* to be 0.029 per 100 neutron captures from TP_{total} of 30 (9) and the theoretical internal conversion coefficients. This transition probability of 30 (9) was corrected for the ²⁴⁴Am contribution to derive a TP_{total} of 32 (9) and P_γ(42.96 keV) of 0.030 (9) per 100 disintegrations of ²⁴⁴Am^m.

Multipolarities and Internal Conversion Coefficients

The nuclear level scheme specified by Akovali has been used to define the multipolarities of the gamma transitions on the basis of known spins and parities (2003Ak04). Recommended internal conversion coefficients have been determined from the theoretical tabulations of Band *et al.* (2002Ba25, 2002Ra45) by means of the methodology of Kibédi *et al.* (2008Ki07). Some of these data were judged to be anomalous from the studies of Hoff *et al.* (1984Ho02), and were adjusted accordingly (ICC data for the 1062.95-, 1084.181- and 1105.91-keV gamma transitions).

Beta-particle Emissions

Energies and emission probabilities

The ²⁴⁴Am^m nuclear level was estimated to have an energy of (89 ± 2) keV from S(n) of 5366.5 (17) keV (2003Au03) and a gamma-ray energy of 5277.6 (4) keV from the neutron capture state to ²⁴⁴Am^m (1984Vo07). Energies of the ²⁴⁴Cm nuclear levels adopted from Akovali (2003Ak04), Q_{β⁻} value of (1427.3 ± 1.0) keV from Audi *et al.* (2003Au03), and ²⁴⁴Am^m nuclear level energy of (89 ± 2) keV were used to determine the energies and uncertainties of the beta-particle transitions.

Adopted Nuclear Levels of ²⁴⁴Cm: J^π and Origins (2003Ak04).

Nuclear level number	Nuclear level energy (keV)	J ^π	Origins
0	0.0	0+	²⁴⁴ Bk EC decay, ²⁴⁴ Am β ⁻ decay, ²⁴⁴ Am ^m β ⁻ decay, ²⁴⁸ Cf α decay, Coulomb excitation
1	42.965 ± 0.010	2+	²⁴⁴ Am β ⁻ decay, ²⁴⁴ Am ^m β ⁻ decay, ²⁴⁸ Cf α decay, Coulomb excitation
2	142.348 ± 0.011	4+	²⁴⁴ Am β ⁻ decay, ²⁴⁸ Cf α decay, Coulomb excitation
3	296.211 ± 0.011	6+	²⁴⁴ Am β ⁻ decay
4	501.786 ± 0.012	8+	²⁴⁴ Am β ⁻ decay
5	970 ± 4	(2+, 3-)	Coulomb excitation
6	984.914 ± 0.021	0+	²⁴⁴ Am ^m β ⁻ decay
7	1020.76 ± 0.03	(2+)	²⁴⁴ Am ^m β ⁻ decay
8	1038 ± 6	(2+, 3-)	Coulomb excitation
9	1040.188 ± 0.012	6+	²⁴⁴ Am β ⁻ decay
10	1084.181 ± 0.014	1, 2+	²⁴⁴ Am ^m β ⁻ decay
11	1105.91 ± 0.02	(1, 2-)	²⁴⁴ Am ^m β ⁻ decay

Beta-particle emission probabilities were determined by balancing the proposed decay scheme through consideration of the βγ-population and γ-depopulation of the nuclear levels of daughter ²⁴⁴Cm. The recommended absolute gamma-ray emission probabilities and theoretical internal conversion coefficients derived from Kibédi *et al.* (2008Ki07) were used in this process, with the theoretical internal conversion coefficients adjusted if identified as anomalous on the basis of the measurements by Hoff *et al.*

Beta-particle Emission Probabilities per 100 Disintegrations of ²⁴⁴Am^m.

	²⁴⁴ Cm level energy (keV)	E _β (keV)	P _β	Transition type	log ft
β _{0,11} ⁻	1105.91 ± 0.02	410 ± 3	0.35 ± 0.09	(1st forbidden non-unique)	6.8
β _{0,10} ⁻	1084.181 ± 0.014	432 ± 3	0.56 ± 0.13	(allowed)	6.67
β _{0,7} ⁻	1020.76 ± 0.03	496 ± 3	0.08 ± 0.02	(allowed)	7.7
β _{0,6} ⁻	984.914 ± 0.021	531 ± 3	1.36 ± 0.16	allowed	6.58
β _{0,1} ⁻	42.965 ± 0.010	1473 ± 3	31 ± 9	allowed	6.74
β _{0,0} ⁻	0.0	1516 ± 3	67 ± 9	allowed	6.45

Σ 100.35

EC TransitionEnergy and transition probability

The EC transition energy was assigned a value of (164 ± 9) keV commensurate with Q_{EC} calculated from Audi *et al.* (2003Au03), while the transition probability was adopted from the recommended BF_{EC} of 0.00036 (1).

EC Transition Probability per 100 Disintegrations of ²⁴⁴Am^m.

	E _{EC} (keV)	P _{EC}	Transition type	log ft	P _K	P _L	P _M
EC _{0,0}	164 ± 9	0.036 ± 0.001	allowed	6.37	0.24 (5)	0.53 (4)	0.168 (12)

Atomic Data**K and L X-ray Emission Probabilities per 100 Disintegrations of ²⁴⁴Am^m.**

			Energy keV	Photons per 100 disint.
XL		(Cm)	12.633 – 23.527	12.3 (27)
	XL ₁	(Cm)	12.633	0.43 (8)
	XL _α	(Cm)	14.746 – 14.961	4.6 (11)
	XL _η	(Cm)	17.314	0.13 (4)
	XL _β	(Cm)	17.286 – 19.688	6.0 (14)
	XL _γ	(Cm)	22.735 – 23.527	1.4 (4)
XK _α	XK _{α2}	(Cm)	104.590	0.013 (4)
	XK _{α1}	(Cm)	109.271	0.020 (6)
XK _{β1}	XK _{β3}	(Cm)	122.304)
	XK _{β1}	(Cm)	123.403) 0.0076 (21)
	XK _{β5}	(Cm)	124.124)
XK _{β2}	XK _{β2}	(Cm)	126.889)
	XK _{β4}	(Cm)	127.352) 0.0027 (8)
	XKO _{2,3}	(Cm)	127.970)

The x-ray and Auger-electron data have been calculated using the evaluated gamma-ray data, and atomic data from 1996Sc06, 1998ScZM and 1999ScZX. Both the x-ray and Auger-electron

emission probabilities were determined by means of the EMISSION computer program (version 4.01, 28 January 2003, with the emission.101 database extended to $Z = 96$ to calculate component L x-ray data of daughter Cm). This program incorporates atomic data from 1996Sc06 and the evaluated gamma-ray data.

Electron energies were determined from electron binding energies tabulated by Larkins (1977La19) and the evaluated gamma-ray energies. Absolute electron emission probabilities were calculated from the evaluated absolute gamma-ray emission probabilities and associated internal conversion coefficients.

References

- 1950St61 K. STREET, Jr., A. GHIORSO, G.T. SEABORG, The isotopes of americium, Phys. Rev. 79 (1950) 530-531. [half-life]
- 1954Gh24 A. GHIORSO, S.G. THOMPSON, G.R. CHOPPIN, B.G. HARVEY, New isotopes of americium, berkelium and californium, Phys. Rev. 94 (1954) 1081. [half-life]
- 1955Fi36 P.R. FIELDS, Jr., J.E. GINDLER, A.L. HARKNESS, M.H. STUDIER, J.R. HUIZENGA, A.M. FRIEDMAN, Electron capture decay of Am²⁴⁴ and the spontaneous fission half-life of Pu²⁴⁴, Phys. Rev. 100 (1955) 172-173. [EC/ β^- ratio]
- 1962Va08 S.E. VANDENBOSCH, P. DAY, The decay scheme of 10.1-h Am²⁴⁴, Nucl. Phys. 30 (1962) 177-190. [spin, parity]
- 1964Va04 R. VANDENBOSCH, P.R. FIELDS, S.E. VANDENBOSCH, D. METTA, Search for a spontaneous fission branch in a metastable state of ²⁴⁴Cm, J. Inorg. Nucl. Chem. 26 (1964) 219-224. [spin, ²⁴³Am(n, γ)²⁴⁴Am cross-section ratio]
- 1976Ga31 V.Ya. GABESKIPIYA, A.P. CHETVERIKOV, V.V. GRYZINA, V.V. TIKHOMIROV, Measurement of relative probability of electron capture in ²⁴⁴Am decay, Sov. At. Energy 41 (1976) 1008-1009. [BF_{EC}]
- 1977La19 F.P. LARKINS, Semiempirical Auger-electron energies for elements $10 \leq Z \leq 100$, At. Data Nucl. Data Tables 20 (1977) 311-387. [Auger-electron energies]
- 1984Ho02 R.W. HOFF, T. VON EGIDY, R.W. LOUGHEED, D.H. WHITE, H.G. BÖRNER, K. SCHRECKENBACH, G. BARREAU, D.D. WARNER, Levels of ²⁴⁴Cm populated by the beta decay of 10-h ²⁴⁴Am^g and 26-min ²⁴⁴Am^m, Phys. Rev. C29 (1984) 618-622. [P_γ , multipolarity]
- 1984Vo07 T. VON EGIDY, R.W. HOFF, R.W. LOUGHEED, D.H. WHITE, H.G. BÖRNER, K. SCHRECKENBACH, D.D. WARNER, G. BARREAU, Nuclear structure of ²⁴⁴Am investigated with the (n, γ) reaction, Phys. Rev. C29 (1984) 1243-1267. [spin, parity, nuclear level energy of ²⁴⁴Am^m]
- 1996Sc06 E. SCHÖNFELD, H. JANßEN, Evaluation of atomic shell data, Nucl. Instrum. Meth. Phys. Res. A369 (1996) 527-533. [X_K , X_L , Auger electrons]
- 1998ScZM E. SCHÖNFELD, G. RODLOFF, Tables of the energies of K-Auger electrons for elements with atomic numbers in the range from $Z = 11$ to $Z = 100$, PTB Report PTB-6.11-98-1, October 1998. [Auger electrons]

Comments on evaluation

- 1999ScZX E. SCHÖNFELD, G. RODLOFF, Energies and relative emission probabilities of K X-rays for elements with atomic numbers in the range from $Z = 5$ to $Z = 100$, PTB Report PTB-6.11-1999-1, February 1999. [X_K]
- 2002Ba85 I.M. BAND, M.B. TRZHASKOVSKAYA, C.W. NESTOR, Jr., P.O. TIKKANEN, S. RAMAN, Dirac–Fock internal conversion coefficients, At. Data Nucl. Data Tables 81 (2002) 1-334. [ICC]
- 2002Ra45 S. RAMAN, C.W. NESTOR, Jr., A. ICHIHARA, M.B. TRZHASKOVSKAYA, How good are the internal conversion coefficients now? Phys. Rev. C66 (2002) 044312, 1-23. [ICC]
- 2003Ak04 Y.A. AKOVALI, Nuclear data sheets for $A = 244$, Nucl. Data Sheets 99 (2003) 197-273. [nuclear levels]
- 2003Au03 G. AUDI, A.H. WAPSTRA, C. THIBAUT, The AME2003 atomic mass evaluation (II). Tables, graphs and references, Nucl. Phys. A729 (2003) 337-676. [Q-values]
- 2008Ki07 T. KIBÉDI, T.W. BURROWS, M.B. TRZHASKOVSKAYA, P.M. DAVIDSON, C.W. NESTOR, Jr., Evaluation of theoretical conversion coefficients using BrIcc, Nucl. Instrum. Methods Phys. Res. A589 (2008) 202-229. [ICC]

²⁴⁴Cm - Comments on evaluation of decay data by V. P. Chechev

This evaluation was completed in February 2005 (see 2006Ch34) and then corrected in October 2009 with a literature cut-off by the same date.

1 Decay Scheme

The decay scheme is based on the evaluation of 2004Ch64. It can be considered essentially complete although some weak gamma-ray transitions have not been observed in ²⁴⁴Cm alpha decay. Such gamma-rays were taken from the ²⁴⁰Np β⁻-decay and the ²⁴⁰Am electron capture and have been included in the decay scheme.

2 Nuclear Data

Q(α) value is from 2003Au03.

The evaluated half-life of ²⁴⁴Cm is based on the experimental values given in Table 1.

Table 1. Experimental values of the ²⁴⁴Cm half-life (in years).

Reference	Author(s)	Value	Measurement method
1954Fr19	Friedman et al.	17.9 (5)	α-activity relative to ²⁴² Cm
1954St33	Stevens et al.	19.2 (6)	α-activity relative to ²⁴² Cm
1961Cao1	Carnall et al.	17.59 (6)	Specific activity
1968Be26	Bentley	18.099 (32) ^a	2π α-counting
1972Ke29	Kerrigan and Dorsett	18.13 (4)	Calorimetry
1982Po14	Polyukhov et al.	18.24 (25)	Specific activity

^a Revised value, recalculated in 2000Ho27.

The EV1NEW program has led to successive rejections of values from 1961Ca01 and 1954St33 due to their too large contribution to χ²-value (more than 80 %). The LRSW method has increased 1.03 times the uncertainty of the value from 1968Be26. The weighted mean of the data set including only the four remaining values is 18.115, with the internal uncertainty 0.028 and χ²/ν = 0.25. The smallest experimental uncertainty is 0.032, thus the recommended value of ²⁴⁴Cm half-life is **18.11 (3) a**.

The recommended spontaneous fission partial half-life of ²⁴⁴Cm is based on the experimental values given in Table 2.

Table 2. Experimental values of the ²⁴⁴Cm spontaneous fission half-life (in 10⁷ years).

Reference	Author(s)	Value	Measurement method
1952Gh27	Ghiorso et al.	1.4 (2) ^a	Ionization chamber
1963Ma56	Malkin et al.	1.46 (6)	Gas scintillator
1965Me02	Metta et al.	1.345 (8) ^a	α/SF counting, α with low geometry counter, SF with 2π parallel plate chamber
1967Ar09	Armani and Gold	1.33 (3)	Fission neutron counting, LiI detector
1970Ba11	Barton and Koontz	1.250 (7)	Low geometry fission fragment counting
1972Ha80	Hastings and Strohm	1.343 (6) ^a	α/SF counting, Si(Au) detector
1993Pa29	Pandey et al.	1.263 (5)	α/SF counting by sequential etching of alpha and fission tracks

^a Revised value, recalculated in 2000Ho27.

The data set in Table 2 is discrepant. The LWEIGHT computer program has chosen the unweighted mean of 1.342 and expanded the uncertainty to 0.079 so its range includes the most precise value of 1993Pa29.

The recommended value of ²⁴⁴Cm spontaneous fission half-life is $1.34 (8) \times 10^7$ years.

2.1 α Transitions

The energies of the alpha transitions have been obtained from the Q value and the ²⁴⁰Pu level energies given in Table 3 from 2004Ch64.

Table 3. ²⁴⁰Pu levels populated in the ²⁴⁴Cm α -decay.

Level number	Energy (keV)	Spin and parity	Half-life	Probability of α -transition (%)
0	0.0	0 ⁺	6561 (7) a	76.7 (4)
1	42.824 (8)	2 ⁺	164 (5) ps	23.3 (4)
2	141.690 (15)	4 ⁺		0.0204 (15)
3	294.319 (24)	6 ⁺		0.00352 (18)
4	497.6 ^a	8 ⁺		4×10^{-5}
5	597.34 (4)	1 ⁻		$5.5 (9) \times 10^{-5}$
6	648.85 (4)	3 ⁻		$4.2 (30) \times 10^{-6}$ b
7	860.71(7)	0 ⁺		$1.49 (16) \times 10^{-4}$
8	900.32 (4)	2 ⁺		$5.0 (5) \times 10^{-5}$
9	938.06 (6)	(1 ⁻)		$4.7 (11) \times 10^{-6}$ b

^a Energy has been taken from ²³⁸U(α , 2n γ)-reaction measurements of 1972Sp06.

^b Deduced from P(γ +ce) decay-scheme probability balances.

The probabilities of the transitions $\alpha_{0,i}$ ($i = 0, 1, 2, 3, 7$) have been obtained by averaging experimental data (Table 4). The experimental results from 1998Ga19 agree well with the evaluated probabilities of the most intense alpha-transitions. The probabilities of the remaining α -transitions have been deduced using the experimental values and the values obtained from P(γ +ce) decay-scheme balances (see footnotes).

Table 4. Experimental and recommended α -transition probabilities (%) in the ²⁴⁴Cm decay.

	α -energy (keV)	1956 Hu96	1960 As11, 1984 Asaro	1963 Dz07	1966 Ba07	1984 BuZJ	1996 Bu50	1996 Sa24	1997 Ka59	1998 Ga19	1998 Ya17	2002 Da21	Recommended
$\alpha_{0,0}$	5805	76.7 (6)	-	76.2 (20)	76.4 (20) ^a	76.98 (5)	76.8 (7)	76.9 (5)	-	76.63 (18)	76.31 (5)	77.16 (11)	76.7 (4) ^b
$\alpha_{0,1}$	5763	23.3 (6)	-	23.8 (9)	23.6 (9) ^a	23.00 (5)	23.2 (5)	23.1 (5)	-	23.34 (18)	23.69 (6)	22.80 (5)	23.3 (4) ^c
$\alpha_{0,2}$	5664	0.017 (3)	0.023 (2)	0.021 (2)	0.02	0.0163 (7)	-	0.0135 (2)	-	0.0205 (15)	-	0.020 (1)	0.0204 (15) ^d
$\alpha_{0,3}$	5515	-	0.0036 (3)	0.003 (1)	0.0034	-	-	-	0.003 42 (9)	0.0038 (5)	-	0.012 (1)	0.003 52 (18) ^e
$\alpha_{0,4}$	5315	-	$\sim 1.5 \cdot 10^{-4}$	-	$\sim 4 \cdot 10^{-5}$	-	-	-	-	-	-	-	$4 \cdot 10^{-5}$ ^f
$\alpha_{0,5}$	5215	-	$1.5 \cdot 10^{-4}$	-	$1 \cdot 10^{-4}$	-	-	-	$4.2 (9) \cdot 10^{-5}$	-	-	-	$5.5 (9) \cdot 10^{-5}$ ^g
$\alpha_{0,7}$	4960	-	$1.55 (16) \cdot 10^{-4}$	-	$3 \cdot 10^{-4}$	-	-	-	$1.42 (16) \cdot 10^{-4}$	-	-	-	$1.49 (16) \cdot 10^{-4}$ ^h
$\alpha_{0,8}$	4920	-	$5.0 (5) \cdot 10^{-5}$	-	$1.3 \cdot 10^{-4}$	-	-	-	$4.9 (8) \cdot 10^{-5}$	-	-	-	$5.0 (5) \cdot 10^{-5}$ ⁱ

^a No uncertainties are quoted by the authors. The uncertainties have been adopted by the evaluator based on the analogy of the spectra obtained with magnetic spectrometers in 1963Dz07 and 1966Ba07.

^b This set of experimental values is discrepant. The LWEIGHT computer program has recommended a weighted average and expanded the uncertainty so the range includes the most precise value from 1998Ya17.

^c Obtained from the relation $P(\alpha_{0,1}) = 100 - P(\alpha_{0,0})$ per 100 disintegrations. An unweighted average of the discrepant set of the experimental values is 23.31, a weighted average is 23.11.

^d Weighted average of the values from 1956Hu96, 1960As11, 1963Dz07, 1998Ga19 and 2002Da21. The lower values from 1984BuZJ and 1996Sa24 have been omitted as outliers. These values conflict greatly with the ratio $P(\gamma_{2,1})/P(\gamma_{1,0}) = 0.067 (7)$ measured in 1972Sc01. The uncertainty of the evaluated $\alpha_{0,2}$ probability has been adopted from the experimental result of 1998Ga19.

^e Average of values from 1960As11, 1963Dz07, 1997Ka59 and 1998Ga19. The EV1NEW computer program using a limitation of relative statistical weights of 0.5 has expanded the uncertainty from 1997Ka59 to 0.00025 and recommended a weighted average and an internal uncertainty.

^f Adopted from 1966Ba07.

^g Deduced from the P(γ +ce)-probability balance at the 597-keV level (“5”).

^h Weighted average of values from 1960As11, 1997Ka59.

ⁱ Weighted average of values from 1960As11, 1997Ka59 and a value of $5.2 (7) \times 10^{-5}$, calculated from P(γ +ce)-probability balance at the 900-keV level (“8”). The uncertainty is the smallest experimental one.

2.2 γ Transitions

The evaluated energies of gamma-ray transitions are virtually the same as the photon energies because nuclear recoil is negligible.

The probabilities, P(γ +ce), for gamma-ray transitions of 42.8-keV ($\gamma_{1,0}$), 98.9-keV ($\gamma_{2,1}$), 152.6-keV ($\gamma_{3,2}$), and 202-keV ($\gamma_{4,3}$) have been deduced from intensity balances, using the probabilities of α -particle transitions evaluated directly from experimental data.

For the 861-keV ($\gamma_{7,0}$) E0 transition its P(ce) value has been obtained from the (α -ce)-coincidence measurement of 1963Bj03: $P(\text{ce } \gamma_{7,0}) + P(\text{ce } \gamma_{7,1}) = 9.5 (20) \times 10^{-6}$ per 100 disintegrations.

The remaining P(γ +ce) values have been calculated from the gamma-ray emission probabilities and the total internal conversion coefficients (ICC's). The ICC's have been interpolated using the BrIcc package with the so called “*Frozen Orbital*” approximation (2008Ki07). The fractional uncertainties of α_K , α_L , α_M , α_T for pure multiplicities have been taken as 2 %.

Multipolarities are from 2004Ch64. These are based on conversion electron measurements of 1956Sm18, 1963Bj03, 1968Du06 and 1990Pe03.

3 Atomic Data

3.1. Fluorescence yields

The fluorescence yields are from 1996Sc06 (Schönfeld and Janßen).

3.2 X radiations

The Pu KX-ray energies and relative emission probabilities are from 1999Schönfeld, where the calculated energy values are based on X-ray wavelengths from 1967Be65 (Bearden). In Table 5 the recommended values of U KX-ray energies are compared with experimental values.

Table 5. Experimental and recommend values of Pu KX-ray energies (keV).

	1980Di13	1982Ba56	Recommended
K α_2	99.55 (3)	99.530 (2)	99.525
K α_1	103.76 (3)	103.741 (2)	103.734
K β_3	116.27	116.242 (2)	116.244
K β_1	117.26	117.233 (2)	117.228
K $\beta_{2,4}$	120.60 (15)	-	120.553
KO $_{2,3}$	121.55 (6)	-	121.543

In 1980Di13 the Pu KX-ray energies were measured in the alpha decay of ²⁴⁵Cm. The relative emission probabilities of KX-rays were obtained as:

$$K\alpha_2 : K\alpha_1 : K\beta_3 : K\beta_1 : K\beta_{2,4} = 64.7 (23) : 100.0 (33) : 12.9 (7) : 23.1 (10) : 8.9 (5).$$

3.3. Auger Electrons

The energies of Auger electrons have been calculated from atomic electron binding energies.

The P(KLX)/P(KLL), P(KXY)/P(KLL) ratios have been taken from 1996Sc06.

4 α Emissions

The energy of alpha particles to the ground state of ²⁴⁰Pu, E($\alpha_{0,0}$), are from the absolute measurement of 1971Gr17 but including the correction of -0.19 keV recommended by A. Rytz in 1991Ry01.

The energies of all other α -particles have been deduced from Q $_{\alpha}$ and ²⁴⁰Pu level energies including the recoil energy corrections.

In Table 6 the recommended values of α -particle energies are compared with experimental results obtained with magnetic alpha spectrometers.

Table 6. Experimental^a and evaluated α -particle energies in the decay of ²⁴⁴Cm (keV).

	1960 As11	1963 Dz07	1966 Ba07	1971 Gr17	1992 Fr04	1998 Ga19	Recommended
$\alpha_{0,0}$	5805	5805 (3)	5805 (1)	5804.77 (5)	5803.6 (22)	-	5804.77 (5)
$\alpha_{0,1}$	5763	5762	5763 (1)	5762.16 (3)	-	-	5762.65 (5)
$\alpha_{0,2}$	5666	5665	5664 (3)	-	-	5664 (2)	5665.41 (5)
$\alpha_{0,3}$	5514	5514	5513 (3)	-	-	5515 (3)	5515.29 (6)
$\alpha_{0,4}$	5316	-	5313	-	-	-	5315.3
$\alpha_{0,5}$	5215	-	5215 (3)	-	-	-	5217.24 (7)
$\alpha_{0,7}$	4956	-	4960 (3)	-	-	-	4958.20 (9)
$\alpha_{0,8}$	4916	-	4920 (3)	-	-	-	4919.24 (7)

^a Authors' values have been adjusted for changes in calibration energies (see 1991Ry01).

5 Electron emissions

The energies of conversion electrons have been obtained from gamma transition energies and relevant electron binding energies. The emission probabilities of conversion electrons have been deduced from the evaluated P(γ) and ICC values.

The absolute emission probabilities of K and L Auger electrons have been calculated using the EMISSION computer program (2000Schönfeld).

6 Photon emissions

6.1 X-ray emissions

The absolute emission probabilities of U KX- and U LX-rays in decay of ²⁴²Pu have been calculated using the EMISSION computer program (2000Schönfeld).

The calculated total absolute emission probability of LX-rays P(XL)= 8.92 (23) % agrees with the experimental value of 8.77 (6) % from 1995Jo23.

In 1990Po14 the relative LX-ray emission probabilities in ²⁴⁴Cm α -decay were measured:

$$[5.3 (8) : Ll; 72 (7) : L\alpha; 100: L\eta\beta; 22.4 (23) : L\gamma].$$

These values agree with the recommended ones with the exception of the (L α /L $\eta\beta$)-ratio.

6.2 Gamma-ray emissions

6.2.1. Gamma-ray energies

The energies of the 43-keV ($\gamma_{1,0}$), 99-keV ($\gamma_{2,1}$), and 153-keV ($\gamma_{3,2}$) gamma rays are from ²⁴⁴Cm α -decay (1972Sc01). Other, less accurate measurements of ²⁴⁴Cm α -decay (1956Sm18), ²⁴⁰Np β^- -decay (1981Hs02) and ²⁴⁰Am ε -decay (1972Ah07) agree with data from 1972Sc01.

The energies of remaining gamma rays have been obtained from the adopted ²⁴⁰Pu level energies. In Table 7 the recommended gamma ray energies are compared with the available experimental data.

Table 7. Experimental and recommended gamma-ray energies (keV).

	1967Lederer (1978LeZA)	1972Ah07	1972Sc01	1981Hs02	Recommended
$\gamma_{1,0}$		42.9 (1)	42.824 (8)	-	42.824 (8)
$\gamma_{2,1}$	-	98.9 (1)	98.860 (13)	-	98.860 (13)
$\gamma_{3,2}$	-	-	152.630 (20)	-	152.630 (20)
$\gamma_{8,6}$	251.20 (20)	-	-	251.5 (1)	251.47 (6)
$\gamma_{7,5}$	263.34 (15)	-	-	263.4 (1)	263.37 (8)
$\gamma_{8,5}$	302.99 (15)	-	-	303.0 (1)	302.98 (6)
$\gamma_{6,2}$	506.9 (3)	-	-	507.2 (1)	507.16 (5)
$\gamma_{5,1}$	554.5 (2)	-	-	554.6 (1)	554.52 (4)
$\gamma_{5,0}$	597.2 (2)	-	-	597.4 (1)	597.34 (4)
$\gamma_{6,1}$	605.8 (2)	-	-	606.1 (1)	606.03 (4)
$\gamma_{8,2}$	758.6 (2)	-	-	758.6 (1)	758.63 (5)
$\gamma_{7,1}$	817.8 (2)	-	-	817.9 (1)	817.89 (7)
$\gamma_{8,1}$	857.5 (2)	-	-	857.5 (1)	857.50 (4)
$\gamma_{9,1}$	894.7 (5)	-	-	895.3 (1)	895.24 (6)
$\gamma_{8,0}$	900.1 (5)	-	-	900.3 (1)	900.32 (4)
$\gamma_{9,0}$	937.6 (10)	-	-	938.0 (1)	938.06 (6)

6.2.2. Gamma-Ray Emission Probabilities

The absolute emission probabilities for gamma rays of 43-keV ($\gamma_{1,0}$), 99-keV ($\gamma_{2,1}$), 153-keV ($\gamma_{3,2}$) and 202-keV ($\gamma_{4,3}$) have been deduced from intensity balances, using the experimental α -particle probabilities. The relative emission probabilities for the first three gamma rays were measured in 1972Sc01 as [100 - $\gamma_{1,0}$, 6.7 (7) - $\gamma_{2,1}$, and 4.1 (1) - $\gamma_{3,2}$]. The measured $P(\gamma_{2,1})/P(\gamma_{1,0}) \times 100$ ratio disagrees with the evaluated 5.3 (4), and the measured $P(\gamma_{3,2})/P(\gamma_{1,0}) \times 100$ ratio agrees with the evaluated 3.95 (23).

The recommended relative emission probabilities of gamma rays with energies greater than 150-keV, obtained by averaging the experimental data from 1967Lederer (1978LeZA) and 1969Sc18 (1970Sc39), are given in Table 8.

Table 8. Experimental and recommended relative emission probabilities of > 150-keV gamma rays from the decay of ²⁴⁴Cm.

	Energy (keV)	1967Lederer 1978LeZA	1969Sc18 1970Sc39	Recommended
$\gamma_{3,2}$	152.6	-	1240 (150)	1170 (160) ^a
$\gamma_{8,6}$	251.5	14 (3)	12.7 (20)	13.1 (20) ^b
$\gamma_{7,5}$	263.4	73 (5)	68 (6)	71 (5) ^b
$\gamma_{8,5}$	303.0	23 (4)	21.0 (20)	21.4 (20) ^b
$\gamma_{6,2}$	507.2	10 (3)	-	10 (3) ^c
$\gamma_{5,1}$	554.5	100	100	100
$\gamma_{5,0}$	597.3	61 (2)	62 (4)	61 (2) ^b
$\gamma_{6,1}$	606.0	10 (2)	9.1 (11)	9.3 (20) ^b
$\gamma_{8,2}$	758.6	15.6 (8)	18.3 (21)	15.9 (8) ^b
$\gamma_{7,1}$	817.9	75 (4)	91 (8)	78 (4) ^b
$\gamma_{8,1}$	857.5	6.6 (4)	< 7.5	6.6 (4) ^c
$\gamma_{9,1}$	895.2	2.1 (6)	< 1.3	2.1 (6) ^c
$\gamma_{8,0}$	900.3	1.5 (6)	< 0.4	1.5 (6) ^c
$\gamma_{9,0}$	938.1	0.5 (5)	< 0.75	0.5 (5) ^c

^a Deduced from the evaluated absolute emission probabilities P(γ 153 keV) and P(γ 555 keV).
^b Weighted average, uncertainty is the smallest experimental value reported.
^c Adopted from 1967Lederer (1978LeZA).

The deduced absolute emission probabilities of gamma-rays with energies greater than 250 keV are based on our recommended relative gamma-ray emission probabilities P(γ)/P(γ 555 keV) in Table 8 and a normalization factor obtained from decay scheme.

The absolute gamma-ray emission probability P⁽¹⁾(γ 555 keV) = 9.1 (11) × 10⁻⁵ per 100 disintegrations (used for decay-scheme normalization) has been obtained from the intensity balance at the 861-keV level (“7”) using the alpha-transition probability P($\alpha_{0,7}$) = 1.49 (16) × 10⁻⁴ per 100 disintegrations, deduced from the experimental data of 1960As11 and 1997Ka59:

$$P(\gamma \text{ 555 keV}) = [P(\alpha_{0,7}) - P(\text{ce 861 keV})] / [P'(\gamma \text{ 263 keV}) \times (1 + \alpha_T^{263}) + P'(\gamma \text{ 818 keV}) \times (1 + \alpha_T^{818})],$$

where P'(γ) is a gamma-ray emission probability relative to that of the 555-keV transition (i.e., P(γ)/P(γ 555 keV)).

Another way of deducing a normalization factor is by using the relative gamma-ray emission probability P(γ 153 keV)/P(γ 555 keV) = 12.4 (15) measured in 1969Sc18 (1970Sc39) and the absolute probability P(γ 153 keV) obtained from the intensity balance for the level 294-keV level (“3”):

$$P^{(2)}(\gamma \text{ 555 keV}) = 8.2 (11) \times 10^{-5} \text{ per 100 disintegrations.}$$

The average of the two P(γ 555 keV) values, 8.7 (11) × 10⁻⁵ per 100 disintegrations, was used as a normalization factor for calculating absolute emission probabilities of gamma-rays with energy greater than 250 keV.

The absolute emission probabilities for the 289-keV ($\gamma_{9,6}$) and 341-keV ($\gamma_{9,5}$) gamma rays have been deduced using the ratios P(γ 895 keV)/P(γ 289 keV) = 3.6 (15) and P(γ 895 keV)/P(γ 341 keV) = 1.0 (3) measured in ²⁴⁰Np β^- -decay (1981Hs02, 2004Ch64).

The absolute emission probability of the 202-keV ($\gamma_{4,3}$) gamma ray has been obtained using the adopted $\alpha_{0,4}$ -transition probability. The 202-keV E2-gamma-ray transition was not observed in the ²⁴⁴Cm alpha decay; however, it is expected from theoretical considerations and by analogy with the ²⁴²Cm decay scheme.

7 Consistency of recommended data

The most accurate Q value, Q(M), is taken from the atomic mass adjustment table of Audi et al. (2003Au03). Comparison of Q(eff)(deduced as the sum of average energies per disintegration ($\sum E_i \times P_i$) for all emissions accompanying ²⁴⁴Cm α - decay) with the tabulated decay energy Q(M) allows to check a consistency of the recommended decay-scheme parameters obtained in this evaluation.

Here E_i and P_i are the evaluated energies and emission probabilities of the i-th alpha particle, beta particle, gamma ray, X-ray, etc. Consistency (percentage deviation) is determined by $\{[Q(M) - Q(\text{eff})] / Q(M)\} \times 100$. "Percentage deviations above 5 % would be regarded as high and imply a poorly defined decay scheme; a value of less than 5 % indicates the construction of a reasonably consistent decay scheme" (quoted from the article by A.L. Nichols in Appl. Rad. Isotopes 55 (2001) 23-70).

For the above ²⁴⁴Cm decay data evaluation we have Q(M) = 5901.74 (5) keV and Q(eff) = 5903 (33) keV, i.e. consistency of (0.02 ± 0.56) % is not superior, but better than 0.6 %.

8 References

- 1952Gh27 A. Ghiorso, G. H. Higgins, A. E. Larsh, G. T. Seaborg, S. G. Thompson, Phys. Rev. 87, 163 (1952) (SF half-life)
- 1954Fr19 A. M. Friedman, A. L. Harkness, P. R. Fields, M. H. Studier, J. R. Huizenga, Phys. Rev. 95, 1501 (1954) (Half-life)
- 1954St33 C. M. Stevens, M. H. Studier, P. R. Fields, J. F. Mech, P. A. Sellers, A. M. Friedman, H. Diamond, J. R. Huizenga, Phys. Rev. 94, 974 (1954) (Half-life)
- 1956Hu96 J. P. Hummel, Thesis, Univ. California (1956), UCRL-3456 (1956) (α -transition probabilities)
- 1956Sm18 W. G. Smith, J. M. Hollander, Phys. Rev. 101, 746 (1956) (Conversion electron measurements, gamma ray multipolarities)
- 1960As11 F. Asaro, I. Perlman, Priv. Comm. (1960), quoted by E. K. Hyde et al. in: The Nuclear Properties of the Heavy Elements, Vol. II, Prentice-Hall, Englewood Cliffs, New Jersey, 1964, p. 880 (α -transition probabilities, α -emission energies)
- 1961Ca01 W. T. Carnall, S. Fried, A. L. Harkness, J. Inorg. Nuclear Chem., 17, 12 (1961) (Half-life)
- 1963Bj03 S. Bjornholm, C. M. Lederer, F. Asaro, I. Perlman, Phys. Rev. 130, 2000 (1963) (E0 gamma and alpha transition probabilities)
- 1963Dz07 B. S. Dzhelepov, R. B. Ivanov, V. G. Nedovesov, V. P. Chechev, Zh. Eksperim. i Teor. Fiz. 45, 1360 (1963); Soviet Phys. JETP 18, 937 (1964) (α -transition probabilities, α -emission energies)
- 1963Ma56 L. Z. Malkin, I. D. Alkhozov, A. S. Krivokhatskii, K. A. Petrzhak, L. M. Belov, At. Energ. USSR 16, 148 (1964); Soviet J. At. Energy 16, 170 (1964) (SF half-life)
- 1965Me02 D. Metta, H. Diamond, R. F. Barnes, J. Milsted, J. Gray, Jr, D. J. Henderson, C. M. Stevens, J. Inorg. Nucl. Chem. 27, 33 (1965) (SF half-life)
- 1966Ba07 S. A. Baranov, Y. F. Rodionov, V. M. Kulakov, V. M. Shatinskii, Yadern. Fiz. 4, 1108 (1966); Soviet J. Nucl. Phys. 4, 798 (1967) (α -transition probabilities, α -emission energies)
- 1967Ar09 R. J. Armani, R. Gold, Proc Symp Standardization of Radionuclides, Vienna, Austria (1966), Intern At Energy Agency, Vienna, p 621 (1967) (SF half-life)
- 1967Be65 J.A. Bearden, Rev. Mod. Phys. 39, 78(1967) (X-ray energies)
- 1967Lederer C. M. Lederer, Priv. Comm. (1967), quoted in 1978LeZA (Gamma ray energies and probabilities)
- 1968Be26 W. C. Bentley, J. Inorg. Nucl. Chem. 30, 2007 (1968) (Half-life)
- 1968Du06 C. L. Duke, W. L. Talbert, Jr, Phys. Rev. 173, 1125 (1968) (Conversion electron measurements, gamma ray multipolarities)
- 1969Sc18 M. R. Schmorak et al., Int. Conf. Radioactivity in Nucl. Spectroscopy Tech. and Appl., Nashville, p. 22. Priv. Comm. (1969) quoted in M. R. Schmorak, Nucl. Data Sheets B4, 661 (1970) (Gamma-ray energies and probabilities)
- 1970Ba11 D. M. Barton, P. G. Koontz, J. Inorg. Nucl. Chem. 32, 769 (1970) (SF half-life)
- 1970Sc39 M. R. Schmorak, Nucl. Data Sheets B4, 661 (1970) (Gamma-ray energies and probabilities)

- 1971Gr17 B. Grennberg, A. Rytz, *Metrologia*. 7, 65 (1971) (Alpha-particle energies)
- 1972Ah07 I. Ahmad, R. F. Barnes, R. K. Sjoblom, P. R. Fields, *J. Inorg. Nucl. Chem.* 34, 3335 (1972) (Gamma ray energies)
- 1972Ha80 J. D. Hastings, W. W. Strohm, *J. Inorg. Nucl. Chem.* 34, 3597 (1972) (SF half-life)
- 1972Ke29 W. J. Kerrigan, R. S. Dorsett, *J. Inorg. Nucl. Chem.* 34, 3603 (1972) (Half-life)
- 1972Sc01 M. Schmorak, C. E. Bemis, Jr., M. J. Zender, N. B. Gove, P. F. Dittner, *Nucl. Phys.* A178, 410 (1972) (Gamma ray energies and probabilities)
- 1972Sp06 H. J. Specht, J. Weber, E. Konecny, D. Heunemann, *Phys. Lett.* 41B, 43 (1972) (Level energies)
- 1978LeZA C. M. Lederer, V. S. Shirley, E. Browne, J. M. Dairiki, R. E. Doebler, A. A. Shihab-Eldin, L. J. Jardine, J. K. Tuli, A. B. Buyrn, *Table of Isotopes*, 7th Ed., John Wiley and Sons, Inc., New York (1978) (Gamma ray energies and probabilities)
- 1980Di13 J. K. Dickens, J. W. McConnell, *Phys. Rev.* C22, 1344 (1980) (Experimental X-ray energies)
- 1981Hs02 H. -C. Hseuh, E. -M. Franz, P. E. Haustein, S. Kateoff, L. K. Peker, *Phys. Rev.* C23, 1217 (1981) (Gamma ray energies and probabilities)
- 1982Ba56 G. Barreau, H. G. Borner, T. von Egidy, R. W. Hoff, *Z. Phys.* A308, 209 (1982) (Experimental X-ray energies)
- 1982Po14 V. G. Polyukhov, G. A. Timofeev, V. V. Kalygin, P. A. Privalova, *Sov. Radiochem.* 24, 408 (1982); *Radiokhimiya* 24, 490 (1982) (Half-life)
- 1984BuZJ P. A. Burns, P. N. Johnston, J. R. Moroney, *Priv. Comm.* (1984), quoted in 1986LoZT, p.147 (α -transition probabilities)
- 1990Pe03 J. Pearcey, S. A. Woods, P. Christmas, *Nucl. Instrum. Methods Phys. Res.* A286, 563 (1990) (Conversion electron measurements, gamma ray multipolarities)
- 1990Po14 Yu. S. Popov, I. B. Makarov, D. Kh. Srurov, E. A. Erin, *Radiokhimiya* 32, 2 (1990); *Sov. J. Radiochemistry* 32, 425 (1990) (Experimental relative LX-ray emission probabilities)
- 1991Ry01 A. Rytz, *At. Data Nucl. Data Tables.* 47, 205 (1991) (Alpha-emission energies)
- 1992Fr04 E. A. Frolov, *Appl. Radiat. Isot.* 43, 211 (1992) (α -emission energies)
- 1993Pa29 A. K. Pandey, R. C. Sharma, P. C. Kalsi, R. H. Iyer, *Nucl. Instrum. Methods Phys. Res.* B82, 151 (1993) (SF half-life)
- 1995Jo23 P. N. Johnston, P. A. Burns, *Nucl. Instrum. Methods Phys. Res.* A361, 229 (1995) (LX-ray emission probabilities)
- 1996Bu50 C. C. Bueno, J. A. C. Goncalves, M. Damy de S. Santos, *Nucl. Instrum. Methods Phys. Res.* A371, 460 (1996) (α -transition probabilities)
- 1996Sa24 A. M. Sanchez, P. R. Montero, F. V. Tome, *Nucl. Instrum. Methods Phys. Res.* A369, 593 (1996) (α -transition probabilities)
- 1996Sc06 E. Schönfeld, H. Janßen, *Nucl. Instrum. Methods Phys. Res.* A369, 527 (1996) (Atomic data)
- 1997Ka59 J. Kasagi, H. Yamazaki, N. Kasajima, T. Ohtsuki, H. Yuki, *J. Phys.(London)* G23, 1451 (1997) (α -transition probabilities)
- 1998Ga19 E. Garcia-Torano, *Appl. Radiat. Isot.* 49, 1325 (1998) (α -transition probabilities, α -emission energies)
- 1998Ya17 J. Yang, *J. Ni Nucl. Instrum. Methods Phys. Res.* A413, 239 (1998) (α -transition probabilities)
- 1999Schönfeld E. Schönfeld, G. Rodloff - PTB-6.11-1999-1999-1, Braunschweig, February 1999 (KX ray energies and relative emission probabilities)
- 2000Ho27 N. E. Holden, D. C. Hoffman, *Pure Appl. Chem.* 72, 1525 (2000) (Spontaneous fission half-life)
- 2000Schönfeld E. Schönfeld, H. Janßen, *Appl. Rad. Isot.* 52, 595 (2000) (X-ray and Auger electron emission probabilities, EMISSION code)
- 2002Da21 F. Dayras, *Nucl. Instrum. Methods Phys. Res.* A490, 492 (2002) (α -transition probabilities)
- 2003Au03 G. Audi, A. H. Wapstra, C. Thibault, *Nucl. Phys.* A729, 337 (2003) (Q value)
- 2004Ch64 F. E. Chukreev, Balraj Singh, *Nuclear Data Sheets* 103, 325 (2004) (Decay scheme, ²⁴⁰Pu level energies, gamma ray multipolarities and probabilities)
- 2006Ch34 V. P. Chechev, *Phys. Atomic Nuclei* 69, 1188 (2006) (²⁴⁴Cm decay data evaluation-2005)
- 2008Ki07 T. Kibedi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, C. W. Nestor, Jr., *Nucl. Instrum. Methods Phys. Res.* A589, 202 (2008) (Theoretical ICC)

**²⁴⁵Cm -Comments on evaluation of decay data
by V.P. Chechev**

Evaluated in November 2010 with a literature cut-off by the same date.

1. DECAY SCHEME

²⁴⁵Cm decays 100 % to levels of ²⁴¹Pu by emission of α particles and, with a very small branch of $5.9(9) \times 10^{-7}$ % by spontaneous fission. The adopted ²⁴¹Pu levels populated in the ²⁴⁵Cm α decay are based generally on the evaluation by Martin (2005Ma88). Questionable ²⁴¹Pu levels with energies of 260.5 and ≈ 376 keV as reported from α spectrometric measurements of 1975Ba65 were not included into the current evaluation. The 260.5-keV nuclear level was judged in 1975Ba65 as belonging possibly to ²⁴³Am α decay, while the 376-keV nuclear level was not identified by 1975Ba65 and may belong to ²³⁹Pu α decay along with the ≈ 384 -keV energy level. However, the latter has been identified as 13/2⁺ belonging to the ²⁴¹Pu 7/2 [624] rotational band populated in the ²⁴⁵Cm α decay (1975Ba65), and therefore has been included in the proposed decay scheme.

The decay scheme overall consistency is supported by the agreement between $Q(\text{calc}) = 5640(30)$ keV, deduced from the evaluated average energies and intensities of all emissions, and $Q(\alpha) = 5622.3(5)$ keV, deduced from measured α -particle energies. Percentage deviation of $Q(\text{calc})$ from the adopted $Q(\alpha)$ and the $Q(\alpha)$ value of Audi *et al.* (2003Au03) is $-(0.3 \pm 0.5)$ %.

2. NUCLEAR DATA

$Q(\alpha)$ value has been deduced from the five alpha transition energies obtained from the α particle energies measured in 1975Ba65 and adjusted for changes in calibration energies by Rytz (1991Ry01). This approach (Table 1) similar to the evaluation by Martin (2005Ma88) is due to absence of an adjusted $Q(\alpha)$ value for ²⁴⁵Cm in 2003Au03. Audi *et al.* (2003) chose $Q(\alpha) = 5623(1)$ keV reported in 1975Ba65.

Table 1. ²⁴⁵Cm $Q(\alpha)$ values deduced from α -transition energies

Energy of ²⁴¹ Pu level (keV)	Energy of α particles (keV) (experimental)	Energy of α -transition (keV)	Deduced $Q(\alpha)$ value (keV)
41.9722 (9)	$\alpha_{0,1}$ 5488.5 (5)	5579.7 (5)	5621.7 (5)
95.7795 (12)	$\alpha_{0,2}$ 5436.1 (5)	5526.4 (5)	5622.2 (5)
175.0523 (14)	$\alpha_{0,5}$ 5361.8 (12)	5450.9 (12)	5625.9 (12)
231.935 (9)	$\alpha_{0,6}$ 5303.6 (12)	5391.7 (12)	5623.6 (12)
301.172 (16)	$\alpha_{0,7}$ 5234.4 (12)	5321.4 (12)	5622.5 (12)

The weighted average of the deduced $Q(\alpha)$ data set is 5622.3 keV, the internal uncertainty is 0.31, the external uncertainty is 0.55, $\chi^2/\nu = 3.05$, χ^2/ν (critical) = 3.30. The smallest value of experimental uncertainties is ± 0.5 keV. The recommended $Q(\alpha)$ value is **5622.3 (5) keV**.

The ²⁴⁵Cm half-life is based on the experimental results given in Table 2.

Table 2. Experimental values of ²⁴⁵Cm half-life (in 10³ years)

Reference	Author(s)	Original value	Re-estimated value	Comments
1954Hu50	Hulet <i>et al.</i>	≈ 20		Not used
1954Fr19	Friedman <i>et al.</i>	11.5 (50)	11.3 (50) ^a	Relative specific activity to ²⁴⁴ Cm - not used.
1955Br02	Browne <i>et al.</i>	14.3 (29)		α counting - not used.
1957Hu76	Huizenga <i>et al.</i>	7.5 (19)		H. Diamond, Priv. Com. no details - outlier.
1961Ca01	Carnall <i>et al.</i>	9.32 (28)	9.60 (29) ^a	Relative specific activity to ²⁴⁴ Cm - outlier.
1969Me01	Metta <i>et al.</i>	8.265 (180)	8.270 (180) ^a	Relative specific activity to ²⁴⁴ Cm
1971Ma32	MacMurdo <i>et al.</i>	8.532 (53)	8.537 (71) ^{a,b}	Relative specific activity to ²⁴⁴ Cm
1982Po14	Polyukhov <i>et al.</i>	8.445 (100)	8.450 (100) ^a	Relative specific activity to ²⁴⁴ Cm
2009KoZV	Kondev <i>et al.</i>	8.245 (70)		Daughter in-growth from ²⁴⁹ Cf sample

^a Re-estimated by the evaluator on the basis of the recommended ²⁴⁴Cm half-life of 18.11 (3) years.

^b Uncertainty has been revised in 1989Ho24.

In the six values adopted in the data analysis, the LWEIGHT computer program identified two outliers (1957Hu76 and 1961Ca01), and indicated that the four remaining experimental values are discrepant: there are two separate groups of measured values at 8.5×10^3 and 8.25×10^3 years. A similar situation for measurements of ²³⁹Pu half-life by the specific activity method (24 400 and 24 100 years) was resolved to the benefit of the lower value on the basis of the detected presence of impurities leading to overestimations of half-life. This method involves the determination of the number of atoms and disintegration rate of the radionuclide with good accuracy, and thereby requires absolute efficiencies (2009KoZV). Daughter growth in a sample in which the parent is shorter lived does not require such efficiencies, and has been successfully adopted recently to determine the ²⁴⁰Pu half-life (2008KoZP). Therefore, for statistical processing the evaluator has chosen two consistent experimental results obtained with different methods: 1969Me01 (re-estimated) and 2009KoZV, and omitted the other two measurements. The weighted average for this limited set of only two measurements is 8.25×10^3 years with an internal uncertainty of 0.065 and external uncertainty of 0.0084 ($\chi^2/\nu = 0.02$).

The recommended value for the ²⁴⁵Cm half-life is **8.25 (7) $\times 10^3$ years**.

A value of $1.4 (2) \times 10^{12}$ years was adopted for ²⁴⁵Cm spontaneous fission (SF) half-life from the measurement of 1985Dr10. SF branching of $5.9 (9) \times 10^{-7}$ % has been derived from the adopted SF half-life and total half-life of $8.25 (7) \times 10^3$ years.

2.1. Alpha Transitions

The energies of alpha transitions $\alpha_{0,1}$, $\alpha_{0,2}$, $\alpha_{0,5}$, $\alpha_{0,6}$ and $\alpha_{0,7}$ have been obtained from the experimental α particle energies taking into account the recoil energies for ²⁴¹Pu (Table 1). The energies of the remaining alpha transitions have been obtained from Q(α) value and ²⁴¹Pu level energies given in Table 3 from the Adopted Levels, Gammas of 2005Ma88 where they were deduced from a least-squares fit to gamma-ray energies.

Table 3. ²⁴¹Pu levels populated in ²⁴⁵Cm α -decay

Level	Energy (keV)	Spin and parity	Half-life	Probability of α - transition (%)
0	0.0	5/2+	14.33 (4) a	0.58
1	41.9722 (9)	7/2+		0.83
2	95.7795 (12)	9/2+		0.04
3	161.314 (4)	11/2+		0.39 (22)
4	161.6852 (9)	1/2+	0.88 (5) μ s	0.0210 (9)
5	175.0523 (14)	7/2+		93.2 (5)
6	231.935 (9)	9/2+		5.0 (1)
7	301.172 (16)	11/2+		0.32
8	385 (3)	(13/2+)		≤ 0.005

The experimental values for the α -transition probabilities of ²⁴⁵Cm from spectrometric measurements are presented in Table 4. Uncertainties were not reported in the cited references, but these for the most intense α -transitions $\alpha_{0,5}$ (5362 keV) and $\alpha_{0,6}$ (5304 keV) observed in 1975Ba65 were estimated in 1976BaZZ. The data of 1966Ba07 for ²⁴⁵Cm are not given in Table 4 as those were superseded in 1975Ba65 by the same group. The probabilities of the α -transitions $\alpha_{0,3}$ and $\alpha_{0,4}$ (observed as a doublet with an energy of ~ 5370 keV) have been deduced from intensity balances at the ²⁴¹Pu levels “3” (161.3 keV) and “4” (161.7 keV), respectively. Probabilities of the remaining alpha transitions have been adopted from the magnetic spectrometer measurements of 1975Ba65.

Table 4. Experimental and recommended probabilities (per 100 decays) of alpha transitions observed in ²⁴⁵Cm α decay

	α -particle energy	1960As11	1963Dz07	1966Fr03	1975Ba65	Deduced from P(γ +ce) balance	Recommended
$\alpha_{0,0}$	5529			1.1	0.58		0.58
$\alpha_{0,1}$	5488			0.9	0.83		0.83
$\alpha_{0,2}$	5436			0.2	0.04		0.04
$\alpha_{0,3}$	5372					0.39 (22)	0.39 (22)
$\alpha_{0,4}$	5371					0.0210 (9)	0.0210 (9)
$\alpha_{0,5}$	5362	93	90	91	93.2 (5)	95.3 (21)	93.2 (5)
$\alpha_{0,6}$	5303	7	7	6.2	5.0 (1)		5.0 (1)
$\alpha_{0,7}$	5234		2	0.5	0.32		0.32
$\alpha_{0,8}$	5152				≤ 0.005		≤ 0.005

The α decay hindrance factors have been calculated using the ALPHAD computer program from the ENSDF evaluation package with $r_0(^{241}\text{Pu}) = 1.4969$ (12) fm (2005Ma88).

2.2. Gamma Transitions and Internal Conversion Coefficients

The recommended energies of the gamma-ray transitions are the same as those of the gamma-ray energies with correction to the minor nuclear recoil for ²⁴¹Pu.

The gamma-ray transition probabilities ($P_{\gamma+ce}$) have been deduced from their evaluated gamma-ray emission probabilities (P_γ) and total internal conversion coefficients (ICCs), apart from $P_{\gamma+ce}$ values for the gamma-ray transitions $\gamma_{6,5}$ (56.9 keV) and $\gamma_{7,6}$ (69.2 keV). The latter values have been deduced directly from intensity balances at the ²⁴¹Pu levels “6” (231.9 keV) and “7” (301.2 keV), respectively.

ICCs have been interpolated using the BrIcc computer program, version v2.2a, data set BrIccFO (2008Ki07). Multipolarities of the gamma-ray transitions and E2/M1 mixing ratios (δ) are based on the measurements of conversion electrons (ce) in the ²⁴⁰Pu (n, γ)-reaction and have been taken from 2005Ma88, except as noted below.

The δ values for $\gamma_{6,5}$ (56.9 keV) and $\gamma_{7,6}$ (69.2 keV) have been obtained from the total ICC deduced using the expression $1 + \alpha_T = P_{\gamma+ce} / P_\gamma$.

3. ATOMIC DATA

The fluorescence yields, X-ray energies and relative probabilities, and Auger electrons energies and relative probabilities are from the SAISINUC software.

4. ALPHA EMISSIONS

The recommended energies of alpha particles for the five transitions ($\alpha_{0,1}$, $\alpha_{0,2}$, $\alpha_{0,5}$, $\alpha_{0,6}$, $\alpha_{0,7}$) used for obtaining $Q(\alpha)$ have been adopted from the most precise measurements of 1975Ba65. The remaining α particle energies have been deduced from the $Q(\alpha)$ value, taking into account the recoil energies for ²⁴¹Pu.

The recommended α -particle energies are compared in Table 5 with the experimental results from spectrometric measurements (1960As11, 1963Dz07, 1966Fr03 and 1975Ba65).

Table 5. Experimental and recommended α -particle energies (keV) in the decay of ²⁴⁵Cm^a

	1960As11	1963Dz07	1966Fr03	1975Ba65	Recommended
$\alpha_{0,0}$			5530 (3)	5529.0 (5)	5530.4 (5)
$\alpha_{0,1}$			5497 (5)	5488.5 (5)	5488.5 (5)
$\alpha_{0,2}$			5447 (5)	5436.1 (5)	5436.1 (5)
$\alpha_{0,3}$				5370	5371.7 (5)
$\alpha_{0,4}$				5370	5371.4 (5)
$\alpha_{0,5}$	5360	5361	5359 (2)	5361.8 (12)	5361.8 (12)
$\alpha_{0,6}$	5305	5305	5306 (2)	5303.6 (12)	5303.6 (12)
$\alpha_{0,7}$		5245	5239 (3)	5234.4 (12)	5234.4 (12)
$\alpha_{0,8}$				\approx 5151	5152 (3)

^a Authors' experimental values have been adjusted for changes in calibration energies, as suggested in 1991Ry01.

5. ELECTRON EMISSIONS

The energies of the conversion electrons have been obtained from the gamma-ray transition energies and the atomic electron binding energies from 1977La19.

The emission probabilities of the conversion electrons have been deduced using the evaluated P_γ and ICC values. Measurements of the ²⁴¹Pu conversion electrons following thermal neutron capture in ²⁴⁰Pu were carried out by 1998Wh01.

The total absolute emission probabilities of K and L Auger electrons have been calculated using the EMISSION computer program (1996Sc06, 2000Sc47).

6. PHOTON EMISSIONS

6.1 X - Ray emissions

The absolute emission probabilities of Pu KX- and LX-rays were calculated using the EMISSION computer program (Table 6). In 1980Di13 the emission probabilities of Pu KX-rays were measured relatively to P_γ ($\gamma_{5,0}$ 175.0 keV). The experimental absolute P(KX) values are given in Table 6 using the evaluated P_γ ($\gamma_{5,0}$ 175.0 keV) = 9.83 (22) %.

Table 6. Experimental (1980Di13) and calculated absolute Pu KX-ray emission probabilities (%)

	1980Di13	Calculated
K α_2 (Pu)	20.0 (7)	19.0 (5)
K α_1 (Pu)	30.9 (11)	30.1 (7)
K β'_1 (Pu)	11.1 (5)	11.1 (3)
K β'_2 (Pu)	3.7 (2)	3.84 (12)

The good agreement between measured and calculated KX-ray emission probabilities supports the recommended γ -ray emission probabilities and assigned multiplicities.

6.2. Gamma emissions

6.2.1. Gamma-ray energies

The gamma-ray energies (E_γ) have been taken from 2005Ma88 (²⁴¹Pu, Adopted Levels, Gammas). They are based mainly on measurements of γ rays from ²⁴⁵Cm α decay by 1980Di13, 1992Daniels, 1994Sh31, 1998Wh01 and γ rays from thermal neutron capture in ²⁴⁰Pu by 1998Wh01 (Table 7). The gamma-ray energies for $\gamma_{7,6}$ (69.2 keV), $\gamma_{5,2}$ (79.3 keV), $\gamma_{2,0}$ (95.8 keV), $\gamma_{6,2}$ (136.1 keV), $\gamma_{7,2}$ (205.4 keV), and $\gamma_{6,0}$ (231.9 keV) have been deduced directly from the adopted ²⁴¹Pu level energies. Other, less accurate measurements of E_γ can be found in 1955Pe32, 1966Ba07, 1991Po17.

6.2.2. Gamma-ray emission probabilities

The relative gamma-ray emission probabilities (I_γ) are weighted averages of the experimental values from 1980Di13 and 1992Daniels, except as noted otherwise in Table 7. The LWEIGHT computer program was used for statistical processing, with the uncertainty assigned to the average value always greater than or equal to the smallest uncertainty of the values used to calculate the average.

The normalization factor (N) was obtained from the intensity balance to the ground state of ^{241}Pu :

$$\Sigma(1+\alpha_T)I_\gamma(\gamma_{1,0}, \gamma_{2,0}, \gamma_{4,0}, \gamma_{5,0}, \gamma_{6,0}) + P(\alpha_{0,0}) = 1$$

$$\text{assuming } P(\alpha_{0,0}) = 0.006(1) \quad (1975\text{Ba65}, 2005\text{Ma88}),$$

$$N = P_\gamma(175.0 \text{ keV}) = 0.0983(22).$$

This adopted value agrees with the directly measured $P_\gamma(175.0 \text{ keV})$ of 0.095(7) (1980Di13) and 0.101(1) (1992Daniels).

The absolute gamma-ray emission probabilities (P_γ) have been deduced from the evaluated relative gamma-ray emission probabilities (Table 7) using the derived normalization factor of 0.0983(22).

Table 7. Experimental (E_γ^{exp}) and recommended gamma-ray energies and experimental and evaluated relative emission probabilities (I_γ^{exp}) in the decay of ^{245}Cm

	E_γ^{exp} from ^{245}Cm decay	Recommended E_γ (keV)	I_γ^{exp} (1980Di13)	I_γ^{exp} (1992Daniels)	I_γ^{exp} (1998Wh01)	Evaluated I_γ
$\gamma_{1,0}$	41.93 (3) ^a	41.972 (1)	3.68 (18)	4.10 (39)		3.75 (18) ^a
$\gamma_{2,1}$	53.72 (4) ^a	53.807 (1)	0.70 (4)	0.77 (4)		0.74 (4) ^a
$\gamma_{6,5}$	56.89 (3) ^b	56.89 (3)	0.38 (2)	0.325 (32)		0.365 (20) ^a
$\gamma_{3,2}$	65.44 (8) ^a	65.535 (3)	0.12 (4)	0.20 (2)		0.18 (2) ^a
$\gamma_{7,6}$	69.17 (6) ^c	69.237 (18)	0.07 (3)	-		0.07 (3)
$\gamma_{5,2}$	79.27 (4) ^a	79.2728 (18)	1.58 (9)	1.22 (7)		1.22 (7) ^d
$\gamma_{2,0}$	95.786 (3) ^{b, d}	95.7795 (12)				0.111 (23) ^c
$\gamma_{7,5}$	126.09 (4) ^b	126.09 (4)			0.07 (2)	0.07 (2) ^b
$\gamma_{5,1}$	133.05 (8) ^a	133.081 (2)	29.2 (15)	28.6 (4)		28.6 (4) ^a
$\gamma_{6,2}$	136.127 (20) ^b	136.156 (9)	1.18 (7)	1.15 (3)		1.15 (3) ^a
$\gamma_{7,3}$	139.87 (4) ^b	139.858 (16)	0.06 (2)	0.06 (3)	0.09 (1)	0.08 (9) ^f
$\gamma_{4,0}$	161.72 (8) ^a	161.685 (1)	0.09 (4)	0.067 (2)		0.072 (2) ^a
$\gamma_{5,0}$	175.01 (9) ^a	175.0523 (14)	100	100	100	100
$\gamma_{6,1}$	189.965 (10) ^b	189.965 (10)	2.03 (13)	2.07 (4)		2.07 (4) ^a
$\gamma_{7,2}$	205.404 (20) ^a	205.393 (16)	-	0.115 (19)	0.08 (1)	0.09 (1) ^g
$\gamma_{6,0}$	231.96 (3) ^b	231.935 (9)	0.16 (4)	0.117 (18)	0.11 (2)	0.119 (18) ^h
$\gamma_{-1,1}$	388.16 (5) ^b	388.16 (5)			0.19 (1)	0.19 (1) ^b

^a Weighted averages of experimental values from 1980Di13 and 1992Daniels (see also 1994Sh31, 2005Ma88).

^b Experimental value from 1998Wh01.

^c Reported only in 1980Di13, but also adopted in the level scheme of 1994Sh31.

^d From 1992Daniels; higher value leads to a large intensity imbalance at the 96-keV and 175-keV levels

^e Obscured by the Pu $K\alpha_2$ X-ray; I_γ is from $I_\gamma/I_\gamma(53.8 \text{ keV}) = 0.15(3)$ in Adopted Gammas (2005Ma88).

^f Weighted averages of experimental values from 1980Di13, 1992Daniels and 1998Wh01.

^g Weighted averages of experimental values from 1992Daniels and 1998Wh01.

^h Weighted averages of experimental values from 1980Di13, 1992Daniels and 1998Wh01.

7. REFERENCES

- 1954Fr19 A.M. Friedman, A.L. Harkness, P.R. Fields, M.H. Studier, J.R. Huizenga, Phys. Rev. 95, 1501 (1954)
(Half-life)
- 1954Hu50 E.K. Hulet, S.G. Thompson, A. Ghiorso, Phys. Rev. 95, 1703 (1954)
(Half-life)
- 1955Br02 C.I. Browne, D.C. Hoffman, W.T. Crane, J.P. Balagna, G.H. Higgins, J.W. Barnes, R.W. Hoff, H.L. Smith, J.P. Mize, M.E. Bunker, J. Inorg. Nucl. Chem. 1, 254 (1955)
(Half-life)
- 1955Pe32 I. Perlman, F. Asaro, F.S. Stephens, J.P. Hummel, UCRL-2932, p.59 (1955)
(γ -ray energies)
- 1957Hu76 J.R. Huizenga, H. Diamond, Phys. Rev. 107, 1087 (1957)
(Half-life)
- 1960As11 F. Asaro, I. Perlman, Private Communication (1960), quoted by 1964Hy02
(α -particle energies and emission probabilities)
- 1961Ca01 W.T. Carnall, S. Fried, A.L. Harkness, J. Inorg. Nucl. Chem. 17, 12 (1961)
(Half-life)
- 1963Dz07 B.S. Dzhelapov, R.B. Ivanov, V.G. Nedovesov, V.P. Chechev, Zh. Eksperim. i Teor. Fiz. 45, 1360 (1963); Sov. Phys. JETP 18, 937 (1964)
(α -particle energies and emission probabilities)
- 1964Hy02 E.K. Hyde, I. Perlman, G.T. Seaborg, The Nuclear Properties of the Heavy Elements, Vol. II, Prentice-Hall, Inc., Englewood Cliffs, N.J. (1964)
(α -particle energies and emission probabilities)
- 1966Ba07 S.A. Baranov, Y.F. Rodionov, V.M. Kulakov, V.M. Shatinskii, Yad. Fiz. 4, 1108 (1966); Sov. J. Nucl. Phys. 4, 798 (1967)
(α -particle energies and emission probabilities)
- 1966Fr03 A.M. Friedman, J. Milsted, Phys. Lett. 21, 179 (1966)
(α -particle energies and emission probabilities)
- 1969Me01 D.N. Metta, H. Diamond, F.R. Kelly, J. Inorg. Nucl. Chem. 31, 1245 (1969)
(Half-life)
- 1971Ma32 K.W. MacMurdo, R.M. Harbour, R.W. Benjamin, J. Inorg. Nucl. Chem. 33, 1241 (1971)
(Half-life)
- 1975Ba65 S.A. Baranov, V.M. Shatinsky, Yad. Fiz. 22, 670 (1975); Sov. J. Nucl. Phys. 22, 346 (1976)
(α -particle energies and emission probabilities)
- 1976BaZZ S.A. Baranov, A.G. Zelenkov, V.M. Kulakov, Proc. Advisory Group Meeting on Transactinium Nuclear Data, Karlsruhe, IAEA technical report IAEA-186, Vol. III, p.249 (1976)
(α -particle energies and emission probabilities)
- 1977La19 F.P. Larkins, At. Data Nucl. Data Tables 20, 311 (1977)
(Atomic electron binding energies)
- 1980Di13 J.K. Dickens, J.W. McConnell, Phys. Rev. C22, 1344 (1980)
(γ -ray energies and emission probabilities)
- 1982Po14 V.G. Polyukhov, G.A. Timofeev, V.V. Kalygin, P.A. Privalova, Radiokhimiya 24, 490 (1982); Sov. Radiochemistry 24, 408 (1982)
(Half-life)
- 1985Dr10 A.A. Druzhinin, V.N. Polynov, S.P. Vesnovsky, A.M. Korochkin, A.A. Lbov, E.A. Nikitin, Dok. Akad. Nauk SSSR 280, 1351 (1985)
(SF half-life)

- 1989Ho24 N.E. Holden, Pure Appl.Chem. 61, 1483 (1989)
(Half-life)
- 1991Po17 Yu.S. Popov, D.Kh. Srurov, I.B. Makarov, E.A. Erin, G.A. Timofeev, Radiokhimiya 33, 3 (1991); Sov. J. Radiochemistry 33, 1 (1991)
(γ -ray energies and emission probabilities)
- 1991Ry01 A. Rytz, At. Data Nucl. Data Tables 47, 205 (1991)
(α -particle energies and emission probabilities)
- 1992Daniels R. Daniels, PhD thesis, University of Manchester, 1992, quoted by 1994Sh31
(γ -ray energies and emission probabilities)
- 1994Sh31 J.A. Shannon, W.R. Phillips, B.J. Varley, I. Ahmad, L.R. Morss, Nucl. Instrum. Methods Phys. Res. A339, 183 (1994)
(γ -ray energies and emission probabilities)
- 1996Sc06 E. Schönfeld, H. Janssen, Nucl. Instrum. Methods Phys. Res. A369, 527 (1996)
(Atomic data)
- 1998Wh01 D.H. White, R.W. Hoff, H.G. Borner, K. Schreckenbach, F. Hoyler, G. Colvin, I. Ahmad, A.M. Friedman, J.R. Erskine, Phys. Rev. C57, 1112 (1998)
(γ -ray energies and emission probabilities)
- 2000Sc47 E. Schönfeld, H. Janssen, Appl. Radiat. Isot. 52, 595 (2000)
(Calculation of emission probabilities of X-rays and Auger electrons)
- 2003Au03 G. Audi, A.H. Wapstra, and C. Thibault, Nucl. Phys. A729, 337 (2003)
(Q value)
- 2005Ma88 M.J. Martin, Nucl. Data Sheets 106, 89 (2005)
(²⁴⁵Cm α -decay scheme, ²⁴¹Pu levels, γ -ray energies and multiplicities)
- 2008Ki07 T. Kibédi, T.W. Burrows, M.B. Trzhaskovskaya, P.M. Davidson, C.W. Nestor, Jr., Nucl. Instrum. Methods Phys. Res. A589, 202 (2008)
(Band-Raman ICC for γ -ray transitions)
- 2008KoZP F.G. Kondev, M.A. Kellett, I. Ahmad, J.P. Greene, A.L. Nichols, Proc. Int. Conf. Nuclear Data for Science and Technology, Nice, France, 22-27 April 2007, O. Bersillon, F. Gunsing, E. Bauge, R. Jacqmin, and S. Leray, Eds., p.93 (2008) EDP Sciences
(Half-life)
- 2009KoZV F.G. Kondev, I. Ahmad, M.P. Carpenter, C.J. Chiara, J.P. Greene, R.V.F. Janssens, M.A. Kellett, T.L. Khoo, T. Lauritsen, C.J. Lister, E.F. Moore, A.L. Nichols, D. Seweryniak, S. Zhu, Proc. 13th Int. Symposium on Capture Gamma-Ray Spectroscopy and Related Topics, Cologne, Germany, 25-29 August 2008, J. Jolie, A. Zilges, N. Warr, A. Blazhev, Eds., AIP Conf. Proc. 1090, p.199 (2009)
(Half-life)

**²⁴⁶Cm - Comments on evaluation of decay data
by F.G. Kondev**

This evaluation was completed in December 2006 with a literature cut off by the same date. The Saisinuc software (2002BeXX) and associated supporting programs were used in assembling the data following the established protocol within the DDEP collaboration.

1. Decay Scheme

The deformed ²⁴⁶Cm nucleus disintegrates by α emissions and spontaneous fission. The strongest α -decay branch populates the ground state of the daughter nuclide ²⁴²Pu, which is also deformed. The level schemes of ²⁴²Pu and ²⁴⁶Cm are based on the evaluations of Akovali (2002Ak06) and Artna -Cohen (1998Ar12), respectively. The recent experimental work of Kondev *et al.* (2007Ko01) reported a weak α -decay branch to the 4⁺ level of the ground-state band of ²⁴²Pu.

2. Nuclear Data

$Q(\alpha)$ value is obtained from the adopted $E_{\alpha,0}$ energy (see section 2.1 for details) and by taking into account the relevant recoil energy. This value differs from that of 5475.1 (9) keV (2003Au03), deduced as a weighted mean of $Q(\alpha) = 5475.2$ (10) keV and 5474.9 (20) keV, which were determined from the $E_{\alpha,0}$ energies of 1984Sh31 and 1966Ba07, respectively. It should be noted that no uncertainty to the $E_{\alpha,0}$ value was reported in the original publication of 1966Ba07, but it was assigned by 2003Au03.

The experimental data on α/SF and $T_{1/2 SF}$, together with results from the earlier evaluation of Holden (2000Ho27), are presented in Table 1.

Table 1. Experimental and evaluated data for the α/SF ratio and the SF half-life of ²⁴⁶Cm

Author	α/SF	$T_{1/2SF}$, (10 ⁷ a)	Method	Used in the evaluation
1956Fi11	2740 (140)	> 1.24	From α/SF	No
1956FrXX		2.0 (8)	relative to ²⁴⁶ Pu weight and the α -counting technique	No
1965Me02	0.139 (9) 10 ^{6 a)}	1.66 (10)	relative to ²⁴⁴ Cm α -decay data ^{b)}	No
1969Me01	3822 (10)	1.80 (1)	From α/SF	Yes
1971Ma32	3833 (32)	1.85 (2)	From α/SF	Yes
2000Ho27		1.81 (2)	Evaluated value	No

a) Net (²⁴⁶Cm fissions)/(²⁴⁴Cm α -disintegrations).

b) Using $T_{1/2,\alpha}(\text{}^{244}\text{Cm}) = 18.11$ (7) a, mole ratio (²⁴⁴Cm/²⁴⁶Cm) = 7.82 (9) and (²⁴⁶Cm fissions)/(²⁴⁴Cm α -disintegrations) = 0.139 (9) 10⁶.

The % α and %SF values were deduced using $\alpha/SF = 3823$ (10), a weighted mean of 3822 (10) (1969Me01) and 3833 (32) (1971Ma32):

$$\%SF = \frac{1}{1 + \alpha / SF} \times 100, \text{ with } \%a = 100 - \%SF \quad (1)$$

Then %SF = 0.02615 (7) % and % α = 99.97385 (7) %

The mean number of neutrons emitted by spontaneous fission is: 2.948 (from ENDF/B-VII)

The recommended partial SF half-life of $T_{1/2 SF} = 1.81 (2) 10^7$ a, was determined as a weighted mean of 1.80 (1) 10^7 a (1969Me01) and 1.85 (2) 10^7 a (1971Ma32).

The experimental data for the partial α -decay half-life of ²⁴⁶Cm are presented in Table 2.

Table 2. Experimental data for the partial α -decay half-life of ²⁴⁶Cm

Author	Method ^{a)}	$T_{1/2 a, (a)}$ ^{b)}	$T_{1/2 a, (a)}$ ^{c)}	$T_{1/2 a, (a)}$ ^{d)}	Used in the evaluation
1954Fr19	RSA to ²⁴⁴ Cm	4000 (600)	18.44 (5)	3928 (589)	No
1955Br02	IA to ²⁴⁶ Pu	2300 (460)			No
1956Bu91	IA to ²⁵⁰ Cf	6620 (320)	9.3 (9)	9311 (623)	No
1961Ca01	RSA to ²⁴⁴ Cm	5480 (170)	17.59 (6)	5642 (175)	No
1969Me01	RSA to ²⁴⁴ Cm	4711 (22)	18.099 (15)	4714 (22)	Yes
1971Mc19	ASA	4654 (40)			Yes
1971Ma32	RSA to ²⁴⁴ Cm	4820 (20)	18.099 (15)	4823 (20)	Yes
1977Po20	RSA to ²⁴⁴ Cm	4852 (76)	18.099 (15)	4855 (76)	Yes
2007Ko01	IA to ²⁵⁰ Cf	4706 (40)	13.08 (9)		Yes

^{a)} RSA-relative specific activity method; ASA – absolute specific activity method; IA in -growth activity method.

^{b)} Value reported in the original publication.

^{c)} Half-life value for the reference ²⁴⁴Cm or ²⁵⁰Cf nuclide used in the original publication.

^{d)} Corrected ²⁴⁶Cm half-life values using $T_{1/2}(\text{²⁴⁴Cm}) = 18.11 (3) \text{ a}$ (2005ChXX) and $T_{1/2}(\text{²⁵⁰Cf}) = 13.08 (9) \text{ a}$ (2001Ak11)

Since in all cases, except 1971Mc19, relative methods were used to deduce $T_{1/2 \alpha}$, the values reported in the original publications were corrected using the most recently adopted $T_{1/2 \alpha}$ of the reference nuclides ²⁴⁴Cm and ²⁵⁰Cf, as summarized in Table 2. Results from the early work of 1954Fr19, 1955Br02, 1956Bu91 and 1961Ca01 are inaccurate and discrepant (with half-life values spanning between 2300 (460) a and 9311 (623) a), and hence, these data were excluded from the present analysis.

Although the remaining five $T_{1/2 \alpha}$ values have better accuracy, these data are also discrepant. For example, while the data of 1969Me01, 1971Mc19 and 2007Ko01 give a weighted mean of $T_{1/2 \alpha} = 4701 (17) \text{ a}$, the results of 1971Ma32 and 1977Po20 are clustered around the weighted mean value of $T_{1/2 \alpha} = 4825 (19) \text{ a}$. In the present work, detailed evaluations of $T_{1/2 \alpha}$ were carried out using specially developed techniques that deal with discrepant data (see references 1992Ra08, 1994Ka08 and 2004MaXX for example) and the results are presented in Table 3. The weighted mean (WM) value (external uncertainty) is $T_{1/2 \alpha} = 4756 (32) \text{ a}$, but $\chi^2_v = 6.16$ (where $\chi^2_v = \chi^2/N-1$) is larger than the critical value of $\chi^2_{v \text{ crit}} = 3.32$ (99 % confidence level) because the data are discrepant.

The Limitation of Relative Statistical Weight (LRSW) method adopts $T_{1/2 \alpha} = 4756 (67) \text{ a}$, which is the WM value, but the uncertainty is extended in order to include “the most precise” value of 4823 (20) a (1971Ma32) (uncertainty of 0.41 %). It should be noted, however, that the determined by the LRSW method “the most precise” value is as accurate as that of 4714 (22) a (1969Me01) (uncertainty of 0.47 %). Hence, if the value from 1969Me01 is adopted as “the most precise” one, then the LRSW would give $T_{1/2 \alpha} = 4756 (42) \text{ a}$. In the LRSW case, χ^2_v is also larger than $\chi^2_{v \text{ crit}}$. The Normalized Residual Method (NRM) evaluates a value of $T_{1/2 \alpha} = 4723 (27) \text{ a}$, while the Rajeval method (RM) adopts $T_{1/2 \alpha} = 4713 (17) \text{ a}$. In both cases χ^2_v is smaller than $\chi^2_{v \text{ crit}}$.

Table 3. Evaluated values of the half-life of ²⁴⁶Cm.

Method/Author ^{a)}	Evaluated T _{1/2} , (a)	c ² /N-1	
UWM	4750 (38)	6.21	
WM (external)	4756 (32)	6.16	
LRSW	4756 (67)	6.16	
NRM	4723 (27)	2.78	Adopted
RM	4713 (17)	1.24	
1989Ho24	4760 (40)	7.48	
1998Ar12	4760 (40) ^{b)}		

^{a)} UWM – Unweighted Mean; WM – Weighted Mean; LRSW – Limitation of Relative Statistical Weight; NRM – Normalized Residual; RM – Rajeval.

^{b)} Value adopted from 1989Ho24

The NRM value is recommended in the present evaluation since the relative statistical weights of the uncertainties (note that only the uncertainty reported in 1971Ma32 has been adjusted by this method) are less than 50 %, while the RM value (uncertainties of 1971Ma32, 1971Mc19 and 1977Po20 were adjusted by this method) is biased towards that of T_{1/2α} = 4714 (22) a (1969Me01) (with a relative statistical weight of 62 %).

2.1 Alpha Transitions

The ²⁴²Pu level energies were deduced by a least-square fit to the adopted γ-ray energies (see section 2.2 and Table 7 for details) using the computer program GTOL from the ENSDF evaluation package. The α_{0,0} energy was taken from the evaluation of Rytz (1991Ry01), while the α_{0,1} and α_{0,2} energies were obtained from the adopted E_{α0,0} = 5387.5 (9) keV, the 2⁺ and 4⁺ level energies of ²⁴²Pu, respectively, and by taking into account the relevant recoil energies.

Table 4. Experimental and evaluated values of the α-particle energies in decay of ²⁴⁶Cm

Authors	E _{a0,0} , (keV)	E _{a0,1} , (keV)	E _{a0,2} , (keV)	Comment ^{a)}
1963Be48	5387	5345		MS
1963Dz07	5387 (4)	5345 (4)		MS
1966Ba07	5385	5342		MS
1984Sh31	5386.5 (10)	5343.5 (10)		MS
2007Ko01	5386 (3)	5342 (3)	5242 (3)	SD
1991Ry01	5387.5 (9)	5342.7 (9)		evaluated
Adopted	5387.5 (9)	5343.7 (9)	5242.5 (10)	Evaluated

a) MS – magnetic α-spectrometer; SD – semiconductor detector

The experimental values for the α-transition probabilities of ²⁴⁶Cm are presented in Table 5. It should be noted that uncertainties were not reported in the work of 1963Be48 and 1966Ba07, but these were estimated by Rytz (1991Ry01).

Table 6 contains the evaluated P_{α0,0} values using two different data sets, one that excludes values reported without uncertainty in the original publications (“limited data”) and the second that includes all experimental values with uncertainties estimated by Rytz (1991Ry01) in cases where those were missing in the original publications (“all data”). The evaluated values deduced using both data sets are consistent and the WM value from the so-called “all data” set is recommended (χ²_v = 1.69 is smaller than the critical value of χ²_{v crit} = 3.32 (99 % confidence level)). The recommended P_{α0,2} value was deduced using the branching ratios of 2007Ko01 and the adopted here P_{α0,0} = 79.17 (22) %. The P_{α0,1} value was determined as:

$$P_{a0,1} = 100 - P_{a0,0} - P_{a0,2} \quad (2)$$

Table 5. Experimental and evaluated α -transition probabilities in decay of ²⁴⁶Cm.

Authors	P _{a0,0} , (%)	P _{a0,1} , (%)	P _{a0,2} , (%)	Comment ^{a)}
1963Be48	78	22		MS
1963Dz07	78 (5)	22 (5)		MS
1966Ba07	79	21		MS
1984Sh31	82.2 (12)	17.8 (12)		MS
2007Ko01	79.08 (22)	20.9 (4)	0.020 (2)	SD
1991Ry01	80.7 (11) ^{b)}	19.3 (11) ^{b)}		evaluated
Adopted	79.17 (22)	20.81 (22)	0.020 (2)	Evaluated

^{a)} MS – magnetic α -spectrometer; SD – semiconductor detector

^{b)} Rytz (1991Ry01) assigned uncertainties to the original 1963Be48 and 1966Ba07 values as follow: P _{α ,0} = 78 (3) and P _{α ,1} = 22 (3) (1963Be48) and P _{α ,0} = 79 (2) and P _{α ,1} = 21 (2) (1966Ba07).

The α -decay hindrance factors were calculated using the computer program ALPHAD from the ENSDF evaluation package with r₀ = 1.4954 (10) fm.

Table 6. Evaluated P _{α ,0} values in the α -decay of ²⁴⁶Cm

Method/Author ^{a)}	“limited data”		“all data”	
	P _{a0,0} , (keV)	c ² /N-1	P _{a0,0} , (keV)	c ² /N-1
UWM	79.8 (13)		79.26 (78)	
WM	79.18 (22)	3.30	79.17 (22)	1.69
LRSW	79.18 (22)	3.30	79.17 (22)	1.69
NRM	79.15 (22)	2.31	79.17 (22)	1.69
RM	79.10 (22)		79.10 (22)	
1991Ry01			80.7 (11)	

^{a)} UWM – Unweighted Mean; WM – Weighted Mean; LRSW – Limitation of Relative Statistical Weight; NRM – Normalized Residual; RM – Rajeval.

2.2 Gamma-Ray Transitions and Electron Internal Conversion Coefficients

The energy of the 2⁺ → 0⁺ ground state band γ -ray transition of ²⁴²Pu was taken from 1972Sc01. The 4⁺ → 2⁺ γ -ray transition was not observed in the α -decay of ²⁴⁶Cm and its energy was taken from the Coulomb excitation data of 1983Sp03 (note that the uncertainty in this value comes from the work of 1971EiZS). Gamma-ray transition multipolarities were taken from the ENSDF evaluation of 1998Ar12. Since absolute γ -ray emission probabilities were not measured directly for any of the γ -ray transitions that follow α -decay of ²⁴⁶Cm, the absolute transition probabilities, P _{γ +ce}, were deduced from the relative α -transition probabilities, presented in Table 5, after a correction for the α -decay branching was applied:

$$P_{g+ce}(g_{2,0}) = \frac{\%a}{100} \times P_{a0,2} \text{ and } P_{g+ce}(g_{1,0}) = \frac{\%a}{100} \times (P_{a0,1} + P_{a0,2}) \quad (3)$$

The electron internal conversion coefficients were calculated by a program supplied with the Saisinuc software (2002BeXX) that uses interpolated values of Band *et al.* (2002Ba85) with the hole being taken into account.

Table 7. Energies, multipolarities and electron internal conversion coefficients for γ -ray transitions following α -decay of ²⁴⁶Cm

	Energy, (keV)	Multipolarity	α_K	α_L	α_M	α_N	α_O	α_T
$\gamma_{1,0}$	44.545 (9)	E2	-	542 (16)	152 (5)	41.6 (12)	9.8 (3)	746 (22)
$\gamma_{2,1}$	102.8 (1)	E2	-	10.1 (3)	2.82 (8)	0.775 (23)	0.183 (5)	13.9 (4)

3. Atomic Data

The Atomic data (Fluorescence yields, X-Ray energies and Relative probabilities, and Auger electrons energies and Relative probabilities) were provided by the Saisinuc software (2002BeXX). Details regarding the origin of these data can be found in 1996Sc06, 1998ScZM, 1999ScZX, 2000ScXX and 2003DeXX.

4. Alpha Emissions

Details are given in section 2.1. The number of alphas per 100 disintegrations was obtained by multiplying the corresponding α -transition probabilities that are presented in Table 5 by the α -decay branching ratio of 0.999 738 5 (7).

5. Photon Emissions

5.1 X-Ray Emissions

The X-ray emissions per 100 disintegrations were calculated using the computer program EMISSION (2000ScXX).

	Energy, (keV)	(%)
L α	12.125	0.195 (8)
L β	14.083 – 14.279	3.03 (11)
L γ	16.334	0.082 (4)
L η	16.499 – 19.331	3.76 (14)
L ζ	20.708 – 21.984	0.87 (4)

5.2 Gamma-Ray Emissions

The number of γ rays per 100 disintegrations was obtained from the $P_{\gamma+ce}(\gamma_{i,k})$ values, described in section 2.2, and the total electron internal conversion coefficients, $\alpha_T(\gamma_{i,k})$ that are presented in Table 7:

$$P_g(\mathbf{g}_{i,k}) = \frac{P_{\gamma+ce}(\mathbf{g}_{i,k})}{1 + \alpha_T(\mathbf{g}_{i,k})} \quad (4)$$

6. Electron Emissions

The energies of the conversion electrons have been calculated from the γ -ray transition energies presented in Table 7 and the corresponding electron shell binding energies (1977La19). The number of conversion electrons of type $x=T,L,M,N$ and O , where T stands for total, L for L -shell electrons, etc., per 100 disintegrations have been determined from the evaluated numbers of photons per 100 disintegrations, $P_\gamma(\gamma_{i,k})$, and the corresponding electron internal conversion coefficients, $\alpha_x(\gamma_{i,k})$

$$ec_{i,kx} = P_\gamma(\mathbf{g}_{i,k}) \times \alpha_x(\mathbf{g}_{i,k}) \quad (5)$$

The number of L Auger electrons per 100 disintegrations was obtained from the computer program EMISSION (2000ScXX).

7. References

- A.M. Friedman, A.L. Harkness, P.R. Fields, M.H. Studier, J.H. Huizenga – Phys. Rev. 95 (1954) 1501.
- C.I. Browne, D.C. Hoffman, W.T. Crane, J.P. Balagna, G.H. Higgins, J.W. Barnes, R.W. Hoff, H.L. Smith, J.P. Mize, M.E. Bunker – J. Inorg. Nucl. Chem. 1 (1955) 254.
- J.P. Butler, T.A. Eastwood, H.G. Jackson, R.P. Schuman – Phys. Rev. 103 (1956) 965.
- P.R. Fields, M.H. Studier, H. Diamond, J.F. Mech, M.G. Inghram, G.L. Pyle, C.M. Stevens, S. Fried, W.M. Manning, A. Ghiorso, S.G. Thompson, G.H. Higgins and G.T. Seaborg – Phys. Rev. 102 (1956) 180.

- S.M. Fried, G.L. Pyle, C.M. Stevens, J.R. Huizenga – J. Inorg. Nucl. Chem. 2 (1956) 415.
- W.T. Carnall, S. Fried, A.L. Harkness – J. Inorg. Nucl. Chem. 17 (1961) 12.
- L.M. Belov, B.S. Dzheleпов, R.B. Ivanov, A.S. Krivokhatskii, V.G. Nedovesov, V.P. Chechev – Sov. J. Radiochem. 5 (1963) 362.
- B.S. Dzheleпов, R.B. Ivanov, V.G. Nedovesov, V.P. Chechev – Sov. Phys. JETP 18 (1963) 937.
- D. Metta, H. Diamond, R.F. Barnes, J. Milsted, J. Gray, Jr., D.J. Henderson, C.M. Stevens – J. Inorg. Nucl. Chem. 27 (1965) 33.
- S.A. Baranov, Yu.P. Radionov, V.M. Kulakov, V.M. Shatinskii – Sov. J. Nucl. Phys. 4 (1967) 798.
- D.N. Metta, H. Diamond, F.R. Kelly – J. Inorg. Nucl. Chem. 31 (1969) 1245.
- K.W. MacMurdo, R.M. Harbour, R.W. Benjamin – J. Inorg. Nucl. Chem. 33 (1971) 1241.
- J.E. McCracken, J.R. Stokely, R.D. Baybarz, C.E. Bemis, Jr. and R. Eby – J. Inorg. Nucl. Chem. 33 (1971) 3251.
- E. Eichler, N.R. Johnson, C.E. Bemis, Jr., R.O. Sayer, D.C. Hensley, M.R. Schmorak – ORNL-4706 (1971)
- M. Schmorak, C.E. Bemis, Jr, M.J. Zender, N.B. Gove, P.F. Dittner – Nucl. Phys. A178 (1972) 410.
- V.G. Polyukhov, G.A. Timofeev, P.A. Privalova, V.Ya. Garbeskiriya and A.P. Chetverikov – Sov. J. Radiochem. 19 (1977) 414.
- F.P. Larkins – Atomic Data and Nuclear Data Tables. 20 (1977) 313.
- W. Spreng, F. Azgui, H. Emling, E. Grosse, R. Kulesa, Ch. Michel, D. Schwalm, R.S. Simon, H.J. Wollersheim, M. Mutterer, J.P. Theobald, M.S. Moore, N. Trautmann, J.L. Egido and P. Ring – Phys. Rev. Lett. 51 (1983) 1522.
- V.M. Shatinskii – Sov. J. At. Energy 56 (1984) 282.
- N.E. Holden – Pure Appl. Chem. 61 (1989) 1483.
- Rytz – At. Data Nucl. Data Tables 47 (1991) 205.
- M.U. Rajput, T.D. MacMahon – Nucl. Instrum. Methods Phys. Res. A312 (1992) 289.
- S.I. Kafala, T.D. MacMahon, P.W. Gray – Nucl. Instrum. Methods Phys. Res. A339 (1994) 151.
- E. Schönfeld, H. Janßen – Nucl. Instrum. Methods Phys. Res. A369 (1996) 527.
- E. Schönfeld, G. Rodloff – PTB-6.11-98-1 Braunschweig (1998) .
- Artna-Cohen. Nucl. Data Sheets 84 (1998) 901.
- E. Schönfeld, G. Rodloff – PTB-6.11-1999-1 Braunschweig (1999) .
- N.E. Holden, D.C. Hoffman – Pure Appl. Chem. 72 (2000) 1525.
- E. Schönfeld, H. Janßen – Appl. Rad. Isot. 52 (2000) 595.
- Y.A. Akovali – Nucl. Data Sheets 94 (2001) 131.
- M.-M. Bé, R. Helmer, V. Chisté – J. Nucl. Scien. and Techn. suppl. 2 (2002) 481.
- I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, P.O. Tikkanen, S. Raman – At. Data Nucl. Data Tables. 91 (2002) 1.
- Y.A. Akovali – Nucl. Data Sheets 96 (2002) 177.
- R.D. Deslattes, E.G. Kessler, P. Indelicato, L. De Billy, E. Lindroth, J. Anton – Rev. Mod. Phys. 77 (2003) 35.
- G. Audi, A.H. Wapstra – Nucl. Phys. A729 (2003) 337.
- D. MacMahon, A. Pearce and P. Harris – Appl. Rad. Isot. 60 (2004) 275.
- V.P. Chechev – http://www.nucleide.org/DDEP_WG/DDEPdata.htm (2006) .
- F.G. Kondev, I. Ahmad, J.P. Greene, M.A. Kellett and A.L. Nichols – Appl. Radiat. Isot. 65 (2007) 335.

**²⁵²Cf - Comments on evaluation of decay data
by M.M. Bé and V. Chisté**

This evaluation was completed in November 2007. The literature available by October 2007 was included.

1 Decay Scheme

²⁵²Cf disintegrates by α emissions mainly to the ²⁴⁸Cm ground state level, and by spontaneous fission for 3,086 (8) %.

In the Tables part, the data are then normalized to 96,914 (3) alpha decays (see §2.2).

The calculated Q value of 6217(26) keV deduced from the decay scheme data, for the α decay, is in agreement with the value of 6216,87 (4) keV from Audi *et al.* (2003Au03).

2 Nuclear Data

The Q value is from the atomic mass evaluation of Audi *et al.* (2003Au03).

The level energies, spins and parities are based on the evaluation of Y.A. Akevali (1999Ak02).

2.1 Total half-life

A theoretical calculation of the α -decay half-life of Cf-252, by M. Balasubramaniam *et al.* (1999Ba03) leads to a value of 2,592 a.

The measured half-life are, in years:

Reference	half-life	Uc	Comments
Mehta (1965Me02)	2,646	0,004	
De Volpi (1969De23)	2,621	0,006	Rejected by Chauvenet criterion
Mijnheer (1973Mi05)	2,659	0,010	Rejected by Chauvenet criterion
V.Spiegel (1974Sp02)	2,638	0,007	
V.T. Shchebolev (1974Sh15)	2,628	0,010	Superseded by 1992Sh33
Mozhaev (1976Mo30)	2,637	0,005	
Lagoutine (1982La25)	2,639	0,007	
J.R.Smith (1984SmZW)	2,651	0,003	
W.G.Alberts (1983Al**)	2,648	0,002	
E.J. Axton (1985Ax**)	2,6503	0,0031	
Chen Keliang (1988Ke**)	2,64	0,13	
V.T. Shchebolev (1992Sh33)	2,645	0,003	
Weighted mean	2,6470	0,0014	$\chi^2 = 1,3$; χ^2 crit = 2,5

(See also 1994Ka08, 1994KhZW for previous evaluated values.)

In the set of data listed above, two values were rejected in application of the Chauvenet's criterion. A value from 1974Sh15 has been superseded by a more recent one by the same author (1992Sh33). The remaining set of 9 values is consistent with a reduced χ^2 of 1,3. Then the weighted mean is 2,6470 with an external uncertainty of 0,0014. The largest contribution to the statistical weight (35 %) is from Alberts ; Axton,

Shchebolev and Smith give about 15 % each.

However, in the references listed above the uncertainty budget, in most cases, was not given. Some of them include the statistical part of the uncertainty only and did not take into account the systematic components as the associated presence of Cf-250 for example. So, as recommended in the study of Kharitonov (1994KhZW) an uncertainty of 0,1 % has been applied on the final result.

The adopted value is 2,6470 (26) a.

2.2 Spontaneous fission half-life

The spontaneous fission decay constant λ_{sf} is determined by :

$$\lambda_{sf} = \lambda / [(N\alpha/N_{sf}) + 1]$$

where $(N\alpha/N_{sf})$ is the ratio between the number of α -decays and N_{sf} the number of spontaneous fission events and, λ is the total ²⁵²Cf decay constant.

Measured values of the ratio $N\alpha/N_{sf}$:

Reference	Value	Uc
D.Mehta (1965Me02)	31,3	0,2
B.M.Aleksandrov (1970Al23)	31,39	0,26
J.D.Hastings (1971Ha**)	31,5	0,2
A.K. Pandey (1993Pa29)	31,56	0,35
Y.S.Popov (1990Po24)	31,38	0,12
Weighted mean	31,40	0,08

The 5 data sets given above are consistent (reduced $\chi^2 = 0,2$).

From this value and the total half -life above (§ 2.1), a **spontaneous fission half-life of 85,76 (23) a** is deduced.

From $N\alpha/N_{sf} = 31,40$ (8) and $N\alpha + N_{sf} = 100$ Cf-252 decays, the **percentage of spontaneous fissions in the decay of Cf-252 is 3,086 (8) %**.

Then the percentage of alpha transitions is: 96,914 (8) %.

2.3 Average number of neutrons

The average number of neutrons $\bar{\nu}$ emitted by spontaneous fission is:

$$\bar{\nu} = 3,7675 (40)$$

as evaluated in the study of M. Divadeenam *et al.* (1984Di08) where relevant experimental data are taken into account and a least-squares fitting program was used to obtain an overall fit.

The average number of neutrons emitted per 100 disintegrations is:

$$n = 3,086 (8) \times 3,7675 (40) = 11,627 (33) \%$$

2.4 a Transitions

See Alpha-particle emissions (§ 4)

2.5 g Transitions

Multipolarities of these γ -ray transitions are from 1999Ak02.

The internal conversion coefficients for the 43 and 100-keV gamma transitions were calculated with the BrIcc code for the Frozen Orbital approximation (2005KiZW).

3 Atomic Data

Atomic values, ω_K , ω_L and n_K , are from Schönfeld and Janßen (1996Sc33).

4 α -Particle Emissions

4.1 α -Particle Energies

From the measured values of Rytz (1986Ry04) and Baranov (1976BaZZ, 1971Ba10, 1970 Ba18), Rytz (1991Ry01) made some adjustments taking into account variations in the energies used as calibration standards. This leads, for the two main groups, to the recommended values of :
6118,10 (10) keV and 6075,64 (11) keV

The other energies : 5976,6 ; 5826,3 and 5615,6-keV are from Baranov (1970Ba18 and 1971Ba10)

Recorded spectra are also shown in Glover (1984Gl03) and Wiltshire (1985Wi14).

4.2 α -Particle Intensities

Measured alpha intensities, per 100 alpha decays :

Energy (keV)	Reference	Intensity (%)	Uc	Comments
6118,10	Asaro (1955As42)	84,5		
	Baranov (1976BaZZ)	84,1	0,4	See also 1970Ba18
	Adopted	84,3	0,3	Unweighted mean
6075,64	Asaro (1955As42)	15,5		
	Baranov (1976BaZZ)	15,8	0,1	See also 1970Ba18
	Adopted	15,6	0,3	Unweighted mean
5976,6	Baranov (1970Ba18)	0,2		See also 1985Wi14
	Asaro (1958As64)	0,28		
	Adopted	0,24	0,04	Unweighted mean
5826,3	Baranov (1970Ba18)	$2 \cdot 10^{-3}$		
5616	Baranov (1970Ba18)	$\sim 6 \cdot 10^{-5}$		

The number of measurements is very scarce moreover the results given by Asaro are without uncertainties. To try to make the most of this limited data, the unweighted mean is adopted, for the 6118 -, 6075-, 5976-keV groups, with uncertainty covering the two existing values.

The intensity of the 5826-keV group is from Baranov (1970Ba18).

The weak group with energy 5615-keV, possibly feeding a 505-keV level, is not adopted, because no photons depopulating this level have been observed in the Cf-252 decay.

In the Tables part, these data are normalized to 96,914 (8) alpha decays (see §2.2).

5 Photon Emissions

5.1 g-Ray Emissions

Measured gamma-ray intensities, per 100 alpha decays :

Energy (keV)	Reference	Intensity (%)	Uc	Comments
42	Asaro (1955As42)	0,014		
43,399 (25)	Watson (1971Wa28)	0,0153	0,0009	
	Adopted	0,0157	0,0004	From decay scheme
100,2 (4)	Asaro (1955As42)	0,013		Adopted E _γ (1999Ak02)
	Adopted	0,0123	0,0021	From decay scheme
154,5 (2)	Piercey (1993Pi07)			Adopted E _γ (1993Pi07)
	Adopted	0,00053	0,00001	From decay scheme

The gamma ray intensities were deduced from the gamma -ray transition probabilities (see §2.5) and the theoretical ICC values.

In the Tables, these data are normalized to 100 decays of Cf-252 (see §2.2).

5.2 X-ray emissions

Asaro (1955As42) measured a K X-ray intensity of 0,007 %. This value disagrees with an expected KX-ray intensity of 0,000 086 % from the internal conversion electrons of the 154,5-keV gamma ray.

Relative intensities were measured by Popov *et al.* (1990Po14).

Total L X-ray intensity following the Cf-252 decay to Cm -248 was measured by Watson (1971Wa28) as 7,83 (40) % per 100 alpha decays.

The L X-ray total intensity calculated from the decay scheme data is 6,26 (14) % per 100 alpha decays. This result is in reasonable agreement with the measured value of Watson.

6 References

- 1955As42 - F.Asaro, et al. Phys. Rev. 100, 1 (1955) 137.
 1963Bj03 - S.Bjornholm, et al. Phys. Rev. 130, 5 (1963) 2000.
 1965Me02 - D.Mehta, et al. J.Inorg.Nucl.Chem. 27 (1965) 33.
 1969De23 - A.De Volpi, et al. Inorg. Nucl. Chem. Letters;5 (1969) 699.
 1970Ba18 - S.A.Baranov, et al. Sov. J. Nucl. Phys. 11,3 (1970) 393.
 1970Al23 - B.M.Aleksandrov, et al. Sov. Atomic Energy 28 (1970) 462.
 1971Wa28 - R.L.Watson, T.K.Li. Nucl. Phys. A178 (1971) 201.
 1971Ha** - J.D.Hastings, et al. Mound Laboratory report- MLM-1845 ; TID-4500 ; UC-4; (1971) 1.
 1971Ba10- S.A.Baranov, et al. Sov. J. Nucl. Phys;14,5 (1972) 614.
 1973Mi05 - B.J.Mijnheer. Int. J. Appl. Radiat. Isotop. 24 (1973) 185.
 1974Sp02 - V.Spiegel. Nucl. Sci. Eng. 53 (1974) 327.
 1974Sh15 - V.T.Shchebolev, et al. Sov. Atomic Energy 36 (1974) 507.
 1976BaZZ - S.A.Baranov, et al. Proc. Advisory Group meeting Transactinium Nucl.Data, Karlsruhe, IAEA 186; B6 (1976) 249.
 1976Mo30 - V.K.Mozhaev. Sov.Atomic Energy 40 (1976) 200.
 1982La25 - F.Lagoutine, et al. Int. J. Appl. Radiat. Isotop. 33 (1982) 711.

- 1983Al** - W.G.Alberts, et al. Report PTB-Mitteilungen 93 (1983) 315.
 1984G103 – K.M. Glover. Int. J. Appl. Radiat. Isot. 35, 4 (1984) 239.
 1984Di08 - M.Divadeenam, et al. Ann. Nucl. Energy 11, 8 (1984) 375.
 1985Wi14 - R.A.P.Wiltshire. Nucl. Instrum. Methods Phys. Res. A236 (1985) 514.
 1985Ax** - E.J.Axton, et al. Metrologia 21 (1985) 59.
 1986Ry04 - A.Rytz, et al. Nucl. Instrum. Methods Phys. Res. A253 (1986) 47.
 1984SmZV - J.R.Smith. Report EPRI NP-3436 (1984).
 1987Sh** - E.A.Shlyamin, et al. IAEA TECDOC-410 (1987) 225.
 1988Ke** - Chen Keliang, et al. China Nucl. Information Centre Beijing, CNIC-I-004; (1988) 23.
 1990Po24 - Yu.S.Popov, et al. Sov. J. Radiochemistry 32 (1990) 425.
 1991Ry01 - A.Rytz. At. Data. Nucl. Data Tables 47,2 (1991) 229.
 1992Sh33 - V.T.Shchebolev, et al. Sov. Atomic Energy 73, 6 (1992) 1015.
 1993Pa29 - A.K.Pandey, et al. Nucl. Instrum. Methods Phys. Res. B82 (1993) 151.
 1994Ka08 - S.I.Kafala, T.D.MacMahon, P.W.Gray, Nucl.Instrum.Methods Phys.Res. A339 (1994) 151
- Testing of Data Evaluation Methods*
- 1994KhZW – I.A.Kharitonov. Report IAEA INDC(CCP)-362 (1994)
 1996Sc33 - E.Schönfeld, et al. Nucl. Instrum. Methods Phys. Res. A369 (1996) 527.
 1999Ba03 - M.Balasubramaniam, et al. Phys. Rev. C60 (1999) 064316.
 1999Ak02 - Y.A.Akovali. Nucl. Data Sheets 87 (1999) 257.
 1999Po - Yu.S.Popov, et al. Radiochemistry 41, 1 (1999) 43.
 2002Ba85 – I.M. Band, M.B. Trzhaskovskaya, C.W. Nestor, Jr., P.O. Tikkanen, S. Raman, Atomic Data and Nuclear Data Tables 81(2002)1 [ICC].
 2003Au03 - G. Audi, A.H. Wapstra, C. Thibault. Nucl. Phys. A729 (2003) 337.
 2005KiZW - T. Kibedi, T.W. Burrows, M.B. Trzhaskovskaya, C.W. Nestor Proc. Intern. Conf. Nuclear Data for Science and Technology, Santa Fe, New Mexico, 26 September-1 October; 769 (2005) 268.

